

Carbon 13 isotopes reveal limited ocean circulation changes between interglacials of the last 800 ka

N. Bouttes^{1,2}, N. Vazquez Riveiros³, A. Govin¹, D. Swingedouw², M. F. Sanchez-Goni^{2,4},
X. Crosta², D. M. Roche^{1,5}

¹ Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91190 Gif-sur-Yvette, France.

² Université de Bordeaux, EPOC, UMR 5805, F-33615 Pessac, France.

³ IFREMER, *Unité de Géosciences Marines*, Pointe du Diable, 29200, Plouzané, France.

⁴ Ecole Pratique des Hautes Etudes (EPHE, PSL University), F-33615 Pessac, France

⁵ Earth and Climate Cluster, Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands.

Contents of this file

Figures S1 to S2

Introduction

We present two additional figures with $\delta^{13}\text{C}$ distribution (permil) in the Atlantic, Pacific and Indian Oceans for the interglacials considered in this study.




Figure S1. Data $\delta^{13}\text{C}$ distribution (permil) in the Atlantic Ocean for the nine considered interglacials.




Figure S2. Data $\delta^{13}\text{C}$ distribution (permil) in the Pacific (squares) and Indian (circles) Oceans for the nine considered interglacials.