



# Taxonomic revision of the genus *Xenogryllus* Bolívar, 1890 (Orthoptera, Gryllidae, Eneopterinae, *Xenogryllini*)

Ranjana Jaiswara, Jiajia Dong, Libin Ma, Haisheng Yin, Tony Robillard

## ► To cite this version:

Ranjana Jaiswara, Jiajia Dong, Libin Ma, Haisheng Yin, Tony Robillard. Taxonomic revision of the genus *Xenogryllus* Bolívar, 1890 (Orthoptera, Gryllidae, Eneopterinae, *Xenogryllini*). Zootaxa, 2019, 4545 (3), pp.301. 10.11646/zootaxa.4545.3.1 . hal-02866078

HAL Id: hal-02866078

<https://hal.science/hal-02866078>

Submitted on 12 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1   **Taxonomic revision of the genus *Xenogryllus* (Orthoptera, Gryllidae,  
2   Eneopterinae)**

3

4   **Page headlines:** Revision of *Xenogryllus*

5

6   *Ranjana Jaiswara\** (1,2)

7   *Jiajia Dong\** (1,3)

8   *Libin Ma* (3)

9   *Haisheng Yin* (4)

10   *Tony Robillard* (1)

11

12   *1\_Institut de Systématique, Evolution et Biodiversité (ISYEB), Muséum national  
13   d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, 57 rue Cuvier, CP 50, 75231  
14   Paris Cedex 05, France*

15   *2\_Department of Zoology, Panjab University, Chandigarh, 160014, India*

16   *3\_College of Life Science, Shaanxi Normal University, 710119, Xi'an, Shaanxi, P.R.  
17   China*

18   *4\_Shanghai Entomological Museum, Institute of Plant Physiology and Ecology,  
19   Chinese Academy of Sciences, Shanghai 200032, China*

20   *\* Equal contribution*

21

22   *Corresponding author: Tony Robillard, tony.robillard@mnhn.fr*

23    **Abstract**

24    Subfamily Eneopterinae has been studied for its diversified acoustic modalities and  
25    disjunct distribution. The genus *Xenogryllus* Bolívar, 1890 is one of the oldest genera of  
26    the subfamily, and the first genus of the tribe Xenogryllini. *Xenogryllus* is known for its  
27    low-frequency calling songs and wide distribution across Africa and Asia. It is known  
28    from five species and has never been subject to formal taxonomic revision and  
29    description of acoustic features of their calling songs is lacking. Therefore, this study  
30    consists of a detailed taxonomic revision of all the species. We re-describe or append  
31    their taxonomic features using external morphological features and internal genitalic  
32    structures. We update the list of *Xenogryllus* by describing three new species from  
33    Africa. In addition, we provide keys to identify species and describe the calling songs of  
34    five species.

35

36      **Introduction**

37      The crickets of the subfamily Eneopterinae have been recently studied for the  
38      diversity of their communication signals and related structures (e.g., Robillard &  
39      Desutter-Grandcolas 2004a, b; Robillard *et al.* 2013; ter Hofstede *et al.*, 2015;  
40      Schneider *et al.*, 2017), and for their biogeographical patterns (Nattier *et al.* 2011;  
41      Vicente *et al.* 2017; Dong *et al.* 2018). To investigate how this diversity occurred and  
42      how the original traits evolved, it is necessary to understand the context in which it  
43      happened. The tribe Xenogryllini is the sister group of the most diversified Lebinthini  
44      (Vicente *et al.* 2017), therefore, it serves as the best “control lineage” to understand the  
45      evolution of unique features in the Lebinthini. A taxonomic, bioacoustics and  
46      phylogenetic focus on the tribe Xenogryllini is thus particularly needed.

47      The Xenogryllini are currently composed of two genera, *Xenogryllus* Bolívar,  
48      1890 (six species) and *Pseudolebinthus* Robillard, 2006 (two species) according to  
49      Orthoptera Species File Online data base (Cigliano *et al.* 2018), but a taxonomic  
50      revision was presented as necessary by Robillard (2006). The members of the  
51      Xenogryllini produce calling songs with low dominant frequencies (Robillard &  
52      Desutter-Grandcolas 2011). However, their spectral and temporal features were never  
53      formally documented.

54      In the present study, we conduct a complete taxonomic revision of the genus  
55      *Xenogryllus* (Table 1) based on freshly collected specimens or procured from 17  
56      natural history repositories. We evaluate the status, redescribe or amend description of  
57      the existing species, describe three new species, *Xenogryllus lamottei* Robillard n. sp.,  
58      *X. maniema* Robillard & Jaiswara n. sp. and *X. mozambicus* Robillard n. sp., and  
59      provide and a key to species.

60

61      **Material and methods**

62      **Material**

63            The material examined belongs to the collections of 17 institutions (see Appendix  
64       S1 for lists of material and abbreviations). Newly collected material comes from several  
65       field work expeditions in Mozambique (2009), India (2015, 2017) and China (2012,  
66       2014, 2017). Newly collected specimens are deposited in the collections of Zoological  
67       Survey of India, Kolkata, Muséum national d'Histoire naturelle, Paris and Shaanxi  
68       Normal University, Xi'an. Political districts are mentioned for each specimen within lists  
69       of materials, and square brackets are used for additional information not mentioned on  
70       specimen labels or for translation of key information present on labels in languages  
71       other than English.

72

73      **Taxonomic description**

74            Description of studied species follows terminologies as proposed by Robillard *et*  
75       *al.* (2014). Observations of external morphological characters and dissection of male  
76       and female genitalia were performed using Leica stereomicroscopes MZ16 (at MNHN,  
77       Paris) and M205C (at IISER, Mohali). SEM observations were performed at the  
78       Plateforme de Microscopie électronique of the MNHN, using a JEOL-JSM 840  
79       electronic microscope (7kV), after a 60 s gold-coating. Terminologies for male FW  
80       venation follow Ragge (1995) and Robillard & Desutter-Grandcolas (2004b). Male and  
81       female genitalia were dissected either from dry preserved or freshly killed specimens.  
82       Male genitalia were dissected by making a small slit between paraproct and subgenital  
83       plate. Female copulatory papilla was dissected out by cutting the membrane between  
84       ovipositor and subgenital plate. Dissected genitalia were cleared in 10% cold KOH  
85       solution and preserved in glass vials containing glycerine. Terminologies for genitalia  
86       follow Desutter (1987), modified in Desutter-Grandcolas (2003) and Robillard &

87 Desutter-Grandcolas (2004a). Imaging of male and female genitalia were made using  
88 an AmScope MU1000 digital camera () or with a Canon EOS 40D Digital SLR camera  
89 (at MNHN), or with a MC120 HD, Leica, Germany (at IISER, Mohali). To highlight the  
90 structural components of male and female genitalia, water solution containing a drop of  
91 JBL Punktol was used. To fix orientations and stabilization of genitalia for photography,  
92 a clear and viscous Power Plast Hand Sanitizer was used following Su (2016).

93

94 **Abbreviations used in taxonomic descriptions**

95 General morphology: FI, FII, FIII, fore, median, hind femur; FW, forewing; TI, TII,  
96 TIII, fore, median, hind tibia; Tarsomere I/II/III-1: basal segment of fore, median and  
97 hind leg tarsomere.

98 Tegminal venation: 1A-4A, first to fourth anal veins; CuA, anterior cubitus; CuA1,  
99 CuA2, ... first, second, ... branches of CuA; CuP, posterior cubitus; M, median vein; R,  
100 radial vein; Sc, subcostal vein and its branches; c1-3, first to third cells of C alignment;  
101 d1 cell (mirror), first cell(s) of D alignment; d2, second cell of D alignment; e1, first cell  
102 of E alignment.

103 Species abbreviations used in figures: ene, *Xenogryllus enoopteroides*; lam,  
104 *Xenogryllus lamottei* n. sp.; mai, *Xenogryllus maichauensis*; man, *Xenogryllus*  
105 *maniema* n. sp.; mar, *Xenogryllus marmoratus*; moz, *Xenogryllus mozambicus* n. sp.;  
106 tra, *Xenogryllus transversus*; ulu, *Xenogryllus ululiu*.

107

108 **Abbreviations of Institutions used in the lists of material**

109

110 AMNH: American Museum of Natural History, New York, USA.

111 ECNU: East China Normal University, Shanghai, China.

112 DEI: Deutsches Entomologisches Institut, Müncheberg, Germany.

113 NHMUK: Natural History Museum (formerly British Museum of Natural History),  
114 London, United Kingdom.  
115 IISERM: Indian Institute of Science Education and Research Mohali, Punjab, India.  
116 MNCN: Museo Nacional de Ciencias Naturales, Madrid, Spain.  
117 MNHN: Muséum national d'Histoire naturelle, Paris, France.  
118 MRAC: Musée Royal de l'Afrique Centrale, Tervuren, Belgium.  
119 NHMW: Naturhistorisches Museum, Vienna, Austria.  
120 NWAFU: Entomological Museum, Northwest A & F University, Yangling, China.  
121 RMNH: National Natuurhistorisch Museum (formerly Rijksmuseum van Natuurlijke  
122 Historie), Leiden, The Netherlands.  
123 SNNU: College of Life Sciences, Shaanxi Normal University, Xi'an, China.  
124 SIPPE: Institute of Plant Physiology and Ecology, Shanghai Institutes for Biological  
125 Sciences, Chinese Academy of Sciences, China.  
126 RBINS: Royal Belgian Institute of Natural Sciences, Brussels, Belgium.  
127 ZFMK: Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn,  
128 Germany.  
129 ZIN: Zoological Institute, Russian Academy of Sciences, S. Petersburg, Russia.  
130 ZSI: Zoological Survey of India, Kolkata, India.  
131

132 **Measurements**

133 BL, body length in dorsal view, from fastigium to apex of abdomen; FIIIL, length  
134 of FIII; FIIIW, width of FIII; TIIIL, length of TIII; FWL, forewing length; FWW, forewing  
135 width (at the level of maximal width at about 1/3 of FWL); HWT, hind wing tail length  
136 (part of the hind wings longer than the FWs); las, inner spines on TIII dorsal side above  
137 the spurs; lbs, inner spines on TIII dorsal side between the spurs; Oas, outer spines on  
138 TIII dorsal side above the spurs; Obs, outer spines on TIII dorsal side between the  
139 spurs; OL, ovipositor length; PronL, pronotum length; PronW, pronotum width; TallIs,

140 spines of third hind tarsomere, not including the apical spines: Ids, inner dorsal spines;  
141 Ods, outer dorsal spines; Ols, outer spines on lateral side of TaliI.

142

### 143 **Acoustic analyses**

144 We document the temporal and spectral measurements of calling songs of  
145 *Xenogryllus enoopteroides*, *X. maichauensis*, *X. marmoratus*, *X. mozambicus*, *X.*  
146 *transversus* and *X. ululiu*. The calling songs of *X. lamottei* n. sp. and *X. maniema* n. sp.  
147 are unknown.

148 The basic cricket song terminology follows Ragge & Reynolds (1998). A basic  
149 song unit is called syllable and corresponds to one opening-closure cycle of the male  
150 forewings. The calling song of *X. transversus* was recorded using TASCAM DR07 MKII  
151 at 44.1 k-samples/s sampling rate (RJ). Males of the species *X. mozambicus* n. sp. and  
152 *X. marmoratus* were recorded with a modified Condenser Microphone Capsule CM16  
153 (Avisoft Bioacoustics, Berlin), with a flat frequency response from 3 to 150 kHz (R.  
154 Specht pers. comm.), connected to a Tascam HD-P2 digital recorder (96 kHz sampling  
155 frequency, 16 bit) in the field, or using Avisoft Triggering Harddisk Recorder version  
156 2.97 and an 8-Pre MOTU sound card at a sampling frequency of 96 k-samples/s (16  
157 bit) (TR). The calling songs of *X. maichauensis* and *X. ululiu* were obtained from the  
158 Orthoptera Species File Online (Cigliano *et al.* 2018) database and were recorded by  
159 Sigfrid Ingrish with a Sony WM-D3 recorder with a Universum mono microphone.

160 Acoustic analyses were performed using the computer softwares Avisoft-SASLab  
161 Pro version 4.40 (Specht 2009) and Audacity version 2.0.0. Acoustic features were  
162 measured either manually on Audacity or using the automatic commands of Avisoft-  
163 SASLab Pro. Newly recording files are deposited in the Sound Library of the MNHN  
164 (inventory numbers mentioned as MNHN-SO\*\*\* in lists of material), and in the  
165 Orthoptera Sound Library of Indian Institute of Science Education and Research Mohali,  
166 Punjab, India.

167

168   **Systematic part**

169   **Orthoptera**

170   **Gryllidae**

171   **Eneopterinae Saussure, 1874**

172   **Xenogryllini Robillard, 2004**

173   *Emended diagnosis.* Crickets of contrasted sizes, generally light brown; head higher  
174   than wide in facial view; fastigium rather wide; eyes of variable sizes, usually  
175   ornamented with thin dorso-ventral dark stripes. TI with two tympana, inner one  
176   covered by a sclerotized expansion, its membrane visible along a longitudinal slit only;  
177   outer tympanum oval in shape, its surface smooth. TIII spurs characteristic of the  
178   subfamily, with four pairs of subapical spurs and three apical spurs on each side, the  
179   median the longest; subapical spurs rather straight, with curved apex as in Nisitrini.  
180   Wings variable within the tribe in males and females. *Male.* Metanotum with  
181   characteristic glandular structures, with a dense bunch of very long setae on scutum  
182   anterior edge. Dorsal margin of subgenital plate with baso-lateral glandular structure or  
183   dorsal edge with a median invagination. *Male FW venation:* 2A and 3A veins narrowly  
184   coupled at plectrum level; harp veins variable; mirror generally well developed and  
185   rounded; bases of CuA2 and CuA3 fused. *Male genitalia:* Pseudepiphalllic lophi well-  
186   developed, long and sclerotized, usually with an apical hook-like expansion;  
187   pseudepiphallus with lateral membranous lobes more or less developed; rami fused  
188   with pseudepiphalllic sclerite, rather short with convergent apex; ectophalllic arc faintly  
189   or not sclerotized; endophalllic sclerite with long latero-posterior arms connected to  
190   ectophalllic fold, and short anterior region; endophalllic apodeme with wide lateral  
191   lamellas. Female ovipositor apex not denticulate, variably pointed.

192     *Distribution.* Asia (Japan, China, India, Thailand, Myanmar, Vietnam, Pakistan,  
193     Bangladesh) and Sub-Saharan Africa.

194

195     ***Xenogryllus* Bolívar, 1890**

196     Type species: *Xenogryllus eneopteroides* Bolívar, 1890.

197     *Xenogryllus* Bolívar, 1890: 232; Kirby 1906: 106; Chopard 1968: 349; 1969: 307;  
198     Vasanth 1993: 130; Robillard & Desutter-Grandcolas 2008: 67; Cigliano *et al.* 2018  
199     (Orthoptera Species File Online).

200     Synonym names:

201     *Dindymus* Kirby, 1906: 88 - incorrect spelling of *Dionymus*.

202     *Dionymus* Brunner von Wattenwyl, 1893: 213 – Chopard 1968: 349.

203

204     *Emended diagnosis.* Genus of average to large size, differing from *Pseudolebinthus* by  
205     FWs as long or longer than abdomen in both sexes (shorter in *Pseudolebinthus*), hind  
206     wings longer than FWs, forming a short tail posterior to FWs. Eyes small and little  
207     prominent, located on face. Face with a whitish or yellowish mask with dark spots. FIII  
208     long and thin. *Male.* Dorsal disc of pronotum forming a wide trapezoid. FWs almost  
209     completely overlapping, widened basally, usually forming a characteristic box around  
210     abdomen, twice as wide as abdomen. FW cells with thin longitudinal wrinkles, including  
211     harp and mirror, absent in *Pseudolebinthus*. Left and right FWs similar in sclerotization  
212     and coloration (asymmetrical in *Pseudolebinthus*). *Male genitalia:* Pseudepiphalllic  
213     sclerite longer than rami; lophi long and sclerotized (shorter in *Pseudolebinthus*). Lophi  
214     with apical hook-like inner expansions. Pseudepiphalllic sclerite with lateral  
215     membranous lobes. Rami strong and short, their apex convergent and hook-like.  
216     *Female.* Dorsal disc of pronotum almost rectangular. FWs as long as in male, not  
217     widened, reaching apex of abdomen. Ovipositor flattened longitudinally as in other

218 eneopterine genera, its apex rounded and smooth. *Female copulatory papilla*: long,  
219 thin and conical.

220 *Redescription*. Genus of average to large size (Figs 1-2), generally light brown to  
221 ochre, with a few dark patterns on FWs. Vertex with five wide longitudinal dark bands  
222 (Fig. 4), including a wide median one prolonged on fastigium, two lateral bands more or  
223 less marked, made of aggregates of brown dots, and two thin stripes posterior to eyes.  
224 Eyes rather small, little prominent, ornamented with thin dorso-ventral dark stripes.  
225 Fastigium forming a wide rectangular rostrum slightly prominent on face. Scapes yellow  
226 with brown patterns, antennae light brown. Face with a whitish mask with dark spots  
227 (Fig. 3). Lateral part of head and lateral lobes of pronotum almost homogeneously  
228 brown. Legs usually homogeneously brown. TI with two tympana; inner tympanum  
229 covered by a sclerotized expansion, its membrane visible along a small longitudinal slit  
230 only; outer tympanum oval in shape, its surface smooth. TI with three apical spurs;  
231 outer dorsal apical spur absent. TII with four apical spurs (two dorsal and two ventral).  
232 FIII long and thin; TIII with four outer and four inner sub-apical spurs.

233 *Male*. Dorsal disc of pronotum forming a wide trapezoid FWs almost completely  
234 overlapping, widened basally, forming a wide box around abdomen (Fig. 5), with  
235 glandular structures on metanotum (Fig. 6). FW cells with thin longitudinal wrinkles.  
236 Left and right FWs similar in sclerotization and coloration. FWs light brown, translucent,  
237 with the following pattern of black spots on dorsal field: wide black transverse band  
238 anterior to transverse part of 1A, including file angle in some species; four corners or  
239 mirror and median area of chords with faint dark markings (Fig. 5). Harp with two  
240 complete parallel oblique veins and one incomplete oblique vein between diagonal vein  
241 and first complete oblique; harp distal angle concave. Mirror large and rounded,  
242 separated in two sub-equal parts by a sinuous accessory vein. Cell d2 crescent-like,  
243 thin and underlying mirror. Cell e1 very long and curved, underlying posterior margin of  
244 mirror. Apical field well developed, triangular, including 4-5 cell alignments. Lateral field

245 crossed by numerous parallel projections of Sc. *Male genitalia* (Fig.7): Pseudepiphalllic  
246 sclerite longer than rami; pseudepiphalllic lophi very long and sclerotized, with apical  
247 hook-like inner expansions (Fig. 8). Pseudepiphalllic sclerite with lateral membranous  
248 lobes. Membrane at bases of pseudepiphallus forming a sclerotized plate; basal margin  
249 with a ventral reinforcement. Rami usually strong with convergent hook-like apex; most  
250 often with a ventral posterior expansion reaching base of pseudepiphalllic membranous  
251 lateral lobes. Pseudepiphalllic parameres with one strong rectangular ventral lobe  
252 covered with scale-like sculptures, with a basal lobe mostly membranous. Ectophallic  
253 arc nearly membranous. Ectophallic apodemes usually strong. Ectophallic ventral  
254 expansions well developed and sclerotized. Ectophallic fold entirely sclerotized  
255 ventrally, its lateral expansions partly fused with lateral arms of endophallic sclerite.  
256 Endophallic sclerite forming a flat plate with long latero-posterior arms and a short  
257 medio-posterior triangular expansion. Endophallic apodeme with two wide lateral  
258 lamellas and no dorsal crest.

259 *Female*. Pronotum dorsal disc almost rectangular. FWs as long as in male, reaching  
260 apex of abdomen or slightly longer, not widened as in male; light brown with a dark  
261 spot of variable size between veins CuP and CuA, near lateral angle, at  $\frac{1}{4}$  of FW  
262 length; with strong longitudinal veins and faint transverse ones. Lateral field crossed by  
263 numerous parallel projections of Sc; area between R and Sc most often with a thin dark  
264 brown band. Subgenital plate variably indented apically (Fig. 9). Ovipositor flattened  
265 longitudinally, its apex rounded and smooth (Fig. 10). *Female copulatory papilla*: long,  
266 thin and conical, its sclerotization variable (Fig. 11).

267 *Distribution*. Japan, Southern China, Taiwan, India, Sri Lanka, Thailand, Myanmar,  
268 Vietnam, Bangladesh and Sub-Saharan Africa.

269 *Natural history*. *Xenogryllus* species live in open areas, in grassland and savannah.  
270 Males sing at night from low bushy vegetation.

271

272 Key to species of *Xenogryllus*

- 273 1. Dorsal disc of pronotum carinated laterally. Eyes higher than long in lateral view,  
274 occupying half of head height  
275 ..... 2
- 276 - Dorsal disc of pronotum not carinated laterally. Eyes as long as high in lateral view,  
277 restricted to dorsal third of head  
278 ..... 4
- 279
- 280 2. Face almost flat in lateral view ..... *X. eneopteroides* Bolívar,  
281 1890
- 282 - Face rounded in lateral view  
283 ..... 3
- 284
- 285 3. Face bulbous in lateral view. Size rather larger. Male genitalia: Pseudepiphalllic lophi  
286 (Fig. 8F) ended by a small lamella, their inner margin almost straight,  
287 membranous..... *X. mozambicus* Robillard  
288 n. sp.
- 289 - Face less rounded in lateral view. Size rather shorter. Male genitalia: Pseudepiphalllic  
290 lophi (Fig. 8E) ended by a long conical apex and with a widened inner membranous  
291 margin ..... *X. maniema* Robillard &  
292 Jaiswara n. sp.
- 293
- 294 4. Male FWs well widened basally, forming a wide box around  
295 abdomen..... 5

- 296 - Male FWs not widened, as wide as abdomen..... *X. lamottei* Robillard
- 297 n. sp.
- 298
- 299 5. Size larger (body length > 20 mm). Male genitalia: pseudepiphalllic lophi strong,  
300 ended by a sharp apex with a dorsal preapical pointed expansion. Female subgenital  
301 plate posterior margin clearly  
302 indented..... 6
- 303 - Size small (body length <20 mm). Male genitalia: pseudepiphalllic lophi very long and  
304 thin, ended by a lamella, without dorsal preapical pointed expansion. Female  
305 subgenital plate posterior margin not indented (Fig. 9B)..... *X.  
306 marmoratus* (Haan, 1844)
- 307
- 308 6. Male genitalia: Pseudepiphalllic lophi with pseudepiphalllic lophi forming curved  
309 elongate hooks widened apically (Fig. 8D). Female subgenital plate posterior margin  
310 indentation rounded laterally, as wide as long (Fig. 9C)..... *X. maichauensis*  
311 Gorochov, 1992
- 312 - Male genitalia: Pseudepiphalllic lophi straight apically. Female subgenital plate  
313 posterior margin indentation pointed laterally  
314 ..... 7
- 315
- 316 7. Size larger. Male genitalia with larger pseudepiphalllic sclerite; lophi proportionally  
317 smaller (Fig. 8I), fused only basally, separated by a deep indentation. Female  
318 subgenital plate posterior margin indentation very deep, longer than wide (Fig.  
319 9G)..... *X. transversus* (Walker,  
320 1869)

321 - Size smaller. Male genitalia with smaller pseudepiphallic sclerite; lophi fused on half  
322 of their length (Fig. 8H). Female subgenital plate posterior margin indentation weak,  
323 wider than long (Fig. 9E).....X.  
324 *ululiu* Gorochov, 1990

325

326 ***Xenogryllus eneopteroides* Bolívar, 1890**

327 (Figs 1A-C; 3A-B; 4A; 5A; 6; 7A-C; 8A-B; 9A; 10A; 11A-B)

328

329 *Xenogryllus eneopteroides* Bolívar, 1890: 232 – Karsch 1893: 204; Kirby 1906: 106;  
330 Giglio-Tos 1907: 26; Bolívar 1910: 544; Chopard 1954: 80; Chopard 1968: 349;  
331 Chiffaud & Gillon 1984 (life history); Paris 1994[1993]: 69 (lectotype designated);  
332 Robillard & Desutter-Grandcolas 2004a: 578; 2004b: 273 (morphological phylogeny);  
333 2006: 644; 2008: 67; 2011: 637; Robillard 2006: 673; Robillard *et al.* 2007: 1267 (song  
334 frequency); Nattier *et al.* 2011: 2201 (molecular phylogeny); Chintauan-Marquier *et al.*  
335 2016: 62 (molecular phylogeny); Vicente *et al.* 2017: 2203 (historical biogeography);  
336 Cigliano *et al.* 2018 (Orthoptera Species File Online).

337 *Xenogryllus eucopterooides* - Kevan & Knipper 1961: 368 (misspelling); Chopard 1968:  
338 349.

339

340 *Type material. Lectotype*, ♂, **Angola**: Duque de Bragança, Bayão (Paris, 1994)  
341 [MNCN, examined on photograph].

342 *Additional material examined. Benin*: Bas-Dahomey, plateaux de Zagnamado et de  
343 Ketou, forêt d'Aqua-queré (saison sèche), 1910 P. Ducorps, 1♀ (MNHN). **Cameroun**:  
344 [L.] Conradt, 1♂, identified *Xenogryllus eneopteroides* Bol. (MNHN). Reg[ion] de Kribi,  
345 1925, littoral, saison humide, Dr. Gromier, 1♂ (MNHN). Kamerun [Cameroun], Pipindi  
346 [Bipindi], Eitel ded. 1♀, identified *Xenogryllus eneopteroides* Bol. (MNHN). 10 km N. of

347 Yaoundé, E. of Agronomy Station, Nkolbisson, 10.vii.1975, N. D. Jago, 1♂, identified  
348 *Xenogryllus eneopteroides* Bolivar by B. C. Townsend, 1976 (NHMUK 010926525).  
349 **Central African Republic:** République Centrafricaine, 75 km WNW Mbaiki, N'Gotto  
350 [Ngoto], W Lobaye, savane arbustive, 7.iv.1995, nuit [night], L. Desutter-Grandcolas, 1  
351 juvenile, n°14, molecular sample Xen-CA (MNHN-E0-ENSIF3159); 1♂, #1, sur plant  
352 [on plant] (MNHN). **Democratic Republic of the Congo:** Congo, 20 km from L. Tumba  
353 (towards Coquilhatville) [Mbandaka], 5.iii.1964, from grassy clearing in forest, 1♂, 1♀,  
354 Tyson Roberts, Brit. Mus. 1982-71 (NHMUK 010926531, 010926572). Kalembe,  
355 15.ix.1947, Miss. Tanganika, 1♀ (RBINS). Lukafa [Lukafu], Congo Belge, 6.xii.1938,  
356 H.J. Brédo, 1♀, R.Mus.Nat. Belg.I.G.12.204 (RBINS). Belgian Congo, Aru, iii.1936, 5♂,  
357 2♀, H.J. Bredo, dry Acacia bush (NHMUK 010926536, 010926527, 010926578,  
358 010926580, 010926539, 010926566, 010926569). Niangara, v-vi-1915, 28°0E 3°40"N,  
359 1♂ (AMNH). Haut-Ituri: Faradje, 1♂, Blommaert, Musée du Congo, identified  
360 *Xenogryllus eneopteroides* by L. Chopard, R. Det. N 2814 (MRAC). S.E. Katanga:  
361 Ngaye: xi-xii.1931, 1♂, R.P. Claquin, Musée du Congo, identified *Xenogryllus*  
362 *eneopteroides* by L. Chopard, R. Det. N 2814 (MRAC); 1932, 1♂, 1♀, R.P. Claquin,  
363 Musée du Congo, identified *Xenogryllus eneopteroides* by L. Chopard, R. Det. DD3509  
364 (MRAC). Haut-Uele, vi.1925, 1♀, Musée du Congo, identified *Xenogryllus*  
365 *eneopteroides* by L. Chopard, R. Det. N 2814 (MRAC). Kibali-Ituri: Geti, ii-iv.1937, 1♂,  
366 Ch. Scops, Musée du Congo, identified *Xenogryllus eneopteroides* by L. Chopard, R.  
367 Det. B 5318 (MRAC). **Gabon:** Plateaux Batéké, piste Ekalla, 30 km W Leconi [Lekoni],  
368 milieu herbeux, 19.vi.1994L. Desutter-Grandcolas: 1 juvenile, n°61, jour [day], litière  
369 [leaf litter], molecular sample Xen-GA (MNHN-EO-ENSIF3442); 1♂, n°3, nuit [night],  
370 sur plante – base tronc [on plant, low on tree trunk] (MNHN). Bas Ogooué, 1♂,  
371 identified *Xenogryllus eneopteroides* Bol. by L. Chopard (MNHN). Ntoum, 19.vii.[19]85,  
372 A. Pauly rec, 2♂, 2♀, 2 juveniles, lumière (RBINS). **Ghana:** Eastern region,  
373 Hansowodze, 2 mls from Kade A.R.S. [Agricultural Research Station], 19.vii.1963, I. K.  
374 Bacheampong, 4♂ (NHMUK 010926524, 010926567, 010926528, 010926571).

375 Ghana, Togoland plateau, grassy, summit above [Nkonya] Wurupong, 23.xii.1959, N.D.  
376 Jago, 1♂, Brit. Mus. 1965-474 (NHMUK 010926535). Eastern region, Kade Agr. Res.  
377 Stn [Agricultural Research Station], 11.vii.1963, I. K. Bacheampong, 2♂, Brit. Mus.  
378 1965-474 (NHMUK 010926526, 010926538). Transvolta, Togoland, Amedzofe:  
379 3.v.1959, (montane grass), 1♀, N.D. Jago, Brit. Mus. 1965-474 (NHMUK 010926529);  
380 25.v.[19]59, N.D. Jago, Brit. Mus. 1965-474 (NHMUK 010926579). Western Reg.[ion]  
381 mls E of Elmina, 1°20W 5°05.5N, xii.1960, 1♂, 1♀, N.D. Jago, identified *Xenogryllus*  
382 *eneopteroides* by P.C. Tinning, 1966, Brit. Mus. 1965-474 (NHMUK 010926509,  
383 010926511). T.v.T [Trans-Volta Togo], Ho-Hohoe rd. 10 miles from Volta bridge,  
384 iii.1960, N.D. Jago, Brit. Mus. 1965-474 (NHMUK 010926553). Western Reg.  
385 Asankragua-Enchi rd., Tano R. Ferry, 24.ix.1962, 1♂, N.D. Jago, Brit. Mus. 1965-474  
386 (NHMUK 010926510). **Guinea:** Kaoulenta [Kéouagenta], [Mount] Nimba (Guinée), ii-  
387 vi.[19]42, M. Lamotte, 1♂, 1♀, 2 juveniles (MNHN). Nimba (Guinée), [Mount] Pierre  
388 Richaud, 900 m, M. Lamotte ii-iv.[19]42, 1♀, identified *Xenogryllus eneopteroides* Bol.  
389 by L. Chopard; 1♀ (MNHN). Nimba (Guinée), Lamotte, Amiet, Vanderplaeften, xii.56-  
390 v.57: Ziéla, 17.ii.[19]57, 1 juvenile; Ziéla, 20.iv.[19]57, 1 juvenile; sav. [savannah] entre  
391 Ziéla et Gbakbré, 9.xii.1956, 1 juvenile; Zgpo. [Zouguépo], 900 m, androp, 18.iv.[1957],  
392 2♀; sav. [savannah] Ziéla, 3.ii.[19]57, 1 juvenile; sav. [savannah] Ziéla, 20-  
393 25.xii.[19]56, 4 juveniles; sav. [savannah] Kéouagenta, 5.xii.[19]56, 2 juveniles; forêt  
394 Ziéla, 19.ii.[19]57, 1 juvenile; 1 juvenile, A.V. n°23, identified *Xenogryllus eneopteroides*  
395 Bol. larva by L. Chopard; 1 juvenile, A.V. n°38; 1 juvenile, 87CD; 14.iii.[19]57, 1  
396 juvenile, fge n°47 (MNHN). Mt. Nimba, Mt. Leclerc [Mount Jean Charles Leclerc], 1500  
397 m, 23.iii.1991. M. Lamotte, n°319, 1♀ (MNHN); 1♀, molecular sample X22-XenGU  
398 (MNHN-EO-ENSIF1494). Nimba (Guinée), M. Lamotte, xii.56-v.57, 29.x.[1956], 1♂,  
399 herbes A2 Ziela, identified *Xenogryllus eneopteroides* Bol. by R. Roy, 1959 (MNHN).  
400 Guinée Française, Diéké [Diecke], P. Chabamaud, 1920, 1♀, identified *Xenogryllus*  
401 *eneopteroides* Bol. by L. Chopard (MNHN). **Ivory Coast:** Lamto (Toumodi), Coll.  
402 E.N.S. Paris, Lamotte et collab.: 1-10.ii.[19]62, 3♂, 2♀; 1963, 1♂, Xen-Robillard-

403 morpho; 25-31.xi.[19]62, 1♀; 15-20.viii.[19]62, 1 juvenile; iv.[19]62, 1♀; 7.xi.[19]62, 1♀;  
404 20.xi.[19]62, 1♀; vi.[19]62, 1♀; 4.vii.[19]62, 1♀, mare sav. inondée [inundated  
405 savannah] (MNHN). Forêt à 15 km de Lamto, 10.ix.1968, C. Girard, 1♀, 1 juvenile  
406 (MNHN). Forêt de Yape, 1♂ (MNHN). Toumodi, xii.1930-iv.1931, Ch. Alluaud & P. A.  
407 Chapuis, 1♂ (MNHN). Danané, i.1939, 1♀ (MNHN). Lamto (Toumodi), C. Girard:  
408 12.ii.1968, savane [savannah], lisière forêt galerie [hedge of gallery forest] 6♂, 3♀, 5  
409 juveniles; 1♂, identified *Xenogryllus eneopteroides* Bol. by R. Roy, 1974; 12.iii.1968,  
410 lisière, forêt, savane, 1♂, molecular sample XenCI (MNHN-EO-ENSIF1481); 8♂, 2♀, 7  
411 juveniles; 1♂, photos MEB [SEM photos] (MNHN-EO-ENSIF1518); 15-30.vii.1968,  
412 savane [savannah], lisière forêt galerie [hedge of gallery forest] 1♂, 2♀, 3 juveniles; 27-  
413 28.ii.1968, 2♀, 1 juvenile, à la lumière; 10-20.iv.1968, 1♂, à la lumière (MNHN). Lamto  
414 (Toumodi), Paul Planquette: 1♀, PNB 18; 1 juvenile, PNB 13B; 15.i.[19]64, 1♀, PNB 64  
415 (MNHN). Lamto (Toumodi), 30.vi.[19]62, Coll. E.N.S. Paris, Lamotte *et al.* 1♂, identified  
416 *Xenogryllus eneopteroides* by L. Chopard (MNHN). Nimba [Mount]: Lamotte, 1946, Z4,  
417 1♀, identified *Xenogryllus eneopteroides* by R. Roy, 1967 (MNHN). Lamto, 1963, 1♂  
418 (MNHN). Assinie, 3295-85, Chaper, 1♂ (MNHN). Réserve du Banco [Banco national  
419 park], R. Paulian & G. Delamare, 2♂, 2♀, 2 juveniles (MNHN). Bouaké (C.I.),  
420 22.v.[19]62, coll. E.N.S. Paris, Lamotte et collab., 1♂ (MNHN). Taï, 18.viii.1978, G.  
421 Couturier, 1♀ (MNHN). Lamto, 1962, P. Le Gall, 2♂, 2♀ (MNHN). **Kenya:** CRS  
422 [Catholic Relief Service] Tebere, 0039.5S 3723E, 10.i.[19]72, I.A.D.R. 72/58, 1♀, Brit.  
423 Mus. 1982-71 (NHMUK 010926573). **Nigeria:** Lagos, Ikoyi, viii. 1951, L. Bala, 1♂,  
424 identified *Xeneogryllus eneopteroides* by L. Chopard (MNHN). U.C. [University College]  
425 Ibadan, 5.ix.1952, Tephrosia, leaf, Teph-66 (MNHN) coll. G.H. Caswell, 1♀ (MNHN).  
426 Ibadan, 6.vi.[19]23, 1♂, 1605 (NHMUK 010926581). Calabar, Nigeria, 10.iv.[18]82,  
427 10.30 A.M., W. edge of [?] Kwa swamp, E. edge of UNICAL staff quarters; about 50  
428 cm up on base of tangle of turns e.g. *Clitonia ? rubiginosa*, notes 72, 1♀, C.I.E. coll  
429 A.156 76 Pres by Comm Inst Ent B.M. 1985-1(NHMUK 010926537). **Republic of**  
430 **Congo:** Sibiti, xi-1963, mission A. Descarpentries et A. Villiers 1963-1964, 2♂, 1♀

431 (MNHN). Baie de Lagoa, envoi H. Deyrolle, 1♀ (MNHN). Mayumbe, Luki, L. Tiebers [?],  
432 1♀, Musée du Congo (MNHN). Odzala, x.1963, 3♂, 3♀, 1 juvenile (MNHN). Congo,  
433 1963, 1♀ (MNHN). Ile [island] M'Bamou, 26.vii.1968, Fidèle, 1♀ (MNHN). Boma, Sundi,  
434 P. Rolin, 1♂ (RBINS). Banana, F. Busschot, 4♀ (RBINS). **Sierra Leone:** Piste  
435 Bandankoro, 29.v.1963, savane de plaine ayant brûlé [burnt lowland savannah],  
436 mission ENS-IFAN aux Monts Loma, 1♂, identified *Xenogryllus eneopteroides* by R.  
437 Roy, 1992; 2♂, 3♀ (MNHN). Piste Keimadugu, savane [savannah], 520 m, 29.v.1963,  
438 mission ENS-IFAN aux Monts Loma, 2♂, 1♀ (MNHN). Firawa, savane [savannah],  
439 1.vi.1963, mission ENS-IFAN aux Monts Loma, 1♂, 1♀ (MNHN). Njala, E. Hargreaves:  
440 21.vii.[19]32, B.M. 1938-149, identified *X. eneopteroides* Bol. by L. Chopard, pres Imp.  
441 Inst. Ent. Brit. Mus. 1965-474 (NHMUK 010926552); x.1936, 1♂, pres Imp. Inst. Ent.  
442 Brit. Mus. 1965-474, 1♂, identified *Xenogryllus eneopteroides* by B. Uvarov (NHMUK  
443 010926534); 23.v.[19]25, 1♂, identified *X. eneopteroides* Bol. by L. Chopard, B. M.  
444 1938-149 (NHMUK 010926530); xi.1935, 1♂, pres Imp. Inst. Ent. Brit. Mus. 1936-252,  
445 1♂, identified *Xenogryllus eneopteroides* by B. Uvarov (NHMUK 010926576). **Uganda:**  
446 Entebbe, 20.xi.1970, W.J. Bailey, B.M.1977-246, identified *Xenogryllus eneopteroides*  
447 by B. C. Townsend, 1978 (NHMUK 010926570). Kisaru, 19.vii.1983, 1♂, H.B. Johnson,  
448 at light (NHMUK 010926551). **Zambia:** Lusaka, about fifty metres from the Great East  
449 Road in Chalimbana, Chongwe Distr., 19.xii.2013, William van Niekerk, 1♂, identified  
450 *X. eneopteroides* by William van Niekerk [online photograph]. Kapiri, 13.xi.1913, 2♀, L.  
451 Charliers, Musée du Congo, identified *Xenogryllus eneopteroides* by L. Chopard, R.  
452 Det. N 2814 (MRAC).

453 *Type locality.* Angola, Duque de Bragança, Bayão (=Kalandula).

454 *Distribution.* Sub-Saharan Africa: Angola, Benin, Cameroun, Central African Republic,  
455 Democratic Republic of the Congo, Gabon, Ivory Coast, Kenya (one large female  
456 specimen with unclear identification), Nigeria, Republic of Congo, Sierra Leone,  
457 Uganda, Zambia.

458     *Emended diagnosis.* Species of average size, close to *X. mozambicus* n. sp. and *X.*  
459     *maniema* n. sp., from which it differs by face almost flat in lateral view (more rounded in  
460     other species), shorter male FWs (short apical field), and shape of lophi in male  
461     genitalia. Differing from *X. lamottei* n. sp. and Asian species by following characters:  
462     well-carinated lateral angles of dorsal disc of pronotum (also carinated in *X.*  
463     *mozambicus* and *X. maniema*), very short ovipositor and male genitalia.

464     *Redescription.* Species of average size (Fig. 1A-C), coloration gray brown little  
465     contrasted. Eyes large, lateral, occupying almost half of head height in lateral view (Fig.  
466     3A-B). Face almost flat in lateral view, with typical whitish mask underlined by a black  
467     line below eyes and on mandibles. Pronotum dorsal disc strongly carinated laterally  
468     (Fig. 3B), coloration dark brown or brown, most often with a median dark brown band  
469     extended laterally near anterior margin, forming a T shape; lateral lobes almost  
470     homogeneously dark brown. First article of antennae dark brown.

471     *Male.* FWs very wide, dark coloration anterior to 1A including angle of 1A (Fig. 5A). File  
472     with 658 stridulatory teeth (n=2) on transverse part of 1A. Harp longer than wide. Cell  
473     c1 narrowed posteriorly, its fusion with cell b1 slightly shorter than its individualised  
474     part. Cell c2 large, d2 long and narrow. Mirror large, wider than long and little rounded,  
475     its inner angle forming an angle. Apical field short, wider than long, including four or  
476     five cell alignments.

477     *Male genitalia* (Figs 7A-C, 8A-B). Pseudepiphalllic lophi as long as rest of  
478     pseudepiphalllic sclerite, forming a long rectangle, sometimes slightly widened  
479     preapically; lophi with a narrow membranous inner margin setose basally (Fig. 8A-B);  
480     their bases fused until mid-length; apex pointed and divergent, ended by a thin lamella  
481     without hook-like inner dorsal expansion (in dorsal view, anterior base of lamella  
482     suggests that lamella may partly correspond to modified inner dorsal expansion of  
483     other species); ventral blade of lophi without longitudinal wrinkles, but with a strong  
484     ventral transverse carina, as in *X. mozambicus* n. sp. Pseudepiphalllic parameres with

485 a strong rectangular ventral lobe and a basal membranous lobe. Rami strong, their  
486 apex forming wide convergent hooks. Ectophallic apodemes strong, not lamellate.  
487 Ectophallic lateral expansions, lateral sclerites of ectophallic fold and endophallic  
488 sclerite partly fused, forming a wide ventral sclerotized plate, trifid apically; endophallic  
489 apodemes made of wide lateral lamellas and a narrow dorsal crest.

490 *Female* (Fig. 1C). Dorsal disc of pronotum almost rectangular, its posterior margin  
491 slightly bisinuate. FWs dorsal field with 7-10 longitudinal veins ( $m=8$ ,  $n=10$ ). Subgenital  
492 plate with a shallow apical indentation (Fig. 9A). Ovipositor (Fig. 10A) very short, about  
493 one third of FIII length.

494 *Female genitalia*. Copulatory papilla (Fig. 11A-B) conical and narrow, its apex rounded;  
495 well-sclerotized except base and apex.

496 *Life history traits*. According to Chiffaud & Gillon (1984), *X. eneopterooides* is found in  
497 Ivory Coast at the end of the dry season, in parts of savannah protected from fires.  
498 Their study revealed that the species feeds only on leaves and flowers of dicotyledones  
499 which are specific to savannah.

500 *Calling song*. According to the sonogram showed in Desutter (1983), analysed by  
501 Robillard & Desutter-Grandcolas (2004a, 2011), the calling song of *X. eneopterooides* is  
502 a short echeme made of five long syllables (syllable duration = 65 ms, period = 106.5  
503 ms. Echeme duration = 611 ms, echeme period = 1062 ms. The dominant frequency is  
504 low (3.6 kHz) and corresponds to the first peak of the spectrum.

505 *Measurements*. See Table 2.

506 *Taxonomic discussion*. The material examined reveals numerous slight differences in  
507 male genitalia (Fig. 8A-B) and size along the wide geographic distribution of the  
508 species. In particular, specimens from Congo differ from the material from the western  
509 coast of Africa. However, these differences are not sufficiently clear or are based on

510 too few individuals to delimit different species yet. More information and sampling will  
511 be necessary to improve the study *Xenogryllus* in Africa.

512

513 ***Xenogryllus carmichaeli* (Chopard, 1928)**

514

515 *Dionymus carmichaeli* Chopard, 1928: 30.

516 *Xenogryllus carmichaeli* – Chopard 1968: 350; Yin & Liu 1995; Cigliano *et al.* 2018  
517 (Orthoptera Species File Online).

518 *Madasumma carmichaeli* – Chopard 1969: 311.

519 *Nomen dubium* – this study.

520

521 *Taxonomic discussion.* According to the description of Chopard (1928), the female of  
522 *X. carmichaeli* is similar to *X. marmoratus* in terms of shape, size and coloration.  
523 However, the male specimen described by Chopard possesses five oblique veins,  
524 which is not consistent with any other existing species of *Xenogryllus*, as noted by  
525 Chopard himself. It is probable that the type series was heterogeneous, the male  
526 belonging to another subfamily than Eneopterinae. The male type may belong to the  
527 genus *Madasumma*, according the subsequent identification of the species by Chopard  
528 (1969), while the female could belong to *X. marmoratus*. The type locality, “India,  
529 Darjeeling District, alt. 100-300 ft, West Bengal”, could fit with *X. marmoratus*, but also  
530 *X. transversus* or *X. maichauensis*. Given these problems and the fact that the type  
531 series is missing in ZSI, Kolkata (RJ, pers. obs.), we consider this species as *nomen  
532 dubium*.

533

534    ***Xenogryllus lamottei* Robillard n. sp.**

535    (Figs 1H-I; 3C-D; 4B; 5B; 7D-E; 8C)

536

537    *Type material. Holotype*, ♂, **Guinea**: Simandou [Mount], Guinée [Guinea], Cpt.[camp]  
538    Fon Yenfédou, Ifan [Institut Français d'Afrique Noire], ix.1951, [M.] Lamotte (MNHN-  
539    EO-ENSIF10685).

540    *Type locality*: Mount Simandou, Guinea.

541    *Distribution*. Species only known from the type locality in Guinea.

542    *Etymology*. The species is dedicated to the great French entomologist Marcel Lamotte  
543    who collected the type specimen.

544    *Diagnosis*. Species of average size, characterized by male FWs not widened as in  
545    other species of the genus; general morphology differing from the other African species  
546    (*X. eneopteroides*, *X. mozambicus* n. sp. and *X. maniema* n. sp.) and more similar to  
547    the Asian species (*X. marmoratus*, *X. transversus*, *X. ululiu* and *X. maichauensis*) by  
548    the following characters: pronotum not carinated laterally (Fig. 3D); wrinkles on surface  
549    of male FWs weak; eyes small, restricted to dorsal quarter of head (reaching ½ of head  
550    in other African species); face almost flat in lateral view; male genitalia with short  
551    pseudepiphalllic lophi (Fig. 8C), close to that of *X. ululiu* (Fig. 8H), their apex short and  
552    bifid, with a pre-apical dorsal hook-like expansion (absent in other African species);  
553    ectophalllic fold and latero-ventral expansions shorter than in other species.

554    *Description*. In addition to the characters of the genus, species of average size (Fig.  
555    1H-I), coloration gray brown little contrasted. Fastigium longer than wide, thinner than  
556    in other species (Fig.4B), slightly widened apically. Face almost flat in lateral view (Fig.  
557    3D), with typical pale mask with black spots underlined by a thick black line below eyes  
558    (Fig. 3C); mandibles dark brown; clypeus and labrum mottled with yellow and dark  
559    brown; maxillary palpi dark brown. Scapes and first article of antennae dark brown,

560 flagellum light brown. Eyes rounded and restricted to posterior quarter of head in lateral  
561 view. Pronotum dorsal disc not carinated laterally (Fig. 3D), brown, with lateral margins  
562 underlined by a yellow line, with a median dark brown band, ticker posteriorly; posterior  
563 margin almost straight. FIII narrow, ended by a long linear region. Abdomen slightly  
564 shorter than FWs. Cerci dark brown.

565 *Male*. FWs as wide as abdomen (Fig. 5B), not widened as in other species of the  
566 genus; light brown, translucent, with weak wrinkles on surface. Dark coloration anterior  
567 to 1A not including angle of file. CuP visible posteriorly until angle of 1A. Angle of 1A  
568 straight. Harp wide, with two straight parallel oblique veins and a partial one, straight  
569 and reaching harp mid-length. Cell c1 not narrowed posteriorly. Mirror almost rounded,  
570 its inner limit forming a curve; underlined posteriorly by cells e1 and d2 fused together.  
571 Apical field forming a narrow triangle made of four cell alignments (E-H). CuA thin and  
572 brown; M thick, whitish; R and Sc almost fused, brown; M-R area dark mostly brown, its  
573 dorsal margin whitish; lateral field translucent brown crossed by 22 projections of Sc.

574 *Male genitalia* (Fig. 7D-E). Pseudepiphallic lophi forming a short rectangle posterior to  
575 wide base of pseudepiphallic sclerite; lophi (Fig. 8C) with a narrow membranous inner  
576 margin, setose basally; apex with a black hook-like inner dorsal expansion; sclerotized  
577 ventral blades of lophi with longitudinal wrinkles as in *X. transversus* and *X. ululiu*, but  
578 without strong ventral carina as in *X. eneopteroides* and *X. mozambicus*.  
579 Pseudepiphallic lateral membranous lobes small. Pseudepiphallic basal margin  
580 reinforcement weak. Rami thin and straight, their apex narrowed but not hooked  
581 innerly. Pseudepiphallic parameres ventral plate r-shaped,. Ectophallic arc not  
582 sclerotized. Ectophallic apodemes thin and divergent. Ectophallic fold and latero-dorsal  
583 expansions short. Endophallic sclerite and apodeme little differentiated.

584 *Female*. Unknown.

585 *Life history traits*. Unknown.

586 *Calling song*. Unknown.

587 *Measurements*. See Table 3.

588

589 ***Xenogryllus maichauensis Gorochov, 1992***

590 (Figs 1L-O; 3E-F; 4C; 5C; 7L-N; 8D; 9C; 10B; 11E-G; 12A; 13)

591

592 *Xenogryllus maichauensis* Gorochov, 1992: 9 – Kim & Pham 2014: 60; Cigliano *et al.*

593 2018 (Orthoptera Species File Online).

594 *Xenogryllus transversus* – Yin & Liu 1995: 96; Robillard *et al.* 2007: 1267 (song

595 frequency, wrong identification); He 2018: 516; Cigliano *et al.* 2018 (Orthoptera

596 Species File Online) (wrong identification).

597

598 *Type material. Holotype*, ♂, **Vietnam: Mai Châu** (ZIN) [examined through photographs

599 sent by A. Gorochov].

600 *Additional material examined. China: Yunnan*, Mengla, E101°33' N21°28', 3.ix.1991,

601 Zuyao Liu, Tianqi Wang & Heisheng Yin: 1♂ (14062446), 2♀ (14062547, 14062622)

602 (SIPPE). Mengla Distr., 12-23.x.2014, coll. Zhang Tao, 14♂, 2♀, (SNNU). San Chahe,

603 Mongyang, E98°22' N25°23', 31.vii.1995, Xianwei Liu, Weinian Zhang, Xingbao Jin, 1♂

604 (14062620) (SIPPE). Xi Shuang Ban Na, 580 m, 10.ix.1993, Xinyue Chen, 3♂,

605 (14064963, 14064964, 14064966); 2♂, 7.ix.1993 (14064968, no number); 1♂,

606 5.ix.1993 (14064967) (SIPPE). Monglun Xi Shuang Ban Na, E101°15' N21°56',

607 5.ix.1993, Longlong Yang, 2♀, (14064969, 14064971); 7.ix.1993, Xinyae Chen, 1♀

608 (14064970) (SIPPE). Jinhong, E101°15' N21°56°, Zuyao Liu, Tianqi Wang, Heisheng

609 Yin: 6.ix.1991, 3♂ (14062619, 14062546, 14062448); 8.ix.1991, 1♂ (14062447)

610 (SIPPE). Jinghong, [N22°0'6.97" 100°46'25.07"E], 9.ix.1991, Zuyao Liu, Tianqi Wang,

611 Heisheng Yin, 1♀ (14062449) (SIPPE). Mohan Vill., 950 m, 3-ix.2005, coll. Xue Guoxi,  
612 2♂ (NWAFU). **Guangdong**, Shen zhen, [22°33'59.69"N, 114° 2'25.53"E], 9.x.2012,  
613 Zhang Tao [online photograph]. **Thailand: Tak**, Doi Musoe, 700 m, Agric. Res. Stn at  
614 night, 9.x.1990, S. Ingrish: 1♀, 1♂, molecular sample XtrTh, identified *Xenogryllus*  
615 *transversus* by S. Ingrish (ZFMK). Mae Salid, Monkrathing, 17° 30' N, 98° 5' E, 700 m,  
616 19.ix.1989-21.ix.1989, S. Ingrisch, 1♂, mountain forest and agricultural land [day  
617 collecting and night collecting following stridulation, Stridulation recorded by S.Ingrisch  
618 0295DXEN.WAV [old ID CIGxentraSW03], identified *Xenogryllus transversus* by S.  
619 Ingrish (ZFMK). Umphang, 1-6 km S., 15° 59' N, 98° 50' E, 16.x.1991, S. Ingrisch, 1♂,  
620 bamboo forest, along road and trails, stridulation recorded by S. Ingrisch  
621 0510XENO.WAV, identified *Xenogryllus transversus* by S. Ingrish (ZFMK). **Chiang**  
622 **Mai**, Roadside bw Samoeng and Mae Rim, 18° 50' N, 99° 0', 1♂, 11.x.1991, S.  
623 Ingrisch, 1♂, stridulation recorded by S. Ingrisch 0502XENO.WAV, [old ID  
624 CIGxentraSW05], roadside at night, identified *Xenogryllus transversus* by S. Ingrish  
625 (ZFMK). **Petchabun**, Nam Nao, 16° 47' N, 101° 27' O, 1000 m, S. Ingrisch; 13.ix.1989-  
626 14.ix.1989, S. Ingrisch, 1♂, stridulation recorded by S. Ingrisch 0286XENO.WAV [old  
627 ID CIGxentraSW01], mixed Oak-Pine forest, grassy undergrowth, Bamboo thicket [day  
628 collecting and night collecting following stridulation], identified *Xenogryllus transversus*  
629 by S. Ingrish (ZFMK).

630 *Type locality*. Mai Châu, Vietnam.

631 *Distribution*. Southern China, Northern Vietnam and Northern Thailand.

632 *Emended diagnosis*. Species of large size, similar in size, venation and coloration to *X.*  
633 *transversus*, from which it differs mostly by male genitalia, with pseudopiphalllic lophi  
634 forming curved elongate hooks widened apically (Fig.8D), while lophi are straight in *X.*  
635 *transversus* (Fig. 8I).

636 *Redescription*. In addition to the characters of the genus, *X. maichauensis* has a large  
637 size and a light brown or golden coloration (Fig. 1L-O). Fastigium slightly widened

638 apically as in *X. ululiu* and *X. transversus* (Fig. 4C). Eyes rather small, located on face,  
639 restricted to the dorsal third of head in lateral view (Fig. 3E-F). Lateral angle of dorsal  
640 disc of pronotum not carinated, with a thin yellow band underlined by a thin black line  
641 anteriorly. Hind wings tail gray brown, twice as long a pronotum.

642 *Male.* Pronotum dorsal disc more rectangular than in *X. transversus*, with a wide  
643 median black longitudinal band, its posterior margin slightly bisinuate. FW venation  
644 (Fig. 5C): 1A forming a straight angle, with 299 stridulatory teeth (n=1) located on  
645 transverse part of 1A. Dark coloration anterior to 1A not including angle of file. Harp  
646 wide. Mirror large, well-rounded, its inner limit curved. Apical field forming a triangle  
647 longer than wide, with six cell alignments.

648 *Male genitalia* (Fig. 7L-N). Characteristic Y-shaped pseudoeiphallic lophi forming  
649 curved elongate hooks, widened apically, with a sharp dorsal preapical expansion (Fig.  
650 8D). Rami strong, convergent apically, with a ventral posterior expansion reaching  
651 base of pseudoeiphallic lateral membranous lobes. Ectophallic apodemes long and  
652 thin. Ectophallic fold and endophallic sclerite almost fused, forming a wide sclerotized  
653 plate, trifurcate posteriorly; endophallic apodemes made of wide lateral lamellas.

654 *Female.* Head slightly wider than pronotum, with strong jaws visible from dorsal view  
655 (Fig. 1N-O). Dorsal disc of pronotum almost rectangular, its posterior margin slightly  
656 bisinuate. FWs light brown, anterior dark spot larger than in *X. ululiu*, *X. transversus*  
657 and *X. marmoratus*. Dorsal field with nine strong longitudinal veins. Subgenital plate  
658 with a V-shaped apical indentation with rounded edges (Fig. 9C). Ovipositor (Fig. 10B)  
659 as long as FIII.

660 *Female genitalia.* Copulatory papilla (Fig. 11E-F) conical, its apex rounded and  
661 sclerotized.

662 *Life history traits.* In Vietnam, the type specimens were found in forest, on leaves of  
663 bushes near a stream (A. Gorochov, pers. comm.). According to the information

664 associated with the recordings made by S. Ingrish in Thailand (available on the  
665 Orthoptera Species Files online under the name *X. transversus* (Cigliano *et al.* 2018),  
666 the species was found in mixed oak-pine forest on grassy undergrowth with bamboo  
667 thicket. Males call at night on vegetation.

668 *Calling song.* (Figs 12A, 13) *X. maichauensis* was recorded in Thailand by S. Ingrish  
669 (Cigliano *et al.* 2018). At 24.5°C, the calling song is made of short echemes quickly  
670 repeated and composed of 2-4 long syllables ( $m = 3.02 \pm 0.63$ ), lasting for  $179.5 \pm 41.6$   
671 ms, with a period of  $502 \pm 42$  ms. Within echemes, the first syllable has a lower  
672 amplitude than the next ones. Syllables are rather long (duration =  $46.5 \pm 1.7$  ms) with  
673 a syllable period of  $65.1 \pm 1.7$  ms (syllable duty cycle = 71%). The frequency spectrum  
674 shows a dominant frequency at  $4.91 \pm 0.07$  kHz followed by two powerful harmonics.

675 *Measurements.* See Table 4.

676 *Taxonomic discussion.* The male genitalia of the specimen from Thailand shows  
677 slightly different pseudopiphalllic lophi compared to the holotype from Vietnam and  
678 examined males from China, with more straight apical branches and less globular apex  
679 (Fig. 7L-M). Such differences are however based on too few observations to  
680 characterise a new species.

681

682 ***Xenogryllus maniema Robillard & Jaiswara n. sp.***

683 (Figs 1J-K; 3G-H; 4D; 5D; 7F-G; 8E)

684

685 *Type material. Holotype,* ♂, **Democratic Republic of the Congo:** Lokandu, île [island]  
686 Biawa, vii.1939, Lt. Vissers, coll. Mus. Congo (MNHN-EO-ENSIF10686). *Paratypes*  
687 (5♂), **Democratic Republic of the Congo:** same information as holotype; 3♂,  
688 identified *Xenogryllus eneopteroides* by L. chopard, R. Det. B.5318 (MRAC). Katanga:  
689 Kafakumba, ix.1924, 1♂, G. F. Overlaet, coll. Musée du Congo, identified *Xenogryllus*

690   *eneopteroides* by L. Chopard, R. Det. N 2814 (MRAC). Congo Belge, Musosa  
691   [Mususa], ix.1939, H. J. Brédo, 1♂, I.G.13.212 (RBINS).

692   *Type locality.* Democratic Republic of the Congo, Lokandu, island Biawa.

693   *Distribution.* Species only known from forested areas in the eastern part of the  
694   Democratic Republic of the Congo.

695   *Etymology.* The species is named after the type locality. Maniema, which means  
696   “jungle” or “rain forest” in Kibangubangu dialect, is one of 26 provinces of the  
697   Democratic Republic of the Congo.

698   *Diagnosis.* Species of average size, closer to *X. mozambicus* n. sp., from which it  
699   differs by less rounded face in lateral view (Fig. 3H), pseudopiphallic lophi (Fig. 8E)  
700   ended by a long conical apex and with a widened inner membranous margin  
701   resembling that of in *X. maichauensis*. From *X. eneopteroides* and *X. mozambicus*, *X.*  
702   *maniema* differs by absence of T-shaped median band on pronotum, absence of  
703   transverse carina on ventral face of lophi, and by rami weak, without convergent hook-  
704   like apex. Differing from *X. lamottei* n. sp. and Asian species by strongly carinated  
705   lateral angles of pronotum dorsal disc (also carinated in *X. mozambicus* and *X.*  
706   *eneopteroides*).

707   *Description.* Species of average size, coloration yellow brown little contrasted (Fig. 1J-  
708   K). Eyes large, lateral, higher than long, occupying almost half of head height in lateral  
709   view (Fig. 3G-H). Face well-rounded in lateral view (less than in *X. mozambicus*), with  
710   typical whitish mask underlined by a black line below eyes and on mandibles.  
711   Pronotum dorsal disc strongly carinated laterally (Fig. 3H), coloration light brown, with a  
712   median dark brown band, not extended laterally near anterior margin; lateral lobes  
713   almost homogeneously brown. First article of antennae dark brown.

714   *Male.* FWs very wide (Fig. 5D), longer than abdomen; dark coloration anterior to 1A  
715   including angle of 1A. FW venation as in *X. eneopteroides*; apical field longer, forming

716 a long triangle made of five (n=2) cell alignments.

717 *Male genitalia* (Fig. 7F-G). Pseudepiphallic lophi twice longer than rest of  
718 pseudepiphallic sclerite; with a wide membranous inner margin (Fig. 8E); their bases  
719 fused until mid-length. Apex of lophi long and pointed, slightly convergent and ended  
720 by a thin lamella, without inner dorsal expansions. Ventral blade of lophi with faint  
721 longitudinal wrinkles, but without strong transverse carina as in *X. eneopteroides*.  
722 Pseudepiphallic parameres with a strong rectangular ventral lobe. Rami rather weak,  
723 their apex almost straight, not forming convergent hooks. Ectophallic apodemes strong,  
724 not lamellate. Ectophallic lateral expansions, lateral sclerites of ectophallic fold and  
725 endophallic sclerite partly fused, forming a wide ventral sclerotized plate, trifid apically;  
726 endophallic apodemes made of wide lateral lamellas and a narrow dorsal crest.

727 *Female*. Unknown.

728 *Life history traits*. Unknown.

729 *Calling song*. Unknown.

730 *Measurements*. See Table 5.

731

732 ***Xenogryllus marmoratus* (Haan, 1844)**

733 (Figs 1D-G; 3I-J; 4E; 5E; 7H-I; 8G; 9B; 11CD; 12B; 14)

734

735 *Gryllus (Phalangopsis) marmoratus* Haan, 1844: 235.

736 *Calyptotrypus marmoratus* – Saussure 1878: 714 (the figures do not correspond to the  
737 species *X. marmoratus*, which is consistent with the surprising distribution “Java, Iles  
738 de la Sonde” proposed by Saussure, where the species is not distributed); Bolívar  
739 1900[1899]: 805.

740 *Madasumma marmorata* – Chopard 1924: 56; Kirby 1906: 93.

741    *Xenogryllus marmoratus* – Chopard 1968: 350; Yin & Liu 1995: 96; Oshiro 1995: 43;  
742    Walker 2010: 27; Robillard & Desutter-Grandcolas 2004a: 578 (calling song); 2004b:  
743    273 (morphological phylogeny); 2006: 644 (molecular phylogeny); 2008: 67; 2011: 637;  
744    Anso *et al.* 2016: 9 (molecular phylogeny); Chintauan-Marquier *et al.* 2016: 62  
745    (molecular phylogeny); Vicente *et al.* 2017: 2203 (historical biogeography); Schneider  
746    *et al.* 2017 (tympanum morphology).

747    *Xenogryllus marmoratus marmoratus* Gorochov, 1992, in Gorochov & Belokobylskij  
748    1992: 11 (nominal subspecies); Ichikawa *et al.* 2000: 276; 2006: 180; Storozhenko &  
749    Paik 2007; Storozhenko *et al.* 2015: 145; Cigliano *et al.* 2018 (Orthoptera Species File  
750    Online); Olivero & Robillard 2017: 1 (mating behavior).

751    *Xenogryllus marmoratus unipartitus* (Karny, 1915) – Gorochov & Belokobylskij 1992:  
752    10 (subspecies).

753    Synonym names:

754    *Heterotrypus unipartitus* Karny, 1915 – Chopard, 1968: 350.

755    Common names: Pine cricket (English), matsumushi (Japanese = pine insect, waiting  
756    insect); bao ta ling (Chinese = pagoda bell).

757

758    *Type material.* Neotype, ♂ [new designation], Japan: Honshu, collection Finot,  
759    identified *Calyptotrypus* species nova by A. Finot (MNHN-EO-ENSIF1592).

760    *Additional material examined.* Japan: Honshu, Mie, ix.1957, F. Ohmachi, 1♂, 1♀,  
761    identified *Xenogryllus marmoratus* de Haan, Xma Robillard morpho (MNHN). Japon,  
762    1♀, Exposition Universelle 1869, #1276-69, identified *Dionymus marmoratus* by L.  
763    Chopard (MNHN-EO-ENSIF1593). Japon [Japan, no precise locality and date], 1♂, 1♀,  
764    identified *Dionymus marmoratus* by L. Chopard, Xma Robillard morpho; 1♂, 2♀  
765    (MNHN); 1♀ (NHMW); 1♂, 20.679, identified *X. marmoratus unipartitus* by Karny  
766    (NHMW); 4♀, identified *Calyptotrypus* species nova, collection Finot (MNHN). Kobe, H.

767 Fruhstorfer, 1♀, 24.125 (NHMW). Kyoto, I. Yamashiro, 1♂, Xma Robillard morpho  
768 MEB, MEB Ziegler; 1♀; 1♀, molecular sample T. Robillard 2004 (MNHN). Kioto  
769 [Kyoto], Y. Hirase, 1♂, identified *Madasumma marmorata* (Haan) and *Calyptotrypus*  
770 *marmoratus* (Haan) by Hebard, 1924 (MNHN). Kanagawa Prefecture, Kanate, Ooi-Cho  
771 Ashigara-kami-gun [35° 23' 00"N, 139° 08' 00"E], H. Sakai, 14.viii.2010, 1♂, 2♀, 1  
772 juvenile, identified *Xenogryllus marmoratus* by A. Ichikawa (MNHN); 1♂,  
773 enregistrement appel TR-male2 [call recording - MNHN-SO-2016-14364], identified  
774 *Xenogryllus marmoratus* by A. Ichikawa (MNHN-EO-ENSIF1704); 1♂, enregistrement  
775 appel TR-male1 [call recording, MNHN-SO-2016-14365], identified *Xenogryllus*  
776 *marmoratus* by A. Ichikawa (MNHN-EO-ENSIF1598). Japon, environs de Tokyo, J.  
777 Harmand, 1906 (MNHN). Kinki Distr.[ict], Wakayama Pref.[ecture], Hashimoto city,  
778 19.x.1986, A. Ichikawa, 1♂, identified *X. marmoratus* by A. Ichikawa (RBINS).  
779 **Tsushima**, H. Fruhstorfer, sept-oct [ix-x], 24.123 (NHMW). **China: Guanxi**, Jin Xiu,  
780 10.x.1981, E110°11' N24°07', 1♀ (14062635) (SIPPE). Longzhou, 1995-viii18/23,  
781 Xianwei Liu, Weinian Zhang, Xinbao Jin, 1♀ (14062635) (SIPPE). **Chongqing**, Beibei,  
782 E106°23' N29°48', 1.x.2000, Zhou, 1♂ (14080765) (SIPPE). **Anhui**, Huangshan,  
783 E118°19' N29°43' [Shanghai market], viii.2012, T. Robillard, 1♂ (TR40),  
784 enregistrement appel [call recording MNHN-SO-2018-36] (MNHN-EO-ENSIF1702); 1♂  
785 (TR2), enregistrement appel [call recording MNHN-SO-2018-52], molecular sample  
786 X12 (MNHN-EO-ENSIF1594). **Jiangsu**, Zhenjiang, 32°12'0.00"N 119°27'0.00"E  
787 [Shanghai market], viii.2012, T. Robillard, 1♂ (TR41), enregistrement appel [call  
788 recording, MNHN-SO-2018-38] (MNHN-EO-ENSIF1707); 1♂ (TR3), enregistrement  
789 appel [call recording MNHN-SO-2018-], molecular sample X13 (MNHN-EO-  
790 ENSIF3562). **Shanghai**, Padang Shanghai, E121° 32' N31° 13' [Shanghai market],  
791 viii.2012, T. Robillard, 1♂, molecular sample X11, enregistrement appel [call recording,  
792 MNHN-SO-2018-37] (MNHN-EO-ENSIF1599); 1♂ (MNHN). Shanghai, Prov. Klange[?],  
793 Musée Meude, O. Piel, 30.viii.[19]30, 1♀, #1623, 1♂, 6.ix.[19]30 (MNHN).  
794 **Guangdong**, Shenzhen Distr., Xichong, 27.ix.2014, coll. Zhang Tao, 4♂, 2♀ (SNNU).

795   **Henan**, Xinxian Distr., 4-8.ix.2014, coll. Ma Libin, 2♀ (SNNU). **Hainan**, Wuzhishan Mt.,  
796   12;viii.2010, coll. Jiang Chaozhong, 1♂ (NWAFU). **Zhejiang**, Tianlongshan Mt.,  
797   8.vii.2009, coll. He Zhuqing, 1♂ (ECNU). **Shaanxi**, Ankang, 10.viii.2017, coll. Ma Libin,  
798   1♂ (SNNU). **Taiwan**, Takao, Formosa, Sauter XI.[19]07, type of *Heterotrypus*  
799   *unipartitus*, det. Karny, identified *Dionymus formosanus* n. sp. Type by T. Shiraki, (DEI,  
800   Eberswalde), identified *Xenogryllus marmoratus unipartitus* (Karny) by A. V. Gorochov  
801   (DEI) [examined on photo]. **Sri Lanka**: Ceylon, Kandy, H. Rolle Berlin SW.11, 1♂  
802   (MNHN). **South Korea**: data from website “South Korea: Orthopteroids of Korea”  
803   (<http://www.jasa.pe.kr/pulmuchi/korthoptera/Xenogryllus-marmoratus.htm>).  
  
804   *Distribution*. Japan, China (including Taiwan and Hainan), South Korea  
805   (<http://www.jasa.pe.kr/pulmuchi/korthoptera/Xenogryllus-marmoratus.htm>), India  
806   (specimens observed by Chopard (1969)), Sri Lanka (1 male specimen).  
  
807   *Emmended diagnosis*. Species characterised by its small size (Fig. 1E-G), head dorsal  
808   coloration with a faint median dark band on vertex, and male genitalia with very long  
809   and thin pseudepiphalllic lophi (Fig. 8G); in male FWs (Fig. 5E), mirror wider than long,  
810   less rounded than in *X. maichauensis*, *X. ululiu* and *X. transversus*.  
  
811   *Redescription*. Species of small to average size, the smallest for the genus (Fig. 1E-G).  
812   Coloration almost homogeneously light ochre. Median dark band on head dorsum faint  
813   on vertex (Fig. 4E), prolonged anteriorly by black fastigium, with three narrow lateral  
814   bands on each side. Fastigium straight, not widened anteriorly. Antennae light brown,  
815   first segment darker. Face relatively flat in lateral view (Fig. 3J). Pronotum dorsal disc  
816   light brown, with a median brown band variably marked prolonging head coloration;  
817   lateral margins not carinated, with a narrow yellow line; lateral lobes ochre,  
818   progressively lighter ventrally.  
  
819   *Male*. FWs light brown, translucent (Fig. 5E); dark coloration anterior to 1A sometimes  
820   extended on angle of 1A. Transverse part of file almost straight, with 435 stridulatory  
821   teeth (n=1) on transverse part of 1A. Harp as wide as long. Fused part of cells c1 and

822 b1 as long as individualised part of c1. Cell d2 wider than in other species. Mirror wider  
823 than long, little rounded, its inner limit forming a wide angle. Apical field short, including  
824 five cell alignments.

825 *Male genitalia* (Fig. 7H-I). Pseudepiphalllic sclerite longer than rami; pseudepiphalllic  
826 lophi very long (Fig. 8G), narrow and sclerotized, slightly convergent, ended by an  
827 apical lamella curved dorsally and forming a small notch. Pseudepiphalllic sclerite with  
828 wide lateral membranous lobes. Basal reinforcement of pseudepiphalllic sclerite strong.  
829 Rami short with convergent hook-like apex. Pseudepiphalllic parameres with a ventral  
830 transverse crest absent in other species; inner apex of of pseudepiphalllic parameres  
831 sharp, slightly sclerotized apically. Ectophalllic apodemes strong, not lamellate.  
832 Ectophalllic lateral arms strong, fused to lateral arms of ectophalllic fold, and with lateral  
833 arms of endophalllic sclerite, forming together a wide gutter. Ectophalllic fold entirely  
834 sclerotized ventrally. Endophalllic sclerite small, apodeme made of a thin anterior dorsal  
835 crest and narrow lateral lamellas.

836 *Female*. FWs dorsal field with 8-9 strong longitudinal veins ( $m = 8$ ,  $n = 10$ ) and faint  
837 transverse ones. Subgenital plate not indented apically (Fig. 9B). Ovipositor longer  
838 than FIII, its apex rounded and smooth.

839 *Female genitalia*. Copulatory papilla (Fig. 11C-D) thin and conical, widened basally;  
840 sclerotization strong basally and apically, its membrane plicate.

841 *Life history traits*. *X. marmoratus* lives on low vegetation in bushes and grass in  
842 secondary habitats and along roads. The species has a nocturnal activity; males call at  
843 night from the vegetation. See Olivero & Robillard (2017) for description of peculiar  
844 mating behaviours.

845 *Calling song* (Figs 12B, 14). At 26.5°C, the calling song of *X. marmoratus* is made of  
846 relatively long syllables, each corresponding to one FW closure. Syllables form a two-  
847 part echeme: 1-3 singleton syllables ( $m = 2$ ,  $n = 10$ ), followed by a chirp made of three

848 syllables, sometimes duplicated. The amplitude profile of the call shows that the  
849 singletons are usually less loud than the chirp. Total echeme duration is  $711 \pm 33$  ms,  
850 for an echeme period of  $6776 \pm 368$  ms (echeme duty cycle = 10.5%). The syllables  
851 have the following characteristics: 1<sup>st</sup> syllable duration =  $51.8 \pm 1.5$  ms (period = 318.3  
852  $\pm 35.6$  ms); 2<sup>nd</sup> syllable duration =  $50.3 \pm 1$  ms; chirp duration =  $183.8 \pm 1$  ms; chirp  
853 syllable duration =  $49.5 \pm 1.9$  ms (period =  $68 \pm 0.8$  ms). The frequency spectrum  
854 shows a pure tone fundamental peak corresponding to the dominant frequency at ca.  
855 5.9  $\pm 0.2$  kHz, with a clear harmonic series, the third harmonic (ca. 18 kHz) being  
856 almost as powerful as the fundamental peak.

857 *Measurements.* See Table 6.

858 *Taxonomic discussion.* The types of De Haan, supposed to be in Leiden, were not  
859 found in RMNH in 2006 (TR, pers. obs.), with no record mentioning them as loaned  
860 (Rob De Vries, curator of the Orthopteran collection in RMNH, pers. com. 2006;  
861 confirmation by Luc Willemse, current curator of the Orthopteran collection in RMNH,  
862 pers. com. 2018). Specimens possibly belonging to the original type series were also  
863 searched for in other European museums, but could not be found. We thus designate a  
864 neotype from Japan that will serve as a reference for future systematic works.

865 Gorochov (1992) distinguished two subspecies, *X. m. marmoratus* (Haan, 1844) and *X.*  
866 *m. unipartitus* (Karny, 1915). In his study, he considered *Gryllus (Phalangopsis)*  
867 *marmoratus* Haan, 1844 (type from Japan), as the nominal subspecies *Xenogryllus*  
868 *marmoratus marmoratus*, while he established the subspecies *X. m. unipartitus* based  
869 on the species *Heterotrypus unipartitus* Karny, 1915, which was described from one  
870 female from Taiwan (Takao, examined on photo). The type of *H. unipartitus* clearly  
871 corresponds to a female of *Xenogryllus*, however, given the continuous variation  
872 observed across the wide distribution of *X. marmoratus*, and the limited information  
873 available based on female morphology, there is no reason to maintain these two  
874 subspecies. We thus synonymise the subspecies under *X. marmoratus*, and *H.*

875 *unipartitus* becomes a junior synonym of *X. marmoratus*, as previously proposed by  
876 Chopard (1968).

877

878 ***Xenogryllus mozambicus* Robillard n. sp.**

879 (Figs 2A-D; 3K-L; 4H; 5F; 7J-K; 8F; 9D; 11H-J; 12C; 15; 16)

880

881 *Xenogryllus eniopterooides* (wrong spelling of *X. eneopterooides*) – Toms 1984: 309.

882

883 *Type material. Holotype*, ♂, **Mozambique: Cabo Delgado**, Pantanos de Nhica, zone  
884 herbacée à l'Est du camp [herbaceous area E of camp], 10°45'19,1"S 40°13'00"E, 122  
885 m, 29.xi.2009, T. Robillard, nuit, enregistrement appel [call recording MNHN-SO-2018-  
886 140] (TR475) (MNHN-EO-ENSIF1517). *Allotype*, ♀, **Mozambique: Cabo Delgado**,  
887 mare SE de Nhica, bras mort de la Rovuma [pond SE Nhica, dead arm of Rovuma  
888 river], 10°45'41,9"S 40°13'31,7"E, 124 m, 27.xi.2009, T. Robillard, mort en élevage  
889 [dead in captivity], (MNHN-EO-ENSIF1539). *Paratypes* (8♂, 4♀): **Mozambique: Cabo**  
890 **Delgado**, same information as holotype: 1♂, 1♀, morts en élevage [dead in captivity]  
891 (MNHN-EO-ENSIF1576-ENSIF1556). Same information as allotype: 2♂, mort en  
892 élevage [dead in captivity] (MNHN-EO- ENSIF1557, ENSIF1508). Mare SE de Nhica,  
893 bras mort de la Rovuma [pond SE Nhica, dead arm of Rovuma river], 10°45'41,9"S  
894 40°13'31,7"E, 124 m, nuit, zone herbacée en bord de piste [herbaceous area near  
895 track] T. Robillard: 22.xi.2009, 1♂ (MNHN-EO-ENSIF1554), 3♀ (MNHN-EO-  
896 ENSIF1502, ENSIF1503, ENSIF1505); 23.xi.2009, 1♂ (TR208), enregistrement appel  
897 Take 224 [call recording MNHN-SO-2018-133] (MNHN-EO-ENSIF3079); 1♂ (TR206),  
898 enregistrement appel Take 222 [call recording MNHN-SO-2018-134], molecular sample  
899 X7-XenMoz (MNHN-EO-ENSIF1515). Mare SW de Nhica, bras mort de la Rovuma

900 [pond SW Nhica, dead arm of Rovuma river], 10°52'09,5"S 40°06'47,1"E, 122 m,  
901 24.xi.2009, nuit, zone herbacée en bord de piste [herbaceous area near track] T.  
902 Robillard: 1♂ (TR224), hautes herbes (h=1.3 m) [on high grass], enregistrement appel  
903 Take 228 [call recording MNHN-SO-2018-135], molecular sample X23 (MNHN-EO-  
904 ENSIF1559); 1♂, mort en élevage (MNHN-EO-ENSIF1581).

905 *Additional material examined.* **Mozambique:** Delagoa [Maputo Bay], xii.1898, 1 ♂,  
906 identified *Phormincter* species nova by A. Finot (MNHN). **South Africa:** Zululand,  
907 Mtunzini, 8.i.1952, R. Toms, 1♂, R146, identified *Xenogryllus eneopteroides* by B. C.  
908 Townsend, 1984, B.M.1983-166 (NHMUK 010926568). **Malawi:** Fish Eagle Bay Lodge,  
909 herbaceous area near lake Malawi (Mal7), S13°02'21.1" E34°19'34.8", 503 m,  
910 6.x.2018, night, 2♂, call recording, 1♀, T. Robillard, K. Salazar & R. Murphy (MNHN).

911 *Type locality.* Mozambique, Cabo Delgado, Pantanos of Nhica, 10°45'19,1"S  
912 40°13'00"E.

913 *Distribution.* Mozambique, South Africa, Malawi.

914 *Etymology.* Species named after the type locality.

915 *Diagnosis.* Species of average to large size, close to *X. eneopteroides* and *X. maniema*  
916 n. sp. From *X. eneopteroides*, the new species differs by rather larger size, face more  
917 rounded, almost globular in lateral view, absence of T-shaped median band on  
918 pronotum (also absent in *X. maniema*) larger mirror, and slight differences in male  
919 genitalia. From *X. maniema* n. sp., *X. mozambicus* mostly differs by male genitalia with  
920 larger lophi (Fig. 8F) and presence of transverse carina on ventral face of lophi (also  
921 present in *X. eneopteroides*). As *X. maniema* and *X. eneopteroides*, *X. mozambicus*  
922 differs from *X. lamottei* n. sp. and Asian species by strongly carinated lateral angles of  
923 pronotum.

924 *Description.* Species of average size (Fig. 2A-D), coloration light brown little contrasted.  
925 Eyes little prominent laterally (Fig. 3K), higher than long, occupying almost half of head

926 height in lateral view. Face globular in lateral view (Fig. 3L), with typical whitish mask  
927 underlined by a black line below eyes and on mandibles. Pronotum dorsal disc strongly  
928 carinated laterally, coloration light brown, with a median dark brown band, not extended  
929 laterally near anterior margin; lateral lobes almost homogeneously dark brown. First  
930 article of antennae dark brown.

931 *Male*. FWs very wide, longer than abdomen; dark coloration anterior to 1A including  
932 angle of 1A (Fig. 5F). Stridulatory file with 505 teeth (n=1) on transverse part of 1A. FW  
933 venation as in *X. eneopteroides*, mirror wider and apical field longer, forming a long  
934 triangle including 5-6 (n=8) cell alignments.

935 *Male genitalia* (Fig. 7J-K). Similar to *X. eneopteroides*, except longer pseudepiphalllic  
936 lophi (Fig 8F); transverse carina on ventral face of lophi present but weak.

937 *Female* (Fig. 2C-D). FWs dorsal field elongate, with 8-9 longitudinal veins (m = 9, n =  
938 6). Subgenital plate with a shallow apical indentation (Fig. 9D). Ovipositor short, about  
939 0.6 times FIII length.

940 *Female genitalia*. Copulatory papilla (Fig. 11H-J) as in *X. eneopteroides*, conical and  
941 narrow, well-sclerotized except base and apex.

942 *Life history traits*. *X. mozambicus* lives in areas of wet savannah (Fig. 15). Individuals  
943 of both sexes are found only at night in herbaceous vegetation, and males sing at night  
944 from the vegetation (ca. 30-80 cm high). Toms (1984) demonstrated that this species  
945 have directional calls and turn while calling, giving rise to changes in sound intensity.

946 *Calling song* (Figs 12C, 16). At 21.5 °C, males of *X. mozambicus* emit almost  
947 continuously long bouts of highly musical calling songs. After a warming phase, call  
948 bouts are made of successions of echemes of  $23 \pm 2$  syllables, barely separated from  
949 each other (echeme duration =  $3.13 \pm 0.29$  s; echeme period  $3.21 \pm 0.30$  s, echeme  
950 duty cycle = 97.6 %), with a regular amplitude profile, except for the last syllable, which  
951 is more intense. Within echemes, syllables are initially organized in  $4 \pm 1$  doublets,

952 which are followed by a series of similar syllables. Syllables are very long (syllable  
953 duration =  $126 \pm 5$  ms), with a relatively short syllable period of  $137 \pm 8$  ms (syllable  
954 duty cycle = 92%). All syllables are characterized by a large amplitude modulation: in  
955 initial doublets within echemes, the first syllables are less intense than the second ones  
956 and they start with a lower amplitude than their end (initial amplitude < ending  
957 amplitude), contrary to the second syllables of the doublets (initial amplitude > ending  
958 amplitude); the rest of the syllables have a higher starting amplitude, except for the last  
959 syllable of each echeme, which has a higher ending amplitude. The frequency  
960 spectrum shows a pure-tone dominant frequency at  $3.3 \pm 0.01$  kHz, followed by a rich  
961 harmonic content including three powerful harmonics at ca. 6.6, 9.9, 13.2 kHz,  
962 especially visible in ending syllables.

963 *Measurements*. See Table 7.

964 *Taxonomic discussion*. The specimens observed by Toms (1984) and identified  
965 *Xenogryllus eniopterooides* (wrong spelling for *X. eneopterooides*) from Zululand  
966 (Mtunzini and Eastern Transvaal (Clanor)), South Africa, belong to *X. mozambicus* n.  
967 sp. according to the male specimen observed from the series of Toms (NHMUK  
968 010926568).

969

970 ***Xenogryllus transversus* (Walker, 1869)**

971 (Figs 2E-H; 3O-P; 4E; 5G; 7O-P; 8I; 9G; 11M-O; 12D; 17)

972

973 *Platydactylus transversus* Walker, 1869: 88.

974 *Cardiodactylus?* *transversus* – Saussure 1878: 665.

975 *Madasumma transversa* – Kirby 1906: 94; Chopard 1968: 350.

976 *Xenogryllus transversus* – Chopard 1969: 310; Bhowmik 1976: 46; 1985: 67; Saeed *et*  
977 *al.* 1990: 309 (redescription); Yin & Liu 1995: 96; Nattier *et al.* 2011: 2199 (molecular  
978 phylogeny); Vicente *et al.* 2017: 2203 (historical biogeography); He 2018: 516; Cigliano  
979 *et al.* 2018 (Orthoptera Species File Online).

980 Synonym names:

981 *Dionymus calcaratus* Brunner von Wattenwyl, 1893: 213 – Chopard 1968: 350.

982 *Calyptotrypus roonwali* Bhowmik, 1977: 35 – Vasanth 1993: 131.

983

984 *Type material. Holotype*, ♂, **Bangladesh**: Silhat [= Silhet / Sylhet] (NHMUK010362937)  
985 [examined].

986 *Additional examined material. India: Meghalaya*, Jaintia Hills, 1♀, Narpuh R.F.  
987 Umprysung - off R. Lubha Sunapor Town, N25°06.683' E92°21.559', 149 ft.  
988 N25°06.983' E92°21.559', Elev.149ft, 19.x.2004, molecular sample Xtr-X2, Svenson  
989 (MNHN-EO-ENSIF87). India, 1♀, #6752 (MNHN). **Sikkim**, 1♂, identified *Dionymus*  
990 *calcaratus* Br. by L. Chopard (MNHN-EO-ENSIF1516). Lebong, 3000 ft, IX.1908, 1♀,  
991 H.M.L., identified *Dionymus calcaratus* Br. (NHMUK010362936). **Manipur**, Imphal  
992 Valley, alt 715 m., 11.x.1945, 1♂, HT of *Calyptotrypus roonwali* Bhowmik (ZSI).  
993 **Jammu**, Tawi river, 32.715736° 74.860403°, secondary area, 30.VII, 11 pm, 1♂,  
994 chorus, molecular sample X26 (MNHN-EOENSIF4395), 1♂ (MNHN). **Punjab**, IISER  
995 Mohali, 30.663169° 76.724212°, 6♂ (MJO\_765, 766, 411, 715, 413), 2 ♀ (MJO\_36,  
996 MJO\_726), molecular sample Xtr-715, 765, 766, R. Jaiswara (IISER Mohali).  
997 **Myanmar**: Bhamo, Birmania, Fea ix.1886, Museo Civ. Genova, 19.176, syntype of  
998 *Dionymus calcaratus* (NHMW). **Pakistan**: Mansehra, 34.333882° 73.201062°, 27.ix,  
999 1♂, W. A. Panhwar, molecular sample X25 (MNHN-EO-ENSIF4388).

1000 *Type locality*. Bangladesh, Silhet.

1001 *Distribution*. Myanmar, India (North and East), Bangladesh, Pakistan.

1002     *Emended diagnosis.* Species of large size, similar in size, venation and coloration to *X.*  
1003     *maichauensis*, from which it differs by male genitalia, with long sclerotized  
1004     pseudepiphalllic lophi ended by a sharp apex with a dorsal preapical pointed expansion  
1005     (Fig. 8I); from *X. ululiu*, *X. transversus* differs by its larger size and by larger  
1006     pseudepiphalllic sclerite in male genitalia, with lophi proportionally smaller, separated  
1007     by a deep indentation (lophi basally fused in *X. ululiu*).  
  
1008     *Redescription.* In addition to the characters of the genus, *X. transversus* has a large  
1009     size and a golden coloration (Fig. 2E-H). Fastigium 1.5 times wider than scape, slightly  
1010     widened apically as in *X. ululiu* and *X. maichauensis*. Eyes rather small, located on  
1011     face, restricted to the dorsal third of head in lateral view (Fig. 3O-P). Lateral angle of  
1012     dorsal disc of pronotum not carinated, as in *X. maichauensis* and *X. ululiu*, with a thin  
1013     yellow band underlined by a black line anteriorly. Hind wings tail gray brown, nearly as  
1014     long as long as pronotum.  
  
1015     *Male.* Pronotum dorsal disc forming a wide trapezoid, with a wide median black  
1016     longitudinal band prolonging vertex coloration; posterior margin slightly bisinuate. FW  
1017     venation (Fig. 5G): 1A forming a 90° angle, with 265 stridulatory teeth (n=1) on  
1018     transverse part of 1A. Dark coloration anterior to 1A including angle of file. Harp wide.  
1019     Mirror large, well-rounded, its inner limit forming a wide curve. Apical field forming a  
1020     triangle longer than wide, with six cell alignments.  
  
1021     *Male genitalia* (Fig. 7O-P). Pseudepiphalllic sclerite elongate, with a wide basis; lophi  
1022     separated by a large indentation (Fig. 8I), similar to that of *X. ululiu*, with a sharp apex  
1023     and a strong dorsal preapical expansion. Rami strong, convergent apically. Ectophalllic  
1024     apodemes strong, lamellate apically. Ectophalllic fold and endophalllic sclerite almost  
1025     fused, forming a long ventral gutter, wider than in *X. ululiu*, trifurcate posteriorly.  
1026     Endophalllic apodemes made of lateral lamellas.  
  
1027     *Female* (Fig. 2G-H). Head slightly wider than pronotum. Dorsal disc of pronotum almost  
1028     rectangular, its posterior margin slightly bisinuate. FWs slightly longer than abdomen,

1029 light brown, with anterior dark spot very small; dorsal field with 10-11 (n = 3) strong  
1030 longitudinal veins. Subgenital plate with a deep V-shaped apical indentation with sharp  
1031 edges (Fig. 9G). Ovipositor as long as FIII.

1032 *Female genitalia.* Copulatory papilla conical (Fig. 11M-O), wider than in *X.*  
1033 *maichauensis*, its apex rounded and sclerotized.

1034 *Life history traits.* *X. transversus* lives in open secondary habitats, where males usually  
1035 call in chorus from 120-180 cm height dense shrubs. In IISER Mohali (Northern India),  
1036 two males were seen calling from a bamboo plantation on one occasion, from a height  
1037 of 60 cm, where one male was probably trying to mount the other (RJ personal obs.). In  
1038 Jammu (Northern India), males were found calling from cannabis plant. Calling activity  
1039 starts late in the evening, almost around 22:00 hrs and continues until 03:00 hrs in the  
1040 morning. Spacing between calling males of this species is quite variable, the closest  
1041 males being spaced by approximately one meter horizontally.

1042 *Calling song.* (Figs 12D, 17). The calling songs of three males were recorded from  
1043 IISER Mohali campus. At 25-27 °C, the calling song of *X. transversus* is made of short  
1044 echemes quickly repeated and lasting for  $197 \pm 4$  ms (echeme period =  $873 \pm 13$  ms),  
1045 and composed of 4-5 long syllables ( $m = 4.5 \pm 0.5$ ). Within echemes, the syllables  
1046 usually show increasing amplitudes. Syllables are rather long (duration =  $31 \pm 4$  ms)  
1047 with a syllable period of  $46 \pm 3$  ms (syllable duty cycle = 71%). The frequency spectrum  
1048 shows a pure-tone dominant frequency at  $4.5 \pm 0.5$  kHz followed by two powerful  
1049 harmonics.

1050 *Measurements.* See Table 8.

1051 *Taxonomic discussion.* We confirm the status of junior synonyms of *Calyptotrypus*  
1052 *roonwali* and *Dionymus calcaratus* according to re-examination of type specimens (HT  
1053 of *C. roonwali* and one male ST of *D. calcaratus*).

1054

- 1055    ***Xenogryllus ululiu* Gorochov, 1990**
- 1056    (Figs 2I-L; 3M-N; 4G; 5H; 7Q-R; 8H; 9E; 11K-L; 12E; 18)
- 1057
- 1058    *Xenogryllus ululiu* Gorochov, 1990: 14 – Gorochov 1992: 10; Kim & Pham 2014: 60;
- 1059    Robillard *et al.* 2007: 1267 (song frequency); Cigliano *et al.* 2018 (Orthoptera Species
- 1060    File Online).
- 1061    *Type material. Holotype*, ♂, **Vietnam**: **Gia Lai**, Kannack, A.V. Gorochov (ZIN) [not  
examined]. *Paratype* (1♂), **Vietnam**: **Gia Lai**, Kannack, A.V. Gorochov: 8-16.ii.1988,  
1063    1♂ (MNHN-EO-ENSIF2809) [examined].
- 1064    *Additional material examined. Vietnam*: **Gia Lai**, 2.ii.1993, 1♀, identified *Xenogryllus*  
1065    *ululiu* by A.V. Gorochov, molecular sample XulV (MNHN-EO-ENSIF2810); 7.XI.1993,  
1066    1♂, molecular sample XulV2, A.V. Gorochov. **Thailand**: **Petchabun**, Nam Nao, 16° 47'  
1067    N, 101° 27' O, 1000 m, S. Ingrisch; 13.ix.1989-14.ix.1989, S. Ingrisch, 1♂, Stridulation  
1068    recorded by S. Ingrisch 0291XENO.WAV, mixed Oak-Pine forest, grassy undergrowth,  
1069    Bamboo thicket [day collecting and night collecting following stridulation], identified  
1070    *Xenogryllus ululiu* by S. Ingrish, molecular sample XulTh (ZFMK). **Loei**, NamNao NP at  
1071    night, 13(14.ix.1989, S. Ingrisch, 1♂, identified *Xenogryllus ululiu* by S. Ingrish,  
1072    molecular sample XulTh (ZFMK). **Cambodia**: **Kampong Spoe**, northern part of Elefan  
1073    mts [Dâmrei Mountains], Kiri-Rom National Park, 130 km NNE Sihanoukville, 300-  
1074    500m, 27.ix-1.x.2003, A. Gorochov, M. Berezin, 1♂, molecular sample X21-XulCam2,  
1075    1♀ (ZIN). **Kaoh Kong**, central part of Elefan mountains [Dâmrei Mountains], vill.  
1076    Styeng-Chkhral (100 km NE of Sihanoukville), 300-500m, 27.viii-6.ix.2003, A.  
1077    Gorochov, L. Anisyutkin, 1♂, molecular sample X20-XulCam1 (MNHN-EO-  
1078    ENSIF4385), 1♂ (ZIN). **Campot**, environs of Sihanoukville (=Kampong Som), 14-  
1079    21.ii.1998, A.V. Gorochov, 1♀ (ZIN). Sihanoukville, citi near Siam bay, environs, 22-  
1080    26.VIII.2003, A. Gorochov, L. Anisyutkin, 1♂, molecular sample X19-XulSi (ZIN).

1081   **China: Guangdong**, Shenzhen District, Wutongshan Mt., 21.ix.2014, coll. Zhang Tao,  
1082   5♂, 3♀, (SNNU).

1083   *Type locality*. Vietnam, Gia Lai, Kannack.

1084   *Distribution* (Fig. 16). Vietnam, Thailand, Cambodia, Southern China.

1085   *Emended diagnosis*. Species average to large size, close to *X. transversus* and *X.*  
1086   *maichauensis* in terms of shape and coloration, from which it differs by smaller size,  
1087   mirror slightly wider than long (less rounded than in *X. transversus* and *X.*  
1088   *maichauensis*), and male genitalia looking like a stockier version of that of *X.*  
1089   *transversus*, with pseudoeiphallic lophi basally fused together (Fig. 8H), with a sharp  
1090   posterior apex and a strong dorsal hook-like expansion. In females, head larger than in  
1091   other species, wider in dorsal view than rest of body.

1092   *Redescription*. In addition to the characters of the genus, *X. ululiu* has an average size  
1093   and a light brown or golden brown coloration (Fig. 2I-L). Head (Fig. 3M-N): Fastigium  
1094   slightly widened apically, as in *X. transversus* and *X. maichauensis* (Fig. 4G); eyes  
1095   rather small, restricted to the dorsal third of head in lateral view (Fig. 3N). Pronotum  
1096   dorsal disc with a wide median black longitudinal band, its lateral angles not carinated,  
1097   with a thin yellow band underlined by a black line ventrally. Hind wings tail gray brown,  
1098   near twice as long a pronotum.

1099   *Male*. FW venation (Fig. 5H): 1A forming a straight angle. Dark coloration anterior to 1A  
1100   not including angle of file. Harp wide, with one short incomplete oblique vein and two  
1101   complete ones slightly bisinuate. Mirror large, slightly wider than long, less rounded  
1102   than in *X. maichauensis* and *X. transversus*, its inner margin curved as in these  
1103   species. Apical field forming a triangle longer than wide, with 6-7 (n = 5) cell  
1104   alignments.

1105   *Male genitalia* (Fig. 7Q-R). Pseudoeiphallic lophi stocky, basally fused together, with a  
1106   sharp apex and a strong dorsal preapical expansion (Fig. 8H). Rami strong, convergent

1107 apically. Ectophallic apodemes strong, lamellate apically. Ectophallic fold and  
1108 endophallic sclerite almost fused, forming a long ventral gutter, trifurcate posteriorly  
1109 and extended anteriorly between rami. Endophallic apodemes made of anterior lateral  
1110 lamellas.

1111 *Female*. Head larger than in male (Fig. 2K-L), wider than rest of body in dorsal view.  
1112 Dorsal disc of pronotum almost rectangular, its posterior margin slightly bisinuate. FWs  
1113 light brown, anterior dark spot small. Dorsal field with nine strong longitudinal veins.  
1114 Subgenital plate with a wide, apical U-shaped indentation with sharp edges (Fig. 9E).  
1115 Ovipositor as long as FIII.

1116 *Female genitalia*. Copulatory papilla small, conical, its apex rounded and sclerotized  
1117 (Fig. 11K-L).

1118 *Life history traits*. According to S. Ingrish who recorded the species' call in Thailand,  
1119 males of *X. ululiu* sing at night in grassy undergrowth of mixed Oak-Pine forest or  
1120 bamboo thicket (Cigliano *et al.* 2018).

1121 *Calling song* (Figs 12E, 18). One male from Thailand was recorded by S. Ingrish at  
1122 21°C. The song consists of echemes during  $566 \pm 57$  ms, with a period of  $2041 \pm 358$   
1123 ms. Echemes are made of 4 or 5 syllables ( $m=5$ ), with syllable duration =  $108 \pm 3.8$  ms  
1124 and syllable period =  $120.3 \pm 3$  ms. Within echemes, the three starting syllables have a  
1125 lower amplitude than the final 1-2 syllables. Dominant frequency is  $4.8 \pm 0.4$  kHz and  
1126 corresponds to the fundamental frequency of the spectrum.

1127 *Measurements*. See Table 9.

1128

## 1129 **Discussion**

1130 The taxonomic revision of *Xenogryllus* is based on the thorough study of  
1131 specimens from natural history collections belonging to 17 institutions and on field work  
1132 in Mozambique, China and India. Our study reveals that the genus is relatively little

1133 diversified compared to genera of other tribes of the subfamily Eneopterinae. The  
1134 revision clarifies the status of all previously described species and subspecies (Table  
1135 1), and added three new species from Africa.

1136 The large geographical coverage of the available data for *Xenogryllus* suggest  
1137 that the most common species of the genus are now clearly defined, formally described  
1138 or redescribed and included in identification keys. However two of the three new  
1139 species described here are based on few specimens only (six males from three  
1140 localities in Congo for *X. maniema* n. sp. and a unique male from Guinea for *X. lamottei*  
1141 n. sp.). As a consequence, while some species of *Xenogryllus* have wider distributions  
1142 across Asia and Africa, where they occur mostly in savannahs and in open shrubby  
1143 areas, other species may have more limited distributions, perhaps in association with  
1144 more constrained and limited forested habitats. This would involve an undefined  
1145 proportion of endemic taxa yet to discover.

1146

## 1147 **Acknowledgements**

1148 We acknowledge the Indo-French Centre for Promotion of Advanced Research  
1149 (CEFIPRA) for funding this project (PIs: Rohini Balakrishnan & Laure Desuter-  
1150 Grandcolas), and the National Biodiversity Authority of the Government of India for  
1151 providing the necessary permits to allow us to carry out the fieldwork. The collaboration  
1152 between France and China was facilitated by CAMPUS FRANCE and the program  
1153 PHC XU GUANGQI (2012). This work was conducted in the context of the PhD thesis  
1154 of JD, which was funded by China Scholarship Council (CSC), the Innovation Funds of  
1155 Graduate Programs of Shaanxi Normal University [2012CXB019], the National Science  
1156 Foundation of China [Grant No. 31402006] and a supporting grant from “La Société  
1157 des Amis du Muséum” and the Institut de Systématique, Evolution, Biodiversité  
1158 (ISYEB). We thank Jérôme Constant and Carole Paleco (RBINS) for their help during  
1159 the study of the collections of the Royal Belgian Institute of Natural sciences, Brussels,

1160 funded by the SYNTHESYS European program (BE-TAF-6640); Andrej Gorochov  
1161 (ZIN) for providing samples from the collections of Saint Petersburg; Rob de Vries,  
1162 Caroline Pepermans and Luc Willemse (RMNH) for their help with the Leiden  
1163 collections, and the following persons for additional loans of material or photos:  
1164 Merdedes Paris (MNCN), Stéphane Hanot (MRAC), Stephan Blank (Müncheberg) and  
1165 Sigfrid Ingrisih (ZFMK). We also thank Simon Poulain (MNHN) for his help  
1166 photographing the specimens, and Karen Salazar (MNHN) for her help with line  
1167 drawing figures. We also thank Rohini Balakrishnan (CES, IISc Bangalore) and Manjari  
1168 Jain, (IISER Mohali) for supporting RJ with logistics and lab space to carry out part of  
1169 this work in India. We also thank the Orthopterist Society for supporting RJ with one-  
1170 month project grant to visit MNHN in 2017. We thank Jean-Yves Rasplus (CBGP,  
1171 Montpellier) and Oliver Pascal for organising the 2009 expedition in Mozambique; this  
1172 expedition was made possible by generous contributions from the Prince Albert II of  
1173 Monaco Foundation and the Stavros Niarchos Foundation; we would like to thank Pro-  
1174 Natura International and the Muséum national d'Histoire naturelle, Paris, for organising  
1175 the expedition under their Our Planet Reviewed programme, and the Instituto de  
1176 Investigação Agrária de Moçambique for helping with the administration and the  
1177 collecting permits issue.

1178

## 1179 **References**

1180 Anso, J., Barrabe, L., Desutter-Grandcolas, L., Jourdan, H., Grandcolas, P., Dong, J. &  
1181 Robillard, T. (2016) *Pixibinthus*, a new cricket genus endemic in New Caledonia:  
1182 phylogenetic study of a potential relict lineage from maquis minier vegetation.  
1183 *PLoS One*, 11, e0150920. <https://doi.org/10.1371/journal.pone.0150920>.

1184 Audacity Team (2016) Audacity(R): Free Audio Editor and Recorder [Computer  
1185 application]. Version 2.0.0 retrieved 12 January 2016 from  
1186 <https://audacityteam.org/>.

- 1187 Bhowmik, H.K. (1976) On the gryllid fauna (Gryllidae: Orthoptera) of the districts of the  
1188 North Bengal, with description of three new species. *Indian Museum Bulletin*, 11,  
1189 42–48.
- 1190 Bhowmik, H.K. (1977) Studies on Indian Crickets (Orthoptera: Insecta) with  
1191 Descriptions of Two New Species. *Records of Zoological Survey of India*, 73,  
1192 368–382.
- 1193 Bhowmik, H.K. (1985) Contribution to the gryllid fauna of the Western Himalayas  
1194 (Orthoptera: Gryllidae). *Records of the Zoological Survey of India*, 73, 1–74.
- 1195 Bolívar, I. (1890) Ortópteros de África del Museo de Lisboa. *Jornal de Ciencias  
1196 Mathematicas, Physicas e Naturaes*, Series 2, 73–232.
- 1197 Bolívar, I. (1900[1899]) Les Orthoptères de St-Joseph's College à Trichinopoly (Sud de  
1198 l'Inde). *Annales de la Société Entomologique de France*, 68, 761–812.
- 1199 Bolívar, I. (1910) Fauna de la Guinea Espanola. *Memorias de la Real Sociedad  
1200 Espanola de Historia Natural*, Tomo 1, 525–544.
- 1201 Brunner von Wattenwyl, C. (1893) Révision du système des Orthoptères et description  
1202 des espèces rapportées par M. Leonardo Fea de Birmanie. *Annali del Museo  
1203 civico di storia naturale di Genova*, 2, 13, 5–230.
- 1204 Chintauan-Marquier, I.C., Legendre, F., Hugel, S., Robillard, T., Grandcolas, P., Nel,  
1205 A., Zuccon, D. & Desutter-Grandcolas, L. (2016) Laying the foundations of  
1206 evolutionary and systematic studies in crickets (Insecta, Orthoptera): a multilocus  
1207 phylogenetic analysis. *Cladistics*, 32, 54–81. <https://dx.doi.org/10.1111/cla.12114>
- 1208 Chopard, L. (1928) Revision of the Indian Gryllidae. *Records of Indian Museum*, 30, 1–  
1209 36.
- 1210 Chopard, L. (1954) La Réserve naturelle intégrale du Mt. Nimba, III. Orthoptères  
1211 Ensifères, *Mémoires de l'Institut Français d'Afrique Noire*, 40, 25–97.

- 1212 Chopard L. (1968) Gryllides. Fam. Gryllidae: Subf. Mogoplistinae, Myrmecophilinae,  
1213 Scleropterinae, Cachoplistinae Pteroplistinae, Pentacenturinae, Phalangopsinae,  
1214 Trigonidinae, Eneopierinae; Fam.Oecanthidae, Gryllootalpidae. *In:* Beier, M. (Ed.),  
1215 *Orthopterorum Catalogus. Vol. 12.* Uitgeverij Dr. W. Junk N.V.'s, Gravenhage,  
1216 pp. 213–500.
- 1217 Chopard, L. (1969) *The fauna of India and adjacent countries. Orthoptera. Vol. 2.*  
1218 *Grylloidea.* Zoological Survey of India, Calcutta: Baptist Mission Press, 421 pp.
- 1219 Cigliano, M.M., H. Braun, D.C. Eades & Otte, D. (2018) Orthoptera Species File.  
1220 Version 5.0/5.0. [Retrieved 10 May 2018]. <http://Orthoptera.SpeciesFile.org>
- 1221 Desutter, L. (1987) Structure et évolution du complexe phallique des Gryllidea  
1222 (Orthoptera) et classification des genres néotropicaux de Grylloidea. 1re partie.  
1223 *Annales de la Société entomologique de France, Nouvelle Série,* 23, 213–239.
- 1224 Desutter-Grandcolas, L. (2003) Phylogeny and the evolution of acoustic  
1225 communication in extant Ensifera (Insecta, Orthoptera). *Zoologica Scripta,* 32,  
1226 525–561. <https://doi.org/10.1046/j.1463-6409.2003.00142.x>
- 1227 Dong, J., Kergoat, G.J., Vicente, N., Rahmadi, C., Xu, S. & Robillard, T. (2018)  
1228 Biogeographic patterns and diversification dynamics of the genus Cardiodactylus  
1229 Saussure (Orthoptera, Grylloidea, Eneopterinae) in Southeast Asia. *Molecular*  
1230 *Phylogenetics and Evolution,* 129 (2018) 1–14  
  
1231 <https://doi.org/10.1016/j.ympev.2018.06.001>
- 1232 Giglio-Tos, E. (1907) Ortotteri africani. Parte II, Blattodea, Mantodea, Phasmodea,  
1233 Locustodea, Gryllodea. *Bulletino dei Musei di Zoologia ed Anatomia comparata,*  
1234 *della R. Universita di Torino,* 22, 563, 1–26.
- 1235 Gorochov, A.V. (1990) New and insufficiently studied crickets (Orthoptera, Gryllidae)  
1236 from Vietnam and some other territories. *In:* Gorochov, A.V. [Ed.]. *News of*  
1237 *systematics and faunistics of Vietnam insects.* Part 1. Trudy Zoologicheskogo

- 1238       Instituta. Akademiiia nauk SSSR. Leningrad [= Proceedings of the Zoological  
1239       Institute, USSR Academy of Sciences, Leningrad] 209, pp. 3–28.
- 1240       Gorochov, A.V. (1992) New and little known crickets (Orthoptera, Gryllidae) from  
1241       Vietnam. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR*, 245, 3–16.
- 1242       Haan, D. (1844) Verhandelingen over de Natuurlijke Geschiedenis der Nederlansche  
1243       Overzeesche Bezittingen, 24, 235. <https://doi.org/10.5962/bhl.title.114730>
- 1244       Ichikawa, A. Murai, T. Honda, E. (2000) Monograph of Japanese crickets (Orthoptera;  
1245       Grylloidea). *Bulletin of the Hoshizaki Green Foundation*, 4, 257–332.
- 1246       Ichikawa, M. & Honda, E. (2000) Bulletin of the Hoshizaki Green Foundation, 4, 276.
- 1247       Karny, V.H. (1915) Orthoptera et Ootbecaria. *Supplementa Entomologica*, 4, 56–108.
- 1248       Karsch, F. (1893) Verzeichniss der von Herrn Dr. Paul Preuss in Kamerun  
1249       gesammelten Grillen. *Entomologische Nachrichten*, 19, 199–208.
- 1250       Kevan, D.K. & Knipper, H. (1961) Geraadflügler aus Ostafrika. *Beiträge zur  
1251       Entomologie*, 11, 356–413.
- 1252       Kirby, W. F. (1906) *A synonymic catalogue of Orthoptera* (Vol. 2). *Orthoptera  
1253       Saltatoria. Part I. (Achetidae et Phasgonuridae)*. The trustees of the British  
1254       Museum, London, pp. viii+562
- 1255       Nattier, R., Robillard, T., Desutter-Grandcolas, L., Couloux, A., & Grandcolas, P. (2011)  
1256       Older than New Caledonia emergence? A molecular phylogenetic study of the  
1257       eneopterine crickets (Orthoptera: Grylloidea). *Journal of Biogeography*, 38,  
1258       2195–2209. <https://doi.org/10.1111/j.1365-2699.2011.02563.x>
- 1259       Oki, S., Hayashi, I., Iwatsuki, N. & Morikawa, K. (2002) Study on sound generating  
1260       mechanism of *Xenogryllus marmoratus*. *Proceedings of Dynamics & Design  
1261       Conference*, 95.

- 1262 Olivero, P. & Robillard, T. (2017) Same-sex sexual behavior in *Xenogryllus*  
1263 *marmoratus* (Haan, 1844) (Grylloidea: Gryllidae: Eneopterinae): Observation in  
1264 the wild from YouTube. *Journal of Orthoptera Research*, 26, 1–5.  
1265 <https://doi.org/10.3897/jor.26.14569>
- 1266 Oshiro, Y. (1995) *Studies on the Gryllidae from the Ryukyu Islands*. Nakimushikai,  
1267 Okinawa, 131 pp.
- 1268 Ragge, D.R. (1955) *The wing-venation of the Orthoptera Saltatoria, with notes on*  
1269 *Dictyopteran wing-venation*. British Museum (Natural History), London, 159 pp.
- 1270 Ragge, D.R., Reynolds, W.J. & Natural History Museum (London, England) (1998).  
1271 The Songs of the Grasshoppers and Crickets of Western Europe. Harley Books,  
1272 Colchester, England, 612 pp. <https://doi.org/10.2307/3503528>
- 1273 Robillard, T. & Desutter-Grandcolas, L. (2004a) High-frequency calling in Eneopterinae  
1274 crickets (Orthoptera, Grylloidea, Eneopteridae): adaptive radiation revealed by  
1275 phylogenetic analysis. *Biological Journal of the Linnean Society*, 83, 577–584.  
1276 <https://doi.org/10.1111/j.1095-8312.2004.00417.x>
- 1277 Robillard, T. & Desutter-Grandcolas, L. (2004b) Phylogeny and the modalities of  
1278 acoustic diversification in extant Eneopterinae (Insecta, Orthoptera, Grylloidea,  
1279 Eneopteridae). *Cladistics*, 20, 271–293. <https://doi.org/10.1111/j.1096-0031.2004.00025.x>
- 1281 Robillard, T. & Desutter-Grandcolas, L. (2006) Phylogeny of the cricket subfamily  
1282 Eneopterinae (Orthoptera, Grylloidea, Eneopteridae) based on four molecular loci  
1283 and morphology. *Molecular Phylogenetics and Evolution*, 40, 643–661.  
1284 <https://doi.org/10.1016/j.ympev.2005.10.019>
- 1285 Robillard, T. (2006) Phylogenetic systematics of *Pseudolebinthus*, a new genus of  
1286 Eneopterinae crickets (Orthoptera, Grylloidea, Eneopteridae) from south-east

- 1287 Africa. *Systematic Entomology*, 31, 671–683. <https://doi.org/10.1111/j.1365->
- 1288 3113.2006.00347.x
- 1289 Robillard, T. & Desutter-Grandcolas, L. (2008) Clarification of the taxonomy of extant  
1290 crickets of the subfamily Eneopterinae (Orthoptera: Grylloidea; Gryllidae).  
1291 *Zootaxa*, 1789, 66–68.
- 1292 Robillard, T. & Desutter-Grandcolas, L. (2011) Evolution of calling songs as  
1293 multicomponent signals in crickets (Orthoptera: Grylloidea: Eneopterinae).  
1294 *Behaviour*, 148, 627–672.
- 1295 Robillard, T., Grandcolas, P. & Desuter-Grandcolas, L. (2007) A shift toward harmonics  
1296 for high-frequency spectra in Eneopterinae crickets (Orthoptera, Grylloidea,  
1297 Eneopteridae). *Canadian Journal of Zoology*, 85, 1264–1275.  
1298 <https://doi.org/10.1139/z07-106>
- 1299 Robillard T, Montealegre-Z, F., Desutter-Grandcolas, L., Grandcolas, P. & Robert, D.  
1300 (2013) Mechanisms of high-frequency song generation in brachypterous crickets  
1301 and the role of ghost frequencies. *Journal of Experimental Biology*, 216, 2001–  
1302 2011.
- 1303 Saeed, A. & Yousuf, M. (1990) New record of family Eneopteridae (Grylloidea:  
1304 Orthoptera) from Pakistan. *Pakistan Journal of Zoology*, 22, 309.
- 1305 Saussure, H. D. (1878) Mélanges Orthoptérolgiques. Memoires de la Société de  
1306 physique et d'histoire naturelle de Genève, 25, 1–836.
- 1307 Schneider, E.S., Römer, H., Robillard, T. & Schmidt, A.K.D. (2017) Hearing with  
1308 exceptionally thin tympana: Ear morphology and tympanal membrane vibrations  
1309 in eneopterine crickets. *Scientific Reports*, 7, 15266.  
1310 <https://doi.org/10.1038/s41598-017-15282-z>
- 1311 Storozhenko, S.Y. & Paik, J.C. (2007) *Orthoptera of Korea*. Dalnauka, Vladivostok, 232  
1312 pp.

- 1313 Storozhenko, S.Y., Kim, T.W. & Jeon, M.J. (2015) *Monograph of Korean Orthoptera*.  
1314 National Institute of Biological Resources, 145 pp.
- 1315 Su, Y.N. (2016) A simple and quick method of displaying liquid-preserved  
1316 morphological structures for microphotography. *Zootaxa*, 4208, 592–593.  
1317 <https://doi.org/10.11646/zootaxa.4208.6.6>
- 1318 ter Hofstede, H.M., Schoneich, S., Robillard, T. & Hedwig, B. (2015) Evolution of a  
1319 communication system by sensory exploitation of startle behavior. *Current  
1320 Biology*, 25, 1–8. <https://doi.org/10.1016/j.cub.2015.10.064>
- 1321 Vasanth, M. (1993) Studies on crickets (Insecta: Orthoptera: Gryllidae) of northeast  
1322 India. *Records of the Zoological Survey of India*, Occasional Paper, Calcutta,  
1323 132, 1–178.
- 1324 Vicente, N., Kergoat, G.J., Dong, J., Yotoko, K., Legendre, F., Nattier, R. & Robillard,  
1325 T. (2017) In and out of the Neotropics: historical biogeography of Eneopterinae  
1326 crickets. *Journal of Biogeography*, 44, 2199–2210.  
1327 <https://doi.org/10.1111/jbi.13026>
- 1328 Walker, F. (1869) *Catalogue of the Specimens of Dermaptera Saltatoria and  
1329 Supplement to the Blattariae in the Collection of the British Museum*, 224 pp.  
1330 <https://doi.org/10.5962/bhl.title.8149>
- 1331 Walker, B.L. (2010) *Toxic Archipelago: a history of industrial disease in Japan*. The  
1332 University of Washington press, 352 pp.
- 1333 Yin, H. & Liu, X.W. (1995) *Synopsis on the classification of Grylloidea and  
1334 Gryllotalpoidea from China*. Shanghai Science and Technology and Literature  
1335 Press, Shanghai.
- 1336

1337 **Tables**

1338

1339 **Table 1.**

1340 List of species and revised status of *Xenogryllus* species based on Robillard (2006)  
1341 and Orthoptera Species File Online (Cigliano *et al.* 2018). Junior synonym names for  
1342 which the status is not updated are not considered in this list but are presented in the  
1343 revision under the senior name.

Taxa	Status
<i>Xenogryllus</i> Bolívar 1890	valid genus (emended diagnosis, redescription)
<i>X. eneopteroides</i> Bolívar, 1890	valid species (redescription, type species)
<i>X. carmichaeli</i> (Chopard, 1928)	<i>nomen dubium</i> (new status)
<i>X. lamottei</i> Robillard n. sp.	new species
<i>X. maichauensis</i> Gorochov, 1992	valid species (redescription)
<i>X. maniema</i> Robillard & Jaiswara n. sp.	new species
<i>X. marmoratus</i> (Haan, 1844)	valid species (redescription, neotype, synonymy of subspecies)
<i>X. mozambicus</i> Robillard n. sp.	new species
<i>X. transversus</i> (Walker, 1869)	valid species (redescription)
<i>X. ululiu</i> Gorochov, 1990	valid species (redescription)

1344

1345

1346

1347

**Table 2.** Measurements of *Xenogryllus eneopteroides*

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>		
	Males (n=6)	20-27.1 (22.7)	3.4-3.9 (3.6)	6.1-7 (6.7)	21.5-24.2 (23.0)	8-8.3 (8.2)	7.4-8.3 (7.8)	20.0-22.4 (21.1)	3.1-3.6 (3.4)	
	Females (n=5)	16.9-22.2 (19.4)	3.8-4.1 (4)	5.8-6.0 (5.9)	20.0-20.9 (20.5)	4.9-5.3 (5.1)	7.3-8.4 (7.9)	22.1-22.6 (22.4)	2.2-3.6 (3)	
	TIIIL		TIIIs				Tallls	OL		
		las	Ibs	Oas	Obs		lds	Ods	Ols	
	Males (n=6)	20.0-21.5 (20.2)	9-13 (10.5)	9-11 (9.8)	18-21 (19.3)	8-13 (10.8)	0	6-7 (6)	3-6 (5)	-
	Females (n=5)	21.6-25.5 (23.6)	9-10 (9.5)	10-13 (11.5)	18-20 (19.0)	13-14 (13.5)	0	6-7 (6)	3-6 (5)	8-9 (8.42)

1348

1349

**Table 3.** Measurements of *Xenogryllus lamottei* n. sp.

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>	
Holotype male	24.2	3.1	4.6	20	6.6	5.7	19.5	3.3	
	TIIIL		TIIIs				Tallls	OL	
		las	Ibs	Oas	Obs		lds	Ods	Ols
Holotype male	?	?	?	?	?	?	?	?	-

1350

1351

**Table 4.** Measurements of *Xenogryllus maichauensis*

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>	
<b>Thailand</b>									
Male	33.3	3.8	5.7	24	8.2	8.4	22.9	4.1	
Female	23.7	3.7	5	24.6	5.5	9.4	22.5	4.2	
<b>China</b>									
Males (n=5)	30.6-32.8 (31.4)	3.3-3.9 (3.7)	4.8-5.1 (5)	21.3-24 (22.7)	7.7-8.1 (7.8)	4-5.8 (5.2)	21-22 (21.6)	3.5-4.6 (3.9)	
Females (n=2)	24-27.4 (25.7)	3.4-3.9 (3.6)	4.8-5.3 (5)	15.5-18 (18.5)	4.4-5.5 (5.1)	5-8.5 (6.2)	22.1-22 (22)	3.7-5.5 (4.9)	
	TIIIL		TIIIs				Tallls	OL	
		las	Ibs	Oas	Obs		lds	Ods	Ols
<b>Thailand</b>									
Male	24.4	12	11	19	13	0	3	6	-
Female	23	10	12	20	17	0	4	4	20.7
<b>China</b>									
Males (n=5)	20-21 (20.5)	11-15 (12.3)	10-12 (11)	15 (15)	12-13 (12)	0 (0)	3 (3)	3 (3)	-
Females (n=2)	21-22 (21.3)	8-11 (9)	10-12 (11)	14-20 (17)	14-17 (16)	0 (0)	4-5 (5)	3-4 (4)	23.6-24.2 (23.9)

1352

1353

**Table 5.** Measurements of *Xenogryllus maniema* n. sp.

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>	
Holotype male	21.8	2.9	4	18.3	6.6	6.4	?	?	
Males (n=5)	19.8-22.9 (21.3)	2.7-3 (2.9)	3.8-4.4 (4.1)	18.5-21.3 (19.3)	6.4-7.6 (6.9)	5.6-6.4 (5.8)	18.3-19.1 (18.8)	3.1-3.2 (3.1)	
	TIIIL		TIIIs				Tallls	OL	
		las	Ibs	Oas	Obs		lds	Ods	Ols
Holotype male	?	?	?	?	?	?	?	?	-
Males (n=5)	18.3-19.1 (18.7)	7-13 (10)	8-15 (12)	16-20 (18)	11-16 (14)	0 (0)	5-9 (7)	7-10 (8)	-

1354

1355

1356

**Table 6.** Measurements of *Xenogryllus marmoratus*

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>
Neotype male	19.4	2.9	4.1	15.5	6.7	3.5	15.3	2.8
Males (n=4)	18.6-22	2.5-3	3.9-4.8	15.5-17	6.4-6.7	3.1-4.3	14.2-15.8	2.6-3.4
(Male mean)	(20.4)	(2.9)	(4.3)	(16.5)	(6.5)	(3.4)	(15.3)	(3)
Females (n=5)	18.8-20	2.1-3.1	3.1-4.3	11.5-15	2.8-4	1.2-6.3	11.8-16.3	2.2-3.2
(Female mean)	(19.1)	(2.7)	(3.7)	(13.4)	(3.3)	(3.4)	(14.4)	(2.7)
	TIIIL	TIIIs				Tallis		OL
		<b>las</b>	<b>Ibs</b>	<b>Oas</b>	<b>Obs</b>	<b>lds</b>	<b>Ods</b>	<b>Ols</b>
Neotype male	11.1	9	11	14	16	0	4	5
Males (n=4)	11.1-15.8	5-12	10-12	11-20	11-16	0-2	3-7	5-7
(Male mean)	(14.1)	(9)	(11)	(16)	(5)	(1)	(5)	(5)
Females (n=5)	12.1-14.9	4-10	7-12	13-18	9-13	0-2	2-5	5-10
(Female mean)	(13.3)	(8)	(9)	(16)	(11)	(1)	(3)	(7)

1357

**Table 7.** Measurements of *Xenogryllus mozambicus* n. sp.

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>
Holotype male	25.5	3.3	4.5	20.5	7.3	7.5	21	3.6
Males (n=5)	22.5-25.5	2.9-3.3	4.3-4.5	21-22	7-7.2	7-7.5	19-20	3.1-3.9
(Male mean)	(23.8)	(3.2)	(4.4)	(21.2)	(7.2)	(7.3)	(20)	(3.5)
Female allotype	21.9	2.9	4.4	17.5	3.8	9	20.5	4
Females (n=4)	21.5-22.1	3-3.3	4.2-4.5	18-21.5	4-7.7	7.5-8.5	19-20	3.5-4
(Female mean)	(21.8)	(3)	(4.3)	(19.2)	(5.3)	(7.9)	(19.9)	(3.7)
	TIIIL	TIIIs				Tallis		OL
		<b>las</b>	<b>Ibs</b>	<b>Oas</b>	<b>Obs</b>	<b>lds</b>	<b>Ods</b>	<b>Ols</b>
Holotype male	20	13	10	15	19	1	8	4
Males (n=5)	18-20	9-13	9-14	11-16	16-21	1-2	8-10	4-5
(Male mean)	(18.2)	(11)	(12)	(14)	(18)	(1)	(9)	(5)
Female allotype	20	9	12	14	17	0	7	6
Females (n=4)	19-20	9-15	9-12	14-18	13-18	0-2	6-8	4-6
(Female mean)	(19.7)	(16)	(10)	(16)	(17)	(1)	(7)	(5)

**Table 8.** Measurements of *Xenogryllus transversus*

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>
Males (n=6)	27.8-34.6	3.4-3.9	6.1-7	21.5-24.2	8-8.3	7.4-8.3	20.0-22.4	3-4
(Male mean)	(31.4)	(3.6)	(6.7)	(23.0)	(8.2)	(7.8)	(21.1)	(3.7)
Females (n=2)	28.5-30	3.8-4.1	5.8-6.0	20.0-20.9	4.9-5.3	7.3-8.4	22.1-22.6	4.2-4.8
(Female mean)	(29.3)	(4)	(5.9)	(20.5)	(5.1)	(7.9)	(22.4)	(4.5)
	TIIIL	TIIIs				Tallis		OL
		<b>las</b>	<b>Ibs</b>	<b>Oas</b>	<b>Obs</b>	<b>lds</b>	<b>Ods</b>	<b>Ols</b>
Males (n=6)	20.0-21.5	9-13	9-11	18-21	8-13	0	2-5	2-3
(Male mean)	(20.2)	(10.5)	(9.8)	(19.3)	(10.8)	(0)	(3)	(3)
Females (n=2)	21.6-25.5	9-10	10-13	18-20	13-14	0	2-3	4-5
(Female mean)	(23.6)	(9.5)	(11.5)	(19.0)	(13.5)	(0)	(3)	(5)

1358

1359

**Table 9.** Measurements of *Xenogryllus ululiu*

	<b>BL</b>	<b>PronL</b>	<b>PronW</b>	<b>FWL</b>	<b>FWW</b>	<b>HWT</b>	<b>FIIIL</b>	<b>FIIIW</b>
Male paratype	28.2	3.5	4.3	22.2	7.9	7.1	22	3.6
Males (n=5)	23.3-28.2	3.3-3.5	4.2-4.8	20-22.2	7.1-8.1	3.3-7.9	19-22.7	3.3-4
(Male mean)	(26)	(3.4)	(4.5)	(21.5)	(7.7)	(5.8)	(21.3)	(3.7)
Females (n=3)	23.7-25.9	3.4-4	4.5-5	17.2-23.1	3.3-4	5.1-8.2	20-21.5	3.4-4.3
(Female mean)	(24.5)	(3.7)	(4.8)	(20.2)	(3.8)	(6.9)	(20.5)	(3.8)
	TIIIL	TIIIs				Tallis		OL
		<b>las</b>	<b>Ibs</b>	<b>Oas</b>	<b>Obs</b>	<b>lds</b>	<b>Ods</b>	<b>Ols</b>
Male paratype	21.8	12	10	20	14	1	4	6
Males (n=5)	19.9-21.8	11-12	10-11	16-20	14-16	0-1	4-7	5-8
(Male mean)	(20.8)	(11)	(10)	(18)	(15)	(0)	(5)	(6)
Females (n=3)	19.8-20.7	7-10	10-11	12-16	14-16	0	3-5	5-7
(Female mean)	(20.1)	(9)	(11)	(14)	(15)	(0)	(4)	(6)

1360 **Legends of figures**

1361  
1362 **Figure 1.** (A-C) *Xenogryllus eneopteroides*: male in dorsal (A) and lateral (B) views,  
1363 female in dorsal view (C); (D-G) *Xenogryllus marmoratus* male in dorsal (D) and  
1364 lateral (E) views, female in dorsal (F) and lateral (G) views; (H-I) *Xenogryllus*  
1365 *lamottei* n. sp.: male holotype in dorsal (H) and lateral (I) views; (J-K)  
1366 *Xenogryllus maniema* n. sp.: male paratype in dorsal (J) and lateral (K) views;  
1367 (L-O) *Xenogryllus maichauensis*: male in dorsal (L) and lateral (M) views,  
1368 female in dorsal (N) and lateral (O) views. Scale bar: 1 cm.

1369 **Figure 2.** (A-D) *Xenogryllus mozambicus* n. sp.: male in dorsal (A) and lateral (B)  
1370 views, female in dorsal (C) and lateral (D) views; (E-H) *Xenogryllus transversus*:  
1371 male in dorsal (E) and lateral (F) views, female in dorsal (G) and lateral (H)  
1372 views; (I-L) *Xenogryllus ululiu*: male in dorsal (I) and lateral (J) views, female in  
1373 dorsal (K) and lateral (L) views. Scale bar: 1 cm.

1374 **Figure 3.** Heads of *Xenogryllus* in facial (left panel) and lateral views (right panel): (A-  
1375 B) *Xenogryllus eneopteroides*; (C-D) *Xenogryllus lamottei* n. sp.; (E-F)  
1376 *Xenogryllus maichauensis*; (G-H) *Xenogryllus maniema* n. sp.; (I-J) *Xenogryllus*  
1377 *marmoratus*. (K-L); *Xenogryllus mozambicus* n. sp.; (M-N) *Xenogryllus ululiu*;  
1378 (O-P) *Xenogryllus transversus*. Scale bar: 1 mm.

1379 **Figure 4.** Heads of *Xenogryllus* in dorsal views: (A) *Xenogryllus eneopteroides*; (B)  
1380 *Xenogryllus lamottei* n. sp.; (C) *Xenogryllus maichauensis*; (D) *Xenogryllus*  
1381 *maniema* n. sp.; (E) *Xenogryllus marmoratus*; (F) *Xenogryllus transversus*; (G)  
1382 *Xenogryllus ululiu*; (H) *Xenogryllus mozambicus* n. sp. Scale bar: 1 mm.

1383 **Figure 5.** Male forewing venation in dorsal view: (A) *Xenogryllus eneopteroides*; (B)  
1384 *Xenogryllus lamottei* n. sp.; (C) *Xenogryllus maichauensis*; (D) *Xenogryllus*  
1385 *maniema* n. sp.; (E) *Xenogryllus marmoratus*; (F) *Xenogryllus mozambicus* n.  
1386 sp.; (G) *Xenogryllus transversus*; (H) *Xenogryllus ululiu*. Scale bar: 5 mm.

1387 **Figure 6.** Male metanotal glandular structures in *Xenogryllus eneopteroides*: (A)  
1388 glandular pit (drawing by G. Hodebert (MNHN) modified from Robillard &  
1389 Desutter-Grandcolas, 2004b), (B-C), SEM view of glandular pores on scutum.  
1390 Scale bars: 1 mm (A), 10 µm (B), 1 µm (C).

1391 **Figure 7.** Male genitalia: (A-C) *Xenogryllus eneopteroides*, dorsal (A), ventral (B) and  
1392 lateral (C) views; (D-E) *Xenogryllus lamottei* n. sp., ventral (D), lateral (E) views;  
1393 (F-G) *Xenogryllus maniema* n. sp., ventral (F), lateral (G) views; (H-I)  
1394 *Xenogryllus marmoratus*, ventral (H), lateral (I) views; (J-K) *Xenogryllus*

1395       *mozambicus* n. sp., ventral (J), lateral (K) views; (L-N) *Xenogryllus*  
1396       *maichauensis*, pseudepiphalllic lophi in ventral (L-M) and lateral (N) views, male  
1397       from South China (L), male from Thailand (M, N); (O-P) *Xenogryllus*  
1398       *transversus*, dorsal (O), ventral (P) views; (Q-R) *Xenogryllus ululiu*,  
1399       pseudepiphalllic lophi in dorsal (Q) and ventral (R) views. Scale bar: 1 mm.

1400       **Figure 8.** Right pseudepiphalllic lophi of male genitalia in dorsal view: (A-B) *Xenogryllus*  
1401       *eneopteroides*, male from Central African Republic (A), male from Ivory Coast  
1402       (B); (C) *Xenogryllus lamottei* n. sp.; (D) *Xenogryllus maichauensis*; (E)  
1403       *Xenogryllus maniema* n. sp.; (F) *Xenogryllus mozambicus* n. sp.; (G)  
1404       *Xenogryllus marmoratus*; (H) *Xenogryllus ululiu*; (I) *Xenogryllus transversus*.  
1405       Symbols: Preapical hook-like expansion figured in gray or with a black arrow;  
1406       dotted parts represent membranous areas; thin lines represent folds and reliefs  
1407       in sclerites. Scale bar: 1 mm.

1408       **Figure 9.** Apex of female subgenital plate and base of ovipositor in ventral view: (A)  
1409       *Xenogryllus eneopteroides*; (B) *Xenogryllus marmoratus*; (C) *Xenogryllus*  
1410       *maichauensis*; (D) *Xenogryllus mozambicus*; (E) *Xenogryllus transversus*; (F)  
1411       *Xenogryllus ululiu*. Scale bar: 1 mm.

1412       **Figure 10.** Apex of female ovipositor: (A) *Xenogryllus eneopteroides*; (B) *Xenogryllus*  
1413       *maichauensis*. Scale bar: 1 mm.

1414       **Figure 11.** Female copulatory papilla: (A-B) *Xenogryllus eneopteroides*, in ventral (A)  
1415       and lateral (B) views; (C-D) *Xenogryllus marmoratus*, in ventral (C) and lateral  
1416       (D) views; (E-G) *Xenogryllus maichauensis*, in ventral (E), dorsal (F) and lateral  
1417       (G) views; (H-J) *Xenogryllus mozambicus* n. sp., in ventral (H), dorsal (I) and  
1418       lateral (J) views; (K-L) *Xenogryllus ululiu*, in ventral (K) and lateral (L) views; (M-  
1419       O) *Xenogryllus transversus*, in ventral (M), dorsal (N) and lateral (O) views.  
1420       Scale bar: 0.5 mm.

1421       **Figure 12.** Comparison of calling songs, oscillogram of five echemes: (A) *Xenogryllus*  
1422       *maichauensis*; (B) *Xenogryllus marmoratus*; (C) *Xenogryllus mozambicus*; (D)  
1423       *Xenogryllus transversus*; (E) *Xenogryllus ululiu*. Scale bar: 1 s.

1424       **Figure 13.** Calling song of *Xenogryllus maichauensis*: (A) oscillogram of 5 echemes;  
1425       (B) detailed oscillogram (upper panel) and sonogram (lower panel) of 1  
1426       echeme; (C) frequency spectrum.

1427       **Figure 14.** Calling song of *Xenogryllus marmoratus*: (A) oscillogram of 5 echemes; (B)  
1428       detailed oscillogram (upper panel) and sonogram (lower panel) of 1 echeme;  
1429       (C) detailed oscillogram of one syllable; (D) frequency spectrum.

1430      **Figure 15.** (A-D) *Xenogryllus mozambicus* n. sp.: Savannah habitat in Mozambique  
1431                (A), male (B) and female (C) in bushy vegetation at night; male (D) predated by  
1432                Reduviidae sp.

1433      **Figure 16.** Calling song of *Xenogryllus mozambicus*: (A) oscillogram of 5 echemes; (B)  
1434                detailed oscillogram (upper panel), amplitude envelope (middle panel) and  
1435                sonogram (lower panel) of 1 echeme; (C) detailed oscillogram (upper panels)  
1436                and sonograms (lower panels) of one the two initial syllables (left) and of the  
1437                final syllable (right); (D) frequency spectrum.

1438      **Figure 17.** Calling song of *Xenogryllus transversus*: (A) oscillogram of 5 echemes; (B)  
1439                detailed oscillogram (upper panel) and sonogram (lower panel) of 1 echeme;  
1440                (C) frequency spectrum.

1441      **Figure 18.** Calling song of *Xenogryllus ululiu*: (A) oscillogram of 5 echemes; (B)  
1442                detailed oscillogram (upper panel) and sonogram (lower panel) of 1 echeme;  
1443                (C) frequency spectrum.


1444

1445



1446


1447



1448


60

1449   **Fig.3**  
1450




1451  
1452  
1453

1454 Fig. 4




1455  
1456

1457 **Fig. 5**




1458  
1459

1460 **Fig. 6**




1461  
1462

1463 **Fig. 7**  
1464




1465  
1466

1467    **Fig. 8**  
1468



1469  
1470  
1471

1472      **Fig.9**  
1473



1474  
1475  
1476

1477      ***Fig. 10***  
1478




1479  
1480

1481 **Fig. 11**

1482


1483



1484


1485  
1486

**Fig. 12**



1487  
1488


1489 **Fig. 13**  
1490



1491

1492 **Fig. 14**

1493



1494

1495 ***Fig. 15***

1496




1497

1498

1499


1500 Fig. 16



1501

1502


1503 **Fig. 17**



1504

1505

1506 **Fig. 18**



1507

1508

1509