



HAL
open science

CONFORMAL BOOTSTRAP IN LIOUVILLE THEORY

Colin Guillarmou, Antti Kupiainen, Rémi Rhodes, Vincent Vargas

► **To cite this version:**

Colin Guillarmou, Antti Kupiainen, Rémi Rhodes, Vincent Vargas. CONFORMAL BOOTSTRAP IN LIOUVILLE THEORY. *Acta Mathematica*, 2024, 233 (1), pp.33-194. 10.4310/acta.2024.v233.n1.a2 . hal-02866510v2

HAL Id: hal-02866510

<https://hal.science/hal-02866510v2>

Submitted on 24 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONFORMAL BOOTSTRAP IN LIOUVILLE THEORY

COLIN GUILLARMOU, ANTTI KUPIAINEN, RÉMI RHODES, AND VINCENT VARGAS

ABSTRACT. The conformal bootstrap hypothesis is a powerful idea in theoretical physics which has led to spectacular predictions in the context of critical phenomena. It postulates an explicit expression for the correlation functions of a conformal field theory in terms of its 3-point correlation functions. In this paper we give the first mathematical proof of the conformal bootstrap hypothesis in the context of Liouville theory, a 2-dimensional conformal field theory studied since the eighties in theoretical physics and constructed recently by F. David and the three last authors using probability theory. The proof is based on a probabilistic construction of the Virasoro algebra highest weight modules through spectral analysis of an associated self adjoint operator akin to harmonic analysis on non compact Lie groups but in an infinite dimensional setup.

“We developed a general approach to CFTs, something like complex analysis in the quantum domain. It worked very well in the various problems of statistical mechanics but the Liouville theory remained unsolved.”

— Alexander Polyakov, *From Strings to Quarks* (2008)

CONTENTS

1. Introduction	2
1.1. Overview	2
1.2. Probabilistic approach of Liouville CFT	4
1.3. DOZZ formula and Conformal Bootstrap.	6
1.4. Main result on conformal bootstrap	7
1.5. Conformal blocks and relations with the AGT conjecture.	8
2. Outline of the proof	9
2.1. Reflection positivity	9
2.2. Hilbert space of LCFT	9
2.3. Hamiltonian of Liouville theory	10
2.4. Spectral resolution of the Hamiltonian	11
2.5. Analytic continuation	11
2.6. Conformal Ward Identities	12
2.7. Organization of the paper	13
2.8. Notations and conventions	14
3. Reflection positivity	14
3.1. Gaussian Free Fields	14
3.2. Reflection positivity	16
3.3. Dilation Semigroup	18
4. Gaussian Free Field: dynamics and CFT aspects	20
4.1. Fock space and harmonic oscillators	20
4.2. Quadratic form	21
4.3. Dynamics of the GFF	22
4.4. Diagonalization of the free Hamiltonian using the Virasoro algebra	23
5. Liouville CFT: dynamics and quadratic form	26
5.1. Feynman-Kac formula	27
5.2. Quadratic forms and the Friedrichs extension of \mathbf{H}	28

6. Scattering of the Liouville Hamiltonian	33
6.1. The operators $\mathbf{H}, \mathbf{H}^0, e^{\gamma c} V$	34
6.2. Resolvent of \mathbf{H}	36
6.3. The Poisson operator	57
6.4. The Scattering operator	61
6.5. Holomorphic parametrization of the eigenstates	64
7. Probabilistic representation of the Poisson operator	64
7.1. Highest weight states	65
7.2. Descendant states	65
7.3. Stress Energy Field	66
7.4. Conformal Ward Identities	70
7.5. Computing the 3-point correlation functions of descendant states	72
8. Proof of Theorem 1.1	74
8.1. Definition of Conformal Blocks	74
8.2. Proof of Theorem 1.1	74
9. Proof of Proposition 7.11	76
9.1. Preliminary remarks	76
9.2. Gaussian integration by parts	77
9.3. Proof of Proposition 9.3	79
Appendix A. Elementary lemmas on the Virasoro algebra	85
Appendix B. The DOZZ formula	86
Appendix C. Integration by parts calculations	87
C.1. Analyticity of the vertex operators	88
References	89

1. INTRODUCTION

1.1. **Overview.** There are essentially two approaches to Quantum Field Theory (QFT) in the physics literature. In the first approach the quantum fields are (generalized) functions $\hat{\mathcal{O}}(t, x)$ on the space-time \mathbb{R}^{d+1} ($d = 3$ for the Standard Model of physics) taking values in operators acting in a Hilbert space \mathcal{H} of physical states. Matrix elements of products of quantum fields at different points $\langle \psi | \prod_{k=1}^N \hat{\mathcal{O}}_k(t_k, x_k) | \psi \rangle$ where $\psi \in \mathcal{H}$ (the “vacuum” state) are (generalized) functions on $\mathbb{R}^{N(d+1)}$. Physical principles (positivity of energy) imply that these matrix elements should have an analytic continuation to the *Euclidean* domain where $t_k \in i\mathbb{R}$ and be given there as correlation functions of *random fields* $\mathcal{O}(y)$ defined on $y \in \mathbb{R}^n$ where $n = d + 1$. In the second approach, based on the so-called path integral approach due to Feynman, these correlation functions are formally given as integrals over a space of generalized functions on \mathbb{R}^n , called fields, with the formal integration measure given explicitly as a Gibbs measure with potential a functional of the fields. The Euclidean formulation serves also as a setup for the theory of second order phase transitions in statistical mechanics systems where now $n \leq 3$. In this case one expects the correlation functions to possess an additional symmetry under the conformal transformations of \mathbb{R}^n and the QFT is now a Conformal Field Theory (CFT).

In practice most of the information on QFT obtained by physicists has been perturbative, namely given in terms of a formal power series expansion in parameters perturbing a Gaussian measure (and pictorially described by Feynman diagrams). In CFT however there is another, nonperturbative, approach going under the name of *Conformal bootstrap*. In this approach one postulates a set of special *primary* fields $\mathcal{O}_\alpha(y)$ (or operators $\mathcal{O}_\alpha(t, x)$ in the Hilbert space formulation) whose correlation functions transform as conformal tensors under the action of the conformal group. Furthermore one postulates a rule called the *operator product expansion* allowing to expand the product of two primary fields inside a correlation function (or a product of two quantum fields in the Hilbert space formulation) as a sum running over a subset of primary fields called the *spectrum* with explicit coefficients depending on the three point correlation functions, the

so called *structure constants* of the CFT (see Section 1.3). Hence the correlation functions of a CFTs are determined in the bootstrap approach if one knows its spectrum and the structure constants.

In the case of two dimensional conformal field theories ($d = 1$ or $n = 2$ above) the conformal symmetry constrains the possible CFTs particularly strongly and Belavin, Polyakov and Zamolodchikov [BPZ84] (BPZ from now on) showed the power of the bootstrap hypothesis by producing explicit expressions for the correlation functions of a large family of CFTs of interest to statistical physics among which the CFT that is believed to coincide with the scaling limit of critical Ising model. In a nutshell, BPZ argued that one could parametrize CFTs by a unique parameter c called the central charge and they found the correlation functions for certain rational values of c where the number of primary fields is finite (the minimal models). During the last decade the bootstrap approach has also led to spectacular predictions of critical exponents in the three dimensional case [PoRyVi]. The reader may consult [Ga] for some background on 2d CFTs.

Giving a rigorous mathematical meaning to these two approaches and relating them has been a huge challenge for mathematicians. On the axiomatic level the transition from the operator theory on Hilbert space to the Euclidean probabilistic theory was understood early on and for the converse the crucial concept of *reflection positivity* was isolated [OsSh73, OsSh75]. Reflection positivity is a property of the probability law underlying the random fields that allows for a construction of a canonical Hilbert space where operators representing the symmetries of the theory act. Reflection positivity is one of the crucial inputs in the present paper.

However on a more concrete level of explicit examples of QFTs mathematical progress has been slower. The (Euclidean) path integral approach was addressed by constructive field theory in dimensions $d + 1 \leq 4$ using probabilistic methods but detailed information has been restricted to the cases that are small perturbations of a Gaussian measure. In particular the 2d CFTs have been beyond this approach so far. A different probabilistic approach to conformal invariance has been developed during the past twenty years following the introduction by Schramm [Sch00] of random curves called Schramm-Loewner evolution (SLE). This approach, centred around the geometric description of critical models of statistical physics, has led to exact statements on the interfaces of percolation or the critical Ising model; following the introduction of SLE and the work of Smirnov, probabilists also managed to justify and construct the CFT correlation functions of the scaling limit of the 2d Ising model [ChSm12, CHI15] (see also the review [Pel19] for the construction of CFT correlations via SLE observables).

Making a mathematical theory of the BPZ approach triggered in the 80's and 90's intense research in the field of Vertex Operator Algebras (VOA for short) introduced by Borchers [Bo86] and Frenkel-Lepowsky-Meurman [FLM89] (see also the book [Hu97] and the article [HuKo07] for more recent developments on this formalism). Even if the theory of VOA was quite successful to rigorously formalize numerous CFTs, the approach suffers certain limitations at the moment. First, correlations are defined as formal power series (convergence issues are not tackled in the first place and are often difficult); second, many fundamental CFTs have still not been formalized within this approach, among which the CFTs with uncountable collections of primary fields and in particular Liouville conformal field theory (LCFT in short) studied in this paper. Moreover, the theory of VOA, which is based on axiomatically implementing the operator product expansion point of view of physics, does not elucidate the link to the the path integral approach or to the models of statistical physics at critical temperature (if any).

In their seminal work, BPZ were in fact motivated by the quest to compute the correlation functions in LCFT, which had been introduced a few years before by Polyakov under the form of a path integral in his approach to bosonic string theory [Po81]. Although BPZ failed to carry out the bootstrap program for LCFT¹, this was successfully implemented later in the physics literature by Dorn, Otto, Zamolodchikov and Zamolodchikov [DoOt94, ZaZa96]. Since then, LCFT has appeared in the physics literature in a wide range of fields including random planar maps (discrete quantum gravity, see the review [Ko11]) and the supersymmetric Yang-Mills theory (via the AGT correspondence [AGT10]). Recently, there has been a large effort in probability theory to make sense of Polyakov's path integral formulation of LCFT within the framework of random conformal geometry and the scaling limit of random planar maps: see [LeG13, Mier13, MS20a, MS20b, MS20c, DiDuDuFa, DuFaGwPfs, MiGw] for the construction of a random metric

¹See Polyakov's citation [Po08] at the beginning of this paper.

space describing (at least at the conjectural level) the limit of random planar maps and [DMS14, NS] for exact results on their link with LCFT². In particular, the three last authors of the present paper in collaboration with F. David [DKRV16] have constructed the path integral formulation of LCFT on the Riemann sphere using probability theory. This was extended to higher genus surfaces in [DRV16, GRV19]. In this paper, we will be concerned with LCFT on the Riemann sphere.

LCFT depends on two parameters $\gamma \in (0, 2)$ and $\mu > 0$ ³. In this paper we prove that for all $\gamma \in (0, 2)$ and $\mu > 0$ the probabilistic construction of LCFT satisfies the hypothesis of the bootstrap approach envisioned in [BPZ84]. In particular we determine the spectrum of LCFT and prove the bootstrap formula for the 4-point function in terms of the structure constants that were identified by the last three authors in [KRV20]. LCFT for $\gamma \in (0, 2)$ is a highly nontrivial CFT with an uncountable family of primary fields and a nontrivial OPE and we believe the proof of conformal bootstrap in this case provides the first nontrivial test case where a mathematical justification for this beautiful idea from physics has been achieved. Let us emphasize that determining the spectrum and the structure constants of LCFT is the cornerstone of the deep link between LCFT and representation theory via exact formulae for correlation functions. Indeed they lead to exact formulae for n point correlation functions on the sphere and pave the way towards understanding bootstrap formulae for higher genus surfaces [GKRV].

In a nutshell, our proof uses reflection positivity to construct a representation of the semigroup of dilations in \mathbb{C} on an explicit Hilbert space associated to LCFT. Its generator, called the *Hamiltonian* of LCFT, is a self-adjoint unbounded operator. It has the form of a Schrödinger operator acting in the L^2 -space of an infinite dimensional Gaussian measure with a non trivial potential which is a positive function for $\gamma \in (0, \sqrt{2})$ and a measure for $\gamma \in [\sqrt{2}, 2)$. We perform the spectral analysis of this operator to determine the spectrum of LCFT⁴ and show that the bootstrap formula for the 4 point correlations functions can be seen as a Plancherel formula with respect to its spectral decomposition. We obtain exact expressions for the generalized eigenfunctions of the Hamiltonian by deriving conformal Ward identities for LCFT correlation functions which reflect the underlying symmetry algebra of LCFT (i.e. the Virasoro algebra).

1.2. Probabilistic approach of Liouville CFT. Let us start with the physicists formulation of LCFT. It is a theory of a random (generalized) function $\phi : \mathbb{C} \rightarrow \mathbb{R}$ called the Liouville field. One is interested in averages of functionals F of ϕ formally given by the path integral

$$(1.1) \quad \langle F \rangle_{\gamma, \mu} := \int F(\phi) e^{-S_L(\phi)} \mathbb{D}\phi,$$

where S_L is the Liouville action functional

$$S_L(\phi) := \frac{1}{\pi} \int_{\mathbb{C}} (|\partial_z \phi(z)|^2 + \pi \mu e^{\gamma \phi(z)}) dz$$

and where dz denotes the Lebesgue measure on \mathbb{C} . It depends on two parameters $\gamma \in (0, 2)$ and $\mu > 0$ (called cosmological constant). The notation $\mathbb{D}\phi$ refers to a formal “Lebesgue measure” on the space of functions $\phi : \mathbb{C} \rightarrow \mathbb{R}$ obeying the asymptotic

$$\phi(z) \underset{|z| \rightarrow \infty}{\sim} -2Q \ln |z|$$

with $Q = \frac{2}{\gamma} + \frac{\gamma}{2}$. This asymptotic is formally required by conformal invariance whereby the field ϕ should be thought to be defined on the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. The physically interesting expectations in LCFT are the n -point correlation functions

$$(1.2) \quad \left\langle \prod_{i=1}^n V_{\alpha_i}(z_i) \right\rangle_{\gamma, \mu}$$

²It is beyond the scope of this introduction to state and comment all the exciting results that have been obtained recently in this flourishing field of probability theory.

³The case $\mu = 0$ is different and corresponds to Gaussian Free Field theory; for the study of a related model, see Kang-Makarov [KM13].

⁴The importance of understanding the spectral analysis of the Hamiltonian of LCFT was stressed by Teschner in [Te01] which was an inspiration for us.

of the exponentials of the Liouville field $V_\alpha(z) = e^{\alpha\phi(z)}$ called “vertex operators” in physics. In (1.2) the points $z_i \in \mathbb{C}$ are distinct and $\alpha_i \in \mathbb{C}$.

The recent work [DKRV16] gave a rigorous mathematical meaning to the correlation functions (1.2) via probability theory as we now describe. As usual the quadratic part of the action functional can be interpreted in terms of a Gaussian Free Field (GFF). We consider the GFF X on \mathbb{C} with the following covariance kernel

$$(1.3) \quad \mathbb{E}[X(z)X(z')] = \ln \frac{1}{|z-z'|} + \ln |z|_+ + \ln |z'|_+ := G(z, z')$$

with $|x|_+ = \max(|x|, 1)$. This GFF can be defined as a random generalized function on a suitable probability space $(\Omega, \Sigma, \mathbb{P})$ (with expectation $\mathbb{E}[\cdot]$): see Section 3.1. As is readily checked with the covariance (1.3) $X(z) \stackrel{law}{=} X(1/z)$ so that the GFF is naturally defined on the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. Furthermore it defines \mathbb{P} almost surely an element in the space of distributions $\mathcal{D}'(\hat{\mathbb{C}})$.

The second ingredient needed for making sense of the path integral is the following Gaussian multiplicative chaos measure (GMC, originally introduced by Kahane [Ka85])

$$(1.4) \quad M_\gamma(dz) := \lim_{\epsilon \rightarrow 0} e^{\gamma X_\epsilon(z) - \frac{\gamma^2}{2} \mathbb{E}[X_\epsilon(z)^2]} \frac{dz}{|z|_+^4}$$

where $X_\epsilon = X * \theta_\epsilon$ is the mollification of X with an approximation $(\theta_\epsilon)_{\epsilon>0}$ of the Dirac mass δ_0 ; indeed, one can show that the limit (1.4) exists in probability in the space of Radon measures on $\hat{\mathbb{C}}$ and that the limit does not depend on the mollifier θ_ϵ : see [RoVa10, RhVa14, Be17] for example. The condition $\gamma \in (0, 2)$ stems from the fact that the random measure M_γ is different from zero if and only if $\gamma \in (0, 2)$. With these definitions the rigorous definition of the Liouville field ϕ is

$$(1.5) \quad \phi(z) = c + X(z) - 2Q \ln |z|_+$$

and the expectation (1.1) for F continuous and non negative on $\mathcal{D}'(\hat{\mathbb{C}})$ is defined as ⁵

$$(1.6) \quad \langle F(\phi) \rangle_{\gamma, \mu} = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}[F(c + X - 2Q \ln |\cdot|_+) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{C})}] dc.$$

The variable $c \in \mathbb{R}$ stems from the fact that in the path integral (1.1) one wants to include also the constant functions on \mathbb{C} which are not captured by the GFF. Indeed, it is needed to ensure conformal invariance of LCFT [DKRV16]. We remark also that the expectation $\langle \cdot \rangle_{\gamma, \mu}$ is *not* a probability measure as $\langle 1 \rangle_{\gamma, \mu} = \infty$ [DKRV16].

Following [DKRV16], the n -point correlations (1.2) can be defined for *real* valued α_i via the following limit

$$(1.7) \quad \langle \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_{\gamma, \mu} := \lim_{\epsilon \rightarrow 0} \langle \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_{\gamma, \mu}$$

where $z_1, \dots, z_n \in \mathbb{C}$ are distinct,

$$(1.8) \quad V_{\alpha, \epsilon}(z) = |z|_+^{-4\Delta_\alpha} e^{\alpha c} e^{\alpha X_\epsilon(z) - \frac{\alpha^2}{2} \mathbb{E}[X_\epsilon(z)^2]}$$

and Δ_α is called the *conformal weight* of V_α

$$(1.9) \quad \Delta_\alpha = \frac{\alpha}{2} \left(Q - \frac{\alpha}{2} \right), \quad \alpha \in \mathbb{C}.$$

The limit (1.7) exists and is non trivial if and only if the following bounds hold

$$(1.10) \quad \sum_{i=1}^n \alpha_i > 2Q, \quad \alpha_i < Q, \quad \forall i = 1, \dots, n \quad \textbf{(Seiberg bounds)}.$$

One of the main results of [DKRV16] is that the limit (1.7) admits the following representation in terms of the moments of GMC

$$(1.11) \quad \langle \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_{\gamma, \mu} = \gamma^{-1} \left(\prod_{1 \leq j < j' \leq n} \frac{1}{|z_j - z_{j'}|^{\alpha_j \alpha_{j'}}} \right) \mu^{-s} \Gamma(s) \mathbb{E}[Z^{-s}]$$

⁵In the recent paper [KRV20], the authors used a convention where the RHS is multiplied by 2.

where $s = \frac{\sum_{i=1}^n \alpha_i - 2Q}{\gamma}$, Γ is the standard Gamma function and (recall that $|x|_+ = \max(|x|, 1)$)

$$Z = \int_{\mathbb{C}} \left(\prod_{i=1}^n \frac{|x|_+^{\gamma \alpha_i}}{|x - z_i|^{\gamma \alpha_i}} \right) M_{\gamma}(dx).$$

We stress that the formula (1.11) is valid for correlations with $n \geq 3$, which can be seen at the level of the Seiberg bounds (1.10)⁶. Also it was proved in [DKRV16, Th 3.5] that these correlation functions are *conformally covariant*. More precisely, if z_1, \dots, z_n are n distinct points in \mathbb{C} then for a Möbius map $\psi(z) = \frac{az+b}{cz+d}$ (with $a, b, c, d \in \mathbb{C}$ and $ad - bc = 1$)

$$(1.12) \quad \left\langle \prod_{i=1}^n V_{\alpha_i}(\psi(z_i)) \right\rangle_{\gamma, \mu} = \prod_{i=1}^n |\psi'(z_i)|^{-2\Delta_{\alpha_i}} \left\langle \prod_{k=1}^n V_{\alpha_k}(z_k) \right\rangle_{\gamma, \mu}.$$

Because of relation (1.12), the vertex operators are *primary fields* in the language of CFT. The Möbius covariance implies in particular that the three point functions are determined up to a constant, called the *structure constant*, which we write as

$$(1.13) \quad \langle V_{\alpha_1}(0)V_{\alpha_2}(1)V_{\alpha_3}(\infty) \rangle_{\gamma, \mu} := \lim_{|u| \rightarrow \infty} |u|^{4\Delta_{\alpha_3}} \langle V_{\alpha_1}(0)V_{\alpha_2}(1)V_{\alpha_3}(u) \rangle_{\gamma, \mu}.$$

Similarly the four point function can be reduced to a function of one complex variable defined by

$$(1.14) \quad \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma, \mu} := \lim_{|u| \rightarrow \infty} |u|^{4\Delta_{\alpha_4}} \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(u) \rangle_{\gamma, \mu}.$$

Let us now turn to the conformal bootstrap approach to LCFT.

1.3. DOZZ formula and Conformal Bootstrap. In theoretical physics conformal field theory is a quantum field theory with conformal group symmetry. In particular one then postulates existence of primary fields whose correlation functions transform covariantly under conformal maps (Möbius transformations in the two dimensional case). The basic physical axiom of conformal field theory apart from this covariance is the *operator product expansion* (OPE for short). Denoting the primary fields still as $V_{\alpha}(x)$ for $x \in \mathbb{R}^n$ then, in very loose terms and cutting many corners, the OPE is the identity

$$(1.15) \quad V_{\alpha}(x)V_{\alpha'}(x') = \sum_{\beta \in \mathcal{S}} C_{\alpha\alpha'}^{\beta}(x, x', \nabla_x) V_{\beta}(x)$$

where \mathcal{S} labels a special set of primary fields called the *spectrum* of the CFT and the $C_{\alpha\alpha'}^{\beta}$ are differential operators *completely determined* by the conformal weights of the fields $V_{\alpha}, V_{\alpha'}, V_{\beta}$ and linear in the structure constants $C_{\alpha\alpha'\beta}$ (defined in general analogously to (1.13)). The identity (1.15) is assumed to hold once inserted in the correlation functions. Obviously a repeated application of the OPE allows to express an n -point function of the CFT as a sum of products of the structure constants. Hence to “solve a CFT” one needs to find its structure constants and spectrum. To find these, the following approach where the 4-point function⁷ (1.14) plays a fundamental role has been extremely fruitful. Applying the OPE (1.15) in (1.14) to α_1 and α_2 leads to a quadratic expression in the structure constants. Applying the OPE instead to α_2 and α_3 yields another quadratic expression and equating the two produces a quadratic equation for the structure constants. Varying $\alpha_1, \dots, \alpha_4$ results in a set of quadratic equations labeled by 4-tuples of $\alpha_i \in \mathcal{S}$ (for LCFT see (1.17)). These 4-point bootstrap equations pose strong constraints on the spectrum and the structure constants and have been used to great effect e.g. in the case of the 3-dimensional Ising model [EPPR12a, EPPR12b]. In two dimensions their study in the case when one of the fields V_{α_i} is a so-called degenerate field led in [BPZ84] to the discovery of the *minimal models* (2d Ising model among them) and their spectra and structure constants.

The motivation for the BPZ paper [BPZ84] was to solve the bootstrap equations for LCFT but in this they were unsuccessful [Po08]. The spectrum of LCFT was conjectured in [CuTh, BCT82, GeNe84] to be *continuous* $\mathcal{S} = Q + i\mathbb{R}_+$ and an explicit formula (see Appendix B) for the structure constants, the DOZZ formula, was postulated by Dorn, Otto, Zamolodchikov and Zamolodchikov [DoOt94, ZaZa96]. A

⁶In fact, one can extend the probabilistic construction (1.11) a bit beyond the Seiberg bounds but the extended bounds also imply $n \geq 3$. We will not discuss these extended bounds in this paper.

⁷In $d > 2$ the four point function is a function of two complex variables.

derivation based on the degenerate 4-point bootstrap equations was subsequently given by Teschner [Te95] and further evidence for the formula was given in [Te01], [Te04]. The DOZZ expression which we denote by $C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, \alpha_3)$ is analytic in the variables $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{C}$ (with a countable number of poles) and one is led to expect that it coincides with the probabilistic expression (1.13) on the domain of validity of the probabilistic construction, i.e. for real $\alpha_1, \alpha_2, \alpha_3$ satisfying the Seiberg bounds (1.10). This is indeed the case: in a recent work [KRV20], the last three authors proved that the probabilistically constructed 3-point correlation functions satisfy the DOZZ formula:

$$\langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(\infty) \rangle_{\gamma,\mu} = \frac{1}{2}C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, \alpha_3)^8.$$

Given that the spectrum is $\mathcal{S} = Q + i\mathbb{R}_+$, the formula (1.15) (where the sum becomes an integral) leads formally to the following bootstrap conjecture for the 4 point correlation functions

$$(1.16) \quad \begin{aligned} & \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma,\mu}^{\text{Boot}} \\ &= \frac{1}{8\pi} \int_0^\infty C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)C_{\gamma,\mu}^{\text{DOZZ}}(Q + iP, \alpha_3, \alpha_4)|z|^{2(\Delta_{Q+iP} - \Delta_{\alpha_1} - \Delta_{\alpha_2})}|\mathcal{F}_P(z)|^2 dP \end{aligned}$$

where \mathcal{F}_P are holomorphic functions in z called (spherical) *conformal blocks*. The conformal blocks are universal in the sense that they only depend on the conformal weights Δ_{α_i} and Δ_{Q+iP} and the central charge $c_L = 1 + 6Q^2$ of LCFT, i.e. $\mathcal{F}_P(z) = \mathcal{F}(c_L, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4}, \Delta_{Q+iP}, z)$; see Subsection 8.1 for the exact definition.

The second hypothesis on the bootstrap approach i.e. the fact that the OPE may be applied in the two ways explained above goes under the name of *crossing symmetry*. More specifically, it is the conjecture that the following identity holds for real $z \in (0, 1)$

$$(1.17) \quad \begin{aligned} & \int_{\mathbb{R}^+} C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_3, \alpha_4, Q + iP)|z|^{2(\Delta_{Q+iP} - \Delta_{\alpha_1} - \Delta_{\alpha_2})}|\mathcal{F}_P(z)|^2 dP \\ &= \int_{\mathbb{R}^+} C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_3, \alpha_2, Q - iP)C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_4, Q + iP)|1 - z|^{2(\Delta_{Q+iP} - \Delta_{\alpha_3} - \Delta_{\alpha_2})}|\tilde{\mathcal{F}}_P(1 - z)|^2 dP \end{aligned}$$

where $\tilde{\mathcal{F}}_P$ is obtained from \mathcal{F}_P by flipping the parameter α_1 with α_3 . As explained above this identity is a very strong constraint in LCFT.

1.4. Main result on conformal bootstrap. In this paper, we justify rigorously the bootstrap approach to LCFT by constructing the spectral representation of LCFT; as an output, we prove the bootstrap and crossing formulas described in the previous subsection.

To understand our approach to the spectrum of LCFT it is useful to draw the analogy with the harmonic analysis on a Lie group G and in particular the Plancherel identity on $L^2(G)$. For a compact G , $L^2(G)$ is decomposed into a direct sum of irreducible (highest weight) representations of G whereas in the non-compact case also a continuous family of representations (a direct integral) appears. This decomposition is related to the spectral decomposition of a self-adjoint operator (the Laplacian) acting on the Hilbert space $L^2(G)$. In 2d CFT, Osterwalder-Schrader's method of reflection positivity provides a canonical Hilbert space where the symmetry algebra of 2d CFT, the *Virasoro algebra*, acts. The role of the Laplacian is played by a self adjoint operator, a special element in the Virasoro algebra called the Hamiltonian of the CFT. In the case of "compact CFTs" (examples being the minimal models of BPZ) the spectrum of this operator is discrete whereas in case of "non-compact CFTs" the spectrum is continuous. Our proof is based on finding the spectral resolution of this operator using scattering theory and representation theory of the Virasoro algebra, leading to a Plancherel type identity as a rigorous version of the OPE.

The main result of this paper is the following theorem proving that the conformal bootstrap formula (1.16) holds for the probabilistic construction of the 4-point function:

Theorem 1.1. *Let $\gamma \in (0, 2)$ and $\alpha_i < Q$ for all $i \in \llbracket 1, 4 \rrbracket$. Then the following identity holds for $\alpha_1 + \alpha_2 > Q$ and $\alpha_3 + \alpha_4 > Q$*

$$(1.18) \quad \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma,\mu} = \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma,\mu}^{\text{Boot}}.$$

⁸The $\frac{1}{2}$ factor here is a general global constant which can be absorbed in the definition of the probabilistic construction: see footnote associated to (1.6).

The conditions $\alpha_1 + \alpha_2 > Q$ and $\alpha_3 + \alpha_4 > Q$ are essential. Indeed, if $\alpha_1 + \alpha_2 < Q$ the analytic continuation of (1.16) from $\alpha_i \in \mathcal{S}$ requires adding an extra term (cf. the discussion on the so-called discrete terms in [ZaZa96]). The second main input to the bootstrap hypothesis, namely the crossing symmetry conjecture (1.17), follows directly from our work since one has by conformal covariance (1.12) of the probabilistic construction of the correlations

$$\langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma,\mu} = \langle V_{\alpha_3}(0)V_{\alpha_2}(1-z)V_{\alpha_1}(1)V_{\alpha_4}(\infty) \rangle_{\gamma,\mu}$$

whereby we get the following corollary:

Corollary 1.2. *The bootstrap construction of LCFT satisfies crossing symmetry for $\gamma \in (0, 2)$.*

This result seems to be very hard to prove directly, however let us also mention that Teschner has given strong arguments in [Te01] in that direction.

Remark 1.3. *We have stated the bootstrap conjecture as the statements (1.16) and (1.17) since these are the crucial relations following from the OPE axiom (1.15) used by physicists to study CFTs. However, as explained above the OPE axiom (1.15) leads also to a recursive computation of the n -point correlation functions for all n . Likewise, the approach in this paper can be extended to $n > 4$ by a $n - 3$ -fold application of the Plancherel identity. Hence the spectral analysis of the LCFT Hamiltonian is the crucial result of this paper. In order to keep the length of this paper reasonable, we will discuss these generalisations elsewhere, in particular for the case of LCFT on the complex torus where the n -point formulae are mathematically quite appealing [GKRV].*

1.5. Conformal blocks and relations with the AGT conjecture. Let us mention that it is not at all obvious that the bootstrap definition of the four point correlation function, i.e. the right hand side of (1.18), exists for real α_i satisfying the condition $\alpha_i < Q$ for all $i \in \llbracket 1, 4 \rrbracket$ along with $\alpha_1 + \alpha_2 > Q$ and $\alpha_3 + \alpha_4 > Q$. Indeed, first the conformal blocks are defined via a series expansion

$$(1.19) \quad \mathcal{F}_P(z) = \sum_{n=0}^{\infty} \beta_n z^n$$

where the coefficients β_n , which have a strong representation theoretic content, are non explicit : see (8.2) for the exact definition of β_n . Hence, it is not obvious that the series (1.19) converges for $|z| < 1$. Second, it is not clear that the integral in $P \in \mathbb{R}^+$ of expression (1.16) is convergent. As a matter of fact, in the course of the proof of Theorem 1.1, we establish both that the radius of convergence⁹ of (1.19) is 1 for almost all P and that the integral (1.16) makes sense. To the best of our knowledge, the proof of the convergence of the conformal blocks is new and we expect that the result holds for all P , although we do not need such a strong statement for our purpose. Let us mention here the recent work [GRSS20] which establishes a probabilistic formula involving moments of a GMC type variable for the conformal blocks of LCFT on the complex torus thereby proving the existence of the torus blocks for all values of the relevant parameters. Convergence of conformal blocks defining series is also topical in physics, see [PRER12, HoRy13, KQR20].

The AGT correspondence [AGT10] between $4d$ supersymmetric Yang-Mills theory and the bootstrap construction of LCFT conjectures that $\mathcal{F}_P(z)$ coincides with special cases of Nekrasov's partition function [Ne04]. In particular this leads to an explicit formula for β_n in (1.19). However, even admitting this conjecture, it remains difficult to show that the radius of convergence in (1.19) is 1: see for instance [FLM18]. The AGT conjecture has been proved as an identity between formal power series in the case of the torus in [Ne16] following the works [MaOk12, ScVa13] but this does not address the issue of convergence. See also [FL10, AFLT11] for arguments in the physics literature which support the AGT conjecture on the torus or the Riemann sphere.

Acknowledgements. C. Guillarmou acknowledges that this project has received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme, grant agreement No 725967. A. Kupiainen is supported by the Academy of Finland and ERC Advanced Grant 741487. R. Rhodes is partially supported by the Institut Universitaire de France (IUF). The authors

⁹We acknowledge here an argument that was given to us by Slava Rychkov in private communication.

wish to thank Zhen-Qing Chen, Naotaka Kajino for discussions on Dirichlet forms, Ctirad Klimcik and Yi-Zhi Huang for explaining the links with Vertex Operator Algebras, Slava Rychkov for fruitful discussions on the convergence of conformal blocks, Alex Strohmaier, Tanya Christiansen and Jan Dereziński for discussions on the scattering part and Baptiste Cerclé for comments on earlier versions of this manuscript.

2. OUTLINE OF THE PROOF

In this section we give an informal summary of the proof of the bootstrap formula with pointers to precise definitions and statements.

2.1. Reflection positivity. The LCFT expectation (1.6) is an expectation in a positive measure but it has another positivity property called reflection positivity (or Osterwalder-Schrader positivity) that allows us to express the correlation functions of LCFT in terms of a scalar product in the *physical Hilbert space* of LCFT. This task is carried out in Section 3 and outlined now. The construction of the Hilbert space is based on an involution acting on observables F . For this, we consider the reflection at the unit circle $\theta: \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$

$$(2.1) \quad \theta(z) = 1/\bar{z}$$

which maps the unit disk \mathbb{D} to its complement \mathbb{D}^c . We promote it to an operator Θ acting on observables $F \mapsto \Theta F$ by

$$(2.2) \quad (\Theta F)(\phi) := F(\phi \circ \theta - 2Q \ln |\cdot|).$$

This allows us to define a sesquilinear form acting on a set $\mathcal{F}_{\mathbb{D}}$ of observables F that depend only on the restriction $\phi|_{\mathbb{D}}$ of ϕ to the unit disk (i.e. $F(\phi) = F(\phi|_{\mathbb{D}})$) by

$$(2.3) \quad (F, G)_{\mathbb{D}} := \langle \Theta F(\phi) \overline{G(\phi)} \rangle_{\gamma, \mu}, \quad F, G \in \mathcal{F}_{\mathbb{D}}.$$

Note that ΘF is an observable depending on the restriction $\phi|_{\mathbb{D}^c}$ of ϕ to the complementary disc \mathbb{D}^c so that the scalar product of two observables on the disk \mathbb{D} is given by the LCFT expectation of their product when one of them is reflected to \mathbb{D}^c . Reflection positivity is the statement that the sesquilinear form (2.3) is nonnegative:

$$(F, F)_{\mathbb{D}} \geq 0$$

see Proposition 3.3. The canonical Hilbert space $\mathcal{H}_{\mathbb{D}}$ of LCFT is defined as the completion of $\mathcal{F}_{\mathbb{D}}$, quotiented out by the null set $\mathcal{N}_0 = \{F \in \mathcal{F}_{\mathbb{D}} \mid (F, F)_{\mathbb{D}} = 0\}$, with respect to the sesquilinear form (2.3). This space can be realized in more concrete terms as follows.

2.2. Hilbert space of LCFT. The space $\mathcal{H}_{\mathbb{D}}$ can be realized as an L^2 space on a set of fields on the equatorial circle $\mathbb{T} = \partial\mathbb{D}$ using the domain Markov property of the GFF. To do this let $\varphi = X|_{\mathbb{T}}$ be the restriction of the GFF to the unit circle. φ can be realized (see Section 3.1) as a (real valued) random Fourier series

$$(2.4) \quad \varphi(\theta) = \sum_{n \neq 0} \varphi_n e^{in\theta}$$

with $\varphi_n = \frac{1}{2\sqrt{n}}(x_n + iy_n)$ for $n > 0$ where x_n, y_n are i.i.d. unit Gaussians. φ can be understood as a random element in a Sobolev space $W^s(\mathbb{T})$ with $s < 0$ (see (3.3)) and as the coordinate function in the probability space $\Omega_{\mathbb{T}} = (\mathbb{R}^2)^{\mathbb{N}^*}$ equipped with a cylinder set sigma algebra and a Gaussian probability measure \mathbb{P} (see (3.6)). The GFF X (1.3) can now be decomposed (Section 3.1) as an independent sum

$$(2.5) \quad X \stackrel{\text{law}}{=} P\varphi + X_{\mathbb{D}} + X_{\mathbb{D}^c}$$

where $P\varphi$ is the harmonic extension of φ to $\hat{\mathbb{C}}$ and $X_{\mathbb{D}}, X_{\mathbb{D}^c}$ are two independent GFFs on \mathbb{D} and \mathbb{D}^c with Dirichlet boundary conditions. In Proposition 3.3 we show that there is a unitary map $U: \mathcal{H}_{\mathbb{D}} \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ given by

$$(2.6) \quad (UF)(c, \varphi) = e^{-Qc} \mathbb{E}_{\varphi} [F(c + X) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D})}],$$

where \mathbb{E}_φ is expectation over $X_{\mathbb{D}}$ in the decomposition $X|_{\mathbb{D}} = X_{\mathbb{D}} + P\varphi$. Hence $\mathcal{H}_{\mathbb{D}}$ can be identified with $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. The operator U then allows us to write the 4 point function in terms of the scalar product $\langle \cdot | \cdot \rangle_2$ in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, namely, we have for $|z| < 1$

$$(2.7) \quad \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma, \mu} = \langle U(V_{\alpha_1}(0)V_{\alpha_2}(z)) | U(V_{\alpha_3}(0)V_{\alpha_4}(1)) \rangle_2.$$

The bootstrap formula is then obtained by expanding this scalar product along the eigenstates of a self adjoint operator \mathbf{H} on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, the LCFT Hamiltonian to which we now turn.

2.3. Hamiltonian of Liouville theory. For $q \in \mathbb{C}$ with $|q| \leq 1$, the dilation map $z \in \mathbb{C} \rightarrow s_q(z) = qz$ maps the unit disc to itself and it gives rise to a map $S_q : \mathcal{F}_{\mathbb{D}} \rightarrow \mathcal{F}_{\mathbb{D}}$ by

$$(2.8) \quad S_q F(\phi) := F(\phi \circ s_q + Q \ln |q|)$$

for $F \in \mathcal{F}_{\mathbb{D}}$. In Proposition 3.6 we show that these operators satisfy $S_q S_{q'} = S_{qq'}$ and $q \mapsto S_q$ descends to a strongly continuous semigroup on $\mathcal{H}_{\mathbb{D}}$. Taking $q = e^{-t}$ for $t \geq 0$, we get a contraction semigroup on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ via the unitary map U in (2.6), given by the relation

$$(2.9) \quad e^{-t\mathbf{H}} = U S_{e^{-t}} U^{-1}.$$

The generator of the semi-group is a positive self-adjoint operator \mathbf{H} with domain $\mathcal{D}(\mathbf{H}) \subset L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, called the *Hamiltonian of Liouville theory*.

To get a more concrete representation for \mathbf{H} we use a probabilistic Feynman-Kac representation for $e^{-t\mathbf{H}}$. It is based on the well known fact that the GFF $X(e^{-t+i\theta})$ for $t \geq 0$ can be realized as a continuous Markov process

$$t \mapsto X(e^{-t+i\cdot}) = (B_t, \varphi_t(\cdot)) \in W^s(\mathbb{T})$$

with $s < 0$. Here we decomposed $W^s(\mathbb{T}) = \mathbb{R} \oplus W_0^s(\mathbb{T})$ where $W_0^s(\mathbb{T})$ has the $n \neq 0$ Fourier components. In this decomposition B_t is a standard Brownian motion and the process φ_t gives rise to an Ornstein-Uhlenbeck semigroup on $W_0^s(\mathbb{T})$ (i.e. the harmonic components evolve as independent OU processes), whose generator is a positive self-adjoint operator \mathbf{P} . The operator \mathbf{P} is essentially an infinite sum of harmonic oscillators (see (4.4) for the exact definition). This leads to the expression for the Hamiltonian when $\mu = 0$

$$(2.10) \quad \mathbf{H}^0 := \mathbf{H}|_{\mu=0} = -\frac{1}{2}\partial_c^2 + \frac{1}{2}Q^2 + \mathbf{P}$$

defined on an appropriate domain where the first two terms come from the Brownian motion, see Section 4 for further details.

Using the formula (2.6) we then deduce a Feynman-Kac formula for the LCFT semigroup in Proposition 5.1) which then formally gives \mathbf{H} as a "Schrödinger operator"

$$(2.11) \quad \mathbf{H} = \mathbf{H}^0 + \mu e^{\gamma c} V(\varphi)$$

where the potential is formally given by

$$(2.12) \quad V(\varphi) = \int_0^{2\pi} e^{\gamma\varphi(\theta) - \frac{\gamma^2}{2}\mathbb{E}[\varphi(\theta)^2]} d\theta.$$

In Section 5.2, we give the rigorous definition of (2.11) which actually is quite subtle and nonstandard. The operator (2.11) is defined as the Friedrichs extension of an associated quadratic form. For $\gamma \in [0, \sqrt{2})$ V is a well defined random variable defined as the total mass of a GMC measure on \mathbb{T} associated to φ . For $\gamma \in [\sqrt{2}, 2)$ however, this GMC measure vanishes identically and V can no longer be considered as a multiplication operator: it has to be understood as a measure on some subspace of $L^2(\Omega_{\mathbb{T}})$ but it still gives rise to a positive operator.

We mention here that the operator (2.11) in the absence of the c -variable, i.e. the operator $\mathbf{P} + \mu V$, was first studied in [Ho71] and was shown to be essentially self-adjoint on an appropriate domain provided $\gamma \in (0, 1)$, condition under which V is a random variable in $L^2(\Omega_{\mathbb{T}})$.

2.4. Spectral resolution of the Hamiltonian. One of the main mathematical inputs in our proof comes from stationary *scattering theory*, see Section 6. It is based on the observation that the potential $e^{\gamma c}V$ vanishes as $c \rightarrow -\infty$ so that \mathbf{H} -eigenstates should be reconstructed from their asymptotics at $c \rightarrow -\infty$, region over which they should behave like \mathbf{H}^0 -eigenstates.

The spectral analysis of \mathbf{H}^0 is simple. The spectrum $\{0 < \lambda_1 < \dots < \lambda_k < \dots\}$ of the operator \mathbf{P} is given by the natural numbers $\lambda_k = k$ and each eigenvalue has finite multiplicity, see Section 4.1. The corresponding eigenspace $\ker(P - \lambda_k)$ is spanned by a family $(h_{jk})_{j=0, \dots, J(k)}$ of finite products of Hermite polynomials in the Fourier components $(\varphi_n)_{n \in \mathbb{Z}^*}$ in (2.4), providing an orthonormal basis $\{h_{jk}\}_{k \in \mathbb{N}, j \leq J(k)}$ of $L^2(\Omega_{\mathbb{T}})$. The spectrum of \mathbf{H}^0 is then readily seen to be absolutely continuous and given by the half-line $[\frac{Q^2}{2}, +\infty)$ with a complete set of generalized eigenstates

$$(2.13) \quad \Psi_{Q+iP, jk}^0(c, \varphi) = e^{iPc} h_{jk}(\varphi),$$

with $P \in \mathbb{R}$, $k \in \mathbb{N}$, $j \leq J(k)$ and eigenvalue $\frac{1}{2}(Q^2 + P^2) + \lambda_k = 2\Delta_{Q+iP} + \lambda_k$ where Δ_{Q+iP} is the conformal weight (1.9).

In physics, the spectrum of \mathbf{H} was first elaborated by Curtright and Thorn [CuTh] (see also Teschner [Te01] for a nice discussion of the scattering picture). To explain the intuition, let us consider a toy model where only the c -variable enters, namely the operator $\frac{1}{2}(-\partial_c^2 + Q^2) + \mu e^{\gamma c}$ on $L^2(\mathbb{R})$. It is a Schrödinger operator with potential tending to 0 as $c \rightarrow -\infty$ and to $+\infty$ as $c \rightarrow +\infty$. Thus for $c \rightarrow -\infty$ the eigenfunctions should tend to eigenfunctions of the $\mu = 0$ problem i.e. to a linear combination of $e^{\pm iPc}$. Indeed, the operator has a complete set of generalized eigenfunctions f_P , $P \in \mathbb{R}^+$, with asymptotics

$$f_P(c) \sim e^{iPc} + R(P)e^{-iPc} \quad \text{as } c \rightarrow -\infty$$

and $f_P(c) \rightarrow 0$ as $c \rightarrow +\infty$. These eigenfunctions describe scattering of waves from a wall (the exponential potential $e^{\gamma c}$ acts as a wall when $c \rightarrow +\infty$) and $R(P)$ is an explicit coefficient called the reflection coefficient¹⁰.

For the operator \mathbf{H} we expect the same intuition to hold: an eigenfunction $\Psi(c, \varphi)$ of \mathbf{H} should, as $c \rightarrow -\infty$, tend to an eigenfunction of \mathbf{H}^0 of the same eigenvalue. Indeed we prove (see Theorem 6.25) which uses a different labelling) that to each \mathbf{H}^0 eigenvector $\Psi_{Q+iP, jk}^0$ in (2.13) there is a corresponding \mathbf{H} eigenvector $\Psi_{Q+iP, jk}$ with the same eigenvalue

$$(2.14) \quad \mathbf{H}\Psi_{Q+iP, jk} = (2\Delta_{Q+iP} + \lambda_k)\Psi_{Q+iP, jk},$$

and these form a complete set for $P \in \mathbb{R}_+$, $k \in \mathbb{N}$, $j \leq J(k)$. Applying the spectral decomposition of \mathbf{H} to (2.7) using this basis leads to

$$(2.15) \quad \langle V_{\alpha_1}(0)V_{\alpha_2}(z)V_{\alpha_3}(1)V_{\alpha_4}(\infty) \rangle_{\gamma, \mu} = \frac{1}{2\pi} \sum_{k \in \mathbb{N}} \sum_{j=0}^{J(k)} \int_0^\infty \langle U(V_{\alpha_1}(0)V_{\alpha_2}(z)) | \Psi_{Q+iP, jk} \rangle_2 \langle \Psi_{Q+iP, jk} | U(V_{\alpha_4}(0)V_{\alpha_3}(1)) \rangle_2 dP.$$

Proving this spectral resolution for \mathbf{H} is the technical core of the paper and involves a considerable amount of work (the whole Section 6). The main difficulty comes from the fact that the potential V appearing in \mathbf{H} acts on the L^2 -space of an infinite dimensional space $\mathbb{R} \times \Omega_{\mathbb{T}}$ and moreover V is not even a function for $\gamma \in [\sqrt{2}, 2)$ as discussed in the previous subsection. This weak regularity and unboundedness of the potential make the problem quite non-standard.

2.5. Analytic continuation. The spectral resolution (2.15) is not yet the bootstrap formula. Indeed, it holds under quite general assumptions on V . To get the bootstrap formula from (2.15) we need to connect the scalar products in (2.15) to the DOZZ 3-point functions $C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)$ and $C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_3, \alpha_4, Q + iP)$ respectively. The probabilistic 3-point function $C_{\gamma, \mu}(\alpha_1, \alpha_2, \alpha)$ is defined only for α real and satisfying the Seiberg bounds $\alpha_1 < Q$, $\alpha_2 < Q$, $\alpha < Q$, $\alpha_1 + \alpha_2 + \alpha > 2Q$ and $C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)$ is an analytic continuation of this probabilistic expression. Our strategy then is to analytically continue the eigenfunction $\Psi_{Q+iP, jk}$ to real values of $Q + iP$ so that we can use the probabilistic construction of the LCFT and in particular its conformal invariance to derive identities that allow to determine the above scalar products.

¹⁰This coefficient is a simplified version of the (quantum) reflection coefficient which appears in the proof of the DOZZ formula [KRV19, KRV20].

In Section, 6 we show that the eigenfunctions can be analytically continued to $\Psi_{\alpha,jk}$ which are analytic in α in a connected region which contains both the spectrum line $\alpha \in Q + i\mathbb{R}_+$ and a half-line $]-\infty, A_k] \subset \mathbb{R}$ for some $A_k < Q$. This analytic continuation still satisfies the eigenvalue problem (2.14) (with $Q + iP$ replaced by α) but it has exponential growth in the c variable as $c \rightarrow -\infty$ and it does not contribute to the spectral resolution (compare with eigenfunctions e^{ac} of the operator $-\partial_c^2$ which are associated to the L^2 spectrum only for $a \in i\mathbb{R}$).




FIGURE 1. Analytic continuation of eigenstates and probabilistic region.

The crucial point we show is that the analytically continued \mathbf{H} -eigenfunctions $\Psi_{\alpha,jk}$ can be obtained by *intertwining* the \mathbf{H}^0 -eigenfunctions for α in some complex neighborhood of the half-line $]-\infty, A_k]$ that we call *the probabilistic region* (see Figure 1):

$$(2.16) \quad \Psi_{\alpha,jk} = \lim_{t \rightarrow \infty} e^{(2\Delta_\alpha + \lambda_k)t} e^{-t\mathbf{H}} \Psi_{\alpha,jk}^0 \quad (\text{Intertwining})$$

where $\Psi_{\alpha,jk}^0 = e^{(\alpha-Q)c} h_{jk}$ are the analytically continued eigenfunctions of \mathbf{H}^0 (see Proposition 7.2 which has a different labelling). Furthermore, combining (2.16) with the Feynman-Kac formula one finds (Section 7.1) for $\lambda_k = 0$ (then $J(0) = 0$, $h_{00}(\varphi) = 1$ and we denote $\Psi_{\alpha,0}$ for $\Psi_{\alpha,00}$) that

$$(2.17) \quad \Psi_{\alpha,0} = U(V_\alpha(0))$$

where U is the unitary map (2.6). Hence the “non-spectral” eigenfunctions $\Psi_{\alpha,0}$ for $\alpha \in \mathbb{R}$ have a probabilistic interpretation in LCFT whereas the “spectral” eigenfunctions $\Psi_{Q+iP,0}$ do not have one. In the physical literature the correspondence between local fields and states in the Hilbert space is called the state-operator correspondence and in LCFT this correspondence is broken. The spectral states $\Psi_{Q+iP,0}$ do not correspond to local fields and they are called *macroscopic states* whereas the non-normalizable states $\Psi_{\alpha,0}$ correspond to the local field V_α and are called *microscopic states*, see [Se90] for a lucid discussion.

The relation (2.17) leads to

$$(2.18) \quad \langle U(\prod_{i=1}^n V_{\alpha_i}(z_i)) | \Psi_{\alpha,0} \rangle_2 = \langle (\prod_{i=1}^n V_{\alpha_i}(z_i)) V_\alpha(\infty) \rangle_{\gamma,\mu}$$

for $\{\alpha_i\}$, α satisfying the Seiberg bounds (the scalar product in (2.18) still makes sense since the vector on the left has sufficient decay in the c -variable to counter the exponential increase of $\Psi_{\alpha,0}$). Note in particular that (2.18) is defined only if n is large enough since the Seiberg bound requires $\sum_i \alpha_i > 2Q - \alpha > 2Q - A_0$. To understand the case $k > 0$ we need to discuss the conformal symmetry of LCFT.

2.6. Conformal Ward Identities. The symmetries of conformal field theory are encoded in an infinite dimensional Lie algebra, the Virasoro algebra. In the case of GFF this means that the Hilbert space $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ carries a representation of two commuting Virasoro algebras with generators $\{\mathbf{L}_n^0\}_{n \in \mathbb{Z}}$ and $\{\tilde{\mathbf{L}}_n^0\}_{n \in \mathbb{Z}}$ (see Section 4.4). In particular the GFF Hamiltonian is given by $\mathbf{H}^0 = \mathbf{L}_0^0 + \tilde{\mathbf{L}}_0^0$ and the \mathbf{H}^0 eigenstates $\Psi_{Q+iP,jk}^0$ for fixed P can be organized to a highest weight representation of these algebras. In concrete terms the highest weight state $\Psi_{Q+iP,0}^0$ satisfies

$$(2.19) \quad \mathbf{L}_0^0 \Psi_{Q+iP,0}^0 = \tilde{\mathbf{L}}_0^0 \Psi_{Q+iP,0}^0 = \Delta_{Q+iP} \Psi_{Q+iP,0}^0, \quad \mathbf{L}_n^0 \Psi_{Q+iP,0}^0 = \tilde{\mathbf{L}}_n^0 \Psi_{Q+iP,0}^0 = 0, \quad n > 0,$$

and given two non-increasing sequences of positive integers $\nu = (\nu_1, \dots, \nu_k)$ and $\tilde{\nu} = (\tilde{\nu}_1, \dots, \tilde{\nu}_l)$, $k, l \in \mathbb{N}$ and setting $\mathbf{L}_{-\nu}^0 = \mathbf{L}_{-\nu_k}^0 \cdots \mathbf{L}_{-\nu_1}^0$ and $\tilde{\mathbf{L}}_{-\tilde{\nu}}^0 = \tilde{\mathbf{L}}_{-\tilde{\nu}_l}^0 \cdots \tilde{\mathbf{L}}_{-\tilde{\nu}_1}^0$ the states

$$(2.20) \quad \Psi_{Q+iP, \nu, \tilde{\nu}}^0 := \mathbf{L}_{-\nu}^0 \tilde{\mathbf{L}}_{-\tilde{\nu}}^0 \Psi_{Q+iP, 0},$$

are eigenstates of \mathbf{H}^0 of eigenvalue $E = 2\Delta_{Q+iP} + \sum_i \nu_i + \sum_j \tilde{\nu}_j$ and, for fixed E , span that eigenspace. Thus at each eigenspace there is a nonsingular matrix $M(Q+iP)$ relating the vectors $\Psi_{Q+iP, \nu, \tilde{\nu}}^0$ and $\Psi_{Q+iP, jk}^0$ and furthermore we show (Proposition 4.9) that this matrix is analytic in the variable $\alpha = Q+iP$. Setting

$$\Psi_{\alpha, \nu, \tilde{\nu}} := \sum_{k, j} M(\alpha)_{\nu, \tilde{\nu}; jk} \Psi_{\alpha, jk}$$

the vectors $\Psi_{Q+iP, \nu, \tilde{\nu}}$ are a complete set of generalized eigenvectors of \mathbf{H} and they can be used in the identity (2.15) (with appropriate Gram matrices), see Section 8. Furthermore $\Psi_{\alpha, \nu, \tilde{\nu}}$ provides an analytic continuation of $\Psi_{Q+iP, \nu, \tilde{\nu}}$ to the half line $\alpha < Q - A$ for some $A > 0$ and it is intertwined there with the corresponding vector $\Psi_{\alpha, \nu, \tilde{\nu}}^0$ by the relation (2.16).

The bootstrap formula (1.16) is a consequence of the following fundamental identity (see Proposition 7.12)

$$(2.21) \quad \langle \Psi_{Q+iP, \nu, \tilde{\nu}} | U(V_{\alpha_1}(0)V_{\alpha_2}(z)) \rangle_2 = d(\alpha_1, \alpha_2, \nu, \tilde{\nu}) C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q+iP) \bar{z}^{|\nu|} z^{|\tilde{\nu}|} |z|^{2(\Delta_{Q+iP} - \Delta_{\alpha_1} - \Delta_{\alpha_2})}$$

where the function d is an explicit function of the parameters that will contribute to the conformal blocks. To prove (2.21) we consider the scalar product $\langle \Psi_{Q+iP, \nu, \tilde{\nu}} | U(\prod_{i=1}^n V_{\alpha_i}(z_i)) \rangle_2$ where $z_i \in \mathbb{D}$ and $\sum_i \alpha_i > Q+A$ and analytically continue it from $\alpha = Q+iP$ ($P \in \mathbb{R}_+$) to $\alpha \in (2Q - \sum_i \alpha_i, Q - A)$. For such α we prove the Conformal Ward Identity (see Proposition 7.11)

$$(2.22) \quad \langle U(\prod_{i=1}^n V_{\alpha_i}(z_i)) | \Psi_{\alpha, \nu, \tilde{\nu}} \rangle_2 = \mathcal{D}(\alpha, \alpha, \nu, \tilde{\nu}) \langle \prod_{i=1}^n V_{\alpha_i}(z_i) V_{\alpha}(\infty) \rangle_{\gamma, \mu}$$

where $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\mathcal{D}(\alpha, \alpha, \nu, \tilde{\nu})$ is an explicit partial differential operator in the variables z_i . Using (2.18), continuing back to $\alpha = Q+iP$ and taking $\alpha_i \rightarrow 0$ for $i > 2$ we then deduce (the complex conjugate of) (2.21).

The proof of (2.22) occupies the whole Section 7. It is based on a representation of the states $\Psi_{\alpha, \nu, \tilde{\nu}}^0$ in terms of $V_{\alpha}(0)$ and the Stress-Energy-Tensor (SET) in Proposition 7.23. Let us briefly explain this for $\nu = n, \tilde{\nu} = \emptyset$ ie for the state $\mathbf{L}_{-n}^0 \Psi_{\alpha, 0}^0$. We have $\Psi_{\alpha, 0}^0 = U_0 V_{\alpha}(0)$ where U_0 is the map 2.6 for $\mu = 0$. The SET is given in the GFF theory by the field

$$(2.23) \quad T(z) := Q \partial_z^2 X(z) - (\partial_z X(z))^2 + \mathbb{E}[(\partial_z X(z))^2],$$

defined through regularization and limit. Then we prove

$$\mathbf{L}_{-n}^0 \Psi_{\alpha, 0}^0 = \frac{1}{2\pi i} \oint z^{1-n} U_0(T(z)V_{\alpha}(0)) dz$$

where the integration contour circles the origin in \mathbb{D} . Plugging this identity to the intertwining relation (2.16) and using the Feynman-Kac formula one ends up with a contour integral of the SET insertion in LCFT correlation function. This is analyzed by Gaussian integration by parts and results in the Ward identity.

2.7. Organization of the paper. The paper is organized as follows. In Section 3, we will introduce the relevant material on the Gaussian Free Field and explain the construction of the Hilbert space as well as the quantization of dilations; the Liouville Hamiltonian is then defined as the generator of dilations. This section uses the concept of reflection positivity. In Section 4, we study the dynamics induced by dilations in the GFF theory (i.e. $\mu = 0$) and recall the basics of representation theory related to the GFF. In Section 5, we study in more details the Liouville Hamiltonian: we establish the Feynman-Kac formula for the associated semigroup and identify its quadratic form, which allows us to use scattering theory to diagonalize the Liouville Hamiltonian in Section 6. In Sections 7 and 9, we will prove the Conformal Ward identities for the correlations of LCFT: in a way, this can be seen as an identification of the eigenstates of the Liouville Hamiltonian. In Section 8, we will prove the main result of the paper Theorem 1.1 using the material proved in the other sections. Finally, in the appendix, we will recall the DOZZ formula and gather auxiliary results (analyticity of vertex operators).

2.8. Notations and conventions. We gather here the frequently used notations:

$\langle \cdot \rangle_{\gamma, \mu}$ denotes the LCFT expectation (1.6).

X, ϕ, φ denote respectively the GFF (1.3), the Liouville field (1.5) and the GFF on \mathbb{T} (2.4).

$(\Omega_{\mathbb{T}}, \mathbb{P}_{\mathbb{T}})$ denotes the probability space (3.5) with measure (3.6).

$L^p(\Omega_{\mathbb{T}})$ for $p \geq 1$: complex valued functions $\psi(\varphi)$ with norm $\|\cdot\|_{L^p(\Omega_{\mathbb{T}})}$.

$L^p(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $p \geq 1$: complex valued functions $\psi(c, \varphi)$ with norm $\|\cdot\|_p$.

$\langle \cdot | \cdot \rangle_{L^2(\Omega_{\mathbb{T}})}$: scalar product in $L^2(\Omega_{\mathbb{T}})$.

$\langle \cdot | \cdot \rangle_2$: scalar product in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$.

$e^{\alpha\rho(c)}L^p(\mathbb{R} \times \Omega_{\mathbb{T}})$. Weighted L^p -space equipped with the norm $\|f\|_{e^{\alpha\rho(c)}L^p} := \|e^{-\alpha\rho(c)}f\|_p$.

$C^\infty(D)$ denotes the set of smooth functions on the domain D .

$C_c^\infty(D)$ denotes the set of smooth functions with compact support in D .

$(\cdot, \cdot)_{\mathbb{D}}$ denotes the scalar product (3.16) associated to reflexion positivity

$\mathcal{H}_{\mathbb{D}}$: Hilbert space associated to $(\cdot, \cdot)_{\mathbb{D}}$.

$\langle f, g \rangle_{\mathbb{T}} := \int_0^{2\pi} f(\theta)g(\theta)d\theta$ denotes the scalar product on $L^2(\mathbb{T})$.

$\langle f, g \rangle_{\mathbb{D}} := \int_{\mathbb{D}} f(x)g(x)dx$ denotes the scalar product in $L^2(\mathbb{D})$.

All sesquilinear forms are linear in their first argument, antilinear in their second one.

For integral operators on some measure space (M, μ) we use the notation $(Gf)(x) = \int G(x, y)f(y)\mu(dy)$.

GFF quantities as opposed to corresponding LCFT ones will carry a subscript or superscript 0: for example the Hamiltonians are \mathbf{H}^0 (GFF) and \mathbf{H} (LCFT).

3. REFLECTION POSITIVITY

In this section we prove the reflection positivity of the LCFT and explain the isometry U (2.6) mapping the LCFT observables to states in the Hilbert space $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ as well as the semigroup in (2.9). We start by a discussion of the various GFF's.

3.1. Gaussian Free Fields. We will now define the fields entering the decomposition (2.5).

GFF on the unit circle \mathbb{T} . Given two independent sequences of i.i.d. standard Gaussians $(x_n)_{n \geq 1}$ and $(y_n)_{n \geq 1}$, the GFF on the unit circle is the random Fourier series

$$(3.1) \quad \varphi(\theta) = \sum_{n \neq 0} \varphi_n e^{in\theta}$$

where for $n > 0$

$$(3.2) \quad \varphi_n := \frac{1}{2\sqrt{n}}(x_n + iy_n), \quad \varphi_{-n} := \bar{\varphi}_n.$$

Let $W^s(\mathbb{T}) \subset \mathbb{C}^{\mathbb{Z}}$ be the set of sequences s.t.

$$(3.3) \quad \|\varphi\|_{W^s(\mathbb{T})}^2 := \sum_{n \in \mathbb{Z}} |\varphi_n|^2 (|n| + 1)^{2s} < \infty.$$

One can easily check that $\mathbb{E}[\|\varphi\|_{W^s(\mathbb{T})}^2] < \infty$ for any $s < 0$ so that the series (3.1) defines a random element in $W^s(\mathbb{T})$. Moreover, by a standard computation, one can check that it is a centered Gaussian field with covariance kernel given by

$$(3.4) \quad \mathbb{E}[\varphi(\theta)\varphi(\theta')] = \ln \frac{1}{|e^{i\theta} - e^{i\theta'}|}.$$

We will view φ as the coordinate function of the probability space

$$(3.5) \quad \Omega_{\mathbb{T}} = (\mathbb{R}^2)^{\mathbb{N}^*}$$

which is equipped with the cylinder sigma-algebra $\Sigma_{\mathbb{T}} = \mathcal{B}^{\otimes \mathbb{N}^*}$, where \mathcal{B} stands for the Borel sigma-algebra on \mathbb{R}^2 and the product measure

$$(3.6) \quad \mathbb{P}_{\mathbb{T}} := \bigotimes_{n \geq 1} \frac{1}{2\pi} e^{-\frac{1}{2}(x_n^2 + y_n^2)} dx_n dy_n.$$

Here $\mathbb{P}_{\mathbb{T}}$ is supported on $W^s(\mathbb{T})$ for any $s < 0$ in the sense that $\mathbb{P}_{\mathbb{T}}(\varphi \in W^s(\mathbb{T})) = 1$.

Harmonic extension of the GFF on \mathbb{T} . The next ingredient we need for the decomposition of the GFF (2.5) is the harmonic extension $P\varphi$ of the circle GFF defined on $z \in \mathbb{D}$ by

$$(3.7) \quad (P\varphi)(z) = \sum_{n \geq 1} (\varphi_n z^n + \bar{\varphi}_n \bar{z}^n)$$

and on $z \in \mathbb{D}^c$ by $(P\varphi)(1/\bar{z})$ so that we have

$$P\varphi = (P\varphi) \circ \theta$$

where θ is the reflection in the unit circle (2.1). $P\varphi$ is a.s. a smooth field in the complement of the unit circle with covariance kernel given for $z, u \in \mathbb{D}$

$$\mathbb{E}[(P\varphi)(z)(P\varphi)(u)] = \frac{1}{2} \sum_{n>0} \frac{1}{n} ((z\bar{u})^n + (\bar{z}u)^n) = -\ln|1 - z\bar{u}|$$

and for $z \in \mathbb{D}, u \in \mathbb{D}^c$

$$\mathbb{E}[(P\varphi)(z)(P\varphi)(u)] = -\ln|1 - z/u|.$$

Dirichlet GFF on the unit disk. The Dirichlet GFF $X_{\mathbb{D}}$ on the unit disk \mathbb{D} is the centered Gaussian distribution (in the sense of Schwartz) with covariance kernel $G_{\mathbb{D}}$ given by

$$(3.8) \quad G_{\mathbb{D}}(x, x') := \mathbb{E}[X_{\mathbb{D}}(x)X_{\mathbb{D}}(x')] = \ln \frac{|1 - x\bar{x}'|}{|x - x'|}.$$

Here, $G_{\mathbb{D}}$ is the Green function of the negative of the Laplacian $\Delta_{\mathbb{D}}$ with Dirichlet boundary condition on $\mathbb{T} = \partial\mathbb{D}$ and $X_{\mathbb{D}}$ can be realized as an expansion in eigenfunctions of $\Delta_{\mathbb{D}}$ with Gaussian coefficients. However, it will be convenient for us to use another realization based on the following observation. Let, for $n \in \mathbb{Z}$

$$X_n(t) = \int_0^{2\pi} e^{-in\theta} X_{\mathbb{D}}(e^{-t+i\theta}) \frac{d\theta}{2\pi}.$$

Then we deduce from (3.8)

$$(3.9) \quad \mathbb{E}[X_n(t)X_m(t')] = \begin{cases} \frac{1}{2|n|} \delta_{n,-m} (e^{-|t-t'||n|} - e^{-(t+t')|n|}) & n \neq 0 \\ t \wedge t' & n = m = 0 \end{cases}.$$

Thus $\{X_n\}_{n \geq 0}$ are independent Gaussian processes with $X_{-n} = \bar{X}_n$ and X_0 is Brownian motion. We can and will realize them in a probability space $(\Omega_{\mathbb{D}}, \Sigma_{\mathbb{D}}, \mathbb{P}_{\mathbb{D}})$ s.t. $X_n(t)$ have continuous sample paths. Then for fixed t

$$(3.10) \quad X_{\mathbb{D}}(e^{-t+i\theta}) = \sum_{n \in \mathbb{Z}} X_n(t) e^{in\theta}.$$

takes values in $W^s(\mathbb{T})$ for $s < 0$ a.s. (defined in (3.3)) and we can take the map $t \in \mathbb{R}^+ \mapsto X_{\mathbb{D}}(e^{-t+i\cdot}) \in W^s(\mathbb{T})$ to be continuous a.s. in $\Omega_{\mathbb{D}}$. Decompose $W^s(\mathbb{T}) = \mathbb{R} \oplus W_0^s(\mathbb{T})$ where $f \in W_0^s(\mathbb{T})$ has zero average: $\int f(\theta) d\theta = 0$. Then

$$X_{\mathbb{D}}(e^{-t+i\cdot}) = (B_t, Y_t)$$

where B_t is Brownian motion and $Y_t(\theta) = \sum_{n \neq 0} Y_n(t) e^{in\theta}$ is a continuous process in $W_0^s(\mathbb{T})$, independent of B_t . The Dirichlet GFF $X_{\mathbb{D}^c}$ on the complement \mathbb{D}^c of \mathbb{D} can be constructed in the same way (in a probability space $(\Omega_{\mathbb{D}^c}, \Sigma_{\mathbb{D}^c}, \mathbb{P}_{\mathbb{D}^c})$) and we have the relation in law

$$(3.11) \quad X_{\mathbb{D}^c} \stackrel{\text{law}}{=} X_{\mathbb{D}} \circ \theta$$

or in other words $X_{\mathbb{D}^c}(e^{t+i\cdot}) \stackrel{\text{law}}{=} X_{\mathbb{D}}(e^{-t+i\cdot})$, $t \geq 0$.

GFF on the Riemann sphere. One can check that adding the covariances in the previous subsections we get that the field X defined by (2.5) has the covariance

$$(3.12) \quad \mathbb{E}[X(x)X(y)] = \ln \frac{1}{|x-y|} + \ln |x|_+ + \ln |y|_+$$

which coincides with (1.3). In the sequel, we suppose that the GFF on the Riemann sphere X is defined on a probability space $(\Omega, \Sigma, \mathbb{P})$ (with expectation $\mathbb{E}[\cdot]$) where $\Omega = \Omega_{\mathbb{T}} \times \Omega_{\mathbb{D}} \times \Omega_{\mathbb{D}^c}$, $\Sigma = \Sigma_{\mathbb{T}} \otimes \Sigma_{\mathbb{D}} \otimes \Sigma_{\mathbb{D}^c}$ and \mathbb{P} is a product measure $\mathbb{P} = \mathbb{P}_{\mathbb{T}} \otimes \mathbb{P}_{\mathbb{D}} \otimes \mathbb{P}_{\mathbb{D}^c}$. At the level of random variables, the GFF decomposes as the sum of three independent variables

$$(3.13) \quad X = P\varphi + X_{\mathbb{D}} + X_{\mathbb{D}^c}$$

where $P\varphi$ is the harmonic extension of the GFF restricted to the circle $\varphi = X|_{\mathbb{T}}$ defined on $(\Omega_{\mathbb{T}}, \Sigma_{\mathbb{T}}, \mathbb{P}_{\mathbb{T}})$ and $X_{\mathbb{D}}, X_{\mathbb{D}^c}$ are two independent GFFs on \mathbb{D} and \mathbb{D}^c with Dirichlet boundary conditions defined respectively on the probability spaces $(\Omega_{\mathbb{D}}, \Sigma_{\mathbb{D}}, \mathbb{P}_{\mathbb{D}})$ and $(\Omega_{\mathbb{D}^c}, \Sigma_{\mathbb{D}^c}, \mathbb{P}_{\mathbb{D}^c})$ ¹¹. We will write $\mathbb{E}_{\varphi}[\cdot]$ for conditional expectation with respect to the GFF on the circle φ (instead of $:= \mathbb{E}[\cdot | \Sigma_{\mathbb{T}}]$). We will view X in two ways in what follows: as a process $X_t \in W^s(\mathbb{T})$ ($s < 0$)

$$(3.14) \quad X_t(\theta) = X_{\mathbb{D}}(e^{-t+i\theta})1_{t>0} + X_{\mathbb{D}^c}(e^{-t+i\theta})1_{t<0} + (P\varphi)(e^{-|t|+i\theta}).$$

and as a random element in $\mathcal{D}'(\hat{\mathbb{C}})$. In the sequel, we will denote for $t \geq 0$:

$$(3.15) \quad \varphi_t(\theta) := P\varphi(e^{-t+i\theta}) + Y_t(\theta).$$

3.2. Reflection positivity. Let $\mathbb{D} = \{|z| < 1\}$ be the unit disk. Recall the definition of the Liouville field (1.5) which is given on \mathbb{D} by $\phi(z) = c + X(z) = c + X_{\mathbb{D}}(z) + P\varphi(z)$. Let $\mathcal{A}_{\mathbb{D}}$ be the sigma-algebra on $\mathbb{R} \times \Omega$ generated by the maps $\phi \mapsto \langle \phi, g \rangle_{\mathbb{D}}$ for $g \in C_0^{\infty}(\mathbb{D})$ and we recall the notation $\langle \phi, g \rangle_{\mathbb{D}} = \int_{\mathbb{D}} g(z)\phi(z)dz$. Let $\mathcal{F}_{\mathbb{D}}$ be the set of $\mathcal{A}_{\mathbb{D}}$ -measurable functions with values in \mathbb{R} .

For $F, G \in \mathcal{F}_{\mathbb{D}}$ such that the following quantities make sense (see below), we define (recall (2.2))

$$(3.16) \quad (F, G)_{\mathbb{D}} := \langle \Theta F(\phi) \overline{G(\phi)} \rangle_{\gamma, \mu}.$$

Reflection positivity is the statement that this bilinear form is non-negative, namely $(F, F)_{\mathbb{D}} \geq 0$. In what follows, we will study this statement separately for the GFF theory ($\mu = 0$) and for LCFT.

Reflection positivity of the GFF. Here we assume $\mu = 0$. Let $F, G \in \mathcal{F}_{\mathbb{D}}$ be nonnegative. The sesquilinear form (3.16) becomes at $\mu = 0$

$$(3.17) \quad (F, G)_{\mathbb{D}, 0} = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}[(\Theta F)(\phi) \overline{G(\phi)}] dc = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}[F(c + X^{(2)}) \overline{G(c + X^{(1)})}] dc$$

where we denoted $X^{(i)} = X_{\mathbb{D}}^{(i)} + P\varphi$ with $X_{\mathbb{D}}^{(1)} = X_{\mathbb{D}}$ and $X_{\mathbb{D}}^{(2)} = X_{\mathbb{D}^c} \circ \theta$ which are two independent GFFs in the unit disk. Hence by independence of $X_{\mathbb{D}}^{(i)}$

$$(3.18) \quad \begin{aligned} (F, G)_{\mathbb{D}, 0} &= \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}[\mathbb{E}_{\varphi}[F(c + X_{\mathbb{D}}^{(2)} + P\varphi)] \overline{\mathbb{E}_{\varphi}[G(c + X_{\mathbb{D}}^{(1)} + P\varphi)}] dc \\ &= \langle U_0 F | U_0 G \rangle_2 \end{aligned}$$

where the map U_0 is defined by

$$(3.19) \quad (U_0 F)(c, \varphi) = e^{-Qc} \mathbb{E}_{\varphi}[F(c + X_{\mathbb{D}} + P\varphi)]$$

and we recall \mathbb{E}_{φ} denotes expectation over $X_{\mathbb{D}}$. Such a map is well defined on nonnegative $F \in \mathcal{F}_{\mathbb{D}}$ and extended to $\mathcal{F}_{\mathbb{D}}^{0, \infty}$, which is defined as the space of $F \in \mathcal{F}_{\mathbb{D}}$ such that $U_0|F| < \infty$ $dc \otimes \mathbb{P}_{\mathbb{T}}$ -almost everywhere. Let $\mathcal{F}_{\mathbb{D}}^{0, 2} = \{F \in \mathcal{F}_{\mathbb{D}}^{0, \infty} \mid \|U_0 F\|_2 < \infty\}$.

¹¹With a slight abuse of notations, we will assume that these spaces are canonically embedded in the product space (Ω, Σ) and we will identify them with the respective images of the respective embeddings.

Proposition 3.1. *The sesquilinear form (3.17) extends to $\mathcal{F}_{\mathbb{D}}^{0,2}$. This extension is non negative*

$$(3.20) \quad \forall F \in \mathcal{F}_{\mathbb{D}}^{0,2}, \quad (F, F)_{\mathbb{D},0} \geq 0.$$

Let $\overline{\mathcal{F}_{\mathbb{D}}^{0,2}/\mathcal{N}_0^0}$ be the Hilbert space completion of the pre-Hilbert space $\mathcal{F}_{\mathbb{D}}^{0,2}/\mathcal{N}_0^0$ with $\mathcal{N}_0^0 := \{F \in \mathcal{F}_{\mathbb{D}}^{0,2} \mid (F, F)_{\mathbb{D},0} = 0\}$. The map U_0 in (3.19) descends to a unitary map $U_0 : \overline{\mathcal{F}_{\mathbb{D}}^{0,2}/\mathcal{N}_0^0} \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$.

Proof. By (3.18) U_0 descends to an isometry on $\mathcal{F}_{\mathbb{D}}^{0,2}/\mathcal{N}_0^0$ so we need to show it is onto. We take F of the form

$$(3.21) \quad F(c + X) = \rho(\langle c + X, g \rangle_{\mathbb{D}}) e^{\langle c + X, f \rangle_{\mathbb{D}} - \frac{1}{2} \langle f, G_{\mathbb{D}} f \rangle_{\mathbb{D}}}$$

with $\rho \in C_0^\infty(\mathbb{R})$ and $g, f \in C_0^\infty(\mathbb{D})$ with the further conditions that g is rotation invariant i.e. $g(re^{i\theta}) = g(r)$, and that $\int_0^{2\pi} f(re^{i\theta}) d\theta = 0$ for all $r \in [0, 1]$. Then $\langle c, f \rangle_{\mathbb{D}} = 0$ and $\langle P\varphi, g \rangle_{\mathbb{D}} = 0$ and we get

$$(3.22) \quad \begin{aligned} (U_0 F)(c, \varphi) &= e^{-Qc} e^{\langle P\varphi, f \rangle_{\mathbb{D}}} \mathbb{E}[\rho(c + \langle X_{\mathbb{D}}, g \rangle_{\mathbb{D}}) e^{\langle X_{\mathbb{D}}, f \rangle_{\mathbb{D}} - \frac{1}{2} \langle f, G_{\mathbb{D}} f \rangle_{\mathbb{D}}}] \\ &= e^{-Qc} e^{\langle P\varphi, f \rangle_{\mathbb{D}}} \mathbb{E}[\rho(c + \langle X_{\mathbb{D}}, g \rangle_{\mathbb{D}})] \end{aligned}$$

where we observed that $\langle X_{\mathbb{D}}, g \rangle_{\mathbb{D}}$ and $\langle X_{\mathbb{D}}, f \rangle_{\mathbb{D}}$ are independent as their covariance vanishes. Indeed, by rotation invariance of g , the function $\mathcal{O}(r, \theta) := \int_{\mathbb{D}} g(x) G_{\mathbb{D}}(x, re^{i\theta}) dx$ does not depend on θ hence

$$\begin{aligned} \mathbb{E}[\langle X_{\mathbb{D}}, g \rangle_{\mathbb{D}} \langle X_{\mathbb{D}}, f \rangle_{\mathbb{D}}] &= \int_{\mathbb{D}} \int_{\mathbb{D}} G_{\mathbb{D}}(x, y) g(x) f(y) dy \\ &= \int_0^1 r \int_0^{2\pi} f(re^{i\theta}) \mathcal{O}(r, \theta) d\theta dr \\ &= \int_0^1 r \mathcal{O}(r, 0) \int_0^{2\pi} f(re^{i\theta}) d\theta dr = 0. \end{aligned}$$

Let $h \in C^\infty(\mathbb{T})$, $f_\epsilon \in C_0^\infty(\mathbb{D})$ and g_ϵ be given by $g_\epsilon(re^{i\theta}) = \epsilon^{-1} \eta(\frac{1-r}{\epsilon})$, $f_\epsilon = h g_\epsilon$ where η is a smooth bump with support on $[1, 2]$ and total mass one. Then $\lim_{\epsilon \rightarrow 0} \langle P\varphi, f_\epsilon \rangle_{\mathbb{D}} = \langle \varphi, h \rangle_{\mathbb{T}}$ and $\lim_{\epsilon \rightarrow 0} \mathbb{E}(\langle X_{\mathbb{D}}, g_\epsilon \rangle_{\mathbb{D}}^2) = 0$ so that

$$\lim_{\epsilon \rightarrow 0} (U_0 F_\epsilon)(c, \varphi) = e^{-Qc} \rho(c) e^{\langle \varphi, h \rangle_{\mathbb{T}}}$$

where the convergence is in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. Thus the functions $e^{-Qc} \rho(c) e^{\langle \varphi, h \rangle_{\mathbb{T}}}$ are in the image of U_0 for all $\rho \in C_0^\infty(\mathbb{R})$ and $h \in C^\infty(\mathbb{T})$. Since the linear span of these is dense in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ the claim follows. \square

Remark 3.2. *Note that this argument shows that U_0 extends from $\mathcal{F}_{\mathbb{D}}^{0,2}$ to functionals of form $F(c + X_{|\mathbb{T}})$ and then*

$$(U_0 F)(c, \varphi) = e^{-Qc} F(c + \varphi).$$

Reflection positivity of LCFT. Next we want to show reflection positivity for the LCFT expectation (1.6) with $\mu > 0$. The GMC measure M_γ defined in (1.4) can also be constructed as the martingale limit

$$(3.23) \quad M_\gamma(dx) = \lim_{N \rightarrow \infty} e^{\gamma X_N(x) - \frac{\gamma^2}{2} \mathbb{E}[X_N(x)^2]} |x|_+^{-4} dx.$$

where in X_N we cut off the series (3.10) and (3.7) defining $X_{\mathbb{D}}^{(i)}$ and $P\varphi$ respectively to finite number of terms $|n| \leq N$. We claim that

$$M_\gamma(\hat{\mathbb{C}}) = M_\gamma^{(1)}(\mathbb{D}) + M_\gamma^{(2)}(\mathbb{D})$$

where $M_\gamma^{(i)}$ are the GMC measures of the fields $X^{(i)} = X_{\mathbb{D}}^{(i)} + P\varphi$, $i = 1, 2$. Indeed, we take the limit $N \rightarrow \infty$ in

$$\int_{\mathbb{D}^c} e^{\gamma X_N(x) - \frac{\gamma^2}{2} \mathbb{E}[X_N(x)^2]} |x|^{-4} dx = \int_{\mathbb{D}^c} e^{\gamma X_N^{(2)}(\frac{1}{\mathbb{R}}) - \frac{\gamma^2}{2} \mathbb{E}[X_N^{(2)}(\frac{1}{\mathbb{R}})^2]} |x|^{-4} dx = \int_{\mathbb{D}^c} e^{\gamma X_N^{(2)}(x) - \frac{\gamma^2}{2} \mathbb{E}[X_N^{(2)}(x)^2]} dx.$$

Thus, for nonnegative $F, G \in \mathcal{F}_{\mathbb{D}}$

$$(F, G)_{\mathbb{D}} = \langle \Theta F \overline{G} \rangle_{\gamma, \mu} = \langle U_0(IF) \mid U_0(IG) \rangle_2$$

where

$$I = e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D})}.$$

Let $\mathcal{F}_\mathbb{D}^\infty$ be the space of $F \in \mathcal{F}_\mathbb{D}$ such that $U_0(|F|I) < \infty$ $dc \otimes \mathbb{P}_\mathbb{T}$ -almost everywhere. Let $\mathcal{F}_\mathbb{D}^2 = \{F \in \mathcal{F}_\mathbb{D}^\infty \mid \|U_0(FI)\|_2 < \infty\}$. From the above considerations, we arrive at:

Proposition 3.3. *The sesquilinear form (3.16) extends to $\mathcal{F}_\mathbb{D}^2$, is nonnegative and given by*

$$(3.24) \quad (F, G)_\mathbb{D} = \langle UF \mid UG \rangle_2$$

for all $F, G \in \mathcal{F}_\mathbb{D}^2$ where

$$(3.25) \quad (UF)(c, \varphi) = (U_0(FI))(c, \varphi) = e^{-Qc} \mathbb{E}_\varphi[F(c+X)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D})}],$$

$X = X_\mathbb{D} + P\varphi$ and M_γ is its GMC measure. Define $\mathcal{N}_0 := \{F \in \mathcal{F}_\mathbb{D}^2 \mid (F, F)_\mathbb{D} = 0\}$. Then U descends to a unitary map

$$U : \mathcal{H}_\mathbb{D} \rightarrow L^2(\mathbb{R} \times \Omega_\mathbb{T})$$

with $\mathcal{H}_\mathbb{D} := \overline{\mathcal{F}_\mathbb{D}^2 / \mathcal{N}_0}$ (the completion with respect to $(\cdot, \cdot)_\mathbb{D}$).

Proof. We need to show U is onto. This follows from $U(I^{-1}F) = U_0F$ and the fact that U_0 is onto. \square

Remark 3.4. *From Remark 3.2 we conclude that U extends from $\mathcal{F}_\mathbb{D}$ to functionals $F(c + X_\mathbb{T})$ for which*

$$(UF)(c, \varphi) = F(c + \varphi) \times (U1)(c + \varphi)$$

or, in other words, for $f \in L^2(\mathbb{R} \times \Omega_\mathbb{T})$

$$(3.26) \quad U^{-1}f = (U1)^{-1}f.$$

3.3. Dilation Semigroup. Recall the action of the dilation map (2.8) on $\mathcal{F}_\mathbb{D}$. The reason for the $Q \ln |q|$ -factor is the Möbius invariance property of LCFT [DKRV16]

Proposition 3.5. *Let $\psi : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ be a Möbius map and let F be a functional on $\mathcal{D}'(\hat{\mathbb{C}})$ so that $\langle |F(\phi)| \rangle_{\gamma, \mu} < \infty$. Then*

$$(3.27) \quad \langle F(\phi \circ \psi + Q \ln |\psi'|) \rangle_{\gamma, \mu} = \langle F(\phi) \rangle_{\gamma, \mu}.$$

We have then:

Proposition 3.6. *The map S_q descends to a contraction $S_q : \mathcal{H}_\mathbb{D} \rightarrow \mathcal{H}_\mathbb{D}$:*

$$(3.28) \quad \forall F \in \mathcal{H}_\mathbb{D}, \quad (S_q F, S_q F)_\mathbb{D} \leq (F, F)_\mathbb{D}.$$

The adjoint of S_q is $S_q^* = S_{\bar{q}}$ i.e. for all $F, G \in \mathcal{H}_\mathbb{D}$

$$(3.29) \quad (S_q F, G)_\mathbb{D} = (F, S_{\bar{q}} G)_\mathbb{D}.$$

Finally the map $q \in \mathbb{D} \mapsto S_q$ is strongly continuous and satisfies the group property

$$(3.30) \quad S_q S_{q'} = S_{qq'}$$

so that $q \in \mathbb{D} \mapsto S_q$ is a strongly continuous contraction semigroup.

Proof. Let us start with (3.29). It suffices to consider $F, G \in \mathcal{F}_\mathbb{D}^2$ real. By definition

$$(S_q F, G)_\mathbb{D} = \langle F(\phi \circ \theta \circ s_q + Q \ln |q| - 2Q\ell \circ s_q) G(\phi) \rangle_{\gamma, \mu} := \langle \tilde{F}(\phi) G(\phi) \rangle_{\gamma, \mu}$$

where $\ell(z) := \ln |z|$. Applying Proposition 3.5 with $\psi = s_{\bar{q}}$ we get

$$\langle \tilde{F}(\phi) G(\phi) \rangle_{\gamma, \mu} = \langle \tilde{F}(\phi \circ s_{\bar{q}} + Q \ln |q|) G(\phi \circ s_{\bar{q}} + Q \ln |q|) \rangle_{\gamma, \mu}$$

But

$$\tilde{F}(\phi \circ s_{\bar{q}} + Q \ln |q|) = F(\phi \circ s_{\bar{q}} \circ \theta \circ s_q + 2Q \ln |q| - 2Q\ell \circ s_q) = F(\phi \circ \theta - 2Q\ell)$$

and therefore $\langle \tilde{F}(\phi) G(\phi) \rangle_{\gamma, \mu} = (F, S_{\bar{q}} G)_\mathbb{D}$ as claimed.

The group property (3.30) is obvious.

To prove the contraction, denote for $F \in \mathcal{F}_{\mathbb{D}}$, the seminorm $\|F\|_{\mathbb{D}} := (F, F)_{\mathbb{D}}^{\frac{1}{2}}$. Then we have

$$\|S_q F\|_{\mathbb{D}} = (S_q F, S_q F)_{\mathbb{D}}^{\frac{1}{2}} = (F, S_{|q|^2} F)_{\mathbb{D}}^{\frac{1}{2}} \leq \|F\|_{\mathbb{D}}^{\frac{1}{2}} \|S_{|q|^2} F\|_{\mathbb{D}}^{\frac{1}{2}}.$$

Iterating this inequality we obtain

$$\|S_q F\|_{\mathbb{D}} \leq \|F\|_{\mathbb{D}}^{1-2^{-k}} \|S_{|q|^{2k}} F\|_{\mathbb{D}}^{2^{-k}}.$$

Recall that

$$(G, G)_{\mathbb{D}} = \langle U_0(IG) | U_0(IG) \rangle_2 = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}[\mathbb{E}_{\varphi}[IG]^2] dc$$

and then by Cauchy-Schwartz applied to $\mathbb{E}_{\varphi}[\cdot]$

$$\mathbb{E}[\mathbb{E}_{\varphi}[IG]^2] = \mathbb{E}[\mathbb{E}_{\varphi}[I^{\frac{1}{2}} G I^{\frac{1}{2}}]^2] \leq \mathbb{E}[\mathbb{E}_{\varphi}[IG^2] \mathbb{E}_{\varphi}[I]]$$

so that

$$(G, G)_{\mathbb{D}} \leq \langle U_0(IG^2) | U_0 I \rangle_2 = \langle UG^2 | U1 \rangle_2 = \langle G^2 \rangle_{\gamma, \mu}.$$

Hence

$$\|S_q F\|_{\mathbb{D}} \leq \|F\|_{\mathbb{D}}^{1-2^{-k}} \langle (S_{|q|^{2k}} F)^2 \rangle_{\gamma, \mu}^{2^{-k-1}} = \|F\|_{\mathbb{D}}^{1-2^{-k}} \langle F^2 \rangle_{\gamma, \mu}^{2^{-k-1}}$$

where we used again the Möbius invariance of $\langle \cdot \rangle_{\gamma, \mu}$. Taking $k \rightarrow \infty$ we conclude $\|S_q F\|_{\mathbb{D}} \leq \|F\|_{\mathbb{D}}$ for $F \in \mathcal{F}_{\mathbb{D}}$ which satisfy $\langle F^2 \rangle_{\gamma, \mu} < \infty$. Such F form a dense set in $\mathcal{F}_{\mathbb{D}}$. Indeed, let $F \in \mathcal{F}_{\mathbb{D}}$ with $\|F\|_{\mathbb{D}} < \infty$ and let $F_R = F 1_{|F| < R}$. Then $\langle F_R^2 \rangle_{\gamma, \mu} < \infty$ and

$$\|F - F_R\|_{\mathbb{D}}^2 = \|F 1_{|F| \geq R}\|_{\mathbb{D}}^2 \leq \|F\|_{\mathbb{D}}^2 \|1_{|F| \geq R}\|_{\mathbb{D}}^2$$

and $\|1_{|F| \geq R}\|_{\mathbb{D}}^2 = \langle 1_{F \geq R} \Theta 1_{|F| \geq R} \rangle \leq \frac{1}{R^2} \langle |F| \theta |F| \rangle \rightarrow 0$ as $R \rightarrow \infty$.

Hence (3.28) holds for all $F \in \mathcal{F}_{\mathbb{D}}$ with $\|F\|_{\mathbb{D}} < \infty$. This implies S_q maps the null space \mathcal{N}_0 to \mathcal{N}_0 and thus S_q extends to $\mathcal{H}_{\mathbb{D}}$ so that (3.28) holds.

Finally to prove strong continuity, by the semigroup property it suffices to prove it at $q = 1$ and by the contractive property we need to prove it only on a dense set. Since

$$\|S_q F - F\|_{\mathbb{D}}^2 = \|S_q F\|_{\mathbb{D}}^2 + \|F\|_{\mathbb{D}}^2 - (S_q F, F)_{\mathbb{D}} - (F, S_q F)_{\mathbb{D}} \leq 2\|F\|_{\mathbb{D}}^2 - (S_q F, F)_{\mathbb{D}} - (F, S_q F)_{\mathbb{D}}$$

it suffices to prove $(S_q F, F)_{\mathbb{D}} \rightarrow (F, F)_{\mathbb{D}}$ as $q \rightarrow 1$ on a dense set of F . Take $F = GI^{-1}$ so that $UF = U_0 G$. Then

$$(F, S_q F)_{\mathbb{D}} = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E}(\Theta G G e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D} \setminus |q|\mathbb{D})}) dc$$

which converges as $q \rightarrow 1$ to $(F, F)_{\mathbb{D}}$ (use $\mathbb{P}(M_{\gamma}(\mathbb{D} \setminus |q|\mathbb{D}) > \epsilon) \rightarrow 0$ as $q \rightarrow 1$). \square

In particular we can form two one-parameter (semi) groups from S_q . Taking $q = e^{-t}$ we define $T_t = S_{e^{-t}}$. Then $T_{t+s} = T_t T_s$ so T_t is a strongly continuous contraction semigroup on the Hilbert space $\mathcal{H}_{\mathbb{D}}$. Hence by the Hille-Yosida theorem

$$(3.31) \quad U S_{e^{-t}} U^{-1} = e^{-t \mathbf{H}_*}$$

where the generator \mathbf{H}_* (in the case $\mu = 0$, we will write \mathbf{H}_*^0) is a positive self-adjoint operator with domain $\mathcal{D}(\mathbf{H}_*)$ consisting of $\psi \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ such that $\lim_{t \rightarrow 0} \frac{1}{t} (e^{-t \mathbf{H}_*} - 1)\psi$ exists in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. The operator \mathbf{H}_* is the *Hamiltonian* of LCFT. Taking $q = e^{i\alpha}$ we get that $\alpha \mapsto S_{e^{i\alpha}}$ is a strongly continuous unitary group so that by Stone's theorem

$$U S_{e^{i\alpha}} U^{-1} = e^{i\alpha \Pi_*}$$

where Π_* is the self adjoint *momentum* operator of LCFT. As we will have no use for Π_* in this paper we will concentrate on \mathbf{H}_* from now on. Let us emphasize here that it is defined in the full range $\gamma \in (0, 2)$. One of our next tasks will be to show that for $\gamma \in (0, 2)$: $\mathbf{H}_* = \mathbf{H}$, where the Hamiltonian \mathbf{H} will be defined as the Friedrichs extension of (2.11).

4. GAUSSIAN FREE FIELD: DYNAMICS AND CFT ASPECTS

4.1. Fock space and harmonic oscillators. The Hilbert space $L^2(\Omega_{\mathbb{T}}, \mathbb{P}_{\mathbb{T}})$ (denoted from now on by $L^2(\Omega_{\mathbb{T}})$) has the structure of Fock space. Let $\mathcal{P} \subset L^2(\Omega_{\mathbb{T}})$ (resp. $\mathcal{S} \subset L^2(\Omega_{\mathbb{T}})$) be the linear span of the functions of the form $F(x_1, y_1, \dots, x_N, y_N)$ for some $N \geq 1$ where F is a polynomial on \mathbb{R}^{2N} (resp. $F \in C^\infty((\mathbb{R}^2)^N)$ with at most polynomial growth at infinity for F and its derivatives). Obviously $\mathcal{P} \subset \mathcal{S}$ and they are both dense in $L^2(\Omega_{\mathbb{T}})$.

On \mathcal{S} we define the annihilation and creation operators

$$(4.1) \quad \mathbf{X}_n = \partial_{x_n}, \quad \mathbf{X}_n^* = -\partial_{x_n} + x_n,$$

$$(4.2) \quad \mathbf{Y}_n = \partial_{y_n}, \quad \mathbf{Y}_n^* = -\partial_{y_n} + y_n.$$

They are formally adjoint of each other (see e.g. [ReSi1, VIII.11] for more about the closure of these operators, which we will not need here) and form a representation of the algebra of canonical commutation relations on \mathcal{S} :

$$(4.3) \quad [\mathbf{X}_n, \mathbf{X}_m^*] = \delta_{nm} = [\mathbf{Y}_n, \mathbf{Y}_m^*]$$

with other commutators vanishing. The operator \mathbf{P} is then given on \mathcal{S} as

$$(4.4) \quad \mathbf{P} = \sum_{n=1}^{\infty} n(\mathbf{X}_n^* \mathbf{X}_n + \mathbf{Y}_n^* \mathbf{Y}_n)$$

(only finite number of terms in the sum contributes when acting on \mathcal{S}) and extends uniquely to an unbounded self-adjoint positive operator on $L^2(\Omega_{\mathbb{T}})$: this follows from the fact that we can find a complete system of eigenfunctions in \mathcal{P} , as described now. Let \mathcal{N} be the set of non-negative integer valued sequences with only a finite number of non null integers, namely $\mathbf{k} = (k_1, k_2, \dots) \in \mathcal{N}$ iff $\mathbf{k} \in \mathbb{N}^{\mathbb{N}^+}$ and $k_n = 0$ for all n large enough. For $\mathbf{k}, \mathbf{l} \in \mathcal{N}$ define the polynomials (here $1 \in L^2(\Omega_{\mathbb{T}})$ is the constant function)

$$(4.5) \quad \hat{\psi}_{\mathbf{k}\mathbf{l}} = \prod_n (\mathbf{X}_n^*)^{k_n} (\mathbf{Y}_n^*)^{l_n} 1 \in \mathcal{P}.$$

Equivalently, $\hat{\psi}_{\mathbf{k}\mathbf{l}} = \prod_n \text{He}_{k_n}(x_n) \text{He}_{l_n}(y_n)$ where $(\text{He}_k)_{k \geq 0}$ are the standard Hermite polynomials. Then, using (4.3), one checks that these are eigenstates of \mathbf{P} :

$$(4.6) \quad \mathbf{P} \hat{\psi}_{\mathbf{k}\mathbf{l}} = (|\mathbf{k}| + |\mathbf{l}|) \hat{\psi}_{\mathbf{k}\mathbf{l}} = \lambda_{\mathbf{k}\mathbf{l}} \hat{\psi}_{\mathbf{k}\mathbf{l}}.$$

where we use the notations

$$(4.7) \quad |\mathbf{k}| := \sum_{n=1}^{\infty} n k_n, \quad \lambda_{\mathbf{k}\mathbf{l}} := |\mathbf{k}| + |\mathbf{l}|$$

for $\mathbf{k}, \mathbf{l} \in \mathcal{N}$. It is also well known that the family $\{\psi_{\mathbf{k}\mathbf{l}} = \hat{\psi}_{\mathbf{k}\mathbf{l}} / \|\hat{\psi}_{\mathbf{k}\mathbf{l}}\|_{L^2(\Omega_{\mathbb{T}})}\}$ (where $\|\cdot\|_{L^2(\Omega_{\mathbb{T}})}$ is the standard norm in $L^2(\Omega_{\mathbb{T}})$) forms an orthonormal basis of $L^2(\Omega_{\mathbb{T}})$. Finally we claim

Proposition 4.1. *The operator \mathbf{P} generates a strongly continuous semigroup of self-adjoint contractions $(e^{-t\mathbf{P}})_{t \geq 0}$ on $L^2(\Omega_{\mathbb{T}})$ with probabilistic representation, for $t \geq 0$,*

$$\forall f \in L^2(\Omega_{\mathbb{T}}), \quad e^{-t\mathbf{P}} f = \mathbb{E}_{\varphi}[f(\varphi_t)]$$

with $(\varphi_t)_{t \geq 0}$ the process defined by (3.15).

Proof. The fact that \mathbf{P} generates a strongly continuous semigroup of self-adjoint contractions results from the fact that \mathbf{P} is self-adjoint and nonnegative. Since $(\psi_{\mathbf{k}\mathbf{l}})_{\mathbf{k}, \mathbf{l}}$ form an orthonormal basis of $L^2(\Omega_{\mathbb{T}})$, it suffices to study the semigroup on this basis. Obviously $e^{-t\mathbf{P}} \psi_{\mathbf{k}\mathbf{l}} = e^{-\lambda_{\mathbf{k}\mathbf{l}} t} \psi_{\mathbf{k}\mathbf{l}}$. Furthermore the decomposition (3.14) together with the covariance structure (3.9) (and recalling the decomposition (3.1)+(3.2) of the field φ) entails that the law of φ_t (see (3.15)) conditionally on φ is given by

$$\varphi_t(\theta) = \sum_{n>0} \frac{x_n(t) + iy_n(t)}{2\sqrt{n}} e^{in\theta} + \sum_{n<0} \frac{x_{-n}(t) - iy_{-n}(t)}{2\sqrt{-n}} e^{in\theta}$$

where $x_n(t), y_n(t)$ are independent Ornstein-Uhlenbeck processes. In particular for each fixed t , there are two independent sequences of independent standard Gaussians $(\bar{x}_n)_n$ and $(\bar{y}_n)_n$ such that, for $n \geq 1$

$$x_n(t) \stackrel{\text{law cond. on } \varphi}{=} e^{-nt}x_n + \sqrt{1 - e^{-2tn}}\bar{x}_n, \quad y_n(t) \stackrel{\text{law cond. on } \varphi}{=} e^{-nt}y_n + \sqrt{1 - e^{-2tn}}\bar{y}_n.$$

Finally, we recall the following elementary result: given Y a standard Gaussian random variable, the standard Hermite polynomials $(\text{He}_k)_{k \geq 0}$ on \mathbb{R} , $x \in \mathbb{R}$ and $u, v \geq 0$ such that $u^2 + v^2 = 1$ then

$$(4.8) \quad \mathbb{E}[\text{He}_k(ux + vY)] = u^k \text{He}_k(x).$$

Using this lemma and our description of the law of X_t , it is then plain to deduce that

$$e^{-t\mathbf{P}}\psi_{\mathbf{k}\mathbf{l}} = \mathbb{E}_\varphi[\psi_{\mathbf{k}\mathbf{l}}(\varphi_t)] = e^{-\lambda_{\mathbf{k}\mathbf{l}}t}\psi_{\mathbf{k}\mathbf{l}}.$$

Hence our claim. \square

Remark 4.2. List the eigenvalues $\lambda = |\mathbf{k}| + |\mathbf{l}|$ of \mathbf{P} in increasing order $\lambda_1 < \lambda_2 < \dots$ and let P_i be the corresponding spectral projectors. Since each λ_i is of finite multiplicity and $\lambda_i \rightarrow \infty$ as $i \rightarrow \infty$ the semigroup $e^{-t\mathbf{P}} = \sum_i e^{-t\lambda_i}P_i$ and the resolvent $(z - \mathbf{P})^{-1} = \sum_i (z - \lambda_i)^{-1}P_i$ are compact if $t > 0$ and $\Im z \neq 0$ since they are norm convergent limits of finite rank operators.

4.2. Quadratic form. Introduce the bilinear form (with associated quadratic form still denoted by \mathcal{Q}_0)

$$(4.9) \quad \forall u, v \in \mathcal{C}, \quad \mathcal{Q}_0(u, v) := \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(\partial_c u \partial_c \bar{v} + Q^2 u \bar{v} + 2(\mathbf{P}u)\bar{v} \right) dc$$

with

$$(4.10) \quad \mathcal{C} = \text{Span}\{\psi(c)F \mid \psi \in C_c^\infty(\mathbb{R}) \text{ and } F \in \mathcal{S}\}.$$

We claim

Proposition 4.3. The quadratic form (4.9) is closable (and we still denote its closure by \mathcal{Q}_0 with domain $\mathcal{D}(\mathcal{Q}_0)$) and lower semibounded: $\mathcal{Q}(u) \geq Q^2 \|u\|_2^2/2$. It determines uniquely a self-adjoint operator \mathbf{H}^0 , called the Friedrichs extension, with domain denoted by $\mathcal{D}(\mathbf{H}^0)$ such that:

$$\mathcal{D}(\mathbf{H}^0) = \{u \in \mathcal{D}(\mathcal{Q}_0) \mid \exists C > 0, \forall v \in \mathcal{D}(\mathcal{Q}_0), |\mathcal{Q}_0(u, v)| \leq C \|v\|_2\}$$

and for $u \in \mathcal{D}(\mathbf{H}^0)$, $\mathbf{H}^0 u$ is the unique element in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ satisfying

$$\mathcal{Q}_0(u, v) = \langle \mathbf{H}^0 u | v \rangle_2.$$

Proof. Recall that the closability of the quadratic form means that its completion with respect to the \mathcal{Q}_0 -norm embeds continuously and injectively in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. Its completion is the vector space consisting of equivalence classes of Cauchy sequences of \mathcal{C} for the \mathcal{Q}_0 -norm under the equivalence relation $u \sim v$ iff $\mathcal{Q}_0(u_n - v_n) \rightarrow 0$ as $n \rightarrow \infty$. This space is a Hilbert space. Let us show that it embeds injectively and continuously in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ by the map $j : [u] \mapsto \lim_{n \rightarrow \infty} u_n$. Indeed, u_n is Cauchy for $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ since $\|u_n - u_m\|_2^2 \leq 2Q^{-2} \mathcal{Q}_0(u_n - v_n)$, it thus converges in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. Moreover $\|\lim_n u_n\|_2^2 \leq 2Q^{-2} \lim_n \mathcal{Q}_0(u_n)$ thus j is bounded. Finally if $j([u]) = 0$, then for $(u_n)_n$ a representative Cauchy sequence of $[u]$, we have $u_n \rightarrow 0$ in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and using

$$\frac{1}{2} \|\partial_c(u_n - u_m)\|_2^2 + \|\mathbf{P}^{1/2}(u_n - u_m)\|_2^2 \leq \mathcal{Q}_0(u_n - u_m, u_n - u_m),$$

one has the convergence in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ of $\partial_c u_n \rightarrow v$ and $\mathbf{P}^{1/2} u_n \rightarrow w$ for some $v, w \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. For each $\varphi \in \mathcal{C}$, we have as $n \rightarrow \infty$

$$\langle \partial_c u_n, \varphi \rangle_2 = \langle u_n, -\partial_c \varphi \rangle \rightarrow 0, \quad \langle \mathbf{P}^{1/2} u_n, \varphi \rangle_2 = \langle u_n, \mathbf{P}^{1/2} \varphi \rangle_2 \rightarrow 0,$$

thus $v = w = z = 0$ by density of \mathcal{C} in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. This implies that $\mathcal{Q}_0(u_n) \rightarrow 0$ and thus j is injective.

Let us now consider the closure \mathcal{Q}_0 with domain $\mathcal{D}(\mathcal{Q}_0)$. Obviously it is closed and lower semi-bounded $\mathcal{Q}_0(u) \geq Q^2 \|u\|_2^2/2$ so that the construction of the Friedrichs extension then follows from [ReSi1, Theorem 8.15]. \square

If we let $\mathcal{D}(\mathcal{Q}_0)'$ be the dual to $\mathcal{D}(\mathcal{Q}_0)$ (i.e. the space of bounded conjugate linear functionals on $\mathcal{D}(\mathcal{Q}_0)$), the injection $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \subset \mathcal{D}(\mathcal{Q}_0)'$ is continuous and the operator \mathbf{H}^0 can be extended as a bounded isomorphism

$$\mathbf{H}^0 : \mathcal{D}(\mathcal{Q}_0) \rightarrow \mathcal{D}(\mathcal{Q}_0)'.$$

We also have $\mathcal{D}(\mathbf{H}^0) = \{u \in \mathcal{D}(\mathcal{Q}_0) \mid \mathbf{H}^0 u \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})\}$ and $(\mathbf{H}^0)^{-1} : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H}^0)$ is bounded. Furthermore, by the spectral theorem, it generates a strongly continuous contraction semigroup of self-adjoint operators $(e^{-t\mathbf{H}^0})_{t \geq 0}$ on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$.

4.3. Dynamics of the GFF. The goal of this subsection is to prove the relation $\mathbf{H}_*^0 = \mathbf{H}^0$, i.e. we want to show

Proposition 4.4. *For all $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and all $t \geq 0$*

$$(4.11) \quad U_0 S_{e^{-t}} U_0^{-1} f = e^{-t\mathbf{H}^0} f = e^{-\frac{Q^2 t}{2}} \mathbb{E}_{\varphi} [f(c + B_t, \varphi_t)]$$

Proof. Recalling (3.10), we have the independent sum

$$X_{\mathbb{D}}(e^{-t+i\theta}) = B_t + Y_t(\theta)$$

where B_t is a Brownian motion and Y_t has zero average on the circle. We then have

$$\begin{aligned} (U_0 S_{e^{-t}} U_0^{-1} f)(c, \varphi) &= e^{-Qc} \mathbb{E}_{\varphi} [e^{Q(c+B_t-Qt)} f(c + B_t - Qt, P\varphi(e^{-t+i\cdot}) + Y_t(\cdot))] \\ &= e^{-\frac{Q^2}{2}t} \mathbb{E}_{\varphi} [f(c + B_t, P\varphi(e^{-t+i\cdot}) + Y_t(\cdot))] \end{aligned}$$

where we have used the Girsanov transform to obtain the last equality. Since B_t and Y_t are independent conditionally on φ , this last quantity is also equal to $e^{-t(\frac{Q^2}{2} - \frac{\partial_c^2}{2})} e^{-t\mathbf{P}} f$ by using Proposition 4.1. Furthermore, for $f \in \mathcal{C}$, it is plain to see that the mapping $t \mapsto e^{-t(\frac{Q^2}{2} - \frac{\partial_c^2}{2})} e^{-t\mathbf{P}} f$ solves the Cauchy problem $\partial_t u = -\mathbf{H}^0 u$ with $u(0) = f$. Hence $e^{-t(\frac{Q^2}{2} - \frac{\partial_c^2}{2})} e^{-t\mathbf{P}} f = e^{-t\mathbf{H}^0} f$. \square

Finally, we have the simple:

Proposition 4.5. *The following properties hold:*

- (1) *The measure $dc \times \mathbb{P}_{\mathbb{T}}$ is invariant for $e^{\frac{Q^2 t}{2}} e^{-t\mathbf{H}^0}$.*
- (2) *$e^{-t\mathbf{H}^0}$ extends to a continuous semigroup on $L^p(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $p \in [1, +\infty]$ with norm $e^{-\frac{Q^2}{2}t}$ and it is strongly continuous for $p \in [1, +\infty)$.*
- (3) *$e^{-t\mathbf{H}^0}$ extends to a strongly continuous semigroup on $e^{-\alpha c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $\alpha \in \mathbb{R}$ with norm $e^{(\frac{\alpha^2}{2} - \frac{Q^2}{2})t}$.*

Proof. 1) This is a consequence of (4.11): indeed the processes B and Y are independent and describe two dynamics for which the measures dc and \mathbb{P} are respectively invariant. 2) follows from (4.11), Jensen's inequality and the fact that $dc \otimes \mathbb{P}_{\mathbb{T}}$ is invariant for \mathbf{H}^0 . 3) The map $K : f \mapsto e^{-\alpha c} f$ is unitary from $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\alpha c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. We have $K e^{-t\mathbf{H}^0} K^{-1} = e^{t(\frac{\alpha^2}{2} - \alpha \partial_c)} e^{-t\mathbf{H}^0}$ which implies the claim. \square

Remark 4.6. *Using the decomposition $L^2(\Omega_{\mathbb{T}}) = \bigoplus_{\mathbf{k}, \mathbf{l}} \ker(\mathbf{P} - \lambda_{\mathbf{k}, \mathbf{l}})$, the operator \mathbf{H}^0 is unitarily equivalent to the direct sum $\bigoplus_{\mathbf{k}, \mathbf{l}} (-\frac{1}{2}\partial_c^2 + \frac{Q^2}{2} + \lambda_{\mathbf{k}, \mathbf{l}})$, each of these operators being a shifted Laplacian on the real line \mathbb{R} . Consequently (using Fourier transform in c), \mathbf{H}^0 has no L^2 -eigenvalue, its spectrum is absolutely continuous and the family $(e^{iPc} \psi_{\mathbf{k}, \mathbf{l}})_{P, \mathbf{k}, \mathbf{l}}$ form a complete family of generalized eigenstates diagonalizing \mathbf{H}^0 .*

4.4. Diagonalization of the free Hamiltonian using the Virasoro algebra. We start by explaining the diagonalization of the free (i.e. non interacting) Hamiltonian \mathbf{H}^0 which corresponds to the case $\mu = 0$ in (2.11). As explained in Remark 4.6, that can be done directly by using the orthonormal basis of Hermite polynomials $\psi_{\mathbf{k}\mathbf{l}}$ of $L^2(\Omega_{\mathbb{T}})$ combined with the Plancherel formula for the Fourier transform on the real line: for each $u_1, u_2 \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, one has

$$(4.12) \quad \langle u_1 | u_2 \rangle_2 = \frac{1}{2\pi} \sum_{\mathbf{k}, \mathbf{l} \in \mathcal{N}} \int_{\mathbb{R}} \langle u_1 | e^{iPc} \psi_{\mathbf{k}\mathbf{l}} \rangle_2 \langle e^{iPc} \psi_{\mathbf{k}\mathbf{l}} | u_2 \rangle_2 dP.$$

It will be useful however to use another basis for $L^2(\Omega_{\mathbb{T}})$ which respects its underlying complex analytic structure; this new basis, made up of \mathbf{H}^0 -eigenstates, will be generated by the action on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ of two commuting unitary representations of the Virasoro algebra (as motivated in the end of Subsection 2.4). We follow below the Segal-Sugawara construction for the Fock representation of the Heisenberg algebra. Let us emphasize that the material we introduce here is standard; to keep the paper self-contained, we recall the main properties of the construction and just give sketches of the proofs (see for instance [Go09, KaRa] for more details).

Fock representation of the Heisenberg algebra. We will work on the vector space

$$(4.13) \quad \mathcal{C}_{\infty} := \text{Span}\{\psi(c)F \mid \psi \in C^{\infty}(\mathbb{R}) \text{ and } F \in \mathcal{S}\}.$$

(not to be confused with \mathcal{C} which is a subset of \mathcal{C}_{∞}) and use the complex coordinates (3.2), i.e. we denote for $n > 0$

$$\partial_n := \frac{\partial}{\partial \varphi_n} = \sqrt{n}(\partial_{x_n} - i\partial_{y_n}) \quad \text{and} \quad \partial_{-n} := \frac{\partial}{\partial \varphi_{-n}} = \sqrt{n}(\partial_{x_n} + i\partial_{y_n}).$$

We define on \mathcal{C}_{∞} the following operators for $n > 0$:

$$\begin{aligned} \mathbf{A}_n &= \frac{i}{2} \partial_n, & \mathbf{A}_{-n} &= \frac{i}{2} (\partial_{-n} - 2n\varphi_n) \\ \tilde{\mathbf{A}}_n &= \frac{i}{2} \partial_{-n}, & \tilde{\mathbf{A}}_{-n} &= \frac{i}{2} (\partial_n - 2n\varphi_{-n}) \\ \mathbf{A}_0 &= \tilde{\mathbf{A}}_0 = \frac{i}{2} (\partial_c + Q). \end{aligned}$$

Their restrictions to \mathcal{C} are closable operators in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ satisfying (on their closed extension)

$$(4.14) \quad \mathbf{A}_n^* = \mathbf{A}_{-n}, \quad \tilde{\mathbf{A}}_n^* = \tilde{\mathbf{A}}_{-n}.$$

Furthermore $\mathbf{A}_n \mathbf{1} = 0$ and $\tilde{\mathbf{A}}_n \mathbf{1} = 0$ for $n > 0$. It is easy to see that the space \mathcal{C}_{∞} is stable by the operators $\mathbf{A}_n, \tilde{\mathbf{A}}_n$ and we have the commutation relations on \mathcal{C}_{∞}

$$(4.15) \quad [\mathbf{A}_n, \mathbf{A}_m] = \frac{n}{2} \delta_{n, -m} = [\tilde{\mathbf{A}}_n, \tilde{\mathbf{A}}_m], \quad [\mathbf{A}_n, \tilde{\mathbf{A}}_m] = 0.$$

Thus \mathbf{A}_n and $\tilde{\mathbf{A}}_n$ ($n > 0$) are annihilation operators and $\mathbf{A}_{-n}, \tilde{\mathbf{A}}_{-n}$ creation operators. By identifying canonically \mathcal{S} (defined in the beginning of subsection 4.1) as a subspace of \mathcal{C}_{∞} , it is plain to check that \mathcal{S} is stable under the action of these operators. As before, let $\mathbf{k}, \mathbf{l} \in \mathcal{N}$ and define the polynomials

$$(4.16) \quad \hat{\pi}_{\mathbf{k}\mathbf{l}} = \prod_{n>0} \mathbf{A}_{-n}^{k_n} \tilde{\mathbf{A}}_{-n}^{l_n} \mathbf{1}.$$

Then $\hat{\pi}_{\mathbf{k}\mathbf{l}}$ and $\hat{\pi}_{\mathbf{k}'\mathbf{l}'}$ are orthogonal if $\mathbf{k} \neq \mathbf{k}'$ or $\mathbf{l} \neq \mathbf{l}'$ and the $\hat{\pi}_{\mathbf{k}\mathbf{l}}$'s with $|\mathbf{k}| + |\mathbf{l}| = N$ span the eigenspace of \mathbf{P} in $L^2(\Omega_{\mathbb{T}})$ with eigenvalue N ¹². We denote $\pi_{\mathbf{k}\mathbf{l}} := \hat{\pi}_{\mathbf{k}\mathbf{l}} / \|\hat{\pi}_{\mathbf{k}\mathbf{l}}\|_{L^2(\Omega_{\mathbb{T}})}$ the normalized eigenvectors.

¹²Explicitly: $\hat{\pi}_{\mathbf{k}\mathbf{l}} = \prod_{n>0} (-in)^{k_n+l_n} \varphi_n^{k_n} \varphi_{-n}^{l_n} + P(\varphi_n, \varphi_{-n})$ where P is a polynomial in φ_n, φ_{-n} spanned by monomials of the form $\prod_{n>0} \varphi_n^{k'_n} \varphi_{-n}^{l'_n}$ with $k'_n \leq k_n, l'_n \leq l_n$ and $\sum_{n>0} k'_n + l'_n < \sum_{n>0} k_n + l_n$.

Segal-Sugawara construction. Now we use the Fock representation of the Heisenberg algebra to construct the Virasoro representation. We define the *normal ordered product* on \mathcal{C}_∞ by $:\mathbf{A}_n\mathbf{A}_m:=\mathbf{A}_n\mathbf{A}_m$ if $m > 0$ and $\mathbf{A}_m\mathbf{A}_n$ if $n > 0$ (i.e. annihilation operators are on the right) and then for all $n \in \mathbb{Z}$

$$(4.17) \quad \mathbf{L}_n^0 := -i(n+1)Q\mathbf{A}_n + \sum_{m \in \mathbb{Z}} : \mathbf{A}_{n-m}\mathbf{A}_m :$$

$$(4.18) \quad \tilde{\mathbf{L}}_n^0 := -i(n+1)Q\tilde{\mathbf{A}}_n + \sum_{m \in \mathbb{Z}} : \tilde{\mathbf{A}}_{n-m}\tilde{\mathbf{A}}_m : .$$

These operators are well defined on \mathcal{C}_∞ (since only a finite number of terms contribute) and their restrictions to \mathcal{C} are closable operators satisfying (on their closed extensions)

$$(4.19) \quad (\mathbf{L}_n^0)^* = \mathbf{L}_{-n}^0, \quad (\tilde{\mathbf{L}}_n^0)^* = \tilde{\mathbf{L}}_{-n}^0.$$

Furthermore the vector space \mathcal{C}_∞ is stable under \mathbf{L}_n^0 and $\tilde{\mathbf{L}}_n^0$ for all $n \in \mathbb{Z}$; on \mathcal{C}_∞ the \mathbf{L}_n^0 satisfy the commutation relations of the *Virasoro Algebra* (see [KaRa, Prop 2.3]):

$$(4.20) \quad [\mathbf{L}_n^0, \mathbf{L}_m^0] = (n-m)\mathbf{L}_{n+m}^0 + \frac{c_L}{12}(n^3-n)\delta_{n,-m}$$

where the central charge is

$$c_L = 1 + 6Q^2.$$

These commutation relations can be checked by using the fact that, on \mathcal{C}_∞ , only finitely many terms contribute in (4.20) and using the commutation relation (4.15). $\tilde{\mathbf{L}}_n^0$ satisfy the same commutation relations (4.20) and commute with the \mathbf{L}_n^0 's. Note also that

$$(4.21) \quad \mathbf{L}_0^0 = \frac{1}{4}(-\partial_c^2 + Q^2) + 2 \sum_{n>0} \mathbf{A}_{-n}\mathbf{A}_n$$

$$(4.22) \quad \tilde{\mathbf{L}}_0^0 = \frac{1}{4}(-\partial_c^2 + Q^2) + 2 \sum_{n>0} \tilde{\mathbf{A}}_{-n}\tilde{\mathbf{A}}_n,$$

so that one can easily check that the $\mu = 0$ Hamiltonian $\mathbf{H}^0 := -\frac{1}{2}\partial_c^2 + \frac{1}{2}Q^2 + \mathbf{P}$ has the following decomposition when restricted on \mathcal{C}_∞

$$\mathbf{H}^0 = \mathbf{L}_0^0 + \tilde{\mathbf{L}}_0^0.$$

Remark 4.7. *In the terminology of representation theory, we have a unitary representation of two commuting Virasoro Algebras on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ (unitary in the sense that (4.19) holds) and this representation is reducible as we will see below by constructing stable sub-representations.*

Diagonalizing \mathbf{H}^0 using the Virasoro representation. Now we explain how to construct the generalized eigenstates of the free Hamiltonian \mathbf{H}^0 using the families of operators $(\mathbf{L}_n^0)_n$ and $(\tilde{\mathbf{L}}_n^0)_n$. Recall that, for $\alpha \in \mathbb{C}$, we have defined the function

$$(4.23) \quad \Psi_\alpha^0(c, \varphi) := e^{(\alpha-Q)c} \in \mathcal{C}_\infty.$$

For $\alpha \in \mathbb{C}$, these are generalized eigenstates of \mathbf{H}^0 : they never belong to $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ but rather to some weighted spaces $e^{\beta|c|}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > |\operatorname{Re}(\alpha) - Q|$, hence their name ‘‘generalized eigenstates’’. We have

$$(4.24) \quad \begin{aligned} \mathbf{L}_0^0\Psi_\alpha^0 &= \tilde{\mathbf{L}}_0^0\Psi_\alpha^0 = \Delta_\alpha\Psi_\alpha^0 \\ \mathbf{L}_n^0\Psi_\alpha^0 &= \tilde{\mathbf{L}}_n^0\Psi_\alpha^0 = 0, \quad n > 0, \end{aligned}$$

where Δ_α is the conformal weight (1.9). In the language of representation theory (or in the CFT terminology), Ψ_α^0 is called *highest weight state* with highest weight Δ_α for both algebras. Before defining the so-called descendants of Ψ_α^0 , we introduce the following definition:

Definition 4.8. *A sequence of integers $\nu = (\nu_i)_{i \geq 0}$ is called a Young diagram if the mapping $i \mapsto \nu_i$ is non-increasing and if $\nu_i = 0$ for i sufficiently large. We denote by \mathcal{T} the set of all Young diagrams. We will sometimes write $\nu = (\nu_i)_{i \in [1, k]}$ where k is the last integer i such that $\nu_i > 0$ and denote by $|\nu| := \sum_{i \geq 1} \nu_i$ the length of the Young diagram¹³. We set $\mathcal{T}_j := \{\nu \in \mathcal{T} \mid |\nu| = j\}$ the set of Young diagrams of length j .*

¹³This length should not be confused with the length (4.7) of a sequence of integers.

Given two Young diagrams $\nu = (\nu_i)_{i \in [1, k]}$ and $\tilde{\nu} = (\tilde{\nu}_i)_{i \in [1, j]}$ we denote

$$\mathbf{L}_{-\nu}^0 = \mathbf{L}_{-\nu_k}^0 \cdots \mathbf{L}_{-\nu_1}^0, \quad \tilde{\mathbf{L}}_{-\tilde{\nu}}^0 = \tilde{\mathbf{L}}_{-\tilde{\nu}_j}^0 \cdots \tilde{\mathbf{L}}_{-\tilde{\nu}_1}^0$$

and define

$$(4.25) \quad \Psi_{\alpha, \nu, \tilde{\nu}}^0 = \mathbf{L}_{-\nu}^0 \tilde{\mathbf{L}}_{-\tilde{\nu}}^0 \Psi_{\alpha}^0,$$

with the convention that $\Psi_{\alpha, \emptyset, \emptyset}^0 = \Psi_{\alpha}^0$. The vectors $\Psi_{\alpha, \nu, \tilde{\nu}}^0$ are called the *descendants* states of Ψ_{α}^0 . We gather in the following proposition their main properties

Proposition 4.9. *The following holds:*

1) *For each pair of Young diagrams $\nu, \tilde{\nu} \in \mathcal{T}$, the descendant state $\Psi_{\alpha, \nu, \tilde{\nu}}^0$ can be written as*

$$(4.26) \quad \Psi_{\alpha, \nu, \tilde{\nu}}^0 = \mathcal{Q}_{\alpha, \nu, \tilde{\nu}} \Psi_{\alpha}^0$$

where $\mathcal{Q}_{\alpha, \nu, \tilde{\nu}} \in \mathcal{P}$ is a polynomial.

2) *for all $\alpha \in \mathbb{C}$*

$$\mathbf{L}_0^0 \Psi_{\alpha, \nu, \tilde{\nu}}^0 = (\Delta_{\alpha} + |\nu|) \Psi_{\alpha, \nu, \tilde{\nu}}^0, \quad \tilde{\mathbf{L}}_0^0 \Psi_{\alpha, \nu, \tilde{\nu}}^0 = (\Delta_{\alpha} + |\tilde{\nu}|) \Psi_{\alpha, \nu, \tilde{\nu}}^0$$

and thus since $\mathbf{H}^0 = \mathbf{L}_0^0 + \tilde{\mathbf{L}}_0^0$

$$\mathbf{H}^0 \Psi_{\alpha, \nu, \tilde{\nu}}^0 = (2\Delta_{\alpha} + |\nu| + |\tilde{\nu}|) \Psi_{\alpha, \nu, \tilde{\nu}}^0.$$

3) **Completeness:** *the inner products of the descendant states obey*

$$(4.27) \quad \langle \mathcal{Q}_{2Q-\bar{\alpha}, \nu, \tilde{\nu}} | \mathcal{Q}_{\alpha, \nu', \tilde{\nu}'} \rangle_{L^2(\Omega_{\mathbb{T}})} = \delta_{|\nu|, |\nu'|} \delta_{|\tilde{\nu}|, |\tilde{\nu}'|} F_{\alpha}(\nu, \nu') F_{\alpha}(\tilde{\nu}, \tilde{\nu}')$$

where each coefficient $F_{\alpha}(\nu, \nu')$ is a polynomial in α , called the *Schapovalov form*. The functions $(\mathcal{Q}_{\alpha, \nu, \tilde{\nu}})_{\nu, \tilde{\nu} \in \mathcal{T}}$ are linearly independent for

$$\alpha \notin \{\alpha_{r,s} \mid r, s \in \mathbb{N}^*, rs \leq \max(|\nu|, |\tilde{\nu}|)\} \quad \text{with} \quad \alpha_{r,s} = Q - r \frac{\gamma}{2} - s \frac{2}{\gamma}.$$

4) *the following holds*

$$(4.28) \quad \mathcal{Q}_{\alpha, \nu, \tilde{\nu}} := \sum_{\mathbf{k}, \mathbf{l}, |\mathbf{k}|+|\mathbf{l}|=N} M_{\alpha, \mathbf{k}, \mathbf{l}, \nu, \tilde{\nu}}^N \psi_{\mathbf{k}, \mathbf{l}},$$

for some coefficients $M_{\alpha, \mathbf{k}, \mathbf{l}, \nu, \tilde{\nu}}^N$ polynomial in $\alpha \in \mathbb{C}$.

5) **Spectral decomposition:** *if $u_1, u_2 \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ then*

$$(4.29) \quad \langle u_1 | u_2 \rangle_2 = \frac{1}{2\pi} \sum_{\nu, \tilde{\nu}, \nu', \tilde{\nu}' \in \mathcal{T}} \int_{\mathbb{R}} \langle u_1 | \Psi_{Q+iP, \nu', \tilde{\nu}'}^0 \rangle_2 \langle \Psi_{Q+iP, \nu, \tilde{\nu}}^0 | u_2 \rangle_2 F_{P+iQ}^{-1}(\nu, \nu') F_{P+iQ}^{-1}(\tilde{\nu}, \tilde{\nu}') dP.$$

Proof. The decomposition (4.26) can be obtained from the definition (4.25) by using that $[\mathbf{A}_n, e^{(\alpha-Q)c}] = [\tilde{\mathbf{A}}_n, e^{(\alpha-Q)c}] = 0$ for all $n \neq 0$, $[\partial_c, e^{(\alpha-Q)c}] = e^{(\alpha-Q)c}(\alpha - Q)\text{Id}$ and that finitely many applications of \mathbf{A}_n and $\tilde{\mathbf{A}}_n$ to $\mathbf{1}$ is a polynomial in \mathcal{P} . Actually, one can even show that for ν a Young diagram

$$\mathbf{L}_{-\nu}^0 \Psi_{\alpha}^0 = (\mathbf{L}_{-\nu}^{0, \alpha} \mathbf{1}) \Psi_{\alpha}^0$$

where, for a Young diagram $\nu = (\nu_1, \dots, \nu_k)$, we define $\mathbf{L}_{\nu}^{0, \alpha} := \mathbf{L}_{\nu_1}^{0, \alpha} \dots \mathbf{L}_{\nu_k}^{0, \alpha}$ and the operators $\mathbf{L}_n^{0, \alpha}$, $n \in \mathbb{Z}$ act in $L^2(\Omega_{\mathbb{T}})$ and are given by the expression (4.17) where we replace \mathbf{A}_0 by $\frac{i}{2}(\alpha + Q)$ i.e.

$$\mathbf{L}_n^{0, \alpha} := \begin{cases} i(\alpha - Q - nQ) \mathbf{A}_n + \sum_{m \neq n, 0} \mathbf{A}_{n-m} \mathbf{A}_m & n \neq 0 \\ \frac{\alpha}{2}(Q - \frac{\alpha}{2}) + 2 \sum_{m > 0} \mathbf{A}_{-m} \mathbf{A}_m & n = 0 \end{cases}.$$

Hence on their closed extensions we have

$$(4.30) \quad (\mathbf{L}_n^{0, \alpha})^* = \mathbf{L}_{-n}^{0, 2Q-\bar{\alpha}}$$

and $\mathbf{L}_n^{0, \alpha}$ satisfies (4.20). The reader should notice here that the order in which the ν_j appear in previous definition ensures that $(\mathbf{L}_{\nu}^{0, 2Q-\bar{\alpha}})^* = \mathbf{L}_{-\nu}^{0, \alpha}$. The operators $(\tilde{\mathbf{L}}_n^{0, \alpha})_n$ are defined in a similar fashion and commute with $(\mathbf{L}_n^{0, \alpha})_n$. Therefore $\mathcal{Q}_{\alpha, \nu, \tilde{\nu}} = \mathbf{L}_{-\nu}^{0, \alpha} \tilde{\mathbf{L}}_{-\tilde{\nu}}^{0, \alpha} \mathbf{1}$. Then 2) results from (4.24) and the commutation relations (4.20) with $n = 0$.

Since $\mathbf{L}_\nu^{0,\alpha}$ and $\tilde{\mathbf{L}}_{\tilde{\nu}}^{0,\alpha}$ commute we have

$$(4.31) \quad \langle \mathcal{Q}_{2Q-\bar{\alpha},\nu,\tilde{\nu}} | \mathcal{Q}_{\alpha,\nu',\tilde{\nu}'} \rangle_{L^2(\Omega_\mp)} = \langle 1 | \tilde{\mathbf{L}}_{\tilde{\nu}}^{0,\alpha} \tilde{\mathbf{L}}_{-\tilde{\nu}'}^{0,\alpha} \mathbf{L}_\nu^{0,\alpha} \mathbf{L}_{-\nu'}^{0,\alpha} 1 \rangle_{L^2(\Omega_\mp)}.$$

Let us now compute the right-hand side. By Lemma A.1, one has for arbitrary $t_1, \dots, t_k \in \mathbb{Z}$ such that $t_1 + \dots + t_k > 0$

$$(4.32) \quad \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 = 0, \quad \tilde{\mathbf{L}}_{t_1}^{0,\alpha} \dots \tilde{\mathbf{L}}_{t_k}^{0,\alpha} 1 = 0.$$

Therefore, if $|\nu| > |\nu'|$ or $|\tilde{\nu}| > |\tilde{\nu}'|$ then we get that (4.31) is equal to 0 by using (4.32). The case $|\nu| < |\nu'|$ or $|\tilde{\nu}| < |\tilde{\nu}'|$ can be dealt similarly and yields 0 also. Hence in what follows we suppose that $|\nu| = |\nu'|$ and $|\tilde{\nu}| = |\tilde{\nu}'|$.

Lemma A.2 establishes that for $|\nu| = |\nu'|$

$$(4.33) \quad \mathbf{L}_\nu^{0,\alpha} \mathbf{L}_{-\nu'}^{0,\alpha} 1 = \sum_{k \geq 0} a_k (\mathbf{L}_0^{0,\alpha})^k 1$$

where the coefficients a_k are determined by the algebra (4.20) and are independent of α . Since $(\mathbf{L}_0^{0,\alpha})^k 1 = \Delta_\alpha^k$ and $\Delta_\alpha = \frac{\alpha}{2}(Q - \frac{\alpha}{2})$ we conclude

$$\mathbf{L}_\nu^{0,\alpha} \mathbf{L}_{-\nu'}^{0,\alpha} 1 = F_\alpha(\nu, \nu')$$

where $F_\alpha(\nu, \nu')$ is a polynomial in Δ_α and thus in α . Repeating the argument for $\tilde{\mathbf{L}}_{\tilde{\nu}}^{0,\alpha} \tilde{\mathbf{L}}_{-\tilde{\nu}'}^{0,\alpha} 1$ yields (4.27).

The determinant of the matrix $(F_\alpha(\nu, \nu'))_{|\nu|=|\nu'|=N}$ is given by the Kac determinant formula (see Feigin-Fuchs [FF84])

$$(4.34) \quad \det(F_\alpha(\nu, \nu'))_{|\nu|=|\nu'|=N} = \kappa_N \prod_{r,s=1; rs \leq N}^N (\Delta_\alpha - \Delta_{\alpha_{r,s}})^{p(N-rs)}$$

where κ_N does not depend on α or c_L , $p(M)$ is the number of Young Diagrams of length M and

$$\alpha_{r,s} = Q - r \frac{\gamma}{2} - s \frac{2}{\gamma}.$$

Now we turn to 4). The polynomials $\mathcal{Q}_{\alpha,\nu,\tilde{\nu}}$ with $|\nu| + |\tilde{\nu}| = N$ belong to the space spanned by the basis $\psi_{\mathbf{k}\mathbf{l}}$ (4.6) (or equivalently the basis $\pi_{\mathbf{k}\mathbf{l}}$ (4.16)) above with $|\mathbf{k}| + |\mathbf{l}| = N$. Let us also stress that item 3) entails that the space spanned by $\mathcal{Q}_{\alpha,\nu,\tilde{\nu}}$ with $|\nu| + |\tilde{\nu}| = N$ is exactly the same as $\psi_{\mathbf{k}\mathbf{l}}$ with $|\mathbf{k}| + |\mathbf{l}| = N$ as soon as $\alpha \notin Q - \frac{\gamma}{2}\mathbb{N}^* - \frac{2}{\gamma}\mathbb{N}^*$. Hence the existence of a decomposition of the type (4.28). The fact that the coefficients of this decomposition are polynomials in α can be more easily seen in the basis $\pi_{\mathbf{k}\mathbf{l}}$: in that case, the coefficients are given by the scalar products $\langle \mathcal{Q}_{\alpha,\nu,\tilde{\nu}} | \pi_{\mathbf{k}\mathbf{l}} \rangle_{L^2(\Omega_\mp)}$ and one can use the expression (4.16) together with the commutation relations

$$[\mathbf{A}_n, \mathbf{L}_m^{0,\alpha}] = n \mathbf{A}_{n+m}^\alpha - \frac{i}{2} n(n+1) Q \delta_{n,-m}$$

where $\mathbf{A}_0^\alpha = \frac{i}{2}(\alpha + Q)$ and $\mathbf{A}_n^\alpha = \mathbf{A}_n$ for $n \neq 0$ and $\mathbf{L}_0^{0,\alpha} 1 = \Delta_\alpha$. 4) follows.

Now, we specialise the above considerations to the case $\alpha = Q + iP$ with $P \in \mathbb{R}$. In this case, one has

$$(4.35) \quad \langle \mathcal{Q}_{Q+iP,\nu,\tilde{\nu}} | \mathcal{Q}_{Q+iP,\nu',\tilde{\nu}'} \rangle_{L^2(\Omega_\mp)} = \delta_{|\nu|,|\nu'|} \delta_{|\tilde{\nu}|,|\tilde{\nu}'|} F_{Q+iP}(\nu, \nu') F_{Q+iP}(\tilde{\nu}, \tilde{\nu}')$$

where the matrices $(F_{Q+iP}(\nu, \nu'))_{\nu,\nu' \in \mathcal{T}_j}$ are positive definite (hence invertible) for each $j \in \mathbb{N}$ and they depend on $P + iQ$: more precisely they are polynomials in the weight $\Delta_{Q+iP} = \frac{1}{4}(P^2 + Q^2)$ and in the central charge c_L . Item 5) then follows from the representation (4.12) and decomposing the basis $\psi_{\mathbf{k}\mathbf{l}}$ in terms of the new basis $\mathcal{Q}_{Q+iP,\nu,\nu'}$. \square

5. LIOUVILLE CFT: DYNAMICS AND QUADRATIC FORM

The goal of this section is to construct explicitly the quadratic form associated to \mathbf{H}_* and to obtain a probabilistic representation of the semigroup. This probabilistic representation, i.e. a Feynman-Kac type formula, easily follows from the definition of \mathbf{H}_* , this is done in subsection 5.1 just below. The construction of the quadratic form is more subtle, especially in the case $\gamma \in (\sqrt{2}, 2)$. Indeed we will construct the Friedrichs extension of the operator (2.11) (restricted to appropriate domain) and we call it \mathbf{H} . Then we will show that $\mathbf{H}_* = \mathbf{H}$. The main reason why the difficulty to construct the quadratic form increases with γ is due to

the interpretation of the term V in (2.11). For $\gamma \in (0, \sqrt{2})$ only, it makes sense as a GMC random variable. Indeed, define the regularized field for $k \geq 0$

$$(5.1) \quad \varphi^{(k)}(\theta) = \sum_{|n| \leq k} \varphi_n e^{in\theta}$$

which is a.s. a smooth function. Then the GMC random variable V can be defined as the following limit

$$(5.2) \quad V := \lim_{k \rightarrow \infty} V^{(k)}, \quad V^{(k)} := \int_0^{2\pi} e^{\gamma \varphi^{(k)}(\theta) - \frac{\gamma^2}{2} \mathbb{E}[\varphi^{(k)}(\theta)^2]} d\theta$$

where the above limit exists $\mathbb{P}_{\mathbb{T}}$ -almost surely and is non trivial for $\gamma \in (0, \sqrt{2})$, see [Ka85, RhVa14, Be17] for instance on the topic, in which case $V \in L^p(\Omega_{\mathbb{T}})$ for all $p < \frac{2}{\gamma^2}$. For $\gamma \in [\sqrt{2}, 2)$, the limit V vanishes, in which case we will rather make sense of the multiplication operator V as a measure (singular with respect to $\mathbb{P}_{\mathbb{T}}$): this case is more problematic, see Subsection 5.2.

5.1. Feynman-Kac formula. We consider the semigroup $e^{-t\mathbf{H}^*}$ and start by writing some form of Feynman-Kac formula for this semigroup.

Proposition 5.1. (Feynman-Kac formula) *For $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ we have*

$$(5.3) \quad e^{-t\mathbf{H}^*} f = e^{-\frac{Q^2 t}{2}} \mathbb{E}_{\varphi} [f(c + B_t, \varphi_t) e^{-\mu e^{\gamma c} \int_{\mathbb{D}_t^c} |z|^{-\gamma Q} M_{\gamma}(dz)}]$$

where $(c + B_t, \varphi_t)$ is the process on $W^s(\mathbb{T})$ defined in subsection 3.1 and $\mathbb{D}_t^c := \{z \in \mathbb{D} \mid |z| > e^{-t}\}$. In particular, for $\gamma \in (0, \sqrt{2})$

$$(5.4) \quad e^{-t\mathbf{H}^*} f = e^{-\frac{Q^2 t}{2}} \mathbb{E}_{\varphi} [f(c + B_t, \varphi_t) e^{-\mu \int_0^t e^{\gamma(c+B_s)} V(\varphi_s) ds}].$$

where

$$V(\varphi_s) := \int_0^{2\pi} e^{\gamma \varphi_s(\theta) - \frac{\gamma^2}{2} \mathbb{E}[\varphi_s(\theta)^2]} d\theta.$$

Proof. Let $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. By definition we have $e^{-t\mathbf{H}^*} f = U S_{e^{-t}} U^{-1} f$, where, from (3.26), we have $U^{-1} f = (U1)^{-1} f$ with

$$(U1)(c, \varphi) = (U_0 e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D})})(c, \varphi).$$

Now we claim that we have for $t > 0$

$$(5.5) \quad \mathbb{E}[e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_t)} | \mathcal{F}_{\mathbb{D}_t^c}] = S_{e^{-t}}(e^{Qc} U1)$$

where $\mathbb{D}_t = \{z \in \mathbb{D} \mid |z| < e^{-t}\}$. Indeed, by the domain Markov property of the GFF and conditionally on $\mathcal{F}_{\mathbb{D}_t^c}$, the law of X inside \mathbb{D}_t is given by the independent sum

$$X \stackrel{\text{law}}{=} P(s_{e^{-t}} X)(e^t) + X_{\mathbb{D}}(e^t).$$

As a consequence and using a change of variables by dilation, we get the relation conditionally on $\mathcal{F}_{\mathbb{D}_t^c}$

$$M_{\gamma}(\mathbb{D}_t) \stackrel{\text{law}}{=} S_{e^{-t}} M_{\gamma}(\mathbb{D}),$$

which gives (5.5). Next, using (5.5), we get

$$\begin{aligned} e^{-t\mathbf{H}^*} f &= U S_{e^{-t}} U^{-1} f \\ &= e^{-Qc} \mathbb{E}_{\varphi} [S_{e^{-t}}(U^{-1} f) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D})}] \\ &= e^{-Qc} \mathbb{E}_{\varphi} [\mathbb{E}[S_{e^{-t}}(U^{-1} f) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D})} | \mathcal{F}_{\mathbb{D}_t^c}]] \\ &= e^{-Qc} \mathbb{E}_{\varphi} [S_{e^{-t}}(U^{-1} f) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_t^c)} \mathbb{E}[e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_t)} | \mathcal{F}_{\mathbb{D}_t^c}]] \\ &= e^{-Qc} \mathbb{E}_{\varphi} [S_{e^{-t}}(e^{Qc} f) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_t^c)}]. \end{aligned}$$

We complete the argument by applying the Cameron-Martin formula to $S_{e^{-t}}(e^{Qc})$ to get (5.3).

In the case when $\gamma \in (0, \sqrt{2})$, we write the chaos measure in terms of the process $Z_t = (c + B_t, \varphi_t)$. The function $V : W^s(\mathbb{T}) \rightarrow \mathbb{R}^+$ given by

$$V(\varphi) = \lim_{k \rightarrow \infty} \int_{\mathbb{T}} e^{\gamma \varphi^{(k)}(\theta) - \frac{\gamma^2}{2} \mathbb{E}[\varphi^{(k)}(\theta)^2]} d\theta$$

is measurable and we get that conditionally on $\varphi = \varphi_0$ (by making the change of variables $dx = r dr d\theta = e^{-2s} ds d\theta$ with $r = |x| = e^{-s}$ and $-\frac{\gamma^2}{2} \mathbb{E}[B_s^2] - 2s = -\gamma Q_s$)

$$e^{\gamma c} M_{\gamma}(\mathbb{D}_t^c) \stackrel{law}{=} \int_0^t e^{\gamma(c+B_s-Q_s)} V(\varphi_s) ds. \quad \square$$

As a consequence of this formula and similarly to \mathbf{H}^0 we have

Proposition 5.2. *The following properties hold:*

- (1) $e^{-t\mathbf{H}^*}$ extends to a continuous semigroup on $L^p(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $p \in [1, +\infty]$ with norm $e^{-\frac{Q^2}{2}t}$ and it is strongly continuous for $p \in [1, +\infty)$.
- (2) $e^{-t\mathbf{H}^*}$ extends to a strongly continuous semigroup on $e^{-\alpha c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $\alpha \in \mathbb{R}$ with norm $e^{(\frac{\alpha^2}{2} - \frac{Q^2}{2})t}$.

Proof. Using in turn (5.4) and $V \geq 0$, we see that $\|e^{-t\mathbf{H}^*} f\|_p \leq \|e^{-t\mathbf{H}^0} |f|\|_p$ so that the claim 1) follows from Proposition 4.5. The same argument works for 2). \square

5.2. Quadratic forms and the Friedrichs extension of \mathbf{H} . Here we construct the quadratic forms associated to (2.11) and the Friedrichs extension of \mathbf{H} in the case $\gamma \in (0, 2)$.

Recall that the underlying measure on the space $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ is $dc \times \mathbb{P}_{\mathbb{T}}$, with $\Omega_{\mathbb{T}} = (\mathbb{R}^2)^{\mathbb{N}^*}$, $\Sigma_{\mathbb{T}} = \mathcal{B}^{\otimes \mathbb{N}^*}$ (where \mathcal{B} stands for the Borel sigma-algebra on \mathbb{R}^2) and the probability measure $\mathbb{P}_{\mathbb{T}}$ defined by (3.6). Also, recall the GFF on the unit circle $\varphi : \Omega \rightarrow W^s(\mathbb{T})$ (with $s < 0$) defined by (3.1). Finally recall that \mathcal{S} is the set of smooth functions depending on finitely many coordinates, i.e. of the form $F(x_1, y_1, \dots, x_n, y_n)$ with $n \geq 1$ and $F \in C^\infty((\mathbb{R}^2)^n)$, with at most polynomial growth at infinity for F and its derivatives. \mathcal{S} is dense in $L^2(\Omega_{\mathbb{T}})$.

We will construct the quadratic form associated to \mathbf{H} as a limit (in a suitable sense) of regularized quadratic forms associated with the regularized potential $V^{(k)}$.

For $k \geq 1$ and $\mu > 0$, we introduce the bilinear form (with associated quadratic form still denoted by \mathcal{Q}_k)

$$(5.6) \quad \mathcal{Q}^{(k)}(u, v) := \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(\partial_c u \partial_c \bar{v} + Q^2 u \bar{v} + 2(\mathbf{P}u) \bar{v} + 2\mu e^{\gamma c} V^{(k)} u \bar{v} \right) dc.$$

Here u, v belong to the domain $\mathcal{D}(\mathcal{Q}^{(k)})$ of the quadratic form, namely the completion for the $\mathcal{Q}^{(k)}$ -norm in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ of the space \mathcal{C} defined by (4.10). Also, it is clear that $\mathcal{D}(\mathcal{Q}^{(k)})$ embeds continuously and injectively in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ (same argument as in the proof of Prop 4.3). Also, since $V^{(k)} \geq 0$ it is clearly lower semi-bounded $\mathcal{Q}^{(k)}(u) \geq Q^2 \|u\|_2^2 / 2$ so that the construction of the Friedrichs extension then follows from [ReSi1, Theorem 8.15]. It determines uniquely a self-adjoint operator $\mathbf{H}^{(k)}$, called the *Friedrichs extension*, with domain denoted by $\mathcal{D}(\mathbf{H}^{(k)})$ such that:

$$\mathcal{D}(\mathbf{H}^{(k)}) = \{u \in \mathcal{D}(\mathcal{Q}^{(k)}); \exists C > 0, \forall v \in \mathcal{D}(\mathcal{Q}^{(k)}), \mathcal{Q}^{(k)}(u, v) \leq C \|v\|_2\}$$

and for $u \in \mathcal{D}(\mathbf{H}^{(k)})$, $\mathbf{H}^{(k)}u$ is the unique element in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ satisfying

$$\mathcal{Q}^{(k)}(u, v) = \langle \mathbf{H}^{(k)}u | v \rangle_2.$$

We also denote by $\mathbf{R}_{\lambda}^{(k)}$ the associated resolvent family. The following result is rather standard but we found no reference corresponding exactly to our context so that we give a short proof.

Proposition 5.3. *The strongly continuous contraction semigroup $(e^{-t\mathbf{H}^{(k)}})_{t \geq 0}$ of self-adjoint operators on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ obeys the Feynman-Kac formula*

$$(5.7) \quad e^{-t\mathbf{H}^{(k)}} f = e^{-\frac{Q^2 t}{2}} \mathbb{E}_{\varphi} \left[f(c + B_t, \varphi_t) e^{-\mu \int_0^t e^{\gamma(c+B_s)} V^{(k)}(\varphi_s) ds} \right],$$

where we have, after decomposing the field φ_s along its harmonics i.e. $\varphi_s(\theta) := \sum_{n \neq 0} \varphi_{s,n} e^{in\theta}$,

$$V^{(k)}(\varphi) = \int_0^{2\pi} e^{\gamma \varphi_s^{(k)}(\theta) - \frac{\gamma^2}{2} \mathbb{E}[\varphi_s^{(k)}(\theta)^2]} d\theta \quad \text{with } \varphi_s^{(k)}(\theta) := \sum_{|n| \leq k, n \neq 0} \varphi_{s,n} e^{in\theta}.$$

Proof. We use Kato's strong Trotter product formula (see [ReSi1, Theorem S.21 page 379]) applied to the self-adjoint operators \mathbf{H}_0 and $e^{\gamma c V^{(k)}}$: since the domain $\mathcal{D}(\mathcal{Q}^{(k)})$ of the quadratic form $\mathcal{Q}^{(k)}$ is dense in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and satisfies $\mathcal{D}(\mathcal{Q}^{(k)}) = \mathcal{D}(\mathcal{Q}_0) \cap \mathcal{D}(\mathcal{Q}_{V^{(k)}})$, where $\mathcal{D}(\mathcal{Q}_{V^{(k)}}) = \{f \mid \|e^{\gamma c/2} V^{(k)}\}^{1/2} f\|_2 < +\infty\}$ is the domain of the quadratic form associated to the operator of multiplication by $e^{\gamma c V^{(k)}}$, we have the identity

$$\lim_{n \rightarrow \infty} (e^{-\frac{t}{n} \mathbf{H}_0} e^{-\frac{t}{n} \mu e^{\gamma c V^{(k)}}})^n = e^{-t \mathbf{H}^{(k)}}$$

where the limit is understood in the strong sense (i.e. convergence in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ when this relation is applied to $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$). Now we compute the limit in the left-hand side. For $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ we have

$$(5.8) \quad (e^{-\frac{t}{n} \mathbf{H}_0} e^{-\frac{t}{n} \mu e^{\gamma c V^{(k)}}})^n f = e^{-Q^2 t/2} \mathbb{E}_{\varphi} [f(c + B_t, \varphi_t) e^{-\mu R_t^{n,(k)}}]$$

with $R_t^{n,(k)}$ the Riemann sum

$$R_t^{n,(k)} := \frac{t}{n} \sum_{j=1}^n e^{\gamma(c+B_{jt/n})} V^{(k)}(\varphi_{jt/n}).$$

The right-hand side of (5.8) converges in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ towards the same expression with $R_t^{n,(k)}$ replaced by $\int_0^t e^{\gamma(c+B_s)} V^{(k)}(\varphi_s) ds$: indeed this can be established by using Jensen and the fact that, almost surely, the Riemann sum $R_t^{n,(k)}$ converges almost surely (for all fixed c) towards the integral $\int_0^t e^{\gamma(c+B_s)} V^{(k)}(\varphi_s) ds$ since the process $s \mapsto e^{\gamma(c+B_s)} V^{(k)}(\varphi_s)$ is continuous. This provides the Feynman-Kac representation as claimed. \square

Now our main goal is to construct a quadratic form corresponding to the limit $k \rightarrow \infty$ of the quadratic forms $(\mathcal{Q}^{(k)}, \mathcal{D}(\mathcal{Q}^{(k)}))$. For $u, v \in \mathcal{C}$, we define

$$(5.9) \quad \mathcal{Q}(u, v) := \lim_{k \rightarrow \infty} \mathcal{Q}^{(k)}(u, v).$$

Of course, when $\gamma \in (0, \sqrt{2})$, existence of the limit is trivial as $(V^{(k)})$ converges towards V in $L^p(\Omega_{\mathbb{T}})$ for $p < 2/\gamma^2$ and $u\bar{v} \in C_c^\infty(\mathbb{R}; L^q(\Omega_{\mathbb{T}}))$ for any $q > 1$. But treating the case $\gamma \in (\sqrt{2}, 2)$ as well requires another argument: the existence of the limit is guaranteed by the Girsanov transform, namely that for $u = u(x_1, y_1, \dots, x_n, y_n)$, $v = v(x_1, y_1, \dots, x_n, y_n)$ and $k \geq n$ the term involving $V^{(k)}$ in $\mathcal{Q}^{(k)}$ can be rewritten as

$$(5.10) \quad \mathbb{E} \int_{\mathbb{R}} e^{\gamma c} V^{(k)} u \bar{v} dc = \mathbb{E} \int_{\mathbb{R}} \int_0^{2\pi} e^{\gamma c} u_{\text{shift}} \bar{v}_{\text{shift}} dc d\theta$$

where the function u_{shift} (and similarly for v) is defined by

$$u_{\text{shift}}(\theta, c, x_1, y_1, \dots, x_n, y_n) := u\left(c, x_1 + \gamma \cos(\theta), y_1 - \gamma \sin(\theta), \dots, x_n + \frac{\gamma}{\sqrt{n}} \cos(n\theta), y_n - \frac{\gamma}{\sqrt{n}} \sin(n\theta)\right).$$

Hence the term $\mathbb{E} \int_{\mathbb{R}} e^{\gamma c} V^{(k)} u \bar{v} dc$ does not depend on $k \geq n$. We also denote by $u_{\text{shift}(N)}$ the same as above except that the first N variables $(x_1, y_1, \dots, x_N, y_N)$ are shifted and other variables remain unchanged.

Also, we denote by $\mathbf{R}_{*,\lambda}$ the resolvent family associated with the Feynman-Kac semigroup $e^{-t\mathbf{H}_*}$. Let \mathcal{Q}_* denote the quadratic form associated to \mathbf{H}_* with domain $\{u \in L^2; \lim_{t \rightarrow 0} \langle u, \frac{u - e^{-t\mathbf{H}_*} u}{t} \rangle_2 < \infty\}$. Using [Sz98, Section 1.4], \mathcal{Q}_* is closed. The following lemma is fundamental: though the limit (5.9) makes sense for any value of $\gamma \in \mathbb{R}$, the fact that it can be related to the quadratic form \mathcal{Q}_* deeply relies on GMC theory, hence on the fact that $\gamma \in (0, 2)$.

Lemma 5.4. *For $u, v \in \mathcal{C}$, we have $\mathcal{Q}_*(u, v) = \mathcal{Q}(u, v)$. In particular, \mathcal{Q} is closable.*

Proof. Let $u, v \in \mathcal{C}$, say depending on the first n -th harmonics. From (5.3), we have to show that, as $t \rightarrow 0$,

$$D_t := e^{-\frac{Q^2 t}{2}} \int \mathbb{E} \left[u(c + B_t, \varphi_t) v(c, \varphi) e^{-\mu e^{\gamma c} \int_{\mathbb{D}_t^c} |z|^{-\gamma Q} M_\gamma(dz)} \right] dc = \langle u, v \rangle_2 - t \mathcal{Q}(u, v) + o(t).$$

For notational simplicity and only in this proof, let us denote $V_t := \int_{\mathbb{D}_t^c} |z|^{-\gamma Q} M_\gamma(dz)$. Then

$$\begin{aligned} D_t - \langle u, v \rangle_2 &= \langle e^{-t \mathbf{H}_0} u - u, v \rangle_2 + (1 + o(1)) \int \mathbb{E} \left[u(c + B_t, \varphi_t) v(c, \varphi) (e^{-\mu e^{\gamma c} V_t} - 1) \right] dc \\ &= -t \mathcal{Q}_0(u, v) + o(t) + (1 + o(1)) \int \mathbb{E} \left[(u(c + B_t, \varphi_t) - u(c, \varphi)) v(c, \varphi) (e^{-\mu e^{\gamma c} V_t} - 1) \right] dc \\ &\quad - (1 + o(1)) \int \mathbb{E} \left[u(c, \varphi) v(c, \varphi) (1 - e^{-\mu e^{\gamma c} V_t}) \right] dc \\ &=: -t \mathcal{Q}_0(u, v) + o(t) + (1 + o(1)) D_t^1 - (1 + o(1)) D_t^2. \end{aligned}$$

Let us show that

$$(5.11) \quad D_t^2 = t(1 + o(1)) \mu \int e^{\gamma c} \int_0^{2\pi} \mathbb{E} \left[u_{\text{shift}}(c, \varphi) v_{\text{shift}}(c, \varphi) \right] dcd\theta.$$

For this and even if it means decomposing u, v into their positive/negative parts, we may assume that u, v are non-negative. First, we use the inequality $1 - e^{-x} \leq x$ for $x \geq 0$ to get

$$\begin{aligned} D_t^2 &\leq \int e^{\gamma c} \mathbb{E} \left[u(c, \varphi) v(c, \varphi) \mu e^{\gamma c} V_t \right] c \\ &= \mu \int e^{\gamma c} \int_0^{2\pi} \int_{e^{-t}}^1 \mathbb{E} \left[e^{\gamma P \varphi(r e^{i\theta}) - \frac{\gamma^2}{2} \mathbb{E}[P \varphi(r e^{i\theta})^2]} u(c, \varphi) v(c, \varphi) \right] dcr^{1-\gamma Q} drd\theta \\ &= \mu \int e^{\gamma c} \int_0^{2\pi} \int_{e^{-t}}^1 \mathbb{E} \left[u_{\text{shift}}^r(c, \varphi) v_{\text{shift}}^r(c, \varphi) \right] dcr^{1-\gamma Q} drd\theta \end{aligned}$$

where we have used the Girsanov transform in the last line and the function u_{shift}^r is defined by

$$u_{\text{shift}}(\theta, c, x_1, y_1, \dots, x_n, y_n) := u(c, x_1 + a_1, y_1 + b_1, \dots, x_n + a_n, y_n + b_n),$$

with $a_k := \frac{\gamma}{\sqrt{k}} r^k \cos(k\theta)$ and $b_k := -\frac{\gamma}{\sqrt{k}} r^k \sin(k\theta)$. Because $u, v \in \mathcal{C}$, it is then plain to deduce that $D_t^2 \leq t(1 + o(1)) \mu \int e^{\gamma c} \int_0^{2\pi} \mathbb{E} \left[u_{\text{shift}}(c, \varphi) v_{\text{shift}}(c, \varphi) \right] dcd\theta$. Second, we use the inequality $1 - e^{-x} \geq x e^{-x}$ for $x \geq 0$ to get, using again the Girsanov transform,

$$\begin{aligned} D_t^2 &\geq \int e^{\gamma c} \mathbb{E} \left[u(c, \varphi) v(c, \varphi) \mu e^{\gamma c} V_t e^{-\mu e^{\gamma c} V_t} \right] c \\ &= \mu \int e^{\gamma c} \int_0^{2\pi} \int_{e^{-t}}^1 \mathbb{E} \left[u_{\text{shift}}^r(c, \varphi) v_{\text{shift}}^r(c, \varphi, r e) e^{-\mu e^{\gamma c} V_t^{\text{shift}}} \right] dcr^{1-\gamma Q} drd\theta \end{aligned}$$

with $V_t^{\text{shift}}(r e^{i\theta}) := \int_{\mathbb{D}_t^c} |z|^{-\gamma Q} |z - r e^{i\theta}|^{-\gamma^2} M_\gamma(dz)$. It remains to get rid of the exponential term in the last expectation. So we split the above expectation in two parts by writing $e^{-\mu e^{\gamma c} V_t^{\text{shift}}} = 1 + (e^{-\mu e^{\gamma c} V_t^{\text{shift}}} - 1)$. The first part produces $t(1 + o(1)) \mu \int e^{\gamma c} \int_0^{2\pi} \mathbb{E} \left[u_{\text{shift}}(c, \varphi) v_{\text{shift}}(c, \varphi) \right] dcd\theta$ similarly as above. For the second part corresponding to $(e^{-\mu e^{\gamma c} V_t^{\text{shift}}} - 1)$, we want to show that it is neglectable. For this, we write

$$(5.12) \quad \begin{aligned} &\mu \int e^{\gamma c} \int_0^{2\pi} \int_{e^{-t}}^1 \mathbb{E} \left[u_{\text{shift}}^r(c, \varphi) v_{\text{shift}}^r(c, \varphi) (1 - e^{-\mu e^{\gamma c} V_t^{\text{shift}}}) \right] dcr^{1-\gamma Q} drd\theta \\ &\leq \mu \int e^{\gamma c} \int_0^{2\pi} \int_{e^{-t}}^1 \mathbb{E} \left[u_{\text{shift}}^r(c, \varphi) v_{\text{shift}}^r(c, \varphi) \right]^p \mathbb{E} \left[|1 - e^{-\mu e^{\gamma c} V_t^{\text{shift}}(r e^{i\theta})}|^q \right]^{1/q} dcr^{1-\gamma Q} drd\theta \end{aligned}$$

where we have used Hölder's inequality in the last line for any fixed conjugate exponents p, q . The first expectation $c \mapsto \mathbb{E} \left[u_{\text{shift}}^r(c, \varphi) v_{\text{shift}}^r(c, \varphi) \right]^p$ is bounded uniformly with respect to r, c and has fixed compact support in c (because v has so), say $[-A, A]$ for some $A > 0$. Now we claim that the family of functions $(c, r) \mapsto f_t(c, r) := \mathbb{E} \left[|1 - e^{-\mu e^{\gamma c} V_t^{\text{shift}}(r e^{i\theta})}|^q \right]^{1/q}$ (this quantity does not depend on θ by invariance in law under rotation) converges uniformly towards 0 as $t \rightarrow 0$, from which one can easily deduce that the quantity (5.12) is $o(t)$. Since f_t is increasing in c hence monotonic for c in $[-A, A]$ it is enough to prove this claim

for a fixed c . Next, recall [DKRV16, Lemma 3.10] the condition of finiteness for integrals of GMC with singularities: for any $\alpha \in (0, Q)$, $p > 0$ and $z_0 \in \mathbb{D}$

$$(5.13) \quad \mathbb{E}\left[\left(\int_{\mathbb{D}} \frac{1}{|z - z_0|^{\gamma\alpha}} M_\gamma(dz)\right)^p\right] < +\infty \Leftrightarrow p < \frac{2}{\gamma}(Q - \alpha).$$

Taking $\alpha = \gamma$ and $z_0 = re^{i\theta}$, the finiteness of the expectation above implies that for fixed $r \in [\frac{1}{2}, 1]$ (and fixed c) $\lim_{t \rightarrow 0} f_t(c, r) = 0$ by using dominated convergence. Now we claim that for fixed t and c , the mapping $r \mapsto f_t(c, r)$ is continuous. To see this, for each $\delta > 0$ let us introduce the mapping

$$r \in [\frac{1}{2}, 1] \mapsto f_{t,\delta}(c, r) := \mathbb{E}\left[|1 - e^{-\mu e^{\gamma c} V_{t,\delta}^{\text{shift}}(re^{i\theta})}|^q\right]^{1/q}$$

where $V_{t,\delta}^{\text{shift}}$ is the potential with regularized singularity at scale δ

$$V_{t,\delta}^{\text{shift}}(re^{i\theta}) := \int_{\mathbb{D}_c^c} |z|^{-\gamma Q} (|z - re^{i\theta}| \vee \delta)^{-\gamma^2} M_\gamma(dz).$$

Obviously, $f_{t,\delta}(c, r)$ is a continuous function of the variable r . We want to show that the family $(f_{t,\delta})_\delta$ converges uniformly towards f_t as $\delta \rightarrow 0$. By using the triangular inequality

$$\begin{aligned} |f_{t,\delta}(c, r) - f_t(c, r)| &\leq \mathbb{E}\left[|e^{-\mu e^{\gamma c} V_{t,\delta}^{\text{shift}}(re^{i\theta})} - e^{-\mu e^{\gamma c} V_t^{\text{shift}}(re^{i\theta})}|^q\right]^{1/q} \\ &\leq \mu e^{\alpha\gamma c} \mathbb{E}\left[|V_t^{\text{shift}}(re^{i\theta}) - V_{t,\delta}^{\text{shift}}(re^{i\theta})|^{\alpha q}\right]^{1/q}. \end{aligned}$$

where we have used the inequality $|e^{-x} - e^{-y}| \leq |x - y|^\alpha$ for any arbitrary $\alpha \in (0, 1]$ and $x, y \geq 0$. We fix α such that $\alpha q < \frac{2}{\gamma}(Q - \gamma)$ to make sure that the expectation is finite using the criterion (5.13). By invariance in law of M_γ under translation, this quantity does not depend on r and is less than (in fact this is true when the singularity is at positive distance from the boundary: when the singularity approaches the boundary, the above quantity is strictly less than the bound below)

$$(5.14) \quad C\mathbb{E}\left[\left(\int_{|z| \leq \delta} |z|^{-\gamma^2} M_\gamma(dz)\right)^{\alpha q}\right]^{1/q}.$$

By multifractal scaling (see the proof of [DKRV16, Lemma 3.10]), (5.14) is equal $C\delta^{\gamma(Q-\gamma)\alpha - \frac{\gamma^2}{2}\alpha^2 q}$ and one can check that the exponent is positive under the condition $\alpha q < \frac{2}{\gamma}(Q - \gamma)$. This establishes the uniform convergence of $(f_{t,\delta})_\delta$ towards f_t as $\delta \rightarrow 0$ over $r \in [\frac{1}{2}, 1]$. In conclusion, for fixed c , the family $(f_t(c, \cdot))_{t>0}$ is a family of continuous functions that decrease pointwise towards 0 as $t \rightarrow 0$. Hence the convergence is uniform by the Dini theorem. So we have proved (5.11).

Similar arguments can be used to show that $D_t^1 = o(t)$. Indeed, we use again the inequality $1 - e^{-x} \leq x$ to get the bound

$$|D_t^1| \leq \mu \int e^{\gamma c} \mathbb{E}\left[|u(c + B_t, \varphi_t) - u(c, \varphi)| |v(c, \varphi)| V_t\right] dc$$

and then the Girsanov transform and Hölder to obtain

$$\begin{aligned} |D_t^1| &\leq \mu \int_0^{2\pi} \int_{e^{-t}}^1 \int e^{\gamma c} \mathbb{E}\left[|u^{\text{shift}}(t, c + B_t, \theta, \varphi_t) - u^{\text{shift}}(0, c, \theta, \varphi)| |v^{\text{shift}}(0, c, \theta, \varphi)|\right] d\theta dr dc \\ &\leq \mu \int_0^{2\pi} \int_{e^{-t}}^1 \int e^{\gamma c} \mathbb{E}\left[|u^{\text{shift}}(t, c + B_t, \theta, \varphi_t) - u^{\text{shift}}(0, c, \theta, \varphi)|^p\right]^{1/p} \mathbb{E}\left[|v^{\text{shift}}(0, c, \theta, \varphi)|^q\right]^{1/q} d\theta dr dc \end{aligned}$$

with

$$u^{\text{shift}}(t, c, \theta, x_1, y_1, \dots, x_n, y_n) := u\left(c, x_1 + a_1, y_1 + b_1, \dots, x_n + a_n, y_n + b_n\right)$$

and

$$a_k := \frac{\gamma k^{1/2}}{\pi} \int_0^{2\pi} \ln \frac{1}{|e^{-t+i\theta'} - e^{-r+i\theta}|} \cos(k\theta') d\theta', \quad b_k := \frac{\gamma k^{1/2}}{\pi} \int_0^{2\pi} \ln \frac{1}{|e^{-t+i\theta'} - e^{-r+i\theta}|} \sin(k\theta) d\theta.$$

Again, the second expectation has compact support in c whereas the second satisfies $\sup_{r \leq t} \mathbb{E} \left[|u^{\text{shift}}(t, c + B_t, \theta, \varphi_t) - u^{\text{shift}}(0, c, \theta, \varphi)|^p \right]^{1/p} \rightarrow 0$ as $t \rightarrow 0$ since $u \in \mathcal{C}$ and B_t and the first n -th harmonics of the field $(\varphi_t)_t$ are continuous. One can then easily conclude. \square

From the above lemma, the quadratic form $(\mathcal{C}, \mathcal{Q})$ is closable. Let $\mathcal{D}(\mathcal{Q})$ be the completion of \mathcal{C} for the \mathcal{Q} -norm. The completion is the vector space consisting of equivalence classes of Cauchy sequences of \mathcal{C} for the norm $\|u\|_{\mathcal{Q}} := \sqrt{\mathcal{Q}(u, u)}$ under the equivalence relation $u \sim v$ iff $\|u_n - v_n\|_{\mathcal{Q}} \rightarrow 0$ as $n \rightarrow \infty$. This space is a Hilbert space. It can be identified with the closure of the space of quadruples $\{(u, \partial_c u, \mathbf{P}^{1/2} u, u_{\text{shift}}); u \in \mathcal{C}\}$ in $(L^2)^3 \times L^2(\mathbb{R} \times \Omega_{\mathbb{T}} \times [0, 2\pi], e^{\gamma c} dc \otimes \mathbb{P}_{\mathbb{T}} \otimes d\theta)$. $\mathcal{D}(\mathcal{Q})$ embeds injectively in L^2 as it is closed. Now we want to show that

Proposition 5.5. *For $\gamma \in (0, 2)$, the quadratic form $(\mathcal{D}(\mathcal{Q}), \mathcal{Q})$ defines an operator \mathbf{H} (the Friedrichs extension), which satisfies $\mathbf{H}_* = \mathbf{H}$.*

Proof. Take $F \in \mathcal{C}$ and consider $u := \mathbf{R}_{*, \lambda} F$ (resp. $u_k := \mathbf{R}_{\lambda}^{(k)} F$) for $\lambda > 0$. We will repeatedly use below the fact that for k large enough $u_{k, \text{shift}} = u_{k, \text{shift}(k)}$ since $F \in \mathcal{C}$. Indeed, this follows from the Feynman-Kac formula (by Prop. 5.3)

$$(5.15) \quad u_k = \int_0^\infty e^{-\lambda t} e^{-t\mathbf{H}^{(k)}} f dt = \int_0^\infty e^{-(\lambda + \frac{Q^2}{2})t} \mathbb{E}_\varphi [f(c + B_t, \varphi_t) e^{-\mu \int_0^t e^{\gamma(c+B_s)} V^{(k)}(\varphi_s) ds}] dt.$$

With this expression, one can see that if f depends on the first N harmonics of the field φ then for $k \geq N$ we have $u_{k, \text{shift}} = u_{k, \text{shift}(k)}$.

The first step of the proof is to observe that

$$u_k \rightarrow u \quad \text{as } k \rightarrow \infty \quad \text{in } L^2(\mathbb{R} \times \Omega_{\mathbb{T}}).$$

This follows from the Feynman-Kac representation (5.15)+(5.3) and standard GMC theory that ensures that $\int_0^t e^{\gamma(c+B_s)} V^{(k)}(\varphi_s) ds$ converges almost surely towards $e^{\gamma c} \int_{\mathbb{D}_c^+} |z|^{-\gamma Q} M_\gamma(dz)$ for $\gamma \in (0, 2)$. Furthermore, since

$$\lambda \|u_k\|_2^2 + \mathcal{Q}^{(k)}(u_k) = \langle F, u_k \rangle_2$$

we deduce using Cauchy-Schwartz in the r.h.s. that

$$(5.16) \quad \sup_k \mathcal{Q}^{(k)}(u_k) < +\infty.$$

This entails that the sequences $(\partial_c u_k)_k$ and $(\mathbf{P}^{1/2} u_k)_k$ weakly converge up to subsequences in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and, by (5.10), that $(u_{k, \text{shift}})_k$ weakly converges in $e^{-\gamma c/2} L^2([0, 2\pi] \times \mathbb{R} \times \Omega_{\mathbb{T}})$. Strong convergence of $(u_k)_k$ towards u and weak convergence of $(\partial_c u_k)_k$ and $(\mathbf{P}^{1/2} u_k)_k$ implies that their respective weak limits must be $\partial_c u$ and $\mathbf{P}^{1/2} u$. The resolvent equation associated to u_k reads for $v \in \mathcal{C}$

$$(5.17) \quad \lambda \langle u_k, v \rangle_2 + \mathcal{Q}^{(k)}(u_k, v) = \langle F, v \rangle_2.$$

Denote by z a possible weak limit of $(u_{k, \text{shift}})_k$ in $e^{-\gamma c/2} L^2([0, 2\pi] \times \mathbb{R} \times \Omega_{\mathbb{T}})$. Passing to the limit in $k \rightarrow \infty$ (up to appropriate subsequence) produces

$$(5.18) \quad \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(\partial_c u \partial_c \bar{v} + (Q^2 + 2\lambda) u \bar{v} + 2(\mathbf{P}^{1/2} u) \overline{\mathbf{P}^{1/2} v} + 2\mu \int_0^{2\pi} e^{\gamma c} z \bar{v}_{\text{shift}} d\theta \right) dc = \langle F, v \rangle_2.$$

Taking $v = u_k$ and passing to the limit as $k \rightarrow \infty$, we get

$$(5.19) \quad \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(|\partial_c u|^2 + (Q^2 + 2\lambda) |u|^2 + 2|\mathbf{P}^{1/2} u|^2 + 2\mu \int_0^{2\pi} e^{\gamma c} |z|^2 d\theta \right) dc = \langle F, u \rangle_2.$$

By weak limit we have

$$\begin{aligned} & \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(|\partial_c u|^2 + (Q^2 + 2\lambda) |u|^2 + 2|\mathbf{P}^{1/2} u|^2 + 2\mu \int_0^{2\pi} e^{\gamma c} |z|^2 d\theta \right) dc \\ & \leq \liminf_k \frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(|\partial_c u_k|^2 + (Q^2 + 2\lambda) |u_k|^2 + 2|\mathbf{P}^{1/2} u_k|^2 + 2\mu \int_0^{2\pi} e^{\gamma c} |u_{k, \text{shift}}|^2 d\theta \right) dc. \end{aligned}$$

Also

$$\frac{1}{2} \mathbb{E} \int_{\mathbb{R}} \left(|\partial_c u_k|^2 + (Q^2 + 2\lambda) |u_k|^2 + 2|\mathbf{P}^{1/2} u_k|^2 + 2\mu \int_0^{2\pi} e^{\gamma c} |u_{k,\text{shift}}|^2 d\theta \right) dc = \langle F, u_k \rangle_2 \rightarrow \langle F, u \rangle_2$$

as $k \rightarrow \infty$. This shows that the weak convergence of the sequence $(u_k, \partial_c u_k, \mathbf{P}^{1/2} u_k, u_{k,\text{shift}})_k$ actually holds in the strong sense. Also, it is easy to check from (5.18) that the limit is unique. This implies the convergence of $(u_k)_k$ in \mathcal{Q} -norm toward u , which thus belongs to $\mathcal{D}(\mathcal{Q})$. Hence $\mathbf{R}_{*,\lambda}$ maps \mathcal{C} into $\mathcal{D}(\mathcal{Q}) \subset L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and it coincides on \mathcal{C} with the resolvent associated to \mathcal{Q} . Since \mathcal{C} is dense in $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, this shows that both resolvent families coincide, hence their semigroups, quadratic forms and generators too. \square

Again, we stress that $\mathcal{D}(\mathbf{H}) = \{u \in \mathcal{D}(\mathcal{Q}) \mid \mathbf{H}u \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})\}$ and $\mathbf{H}^{-1} : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H})$ is bounded. Furthermore, by the spectral theorem, \mathbf{H} generates a strongly continuous contraction semigroup of self-adjoint operators $(e^{-t\mathbf{H}})_{t \geq 0}$ on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$.

If we let $\mathcal{D}(\mathcal{Q})'$ be the dual to $\mathcal{D}(\mathcal{Q})$ (i.e. the space of bounded conjugate linear functionals on $\mathcal{D}(\mathcal{Q})$), the injection $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \subset \mathcal{D}(\mathcal{Q})'$ is continuous and the operator \mathbf{H} can be extended as a bounded isomorphism

$$\mathbf{H} : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})'.$$

Remark 5.6. *By adapting some argument in [ReSi1, ReSi2] one can prove that \mathbf{H} is essentially self-adjoint on $\mathcal{D}(\mathbf{H}^0) \cap \mathcal{D}(e^{\gamma c} V)$ for $\gamma \in (0, 1)$, condition that ensures that the potential V is in $L^2(\Omega_{\mathbb{T}})$.*

6. SCATTERING OF THE LIOUVILLE HAMILTONIAN

In this section, we develop the scattering theory for the operator \mathbf{H} on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ with underlying measure $dc \otimes \mathbb{P}_{\mathbb{T}}$ (where \mathbf{H} is the generator of the dilation semigroup studied above). This operator has continuous spectrum and cannot be diagonalized with a complete set of $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ -eigenfunctions. We will rather use a stationary approach for this operator, in a way similar to what has been done in geometric scattering theory for manifolds with cylindrical ends in [Gu89, Me93]. The goal is to obtain a spectral resolution for \mathbf{H} in terms of generalized eigenfunctions, which will be shown to be analytic in the spectral parameter. In other words, we search to write the spectral measure of \mathbf{H} using these generalized eigenfunctions, which are similar to plane waves $(e^{i\lambda\omega \cdot x})_{\lambda \in \mathbb{R}, \omega \in S^{n-1}}$ in Euclidean scattering for the Laplacian Δ_x on \mathbb{R}^n . In our case, the generalized eigenfunctions will be functions in weighted spaces of the form $e^{-\beta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > 0$ with particular asymptotic expansions at $c = -\infty$. Let us explain briefly the simplest one, corresponding to the functions $\Psi_{\alpha} := \Psi_{\alpha,0,0} = U(V_{\alpha}(0))$ defined in the Introduction and represented probabilistically by (7.5) using the unitary map (3.25) when $\alpha < Q$ is real. For $\alpha \in (Q - \gamma/2, Q)$, they will be the only eigenfunctions of \mathbf{H} in $e^{-\beta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > Q - \alpha$ satisfying

$$(\mathbf{H} - 2\Delta_{\alpha})\Psi_{\alpha} = 0, \quad \Psi_{\alpha} = e^{(\alpha-Q)c} + u, \quad \text{with } u \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}}).$$

Similarly, we will construct (in Section 6.5) some eigenfunctions $\Psi_{\alpha,\mathbf{k},\mathbf{l}}$ of \mathbf{H} with eigenvalue $2\Delta_{\alpha} + \lambda_{\mathbf{k},\mathbf{l}}$, with $\Psi_{\alpha,\mathbf{k},\mathbf{l}}|_{\mathbb{R}^+} \in L^2$ and asymptotic (for all $\beta \in (0, \gamma/2)$)

$$\Psi_{\alpha,\mathbf{k},\mathbf{l}}|_{\mathbb{R}^-} = e^{(\alpha-Q)c} \psi_{\mathbf{k},\mathbf{l}} + u, \quad \text{with } u \in e^{(\alpha-Q+\beta)c} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}),$$

where we recall that $\lambda_{\mathbf{k},\mathbf{l}} \in \mathbb{N}$ are the eigenvalues of P defined in (4.7) and the corresponding eigenfunctions $\psi_{\mathbf{k},\mathbf{l}}$ of P appear in (4.5). One way to construct the Ψ_{α} , and similarly for $\Psi_{\alpha,\mathbf{k},\mathbf{l}}$, is to take the limit (see Proposition 6.9)

$$\Psi_{\alpha} = \lim_{t \rightarrow \infty} e^{-t\mathbf{H}} (e^{2t\Delta_{\alpha}} e^{(\alpha-Q)c}),$$

where we observe that $e^{-t\mathbf{H}^0} e^{(\alpha-Q)c} = e^{-2t\Delta_{\alpha}} e^{(\alpha-Q)c}$ so that, formally speaking, Ψ_{α} is the limit of the intertwining $e^{-t\mathbf{H}} e^{t\mathbf{H}^0} (e^{(\alpha-Q)c})$ as $t \rightarrow +\infty$. An alternative expression is to write them as

$$(6.1) \quad \Psi_{\alpha} = e^{(\alpha-Q)c} \chi(c) - (\mathbf{H} - 2\Delta_{\alpha})^{-1} (\mathbf{H} - 2\Delta_{\alpha}) (e^{(\alpha-Q)c} \chi(c))$$

where $\chi \in C^{\infty}(\mathbb{R})$ equal to 1 near $-\infty$ and 0 near $+\infty$ (see (6.20) and Lemma 6.7); here we notice that $e^{(\alpha-Q)c} \chi(c)$ is not $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ but one can check that $(\mathbf{H} - 2\Delta_{\alpha}) (e^{(\alpha-Q)c} \chi(c)) \in L^2$ if $\gamma > Q - \alpha$ (and $\gamma < 1$) so that we can apply the resolvent $\mathbf{R}(\alpha) := (\mathbf{H} - 2\Delta_{\alpha})^{-1}$ to it, and $(\mathbf{H} - 2\Delta_{\alpha})^{-1} (\mathbf{H} - 2\Delta_{\alpha}) (e^{(\alpha-Q)c} \chi(c)) \in L^2$ is not equal to $(e^{(\alpha-Q)c} \chi(c))$. Our goal will be to extend analytically these Ψ_{α} (and actually the whole

family of generalized eigenstates denoted by $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ to $\operatorname{Re}(\alpha) \leq Q$ and in particular to the line $\alpha \in Q + i\mathbb{R}$ corresponding to the spectrum of \mathbf{H} . To perform this, we see that we need to extend analytically the resolvent operator $\mathbf{R}(\alpha)$ to $\operatorname{Re}(\alpha) \leq Q$, which will be the main part of this section. In fact, we shall show that $\mathbf{R}(\alpha)$ extends analytically on an open set of a Riemann surface covering the complex plane, containing the real half-plane $\operatorname{Re}(\alpha) \leq Q$. We note that the functions $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ will be expressed as the elements in the range of some Poisson operator denoted $\mathcal{P}(\alpha)$, mapping (some subspaces of) $L^2(\Omega_{\mathbb{T}})$ to weighted spaces $e^{-\beta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for some $\beta > 0$ depending on α . The results proved here hold in some cases for geometric scattering in finite dimension [Gu89, Me93], but we are not aware of some results of this type in quantum field theory where the base space is infinite dimensional. The main difficulty will be to deal with the fact that the perturbation V is quite singular (even for $\gamma < \sqrt{2}$ where V is a potential, it is not bounded) and the fact that the eigenfunctions of \mathbf{P} (Hermite polynomials) have $L^p(\Omega_{\mathbb{T}})$ norms which blow up very fast in terms of their eigenvalues.

In this section, we shall start by describing the resolvent of \mathbf{H} in the *probabilistic region* $\{\operatorname{Re}((\alpha - Q)^2) > \beta^2\}$ acting on weighted spaces $e^{\beta c} L^2$ for $\beta \in \mathbb{R}$ and deduce the construction of the $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ in this region. Next, we will show that the resolvent $\mathbf{R}(\alpha)$ admits an analytic extension in a neighborhood of $\{\operatorname{Re}(\alpha) \leq Q\}$ (for α on some Riemann surface Σ). We shall use these results to prove the analytic continuation of the $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ to $\operatorname{Re}(\alpha) \leq Q$ and we shall finally construct the scattering operator $\mathbf{S}(\alpha)$ in Section 6.4 and write the spectral decomposition of \mathbf{H} in terms of the $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ in Theorem 6.25 (written in terms of Poisson operator in this section).

In what follows, we will mostly consider the L^2 (or L^p) spaces on $\Omega_{\mathbb{T}}$ or on $\mathbb{R} \times \Omega_{\mathbb{T}}$ respectively equipped with the measure $\mathbb{P}_{\mathbb{T}}$ or $dc \otimes \mathbb{P}_{\mathbb{T}}$, which we will denote by $L^2(\Omega_{\mathbb{T}})$ or $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for short. When the space is omitted, i.e. we simply write L^2 , this means that we consider $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$: this will relieve notations in some latter part of the paper. Recall that we denote by $\langle \cdot | \cdot \rangle_2$ the standard scalar product associated to $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}, dc \otimes \mathbb{P}_{\mathbb{T}})$ and $\| \cdot \|_2$ the associated norm; in general our scalar products will always be complex linear in the left component and anti-linear in the right component. Given two normed vector space E and F , the space of continuous linear mappings from E into F will be denoted by $\mathcal{L}(E, F)$ and when $E = F$ we will simply write $\mathcal{L}(E)$. The corresponding operator norms will be denoted by $\| \cdot \|_{\mathcal{L}(E, F)}$ or $\| \cdot \|_{\mathcal{L}(E)}$.

6.1. The operators $\mathbf{H}, \mathbf{H}^0, e^{\gamma c} V$. The operator \mathbf{H} is made up of several pieces. The first piece is the operator \mathbf{P} defined in (4.4), which is a self-adjoint non-negative unbounded operator on $L^2(\Omega_{\mathbb{T}})$. It has discrete spectrum $(\lambda_{\mathbf{k}, \mathbf{l}})_{\mathbf{k}, \mathbf{l} \in \mathcal{N}}$, but to simplify the indexing we shall order them in increasing order (without counting multiplicity) and denote them by

$$\sigma(\mathbf{P}) = \{\lambda_j \mid j \in \mathbb{N}, \lambda_j < \lambda_{j+1}\}.$$

We denote by

$$(6.2) \quad E_k := \{F \in L^2(\Omega_{\mathbb{T}}) \mid 1_{[0, \lambda_k]}(\mathbf{P})F = F\} = \bigoplus_{j \leq k} \ker(\mathbf{P} - \lambda_j)$$

the sum of eigenspaces with eigenvalues less or equal to λ_k and $\Pi_k : L^2(\Omega_{\mathbb{T}}) \rightarrow E_k$ the orthogonal projection. We also will use the important fact

$$(6.3) \quad E_k \subset L^p(\Omega_{\mathbb{T}}), \quad \forall p < \infty.$$

The quadratic form associated to \mathbf{H} is the form \mathcal{Q} defined in Subsection 5.2, with domain $\mathcal{D}(\mathcal{Q})$. We will consider the self-adjoint extension associated of \mathbf{H} obtained using the quadratic form \mathcal{Q} . The operator $\mathbf{H}^0 : \mathcal{D}(\mathbf{H}^0) \rightarrow L^2$ corresponds to the case $\mu = 0$, its quadratic form has domain $\mathcal{D}(\mathcal{Q}_0)$ containing $\mathcal{D}(\mathcal{Q})$. We recall that

$$\mathbf{H} : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q}), \quad \mathbf{H}^0 : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$$

if $\mathcal{D}'(\mathcal{Q})$ is the dual of $\mathcal{D}(\mathcal{Q})$. We then define $e^{\gamma c} V : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$ by the equation

$$(6.4) \quad \mathbf{H} = \mathbf{H}^0 + e^{\gamma c} V = -\frac{1}{2} \partial_c^2 + \frac{1}{2} Q^2 + \mathbf{P} + e^{\gamma c} V.$$

We will define the associated quadratic form

$$\mathcal{Q}_{e^{\gamma c}V} := \mathcal{Q} - \mathcal{Q}_0$$

defined on $\mathcal{D}(\mathcal{Q})$ and remark that

$$\mathcal{Q}_{e^{\gamma c}V}(u, u) = \int_{\mathbb{R}} e^{\gamma c} \mathcal{Q}_V(u, u) dc$$

for some well-defined quadratic form \mathcal{Q}_V , defined on a domain $\mathcal{D}(\mathcal{Q}_V) \subset L^2(\Omega_{\mathbb{T}})$ containing E_k for each $k \geq 0$.

For our study of the resolvent of \mathbf{H} , we need to introduce the orthogonal projection

$$\Pi_k : L^2(\Omega_{\mathbb{T}}) \rightarrow E_k,$$

extended trivially in the c variable as $\Pi_k : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R}; E_k)$.

Lemma 6.1. *Let $\chi \in L^\infty(\mathbb{R}) \cap C^\infty(\mathbb{R})$ with support in $(-\infty, A)$ for some $A \in \mathbb{R}$ and $\chi' \in L^\infty(\mathbb{R})$. Then, for all $\beta \geq -\gamma/2$ and $\beta' \in \mathbb{R}$, the following operators are bounded*

$$\begin{aligned} \chi \Pi_k &: \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q}), & \chi \Pi_k &: \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q}) \\ \chi e^{\gamma c} V \Pi_k &: e^{\beta' c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{(\beta' - \beta)c} \mathcal{D}'(\mathcal{Q}), \\ \chi e^{\gamma c} \Pi_k V &: e^{(\beta - \beta')c} \mathcal{D}(\mathcal{Q}) \rightarrow e^{-\beta' c} L^2(\mathbb{R}; E_k), \\ \chi \Pi_k e^{\gamma c} V \Pi_k &: e^{\beta' c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{(\beta' - \beta)c} L^2(\mathbb{R}; E_k). \end{aligned}$$

If $\beta > -\gamma/2$ then one also has

$$\chi e^{\gamma c} V \Pi_k : e^{\beta' c} L^\infty(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{(\beta' - \beta)c} \mathcal{D}'(\mathcal{Q}).$$

Proof. For the first operator $\chi \Pi_k$, it suffices to check that for $u \in L^2((-\infty, A); E_k)$, $\mathcal{Q}_{e^{\gamma c}V}(u) \leq C_k \|u\|_{L^2}^2$. Since there is $C_k > 0$ depending on k such that for all $F \in E_k$, $\mathcal{Q}_V(F) \leq C_k \|F\|_{L^2(\Omega_{\mathbb{T}})}^2$, we have

$$\mathcal{Q}_{e^{\gamma c}V}(u) \leq C_k \int_{-\infty}^A e^{\gamma c} \|u\|_{L^2(\Omega_{\mathbb{T}})}^2 dc \leq C_{k,A} \|u\|_{L^2(\mathbb{R} \times \Omega_{\mathbb{T}})}^2.$$

The extension of $\chi \Pi_k$ to $\mathcal{D}'(\mathcal{Q})$ follows by duality, using that $\Pi_k^* = \Pi_k$ on L^2 . Next, we analyze $\chi e^{\gamma c} V \Pi_k$. It suffices to deal with the case $\beta' = 0$ since $e^{\beta' c}$ commutes with $e^{\gamma c} V$. Using that for $F \in L^2(\mathbb{R}; E_k)$ and $\chi \in L^\infty(\mathbb{R})$ with support in \mathbb{R}^- , we have by Cauchy-Schwarz that for all $u \in \mathcal{D}(\mathcal{Q})$

$$\begin{aligned} |\langle \chi(c) e^{(\gamma + \beta)c} V F, u \rangle| &\leq \int_{-\infty}^A \chi(c) e^{(\gamma + \beta)c} |\mathcal{Q}_V(F, u)| dc \leq \|\chi\|_{L^\infty} \left(\int_{-\infty}^A e^{\gamma c} \mathcal{Q}_V(u) dc \right)^{\frac{1}{2}} \left(\int_{-\infty}^A e^{(\gamma + 2\beta)c} \mathcal{Q}_V(F) dc \right)^{\frac{1}{2}} \\ &\leq C_{k,A} \|\chi\|_{L^\infty} \|F\|_{L^2} \mathcal{Q}(u)^{1/2} \end{aligned}$$

for some constant $C_{k,A} > 0$, provided $\gamma + 2\beta \geq 0$. The same argument also holds for $F \in L^\infty(\mathbb{R}; E_k)$. Finally, $\chi \Pi_k e^{\gamma c} V = \chi e^{\gamma c} \Pi_k V$ makes sense as a map $e^{\beta c} \mathcal{D}(\mathcal{Q}) \rightarrow e^{\beta c} \mathcal{D}'(\mathcal{Q})$ using the boundedness $\Pi_k : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$. To prove that it actually maps to L^2 if $\gamma + 2\beta > 0$, we note that for all $u' \in \mathcal{C}$

$$|\langle \chi \Pi_k e^{\gamma c} V u, u' \rangle| = |\langle \chi e^{\gamma c} V u, \Pi_k u' \rangle| = \left| \int_{-\infty}^A \chi(c) e^{\gamma c} \mathcal{Q}_V(u, \Pi_k(u')) dc \right| \leq C_{k,A} \|\chi\|_{L^\infty} \mathcal{Q}(e^{-\beta c} u)^{1/2} \|u'\|_{L^2}$$

where the last bound is obtained as above by Cauchy-Schwarz and the bounds on \mathcal{Q}_V on E_k . For the last term $\chi \Pi_k V \Pi_k$, we take $u \in L^2(\mathbb{R} \times \Omega)$ and note that for $u' \in \mathcal{D}(\mathcal{Q})$

$$\begin{aligned} |\langle e^{\beta c} \Pi_k V \Pi_k u, u' \rangle| &= |\langle \chi e^{\beta c} \Pi_k V \Pi_k u, \Pi_k u' \rangle| \leq \left| \int_{-\infty}^A \chi(c) e^{(\gamma + \beta)c} \mathcal{Q}_V(\Pi_k u, \Pi_k u') dc \right| \\ &\leq \left| \int_{-\infty}^A \chi(c) e^{(\gamma + \beta)c} \mathcal{Q}_V(\Pi_k u)^{\frac{1}{2}} \mathcal{Q}_V(\Pi_k u')^{\frac{1}{2}} dc \right| \leq C_{k,A} \|u\|_{L^2} \|u'\|_{L^2} \end{aligned}$$

showing that $\|e^{\beta c} \Pi_k V \Pi_k u\|_2 \leq C_{k,A} \|u\|_2$. \square

First, we show a useful result for the spectral decomposition.

Lemma 6.2. *The operator \mathbf{H} does not have non-zero eigenvectors $u \in \mathcal{D}(\mathbf{H})$. If $\gamma \in (0, 1)$, the spectrum of \mathbf{H} is given by $\sigma(\mathbf{H}) = [\frac{Q^2}{2}, \infty)$ and consists of essential spectrum.*

Proof. In the case $\gamma \in (0, 1)$, the space \mathcal{C} is included in $\mathcal{D}(\mathbf{H})$. It is then easy to check that $\sigma(\mathbf{H}) = [\frac{Q^2}{2}, \infty)$ consists only of essential spectrum by using Weyl sequences $(e^{ipc}\chi(2^{-n}c)/\omega_n)_n \in \mathbb{N}$ where $\chi \in C_c^\infty(\mathbb{R})$ have support in $[-\frac{3}{2}, -1]$ and equal to 1 on some interval, with $\omega_n = \|\chi(2^{-n}\cdot)\|_{L^2(\mathbb{R})}$.

Let $u \in \mathcal{D}(\mathbf{H})$ such that $\mathbf{H}u = \lambda u$ with $\lambda \in [\frac{Q^2}{2}, \infty)$. Then $u \in \mathcal{D}(\mathcal{Q})$ (hence $\partial_c u \in L^2$), and it satisfies $\mathcal{Q}(u, v) = \langle \lambda u | v \rangle_2$ for all $v \in \mathcal{D}(\mathcal{Q})$. Now we claim

Lemma 6.3. *Assume we are given $f \in L^2$ such that $\partial_c f \in L^2$. Consider $u \in \mathcal{D}(\mathcal{Q})$ such that*

$$(6.5) \quad \mathcal{Q}(u, v) = \langle f | v \rangle_2, \quad \forall v \in \mathcal{D}(\mathcal{Q}).$$

Then $\partial_c u \in \mathcal{D}(\mathcal{Q})$ and

$$(6.6) \quad \mathcal{Q}(\partial_c u, v) = \langle \partial_c f | v \rangle_2 - \gamma \int e^{\gamma c} \int_0^{2\pi} \mathbb{E}[u_{\text{shift}} v_{\text{shift}}] d\theta dc, \quad \forall v \in \mathcal{D}(\mathcal{Q}).$$

We postpone the proof of this lemma and conclude first. Consider next (6.5) with $f = \lambda u$ and choose $v = \partial_c u \in \mathcal{D}(\mathcal{Q})$ to obtain $\mathcal{Q}(u, \partial_c u) = \langle \lambda u | \partial_c u \rangle_2 = 0$. Also, choosing $v = u$ in (6.6) we obtain $\mathcal{Q}(\partial_c u, u) = \langle \lambda \partial_c u | u \rangle_2 - \gamma \int e^{\gamma c} \int_0^{2\pi} \mathbb{E}|u_{\text{shift}}|^2 d\theta dc$. These relations imply $\gamma \int e^{\gamma c} \int_0^{2\pi} \mathbb{E}|u_{\text{shift}}|^2 d\theta dc = 0$. In the case when $\gamma \in (0, \sqrt{2})$ then V exists as a fairly defined function and this relation translates into $\|e^{\gamma c/2} V^{1/2} u\|_2 = 0$. Hence $u = 0$ as $V > 0$ almost surely. In the general situation $\gamma \in (0, 2)$, the argument is as follows: the relation $\gamma \int e^{\gamma c} \int_0^{2\pi} \mathbb{E}|u_{\text{shift}}|^2 d\theta dc = 0$ implies that $\mathcal{Q}(u, v) = \mathcal{Q}_0(u, v)$ for all $v \in \mathcal{C}$. Therefore, $u \in \mathcal{D}(\mathbf{H}_0)$ and u is an eigenfunction \mathbf{H}_0 , hence $u = 0$ (cf. Remark 4.6). \square

Proof of Lemma 6.3. For $h > 0$, introduce the translation operator $T_h : L^2 \rightarrow L^2$ by $T_h v := v(c + h, \cdot)$ and the discrete derivative operator $D_h : L^2 \rightarrow L^2$ by $D_h v := (T_h v - v)/h$. Note that T_h maps $\mathcal{D}(\mathcal{Q})$ into itself, that $\|D_h v\|_2 \leq \|\partial_c v\|_2$, $D_h u_{\text{shift}} = (D_h u)_{\text{shift}}$ and we have the discrete IPP $\langle D_h u | v \rangle_2 = -\langle u | D_{-h} v \rangle_2$ for all $u, v \in L^2$. Now we can replace v by $D_{-h} v$ in (6.5) and use discrete IPP to obtain

$$(6.7) \quad \mathcal{Q}(D_h u, v) = \langle D_h f, v \rangle_2 - \int \int_0^{2\pi} \mathbb{E}[(T_h e^{\gamma c} - e^{\gamma c}) D_h u_{\text{shift}} v_{\text{shift}}] d\theta dc$$

$$(6.8) \quad - \int \int_0^{2\pi} \mathbb{E}[D_h(e^{\gamma c}) u_{\text{shift}} v_{\text{shift}}] d\theta dc, \quad \forall v \in \mathcal{D}(\mathcal{Q}).$$

Next we choose $v = D_h u$ and, using the inequality $ab \leq \frac{\epsilon}{2} a^2 + \frac{1}{2\epsilon} b^2$ for arbitrary $\epsilon > 0$, we obtain the a priori estimate (for some constant $C > 0$ depending only on γ)

$$(6.9) \quad \begin{aligned} \mathcal{Q}(D_h u, D_h u) &\leq \frac{1}{2} (\|D_h f\|_2^2 + \|D_h u\|_2^2) - (e^{\gamma h} - 1) \int \int_0^{2\pi} \mathbb{E}(e^{\gamma c} |D_h u_{\text{shift}}|^2) d\theta dc \\ &\quad + \frac{1}{2} \left| \frac{e^{\gamma h} - 1}{h} \right| \left(\epsilon^{-1} \int \int_0^{2\pi} \mathbb{E}(e^{\gamma c} |u_{\text{shift}}|^2) d\theta dc + \epsilon \int \int_0^{2\pi} \mathbb{E}(e^{\gamma c} |D_h u_{\text{shift}}|^2) d\theta dc \right) \\ &\leq (C + \epsilon^{-1}) (\|\partial_c f\|_2^2 + \mathcal{Q}(u, u)) + C(\epsilon + h) \mathcal{Q}(D_h u, D_h u) \end{aligned}$$

for all $h > 0$ small, and therefore the last term can be absorbed in the left hand side if $\epsilon, h > 0$ are small enough. Then, writing (6.7) for h and h' , subtracting and then choosing $v = D_h u - D_{h'} u$, we find

$$\mathcal{Q}(D_h u - D_{h'} u, D_h u - D_{h'} u) \leq C (\|D_h f - D_{h'} f\|_2^2 + |h - h'| \mathcal{Q}(u, u) + |h - h'| \mathcal{Q}(D_{(h-h')} u, D_{(h-h')} u)).$$

Using (6.9) with h replaced by $h - h'$ to bound the last term, we obtain that the sequence $(D_h u)_h$ is Cauchy for the \mathcal{Q} -norm. Hence the limit $\partial_c u$ belongs to $\mathcal{D}(\mathcal{Q})$. \square

Remark 6.4. *When $\gamma \in [1, 2)$ the spectrum is also $[Q^2/2, \infty)$ and is made of essential spectrum, and this will follow from our analysis of the resolvent: in fact we will show for $\gamma \in (0, 2)$ that the spectrum of \mathbf{H} is absolutely continuous.*

6.2. Resolvent of \mathbf{H} . To describe the spectral measure of \mathbf{H} and construct its generalized eigenfunctions, the main step is to understand the resolvent of \mathbf{H} as a function of the spectral parameter, in particular when the spectral parameter approach the spectrum. Due to the fact that the spectrum of \mathbf{H} starts at $Q^2/2$, it is convenient to use the spectral parameter $2\Delta_\alpha$ where $\Delta_\alpha = \frac{\alpha}{2}(Q - \frac{\alpha}{2})$ and $\alpha \in \mathbb{C}$. That way we have with $\alpha = Q + ip$

$$\mathbf{H} - 2\Delta_\alpha = -\frac{1}{2} \partial_c^2 + \mathbf{P} + e^{\gamma c} V - \frac{1}{2} p^2$$

where $p \in \mathbb{R}$ plays the role of a frequency: in particular $2\Delta_\alpha \in [Q^2/2, \infty)$ if and only if $p \in \mathbb{R}$. The half-plane $\{\alpha \in \mathbb{C} \mid \operatorname{Re}(\alpha) < Q\}$ is mapped by Δ_α to the resolvent set $\mathbb{C} \setminus [Q^2/2, \infty)$ of \mathbf{H} and will be called the *physical sheet*. By the spectral theorem

$$\mathbf{R}(\alpha) = (\mathbf{H} - 2\Delta_\alpha)^{-1} : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H})$$

is bounded if $\operatorname{Re}(\alpha) < Q$. Our goal is to extend this resolvent up to the line $\operatorname{Re}(\alpha) = Q$ analytically, and we will actually do it in an even larger region. The price to pay is that $\mathbf{R}(\alpha)$ will not be bounded on L^2 but rather on certain weighted L^2 spaces, where the weights are $e^{\beta c}$ in the region $c \leq -1$, with $\beta \in \mathbb{R}$ tuned with respect to α .

Resolvent and propagator on weighted spaces in the probabilistic region. Our first task is to understand the resolvent on weighted spaces in a subregion of $\operatorname{Re}(\alpha) < Q$, that we call the *probabilistic region* due to the fact that the resolvent can be written in terms of the semigroup $e^{-t\mathbf{H}}$.

Let $\rho : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth non-decreasing function satisfying

$$\rho(c) = c + a \text{ for } c \leq -1, \quad \rho(c) = 0 \text{ for } c \geq 0, \quad 0 \leq \rho' \leq 1$$

for some $a \in \mathbb{R}$. We have for $\beta \geq 0$ the inclusion of weighted spaces

$$e^{\beta\rho(c)} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \subset L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \subset e^{-\beta\rho(c)} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}).$$

The weighted spaces $e^{\beta\rho(c)} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ are obviously Hilbert spaces with product $\langle u, v \rangle_{e^{\beta\rho} L^2} := \langle e^{-\beta\rho} u, e^{-\beta\rho} v \rangle_2$.

Lemma 6.5. *Let $\beta \in \mathbb{R}$. If $\operatorname{Re}((\alpha - Q)^2) > \beta^2$ and $\operatorname{Re}(\alpha) < Q$, the resolvent $\mathbf{R}(\alpha) = (\mathbf{H} - 2\Delta_\alpha)^{-1}$ extends to a bounded operator*

$$\begin{aligned} \mathbf{R}(\alpha) : e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) &\rightarrow e^{-\beta\rho} \mathcal{D}(\mathbf{H}), \\ \mathbf{R}(\alpha) : e^{-\beta\rho} \mathcal{D}'(\mathcal{Q}) &\rightarrow e^{-\beta\rho} \mathcal{D}(\mathcal{Q}) \end{aligned}$$

which is analytic in α in this region. The operator $\mathbf{H} : e^{-\beta\rho} L^2 \rightarrow e^{-\beta\rho} L^2$ is closed with domain $e^{-\beta\rho} \mathcal{D}(\mathbf{H})$, it is a bijective mapping $e^{-\beta\rho} \mathcal{D}(\mathbf{H}) \rightarrow e^{-\beta\rho} L^2$ with inverse $\mathbf{R}(\alpha)$. Moreover, for $\alpha \in (-\infty, Q)$ and $0 \leq \beta < Q - \alpha$, the resolvent is bounded with norm $\|\mathbf{R}(\alpha)\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq 2((\alpha - Q)^2 - \beta^2)^{-1}$ and is equal to the integral

$$(6.10) \quad \mathbf{R}(\alpha) = \int_0^\infty e^{-t\mathbf{H} + t2\Delta_\alpha} dt$$

where $e^{-t\mathbf{H}}$ is the semigroup on $e^{-\beta\rho} L^2$ obtained by Hille-Yosida theorem with norm

$$(6.11) \quad \forall t \geq 0, \quad \|e^{-t\mathbf{H}}\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq e^{-t\frac{Q^2 - \beta^2}{2}}.$$

The integral (6.10) converges in $\mathcal{L}(e^{-\beta\rho} L^2)$ operator norm and $e^{-t\mathbf{H}} : e^{-\beta\rho} L^2 \rightarrow e^{-\beta\rho} L^2$ extends the semigroup defined in (3.31). Finally, $e^{-t\mathbf{H}} : e^{\beta\rho} \mathcal{D}(\mathcal{Q}) \rightarrow e^{\beta\rho} \mathcal{D}(\mathcal{Q})$ and $e^{-t\mathbf{H}} : e^{\beta\rho} \mathcal{D}'(\mathcal{Q}) \rightarrow e^{\beta\rho} \mathcal{D}'(\mathcal{Q})$ are bounded and for each $\varepsilon > 0$ there is some constant $C_\varepsilon > 0$ such that for all $t > 0$

$$(6.12) \quad \|e^{-t\mathbf{H}}\|_{\mathcal{L}(e^{\beta\rho} \mathcal{D}(\mathcal{Q}))} + \|e^{-t\mathbf{H}}\|_{\mathcal{L}(e^{\beta\rho} \mathcal{D}'(\mathcal{Q}))} \leq C_\varepsilon e^{-t(\frac{Q^2 - \beta^2}{2} - \varepsilon)}.$$

Proof. Consider the operator for $\beta \in \mathbb{R}$, acting on the space \mathcal{C} (defined in (4.10)) and with value in $e^{\beta\rho} \mathcal{D}'(\mathcal{Q})$,

$$\mathbf{H}_\beta := e^{\beta\rho(c)} \mathbf{H} e^{-\beta\rho(c)} = \mathbf{H} - \frac{\beta^2}{2} (\rho'(c))^2 + \frac{\beta}{2} \rho''(c) + \beta \rho'(c) \partial_c.$$

Let $u \in \mathcal{C}$, then we have (using integration by parts)

$$\begin{aligned} \operatorname{Re}(\mathbf{H}_\beta u | u)_2 &= \mathcal{Q}(u) - \frac{\beta^2}{2} \|\rho' u\|_2^2 + \frac{\beta}{2} \langle \rho'' u | u \rangle_2 + \beta \operatorname{Re}(\langle \rho' \partial_c u | u \rangle_2) \\ (6.13) \quad &= \mathcal{Q}(u) - \frac{\beta^2}{2} \|\rho' u\|_2^2 \\ &\geq \mathcal{Q}(u) - \frac{\beta^2}{2} \|u\|_2^2 \geq \frac{Q^2 - \beta^2}{2} \|u\|_2^2 + \frac{1}{2} \|\partial_c u\|_2^2 + \|\mathbf{P}^{1/2} u\|_2^2 + \mathcal{Q}_{e^{\gamma c} V}(u, u), \end{aligned}$$

where \mathcal{Q} was defined in Section 5.2. Consider the sesquilinear form $\mathcal{Q}_{\alpha, \beta}(u, v) := \langle (\mathbf{H}_\beta - 2\Delta_\alpha) u | v \rangle_2$ defined on \mathcal{C} . We easily see that if $-\operatorname{Re}(2\Delta_\alpha) > \frac{\beta^2 - Q^2}{2}$, then

$$\{u \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \mid \mathcal{Q}_{\alpha, \beta}(u, u) < \infty\} = \mathcal{D}(\mathcal{Q}).$$

Let $\mathcal{D}'(\mathcal{Q})$ be the dual of $\mathcal{D}(\mathcal{Q})$ (note that $L^2 \subset \mathcal{D}'(\mathcal{Q})$). By Lax-Milgram, if $-\operatorname{Re}(2\Delta_\alpha) > \frac{\beta^2 - Q^2}{2}$, then for each $f' \in \mathcal{D}'(\mathcal{Q})$, there is a unique $u \in \mathcal{D}(\mathcal{Q})$ such that

$$(6.14) \quad \forall v \in \mathcal{D}(\mathcal{Q}), \quad \mathcal{Q}_{\alpha,\beta}(u, v) = f'(v), \quad \mathcal{Q}(u)^{1/2} \leq C' \|f'\|_{\mathcal{D}'(\mathcal{Q})}$$

for all $v \in \mathcal{D}(\mathcal{Q})$, where $C' > 0$ depends only on $\operatorname{Re}(2\Delta_\alpha)$ and β^2 . This holds in particular for the linear form $f' : v \mapsto \langle f | v \rangle_2$ with norm $\|f'\|_{\mathcal{D}'(\mathcal{Q})} \leq C \|f\|_2$ for some $C > 0$ depending on $\operatorname{Re}(2\Delta_\alpha)$ and β^2 . We define $\tilde{\mathbf{R}}(\alpha)(e^{-\beta\rho} f) := e^{-\beta\rho} u$, this gives a bounded linear operator

$$(6.15) \quad \tilde{\mathbf{R}}(\alpha) : e^{-\beta\rho} \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho} \mathcal{D}(\mathcal{Q}) \subset e^{-\beta\rho} L^2$$

inverting the bounded operator $e^{\beta\rho}(\mathbf{H} - 2\Delta_\alpha)e^{-\beta\rho} : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$. Moreover, by (6.13), its weighted L^2 -norm is bounded by

$$(6.16) \quad \|\tilde{\mathbf{R}}(\alpha)\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq 2(\operatorname{Re}((\alpha - Q)^2 - \beta^2))^{-1} = \left(\frac{Q^2 - \beta^2}{2} - 2\operatorname{Re}(\Delta_\alpha)\right)^{-1}$$

Using $\mathcal{D}(\mathcal{Q}) \subset e^{-\beta\rho} \mathcal{D}(\mathcal{Q})$ and the uniqueness property above, this means that for $f \in L^2$, we have $\mathbf{R}(\alpha)f = \tilde{\mathbf{R}}(\alpha)f$ and thus $\tilde{\mathbf{R}}(\alpha)$ is a continuous extension of $\mathbf{R}(\alpha)$ to the Hilbert space $e^{-\beta\rho} L^2$. The analyticity in α comes from Lax-Milgram, but can also alternatively be obtained by Cauchy formula (for $\varepsilon > 0$ small)

$$\tilde{\mathbf{R}}(\alpha)f = \frac{1}{2\pi i} \int_{|z-\alpha|=\varepsilon} \frac{\tilde{\mathbf{R}}(z)f}{z-\alpha} dz$$

which holds for all $f \in \mathcal{C}$ (since $\tilde{\mathbf{R}}(\alpha)f = \mathbf{R}(\alpha)f$ for such f), and can then be extended to $e^{-\beta\rho} L^2$ by density of \mathcal{C} in $e^{-\beta\rho} L^2$. The domain $\mathcal{D}(e^{\beta\rho} \mathbf{H} e^{-\beta\rho}) = \{u \in \mathcal{D}(\mathcal{Q}) \mid e^{\beta\rho} \mathbf{H} e^{-\beta\rho} u \in L^2\}$ of the operator $e^{\beta\rho} \mathbf{H} e^{-\beta\rho}$ is actually equal to $\mathcal{D}(\mathbf{H}) = \{u \in \mathcal{D}(\mathcal{Q}) \mid \mathbf{H}u \in L^2\}$ since

$$(6.17) \quad e^{-\beta\rho} \mathbf{H}(e^{\beta\rho} u) = \mathbf{H}u - \frac{\beta^2}{2}(\rho'(c))^2 u - \frac{\beta}{2}\rho''(c)u - \beta\rho'(c)\partial_c u$$

with $\rho' \in C_c^\infty(\mathbb{R})$ (thus $\rho'(c)\partial_c u \in L^2$ for $u \in \mathcal{D}(\mathcal{Q})$). The operator $\mathbf{H} : e^{-\beta\rho} \mathcal{D}(\mathbf{H}) \rightarrow e^{-\beta\rho} L^2$ is thus closed. By Hille-Yosida theorem, there is an associated bounded semigroup $e^{-t\mathbf{H}}$ on $e^{-\beta\rho} L^2$, and by density of $L^2 \subset e^{-\beta\rho} L^2$ when $\beta \geq 0$, it is an extension of the $e^{-t\mathbf{H}}$ semigroup on L^2 . Let us check that the resolvent can be written as an integral of the propagator. For $f \in \mathcal{C} \subset L^2$, we have

$$\tilde{\mathbf{R}}(\alpha)f = \mathbf{R}(\alpha)f = \int_0^\infty e^{-t\mathbf{H} + t2\Delta_\alpha} f dt.$$

By Hille-Yosida theorem and (6.15), we have $\|e^{-t\mathbf{H}}\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq e^{-t\frac{Q^2 - \beta^2}{2}}$, so that the integral above converges if $Q - \alpha > \beta$ (for $\alpha \in (-\infty, Q)$) as a bounded operator on $e^{-\beta\rho} L^2$, showing the desired claim by density of \mathcal{C} in $e^{-\beta\rho} L^2$.

We conclude with a $e^{\beta\rho} \mathcal{D}'(\mathcal{Q})$ bound for $\tilde{\mathbf{R}}(\alpha)$ and $e^{-t\mathbf{H}}$. First, we note using (6.17) that for $u \in \mathcal{D}(\mathcal{Q})$

$$|\operatorname{Im}(\mathcal{Q}_{\alpha,\beta}(u, u))| \geq (|\operatorname{Im}(2\Delta_\alpha)| - \beta^2) \|u\|_2^2 - \frac{1}{4} \|\partial_c u\|_2^2,$$

which implies

$$|\langle \mathcal{Q}_{\alpha,\beta}(u, u) \rangle| \geq \frac{1}{\sqrt{2}} \left(c_{\Delta_\alpha} \|u\|_2^2 + \frac{1}{4} \|\partial_c u\|_2^2 + \|\mathbf{P}^{1/2} u\|_2^2 + \mathcal{Q}_{e^{\gamma c} V}(u, u) \right).$$

$$c_z := \min \left(-2\operatorname{Re}(z) + 2|\operatorname{Im}(z)| + \frac{Q^2 - 3\beta^2}{2}, -2\operatorname{Re}(z) + \frac{Q^2 - \beta^2}{2} \right).$$

This in turn gives that, provided $\operatorname{Re}(-2\Delta_\alpha) + |\operatorname{Im}(2\Delta_\alpha)| + \frac{Q^2 - 3\beta^2}{2} > 0$, $\tilde{\mathbf{R}}(\alpha) : e^{-\beta\rho} L^2 \rightarrow e^{-\beta\rho} L^2$ is also well-defined, analytic in α and bounded, and moreover satisfies for $|\Delta_\alpha| \gg 1$ and $\operatorname{Re}(\Delta_\alpha) \leq \frac{1}{2} |\operatorname{Im}(\Delta_\alpha)|$

$$\|\tilde{\mathbf{R}}(\alpha)\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq C |\Delta_\alpha|^{-1}$$

where $C > 0$ is a uniform constant. First, for each $\varepsilon > 0$, there is $C_\varepsilon > 0$ such that for all $f \in e^{-\beta\rho} \mathcal{D}(\mathcal{Q})$, $u := \mathbf{R}(\alpha)f$

$$\mathcal{Q}(u, u) \leq C_\varepsilon \operatorname{Re}(\mathcal{Q}_{\alpha,\beta}(u, u)) \leq C_\varepsilon |\langle u | f \rangle_2| \leq \frac{C_\varepsilon}{c_{\Delta_\alpha}} \|f\|_2^2$$




FIGURE 2. The blue region corresponds to the set of parameters $\alpha \in \mathbb{C}$ such that $\operatorname{Re}((\alpha - Q)^2) > \beta^2$, i.e. region of validity of Lemma 6.5 (here $\beta = 1$ on the plot).

if $c_{\Delta_\alpha} > 0$, thus for $c_z > 0$

$$(6.18) \quad \begin{aligned} \|(\mathbf{H} - z)^{-1}\|_{\mathcal{L}(e^{-\beta\rho}\mathcal{D}(\mathcal{Q}))} &\leq C_\varepsilon^{1/2} c_z^{-1/2} \\ \|(\mathbf{H} - z)^{-1}\|_{\mathcal{L}(e^{-\beta\rho}L^2)} &\leq C_\varepsilon c_z^{-1}. \end{aligned}$$

Let us consider a contour $\Gamma = \Gamma_0 \cup \Gamma_+ \cup \Gamma_- \subset \mathbb{C}$ with $a := \frac{Q^2 - \beta^2}{2} - \varepsilon$, $\Gamma_\pm := a \pm iN + e^{\pm 3i\pi/8}\mathbb{R}^+ \subset \mathbb{C}$ and $\Gamma_0 = a + i[-N, N]$ for some $N > 0$ large enough so that $c_z > 0$ on Γ , and where Γ oriented clockwise around $[a, +\infty)$. Using the holomorphic functional calculus, we have

$$e^{-t\mathbf{H}} = \frac{1}{2\pi i} \int_{\Gamma_+ \cup \Gamma_-} e^{-tz} (\mathbf{H} - z)^{-1} dz$$

and the integral converges both in $\mathcal{L}(e^{-\beta\rho}L^2)$ and $\mathcal{L}(e^{-\beta\rho}\mathcal{D}(\mathcal{Q}))$ using (6.18), with bound

$$\|e^{-t\mathbf{H}}\|_{e^{-\beta\rho}\mathcal{D}(\mathcal{Q})} \leq C e^{-ta}$$

for some C depending only on $\varepsilon > 0$. Using duality, this gives (6.12). \square

In what follows, we will always write $\mathbf{R}(\alpha)$ for the resolvent, for both the operator acting on L^2 or acting on $e^{-\beta\rho}L^2$.

Poisson operator in the probabilistic region. For $\ell \in \mathbb{N}$, we shall define the Poisson operator $\mathcal{P}_\ell(\alpha)$ in the resolvent set. This operator is a way to construct the generalized eigenfunctions of \mathbf{H} : it takes an element $F \in E_\ell \subset L^2(\Omega_{\mathbb{T}})$ and produces a function $u = \mathcal{P}_\ell(\alpha)F$ solving $(\mathbf{H} - 2\Delta_\alpha)u = 0$ with a prescribed leading asymptotic in terms of F as $c \rightarrow -\infty$.

First, we explain in details our convention on square roots in \mathbb{C} and since it will be important in the proof and to avoid confusions for the reader. We will denote by $\sqrt{\cdot}$ the square root defined so that $\operatorname{Im}(\sqrt{z}) > 0$ if $z \in \mathbb{C} \setminus \mathbb{R}^+$, i.e. $\sqrt{re^{i\theta}} = \sqrt{r}e^{i\theta/2}$ for $\theta \in (0, 2\pi)$ and $r > 0$. In particular, one has $\sqrt{z^2} = z$ for $\operatorname{Im}(\sqrt{z}) > 0$ and this extends holomorphically to $z \in \mathbb{C}$. For $\lambda \in \mathbb{R}^+$, the function $\sqrt{z^2 - \lambda}$ will be of special interest to us: it is well-defined and analytic in $\{\operatorname{Im}(z) > 0\} \cup (-\sqrt{\lambda}, \sqrt{\lambda})$ (since $z^2 \notin [\lambda, \infty)$ in this region) and it extends analytically in a neighborhood of the half lines $(-\infty, -\sqrt{\lambda})$ and $(\sqrt{\lambda}, +\infty)$, for exemple in $z \in \sqrt{\lambda} + \mathbb{R}^+ e^{-i\theta}$ and $z \in -\sqrt{\lambda} + \mathbb{R}^+ e^{i(\pi+\theta)}$ for $\theta \in [0, \varepsilon)$. With that convention, which will be used all along this Section on scattering for \mathbf{H} , we have $\sqrt{z^2 - \lambda} > 0$ if $z \in (\sqrt{\lambda}, \infty)$ while $\sqrt{z^2 - \lambda} < 0$ if $z \in (-\infty, -\sqrt{\lambda})$. Later we will view these functions as holomorphic functions on a Riemann surface.

We note the following elementary property, which will be useful in the following.

Lemma 6.6. *For $z \in \mathbb{C} \setminus \mathbb{R}^+$, the following map is non-decreasing*

$$x \in \mathbb{R}^+ \mapsto \operatorname{Im} \sqrt{z - x}.$$

Proof. It suffices to differentiate in x . □

Let $\chi \in C^\infty(\mathbb{R})$ be equal to 1 in $(-\infty, a - 1)$ and equal to 0 in $(a, +\infty)$ for some $a \in (0, 1/2)$, then for $\alpha = Q + ip$ with $\operatorname{Im}(p) > 0$ we choose

$$(6.19) \quad \beta_\ell > \max_{j=0, \dots, \ell} \operatorname{Im} \sqrt{p^2 - 2\lambda_j} - \gamma/2 = \operatorname{Im} \sqrt{p^2 - 2\lambda_\ell} - \gamma/2, \quad \text{and } \beta_\ell \geq 0.$$

Then for $\operatorname{Re}((\alpha - Q)^2) > \beta_\ell^2$ we define

$$(6.20) \quad \mathcal{P}_\ell(\alpha) : \begin{cases} E_\ell = \oplus_{j=0}^\ell \ker(\mathbf{P} - \lambda_j) & \rightarrow e^{-(\beta_\ell + \gamma/2)\rho} \mathcal{D}(\mathcal{Q}) \\ F = \sum_{0 \leq j \leq \ell} F_j & \rightarrow \chi F_-(\alpha) - \mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha)(\chi F_-(\alpha)), \end{cases}$$

with $F_-(\alpha) := \sum_{j=0}^\ell F_j e^{ic\sqrt{p^2 - 2\lambda_j}}$.

We will show in the following Lemma that this definition makes sense by using Lemma 6.5. Before going to the proof of it, recall that $\operatorname{Im} \sqrt{p^2 - 2\lambda_j} > 0$ for $\operatorname{Re}(\alpha) < Q$ by Lemma 6.6, and note that the condition $\operatorname{Re}((\alpha - Q)^2) > \beta_\ell^2$ implies that $\operatorname{Im} \sqrt{p^2 - 2\lambda_j} \geq \operatorname{Im}(p) > \beta_\ell$ for all $j = 0, \dots, \ell$. We then emphasize that the main reason for $\mathcal{P}_\ell(\alpha)F$ to be defined and non-trivial is that $\chi F_-(\alpha) \in e^{-(\beta_\ell + \gamma/2)\rho} \mathcal{D}(\mathcal{Q}) \setminus e^{-\beta_\ell \rho} \mathcal{D}(\mathcal{Q})$ and $(\mathbf{H} - 2\Delta_\alpha)(\chi F_-(\alpha)) \in e^{-\beta_\ell \rho} \mathcal{D}'(\mathcal{Q})$ so that $\mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha)(\chi F_-(\alpha))$ is well-defined but not equal to $\chi F_-(\alpha)$.

Lemma 6.7. *For each $\ell \in \mathbb{N}$, let $\beta_\ell \geq 0$, then the operator $\mathcal{P}_\ell(\alpha)$ is well-defined, bounded and holomorphic in the region*

$$(6.21) \quad \left\{ \alpha = Q + ip \in \mathbb{C} \mid \operatorname{Re}(\alpha - Q) < 0, \operatorname{Re}((\alpha - Q)^2) > \beta_\ell^2, \beta_\ell > \operatorname{Im} \sqrt{p^2 - 2\lambda_\ell} - \gamma/2 \right\}$$

and it satisfies in $e^{-(\beta_\ell + \gamma/2)\rho} \mathcal{D}'(\mathcal{Q})$

$$(6.22) \quad (\mathbf{H} - 2\Delta_\alpha) \mathcal{P}_\ell(\alpha) = 0,$$

and in the region $c \leq -1$, one has the asymptotic behaviour, with $F_j := (\Pi_j - \Pi_{j-1})F$,

$$(6.23) \quad \mathcal{P}_\ell(\alpha)F = \sum_{j=0}^\ell F_j e^{ic\sqrt{p^2 - 2\lambda_j}} + F_+(\alpha), \quad F_+(\alpha) \in e^{-\beta_\ell \rho} \mathcal{D}(\mathcal{Q}).$$

Proof. First we observe that $(\mathbf{H}_0 - 2\Delta_\alpha)F_-(\alpha) = 0$ thus

$$(\mathbf{H} - 2\Delta_\alpha)\chi F_-(\alpha) = -\frac{1}{2}\chi''(c)F_-(\alpha) + e^{\gamma c}V\chi F_-(\alpha) - \chi'(c)\partial_c F_-(\alpha).$$

We note that χ', χ'' have compact support in \mathbb{R} , and also for each $u \in \mathcal{D}(\mathcal{Q})$

$$\begin{aligned} |\langle \chi e^{\gamma c + \beta_\ell \rho} V F_-(\alpha), u \rangle| &\leq \int_{\mathbb{R}^-} e^{\gamma c + \beta_\ell \rho} |\mathcal{Q}_V(F_-(\alpha), u)| dc \leq \left(\int_{\mathbb{R}^-} e^{\gamma c} \mathcal{Q}_V(u) dc \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^-} e^{\gamma c} \mathcal{Q}_V(e^{\beta_\ell \rho} F_-) dc \right)^{\frac{1}{2}} \\ &\leq \mathcal{Q}(u)^{1/2} \mathcal{Q}_{e^{\gamma c} V}(e^{\beta_\ell \rho} F_-)^{\frac{1}{2}} \end{aligned}$$

and $\mathcal{Q}_{e^{\gamma c} V}(e^{\beta_\ell \rho} F_-) < \infty$ by using (6.21) and the fact that $\mathcal{Q}_V(\psi) < \infty$ for all $\psi \in E_\ell$. We obtain that

$$\chi F_-(\alpha) \in e^{-(\beta_\ell + \gamma/2)\rho} \mathcal{D}(\mathcal{Q}), \quad (\mathbf{H} - 2\Delta_\alpha)\chi F_-(\alpha) \in e^{-\beta_\ell \rho} \mathcal{D}'(\mathcal{Q}).$$

This shows, using Lemma 6.5, that $\mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha)(\chi F_-(\alpha))$ is well-defined as an element of $e^{-\beta_\ell \rho} \mathcal{D}(\mathcal{Q})$, with holomorphic dependence in α , provided $\operatorname{Re}((\alpha - Q)^2) > \beta_\ell^2$. By construction, it clearly also solves (6.22) in $e^{-(\beta_\ell + \gamma/2)\rho} \mathcal{D}'(\mathcal{Q})$. □

Note that the error term $F_+(\alpha)$ in (6.23) is smaller than the bigger term in $F_-(\alpha)$ but is not necessarily neglectible with respect to all terms of $F_-(\alpha)$.

We also notice that where they are defined, we have for $j \geq 0, \ell \geq 0$

$$(6.24) \quad \mathcal{P}_{\ell+j}(\alpha)|_{E_\ell} = \mathcal{P}_\ell(\alpha)|_{E_\ell}.$$




FIGURE 3. Left picture: regions of definition of the Poisson operator $\mathcal{P}_0(\alpha)$ (for the plot, we take $\gamma = 1/2$ and β_0 is optimized to obtain the largest possible region). Right picture: regions of definition of the Poisson operator $\mathcal{P}_\ell(\alpha)$ with $\ell > 0$ (for the plot we take $\lambda_\ell = 4, \gamma = 1/2$ and β_ℓ optimized).

In (6.20), the definition of the operator $\mathcal{P}_\ell(\alpha)$ seemingly depends on the cutoff function χ . In fact, we can show that this is not the case. We state a lemma below in this direction

Lemma 6.8. *For $\ell \in \mathbb{N}$, β_ℓ satisfying (6.19) and for $\text{Re}((\alpha - Q)^2) > \beta_\ell^2$ the definition of the operator $\mathcal{P}_\ell(\alpha)|_{E_\ell}$ does not depend on χ .*

Proof. Pick two functions $\chi, \hat{\chi}$ satisfying our assumptions and denote by $\mathcal{P}_\ell^\chi(\alpha), \mathcal{P}_\ell^{\hat{\chi}}(\alpha)$ the corresponding Poisson operators. Set $d(\chi) := \chi - \hat{\chi}$. Then observe that for $F \in E_\ell$

$$\mathcal{P}_\ell^\chi(\alpha)F - \mathcal{P}_\ell^{\hat{\chi}}(\alpha)F = d(\chi)F_-(\alpha) - \mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha)(d(\chi)F_-(\alpha)).$$

Now we note that $d(\chi)F_-(\alpha) \in \mathcal{D}(\mathcal{Q})$ since $d(\chi)(c) = 0$ for $c \notin (-1, a)$ for some $a > 0$ and \mathcal{Q}_V is bounded on E_ℓ . Then $\mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha)(d(\chi)F_-(\alpha)) = d(\chi)F_-(\alpha)$ since $(\mathbf{H} - 2\Delta_\alpha) : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$ is an isomorphism, hence $\mathcal{P}_\ell^\chi(\alpha)F - \mathcal{P}_\ell^{\hat{\chi}}(\alpha)F = 0$. \square

We can also rewrite the construction of the Poisson operator using the propagator.

Proposition 6.9. *The following properties hold true:*

1) *Let $\ell \in \mathbb{N}$ and let $F \in L^2(\Omega_{\mathbb{T}}) \cap \ker(\mathbf{P} - \lambda_j)$ for $j \leq \ell$. Then we have the identity*

$$\mathcal{P}_\ell(\alpha)F = \lim_{t \rightarrow +\infty} e^{t2\Delta_\alpha} e^{-t\mathbf{H}} \left(e^{ic\sqrt{p^2 - 2\lambda_j}} \chi(c)F \right)$$

in $e^{-(\beta+\gamma/2)\rho}\mathcal{D}'(\mathcal{Q})$ provided $\text{Re}((\alpha - Q)^2) > \beta^2$ with $\beta > \text{Im}(\sqrt{p^2 - 2\lambda_j}) - \gamma/2$ and $\alpha = Q + ip$. Furthermore if $\text{Im}(\sqrt{p^2 - 2\lambda_j}) > \gamma$ then we can take $\chi = 1$ in the above statement.

2) *Let $F \in L^2(\Omega_{\mathbb{T}}) \cap \ker(\mathbf{P})$. If $\alpha \in \mathbb{R}$ with $\alpha < Q$, then $dc \otimes \mathbf{P}$ almost everywhere*

$$\mathcal{P}_0(\alpha)F = \lim_{t \rightarrow +\infty} e^{t2\Delta_\alpha} e^{-t\mathbf{H}} \left(e^{(\alpha-Q)c} F \right).$$

Proof. We first prove 1). Recall that $\mathbf{H} = \mathbf{H}_0 + e^{\gamma c}V$. We have

$$(\mathbf{H} - 2\Delta_\alpha)(\chi(c)e^{i\nu_j c}F) = \chi(c)e^{(i\nu_j + \gamma)c}VF - \tilde{\chi}(c)F e^{i\nu_j c}$$

where $\nu_j := \sqrt{p^2 - 2\lambda_j}$ and $\tilde{\chi}(c) := \frac{1}{2}\chi''(c) + i\nu_j\chi'(c) \in C_c^\infty(\mathbb{R})$, and this belongs to $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q})$. Using Lemma 6.5,

$$\mathcal{P}_\ell(\alpha)F = \chi(c)e^{i\nu_j c}F - \mathbf{R}(\alpha)(e^{(i\nu_j + \gamma)c}V\chi F - \tilde{\chi}e^{i\nu_j c}F)$$

provided $\operatorname{Re}((\alpha - Q)^2) > \beta^2$ for any $\beta > \operatorname{Im}(\nu_j) - \gamma/2$. Noticing that the bound (6.11) remains valid with $V = 0$, we can make sense of $u(t) := e^{-t\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F)$ as an element in $e^{-\beta_0\rho}L^2(\mathbb{R}; E_j)$ for any $\beta_0 > \operatorname{Im}(\nu_j)$. Then

$$(\partial_t + 2\Delta_\alpha)u(t) = e^{-t\mathbf{H}_0}(-\mathbf{H}_0 + 2\Delta_\alpha)(\chi(c)e^{i\nu_j c}F) = e^{-t\mathbf{H}_0}(e^{i\nu_j c}F\tilde{\chi}(c))$$

thus we get by integrating in t

$$(6.25) \quad \begin{aligned} e^{-t(\mathbf{H}_0 - 2\Delta_\alpha)}(\chi(c)e^{i\nu_j c}F) &= \chi(c)e^{i\nu_j c}F + \int_0^t e^{-s(\mathbf{H}_0 - 2\Delta_\alpha)}(e^{i\nu_j c}F\tilde{\chi}(c))ds \\ &= \chi(c)e^{i\nu_j c}F + (1 - e^{-t(\mathbf{H}_0 - 2\Delta_\alpha)})\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)) \end{aligned}$$

where $\mathbf{R}_0(\alpha) := (\mathbf{H}_0 - 2\Delta_\alpha)^{-1}$ is defined also on $e^{-\beta\rho}L^2$ by taking the proof of Lemma 6.5 in the case of the trivial potential $V = 0$. We also note that $e^{-t\mathbf{H}_0}(e^{i\nu_j c}F\tilde{\chi}(c))$ and $e^{-t\mathbf{H}_0}\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))$ are in $L^2(\mathbb{R}; E_j)$ since $\mathbf{H}_0 F = (Q^2/2 + \lambda_j)F$ (and $\tilde{\chi} \in C_c^\infty(\mathbb{R})$), i.e. all terms above are functions of c with values in E_j .

We next claim that

$$e^{-t\mathbf{H}}(\chi(c)e^{i\nu_j c}F) = e^{-t\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F) - \int_0^t e^{-(t-s)\mathbf{H}}e^{\gamma c}Ve^{-s\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F)ds.$$

Indeed, first, all terms are well-defined: since $e^{-s\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F) \in e^{-\beta_0\rho}L^2(\mathbb{R}; E_j) \cap e^{-\beta_0\rho}\mathcal{D}(\mathcal{Q})$ one has $e^{\gamma c}Ve^{-s\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F) \in e^{-\beta_0\rho}\mathcal{D}'(\mathcal{Q})$, and we can then use (6.12). Then the identity above is obtained since both terms solve $(\partial_t + \mathbf{H})u(t) = 0$ in $e^{-\beta_0\rho}\mathcal{D}'(\mathcal{Q})$ with $u(0) = \chi(c)e^{i\nu_j c}F$ in $e^{-\beta_0\rho}\mathcal{D}(\mathcal{Q})$. By applying twice (6.25), we thus obtain

$$\begin{aligned} e^{-t(\mathbf{H} - 2\Delta_\alpha)}(\chi(c)e^{i\nu_j c}F) &= \chi(c)e^{i\nu_j c}F - e^{t2\Delta_\alpha} \int_0^t e^{-(t-s)\mathbf{H}}e^{\gamma c}Ve^{-s\mathbf{H}_0}(\chi(c)e^{i\nu_j c}F)ds \\ &\quad + (1 - e^{-t(\mathbf{H}_0 - 2\Delta_\alpha)})\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)) \\ &= \chi(c)e^{i\nu_j c}F - \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}(e^{\gamma c}V\chi(c)e^{i\nu_j c}F)ds \\ &\quad + (1 - e^{-t(\mathbf{H}_0 - 2\Delta_\alpha)})\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)) \\ &\quad - \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}e^{\gamma c}V(1 - e^{-s(\mathbf{H}_0 - 2\Delta_\alpha)})\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))ds. \end{aligned}$$

Using (6.11) and (6.12) (applied with both $V \neq 0$ and $V = 0$) and $\operatorname{Re}((\alpha - Q)^2) > \beta^2$, we have as bounded operators on respectively $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q})$ and $e^{-\beta\rho}L^2$

$$\begin{aligned} \lim_{t \rightarrow +\infty} \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}ds &= \lim_{t \rightarrow +\infty} (1 - e^{-t(\mathbf{H} - 2\Delta_\alpha)})\mathbf{R}(\alpha) = \mathbf{R}(\alpha) \\ \lim_{t \rightarrow +\infty} (1 - e^{-t(\mathbf{H}_0 - 2\Delta_\alpha)})\mathbf{R}_0(\alpha) &= \mathbf{R}_0(\alpha). \end{aligned}$$

This gives in particular in $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q})$

$$\lim_{t \rightarrow +\infty} \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}(e^{\gamma c}V\chi(c)e^{i\nu_j c}F)ds = \mathbf{R}(\alpha)(e^{\gamma c}V\chi(c)e^{i\nu_j c}F).$$

Similarly one has in $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q})$

$$\lim_{t \rightarrow +\infty} \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}e^{\gamma c}V\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))ds = \mathbf{R}(\alpha)e^{\gamma c}V\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)).$$

Finally we claim that

$$\lim_{t \rightarrow +\infty} \int_0^t e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}e^{\gamma c}Ve^{-s(\mathbf{H}_0 - 2\Delta_\alpha)}\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))ds = 0.$$

Indeed, we can apply Lebesgue theorem if one can show

$$\|e^{\gamma c}Ve^{-s(\mathbf{H}_0 - 2\Delta_\alpha)}\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))\|_{e^{\beta\rho}\mathcal{D}'(\mathcal{Q})} \leq e^{-\delta s}$$

for some $\delta > 0$, since $\|e^{-(t-s)(\mathbf{H} - 2\Delta_\alpha)}\|_{\mathcal{L}(e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}))} \rightarrow 0$ by (6.12). But this estimate follows again from (6.11) with $V = 0$ and the fact that $\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)) \in e^{-\beta\rho}L^2(\mathbb{R}; E_j)$, which in turn implies that

$$e^{\gamma c}Ve^{-s(\mathbf{H}_0 - 2\Delta_\alpha)}\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)) \in e^{\beta\rho}\mathcal{D}'(\mathcal{Q}).$$

We have thus proved that

$$\begin{aligned} \lim_{t \rightarrow +\infty} e^{-t(\mathbf{H}-2\Delta_\alpha)}(\chi(c)e^{i\nu_j c}F) &= \chi(c)e^{i\nu_j c}F - \mathbf{R}(\alpha)e^{\gamma c}V(e^{i\nu_j c}F\chi(c) + \mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))) \\ &\quad + \mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c)). \end{aligned}$$

We conclude by observing that

$$\mathbf{R}(\alpha)(e^{\gamma c}V\chi e^{i\nu_j c}F - \tilde{\chi}e^{i\nu_j c}F) = -\mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}) + \mathbf{R}(\alpha)e^{\gamma c}V(\chi e^{i\nu_j c}F + \mathbf{R}_0(\alpha)(e^{i\nu_j c}F\tilde{\chi}(c))),$$

which can be established by applying $(\mathbf{H} - 2\Delta_\alpha)$ to this equation and using the injectivity of $(\mathbf{H} - 2\Delta_\alpha)$ on $e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ under our condition on α, β .

Notice that for $\text{Im}(\sqrt{p^2 - 2\lambda_j}) > \gamma$, we have $e^{(i\nu_j + \gamma)c}\mathbf{1}_{(0, +\infty)}(c) \in L^2(\mathbb{R})$ so that the above argument applies with $\chi = 1$.

Now we prove 2). As $F \in L^2(\Omega_{\mathbb{T}}) \cap \ker(\mathbf{P} - \lambda_0)$, we may assume $F = 1$ without loss of generality. With our assumptions, we can write $\alpha = Q + ip$ with $p = i(Q - \alpha)$ and choose $(Q - \alpha) > \beta > (Q - \alpha) - \gamma$. By applying 1), we get that

$$\mathcal{P}_\ell(\alpha)1 = \lim_{t \rightarrow +\infty} e^{t2\Delta_\alpha} e^{-t\mathbf{H}}(e^{(\alpha-Q)c}\chi(c))$$

in $e^{-\beta\rho}L^2$, hence $dc \otimes \mathbb{P}$ almost everywhere (up to extracting subsequence). We have to show that we can replace χ by 1. For this we will use the probabilistic representation (5.4): we have

$$\begin{aligned} e^{t2\Delta_\alpha} e^{-t\mathbf{H}}(e^{(\alpha-Q)c}(1 - \chi(c))) &= e^{-Qc} \mathbb{E}_\varphi[S_{e^{-t}}(e^{\alpha c}(1 - \chi(c)))e^{-\mu M_\gamma(\mathbb{D}_t^c)}] \\ &= e^{t2\Delta_\alpha} e^{(\alpha-Q)c} \mathbb{E}_\varphi[e^{\alpha(X_t(0) - Qt)}(1 - \chi(c + X_t(0) - Qt))]e^{-\mu M_\gamma(\mathbb{D}_t^c)}. \end{aligned}$$

By Girsanov's transform this expression can be rewritten as

$$e^{t2\Delta_\alpha} e^{-t\mathbf{H}}(e^{(\alpha-Q)c}(1 - \chi(c))) = e^{(\alpha-Q)c} \mathbb{E}_\varphi[(1 - \chi(c + X_t(0) - (Q - \alpha)t)) \exp(-\mu \int_{\mathbb{D}_t^c} |z|^{-\gamma\alpha} M_\gamma(dz))].$$

Recalling that $\chi = 1$ on $(-\infty, a - 1)$ and that $t \mapsto X_t(0)$ evolves as a Brownian motion independent of φ , then estimating the exponential term by 1 we obtain

$$|e^{t2\Delta_\alpha} e^{-t\mathbf{H}}(e^{(\alpha-Q)c}(1 - \chi(c)))| \leq e^{(\alpha-Q)c} \mathbb{P}(c + X_t(0) - (Q - \alpha)t \geq a - 1).$$

The result easily follows from this estimate. \square

Meromorphic extension of the resolvent near the L^2 spectrum. We denote by Z the Riemann surface on which the functions $p \mapsto \sqrt{p^2 - 2\lambda_j}$ are single valued for all j . This is a ramified covering of \mathbb{C} with ramification points $\{\pm\sqrt{2\lambda_j} \mid j \in \mathbb{N}\}$, and in which we embed the region $\text{Im}(p) > 0$ that we call the *physical sheet*. We will call $\pi : Z \rightarrow \mathbb{C}$ the projection of the covering. The construction of Z can be done iteratively on j , as explained in Chapter 6.7 of Melrose's book [Me93]. The map $p \mapsto \alpha := Q + ip$ from $\text{Im}(p) > 0$ to $\text{Re}(\alpha) < Q$ (now called the physical sheet for the variable α) extends analytically as a map $Z \rightarrow \Sigma$ where Σ is an isomorphic Riemann surface to Z (it just amounts to a linear change of complex coordinates from p to α). We shall also denote by $\pi : \Sigma \rightarrow \mathbb{C}$ the projection. Finally we choose a specific function χ of the form indicated previously but we further impose that $\chi \in C^\infty(\mathbb{R})$ is equal to 1 in $(-\infty, -1 + \delta)$ and equal to 0 in $(0, +\infty)$ (for some small $\delta > 0$).

First, we recall the notation for the orthogonal projectors

$$\Pi_k = 1_{[0, \lambda_k]}(\mathbf{P}) : L^2(\Omega_{\mathbb{T}}, \mathbb{P}) \rightarrow L^2(\Omega_{\mathbb{T}}, \mathbb{P})$$

and we denote by E_k their range (which are Hilbert spaces) and E_k^\perp the range of $1 - \Pi_k$. The goal of this section is to show the following:

Proposition 6.10. *Assume that $\gamma \in (0, \sqrt{2})$ and let $0 \leq \beta < \gamma/2$. Then the following holds true:*

1) *The resolvent $\mathbf{R}(\alpha) := (\mathbf{H} - 2\Delta_\alpha)^{-1}$ is bounded as a map $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H})$ for $\text{Re}(\alpha) < Q$ and for $k \geq 0$ large enough, the operator $(\mathbf{H} - 2\Delta_{\pi(\alpha)})^{-1}$ admits a meromorphic continuation to the region*

$$\{\alpha = Q + ip \in \Sigma \mid |\pi(p)|^2 \leq \lambda_k^{\frac{1}{4}}, \forall j \leq k, \text{Im}\sqrt{p^2 - 2\lambda_j} > -\min(\beta, \gamma/2 - \beta)\}$$

as a map

$$\mathbf{R}(\alpha) : e^{\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathbf{H}),$$

and the residue at each pole is a finite rank operator. Moreover, for each $\psi \in C^\infty(\mathbb{R}) \cap L^\infty(\mathbb{R})$ satisfying $\psi' \in L^\infty$ and $\text{supp}(\psi) \subset (-\infty, A)$ for some $A \in \mathbb{R}$,

$$\mathbf{R}(\alpha)(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q}).$$

2) If $f \in e^{\beta\rho}\mathcal{D}'(\mathcal{Q})$ is supported in $c \in (-\infty, A)$ for some $A > 0$ and is such that $\Pi_k f \in e^{\beta\rho}L^2(\mathbb{R}; E_k)$ and $(1 - \Pi_k)f \in \mathcal{D}'(\mathcal{Q})$, then for α as above and not a pole, one has in $c \leq 0$

$$(6.26) \quad \mathbf{R}(\alpha)f = \sum_{j=0}^k a_j(\alpha, f) e^{-ic\sqrt{p^2 - 2\lambda_j}} + G(\alpha, f),$$

with $a_j(\alpha, f) \in \ker(\mathbf{P} - \lambda_j)$, and $G(\alpha, f), \partial_c G(\alpha, f) \in e^{\beta\rho}L^2(\mathbb{R}^-; E_k) + L^2(\mathbb{R}^-; E_k^\perp)$, all depending meromorphically in α in the region they are defined.

3) There is no pole for $\mathbf{R}(\alpha)$ in $\{\alpha \in \Sigma \mid \text{Re}(\alpha) \leq Q\} \setminus \cup_{j=0}^\infty \{Q \pm i\sqrt{2\lambda_j}\}$ and $Q \pm i\sqrt{2\lambda_j}$ can be at most a pole of order 1.

To prove this Proposition, we will construct parametrices for the operator $\mathbf{H} - 2\Delta_\alpha = \mathbf{H} - \frac{Q^2 + p^2}{2}$ in several steps and will split the argument. More concretely, we will search for some bounded model operator $\tilde{\mathbf{R}}(\alpha) : e^{\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$, holomorphic in α in the desired region of Σ , such that

$$(\mathbf{H} - 2\Delta_\alpha)\tilde{\mathbf{R}}(\alpha) = 1 - \mathbf{K}(\alpha)$$

where $\mathbf{K}(\alpha) \in \mathcal{L}(e^{\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}))$ is an analytic family of compact operators with $1 - \mathbf{K}(\alpha_0)$ invertible at some α_0 belonging to the physical sheet. Then the Fredholm analytic theorem will imply that $(1 - \mathbf{K}(\alpha))^{-1}$ exists as a meromorphic family and $\mathbf{R}(\alpha) := \tilde{\mathbf{R}}(\alpha)(1 - \mathbf{K}(\alpha))^{-1}$ gives us the desired meromorphic extension of $\mathbf{R}(\alpha)$. Our strategy will be based on that method with slight modifications. The continuous spectrum of \mathbf{H} near frequency $(Q^2 + p^2)/2 \in \mathbb{R}^+$ will come only from finitely many eigenmodes of \mathbf{P} , namely those λ_j for which $2\lambda_j \leq p^2$. This suggests, in order to construct the approximation $\tilde{\mathbf{R}}(\alpha)$ to split the modes of \mathbf{P} depending on the value of $\text{Im}(\alpha - Q)$. The parametrix will be constructed in three steps as follows:

- First, we deal with the large eigenmodes for the operator \mathbf{P} in the region $c \in (-\infty, 0]$ of $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. We will show that this part does not contribute to the continuous spectrum at frequency $(Q^2 + p^2)/2$, and we shall obtain a parametrix for that part by energy estimates.
- Then we consider the region $c \geq -1$ where we shall show that the model operator in that region (essentially \mathbf{H} on $L^2([-1, \infty) \times \Omega_{\mathbb{T}})$ with Dirichlet condition at $c = 0$) has compact resolvent, providing a compact operator for the parametrix of that region.
- Finally, we will deal with the $c \leq 0$ region corresponding to eigenmodes of \mathbf{P} of order $\mathcal{O}(|p|^2)$, where there is *scattering* at $c = -\infty$ for frequency $(Q^2 + p^2)/2$, producing continuous spectrum. The parametrix for this part is basically the exact inverse of $\mathbf{H}_0 - 2\Delta_\alpha$ restricted to finitely many modes of \mathbf{P} .

For $s \geq 0$ and $I \subset \mathbb{R}$ an open interval, we will denote by $H^s(I; L^2(\Omega_{\mathbb{T}}))$ the Sobolev space of order s in the c -variable

$$H^s(I; L^2(\Omega_{\mathbb{T}})) := \{u \in L^2(I \times \Omega_{\mathbb{T}}) \mid \forall j \leq \ell, \xi \mapsto \langle \xi \rangle^s \mathcal{F}(u)(\xi) \in L^2(I \times \Omega_{\mathbb{T}})\}$$

where \mathcal{F} denotes Fourier transform in c , and similarly $H_0^s(I; L^2(\Omega_{\mathbb{T}}))$ will be the completion of $C_c^\infty(I; L^2(\Omega_{\mathbb{T}}))$ with respect to the norm $\|\langle \xi \rangle^s \mathcal{F}(u)\|_{L^2(\mathbb{R}; L^2(\Omega_{\mathbb{T}}))}$.

1) Large \mathbf{P} eigenmodes in the region $c \leq 0$. Let $\chi \in C^\infty(\mathbb{R}, [0, 1])$ which satisfies $\chi(c) = 1$ for $c \leq -1 + \delta$ for some $\delta \in (0, \frac{1}{2})$ and $\chi(c) = 0$ in $[-1/2, \infty)$ and $\tilde{\chi} \in C^\infty(\mathbb{R}, [0, 1])$ such that $\tilde{\chi} = 1$ on the support of χ and $\text{supp}(\tilde{\chi}) \subset \mathbb{R}^-$, and we now view these functions as multiplication operators by $\chi(c)$ on the spaces $e^{\beta\rho}L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})$. We will first show the following

Lemma 6.11. 1) *There is a constant $C > 0$ depending only on $|\tilde{\chi}'|_\infty, |\tilde{\chi}''|_\infty$ such that for each $k \in \mathbb{N}$, there is a bounded operator*

$$\mathbf{R}_k^\perp(\alpha) : L^2(\mathbb{R}; E_k^\perp) \rightarrow L^2(\mathbb{R}^-; E_k^\perp)$$

holomorphic in $\alpha = Q + ip \in \mathbb{C}$ in the region $\{\operatorname{Re}(\alpha) < Q\} \cup \{|\alpha - Q|^2 \leq \lambda_k\}$, with

$$\tilde{\chi} \mathbf{R}_k^\perp(\alpha) \chi : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R}; E_k^\perp) \cap \mathcal{D}(\mathbf{H}), \quad \tilde{\chi} \mathbf{R}_k^\perp(\alpha) \chi : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q}) \cap L^2(\mathbb{R}; E_k^\perp)$$

bounded, so that

$$(6.27) \quad (\mathbf{H} - \frac{Q^2 + p^2}{2}) \tilde{\chi} \mathbf{R}_k^\perp(\alpha) (1 - \Pi_k) \chi = (1 - \Pi_k) \chi + \mathbf{L}_k^\perp(\alpha) + \mathbf{K}_k^\perp(\alpha) \quad \text{and} \quad \tilde{\chi} \mathbf{R}_k^\perp(\alpha) \Pi_k \chi = 0$$

with $\mathbf{L}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}) \rightarrow L^2(\mathbb{R}^-; E_k^\perp)$ and $\mathbf{K}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}) \rightarrow e^{\beta\rho} L^2(\mathbb{R}; E_k)$ bounded and holomorphic in α as above for each $0 < \beta < \gamma/2$. In the region where $|p^2| \leq \lambda_k$, one has the bound

$$(6.28) \quad \|\mathbf{L}_k^\perp(\alpha)\|_{\mathcal{L}(L^2)} \leq C \lambda_k^{-1/2}$$

and $\mathbf{K}_k^\perp(\alpha)$ is compact as a map $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta\rho} L^2(\mathbb{R}; E_k)$.

2) *Let $\beta \in \mathbb{R}$, then in the region $\operatorname{Re}((\alpha - Q)^2) > \beta^2 - 2\lambda_k + 1$, the operator $\mathbf{R}_k^\perp(\alpha) : e^{-\beta\rho} \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho} (L^2(\mathbb{R}^-; E_k^\perp) \cap \mathcal{D}(\mathcal{Q}))$ is a bounded holomorphic family, $\mathbf{K}_k^\perp(\alpha) : e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} L^2(\mathbb{R}^-; E_k)$ is a compact holomorphic family, $\mathbf{L}_k^\perp(\alpha) : e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} L^2(\mathbb{R}^-; E_k^\perp)$ is bounded analytic with norm*

$$\|\mathbf{L}_k^\perp(\alpha)\|_{\mathcal{L}(e^{-\beta\rho} L^2)} \leq \frac{C(1 + |\beta|)}{\sqrt{\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2}}$$

for some $C > 0$ depending only on $|\tilde{\chi}'|_\infty$ and $|\tilde{\chi}''|_\infty$.

Proof. Let us define \mathcal{Q}_k^\perp the restriction of \mathcal{Q} to the closed subspace

$$\mathcal{D}(\mathcal{Q}_k^\perp) := \mathcal{D}(\mathcal{Q}) \cap L^2(\mathbb{R}^-; E_k^\perp) = \mathcal{D}(\mathcal{Q}) \cap \ker \Pi_k \cap \ker r_{\mathbb{R}^+}$$

where $r_{\mathbb{R}^\pm} : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R}^\pm \times \Omega_{\mathbb{T}})$ is the restriction for $c \geq 0$. Note that this is a closed form and is thus the quadratic form of a unique self-adjoint operator \mathbf{H}_k^\perp , which maps $\mathcal{D}(\mathcal{Q}_k^\perp) \rightarrow \mathcal{D}'(\mathcal{Q}_k^\perp)$ and has a domain $\mathcal{D}(\mathbf{H}_k^\perp) \subset \mathcal{D}(\mathcal{Q}_k^\perp)$. For $u \in \mathcal{D}(\mathcal{Q}_k^\perp)$ we have $\|\mathbf{P}^{1/2} u\|_2^2 \geq \lambda_k \|u\|_2^2$, thus for each $\varepsilon \geq 0$

$$(6.29) \quad \mathcal{Q}(u) \geq \frac{1}{2} \|\partial_c u\|_{L^2}^2 + \left(\frac{Q^2}{2} + (1 - \varepsilon)\lambda_k\right) \|u\|_{L^2}^2 + \varepsilon \|\mathbf{P}^{1/2} u\|_{L^2}^2 + \mathcal{Q}_{e^{\gamma c} V}(u, u)$$

and therefore the quadratic form $\mathcal{Q}_k^\perp(u)$ is bounded below by $\frac{\lambda_k}{2} \|u\|_{L^2}^2$ on $\mathcal{D}(\mathcal{Q}_k^\perp)$ (if ε is chosen small enough). There is a natural injection $\iota : \mathcal{D}(\mathcal{Q}_k^\perp) \rightarrow \mathcal{D}(\mathcal{Q})$ which we view as an inclusion. Let $\chi, \tilde{\chi} \in C^\infty(\mathbb{R})$ be equal to 1 in $c < -1/2$ and supported in $(-\infty, 0)$. Let $\tilde{\chi}, \chi$ defined as χ but equal to 1 on the support of χ . We view χ as the operator of multiplication by $\chi(c)$ on $L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, and similarly for $\tilde{\chi}$. Let us check that $\iota \chi = \chi \iota : \mathcal{D}(\mathbf{H}_k^\perp) \rightarrow \mathcal{D}(\mathbf{H})$ and for $u \in \mathcal{D}(\mathbf{H}_k^\perp)$

$$(6.30) \quad \mathbf{H} \iota(\chi u) = \iota(\mathbf{H}_k^\perp \chi u) + \chi e^{\gamma c} \Pi_k(V \iota(u))$$

(note that $\chi u \in \mathcal{D}(\mathbf{H}_k^\perp)$ if $u \in \mathcal{D}(\mathbf{H}_k^\perp)$ since $\mathbf{H}_k^\perp \chi u = \chi \mathbf{H}_k^\perp u - \frac{1}{2} \chi'' u - \chi' \partial_c u$). Here we use Lemma 6.1, which insures that $\chi e^{\gamma c} \Pi_k(V \iota(u))$ makes sense. First, we have for $u \in \mathcal{D}(\mathcal{Q}_k^\perp)$ that $\iota(u) = (1 - \Pi_k) \iota(u)$. For such u and for $v \in \mathcal{D}(\mathcal{Q})$, one gets

$$\begin{aligned} \mathcal{Q}(\iota(\chi u), v) &= \frac{1}{2} \langle \partial_c(1 - \Pi_k) \chi u, \partial_c(1 - \Pi_k) \tilde{\chi} v \rangle_{L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})} + \frac{Q^2}{2} \|(1 - \Pi_k) \chi u, (1 - \Pi_k) \tilde{\chi} v\|_{L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})}^2 \\ &\quad + \langle \mathbf{P}^{1/2}(1 - \Pi_k) \chi u, \mathbf{P}^{1/2}(1 - \Pi_k) \tilde{\chi} v \rangle_{L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})} + \int_{\mathbb{R}} \chi(c) e^{\gamma c} (\mathcal{Q}_V(u, (1 - \Pi_k)v) + \mathcal{Q}_V(u, \Pi_k v)) dc \\ &= \mathcal{Q}_k^\perp(\chi u, (1 - \Pi_k) \tilde{\chi} v) + \int_{\mathbb{R}} e^{\gamma c} \mathcal{Q}_V(\chi u, \Pi_k v) dc \\ &= \langle \mathbf{H}_k^\perp \chi u, (1 - \Pi_k) \tilde{\chi} v \rangle_2 + \int_{\mathbb{R}} e^{\gamma c} \mathcal{Q}_V(\chi u, \Pi_k v) dc = \langle \mathbf{H}_k^\perp \chi u, v \rangle_{L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})} + \int_{\mathbb{R}} e^{\gamma c} \mathcal{Q}_V(\chi u, \Pi_k v) dc \end{aligned}$$

where in the last line, $\tilde{v} := (1 - \Pi_k) \tilde{\chi} v$ is viewed as an element in $\mathcal{D}(\mathcal{Q}_k^\perp)$. As in the proof of Lemma 6.1, there is a uniform constant C_k depending only on k, γ such that

$$\left| \int_{\mathbb{R}^-} e^{\gamma c} \mathcal{Q}_V(u, \Pi_k v) dc \right| \leq C_k |\mathcal{Q}_{e^{\gamma c} V}(i(u))|^{1/2} \|v\|_2$$

which implies that $\iota(\chi u) \in \mathcal{D}(\mathbf{H})$ since $|\langle \mathbf{H}_k^\perp \chi u, v \rangle_{L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})}| \leq \|\mathbf{H}_k^\perp \chi u\|_2 \|v\|_2$. Now we also notice that

$$\int_{\mathbb{R}} e^{\gamma c} \mathcal{Q}_V(\chi u, \Pi_k v) dc = \langle e^{\gamma c} V \chi \iota(u), \Pi_k \tilde{\chi} v \rangle_2$$

where the pairing makes sense since $e^{\gamma c} V \chi \iota(u) \in \mathcal{D}'(\mathcal{Q})$ and $\Pi_k \tilde{\chi} v \in \mathcal{D}(\mathcal{Q})$ by Lemma 6.1. Moreover, using $\tilde{\chi} \chi = \chi$ and $\Pi_k^* = \Pi_k$ this term is also equal to

$$\langle e^{\gamma c} V \chi \iota(u), \Pi_k \tilde{\chi} v \rangle_2 = \langle \chi e^{\gamma c} \Pi_k(V \iota(u)), v \rangle_2.$$

This shows the formula (6.30).

The spectrum of \mathbf{H}_k^\perp is contained in $[\frac{Q^2}{2} + \lambda_k, \infty)$ due to (6.29). It will be said to have Dirichlet condition at $c = 0$, by analogy with the Laplacian on finite dimensional manifolds. The resolvent $\mathbf{R}_k^\perp(\alpha) := (\mathbf{H}_k^\perp - \frac{Q^2 + p^2}{2})^{-1}$ (with $\alpha = Q + ip$) is well defined as a bounded map

$$\mathbf{R}_k^\perp(\alpha) : L^2(\mathbb{R}^-; E_k^\perp) \rightarrow \mathcal{D}(\mathbf{H}_k^\perp) \quad \mathbf{R}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}_k^\perp) \rightarrow \mathcal{D}(\mathcal{Q}_k^\perp)$$

if $p \in \mathbb{C}$ is such that $p^2 \notin [2\lambda_k, \infty)$, with L^2 norm

$$(6.31) \quad \|\mathbf{R}_k^\perp(\alpha)\|_{\mathcal{L}(L^2)} \leq 2/\lambda_k \text{ for } |p|^2 \leq \lambda_k.$$

It is also holomorphic in α in $\{\operatorname{Re}(\alpha) < Q\} \cup \{|\alpha - Q|^2 < \lambda_k\}$. We also define the dual map $\iota^T : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q}_k^\perp)$ (which also maps $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R}^-; E_k^\perp)$). We get

$$\iota \chi \mathbf{R}_k^\perp(\alpha) \iota^T (1 - \Pi_k) : L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H}) \cap L^2(\mathbb{R}; E_k^\perp) \quad \iota \mathbf{R}_k^\perp(\alpha) \iota^T : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$$

with the same properties. To avoid heavy notations, we shall now remove the ι, ι^T maps in the notations so that we view $\chi \mathbf{R}_k^\perp(\alpha) (1 - \Pi_k)$ as map $L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H}) \cap L^2(\mathbb{R}; E_k^\perp)$.

Using (6.29) with $u = \mathbf{R}_k^\perp(\alpha) f$ we also obtain that

$$(6.32) \quad \|\partial_c \mathbf{R}_k^\perp(\alpha)\|_{\mathcal{L}(L^2)} \leq \sqrt{\frac{4}{\lambda_k}}, \text{ for } |p|^2 \leq \lambda_k/2.$$

Now we fix $\chi, \tilde{\chi}$ as defined before the Lemma. Thus, using $\Pi_k \mathbf{H}_0 = \mathbf{H}_0 \Pi_k$ and (6.30), we get for $\alpha = Q + ip$

$$\begin{aligned} (\mathbf{H} - \frac{Q^2 + p^2}{2}) \tilde{\chi} \mathbf{R}_k^\perp(\alpha) (1 - \Pi_k) \chi &= (1 - \Pi_k) \chi - \frac{1}{2} [\partial_c^2, \tilde{\chi}] \mathbf{R}_k^\perp(\alpha) (1 - \Pi_k) \chi + e^{\gamma c} \tilde{\chi} \Pi_k V \mathbf{R}_k^\perp(\alpha) \chi (1 - \Pi_k) \\ &=: (1 - \Pi_k) \chi + \mathbf{L}_k^\perp(\alpha) + \mathbf{K}_k^\perp(\alpha). \end{aligned}$$

Since $[\partial_c^2, \tilde{\chi}]$ is a first order operator with compact support in c commuting with Π_k , we notice that $\mathbf{L}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}) \rightarrow L^2(\mathbb{R}^-; E_k^\perp)$ and we can use (6.32), (6.31) to deduce that there is $C > 0$ depending only on $|\tilde{\chi}'|_\infty, |\tilde{\chi}''|_\infty$ such that

$$\|\mathbf{L}_k^\perp(\alpha)\|_{\mathcal{L}(L^2)} \leq C \lambda_k^{-1/2}$$

as long as $|p^2| \leq \lambda_k$. Let us now deal with $\mathbf{K}_k^\perp(\alpha)$. First, notice that $\mathbf{K}_k^\perp(\alpha)$ maps $\mathcal{D}'(\mathcal{Q})$ to $e^{\gamma \rho/2} L^2(\mathbb{R}^-; E_k)$: indeed,

$$\mathbf{R}_k^\perp(\alpha) (1 - \Pi_k) \chi : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q}), \quad e^{\gamma c} \tilde{\chi} \Pi_k V : \mathcal{D}(\mathcal{Q}) \rightarrow e^{\gamma \rho/2} L^2(\mathbb{R}^-; E_k)$$

using Lemma 6.1 with $\beta = \beta' = -\gamma/2$. We would like to prove some regularization property in c to deduce that $\mathbf{K}_k^\perp(\alpha)$ is compact on L^2 (or some weighted L^2 space). First, we have

$$\partial_c \mathbf{H}_k^\perp = \mathbf{H}_k^\perp \partial_c + \gamma e^{\gamma c} (1 - \Pi_k) V (1 - \Pi_k)$$

thus applying $\mathbf{R}_k^\perp(\alpha)$ on the left and right:

$$\mathbf{R}_k^\perp(\alpha) \partial_c = \partial_c \mathbf{R}_k^\perp(\alpha) + \gamma \mathbf{R}_k^\perp(\alpha) e^{\gamma c} V \mathbf{R}_k^\perp(\alpha).$$

Here the last term really is $\mathbf{R}_k^\perp(\alpha) \iota^T e^{\gamma c} V \iota \mathbf{R}_k^\perp(\alpha)$, viewed as a bounded map $L^2(\mathbb{R}^-; E_k^\perp) \rightarrow \mathcal{D}(\mathcal{Q}_k^\perp)$ using $e^{\gamma c} V : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$. Therefore (using $[\partial_c, V] = 0$)

$$\begin{aligned} \partial_c \mathbf{K}_k^\perp(\alpha) &= \gamma \mathbf{K}_k^\perp(\alpha) + e^{\gamma c} \tilde{\chi}' \Pi_k V \mathbf{R}_k^\perp(\alpha) (1 - \Pi_k) \chi \\ &\quad + \tilde{\chi} e^{\gamma c} \Pi_k V \mathbf{R}_k^\perp(\alpha) \partial_c \chi - \gamma e^{\gamma c} \tilde{\chi} \Pi_k V \mathbf{R}_k^\perp(\alpha) e^{\gamma c} V \mathbf{R}_k^\perp(\alpha) \chi. \end{aligned}$$

Just as above, the first two terms map $\mathcal{D}'(\mathcal{Q})$ to $e^{\gamma\rho/2}L^2(\mathbb{R}; E_k)$. Next, $\partial_c\chi : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}'(\mathcal{Q})$, $\mathbf{R}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$ and $\tilde{\chi}e^{\gamma c}\Pi_k V : \mathcal{D}(\mathcal{Q}) \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ are bounded (using again Lemma 6.1), so

$$\tilde{\chi}e^{\gamma c}\Pi_k V \mathbf{R}_k^\perp(\alpha) \partial_c \chi : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\gamma\rho/2}L^2(\mathbb{R}; E_k)$$

is bounded. Finally, by Lemma 6.1

$$\tilde{\chi}e^{\gamma c}\Pi_k V \mathbf{R}_k^\perp(\alpha) : \mathcal{D}'(\mathcal{Q}_k^\perp) \rightarrow e^{\gamma\rho/2}L^2(\mathbb{R}; E_k), \quad e^{\gamma c}V \mathbf{R}_k^\perp(\alpha) \chi : L^2 \rightarrow \mathcal{D}'(\mathcal{Q}_k^\perp)$$

is bounded so

$$e^{\gamma c}\tilde{\chi}\Pi_k V \mathbf{R}_k^\perp(\alpha) e^{\gamma c}V \mathbf{R}_k^\perp(\alpha) \chi : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\gamma\rho/2}L^2(\mathbb{R}; E_k)$$

is also bounded and therefore $\partial_c \mathbf{K}_k^\perp(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\gamma\rho/2}L^2(\mathbb{R}; E_k)$. This shows that

$$\mathbf{K}_k^\perp(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\gamma\rho/2}H_0^1(\mathbb{R}; E_k).$$

It is then easy to see this is a compact operator on L^2 as announced since $e^{\gamma\rho/2}H_0^1(\mathbb{R}; E_k)$ injects compactly into $e^{\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta < \gamma/2$ by using that E_k has finite dimension. We conclude that $\mathbf{K}_k^\perp(\alpha)$ is compact as a map $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ if $\beta < \gamma/2$. Moreover $\mathbf{K}_k^\perp(\alpha)$, $\mathbf{L}_k^\perp(\alpha)$ are holomorphic in $\alpha \in \mathbb{C}$ in the region $\{\operatorname{Re}(\alpha) < Q\} \cup \{|\alpha - Q|^2 < \lambda_k\}$ since $\mathbf{R}_k^\perp(\alpha)$ is. This concludes the proof of 1).

Let us next consider the region $\{\operatorname{Re}(\alpha) \leq Q\}$, and we proceed as in Lemma 6.5. Let $\mathbf{H}_{k,\beta}^\perp := e^{\beta\rho}\mathbf{H}_k^\perp e^{-\beta\rho}$ for $\beta \in \mathbb{R}$ which is also given by

$$\mathbf{H}_{k,\beta}^\perp = \mathbf{H}_k^\perp + (1 - \Pi_k) \left(-\frac{\beta^2}{2}(\rho'(c))^2 + \frac{\beta}{2}\rho''(c) + \beta\rho'(c)\partial_c \right)$$

and the associated sesquilinear form on $\mathcal{D}(\mathcal{Q}_k^\perp)$

$$\mathcal{Q}_{k,\beta}^\perp(u) := \mathcal{Q}_k^\perp(u) - \frac{\beta^2}{2}\|\rho'u\|_2^2 + \frac{\beta}{2}\langle \rho''u, u \rangle_2 + \beta\langle \partial_c u, \rho'u \rangle_2.$$

Note that on $\mathcal{D}(\mathcal{Q}_k^\perp)$, we have

$$\operatorname{Re}(\mathcal{Q}_{k,\beta}^\perp(u)) \geq \mathcal{Q}_k^\perp(u) - \frac{\beta^2}{2}\|u\|_2^2 \geq \frac{1}{2}\|\partial_c u\|_2^2 + \left(\frac{Q^2 - \beta^2}{2} + \lambda_k\right)\|u\|_2^2 + \|e^{\frac{\gamma c}{2}}V^{\frac{1}{2}}u\|_2^2.$$

This implies by Lax-Milgram, just as in the proof of Lemma 6.5, that if $\operatorname{Re}((\alpha - Q)^2) > \beta^2 - 2\lambda_k$, then for each $f \in \mathcal{D}'(\mathcal{Q}_k^\perp)$, there is a unique $u \in \mathcal{D}(\mathcal{Q}_k^\perp)$ such that

$$(6.33) \quad e^{\beta\rho}(\mathbf{H}_k^\perp - 2\Delta_\alpha)e^{-\beta\rho}u = f, \text{ and if } f \in L^2$$

$$(6.34) \quad \|u\|_2 \leq \frac{2\|f\|_2}{\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2}, \quad \|\partial_c u\|_2 \leq \frac{2\|f\|_2}{\sqrt{\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2}}.$$

In particular, this shows that, for $\operatorname{Re}((\alpha - Q)^2) > \beta^2 - 2\lambda_k$, $\mathbf{R}_k^\perp(\alpha)$ extends as a map

$$\mathbf{R}_k^\perp(\alpha) : e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}_k^\perp) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q}_k^\perp)$$

with $\|\mathbf{R}_k^\perp(\alpha)\|_{\mathcal{L}(e^{\beta\rho}L^2)} \leq 2(\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2)^{-1}$. If we further impose that $\operatorname{Re}((\alpha - Q)^2) > \beta^2 - 2\lambda_k + 1$ then, since $e^{\beta\rho}[\partial_c, e^{-\beta\rho}] = -\beta\rho'$ and using (6.34),

$$\|\mathbf{L}_k^\perp(\alpha)\|_{\mathcal{L}(e^{-\beta\rho}L^2)} \leq \frac{2|\chi'|_\infty(1 + |\beta|) + |\chi''|_\infty}{\sqrt{\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2}}.$$

Finally, the same argument as above for $\mathbf{K}_k^\perp(\alpha)$ shows that for $\operatorname{Re}((\alpha - Q)^2) + 2\lambda_k - \beta^2 > 1$, the operator $\mathbf{K}_k^\perp(\alpha)$ is compact from $e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ to $e^{-\beta\rho}L^2(\mathbb{R}^-; E_k)$. \square

Remark 6.12. We notice that the operators $\mathbf{R}_k^\perp(\alpha)$, $\mathbf{K}_k^\perp(\alpha)$, $\mathbf{L}_k^\perp(\alpha)$ lift as holomorphic family of operators to the region $\{\alpha \in \Sigma \mid \operatorname{Re}(\pi(\alpha)) < Q, |\pi(\alpha) - Q|^2 < \lambda_k\}$ by simply composing with the projection $\pi : \Sigma \rightarrow \mathbb{C}$.

2) The region $c \geq -1$. Next, consider the operator $\mathbf{H} - \frac{Q^2 + p^2}{2}$ on $L^2([-1, \infty); L^2(\Omega_{\mathbb{T}}))$ with Dirichlet condition at $c = -1$ (i.e. the extension associated to the quadratic form on functions supported in $c \geq -1$), and $\hat{\chi} \in C^\infty(\mathbb{R}; [0, 1])$ such that $(1 - \hat{\chi}) = 1$ on $\operatorname{supp}(1 - \chi)$ and $1 - \hat{\chi}$ supported in $(-1, \infty)$ (otherwise stated, $\hat{\chi} = 0$ on $(-1 + \delta, +\infty)$ and $\hat{\chi} = 1$ on $(-\infty, -1)$).

We will construct a quasi-compact approximate inverse to \mathbf{H} in $[-1, \infty)$ by using energy estimates and the properties of V , in particular the fact the region where $V > 0$ is small are somehow small. We show the following:

Lemma 6.13. *There is a uniform constant $C > 0$ and a bounded operator, independent of α ,*

$$\mathbf{R}_+ : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow \mathcal{D}(\mathbf{H}),$$

satisfying

$$(1 - \hat{\chi})\mathbf{R}_+(1 - \chi) : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$$

and for $\alpha = Q + ip \in \mathbb{C}$ and $k \geq 1$

$$\left(\mathbf{H} - \frac{Q^2 + p^2}{2}\right)(1 - \hat{\chi})\mathbf{R}_+(1 - \chi) = (1 - \chi) + \mathbf{K}_{+,k}(\alpha) + \mathbf{L}_{+,k}(\alpha)$$

where $\mathbf{K}_{+,k}(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2([-1, \infty) \times \Omega_{\mathbb{T}})$ compact and holomorphic in $\alpha \in \mathbb{C}$, and the operator $\mathbf{L}_{+,k}(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2([-1, \infty) \times \Omega_{\mathbb{T}})$ is bounded and holomorphic in $\alpha \in \mathbb{C}$, such that

$$(6.35) \quad \|\mathbf{L}_{+,k}(\alpha)\|_{\mathcal{L}(L^2)} \leq C(1 + |p|^2 \lambda_k^{-1/2}), \quad \|\mathbf{L}_{+,k}(\alpha)^2\|_{\mathcal{L}(L^2)} \leq C(1 + |p|^2) \lambda_k^{-1/2} + C(1 + |p|^4) \lambda_k^{-1}$$

for some uniform constant C depending only on $\hat{\chi}$. Moreover $\mathbf{L}_{+,k}(\alpha)$ and $\mathbf{K}_{+,k}(\alpha)$ are bounded as maps $\mathcal{D}'(\mathcal{Q}) \rightarrow L^2([-1, \infty) \times \Omega_{\mathbb{T}})$.

Proof. We consider $\mathbf{R}_+ := (\mathbf{H} - \frac{Q^2}{2} + 1)^{-1} : L^2 \rightarrow \mathcal{D}(\mathbf{H})$, which we also view as a map $\mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$. We have

$$(6.36) \quad \frac{1}{2} \|\partial_c \mathbf{R}_+ f\|_2^2 + \|\mathbf{R}_+ f\|_2^2 + \|\mathbf{P}^{1/2} \mathbf{R}_+ f\|_2^2 + \mathcal{Q}_{e^{\gamma c} V}(\mathbf{R}_+ f) \leq \|f\|_2^2,$$

We write

$$(6.37) \quad \begin{aligned} \left(\mathbf{H} - \frac{Q^2 + p^2}{2}\right)(1 - \hat{\chi})\mathbf{R}_+(1 - \chi) &= (1 - \chi) + \frac{1}{2}[\partial_c^2, \hat{\chi}]\mathbf{R}_+(1 - \chi) - \left(\frac{p^2}{2} - 1\right)(1 - \hat{\chi})\mathbf{R}_+(1 - \chi). \\ &=: (1 - \chi) + \mathbf{K}_+^1 + \mathbf{K}_+^2(\alpha). \end{aligned}$$

Notice that $\mathbf{K}_+^1, \mathbf{K}_+^2(\alpha)$ are bounded as maps $\mathcal{D}'(\mathcal{Q}) \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ by using that $[\partial_c^2, \hat{\chi}] : \mathcal{D}(\mathcal{Q}) \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ is bounded. By Lemma 6.14, there is $\beta > 0$ such that have (here $c_+ := c \mathbf{1}_{c > 0}$.)

$$(6.38) \quad \Pi_k \mathbf{R}_+ : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta c_+} L^2(\mathbb{R}; E_k)$$

is bounded. Let $\psi \in C_c^\infty(\mathbb{R})$ non-negative, equal to 1 in $[-A, A]$ and 0 for $|c| > 2A$ and so that $\|\psi\|_\infty + \|\psi'\|_\infty + \|\psi''\|_\infty \leq 1$ (we shall take $A \rightarrow \infty$ later). By using Lemma 6.1, we have

$$\Pi_k \psi : \mathcal{D}(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$$

is bounded: indeed, $\partial_c(\psi u) = \psi' u + \psi \partial_c u \in L^2$ if $u \in \mathcal{D}(\mathcal{Q})$ and

$$\int_{\mathbb{R}} \psi^2(c) e^{\gamma c} \mathcal{Q}_V(u, u) dc \leq C_{\gamma, \psi} \mathcal{Q}(u, u)$$

for some $C_{\gamma, \psi}$ depending on γ and ψ .

We are going to show that for all such ψ ,

$$(6.39) \quad \|(1 - \Pi_k)\psi \mathbf{R}_+\|_{L^2 \rightarrow L^2} \leq 3/\sqrt{\lambda_k},$$

which by taking $A \rightarrow \infty$ shows that $(1 - \Pi_k)\mathbf{R}_+$ has $\mathcal{L}(L^2)$ norm bounded by $3/\sqrt{\lambda_k}$. Let $u = \mathbf{R}_+ f$ with $f \in L^2$, then we claim that

$$(6.40) \quad \begin{aligned} \lambda_k \|(1 - \Pi_k)\psi u\|_2^2 + \frac{1}{2} \|\partial_c(1 - \Pi_k)\psi u\|_2^2 + \frac{1}{2} \mathcal{Q}_{e^{\gamma c} V}((1 - \Pi_k)\psi u) \\ \leq \|(1 - \Pi_k)\psi f\|_2 \|(1 - \Pi_k)\psi u\|_2 + \frac{1}{2} \mathcal{Q}_{e^{\gamma c} V}(\Pi_k \psi u) + 2\|f\|_2^2 \end{aligned}$$

To prove this, we note that, in $\mathcal{D}'(\mathcal{Q})$

$$\left(\mathbf{H} - \frac{Q^2}{2} + 1\right)(1 - \Pi_k)\psi u = \psi(1 - \Pi_k)f + \psi(\Pi_k e^{\gamma c} V - e^{\gamma c} V \Pi_k)u - \frac{1}{2} \psi''(1 - \Pi_k)u - \psi'(1 - \Pi_k)\partial_c$$

thus pairing with $(1 - \Pi_k)\psi u \in \mathcal{D}(\mathcal{Q})$, this gives

$$\begin{aligned} \mathcal{Q}((1 - \Pi_k)\psi u, (1 - \Pi_k)\psi u) &= \langle (1 - \Pi_k)\psi f, (1 - \Pi_k)\psi u \rangle_2 - \mathcal{Q}_{e^{\gamma c V}}(\psi \Pi_k u, \psi(1 - \Pi_k)u) \\ &\quad - \frac{1}{2} \langle \psi''(1 - \Pi_k)u, (1 - \Pi_k)\psi u \rangle_2 - \langle \psi'(1 - \Pi_k)\partial_c u, (1 - \Pi_k)\psi u \rangle_2 \end{aligned}$$

This gives the bound (6.40), using Cauchy-Schwarz, $\|\mathbf{P}^{1/2}(1 - \Pi_k)u\|_{L^2(\Omega_{\mathbb{T}})}^2 \geq \lambda_k \|u\|_{L^2(\Omega_{\mathbb{T}})}^2$ and the fact that $\|u\|_{\mathcal{D}(\mathcal{Q})} \leq \|f\|_2$ (by (6.36)). Now we do the same computation with $(1 - \Pi_k)$ replaced by Π_k :

$$(6.41) \quad \begin{aligned} &\|\Pi_k \psi u\|_2^2 + \frac{1}{2} \|\partial_c \Pi_k \psi u\|_2^2 + \frac{1}{2} \mathcal{Q}_{e^{\gamma c V}}(\Pi_k \psi u) \\ &\leq \|\Pi_k \psi f\|_2 \|\Pi_k \psi u\|_2 + \frac{1}{2} \mathcal{Q}_{e^{\gamma c V}}((1 - \Pi_k)\psi u) + 2\|f\|_2^2. \end{aligned}$$

Combining (6.41) and (6.40) and using $\|u\|_2 \leq \|f\|_2$, we obtain

$$(6.42) \quad \|(1 - \Pi_k)\psi \mathbf{R}_+ f\|_2 = \|(1 - \Pi_k)\psi u\|_2 \leq \frac{3}{\sqrt{\lambda_k}} \|f\|_2$$

Since $[\partial_c^2, \hat{\chi}] = \hat{\chi}'' + 2\hat{\chi}'\partial_c$ and $\hat{\chi}' = 0$ on $\text{supp}(1 - \chi)$, we have $(\mathbf{K}_+^1)^2 = 0$ and $\|\mathbf{K}_+^1\|_{\mathcal{L}(L^2)} \leq C$ (using (6.36)) for some uniform C depending only on $\hat{\chi}$. By combining with (6.42), we deduce that

$$\begin{aligned} &\|(\mathbf{K}_+^1 + (1 - \Pi_k)\mathbf{K}_+^2(\alpha))\|_{\mathcal{L}(L^2)} \leq C(1 + |p|^2 \lambda_k^{-1/2}), \\ &\|(\mathbf{K}_+^1 + (1 - \Pi_k)\mathbf{K}_+^2(\alpha))^2\|_{\mathcal{L}(L^2)} \leq C((1 + |p|^2)\lambda_k^{-1/2} + (1 + |p|^2)^2 \lambda_k^{-1}) \end{aligned}$$

for some uniform C depending only on $\hat{\chi}$. Next we consider the operator $\Pi_k \mathbf{K}_+^2(\alpha)$. Recall that, by (6.36),

$$(6.43) \quad \partial_c \Pi_k \mathbf{R}_+ = \Pi_k \partial_c \mathbf{R}_+ : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$$

is bounded. Now we claim that the injection

$$(6.44) \quad F_k := \{u \in e^{-\beta c} L^2([-1, \infty); E_k) \mid \partial_c u \in L^2([-1, \infty); E_k)\} \rightarrow e^{-\frac{\beta c}{2}} L^2([-1, \infty) \times \Omega_{\mathbb{T}})$$

is compact if we put the norm $\|u\|_{F_k} := \|e^{\beta c} u\|_2 + \|\partial_c u\|_2$ on F_k . Indeed, consider the operator $\eta_T \text{Id} : F_k \rightarrow e^{-\frac{\beta c}{2}} L^2([-1, \infty); E_k)$ where $\eta_T(c) = \eta(c/T)$ if $\eta \in C_0^\infty((-2, 2))$ is equal to 1 on $(-1, 1)$ and $0 \leq \eta \leq 1$. Since E_k has finite dimension, this is a compact operator by the compact embedding $H^1([-1, T]; E_k) \rightarrow e^{-\frac{\beta c}{2}} L^2([-1, \infty); E_k)$, and as $T \rightarrow \infty$ we have for $u \in F_k$

$$\|e^{\frac{\beta c}{2}} (\eta_T u - u)\|_2^2 \leq e^{-\beta T} \int_{-1}^\infty (1 - \eta_T)^2 e^{2\beta c} \|u\|_{L^2(\Omega_{\mathbb{T}})}^2 dc \leq e^{-\beta T} \|u\|_{F_k}^2$$

thus the injection (6.44) is a limit of compact operators for the operator norm topology, therefore is compact.

By (6.38) and (6.43), the operator $\Pi_k \mathbf{R}_+ : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\frac{\beta c}{2} c} L^2([-1, \infty) \times \Omega_{\mathbb{T}})$ is compact. We get that

$$\Pi_k \mathbf{K}_+^2(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R} \times E_k)$$

is compact. This complete the proof by setting $\mathbf{K}_{+,k}(\alpha) := \Pi_k \mathbf{K}_+^2(\alpha)$ and $\mathbf{L}_{+,k}(\alpha) := \mathbf{K}_+^1 + (1 - \Pi_k)\mathbf{K}_+^2(\alpha)$. The holomorphy in $\alpha \in \mathbb{C}$ is clear since $\mathbf{K}_+^2(\alpha)$ is polynomial in α . \square

Lemma 6.14. *For $\alpha = Q + ip$ with $p \in i\mathbb{R}^*$, there is $\beta > 0$ such that the resolvent $\Pi_k \mathbf{R}(\alpha) : L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ is bounded.*

Proof. We estimate the norm for $e^{-\beta c} L^2(\mathbb{R}_+ \times \Omega_{\mathbb{T}})$ since the estimate for $e^{-\beta c} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})$ results from Proposition 5.2. Recall (6.10) with the representation for the resolvent $\mathbf{R}(\alpha) = \int_0^\infty e^{-t\mathbf{H} + t^2 \Delta_\alpha} dt$, valid for the range of α we consider. A key observation to establish our lemma is the following estimate on $e^{-t\mathbf{H}}$, based on the Feynman-Kac formula (5.3). Here we write for simplicity $V_t := \int_{\mathbb{D}_t^c} |z|^{-\gamma Q} M_\gamma(dz)$. By using the Markov property, we get that for $f \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and $a \in (0, 1)$,

$$(6.45) \quad \begin{aligned} |e^{-t\mathbf{H}} f| &= |e^{-\frac{Q^2}{2} t} \mathbb{E}_\varphi[f(c + B_t, \varphi_t) e^{-\mu e^{\gamma c} V_t}]| \\ &\leq e^{-\frac{Q^2}{2} t} \mathbb{E}_\varphi[|f(c + B_t, \varphi_t)| e^{-\mu e^{\gamma c} V_{(1-a)t}}] \\ &= e^{-\frac{Q^2}{2} (1-a)t} \mathbb{E}_\varphi[e^{-at\mathbf{H}_0} |f|(c + B_{(1-a)t}, \varphi_{(1-a)t}) e^{-\mu e^{\gamma c} V_{(1-a)t}}]. \end{aligned}$$

Now we want to exploit this relation in the integral representation of the resolvent. Take $q \in [1, 2)$. By using in turn the fact that Π_k has finite dimensional rank (in particular we have equivalence of norms on its rank) and the continuity of the map $\Pi_k : L^q(\Omega_{\mathbb{T}}) \rightarrow L^q(\Omega_{\mathbb{T}})$ (see [Ja, Th 5.14]), we obtain for some constant $C = C(t_0, \alpha, k, q, \mu, \gamma)$ which may change along the lines below

$$\begin{aligned} & \int_0^\infty e^{2\beta c} \mathbb{E} \left[\left(\Pi_k \int_0^\infty e^{-t\mathbf{H}+t2\Delta_\alpha} f dt \right)^2 \right] dc \\ & \leq C \int_0^\infty e^{2\beta c} \mathbb{E} \left[\left(\Pi_k \int_0^\infty e^{-t\mathbf{H}+t2\Delta_\alpha} f dt \right)^q \right]^{2/q} dc \\ & \leq C \int_0^\infty e^{2\beta c} \mathbb{E} \left[\left| \int_0^\infty e^{-t\mathbf{H}+t2\Delta_\alpha} f dt \right|^q \right]^{2/q} dc. \end{aligned}$$

We are going to estimate this quantity by splitting the dt -integral above in two parts, $\int_0^{t_0} \dots$ and $\int_{t_0}^\infty \dots$ for some $t_0 > 0$, which we call respectively A_1 and A_2 .

Let us first focus on the first part. Using the relation (6.45) with $a = 1/2$ produces the bound

$$A_1 \leq C \int_0^\infty e^{2\beta c} \mathbb{E} \left[\left| \int_0^{t_0} \mathbb{E}_\varphi \left[e^{-\frac{t}{2}\mathbf{H}_0} |f|(c + B_{t/2}, \varphi_{t/2}) e^{-\mu e^{\gamma c} V_{t/2}} \right] dt \right|^q \right]^{2/q} dc.$$

Then Jensen gives the estimate

$$A_1 \leq C \int_0^\infty e^{2\beta c} \int_0^{t_0} \mathbb{E} \left[\left(e^{-\frac{t}{2}\mathbf{H}_0} |f|(c + B_{t/2}, \varphi_{t/2}) \right)^q e^{-q\mu e^{\gamma c} V_{t/2}} \right]^{2/q} dt dc.$$

Now we use Hölder's inequality with conjugate exponents $2/q, 2/(2-q)$ to estimate the above quantity by

$$A_1 \leq C \int_0^\infty \int_0^{t_0} e^{2\beta c} \mathbb{E} \left[\left(e^{-\frac{t}{2}\mathbf{H}_0} |f|(c + B_{t/2}, \varphi_{t/2}) \right)^2 \right] \left(\mathbb{E} \left[e^{-\frac{2-q}{2-q}\mu e^{\gamma c} V_{t/2}} \right] \right)^{\frac{2-q}{q}} dt dc.$$

Using the elementary inequality $x^\theta e^{-x} \leq C$ for $\theta > 0$, we deduce

$$A_1 \leq C \int_0^\infty \int_0^{t_0} e^{(2\beta - \theta \frac{2-q}{q})c} \mathbb{E} \left[\left(e^{-\frac{t}{2}\mathbf{H}_0} |f|(c + B_{t/2}, \varphi_{t/2}) \right)^2 \right] \left(\mathbb{E} \left[V_{t/2}^{-\theta} \right] \right)^{\frac{2-q}{q}} dt dc.$$

It remains to estimate the quantity $\mathbb{E} \left[V_{t/2}^{-\theta} \right]$. Notice that $V_{t/2}$ is a GMC over an annulus of radii 1 and $e^{-t/2}$. Hence the quantity to be investigated is less than the same expression with $V_{t/2}$ replaced by a GMC over a ball of diameter $t/2$ contained in this annulus. Standard computations of GMC multifractal spectrum [RhVa14, Th 2.14] (also valid for negative moments) then yields that

$$(6.46) \quad \mathbb{E} \left(V_{t/2}^{-\theta} \right) \leq C t^{\psi(-\theta)}$$

where $\psi(u) := (2 + \frac{\gamma^2}{2})u - \frac{\gamma^2}{2}u^2$ is the multifractal spectrum. Since ψ is continuous and $\psi(0) = 0$, we can find $\theta > 0$ such that $\psi(-\theta) > -1$. Then, for $2\beta = \frac{2-q}{q}\theta\gamma$, we deduce that

$$\begin{aligned} A_1 & \leq C \int_0^\infty \int_0^{t_0} \mathbb{E} \left[\left(e^{-\frac{t}{2}\mathbf{H}_0} |f|(c + B_{t/2}, \varphi_{t/2}) \right)^2 \right] t^{\psi(-\theta)} dt dc \\ & \leq C \int_0^{t_0} \|e^{-\frac{t}{2}\mathbf{H}_0} |f|\|_2^2 t^{\psi(-\theta)} dt \\ & \leq C \|f\|_2^2 \int_0^{t_0} t^{\psi(-\theta)} dt, \end{aligned}$$

which proves our claim for A_1 , namely the part corresponding to $\int_0^{t_0}$.

Concerning A_2 , we start as above

$$\begin{aligned} A_2 & := \int_0^\infty e^{2\beta c} \mathbb{E} \left[\left(\Pi_k \int_{t_0}^\infty e^{-t\mathbf{H}+t2\Delta_\alpha} f dt \right)^2 \right] dc \\ & \leq C \int_0^\infty e^{2\beta c} \left(\mathbb{E} \left[\left(\int_{t_0}^\infty e^{-t\mathbf{H}+t2\Delta_\alpha} f dt \right)^q \right] \right)^{2/q} dc \end{aligned}$$

for $q \in (1, 2)$. For the values of α we consider, we have $2\Delta_\alpha \leq \frac{Q^2}{2} - \frac{|p|^2}{2}$ so that, using Jensen,

$$A_2 \leq C \int_0^\infty e^{2\beta c} \left(\mathbb{E} \left[\int_{t_0}^\infty e^{-\frac{|p|^2}{2}t} \left(\mathbb{E}_\varphi [|f(c + B_t, \varphi_t)| e^{-\mu e^{\gamma c} V_t}] \right)^q dt \right] \right)^{2/q} dc$$

Let us now fix $a \in (0, 1)$ small enough such that $aQ^2 < |p|^2/2$. Then, using (6.45), Jensen and the inequality $x^\theta e^{-x} \leq C$ for $\theta > 0$, we deduce

$$A_2 \leq C \int_0^\infty e^{2(\beta - \gamma\theta/q)c} \left(\mathbb{E} \left[\int_{t_0}^\infty e^{-\frac{|p|^2}{4}t} \mathbb{E}_\varphi \left[\left(e^{-at\mathbf{H}_0} |f|(c + B_{(1-a)t}, \varphi_{(1-a)t}) \right)^q V_{(1-a)t}^{-\theta} \right] dt \right] \right)^{2/q} dc.$$

Now we can proceed similarly as above by using the fact that for $a < 1$ and all $q > 0$ $\sup_{t>t_0} \mathbb{E}[V_{(1-a)t}^{-\theta}] < \infty$ (this quantity is decreasing in t and GMC has negative moments) to get

$$\begin{aligned} A_2 &\leq C \int_0^\infty e^{2(\beta - \gamma\theta/q)c} \left(\int_{t_0}^\infty e^{-\frac{|p|^2}{4}t} \mathbb{E} \left[\left(e^{-at\mathbf{H}_0} |f|(c + B_{(1-a)t}, \varphi_{(1-a)t}) \right)^2 \right]^{q/2} \mathbb{E} \left[V_{(1-a)t}^{-\frac{2\theta}{2-q}} \right]^{\frac{2-q}{2}} dt \right)^{2/q} dc \\ &\leq C \int_0^\infty e^{2(\beta - \gamma\theta/q)c} \left(\int_{t_0}^\infty e^{-\frac{|p|^2}{4}t} \mathbb{E} \left[\left(e^{-at\mathbf{H}_0} |f|(c + B_{(1-a)t}, \varphi_{(1-a)t}) \right)^2 \right]^{q/2} dt \right)^{2/q} dc. \end{aligned}$$

For $\beta = \theta\gamma/q$ and using Jensen, the above quantity is less than

$$C \int_{t_0}^\infty e^{-\frac{|p|^2}{4}t} \|e^{-at\mathbf{H}_0} |f|\|_2^2 dt \leq C \|f\|_2^2 \int_{t_0}^\infty e^{-\frac{|p|^2}{4}t} dt.$$

Hence the result. \square

Remark 6.15. *As above, the operators \mathbf{R}_+ , $\mathbf{K}_{+,k}(\alpha)$ and $\mathbf{L}_{+,k}(\alpha)$, lift as holomorphic family of operators to Σ .*

3) Small \mathbf{P} eigenmodes in the region $c \leq 0$, where there is scattering. We will view the operator \mathbf{H} as a perturbation of the free Hamiltonian $\mathbf{H}_0 := -\frac{1}{2}\partial_c^2 + \frac{Q^2}{2} + \mathbf{P}$ on $L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}})$ with Dirichlet condition at $c = 0$. We show (recall that $\pi : \Sigma \rightarrow \mathbb{C}$ is the covering map)

Lemma 6.16. *1) Fix k and $0 < \beta < \gamma/2$. The operators*

$$(6.47) \quad \mathbf{R}_k(\alpha) := \left(\mathbf{H}_0 - \frac{Q^2 + p^2}{2} \right)^{-1} \Pi_k : e^{\beta\rho} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} L^2(\mathbb{R}^-; E_k)$$

defined for $\text{Im}(p) > 0$ can be holomorphically continued to the region

$$(6.48) \quad \{ \alpha = Q + ip \in \Sigma \mid \forall j \leq k, \text{Im}\sqrt{p^2 - 2\lambda_j} > -\beta \}.$$

This continuation, still denoted $\mathbf{R}_k(\alpha) : e^{\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} L^2(\mathbb{R}^-; E_k)$, satisfies

$$\widetilde{\chi} \mathbf{R}_k(\alpha) \chi : e^{\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} (L^2(\mathbb{R}^-; E_k) \cap \mathcal{D}(\mathcal{Q})),$$

$$\left(\mathbf{H} - \frac{Q^2 + p^2}{2} \right) \widetilde{\chi} \mathbf{R}_k(\alpha) \chi = \Pi_k \chi + \mathbf{K}_{k,1}(\alpha) + \mathbf{K}_{k,2}(\alpha)$$

where $\mathbf{K}_{k,1}(\alpha)$, $\mathbf{K}_{k,2}(\alpha)$ are such that for $\text{Im}\sqrt{p^2 - 2\lambda_j} > -\min(\beta, \gamma/2 - \beta)$

$$\mathbf{K}_{k,1}(\alpha) : e^{\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$$

$$\mathbf{K}_{k,2}(\alpha) : e^{\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta\rho} \mathcal{D}'(\mathcal{Q})$$

are holomorphic families of compact operators in (6.48), and we have $\mathbf{K}_{k,i}(\alpha)(1 - \Pi_k) = 0$ for $i = 1, 2$, $(1 - \Pi_k)\mathbf{K}_{k,1}(\alpha) = 0$ and $\Pi_k\mathbf{K}_{k,2}(\alpha) = 0$.

2) If $f \in e^{\beta\rho} L^2$, then there is $C_k > 0$ depending on k , some $a_j(\alpha, f)$ and $G(\alpha, f) \in H^1(\mathbb{R}; E_k)$ depending linearly on f and holomorphic in $\{ \alpha = Q + ip \in \Sigma \mid \forall j \leq k, \text{Im}\sqrt{p^2 - 2\lambda_j} > -\min(\beta, \gamma/2 - \beta) \}$ such that in the region $c \leq 0$

$$(6.49) \quad (\mathbf{R}_k(\alpha)f) = \sum_{\lambda_j \leq \lambda_k} a_j(\alpha, f) e^{-ic\sqrt{p^2 - 2\lambda_j}} + G(\alpha, f),$$

$$\|G(\alpha, f)(c)\|_{L^2(\Omega_{\mathbb{T}})} + \|\partial_c G(\alpha, f)(c)\|_{L^2(\Omega_{\mathbb{T}})} \leq C_k e^{\beta\rho} \|e^{-\beta\rho} \Pi_k f\|_2.$$

3) For each $\beta \in \mathbb{R}$, the operator $\mathbf{R}_k(\alpha)$ extends as a bounded analytic family

$$\mathbf{R}_k(\alpha) : e^{-\beta\rho} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} (L^2(\mathbb{R}^-; E_k) \cap \mathcal{D}(\mathcal{Q}))$$

in the region $\text{Im}(p) > |\beta|$ and $\mathbf{K}_{k,1}(\alpha) : e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$, $\mathbf{K}_{k,2}(\alpha) : e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho} \mathcal{D}'(\mathcal{Q})$ are compact analytic families in that same region.

Proof. We first consider \mathbf{H}_0 on $(-\infty, 0]$ with Dirichlet condition at $c = 0$. Using the diagonalisation of \mathbf{P} on E_k , we can compute the resolvent $(\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1}$ on E_k by standard ODE methods (Sturm-Liouville theory): for $\text{Im}(p) > 0$, this is the diagonal operator given for $j \leq k$ and $f \in L^2(\mathbb{R}^-)$ and $\phi_j \in \ker(\mathbf{P} - \lambda_j)$

$$\begin{aligned} (\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} f(c) \phi_j &= \frac{2}{\sqrt{p^2 - 2\lambda_j}} \phi_j \left(\int_{-\infty}^c \sin(c\sqrt{p^2 - 2\lambda_j}) e^{-ic'\sqrt{p^2 - 2\lambda_j}} f(c') dc' \right. \\ &\quad \left. + \int_c^0 e^{-ic'\sqrt{p^2 - 2\lambda_j}} \sin(c'\sqrt{p^2 - 2\lambda_j}) f(c') dc' \right) \end{aligned}$$

where our convention is that \sqrt{z} is defined with the cut on \mathbb{R}^+ , so that $\sqrt{p^2} = p$ if $\text{Im}(p) > 0$. For $j = 0$, that is $\phi_0 = 1$, for each $\beta > 0$ the resolvent restricted to E_0 admits an analytic continuation from $\text{Im}(p) > 0$ to $\text{Im}(p) > -\beta$, as a map

$$(\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} \Pi_0 : e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} L^2(\mathbb{R}^-; E_0).$$

This is easy to see by using Schur's lemma and the analyticity in p for the Schwartz kernel

$$\kappa_0(c, c') := \mathbf{1}_{\{c \geq c'\}} e^{-\beta(|c|+|c'|)} \sin(cp) e^{-ic'p} + \mathbf{1}_{\{c' \geq c\}} e^{-\beta(|c'|+|c|)} \sin(c'p) e^{-icp}$$

of the operator $\Pi_0 e^{-\beta|c|} (\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} e^{-\beta|c|} \Pi_0$ that we view as an operator on $L^2(\mathbb{R}^-)$. Moreover, one directly also obtains that it maps $e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta|c|} H^2(\mathbb{R}^-; E_0) \cap H_0^1(\mathbb{R}^-; E_0)$. Similarly, the operators

$$(\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} \phi_j \langle \phi_j, \cdot \rangle : e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} L^2(\mathbb{R}^-; \mathbb{C} \phi_j)$$

are analytic in p , which implies that

$$\mathbf{R}_k(\alpha) := (\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} \Pi_k : e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} H^2(\mathbb{R}^-; E_k) \cap H_0^1(\mathbb{R}^-; E_k)$$

admits an analytic extension in p to the region $\{p \in Z \mid \forall j \geq 0, \text{Im}(\sqrt{p^2 - 2\lambda_j}) > -\beta\}$ of the ramified Riemann surface Z . The proof of Lemma 6.1 also shows that $\mathcal{Q}_{e^{\gamma c} V}(\tilde{\chi} \Pi_k u) < \infty$ if $u \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and clearly also $\mathbf{P}^{\frac{1}{2}} \Pi_k \in \mathcal{L}(L^2(\Omega_{\mathbb{T}}))$, thus we deduce that

$$\tilde{\chi} \mathbf{R}_k(\alpha) \chi : e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} (L^2(\mathbb{R}^-; E_k) \cap \mathcal{D}(\mathcal{Q}))$$

is bounded. We have (using $\Pi_k \mathbf{R}_k(\alpha) = \mathbf{R}_k(\alpha)$ and Lemma 6.1)

$$\begin{aligned} (\mathbf{H} - \frac{Q^2+p^2}{2}) \tilde{\chi} \mathbf{R}_k(\alpha) \chi &= \Pi_k \chi - \frac{1}{2} [\partial_c^2, \tilde{\chi}] \Pi_k \mathbf{R}_k(\alpha) \chi + e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi \\ &= \Pi_k \chi + \frac{1}{2} [\partial_c^2, \tilde{\chi}] \Pi_k \mathbf{R}_k(\alpha) \chi + \Pi_k e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi + \tilde{\chi} (1 - \Pi_k) e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi \\ &= \Pi_k \chi + \mathbf{K}_{k,1}(\alpha) + \mathbf{K}_{k,2}(\alpha) \end{aligned}$$

where $\mathbf{K}_{k,2}(\alpha) := \tilde{\chi} (1 - \Pi_k) e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi$ satisfies $\Pi_k \mathbf{K}_{k,2}(\alpha) = 0$. The operator $[\partial_c^2, \tilde{\chi}] \Pi_k \mathbf{R}_k(\alpha) \chi$ is compact on $e^{-\beta|c|} L^2(\mathbb{R}; L^2(\Omega_{\mathbb{T}}))$ since $e^{\beta|c|} [\partial_c^2, \tilde{\chi}]$ is a compactly supported first order operator in c , $E_k = \text{Im}(\Pi_k)$ is finite dimensional in $L^2(\Omega_{\mathbb{T}})$ and $\mathbf{R}_k(\alpha) : e^{-\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} H^2(\mathbb{R}^-; E_k)$ (this amounts to the compact injection $H^2([-1, 0]; E_k) \rightarrow H^1(\mathbb{R}^-; E_k)$).

The operator $e^{(\beta-\gamma/2)|c|} \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi e^{-\beta|c|}$ is also compact as maps $L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow L^2(\mathbb{R}; E_k)$ and $L^2 \rightarrow \mathcal{D}(\mathcal{Q})$ by using the same type of argument as for proving the compact injection (6.44): indeed, one has the pointwise bound on its Schwartz kernel restricted to $\ker(\mathbf{P} - \lambda_j)$

$$\begin{aligned} |\kappa_j(c, c')| &\leq C e^{(\beta-\gamma/2)|c| - \beta|c'|} (e^{\text{Im}(\sqrt{p^2 - 2\lambda_j})(|c| - |c'|)} + e^{-\text{Im}(\sqrt{p^2 - 2\lambda_j})(|c| + |c'|)}) \mathbf{1}_{|c'| \geq |c|} \\ &\quad + C e^{(\beta-\gamma/2)|c'| - \beta|c|} (e^{\text{Im}(\sqrt{p^2 - 2\lambda_j})(|c'| - |c|)} + e^{-\text{Im}(\sqrt{p^2 - 2\lambda_j})(|c| + |c'|)}) \mathbf{1}_{|c| \geq |c'|}. \end{aligned}$$

We see that for $\text{Im}(\sqrt{p^2 - 2\lambda_j}) \geq 0$, if $0 < \beta < \gamma/2$, this is bounded by $C \max(e^{(\beta-\gamma/2)|c|-\beta|c'|}, e^{(\beta-\gamma/2)|c'|-\beta|c|})$, and is thus the integral kernel of a compact operator on $L^2(\mathbb{R}^-)$ since it is Hilbert-Schmidt (the kernel being in $L^2(\mathbb{R} \times \mathbb{R})$). If now $\text{Im}(\sqrt{p^2 - 2\lambda_j}) < 0$, the same argument shows that a sufficient condition to be compact is that

$$\text{Im}\sqrt{p^2 - 2\lambda_j} > -\beta \text{ and } \text{Im}\sqrt{p^2 - 2\lambda_j} > \beta - \gamma/2.$$

Combining with Lemma 6.1, we deduce that if $0 < \beta < \gamma/2$, then

$$e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi : e^{\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta \rho} \mathcal{D}'(\mathcal{Q})$$

$$\text{and } \Pi_k e^{\gamma c} V \Pi_k \tilde{\chi} \mathbf{R}_k(\alpha) \chi : e^{\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta \rho} L^2(\mathbb{R}^-; E_k)$$

are compact operators. Thus $\mathbf{K}_{k,1}(\alpha) : e^{\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta \rho} L^2(\mathbb{R}^-; E_k)$ is also compact. This proves 1).

Now, if $f \in e^{\beta \rho} L^2(\mathbb{R}; E_k)$ we have for $c \leq 0$ and writing $f = \sum_{j \leq k} f_j$ with $f_j \in \ker(\mathbf{P} - \lambda_j)$

$$(6.50) \quad \begin{aligned} (\mathbf{R}_k(\alpha) \chi f)(c) &= 2 \sum_{j \leq k} \frac{e^{-ic\sqrt{p^2-2\lambda_j}}}{\sqrt{p^2-2\lambda_j}} \int_{-\infty}^0 \sin(c' \sqrt{p^2-2\lambda_j}) \chi(c') f_j(c') dc' \\ &\quad + 2 \sum_{j \leq k} \frac{e^{-ic\sqrt{p^2-2\lambda_j}}}{\sqrt{p^2-2\lambda_j}} \int_{-\infty}^c \sin((c-c') \sqrt{p^2-2\lambda_j}) f_j(c') dc' \end{aligned}$$

and the term in the second line, denoted $G(c)$, satisfies for $c < -1$

$$\begin{aligned} \|G(c)\|_{L^2(\Omega_{\mathbb{T}})} &\leq 2 \sum_{j \leq k} \int_{-\infty}^c e^{|\text{Im}\sqrt{p^2-2\lambda_j}|(c-c')+\beta c'} \|e^{-\beta \rho(c')} f_j(c')\|_{L^2(\Omega_{\mathbb{T}})} dc' \\ &\leq C_k \|e^{-\beta \rho} f\|_2 e^{-\beta|c|}. \end{aligned}$$

and the same bounds hold for $\|\partial_c G(c)\|_{L^2(\Omega_{\mathbb{T}})}$; this completes the proof of 2).

We remark that if $\text{Im}(p) > 0$, then the operator

$$(\mathbf{H}_0 - \frac{Q^2+p^2}{2})^{-1} \Pi_k : e^{\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} L^2(\mathbb{R}^-; E_k)$$

is bounded and analytic in p , provided $0 < \beta < \min_{j \leq k} \text{Im}(\sqrt{p^2 - 2\lambda_j}) = \text{Im}(p)$. Indeed, this follows again from Schur's lemma applied to the Schwartz kernel

$$\mathbf{1}_{\{c \geq c'\}} e^{-\beta(|c|-|c'|)} \sin(c\sqrt{p^2-2\lambda_j}) e^{-ic'\sqrt{p^2-2\lambda_j}} + \mathbf{1}_{\{c' \geq c\}} e^{-\beta(|c'|-|c|)} \sin(c'\sqrt{p^2-2\lambda_j}) e^{-ic\sqrt{p^2-2\lambda_j}}.$$

Using again that $\Pi_k : E_k \rightarrow L^2(\Omega_{\mathbb{T}})$ is bounded, this implies that

$$\mathbf{R}_k(\alpha) : e^{\beta|c|} L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{\beta|c|} (L^2(\mathbb{R}^-; E_k) \cap \mathcal{D}(\mathcal{Q}))$$

is an analytic bounded family in p in the region $0 < \beta < \text{Im}(p)$. The same argument works with $0 < -\beta < \text{Im}(p)$ in case $\beta < 0$. The operator $\mathbf{K}_{k,1}(\alpha) : e^{-\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ is compact by the same argument as above since it is Hilbert-Schmidt and $\mathbf{K}_{k,2}(\alpha) : e^{-\beta \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta \rho} \mathcal{D}'(\mathcal{Q})$ is compact. \square

4) Proof of Proposition 6.10. For $\beta \in \mathbb{R}$, define the Hilbert space for fixed k

$$\mathcal{H}_{k,\beta} := e^{\beta \rho} L^2(\mathbb{R}; E_k) \oplus L^2(\mathbb{R}; E_k^\perp)$$

with scalar product

$$\langle f, f' \rangle_{\mathcal{H}_{k,\beta}} := \int_{\mathbb{R}} e^{-2\beta \rho(c)} \langle \Pi_k f, \Pi_k f' \rangle_{L^2(\Omega_{\mathbb{T}})} dc + \langle (1 - \Pi_k) f, (1 - \Pi_k) f' \rangle_{L^2(\mathbb{R} \times \Omega_{\mathbb{T}})}.$$

We now fix β and β' as in the statement of Proposition 6.10. We now use the operators of Lemma 6.11, Lemma 6.13 and Lemma 6.16: let $\chi; \tilde{\chi}; \hat{\chi}$ be the cutoff functions of these Lemmas and let $\check{\chi} \in C^\infty(\mathbb{R})$ equal to 1 on $\text{supp}(\tilde{\chi})$ and supported in \mathbb{R}^- . We define

$$\tilde{\mathbf{R}}(\alpha) := \tilde{\chi} \mathbf{R}_k^\perp(\pi(\alpha)) \chi + (1 - \hat{\chi}) \mathbf{R}_+(1 - \chi) + \tilde{\chi} \mathbf{R}_k(\alpha) \chi - \check{\chi} \mathbf{R}_k^\perp(\alpha) \mathbf{K}_{k,2}(\alpha)$$

which in $\{\alpha = Q + ip \in \Sigma \mid \forall j \leq k, \text{Im}\sqrt{p^2 - 2\lambda_j} > -\beta\}$ is bounded and holomorphic (in α) as a map $\tilde{\mathbf{R}}(\alpha) : \mathcal{H}_{k,\beta} \rightarrow \mathcal{H}_{k,-\beta} \cap e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$. It moreover satisfies the identity

$$\begin{aligned} (\mathbf{H} - 2\Delta_{\pi(\alpha)})\tilde{\mathbf{R}}(\alpha) &= 1 + \mathbf{L}_k^\perp(\pi(\alpha)) + \mathbf{K}_k^\perp(\pi(\alpha)) + \mathbf{K}_{+,k}(\pi(\alpha)) + \mathbf{L}_{+,k}(\pi(\alpha)) + \mathbf{K}_{k,1}(\alpha) \\ &\quad - (\check{\mathbf{L}}_k^\perp(\pi(\alpha)) + \check{\mathbf{K}}_k^\perp(\pi(\alpha)))\mathbf{K}_{k,2}(\alpha) \end{aligned}$$

where $\check{\mathbf{L}}_k^\perp(\pi(\alpha))$ and $\check{\mathbf{K}}_k^\perp(\pi(\alpha))$ are the operators of Lemma 6.11 with $\tilde{\chi}$ (resp. χ) is replaced by $\check{\chi}$ (resp. $\tilde{\chi}$). Let us define

$$(6.51) \quad \tilde{\mathbf{K}}_k(\alpha) := \mathbf{K}_{k,1}(\alpha) + \mathbf{K}_{+,k}(\pi(\alpha)) + \mathbf{K}_k^\perp(\pi(\alpha)) - (\check{\mathbf{L}}_k^\perp(\pi(\alpha)) + \check{\mathbf{K}}_k^\perp(\pi(\alpha)))\mathbf{K}_{k,2}(\alpha).$$

By Lemma 6.11, Lemma 6.13 and Lemma 6.16, $\tilde{\mathbf{K}}_k(\alpha) : \mathcal{H}_{k,\beta} \rightarrow \mathcal{H}_{k,\beta}$ is compact and holomorphic in α (recall that $\mathbf{K}_{k,j}(\alpha)(1 - \Pi_k) = 0$). We claim that for each $\psi \in C^\infty(\mathbb{R}) \cap L^\infty(\mathbb{R})$ satisfying $\psi' \in L^\infty$ and $\text{supp}(\psi) \subset (-\infty, A)$ for some $A \in \mathbb{R}$,

$$(6.52) \quad \tilde{\mathbf{K}}_k(\alpha)(1 - \Pi_k) : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{H}_{k,\beta} \quad \text{and} \quad \tilde{\mathbf{R}}(\alpha)(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$$

are bounded. Indeed, Lemma 6.11 and Lemma 6.13 show that $\mathbf{K}_{+,k}(\pi(\alpha)) + \mathbf{K}_k^\perp(\pi(\alpha))$ is bounded as a map $\mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{H}_{k,\beta}$ and that $\tilde{\chi}\mathbf{R}_k^\perp(\pi(\alpha))\chi + (1 - \tilde{\chi})\mathbf{R}_+(1 - \chi)$ is bounded as a map $\mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}(\mathcal{Q})$, while Lemma 6.1 shows that $(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$.

Now if $|\pi(\alpha) - Q|^2 \leq \lambda_k^{1/4}$ and if k is large enough, the operator $\tilde{\mathbf{L}}_k(\alpha) := \mathbf{L}_k^\perp(\pi(\alpha)) + \mathbf{L}_{+,k}(\alpha)$ is bounded as map

$$(6.53) \quad \tilde{\mathbf{L}}_k(\alpha) : \mathcal{H}_{k,\beta} \rightarrow L^2(\mathbb{R}; E_k^\perp) \quad \tilde{\mathbf{L}}_k(\alpha) : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{H}_{k,\beta}$$

with holomorphic dependance in α , and with bound (recall (6.28) and (6.35))

$$\|\tilde{\mathbf{L}}_k(\alpha)\|_{\mathcal{H}_{k,\beta} \rightarrow L^2(\mathbb{R}; E_k^\perp)} < 1/2.$$

In particular, $(1 + \tilde{\mathbf{L}}_k(\alpha))(1 - \tilde{\mathbf{L}}_k(\alpha)) = 1 - \tilde{\mathbf{L}}_k(\alpha)^2$ is invertible on $\mathcal{H}_{k,\beta}$ with holomorphic inverse given by the Neumann series $\sum_{j=0}^{\infty} \tilde{\mathbf{L}}_k(\alpha)^{2j}$; we write $(1 + \mathbf{T}_k(\alpha)) := (1 - \tilde{\mathbf{L}}_k(\alpha))(1 - \tilde{\mathbf{L}}_k(\alpha)^2)^{-1}$, with $\mathbf{T}_k(\alpha)$ mapping boundedly $\mathcal{H}_{k,\beta} \rightarrow \mathcal{H}_{k,\beta}$. Moreover we have

$$(\mathbf{H} - 2\Delta_{\pi(\alpha)})\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) = 1 + \tilde{\mathbf{K}}_k(\alpha)(1 + \mathbf{T}_k(\alpha))$$

and the remainder $\hat{\mathbf{K}}(\alpha) := \tilde{\mathbf{K}}_k(\alpha)(1 + \mathbf{T}_k(\alpha))$ is now compact on $\mathcal{H}_{k,\beta}$, and $1 + \hat{\mathbf{K}}(\alpha)$ is thus Fredholm of index 0.

Let $p_0 = iq$ for some $q \gg \beta$, the operator $\mathbf{H} - \frac{Q^2}{2}$ being self-adjoint on its domain $\mathcal{D}(\mathbf{H})$ and non-negative, $\mathbf{H} - \frac{Q^2 + p_0^2}{2}$ is invertible with inverse denoted $\mathbf{R}(\alpha_0)$ if $\alpha_0 = Q + ip_0 = Q - q$. Now, let $(\psi_j)_{j \leq J} \subset \mathcal{H}_{k,\beta}$ be an orthonormal basis of $\ker(1 + \hat{\mathbf{K}}(\alpha_0)^*)$, and $(\varphi_j)_{j \leq J} \subset \mathcal{H}_{k,\beta}$ an orthonormal basis of $\ker(1 + \hat{\mathbf{K}}(\alpha_0))$. For each j , there is $w_j \in \mathcal{D}(\mathbf{H})$ such that $(\mathbf{H} - \frac{Q^2 + p_0^2}{2})w_j = \psi_j$. If $\theta \in C^\infty(\mathbb{R})$ equal 1 in $c \in (-\infty, -1)$ and is supported in $c \in \mathbb{R}^-$, we have in $\mathcal{D}'(\mathcal{Q})$

$$(6.54) \quad (\mathbf{H}_0 - \frac{Q^2 + p_0^2}{2})\theta w_j = \theta\psi_j - \theta e^{\gamma c} V w_j - \frac{1}{2}[\partial_c^2, \theta]w_j$$

and this implies by projecting this relation on E_k with Π_k that, setting $\psi_{j,k} = \Pi_k\psi_j$ and $\psi_{j,k}^\perp = (1 - \Pi_k)\psi_j$, similarly $w_{j,k} = \Pi_k w_j$ and $w_{j,k}^\perp = (1 - \Pi_k)w_j$

$$(6.55) \quad (\mathbf{H}_0 - \frac{Q^2 + p_0^2}{2})\theta w_{j,k} = \theta\psi_{j,k} - \theta\Pi_k(e^{\gamma c} V w_j) - \frac{1}{2}[\partial_c^2, \theta]w_{j,k}.$$

By Lemma 6.1 with $\beta' = \beta = \gamma/2$, we see that $\Pi_k(e^{\gamma c} V w_j) \in e^{\gamma\rho/2}L^2$. Since $[\partial_c^2, \theta]$ is a first order differential operator with compact support, we get $[\partial_c^2, \theta]w_{j,k} \in L^2$ as $\tilde{\theta}w_{j,k} \in \mathcal{D}(\mathcal{Q})$ if $\tilde{\theta} \in C_c^\infty(\mathbb{R})$ by Lemma 6.1. The right hand side of (6.55) is then in $e^{\beta\rho}L^2$. This shows in particular that $\theta w_{j,k} \in H^2(\mathbb{R}^-; E_k) \cap H_0^1(\mathbb{R}^-; E_k)$ and since E_k is finite dimensional, it is direct to check that

$$\mathbf{R}_k(Q + ip_0)(\mathbf{H}_0 - \frac{Q^2 + p_0^2}{2})\theta w_{j,k} = \theta w_{j,k}$$

with $\mathbf{R}_k(Q + ip_0)$ the operator of Lemma 6.16. We obtain in the region $c \in \mathbb{R}^-$

$$\theta w_{j,k} = \mathbf{R}_k(Q + ip_0)(\theta\psi_{j,k} - \theta e^{\gamma c} \Pi_k(V w_j) - \frac{1}{2}[\partial_c^2, \theta]w_{j,k}).$$

Using the properties of $\mathbf{R}_k(\alpha)$ in Part 3) of Lemma 6.16 (applied with $-\beta$ instead of β), we see that for each $0 < \beta < \gamma/2$

$$\theta w_{j,k} \in e^{\beta\rho} L^2(\mathbb{R}^-; E_k)$$

and thus we deduce that

$$w_j \in e^{\beta\rho} L^2(\mathbb{R}; E_k) \oplus L^2(\mathbb{R}; E_k^\perp) = \mathcal{H}_{k,\beta}.$$

If we consider the finite rank operator \mathbf{W} defined by $\mathbf{W}f := \sum_{j=1}^J w_j \langle f, \varphi_j \rangle_{\mathcal{H}_{k,\beta}}$ for $f \in \mathcal{H}_{k,\beta}$, we have

$$\left(\mathbf{H} - \frac{Q^2 + p_0^2}{2}\right) \mathbf{W} = \mathbf{Y} \quad \text{with } \mathbf{Y}f := \sum_{j=1}^J \psi_j \langle f, \varphi_j \rangle_{\mathcal{H}_{k,\beta}}.$$

But now it is direct to check that $1 + \hat{\mathbf{K}}(\alpha_0) + \mathbf{Y}$ is invertible on $\mathcal{H}_{k,\beta}$ and we obtain that

$$(\mathbf{H} - 2\Delta_{\pi(\alpha)}) (\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) + \mathbf{W}) = 1 + \hat{\mathbf{K}}(\alpha) + \mathbf{Y} - 2(\Delta_{\pi(\alpha)} - \Delta_{\alpha_0}) \mathbf{W}.$$

The remainder $\mathbf{K}(\alpha) := \hat{\mathbf{K}}(\alpha) + \mathbf{Y} - 2(\Delta_{\pi(\alpha)} - \Delta_{\alpha_0}) \mathbf{W}$ is compact on $\mathcal{H}_{k,\beta}$, analytic in α in the desired region and $1 + \mathbf{K}(\alpha)$ is invertible for $\alpha = \alpha_0$, therefore we can apply the Fredholm analytic theorem to conclude that the family of operator $(1 + \mathbf{K}(\alpha))^{-1}$ exists as a meromorphic family of bounded operators on $\mathcal{H}_{k,\beta}$ for α in $\{\alpha = Q + ip \in \Sigma \mid |\pi(\alpha) - Q|^2 \leq \lambda_k^{1/4}, \forall j \leq k, \text{Im}\sqrt{p^2 - \lambda_j} > -\beta\}$ except on a discrete set of poles with finite rank polar part. We can thus set

$$(6.56) \quad \mathbf{R}(\alpha) := (\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) + \mathbf{W})(1 + \mathbf{K}(\alpha))^{-1}$$

which satisfies the desired properties. To prove the boundedness of $\mathbf{R}(\alpha)(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ for $\psi \in L^\infty(\mathbb{R}) \cap C^\infty(\mathbb{R})$ with $\psi' \in L^\infty$ and $\text{supp}(\psi) \subset (-\infty, A)$ for $A \in \mathbb{R}$, we write

$$\mathbf{R}(\alpha) = \tilde{\mathbf{R}}(\alpha) - \mathbf{R}(\alpha)(\tilde{\mathbf{K}}_k(\alpha) + \tilde{\mathbf{L}}_k(\alpha))$$

and we have seen in (6.52) and (6.53) that $(\tilde{\mathbf{K}}_k(\alpha) + \tilde{\mathbf{L}}_k(\alpha)) : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{H}_{k,\beta}$ is bounded and $\tilde{\mathbf{R}}(\alpha)(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ is bounded, while $(1 - \Pi_k)\psi : \mathcal{D}'(\mathcal{Q}) \rightarrow \mathcal{D}'(\mathcal{Q})$ by Lemma 6.1. Finally $\mathbf{R}(\alpha) : \mathcal{H}_{k,\beta} \rightarrow e^{\beta\rho}\mathcal{D}(\mathcal{Q})$ and the same holds for \mathbf{W} . This shows the announced property of $\mathbf{R}(\alpha)(1 - \Pi_k)\psi$. We can use the mapping properties of $\tilde{\mathbf{R}}(\alpha)$ and \mathbf{W} , together with (6.49) and (6.56) to deduce (6.26).

We finally need to prove that there is no pole in the half plane $\text{Re}(\alpha) \leq Q$ except possibly at the points $\alpha = Q \pm i\sqrt{2\lambda_j}$. First, by the spectral theorem, one has for each $f \in e^{\beta\rho}L^2 \subset L^2$ with $\beta > 0$ and each α satisfying $\text{Re}(\alpha) < Q$

$$\|\mathbf{R}(\alpha)f\|_{e^{\beta\rho}L^2} \leq C\|\mathbf{R}(\alpha)f\|_2 \leq \frac{C\|f\|_2}{|\text{Im}(\alpha)| \cdot |\text{Re}(\alpha) - Q|}$$

which implies that a pole $\alpha_0 = Q + ip_0$ with $p_0 \notin \{\pm\sqrt{2\lambda_j} \mid j \geq 0\}$ on $\text{Re}(\alpha) = Q$ must be at most of order 1, while at $p_0 = \pm\sqrt{2\lambda_j}$ it can be at most of order 2 on Σ . Since $\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) + \mathbf{W}$ is analytic, a pole of $\tilde{\mathbf{R}}(\alpha)$ can only come from a pole of $(1 + \mathbf{K}(\alpha))^{-1}$, with polar part being a finite rank operator. We now assume that $p_0 \notin \{\pm\sqrt{2\lambda_j} \mid j \geq 0\}$. Let us denote by \mathbf{Z} the finite rank residue $\mathbf{Z} = \text{Res}_{\alpha_0} \mathbf{R}(\alpha)$. Then $(\mathbf{H} - \frac{Q^2 + p_0^2}{2})\mathbf{Z} = 0$, which means that each element in $\text{Ran}(\mathbf{Z})$ is a $w \in e^{-\beta\rho}\mathcal{D}(\mathbf{H})$ such that $(\mathbf{H} - \frac{Q^2 + p_0^2}{2})w = 0$. There are finite rank operators $\mathbf{Z}_0, \dots, \mathbf{Z}_N$ on $\mathcal{H}_{k,\beta}$ so that for $\psi \in C^\infty((-\infty, -2)_c; [0, 1])$

$$\psi\mathbf{Z} = \sum_{n=0}^N \psi \partial_\alpha^n \tilde{\mathbf{R}}(\alpha_0) \mathbf{Z}_n = \sum_{n=1}^N \psi \partial_\alpha^n \mathbf{R}_k(\alpha_0) \chi \mathbf{Z}_n + \psi \tilde{\mathbf{R}}(\alpha_0) \mathbf{Z}_0 + \psi \mathbf{Z}_{L^2}$$

where \mathbf{Z}_{L^2} is a finite rank operator mapping to $\mathcal{H}_{k,\beta} \subset L^2$. For $f \in \mathcal{H}_{k,\beta}$, the expression of $\partial_\alpha^j \mathbf{R}_k(\alpha_0)f$ is explicit from (6.50), and one directly checks by differentiating (6.50) in α that it is of the form (for $c < -2$)

$$(\partial_\alpha^n \tilde{\mathbf{R}}(\alpha_0)f)(c) = \sum_{\lambda_j \leq \lambda_k} \sum_{m \leq n} \tilde{a}_{j,m}(\alpha, f) c^m e^{-ic\sqrt{p_0^2 - 2\lambda_j}} + \tilde{G}(\alpha_0, f)$$

for some $\tilde{a}_{j,m}(\alpha_0, f) \in \ker(\mathbf{P} - \lambda_j)$ and $\tilde{\chi}\tilde{G}(\alpha_0, f) \in \mathcal{H}_{k,\beta}$ satisfying $\mathbf{H}\tilde{\chi}\tilde{G}(\alpha_0, f) \in \mathcal{H}_{k,\beta}$. This implies that, in $c < -2$, $w \in \text{Ran}(\mathbf{Z})$ is necessarily of the form

$$w = \sum_{j \leq k} \sum_{m \leq N} b_{j,m} c^m e^{-ic\sqrt{p_0^2 - 2\lambda_j}} + \hat{G}$$

for some $b_{j,m} \in \ker(\mathbf{P} - \lambda_j)$ and $\hat{G} \in \mathcal{H}_{k,\beta}$ with $\mathbf{H}\hat{G} \in \mathcal{H}_{k,\beta}$. Using that $\tilde{\chi}(c)e^{\gamma c}\Pi_k(Vw) \in e^{\beta c}L^2$, we see from the equation $(\mathbf{H}_0 - \frac{Q^2 + p_0^2}{2})\Pi_k(w) = -e^{\gamma c}\Pi_k(Vw)$ that

$$\left(\mathbf{H}_0 - \frac{Q^2 + p_0^2}{2} \right) \left(\sum_{\lambda_j \leq \lambda_k} \sum_{m \leq N} b_{j,m} c^m e^{-ic\sqrt{p_0^2 - 2\lambda_j}} \right) \Big|_{c \leq 0} \in e^{\beta c} L^2$$

and by using the explicit expression of \mathbf{H}_0 , it is clear that necessarily $b_{j,m} = 0$ for all $m \neq 0$. Then we may apply Lemma 6.18 with $u_1 = u_2 = w$ to deduce that $b_{j,0} = 0$, and therefore $w \in \mathcal{D}(\mathbf{H})$, which implies $w = 0$ by Lemma 6.2.

It remains to show that $\alpha = Q \pm i\sqrt{2\lambda_j}$ is a pole of order at most 1. To simplify, we write the argument for $\alpha = Q$, the proof is the same for all j . The method is basically the same as in the proof of [Me93, Proposition 6.28]: the resolvent has Laurent expansion $\mathbf{R}(\alpha) = (\alpha - Q)^{-2}\mathbf{Q} + (\alpha - Q)^{-1}\mathbf{R}'(\alpha)$ for some holomorphic operator $\mathbf{R}'(\alpha)$ near $\alpha = Q$ and \mathbf{Q} has finite rank, then we also have $\|\mathbf{R}(\alpha)\phi\|_{L^2} \leq |Q - \alpha|^{-2}$ for $\alpha < Q$ and all $\phi \in \mathcal{C}$, thus we can deduce that

$$\mathbf{Q}\phi = \lim_{\alpha \rightarrow Q^-} (\alpha - Q)^2 \mathbf{R}(\alpha)\phi.$$

The limit holds in $e^{-\delta\rho}L^2$ for all $\delta > 0$ small, but the right hand side has actually a bounded L^2 -norm, so $\mathbf{Q}\phi \in L^2$ and thus $\text{Ran}(\mathbf{Q}) \subset L^2$. Since we also have $\mathbf{H}\mathbf{Q} = 0$ from Laurent expanding $(\mathbf{H} - 2\Delta_\alpha)\mathbf{R}(\alpha) = \text{Id}$ at $\alpha = Q$, we conclude that $\mathbf{Q} = 0$ by using Lemma 6.2. \square

The resolvent in the physical sheet on weighted spaces. We shall conclude this section on the resolvent of \mathbf{H} by analyzing its boundedness on weighted spaces $e^{-\beta\rho}L^2$ in the half-plane $\{\text{Re}(\alpha) < Q\}$. We recall that Lemma 6.5 was precisely proving such boundedness but the region of validity in α of this Lemma was not covering the whole physical-sheet, and in particular not the region close to the line $\text{Re}(\alpha) = Q$. Just as in Lemma 6.7, the main application of such boundedness on weighted spaces is to define the Poisson operator $\mathcal{P}_\ell(\alpha)$, and we aim to define it in a large connected region of $\{\text{Re}(\alpha) \leq Q\}$ relating the probabilistic region and the line $\alpha \in Q + i\mathbb{R}$ corresponding to the L^2 -spectrum of \mathbf{H} .

Proposition 6.17. *Let $\beta \in \mathbb{R}$ and $\text{Re}(\alpha) < Q$, then the resolvent $\mathbf{R}(\alpha)$ of \mathbf{H} extends as an analytic family of bounded operators*

$$\mathbf{R}(\alpha) : e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho}\mathcal{D}(Q)$$

in the region $\text{Re}(\alpha) < Q - |\beta|$, and it satisfies for each $\psi \in C^\infty(\mathbb{R}) \cap L^\infty(\mathbb{R})$ such that $\psi' \in L^\infty$ and $\text{supp}(\psi) \subset (-\infty, A)$ for some $A \in \mathbb{R}$,

$$\mathbf{R}(\alpha)(1 - \Pi_k)\psi : e^{-\beta\rho}\mathcal{D}'(Q) \rightarrow e^{-\beta\rho}\mathcal{D}(Q).$$

Proof. We proceed as in the proof of Proposition 6.10: we let for $\text{Re}(\alpha) < Q$

$$\tilde{\mathbf{R}}(\alpha) := \tilde{\chi}\mathbf{R}_k^+(\alpha)\chi + (1 - \hat{\chi})\mathbf{R}_+(1 - \chi) + \tilde{\chi}\mathbf{R}_k(\alpha)\chi - \hat{\chi}\mathbf{R}_k^+(\alpha)\mathbf{K}_{k,2}(\alpha)$$

and we get

$$\left(\mathbf{H} - \frac{Q^2 + p^2}{2} \right) \tilde{\mathbf{R}}(\alpha) = \text{Id} + \tilde{\mathbf{K}}_k(\lambda) + \tilde{\mathbf{L}}_k(\alpha)$$

where we used the operators of the proof of Proposition 6.10 (see (6.51) and (6.53)). Now we take $\text{Re}(\alpha) < Q$ and $|Q - \alpha| < A_0$ for some fixed constant $A_0 > 0$ that can be chosen arbitrarily large, and we let $k > 0$ large enough so that

$$(6.57) \quad A_0^2 + 1 < \min\left(\frac{\lambda_k^{1/2}}{16(1 + C_2)}, \lambda_k^{1/4}\right), \quad \lambda_k > (16^2 C_1^2 (1 + C_2^2) + 1)(|\beta| + 1)^2,$$

where the constant C_1, C_2 above are the constants respectively given in Lemma 6.11 and Lemma 6.13. The conditions in (6.57) ensures both the condition $\text{Re}((\alpha - Q)^2) > \beta^2 - 2\lambda_k + 1$ of Lemma 6.11 is satisfied and

the operator $\tilde{\chi}\mathbf{R}_k^+(\alpha)\chi : e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}(L^2(\mathbb{R}^-; E_k^+) \cap \mathcal{D}(\mathcal{Q}))$ is a bounded holomorphic family, and the norm estimate appearing in 2) of Lemma 6.11 gives

$$(6.58) \quad \|\mathbf{L}_k^+(\alpha)\|_{\mathcal{L}(e^{-\beta\rho}L^2)} \leq \frac{C_1(1+|\beta|)}{\sqrt{\operatorname{Re}((\alpha-Q)^2) + 2\lambda_k - \beta^2}} \leq \frac{1}{16(1+C_2)}.$$

The condition $|\beta| < Q - \operatorname{Re}(\alpha)$ (equivalent to $\operatorname{Im}(p) > |\beta|$) makes sure that we can apply 3) of Lemma 6.16: in particular the operator $\mathbf{R}_k(\alpha) : e^{-\beta\rho}L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho}(L^2(\mathbb{R}^-; E_k) \cap \mathcal{D}(\mathcal{Q}))$ is a bounded holomorphic family. Also, Lemma 6.13 ensures that $(1-\hat{\chi})\mathbf{R}_+(1-\chi) : e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ is a bounded holomorphic family (note that both cutoff functions $(1-\hat{\chi})$ and $(1-\chi)$ kill the $c \rightarrow -\infty$ behaviour and this is why Lemma 6.13 extends to weighted spaces $e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$). Also the first condition in (6.57) ensures the norm estimate (as given by (6.35))

$$(6.59) \quad \|\mathbf{L}_{+,k}(\alpha)\|_{\mathcal{L}(L^2)} \leq 2C_2, \quad \|\mathbf{L}_{+,k}(\alpha)^2\|_{\mathcal{L}(L^2)} \leq \frac{1}{8}.$$

As a consequence

$$\tilde{\mathbf{R}}(\alpha) : e^{-\beta\rho}L^2 \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$$

is bounded and holomorphic in $U := \{\alpha \in \mathbb{C} \mid |Q - \alpha| < A_0, \operatorname{Re}(\alpha) < Q - |\beta|\}$. Furthermore (6.58) and (6.59) provide the estimate

$$\|(\mathbf{L}_{+,k}(\alpha) + \mathbf{L}_k^+(\alpha))^2\|_{\mathcal{L}(e^{-\beta\rho}L^2)} < 1/2.$$

Moreover, 2) of Lemma 6.11, Lemma 6.13 and 3) of Lemma 6.16 also give that $\tilde{\mathbf{K}}_k(\alpha)$ is compact on the Hilbert space $e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. Exactly the same argument as in the proof of Proposition 6.10 gives that

$$\left(\mathbf{H} - \frac{Q^2 + p^2}{2}\right)\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) = 1 + \tilde{\mathbf{K}}_k(\alpha)(1 + \mathbf{T}_k(\alpha))$$

for some \mathbf{T}_k bounded holomorphic on $e^{-\beta\rho}L^2$ in U . Since by Lemma 6.5 we know that $(\mathbf{H} - 2\Delta_\alpha)$ is invertible on $e^{-\beta\rho}L^2$ for some $\alpha_0 \in U$, one can always add a finite rank operator $\mathbf{W} : e^{-\beta\rho}L^2 \rightarrow e^{-\beta\rho}\mathcal{D}(\mathbf{H})$, so that

$$(\mathbf{H} - 2\Delta_\alpha)(\mathbf{R}(\alpha)(1 + \mathbf{T}_k(\alpha)) + \mathbf{W}) = 1 + \mathbf{K}(\alpha)$$

for some compact remainder $\mathbf{K}(\alpha)$ on $e^{-\beta\rho}L^2$, analytic in $\alpha \in U$ in the desired region and $1 + \mathbf{K}(\alpha)$ being invertible for $\alpha = \alpha_0 \in U$. This implies by analytic Fredholm theorem that

$$\mathbf{R}(\alpha) = (\tilde{\mathbf{R}}(\alpha)(1 + \mathbf{T}_k(\alpha)) + \mathbf{W})(1 + \mathbf{K}(\alpha))^{-1} : e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$$

is meromorphic for $\alpha \in U$. Now, using the density of the embeddings $e^{|\beta|\rho}L^2 \subset L^2 \subset e^{-|\beta|\rho}L^2$ and using that $\mathbf{R}(\alpha)$ is holomorphic in U as a bounded operator on L^2 , it is direct to check that $\mathbf{R}(\alpha) : e^{-\beta\rho}L^2 \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ is analytic in U . Since A_0 (and thus U) can be chosen arbitrarily large as long as the constraint $\operatorname{Re}(\alpha) < Q - |\beta|$ is satisfied, we obtain our desired result. To prove that $\mathbf{R}(\alpha)(1 - \Pi_k)\psi$ maps $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q})$ to $e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$, we proceed as in the proof of Proposition 6.10 and write

$$\mathbf{R}(\alpha) = \tilde{\mathbf{R}}(\alpha) - \mathbf{R}(\alpha)(\tilde{\mathbf{K}}_k(\alpha) + \tilde{\mathbf{L}}_k(\alpha)).$$

We have seen that $\tilde{\mathbf{R}}(\alpha)(1 - \Pi_k)\psi : e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$. The same arguments (just as in the proof of Proposition 6.10) also prove that that operators $\tilde{\mathbf{K}}_k(\alpha), \tilde{\mathbf{L}}_k(\alpha)$ are bounded as operators $e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}L^2$, thus we obtain that $\mathbf{R}(\alpha)(1 - \Pi_k)\psi : e^{-\beta\rho}\mathcal{D}'(\mathcal{Q}) \rightarrow e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ is bounded. \square

6.3. The Poisson operator. We have seen in Lemma 6.7 that it is possible to construct a family of Poisson operators $\mathcal{P}_\ell(\alpha)$ in what we called the *probabilistic region*, which contains a half line $(-\infty, Q - c_\ell)$ for some $c_\ell \geq 0$ depending on ℓ . The construction was using the resolvent acting on weighted L^2 -spaces. In this section, we will use Proposition 6.10 and Proposition 6.17 to prove that the Poisson operators extend holomorphically in α in a connected region of $\operatorname{Re}(\alpha) \leq Q$ containing the probabilistic region and the line $Q + i\mathbb{R}$.

Regime close to the continuous spectrum of H . We first start with a technical lemma that allows to define the Poisson operator on the continuous spectrum $Q + i\mathbb{R}$:

Lemma 6.18. *Let $p \in \mathbb{R}$ and for $m = 1, 2$, let $u_m \in e^{-\delta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ with $\delta > 0$ such that:*
 1) *for each $\theta \in C^\infty(\mathbb{R}; [0, 1])$ supported in $(a, +\infty)$ for some $a \in \mathbb{R}$ then $\theta u_m \in \mathcal{D}(\mathcal{Q})$*
 2) *u_m satisfies*

$$(\mathbf{H} - \frac{Q^2 + p^2}{2})u_m = r_m \in e^{\delta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}).$$

Set $k = \max\{j \geq 0 \mid 2\lambda_j \leq p^2\}$. Then u_m has asymptotic behaviour

$$(6.60) \quad u_m = \sum_{j, 2\lambda_j \leq p^2} \left(a_m^j e^{-ic\sqrt{p^2 - 2\lambda_j}} + b_m^j e^{ic\sqrt{p^2 - 2\lambda_j}} \right) + G_m$$

with $a_m^j, b_m^j \in \ker(\mathbf{P} - \lambda_j)$, and both $G_m, \partial_c G_m \in e^{\delta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) + L^2(\Omega_{\mathbb{T}}; E_k^\perp)$. Then we have

$$\langle u_1 \mid r_2 \rangle - \langle r_1 \mid u_2 \rangle = i \sum_{j, 2\lambda_j \leq p^2} \sqrt{p^2 - 2\lambda_j} \left(\langle a_1^j \mid a_2^j \rangle_{L^2(\Omega_{\mathbb{T}})} - \langle b_1^j \mid b_2^j \rangle_{L^2(\Omega_{\mathbb{T}})} \right).$$

Proof. Let $\theta \in C^\infty(\mathbb{R})$ be non-negative satisfying $\theta_T = 1$ on $[-T, \infty)$ and $\text{supp}(\theta_T) \subset [-T - \varepsilon, \infty)$ where $T > 0$ is a large parameter and $\varepsilon > 0$ small, and let $\tilde{\theta}_T(\cdot) = \theta_T(\cdot + 1)$. In particular we have $\tilde{\theta}_T \theta_T = \theta_T$. First, $\tilde{\theta}_T u_m \in H^1(\mathbb{R}; L^2(\Omega_{\mathbb{T}}))$ satisfies

$$(\mathbf{H} - \frac{Q^2 + p^2}{2})(\tilde{\theta}_T u_m) = \tilde{\theta}_T r_m - \frac{1}{2} [\partial_c^2, \tilde{\theta}_T] u_m \in L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$$

thus $\tilde{\theta}_T u_m \in \mathcal{D}(\mathbf{H})$ (we used that $[\partial_c^2, \tilde{\theta}_T]$ is a first order differential operator with compactly supported coefficients). This implies, using $[\mathbf{H}, \tilde{\theta}_T] \theta_T = 0 = \theta_T [\mathbf{H}, \tilde{\theta}_T]$, that

$$(6.61) \quad \begin{aligned} \langle u_1, r_2 \rangle - \langle r_1, u_2 \rangle &= \lim_{T \rightarrow \infty} \langle \theta_T u_1, \mathbf{H}(\tilde{\theta}_T u_2) \rangle - \langle \mathbf{H}(\tilde{\theta}_T u_1), \theta_T u_2 \rangle \\ &= - \lim_{T \rightarrow \infty} \frac{1}{2} \langle [\partial_c^2, \theta_T] u_1, u_2 \rangle. \end{aligned}$$

We write $u_m = u_m^0 + G_m$ by using (6.60). Then we claim that, as $T \rightarrow \infty$,

$$|\langle [\partial_c^2, \theta_T] u_1^0, G_2 \rangle| + |\langle [\partial_c^2, \theta_T] G_1, u_2 \rangle| \rightarrow 0.$$

Indeed, we have $(|u_1^0| + |\partial_c u_1^0|)G_2 \in L^1(\mathbb{R} \times \Omega_{\mathbb{T}})$ and $(|G_1| + |\partial_c G_1|)u_2 \in L^1(\mathbb{R} \times \Omega_{\mathbb{T}})$, and the support of $[\partial_c^2, \theta_T]$ is contained in $[-T - 1, -T]$. We are left in (6.61) to study the limit of $\langle [\partial_c^2, \theta_T] u_1^0, u_2^0 \rangle$. But now we have $[\partial_c^2, \theta_T] u_1^0 = \theta_T'' u_1^0 + 2\theta_T' \partial_c u_1^0$ and for fixed $T > 0$ it is direct to check, using integration by parts and the fact that $(\mathbf{H}_0 - \frac{Q^2 + p^2}{2})u_m^0 = 0$ that

$$\begin{aligned} \langle [\partial_c^2, \theta_T] u_1^0, u_2^0 \rangle &= \int_{-T-1}^{-T} \partial_c (\theta_T \langle \partial_c u_1^0, u_2^0 \rangle_{L^2(\Omega_{\mathbb{T}})} - \theta_T \langle u_1^0, \partial_c u_2^0 \rangle_{L^2(\Omega_{\mathbb{T}})}) dc \\ &= \langle \partial_c u_1^0(-T), u_2^0(-T) \rangle_{L^2(\Omega_{\mathbb{T}})} - \langle u_1^0(-T), \partial_c u_2^0(-T) \rangle_{L^2(\Omega_{\mathbb{T}})}. \end{aligned}$$

A direct computation gives that this is equal to

$$2i \sum_{j, \lambda_j \leq p^2} \sqrt{p^2 - 2\lambda_j} \left(\langle b_1^j, b_2^j \rangle_{L^2(\Omega_{\mathbb{T}})} - \langle a_1^j, a_2^j \rangle_{L^2(\Omega_{\mathbb{T}})} \right).$$

This completes the proof. \square

Now we extend the construction of the Poisson operator (6.20) in a neighborhood of the line spectrum $\alpha \in Q + i\mathbb{R}$.

Proposition 6.19. *Let $0 < \beta < \gamma/2$ and $\ell \in \mathbb{N}$. Then there is an analytic family of operators $\mathcal{P}_\ell(\alpha)$*

$$\mathcal{P}_\ell(\alpha) : E_\ell \rightarrow e^{-\beta\rho} \mathcal{D}(\mathcal{Q})$$

in the region

$$(6.62) \quad \left\{ \alpha \in \mathbb{C} \mid \text{Re}(\alpha) < Q, \text{Im} \sqrt{p^2 - 2\lambda_\ell} < \beta \right\} \cup \left\{ Q + ip \in Q + i\mathbb{R} \mid |p| \in \bigcup_{j \geq \ell} (\sqrt{2\lambda_j}, \sqrt{2\lambda_{j+1}}) \right\},$$

continuous at each $Q \pm i\sqrt{2\lambda_j}$ for $j \geq \ell$, satisfying $(\mathbf{H} - \frac{Q^2+p^2}{2})\mathcal{P}_\ell(\alpha)F = 0$ and

$$(6.63) \quad \mathcal{P}_\ell(\alpha)F = \sum_{j \leq \ell} \left(F_j^- e^{ic\sqrt{p^2-2\lambda_j}} + F_j^+(\alpha) e^{-ic\sqrt{p^2-2\lambda_j}} \right) + G_\ell(\alpha, F)$$

with $F_j^- = \Pi_{\ker(\mathbf{P}-\lambda_j)}F$, $F_j^+(\alpha) \in \ker(\mathbf{P}-\lambda_j)$, and $G_\ell(\alpha, F), \partial_c G_\ell(\alpha, F) \in e^{\frac{\beta}{2}\rho(c)}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) + L^2(\mathbb{R}; E_\ell^\perp)$. In particular, $\mathcal{P}_\ell(\alpha) \in e^{-(\text{Im}(\sqrt{p^2-\lambda_\ell})+\varepsilon)\rho}\mathcal{D}(\mathcal{Q})$ for all $\varepsilon > 0$. Moreover, for each $\theta \in C_c^\infty(\mathbb{R})$, one has $\theta\mathcal{P}_\ell(\alpha)F \in \mathcal{D}(\mathbf{H})$. Such a solution $u \in e^{-\beta\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ to the equation $(\mathbf{H} - \frac{Q^2+p^2}{2})u = 0$ with the asymptotic expansion (6.63) is unique. The operator $\mathcal{P}_\ell(\alpha)$ admits a meromorphic extension to the region

$$(6.64) \quad \left\{ \alpha = Q + ip \in \Sigma \mid \forall j \in \mathbb{N} \cup \{0\}, \text{Im}\sqrt{p^2-2\lambda_j} \in (\beta/2 - \gamma, \beta/2) \right\}$$

and $\mathcal{P}_\ell(\alpha)F$ satisfies (6.63) in that region. Finally, $F_j^+(\alpha)$ depends meromorphically on α in the region above.

Proof. We start by setting $u_-(\alpha) := \sum_{j=0}^\ell F_j^- e^{ic\sqrt{p^2-2\lambda_j}}$, and let $\chi \in C^\infty(\mathbb{R})$ equal to 1 in $(-\infty, -1)$ and with $\text{supp}(\chi) \subset \mathbb{R}^-$. We get

$$(\mathbf{H} - \frac{Q^2+p^2}{2})(\chi u_-(\alpha)) = -\frac{1}{2}\chi''(c)u_-(\alpha) - \chi'(c)\partial_c u_-(\alpha) + e^{\gamma c}V\chi u_-(\alpha).$$

The first two terms are in $e^{N\rho}L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all N , the term $e^{\gamma c}V\chi u_-(\alpha)$ can be decomposed (for each $k \geq \ell$) as

$$\Pi_k(e^{\gamma c}V\chi u_-(\alpha)) + (1 - \Pi_k)(e^{\gamma c}V\chi u_-(\alpha)).$$

Using that $u_-(\alpha) = \Pi_k u_-(\alpha)$ if $k \geq \ell$ together with Lemma 6.1, we see that the first term satisfies $\Pi_k(e^{\gamma c}V\chi u_-(\alpha)) \in e^{\beta\rho}L^2(\mathbb{R}^-; E_k)$ and the second term $(1 - \Pi_k)(e^{\gamma c}V\chi u_-(\alpha)) \in e^{\beta\rho}\mathcal{D}'(\mathcal{Q})$, provided that $\beta < \gamma/2$ and that $\text{Im}\sqrt{p^2-\lambda_j} \leq \gamma/2$ for $j \leq \ell$. We can thus define, using Proposition 6.10 (with $k \gg \ell$ large enough), if $\text{Im}(\sqrt{p^2-2\lambda_j}) \in (-\min(\beta, \frac{\gamma}{2} - \beta), \gamma/2)$ for all $j \leq k$ and $|\pi(p)|^2 \leq \lambda_k^{1/4}$

$$u_+(\alpha) := \mathbf{R}(\alpha)(\mathbf{H} - \frac{Q^2+p^2}{2})(\chi u_-(\alpha)) \in e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$$

so that $u(\alpha) := \chi u_-(\alpha) - u_+(\alpha)$ solves $(\mathbf{H} - \frac{Q^2+p^2}{2})u(\alpha) = 0$ in $e^{-\frac{\gamma}{2}\rho}\mathcal{D}'(\mathcal{Q})$. We use Proposition 6.10 with $k \gg \ell$ large enough, and we see that $u_+(\alpha)$ is of the form, in $c \leq 0$,

$$u_+(\alpha) = \sum_{j \leq k} \tilde{a}_j(\alpha, F) e^{-ic\sqrt{p^2-2\lambda_j}} + G(\alpha, F) = \sum_{j \leq \ell} \tilde{a}_j(\alpha, F) e^{-ic\sqrt{p^2-2\lambda_j}} + G_\ell(\alpha, F)$$

with $\tilde{a}_j(\alpha, F) \in \ker(\mathbf{P}-\lambda_j)$ and $G(\alpha, F), \partial_c G(\alpha, F) \in e^{\beta\rho}L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) + L^2(\mathbb{R}^-; E_k^\perp)$ and $G_\ell(\alpha, F) \in e^{\beta\rho}L^2(\mathbb{R}^- \times \Omega_{\mathbb{T}}) + L^2(\mathbb{R}^-; E_\ell^\perp)$ if $\text{Re}(\alpha) < Q$. Here we have used the fact that $\text{Im}\sqrt{p^2-2\lambda_j} > 0$ (since either $\text{Re}(\alpha) < Q$ or $p^2 < 2\lambda_j$ for all $j > \ell$ if $\alpha = Q + ip$ with $p \in \mathbb{R} \setminus [-\sqrt{2\lambda_\ell}, \sqrt{2\lambda_\ell}]$) to place all terms corresponding to all $\ell < j \leq k$, which belong to $L^2(\mathbb{R}; E_\ell^\perp)$, in the remainder term $G_\ell(\alpha, F)$. This shows that

$$(6.65) \quad \mathcal{P}_\ell(\alpha)F := u(\alpha) = \chi(c) \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} - \mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha) \left(\chi(c) \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \right)$$

satisfies all the required properties. The analyticity in α except possibly at the points $Q \pm i\sqrt{2\lambda_j}$ for $j \in \mathbb{N}_0$ follows from Proposition 6.10, in particular 3) of that Proposition. At the points $Q \pm i\sqrt{2\lambda_j}$, the analyticity on the surface Σ is a consequence of the Lemma 6.20, in particular (6.67) and the fact that $Q \pm i\sqrt{2\lambda_j}$ is at most a pole of order 1 of $\mathbf{R}(\alpha)$ and $a_j(\alpha, \varphi)$. We notice that the expression of $\mathcal{P}_\ell(\alpha)$ is the same as in (6.20), thus when the regions of α considered in Lemma 6.7 and here have an intersection, then this corresponds to the same operator, by analytic continuation.

The uniqueness of the solution with such an asymptotic is direct if $\text{Re}(\alpha) < Q$: the difference of two such solutions would be in $\mathcal{D}(\mathcal{Q})$ and the operator \mathbf{H} has no L^2 eigenvalues (Lemma 6.2), hence the difference is identically 0. For the case $\alpha = Q + ip$ with $p \in \mathbb{R}$, denote by $\hat{u}(\alpha)$ the difference of two such solutions. Then $\hat{u}(\alpha)$ can be written under the form

$$\hat{u}(\alpha) = \sum_{j \leq \ell} \hat{F}_j^+(\alpha) e^{-ic\sqrt{p^2-2\lambda_j}} + \hat{G}_\ell(\alpha, F)$$

where $\hat{F}_j^+(\alpha) \in \ker(\mathbf{P} - \lambda_j)$ and $\hat{G}_\ell(\alpha, F) \in e^{\beta\rho(c)}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) + L^2(\mathbb{R}; E_\ell^1)$. We can split the sum above as $\sum_{j, 2\lambda_j \leq p^2} \dots + \sum_{j, p^2 < j \leq 2\lambda_\ell} \dots$. The sum $\sum_{j, p^2 < j \leq 2\lambda_\ell} \dots$ belongs to some $e^{\delta\rho}L^2$ as well as its ∂_c derivative. We can use Lemma 6.18 to see that $\sum_{j, 2\lambda_j \leq p^2} \|\hat{F}_j^+(\alpha)\|_{L^2(\Omega_{\mathbb{T}})}^2 = 0$, hence again $\hat{u}(\alpha) \in L^2$ and we can conclude as previously.

The meromorphic extension of $\mathcal{P}_\ell(\alpha)$ is a direct consequence of the meromorphic extension of $\mathbf{R}(\alpha)$ in Proposition 6.10. \square

We notice that for $\alpha = Q + ip$ with $p \in \mathbb{R}$, the function $\overline{\mathcal{P}_\ell(\bar{\alpha})F}$ is another solution of $(\mathbf{H} - \frac{Q^2+p^2}{2})u = 0$ satisfying

$$\overline{\mathcal{P}_\ell(\bar{\alpha})F} = \sum_{j \leq \ell} \left(\overline{F_j^-} e^{-ic\sqrt{p^2-\lambda_j}} + \overline{F_j^+(\bar{\alpha})} e^{ic\sqrt{p^2-\lambda_j}} \right) + \overline{G_\ell(\bar{\alpha}, F)}.$$

This implies that for each $F = \sum_{j \leq \ell} F_j^- \in E_\ell$, there is a unique solution $u = \widehat{\mathcal{P}}_\ell(\alpha)F$ to $(\mathbf{H} - \frac{Q^2+p^2}{2})u = 0$ of the form

$$(6.66) \quad \widehat{\mathcal{P}}_\ell(\alpha)F = \sum_{j \leq \ell} \left(F_j^- e^{-ic\sqrt{p^2-\lambda_j}} + \hat{F}_j^+(\alpha) e^{ic\sqrt{p^2-\lambda_j}} \right) + \widehat{G}_\ell(\alpha, F)$$

with $\widehat{G}_\ell(\alpha, F) \in e^{\frac{\beta}{2}\rho(c)}L^2(\mathbb{R} \times \Omega_{\mathbb{T}}) + L^2(\mathbb{R}; E_\ell^1)$ and $\hat{F}_j^+ \in \mathbb{C}\rho_j$, and $\widehat{\mathcal{P}}_\ell(\alpha)$ extends meromorphically on an open set of Σ just like $\mathcal{P}_\ell(\alpha)$.

Lemma 6.20. *Let $\ell \in \mathbb{N}$, $0 < \beta < \gamma$ and α in (6.62), the Poisson operator $\mathcal{P}_\ell(\alpha)$ can be obtained from the resolvent as follows: for $F = \sum_{j \leq \ell} F_j^- \in E_\ell$ and $\varphi \in e^{\beta\rho}L^2$*

$$(6.67) \quad \langle \mathcal{P}_\ell(\alpha)F, \varphi \rangle_2 = i \sum_{j \leq \ell} \sqrt{p^2 - 2\lambda_j} \left\langle F_j^-, a_j(\bar{\alpha}, \varphi) \right\rangle_{L^2(\Omega_{\mathbb{T}})}$$

where $a_j(\alpha, \varphi)$ are the functionals obtained from (6.26), holomorphic in α and linear in φ .

Proof. Let $\alpha = Q + ip$ with $p \in \mathbb{R} \setminus ([-\sqrt{2\lambda_\ell}, \sqrt{2\lambda_\ell}] \cup_{j \geq \ell} \{\pm\sqrt{2\lambda_j}\})$ and let us take $F = \sum_{j \leq \ell} F_j^-$ with $F_j^- \in \text{Ker}(\mathbf{P} - \lambda_j)$ for $j \leq \ell$. Then from the construction of $\mathcal{P}_\ell(\alpha)F$ (with a function $\chi = \chi(c)$) in the proof of Proposition 6.19

$$\begin{aligned} \langle \mathcal{P}_\ell(\alpha)F, \varphi \rangle &= \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \varphi \right\rangle - \left\langle \mathbf{R}(\alpha)(\mathbf{H} - 2\Delta_\alpha) \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \varphi \right\rangle \\ &= \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \varphi \right\rangle - \left\langle (\mathbf{H} - 2\Delta_\alpha) \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \mathbf{R}(\bar{\alpha})\varphi \right\rangle. \end{aligned}$$

Here we used $\mathbf{R}(\alpha)^* = \mathbf{R}(\bar{\alpha}) = \mathbf{R}(2Q - \alpha)$. Let θ_T be as in the proof of Lemma 6.18. We have

$$\begin{aligned} &\lim_{T \rightarrow \infty} \left\langle \theta_T(c)(\mathbf{H} - 2\Delta_\alpha) \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \mathbf{R}(\bar{\alpha})\varphi \right\rangle \\ &= \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \varphi \right\rangle - \frac{1}{2} \lim_{T \rightarrow \infty} \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \theta_T'' \mathbf{R}(\bar{\alpha})\varphi \right\rangle \\ &\quad - \lim_{T \rightarrow \infty} \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \theta_T' \partial_c \mathbf{R}(\bar{\alpha})\varphi \right\rangle. \end{aligned}$$

Using now the asymptotic form (6.26), the last two limits above can be rewritten as

$$\begin{aligned} &\lim_{T \rightarrow \infty} \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \theta_T'' \mathbf{R}(\bar{\alpha})\varphi \right\rangle = \lim_{T \rightarrow \infty} \sum_{j \leq \ell} \left\langle F_j^- e^{ic\sqrt{p^2-2\lambda_j}}, \theta_T'' a_j(\bar{\alpha}, \varphi) e^{ic\sqrt{p^2-2\lambda_j}} \right\rangle \\ &\lim_{T \rightarrow \infty} \left\langle \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2-2\lambda_j}} \chi, \theta_T' \partial_c \mathbf{R}(\bar{\alpha})\varphi \right\rangle = \lim_{T \rightarrow \infty} \sum_{j \leq \ell} \left\langle F_j^- e^{ic\sqrt{p^2-2\lambda_j}}, \theta_T' a_j(\bar{\alpha}, \varphi) \partial_c (e^{ic\sqrt{p^2-2\lambda_j}}) \right\rangle \end{aligned}$$

and this easily yields

$$\langle \mathcal{P}_\ell(\alpha)F, \varphi \rangle = i \sum_{j \leq \ell} \sqrt{p^2 - \lambda_j} \left\langle F_j^-, a_j(\bar{\alpha}, \varphi) \right\rangle_{L^2(\Omega_{\mathbb{T}})}.$$

where a_j are the functional obtained from (6.26). The result then extends holomorphically to the region (6.62) and meromorphically to (6.64) \square

The Poisson operator far from $\text{Re}(\alpha) = Q$. We have seen in Lemma 6.7 that the Poisson operator can be defined far from the spectrum. The problem is that the region of analyticity of $\mathcal{P}_\ell(\alpha)$ in Lemma 6.7 does not intersect (for ℓ large at least) the region of analyticity of $\mathcal{P}_\ell(\alpha)$ from Proposition 6.19. The proposition below extends the construction of the Poisson operator to a region overlapping both regions in Lemma 6.7 and Proposition 6.19 (see figure 4).

Proposition 6.21. *For ℓ fixed, the Poisson operator $\mathcal{P}_\ell(\alpha)$ of Lemma 6.7 extends analytically to the region*

$$(6.68) \quad \left\{ \alpha = Q + ip \in \mathbb{C} \mid \text{Re}(\alpha) < Q, \text{Im}(p) > \text{Im}(\sqrt{p^2 - 2\lambda_\ell}) - \gamma/2 \right\}$$

as an function in $e^{-(\text{Im}(\sqrt{p^2 - 2\lambda_\ell}) + \varepsilon)\rho} \mathcal{D}(\mathcal{Q})$ for all $\varepsilon > 0$.

Proof. As before, for $F = \sum_{j=0}^{\ell} F_j^- \in E_\ell$ with $F_j^- \in \text{Ker}(\mathbf{P} - \lambda_j)$, we set $u_-(\alpha) := \sum_{j \leq \ell} F_j^- e^{ic\sqrt{p^2 - 2\lambda_j}}$, and let $\chi \in C^\infty(\mathbb{R})$ equal to 1 in $(-\infty, -1)$ and with $\text{supp}(\chi) \subset \mathbb{R}^-$. We get

$$(\mathbf{H} - \frac{Q^2 + p^2}{2})(\chi u_-(\alpha)) = -\frac{1}{2}\chi''(c)u_-(\alpha) - \chi'(c)\partial_c u_-(\alpha) + e^{\gamma c} V \chi u_-(\alpha).$$

The first two terms in the right hand side are in $e^{N\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $N > 0$ (indeed, compactly supported in c), while the last term can be decomposed as

$$\Pi_k e^{\gamma c} V \chi u_-(\alpha) + (1 - \Pi_k) \chi e^{\gamma c} V u_-(\alpha) \in e^{(-\text{Im}\sqrt{p^2 - 2\lambda_\ell} + \gamma/2)\rho} L^2(\mathbb{R}; E_k) + e^{(-\text{Im}\sqrt{p^2 - 2\lambda_\ell} + \gamma/2)\rho} \mathcal{D}'(\mathcal{Q})$$

by using Lemma 6.1. Using Proposition 6.17, we can thus define, with the same formula as in Lemma 6.20 and Proposition 6.19, the Poisson operator

$$u_+(\alpha) := \mathbf{R}(\alpha) \left(\mathbf{H} - \frac{Q^2 + p^2}{2} \right) (\chi u_-(\alpha)) \in e^{(-\text{Im}\sqrt{p^2 - 2\lambda_\ell} + \gamma/2)\rho} \mathcal{D}(\mathcal{Q}),$$

$$\mathcal{P}_\ell(\alpha) F := u_-(\alpha) \chi - u_+(\alpha)$$

in the region

$$\text{Im}(p) = \text{Re}(Q - \alpha) > \max_{j \leq \ell} \text{Im}\sqrt{p^2 - 2\lambda_j} - \gamma/2 = \text{Im}\sqrt{p^2 - 2\lambda_\ell} - \gamma/2.$$

□

Remark 6.22. *Notice that this region of holomorphy is non-empty and connected, as for ℓ and $|p| = R \gg \lambda_\ell$*

$$\text{Im}(p) - \text{Im}\sqrt{p^2 - 2\lambda_\ell} + \gamma/2 = \gamma/2 + \mathcal{O}\left(\frac{\lambda_\ell}{R^2}\right) > 0.$$

6.4. The Scattering operator.

Definition 6.23. *Let $\ell \in \mathbb{N}$ and $\alpha = Q + ip$ with $p \in \mathbb{R} \setminus (-\sqrt{2\lambda_\ell}, \sqrt{2\lambda_\ell})$. The scattering operator $\mathbf{S}_\ell(\alpha) : E_\ell \rightarrow E_\ell$ for the ℓ -th layer (also called ℓ -scattering operator) is the operator defined as follows: let $F = \sum_{j \leq \ell} F_j \in E_\ell$ (with $F_j \in \text{Ker}(\mathbf{P} - \lambda_j)$) and let $F_j^- := (p^2 - 2\lambda_j)^{-1/4} F_j$, then we set*

$$\mathbf{S}_\ell(\alpha) F := \begin{cases} \sum_{j \leq \ell} F_j^+(\alpha) (p^2 - 2\lambda_j)^{1/4}, & \text{if } p > \sqrt{2\lambda_\ell}, \\ \sum_{j \leq \ell} \hat{F}_j^+(\alpha) (p^2 - 2\lambda_j)^{1/4}, & \text{if } p < -\sqrt{2\lambda_\ell}. \end{cases}$$

where $F_j^+(\alpha), \hat{F}_j^+(\alpha)$ are the functions in (6.63) and (6.66). We will call more generally

$$\mathbf{S}(\alpha) := \begin{cases} \bigcup_{\ell, 2\lambda_\ell < p^2} \text{ker}(\mathbf{P} - \lambda_j) & \rightarrow \bigcup_{\ell, 2\lambda_\ell < p^2} \text{ker}(\mathbf{P} - \lambda_j) \\ F \in E_\ell & \mapsto \mathbf{S}_\ell(\alpha) F \end{cases}$$

the scattering operator, where we use $\mathbf{S}_\ell(\alpha)|_{E_{\ell'}} = \mathbf{S}_{\ell'}(\alpha)$ for $\ell' < \ell$.

Let us define the map $\omega_\ell : \Sigma \rightarrow \Sigma$ by the following property: if $r_j(\alpha) = \sqrt{p^2 - 2\lambda_j}$ are the analytic functions on Σ used to define this ramified covering (with $\alpha = Q + ip$ in the physical sheet), then $\omega_\ell(\alpha)$ is the point in Σ so that

$$r_j(\omega_\ell(\alpha)) = \begin{cases} -r_j(\alpha) & \text{if } j \leq \ell, \\ r_j(\alpha) & \text{if } j > \ell \end{cases}.$$

As a consequence of Proposition 6.19 and Lemma 6.18, we obtain the




FIGURE 4. The green colored region correspond to the region (6.68) of validity of Lemma 6.21 for the analyticity of the Poisson operator $\mathcal{P}_\ell(\alpha)$ with $\ell > 0$ (for the plot, we take $\lambda_\ell = 4$, $\gamma = 1/2$). It overlaps the probabilistic (6.21) and near spectrum (6.62) regions.

Corollary 6.24. *For each $\ell \in \mathbb{N}$, the ℓ -th scattering operator $\mathbf{S}_\ell(\alpha)$ is unitary on E_ℓ if $\alpha = Q + ip$ ($p \in \mathbb{R}$) is such that $\lambda_\ell < p^2 < \min\{\lambda_j \mid \lambda_\ell < \lambda_j\}$. It also satisfies the functional equation*

$$(6.69) \quad \mathbf{S}_\ell(\alpha)\mathbf{S}_\ell(\omega_\ell(\alpha)) = \text{Id}.$$

Moreover it extends meromorphically in (6.64) if $\beta < \gamma$. It also satisfies the following functional equation for each $F = \sum_{j=0}^{\ell} F_j \in E_\ell$

$$(6.70) \quad \mathcal{P}_\ell(\alpha) \sum_{j=0}^{\ell} (p^2 - 2\lambda_j)^{-1/4} F_j = \mathcal{P}_\ell(\omega_\ell(\alpha)) \mathbf{S}_\ell(\alpha) \sum_{j=0}^{\ell} F_j.$$

Proof. The unitarity of $\mathbf{S}_\ell(\alpha)$ on the line $\text{Re}(\alpha) = Q$ follows directly from Lemma 6.18 applied with $u_\ell = \mathcal{P}_\ell(\alpha)F$. The functional equation (6.69) reads $\mathbf{S}_\ell(\alpha)\mathbf{S}_\ell(2Q - \alpha) = \text{Id}$ on the line $\text{Re}(\alpha) = Q$ and that comes directly from the uniqueness statement in Proposition 6.19 on the line. The extension of $\mathbf{S}(\alpha)$ with respect to α comes directly from the meromorphy of the $a_j(\alpha, F)$ in Proposition 6.10. The functional identity extends meromorphically under the formula (6.69). The functional equation (6.69) also comes from uniqueness of the Poisson operator. \square

Theorem 6.25. *For each $j \in \mathbb{N}$, let $(h_{jk})_{k=1, \dots, k(j)}$ be an orthonormal basis of $\ker_{L^2(\Omega_{\mathbb{T}})}(\mathbf{P} - \lambda_j)$. The spectral resolution holds for all $\varphi, \varphi' \in e^{\beta\rho} L^2(\Omega_{\mathbb{T}} \times \mathbb{R})$ with $\beta > 0$*

$$(6.71) \quad \langle \varphi | \varphi' \rangle_2 = \frac{1}{2\pi} \sum_{j=0}^{\infty} \sum_{k=1}^{k(j)} \int_0^{\infty} \langle \varphi | \mathcal{P}_j(Q + i\sqrt{p^2 + 2\lambda_j}) h_{jk} \rangle \langle \mathcal{P}_j(Q + i\sqrt{p^2 + 2\lambda_j}) h_{jk} | \varphi' \rangle dp.$$

As a consequence the spectrum of \mathbf{H} is absolutely continuous.

Proof. We recall the Stone formula: for $\varphi, \varphi' \in e^{\beta\rho}L^2$ for $\beta > 0$

$$\begin{aligned} \langle \varphi | \varphi' \rangle_2 &= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_0^\infty \langle [(\mathbf{H} - \frac{Q^2}{2} - t - i\varepsilon)^{-1} - (\mathbf{H} - \frac{Q^2}{2} - t + i\varepsilon)^{-1}] \varphi | \varphi' \rangle dt \\ &= \frac{1}{2\pi i} \lim_{\varepsilon \rightarrow 0^+} \int_0^\infty \langle [(\mathbf{H} - \frac{Q^2+p^2}{2} - i\varepsilon)^{-1} - (\mathbf{H} - \frac{Q^2+p^2}{2} + i\varepsilon)^{-1}] \varphi | \varphi' \rangle p dp \\ &= \frac{1}{2\pi i} \int_0^\infty \langle [(\mathbf{R}(Q+ip) - \mathbf{R}(Q-ip))] \varphi | \varphi' \rangle p dp. \end{aligned}$$

Here $\alpha = Q + ip$ (with $p > 0$) has to be viewed as an element in Σ obtained by limit $Q + ip - \varepsilon$ as $\varepsilon \rightarrow 0^+$ and, if ℓ is the largest integer such that $2\lambda_\ell \leq p^2$, we write $\bar{\alpha}$ for the point $\omega_\ell(\alpha)$ on Σ . For $\alpha = Q + ip$ with $p \in \mathbb{R}^+$, we have for $\varphi \in e^{\beta\rho}L^2$ with $\beta > 0$

$$\begin{aligned} \frac{1}{2i} \langle (\mathbf{R}(\alpha) - \mathbf{R}(\bar{\alpha})) \varphi | \varphi \rangle &= \text{Im} \langle \mathbf{R}(\alpha) \varphi | \varphi \rangle \\ &= \text{Im} \langle \mathbf{R}(\alpha) (\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) \varphi | \varphi \rangle \\ &= \text{Im} \langle (\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) \varphi | \mathbf{R}(\bar{\alpha}) \varphi \rangle \end{aligned}$$

Here we have used that $(\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) = \text{Id}$ on $e^{\beta\rho}L^2$ provided $p \in \mathbb{R}$ and that $\langle \mathbf{R}(\bar{\alpha}) \varphi | \varphi' \rangle = \langle \varphi, \mathbf{R}(\alpha) \varphi' \rangle$ for $\varphi | \varphi' \in e^{\beta\rho}L^2$, this last fact coming from the identity $\mathbf{R}(\bar{\alpha}) = \mathbf{R}(\alpha)^*$ for $\text{Re}(\alpha) < Q$ and passing to the limit $\text{Re}(\alpha) \rightarrow Q$. Let $\theta_T(c)$ as in the proof of Lemma 6.18. We have

$$\begin{aligned} \langle (\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) \varphi, \mathbf{R}(\bar{\alpha}) \varphi \rangle &= \lim_{T \rightarrow +\infty} \langle \theta_T (\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) \varphi, \mathbf{R}(\bar{\alpha}) \varphi \rangle \\ &= \lim_{T \rightarrow +\infty} \langle \theta_T \mathbf{R}(\bar{\alpha}) \varphi | \varphi \rangle + \lim_{T \rightarrow +\infty} \frac{1}{2} \langle [\partial_c^2, \theta_T] \mathbf{R}(\bar{\alpha}) \varphi | \mathbf{R}(\bar{\alpha}) \varphi \rangle. \end{aligned}$$

Using (6.26) and arguing as in the proof of Lemma 6.18, as $T \rightarrow \infty$ we get

$$\begin{aligned} \langle (\mathbf{H} - 2\Delta_\alpha) \mathbf{R}(\bar{\alpha}) \varphi, \mathbf{R}(\bar{\alpha}) \varphi \rangle &= \langle \mathbf{R}(\bar{\alpha}) \varphi, \varphi \rangle + \frac{1}{2} \lim_{T \rightarrow \infty} \sum_{j \leq \ell} \|a_j(\bar{\alpha}, \varphi)\|_{L^2(\Omega_T)}^2 \partial_c (e^{ic\sqrt{p^2-2\lambda_j}})|_{c=-T} e^{iT\sqrt{p^2-2\lambda_j}} \\ &\quad - \frac{1}{2} \lim_{T \rightarrow \infty} \sum_{j \leq \ell} \|a_j(\bar{\alpha}, \varphi)\|_{L^2(\Omega_T)}^2 e^{-iT\sqrt{p^2-2\lambda_j}} \partial_c (e^{-ic\sqrt{p^2-2\lambda_j}})|_{c=-T} \\ &= \langle \mathbf{R}(\bar{\alpha}) \varphi, \varphi \rangle - i \sum_{j \leq \ell} \sqrt{p^2-2\lambda_j} \|a_j(\bar{\alpha}, \varphi)\|_{L^2(\Omega_T)}^2. \end{aligned}$$

We conclude that

$$(6.72) \quad -\text{Im} \langle \mathbf{R}(\bar{\alpha}) \varphi, \varphi \rangle = \frac{1}{2} \sum_{j \leq \ell} \sqrt{p^2-2\lambda_j} \|a_j(\bar{\alpha}, \varphi)\|_{L^2(\Omega_T)}^2.$$

By Lemma 6.20, we have

$$(6.73) \quad \mathcal{P}_\ell(\alpha)^* \varphi = -i \sum_{j \leq \ell} a_j(\bar{\alpha}, \varphi) \sqrt{p^2-2\lambda_j}.$$

By polarisation and by denoting Π_j the orthogonal projectors on $\text{Ker}(\mathbf{P} - \lambda_j)$, we deduce from (6.72) and (6.73) that

$$-\text{Im} \langle \mathbf{R}(\bar{\alpha}) \varphi, \varphi \rangle = \frac{1}{2} \sum_{j \leq \ell} \frac{1}{\sqrt{p^2-2\lambda_j}} \langle \Pi_j \mathcal{P}_\ell(\alpha)^* \varphi, \Pi_j \mathcal{P}_\ell(\alpha)^* \varphi \rangle_{L^2(\Omega_T)}.$$

Rewriting $\mathcal{P}_\ell(\alpha)^* \varphi = \sum_{j=0}^\ell \sum_{k=1}^{k(j)} \langle \varphi, \mathcal{P}_\ell(\alpha) h_{jk} \rangle_2 h_{jk}$, we obtain

$$\langle \varphi, \varphi' \rangle_2 = \frac{1}{2\pi} \int_0^\infty \sum_{\ell=0}^\infty \mathbb{1}_{[\sqrt{2\lambda_\ell}, \sqrt{2\lambda_{\ell+1}})}(p) \sum_{j=0}^\ell \sum_{k=1}^{k(j)} \langle \varphi, \mathcal{P}_\ell(Q+ip) h_{jk} \rangle_2 \overline{\langle \varphi', \mathcal{P}_\ell(Q+ip) h_{jk} \rangle_2} \frac{p}{\sqrt{p^2-2\lambda_j}} dp.$$

This finally can be rewritten, using (6.24), as

$$\begin{aligned} \langle \varphi, \varphi' \rangle_2 &= \frac{1}{2\pi} \sum_{j=0}^{\infty} \sum_{\ell \geq j} \int_0^{\infty} \mathbb{1}_{[\sqrt{2\lambda_\ell}, \sqrt{2\lambda_{\ell+1}})}(p) \sum_{k=1}^{k(j)} \langle \varphi, \mathcal{P}_\ell(Q + ip)h_{jk} \rangle \overline{\langle \varphi', \mathcal{P}_\ell(Q + ip)h_{jk} \rangle} \frac{p}{\sqrt{p^2 - 2\lambda_j}} dp \\ &= \frac{1}{2\pi} \sum_{j=0}^{\infty} \int_{\sqrt{2\lambda_j}}^{\infty} \sum_{k=1}^{k(j)} \langle \varphi, \mathcal{P}_j(Q + ip)h_{jk} \rangle \overline{\langle \varphi', \mathcal{P}_j(Q + ip)h_{jk} \rangle} \frac{p}{\sqrt{p^2 - 2\lambda_j}} dp \\ &= \frac{1}{2\pi} \sum_{j=0}^{\infty} \sum_{k=1}^{k(j)} \int_0^{\infty} \langle \varphi, \mathcal{P}_j(Q + i\sqrt{r^2 + 2\lambda_j})h_{jk} \rangle \overline{\langle \varphi', \mathcal{P}_j(Q + i\sqrt{r^2 + 2\lambda_j})h_{jk} \rangle} dr \end{aligned}$$

where we performed the change of variables in the last line $r = \sqrt{p^2 - 2\lambda_j}$. \square

6.5. Holomorphic parametrization of the eigenstates. We conclude this section by defining the generalized eigenfunctions $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ by using the standard orthonormal basis $(\psi_{\mathbf{k}\mathbf{l}})_{\mathbf{k}, \mathbf{l} \in \mathcal{N}}$ of $L^2(\Omega_{\mathbb{T}})$ made of Hermite polynomials and introduced below (4.7). We set

$$\Psi_{\alpha, \mathbf{k}, \mathbf{l}} := \mathcal{P}_j(Q - \sqrt{(Q - \alpha)^2 - 2\lambda_{\mathbf{k}\mathbf{l}}})\psi_{\mathbf{k}\mathbf{l}}$$

where ℓ is defined by $\lambda_\ell = \lambda_{\mathbf{k}\mathbf{l}}$ and \sqrt{z} is used here with the convention that the cut is on \mathbb{R}^- (ie. $\sqrt{Re^{i\theta}} = \sqrt{R}e^{i\theta/2}$ for $\theta \in (-\pi, \pi)$). These are eigenfunctions of \mathbf{H} with eigenvalues $2\Delta_\alpha + \lambda_{\mathbf{k}\mathbf{l}}$. Using (6.62) and (6.68) for the holomorphy of $\mathcal{P}_\ell(\cdot)$, we obtain the

Proposition 6.26. *Let $\ell \geq 0$ such that $\lambda_{\mathbf{k}\mathbf{l}} = |\mathbf{k}| + |\mathbf{l}| = \lambda_\ell$. For each $\varepsilon > 0$, the function $\Psi_{\alpha, \mathbf{k}, \mathbf{l}} \in e^{-\beta\rho}\mathcal{D}(\mathcal{Q})$ is an eigenfunction of \mathbf{H} with eigenvalue $2\Delta_\alpha + \lambda_\ell$ for all $\beta > Q - \text{Re}(\alpha)$, it is holomorphic on the set*

$$(6.74) \quad W_\ell := \left\{ \alpha \in \mathbb{C} \setminus \mathcal{D}_\ell \mid \text{Re}(\alpha) \leq Q, \text{Re}\sqrt{(Q - \alpha)^2 - 2\lambda_\ell} > \text{Re}(Q - \alpha) - \gamma/2 \right\}$$

where $\mathcal{D}_\ell := \cup_{j \geq \ell} \{Q \pm i\sqrt{2(\lambda_j - \lambda_\ell)}\}$ is a discrete set where $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ is continuous in α with square root singularities. The set W_ℓ is a connected subset of the half-plane $\text{Re}(\alpha) \leq Q$, containing $(Q + i\mathbb{R}) \setminus \mathcal{D}_\ell$ and the real half-line $(-\infty, Q - \frac{2\lambda_\ell}{\gamma} - \frac{\gamma}{4})$.

Finally, one has for $P \in \mathbb{R}$, $\Psi_{Q+iP, \mathbf{k}, \mathbf{l}} = \mathcal{P}_j(Q + i\sqrt{P^2 + 2\lambda_{\mathbf{k}\mathbf{l}}})\psi_{\mathbf{k}\mathbf{l}}$ and one can rewrite (6.71) as

$$(6.75) \quad \langle \varphi | \varphi' \rangle_2 = \frac{1}{2\pi} \sum_{\mathbf{k}, \mathbf{l} \in \mathcal{N}} \int_0^{\infty} \langle \varphi | \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} \rangle \langle \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} | \varphi' \rangle dP.$$

7. PROBABILISTIC REPRESENTATION OF THE POISSON OPERATOR

In Section 6, we constructed the generalized eigenstates $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ by means of the Poisson operator (see Prop 6.26) of the Liouville Hamiltonian \mathbf{H} on the spectrum line $\alpha \in Q + i\mathbb{R}$ and showed that these generalized eigenstates can be analytically continued in the parameter α over the region W_ℓ defined by (6.74) for $|\mathbf{k}| + |\mathbf{l}| = \lambda_\ell$. As in the case of the \mathbf{H}^0 -eigenstates, we will need to perform a change of basis. So, similarly to Prop 4.9 item 4, we set for $\nu, \tilde{\nu} \in \mathcal{T}$ Young diagrams, with $N := |\nu| + |\tilde{\nu}|$,

$$(7.1) \quad \Psi_{\alpha, \nu, \tilde{\nu}} := \sum_{\mathbf{k}, \mathbf{l}, |\mathbf{k}| + |\mathbf{l}| = N} M_{\alpha, \mathbf{k}\mathbf{l}, \nu\tilde{\nu}}^N \Psi_{\alpha, \mathbf{k}, \mathbf{l}},$$

with the convention that $\Psi_{\alpha, \emptyset, \emptyset} = \Psi_{\alpha, 0, 0}$, which will be denoted by Ψ_α . The $(\Psi_{\alpha, \nu, \tilde{\nu}})_{\nu, \tilde{\nu}}$ (with $\nu \neq \emptyset$ or $\tilde{\nu} \neq \emptyset$) will be called descendant states. Since the coefficients $M_{\alpha, \mathbf{k}\mathbf{l}, \nu\tilde{\nu}}^N$ are analytic in $\alpha \in \mathbb{C}$ (see Prop 4.9), $\Psi_{\alpha, \nu, \tilde{\nu}}$ satisfy the same holomorphy properties (namely Prop 6.26) as $\Psi_{\alpha, \mathbf{k}, \mathbf{l}}$ provided that $|\mathbf{k}| + |\mathbf{l}| = |\nu| + |\tilde{\nu}|$. The main goal of this section is to compute the correlation functions of these descendant states and our strategy can be summarized as follows.

It turns out that for α real in the physical region, which we will call probabilistic region, we will be able to give a probabilistic representation for the descendant states thanks to the *intertwining* construction based on the descendant states $\Psi_{\alpha, \nu, \tilde{\nu}}^0$ of the GFF theory $\mu = 0$:

$$(7.2) \quad \lim_{t \rightarrow +\infty} e^{t(2\Delta_\alpha + |\nu| + |\tilde{\nu}|)} e^{-t\mathbf{H}} \Psi_{\alpha, \nu, \tilde{\nu}}^0 = \Psi_{\alpha, \nu, \tilde{\nu}}$$

for α real and negative enough (see Proposition 7.2). Indeed, in Subsection 7.3, we express the free descendant states in terms of contour integrals of the Stress Energy Tensor (SET) of the GFF theory, which we plug then in (7.2). Applying the Feynman-Kac formula for the propagator $e^{-t\mathbf{H}}$ in the expression (7.2), we obtain a probabilistic expression for the descendant states $\Psi_{\alpha,\nu,\tilde{\nu}}$ in terms of contour integrals of SET in LCFT¹⁴ (see Corollary 7.10). The Ward identities (Proposition 7.11) then allows us to translate these contour integrals in terms of differential operators: more precisely, correlation functions of descendant states $\Psi_{\alpha,\nu,\tilde{\nu}}$ can be obtained in terms of differential operators acting on the correlation functions of primary states Ψ_α . Finally we will analytically continue these relations from the region $\{\alpha \in \mathbb{R}; \alpha < Q\}$ back to the spectrum line and obtain the consequences in Subsection 7.5 about the structure of 3 point correlation functions involving descendant states.

7.1. Highest weight states. Recall the definition (4.23) of the highest weight state $\Psi_\alpha^0(c, \varphi) = e^{(\alpha-Q)c}$ for $\alpha \in \mathbb{C}$ of the $\mu = 0$ theory. From Proposition 6.9 item 2 (applied with $F = 1$), we know that for $\alpha \in \mathbb{R}$ with $\alpha < Q$, the state

$$(7.3) \quad \Psi_\alpha = \mathcal{P}_0(\alpha)1$$

is given by the large time limit

$$(7.4) \quad \Psi_\alpha = \lim_{t \rightarrow \infty} e^{2t\Delta_\alpha} e^{-t\mathbf{H}} \Psi_\alpha^0, \quad dc \otimes \mathbb{P}_\mathbb{T} \text{ a.e.}$$

where Δ_α denotes the conformal weight (1.9). In physics (or representation theory) terminology the state Ψ_α is the highest weight state corresponding to the *primary field* V_α . Combining (7.4) with the Feynman-Kac formula (5.4) leads to the probabilistic representation for $\alpha < Q$

$$(7.5) \quad \Psi_\alpha(c, \varphi) := e^{(\alpha-Q)c} \mathbb{E}_\varphi \left[\exp \left(-\mu e^{\gamma c} \int_{\mathbb{D}} |x|^{-\gamma\alpha} M_\gamma(dx) \right) \right].$$

We recall here that the integrability of $|x|^{-\gamma\alpha}$ with respect to $M_\gamma(dx)$ is detailed in [DKRV16].

Remark 7.1. *In forthcoming work, we will show that, for $\alpha \in (\frac{2}{\gamma}, Q)$, we have as $c \rightarrow -\infty$*

$$\Psi_\alpha(c, \varphi) = e^{(\alpha-Q)c} + e^{(Q-\alpha)c} R(\alpha) + e^{(Q-\alpha)c} o(1)$$

with $o(1) \rightarrow 0$ in $L^2(\Omega_\mathbb{T})$ as $c \rightarrow -\infty$ with R the reflection coefficient defined in [KRV20], which thus appears as the scattering coefficient of constant functions: for $\ell = 0$ we have $\mathbf{S}_0(\alpha) = R(\alpha)\mathbf{Id}$. More generally, we will show that the scattering matrix is diagonal.

7.2. Descendant states. Recall (Subsection 4.4) that for $\mu = 0$ we have descendant states given by

$$\Psi_{\alpha,\nu,\tilde{\nu}}^0(c, \varphi) = \mathcal{Q}_{\alpha,\nu,\tilde{\nu}}(\varphi) e^{(\alpha-Q)c}$$

where $\mathcal{Q}_{\alpha,\nu,\tilde{\nu}}$ are eigenstates of the operator \mathbf{P} :

$$\mathbf{P} \mathcal{Q}_{\alpha,\nu,\tilde{\nu}} = (|\nu| + |\tilde{\nu}|) \mathcal{Q}_{\alpha,\nu,\tilde{\nu}}$$

so that

$$\mathbf{H}^0 \Psi_{\alpha,\nu,\tilde{\nu}}^0 = (2\Delta_\alpha + |\nu| + |\tilde{\nu}|) \Psi_{\alpha,\nu,\tilde{\nu}}^0.$$

From Proposition 6.9 we infer the following

Proposition 7.2. *Let $\alpha < (Q - \frac{2(|\nu|+|\tilde{\nu}|)}{\gamma} - \frac{\gamma}{4}) \wedge (Q - \gamma)$. Then the limit*

$$(7.6) \quad \lim_{t \rightarrow +\infty} e^{t(2\Delta_\alpha + |\nu| + |\tilde{\nu}|)} e^{-t\mathbf{H}} \Psi_{\alpha,\nu,\tilde{\nu}}^0 = \Psi_{\alpha,\nu,\tilde{\nu}}$$

holds in $e^{-(\beta+\gamma/2)\rho} L^2(\mathbb{R} \times \Omega_\mathbb{T})$ for $\beta > Q - \alpha - \gamma/2$.

¹⁴Recast in the language of CFTs, this is somewhat equivalent to stating that descendant states can be obtained via the action of Virasoro generators on primary states.

Proof. Write $|\nu| + |\tilde{\nu}| = \lambda_j$ for some j . We apply Proposition 6.9 item 1 but we have to make a small notational warning: indeed recall that in Section 6, eigenvalues are parametrized by $2\Delta_\alpha$ whereas here eigenvalues correspond to $2\Delta_\alpha + |\nu| + |\tilde{\nu}| = 2\Delta_\alpha + \lambda_j$. So let us call α' the α in the statement of Proposition 6.9, and write it as $\alpha' = Q + ip$ with $p^2 = 2\lambda_j - (Q - \alpha)^2$ ($p \in i\mathbb{R}$) in such a way that

$$2\Delta_{\alpha'} = 2\Delta_\alpha + \lambda_j, \quad \text{otherwise stated } \sqrt{p^2 - 2\lambda_j} = i(Q - \alpha).$$

In particular $\text{Im}\sqrt{p^2 - 2\lambda_j} = Q - \alpha > \gamma$ so that we can choose $\chi = 1$ in Proposition 6.9 item 1. Let $\ell \geq 1$, $j \leq \ell$ and $F = \mathcal{Q}_{\alpha, \nu, \tilde{\nu}}$. Then the limit (7.6) exists in $e^{-(\beta+\gamma/2)\rho} L^2$ if $\beta > \text{Im}\sqrt{p^2 - 2\lambda_j} - \gamma/2 = Q - \alpha - \gamma/2 > 0$ and $-p^2 > \beta^2$. In conclusion we get $\alpha < Q - \gamma/2 = 2/\gamma$ and $-p^2 > (Q - \alpha - \gamma/2)^2$. Substituting $p^2 = 2\lambda_j - (Q - \alpha)^2$ in the latter, we arrive at the relation $(Q - \alpha)^2 - 2\lambda_j > (Q - \alpha - \gamma/2)^2$, which can be solved to find our condition. \square

7.3. Stress Energy Field. In this section we construct a probabilistic representation for the Virasoro descendants (4.25). This can be done in terms of a local field, the *stress-energy tensor*, formally given for $z \in \mathbb{D}$ by

$$(7.7) \quad T(z) := Q\partial_z^2 X(z) - (\partial_z X(z))^2 + \mathbb{E}[(\partial_z X(z))^2].$$

The stress tensor does not make sense as a random field but can be given sense at the level of correlation functions as the limit of $T_\epsilon(z)$ defined by (7.7) with X replaced by a regularized field X_ϵ which we take in the form

$$(7.8) \quad X_\epsilon(z) := \langle X, f_{\epsilon, z} \rangle_{\mathbb{D}}$$

for suitable test function $f_{\epsilon, z}$. We will use two regularizations in what follows. For the first one we take $f_{\epsilon, z}(u) = \frac{1}{\epsilon^2} \varrho(\frac{z-u}{\epsilon})$ with non-negative $\varrho \in C_c^\infty(\mathbb{C})$ with $\varrho(0) = 1$ and $\int_{\mathbb{C}} \varrho(x) dx = 1$. For this regularization we have for $t \geq 0$ the following scaling relation

$$(7.9) \quad S_{-t} T_\epsilon(z) = e^{-2t} T_{e^{-t}\epsilon}(e^{-t}z).$$

For the second regularization we write $z = e^{-t+i\theta}$ and take $f_{\epsilon, z}(e^{-s+i\theta'}) = \frac{1}{2\pi\epsilon} \rho(\frac{t-s}{\epsilon}) \sum_{|n| < \epsilon^{-1}} e^{in(\theta-\theta')}$ with $\rho \in C_c^\infty(\mathbb{R})$, $\rho(0) = 1$ and $\int_{\mathbb{R}} \rho(x) dx = 1$. This regularization has the property that $T_\epsilon(z)$ depends only on the Fourier components φ_n with $|n| < 1/\epsilon$ when we decompose $X = P\varphi + X_{\mathbb{D}}$. We denote also by $\bar{T}(z)$ the complex conjugate of $T(z)$.

Our goal is to express the action of the Virasoro generators (4.17) and (4.18) on the state $U_0 F$ in terms of the states

$$(7.10) \quad U_0 \left(\prod_{i=1}^k T(u_i) \prod_{j=1}^l \bar{T}(v_j) F \right)$$

which will be defined as limits of regularized expressions. We start by specifying a suitable class of F for which (7.10) makes sense. Let $\delta < 1$. We introduce the set \mathcal{E}_δ defined by

$$(7.11) \quad \mathcal{E}_\delta := \left\{ f \in C_0^\infty(\delta\mathbb{D}) \mid f(e^{-t+i\theta}) = \sum_{n \in \mathbb{Z}} f_n(t) e^{in\theta}, \text{ with } f_n \in C_0^\infty((-\ln \delta, \infty)) \text{ and } f_n = 0 \text{ for } |n| \text{ large enough} \right\}.$$

Define $\mathcal{F}_\delta \subset \mathcal{F}_{\mathbb{D}}$ by

$$(7.12) \quad \mathcal{F}_\delta = \text{span} \left\{ \prod_{i=1}^l \langle g_i, c + X \rangle_{\mathbb{D}} e^{(f, c+X)_{\mathbb{D}}}; l \geq 0, f, g_i \in \mathcal{E}_\delta \right\}.$$

with $l \geq 0$, $f, g_i \in \mathcal{E}_\delta$. We note that $U_0 F \in e^{\beta|c|} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > |f_0 - Q|$ where $f_0 = \langle f, 1 \rangle_{\mathbb{D}}$ and is in the domain of the Virasoro operators $\mathbf{L}_{-\nu}^0 \tilde{\mathbf{L}}_{-\tilde{\nu}}^0$ defined in Subsection 4.4 since it depends on a finite number of φ_n . Let

$$(7.13) \quad \mathcal{O}_\delta = \{(\mathbf{u}, \mathbf{v}) \in \mathbb{C}^{m+n} \mid \delta < |u_j|, |v_j| < 1, \forall j \neq j', |u_j| \neq |u_{j'}|, |v_j| \neq |v_{j'}|, |u_j| \neq |v_{j'}|\}.$$

We have the simple:

Proposition 7.3. *Let $F \in \mathcal{F}_\delta$. Then the functions*

$$(\mathbf{u}, \mathbf{v}) \rightarrow U_0 \left(\prod_{i=1}^k T_{\epsilon_i}(u_i) \prod_{j=1}^l \bar{T}_{\epsilon'_j}(v_j) F \right)$$

are continuous on \mathcal{O}_δ with values in $e^{\beta|c|} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > |f_0 - Q|$ and converge uniformly on compact subsets of \mathcal{O}_δ as $(\epsilon_i)_i$ and $(\epsilon'_j)_j$ tend successively to 0 in whatever order. The limit is independent of the regularization or the order along which limits have been taken and denoted by (7.10) and it defines a function holomorphic in \mathbf{u} and anti-holomorphic in \mathbf{v} in the region \mathcal{O}_δ taking values in $e^{\beta|c|} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$.

Proof. To keep the notation simple we consider only the case $j = 0$ in (7.10) and the case where $F \in \mathcal{F}_\delta$ is of the form $F = e^{\langle f, c+X \rangle_{\mathbb{D}}}$ with $f \in \mathcal{E}_\delta$. By replacing f by $f + \sum_i \lambda_i g_i$ with $g_i \in \mathcal{E}_\delta$ and differentiating at $\lambda_i = 0$ we can deduce the result for general F . For such F we have

$$(7.14) \quad (U_0 F)(c, \varphi) = e^{(f_0 - Q)c} e^{\langle P\varphi, f \rangle_{\mathbb{D}}} e^{\frac{1}{2} \langle f, G_{\mathbb{D}} f \rangle_{\mathbb{D}}}.$$

By Gaussian integration by parts (see (9.4)) the right-hand side of (7.10) is a sum of terms

$$(7.15) \quad \text{const} \times \prod_i (\partial^{a_i} P\varphi_{\epsilon_i}(u_i))^{b_i} \prod_{j < k} (\partial_{u_j}^{b_{jk}} \partial_{u_k}^{c_{jk}} (f_{\epsilon_j, u_j}, G_{\mathbb{D}} f_{\epsilon_k, u_k})_{\mathbb{D}})^{d_{jk}} \prod_l \partial_{u_l}^{d_l} (f_{\epsilon_l, u_l}, G_{\mathbb{D}} f)_{\mathbb{D}}^{e_l} U_0 F$$

where $a_i, b_{jk}, c_{jk} \in \{1, 2\}$, $b_i, d_{jk}, e_l \in \{0, 1, 2\}$, $d_l \in \{1, 2\}$ and $G_{\mathbb{D}}$ is the Dirichlet Green function (3.8). The functions $(f_{\epsilon_j, u_j}, G_{\mathbb{D}} f_{\epsilon_k, u_k})_{\mathbb{D}}$ and $(f_{\epsilon_l, u_l}, G_{\mathbb{D}} f)_{\mathbb{D}}$ converge uniformly on compacts of \mathcal{O}_δ to smooth functions. From (3.8) we get

$$(7.16) \quad \partial_z G_{\mathbb{D}}(z, u) = -\frac{1}{2} \left(\frac{1}{z - u} - \frac{1}{z - \frac{1}{\bar{u}}} \right).$$

Hence the limit of the second product in (7.15) is holomorphic since $b_{jk}, c_{jk} > 0$. For the third product, we get convergence to terms of the form

$$\int_{\mathbb{D}} \partial_u G_{\mathbb{D}}(u, z) f(z) dz = -\frac{1}{2} \sum_{n=0}^{\infty} (f_n u^{-n-1} + f_{-n-1} u^n)$$

or its ∂_u derivative where $f_n = \langle f, u^n \rangle_{\mathbb{D}}$ and $f_{-n} = \langle f, \bar{u}^n \rangle_{\mathbb{D}}$ and the sum is finite and holomorphic. Finally, recalling (3.7) it is easy to check that the first product converges in $L^2(\Omega_{\mathbb{T}})$ uniformly on compacts of \mathcal{O}_δ to a holomorphic function $g(\mathbf{u})$ and $g(\mathbf{u}) U_0 F \in e^{\beta|c|} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. \square

Now we will consider contour integrals of observables of the type (7.10), for which we use the following notation: for $f : \mathbb{D} \rightarrow \mathbb{C}$ and $\delta > 0$

$$\oint_{|z|=\delta} f(z) dz := i\delta \int_0^{2\pi} f(\delta e^{i\theta}) e^{i\theta} d\theta, \quad \oint_{|z|=\delta} f(z) d\bar{z} := i\delta \int_0^{2\pi} f(\delta e^{i\theta}) e^{-i\theta} d\theta.$$

Then we have

Lemma 7.4. *Let $F \in \mathcal{F}_\delta$ and $\delta' > \delta$. Then for all $n > 0$*

$$(7.17) \quad \frac{1}{2\pi i} \oint_{|z|=\delta'} z^{1-n} U_0(T(z)F) dz = \mathbf{L}_{-n}^0 U_0 F, \quad \frac{1}{2\pi i} \oint_{|z|=\delta'} \bar{z}^{1-n} U_0(\bar{T}(z)F) d\bar{z} = \tilde{\mathbf{L}}_{-n}^0 U_0 F.$$

Proof. Here again (for simplicity) we consider the case where $F \in \mathcal{F}_{\delta\mathbb{D}}$ is of the form $F = e^{\langle f, c+X \rangle_{\mathbb{D}}}$ with $f \in \mathcal{E}_\delta$. Let us write the integration by parts terms explicitly. First

$$U_0(\partial_z X(z)F) = \partial_z P\varphi(z) U_0 F + \int_{\mathbb{D}} \partial_z G_{\mathbb{D}}(z, u) f(u) du U_0 F.$$

From (7.16) we get

$$(7.18) \quad \partial_z G_{\mathbb{D}}(z, u) = -\frac{1}{2} \sum_{n=0}^{\infty} (u^n z^{-n-1} + \bar{u}^{n+1} z^n)$$

which converges since $|u| < \delta < \delta' = |z| < 1$. Therefore

$$\int_{\mathbb{D}} \partial_z G_{\mathbb{D}}(z, u) f(u) du = -\frac{1}{2} \sum_{n=0}^{\infty} (f_n z^{-n-1} + f_{-n-1} z^n)$$

where $f_n = \langle f, u^n \rangle_{\mathbb{D}}$ and $f_{-n} = \langle f, \bar{u}^n \rangle_{\mathbb{D}}$ for $n \geq 0$. Recalling (3.7), we have obtained

$$(7.19) \quad \begin{aligned} U_0(\partial_z X(z)F) &= \sum_{n \geq 0} (n\varphi_n z^{n-1} - \frac{1}{2}(f_n z^{-n-1} + f_{-n-1} z^n)) U_0 F \\ &= \sum_{n \in \mathbb{Z}} z^{n-1} (n\varphi_n 1_{n>0} - \frac{1}{2} f_{-n}) U_0 F. \end{aligned}$$

By (7.14)

$$(7.20) \quad (U_0 F)(c, \varphi) = e^{(f_0 - Q)c} e^{\sum_{n>0} (\varphi_n f_n + \varphi_{-n} f_{-n})} e^{\frac{1}{2} \langle f, G_{\mathbb{D}} f \rangle_{\mathbb{D}}}$$

so that

$$(7.21) \quad \begin{aligned} U_0(\partial_z X(z)F) &= (-\frac{1}{2} f_0 z^{-1} + \sum_{n \neq 0} z^{n-1} (n\varphi_n 1_{n>0} - \frac{1}{2} \partial_{-n})) U_0 F \\ &= i \sum_{n \in \mathbb{Z}} z^{n-1} \mathbf{A}_{-n} U_0 F \end{aligned}$$

where we recall that $\mathbf{A}_0 = \frac{i}{2}(\partial_c + Q)$. Hence

$$(7.22) \quad U_0(Q \partial_z^2 X(z)F) = -iQ \sum_{n \in \mathbb{Z}} (n+1) z^{-n-2} \mathbf{A}_n U_0 F.$$

Next consider the quadratic terms in T and use Gaussian integrating by parts twice to get

$$\begin{aligned} U_0(((\partial_z X(z))^2 - \mathbb{E}[(\partial_z X(z))^2])F) &= \left(\partial_z P \varphi(z) + \int_{\mathbb{D}} \partial_z G_{\mathbb{D}}(z, u) f(u) du \right)^2 U_0 F \\ &= \left(\sum_{n \in \mathbb{Z}} z^{n-1} (n\varphi_n 1_{n>0} - \frac{1}{2} f_{-n}) \right)^2 U_0 F \\ &= \sum_{n, m} z^{n+m-2} (-\mathbf{A}_{-n} \mathbf{A}_{-m} + \frac{m}{2} \delta_{m, -n} 1_{m>0}) U_0 F \\ &= - \sum_{n, m} z^{n+m-2} : \mathbf{A}_{-n} \mathbf{A}_{-m} : U_0 F \end{aligned}$$

where we used (7.19) in the second step, (7.21) in the third step and $\mathbf{A}_m \mathbf{A}_{-m} = \mathbf{A}_{-m} \mathbf{A}_m + \frac{m}{2}$ for $m > 0$ in the last step. The sum converges in $e^{\beta|c|} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$. Combining this with (7.22) the claim (7.17) follows upon doing the contour integral. The claim for \bar{T} is proved in the same way. \square

Let us now introduce some notation for the general correlations (7.10). Denote $\mathbf{u} = (u_1, \dots, u_k) \in \mathbb{D}^k$. We define nested contour integrals for $f : \mathbb{D}^k \times \mathbb{D}^j \rightarrow \mathbb{C}$ by

$$(7.23) \quad \oint_{|\mathbf{u}|=\delta} \oint_{|\mathbf{v}|=\tilde{\delta}} f(\mathbf{u}, \mathbf{v}) d\bar{\mathbf{v}} d\mathbf{u} := \oint_{|u_k|=\delta_k} \dots \oint_{|u_1|=\delta_1} \oint_{|v_j|=\tilde{\delta}_j} \dots \oint_{|v_1|=\tilde{\delta}_1} f(\mathbf{u}, \mathbf{v}) d\bar{v}_1 \dots d\bar{v}_j du_1 \dots du_k.$$

where $\delta := (\delta_1, \dots, \delta_k)$ with $0 < \delta_1 < \dots < \delta_k < 1$ and similarly for $\tilde{\delta}$. Furthermore we always suppose $\delta_i \neq \tilde{\delta}_j$ for all i, j . Next, for $\epsilon \in (\mathbb{R}^+)^k$ and $\epsilon' \in (\mathbb{R}^+)^j$ we set $T_{\epsilon}(\mathbf{u}) := \prod_{i=1}^k T_{\epsilon_i}(u_i)$ (and similarly for the anti-holomorphic part) and given two Young diagrams $\nu = (\nu_i)_{1 \leq i \leq k}$, $\tilde{\nu} = (\tilde{\nu}_i)_{1 \leq i \leq j}$, we denote $\mathbf{u}^{1-\nu} = \prod u_i^{1-\nu_i}$ and $\bar{\mathbf{v}}^{1-\tilde{\nu}} = \prod \bar{v}_i^{1-\tilde{\nu}_i}$. With these notations we have:

Proposition 7.5. *Let $F \in \mathcal{F}_{\delta}$ and $\delta < \delta_1 \wedge \tilde{\delta}_1$. Then*

$$(7.24) \quad (2\pi i)^{-k-j} \oint_{|\mathbf{u}|=\delta} \oint_{|\mathbf{v}|=\tilde{\delta}} \mathbf{u}^{1-\nu} \bar{\mathbf{v}}^{1-\tilde{\nu}} U_0(T(\mathbf{u})\bar{T}(\mathbf{v})F) d\bar{\mathbf{v}} d\mathbf{u} = \mathbf{L}_{-\nu}^0 \mathbf{L}_{-\tilde{\nu}}^0 U_0 F.$$

Proof. For simplicity consider again the case with only T insertions. We proceed by induction in k . By Lemma 7.4 the claim holds for $k = 1$. Suppose it holds for $k - 1$. We use the second regularization introduced above. This entails that $T_{\epsilon}(u_i) \in \mathcal{F}_{\delta_{i+1}}$ for ϵ small enough. By Proposition 7.3 we have

$$\begin{aligned} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} U_0(T(\mathbf{u})F) d\mathbf{u} &= \lim_{\epsilon^{(k)} \rightarrow 0} \lim_{\epsilon_k \rightarrow 0} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} U_0(T_{\epsilon_k}(u_k) T_{\epsilon^{(k)}}(\mathbf{u}^{(k)}) F) d\mathbf{u} \\ &= \lim_{\epsilon^{(k)} \rightarrow 0} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} U_0(T(u_k) T_{\epsilon^{(k)}}(\mathbf{u}^{(k)}) F) d\mathbf{u} \end{aligned}$$

where we introduced the notation $\epsilon^{(k)} = (\epsilon_1, \dots, \epsilon_{k-1})$, $\mathbf{u}^{(k)} = (u_1, \dots, u_{k-1})$. We have, for $\epsilon^{(k)}$ small enough that $T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F \in \mathcal{F}_{\delta_k}$. Hence by Lemma 7.4

$$\frac{1}{2\pi i} \oint_{|u_k|=\delta_k} u_k^{1-\nu_k} U_0(T(u_k)T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F) du_k = \mathbf{L}_{-\nu_k} U_0(T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F).$$

From (7.15) we infer that $U_0(T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F) = P_k(\epsilon^{(k)}, \varphi) U_0 F$ where $P_k(\epsilon^{(k)}, \varphi)$ is a polynomial in finitely many variables φ_n (depending on k) with coefficients continuous in $\mathbf{u}^{(k)}$ (and depending on $\epsilon^{(k)}$). Therefore we may commute $\mathbf{L}_{-\nu_k}$ and the integration to get

$$(2\pi i)^{-k} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} U_0(T(u_k)T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F) d\mathbf{u} = \mathbf{L}_{-\nu_k} \left((2\pi i)^{-k+1} \oint_{|\mathbf{u}^{(k)}|=\delta^{(k)}} (\mathbf{u}^{(k)})^{1-\nu^{(k)}} U_0(T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F) d\mathbf{u}^{(k)} \right).$$

By the induction hypothesis the term

$$(7.25) \quad I_{\epsilon^{(k)}} := (2\pi i)^{-k+1} \oint_{|\mathbf{u}^{(k)}|=\delta^{(k)}} (\mathbf{u}^{(k)})^{1-\nu^{(k)}} U_0(T_{\epsilon^{(k)}}(\mathbf{u}^{(k)})F) d\mathbf{u}^{(k)}$$

converges to $\mathbf{L}_{-\nu^{(k)}} U_0 F$ as $\epsilon^{(k)}$ goes to 0. Moreover we claim that, for all fixed k , there exists $M, N < \infty$ s.t. $I_{\epsilon^{(k)}} = Q_k(\epsilon^{(k)}, \varphi) U_0 F$ where $Q_k(\epsilon^{(k)}, \varphi)$ is a polynomial in φ_n with $|n| < N$ and of degree less than M . Furthermore the coefficients of $Q_k(\epsilon^{(k)}, \varphi)$ converge as $\epsilon^{(k)} \rightarrow 0$. Therefore we can commute $\mathbf{L}_{-\nu_k}$ and the limit to get

$$\lim_{\epsilon^{(k)} \rightarrow 0} \mathbf{L}_{-\nu_k} I_{\epsilon^{(k)}} = \mathbf{L}_{-\nu_k} \lim_{\epsilon^{(k)} \rightarrow 0} I_{\epsilon^{(k)}} = \mathbf{L}_{-\nu_k} \mathbf{L}_{-\nu^{(k)}} U_0(F) = \mathbf{L}_{-\nu} U_0(F)$$

which completes the induction step.

To prove the above claim we use (7.15). Let \mathcal{G} be the graph with vertex set $\{1, \dots, k\}$ and edges $\{i, j\}$ with $d_{ij} > 0$. Let Γ be a connected component of \mathcal{G} . Then the number of vertices i in Γ s.t. $b_i > 0$ is no more than two. Consider a connected component for which the number is two. This corresponds to a subproduct in (7.15) of the form

$$(7.26) \quad f_{\epsilon}(u_{i_1}, \dots, u_{i_l}) = \partial P \varphi_{\epsilon_{i_1}}(u_{i_1}) \partial P \varphi_{\epsilon_{i_l}}(u_{i_l}) \prod_{a=1}^{l-1} \partial_{u_{i_a}} \partial_{u_{i_{a+1}}} (f_{\epsilon_{i_a}, u_{i_a}}, G_{\mathbb{D}} f_{\epsilon_{i_{a+1}}, u_{i_{a+1}}})_{\mathbb{D}}.$$

We have

$$\partial P \varphi_{\epsilon}(u) = \sum_{n=1}^{M_{\epsilon}} a_n(\epsilon) \varphi_n u^{n-1}$$

where $M_{\epsilon} < \infty$ if $\epsilon > 0$ and $a_n(\epsilon)$ converges as $\epsilon \rightarrow 0$. Similarly, for $|u| < |u'|$,

$$\partial_u \partial_{u'} (f_{\epsilon, u}, G_{\mathbb{D}} f_{\epsilon', u'})_{\mathbb{D}} = \sum_{m=0}^{N_{\epsilon, \epsilon'}} b_m(\epsilon, \epsilon') u^m u'^{-m-2}$$

where $N_{\epsilon, \epsilon'} < \infty$ if $\epsilon, \epsilon' > 0$ and $b_m(\epsilon, \epsilon')$ converge as $\epsilon, \epsilon' \rightarrow 0$. Insert these to (7.26). To simplify notation denote $u_{i_a} = v_a$, $\epsilon_{i_a} = \varepsilon_a$, $\delta_{i_a} = d_a$ and $\nu_{i_a} = \eta_a$. The contour integral of f_{ϵ} becomes

$$\oint_{|\mathbf{v}|=d} \mathbf{v}^{1-\boldsymbol{\eta}} f_{\boldsymbol{\varepsilon}}(\mathbf{v}) d\mathbf{v} = \sum_{n_1, n_l} \varphi_{n_1} \varphi_{n_l} a_{n_1, n_l}(\boldsymbol{\varepsilon})$$

where

$$a_{n_1, n_l}(\boldsymbol{\varepsilon}) = \sum_{m_1, \dots, m_{l-1} > 0} a_{n_1, n_l}(\mathbf{m}, \boldsymbol{\varepsilon}) \int_{[0, 2\pi]^l} e^{i(n_1 \theta_1 + n_l \theta_l + \sum_{i=1}^{l-1} \alpha_i (m_i \theta_i - (m_i + 2) \theta_{i+1}) + \sum_{i=1}^l (1 - \eta_i) \theta_i)} d\boldsymbol{\theta}$$

and $\alpha_i = \pm 1$ depending on whether $|v_a| < |v_{a+1}|$ or the opposite. The $\boldsymbol{\theta}$ integral gives the constraints $n_1 + \alpha_1 m_1 + 1 - \eta_1 = 0$, $n_l - \alpha_{l-1} (m_{l-1} + 2) + 1 - \eta_l = 0$ and $\alpha_i m_i - \alpha_{i-1} (m_{i-1} + 2) + (1 - \eta_i) = 0$ for $i = 2, \dots, l-1$. The last two imply $n_l - \alpha_1 m_1 + C(\boldsymbol{\alpha}, \boldsymbol{\eta}) = 0$ and combining this with the first constraint we get $a_{n_1, n_l} = 0$ if $|n_1|, |n_l| > C(\boldsymbol{\alpha}, \boldsymbol{\eta})$, where $C(\boldsymbol{\alpha}, \boldsymbol{\eta})$ denotes a generic constant depending only on $\boldsymbol{\alpha}, \boldsymbol{\eta}$. Convergence of the $a_{n_1, n_l}(\boldsymbol{\varepsilon})$ as $\boldsymbol{\varepsilon} \rightarrow 0$ then follows from the convergence of the $(a_n)_n$'s and (b_m) 's and the fact that $a_{n_1, n_l}(\boldsymbol{\varepsilon})$ is a polynomial of these coefficients. This finishes the proof for the case Γ has two vertices with $b_i > 0$. The two other cases are similar. \square

In the sequel we will apply Proposition 7.5 to the function

$$F = S_{e^{-t}} U_0^{-1} \Psi_\alpha^0 = e^\alpha \int_0^{2\pi} (c + X(e^{-t+i\theta})) \frac{d\theta}{2\pi} - \alpha Q t$$

where $\Psi_\alpha^0(c, \varphi) = e^{(\alpha-Q)c}$ (in this case integration against a function $f \in \mathcal{E}_\delta$ is replaced by an average on a circle but the previous considerations apply also). Then $U_0 F = e^{-t\mathbf{H}} \Psi_\alpha^0 = e^{-2t\Delta_\alpha} \Psi_\alpha^0$. Thus we arrive to the representation for the Virasoro descendants

$$(7.27) \quad \Psi_{\alpha, \nu, \tilde{\nu}}^0 = e^{2t\Delta_\alpha} (2\pi i)^{-k-j} \oint_{|\mathbf{u}|=\delta} \oint_{|\mathbf{v}|=\tilde{\delta}} \mathbf{u}^{1-\nu} \bar{\mathbf{v}}^{1-\tilde{\nu}} U_0(T(\mathbf{u})\bar{T}(\mathbf{v})S_{e^{-t}}U_0^{-1}\Psi_\alpha^0) d\bar{\mathbf{v}} d\mathbf{u}$$

where now $e^{-t} < \delta_1 \wedge \tilde{\delta}_1$.

7.4. Conformal Ward Identities. In this section we state the main identity relating a LCFT correlation function with a V_α insertion to a scalar product with the descendant states $\Psi_{\alpha, \nu, \tilde{\nu}}$ given by (7.1). We have

Lemma 7.6. *We have*

$$e^{-t\mathbf{H}} U_0(T(\mathbf{u})\bar{T}(\mathbf{v})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0) = \lim_{\epsilon \rightarrow 0} \lim_{\epsilon' \rightarrow 0} e^{-t\mathbf{H}} U_0(T_\epsilon(\mathbf{u})\bar{T}_{\epsilon'}(\mathbf{v})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0)$$

where the limit is in $e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $\beta > Q - \alpha$, uniformly in $(\mathbf{u}, \mathbf{v}) \in \mathcal{O}_{e^{-s}}$ (recall (7.13)) and the LHS is analytic in \mathbf{u} and anti-analytic in \mathbf{v} on $\mathcal{O}_{e^{-s}}$.

Proof. This follows from Proposition 7.3 and (6.11) applied with $\beta > Q - \alpha$. \square

The following lemma gives a probabilistic expression for $e^{-t\mathbf{H}} \Psi_{\alpha, \nu, \tilde{\nu}}^0$ (recall our convention for contour integrals in (7.23)):

Lemma 7.7. *Let $\delta_k \wedge \tilde{\delta}_k < e^{-t}$. Then*

$$e^{-t\mathbf{H}} \Psi_{\alpha, \nu, \tilde{\nu}}^0 = \frac{e^{-(2\Delta_\alpha + |\nu| + |\tilde{\nu}|)t}}{(2\pi i)^{k+j}} \oint_{|\mathbf{u}|=\delta} \oint_{|\mathbf{v}|=\tilde{\delta}} \mathbf{u}^{1-\nu} \bar{\mathbf{v}}^{1-\tilde{\nu}} e^{-Qc} \mathbb{E}_\varphi(T(\mathbf{u})\bar{T}(\mathbf{v})V_\alpha(0)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)}) d\bar{\mathbf{v}} d\mathbf{u}$$

where

$$\mathbb{E}_\varphi(T(\mathbf{u})\bar{T}(\mathbf{v})V_\alpha(0)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)}) := \lim_{\epsilon \rightarrow 0} \lim_{\epsilon' \rightarrow 0} \mathbb{E}_\varphi(T_\epsilon(\mathbf{u})\bar{T}_{\epsilon'}(\mathbf{v})V_\alpha(0)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)})$$

and the limit exists in $e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for all $\beta > Q - \alpha$ and is analytic in \mathbf{u} and anti-analytic in \mathbf{v} in the region $\mathcal{O}_{e^{-t}}$.

Proof. For the sake of readability we write the proof in the case when $\tilde{\nu} = 0$. Thus, consider

$$(7.28) \quad \Psi_{\alpha, \nu, 0}^0 = \frac{e^{2\Delta_\alpha s}}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} U_0(T(\mathbf{u})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0) d\mathbf{u}.$$

We use the regularisation T_ϵ where (7.9) holds. By Lemma 7.6

$$\begin{aligned} e^{-t\mathbf{H}} U_0(T(\mathbf{u})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0) &= \lim_{\epsilon \rightarrow 0} e^{-t\mathbf{H}} U_0(T_\epsilon(\mathbf{u})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0) \\ &= \lim_{\epsilon \rightarrow 0} e^{-t\mathbf{H}} U(T_\epsilon(\mathbf{u})(S_{e^{-s}}U_0^{-1}\Psi_\alpha^0)e^{\mu e^{\gamma c} M_\gamma(\mathbb{D})}) \\ &= \lim_{\epsilon \rightarrow 0} U(S_{e^{-t}}(T_\epsilon(\mathbf{u})(S_{e^{-s}}U_0^{-1}\Psi_\alpha^0)e^{\mu e^{\gamma c} M_\gamma(\mathbb{D})})) \\ &= \lim_{\epsilon \rightarrow 0} e^{-2t} U(T_{e^{-t}\epsilon}(e^{-t}\mathbf{u})(S_{e^{-s-t}}U_0^{-1}\Psi_\alpha^0)S_{e^{-t}}(e^{\mu e^{\gamma c} M_\gamma(\mathbb{D})})) \\ &= e^{-2t} e^{-Qc} \lim_{\epsilon \rightarrow 0} \mathbb{E}_\varphi(T_{e^{-t}\epsilon}(e^{-t}\mathbf{u})(S_{e^{-s-t}}U_0^{-1}\Psi_\alpha^0)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)}) \\ &:= e^{-2t} e^{-Qc} \mathbb{E}_\varphi(T(e^{-t}\mathbf{u})(S_{e^{-s-t}}U_0^{-1}\Psi_\alpha^0)e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)}) \end{aligned}$$

where we used (7.9) in the fourth identity. By Lemma 7.6 the last expression is analytic in \mathbf{u} and since $e^{2\Delta_\alpha s} e^{-t\mathbf{H}} U_0(T(\mathbf{u})S_{e^{-s}}U_0^{-1}\Psi_\alpha^0)$ is independent on s we can take the limit $s \rightarrow \infty$. For this we note that

$$S_{e^{-s-t}} U_0^{-1} \Psi_\alpha^0 = e^{\alpha(c-(t+s)Q)} e^{\alpha(1, X(e^{-t-s} \cdot))_{\mathbb{T}}} = e^{-2\Delta_\alpha(s+t)} e^{\alpha c} e^{\alpha(1, X(e^{-t-s} \cdot))_{\mathbb{T}}} - \frac{1}{2} \alpha^2 \mathbb{E}(1, X(e^{-t-s} \cdot))_{\mathbb{T}}^2$$

so that

$$e^{2\Delta_\alpha s} \mathbb{E}_\varphi \left(T(e^{-t} \mathbf{u}) (S_{e^{-s-t}} U_0^{-1} \Psi_\alpha^0) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)} \right) = e^{-2\Delta_\alpha t} \mathbb{E}_\varphi \left(T(e^{-t} \mathbf{u}) V_\alpha(0) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)} \right)$$

and the last expression is analytic in \mathbf{u} . Hence by a change of variables in the \mathbf{u} -integral

$$\begin{aligned} e^{-t\mathbf{H}} \Psi_{\alpha, \nu, 0}^0 &= \frac{e^{2\Delta_\alpha s}}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} e^{-t\mathbf{H}} U_0 \left(T(\mathbf{u}) S_{e^{-s}} U_0^{-1} \Psi_\alpha^0 \right) d\mathbf{u} \\ &= \frac{e^{-(2\Delta_\alpha + |\nu|)t}}{(2\pi i)^k} \oint_{|\mathbf{u}|=e^{-t}\delta} \mathbf{u}^{1-\nu} e^{-Qc} \mathbb{E}_\varphi \left(T(\mathbf{u}) V_\alpha(0) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)} \right) d\mathbf{u} \\ &= \frac{e^{-(2\Delta_\alpha + |\nu|)t}}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta} \mathbf{u}^{1-\nu} e^{-Qc} \mathbb{E}_\varphi \left(T(\mathbf{u}) V_\alpha(0) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D} \setminus \mathbb{D}_t)} \right) d\mathbf{u} \end{aligned}$$

where in the last step we used analyticity to move the contours to $|\mathbf{u}| = \delta$. \square

In what follows, for fixed $n \geq 1$, we will denote

$$(7.29) \quad \mathcal{Z} := \{ \mathbf{z} = (z_1, \dots, z_n) \mid \forall i \neq j, z_i \neq z_j \text{ and } \forall i, |z_i| < 1 \}.$$

Denoting $\theta(\mathbf{z}) = (\theta(z_1), \dots, \theta(z_n)) \in \mathbb{C}^n$ we have $\theta\mathcal{Z} = \{(z_1, \dots, z_n) \mid \forall i \neq j, z_i \neq z_j \text{ and } \forall i, |z_i| > 1\}$.

For $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$ such that $\alpha_i < Q$ for all i we define the function $U_\alpha(\mathbf{z}) : \mathbb{R} \times \Omega_{\mathbb{T}} \rightarrow \mathbb{R}$ by

$$(7.30) \quad U_\alpha(\mathbf{z}, c, \varphi) := \lim_{\epsilon \rightarrow 0} e^{-Qc} \mathbb{E}_\varphi \left[\left(\prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \right) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D})} \right], \quad \text{for } \mathbf{z} \in \mathcal{Z}$$

where $V_{\alpha_i, \epsilon}$ stands for the regularized vertex operator (1.8). Let us set

$$(7.31) \quad s := \sum_{i=1}^n \alpha_i.$$

Remark 7.8. *It follows directly from the construction of correlation functions that for $\mathbf{z} \in \theta\mathcal{Z}$*

$$\langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_{\gamma, \nu} = \left(\prod_{i=1}^n |z_i|^{-4\Delta_{\alpha_i}} \right) \langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2$$

and these expressions are finite if $\alpha + s > 2Q$ and $\alpha, \alpha_i < Q$.

Lemma 7.9. *Let $\mathbf{z} \in \theta\mathcal{Z}$. Then almost everywhere in c, φ and for all $R > 0$*

$$U_\alpha(\theta(\mathbf{z}))(c, \varphi) \leq e^{(s-Q)(c \wedge 0) - R(c \vee 0)} A(\varphi)$$

where $A \in L^2(\Omega_{\mathbb{T}})$.

Proof. Let $r = \max_i |\theta(z_i)|$ and $\iota(\varphi) = \inf_{x \in \mathbb{D}_r} P\varphi(x)$ and $\sigma(\varphi) = \sup_{x \in \mathbb{D}_r} P\varphi(x)$ with \mathbb{D}_r the disk centered at 0 with radius r . Then

$$U_\alpha(\theta(\mathbf{z}))(c, \varphi) \leq C e^{-Qc} e^{(c+\sigma(\varphi))s} \mathbb{E} e^{-\mu e^{\gamma(c+\iota(\varphi))} Z}$$

where the expectation is over the Dirichlet GFF $X_{\mathbb{D}}$ and

$$Z = \int_{\mathbb{D}_r} (1 - |z|^2)^{\frac{\gamma}{2}} e^{\sum_i \gamma \alpha_i G_{\mathbb{D}}(z, \theta(z_i))} M_{\gamma, \mathbb{D}}(dz)$$

where $M_{\gamma, \mathbb{D}}$ is the GMC of $X_{\mathbb{D}}$. For $c < 0$ we use the trivial bound

$$U_\alpha(\theta(\mathbf{z}))(c, \varphi) \leq C e^{(s-Q)c} e^{\sigma(\varphi)s}$$

and for $c > 0$ we note that Z has all negative moments so that for $a > 0$

$$\mathbb{E} e^{-aZ} = \mathbb{E} (aZ)^{-n} (aZ)^n e^{-aZ} \leq n! \mathbb{E} (aZ)^{-n} \leq C_n a^{-n}$$

implying

$$U_\alpha(\theta(\mathbf{z}))(c, \varphi) \leq C_n e^{c(s-Q-\gamma n)} e^{s\sigma(\varphi) - n\gamma\iota(\varphi)}$$

Since $e^{s\sigma(\varphi) - n\gamma\iota(\varphi)}$ is in $L^2(\mathbb{P})$ for all s, n the claim follows. \square

Define now the modified Liouville expectation (with now \mathbb{D}_t the disk centered at 0 with radius e^{-t})

$$(7.32) \quad \langle F \rangle_t = \int_{\mathbb{R}} e^{-2Qc} \mathbb{E} \left[F(c, X) e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{C} \setminus \mathbb{D}_t)} \right] dc.$$

Also, in the contour integrals below, for vectors $\boldsymbol{\delta}, \tilde{\boldsymbol{\delta}}$ defining the radii of the respective contours, we will put a subscript t when these variables are multiplied by e^{-t} , namely $\boldsymbol{\delta}_t := e^{-t} \boldsymbol{\delta}$ and similarly for $\tilde{\boldsymbol{\delta}}_t$. Then we get

Corollary 7.10. *Let $\mathbf{z} \in \theta\mathcal{Z}$. For $\alpha \in \mathbb{R}$ such that $\alpha < (Q - \frac{2(|\nu|+|\bar{\nu}|)}{\gamma} - \frac{\gamma}{4}) \wedge (Q - \gamma)$ and $\alpha + \sum_i \alpha_i > 2Q$, we have*

$$\begin{aligned} & \langle \Psi_{\alpha, \nu, \nu'} | U_{\alpha}(\theta(\mathbf{z})) \rangle_2 \\ &= \left(\prod_{i=1}^n |z_i|^{4\Delta_{\alpha_i}} \right) \times \frac{1}{(2\pi i)^{k+j}} \lim_{t \rightarrow \infty} \oint_{|\mathbf{u}|=\boldsymbol{\delta}_t} \oint_{|\mathbf{v}|=\tilde{\boldsymbol{\delta}}'_t} \mathbf{u}^{1-\nu} \bar{\mathbf{v}}^{1-\nu'} \langle T(\mathbf{u}) \bar{T}(\mathbf{v}) V_{\alpha}(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\bar{\mathbf{v}} d\mathbf{u}. \end{aligned}$$

Proof. Combining Proposition 7.2 and Lemma 7.9 the existence of the limit

$$\langle \Psi_{\alpha, \nu, \nu'} | U_{\alpha}(\theta(\mathbf{z})) \rangle_2 = \lim_{t \rightarrow \infty} e^{(2\Delta_{\alpha} + |\nu| + |\nu'|)t} \langle e^{-t\mathbf{H}} \Psi_{\alpha, \nu, \nu'}^0 | U_{\alpha}(\theta(\mathbf{z})) \rangle_2$$

follows. By Lemma 7.7 the RHS is given by the RHS of (7.10). \square

Here is the main result of this section:

Proposition 7.11. *Let $\mathbf{z} \in \theta\mathcal{Z}$. For $\alpha \in \mathbb{R}$ such that $\alpha < (Q - \frac{2(|\nu|+|\bar{\nu}|)}{\gamma} - \frac{\gamma}{4}) \wedge (Q - \gamma)$ and $\alpha + \sum_i \alpha_i > 2Q$ and for all i $\alpha_i < Q$, we have in the distributional sense*

$$\langle \Psi_{\alpha, \nu, \bar{\nu}} | U_{\alpha}(\theta(\mathbf{z})) \rangle_2 = \left(\prod_{i=1}^n |z_i|^{4\Delta_{\alpha_i}} \right) \times \mathbf{D}_{\nu} \tilde{\mathbf{D}}_{\bar{\nu}} (V_{\alpha}(0) \prod_{i=1}^n V_{\alpha_i}(z_i))_{\gamma, \mu}$$

where the differential operators $\mathbf{D}_{\nu}, \tilde{\mathbf{D}}_{\bar{\nu}}$ are defined by

$$(7.33) \quad \mathbf{D}_{\nu} = \mathbf{D}_{\nu_k} \dots \mathbf{D}_{\nu_1} \quad \text{and} \quad \tilde{\mathbf{D}}_{\bar{\nu}} = \tilde{\mathbf{D}}_{\bar{\nu}_j} \dots \tilde{\mathbf{D}}_{\bar{\nu}_1}$$

where for $n \in \mathbb{N}$

$$(7.34) \quad \mathbf{D}_n = \sum_{i=1}^n \left(-\frac{1}{z_i^{n-1}} \partial_{z_i} + \frac{(n-1)}{z_i^n} \Delta_{\alpha_i} \right)$$

$$(7.35) \quad \tilde{\mathbf{D}}_n = \sum_{i=1}^n \left(-\frac{1}{\bar{z}_i^{n-1}} \partial_{\bar{z}_i} + \frac{(n-1)}{\bar{z}_i^n} \Delta_{\alpha_i} \right)$$

Proof. Section 9 will be devoted to the proof of this proposition. \square

7.5. Computing the 3-point correlation functions of descendant states. Now we exploit Proposition 7.11 to give *exact* analytic expressions for the 3-point correlation functions of descendant states. For this, we first need to introduce some notation: for $\nu = (\nu_i)_{i \in [1, k]}$ a Young diagram and some real $\Delta, \Delta', \Delta''$ we set

$$(7.36) \quad v(\Delta, \Delta', \Delta'', \nu) := \prod_{j=1}^k (\nu_j \Delta' - \Delta + \Delta'' + \sum_{u < j} \nu_u).$$

With this notation, we can state the following key result:

Proposition 7.12 (Conformal Ward identities). *Assume $\alpha_1, \alpha_2 < Q$ with $\alpha_1 + \alpha_2 > Q$ and $|z| < 1$. For all $P > 0$, the scalar product $\langle \Psi_{Q+iP, \nu, \bar{\nu}} | U_{\alpha_1, \alpha_2}(0, z) \rangle_2$ is explicitly given by the following expression*

$$(7.37) \quad \begin{aligned} & \langle \Psi_{Q+iP, \nu, \bar{\nu}} | U_{\alpha_1, \alpha_2}(0, z) \rangle_2 = v(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{Q+iP}, \bar{\nu}) v(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{Q+iP}, \nu) \\ & \quad \times \frac{1}{2} C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q+iP) \bar{z}^{|\nu|} z^{|\bar{\nu}|} |z|^{2(\Delta_{Q+iP} - \Delta_{\alpha_1} - \Delta_{\alpha_2})} \end{aligned}$$

where Δ_{α} are conformal weights (1.9) and $C_{\gamma, \mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q+iP)$ are the DOZZ structure constants (see appendix B for the definition).

The remaining part of this subsection is devoted to the proof of this proposition. We first need the following lemma concerning analyticity of $U_\alpha(\mathbf{z})$ in the parameter $\alpha \in \mathbb{C}^n$, proved in Appendix C.1.

Lemma 7.13. *For fixed $\mathbf{z} \in \mathcal{Z}$ the mapping $\alpha \mapsto U_\alpha(\mathbf{z}) \in e^{\beta' \rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ extends analytically to a complex neighborhood \mathcal{A}_U^n in \mathbb{C}^n of the set $\{\alpha \in \mathbb{R}^n \mid \forall i, \alpha_i < Q\}$, for arbitrary $\beta' < \text{Re}(s) - Q$ (recall (7.31)). This analytic extension is continuous in $(\alpha, \mathbf{z}) \in \mathcal{A}_U^n \times \mathcal{Z}$.*

The first conclusion we want to draw is the fact that the pairing of Ψ_α with $U_\alpha(\theta(\mathbf{z}))$ (in the case $n = 2$) is related to the DOZZ formula when α is on the spectrum line $Q + i\mathbb{R}$.

Lemma 7.14. *Here we fix $n = 2$ and we consider $\mathbf{z} \in \theta\mathcal{Z}$. The mapping $(\alpha, \alpha) \mapsto \langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2$ is continuous in the set*

$$\Xi := \{(\alpha, \alpha) \in \mathbb{C} \times \mathcal{A}_U^2 \mid \text{Re}(\alpha) \leq Q, \text{Re}(\alpha + \alpha_1 + \alpha_2) > 2Q\}$$

and analytic in the set $\Xi \cap \{\alpha \in \mathbb{C} \setminus \mathcal{D}_0\}$. Moreover, in the set Ξ , we have the relation

$$\langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2 = |z_1|^{2\Delta_{\alpha_2} - 2\Delta_\alpha + 2\Delta_{\alpha_1}} |z_1 - z_2|^{2\Delta_\alpha - 2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} |z_2|^{2\Delta_{\alpha_1} - 2\Delta_\alpha + 2\Delta_{\alpha_2}} \frac{1}{2} C_{\gamma, \mu}^{\text{DOZZ}}(\alpha, \alpha_1, \alpha_2).$$

In particular this relation holds for $\alpha = Q + iP$ with $P \in (0, +\infty)$ and $\alpha_1, \alpha_2 \in (-\infty, Q)$ with $\alpha_1 + \alpha_2 > Q$.

Proof. By Proposition 6.26 the mapping $\alpha \mapsto \Psi_\alpha \in e^{-\frac{\beta}{2}\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ is analytic on W_0 , i.e. in the region $\{\alpha \in \mathbb{C} \setminus \mathcal{D}_0 \mid Q - \text{Re}(\alpha) < \beta\}$ and continuous on \mathcal{D}_0 . Combining with Lemma 7.13 (with $n = 2$) produces directly the region of analyticity/continuity we claim. Furthermore when all the parameters $\alpha, \alpha_1, \alpha_2$ are real, by Remark 7.8 we have

$$\langle V_\alpha(0) V_{\alpha_1}(z_1) V_{\alpha_2}(z_2) \rangle_{\gamma, \nu} = \left(\prod_{i=1}^2 |z_i|^{-4\Delta_{\alpha_i}} \right) \langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2.$$

Also, for real parameters, the LHS coincide with the DOZZ formula [KRV20], namely

$$\langle V_\alpha(0) V_{\alpha_1}(z_1) V_{\alpha_2}(z_2) \rangle_{\gamma, \nu} = |z_1|^{2\Delta_{\alpha_2} - 2\Delta_\alpha - 2\Delta_{\alpha_1}} |z_1 - z_2|^{2\Delta_\alpha - 2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} |z_2|^{2\Delta_{\alpha_1} - 2\Delta_\alpha - 2\Delta_{\alpha_2}} \frac{1}{2} C_{\gamma, \mu}^{\text{DOZZ}}(\alpha, \alpha_1, \alpha_2).$$

This proves the claim. \square

Now we would like to use the Ward identities, i.e. Proposition 7.11, to express the correlations of descendant fields with two insertions $\langle \Psi_{\alpha, \nu, \tilde{\nu}} | U_\alpha(\theta(\mathbf{z})) \rangle_2$ (here with $n = 2$) in terms of differential operators applied to correlation of primaries $\langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2$ when the parameter α is close to the spectrum line $\alpha \in Q + i\mathbb{R}$. This is not straightforward because Proposition 7.11 is not only restricted to real values of the parameter but also because the constraint on α , which forces it to be negatively large, implies to have n large in order for the global Seiberg bound $\alpha + \sum_i \alpha_i > 2Q$ to be satisfied. Transferring Ward's relations close to the spectrum line is thus our next task.

For this, fix a pair of Young diagrams $\nu, \tilde{\nu}$ and ℓ such that $\lambda_\ell = |\nu| + |\tilde{\nu}|$. By Proposition 6.26 there exists a connected open set $\mathcal{A}_{\nu, \tilde{\nu}} := W_0 \cap W_\ell \subset \mathbb{C}$ such that the mappings $\alpha \mapsto \Psi_{\alpha, \nu, \tilde{\nu}} \in e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ and $\alpha \mapsto \Psi_\alpha \in e^{-\beta\rho} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for $\beta > Q - \text{Re}(\alpha)$ are analytic on $\mathcal{A}_{\nu, \tilde{\nu}}$ and furthermore

- $\mathcal{A}_{\nu, \tilde{\nu}}$ contains a complex neighborhood of the spectrum line $\{\alpha = Q + iP \mid P \in (0, +\infty)\}$, with the discrete set $\mathcal{D}_0 \cup \mathcal{D}_\ell$ removed and the mappings extend continuously to $\mathcal{D}_0 \cup \mathcal{D}_\ell$.
- $\mathcal{A}_{\nu, \tilde{\nu}}$ contains a complex neighborhood of the real half-line $(-\infty, Q - \frac{2\lambda_\ell}{\gamma} - \frac{\gamma}{4})$

Therefore, for arbitrary fixed n and $\mathbf{z} \in \theta\mathcal{Z}$, the pairings $(\alpha, \alpha) \mapsto \langle \Psi_{\alpha, \nu, \tilde{\nu}} | U_\alpha(\theta(\mathbf{z})) \rangle_2$ and $(\alpha, \alpha) \mapsto \langle \Psi_\alpha | U_\alpha(\theta(\mathbf{z})) \rangle_2$ are holomorphic in the region $\mathcal{A}_{\nu, \tilde{\nu}} \odot \mathcal{A}_U^n := \{(\alpha, \alpha) \in \mathcal{A}_{\nu, \tilde{\nu}} \times \mathcal{A}_U^n \mid 2Q < \text{Re}(s + \alpha)\}$. Let us consider the subsets

$$\mathcal{S} := \{(\alpha, \alpha) \mid \forall i \alpha_i \in \mathbb{R} \text{ and } \alpha_i < Q, s - Q > 0, \alpha \in Q + i(0, \infty)\}, \quad \mathcal{S}_* = \mathcal{S} \cap \{(\alpha, \alpha) \mid \alpha \notin \mathcal{D}_0 \cup \mathcal{D}_\ell\}$$

and

$$\mathcal{R}_{\nu, \tilde{\nu}} := \{(\alpha, \alpha) \mid \forall i \alpha_i \in \mathbb{R} \text{ and } \alpha_i < Q, \alpha + s - 2Q > 0, \alpha \in \mathbb{R}, \alpha < (Q - \frac{2\lambda_\ell}{\gamma} - \frac{\gamma}{4}) \wedge (Q - \gamma)\}.$$

They are both in the same connected component of $\mathcal{A}_{\nu, \tilde{\nu}} \odot \mathcal{A}_U^n$. Furthermore, \mathcal{S}_* is obviously non-empty whereas the condition $(n-2)Q > -(Q - \frac{2\lambda_\ell}{\gamma} - \frac{\gamma}{4}) \wedge (Q - \gamma)$ ensures that $\mathcal{R}_{\nu, \tilde{\nu}}$ is non-empty, which we assume from now on.

Now we exploit the Ward identities, valid on $\mathcal{R}_{\nu, \tilde{\nu}}$. Let us consider a smooth compactly supported function φ on $\theta\mathcal{Z}$. The mapping

$$(\alpha, \alpha) \in \mathcal{A}_{\nu, \tilde{\nu}} \circ \mathcal{A}_U^n \mapsto \int \langle \Psi_{\alpha, \nu, \tilde{\nu}} | U_{\alpha}(\theta(\mathbf{z})) \rangle_2 \bar{\varphi}(\mathbf{z}) \, d\mathbf{z}$$

is thus analytic. Furthermore on $\mathcal{R}_{\nu, \tilde{\nu}}$ and by Proposition 7.11, it coincides with the mapping

$$\begin{aligned} (\alpha, \alpha) \in \mathcal{A}_{\nu, \tilde{\nu}} \circ \mathcal{A}_U^n &\mapsto \int \langle V_{\alpha}(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_{\gamma, \mu} \overline{\tilde{\mathbf{D}}_{\tilde{\nu}}^* \mathbf{D}_{\nu}^* \left(\varphi(\mathbf{z}) \prod_{i=1}^n |z_i|^{4\Delta_{\alpha_i}} \right)} \, d\mathbf{z} \\ &= \int \left(\prod_{i=1}^n |z_i|^{-4\Delta_{\alpha_i}} \right) \langle \Psi_{\alpha} | U_{\alpha}(\theta(\mathbf{z})) \rangle_2 \overline{\tilde{\mathbf{D}}_{\tilde{\nu}}^* \mathbf{D}_{\nu}^* \left(\varphi(\mathbf{z}) \prod_{i=1}^n |z_i|^{4\Delta_{\alpha_i}} \right)} \, d\mathbf{z} \end{aligned}$$

where we have introduced the (adjoint) operator \mathbf{D}_{ν}^* by

$$\int_{\mathbb{C}^n} \mathbf{D}_{\nu} f(\mathbf{z}) \bar{\varphi}(z) \, d\mathbf{z} = \int_{\mathbb{C}^n} f(\mathbf{z}) \overline{\mathbf{D}_{\nu}^* \varphi}(z) \, d\mathbf{z}$$

for all functions f in the domain of \mathbf{D}_{ν} and all smooth compactly supported functions φ in \mathbb{C}^n (and similarly for $\tilde{\mathbf{D}}_{\tilde{\nu}}$). Therefore both mappings are analytic and coincide on $\mathcal{R}_{\nu, \tilde{\nu}}$, thus on the connected component of $\mathcal{A}_{\nu, \tilde{\nu}} \circ \mathcal{A}_U$ containing $\mathcal{R}_{\nu, \tilde{\nu}}$, therefore on \mathcal{S}_* and finally on \mathcal{S} by continuity. Notice that, on \mathcal{S} , we can take all the α_i 's equal to 0 but the first two of them provided they satisfy $\alpha_1 + \alpha_2 > Q$. This fact being valid for all test function φ , we deduce that the relation

$$\begin{aligned} &\langle \Psi_{Q+iP, \nu, \tilde{\nu}} | U(V_{\alpha_1}(\theta(z_1)) V_{\alpha_2}(\theta(z_2))) \rangle_2 \\ &= \frac{1}{2} C_{\gamma, \mu}^{\text{DOZZ}}(Q+iP, \alpha_1, \alpha_2) |z_1|^{4\Delta_{\alpha_1}} |z_2|^{4\Delta_{\alpha_2}} \\ &\quad \times \mathbf{D}_{\nu} \tilde{\mathbf{D}}_{\tilde{\nu}} \left(|z_1|^{2\Delta_{\alpha_2} - 2\Delta_{Q+iP} - 2\Delta_{\alpha_1}} |z_1 - z_2|^{2\Delta_{Q+iP} - 2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} |z_2|^{2\Delta_{\alpha_1} - 2\Delta_{Q+iP} - 2\Delta_{\alpha_2}} \right) \end{aligned}$$

holds for almost every z_1, z_2 and $\alpha_1, \alpha_2 < Q$ such that $\alpha_1 + \alpha_2 > Q$, and thus for every $z_1, z_2 \in \theta\mathcal{Z}$ as both sides are continuous in these variables. From this relation and after some elementary algebra to compute the last term, sending $z_2 \rightarrow \infty$, we end up with the claimed relation. \square

8. PROOF OF THEOREM 1.1

8.1. Definition of Conformal Blocks. Before proving Theorem 1.1, we give the definition of the conformal blocks (1.19) which is based on material introduced in subsections 4.4 and 7.5. The conformal blocks are defined as the formal power series

$$(8.1) \quad \mathcal{F}_{Q+iP}(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4} z) = \sum_{n=0}^{\infty} \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4}) z^n$$

with

$$(8.2) \quad \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4}) := \sum_{|\nu|, |\nu'|=n} v(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{Q+iP}, \nu) F_{Q+iP}^{-1}(\nu, \nu') v(\Delta_{\alpha_4}, \Delta_{\alpha_3}, \Delta_{Q+iP}, \nu').$$

where the matrix $(F_{Q+iP}^{-1}(\nu, \nu'))_{|\nu|, |\nu'|=n}$ is the inverse of the scalar product matrix (4.27) and the function v is explicitly defined by expression (7.36). The definition of β_n is not explicit because there is no known explicit formula for the inverse matrix F_{Q+iP}^{-1} and the convergence of (8.1) is an open problem. Below we'll prove the series converges in the unit disc a.e. in P .

8.2. Proof of Theorem 1.1. We start with a Lemma on the spectral decomposition:

Lemma 8.1. *Let $u_1, u_2 \in e^{\delta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ with $\delta > 0$. Then*

$$\langle u_1 | u_2 \rangle_2 = \frac{1}{2\pi} \lim_{N, L \rightarrow \infty} \sum_{|\nu|=|\nu'| \leq N} \sum_{|\tilde{\nu}'|=|\tilde{\nu}| \leq N} \int_0^L \langle u_1 | \Psi_{Q+iP, \nu, \tilde{\nu}} \rangle_2 \langle \Psi_{Q+iP, \nu', \tilde{\nu}'} | u_2 \rangle_2 F_{Q+iP}^{-1}(\nu, \nu') F_{Q+iP}^{-1}(\tilde{\nu}, \tilde{\nu}') \, dP.$$

Proof. We write the spectral representation (6.75) as

$$(8.3) \quad \langle u_1 | u_2 \rangle_2 = \lim_{N, L \rightarrow \infty} \frac{1}{2\pi} \sum_{|\mathbf{k}|+|\mathbf{l}| \leq N} \int_0^L \langle u_1 | \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} \rangle_2 \langle \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} | u_2 \rangle_2 dP.$$

Let $F_{Q+iP}^{-1/2}(\nu, \nu')$ be the square root of the positive definite matrix $(F_{Q+iP}^{-1}(\nu, \nu'))_{|\nu|=|\nu'|=N}$ and set for $|\nu| = N$, $|\tilde{\nu}| = N'$

$$H_{Q+iP, \nu, \tilde{\nu}} := \sum_{|\nu_1|=N, |\nu_2|=N'} F_{Q+iP}^{-1/2}(\nu, \nu_1) F_{Q+iP}^{-1/2}(\tilde{\nu}, \nu_2) \Psi_{Q+iP, \nu_1, \nu_2}.$$

where $\Psi_{Q+iP, \nu_1, \nu_2}$ are defined in (7.1). We get the identity

$$\sum_{|\mathbf{k}|+|\mathbf{l}|=N} \langle u_1 | \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} \rangle_2 \langle \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} | u_2 \rangle_2 = \sum_{|\nu|+|\tilde{\nu}|=N} \langle u_1 | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2 \langle H_{Q+iP, \nu, \tilde{\nu}} | u_2 \rangle_2.$$

Hence we have

$$(8.4) \quad \sum_{|\nu|+|\tilde{\nu}| \leq N} \int_0^L \langle u_1 | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2 \langle H_{Q+iP, \nu, \tilde{\nu}} | u_2 \rangle_2 dP = \sum_{|\mathbf{k}|+|\mathbf{l}| \leq N} \int_0^L \langle u_1 | \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} \rangle_2 \langle \Psi_{Q+iP, \mathbf{k}, \mathbf{l}} | u_2 \rangle_2 dP.$$

On the other hand

$$(8.5) \quad \begin{aligned} & \sum_{|\nu|, |\tilde{\nu}| \leq N} \int_0^L \langle u_1 | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2 \langle H_{Q+iP, \nu, \tilde{\nu}} | u_2 \rangle_2 dP \\ &= \sum_{|\nu_1|=|\nu_3| \leq N} \sum_{|\nu_2|=|\nu_4| \leq N} \int_0^L \langle u_1 | \Psi_{Q+iP, \nu_1, \nu_2} \rangle_2 \langle \Psi_{Q+iP, \nu_3, \nu_4} | u_2 \rangle_2 F_{Q+iP}^{-1}(\nu_1, \nu_3) F_{Q+iP}^{-1}(\nu_2, \nu_4) dP. \end{aligned}$$

Let $u_1 = u_2 = u$. Then

$$(8.6) \quad \int_0^L \sum_{|\nu|+|\tilde{\nu}| \leq N} |\langle u | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2|^2 dP \leq \int_0^L \sum_{\substack{|\nu| \leq N, \\ |\tilde{\nu}| \leq N}} |\langle u | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2|^2 dP \leq \int_0^L \sum_{|\nu|+|\tilde{\nu}| \leq 2N} |\langle u | H_{Q+iP, \nu, \tilde{\nu}} \rangle_2|^2 dP.$$

By (8.4) and (8.3) the limits as $L, N \rightarrow \infty$ of all the three sums in (8.6) are $2\pi \langle u | u \rangle_2$. Hence by polarisation the limits as $L, N \rightarrow \infty$ of the left hand sides of (8.5) and (8.4) exist and equal $2\pi \langle u_1 | u_2 \rangle_2$. This proves the claim. \square

Let $|z| < 1$, $t \in (0, 1)$ and $|u| > 1$ with $u \neq t^{-1}$. The 4-point correlation function is given as a scalar product

$$(8.7) \quad \langle V_{\alpha_1}(0) V_{\alpha_2}(z) V_{\alpha_3}(t^{-1}) V_{\alpha_4}(u) \rangle_{\gamma, \mu} = t^{4\Delta_{\alpha_3}} |u|^{-4\Delta_{\alpha_4}} \left\langle U_{\alpha_1, \alpha_2}(0, z) | U_{\alpha_4, \alpha_3}(\bar{u}^{-1}, t) \right\rangle_2$$

where U_{α_1, α_2} is defined in (7.30). As in (1.14) we have

$$\langle V_{\alpha_1}(0) V_{\alpha_2}(z) V_{\alpha_3}(t^{-1}) V_{\alpha_4}(\infty) \rangle_{\gamma, \mu} = t^{4\Delta_{\alpha_3}} \left\langle U_{\alpha_1, \alpha_2}(0, z) | U_{\alpha_4, \alpha_3}(0, t) \right\rangle_2.$$

Recall from Lemma 7.9 that for $|z| < 1$ the vector $U_{\alpha_1, \alpha_2}(0, z) \in e^{\delta c} L^2(\mathbb{R} \times \Omega_{\mathbb{T}})$ for some $\delta > 0$ if $\alpha_1 + \alpha_2 > Q$, and the same holds for $U_{\alpha_4, \alpha_3}(0, t)$ if $\alpha_3 + \alpha_4 > Q$. Using Lemma 8.1 we get

$$(8.8) \quad \left\langle U_{\alpha_1, \alpha_2}(0, z) | U_{\alpha_4, \alpha_3}(0, t) \right\rangle_2 = \frac{1}{2\pi} \lim_{N, L \rightarrow \infty} I_{N, L}(\alpha, z, t)$$

where

$$(8.9) \quad I_{N, L}(\alpha, z, t) = \sum_{\substack{|\nu'|=|\nu| \leq N, \\ |\tilde{\nu}'|=|\tilde{\nu}| \leq N}} \int_0^L \langle U_{\alpha_1, \alpha_2}(0, z) | \Psi_{Q+iP, \nu, \tilde{\nu}} \rangle_2 \langle \Psi_{Q+iP, \nu', \tilde{\nu}'} | U_{\alpha_4, \alpha_3}(0, t) \rangle_2 F_{Q+iP}^{-1}(\nu, \nu') F_{Q+iP}^{-1}(\tilde{\nu}, \tilde{\nu}') dP.$$

Using Proposition 7.12 for the scalar products we can write this as

$$\begin{aligned}
I_{N,L}(\alpha, z, t) &= \frac{1}{4} t^{4\Delta_{\alpha_3}} t^{-2\Delta_{\alpha_3} - 2\Delta_{\alpha_4}} |z|^{-2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} \int_0^L \overline{C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q + iP)} C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_4, \alpha_3, Q + iP) \\
&\quad |tz|^{2\Delta_{Q+iP}} \sum_{|\nu'|=|\nu| \leq N} z^{|\nu|} t^{|\nu'|} v(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{Q+iP}, \nu) F_{Q+iP}^{-1}(\nu, \nu') v(\Delta_{\alpha_4}, \Delta_{\alpha_3}, \Delta_{Q+iP}, \nu') \\
&\quad \sum_{|\tilde{\nu}'|=|\tilde{\nu}| \leq N} \tilde{z}^{|\tilde{\nu}|} t^{|\tilde{\nu}'|} v(\Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{Q+iP}, \tilde{\nu}) F_{Q+iP}^{-1}(\tilde{\nu}, \tilde{\nu}') v(\Delta_{\alpha_4}, \Delta_{\alpha_3}, \Delta_{Q+iP}, \tilde{\nu}') dP \\
&= \frac{1}{4} t^{2\Delta_{\alpha_3} - 2\Delta_{\alpha_4}} |z|^{-2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} \int_0^L \overline{C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q + iP)} C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_4, \alpha_3, Q + iP) |tz|^{2\Delta_{Q+iP}} \\
&\quad \left| \sum_{n=1}^N \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4})(zt)^n \right|^2 dP
\end{aligned}$$

where we noted that the v factors are real. The coefficients β_n are given by (8.1) so formally the bootstrap formula follows now by taking $N, L \rightarrow \infty$. Yet, rigorously, there is still a gap to bridge since we do not know the convergence of the series (8.1). We adapt here a Cauchy-Schwarz type argument¹⁵ to establish the convergence a.s. in P . For this, we take first $\alpha_3 = \alpha_2$ and $\alpha_1 = \alpha_4$, with $\alpha_1 + \alpha_2 > Q$. Then

$$\begin{aligned}
I_{N,L}(\alpha_1, \alpha_2, \alpha_2, \alpha_1, z, t) &= \frac{1}{4} t^{4\Delta_{\alpha_2}} |tz|^{-2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} \\
&\quad \int_0^L |C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)|^2 |tz|^{2\Delta_{Q+iP}} \left| \sum_{n=0}^N \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_2}, \Delta_{\alpha_1})(zt)^n \right|^2 dP
\end{aligned}$$

If $z \in (0, 1)$ this expression is increasing in the variables N, L and zt since $\beta_n \geq 0$; moreover, it converges as N, L goes to infinity. This implies that the series

$$\sum_{n=0}^{\infty} z^n \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_2}, \Delta_{\alpha_1})$$

is absolutely converging for $|z| < 1$ for almost all $P \geq 0$ and

$$\begin{aligned}
&\langle V_{\alpha_1}(0) V_{\alpha_2}(z) V_{\alpha_2}(t^{-1}) V_{\alpha_1}(\infty) \rangle_{\gamma,\mu} = \\
&\frac{1}{8\pi} t^{4\Delta_{\alpha_2}} |tz|^{-2\Delta_{\alpha_1} - 2\Delta_{\alpha_2}} \int_0^{\infty} |C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, Q - iP)|^2 |tz|^{2\Delta_{Q+iP}} \left| \sum_{n=0}^{\infty} \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_2}, \Delta_{\alpha_1})(zt)^n \right|^2 dP.
\end{aligned}$$

This leads to the formula (1.18) for the case $\alpha_3 = \alpha_2$ and $\alpha_1 = \alpha_4$ by using the identity

$$\langle V_{\alpha_1}(0) V_{\alpha_2}(tz) V_{\alpha_2}(1) V_{\alpha_1}(\infty) \rangle_{\gamma,\mu} = t^{-4\Delta_{\alpha_2}} \langle V_{\alpha_1}(0) V_{\alpha_2}(z) V_{\alpha_2}(t^{-1}) V_{\alpha_1}(\infty) \rangle_{\gamma,\mu}.$$

For the general case we use Cauchy-Schwartz

$$\begin{aligned}
|\beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_3}, \Delta_{\alpha_4})| &\leq \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_2}, \Delta_{\alpha_1})^{\frac{1}{2}} \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_4}, \Delta_{\alpha_3}, \Delta_{\alpha_3}, \Delta_{\alpha_4})^{\frac{1}{2}} \\
&\leq \frac{1}{2} (\beta_n(\Delta_{Q+iP}, \Delta_{\alpha_1}, \Delta_{\alpha_2}, \Delta_{\alpha_2}, \Delta_{\alpha_1}) + \beta_n(\Delta_{Q+iP}, \Delta_{\alpha_4}, \Delta_{\alpha_3}, \Delta_{\alpha_3}, \Delta_{\alpha_4})).
\end{aligned}$$

so that the general case follows from the case $\alpha_1 = \alpha_4$ and $\alpha_2 = \alpha_3$. \square

9. PROOF OF PROPOSITION 7.11

9.1. Preliminary remarks. Before proceeding to computations, we stress that the reader should keep in mind that the SET field $T(u)$ is not a proper random field. In particular the expectation in (7.10) is a notation for the object constructed in the limit as $\epsilon \rightarrow 0$ and $t \rightarrow \infty$. In LCFT the construction of the correlation functions of the SET is subtle. This was done in [KRV19] only for one or two SET insertions. However the situation is here much simpler and we will not have to rely on [KRV19]. The reason is that we need to deal with correlation functions with a regularized LCFT expectation $\langle \cdot \rangle_t$ where we have replaced

¹⁵Communicated to us by S. Rychkov.

$M_\gamma(\mathbb{C})$ by $M_\gamma(\mathbb{C} \setminus \mathbb{D}_t)$ in (7.32), and all the SET insertions that we will consider are located in \mathbb{D}_t which as we will see makes them much more regular than in the full LCFT.

The regularized SET field $T_\epsilon(u)$ is a proper random field and its correlation functions in the presence of the vertex operators are defined as limits of the corresponding ones with regularized vertex operators

$$(9.1) \quad \langle T_\epsilon(\mathbf{u}) \bar{T}_{\epsilon'}(\mathbf{v}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = \lim_{\epsilon'' \rightarrow 0} \langle T_\epsilon(\mathbf{u}) \bar{T}_{\epsilon'}(\mathbf{v}) V_{\alpha, \epsilon''}(0) \prod_{i=1}^n V_{\alpha_i, \epsilon''}(z_i) \rangle_t.$$

The existence of this limit follows from the representation of the expectation on the RHS as a GFF expectation of an explicit function of a GMC integral [DKRV16]. And in particular the limit is independent of the regularization procedure used for the vertex operators. For simplicity we will use in this section the following:

$$(9.2) \quad V_{\alpha, \epsilon}(x) = \epsilon^{\frac{\alpha^2}{2}} e^{\alpha \phi_\epsilon(x)} = |x|_+^{-4\Delta_\alpha} e^{\alpha c} e^{\alpha X_\epsilon(x) - \frac{\alpha^2}{2} \mathbb{E} X_\epsilon(x)^2} (1 + \mathcal{O}(\epsilon))$$

where X_ϵ is the same regularization as in T_ϵ . The $\mathcal{O}(\epsilon)$ will drop out from all terms in the $\epsilon \rightarrow 0$ limit and will not be displayed below.

The proof of Proposition 7.11 consists of using Gaussian integration by parts to the the $T_\epsilon(\mathbf{u})$ and $\bar{T}_{\epsilon'}(\mathbf{v})$ factors in (9.1) to which we now turn.

9.2. Gaussian integration by parts. (9.1) is analysed using Gaussian integration by parts. For a centered Gaussian vector (X, Y_1, \dots, Y_N) and a smooth function f on \mathbb{R}^N , the Gaussian integration by parts formula is

$$\mathbb{E}[X f(Y_1, \dots, Y_N)] = \sum_{k=1}^N \mathbb{E}[X Y_k] \mathbb{E}[\partial_k f(Y_1, \dots, Y_N)].$$

Applied to the LCFT this leads to the following formula. Let $\phi = c + X - 2Q \log |z|_+$ be the Liouville field and F a smooth function on \mathbb{R}^N . Define for $u, v \in \mathbb{C}$

$$(9.3) \quad C(u, v) = -\frac{1}{2} \frac{1}{u-v}, \quad C_{\epsilon, \epsilon'}(u, v) = \int \rho_\epsilon(u-u') \rho_{\epsilon'}(v-v') C(u', v') du' dv'$$

with $(\rho_\epsilon)_\epsilon$ a mollifying family of the type $\rho_\epsilon(\cdot) = \epsilon^{-2} \rho(|\cdot|/\epsilon)$. Then for $z, x_1, \dots, x_N \in \mathbb{C}$

$$(9.4) \quad \langle \partial_z \phi_\epsilon(z) F(\phi_{\epsilon'}(x_1), \dots, \phi_{\epsilon'}(x_N)) \rangle_t = \sum_{k=1}^N C_{\epsilon, \epsilon'}(z, x_k) \langle \partial_k F(\phi_{\epsilon'}(x_1), \dots, \phi_{\epsilon'}(x_N)) \rangle_t \\ - \mu \gamma \int_{\mathbb{C}_t} C_{\epsilon, 0}(z, x) \langle V_\gamma(x) F(\phi_{\epsilon'}(x_1), \dots, \phi_{\epsilon'}(x_N)) \rangle_t dx$$

where F in the applications below is such that all the terms here are well defined. Note that $\partial_u G(u, v) = C(u, v) + \frac{1}{2} u^{-1} 1_{|u|>1}$. The virtue of the Liouville field is that the annoying metric dependent terms $u^{-1} 1_{|u|>1}$ drop out from the formulae. This fact is nontrivial and it was proven in [KRV19, Subsection 3.2], for the case $t = \infty$ with F corresponding to product of vertex operators. The proof goes the same way to produce (9.4) with a finite t .

The first application of this formula is a direct proof of the existence of the $\epsilon, \epsilon' \rightarrow 0$ limit of (9.1) which will be useful also later in the proof. We have

Proposition 9.1. *The functions (9.1) converge uniformly on compact subsets of $(\mathbf{u}, \mathbf{v}, \mathbf{z}) \in \mathbb{D}_t \times \mathbb{D}_t \times \theta \mathcal{Z}$*

$$(9.5) \quad \lim_{\epsilon, \epsilon' \rightarrow 0} \langle T_\epsilon(\mathbf{u}) \bar{T}_{\epsilon'}(\mathbf{v}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t := \langle T(\mathbf{u}) \bar{T}(\mathbf{v}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$$

where the limit is analytic in $\mathbf{u} \in \mathcal{O}_{e^{-t}}$ and anti-analytic in $\mathbf{v} \in \mathcal{O}_{e^{-t}}$.

Proof. We consider for simplicity only the case $\tilde{v} = 0$. The LHS is defined as the limit $\epsilon'' \rightarrow 0$ in (9.1) but we will for clarity work directly with $\epsilon'' = 0$ as it will be clear below that this limit trivially exists. Indeed, the functions $C_{\epsilon, \epsilon'}(u, v)$ are smooth for $\epsilon, \epsilon' > 0$ and they converge together with their derivatives uniformly on $|u| < e^{-s}$, $|v| \geq e^{-t}$ for all $s > t$ to the derivatives of $C(u, v)$.

We will now apply (9.4) to all the ϕ_ϵ factors in the SET tensors in (9.5) one after the other. To make this systematic let us introduce the notation

$$(\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3) := (\partial_z \phi_\epsilon, \partial_z^2 \phi_\epsilon, (\partial_z \phi_\epsilon)^2 - \mathbb{E}(\partial_z \phi_\epsilon)^2).$$

Applying the integration by parts formula to $\partial_z \phi_\epsilon(u_k)$ (or to $\partial_z^2 \phi_\epsilon(u_k)$ if $i_k = 2$ below) we obtain

$$\begin{aligned} \langle \prod_{j=1}^k \mathcal{O}_{i_j}(u_j) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t &= \sum_{l=1}^{k-1} \langle \{\mathcal{O}_{i_k}(u_k) \mathcal{O}_{i_l}(u_l)\} \prod_{j \neq k, l} \mathcal{O}_{i_j}(u_j) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad + \sum_{l=0}^n \alpha_l \langle \{\mathcal{O}_{i_k}(u_k) \phi(z_l)\} \prod_{j \neq k} \mathcal{O}_{i_j}(u_j) V_\alpha(0) \prod_{i \neq l} V_{\alpha_i}(z_i) \rangle_t \\ &\quad - \mu \gamma \int_{\mathbb{C}_t} \langle \{\mathcal{O}_{i_k}(u_k) \phi(x)\} \prod_{j \neq k} \mathcal{O}_{i_j}(u_j) V_\alpha(0) V_\gamma(x) \prod_{i \neq l} V_{\alpha_i}(z_i) \rangle_t dx \end{aligned}$$

where $\alpha_0 = \alpha$ and $z_0 = 0$ and we introduced the following notation

$$\begin{aligned} \{\mathcal{O}_1(u) \mathcal{O}_1(v)\} &:= \partial_v C_{\epsilon, \epsilon}(u, v) & \{\mathcal{O}_2(u) \mathcal{O}_2(v)\} &:= \partial_u \partial_v^2 C_{\epsilon, \epsilon}(u, v) \\ \{\mathcal{O}_1(u) \mathcal{O}_3(v)\} &:= 2\mathcal{O}_1(v) \partial_v C_{\epsilon, \epsilon}(u, v) & \{\mathcal{O}_1(u) \mathcal{O}_2(v)\} &:= \partial_v^2 C_{\epsilon, \epsilon}(u, v) \\ \{\mathcal{O}_3(u) \mathcal{O}_1(v)\} &:= \mathcal{O}_1(u) \partial_v C_{\epsilon, \epsilon}(u, v) & \{\mathcal{O}_2(u) \mathcal{O}_1(v)\} &:= \partial_u \partial_v C_{\epsilon, \epsilon}(u, v) \\ \{\mathcal{O}_3(u) \mathcal{O}_3(v)\} &:= 2\mathcal{O}_1(u) \mathcal{O}_1(v) \partial_v C_{\epsilon, \epsilon}(u, v) & \{\mathcal{O}_2(u) \mathcal{O}_3(v)\} &:= 2\mathcal{O}_1(v) \partial_u \partial_v C_{\epsilon, \epsilon}(u, v) \\ \{\mathcal{O}_3(u) \mathcal{O}_2(v)\} &:= \mathcal{O}_1(u) \partial_v^2 C_{\epsilon, \epsilon}(u, v) \end{aligned}$$

etc. Similarly

$$\{\mathcal{O}_1(u) \phi(x)\} := C_{\epsilon, 0}(u, x), \quad \{\mathcal{O}_2(u) \phi(x)\} := \partial_u C_{\epsilon, 0}(u, x), \quad \{\mathcal{O}_3(u) \phi(x)\} := \mathcal{O}_1(u) C_{\epsilon, 0}(u, x).$$

Since $T_\epsilon = Q\mathcal{O}_2 - \mathcal{O}_3$ we can iterate the integration by parts formula to obtain an expression for $\langle T_\epsilon(\mathbf{u}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$ as a sum of terms of the form

$$(9.6) \quad C \prod_{\alpha, \beta} \partial^{i\alpha\beta} C_{\epsilon, \epsilon}(u_\alpha, u_\beta) \prod_{\alpha, l} \partial^{j\alpha, l} C_{\epsilon, 0}(u_\alpha, z_l) \int_{\mathbb{C}_t^m} \prod_{\alpha, k} \partial^{l\alpha, k} C_{\epsilon, 0}(u_\alpha, x_k) \langle V_\alpha(0) \prod_{j=1}^m V_\gamma(x_j) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx$$

where the products run through some subsets of the index values. We have for all compact $K \subset \mathbb{D}_t$

$$\sup_{u \in K, x \in \mathbb{C}_t} \sup_{\epsilon} |\partial_u^l \partial_u^l C_{\epsilon, 0}(u, x)| < \infty.$$

Hence the expression (9.6) is bounded together with all its derivatives in \mathbf{u} uniformly in ϵ by

$$(9.7) \quad C \int_{\mathbb{C}_t^m} \langle V_\alpha(0) \prod_{j=1}^m V_\gamma(x_j) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx.$$

which by the lemma below is finite. The limit of (9.6) and all its derivatives exist by dominated convergence. Clearly the ∂_u derivatives vanish so the limit is analytic in \mathbf{u} . \square

We need the following KPZ identity for the a priori bound (9.7) (recall (7.31))

Lemma 9.2. *The functions $\mathbf{x} \in \mathbb{C}_t^n \rightarrow \langle V_\alpha(0) \prod_{j=1}^m V_\gamma(x_j) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$ are integrable and*

$$(9.8) \quad \int_{\mathbb{C}_t^m} \langle V_\alpha(0) \prod_{j=1}^m V_\gamma(x_j) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx = C \int_{\mathbb{C}_t^m} \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx$$

where $C = (\mu\gamma)^{-m} \prod_{l=0}^{m-1} (\alpha + s + \gamma l - 2Q)$.

Proof. See [KRV19, Lemma 3.3]. Briefly,

$$\langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = \mu^{\frac{2Q-s-\alpha}{\gamma}} \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \Big|_{\mu=1}$$

and the LHS of (9.8) equals $(-\partial_\mu)^m \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$. \square

The representation (9.6) will not be useful for a direct proof of the Ward identity due to the V_γ insertions. We will rather use the integration by parts inductively, first to $T_\epsilon(u_k)$ which corresponds to the largest contour in the contour integrals in expression (7.10), and by showing then that at each step the V_γ insertions give rise to the derivatives in the Proposition 7.11. We give now the inductive step to prove this claim, stated for simplicity for the case $\tilde{\nu} = 0$. For this, we introduce the (adjoint) operator \mathbf{D}_n^* defined (by duality) by

$$\int_{\mathbb{C}^n} \mathbf{D}_n f(\mathbf{z}) \bar{\varphi}(z) d\mathbf{z} = \int_{\mathbb{C}^n} f(\mathbf{z}) \overline{\mathbf{D}_n^* \varphi}(z) d\mathbf{z}$$

for all functions f in the domain of \mathbf{D}_n and all smooth compactly supported functions φ in \mathbb{C}^n . Then

Proposition 9.3. *Let $\varphi \in C_0^\infty(\theta\mathcal{Z})$ be a smooth compactly supported function in $\theta\mathcal{Z} \subset \mathbb{C}^n$ and define*

$$\mathcal{T}_t(\nu, \varphi) := \int \left(\frac{1}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} \langle T(\mathbf{u}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z}.$$

Then for $\nu^{(k)} = (\nu_1, \dots, \nu_{k-1})$

$$\mathcal{T}_t(\nu, \varphi) = \mathcal{T}_t(\nu^{(k)}, \mathbf{D}_{\nu_k}^* \varphi) + \mathcal{B}_t(\nu, \varphi)$$

where

$$(9.9) \quad |\mathcal{B}_t(\nu, \varphi)| \leq C e^{(\alpha+|\nu|-2)t} \int \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t |\varphi(\mathbf{z})| d\mathbf{z}.$$

Proof of Proposition 7.11. Iterating Proposition 9.3 we get

$$\mathcal{T}_t(\nu, \varphi) = \int \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \overline{\mathbf{D}_\nu^* \varphi}(\mathbf{z}) d\mathbf{z} + \mathcal{B}_t(\varphi)$$

where $\mathcal{B}_t(\varphi)$ satisfies

$$\mathcal{B}_t(\varphi) \leq C e^{(\alpha+|\nu|-2)t} \sum_{\ell=1}^k \int \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t |\mathbf{D}_{\nu_{\ell+1}}^* \cdots \mathbf{D}_{\nu_k}^* \varphi(\mathbf{z})| d\mathbf{z}$$

where by convention $\mathbf{D}_{\nu_{\ell+1}}^* \cdots \mathbf{D}_{\nu_k}^* \varphi = \varphi$ if $\ell = k$. The functions $\mathbf{z} \rightarrow \langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$ are continuous on $\theta\mathcal{Z}$ and converge uniformly as $t \rightarrow \infty$ on compact subsets of $\theta\mathcal{Z}$ to the function $\langle V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle$. Hence $\mathcal{T}_t(\nu, \cdot)$ converges in the Frechet topology of $\mathcal{D}'(\theta\mathcal{Z})$ to the required limit since $\mathcal{B}_t(\varphi)$ goes to 0 as t goes to infinity (recall that $\alpha + |\nu| - 2 < 0$). \square

9.3. Proof of Proposition 9.3. We start the proof of Proposition 9.3 by applying Gaussian integration by parts formula twice to the $T(u_k)$. This produces plenty of terms which we group in four contributions:

$$(9.10) \quad \langle T(\mathbf{u}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = R(\mathbf{u}, \mathbf{z}) + M(\mathbf{u}, \mathbf{z}) + N(\mathbf{u}, \mathbf{z}) + D(\mathbf{u}, \mathbf{z}).$$

In $R(\mathbf{u})$ we group all the contractions with $C(u_k, u_l)$ between u_k and u_l , $l < k$ and u_k and 0. These terms do not contribute to the contour integral of the u_k variable since they give rise to integrals of the form

$$\oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} (u_k - v)^{-a} (u_k - w)^{-b} du_k$$

where $v, w \in \{0, u_1, \dots, u_{k-1}\}$ and $a + b \geq 2$. Since $|v|, |w| < e^{-t}\delta_k$ and $\nu_k > 0$, this integral vanishes. We conclude

$$(9.11) \quad \oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} R(\mathbf{u}, \mathbf{z}) du_k = 0.$$

For the benefit of the reader we display all the terms in $R(\mathbf{u}, \mathbf{z})$ in the Appendix C.

Let us now introduce the notations $\mathbf{u}^{(k)} := (u_1, \dots, u_{k-1})$ and $\mathbf{u}^{(k, \ell)} := (u_1, \dots, u_{\ell-1}, u_{\ell+1}, \dots, u_{k-1})$. The second contribution in (9.10) collects the contractions hitting only one V_{α_p} :

$$(9.12) \quad M(\mathbf{u}, \mathbf{z}) = \sum_{p=1}^n \left(\frac{Q_{\alpha_p}}{2} - \frac{\alpha_p^2}{4} \right) \frac{1}{(u_k - z_p)^2} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t.$$

We can then do the u_k integral explicitly to obtain

$$(9.13) \quad \int \left(\frac{1}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} \mathbf{M}(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z}$$

$$(9.14) \quad = \int \left(\sum_{p=1}^n \frac{\nu_k - 1}{z_p^{\nu_k}} \Delta_{\alpha_p} \right) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \bar{\varphi}(\mathbf{z}) d\mathbf{z}$$

Note that this term contains the constant part of the differential operator \mathbf{D}_{ν_k} .

The third contribution is given by terms where all contractions hit V_γ :

$$(9.15) \quad \begin{aligned} N(\mathbf{u}, \mathbf{z}) &= \left(\frac{\mu\gamma^2}{4} - \frac{\mu\gamma Q}{2} \right) \int_{\mathbb{C}_t} \frac{1}{(u_k - x)^2} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &= -\mu \int_{\mathbb{C}_t} \frac{1}{(u_k - x)^2} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx. \end{aligned}$$

Finally D gathers all the other terms

$$(9.16) \quad D(\mathbf{u}, \mathbf{z}) = \sum_{i=1}^9 T_i(\mathbf{u}, \mathbf{z})$$

with

$$(9.17) \quad \begin{aligned} T_1(\mathbf{u}, \mathbf{z}) &= - \sum_{\ell=1}^{k-1} \sum_{p=1}^n \frac{Q\alpha_p}{(u_k - u_\ell)^3 (u_k - z_p)} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ T_2(\mathbf{u}, \mathbf{z}) &= \sum_{\ell=1}^{k-1} \sum_{p=1}^n \frac{\alpha_p}{(u_k - u_\ell)^2 (u_k - z_p)} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ T_3(\mathbf{u}, \mathbf{z}) &= - \sum_{p=1}^n \frac{\alpha\alpha_p}{2} \frac{1}{(u_k - z_p) u_k} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ T_4(\mathbf{u}, \mathbf{z}) &= \frac{\mu\gamma}{2} \sum_{p=1}^n \alpha_p \int_{\mathbb{C}_t} \frac{1}{(u_k - z_p)(u_k - x)} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ T_5(\mathbf{u}, \mathbf{z}) &= - \sum_{p \neq p'=1}^n \frac{\alpha_p \alpha_{p'}}{4} \frac{1}{(u_k - z_p)(u_k - z_{p'})} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ T_6(\mathbf{u}, \mathbf{z}) &= \mu Q \gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(u_k - u_\ell)^3 (u_k - x)} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ T_7(\mathbf{u}, \mathbf{z}) &= -\mu\gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(u_k - u_\ell)^2 (u_k - x)} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ T_8(\mathbf{u}, \mathbf{z}) &= \frac{\mu\gamma\alpha}{2} \int_{\mathbb{C}_t} \frac{1}{u_k (u_k - x)} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ T_9(\mathbf{u}, \mathbf{z}) &= -\frac{\mu^2\gamma^2}{4} \int_{\mathbb{C}_t} \int_{\mathbb{C}_t} \frac{1}{(u_k - x)(u_k - x')} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) V_\gamma(x') \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx. \end{aligned}$$

We need to show that N and D will give rise (after contour integration) to the ∂_{z_i} -derivatives in the expression $\mathbf{D}_{\nu_k} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$. To show this we need to analyse N further.

Regularizing the vertex insertions (beside the V_γ insertion, we also regularize the V_{α_i} 's for later need) in $N(\mathbf{u}, \mathbf{z})$ given by (9.15), and performing an integration by parts (Green formula) in the x integral we get

$$\begin{aligned} N(\mathbf{u}, \mathbf{z}) &= -\mu \lim_{\epsilon \rightarrow 0} \int_{\mathbb{C}_t} \partial_x \frac{1}{u_k - x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_{\gamma, \epsilon}(x) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t dx \\ &= B_t(\mathbf{u}, \mathbf{z}) + \mu\gamma \lim_{\epsilon \rightarrow 0} \int_{\mathbb{C}_t} \frac{1}{u_k - x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \partial_x \phi_\epsilon(x) V_{\gamma, \epsilon}(x) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t dx \\ &=: B_t(\mathbf{u}, \mathbf{z}) + \tilde{N}(\mathbf{u}, \mathbf{z}) \end{aligned}$$

where the boundary term appearing in Green formula has $\epsilon \rightarrow 0$ limit given by

$$(9.18) \quad B_t(\mathbf{u}, \mathbf{z}) := i\mu \oint_{|x|=e^{-t}} \frac{1}{u_k - x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\bar{x}$$

and we used (1.8) to write

$$\partial_x V_{\gamma, \epsilon}(x) = \alpha \partial_x \phi_\epsilon(x) V_{\gamma, \epsilon}(x).$$

In $\tilde{N}(\mathbf{u}, \mathbf{z})$ we integrate by parts the $\partial_x \phi_\epsilon(x)$ and end up with

$$(9.19) \quad \begin{aligned} \tilde{N}(\mathbf{u}, \mathbf{z}) &= -\mu Q \gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(u_k - x)(x - u_\ell)^3} \langle T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad + \mu \gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(u_k - x)(x - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad - \frac{\mu \gamma \alpha}{2} \int_{\mathbb{C}_t} \frac{1}{(u_k - x)x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad + \frac{\mu^2 \gamma^2}{2} \int_{\mathbb{C}_t} \int_{\mathbb{C}_t} \frac{1}{(u_k - x)(x - x')} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) V_\gamma(x') \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad + \mu \gamma \lim_{\epsilon \rightarrow 0} \sum_{p=1}^n \int_{\mathbb{C}_t} \frac{\alpha_p}{u_k - x} C_{\epsilon, 0}(x, z_p) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t dx \\ &=: T'_6(\mathbf{u}, \mathbf{z}) + T'_7(\mathbf{u}, \mathbf{z}) + T'_8(\mathbf{u}, \mathbf{z}) + T'_9(\mathbf{u}) + T'_4(\mathbf{u}, \mathbf{z}) \end{aligned}$$

where again we took the $\epsilon \rightarrow 0$ limit in the terms where it was obvious. In particular this identity proves that the limit on the RHS, denoted by $T'_4(\mathbf{u}, \mathbf{z})$, exists. The numbering of these terms and the ones below will be used when comparing with (9.17).

Derivatives of correlation functions. We want to compare the expression (9.10) to derivatives of the function $\langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$. We need to treat separately the cases $\nu_k \geq 2$ and $\nu_k = 1$.

Case $\nu_k \geq 2$. We have

Lemma 9.4. *Let*

$$I_\epsilon(\mathbf{u}, \mathbf{z}) := \sum_{p=1}^n \frac{1}{u_k - z_p} \partial_{z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t.$$

Then $\lim_{\epsilon \rightarrow 0} I_\epsilon(\mathbf{u}, \mathbf{z}) := I(\mathbf{u}, \mathbf{z})$ exists and defines a continuous function in $\mathbf{z} \in \theta \mathcal{Z}$ satisfying

$$(9.20) \quad \int \left(\frac{1}{(2\pi i)^k} \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} I(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} = \mathcal{T}_t(\nu^{(k)}, \hat{\mathbf{D}}_{\nu_k} \varphi)$$

for all $\varphi \in C_0^\infty(\theta \mathcal{Z})$ with $\hat{\mathbf{D}}_n = \mathbf{D}_n^* - (n-1) \sum_i \Delta_{\alpha_i} z_i^{-n}$.

Proof. We have

$$I_\epsilon(\mathbf{u}, \mathbf{z}) = \sum_{p=1}^n \frac{\alpha_p}{u_k - z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \partial_{z_p} \phi_\epsilon(z_p) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t = K_\epsilon(\mathbf{u}, \mathbf{z}) + L_\epsilon(\mathbf{u}, \mathbf{z})$$

where we integrate by parts the $\partial_{z_p} \phi_\epsilon(z_p)$ and $K_\epsilon(\mathbf{u}, \mathbf{z})$ collects the terms with an obvious $\epsilon \rightarrow 0$ limit $K(\mathbf{u}, \mathbf{z})$:

$$\begin{aligned} K(\mathbf{u}, \mathbf{z}) &= - \sum_{p=1}^n \sum_{\ell=1}^{k-1} \frac{Q\alpha_p}{(u_k - z_p)(z_p - u_\ell)^3} \langle T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad + \sum_{p=1}^n \sum_{\ell=1}^{k-1} \frac{\alpha_p}{(u_k - z_p)(z_p - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad - \sum_{p=1}^n \frac{\alpha_p \alpha}{2} \frac{1}{(u_k - z_p) z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad - \sum_{p \neq p'=1}^n \frac{\alpha_p \alpha_{p'}}{2} \frac{1}{(u_k - z_p)(z_p - z_{p'})} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &=: D_1(\mathbf{u}, \mathbf{z}) + D_2(\mathbf{u}, \mathbf{z}) + D_3(\mathbf{u}, \mathbf{z}) + D_5(\mathbf{u}, \mathbf{z}), \end{aligned}$$

whereas

$$L_\epsilon(\mathbf{u}, \mathbf{z}) = -\mu\gamma \sum_{p=1}^n \alpha_p \int_{\mathbb{C}_t} \frac{1}{u_k - z_p} C_{\epsilon,0}(z_p, x) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t dx.$$

Since $C_{0,0}(z_p, x) = -\frac{1}{2} \frac{1}{z_p - x}$ and since it is not clear that $\frac{1}{z_p - x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$ is integrable the $\epsilon \rightarrow 0$ limit of L_ϵ is problematic¹⁶. However, we can compare it with the term T'_4 in (9.19). Writing

$$\frac{1}{u_k - z_p} = \frac{1}{u_k - x} + \frac{z_p - x}{(u_k - z_p)(u_k - x)},$$

we conclude that L_ϵ converges:

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} L_\epsilon(\mathbf{u}, \mathbf{z}) &= -\mu\gamma \lim_{\epsilon \rightarrow 0} \sum_{p=1}^n \alpha_p \left(\int_{\mathbb{C}_t} \frac{1}{u_k - x} C_{\epsilon,0}(z_p, x) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t dx \right. \\ &\quad \left. + \int_{\mathbb{C}_t} \frac{z_p - x}{(u_k - z_p)(u_k - x)} C_{\epsilon,0}(z_p, x) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \right) \\ &=: T'_4(\mathbf{u}, \mathbf{z}) + T_4(\mathbf{u}, \mathbf{z}). \end{aligned}$$

Indeed, setting $z = z_p - x$ the function

$$(z_p - x) C_{\epsilon,0}(z_p, x) = -\frac{1}{2} \int \rho_\epsilon(y) \frac{z}{z+y} dy = \frac{i}{2\epsilon^2} \int_{\mathbb{R}^+} \rho\left(\frac{r}{\epsilon}\right) \oint_{|u|=1} \frac{z}{z+ru} \frac{du}{u} r dr = -\pi \int_{\mathbb{R}^+} \rho(r) 1_{r < |z|/\epsilon} r dr$$

is uniformly bounded and converges almost everywhere to $-\frac{1}{2}$.

The same argument can be repeated to the smeared functions to show that (because convergence is uniform over compact subsets of $\theta\mathcal{Z}$)

$$\lim_{\epsilon \rightarrow 0} \sum_{p=1}^n \int \left(\frac{\alpha_p}{u_k - z_p} \partial_{z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} := \ell(\nu, \varphi)$$

exists. Then integrating ∂_{z_p} by parts and using that $\lim_{\epsilon \rightarrow 0} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t$ exists we conclude

$$\ell(\nu, \varphi) = - \int \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i, \epsilon}(z_i) \rangle_t \sum_{p=1}^n \partial_{z_p} \left(\frac{\alpha_p}{u_k - z_p} \varphi(\mathbf{z}) \right) d\mathbf{z}$$

which proves (9.20). \square

We have obtained the relation

$$(9.21) \quad N(\mathbf{u}, \mathbf{z}) - I(\mathbf{u}, \mathbf{z}) = B_t(\mathbf{u}, \mathbf{z}) + \sum_{i=6}^9 T_i(\mathbf{u}, \mathbf{z}) - \sum_{i=1}^3 D_i(\mathbf{u}, \mathbf{z}) - D_5(\mathbf{u}, \mathbf{z}) - T_4(\mathbf{u}, \mathbf{z}).$$

¹⁶Actually, this fact was shown in [KRV20] without the SET insertions and could be proven here as well but we will follow another route because the recursion to prove this extension is painful.

Let us consider the expression

$$\begin{aligned} K(\mathbf{u}, \mathbf{z}) &:= \langle T(\mathbf{u})V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t - I(\mathbf{u}, \mathbf{z}) - M(\mathbf{u}, \mathbf{z}) \\ &= R(\mathbf{u}, \mathbf{z}) + N(\mathbf{u}, \mathbf{z}) + D(\mathbf{u}, \mathbf{z}) - I(\mathbf{u}, \mathbf{z}). \end{aligned}$$

By (9.13) and (9.20) we have

$$(9.22) \quad \int \left(\oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} K(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \varphi(\mathbf{z}) d\mathbf{z} = \mathcal{T}(\nu, \varphi) - \mathcal{T}(\nu^{(k)}, \mathbf{D}_{\nu_k}^* \varphi).$$

On the other hand combining (9.10), (9.16) and (9.21) we obtain

$$(9.23) \quad K = R + \sum_{i=1}^3 (T_i - D_i) + T_5 - D_5 + \sum_{i=6}^9 (T_i + T'_i) + B_t.$$

Now some simple algebra, see Appendix C gives:

$$(9.24) \quad T'_9 = -T_9, \quad T_5 = D_5$$

$$(9.25) \quad \oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} (T'_i(\mathbf{u}, \mathbf{z}) + T'_i(\mathbf{u}, \mathbf{z})) du_k = 0 \quad i = 6, 7, 8,$$

$$(9.26) \quad \oint_{|u_k|=e^{-t}\epsilon_k} u_k^{1-\nu_k} (T_i(\mathbf{u}, \mathbf{z}) - D_i(\mathbf{u}, \mathbf{z})) du_k = 0 \quad i = 1, 2, 3.$$

Hence using these relations and (9.11) we conclude

$$\int \left(\oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} K(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} = \int \left(\oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} B_t(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} =: B_t(\nu, \varphi).$$

Thus to prove Proposition 7.11 for $\nu_k \geq 2$ we need to prove the bound (9.9) for $B_t(\nu, \varphi)$. Recalling (9.18) we get by residue theorem

$$\oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} B_t(\mathbf{u}, \mathbf{z}) d\mathbf{u} = -2\pi \oint_{|\mathbf{u}^{(k)}|=\delta_t^{(k)}} (\mathbf{u}^{(k)})^{1-\nu^{(k)}} \oint_{|x|=e^{-t}} x^{1-\nu_k} \langle T(\mathbf{u}^{(k)})V_\alpha(0)V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\bar{x} d\mathbf{u}.$$

By (9.6) (at $\epsilon = 0$ and an extra $V_{\alpha_{n+1}}(z_{n+1}) = V_\gamma(x)$) the expectation on the RHS is a sum of terms of the form

$$\int_{\mathbb{C}_t^m} I(\mathbf{u}^{(k)}, \mathbf{x}, \mathbf{z}, x) \langle V_\alpha(0)V_\gamma(x) \prod_{\ell=1}^m V_\gamma(x_\ell) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\mathbf{x}$$

where

$$I(\mathbf{u}^{(k)}, \mathbf{x}, \mathbf{z}, x) = C \prod_{\alpha, \beta} \frac{1}{(u_\alpha - u_\beta)^{k_{\alpha\beta}}} \prod_{\alpha, i} \frac{1}{(u_\alpha - z_i)^{l_{\alpha i}}} \prod_{\alpha, \ell} \frac{1}{(u_\alpha - x_\ell)^{m_{\alpha\ell}}} \prod_{\alpha} \frac{1}{(u_\alpha - x)^{n_\alpha}}$$

where $\sum k_{\alpha\beta} + \sum l_{\alpha k} + \sum m_{\alpha\ell} + \sum n_\alpha = 2(k-1)$. Performing the u -integrals in the order u_{k-1}, u_{k-2}, \dots by the residue theorem we get

$$\oint_{|\mathbf{u}^{(k)}|=\delta_t^{(k)}} (\mathbf{u}^{(k)})^{1-\nu^{(k)}} I(\mathbf{u}^{(k)}, \mathbf{x}, \mathbf{z}, x) d\mathbf{u} = \sum C(a, \mathbf{b}, \mathbf{c}) x^{-a} \prod_{\ell} x_\ell^{-b_\ell} \prod_i z_i^{-c_i}$$

with $a + \sum b_\ell + \sum c_i = |\nu^{(k-1)}|$. Since $|x_\ell| \geq e^{-t}$ and $|x| = e^{-t}$ we conclude

$$\left| \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} B_t(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right| \leq C e^{t(|\nu|-2)} \max_{m \leq 2(k-1)} \sup_{|x|=e^{-t}} \int_{\mathbb{C}_t^m} \langle V_\alpha(0)V_\gamma(x) \prod_{\ell=1}^m V_\gamma(x_\ell) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\mathbf{x}.$$

By Lemma 9.2

$$\int_{\mathbb{C}_t^m} \langle V_\alpha(0)V_\gamma(x) \prod_{\ell=1}^m V_\gamma(x_\ell) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\mathbf{x} = C \langle V_\alpha(0)V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \leq C|x|^{-\gamma\alpha} = C e^{t\gamma\alpha}.$$

where we used the formula (1.11) and this estimate is uniform over the compact subsets of $\mathbf{z} \in \theta\mathcal{Z}$. Hence

$$\left| \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} B_t(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right| \leq C e^{t(|\nu|+\alpha-2)}$$

as claimed.

Case $\nu_k = 1$. Here we need to regularize also the Liouville expectation: let $\langle - \rangle_{t,\epsilon}$ be as in (7.32) except we replace $e^{\gamma c} M_\gamma(\mathbb{C}_t)$ by the regularized version $\int_{\mathbb{C}_t} V_{\gamma,\epsilon}(x) dx$. We use following variant of Lemma 9.4.

Lemma 9.5. *Let $I'_\epsilon(\mathbf{u}, \mathbf{z}) := \frac{1}{u_k} \sum_{p=1}^n \partial_{z_p} \langle T(\mathbf{u}^k) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i,\epsilon}(z_i) \rangle_{t,\epsilon}$. Then $\lim_{\epsilon \rightarrow 0} I'_\epsilon(\mathbf{u}, \mathbf{z}) := I'(\mathbf{u}, \mathbf{z})$ exists and defines a continuous function in $\mathbf{z} \in \theta \mathcal{Z}$ satisfying*

$$(9.27) \quad \int \left(\frac{1}{(2\pi i)^{|\nu|}} \oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} I'(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} = \mathcal{T}_t(\nu^{(k)}, \mathbf{D}_{\nu_k}^* \varphi)$$

for all $\varphi \in C_0^\infty(\theta \mathcal{Z})$.

Proof. The proof is similar to Lemma 9.4 but cancellations occur for other reasons and we explain how. First, the integration by parts gives

$$I'_\epsilon(\mathbf{u}, \mathbf{z}) = K'_\epsilon(\mathbf{u}, \mathbf{z}) + L'_\epsilon(\mathbf{u}, \mathbf{z})$$

where $\lim_{\epsilon \rightarrow 0} K'_\epsilon = K'$ exists and is given by

$$\begin{aligned} K'(\mathbf{u}, \mathbf{z}) &= - \sum_{p=1}^n \sum_{\ell=1}^{k-1} \frac{Q\alpha_p}{u_k(z_p - u_\ell)^3} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad + \sum_{p=1}^n \sum_{\ell=1}^{k-1} \frac{\alpha_p}{u_k(z_p - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad - \sum_{p=1}^n \frac{\alpha_p \alpha}{2} \frac{1}{u_k z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &\quad - \sum_{p \neq p'=1}^n \frac{\alpha_p \alpha_{p'}}{2} \frac{1}{u_k(z_p - z_{p'})} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\ &=: C_1(\mathbf{u}, \mathbf{z}) + C_2(\mathbf{u}, \mathbf{z}) + C_3(\mathbf{u}, \mathbf{z}) + C_5(\mathbf{u}, \mathbf{z}) \end{aligned}$$

whereas

$$L'_\epsilon(\mathbf{u}, \mathbf{z}) = - \frac{\mu\gamma}{u_k} \sum_{p=1}^n \alpha_p \int_{\mathbb{C}_t} C_{\epsilon,\epsilon}(z_p, x) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_{\gamma,\epsilon}(x) \prod_{i=1}^n V_{\alpha_i,\epsilon}(z_i) \rangle_{t,\epsilon} dx$$

is the term that needs analysis. Let us define

$$B'_t(\mathbf{u}, \mathbf{z}) := - \frac{i\mu}{u_k} \oint_{|x|=e^{-t}} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t d\bar{x}.$$

Then

$$\begin{aligned} B'_t(\mathbf{u}, \mathbf{z}) &= - \frac{i\mu}{u_k} \lim_{\epsilon \rightarrow 0} \oint_{|x|=e^{-t}} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_{\gamma,\epsilon}(x) \prod_{i=1}^n V_{\alpha_i,\epsilon}(z_i) \rangle_{t,\epsilon} d\bar{x} \\ &= \frac{\mu}{u_k} \lim_{\epsilon \rightarrow 0} \int_{\mathbb{C}_t} \partial_x \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_{\gamma,\epsilon}(x) \prod_{i=1}^n V_{\alpha_i,\epsilon}(z_i) \rangle_{t,\epsilon} dx \\ &= - \frac{\mu Q\gamma}{u_k} \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(x - u_\ell)^3} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad + \frac{\mu\gamma}{u_k} \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(x - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad - \mu \frac{\gamma\alpha}{2u_k} \int_{\mathbb{C}_t} \frac{1}{x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx \\ &\quad - \frac{\mu\gamma}{u_k} \lim_{\epsilon \rightarrow 0} \sum_{p=1}^n \alpha_p \int_{\mathbb{C}_t} C_{\epsilon,\epsilon}(z_p, x) \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_{\gamma,\epsilon}(x) \prod_{i=1}^n V_{\alpha_i,\epsilon}(z_i) \rangle_{t,\epsilon} dx \\ &=: P_6(\mathbf{u}, \mathbf{z}) + P_7(\mathbf{u}, \mathbf{z}) + P_8(\mathbf{u}, \mathbf{z}) + P_4(\mathbf{u}, \mathbf{z}) \end{aligned}$$

This proves the existence of $\lim_{\epsilon \rightarrow 0} L'_\epsilon = L' = P_4$ and furthermore

$$(9.28) \quad I' = \lim_{\epsilon \rightarrow 0} I'_\epsilon = C_1 + C_2 + C_3 + C_5 + B'_t - P_6 - P_7 - P_8.$$

The claim (9.27) follows as in Lemma 9.4. \square

Let us consider the expression

$$\begin{aligned} K'(\mathbf{u}, \mathbf{z}) &:= (T(\mathbf{u})V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i))_t + I'(\mathbf{u}, \mathbf{z}) - M(\mathbf{u}, \mathbf{z}) \\ &= R(\mathbf{u}, \mathbf{z}) + N(\mathbf{u}, \mathbf{z}) + D(\mathbf{u}, \mathbf{z}) + I'(\mathbf{u}, \mathbf{z}). \end{aligned}$$

By (9.13) and (9.27) we have

$$(9.29) \quad \int \left(\oint_{|\mathbf{u}|=\delta_t} \mathbf{u}^{1-\nu} K(\mathbf{u}, \mathbf{z}) d\mathbf{u} \right) \bar{\varphi}(\mathbf{z}) d\mathbf{z} = \mathcal{T}(\nu, \varphi) - \mathcal{T}(\nu^{(k)}, \mathbf{D}_{\nu_k}^* \varphi).$$

On the other hand combining (9.10), (9.16) and (9.21) we obtain

$$(9.30) \quad K = R + N + \sum_{i=1}^9 T_i + \sum_{i=1}^3 C_i + C_5 - \sum_{i=6}^8 P_i + B'_t$$

As before it is easy to check that the following relations hold (see Appendix C)

$$(9.31) \quad \oint_{|u_k|=e^{-t}\epsilon_k} T_i(\mathbf{u}, \mathbf{z}) du_k = 0 \quad \text{for } i = 4, 5, 9$$

$$(9.32) \quad \oint_{|u_k|=e^{-t}\epsilon_k} (T_i(\mathbf{u}, \mathbf{z}) + C_i(\mathbf{u}, \mathbf{z})) du_k = 0 \quad \text{for } i = 1, 2, 3$$

$$(9.33) \quad \oint_{|u_k|=e^{-t}\epsilon_k} (T_i(\mathbf{u}, \mathbf{z}) - P_i(\mathbf{u}, \mathbf{z})) du_k = 0 \quad \text{for } i = 6, 7, 8$$

$$(9.34) \quad \oint_{|u_k|=e^{-t}\epsilon_k} C_5(\mathbf{u}, \mathbf{z}) du_k = 0$$

$$(9.35) \quad \oint_{|u_k|=e^{-t}\epsilon_k} N(\mathbf{u}, \mathbf{z}) du_k = 0.$$

We can now conclude as in the case $\nu_k > 1$. \square

APPENDIX A. ELEMENTARY LEMMAS ON THE VIRASORO ALGEBRA

Lemma A.1. *Let $t_1, \dots, t_k \in \mathbb{Z}$ be such that $t_1 + \dots + t_k > 0$. Then*

$$(A.1) \quad \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 = 0, \quad \widetilde{\mathbf{L}}_{t_1}^{0,\alpha} \dots \widetilde{\mathbf{L}}_{t_k}^{0,\alpha} 1 = 0.$$

Proof. We will prove this relation by recursion on k and work with $\mathbf{L}_n^{0,\alpha}$ (the case of the $\widetilde{\mathbf{L}}_n^{0,\alpha}$ is identical). For $k = 1$, the relation comes from the fact that $\mathbf{L}_t^{0,\alpha} 1 = 0$ for $t > 0$. For $k \geq 2$, let j be the biggest index such that $t_j > 0$. If $j = k$ we are done. Otherwise, we have using (4.20) on $\mathbf{L}_{t_j}^{0,\alpha}, \mathbf{L}_{t_{j+1}}^{0,\alpha}$

$$\begin{aligned} \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 &= \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+1}}^{0,\alpha} \mathbf{L}_{t_j}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 \\ &= (t_j - t_{j+1}) \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_j+t_{j+1}}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 \\ &\quad + \frac{cL}{12} \delta_{t_j=-t_{j+1}} (t_j^3 - t_j) \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1. \end{aligned}$$

The second term in the above equality is equal to 0 by the recursion hypothesis. The third term in the above equality is equal to 0 since if $t_j = -t_{j+1}$ then we can apply the recursion hypothesis on $(t_1, \dots, t_{j-1}, t_{j+2}, \dots, t_k)$ since in this case $t_1 + \dots + t_{j-1} + t_{j+2} + \dots + t_k = t_1 + \dots + t_k$. Therefore by iterating the above procedure with the couple (t_j, t_{j+2}) , etc... we end up with

$$\mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 = \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+1}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} \mathbf{L}_{t_j}^{0,\alpha} 1 = 0. \quad \square$$

Lemma A.2. *We suppose that $t_1, \dots, t_k \in \mathbb{Z}$ are such that $t_1 + \dots + t_k = 0$.*

$$\mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} 1 = \sum_{k \geq 0} a_k (\mathbf{L}_0^{0,\alpha})^k 1$$

where the a_k only depend on the commutation relations and not on α .

Proof. We suppose that for all t_1, \dots, t_r such that $|t_i| \neq 0$ for all i , $t_1 + \dots + t_r = 0$ and $|t_1| + \dots + |t_r| \leq N$ the lemma holds. We consider t_1, \dots, t_k such that $t_1 + \dots + t_k = 0$, $|t_i| \neq 0$ for all i and $|t_1| + \dots + |t_k| \leq N + 1$. We consider j the largest integer such that $t_j > 0$. Hence for all $l > j$ we have $t_l < 0$. We have

$$\begin{aligned} \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_j}^{0,\alpha} \mathbf{L}_{t_{j+1}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} &= \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j+1}}^{0,\alpha} \mathbf{L}_{t_j}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} \\ &\quad + (t_j - t_{j+1}) \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_j+t_{j+1}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} \\ &\quad + \frac{cL}{12} (t_j^3 - t_j) \delta_{t_j = -t_{j+1}} \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha}. \end{aligned}$$

If $t_j = -t_{j+1}$ then the third term is of the desired form by the recursive assumption; if $t_j \neq -t_{j+1}$ then the third term is 0. If $t_j + t_{j+1} \neq 0$ then the second term is of the desired form by the recursive assumption. If not then one can use $\mathbf{L}_n^{0,\alpha} L_0^{0,\alpha} = \mathbf{L}_0^{0,\alpha} \mathbf{L}_n^{0,\alpha} + n \mathbf{L}_n^{0,\alpha}$ to get

$$(A.2) \quad \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_0^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} = (t_1 + \dots + t_{j-1}) \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha} + \mathbf{L}_0^{0,\alpha} \mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j-1}}^{0,\alpha} \mathbf{L}_{t_{j+2}}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha}$$

and one gets the result by the recursive assumption. To conclude the recursion step, one just needs to iterate the above procedure of shifting the $\mathbf{L}_{t_j}^{0,\alpha}$ to the right on the first term $\mathbf{L}_{t_1}^{0,\alpha} \dots \mathbf{L}_{t_{j+1}}^{0,\alpha} \mathbf{L}_{t_j}^{0,\alpha} \dots \mathbf{L}_{t_k}^{0,\alpha}$ and using $\mathbf{L}_{t_j}^{0,\alpha} 1 = 0$.

In the general case (i.e. when some t_i can be equal to 0), one can shift the $L_0^{0,\alpha}$ terms to the left using (A.2). \square

APPENDIX B. THE DOZZ FORMULA

We set $\ell(z) = \frac{\Gamma(z)}{\Gamma(1-z)}$ where Γ denotes the standard Gamma function. We introduce Zamolodchikov's special holomorphic function $\Upsilon_{\frac{\gamma}{2}}(z)$ by the following expression for $0 < \text{Re}(z) < Q$

$$(B.1) \quad \ln \Upsilon_{\frac{\gamma}{2}}(z) = \int_0^\infty \left(\left(\frac{Q}{2} - z \right)^2 e^{-t} - \frac{(\sinh((\frac{Q}{2} - z)\frac{t}{2}))^2}{\sinh(\frac{t\gamma}{4}) \sinh(\frac{t}{\gamma})} \right) \frac{dt}{t}.$$

The function $\Upsilon_{\frac{\gamma}{2}}$ is then defined on all \mathbb{C} by analytic continuation of the expression (B.1) as expression (B.1) satisfies the following remarkable functional relations:

$$(B.2) \quad \Upsilon_{\frac{\gamma}{2}}\left(z + \frac{\gamma}{2}\right) = \ell\left(\frac{\gamma}{2}z\right) \left(\frac{\gamma}{2}\right)^{1-\gamma z} \Upsilon_{\frac{\gamma}{2}}(z), \quad \Upsilon_{\frac{\gamma}{2}}\left(z + \frac{2}{\gamma}\right) = \ell\left(\frac{2}{\gamma}z\right) \left(\frac{\gamma}{2}\right)^{\frac{4}{\gamma}z-1} \Upsilon_{\frac{\gamma}{2}}(z).$$

The function $\Upsilon_{\frac{\gamma}{2}}$ has no poles in \mathbb{C} and the zeros of $\Upsilon_{\frac{\gamma}{2}}$ are simple (if $\gamma^2 \notin \mathbb{Q}$) and given by the discrete set $(-\frac{\gamma}{2}\mathbb{N} - \frac{2}{\gamma}\mathbb{N}) \cup (Q + \frac{\gamma}{2}\mathbb{N} + \frac{2}{\gamma}\mathbb{N})$. With these notations, the DOZZ formula is defined for $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{C}$ by the following formula where we set $\bar{\alpha} = \alpha_1 + \alpha_2 + \alpha_3$

$$(B.3) \quad C_{\gamma,\mu}^{\text{DOZZ}}(\alpha_1, \alpha_2, \alpha_3) = \left(\pi \mu \ell\left(\frac{\gamma^2}{4}\right) \left(\frac{\gamma}{2}\right)^{2-\frac{\gamma^2}{2}} \right)^{\frac{2Q-\bar{\alpha}}{\gamma}} \frac{\Upsilon'_{\frac{\gamma}{2}}(0) \Upsilon_{\frac{\gamma}{2}}(\alpha_1) \Upsilon_{\frac{\gamma}{2}}(\alpha_2) \Upsilon_{\frac{\gamma}{2}}(\alpha_3)}{\Upsilon_{\frac{\gamma}{2}}(\frac{\bar{\alpha}}{2} - Q) \Upsilon_{\frac{\gamma}{2}}(\frac{\bar{\alpha}}{2} - \alpha_1) \Upsilon_{\frac{\gamma}{2}}(\frac{\bar{\alpha}}{2} - \alpha_2) \Upsilon_{\frac{\gamma}{2}}(\frac{\bar{\alpha}}{2} - \alpha_3)}$$

The DOZZ formula is meromorphic with poles corresponding to the zeroes of the denominator of expression (B.3). Note that it is symmetric in $\alpha_1, \alpha_2, \alpha_3$ and real valued when α_j are real.

APPENDIX C. INTEGRATION BY PARTS CALCULATIONS

R-terms. Here we list explicitly the terms in the integration by parts formula in the proof of Proposition 9.3 giving zero contribution to the contour integral:

$$\begin{aligned}
R(\mathbf{u}, \mathbf{z}) := & 3Q^2 \sum_{\ell=1}^{k-1} \frac{1}{(u_k - u_\ell)^4} \langle T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& - 2Q \sum_{\ell=1}^{k-1} \frac{1}{(u_k - u_\ell)^3} \langle T(\mathbf{u}^{(\ell, k)}) \partial_z X(u_\ell) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& + \frac{Q\alpha}{2} \frac{1}{u_k^2} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& - Q^2 \sum_{\ell, \ell' \neq 1}^{k-1} \frac{1}{(u_k - u_\ell)^3 (u_k - u_{\ell'})^3} \langle T(\mathbf{u}^{\ell, \ell', k}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& + 2Q \sum_{\ell, \ell' = 1}^{k-1} \frac{1}{(u_k - u_\ell)^3 (u_k - u_{\ell'})^2} \langle \partial_z X(u_{\ell'}) T(\mathbf{u}^{\ell, \ell', k}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& - Q\alpha \sum_{\ell=1}^{k-1} \frac{1}{(u_k - u_\ell)^3 u_k} \langle T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& - \sum_{\ell, \ell' = 1}^{k-1} \frac{1}{(u_k - u_\ell)^2 (u_k - u_{\ell'})^2} \langle \partial_z X(u_\ell) \partial_z X(u_{\ell'}) T(\mathbf{u}^{\ell, \ell', k}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& + \alpha \sum_{\ell=1}^{k-1} \frac{1}{(u_k - u_\ell)^2 u_k} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t \\
& - \frac{\alpha^2}{4} \frac{1}{u_k^2} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t.
\end{aligned}$$

Proof of (9.24)-(9.26). The claim $T'_9 = -T_9$ results from the relation

$$\frac{1}{(u_k - x)(u_k - x')} = \frac{1}{x - x'} \left(\frac{1}{u_k - x} - \frac{1}{u_k - x'} \right)$$

and the fact that the mapping $(x, x') \mapsto \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) V_\gamma(x') \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$ is symmetric.

Also, $T_5 = D_5$ comes from the relation

$$\frac{1}{(u_k - z_p)(u_k - z_{p'})} = \frac{1}{z_p - z_{p'}} \left(\frac{1}{u_k - z_p} - \frac{1}{u_k - z_{p'}} \right)$$

and a re-indexation of the double sum.

Using

$$\frac{1}{(u_k - u_\ell)^2 (u_k - x)} = \frac{1}{(u_k - u_\ell)^2 (u_\ell - x)} - \frac{1}{(u_k - u_\ell)(x - u_\ell)^2} + \frac{1}{(x - u_\ell)^2 (u_k - x)}$$

we find that

$$T_7 + T'_7 = -\mu\gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \left(\frac{1}{(u_k - u_\ell)^2 (u_\ell - x)} - \frac{1}{(u_k - u_\ell)(x - u_\ell)^2} \right) \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell, k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx$$

satisfies

$$\oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} (T_7(\mathbf{u}, \mathbf{z}) + T'_7(\mathbf{u}, \mathbf{z})) du_k = 0$$

by moving the contour to ∞ ($\nu_k > 1$ is used here).

Using that

$$\frac{1}{u_k(u_k - x)} - \frac{1}{x(u_k - x)} = -\frac{1}{u_k x}$$

and $\oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} \frac{1}{u_k} du_k = 0$ for $\nu_k > 1$ we deduce that

$$\oint_{|u_k|=e^{-t}\delta_k} u_k^{1-\nu_k} (T_8(\mathbf{u}, \mathbf{z}) + T_8'(\mathbf{u}, \mathbf{z})) du_k = 0.$$

The relation

$$\frac{1}{(u_k - u_\ell)^3 (u_k - x)} = \frac{1}{(u_k - u_\ell)^3 (u_\ell - x)} - \frac{1}{(u_k - u_\ell)^2 (x - u_\ell)^2} + \frac{1}{(u_k - u_\ell) (u_\ell - x)^3} - \frac{1}{(u_k - x) (u_\ell - x)^3}$$

entails in the same way (9.25) for $i = 6$.

Finally the relations (9.26) follow by computing residues at the pole $u_k = z_p$.

Proof of (9.31)-(9.35).

The relations (9.31) and (9.35) holds because all the corresponding $T_i(\mathbf{u})$ are holomorphic in $u_k \in \mathbb{D}_t$. For (9.34) we observe that

$$\frac{1}{2\pi i} \oint_{|u_k|=e^{-t}\epsilon_k} C_5(\mathbf{u}) du_k = - \sum_{p \neq p'=1}^n \frac{\alpha_p \alpha_{p'}}{2} \frac{1}{(z_p - z_{p'})} \langle T(\mathbf{u}^k) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t$$

and that this expression is null for antisymmetry reasons. (9.32) and (9.33) follow from the residue at z_p and x respectively

$$\begin{aligned} \oint_{|u_k|=e^{-t}\epsilon_k} T_1(\mathbf{u}, \mathbf{z}) du_k &= 2\pi i \sum_{\ell=1}^{k-1} \sum_{p=1}^n \frac{Q\alpha_p}{(z_p - u_\ell)^3} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = - \oint_{|u_k|=e^{-t}\epsilon_k} C_1(\mathbf{u}, \mathbf{z}) du_k \\ \oint_{|u_k|=e^{-t}\epsilon_k} T_2(\mathbf{u}, \mathbf{z}) du_k &= - 2\pi i \sum_{\ell=1}^{k-1} \sum_{p=1}^n \frac{\alpha_p}{(z_p - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = - \oint_{|u_k|=e^{-t}\epsilon_k} C_2(\mathbf{u}, \mathbf{z}) du_k \\ \oint_{|u_k|=e^{-t}\epsilon_k} T_3(\mathbf{u}, \mathbf{z}) du_k &= 2\pi i \sum_{p=1}^n \frac{\alpha_p}{2} \frac{1}{z_p} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t = - \oint_{|u_k|=e^{-t}\epsilon_k} C_3(\mathbf{u}, \mathbf{z}) du_k, \end{aligned}$$

thus proving (9.32). Finally, we compute

$$\begin{aligned} \frac{1}{2\pi i} \oint_{|u_k|=e^{-t}\epsilon_k} T_6(\mathbf{u}, \mathbf{z}) du_k &= -\mu Q \gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(x - u_\ell)^3} \langle T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx = P_6(\mathbf{u}, \mathbf{z}) \\ \frac{1}{2\pi i} \oint_{|u_k|=e^{-t}\epsilon_k} T_7(\mathbf{u}, \mathbf{z}) du_k &= \mu \gamma \sum_{\ell=1}^{k-1} \int_{\mathbb{C}_t} \frac{1}{(x - u_\ell)^2} \langle \partial_z X(u_\ell) T(\mathbf{u}^{(\ell,k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx = P_7(\mathbf{u}, \mathbf{z}) \\ \frac{1}{2\pi i} \oint_{|u_k|=e^{-t}\epsilon_k} T_8(\mathbf{u}, \mathbf{z}) du_k &= -\frac{\mu \gamma \alpha}{2} \int_{\mathbb{C}_t} \frac{1}{x} \langle T(\mathbf{u}^{(k)}) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx = P_8(\mathbf{u}, \mathbf{z}) \\ \frac{1}{2\pi i} \oint_{|u_k|=e^{-t}\epsilon_k} C_4(\mathbf{u}, \mathbf{z}) du_k &= \mu \gamma \sum_{p=1}^n \frac{\alpha_p}{2} \int_{\mathbb{C}_t} \frac{1}{(z_p - x)} \langle T(\mathbf{u}^k) V_\alpha(0) V_\gamma(x) \prod_{i=1}^n V_{\alpha_i}(z_i) \rangle_t dx = P_4(\mathbf{u}, \mathbf{z}). \end{aligned}$$

C.1. Analyticity of the vertex operators. Recall the definition of $U_\alpha(\mathbf{z})$ in (7.30) for $\mathbf{z} \in \mathcal{Z}$ and real α_i 's such that $\alpha_i < Q$. It is plain to see that $U_\alpha(\mathbf{z})$ agrees with the following slightly different regularization

$$U_\alpha^{\text{holes}}(\mathbf{z}) = \lim_{k \rightarrow \infty} U_{\alpha,k}^{\text{holes}}(\mathbf{z})$$

$$\text{with } U_{\alpha,k}^{\text{holes}}(\mathbf{z}) = e^{\sum_i \alpha_i - Qc} e^{\sum_i \alpha_i P\varphi(z_i)} \mathbb{E}_\varphi \left[\left(\prod_{i=1}^n \epsilon_k^{\frac{\alpha_i^2}{2}} e^{\alpha_i X_{\mathbb{D}, \epsilon_k}(z_i)} \right) e^{-\mu e^{\gamma c} M_\gamma(\mathbb{D}_k)} \right]$$

where $\epsilon_k = 2^{-k}$, $X_{\mathbb{D}, \epsilon_k}$ is the circle average of the Dirichlet GFF and \mathbb{D}_k is the unit disk with small holes removed around each insertion, namely $\mathbb{D}_k := \mathbb{D} \setminus \bigcup_{i=1}^n B(z_i, \epsilon_k)$. Recall that we get the following explicit expression by using the Girsanov theorem:

$$(C.1) \quad U_\alpha^{\text{holes}}(\mathbf{z}) = e^{(\sum_{j=1}^n \alpha_j - Q)c} e^{\sum_i \alpha_i P\varphi(z_i) + \sum_{i < j} \alpha_i \alpha_j G_{\mathbb{D}}(z_i, z_j)} \mathbb{E}_\varphi \left[e^{-\mu e^{\gamma c} \int_{\mathbb{D}} e^{\gamma \sum_{i=1}^n \alpha_i G_{\mathbb{D}}(x, z_i)} M_\gamma(dx)} \right].$$

We fix k_0 such that the open balls $B(z_i, 2^{-k_0})$ are disjoint and included in \mathbb{D} . Set $\mathcal{O}^n := \{\alpha \in \mathbb{R}^n; \alpha_i < Q, \forall i\}$. Then we have the following analyticity result:

Proposition C.1. *The (random) function $\alpha \rightarrow U_{\alpha}^{\text{holes}}(\mathbf{z})$ admits an analytic extension in a complex neighborhood of \mathcal{O}^n such that for all real $\alpha \in \mathcal{O}^n$ there exists some $\epsilon > 0$ (depending on α) and (non random) $C, \tilde{C} > 0$ satisfying*

$$\begin{aligned} & \sup_{\beta \in [-\epsilon, \epsilon]^n} |U_{\alpha+i\beta}^{\text{holes}}(\mathbf{z}) - U_{\alpha+i\beta, k_0}^{\text{holes}}(\mathbf{z})| \\ & \leq C(1 + e^{\gamma c}) e^{(\sum_{j=1}^n \alpha_j - Q)c} e^{\tilde{C} \sup_i \sup_{u \in B(z_i, 2^{-k_0})} P\varphi(u)} \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} \int_{\mathbb{D}_{k_0}} e^{\gamma \sum_{i=1}^n \alpha_i G_{\mathbb{D}}(x, z_i)} M_{\gamma}(dx)} \right]. \end{aligned}$$

Proof. To simplify we will suppose that $n = 1$ and $z_1 = 0$ and set $G_{\mathbb{D}, k}(0, u) = \mathbb{E}[X_{\mathbb{D}, \epsilon k}(0) X_{\mathbb{D}}(u)]$. This is no restriction as the same analysis can be performed around each insertion in case $n > 1$. In this context, we get by using the Markov property of the Dirichlet GFF that:

$$\begin{aligned} & |U_{\alpha+i\beta, k+1}^{\text{holes}} - U_{\alpha+i\beta, k}^{\text{holes}}(\mathbf{z})| \\ & = e^{(\alpha-Q)c} |\mathbb{E}_{\varphi} \left[\epsilon_{k+1}^{(\alpha+i\beta)^2/2} e^{(\alpha+i\beta)X_{\mathbb{D}, \epsilon k+1}(0)} \left(e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_{k+1})} - e^{-\mu e^{\gamma c} M_{\gamma}(\mathbb{D}_k)} \right) \right]| \\ & \leq e^{(\alpha-Q)c} 2^{(k+1)\beta^2/2} |\mathbb{E}_{\varphi} \left[\left(e^{-\mu e^{\gamma c} \int_{\mathbb{D}_{k+1}} e^{\gamma \alpha_1 G_{\mathbb{D}, k+1}(x, 0)} M_{\gamma}(dx)} - e^{-\mu e^{\gamma c} \int_{\mathbb{D}_k} e^{\gamma \alpha_1 G_{\mathbb{D}, k+1}(x, 0)} M_{\gamma}(dx)} \right) \right]| \\ & = e^{(\alpha-Q)c} 2^{(k+1)\beta^2/2} |\mathbb{E}_{\varphi} \left[\left(e^{-\mu e^{\gamma c} (Y_k + \delta Y_k)} - e^{-\mu e^{\gamma c} Y_k} \right) \right]|, \end{aligned}$$

where we have set

$$Y_k = \int_{\mathbb{D}_k} e^{\gamma P\varphi(x)} e^{\gamma \alpha_1 G_{\mathbb{D}, \epsilon k+1}(0, x)} M_{\gamma}(dx) \quad \text{and} \quad \delta Y_k = \int_{\mathbb{D}_k \setminus \mathbb{D}_{k+1}} e^{\gamma P\varphi(x)} e^{\gamma \alpha_1 G_{\mathbb{D}, k+1}(0, u)} M_{\gamma}(dx).$$

Now, we consider the cases $\delta Y_k > 1$ and $\delta Y_k \leq 1$. By FKG inequality for the Dirichlet GFF, we have

$$\begin{aligned} & \mathbb{E}_{\varphi} \left[\mathbf{1}_{\delta Y_k > 1} |e^{-\mu e^{\gamma c} Y_k} - e^{-\mu e^{\gamma c} (Y_k + \delta Y_k)}| \right] \\ & \leq 2 \mathbb{E}_{\varphi} \left[\mathbf{1}_{\delta Y_k > 1} e^{-\mu e^{\gamma c} Y_k} \right] \\ & \leq 2 \mathbb{E}_{\varphi} \left[\mathbf{1}_{\delta Y_k > 1} \right] \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_k} \right] \\ & \leq 2 \mathbb{E}_{\varphi} \left[(\delta Y_k)^{\eta} \right] \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_k} \right]. \end{aligned}$$

Next, we choose $\beta > 0$ and $\eta > 0$ such that $2^{(k+1)\beta^2/2} \mathbb{E}_{\varphi} \left[(\delta Y_k)^{\eta} \right] \leq C e^{\gamma \eta \sup_{|u| \leq \epsilon_k} P\varphi(u)} 2^{-\theta k}$ with $\theta > 0$.

In the case $\delta Y_k \leq 1$, we get (using the inequality $x \mathbf{1}_{\{x \leq 1\}} \leq x^{\eta}$ for $x > 0$ and then FKG for the Dirichlet GFF)

$$\begin{aligned} & 2^{(k+1)\beta^2/2} \mathbb{E}_{\varphi} \left[\mathbf{1}_{\delta Y_k \leq 1} |e^{-\mu e^{\gamma c} Y_k} - e^{-\mu e^{\gamma c} (Y_k + \delta Y_k)}| \right] \\ & \leq 2^{(k+1)\beta^2/2} \mu e^{\gamma c} \mathbb{E}_{\varphi} \left[\mathbf{1}_{\delta Y_k \leq 1} \delta Y_k e^{-\mu e^{\gamma c} Y_k} \right] \\ & \leq 2^{(k+1)\beta^2/2} \mu e^{\gamma c} \mathbb{E}_{\varphi} \left[(\delta Y_k)^{\eta} e^{-\mu e^{\gamma c} Y_k} \right] \\ & \leq 2^{(k+1)\beta^2/2} \mu e^{\gamma c} \mathbb{E}_{\varphi} \left[(\delta Y_k)^{\eta} \right] \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_k} \right] \\ & \leq C 2^{-k\theta} e^{\gamma c} e^{\gamma \eta \sup_{|u| \leq \epsilon_k} P\varphi(u)} \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_k} \right]. \end{aligned}$$

Gathering the above considerations, we get

$$|U_{\alpha+i\beta, k+1}^{\text{holes}} - U_{\alpha+i\beta, k}^{\text{holes}}(\mathbf{z})| \leq C e^{(\alpha-Q)c} 2^{-k\theta} (1 + e^{\gamma c}) e^{\gamma \eta \sup_{|u| \leq \epsilon_n} P\varphi(u)} \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_n} \right]$$

This shows that the (random) analytic function $U_{\alpha+i\beta, k}^{\text{holes}}(\mathbf{z})$ converges as $k \rightarrow \infty$ with probability 1 and for all c towards an analytic function that satisfies

$$|U_{\alpha+i\beta, k+1}^{\text{holes}} - U_{\alpha+i\beta, k_0}^{\text{holes}}(\mathbf{z})| \leq C e^{(\alpha-Q)c} (1 + e^{\gamma c}) e^{\gamma \eta \sup_{|u| \leq \frac{1}{2}} P\varphi(u)} \mathbb{E}_{\varphi} \left[e^{-\mu e^{\gamma c} Y_{k_0}} \right]. \quad \square$$

REFERENCES

- [AFLT11] Alba V.A., Fateev V.A., Litvinov A.V., Tarnopolsky G.M.: On combinatorial expansion of the conformal blocks arising from AGT conjecture, *Lett. Math. Phys.* **98**, 33-64 (2011).
- [AGT10] L. F. Alday, D. Gaiotto, and Y. Tachikawa. Liouville Correlation Functions from Four Dimensional Gauge Theories, *Lett. Math. Phys.* **91**, 167-197 (2010).

- [BPZ84] Belavin A.A., Polyakov A.M., Zamolodchikov A.B., Infinite conformal symmetry in two-dimensional quantum field theory, *Nuclear Physics B* **241** (2), 333-380 (1984).
- [Be17] Berestycki N., An elementary approach to Gaussian multiplicative chaos, *Electronic communications in Probability* **27**, 1-12 (2017).
- [Bo86] Borcherds R., Vertex algebras, Kac-Moody algebras, and the Monster, *Proceedings of the National Academy of Sciences of the United States of America* **83**, 3068-3071 (1986).
- [BCT82] Braaten E., Curtright T., Thorn C.: Phys.Lett., **B118**, 115 (1982).
- [ChSm12] Chelkak D., Smirnov S.: Universality in the 2D Ising model and conformal invariance of fermionic observables, *Inventiones mathematicae* **189**, 515-580 (2012).
- [CHI15] Chelkak D., Hongler C., Izyurov K.: Conformal invariance of spin correlations in the planar Ising model, *Annals of mathematics* **181**, 1087-1138 (2015).
- [CuTh] Curtright T., Thorn C.: Phys.Rev.Lett., **48**, 1309 (1982).
- [DKRV16] David F., Kupiainen A., Rhodes R., Vargas V., Liouville Quantum Gravity on the Riemann sphere, *Communications in Mathematical Physics* **342** (3), 869-907 (2016).
- [DRV16] David F., Rhodes R., Vargas V.: Liouville Quantum Gravity on complex tori, *J. Math. Phys.* **57**, 022302 (2016).
- [DiDuDuFa] Ding J., Dubédat J., Dunlap A., Falconet H.: Tightness of Liouville first passage percolation for $\gamma \in (0, 2)$, [arXiv:1904.08021](https://arxiv.org/abs/1904.08021).
- [DoOt94] Dorn H., Otto H.-J.: Two and three point functions in Liouville theory, *Nuclear Physics B* **429** (2), 375-388 (1994).
- [DuFaGwPfs] Dubédat J., Falconet H., Gwynne E., Pfeffer J., Sun X.: Weak LQG metrics and Liouville first passage percolation, [arXiv:1905.00380](https://arxiv.org/abs/1905.00380)
- [DMS14] Duplantier B., Miller J., Sheffield S.: Liouville quantum gravity as mating of trees, [arXiv:1409.7055](https://arxiv.org/abs/1409.7055).
- [EPPR12a] El-Showk, S., Paulos M.F., Poland D., Rychkov S., Simmons-Duffin D., Vichi A., Solving the 3D Ising model with the conformal bootstrap, *Phys. Rev. D* **86**, 025022 (2012).
- [EPPR12b] El-Showk, S., Paulos M.F., Poland D., Rychkov S., Simmons-Duffin D., Vichi A., Solving the 3D Ising model with the conformal bootstrap II. c-minimization and precise critical exponents, *Phys. Rev. D* **86**, 025022 (2012).
- [FL10] Fateev V.A., Litvinov A.V.: On AGT conjecture, *JHEP*, 1002:014 (2010).
- [FF84] Feigin B.L., Fuchs D.B.: Verma modules over the Virasoro algebra, *General and Algebraic Topology, and Applications Proceedings of the International Topological Conference held in Leningrad, August 23-27, 1982* **1060**, Springer, 230-245 (1984).
- [FLM18] Felder G., Müller-Lennert M., Analyticity of Nekrasov partition functions, *Comm. Math. Phys.* **364** (2), 683-718 (2018).
- [FLM89] Frenkel I., Lepowsky J., Meurman A., Vertex Operator Algebras and the Monster, *Academic Press* (1989).
- [GKRV] Guillarmou C., Kupiainen A., Rhodes R., Vargas V.: in preparation.
- [Ga] Gawędzki K.: Lectures on conformal field theory, [Quantum field theory program at IAS](https://www.maths.ed.ac.uk/~igordon/LA1.pdf).
- [GeNe84] Gervais J.-L. and Neveu A. Nucl.Phys. **B238**, 125 (1984).
- [GRSS20] Ghosal P., Remy G., Sun X., Sun Y., Probabilistic conformal blocks for Liouville CFT on the torus, [arXiv:2003.03802](https://arxiv.org/abs/2003.03802).
- [Go09] Gordon I.: Infinite-dimensional Lie algebras, Lectures available at <https://www.maths.ed.ac.uk/~igordon/LA1.pdf>.
- [GRV19] Guillarmou C., Rhodes R., Vargas V., Polyakov's formulation of 2d bosonic string theory, *Publications mathématiques de l'IHES* **130**, 111-185 (2019).
- [Gu89] L. Guillopé, Théorie spectrale de quelques variétés à bouts, *Ann. Sci. Ecole Norm. Sup.* **22**, 137-160 (1989).
- [Ho71] Hoegh-Krohn R.: A general class of quantum fields without cut-offs in two space-time dimensions, *Commun. Math. Phys.* **21**, 244-255 (1971).
- [NS] Holden N., Sun X.: Convergence of uniform triangulations under the Cardy embedding, [arXiv:1905.13207](https://arxiv.org/abs/1905.13207) [math.PR].
- [HoRy13] Hogervorst M., Rychkov S.: Radial Coordinates for Conformal Blocks, *Phys. Rev. D* **87**, 106004 (2013), [arXiv:1303.1111](https://arxiv.org/abs/1303.1111) [hep-th].
- [Hu97] Huang Y.Z.: -Two-Dimensional Conformal Geometry and Vertex Operator Algebras, *Progress in Mathematics* (1997).
- [HuKo07] Huang Y.-Z., Kong L.: Full field algebras, *Communications in Mathematical Physics* **272**, 345-396 (2007).
- [Ja] Janson S.: Gaussian Hilbert Spaces, Cambridge Tracts in Mathematics, Cambridge, Cambridge University Press (1997).
- [KaRa] Kac V. G. and Raina A. K.: Bombay lectures on highest weight representations of infinite-dimensional Lie algebras, Advanced Series in Mathematical Physics. World Scientific Publishing Co., Inc., Teaneck, NJ, 1987, vol.2, pp.xii+145.
- [Ka85] Kahane, J.-P.: Sur le chaos multiplicatif, *Ann. Sci. Math. Québec*, **9** (2), 105-150 (1985).
- [KM13] Kang N.-G., Makarov N.: Gaussian free field theory and conformal field theory, *Astérisque* **353**, (2013).
- [Ko11] Kostov I.: Two-dimensional quantum gravity, in *The Oxford Handbook of Random Matrix Theory*, Akemann G., Baik J. and Di Francesco P. Eds, Oxford University Press (2011)
- [KQR20] Kravchuk P., Qiao J., Rychkov S.: Distributions in CFT I. Cross-Ratio Space, [arXiv:2001.08778](https://arxiv.org/abs/2001.08778) [hep-th].
- [KRV19] Kupiainen A., Rhodes R., Vargas V., Local conformal structure of Liouville Quantum Gravity, *Communications in Mathematical Physics* **371**, 1005-1069 (2019).
- [KRV20] Kupiainen A., Rhodes R., Vargas V.: Integrability of Liouville theory: proof of the DOZZ formula, *Annals of Mathematics* **191**, 81-166 (2020).
- [LeG13] Le Gall J.-F.: Uniqueness and universality of the Brownian map, *Annals of Probability* **41** (4), 2880-2960 (2013).
- [MaOk12] Maulik D. Okounkov A.: Quantum Groups and Quantum Cohomology, *Asterisque* **408** (2019).

- [Me93] Melrose R.B., The Atiyah-Patodi-Singer Index Theorem, Research Notes in Mathematics, A K Peters/CRC Press (1993).
- [Mier13] Miermont G.: The Brownian map is the scaling limit of uniform random plane quadrangulations, *Acta Mathematica* **210** (2), 319-401 (2013).
- [MiGw] Miller J., Gwynne E.: Existence and uniqueness of the Liouville quantum gravity metric for $\gamma \in (0, 2)$, [arXiv:1905.00383](#).
- [MS20a] Miller J., Sheffield S.: Liouville quantum gravity and the Brownian map I: The QLE(8/3,0) metric, *Inventiones Mathematicae* **219**, 75-152 (2020).
- [MS20b] Miller J., Sheffield S.: Liouville quantum gravity and the Brownian map II: geodesics and continuity of the embedding, [arXiv:1605.03563](#).
- [MS20c] Miller J., Sheffield S.: Liouville quantum gravity and the Brownian map III: The conformal structure is determined, [arXiv:1608.05391](#).
- [Ne16] Negut A.: Exts and the AGT Relations, *Letters in Mathematical Physics* **106**, 1265-1316 (2016).
- [Ne04] Nekrasov N. A. , Seiberg-Witten Prepotential from Instanton Counting, *Adv. Theor. Math. Phys.* **7** (2004) 831-864,
- [OsSh73] Osterwalder, K., Schrader, R., Axioms for Euclidean Green's functions I. *Commun. Math. Phys.* **31**, 83-112 (1973).
- [OsSh75] Osterwalder, K., Schrader, R., Axioms for Euclidean Green's functions II, *Commun. Math. Phys.* **42**, 281-305 (1975).
- [PRER12] Pappadopulo, Rychkov S., Espin J., Rattazzi R.: OPE convergence in Conformal Field Theory, *Phys. Rev. D* **86**, 105043 (2012), [arXiv:1208.6449 \[hep-th\]](#).
- [Pel19] Peltola E.: Towards a conformal field theory for Schramm-Loewner evolutions, *J. Math. Phys.* **60** (10), 103305 (2019).
- [PoRyVi] Poland D, Rychkov S, Vichi A: The Conformal Bootstrap: Theory, Numerical Techniques, and Applications, *Rev. Mod. Phys.* **91**, 15002 (2019)
- [Po81] Polyakov A.M.: Quantum geometry of bosonic strings, *Phys. Lett.* **103B**, 207 (1981).
- [Po08] Polyakov A.M.: From Quarks to Strings, [arXiv:0812.0183](#).
- [ReSi1] Reed M., Simon B.: Methods of Modern Mathematical Physics, Vol 1, revised and enlarged edition, Academic Press (1980).
- [ReSi2] Reed M., Simon B.: Methods of Modern Mathematical Physics, Vol 2, Academic Press (1975).
- [RhVa14] Rhodes R., Vargas, V.: Gaussian multiplicative chaos and applications: a review, *Probability Surveys* **11**, 315-392 (2014).
- [RoVa10] Robert R., Vargas V.: Gaussian multiplicative chaos revisited, *Annals of Probability* **38** (2), 605-631 (2010).
- [ScVa13] Schiffmann O., Vasserot E.: Cherednik algebras, W-algebras and the equivariant cohomology of the moduli space of instantons on A^2 , *Publications mathématiques de l'IHES* **118** (1), 213-342 (2013).
- [Sch00] Schramm O.: Scaling limits of loop-erased random walks and uniform spanning trees, *Israel Journal of Mathematics* **118**, 221-288 (2000).
- [Se90] Seiberg N., Notes on Quantum Liouville Theory and Quantum Gravity, *Progress of Theoretical Physics*, suppl. 102, 1990.
- [Sz98] Sznitman A.-S., Brownian Motion, Obstacles and Random Media, Springer Monographs in Mathematics, Springer-Verlag Berlin Heidelberg, 1998.
- [Te95] Teschner J., On the Liouville three point function, *Phys. Lett. B* **363**, 65-70 (1995).
- [Te01] Teschner J., Liouville Theory Revisited, *Class.Quant.Grav.* **18**, R153-R222 (2001).
- [Te04] Teschner J.: A lecture on the Liouville vertex operators, *Int.J.Mod.Phys. A19S2*, 436-458 (2004)
- [ZaZa95] Zamolodchikov A.B., Zamolodchikov A.B.: Structure constants and conformal bootstrap in Liouville field theory, [arXiv:hep-th/9506136](#).
- [ZaZa96] Zamolodchikov A.B., Zamolodchikov A.B.: Conformal bootstrap in Liouville field theory, *Nuclear Physics B* **477** (2), 577-605 (1996).

UNIVERSITÉ PARIS-SACLAY, CNRS, LABORATOIRE DE MATHÉMATIQUES D'ORSAY, 91405, ORSAY, FRANCE.
E-mail address: colin.guillarmou@math.u-psud.fr

UNIVERSITY OF HELSINKI, DEPARTMENT OF MATHEMATICS AND STATISTICS
E-mail address: antti.kupiainen@helsinki.fi

AIX-MARSEILLE UNIVERSITY, INSTITUT DE MATHÉMATIQUES DE MARSEILLE (I2M), AND INSTITUT UNIVERSITAIRE DE FRANCE (IUF)
E-mail address: remi.rhodes@univ-amu.fr

LAGA UNIVERSITÉ SORBONNE PARIS NORD, CNRS, UMR 7539, F-93430, VILLETANEUSE, FRANCE
E-mail address: vargas@math.univ-paris13.fr