



Characterization of 3D genomic interactions in fetal pig muscle

Maria Eugenia Marti Marimon, Hervé Acloque, Matthias Zytnicki, David Robelin, Sarah Djebali Quelen, Nathalie Villa-Vialaneix, Ole Madsen, Yvette Lahbib Mansais, Diane Esquerré, Florence Mompert, et al.

► To cite this version:

Maria Eugenia Marti Marimon, Hervé Acloque, Matthias Zytnicki, David Robelin, Sarah Djebali Quelen, et al.. Characterization of 3D genomic interactions in fetal pig muscle. 36. International Society for Animal Genetics Conference (ISAG), Jul 2017, Dublin, Ireland. pp.43. hal-02868765

HAL Id: hal-02868765

<https://hal.science/hal-02868765>

Submitted on 15 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: <http://oatao.univ-toulouse.fr/18301>

To cite this version:

Marti Marimon, Maria Eugenia and Acloque, Hervé and Zytnicki, Matthias and Robelin, David and Djebali Quelen, Sarah and Villa-Vialaneix, Nathalie and Madsen, Ole and Lahbib Mansais, Yvette and Esquerré, Diane and Mompert, Florence and Yerle-Bouissou, Martine and Groenen, Martien and Foissac, Sylvain Characterization of 3D genomic interactions in fetal pig muscle. (2017) In: 36th International Society for Animal Genetics Conference, 17 July 2017 - 21 July 2017 (Dublin, Ireland).

Any correspondence concerning this service should be sent to the repository administrator: tech-oatao@listes-diff.inp-toulouse.fr

Characterization of 3D genomic interactions in fetal pig muscle

Maria Marti-Marimon¹, Hervé Acloque¹, Matthias Zytnicki², David Robelin¹, Sarah Djebali¹, Nathalie Villa-Vialaneix², Ole Madsen³, Yvette Lahbib-Mansais¹, Diane Esquerré¹, Florence Mompert¹, Martien Groenen³, Martine Yerle-Bouissou¹, Sylvain Foissac¹.

¹GenPhySE, University of Toulouse, INRA, INPT, ENVT, Castanet Tolosan, France

²MIAT, Université de Toulouse, INRA, Castanet Tolosan, France

³Animal Breeding and Genomics Centre (ABG), Wageningen University, 6708 PB Wageningen, Netherlands.

Genome sequence alone is not sufficient to explain the overall coordination of nuclear activity in a particular tissue. Studies have shown the important role played by nuclear organization on gene expression regulation (Bickmore & Van Steensel 2013). Moreover, genomic long-range intra- and inter-chromosomal interactions are involved in the activation of tissue- specific gene networks (Fanucchi et al. 2013). **Here we present an overview of the pig genome architecture in muscle at two late developmental stages.** The maturation process occurs between the 90th day and the end of gestation (114 days). To characterize this key period for survival at birth, we profiled chromatin interactions genome-wide with in situ Hi-C (High Throughput Chromosome Conformation Capture) in muscle samples collected at 90 and 110 days of gestation, specific moments where a drastic change in gene expression has been reported (Voillet et al. 2014). About 200 million read pairs per library were generated (3 replicates per condition). This allowed to (a) produce an experimental Hi-C protocol optimized on frozen fetal tissues, (b) reveal the first Hi-C contact heatmaps in fetal porcine muscle cells, (c) profile Topologically Associated Domains (TAD) defined as genomic domains with high levels of chromatin interactions (Pombo & Dillon 2015). About 82% of the Hi-C reads could be mapped on the reference genome using the new assembly version *Sus scrofa* v11, from which, after filtering, 49% of valid read pairs have been used to infer the genomic interactions in both developmental stages. In addition, ChIP-seq experiments are being performed to map the binding of the structural protein CTCF, which is known to regulate genome structure by promoting interactions between genes and distal enhancers (Hnisz et al. 2016). By integrating the results from the previous differential analysis of gene expression (Voillet et al. 2014) and our results in nuclear organization (Hi-C data and ChIP-seq data), we will report general differences between both developmental stages in terms of transcription and structure.