



HAL
open science

The square root of a parabolic operator

El Maati Ouhabaz

► **To cite this version:**

| El Maati Ouhabaz. The square root of a parabolic operator. 2020. hal-02871224v1

HAL Id: hal-02871224

<https://hal.science/hal-02871224v1>

Preprint submitted on 17 Jun 2020 (v1), last revised 1 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The square root of a parabolic operator

El Maati Ouhabaz

Abstract

Let $L(t) = -\operatorname{div}(A(x, t)\nabla_x)$ for $t \in (0, \tau)$ be a uniformly elliptic operator with boundary conditions on a domain Ω of \mathbb{R}^d and $\partial = \frac{\partial}{\partial t}$. Define the parabolic operator $\mathcal{L} = \partial + L$ on $L^2(0, \tau, L^2(\Omega))$ by $(\mathcal{L}u)(t) := \frac{\partial u(t)}{\partial t} + L(t)u(t)$. We assume a very little of regularity for the boundary of Ω and assume that the coefficients $A(x, t)$ are measurable in x and piecewise C^α in t for some $\alpha > \frac{1}{2}$. We prove the Kato square root property for $\sqrt{\mathcal{L}}$ and the estimate

$$\|\sqrt{\mathcal{L}}u\|_{L^2(0, \tau, L^2(\Omega))} \approx \|\nabla_x u\|_{L^2(0, \tau, L^2(\Omega))} + \|u\|_{H^{\frac{1}{2}}(0, \tau, L^2(\Omega))} + \left(\int_0^\tau \|u(t)\|_{L^2(\Omega)}^2 \frac{dt}{t} \right)^{1/2}.$$

We also prove L^p -versions of this result.

Keywords: elliptic and parabolic operators, the Kato square root property, maximal regularity, the holomorphic functional calculus, non-autonomous evolution equations.

Home institution:

Institut de Mathématiques de Bordeaux
Université de Bordeaux, CNRS, UMR 5251,
351, Cours de la Libération. 33405 Talence, France.
Elmaati.Ouhabaz@math.u-bordeaux.fr

1 Introduction and the main results

Consider on $L^2(\mathbb{R}^d)$ the differential operator $L(t) = -\operatorname{div}(A(x, t)\nabla_x)$ where the matrix $A(x, t) = (a_{kl}(x, t))_{1 \leq k, l \leq d}$ has complex measurable entries and satisfies the usual ellipticity condition

$$\operatorname{Re}\langle A(x, t)\xi, \xi \rangle \geq \kappa|\xi|^2, \quad |\langle A(x, t)\xi, \zeta \rangle| \leq C|\xi||\zeta| \quad (1.1)$$

for all $\xi, \zeta \in \mathbb{C}^d$, where κ, C are positive constants independent of $(x, t) \in \mathbb{R}^d \times \mathbb{R}$ and $\langle \cdot, \cdot \rangle$ denotes the scalar product of \mathbb{C}^d . We consider the first order differential operator $\partial u = \frac{\partial u(t)}{\partial t}$

for all u in the Sobolev space $H^1(\mathbb{R}, L^2(\mathbb{R}^d))$. One defines the half-order derivative $\partial^{1/2}$ by

$$\partial^{1/2}u(t) = -\frac{1}{2\sqrt{2\pi}} \int_{\mathbb{R}} \frac{1}{|t-s|^{3/2}} (u(t) - u(s)) ds.$$

The following theorem is a parabolic version of the Kato square root property. It is proved by P. Auscher, M. Egert and K. Nyström [5].

Theorem 1.1. *Suppose (1.1). There exists a realization of the parabolic operator $\mathcal{L} := \partial + L$ which is maximal accretive on $L^2(\mathbb{R}^{d+1})$, the domain of its square root $\sqrt{\mathcal{L}}$ coincides with $H^{\frac{1}{2}}(\mathbb{R}, L^2(\mathbb{R}^d)) \cap L^2(\mathbb{R}, H^1(\mathbb{R}^d))$ and*

$$\|\sqrt{\mathcal{L}}u\|_{L^2(\mathbb{R}^{d+1})} \approx \|\nabla_x u\|_{L^2(\mathbb{R}^{d+1})} + \|\partial^{1/2}u\|_{L^2(\mathbb{R}^{d+1})}$$

for all $u \in D(\sqrt{\mathcal{L}})$.

A similar result was proved by K. Nyström [20] in the case where $A(x, t) = A(x)$.

The aim of the present short paper is twofold. We consider the above parabolic Kato square root problem for operators on domains with boundary conditions. Secondly, we investigate the problem on $L^p(0, \tau, L^r(\Omega))$ and not only on $L^2(0, \tau, L^2(\Omega))$. We consider the time variable t in an interval $(0, \tau)$ which is usual for evolution equations rather than the whole set \mathbb{R} . In order to give the precise statements we need some preparation.

Let Ω be an open subset of \mathbb{R}^d with boundary Γ . Consider a closed subspace V of $H^1(\Omega)$ which contains $H_0^1(\Omega)$ and define the sesquilinear form

$$\mathbf{a}(t, u, v) = \int_{\Omega} A(x, t) \nabla_x u \cdot \overline{\nabla_x v} dx$$

with domain V . We assume that the matrix $A(x, t) = (a_{kl}(x, t))_{1 \leq k, l \leq d}$ satisfies the ellipticity condition (1.1) with constants independent of $(x, t) \in \Omega \times (0, \tau)$. The associated operator is formally given by $L(t) = -\operatorname{div}(A(x, t) \nabla_x)$ and subject to the boundary conditions fixed by V . We say that \mathbf{a} is C^α for some $\alpha > 0$ if there exists a positive constant M such that for all $u, v \in V$

$$|\mathbf{a}(t, u, v) - \mathbf{a}(s, u, v)| \leq M|t - s|^\alpha \|u\|_V \|v\|_V.$$

We say that \mathbf{a} is piecewise C^α for some $\alpha > 0$ if there exist $\tau_1 = 0 < \tau_2 < \dots < \tau_N = \tau$ such that on each sub-interval (τ_j, τ_{j+1}) , \mathbf{a} is the restriction of a C^α form on $[\tau_j, \tau_{j+1}]$.

We make the following two assumptions. Suppose that \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$. Observe that this is satisfied if the coefficients a_{kl} , $1 \leq k, l \leq d$ are piecewise C^α in the t -variable for some $\alpha > \frac{1}{2}$.

Next, we assume that for each fixed t , the operator $L(t)$ satisfies the following Kato square root property

$$V \subseteq D\left(\sqrt{L(t)}\right) \quad \text{and} \quad \left\| \sqrt{L(t)} f \right\|_{L^2(\Omega)} \leq C \left[\|\nabla_x f\|_{L^2(\Omega)} + \|f\|_{L^2(\Omega)} \right], \quad f \in V. \quad (1.2)$$

The constant C is independent of t . In this case, a well known duality argument implies that $D\left(\sqrt{L(t)}\right) = V$ and the norms $\left\| \sqrt{L(t)} f \right\|_{L^2(\Omega)} + \|f\|_{L^2(\Omega)}$ and $\|\nabla_x f\|_{L^2(\Omega)} + \|f\|_{L^2(\Omega)}$ are equivalent. In many cases, the homogeneous estimate

$$D\left(\sqrt{L(t)}\right) = V \quad \text{and} \quad \left\| \sqrt{L(t)} f \right\|_{L^2(\Omega)} \approx \|\nabla_x f\|_{L^2(\Omega)} \quad (1.3)$$

holds. The implicit constants in the equivalence \approx are independent of t since they depend only on the ellipticity constants.

The square root property (1.3) is always satisfied if $A(x, t)$ is symmetric. It is satisfied if $\Omega = \mathbb{R}^d$ by the solution of the Kato square root problem (see P. Auscher et al. [4]). The non-homogeneous estimate (1.2) is satisfied if one has in addition terms of lower order. On domains, it is satisfied if the boundary of Ω has a little of regularity (for example Lipschitz is enough) for Dirichlet boundary conditions ($V = H_0^1(\Omega)$), Neumann boundary conditions ($V = H^1(\Omega)$) or even for mixed boundary conditions. For all this we refer to the paper of M. Egert, R. Haller-Dintelmann and P. Tolksdorf [13], the recent preprint of S. Bechtel, M. Egert and R. Haller-Dintelmann [8] and the references therein.

Now we state our first main result.

Theorem 1.2. *Suppose (1.1), (1.2) and suppose that \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$. Then there exists a realization of the parabolic operator $\mathcal{L} := \partial + L$ that is maximal accretive on $L^2(0, \tau, L^2(\Omega))$ and satisfies the Kato square root property*

$$D(\sqrt{\mathcal{L}}) = \left\{ u \in H^{\frac{1}{2}}(0, \tau, L^2(\Omega)) \cap L^2(0, \tau, V), \int_0^\tau \|u(t)\|_{L^2(\Omega)}^2 \frac{dt}{t} < \infty \right\}$$

and

$$\left\| \sqrt{\mathcal{L}} u \right\|_{L^2(0, \tau, L^2(\Omega))} \approx \|\nabla_x u\|_{L^2(0, \tau, L^2(\Omega))} + \|u\|_{H^{\frac{1}{2}}(0, \tau, L^2(\Omega))} + \left(\int_0^\tau \|u(t)\|_{L^2(\Omega)}^2 \frac{dt}{t} \right)^{1/2}$$

for all $u \in D(\sqrt{\mathcal{L}})$.

Our main idea for the proof of this result is to use the maximal regularity for the non-autonomous evolution equation

$$\begin{cases} \frac{\partial u(t)}{\partial t} + L(t) u(t) &= f(t), \quad t \in (0, \tau] \\ u(0) &= u_0. \end{cases} \quad (\text{P})$$

This maximal regularity was proved in an abstract setting by B. Haak and E.M. Ouhabaz [14] under the assumption that the form \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$. It is also proved there that the maximal L^p -regularity holds if there exists a non-decreasing function $\omega : [0, \tau] \rightarrow [0, \infty)$ such that for $u, v \in V$

$$|\mathbf{a}(t, u, v) - \mathbf{a}(s, u, v)| \leq \omega(|t-s|) \|u\|_V \|v\|_V$$

with

$$\int_0^\tau \frac{\omega(t)}{t^{\frac{3}{2}}} dt < \infty. \quad (1.4)$$

See also M. Achache and E.M. Ouhabaz [1] and the references there for an account on recent development on this topic.

The idea of using the maximal regularity in the proof of Theorem 1.2 lies in the fact that we have a precise description of the domain of the maximal accretive operator \mathcal{L} . Then with the help of imaginary powers (or a holomorphic functional calculus) of \mathcal{L} we can

appeal to results on interpolation spaces which in turn give the description of $D(\sqrt{\mathcal{L}})$. Note that the proof of Theorem 1.1 in [5] is very different and it is based on the first order approach initiated by A. McIntosh and his collaborators (see e.g., A. Axelsson, S. Keith and A. McIntosh [6]). One may wonder whether the (piecewise) regularity required in Theorem 1.2 can be removed by adapting the proof in [5] to parabolic operators on domains. This is not known and remains to be done.

Our approach is flexible and applies without any additional effort to other situations such as operators with lower order terms, degenerate operators, systems and operators on weighted spaces. For clarity of exposition we do not search for generality and we keep the setting described above. Instead, we consider another problem which was not studied before in the literature. We study the problem of the square root of \mathcal{L} on $L^p(0, \tau, L^2(\Omega))$ for $p \neq 2$. For this we shall need the following slightly stronger condition than (1.4)

$$\int_0^\tau \frac{\omega(t)}{t^{1+\beta}} dt < \infty \quad (1.5)$$

for some $\beta > \frac{1}{2}$. Clearly, (1.5) is satisfied if the coefficients a_{kl} are C^α in t for some $\alpha > \frac{1}{2}$. We prove the following result.

Theorem 1.3. *Suppose (1.1), (1.2) and (1.5) and let $p \in (1, \infty)$ with $p \neq 2$. There exists a realization of the parabolic operator $\mathcal{L} := \partial + L$ that is maximal accretive on $L^p(0, \tau, L^2(\Omega))$ and satisfies the Kato square root property*

$$\|\sqrt{\mathcal{L}}u\|_{L^p(0, \tau, L^2(\Omega))} \approx \|u\|_{W^{\frac{1}{2}, p}(0, \tau, L^2(\Omega))} + \|\nabla_x u\|_{L^p(0, \tau, L^2(\Omega))}.$$

If $p \in (1, 2)$, the domain of $\sqrt{\mathcal{L}}$ coincides with $W^{\frac{1}{2}, p}(0, \tau, L^2(\Omega)) \cap L^p(0, \tau, V)$. If $p \in (2, \infty)$ this domain coincides with $\{u \in W^{\frac{1}{2}, p}(0, \tau, L^2(\Omega)) \cap L^p(0, \tau, V), u(0) = 0\}$.

The ideas in the proof are similar to the case of $p = 2$ in the sense that we use the maximal regularity of (P) and estimates for imaginary powers ∂^{is} , L^{is} and $(\nu + \mathcal{L})^{is}$ for some constant $\nu \geq 0$. While for $p = 2$, the boundedness of \mathcal{L}^{is} follows from the accretivity of the operator \mathcal{L} on the Hilbert space $L^2(0, \tau, L^2(\Omega))$, the situation for $p \neq 2$ requires some additional work. In order to prove the boundedness of $(\nu + \mathcal{L})^{is}$ we use a perturbation result for the holomorphic functional calculus due to J. Prüss and G. Simonett [24]. The regularity condition (1.5) will be used both to ensure the maximal L^p -regularity and to prove a commutator estimate in order to apply the perturbation theorem in [24]. As a result, we prove that the maximal accretive operator $\nu + \mathcal{L}$ has a bounded holomorphic functional calculus on $L^p(0, \tau, L^2(\Omega))$ for all $p \in (1, \infty)$. This latter result uses only the maximal regularity through the condition (1.5) and not the square root property (1.2). Theorem 1.3 shows that the Kato square root property for the parabolic operator \mathcal{L} holds beyond the Hilbert space setting $L^2(0, \tau, L^2(\Omega))$. A natural question arises whether one might prove a similar result on $L^p(0, \tau, L^r(\Omega))$ for some (or all) $r \neq 2$. We prove such a result for time independent coefficients. The general case is more complicated and remains open unless the coefficients are smooth with respect to the space variable. See the last section of the paper.

Throughout the paper we use $\|\cdot\|_E$ to denote the norm of a given Banach space E . All inessential constants are often denoted by C, C', \dots , the notation $A \approx B$ means that there exists a constant $C > 0$ such that $\frac{1}{C}A \leq B \leq CA$.

Acknowledgements. The author would like to thank Moritz Egert and Sylvie Monniaux for stimulating discussions.

This research is partly supported by the ANR project RAGE, ANR-18-CE-0012-01.

2 Proof of Theorem 1.2

We start by recalling the following maximal regularity result from [14] (Theorem 2 and Corollary 4). It is proved there in an abstract setting of time dependent forms having the same domain. We state it here for the case of elliptic operators as defined in the introduction, so we assume throughout this section that the ellipticity condition (1.1) is satisfied.

Theorem 2.1. *1) Suppose that \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$ and that (1.2) holds. Then the Cauchy problem (P) has maximal L^2 -regularity in $L^2(\Omega)$ for any given $u_0 \in V$. In addition, there exists a positive constant C such that*

$$\|u\|_2 + \left\| \frac{\partial u}{\partial t} \right\|_2 + \|L(\cdot)u(\cdot)\|_2 \leq C [\|f\|_2 + \|u_0\|_V]. \quad (2.1)$$

2) Suppose (1.4). Then (P) with $u_0 = 0$ has maximal L^p -regularity in $L^2(\Omega)$ for all $p \in (1, \infty)$. If in addition ω satisfies the p -Dini condition

$$\int_0^\tau \left(\frac{\omega(t)}{t} \right)^p dt < \infty, \quad (2.2)$$

then (P) has maximal L^p -regularity for all $u_0 \in (L^2(\Omega), D(L(0)))_{1-\frac{1}{p}, p}$.

There exists a positive constant C such that

$$\|u\|_p + \left\| \frac{\partial u}{\partial t} \right\|_p + \|L(\cdot)u(\cdot)\|_p \leq C \left[\|f\|_p + \|u_0\|_{(L^2(\Omega), D(L(0)))_{1-\frac{1}{p}, p}} \right]. \quad (2.3)$$

Recall that (P) has maximal L^p -regularity in $L^2(\Omega)$ if for every $f \in L^p(0, \tau, L^2(\Omega))$ there exists a unique $u \in W^{1,p}(0, \tau, L^2(\Omega))$, $u(t) \in D(L(t))$ for a.e. $t \in (0, \tau)$ and u satisfies (P) for a.e. $t \in (0, \tau)$. We recall that $(L^2(\Omega), D(L(0)))_{1-\frac{1}{p}, p}$ is the real interpolation space and the L^p -norm in the apriori estimates (2.1) and (2.3) is the norm of $L^p(0, \tau, L^2(\Omega))$.

Let us also mention that the maximal L^2 -regularity holds under the slightly weaker regularity property that the map $t \mapsto L(t)$ is piecewise in $H^{\frac{1}{2}}(0, \tau, \mathcal{B}(V, V'))$, where V' is the dual space of V . This is proved in [1] in an abstract setting. Whether the maximal regularity holds for elliptic operators with measurable coefficients in the t -variable (and in the x -variable as we do here) is an important open problem. An example of a family of forms $b(t, \cdot, \cdot)$ such that $t \mapsto b(t, u, v)$ is $C^{\frac{1}{2}}$ in $(0, \tau)$ but the corresponding family of operators does not have the maximal regularity is given in [16]. Note however that these

are not differential operators.

We shall apply the previous theorem in the case where $u(0) = 0$. In this case, we have maximal L^p -regularity for every $p \in (1, \infty)$ provided \mathbf{a} satisfies (1.4). If \mathbf{a} is discontinuous, we assume that it is piecewise C^α for some $\alpha > \frac{1}{2}$ and in addition (1.2) holds. For general forms, the condition $D\left(\sqrt{L(t)}\right) = V$ cannot be removed if \mathbf{a} has (at least) one jump, see [11].

Set $\mathcal{H} = L^2(0, \tau, L^2(\Omega))$ and define $\partial = \frac{\partial}{\partial t}$ with domain

$$D(\partial) = {}_0H^1 := \{u \in H^1(0, \tau, L^2(\Omega)), u(0) = 0\}.$$

Define also the operator L by $(Lu)(t) = L(t)u(t)$ with domain

$$D(L) = \{u \in L^2(0, \tau, L^2(\Omega)), u(t) \in D(L(t)) \text{ a.e. } t \text{ and } L(\cdot)u(\cdot) \in \mathcal{H}\}.$$

Lemma 2.2. *Suppose either (1.4) or \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$ and (1.2) holds. Define the parabolic operator*

$$\mathcal{L} = \partial + L \quad \text{with domain} \quad D(\mathcal{L}) = {}_0H^1 \cap D(L).$$

Then \mathcal{L} is invertible, maximal accretive and has dense domain. The operators $\partial\mathcal{L}^{-1}$ and $L\mathcal{L}^{-1}$ are bounded on \mathcal{H} .

Proof. Integration by parts shows that ∂ is accretive. Then \mathcal{L} is accretive as the sum of two accretive operators. It is invertible on \mathcal{H} by Theorem 2.1. The fact that $\partial\mathcal{L}^{-1}$ and $L\mathcal{L}^{-1}$ are bounded operators on \mathcal{H} is a consequence of the a priori estimate (2.1). A standard duality argument shows that \mathcal{L} is densely defined. \square

Next, for a given $f \in \mathcal{H}$, then $u(t) := \int_0^t f(s) ds$ satisfies $u \in D(\partial)$ and $\partial u = f$. Therefore ∂ is invertible and it is maximal accretive. In particular, this allows us to define its square root $\sqrt{\partial}$ as a maximal accretive operator. Similarly, L is maximal accretive since one checks that $((I + L)^{-1}u)(t) = (I + L(t))^{-1}u(t)$. Therefore, \sqrt{L} is also well defined.

Lemma 2.3. *Suppose either (1.4) or \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$ and (1.2) holds. We have*

$$\|\sqrt{\partial}u\|_{\mathcal{H}} + \|\sqrt{L}u\|_{\mathcal{H}} \leq C\|\sqrt{\mathcal{L}}u\|_{\mathcal{H}} \quad (2.4)$$

for all $u \in D(\sqrt{\mathcal{L}})$. In particular, $D(\sqrt{\mathcal{L}}) \subset D(\sqrt{\partial}) \cap D(\sqrt{L})$.

Proof. Since ∂ and L are maximal accretive it is well known (see e.g. [17]) that they have bounded imaginary powers

$$\|\partial^{is}\|_{\mathcal{B}(\mathcal{H})} \leq e^{\frac{\pi}{2}|s|} \quad \text{and} \quad \|L^{is}\|_{\mathcal{B}(\mathcal{H})} \leq e^{\frac{\pi}{2}|s|}, \quad s \in \mathbb{R}. \quad (2.5)$$

For the same reason, \mathcal{L} also satisfies

$$\|\mathcal{L}^{is}\|_{\mathcal{B}(\mathcal{H})} \leq e^{\frac{\pi}{2}|s|}, \quad s \in \mathbb{R}. \quad (2.6)$$

Define $T(z) := \partial^z \mathcal{L}^{-z}$. Then for $z = is$ with $s \in \mathbb{R}$, it follows from (2.5) and (2.6) that $T(is)$ is bounded on \mathcal{H} with norm bounded by $e^{\pi|s|}$. Using Lemma 2.2, (2.5) and (2.6) we see that $T(1 + is)$ is also bounded on \mathcal{H} with norm bounded by $C e^{\pi|s|}$. This implies that $\partial^{1/2} \mathcal{L}^{-1/2}$ is a bounded operator on \mathcal{H} . Applying the same reasoning with L in place of ∂ shows that $L^{1/2} \mathcal{L}^{-1/2}$ is also bounded on \mathcal{H} . This proves the lemma. \square

Lemma 2.4. *Suppose either (1.4) or \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$ and (1.2) holds. Then there exists a constant $c > 0$ such that*

$$c\|\sqrt{\mathcal{L}}u\|_{\mathcal{H}} \leq \|\sqrt{\partial}u\|_{\mathcal{H}} + \|\sqrt{L}u\|_{\mathcal{H}} \quad (2.7)$$

for all $u \in D(\sqrt{L}) \cap D(\sqrt{\partial})$. In particular, $D(\sqrt{\partial}) \cap D(\sqrt{L}) \subset D(\sqrt{\mathcal{L}})$.

Proof. The proof uses a classical duality argument.

Firstly, one checks easily that the adjoint of ∂ is given by

$$\partial^*v(t) = -\frac{\partial v(t)}{\partial t}, \quad D(\partial^*) = \{v \in H^1(0, \tau, L^2(\Omega)), v(\tau) = 0\}.$$

The adjoint operator L^* is defined similarly to L with $L(t)$ replaced by $L(t)^*$, i.e., $A(x, t)$ is replaced by its adjoint $A^*(x, t)$. On the other hand it is clear that the maximal regularity given by Theorem 2.1 holds for the retrograde problem

$$\begin{cases} -\frac{\partial v(t)}{\partial t} + L(t)^*v(t) &= f(t), \quad t \in (0, \tau] \\ v(\tau) &= 0. \end{cases}$$

Using this we see as above that the operator $\partial^* + L^*$, defined on the intersection of the domains, is invertible and it is maximal accretive. It turns out that this operator is the adjoint of \mathcal{L} . Using the same proof as before, Lemma 2.3 applied to \mathcal{L}^* gives

$$\|\sqrt{\partial^*}v\|_{\mathcal{H}} + \|\sqrt{L^*}v\|_{\mathcal{H}} \leq C\|\sqrt{\mathcal{L}^*}v\|_{\mathcal{H}} \quad (2.8)$$

for all $v \in D(\sqrt{\mathcal{L}^*})$.

Let $u \in D(\mathcal{L})$ and $v \in D(\sqrt{\mathcal{L}^*})$. Then

$$\begin{aligned} \left| (\sqrt{\mathcal{L}}u, \sqrt{\mathcal{L}^*}v)_{\mathcal{H}} \right| &= |(\mathcal{L}u, v)_{\mathcal{H}}| \\ &= \left| (\sqrt{\partial}u, \sqrt{\partial^*}v)_{\mathcal{H}} + (\sqrt{L}u, \sqrt{L^*}v)_{\mathcal{H}} \right| \\ &\leq \left(\|\sqrt{\partial}u\|_{\mathcal{H}} + \|\sqrt{L}u\|_{\mathcal{H}} \right) \left(\|\sqrt{\partial^*}v\|_{\mathcal{H}} + \|\sqrt{L^*}v\|_{\mathcal{H}} \right) \\ &\leq C \left(\|\sqrt{\partial}u\|_{\mathcal{H}} + \|\sqrt{L}u\|_{\mathcal{H}} \right) \left\| \sqrt{\mathcal{L}^*}v \right\|_{\mathcal{H}}. \end{aligned}$$

Note that we have used (2.8) in the last inequality. Since $\sqrt{\mathcal{L}^*}$ is invertible its range is \mathcal{H} . Now we take the supremum over $\left\| \sqrt{\mathcal{L}^*}v \right\|_{\mathcal{H}} \leq 1$ and we obtain (2.7) for $u \in D(\mathcal{L})$. Since $D(\mathcal{L})$ is a core for $\sqrt{\mathcal{L}}$ the inequality holds for all $u \in D(\sqrt{\mathcal{L}})$. \square

Proof of Theorem 1.2. Under the sole assumption (1.4) or if \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$ and (1.2) holds we obtain from the previous lemmas that

$$\|\sqrt{\mathcal{L}}u\|_{\mathcal{H}} \approx \|\sqrt{\partial}u\|_{\mathcal{H}} + \|\sqrt{L}u\|_{\mathcal{H}} \quad (2.9)$$

for all $u \in D(\sqrt{\mathcal{L}}) = D(\sqrt{L}) \cap D(\sqrt{\partial})$. On the other hand since the operator ∂ has bounded imaginary powers it follows that $D(\sqrt{\partial})$ coincides with the complex interpolation space $[_0H^1, \mathcal{H}]_{\frac{1}{2}}$. By [19], p. 68, this interpolation space coincides with

$$\left\{ u \in H^{\frac{1}{2}}(0, \tau, L^2(\Omega)), \int_0^\tau \|u(t)\|_{L^2(\Omega)}^2 \frac{dt}{t} < \infty \right\}.$$

In addition, $\|\sqrt{\partial}u\|_{\mathcal{H}}$ is equivalent to $\|u\|_{H^{\frac{1}{2}}(0,\tau,L^2(\Omega))} + \left(\int_0^\tau \|u(t)\|_{L^2(\Omega)}^2 \frac{dt}{t}\right)^{1/2}$.¹ As mentioned in the introduction, (1.2) implies that the quantities $\|\sqrt{L(t)}u(t)\|_{L^2(\Omega)} + \|u(t)\|_{L^2(\Omega)}$ and $\|\nabla_x u(t)\|_{L^2(\Omega)} + \|u(t)\|_{L^2(\Omega)}$ are equivalent with constants independent of $t \in (0, \tau)$. Therefore, $\|\sqrt{L}u\|_{\mathcal{H}} + \|u\|_{\mathcal{H}}$ and $\|\nabla_x u\|_{\mathcal{H}} + \|u\|_{\mathcal{H}}$ are equivalent. We use this in (2.9) to obtain

$$\|\sqrt{\mathcal{L}}u\|_{\mathcal{H}} + \|u\|_{\mathcal{H}} \approx \|\sqrt{\partial}u\|_{\mathcal{H}} + \|\nabla_x u\|_{\mathcal{H}} + \|u\|_{\mathcal{H}}.$$

From this and the fact that the operators ∂ and $\sqrt{\mathcal{L}}$ are invertible (cf. Lemma 2.2) we obtain the theorem. \square

Remark 2.5. 1-*The proofs of Lemmas 2.3 and 2.4 do not use any specific property of the differential operators $L(t)$. These lemmas are valid in an abstract setting of operators $L(t)$ which are associated with a family of sesquilinear forms*

$$\mathbf{a} : (0, \tau) \times V \times V \rightarrow \mathbb{C}$$

which are quasi-coercive and bounded with uniform constants in t . Here V is a Hilbert space that is densely and continuously embedded into another given Hilbert space H . We define ∂, L and \mathcal{L} as before. Under the sole assumption (1.4) we obtain $D(\sqrt{\mathcal{L}}) = D(\sqrt{L}) \cap D(\sqrt{\partial})$ and

$$\|\sqrt{\mathcal{L}}u\|_{L^2(0,\tau,H)} \approx \|\sqrt{\partial}u\|_{L^2(0,\tau,H)} + \|\sqrt{L}u\|_{L^2(0,\tau,H)}.$$

If \mathbf{a} is piecewise C^α for some $\alpha > \frac{1}{2}$, we assume in addition that (1.2) holds and we obtain the same conclusion.

2- *The ideas used in this section can also be used to describe the domain of any fractional power $D(\mathcal{L}^\alpha)$ for $\alpha \in (0, 1)$.*

3 $L^p(L^2)$ -estimates

During the proofs of the previous section we used the maximal L^2 -regularity given by Theorem 2.1. We take advantage that this latter theorem gives also maximal L^p -regularity for every $p \in (1, \infty)$. We use this in the proof of $L^p(L^2)$ -estimate of Theorem 1.3.

Throughout this section we take the assumptions of Theorem 1.3, that is, we assume (1.1), (1.2) and (1.5).

Fix $p \in (1, \infty)$ with $p \neq 2$. Define on $L^p(0, \tau, L^2(\Omega))$ the operator $\partial = \frac{\partial}{\partial t}$ with domain

$$D(\partial) = {}_0W^{1,p} := \{u \in W^{1,p}(0, \tau, L^2(\Omega)), u(0) = 0\}.$$

It is well known that ∂ has bounded imaginary powers on $L^p(0, \tau, L^2(\Omega))$ (see e.g. [12]). It is not difficult to prove that ∂ is accretive and invertible. Hence, ∂ is maximal accretive.

As in the previous section, we define L by $(Lu)(t) := L(t)u(t)$ with domain

$$D(L) = \{u \in L^p(0, \tau, L^2(\Omega)), u(t) \in D(L(t)) \text{ a.e. } t \text{ and } L(\cdot)u(\cdot) \in L^p(0, \tau, L^2(\Omega))\}.$$

¹Remember that ∂ is invertible, hence the graph norm of $\sqrt{\partial}$ equivalent to $\|\sqrt{\partial}u\|_{\mathcal{H}}$.

Then L is maximal accretive. Since for fixed $t \in (0, \tau)$,

$$\|L(t)^{is}\|_{\mathcal{B}(L^2(\Omega))} \leq e^{\frac{\pi}{2}|s|}$$

and $(L^{is}u)(t) = L(t)^{is}u(t)$,² it follows that the operator L has bounded imaginary powers on $L^p(0, \tau, L^2(\Omega))$. We define $\mathcal{L} = \partial + L$ on the intersection of the domains. It follows from Theorem 2.1 that the operator \mathcal{L} is invertible. In particular, it is maximal accretive. In contrast to the Hilbert space setting of Theorem 1.2, the boundedness of imaginary powers of \mathcal{L} is not a consequence of maximal accretivity. So we have to use a different argument.

Proposition 3.1. *There exists a $\nu \geq 0$ such that the operator $\mathcal{L} + \nu$ has a bounded holomorphic functional calculus on $L^p(0, \tau, L^2(\Omega))$. In particular, $\mathcal{L} + \nu$ has bounded imaginary powers.*

The proof is based on the following perturbation theorem (see Corollary 3.2 in [24]).

Theorem 3.2. *Let A and B be two operators having holomorphic functional calculi with angles ϕ_A and ϕ_B on a Banach space X . Suppose that $0 \in \rho(A)$, B is \mathcal{R} -sectorial and $\phi_A + \phi_B < \pi$. Suppose in addition that for some $0 \leq \alpha < \beta < 1$ the Labbas-Terreni commutator estimate*

$$\|A(\lambda + A)^{-1} [A^{-1}(\mu + B)^{-1} - (\mu + B)^{-1}A^{-1}]\|_{\mathcal{B}(X)} \leq C|\lambda|^{\alpha-1}|\mu|^{-\beta-1} \quad (3.1)$$

holds for all λ and μ with $|\arg(\lambda)| < \pi - \phi_A$ and $|\arg(\mu)| < \pi - \phi_B$. Then there exists a $\nu \geq 0$ such that $\nu + A + B$ has a bounded holomorphic functional calculus on X .

Proof of Proposition 3.1. The operator $\mathcal{L} + \epsilon$ is the sum of (non-commuting) operators $B = \partial$ and $A = L + \epsilon$. Each of these operators has a bounded holomorphic functional calculus on $L^p(0, \tau, L^2(\Omega))$ with angles $\phi_\partial = \frac{\pi}{2} + \epsilon$ and $\phi_L < \frac{\pi}{2}$, respectively. Next, the functional calculus is \mathcal{R} -bounded (for holomorphic functions with modulus ≤ 1). This follows from [15], Theorem 10.3.4 (3) in combination with Proposition 7.5.3 (which shows that $L^p(0, \tau, L^2(\Omega))$ has Pisier's contraction principle since this is the case for the Hilbert space $L^2(\Omega)$). The role of $\epsilon > 0$ above is only to guarantee that $L + \epsilon$ is invertible. For simplicity we forget ϵ and keep in mind that L has to be replaced by $L + \epsilon$ in the sequel. We claim that (3.1) is satisfied with $\alpha = \frac{1}{2}$ and β as in (1.5). Once this is proved we can apply Theorem 3.2 to obtain the proposition.

Let $f \in L^p(0, \tau, L^2(\Omega))$ and set

$$I(t) := \|L(t)(\lambda + L(t))^{-1} [L(t)^{-1}(\mu + \partial)^{-1} - (\mu + \partial)^{-1}L(t)^{-1}f(t)]\|_{L^2(\Omega)}.$$

Since

$$(\mu + \partial)^{-1}f(t) = \int_0^t e^{-\mu(t-s)}f(s) ds \quad (3.2)$$

²one starts from the resolvent formula $((\lambda I + L)^{-1}u)(t) = (\lambda I + L(t))^{-1}u(t)$ and then by integration along an appropriate contour to define the holomorphic functional calculus one obtains such a formula.

and $\operatorname{Re}(\mu) \approx |\mu|$, we have

$$\begin{aligned} I(t) &= \left\| \int_0^t e^{-\mu(t-s)} L(t)(\lambda + L(t))^{-1} [L(t)^{-1} - L(s)^{-1}] f(s) ds \right\|_{L^2(\Omega)} \\ &\leq \int_0^t e^{-c|\mu|(t-s)} \|L(t)(\lambda + L(t))^{-1} [L(t)^{-1} - L(s)^{-1}] f(s)\|_{L^2(\Omega)} ds \end{aligned}$$

for some constant $c > 0$. Now we argue exactly as in [22], p. 1675 to obtain

$$\|L(t)(\lambda + L(t))^{-1} [L(t)^{-1} - L(s)^{-1}] f(s)\|_{L^2(\Omega)} \leq \frac{C}{|\lambda|^{1/2}} \omega(|t-s|) \|f(s)\|_{L^2(\Omega)}.$$

This gives

$$I(t) \leq \frac{C}{|\lambda|^{1/2}} \int_0^t e^{-c|\mu|(t-s)} \omega(|t-s|) \|f(s)\|_{L^2(\Omega)} ds. \quad (3.3)$$

The term $\int_0^t e^{-c|\mu|(t-s)} \omega(|t-s|) \|f(s)\|_{L^2(\Omega)} ds$ can be seen as an operator (acting on $\|f(s)\|_{L^2(\Omega)}$) with kernel

$$K(t, s) = \chi_{(0,t)}(s) e^{-c|\mu|(t-s)} \omega(t-s).$$

Using the assumption (1.5) we have for all $t \in (0, \tau)$

$$\begin{aligned} \int_0^\tau K(t, s) ds &= \frac{1}{|\mu|^{\beta+1}} \int_0^t e^{-c|\mu|(t-s)} (|\mu|(t-s))^{\beta+1} \frac{\omega(t-s)}{(t-s)^{\beta+1}} ds \\ &\leq \frac{C}{|\mu|^{\beta+1}} \int_0^\tau \frac{\omega(r)}{r^{1+\beta}} dr \leq \frac{C'}{|\mu|^{\beta+1}}. \end{aligned}$$

Similarly,

$$\int_0^\tau K(t, s) dt \leq \frac{C'}{|\mu|^{\beta+1}},$$

uniformly in $s \in (0, \tau)$. This implies that the operator with kernel $K(t, s)$ is bounded on $L^p(0, \tau)$ with norm bounded by $\frac{C'}{|\mu|^{\beta+1}}$. It follows from (3.3) that the operator $L(\lambda + L)^{-1} [L^{-1}(\mu + \partial)^{-1} - (\mu + \partial)^{-1} L^{-1}]$ is bounded on $L^p(0, \tau, L^2(\Omega))$ with norm bounded by $\frac{C}{|\lambda|^{1/2} |\mu|^{1+\beta}}$. This is exactly the condition (3.1). \square

We go back to the proof of Theorem 1.3. Since ∂ has imaginary powers, we have $D(\sqrt{\partial}) = [{}_0W^{1,p}, L^p(0, \tau, L^2(\Omega))]_{\frac{1}{2}}$ with equivalent norms. It follows from [2], Theorem 4.7.1 or [10], p. 41 that $[{}_0W^{1,p}, L^p(0, \tau, L^2(\Omega))]_{\frac{1}{2}}$ coincides with $W^{\frac{1}{2},p}(0, \tau, L^2(\Omega))$ if $p < 2$ and with ${}_0W^{\frac{1}{2},p}(0, \tau, L^2(\Omega))$ if $p > 2$.³ Hence

$$\|\sqrt{\partial} u\|_{L^p(0,\tau,L^2(\Omega))} \approx \|u\|_{W^{\frac{1}{2},p}(0,\tau,L^2(\Omega))}. \quad (3.4)$$

³This is stated in [2] and [10] on the interval $(0, \infty)$ instead of $(0, \tau)$. One either uses a similar retraction and coretraction argument used their to deal directly $(0, \tau)$ or use a cut-off argument around the point τ . See also [7] for interpolation results in the scalar case.

By Proposition 3.1, $(\mathcal{L} + \nu)^{is}$ is bounded on $L^p(0, \tau, L^2(\Omega))$, thus we can repeat the proof of Lemma 2.3 and obtain

$$\|\sqrt{\partial} u\|_{L^p(0, \tau, L^2(\Omega))} + \|\sqrt{L} u\|_{L^p(0, \tau, L^2(\Omega))} \leq C \|\sqrt{\mathcal{L} + \nu} u\|_{L^p(0, \tau, L^2(\Omega))}.$$

On the other hand since the operator \mathcal{L} is invertible by Theorem 2.1, we can remove the constant ν in the previous inequality and obtain

$$\|\sqrt{\partial} u\|_{L^p(0, \tau, L^2(\Omega))} + \|\sqrt{L} u\|_{L^p(0, \tau, L^2(\Omega))} \leq C' \|\sqrt{\mathcal{L}} u\|_{L^p(0, \tau, L^2(\Omega))}.$$

Using the same estimate for the adjoint operator on $L^{p'}(0, \tau, L^2(\Omega))$ we argue by duality as in Lemma 2.4 and obtain the the reverse inequality. Therefore,

$$\|\sqrt{\mathcal{L}} u\|_{L^p(0, \tau, L^2(\Omega))} \approx \|\sqrt{\partial} u\|_{L^p(0, \tau, L^2(\Omega))} + \|\sqrt{L} u\|_{L^p(0, \tau, L^2(\Omega))} \quad (3.5)$$

for all $u \in D(\sqrt{\mathcal{L}}) = D(\sqrt{\partial}) \cap D(\sqrt{L})$. Using (3.4) it follows that

$$\|\sqrt{\mathcal{L}} u\|_{L^p(0, \tau, L^2(\Omega))} \approx \|u\|_{W^{\frac{1}{2}, p}(0, \tau, L^2(\Omega))} + \|\sqrt{L} u\|_{L^p(0, \tau, L^2(\Omega))} \quad (3.6)$$

for all $u \in D(\sqrt{\mathcal{L}}) = [{}_0W^{1, p}(0, \tau, L^2(\Omega)), L^p(0, \tau, L^2(\Omega))]_{\frac{1}{2}} \cap D(\sqrt{L})$. Thus we have proved Theorem 1.3.

As we already mentioned before, the method we employed in this paper can be used in other circumstances. For example, the above $L^p(L^2)$ -estimate can be proved for elliptic operators with lower order terms, some degenerate operators as well as parabolic systems. We do not write the details since they are essentially a simple repetition of what is presented above.

4 $L^p(L^r)$ -estimates

In this section we address the question whether the previous results can be extended to $L^p(0, \tau, L^r(\Omega))$ for $r \neq 2$. When reproducing the arguments of the previous sections we face two problems. The first is to have maximal L^p -regularity in $L^r(\Omega)$ since Theorem 2.1 is specific to the $L^2(\Omega)$ case. The second one is to have boundedness of imaginary powers of \mathcal{L} (or $\nu + \mathcal{L}$ for some constant $\nu \geq 0$). The arguments in the proof of Proposition 3.1 use the sesquilinear form setting in order to check (3.1). Note that there are results on maximal regularity outside the Hilbert space (and hence the sesquilinear form) setting. However these results assume the domains of $L(t)$ to be constant. See [3] and the references there. In order to guarantee that the operators $L(t)$ have the same domain on $L^r(\Omega)$ the natural thing to do is to compute this domain and show that it coincides with some Sobolev space. In order to do so one needs some regularity in the x -variable for $A(x, t)$ and more regularity for Ω . In order to stay with non-smooth coefficients in the x -variable we shall concentrate on the case $A(x, t) = A(x)$. We also assume that our elliptic operator is subject to the Dirichlet boundary conditions. With the same notation as before, we have

Theorem 4.1. *Suppose (1.1), (1.2) and (1.5). Suppose that $A(x, t) = A(x)$ has real-valued coefficients. Let $p \in (1, \infty)$ and denote by p' its conjugate. Then for $r \in [\min(p, p'), \max(p, p')]$,*

$$\|\sqrt{\partial} u\|_{L^p(0, \tau, L^r(\Omega))} + \|\sqrt{L} u\|_{L^p(0, \tau, L^r(\Omega))} \approx \|\sqrt{\mathcal{L}} u\|_{L^p(0, \tau, L^r(\Omega))} \quad (4.1)$$

for all $u \in D(\sqrt{\mathcal{L}})$. In addition, for $r \in [\min(p, p'), 2]$, there exists a constant C such

$$\|\sqrt{\partial} u\|_{L^p(0, \tau, L^r(\Omega))} + \|\nabla u\|_{L^p(0, \tau, L^r(\Omega))} \leq C \|\sqrt{\mathcal{L}} u\|_{L^p(0, \tau, L^r(\Omega))}. \quad (4.2)$$

Proof. First since L has real-coefficients and subject to the Dirichlet boundary conditions, the semigroup e^{-tL} is sub-Markovian (cf. [21], Chapter 4). Therefore, by [18], L has maximal L^p -regularity on $L^r(\Omega)$ for all $p, r \in (1, \infty)$. In particular, the operator $\mathcal{L} = \partial + L$ defined on the intersection of the domains ${}_0W^{1,2}(0, \tau, L^r(\Omega))$ and $D(L)$ is maximal accretive (note that both ∂ and L are accretive on $L^p(0, \tau, L^r(\Omega))$). On the other hand, the two maximal accretive operators ∂ and L are generators of positive semigroups. For positivity of e^{-tL} see [21], Chapter 4 and for $e^{-t\partial}$ this follows readily from the positivity of its resolvent (see (3.2)). This and the Trotter product formula give the positivity of the contraction semigroup $e^{-t\mathcal{L}}$ on $L^p(0, \tau, L^r(\Omega))$. Since for $r = p$, $L^p(0, \tau, L^p(\Omega)) \simeq L^p(\Omega \times (0, \tau))$ we may use the transference method [9] to obtain that \mathcal{L} has a bounded holomorphic functional calculus on $L^p(0, \tau, L^p(\Omega))$ (with angle $\phi > \frac{\pi}{2}$). This is also true for $\nu + \mathcal{L}$ for any $\nu \geq 0$. Using this and Proposition 3.1 it follows by interpolation that $\nu + \mathcal{L}$ has a bounded holomorphic functional calculus on $L^p(0, \tau, L^r(\Omega))$ for $r \in [p, 2]$ or $[2, p]$. What we did here for \mathcal{L} is also valid for $\mathcal{L}^* = \partial^* + L^*$ by the same arguments. This gives that $\nu + \mathcal{L}$ has a bounded holomorphic functional calculus on $L^p(0, \tau, L^r(\Omega))$ for all $p \in (1, \infty)$ and $r \in [\min(p, p'), \max(p, p')]$. In particular, the imaginary powers $(\nu + \mathcal{L})^{is}$ are bounded on these spaces. The rest of the proof of (4.1) is exactly the same as for Theorem 1.3.

Suppose now that $r \in [\min(p, p'), 2]$. Then the Riesz transform $\nabla L^{-\frac{1}{2}}$ is bounded on $L^r(\Omega)$ (see [21], Section 7.7). This gives $\|\nabla f\|_{L^r(\Omega)} \leq C \|\sqrt{L} f\|_{L^r(\Omega)}$. Thus, (4.2) follows from (4.1). \square

Remark 4.2. *The idea of using the transference method on $L^p(0, \tau, L^p(\Omega))$ was already used in [23] in the context of parabolic Schrödinger operators.*

References

- [1] M. Achache and E.M. Ouhabaz, Lions' maximal regularity problem with $H^{\frac{1}{2}}$ -regularity in time. *J. Differential Equations* 266 (2019), no. 6, 3654-3678.
- [2] H. Amann, Anisotropic Function spaces and Maximal Regularity for Parabolic Problems. Part 1. Function spaces. *Jindrich Necas Center for Mathematical Modeling Lecture Notes*, 6. Matfyzpress, Prague, 2009. vi+141 pp. ISBN: 978-80-7378-089-0.
- [3] W. Arendt, R. Chill, S. Fornaro, C. Poupaud, L^p -maximal regularity for non-autonomous evolution equations. *J. Differential Equations* 237 no 1 (2007) 1-26.
- [4] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh and Ph. Tchamitchian, The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n . *Ann. of Math.* (2) 156 (2002), no. 2, 633-654.

- [5] P. Auscher, M. Egert and K. Nyström, Boundary value problems for parabolic systems via first order approach. *J. Eur. Math. Soc. (JEMS)*, to appear.
- [6] A. Axelsson, S. Keith and A. McIntosh, Quadratic estimates and functional calculi of perturbed Dirac operators. *Invent. Math.* 163 (2006), no. 3, 455-497.
- [7] S. Bechtel and M. Egert, Interpolation theory for Sobolev functions with partially vanishing trace on irregular open sets. *J. Fourier Anal. Appl.* 25 (2019), no. 5, 2733-2781.
- [8] S. Bechtel, M. Egert and R. Haller-Dintelmann, The Kato square root problem on locally uniform domains. Preprint 2019, available on: <https://arxiv.org/pdf/1902.03957.pdf>.
- [9] R.R. Coifman, G. Weiss, Transference Methods in Analysis, Conference Board of the Mathematical Sciences, Regional Conference Series in Mathematics 31, AMS, Providence (1977).
- [10] R. Denk and M. Kaip, General Parabolic Mixed Order Systems in L_p and Applications. *Operator Theory: Advances and Applications, 239*. Birkhäuser/Springer, Cham, 2013. viii+250 pp. ISBN: 978-3-319-01999-4.
- [11] D. Dier, Non-autonomous Cauchy Problems Governed by Forms: Maximal Regularity and Invariance. PhD Thesis. Universität Ulm, 2014.
- [12] G. Dore and A. Venni, On the closedness of the sum of two closed operators. *Math. Z.* 196 (1987), no. 2, 189-201.
- [13] M. Egert, R. Haller-Dintelmann and P. Tolksdorf, The Kato square root problem for mixed boundary conditions. *J. Funct. Anal.* 267 (2014), no. 5, 1419-1461.
- [14] B. Haak and E.M. Ouhabaz, Maximal regularity for non-autonomous evolution equations. *Math. Ann.* 363 (2015), no. 3-4, 1117-1145.
- [15] T. Hytönen, J. van Neerven, M. Veraar and L. Weis, Analysis in Banach Spaces. Vol. II. Probabilistic Methods and Operator Theory. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics 67*. Springer, Cham, 2017.
- [16] S. Fackler, J.-L. Lions' problem concerning maximal regularity of equations governed by non-autonomous forms. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 34 (2017), no. 3, 699-709.
- [17] T. Kato, Fractional powers of dissipative operators II. *J. Math. Soc. Japan* 14 (1962) 242-248.
- [18] D. Lamberton, Equations d'évolution linéaires associées à des semi-groupes de contractions dans les espaces L^p . *J. Funct. Anal.* 72 (1987), no. 2, 252-262.

- [19] J. L. Lions and E. Magenes, Non-Homogeneous Boundary Value Problems and Applications: Vol. II. *Die Grundlehren der mathematischen Wissenschaften*, Band 182. Springer-Verlag, New York-Heidelberg, 1972. xi+242 pp.
- [20] K. Nyström, Square functions estimates and the Kato problem for second order parabolic operators in \mathbb{R}^{n+1} . *Advances in Mathematics* 293 (2016), 1-36.
- [21] E. M. Ouhabaz, Analysis of Heat Equations on Domains, *London Math. Soc Monographs 31*, Princeton University Press 2005.
- [22] E.M. Ouhabaz and C. Spina, Maximal regularity for non-autonomous Schrödinger type equations. *J. Differential Equations* 248 (2010), no. 7, 1668-1683.
- [23] E.M. Ouhabaz and C. Spina, Riesz transforms of some parabolic operators. *AMSI International Conference on Harmonic Analysis and Applications*, 115?123, *Proc. Centre Math. Appl. Austral. Nat. Univ.*, 45, *Austral. Nat. Univ.*, Canberra, 2013.
- [24] J. Prüss and G. Simonett, H^∞ -calculus for the sum of non-commuting operators. *Trans. Amer. Math. Soc.* 359 (2007), no. 8, 3549-3565.