



**HAL**  
open science

## **Trophic ecology of suspension-feeding bivalves inhabiting a north-eastern Pacific coastal lagoon: Comparison of different biomarkers**

Margaux Mathieu-Resuge, Edouard Kraffe, Fabienne Le Grand, Andy Boens,  
Antoine Bideau, Salvador E. Lluch-Cota, Ilie S. Racotta, Gauthier Schaal

► **To cite this version:**

Margaux Mathieu-Resuge, Edouard Kraffe, Fabienne Le Grand, Andy Boens, Antoine Bideau, et al.. Trophic ecology of suspension-feeding bivalves inhabiting a north-eastern Pacific coastal lagoon: Comparison of different biomarkers. *Marine Environmental Research*, 2019, 145, pp.155-163. 10.1016/j.marenvres.2019.02.016 . hal-02873900

**HAL Id: hal-02873900**

**<https://hal.science/hal-02873900>**

Submitted on 18 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

---

## Trophic ecology of suspension-feeding bivalves inhabiting a north-eastern Pacific coastal lagoon: Comparison of different biomarkers

Mathieu-Resuge Margaux <sup>1,\*</sup>, Kraffe Edouard <sup>1</sup>, Le Grand Fabienne <sup>3</sup>, Boens Andy <sup>1</sup>, Bideau Antoine <sup>1</sup>, Lluch-Cota Salvador E. <sup>2</sup>, Racotta Ilie S. <sup>2</sup>, Schaal Gauthier <sup>1</sup>

<sup>1</sup> Univ Brest, CNRS, IRD, Ifremer, LEMAR, IUEM, F-29280, Plouzane, France

<sup>2</sup> Centro de Investigaciones Biológicas Del Noroeste (CIBNOR), La Paz, BCS, Mexico

\* Corresponding author : Margaux Mathieu-Resuge, email address : [m.mathieuresuge@hotmail.fr](mailto:m.mathieuresuge@hotmail.fr)

---

### Abstract :

Understanding the nature and origins of food sources supporting coastal lagoon-inhabiting organisms is necessary to evaluate the ecological status of such ecosystems. The trophic ecology of a bivalve specie *Spondylus crassisquama* was studied in the Ojo de Liebre lagoon (Baja California, Mexico), combining stable isotope (SI), fatty acid (FA) and sterols analyses along a transect under oceanic influences. The second objective of the study was to investigate if sterols compositions give complementary information to those obtained from FA and SI. Temporal and spatial patterns of the three biomarkers in bivalve tissues suggest oceanic inputs at the mouth of the lagoon, while the inner station was characterized by a contribution of local sources including an important role of micro heterotrophs. This study revealed that the association of lipid biomarkers provide higher taxonomic resolution of the spatiotemporal heterogeneity in nutrient sources influencing the trophic functioning of a large coastal lagoons.

### Highlights

► Stable isotopes fatty acids and sterols were used to characterize the diet of bivalves. ► A transect was sampled in a Pacific lagoon with increasing distance from the mouth. ► Bivalves at the entrance of the lagoon relied on oceanic phytoplankton inputs. ► Bivalves in the inner lagoon relied on local microbial food web. ► Lipid biomarkers and SI were complementary to distinguish the food sources origins.

**Keywords** : coastal zone, Ojo de Liebre, suspension-feeders, multi-tracer analyses, fatty acids, stables isotopes, sterols

## 30           **1. Introduction**

31 Coastal areas are highly productive marine ecosystems (Clavier et al., 2014; Nixon, 1982) that  
32 provide a variety of ecosystem services, such as nutrient recycling, biogeochemical cycles  
33 regulation, or food supply (Gattuso et al., 1998; Martínez et al., 2007). Among coastal  
34 ecosystems, lagoons play key ecological and economic roles, providing habitat to many resident  
35 and migratory species and hosting commercial, artisanal and subsistence fisheries that support  
36 local populations. Their high productivity originates from nutrients inputs of both terrestrial and  
37 marine origins (Carlier et al., 2007; Mann, 1982), as well as from local primary productivity,  
38 enhanced by the shallow and enclosed nature of these areas. Understanding the relative  
39 contribution of autochthonous processes versus allochthonous inputs (lagoon vs oceanic  
40 primary production, respectively) for the functioning of coastal lagoons is therefore critical to  
41 understand the dynamics of these ecosystems, and how they are likely to react when facing  
42 anthropogenic disturbances.

43 Suspension-feeding bivalves are major components of coastal lagoons, where they can  
44 dominate benthic communities and play a key role in the regulation of benthic and pelagic  
45 biogeochemical fluxes (Dame, 1993). Because these organisms are primary consumers, they  
46 directly reflect processes occurring at the lowest level of food webs (Marchais et al., 2013).  
47 Furthermore, suspension-feeders can feed on a wide variety of sestonic particles, hence can be  
48 integrative indicators of coastal oceanic versus lagoon autochthonous inputs into coastal  
49 ecosystems (Coma et al., 2001; Kraak et al., 1991; Lefebvre et al., 2009).

50 Since a few decades, trophic studies have evolved from field or laboratory observational  
51 approaches (*i.e.* stomach contents, food choice experiments) to biomarker-based approaches  
52 (BBA). BBA have been applied to a variety of organisms, from primary consumers (Desvillettes et  
53 al., 1997) to marine mammals (Kelly, 2000), as well as at the community level (Alfaro et al.,  
54 2006; Vander Zanden and Rasmussen, 2001). Every BBA is based on several prerequisites, which  
55 need to be met in order to produce robust trophic information. Uncertainties concerning these  
56 prerequisites are common (*i.e.* concerning turnover, metabolic fate of biomarker compounds)  
57 and result in less reliable data, and stress the need to couple different BBA to fully understand  
58 trophic dynamics of natural environments. Among the most commonly used BBA, stable  
59 isotopes (SI) ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ,  $\delta^{34}\text{S}$ ) have been widely and successfully used to characterize the origin  
60 of food sources and the trophic level of consumers (Peterson and Fry, 1987; Post, 2002).  
61 However, SI analysis requires potential food sources to be discriminated in the two or three  
62 dimensional isotopic space, which is not always the case in high diversity systems. Moreover, SI  
63 analysis provides a limited taxonomic resolution of the food sources assimilated concerning  
64 heterogeneous pools of organic matter (*e.g.* suspended particulate organic matter, sedimentary  
65 organic matter). In this context, the combination with others biomarkers can give useful  
66 information to complement identifications of food sources. Fatty acids allow the identification  
67 of some taxa in the diet of consumers due to the existence of specific FA (Dalsgaard et al., 2003;  
68 Parrish et al., 2000). Although primary producers have characteristic FA compositions, most do  
69 not possess unique FAs which can cause limitations for identifications of food sources (Bec et  
70 al., 2010). In addition, in the specific case of coastal benthic food webs, the important number

71 of potential food sources is a problem for distinguishing their contributions to diets (Kelly and  
72 Scheibling, 2012).

73 Sterols are lipid compounds common in all primary producers. Some of them have been  
74 reported to allow the characterization of classes, genus and in some cases, species of primary  
75 producers that contribute to the diet of filter feeders (Napolitano et al., 1993; Soudant et al.,  
76 1998; Volkman, 1986). As for FAs, the different primary producers may display distinct sterols  
77 composition, which a necessary condition for using these compounds as trophic biomarkers.  
78 Moreover, since the capacity to synthesize *de novo* or modify dietary sterols is absent or  
79 considered very low in marine invertebrates (Holden and Patterson, 1991; Kanazawa, 2001),  
80 sterols have to be acquired from their diet. The role of sterols is still poorly understood, even if  
81 they are recognized as important structural components of cell membranes (Giner et al., 2016)  
82 and necessary for growth and larval development (Soudant et al., 1998). Primary producers that  
83 constitute the major part of the diet of bivalves contain an important diversity of sterols, named  
84 phytosterols (Wasowicz and Rudzinska, 2011), and their presence in consumers could be a  
85 reliable complementary biomarkers to identify and distinguish contribution of specific food  
86 sources to their diet.

87 The present study aims to characterize the trophic ecology of a suspension-feeding bivalve,  
88 *Spondylus crassisquama* (Lamarck, 1819) within the Northeastern Pacific coastal lagoon of Ojo  
89 de Liebre (Baja California, Mexico). By sampling this species at two sites, varying in their  
90 distance to the open ocean, and at two seasons, characterized by different environmental  
91 conditions, we test the hypothesis that oceanic inputs induce a spatial structure in the trophic

92 ecology of suspension-feeding bivalves within the lagoon, including a higher reliance towards  
93 oceanic production in individuals living close to the lagoon mouth. For this purpose, a multi-  
94 tracers approach was chosen, combining stable isotopes, fatty acids and sterols composition in  
95 bivalve digestive glands as well as in their potential food sources (particulate organic matter,  
96 sedimentary organic matter, and seagrass detritus). The potential complementarity among  
97 those different biomarkers is also discussed.  
98

99 **2. Material and methods**100 **2.1 Sampling**

101 The lagoon of Ojo de Liebre (Baja California, Mexico) is one of the largest coastal lagoons in the  
102 north-east Pacific ( $\approx 40$  km long,  $\approx 10$  km wide, 446 km<sup>2</sup>). Its topography is characterized by  
103 deep and narrow channels surrounded by large intertidal flats. The tidal range is about 1 to 2.5  
104 meters and currents are more turbulent in channels (maximum 2.5 knots) than intertidal flats  
105 (0.35 to 1.5 knots) (Phleger and Ewing, 1962). Two stations, located on historically fished banks,  
106 characterized by a depth around 7 to 10 m, were studied into the Ojo de Liebre lagoon (Figure  
107 1). Station 1 (Chocolatero) is a bank located close to the mouth of the lagoon (at 6.1 km from  
108 the entrance), while station 2 (Datil) represented the most inner station (15.5 km from the




Figure 1. Location of the both sampling stations in the Ojo de Liebre lagoon (Baja California Sur, Mexico).

109 mouth) and the southern distribution limit of *Spondylus crassisquama* in this bay.

110 Samples were collected in February and August 2016. For each station, 10 similarly sized  
111 individuals were sampled by divers. Potential food sources available for filters feeders were also  
112 sampled. One sedimentary core was sampled at each station. The upper centimeter, the most  
113 likely to be resuspended by currents and therefore available to suspension-feeders, was  
114 collected. Due to the absence of water column vertical stratification into the lagoon (Phleger  
115 and Ewing, 1962), one surface water sample (3L) was collected for each site and each date.  
116 Because of logistical limitations associated to sampling in the lagoon, we could not replicate our  
117 source sampling within each site. Ojo de Liebre contains one of the largest seagrass bed of  
118 *Zostera marina* in the Pacific (Cabello-Pasini et al., 2003) which represents one of the dominant  
119 primary producer at the bay scale. Because seagrass detritus could contribute to the  
120 composition of particulate organic matter (POM) and potentially as a food source for bivalves,  
121 stranded *Z. marina* detritus were collected from the shore at the eastern part of the sampling  
122 area, and considered as proxies of eelgrass detritus within the lagoon. Indeed, the SI  
123 composition of seagrass is known to be little altered during degradation, in particular for  $\delta^{13}\text{C}$   
124 (Fenton and Ritz, 1988).

125 After collection, samples were placed in cooler boxes containing water and transported to the  
126 laboratory (CIBNOR, Unidad Guerrero Negro). Bivalves were dissected and each tissue of  
127 interest (*i.e.* adductor muscle and DG) was sampled, packed into aluminum and stored in liquid  
128 nitrogen until further analysis. For each station, surface water samples were filtered on pre-



129 combusted GF/F filters (6 h, 450°C). For each filter 1.5 L of sea water was filtered. Immediately  
130 after filtration, filters were frozen in liquid nitrogen. Sediment samples were soaked in local  
131 filtered water (GF/F 0.7 µm), and allowed to settle for a few seconds. The supernatant  
132 containing fine organic particles, was then collected and filtered on pre-combusted (6 h, 450°C)  
133 GF/F filters. Two subsamples of suspended particulate organic matter (POM) (one for SI  
134 analyses and one for FA analyses), and four subsamples of sedimentary organic matter (SOM)  
135 (two for SI and two for FA analyses) were taken at each sampling station/date. All samples were  
136 stored in liquid nitrogen during transportation to the laboratory (CIBNOR, La Paz) and then  
137 stored at -80°C until further preparations for analysis.

## 138 2.2 Stable isotopes analysis

139 For SI analyses, muscle samples were freeze dried for 24 h and ground into homogenized fine  
140 powder using a ball mill. Muscle samples were not previously delipidated because bivalve  
141 muscles generally show a very low content of lipids in this tissue (Lorrain et al., 2002), also C/N  
142 ratios were measured and did not exceed 3.5, suggesting delipidation unnecessary (Post et al.,  
143 2007). A minimum of 0.250 mg of powder were then placed in tin capsules (8x5 mm) for  
144 isotopic analyses. For SOM and POM, filters were scraped, and approximately 1.5 mg was  
145 placed in tin capsules.

146 Samples were analyzed by continuous flow on a Flash EA2000 elemental analyzer coupled to a  
147 Delta V Plus mass spectrometer (Thermo Fisher scientific). Reference gas calibration was  
148 conducted using IAEA-600 Caffeine, IAEA-CH-6 Sucrose and IAEA-N-2 Ammonium

149 Sulphate. Analytical precision, based on repeated analyses of an acetanilide working standard  
150 (Thermo scientific), was  $\pm 0.1$  ‰ for both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ . Results are expressed in  $\delta$ , with respect  
151 to international standards (Vienna Pee Dee Belemnite for  $\delta^{13}\text{C}$  and atmospheric nitrogen for  
152  $\delta^{15}\text{N}$ ), following the equation:  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N} = [(R_{\text{sample}}/R_{\text{standard}})-1] \cdot 10^{-3}$  (expressed in ‰) and  
153 where R is  $^{13}\text{C}/^{12}\text{C}$  or  $^{15}\text{N}/^{14}\text{N}$ .

## 154 2.3 Lipid analysis

### 155 2.3.1 Lipid extraction

156 Samples of bivalve DG and *Z. marina* were ground in liquid nitrogen into a fine homogeneous  
157 powder. Lipids were then extracted, between 0.2 and 0.4 g of material was placed in 6 ml of  
158 chloroform/methanol (2:1; v/v) and sonicated at 4°C during 5 min, to ensure complete  
159 extraction of lipids. Lipid extracts were stored at -20°C under nitrogen atmosphere until further  
160 analysis. Lipids from the other potential food sources (POM and SOM) were extracted following  
161 the same method, by resuspending filters directly in chloroform/methanol (2:1; v/v).

### 162 2.3.2 Separation of neutral lipids

163 Lipids from DG were separated into two fractions (neutral and polar lipids) following the  
164 method described by Le Grand et al. (2014). An aliquot (1/6) of total lipid extract was  
165 evaporated under nitrogen flux, recovered with 3 washes of 0.5 ml of chloroform/methanol  
166 (98:2; v/v) and deposited at the top of a silica gel column (40 mm x 4 mm, silica gel 60A 63-200  
167  $\mu\text{m}$  rehydrated with 6%  $\text{H}_2\text{O}$  (70-230 mesh)). NL were eluted with 10 ml of

168 chloroform/methanol (98/2; v/v). NL fractions were collected in 20 ml glass vial containing 2.3  
169  $\mu\text{g}$  of each internal standards (tricosanoic acid (C23:0), and cholestane). After evaporation to  
170 dryness, NL fraction was recovered with 3 washes of 0.5 ml of chloroform/methanol (98:2; v/v)  
171 and separated in two aliquots (1/2), one for FA analysis and the second for sterol analysis.

### 172 *2.3.3 Methylation of FA*

173 The NL fraction of DG, and an aliquot of the total lipid extract from food sources (POM, SOM,  
174 and eelgrass) were evaporated to dryness under nitrogen flux. Acidic transesterification was  
175 done to obtain fatty acid methyl esters (FAME). Briefly, 0.8 ml of  $\text{H}_2\text{SO}_4$ /methanol were added  
176 (3.4 %; v/v) and lipid samples were then vortexed and heated at 100°C for 10 min. After cooling,  
177 0.8 ml of hexane and 1.5 ml of distilled water saturated in hexane were added. Vials were  
178 homogenized and centrifuged at 738 g, 1 min at 20°C. The aqueous phase (without FAME) was  
179 discarded and the organic phase—containing FAME was washed two more times with distilled  
180 water saturated in hexane.

### 181 *2.3.4 Saponification and silylation of sterols*

182 After evaporation to dryness of NL fraction of DG and of an aliquot of the total lipid extract from  
183 food sources (POM, SOM, and eelgrass), 3 ml of KOH/methanol (5.6:94.4; v/v) were added;  
184 samples were homogenized and placed under nitrogen atmosphere. Samples were then heated  
185 at 100°C during 10 min. After cooling at room temperature, 0.8 ml of hexane and 1.5 ml of  
186 distilled water saturated in hexane were added, followed by a homogenization and

187 centrifugation at 738 g during 1 min at 20°C. The aqueous phase was discarded and the organic  
188 phase—containing sterols was washed two more times with distilled water saturated in hexane.

189 Before GC analysis, sterol samples were silylated to improve resolution for gas chromatography.  
190 For that, organic phases obtained after saponification were evaporated to dryness under  
191 nitrogen flux, then 100 µl of reagent (BSTFA:TMCS; 99:1; v/v) and 100 µl of pyridine were  
192 added. Sterols samples were then heated at 70°C during 30 min. After cooling at room  
193 temperature, 0.8 ml of hexane and 1.5 ml of distilled water saturated in hexane were added,  
194 followed by a homogenization and centrifugation at 738 g during 1 min at 20°C. The aqueous  
195 phase was discarded and the organic phase—containing silylated sterols was washed two more  
196 times with distilled water saturated in hexane.

## 197 *2.4 Lipid analysis by Gas Chromatographs*

### 198 *2.4.1 Fatty acid Gas Chromatography analysis*

199 FAME were analyzed in a Varian CP8400 gas chromatograph, programmed in temperature (from  
200 0°C to 150°C at 50°C min<sup>-1</sup>, then to 170°C at 3.5°C min<sup>-1</sup>, to 185°C at 1.5°C min<sup>-1</sup>, to 225°C at  
201 2.4°C min<sup>-1</sup>, and finally to 250°C at 5.5°C min<sup>-1</sup> and maintained for 15 min) and equipped with  
202 two splitless injectors and two flame-ionization detectors, with hydrogen as vector gas. FAME  
203 were injected and separated simultaneously on two columns, one polar (ZBWAX: 30 m x 0.25  
204 mm ID x 0.2 µm) and one apolar (ZB5HT: 30 m x 0.25 mm ID x 0.2 µm), with two different  
205 polarities we can easily quantify FA in case of co-elution. FAME were identified by comparisons  
206 of their retention time with those of commercial standards (Supelco 37 Component FAME Mix,

207 the PUFA No.1 and No.3, and the Bacterial Acid Methyl Ester Mix from Sigma) and in-house  
208 standard mixtures from marine bivalves and micro- and macroalgae. FA compositions were  
209 expressed as the mass percentage of the total FA content.

#### 210 *2.4.2 Sterol analysis by Gas Chromatography*

211 Sterols were analyzed in a Shimadzu GC-2010 gas chromatograph (GC). Sterols were injected  
212 and separated on a ZB5HT: 30 m x 0.25 mm ID x 0.2  $\mu\text{m}$  (Phenomenex). The GC temperature  
213 program was from 40°C to 100°C at 25°C.min<sup>-1</sup>, then to 250°C at 15°C.min<sup>-1</sup>, and finally to 315°C  
214 at 3°C.min<sup>-1</sup> and maintained for 10 min. GC was equipped of splitless injector slated at 280°C,  
215 and a flame-ionization detector programmed at 250°C, with hydrogen as vector gas and an  
216 interface maintained at 315°C. Sterols were then identified by comparisons of their retention  
217 time with those of commercial standards and an in-house standard mixtures from marine  
218 bivalves. Sterols relative proportions were expressed as the mass percentage of the total sterols  
219 content.

#### 220 *2.5 Statistical Analysis*

221 All data were first tested for normality (Shapiro-Wilks test) and homoscedasticity (Bartlett test).  
222 If conditions were satisfied, parametric analyses were carried out, alternatively, non-parametric  
223 analyses were used. Du to number of replicates, the statistical comparison of food sources were  
224 impossible. SI variations between stations of bivalve muscles were compared by Student's test  
225 (t-tests). In total, 35 FA and 11 sterols having a mass proportion higher than 0.5 % of total FA or  
226 sterol were considered for statistical analyses, and non-identified FA and sterols were grouped

227 as “unknown”. Permutational variance analysis (PERMANOVA) permitted to test the interaction  
228 between periods and stations. As the interaction was significant, we analyzed the global spatial  
229 difference of both FA and sterol compositions by PERMANOVA tests at each periods. Due to the  
230 low proportion of zero in our dataset (less than 5 % for FA and 1.5% for sterols) Euclidean  
231 distances or Euclidean-based methods were used throughout our study. To identify which were  
232 the most discriminant FA and sterols between stations, a similarity of percentages analyses  
233 (SIMPER) was used. Principal component analyses (PCA) were performed to investigate the  
234 variation in lipid profiles between stations, for each species and both periods. FA and sterols  
235 which accounted for more than 80 % of dissimilarity (SIMPER) were represented on PCA. For  
236 each lipid fractions, pairwise comparisons (Wilcoxon tests) were carried out to explore spatial  
237 differences for each sterol, at each periods.

238 Firstly, we considered the consistency between trophic structures derived from FA and sterol  
239 tracers, at the species levels and at the two periods. Coinertia analyses (Dray et al., 2003) were  
240 used to assess whether the same spatial and temporal patterns were retrieved with the two  
241 sets of markers. RV coefficient (between 0 and 1) was used to quantify the degree of adequacy  
242 between the two datasets, with high values indicating strong concordance. Secondly, we  
243 considered the simple relationships among the principal biomarkers using Spearman's rank  
244 correlation (where  $r_s$  is the correlation coefficient). When a correlation was detected, the model  
245 (linear or not) selected by the Akaike Information Criterion (AIC) was adjusted and the normality  
246 of residuals was checked using Q–Q plots.

247 All statistical analyses and graphics were performed with the free software R (Core Team, 2017),  
248 with R Version 3.4.1 (2017 06 30).

ACCEPTED MANUSCRIPT

### 249 3. Results

#### 250 3.1 Spatial differences in biomarker compositions

##### 251 3.1.1 Food sources

252 Spatial and seasonal trends were observed for all biomarker compositions (SI, FA and sterols) of  
253 potential food compartments that contribute to bivalve diets. Nevertheless, precautions should  
254 be taken as the low number of replicates ( $n \leq 2$  per stations), did not permit to carry out  
255 statistical analyses to reveal spatial differences.

256 POM was more  $^{15}\text{N}$ -enriched in February than in August (Figure 2). For both seasons, the highest  
257  $\delta^{15}\text{N}$  values for POM were recorded from station 1, near the lagoon entrance (8.3‰ in February  
258 and 5.5‰ in August) and the lowest at station 2, the furthest site from the bay mouth (5.9‰ in  
259 February and 3.7‰ in August). At the both periods, the  $\delta^{13}\text{C}$  values of POM were close between  
260 both stations but were quite  $^{13}\text{C}$ -depleted compared to SOM (Figure 2). The  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  of  
261 SOM did not seem to differ between stations in February (Figure 2). As for POM, SOM samples  
262 from station 1 were more  $^{15}\text{N}$ -enriched than samples from station 2. The same gradient also  
263 occurred for  $\delta^{13}\text{C}$ , with station 1 which seems more  $^{15}\text{N}$ -enriched than station 2 (Figure 2). SOM  
264  $\delta^{13}\text{C}$  were close to those of *Zostera marina* detritus (Figure 2), between -13.1‰ and -8.6‰, in  
265 February and in August. *Z. marina* detritus did not show big differences in  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$   
266 between seasons with an average  $\delta^{13}\text{C}$  of -11.5‰ in February and August and an average  $\delta^{15}\text{N}$   
267 of 2.8‰ in February and 3.4‰ in August.






Figure 2. Stable Isotope (SI) bi-plot ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ , mean values  $\pm$ SD when  $n \geq 2$ ) of the adductor muscles of *Spondylus crassisquama* (represented by circles) and of their potential food sources (POM, SOM and seag seagrass), in February and August. Samples from station 1 are shown in light blue and from station 2 in green.

268 The main FA found in POM and SOM were 16:0 for saturated FA (SFA), and 16:1n-7, 18:1n-9,  
 269 18:1n-7 for monounsaturated FA (MUFA) (Supplementary Table 1). For polyunsaturated FA  
 270 (PUFA), 20:5n-3 and 22:6n-3 were the FA that exhibited the highest abundances in POM and  
 271 SOM; 18:3n-3 and 18:4n-3 showed highest values in POM and 20:4n-6 in SOM (Supplementary  
 272 Table 1). Station 2 samples had POM enriched in 18:4n-3 (8.3% in February and 5.3% in August)  
 273 and in 22:6n-3 (6.9% in February and 8.8% in August) (Supplementary Table 1). SOM showed  
 274 specifically higher values of 20:4n-6 at station 2 (5.8% in February, and 4.3% in August)

275 (Supplementary Table 1). In February, *Z. marina* was characterized by high levels of 16:3n-3  
276 (6.7%) and 18:3n-3 (9.7%) (Supplementary Table 1). In August, the proportion of 18:3n-3  
277 reached 33.6% of total FA, whereas 16:3n-3 proportion decreased to 3.7%. The proportion of  
278 18:2n-6, which was at low levels in February (1.9%), reached 17.4% in *Z. marina* sampled in  
279 August (Supplementary Table 1).

280 The main sterols found in POM were Cholesterol, Brassicasterol, Norcholesterol, 24-  
281 Methylenecholesterol, Isofucosterol and  $\beta$ -Sitosterol. At both periods, proportions of  
282 Cholesterol in POM collected at station 1 showed higher proportions than station 2 (40.1%  
283 against 34% in February, and 56.1% against 38% in August) (Supplementary Table 2).  
284 Brassicasterol was present at similar levels in POM at both stations in February (15.4% and  
285 15.5%), but was lower at station 1 in August (7.6%) (Supplementary Table 2). The 24-  
286 Methylenecholesterol was more abundant in the POM in February (9.0% at station 1 and 12.8%  
287 at station 2) (Supplementary Table 2). Stigmasterol was not detected in any samples of POM in  
288 February and only 2.9% at station 1 in August, but presented 26.4% of total sterols of POM  
289 sample at station 2 in August. In August, the ratio 24-Methylenecholesterol / Brassicasterol was  
290 more than twice higher in POM at station 1 (0.8) than at station 2 (0.3). Predominant sterols  
291 found in SOM were the same as POM except for Cholestanol, and Campesterol that were  
292 principally found in SOM (only detected in SOM for Campesterol). At both months, SOM seemed  
293 to present a higher abundance of Norcholesterol and Cholestanol at station 1 than at station 2  
294 (Supplementary Table 2). Inversely, Cholesterol, Brassicasterol, and  $\beta$ -Sitosterol were present in  
295 higher proportions in SOM collected at station 2 (Supplementary Table 2). The main eelgrass

296 sterols were  $\beta$ -Sitosterol, Stigmasterol and Cholesterol (Supplementary Table 2). Stigmasterol  
297 showed higher proportions in eelgrass sampled in August (20.9 vs 10.3% in February), whereas  
298 Cholesterol was in much higher proportions in February (16.7% vs 3.8% in August).

### 299 3.1.2 Bivalves

300 Although spatial differences were quantitatively low,  $\delta^{13}\text{C}$  values significantly differed between  
301 stations in February, (-18.4‰ at station 1 and -19.1‰ at station 2) (t-test,  $p < 0.001$ ,  $df = 16.70$ ,  
302  $t = 4.21$ ; Figure 2). In August,  $\delta^{13}\text{C}$  did not differ significantly between stations (t-test,  $p > 0.05$ ,  
303  $df = 15.54$ ,  $t = 0.31$ ), with individuals from station 1 presenting values of -19.1‰ and from  
304 station 2 of -19.2‰ (Figure 2).  $\delta^{15}\text{N}$  differed significantly between stations for both sampling  
305 dates, with  $\delta^{15}\text{N}$  values higher at station 1 than at station 2, this difference being more  
306 pronounced in August. In February,  $\delta^{15}\text{N}$  significantly decreased in muscles along the transect (t-  
307 test,  $p < 0.001$ ,  $df = 17.76$ ,  $t = 7.56$ ), from 8.7‰ at station 1, to 7.6‰ at station 2 (Figure 2). In  
308 August,  $\delta^{15}\text{N}$  significantly differed between station 1 (8.8‰) and station 2 (7.9‰) (t-test,  $p <$   
309  $0.001$ ,  $df = 16.7$ ,  $t = 5.12$ ; Figure 2).

310 Because significant interactions were found for FA proportions between periods and stations  
311 (PERMANOVA,  $p$ -value  $< 0.01$ ,  $df = 1$ ,  $F = 10.14$ ,  $R^2 = 0.13$ ), and because sterol proportions  
312 presented same interactions (PERMANOVA,  $p$ -value  $< 0.01$ ,  $df = 1$ ,  $F = 4.45$ ,  $R^2 = 0.09$ ), we  
313 compared spatial differences at each period.

314 The PCA based on NL FA and sterols composition shows the differences between stations and  
 315 lipid biomarkers responsible for 80% of these differences (Figure 3).




Figure 3. Principal Component Analyses (PCA) of neutral lipids (NL) fatty acids (FA) and sterols compositions (mass %) of *Spondylus crassisquama* digestive gland in February (A and C) and August (B and D). Only FA and sterols that account for > 80% of the contribution of dissimilarity between stations are shown.

316 In February, significant spatial differences were found for both NL FA composition  
 317 (PERMANOVA,  $p < 0.001$ ,  $df = 2$ ,  $F = 5.45$ ,  $R^2 = 0.3$ ), and sterol composition (PERMANOVA,  $p$ -  
 318 value  $< 0.05$ ,  $df = 1$ ,  $F = 10.02$ ,  $R^2 = 0.56$ ). At this period individuals from station 1 were  
 319 characterized by higher levels in 20:4n-6 and 22:6n-3 (Figure 3 and Supplementary table 3) and  
 320 were associated to higher values of  $\beta$ -Sitosterol (Figure 3 and Supplementary table 4).  
 321 Individuals from station 2 tended to be characterized by FA 18:1n-7, 18:3n-3 (Figure 3 and  
 322 Supplementary table 3) and were not associated to any particular phytosterol.

323 In August, NL FA compositions differed among sampling location (PERMANOVA,  $p < 0.001$ ,  $df =$   
324  $2$ ,  $F = 34.11$ ,  $R^2 = 0.74$ ). The same was found for sterol composition (PERMANOVA,  $p$ -value  $<$   
325  $0.01$ ,  $df = 1$ ,  $F = 13.55$ ,  $R^2 = 0.63$ ). Here again, a clear spatial pattern was observed along the first  
326 axis of the PCA (Figure 3). We observed significantly decreasing values of 20:5n-3 and 16:1n-7  
327 from the mouth to the inner bay (Figure 3 and Supplementary table 3). Even if PCA revealed a  
328 contribution of 24-Methylenecholesterol (Figure 3), pairwise comparisons revealed no  
329 significant spatial differences (Supplementary table 4). Individuals from station 2 were  
330 characterized by higher proportions of 18:1n-9 and 18:3n-3 (Figure 3 and Supplementary table  
331 3), and by significantly higher proportions of Campesterol, Stigmasterol, Brassicasterol (Figure 3  
332 and Supplementary table 4).




Figure 4. Boxplot of the ratios of 24-Methylenecholesterol/Brassicasterol ( $n=5$ ) and of 20:5n-3/22:6n-3 (EPA/DHA;  $n=10$ ) of *Spondylus crassisquama* at the both periods. Different letters indicate significant differences (Wilcoxon test, at significant level  $\alpha < 0.05$ ). 24-ME.cholesterol: 24-Methylenecholesterol.

333 We calculated the ratio between 24-Methylenecholesterol and Brassicasterol and the ratio  
 334 between 20:5n-3 and 22:6n-3 (EPA/DHA) (Figure 4). In February, the EPA/DHA was significantly  
 335 higher at the inner site, while no difference was found for the 24-  
 336 Methylenecholesterol/Brassicasterol. In August, the two ratios displayed the same pattern,  
 337 being the highest near the entrance of the lagoon.

338 3.2 Comparisons of different biomarkers

339 We found a robust correlation between the two ordinations of FA and sterols. This relationship  
340 was stronger in February (RV = 0.71) than in August (RV = 0.6). Both co-inertia analyses were  
341 significant (p-value < 0.01). Significant Spearman's rank correlations were found between  
342 EPA/DHA and 24-Me/Bra ( $r_s = 0.7$ , p-values < 0.05; Figure 5). Both ratios were positively  
343 correlated to the 16:1n-7 ( $0.33 < r_s < 0.54$ , p-values < 0.05; Figure 6). Inversely, ratios were  
344 negatively correlated to Campesterol ( $-0.76 < r_s < -0.78$ , p-values < 0.05; Figure 5), as for  
345 Stigmasterol ( $-0.69 < r_s < -0.91$ , p-values < 0.05; Figure 5). Interestingly, Stigmasterol biomarker  
346 was strongly and positively correlated with Campesterol ( $r_s = 0.8$ , p-values < 0.05; Figure 5). In  
347 contrast, it was negatively correlated to 20:5n-3 ( $r_s = -0.48$ , p-values < 0.05; Figure 5). No  
348 correlation was observed between Stigmasterol and 16:1n-7 ( $r_s = -0.39$ ,  $P > 0.05$ ; Figure 5). 24-  
349 Methylenecholetrol was strongly correlated with EPA/DHA ratio ( $r_s = 0.73$ ,  $P < 0.05$ ; Figure 5).

350




Figure 5. Correlations between the three biomarkers (Spearman's test): stable isotope ratios (carbon and nitrogen; ‰), neutral lipid fatty acids, and sterols (mass %) for *Spondylus crassisquama*. The colored circles represent significant correlations between two tracers ( $p$ -value < 0.05). The numbers inside cells are the associated Spearman's correlation.

351



#### 352 4. Discussion

353 In coastal lagoons, investigating the nature of food sources consumed by bivalves can provide a  
354 spatially and temporally integrated overview of nutrient transfers within the system and across  
355 ecosystem boundaries (e.g lagoon primary production vs inputs from coastal ocean). Here, the  
356 coupled approach used revealed quite powerful to unravel spatial patterns in the trophic  
357 ecology of *S. crassiquama*. Our results point out the significant role played by oceanic inputs in  
358 the ecological functioning of coastal lagoons, including the support of bivalve populations. This  
359 study also highlights the role played by remineralization in the food web of the inner lagoon.

##### 360 4.1 Spatial differences in biomarker compositions

361 The similarity of carbon SI composition observed between POM and bivalves isotopic  
362 composition is in accordance with a direct and non-selective feeding (Vander Zanden and  
363 Rasmussen, 2001). Although significant differences were observed between stations especially  
364 for  $\delta^{15}\text{N}$ , the extent of these differences was low, typically below 1.5 ‰ suggesting that no  
365 major contrast occurred in the diet of this species in the lagoon. Individuals sampled close to the  
366 mouth displayed the highest  $\delta^{15}\text{N}$ , while those from the inner lagoon displayed the lowest  $\delta^{15}\text{N}$ .  
367 This pattern matches the one displayed in POM, suggesting that bivalve  $\delta^{15}\text{N}$  directly reflect the  
368 composition of POM, and their diet does not drastically differ from its composition (Coma et al.,  
369 2001). Consequently,  $\delta^{15}\text{N}$  indicates that oceanic inputs affecting the composition of POM are  
370 transferred into the local food web through their consumption by suspension-feeding bivalves  
371 of the lagoon. Ojo de Liebre is located in an arid region, with virtually no freshwater inputs  
372 (Phleger and Ewing, 1962). In contrast, the open ocean facing the lagoon can induce  $^{15}\text{N}$ -

373 enriched nutrient or plankton pulses that could explain the spatial pattern observed here  
374 (Altabet et al., 1999). The higher values at the entrance of the lagoon tend to be close to those  
375 described in the region (McMahon et al., 2013; Olson et al., 2010). It seems therefore that the  
376 lagoon of Ojo de Liebre, at least in its proximal part, is significantly influenced by its nearshore  
377 hydrological conditions, through the entrance of oceanic originating organic matter/nutrients.

378 At both seasons, global FA and sterol compositions differed, revealing spatial differences in the  
379 diet of *S. crassisquama*. In February, even though some biomarkers were significantly different  
380 between stations, the spatial pattern was quite slight, revealing only small dietary differences.  
381 Individuals from the mouth of the lagoon were characterized by higher proportions of 22:6n-3  
382 and  $\beta$ -Sitosterol. The presence of 22:6n-3 suggests a dominance of dinoflagellates in the diet of  
383 *S. crassisquama* (Alfaro et al., 2006; Parrish, 2009). Although  $\beta$ -Sitosterol was the major sterol  
384 found in eelgrass, it is not possible to conclude that  $\beta$ -Sitosterol found in bivalves DG is obtained  
385 from eelgrass detritus, as this sterol was also found in POM and SOM. High proportions of  $\beta$ -  
386 Sitosterol have also been reported in haptophytes and diatoms in the literature (Napolitano et  
387 al., 1993; Soudant et al., 1998). In fact, most phytosterols found in this study were found in all  
388 POM, SOM, and eelgrass samples, making difficult to decipher their origin among potential  
389 foods sources of bivalves.

390 In August, a clear gradient from the mouth to the center of the bay suggested a more important  
391 spatial diet shift. Individuals from the most oceanic stations (station 1) were characterized by  
392 higher levels of 16:1n-7 as well as 20:5n-3, but also by higher levels of 24-Methylenecholesterol.

393 The presence of these lipid biomarker, as well as higher EPA/DHA and 24-Me./Bra ratios,  
394 highlights the importance of diatoms in the diet of bivalves from this station during this period  
395 (Kharlamenko et al., 2001; Volkman, 2003; Alfaro et al., 2006; Lavaud et al., 2018). This supply in  
396 diatoms at the mouth of the lagoon supports the hypothesis of oceanic inputs and intake of  
397 primary production from coastal waters suggested by  $^{15}\text{N}$  enrichment. Du et al. (2015) showed  
398 in the Northern California current that diatoms proliferate during strong upwelling events (from  
399 April to September), when nutrients and turbulence are relatively high. The fact that oceanic  
400 water masses present the same taxonomic dominances supports again the hypothesis that  
401 oceanic organic matter enters the lagoon and supports local the food web. However, the  
402 diminution of diatoms importance at the inner station suggests that oceanic inputs do not  
403 extend to the whole bay. At both seasons, in the center of the bay, *S. crassisquama* was  
404 characterized by high levels of C18 PUFA (18:3n-3) and/or MUFA (18:1n-9), but also by high  
405 proportions of Brassicasterol, Campesterol, or Stigmasterol. These FA can be found in various  
406 primary producers, such as macroalgae (Nelson et al., 2002), higher plants (Valitova et al.,  
407 2016), seagrass (Kharlamenko et al., 2001; Jaffé et al., 2006; Volkman et al., 2008), various  
408 phytoplankton taxa (Soudant et al., 1998; Dalsgaard et al., 2003), including dinoflagellates  
409 (Soudant et al., 1998; Kharlamenko et al., 2001; Hurtado et al., 2012; Nerot et al., 2015) and are  
410 very abundant in various ciliate species, which contribute to the marine microbial food web  
411 (Zhukova and Kharlamenko, 1999). Consequently, bivalves from the center of the bay seem to  
412 rely, at least partly, on microheterotrophs (dinoflagellates, flagellates, ciliates), hence revealing  
413 the influence of the microbial food web in this part of the lagoon in summer. The lower

414 EPA/DHA and 24-Me./Bra ratios also support the hypothesis of the presence of dinoflagellates  
415 in the diet of this species (Alfaro et al., 2006; Kelly and Scheibling, 2012; Lavaud et al., 2018).

416 Campesterol, has been reported in small amount in some species of diatoms like *Chaetoceros*  
417 *calcitrans* (Yamauguchi et al., 1986; Soudant et al., 1998; Mohammady, 2004). The contrast in  
418 the spatial repartition of Campesterol (higher at inner station) and 24-Methylenecholesterol  
419 (higher at entrance station) suggests different diatom assemblages and the contribution of  
420 different diatom species in the diet of bivalves within the lagoon. In addition, Campesterol was  
421 absent from POM samples at both stations and for both seasons, whereas it was present in non-  
422 negligible amounts in SOM. The presence of this sterol in bivalves might therefore suggest a  
423 contribution of SOM in the diet of bivalves, through the resuspension of sediment.

424 The contribution of eelgrass detritus to the diet of bivalves, although already reported as a  
425 secondary dietary source in enclosed coastal systems (Schaal et al., 2008) seems unlikely  
426 because of the lack of  $^{13}\text{C}$ -enrichment in bivalves. However, lipid biomarkers as 18:3n-3 or  
427 Stigmasterol seem to reveal the presence of eelgrass in bivalve diets, and in particular at the  
428 inner station. Seagrass cover a surface of more than 590 km<sup>2</sup> in the Ojo de Liebre lagoon  
429 (Riosmena-Rodríguez et al., 2013), and are known to produce large amounts of detritus, which  
430 can be buried in the sediment, exported to the adjacent nearshore environment, or locally  
431 transferred into the food web (Hyndes and Lavery, 2005; Schaal et al., 2008). Here, SI ratios  
432 measured in sedimentary organic matter revealed strong  $^{13}\text{C}$  enrichment, suggesting that  
433 detritus may be major components of sediment in the Ojo de Liebre lagoon. Although these

434 high  $\delta^{13}\text{C}$  in SOM could also reveal the importance of epiphytes, which commonly display a SI  
435 composition similar to their seagrass host, the highly degraded nature of sediments suggests a  
436 dominance of refractory compounds, which are abundant in seagrass (Fourqurean and Schlau,  
437 2003). In contrast with sediment, POM did not seem to reflect any significant contribution of  
438 seagrass, which would have resulted in more  $^{13}\text{C}$ -enriched SI composition than what was  
439 observed. Only a slight  $^{13}\text{C}$ -enrichment could be found in POM at stations inside the lagoon that  
440 could be due to abundant seagrass detritus in suspension on the water column, but the lack of  
441 replicates prevents any definitive conclusion about seagrass contribution to the POM pool.

442

#### 443 4.2 Comparison of different lipid biomarkers

444 Comparing sterol composition with FA and SI analysis brings some interesting insights into the  
445 potential contribution of benthic primary producers in the diet of bivalve species. The  
446 compositions in FA and sterols were strongly correlated according to the RV coefficient of the  
447 co-inertia analyses. The higher RV coefficient in February could be explained by the fact that the  
448 diet of bivalves was more spatially homogeneous in the winter period. Spearman's correlations  
449 allowed to validate the use of 24Me./Bra as a maker of predominance of diatoms or  
450 dinoflagellates in the bivalve diet. Indeed, this ratio was strongly positively correlated with  
451 EPA/DHA ratio, which is currently used in literature (Le Croizier et al., 2016; Parrish, 2009). Both  
452 ratios were also positively correlated to 16:1n-7, which confirmed their usefulness to unravel  
453 the predominance of diatoms in the diet of bivalves (Dalsgaard et al., 2003; Richoux and  
454 Froneman, 2008). However, Campesterol was negatively correlated to both ratios, which could

455 be surprising for a known diatom biomarkers. In fact, Campesterol is reported to be biomarker  
456 of Bacillariophyceae (Soudant et al., 1998; Mohammady, 2004), which seems to be a group of  
457 diatoms only predominant in the inner lagoon. Besides, Stigmasterol, which could be considered  
458 as biomarker of eelgrass (Jaffé et al., 2006; Volkman et al., 2008), is highly and positively  
459 correlated to 18:3n-3 also known as biomarker of eelgrass (Kharlamenko et al., 2001). This  
460 correlation and high amount of these lipid biomarkers found in seagrass detritus samples,  
461 support their utilization as biomarker of seagrass detritus in the lagoon of Ojo de Liebre.

## 462 **5. Conclusion**

463 Stable isotopes and fatty acids and sterols proved being highly complementary tools to identify  
464 spatio-temporal patterns in the Ojo de Liebre lagoon. The entrance of the lagoon seems to  
465 reflect coastal oceanic processes, supporting the hypothesis of significant oceanic primary  
466 production pulses into the lagoon. Although significant, the influence of oceanic inputs does not  
467 extend to the whole lagoon. The center of the lagoon seems to be more local processes-driven,  
468 with a recycled organic matter-based food web and the role of seagrass suggested by FA and  
469 sterols analysis. Our results revealed strong correlations between FA and sterols in bivalves.  
470 Consequently, sterol compositions could be used as complementary and/or substitute  
471 biomarkers to FA. The approach used in this study brought some interesting new insights on the  
472 potential contribution of primary producers to bivalve diets. This coupling brings new  
473 information about phytoplankton assemblages and diet composition differences at low

474 taxonomic level, which seems to be an improvement compared to the broader discrimination

475 power and SI and FA.

476

ACCEPTED MANUSCRIPT

477 **Acknowledgements**

478 This study was supported by the ECOS-ANUIES program (PROPHYMUS project), the "Laboratoire  
479 d'Excellence" LabexMER (ANR-10-LABX-19) and co-funded by a grant from the French  
480 government under the program "Investissements d'Avenir". MMR's PhD fellowship was  
481 provided by the French Research Ministry and Region Bretagne. We would like to thank the  
482 contribution of the Guerrero Negro unit of CIBNOR for hosting the first steps of sample  
483 preparation after fieldwork, the LIPIDOCEAN analytical facility for hosting all fatty acid analyses,  
484 and the Pôle Spectrométrie Océan (Clément Tanvet) for running stable isotope analyses.

485 We thank the "Pepe" Jose Alfredo Saenz Arce crew for the fieldwork, as well as Rosa Linda  
486 Salgado Garcia, and Laura Guzman for their precious help during sampling and dissections, and  
487 Dr. Elena Palacios and Dr. Kitty Arredondo Vega, for the laboratory facilities provided for the  
488 preparation of samples at CIBNOR La Paz. We thank the Reserva de la biosfera desierto de "el  
489 Vizcaino" for their help and support in carrying out sampling for this study. Authors would like  
490 to thank Aurelien Boye and Olivier Gauthier for their precious help and advices about statistical  
491 analyses.

492



493 **Literature cited**

- 494 Alfaro, A.C., Thomas, F., Sergent, L., Duxbury, M., 2006. Identification of trophic interactions  
495 within an estuarine food web (northern New Zealand) using fatty acid biomarkers and stable  
496 isotopes. *Estuar. Coast. Shelf Sci.*, Applying the Ecohydrology approach to the Guadiana estuary  
497 and coastal areas: lessons learned from dam impacted ecosystems 70, 271–286.  
498 <https://doi.org/10.1016/j.ecss.2006.06.017>
- 499 Altabet, M.A., PilskaIn, C., Thunell, R., Pride, C., Sigman, D., Chavez, F., Francois, R., 1999. The  
500 nitrogen isotope biogeochemistry of sinking particles from the margin of the Eastern North  
501 Pacific. *Deep Sea Res. Part Oceanogr. Res. Pap.* 46, 655–679. <https://doi.org/10.1016/S0967->  
502 [0637\(98\)00084-3](https://doi.org/10.1016/S0967-0637(98)00084-3)
- 503 Bec, A., Perga, M.-E., Desvilettes, C., Bourdier, G., 2010. How well can the fatty acid content of  
504 lake seston be predicted from its taxonomic composition? *Freshw. Biol.* 55, 1958–1972.
- 505 Cabello-Pasini, A., Muñiz-Salazar, R., Ward, D.H., 2003. Annual variations of biomass and  
506 photosynthesis in *Zostera marina* at its southern end of distribution in the North Pacific. *Aquat.*  
507 *Bot.* 76, 31–47. [https://doi.org/10.1016/S0304-3770\(03\)00012-3](https://doi.org/10.1016/S0304-3770(03)00012-3)
- 508 Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Escoubeyrou, K., Desmalades, M., Caparros,  
509 J., Grémare, A., 2007. A seasonal survey of the food web in the Lapalme Lagoon (northwestern  
510 Mediterranean) assessed by carbon and nitrogen stable isotope analysis. *Estuar. Coast. Shelf Sci.*  
511 73, 299–315. <https://doi.org/10.1016/j.ecss.2007.01.012>

- 512 Clavier, J., Chauvaud, L., Amice, E., Lazure, P., Van Der Geest, M., Labrosse, P., Diagne, A.,  
513 Carlier, A., Chauvaud, S., 2014. Benthic metabolism in shallow coastal ecosystems of the Banc  
514 d'Arguin, Mauritania. *Mar. Ecol. Prog. Ser.* 501, 11–23.
- 515 Coma, R., Ribes, M., Gili, J.-M., Hughes, R.N., 2001. The ultimate opportunists: consumers of  
516 seston. *Mar. Ecol. Prog. Ser.* 219, 305–308.
- 517 Core Team, R., 2017. R: A language and environment for statistical computing. R Foundation for  
518 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 519 Dalsgaard, J., St. John, M., Kattner, G., Müller-Navarra, D., Hagen, W., 2003. Fatty acid trophic  
520 markers in the pelagic marine environment. *Adv. Mar. Biol.* 46, 225–340.  
521 [https://doi.org/10.1016/S0065-2881\(03\)46005-7](https://doi.org/10.1016/S0065-2881(03)46005-7)
- 522 Dame, R.F., 1993. The role of bivalve filter feeder material fluxes in estuarine ecosystems, in:  
523 *Bivalve Filter Feeders*. Springer, pp. 245–269.
- 524 Desvillettes, C.H., Bourdier, G., Amblard, C.H., Barth, B., 1997. Use of fatty acids for the  
525 assessment of zooplankton grazing on bacteria, protozoans and microalgae. *Freshw. Biol.* 38,  
526 629–637.
- 527 Dray, S., Chessel, D., Thioulouse, J., 2003. Co-inertia analysis and the linking of ecological data  
528 tables. *Ecology* 84, 3078–3089.

- 529 Du, X., Peterson, W., O'Higgins, L., 2015. Interannual variations in phytoplankton community  
530 structure in the northern California Current during the upwelling seasons of 2001-2010. Mar.  
531 Ecol. Prog. Ser. 519, 75–87.
- 532 Fenton, G.E., Ritz, D.A., 1988. Changes in carbon and hydrogen stable isotope ratios of  
533 macroalgae and seagrass during decomposition. Estuar. Coast. Shelf Sci. 26, 429–436.
- 534 Fourqurean, J.W., Schrlau, J.E., 2003. Changes in nutrient content and stable isotope ratios of C  
535 and N during decomposition of seagrasses and mangrove leaves along a nutrient availability  
536 gradient in Florida Bay, USA. Chem. Ecol. 19, 373–390.
- 537 Gattuso, J.-P., Frankignoulle, M., Wollast, R., 1998. Carbon and carbonate metabolism in coastal  
538 aquatic ecosystems. Annu. Rev. Ecol. Syst. 29, 405–434.
- 539 Giner, J.-L., Zhao, H., Dixon, M.S., Wikfors, G.H., 2016. Bioconversion of <sup>13</sup>C-labeled microalgal  
540 phytosterols to cholesterol by the Northern Bay scallop, *Argopecten irradians irradians*. Comp.  
541 Biochem. Physiol. B Biochem. Mol. Biol. 192, 1–8.
- 542 Holden, M.J., Patterson, G.W., 1991. Absence of sterol biosynthesis in oyster tissue culture.  
543 Lipids 26, 81–82.
- 544 Hurtado, M.A., Racotta, I.S., Arcos, F., Morales-Bojórquez, E., Moal, J., Soudant, P., Palacios, E.,  
545 2012. Seasonal variations of biochemical, pigment, fatty acid, and sterol compositions in female  
546 *Crassostrea corteziensis* oysters in relation to the reproductive cycle. Comp. Biochem. Physiol. B  
547 Biochem. Mol. Biol. 163, 172–183.

- 548 Hyndes, G.A., Lavery, P.S., 2005. Does transported seagrass provide an important trophic link in  
549 unvegetated, nearshore areas? *Estuar. Coast. Shelf Sci.* 63, 633–643.  
550 <https://doi.org/10.1016/j.ecss.2005.01.008>
- 551 Jaffé, R., Rushdi, A.I., Medeiros, P.M., Simoneit, B.R.T., 2006. Natural product biomarkers as  
552 indicators of sources and transport of sedimentary organic matter in a subtropical river.  
553 *Chemosphere* 64, 1870–1884. <https://doi.org/10.1016/j.chemosphere.2006.01.048>
- 554 Kanazawa, A., 2001. Sterols in marine invertebrates. *Fish. Sci.* 67, 997–1007.
- 555 Kelly, J.F., 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian  
556 trophic ecology. *Can. J. Zool.* 78, 1–27.
- 557 Kelly, J.R., Scheibling, R.E., 2012. Fatty acids as dietary tracers in benthic food webs. *Mar. Ecol.*  
558 *Prog. Ser.* 446, 1–22.
- 559 Kharlamenko, V.I., Kiyashko, S.I., Imbs, A.B., Vyshkvartzev, D.I., 2001. Identification of food  
560 sources of invertebrates from the seagrass *Zostera marina* community using carbon and sulfur  
561 stable isotope ratio and fatty acid analyses. *Mar. Ecol. Prog. Ser.* 220, 103–117.
- 562 Kraak, M.H.S., Scholten, M.C.T., Peeters, W.H.M., de Kock, W.C., 1991. Biomonitoring of heavy  
563 metals in the Western European rivers Rhine and Meuse using the freshwater mussel *Dreissena*  
564 *polymorpha*. *Environ. Pollut.* 74, 101–114. [https://doi.org/10.1016/0269-7491\(91\)90107-8](https://doi.org/10.1016/0269-7491(91)90107-8)

- 565 Lavaud, R., Artigaud, S., Le Grand, F., Donval, A., Soudant, P., Flye-Sainte-Marie, J., Strohmeier,  
566 T., Strand, Ø., Leynaert, A., Beker, B., 2018. New insights into the seasonal feeding ecology of  
567 *Pecten maximus* using pigments, fatty acids and sterols analyses. Mar. Ecol. Prog. Ser. 590, 109–  
568 129.
- 569 Le Croizier, G., Schaal, G., Gallon, R., Fall, M., Le Grand, F., Munaron, J.-M., Rouget, M.-L.,  
570 Machu, E., Le Loc'H, F., Laë, R., 2016. Trophic ecology influence on metal bioaccumulation in  
571 marine fish: Inference from stable isotope and fatty acid analyses. Sci. Total Environ. 573, 83–  
572 95.
- 573 Le Grand, F.L., Soudant, P., Siah, A., Tremblay, R., Marty, Y., Kraffe, E., 2014. Disseminated  
574 Neoplasia in the soft-shell Clam *Mya arenaria*: membrane lipid composition and functional  
575 parameters of circulating cells. Lipids 49, 807–818. <https://doi.org/10.1007/s11745-014-3917-4>
- 576 Lefebvre, S., Harma, C., Blin, J.-L., 2009. Trophic typology of coastal ecosystems based on  $\delta^{13}\text{C}$   
577 and  $\delta^{15}\text{N}$  ratios in an opportunistic suspension feeder. Mar. Ecol. Prog. Ser. 390, 27–37.
- 578 Lorrain, A., Paulet, Y.-M., Chauvaud, L., Savoye, N., Donval, A., Saout, C., 2002. Differential  $\delta^{13}\text{C}$   
579 and  $\delta^{15}\text{N}$  signatures among scallop tissues: implications for ecology and physiology. J. Exp. Mar.  
580 Biol. Ecol. 275, 47–61. [https://doi.org/10.1016/S0022-0981\(02\)00220-4](https://doi.org/10.1016/S0022-0981(02)00220-4)
- 581 Mann, K.H., 1982. Ecology of Coastal Waters: A Systems Approach. University of California  
582 Press.

- 583 Marchais, V., Schaal, G., Grall, J., Lorrain, A., Nerot, C., Richard, P., Chauvaud, L., 2013. Spatial  
584 variability of stable isotope Ratios in Oysters (*Crassostrea gigas*) and primary producers along an  
585 estuarine gradient (Bay of Brest, France). *Estuaries Coasts* 36, 808–819.  
586 <https://doi.org/10.1007/s12237-012-9584-x>
- 587 Martínez, M.L., Intralawan, A., Vázquez, G., Pérez-Maqueo, O., Sutton, P., Landgrave, R., 2007.  
588 The coasts of our world: Ecological, economic and social importance. *Ecol. Econ., Ecological*  
589 *Economics of Coastal Disasters* 63, 254–272. <https://doi.org/10.1016/j.ecolecon.2006.10.022>
- 590 McMahon, K.W., Hamady, L.L., Thorrold, S.R., 2013. Ocean ecogeochemistry: a review.  
591 *Oceanogr. Mar. Biol. Annu. Rev.* 51, 327–374.
- 592 Mohammady, N.G., 2004. Total, free and conjugated sterolic forms in three microalgae used in  
593 mariculture. *Z. Für Naturforschung C* 59, 619–624.
- 594 Napolitano, G.E., Ackman, R.G., Silva-Serra, M.A., 1993. Incorporation of dietary sterols by the  
595 sea scallop *Placopecten magellanicus* (Gmelin) fed on microalgae. *Mar. Biol.* 117, 647–654.
- 596 Nelson, M.M., Leighton, D.L., Phleger, C.F., Nichols, P.D., 2002. Comparison of growth and lipid  
597 composition in the green abalone, *Haliotis fulgens*, provided specific macroalgal diets. *Comp.*  
598 *Biochem. Physiol. B Biochem. Mol. Biol.* 131, 695–712. [https://doi.org/10.1016/S1096-](https://doi.org/10.1016/S1096-4959(02)00042-8)  
599 [4959\(02\)00042-8](https://doi.org/10.1016/S1096-4959(02)00042-8)
- 600 Nerot, C., Meziane, T., Schaal, G., Grall, J., Lorrain, A., Paulet, Y.-M., Kraffe, E., 2015. Spatial  
601 changes in fatty acids signatures of the great scallop *Pecten maximus* across the Bay of Biscay

602 continental shelf [WWW Document]. URL <http://ac.els-cdn.com/scd-proxy.univ->  
603 [brest.fr/S0278434315300522/1-s2.0-S0278434315300522-main.pdf?\\_tid=cc01133c-2f39-11e7-](http://ac.els-cdn.com/scd-proxy.univ-brest.fr/S0278434315300522/1-s2.0-S0278434315300522-main.pdf?_tid=cc01133c-2f39-11e7-8950-00000aab0f02&acdnat=1493731274_d207b602cfd94a22be277a4effaa33b7)  
604 [8950-00000aab0f02&acdnat=1493731274\\_d207b602cfd94a22be277a4effaa33b7](http://ac.els-cdn.com/scd-proxy.univ-brest.fr/S0278434315300522/1-s2.0-S0278434315300522-main.pdf?_tid=cc01133c-2f39-11e7-8950-00000aab0f02&acdnat=1493731274_d207b602cfd94a22be277a4effaa33b7) (accessed  
605 5.2.17).

606 Nixon, S.W., 1982. Nutrient dynamics, primary production and fisheries yields of lagoons.  
607 *Oceanol. Acta Spec. Issue* 357–371.

608 Olson, R.J., Popp, B.N., Graham, B.S., López-Ibarra, G.A., Galván-Magaña, F., Lennert-Cody, C.E.,  
609 Bocanegra-Castillo, N., Wallsgrove, N.J., Gier, E., Alatorre-Ramírez, V., Ballance, L.T., Fry, B.,  
610 2010. Food-web inferences of stable isotope spatial patterns in copepods and yellowfin tuna in  
611 the pelagic eastern Pacific Ocean. *Prog. Oceanogr., Climate Impacts on Oceanic TOp Predators*  
612 (CLIOTOP) 86, 124–138. <https://doi.org/10.1016/j.pocean.2010.04.026>

613 Parrish, C.C., 2009. Essential fatty acids in aquatic food webs, in: *Lipids in Aquatic Ecosystems*.  
614 Springer, pp. 309–326.

615 Parrish, C.C., Abrajano, T.A., Budge, S.M., Helleur, R.J., Hudson, E.D., Pulchan, K., Ramos, C.,  
616 2000. Lipid and phenolic biomarkers in marine ecosystems: analysis and applications, in: *Marine*  
617 *Chemistry*. Springer, pp. 193–223.

618 Peterson, B.J., Fry, B., 1987. Stable isotopes in ecosystem studies. *Annu. Rev. Ecol. Syst.* 18, 293–  
619 320.

- 620 Phleger, F.B., Ewing, G.C., 1962. Sedimentology and oceanography of coastal Lagoons in Baja  
621 California, Mexico. Geol. Soc. Am. Bull. 73, 145. [https://doi.org/10.1130/0016-](https://doi.org/10.1130/0016-7606(1962)73[145:SAOOCL]2.0.CO;2)  
622 [7606\(1962\)73\[145:SAOOCL\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1962)73[145:SAOOCL]2.0.CO;2)
- 623 Post, D.M., 2002. Using stable isotopes to estimate trophic position: models, methods, and  
624 assumptions. Ecology 83, 703–718.
- 625 Post, D.M., Layman, C.A., Arrington, D.A., Takimoto, G., Quattrochi, J., Montaña, C.G., 2007.  
626 Getting to the fat of the matter: models, methods and assumptions for dealing with lipids in  
627 stable isotope analyses. Oecologia 152, 179–189. <https://doi.org/10.1007/s00442-006-0630-x>
- 628 Richoux, N.B., Froneman, P.W., 2008. Trophic ecology of dominant zooplankton and macrofauna  
629 in a temperate, oligotrophic South African estuary: a fatty acid approach. Mar. Ecol. Prog. Ser.  
630 357, 121–137.
- 631 Riosmena-Rodríguez, R., Muñiz-Salazar, R., López-Calderón, J., Torre-Cosío, J., Meling, A., Talbo,  
632 S.L., Sage, G.K., Ward, D.H., Cabello-Pasini, A., 2013. Conservation status of *Zostera marina*  
633 populations at Mexican Pacific. Adv. Environ. Res. 27, 35–63.
- 634 Schaal, G., Riera, P., Leroux, C., 2008. Trophic coupling between two adjacent benthic food webs  
635 within a man-made intertidal area: a stable isotopes evidence. Estuar. Coast. Shelf Sci. 77, 523–  
636 534. <https://doi.org/10.1016/j.ecss.2007.10.008>



- 637 Soudant, P., Le Coz, J.-R., Marty, Y., Moal, J., Robert, R., Samain, J.-F., 1998. Incorporation of  
638 microalgae sterols by scallop *Pecten maximus* (L.) larvae. *Comp. Biochem. Physiol. A. Mol.*  
639 *Integr. Physiol.* 119, 451–457.
- 640 Valitova, J.N., Sulkarnayeva, A.G., Minibayeva, F.V., 2016. Plant sterols: diversity, biosynthesis,  
641 and physiological functions. *Biochem. Mosc.* 81, 819–834.
- 642 Vander Zanden, M.J., Rasmussen, J.B., 2001. Variation in  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  trophic fractionation:  
643 implications for aquatic food web studies. *Limnol. Oceanogr.* 46, 2061–2066.  
644 <https://doi.org/10.4319/lo.2001.46.8.2061>
- 645 Volkman, J., 2003. Sterols in microorganisms. *Appl. Microbiol. Biotechnol.* 60, 495–506.
- 646 Volkman, J.K., 1986. A review of sterol markers for marine and terrigenous organic matter. *Org.*  
647 *Geochem.* 9, 83–99.
- 648 Volkman, J.K., Revill, A.T., Holdsworth, D.G., Fredericks, D., 2008. Organic matter sources in an  
649 enclosed coastal inlet assessed using lipid biomarkers and stable isotopes. *Org. Geochem.* 39,  
650 689–710. <https://doi.org/10.1016/j.orggeochem.2008.02.014>
- 651 Wasowicz, E., Rudzinska, M., 2011. Cholesterol and phytosterols. *Chem. Biol. Funct. Asp. Food*  
652 *Lipids* 113–134.
- 653 Yamauguchi, T., Koichi, I.T.O., Mitsuo, H., 1986. Studies on the sterols in some marine  
654 phytoplanktons. *Tohoku J. Agric. Res.* 37, 5–14.

655 Zhukova, N.V., Kharlamenko, V.I., 1999. Sources of essential fatty acids in the marine microbial  
656 loop. *Aquat. Microb. Ecol.* 17, 153–157. <https://doi.org/10.3354/ame017153>

657

ACCEPTED MANUSCRIPT

**Highlights**

- Stable isotopes fatty acids and sterols were used to characterize the diet of bivalves
- A transect was sampled in a Pacific lagoon with increasing distance from the mouth
- Bivalves at the entrance of the lagoon relied on oceanic phytoplankton inputs
- Bivalves in the inner lagoon relied on local microbial food web
- Lipid biomarkers and SI were complementary to distinguish the food sources origins