



**HAL**  
open science

## Draft Genome Sequence of a *Chryseobacterium indologenes* Strain Isolated from a Blood Culture of a Hospitalized Child in Antananarivo, Madagascar

Mamitina Alain Noah Rabenandrasana, Lala Fanomezantsoa Rafetrarivony, Lalainasoa Odile Rivoarilala, Vincent Enouf, Annick Lalaina Robinson, Ando Rakotozanany, Jessica Vanhomwegen, Valérie Caro, Cassandre von Platen, Jean-Claude Manuguerra, et al.

### ► To cite this version:

Mamitina Alain Noah Rabenandrasana, Lala Fanomezantsoa Rafetrarivony, Lalainasoa Odile Rivoarilala, Vincent Enouf, Annick Lalaina Robinson, et al.. Draft Genome Sequence of a *Chryseobacterium indologenes* Strain Isolated from a Blood Culture of a Hospitalized Child in Antananarivo, Madagascar. *Microbiology Resource Announcements*, 2019, 8 (35), pp.e00752-19. 10.1128/MRA.00752-19. hal-02875195

HAL Id: hal-02875195

<https://hal.science/hal-02875195v1>

Submitted on 14 Aug 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Draft Genome Sequence of a *Chryseobacterium indologenes* Strain Isolated from a Blood Culture of a Hospitalized Child in Antananarivo, Madagascar

Mamitina Alain Noah Rabenandrasana,<sup>a</sup> Lala Fanomezantsoa Rafetrarivony,<sup>a</sup> Lalaisoa Odile Rivoarilala,<sup>a</sup> Vincent Enouf,<sup>e</sup> Annick Lalaina Robinson,<sup>d</sup> Ando Rakotozanany,<sup>d</sup> Jessica Vanhomwegen,<sup>b</sup> Valérie Caro,<sup>b</sup> Cassandre Von Platen,<sup>c</sup> Jean-Claude Manuguerra,<sup>b</sup> Jean-Marc Collard<sup>a</sup>

<sup>a</sup>Experimental Bacteriology Unit, Institut Pasteur de Madagascar, Antananarivo, Madagascar

<sup>b</sup>Laboratory for Urgent Response to Biological Threats, Institut Pasteur, Paris, France

<sup>c</sup>Center for Translational Science, Institut Pasteur, Paris, France

<sup>d</sup>Centre Hospitalier Universitaire Mère-Enfant Tsaralalàna (CHUMET), Antananarivo, Madagascar

<sup>e</sup>Pasteur International Bioresources Network (PIBnet), Plateforme de Microbiologie Mutualisée (P2M), Institut Pasteur, Paris, France

**ABSTRACT** We report here the draft genome sequence of a *Chryseobacterium indologenes* strain, isolated from a blood culture of a 2.2-year-old child admitted to the hospital for vomiting and coughing. The genome was composed of 5,063,674 bp and had 37.04% GC content. We detected 4,796 genes with predicted protein-coding functions, including those associated with antibiotic resistance.

**B**acteria of the *Chryseobacterium* genus are nonmotile, chemoorganotrophic, and glucose-nonfermentative Gram-negative rod-shaped bacteria. The most pathogenic species of the genus, *Chryseobacterium meningosepticum*, which causes numerous infections, was reclassified to the genus *Elizabethkingia* (1). *Chryseobacterium indologenes*, although ubiquitous in nature and found mainly in soil and water, is an uncommon human pathogen. However, in rare cases, it can cause serious infections, particularly among immunocompromised hospitalized patients with severe underlying diseases and/or with indwelling catheters (2). These include keratitis, bacteremia (2), pneumonia, cellulitis, and artificial shunt infection (3–6).

*C. indologenes* was isolated from a positive blood culture (time to positivity, 23 h) collected by venipuncture from a 2.2-year-old child hospitalized at the pediatric hospital of Tsaralalàna, Antananarivo, Madagascar. This child was admitted 1 day before blood sampling with fever, coughing, and vomiting. The study was approved by the Ministry of Health and the National Ethics Committee of Madagascar (number 140-MSANP/CE). Informed written consent was obtained from at least one child's parent before sampling.

The strain forms on solid blood agar in typical yellow colonies due to a flexirubin-type pigment. After 24 h, the colonies were identified by matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) as *C. indologenes*, with a score of >2.0. The antimicrobial susceptibility testing was performed according to the standard disc methods described in the CA-SFM guidelines (7), and susceptibility to chloramphenicol and tetracycline was also tested. The isolate was resistant to ticarcillin, piperacillin, ticarcillin plus clavulanate, tazobactam, ceftazidime, aztreonam, imipenem, gentamicin, tobramycin, amikacin, and ciprofloxacin, displayed an intermediate resistance to chloramphenicol, and remained susceptible to cefepime and tetracycline.

DNA extraction was done from an overnight culture using a DNA blood and tissue kit (Qiagen, France). A DNA sequencing library was prepared using a Nextera XT DNA sample prep kit (Illumina, Inc., San Diego, CA), according to the manufacturer's instruc-

**Citation** Rabenandrasana MAN, Rafetrarivony LF, Rivoarilala LO, Enouf V, Robinson AL, Rakotozanany A, Vanhomwegen J, Caro V, Von Platen C, Manuguerra J-C, Collard J-M. 2019. Draft genome sequence of a *Chryseobacterium indologenes* strain isolated from a blood culture of a hospitalized child in Antananarivo, Madagascar. *Microbiol Resour Announc* 8:e00752-19. <https://doi.org/10.1128/MRA.00752-19>.

**Editor** Catherine Putonti, Loyola University Chicago

**Copyright** © 2019 Rabenandrasana et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mamitina Alain Noah Rabenandrasana, [rabalainnoah@gmail.com](mailto:rabalainnoah@gmail.com).

**Received** 28 June 2019

**Accepted** 4 August 2019

**Published** 29 August 2019

**TABLE 1** Antimicrobial resistance genes found in a *C. indologenes* strain isolated in Antananarivo, Madagascar

Resistance gene	Function	Classification
<i>tet(X)</i>	Tetracycline-inactivating enzyme	Antibiotic inactivation enzyme
<i>bla</i> <sub>IND-2a</sub>	Subclass B1 $\beta$ -lactamase	Antibiotic inactivation enzyme
<i>katG</i>	Catalase peroxidase	Antibiotic inactivation enzyme
<i>ant(6)-I</i>	Aminoglycoside 6-nucleotidyltransferase	Antibiotic inactivation enzyme
<i>bla</i> <sub>CIA</sub>	Class A extended-spectrum $\beta$ -lactamase	Antibiotic inactivation enzyme
<i>catB</i>	Chloramphenicol <i>O</i> -acetyltransferase	Antibiotic inactivation enzyme
<i>oxyR</i>	Hydrogen peroxide-inducible gene activator	Regulator modulating expression of antibiotic resistance genes

tions. Genome sequencing was performed using MiSeq Illumina technology ( $2 \times 300$  bp), generating 3,772,882 paired-end reads. FqCleaner version 3.0 was used to eliminate adapter sequences (8, 9), reduce redundant or overrepresented reads (10), correct sequencing errors (11), merge overlapping paired reads, and discard reads with a Phred score (measure of the quality of identification of nucleobases generated by automated DNA sequencing) of  $<20$ . Illumina read *de novo* assembly was performed using SPAdes version 3.10.0 (12, 13), with default parameters. Acquired resistance genes were detected using ResFinder version 3.0 (14). The genome was annotated using the PATRIC Web server version 3.5.39 (15).

In total, 48 assembled contigs with an  $N_{50}$  value of 193,241 bp and an average coverage of  $36.1\times$ , which had 5,063,674 bp and 37.04% GC content, were annotated using the PATRIC Web server annotation pipeline (15–19). The genome harbors 64 tRNA genes, 3 rRNA genes, and 4,796 protein-coding sequences. The antimicrobial resistance (AMR) genes detected in this genome and the corresponding AMR mechanisms are provided in Table 1.

The extended-spectrum  $\beta$ -lactamase gene *bla*<sub>CIA</sub> was responsible for the resistance to ticarcillin, piperacillin, ticarcillin plus clavulanate, tazobactam, and ceftazidime but not to cefepime. A metallo- $\beta$ -lactamase gene, *bla*<sub>IND-2a</sub>, was responsible for the resistance to imipenem. Our phenotypic results showed that the *catB* gene did not confer complete resistance to chloramphenicol and that the *tet(X)* gene was nonfunctional, like that originally found in *Bacteroides* spp. (20–23). The Tet(X) protein showed 62% identity with Tet(X) from the strain AF25-18 (GenBank accession number [RJV36860](#)) and 62% with Tet(X) from *Chryseobacterium oncorhynchi* (GenBank accession number [PWN64762](#)). Studies on hosted bacteriophages, secretion systems, mobility, and ability to form biofilms would allow us to understand the pathogenic potential of our *C. indologenes* isolate and its role in bacteremia.

**Data availability.** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [VCBR00000000](#). The version described in this paper is version VCBR01000000. Raw sequence data for this strain were deposited under SRA accession number [PRJNA551338](#).

## ACKNOWLEDGMENTS

We thank Norohasina Randriamanga, the field investigator, and the staff of pediatric hospital Tsaralàna, Antananarivo, Madagascar.

We thank the Programme Transversal de Recherche (PTR) Child's Play number 471 funded by the Institut Pasteur (Paris, France), which supported this work, and the staff of the Plateforme de Microbiologie Mutualisée (P2M) at Institut Pasteur Paris, where the whole-genome sequencing was performed.

## REFERENCES

- Kim KK, Kim MK, Lim JH, Park HL. 2005. Transfer of *Chryseobacterium meningosepticum* and *Chryseobacterium miricola* to *Elizabethkingia* gen. nov. as *Elizabethkingia meningoseptica* comb. nov. and *Elizabethkingia miricola* comb. nov. *Int J Syst Evol Microbiol* 55:1287–1293. <https://doi.org/10.1099/ijs.0.63541-0>.
- Deng L, Li MF, Li YH, Yang JL, Zhou X. 2015. *Chryseobacterium indolo-*

- genes in four patients with leukemia. *Transpl Infect Dis* 17:583–587. <https://doi.org/10.1111/tid.12400>.
3. Lu PC, Chan JC. 1997. Flavobacterium indologenes keratitis. *Ophthalmologica* 211:98–100. <https://doi.org/10.1159/000310769>.
  4. Kienzle N, Muller M, Pegg S. 2001. Chryseobacterium in burn wounds. *Burns* 27:179–182. [https://doi.org/10.1016/S0305-4179\(00\)00087-5](https://doi.org/10.1016/S0305-4179(00)00087-5).
  5. Doiz O, Llorente MT, Mateo A, Seral C, García C, Rubio MC. 1999. Corneal abscess by Flavobacterium indologenes. A case report. *Enferm Infecc Microbiol Clin* 17:149–150. (In Spanish.)
  6. Green BT, Nolan PE. 2001. Cellulitis and bacteraemia due to Chryseobacterium indologenes. *J Infect* 42:219–220. <https://doi.org/10.1053/jinf.2001.0822>.
  7. Société Française de Microbiologie. 2017. Communiqué 2017. Comité de l'antibiogramme de la Société Française de Microbiologie (CA-SFM)/ European Committee on Antimicrobial Susceptibility Testing (EUCAST), Paris, France.
  8. Criscuolo A, Brisse S. 2013. AlienTrimmer: a tool to quickly and accurately trim off multiple short contaminant sequences from high-throughput sequencing reads. *Genomics* 102:500–506. <https://doi.org/10.1016/j.ygeno.2013.07.011>.
  9. Crusoe MR, Alameldin HF, Awad S, Boucher E, Caldwell A, Cartwright R, Charbonneau A, Constantinides B, Edverson G, Fay S, Fenton J, Fenzl T, Fish J, Garcia-Gutierrez L, Garland P, Gluck J, González I, Guermont S, Guo J, Gupta A, Herr JR, Howe A, Hyer A, Härpfer A, Irber L, Kidd R, Lin D, Lippi J, Mansour T, McA'Nulty P, McDonald E, Mizzi J, Murray KD, Nahum JR, Nanlohy K, Nederbragt AJ, Ortiz-Zuazaga H, Ory J, Pell J, Pepe-Rannek C, Russ ZN, Schwarz E, Scott C, Seaman J, Sievert S, Simpson J, Skennerton CT, Spencer J, Srinivasan R, Standage D, et al. 2015. The khmer software package: enabling efficient nucleotide sequence analysis. *F1000Res* 4:900. <https://doi.org/10.12688/f1000research.6924.1>.
  10. Liu Y, Schröder J, Schmidt B. 2013. Musket: a multistage k-mer spectrum-based error corrector for Illumina sequence data. *Bioinformatics* 29:308–315. <https://doi.org/10.1093/bioinformatics/bts690>.
  11. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. 2012. Predicting the functional effect of amino acid substitutions and indels. *PLoS One* 7:e46688. <https://doi.org/10.1371/journal.pone.0046688>.
  12. Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, Pribelski AD, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, Clingenpeel SR, Woyke T, McLean JS, Lasken R, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling genomes and mini-metagenomes from highly chimeric reads, p 158–170. *In* Deng M, Jiang R, Sun F, Zhang X (ed), *Research in computational molecular biology*. Springer, Berlin, Germany.
  13. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
  14. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
  15. Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, Conrad N, Dietrich EM, Disz T, Gabbard JL, Gerdes S, Henry CS, Kenyon RW, Machi D, Mao C, Nordberg EK, Olsen GJ, Murphy-Olson DE, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Vonstein V, Warren A, Xia F, Yoo H, Stevens RL. 2017. Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource. *Nucleic Acids Res* 45:D535–D542. <https://doi.org/10.1093/nar/gkw1017>.
  16. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res* 44:D457–D462. <https://doi.org/10.1093/nar/gkv1070>.
  17. Davis JJ, Gerdes S, Olsen GJ, Olson R, Pusch GD, Shukla M, Vonstein V, Wattam AR, Yoo H. 2016. PATTyFams: protein families for the microbial genomes in the PATRIC database. *Front Microbiol* 7:118. <https://doi.org/10.3389/fmicb.2016.00118>.
  18. Schomburg I, Chang A, Ebeling C, Gremse M, Heldt C, Huhn G, Schomburg D. 2004. BRENDA, the enzyme database: updates and major new developments. *Nucleic Acids Res* 32:D431–D433. <https://doi.org/10.1093/nar/gkh081>.
  19. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene Ontology: tool for the unification of biology. *Nat Genet* 25:25–29. <https://doi.org/10.1038/75556>.
  20. Markley JL, Wenczewicz TA. 2018. Tetracycline-inactivating enzymes. *Front Microbiol* 9:1058. <https://doi.org/10.3389/fmicb.2018.01058>.
  21. Leski TA, Bangura U, Jimmy DH, Ansumana R, Lizewski SE, Stenger DA, Taitt CR, Vora GJ. 2013. Multidrug-resistant tet(X)-containing hospital isolates in Sierra Leone. *Int J Antimicrob Agents* 42:83–86. <https://doi.org/10.1016/j.ijantimicag.2013.04.014>.
  22. Ghosh S, LaPara TM, Sadowsky MJ. 2014. Draft genome sequence of Sphingobacterium sp. strain PM2-P1-29. *Genome Announc* 2:e00963-14. <https://doi.org/10.1128/genomeA.00963-14>.
  23. Zeng J, Pan Y, Yang J, Hou M, Zeng Z, Xiong W. 2019. Metagenomic insights into the distribution of antibiotic resistance between the gut-associated environments and the pristine environments. *Environ Int* 126:346–354. <https://doi.org/10.1016/j.envint.2019.02.052>.