



HAL
open science

Entropy solutions for a two-phase transition model for vehicular traffic with metastable phase and time depending point constraint on the density flow

Boris Andreianov, Carlotta Donadello, Massimiliano D. Rosini

► To cite this version:

Boris Andreianov, Carlotta Donadello, Massimiliano D. Rosini. Entropy solutions for a two-phase transition model for vehicular traffic with metastable phase and time depending point constraint on the density flow. 2020. hal-02877276v1

HAL Id: hal-02877276

<https://hal.science/hal-02877276v1>

Preprint submitted on 22 Jun 2020 (v1), last revised 12 Mar 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Entropy solutions for a two-phase transition model for vehicular traffic with metastable phase and time depending point constraint on the density flow

Boris Andreianov, Carlotta Donadello, Massimiliano D. Rosini

June 21, 2020

Abstract

We consider a macroscopic two-phase transition model for vehicular traffic flow subject to a point constraint on the density flux. The two phases correspond to light and heavy traffic and their dynamics are described respectively by the Lighthill-Whitham-Richards model and the Aw-Rascle-Zhang model. Their intersection, the so-called metastable phase, is assumed to be non empty. In order to guarantee the control of total variation of solutions, the t -dependent function prescribing the level of the constraint should be piecewise constant. It can be given *a priori* or it can be computed in a time-discrete way from the solution itself, thus giving rise to nonlocal phase transition models with point constraint in the spirit of LWR-based models described in *Andreianov, Donadello, Razafison, Rosini, JMPA (2018)*.

We introduce a new definition of admissible solutions for the Cauchy problem, for which we prove existence and we provide a characterization. In particular, these admissible solutions attain the maximal flux allowed by the constraint whenever it is enforced, which guarantees compatibility of the constructed solutions with the modeling assumption imposed at the level of the Riemann solver. These results rely on the wave front tracking method and on adaptation of the specific entropies and renormalization properties introduced in *Andreianov, Donadello, Rosini, M3AS (2016)* while dealing with the ARZ system.

2010 Mathematical Subject Classification. Primary: 35L65, 90B20, 35L45

Key words. Conservation laws, phase transitions, Lighthill-Whitham-Richards model, Aw-Rascle-Zhang model, point constraint on the density flux, wave-front tracking, entropy conditions.

1 Introduction

Our work enters the framework of so-called phase transition models used in macroscopic description of road traffic. In this paper we prove an existence result for a constrained two-phase transition model of hyperbolic conservation laws introduced in [18]. Such model reproduces the vehicular traffic along a road with pointlike inhomogeneities with limited capacity, such as construction sites, tollgates, traffic lights, etc. Our work is a continuation and an extension of the work [9].

We consider two phases corresponding to the *congested phase* Ω_c and the *free phase* Ω_f . In the congested phase the traffic is governed by a 2×2 system of conservation laws, whereas in the free phase it is governed by a scalar conservation law. The two phases are coupled via *phase transitions*, namely discontinuities between two states belonging to different phases and satisfying the Rankine-Hugoniot conditions.

The first two-phase model has been proposed by Colombo in [15]. Later, Goatin proposed in [22] a two-phase model which couples the Aw, Rascle and Zhang [8, 27] (ARZ) model for the congested phase Ω_c , with the Lighthill, Whitham and Richards [24, 26] (LWR) model for the free-flow phase Ω_f , see also [10] for its generalization.

Both Colombo [15] and Goatin [22] assume that $\Omega_c \cap \Omega_f = \emptyset$. The first two-phase model with a *metastable phase* $\Omega_c \cap \Omega_f \neq \emptyset$ has been introduced in [13]. Differently from [10, 22], for the models in [13, 15] the density flux function vanishes at a unique maximal density.

Existence results for Cauchy problems have already been established in [10, 13, 17, 22] for the case without point constraints, and in [9] for the case with point constraint.

In the present paper we assume that $\Omega_c \cap \Omega_f \neq \emptyset$. For this reason, see [15, Remark 2], we assume that Ω_f is characterized by a unique value V of the velocity. In other words, the scalar conservation law describing the dynamics of the free phase is merely the transport equation with constant velocity V . We consider vehicles having different lengths, hence we allow the density flux function to vanish at different densities. At last, as in [9, 11, 18], we impose that at the interface $x = 0$ the density flux of the solution is lower than a value

F . However, differently from [9, 11, 18], here F is time dependent (restricted to be piecewise constant, with sufficiently high level of passing capacity). In particular our setting includes constraint functions whose values, updated at fixed times, depend on the past evolution of the solution. This allows to model the traffic in presence, for instance, of traffic lights or construction sites operating only during a part of the day, as well as their possible adaptative management.

With respect to [9], our contribution is twofold.

First, we show that the wave-front tracking algorithm gives at the limit an entropy solution; this allows to characterize the flow across the point constraints without technical assumption [9, formula (2.14)]. We strengthen the definition of entropy solution so to include the sharp flux characterization at the constraint. According to our definition, non-classical shocks occur precisely at the level of the flux imposed by the constraint; moreover, the lagrangian marker does not increase across the non-classical shock. Note that these two properties underlied the resolution of the Riemann problems in [9]. Thus, the sharp characterization of admissible solutions adequately reflects the modeling assumptions imposed while prescribing the Riemann solver. To achieve this flux characterization at the constraint, we rely upon a localized version of the renormalization property (cf. [4]) and indagate the value of the limit density flux at the constraint position through a classical application of the Green theorem. Note that the passage to the limit that underlies the existence result exploits the careful choice of the entropy inequalities (or, more precisely, of the contribution of the point constraint to these inequalities) and relies again upon the renormalization property.

Second, our existence result allows to consider time depending constraints under an additional restriction on possible constraint levels. This opens way to the study of modeling situations where a non-local point constraint is imposed at the bottleneck. Analogous models based on Lighthill-Whitham-Richard equation were constructed and analyzed in [2]; here, we restrict our attention to constraints updated at discrete times. Models with continuously varying constraint allows to reproduce capacity drop leading to non-monotone empirical features of real traffic flows such as Faster is Slower and the Braess Paradox (see [1, 3]). Contrarily to [1, 3], here we limit our attention to the situations where the constraint and its variations do not result from the intrinsic desorganization (or, on the contrary, from a self-organization, see [2, 7]) of the flow at high densities. In the models we have in mind in this paper, the constraint and its variations arise from operation of bottlenecks (such as toll gates) at discrete times as a function of data collected non-locally in time upstream the flow. Our restriction to piecewise constant in time constraints F is motivated by technical reasons. In particular, since our existence proof relies on wave front tracking approximations, we need to monitor the increase in total variation of the approximate solutions at any time at which the constraint level changes. In practice this is possible provided changes only occur a locally finite number of times, and the updated constraint level lays in the interval where metastable states exist. In our opinion, the most promising direction to overcome these technical restriction requires different compactness techniques, e.g. of the compensated compactness type; work in this direction is the subject of ongoing research.

The paper is organized as follows. In the next section we give a brief introduction to the model and to the state of the art. In Section 2 we introduce the needed notations, the model and the main result of the paper, which concerns constraint given beforehand. We defer the proofs of our main results to Sections 3 and 4. In the last section we briefly indicate how our results can be applied to a particular class of nonlocal point constraints.

2 Model and main result

In this section we introduce some notations, see Figure 1, recall the two-phase transition model (3), give the definition of entropy solutions in Definition 2.1 and state our main result in Theorem 2.8.

Let $\rho \geq 0$ and $v \geq 0$ be the density and the velocity of the vehicles, respectively. Denote $u \doteq (\rho, v)$ and let $f(u) \doteq v\rho$ be the density flux. If $V > 0$ is the unique velocity in the free-flow phase Ω_f and ρ^+ is the maximal density in Ω_f , then

$$\Omega_f \doteq \{u \in \mathbb{R}_+^2 : \rho \leq \rho^+, v = V\},$$

where $\mathbb{R}_+ \doteq [0, \infty)$. If $[\rho^-, \rho^+] \times \{V\}$ is the metastable phase, i.e. the intersection between Ω_f and the congested phase Ω_c , then

$$\Omega_c \doteq \{u \in \mathbb{R}_+^2 : v \leq V, w^- \leq v + p(\rho) \leq w^+\},$$

where $w^\pm \doteq p(\rho^\pm) + V$. Above $p \in \mathbf{C}^2((0, \infty); \mathbb{R})$ is an anticipation factor, which takes into account drivers' reactions to the state of traffic in front of them. We assume that

$$p'(\rho) > 0, \quad 2p'(\rho) + p''(\rho)\rho > 0 \quad \text{for every } \rho > 0, \quad (1)$$

and

$$v < p'(\rho)\rho \text{ for every } (\rho, v) \in \Omega_c. \quad (2)$$




Figure 1: Notations.

Typical choices for p are $p(\rho) \doteq \rho^\gamma$ with $\gamma > V/(w^- - V)$, see [8], and $p(\rho) \doteq V_{\text{ref}} \ln(\rho/\rho_{\text{max}})$ with $V_{\text{ref}} > V$ and $\rho_{\text{max}} > 0$, see [22]. Denote $f^\pm \doteq V \rho^\pm$ and let $R \doteq p^{-1}(w^+) > 0$ be the maximal density (in the congested phase). Define

$$\Omega \doteq \Omega_f \cup \Omega_c, \quad \Omega_f^- \doteq \{u \in \Omega_f : \rho \in [0, \rho^-]\}, \quad \Omega_f^+ \doteq \{u \in \Omega_f : \rho \in [\rho^-, \rho^+]\}, \quad \Omega_c^- \doteq \Omega_c \setminus \Omega_f^+.$$

Notice that $\Omega_f \cap \Omega_c = \Omega_f^+$ is the metastable phase. The (extended) Lagrangian marker $\mathbf{w}: \Omega \rightarrow [w^- - 1, w^+]$ is defined by

$$\mathbf{w}(u) \doteq \begin{cases} v + p(\rho) & \text{if } u \in \Omega_c, \\ w^- - 1 + \frac{\rho}{\rho^-} & \text{if } u \in \Omega_f^-. \end{cases}$$

Let $\mathbf{W}: \Omega \rightarrow [w^-, w^+]$ be defined by

$$\mathbf{W}(u) \doteq \begin{cases} v + p(\rho) & \text{if } u \in \Omega_c, \\ w^- & \text{if } u \in \Omega_f^-. \end{cases}$$

In this paper we study the constrained Cauchy problem for the phase transition model

$$\begin{array}{ll} \text{Free-flow (LWR)} & \text{Congested flow (ARZ)} \\ \begin{cases} u \in \Omega_f, \\ \rho_t + (\rho V)_x = 0, \\ v = V, \end{cases} & \begin{cases} u \in \Omega_c, \\ \rho_t + (\rho v)_x = 0, \\ (\rho \mathbf{W}(u))_t + (\rho \mathbf{W}(u) v)_x = 0, \end{cases} \end{array} \quad (3a)$$

with initial condition

$$u(0, x) = \bar{u}(x) \quad (3b)$$

and local point constraint on the density flux at $x = 0$

$$f(u(t, 0_\pm)) \leq \mathbf{F}(t), \quad (3c)$$

where $\mathbf{F}: (0, \infty) \rightarrow [0, f^+]$ is a given function. Note that in Section 5 we will explain how to deal with piecewise constant constraint functions \mathbf{F} such that, if t_i and t_{i+1} are two consequent times at which \mathbf{F} is discontinuous, the value of $\mathbf{F}(t)$ for $t \in (t_i, t_{i+1})$ depend on $u|_{[0, t_i] \times \mathbb{R}}$.

For $u \in \Omega$, $k \in [0, V]$ and $F \in [0, f^+]$ we define

$$\mathbf{N}_F^k(u) \doteq \begin{cases} f(u) \mathbf{n}_F^k(\mathbf{W}(u)) & \text{if } F \neq 0, \\ k & \text{if } F = 0, \end{cases} \quad \mathbf{n}_F^k(\mathbf{W}) \doteq \left[\frac{k}{F} - \frac{1}{p^{-1}(\mathbf{W} - k)} \right]_+, \quad (4)$$

and introduce the entropy-entropy flux pair

$$\mathbf{E}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{\rho}{p^{-1}(\mathbf{W}(u) - k)} - 1 & \text{if } v < k, \end{cases} \quad \mathbf{Q}^k(u) \doteq \begin{cases} 0 & \text{if } v \geq k, \\ \frac{f(u)}{p^{-1}(\mathbf{W}(u) - k)} - k & \text{if } v < k, \end{cases} \quad (5)$$

which is obtained by adapting the entropy-entropy flux pair introduced in [4] for the ARZ model.

Definition 2.1. Let $\bar{u} \in \mathbf{BV}(\mathbb{R}; \Omega)$. Let $\mathbf{F} \in \mathbf{L}^\infty((0, \infty); [0, f_+])$. We say that $u \in \mathbf{L}^\infty((0, \infty); \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ is an admissible solution to constrained Cauchy problem (3) if the following holds:

(S.1) Initial condition (3b) holds for a.e. $x \in \mathbb{R}$, namely (bearing in mind the time continuity with $\mathbf{L}_{\text{loc}}^1$ values)

$$u(0, x) = \bar{u}(x) \quad \text{for a.e. } x \in \mathbb{R}.$$

(S.2) The function u provides a weak solution to the mass conservation equation, namely, for any $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ we have

$$\int_0^\infty \int_{\mathbb{R}} (\rho \phi_t + f(u) \phi_x) \, dx \, dt = 0. \quad (6)$$

(S.3) The function w satisfies the weak formulation and the renormalization property away from the constraint, namely, for any $g \in \mathbf{C}([w^-, w^+]; \mathbb{R})$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi(\cdot, 0) \equiv 0$, we have that

$$\int_0^\infty \int_{\mathbb{R}} (\mathbf{E}_g(u) \phi_t + \mathbf{Q}_g(u) \phi_x) \, dx \, dt = 0, \quad (7)$$

where

$$\mathbf{E}_g(u) \doteq \rho g(\mathbf{W}(u)), \quad \mathbf{Q}_g(u) \doteq v \mathbf{E}_g(u).$$

(S.4) Entropy inequalities are satisfied up to the constraint, namely, for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$ we have

$$\int_0^\infty \left(\int_{\mathbb{R}} (\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x) \, dx + \mathbf{N}_{\mathbf{F}(t)}^k(u(t, 0_-)) \phi(t, 0) \right) dt \geq 0. \quad (8)$$

(S.5) The constraint condition (3c) holds for a.e. $t > 0$, namely

$$f(u(t, 0_\pm)) \leq \mathbf{F}(t) \quad \text{for a.e. } t > 0.$$

Remark 2.2. It is possible to prove, using the delicate theory of [25], that as soon as (7) holds with $g = \text{Id}$ (which corresponds to the mere weak formulation) and (6) holds (that is, the field $(\rho, \rho v)$ is divergence-free), the formulation (7) holds with arbitrary g . We include the renormalization property into the definition because of its importance in the derivation of entropy inequalities and also because it is easily proved at the level of approximate solutions constructed with the wave-front tracking algorithm.

Remark 2.3. A solution u to constrained Cauchy problem (3) does not satisfy in general the second Rankine-Hugoniot condition (10) along $x = 0$

$$\rho(t, 0_-) \mathbf{W}(u(t, 0_-)) v(t, 0_-) = \rho(t, 0_+) \mathbf{W}(u(t, 0_+)) v(t, 0_+) \quad \text{for a.e. } t > 0.$$

Indeed the (extended) linearized momentum $\rho \mathbf{W}(u)$ is conserved across (classical) shocks and phase transitions, but in general it is not conserved across non-classical shocks even if they are between states in Ω_c . As a consequence, a solution to (3) taking values in Ω_c is not necessarily a weak solution to the 2×2 system of conservation laws in (3a) for the congested flow. For this reason in (7) we consider test functions ϕ such that $\phi(\cdot, 0) \equiv 0$. This is in the same spirit of the solutions considered in [11, 18–21] for traffic through locations with reduced capacity.

Differently from [9], in this paper we do not require in (8) that $\phi(\cdot, 0) \equiv 0$. This allows us to better characterize the (density) flux at $x = 0$ associated to non-classical shocks. In fact, we can ensure that the flux of the non-classical shocks of any solution is equal to the maximal flux \mathbf{F} allowed by the constraint without requiring the technical assumption [9, (2.14)], as pointed out in Proposition 2.4 below. We can also ensure that the lack of conservativity in the generalized momentum equation is sign-definite, namely $\int_{\mathbb{R}} \rho(t, x) \mathbf{W}(u(t, x)) \, dx$ is non-increasing with t : this is an immediate consequence of (11) below.

In the following proposition we state precisely which discontinuities are admissible for the solutions to (3).

Proposition 2.4. Let u be a solution of constrained Cauchy problem (3) in the sense of Definition 2.1. Assume that \mathbf{F} is a piecewise constant function. Then u has the following properties:

- At any Lipschitz curve of discontinuity $x = \delta(t)$ of u , the first Rankine-Hugoniot jump condition holds for a.e. $t > 0$:

$$\left[\rho(t, \delta(t)_+) - \rho(t, \delta(t)_-) \right] \dot{\delta}(t) = f(u(t, \delta(t)_+)) - f(u(t, \delta(t)_-)), \quad (9)$$

and if $\delta(t) \neq 0$, then it satisfies also the second Rankine-Hugoniot jump condition

$$\begin{aligned} & \left[\rho(t, \delta(t)_+) \mathbb{W}(u(t, \delta(t)_+)) - \rho(t, \delta(t)_-) \mathbb{W}(u(t, \delta(t)_-)) \right] \dot{\delta}(t) \\ &= f(u(t, \delta(t)_+)) \mathbb{W}(u(t, \delta(t)_+)) - f(u(t, \delta(t)_-)) \mathbb{W}(u(t, \delta(t)_-)). \end{aligned} \quad (10)$$

- Any discontinuity of u away from the constraint location $x = 0$ is classical, i.e., it satisfies the Lax entropy inequalities.

- Non-classical discontinuities of u may occur only at $x = 0$, and in this case the (density) flux $f(u(t, 0_{\pm}))$ at $x = 0$ equals the maximal flux \mathbf{F} allowed by the constraint.

Moreover, whatever be the nature of the shock at $x = 0$, there holds

$$f(u(t, 0_{\pm})) \left(\mathbb{W}(u(t, 0_-)) - \mathbb{W}(u(t, 0_+)) \right) \geq 0 \text{ for a.e. } t > 0. \quad (11)$$

The proof is deferred to Section 4.

Note that the assumption of piecewise constancy of \mathbf{F} can be weakened via localisation arguments (cf. [4, Prop. 3.2]); however, we do not go beyond the piecewise constant setting in the rest of this paper, because of the difficulty of controlling the variation of the approximate solutions constructed with wave-front tracking.

Remark 2.5. The characterization of non-classical shocks at $x = 0$ by the constraint saturation condition $f(u(t, 0_{\pm})) = \mathbf{F}$, achieved in Proposition 2.4 for the model at hand, was the cornerstone of the uniqueness results in the LWR point constrained models, [6, 16], and it holds true for the constrained ARZ model, [4]. Note that for phase transition models without the metastable phase, the constraint saturation property can not be true in all situations, see [12].

In order to describe precisely non-classical shocks and define the functional designed to control the total variation of the wave-front tracking solutions, let us introduce $v_F^{\pm} \in [0, V]$ and $w_F \in [w^- - 1, w^+]$ defined by the following conditions, see Figure 2:

$$\begin{aligned} \text{if } F = f^+ : & \quad v_F^+ \doteq V, & \quad v_F^- \doteq V, & \quad w_F \doteq w^+, \\ \text{if } F \in [f^-, f^+) : & \quad v_F^+ \doteq V, & \quad v_F^- + p(F/v_F^-) = w^+, & \quad w_F \doteq p(F/V) + V, \\ \text{if } F \in (0, f^-) : & \quad v_F^+ + p(F/v_F^+) = w^-, & \quad v_F^- + p(F/v_F^-) = w^+, & \quad w_F \doteq w^- - 1 + \frac{F}{f^-}, \\ \text{if } F = 0 : & \quad v_F^+ \doteq 0, & \quad v_F^- \doteq 0, & \quad w_F \doteq w^- - 1. \end{aligned}$$




Figure 2: Geometrical meaning of w_F , v_F^{\pm} and Ξ_F in the case $F \in (0, f^-)$. The curve in the figure on the left is the graph of Ξ_F , which corresponds to the horizontal solid segment in the figure on the right.

For any $F \in (0, f^+)$, let $\Xi_F : [v_F^-, v_F^+] \rightarrow [w^-, w^+]$ be given by $\Xi_F(v) \doteq v + p(F/v)$, see Figure 2. Notice that Ξ_F is strictly decreasing by (2) and is strictly convex by (1).

For any $F \in (0, f^+)$, let $[w^- - 1, w^+] \ni w \mapsto \hat{u}(w, F) = (\hat{x}(w, F), \hat{v}(w, F)) \in \Omega_c$ and $[0, V] \ni v \mapsto \check{u}(v, F) = (\check{x}(v, F), \check{v}(v, F)) \in \Omega$ be defined in the (v, w) -coordinates by

$$\hat{v}(w, F) \doteq \begin{cases} \Xi_F^{-1}(w) & \text{if } w > \max\{w^-, w_F\}, \\ v_F^+ & \text{if } w_F < w \leq w^-, \\ V & \text{if } w \leq w_F, \end{cases} \quad \hat{w}(w, F) \doteq \begin{cases} w & \text{if } w > \max\{w^-, w_F\}, \\ w^- & \text{if } w_F < w \leq w^-, \\ w_F & \text{if } w \leq w_F, \end{cases} \quad (12)$$

$$\check{v}(v, F) \doteq \begin{cases} V & \text{if } v > v_F^+, \\ v & \text{if } v \in [v_F^-, v_F^+], \\ v_F^- & \text{if } v < v_F^-, \end{cases} \quad \check{w}(v, F) \doteq \begin{cases} w_F & \text{if } v > v_F^+, \\ \Xi_F(v) & \text{if } v \in [v_F^-, v_F^+], \\ w^+ & \text{if } v < v_F^-, \end{cases}$$

where $\hat{w} \doteq \mathbf{w} \circ \hat{u}$ and $\check{w} \doteq \mathbf{w} \circ \check{u}$, see Figures 3 and 4.

The following lemma collects some useful properties of the maps defined above. In particular, it explains why we limit our study to the case of F taking values in $[f^-, f^+]$.

Lemma 2.6. 1. For any $F \in (0, f^+)$, $w \in [w^- - 1, w^+]$ and $v \in [0, V]$ we have

$$f(\hat{u}(w, F)) = f(\check{u}(v, F)) = F.$$

2. The maps $w \mapsto \hat{u}(w, F)$ and $v \mapsto \check{u}(v, F)$ are Lipschitz continuous if and only if $F \geq f^-$. The Lipschitz constant is then uniform with respect to $F \in [f^-, f^+]$;
3. If $F < f^-$, then $w \mapsto \hat{u}(w, F)$ and $v \mapsto \check{u}(v, F)$ are only left-continuous;
4. $\hat{w}(w, F) \geq w$ and $\check{v}(v, F) \geq v$.
5. $w \mapsto \hat{w}(w, F)$ and $v \mapsto \check{v}(v, F)$ are non-decreasing, while $w \mapsto \hat{v}(w, F)$ and $v \mapsto \check{w}(v, F)$ are non-increasing.

Proof. We focus on the proof of 2, which is crucial for the analysis of increase of the Glimm-like functionals providing the variation control. The other properties can be assessed analogously, upon examination of the definitions. The uniform in F Lipschitz continuity of \hat{w} and \check{v} is obvious. The uniform Lipschitz regularity of \hat{v} and of \check{w} requires uniform bounds on $\Xi_F^{-1}(w)$ and $(\Xi_F^{-1})'(w)$ for $v \in [v_F^-, v_F^+]$ and $w \geq \max\{w^-, w_F\}$. As $w = \Xi_F(v) = v + p(F/v)$ in these calculations, v cannot lie too close to zero. One readily computes

$$\Xi_F'(v) = 1 - p'(F/v)F/v^2 = \frac{v - p'(\rho_F(v))\rho_F(v)}{v},$$

where $\rho_F(v) = F/v$. By assumption (2) (notice that the inequality $v - p(\rho_F(v))\rho_F(v) > 0$ is strict and can be strengthened to $v - p(\rho_F(v))\rho_F(v) \geq \delta > 0$ since p' is continuous and the domain Ω_c is compact), the claimed uniform bounds follow. \square




Figure 3: Geometrical meaning of \hat{u} and \check{u} defined in (12) in the case $F \in (f^-, f^+)$.

Denote by TV_+ and TV_- the positive and negative total variations, respectively. For any $u: \mathbb{R} \rightarrow \Omega$ and $F \in (0, f^+)$ let

$$\hat{Y}(u, F) \doteq \text{TV}_+(\hat{v}(\mathbf{w}(u), F); (-\infty, 0)) + \text{TV}_-(\hat{w}(\mathbf{w}(u), F); (-\infty, 0)), \quad (13)$$

$$\check{Y}(u, F) \doteq \text{TV}_+(\check{v}(v, F); (0, \infty)) + \text{TV}_-(\check{w}(v, F); (0, \infty)).$$

Remark 2.7. Due to Lemma 2.6, if $F \in [f^-, f^+]$ and $u \in \mathbf{BV}(\mathbb{R}; \Omega)$, then $\hat{Y}(u; F) + \check{Y}(u; F)$ are uniformly bounded by a constant times the total variation of u .




Figure 4: Geometrical meaning of \hat{u} and \tilde{u} defined in (12) in the case $F \in (0, f^-)$.

We can now state our main result. Note that it contains essentially the two cases: constant constraint $F \equiv F(0) \in [0, f_+]$ with an additional restriction on nonlinear variations of \bar{u} , and piecewise constant constraint taking values above the threshold f_- . Note that the assumption $F(t) \geq f_-$ corresponds to the presence of metastable states verifying the imposed constraint.

Theorem 2.8. *Let $\bar{u} \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$. Assume that $F \in \mathbf{PC}((0, \infty); [0, f^+])$ satisfies one of the following conditions:*

- (H.1) F takes its values in $[f^-, f^+]$;
- (H.2) $F(t) \equiv F(0) \in [0, f^-)$ and $\hat{Y}(\bar{u}) + \check{Y}(\bar{u})$ is finite.

Then the approximate solutions u_n constructed in Section 3.4 converge to a solution $u \in \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ of constrained Cauchy problem (3) in the sense of Definition 2.1. Moreover for all $t, s \in \mathbb{R}_+$ the following estimates hold

$$\text{TV}(u(t)) \leq K, \quad \|u(t) - u(s)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L|t - s|, \quad \|u(t)\|_{\mathbf{L}^\infty(\mathbb{R}; \Omega)} \leq R + V, \quad (14)$$

where K and L are constants that depend on \bar{u} and F . Furthermore, non-classical discontinuities of u can occur only at the constraint location $x = 0$, and in this case the (density) flow at $x = 0$ and time $t > 0$ is the maximal flow $F(t)$ allowed by the constraint, moreover, (11) is fulfilled.

As in [4, 10, 17], the proof of the above theorem is based on the wave-front tracking algorithm, see [14, 23] and the references therein. The details of the proof are deferred to Section 3.

Corollary 2.9. *The conclusion of Theorem 2.8 still holds true under the following hypothesis*

- $\bar{u} \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$;
- $F \in \mathbf{PC}((0, \infty); [0, f^+])$ and there exists t_1 such that
 - F takes values in $[f_-, f_+]$ for all $t \geq t_1$;
 - $F(t) \equiv F(0) \in [0, f^-)$ for all $t \in [0, t_1)$, and $\hat{Y}(\bar{u}) + \check{Y}(\bar{u})$ is finite.

3 Proof of Theorem 2.8

In this section we prove Theorem 2.8. Note that the justification of the very particular situation of Corollary 2.9 is immediate, the \mathbf{BV} control being guaranteed by the successive application of the arguments used for the case (H.2) and then for the case (H.1) of Theorem 2.8.

3.1 Lax curves

In the (ρ, f) -plane the Lax curves in Ω_c of the first and second characteristic families passing through $\bar{u} = (\bar{\rho}, \bar{v}) \in \Omega_c$ are respectively described by the graphs of the maps

$$\begin{aligned} [p^{-1}(\mathbb{W}(\bar{u}) - V), p^{-1}(\mathbb{W}(\bar{u}))] \ni \rho &\mapsto \mathfrak{L}_{\mathbb{W}(\bar{u})}(\rho) \doteq f(\rho, \mathbb{W}(\bar{u}) - p(\rho)) \in \mathbb{R}_+, \\ [p^{-1}(w^- - \bar{v}), p^{-1}(w^+ - \bar{v})] \ni \rho &\mapsto \bar{v} \rho \in \mathbb{R}_+. \end{aligned}$$

Conditions (1) and (2) ensure that for any $w \in [w^-, w^+]$ the map $\rho \mapsto \mathfrak{L}_w(\rho) = (w - p(\rho)) \rho$ is strictly concave and strictly decreasing in $[p^{-1}(w - V), p^{-1}(w)]$.

We introduce the following functions, see Figure 1:

$$\begin{aligned}
\omega: \Omega_c &\rightarrow \Omega_f^+, & u = \omega(\bar{u}) &\iff \begin{cases} \mathfrak{w}(u) = \mathfrak{w}(\bar{u}), \\ v = V, \end{cases} \\
\mathfrak{v}^\pm: \Omega &\rightarrow \Omega_c, & u = \mathfrak{v}^\pm(\bar{u}) &\iff \begin{cases} \mathfrak{w}(u) = w^\pm, \\ v = \bar{v}, \end{cases} \\
\mathfrak{u}_*: \Omega^2 &\rightarrow \Omega_c, & u = \mathfrak{u}_*(u_\ell, u_r) &\iff \begin{cases} \mathfrak{w}(u) = \mathfrak{W}(u_\ell), \\ v = v_r, \end{cases} \\
\Lambda: \{(u_\ell, u_r) \in \Omega^2 : \rho_\ell \neq \rho_r\} &\rightarrow \mathbb{R}, & \Lambda(u_\ell, u_r) &\doteq \frac{f(u_r) - f(u_\ell)}{\rho_r - \rho_\ell}.
\end{aligned}$$

Notice that:

- the point $\omega(\bar{u})$ is the intersection of Ω_f^+ and the Lax curve of the first characteristic family passing through \bar{u} ;
- for any $w \in [w^-, w^+]$ the point $(p^{-1}(w), 0)$ is the intersection of the Lax curve of the first characteristic family corresponding to w and the segment $\{(\rho, v) \in \Omega_c : v = 0\}$;
- the point $\mathfrak{v}^\pm(\bar{u})$ is the intersection of the Lax curve of the second characteristic family passing through \bar{u} and $\{u \in \Omega_c : \mathfrak{w}(u) = w^\pm\}$;
- for any $u_\ell, u_r \in \Omega_c$ the point $\mathfrak{u}_*(u_\ell, u_r)$ is the intersection between the Lax curve of the first characteristic family passing through u_ℓ and the Lax curve of the second characteristic family passing through u_r ;
- $\Lambda(u_\ell, u_r)$ is the speed of a discontinuity (u_ℓ, u_r) , that in the (ρ, f) -coordinates coincides with the slope of the segment connecting u_ℓ and u_r .

Observe that by definition $\mathfrak{v}^\pm(\bar{u}) = \mathfrak{u}_*((p^{-1}(w^\pm), 0), \bar{u})$ and $\omega(\bar{u}) = \mathfrak{u}_*(\bar{u}, (0, V))$.

3.2 Riemann solvers

For completeness, we recall the definitions of the Riemann solvers \mathcal{R} and \mathcal{R}_F introduced in [10] and [18], associated to Riemann problem (3a) and to constrained Riemann problem (3a), (3c) with $\mathbf{F} \equiv F$ constant belonging to $[0, f^+]$, respectively, and used in Section 3.3 to define the approximate Riemann solvers \mathcal{R}_n and $\mathcal{R}_{F,n}$.

We recall that Riemann problems for (3a) are Cauchy problems with initial condition of the form

$$u(0, x) = \begin{cases} u_\ell & \text{if } x < 0, \\ u_r & \text{if } x \geq 0. \end{cases} \quad (15)$$

Definition 3.1. *The Riemann solver $\mathcal{R}: \Omega^2 \rightarrow \mathbf{L}^\infty(\mathbb{R}; \Omega)$ associated to Riemann problem (3a), (15) is defined as follows.*

- (R.1) *If $u_\ell, u_r \in \Omega_f$, then $\mathcal{R}[u_\ell, u_r]$ consists of a contact discontinuity (u_ℓ, u_r) with speed of propagation V .*
- (R.2) *If $u_\ell, u_r \in \Omega_c$, then $\mathcal{R}[u_\ell, u_r]$ consists of a 1-wave $(u_\ell, \mathfrak{u}_*(u_\ell, u_r))$ and of a 2-contact discontinuity $(\mathfrak{u}_*(u_\ell, u_r), u_r)$.*
- (R.3) *If $u_\ell \in \Omega_c^-$ and $u_r \in \Omega_f^-$, then $\mathcal{R}[u_\ell, u_r]$ consists of a 1-rarefaction $(u_\ell, \omega(u_\ell))$ and a contact discontinuity $(\omega(u_\ell), u_r)$.*
- (R.4) *If $u_\ell \in \Omega_f^-$ and $u_r \in \Omega_c^-$, then $\mathcal{R}[u_\ell, u_r]$ consists of a phase transition $(u_\ell, \mathfrak{v}^-(u_r))$ and a 2-contact discontinuity $(\mathfrak{v}^-(u_r), u_r)$.*

Since $(t, x) \mapsto \mathcal{R}[u_\ell, u_r](x/t)$ does not in general satisfy constraint condition (3c) with $\mathbf{F} \equiv F$ constant belonging to $[0, f^+]$, we introduce

$$\begin{aligned}
\mathcal{D}_F &\doteq \{ (u_\ell, u_r) \in \Omega \times \Omega : f(\mathcal{R}[u_\ell, u_r](t, 0_\pm)) \leq F \} \\
&= \{ (u_\ell, u_r) \in \Omega_f \times \Omega_f : f(u_\ell) \leq F \} \\
&\quad \cup \{ (u_\ell, u_r) \in \Omega_c \times \Omega : f(\mathfrak{u}_*(u_\ell, u_r)) \leq F \} \\
&\quad \cup \{ (u_\ell, u_r) \in \Omega_f^- \times \Omega_c^- : \min\{f(u_\ell), f(\mathfrak{v}^-(u_r))\} \leq F \}, \\
\mathcal{D}_F^c &\doteq \Omega^2 \setminus \mathcal{D}_F,
\end{aligned}$$

and the constrained Riemann solver \mathcal{R}_F in the following definition.

Definition 3.2. The constrained Riemann solver $\mathcal{R}_F: \Omega^2 \rightarrow \mathbf{L}^\infty(\mathbb{R}; \Omega)$ associated to constrained Riemann problem (3a), (3c), (15) with $\mathbf{F} \equiv F$ constant belonging to $[0, f^+]$ is defined as

$$\mathcal{R}_F[u_\ell, u_r](x) \doteq \begin{cases} \mathcal{R}[u_\ell, u_r](x) & \text{if } (u_\ell, u_r) \in \mathcal{D}_F, \\ \begin{cases} \mathcal{R}[u_\ell, \hat{u}_\ell](x) & \text{if } x < 0, \\ \mathcal{R}[\check{u}_r, u_r](x) & \text{if } x > 0, \end{cases} & \text{if } (u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}, \end{cases}$$

where $\hat{u}_\ell \doteq \hat{u}(w(u_\ell), F) \in \Omega_c$ and $\check{u}_r \doteq \check{u}(v_r, F) \in \Omega$ are defined by (12).




Figure 5: The selection criterion (12) for $\hat{u}_\ell \doteq \hat{u}(w(u_\ell), F)$ and $\check{u}_r \doteq \check{u}(v_r, F)$ exploited in Definition 3.2 in the case $(u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}$ and $F \in (0, f^-)$. In the first picture u_ℓ^1, u_ℓ^2 represent the left state u_ℓ^1, u_ℓ^2 are the corresponding \hat{u}_ℓ . Second and third pictures have analogous meaning, with u_r^1, u_r^2 and $\check{u}_r^1, \check{u}_r^2$.

In Figure 5 we clarify the selection criterion (12) for \hat{u}_ℓ and \check{u}_r in the case $(u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}$ and $F \in (0, f^-)$. We point out that \hat{u}_ℓ and \check{u}_r satisfy the following general properties.

- If $(u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}$, then $w(u_\ell) > w(\check{u}_r)$ and $v_r > \hat{v}_\ell$.
- If $(u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}$ and $u_\ell \in \Omega_f^-$, then $w(\hat{u}_\ell) = w^-$.
- If $(u_\ell, u_r) \in \mathcal{D}_F^{\mathbf{G}}$ and $u_r \in \Omega_f$, then $\check{v}_r = V$.

We recall that both \mathcal{R} and \mathcal{R}_F are $\mathbf{L}_{\text{loc}}^1$ -continuous, see [18, Propositions 2 and 3].

3.3 The approximate Riemann solvers

For simplicity here and in the following we assume that $n \in \mathbb{N}$ is sufficiently large. For any $F \in [0, f^+]$ we introduce below a grid $\mathcal{G}_{F,n}$ in Ω and approximate Riemann solvers $\mathcal{R}_n, \mathcal{R}_{F,n}: \mathcal{G}_{F,n} \times \mathcal{G}_{F,n} \rightarrow \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F,n})$.

The grid

We introduce in Ω a grid $\mathcal{G}_{F,n} \doteq \Omega \cap \mathcal{P}_{F,n}$, see Figure 6, with $\mathcal{P}_{F,n}$ given in the (v, w) -coordinates by

$$\left(\bigcup_{i=0}^{M \cdot 2^n} \{v^i\} \right) \times \left(\bigcup_{i=0}^{N \cdot 2^n} \{w^i\} \right),$$

where M, N, v^i and w^i , are defined as follows:

- If $F = 0$, then we let $M = 1, N = 2$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} & \text{if } i \in \{0, \dots, 2^n\}, \\ w^- + (i - 2^n) 2^{-n} (w^+ - w^-) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \end{cases}$$

and

$$v^i \doteq i 2^{-n} V \quad \text{if } i \in \{0, \dots, 2^n\}.$$

- If $F \in (0, f^-)$, then we let $M = 3, N = 3$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} (w_F - w^- + 1) & \text{if } i \in \{0, \dots, 2^n\}, \\ w_F + (i - 2^n) 2^{-n} (w^- - w_F) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ w^- + (i - 2 \cdot 2^n) 2^{-n} (w^+ - w^-) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}, \end{cases}$$




Figure 6: The grid $\mathcal{G}_{F,n}$ corresponding to $F \in (0, f^-)$ and $n = 2$. The curve in the figure on the left is the support of Ξ_F , which corresponds to (a portion of) the horizontal line in the figure on the right.

and

$$v^i \doteq \begin{cases} i 2^{-n} v_F^- & \text{if } i \in \{0, \dots, 2^n\}, \\ \Xi_F^{-1}(w^{4 \cdot 2^n - i}) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ v_F^+ + (i - 2 \cdot 2^n) 2^{-n} (V - v_F^+) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}. \end{cases}$$

- If $F \in [f^-, f^+]$, then we let $M = 2$, $N = 3$,

$$w^i \doteq \begin{cases} w^- - 1 + i 2^{-n} & \text{if } i \in \{0, \dots, 2^n\}, \\ w^- + (i - 2^n) 2^{-n} (w_F - w^-) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}, \\ w_F + (i - 2 \cdot 2^n) 2^{-n} (w^+ - w_F) & \text{if } i \in \{2 \cdot 2^n + 1, \dots, 3 \cdot 2^n\}, \end{cases}$$

and

$$v^i \doteq \begin{cases} i 2^{-n} v_F^- & \text{if } i \in \{0, \dots, 2^n\}, \\ \Xi_F^{-1}(w^{4 \cdot 2^n - i}) & \text{if } i \in \{2^n + 1, \dots, 2 \cdot 2^n\}. \end{cases}$$

Notice that if $F \in \{f^-, f^+\}$, then we do not necessarily have $w^i \neq w^{i+1}$.

The approximate Riemann solvers

An approximate solution $u_n \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F,n})$ to (3) is constructed by applying the Riemann solvers $\mathcal{R}_n, \mathcal{R}_{F,n} : \mathcal{G}_{F,n} \times \mathcal{G}_{F,n} \rightarrow \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F,n})$, in which rarefactions are replaced by piecewise constant rarefaction fans. More precisely, for any $(u_\ell, u_r) \in \mathcal{G}_{F,n} \times \mathcal{G}_{F,n}$ such that $w_\ell = w_r$ and $v_\ell = v^h < v_r = v^{h+k}$, we let

$$\mathcal{R}_n[u_\ell, u_r](\xi) \doteq \begin{cases} u_\ell & \text{if } \xi \leq \Lambda(u_\ell, u_1), \\ u_j & \text{if } \Lambda(u_{j-1}, u_j) < \xi \leq \Lambda(u_j, u_{j+1}), \quad 1 \leq j \leq k-1, \\ u_r & \text{if } \xi > \Lambda(u_{k-1}, u_r), \end{cases}$$

where $u_0 \doteq u_\ell$, $u_k \doteq u_r$ and $u_j \in \mathcal{G}_{F,n}$ is such that $v_j \doteq v^{h+j}$ and $w_j = w_\ell$. The Riemann solver $\mathcal{R}_{F,n}$ is defined as follows:

1. If $f(\mathcal{R}_n[u_\ell, u_r](0_\pm)) \leq F$, then $\mathcal{R}_{F,n}[u_\ell, u_r] \doteq \mathcal{R}_n[u_\ell, u_r]$.
2. If $f(\mathcal{R}_n[u_\ell, u_r](0_\pm)) > F$, then

$$\mathcal{R}_{F,n}[u_\ell, u_r](\xi) \doteq \begin{cases} \mathcal{R}_n[u_\ell, \hat{u}_\ell](\xi) & \text{if } \xi < 0, \\ \mathcal{R}_n[\tilde{u}_r, u_r](\xi) & \text{if } \xi \geq 0. \end{cases}$$

3.4 The approximate solution

In this section we apply a wave-front tracking algorithm to construct an approximate solution u_n in the space \mathbf{PC} of piecewise constant functions taking finitely many values.

By assumption $F \in \mathbf{PC}((0, \infty); [0, f^+])$. Therefore there exist $F_i \in [0, f^+]$ and $t_i \geq 0$, $i \in \{0, 1, \dots, N\}$, such that

$$F_i \neq F_{i+1}, \quad t_i < t_{i+1}, \quad F(t) = F_i \quad \forall t \in (t_i, t_{i+1}),$$

with $t_0 = 0$ and $t_{N+1} = \infty$.

We recall that in the assumptions of theorem 2.8 we allow $F \equiv F(0) \in [0, f_+]$ when $N = 0$, while for $N \geq 1$ we assume $F_i \in [f_-, f_+]$ for $0 \leq i \leq N$. As stated in Corollary 2.9, one can also consider the intermediate situation in which $N \geq 1$, $F_0 \in [0, f_-]$ and $F_i \in [f_-, f_+]$ for $1 \leq i \leq N$.

An approximate solution $u_n \in \mathbf{PC}(\mathbb{R}_+ \times \mathbb{R}; \cup_{i=0}^N \mathcal{G}_{F_i, n})$ to (3) can be constructed as follows, using at every time step the projection on the corresponding grid. As a first step we consider the grid $\mathcal{G}_{F_0, n}$ and approximate the initial datum \bar{u} with $\bar{u}_n^0 \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_0, n})$ such that

$$\begin{aligned} \|\bar{v}_n^0\|_{\mathbf{L}^\infty} &\leq \|\bar{v}^0\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{v}_n^0) \leq \text{TV}(\bar{v}^0), \quad \|\bar{v}_n^0 - \bar{v}^0\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \\ \|\bar{w}_n^0\|_{\mathbf{L}^\infty} &\leq \|\bar{w}^0\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{w}_n^0) \leq \text{TV}(\bar{w}^0), \quad \|\bar{w}_n^0 - \bar{w}^0\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \end{aligned} \tag{16}$$

where for every compact subset K of \mathbb{R} , $C(K)$ is a constant independent of n . Here

$$\bar{w}_n^0 \doteq \mathfrak{w}(\bar{u}_n^0).$$

The approximate solution u_n^0 is then obtained by gluing together the approximate solutions computed by applying $\mathcal{R}_{F_0, n}$ at $x = 0$ at time $t = 0$ and at any time a wave-front reaches $x = 0$, and by applying \mathcal{R}_n at any discontinuity of \bar{u}_n^0 away from $x = 0$ or at any interaction between wave-fronts taking place away from $x = 0$.

At time $t = t_1$ we restart the above construction by updating the constraint to F_1 and by using $u_n^0(t_1, \cdot)$ as initial datum. More precisely, we consider the grid $\mathcal{G}_{F_1, n}$ and approximate $u_n^0(t_1, \cdot) \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_0, n})$ by $\bar{u}_n^1 \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_1, n})$ such that

$$\begin{aligned} \|\bar{v}_n^1\|_{\mathbf{L}^\infty} &\leq \|v_n^0(t_1, \cdot)\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{v}_n^1) \leq \text{TV}(v_n^0(t_1, \cdot)), \quad \|\bar{v}_n^1 - v_n^0(t_1, \cdot)\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \\ \|\bar{w}_n^1\|_{\mathbf{L}^\infty} &\leq \|w_n^0(t_1, \cdot)\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{w}_n^1) \leq \text{TV}(w_n^0(t_1, \cdot)), \quad \|\bar{w}_n^1 - w_n^0(t_1, \cdot)\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \end{aligned}$$

where

$$\bar{w}_n^1 \doteq \mathfrak{w}(\bar{u}_n^1).$$

The approximate solution u_n^1 is then obtained by gluing together the approximate solutions computed by applying $\mathcal{R}_{F_1, n}$ at $x = 0$ at time $t = t_1$ and at any time a wave-front reaches $x = 0$, and by applying \mathcal{R}_n at any discontinuity of \bar{u}_n^1 away from $x = 0$ or at any interaction between wave-fronts taking place away from $x = 0$.

More in general, at time $t = t_i$ we update the constraint to F_i , consider the grid $\mathcal{G}_{F_i, n}$, approximate $u_n^{i-1}(t_i, \cdot) \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_{i-1}, n})$ with $\bar{u}_n^i \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_i, n})$ such that

$$\begin{aligned} \|\bar{v}_n^i\|_{\mathbf{L}^\infty} &\leq \|v_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{v}_n^i) \leq \text{TV}(v_n^{i-1}(t_i, \cdot)), \quad \|\bar{v}_n^i - v_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \\ \|\bar{w}_n^i\|_{\mathbf{L}^\infty} &\leq \|w_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}^\infty}, \quad \text{TV}(\bar{w}_n^i) \leq \text{TV}(w_n^{i-1}(t_i, \cdot)), \quad \|\bar{w}_n^i - w_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}^1(K)} \leq \frac{C(K)}{2^n}, \end{aligned} \tag{17}$$

where

$$\bar{w}_n^i \doteq \mathfrak{w}(\bar{u}_n^i).$$

The approximate solution u_n^i is then obtained by gluing together the approximate solutions computed by applying $\mathcal{R}_{F_i, n}$ at $x = 0$ at time $t = t_i$ and at any time a wave-front reaches $x = 0$, and by applying \mathcal{R}_n at any discontinuity of \bar{u}_n^i away from $x = 0$ or at any interaction between wave-fronts taking place away from $x = 0$.

By iterating the above procedure we obtain the approximate solution

$$u_n(t, x) = \sum_{i=0}^N u_n^i(t - t_i, x) \cdot \mathbb{1}_{(t_i, t_{i+1}]}(t). \tag{18}$$

As usual, in order to extend the construction globally in time we have to ensure that only finitely many interactions may occur in finite time. In Section 3.5 we prove that $u_n(t, \cdot)$ is well defined for all $t > 0$ and belongs to $\mathbf{PC}(\mathbb{R}_+ \times \mathbb{R}; \cup_{i=0}^N \mathcal{G}_{F_i, n})$. Finally, in Section 3.6 we prove that u_n converges (up to a subsequence) in $\mathbf{L}_{\text{loc}}^1$ to a limit u , which results to be an admissible solution to (3) in the sense of Definition 2.1.

3.5 A priori estimates

In this section we prove the main a priori estimates on the sequence of approximate solutions $\{u_n\}_n$ defined in (18). In Proposition 3.4 we state that $u_n(t, \cdot)$ takes values in $\mathcal{G}_{F_i, n}$ for any $t \in (t_i, t_{i+1}]$ and estimate $\text{TV}(u_n(t, \cdot))$ uniformly in n and t . This together with Proposition 3.6 guarantee that the number of interactions and the number of the discontinuities of u_n are both bounded globally in time.

We choose to study the total variation in the (v, w) -coordinates rather than in the (ρ, v) -coordinates. This choice is convenient to describe the grid, the approximate Riemann solvers and ease the forthcoming analysis, because the total variation of u_n in these coordinates does not increase after any interaction away from $x = 0$. Furthermore, the entropy pairs defined in (5) in the (v, w) -coordinates are well defined, but in the (ρ, v) -coordinates are multi-valued at the vacuum.

Observe that any Contact Discontinuity (CD) has non-negative speed (of propagation), any Shock (S) or Rarefaction Shock (RS) has negative speed, all the Non-classical Shocks (NSs) are stationary and the speed of all the possible Phase Transitions (PTs) ranges in the interval $(-f^-(p^{-1}(w^-) - \rho^-), V)$. Below we say that (u_ℓ, u_r) is a null wave if $u_\ell = u_r$. Notice that if (u_ℓ, u_r) is a PT then $u_\ell \in \Omega_f^-$ and $u_r \in \Omega_c^-$, moreover if (u_ℓ, u_r) is a PT with $w_r > w^-$ then $\rho_\ell = 0$.

Let u_n be an approximate solution of the form (18). For any $t > 0$, let $\sharp_n^i(t)$ and $\sharp_n(t)$ be the number of waves/discontinuities of $u_n^i(t, \cdot)$ and $u_n(t, \cdot)$, respectively. By definition (18) we have

$$\sharp_n(t) = \sum_{i=0}^N \sharp_n^i(t - t_i) \cdot \mathbb{1}_{(t_i, t_{i+1}]}(t). \quad (19)$$

Lemma 3.3. *We have that $\sharp_n : (0, \infty) \rightarrow \mathbb{R}_+$ is uniformly bounded.*

Proof. We have by construction that $\bar{u}_n^i \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_i, n})$ and by assumption that $F_i \in [0, f^+]$. We can therefore apply [9, Proposition 4.1] and obtain that $\sharp_n^i : (0, \infty) \rightarrow \mathbb{R}_+$, $i \in \{1, \dots, N\}$, are uniformly bounded. Hence by (19) the proof is complete. \square

We introduce $\mathcal{T}_n^i, \mathcal{T}_n : (0, \infty) \rightarrow \mathbb{R}_+$ and $\mathcal{T}^0 \geq 0$ defined as

$$\begin{aligned} \mathcal{T}_n^i(t) &\doteq \text{TV}(v_n^i(t, \cdot)) + \text{TV}(w_n^i(t, \cdot)) + 2\hat{\Upsilon}_n^i(t) + 2\check{\Upsilon}_n^i(t), \\ \mathcal{T}_n(t) &\doteq \sum_{i=0}^N \mathcal{T}_n^i(t - t_i) \cdot \mathbb{1}_{(t_i, t_{i+1}]}(t), \\ \mathcal{T}^0 &\doteq \text{TV}(\bar{v}) + \text{TV}(\bar{w}) + 2\hat{\Upsilon}^0 + 2\check{\Upsilon}^0, \end{aligned} \quad (20)$$

where for any $i \in \{0, \dots, N\}$ we define

$$\begin{aligned} \bar{w}_n^i &\doteq \bar{w}(u_n^i), & \hat{\Upsilon}_n^i(t) &\doteq \hat{\Upsilon}(u_n^i(t, \cdot), F_i), & \check{\Upsilon}_n^i(t) &\doteq \check{\Upsilon}(u_n^i(t, \cdot), F_i), \\ \bar{w} &\doteq \bar{w}(\bar{u}), & \hat{\Upsilon}^0(t) &\doteq \hat{\Upsilon}(\bar{u}, F_0), & \check{\Upsilon}^0(t) &\doteq \check{\Upsilon}(\bar{u}, F_0). \end{aligned}$$

Recall that $\hat{\Upsilon}$ and $\check{\Upsilon}$ are defined in (13). Conventionally, we assume that u_n^i , $i \in \{1, \dots, N\}$, are left continuous in time, i.e., $u_n^i(t, \cdot) \equiv u_n^i(t_-, \cdot)$. Then by definition (18) we have that also u_n is left continuous in time. Hence by definition (20) also \mathcal{T}_n is left continuous in time.

The next lemma gives uniform bounds on the total variation of the approximate solution u^n . For convenience, denote by C the uniform Lipschitz constant which existence is claimed in Lemma 2.6. We also need to guarantee that only finite number of fronts is generated by the algorithm in finite time. In order to count emerging fronts, let $\varepsilon_n^i > 0$ be the minimal (v, w) -distance between two points in the grid $\mathcal{G}_{F_i, n}$, namely

$$\varepsilon_n^i \doteq \min_{\substack{u^1, u^2 \in \mathcal{G}_{F_i, n} \\ u^1 \neq u^2}} \left\{ \max \left\{ |v^1 - v^2|, |\bar{w}(u^1) - \bar{w}(u^2)| \right\} \right\},$$

and define $\varepsilon_n \doteq \min \{ \varepsilon_n^i : i \in \{0, \dots, N\} \}$.

Proposition 3.4. *Fix $n \in \mathbb{N}$ sufficiently large. We have that:*

- (a) *for all $i \in \{0, \dots, N\}$, the map $(t_i, t_{i+1}) \ni t \mapsto \mathcal{T}_n(t) \in \mathbb{R}_+$ is non-increasing and decreases by at least ε_n any time the number of waves increases;*
- (b) *$(0, \infty) \ni t \mapsto \mathcal{T}_n(t) \in \mathbb{R}_+$ is uniformly bounded by $(1 + 2C)^N \cdot \mathcal{T}^0$ with respect to $t > 0$ and $n \in \mathbb{N}$;*

(c) for all $i \in \{0, \dots, N\}$, $u_n(t, \cdot) \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_i, n})$ for all $t \in (t_i, t_{i+1}]$.

Proof. By [9, Proposition 4.1] we have that

- (i) the map $(0, \infty) \ni t \mapsto \mathcal{T}_n^i(t) \in \mathbb{R}_+$ is non-increasing and decreases by at least ε_n^i any time the number of waves increases;
- (ii) $u_n^i(t, \cdot) \in \mathbf{PC}(\mathbb{R}; \mathcal{G}_{F_i, n})$ for all $t > 0$.

Therefore by (18) and (20) properties (a) and (c) hold true. Property (b) follows readily from the definition of \mathcal{T} , the requirement that $F_i \in [f_-, f_+]$ for $i \geq 1$, and Lemma 2.6. \square

Remark 3.5. *At this point, let us make apparent the difficulty in extension of our result to the case where F_i need not be restricted to values above the threshold f^- . Consider, e.g., the first re-meshing time t_1 . Under the assumption $F_1 \in [f^-, f^+]$, we control $\hat{\Upsilon}(\bar{u}_n^1, F_1)$, $\check{\Upsilon}(\bar{u}_n^1, F_1)$ by the variation of \bar{u}_n^1 via the uniform Lipschitz constant of the maps \hat{u}, \check{u} (see Lemma 2.6). When $F_1 \in [0, f^-)$, the discontinuity of the maps \hat{u}, \check{u} makes this control impossible. There remains the eventuality of controlling $\hat{\Upsilon}(\bar{u}_n^1, F_1)$, $\check{\Upsilon}(\bar{u}_n^1, F_1)$ by the values $\hat{\Upsilon}(u_n^0(t_1, \cdot), F_0)$, $\check{\Upsilon}(u_n^0(t_1, \cdot), F_0)$, being understood that \bar{u}_n^1 is a projection of $u_n^0(t_1, \cdot)$ due to re-meshing. At this point, it is the change of the constraint level from F_0 to F_1 that creates a major difficulty: we are unable to control, e.g., $\hat{\Upsilon}(u_n^0(t_1, \cdot), F_1)$ by $\hat{\Upsilon}(u_n^0(t_1, \cdot), F_0)$ without artificial restrictions. Note that the technique that was developed for handling the analogous difficulty in the case of the ARZ system (see [4]) does not extend to our case, due to the more complex definition of the interaction potentials $\hat{\Upsilon}, \check{\Upsilon}$ and to the fact that $\mathbb{W}(u)$ may fail to satisfy the conservation equation at $x = 0$.*

Beside the bound on the number of wave-fronts proved in Proposition 3.4, we need to bound also the number of interactions. This is the aim of the next proposition, which together with Proposition 3.4 ensure the global existence of u_n .

Proposition 3.6. *For any fixed $n \in \mathbb{N}$ sufficiently large, we have that the number of interactions occurring in the interval of time $(0, \infty)$ is bounded. In particular u_n is globally defined.*

Proof. By [9, Proposition 3.2] we have that the number of interactions involved in the construction of each u_n^i is bounded. Therefore the statement of the proposition follows directly from the definition (18) of u_n . \square

3.6 Convergence

The convergence is proved by following the traditional method of proving compactness via Helly's theorem.

We observe that

$$|\rho_\ell - \rho_r| \leq L_\rho (|v_\ell - v_r| + |\mathfrak{w}_\ell - \mathfrak{w}_r|),$$

where $L_\rho \doteq \max\{\rho^-, \|1/p'\|_{\mathbf{L}^\infty([p^{-1}(w^-), p^{-1}(w^+)]; \mathbb{R})}\}$, because

$$\rho_{\ell, r} = \begin{cases} p^{-1}(\mathfrak{w}_{\ell, r} - v_{\ell, r}) & \text{if } \mathfrak{w}_{\ell, r} \in [w^-, w^+], \\ (\mathfrak{w}_{\ell, r} + 1 - w^-) \rho^- & \text{if } \mathfrak{w}_{\ell, r} \in [w^- - 1, w^-]. \end{cases}$$

As a consequence $\mathbf{TV}(\rho) \leq L_\rho (\mathbf{TV}(v) + \mathbf{TV}(\mathfrak{w}))$, hence

$$\mathbf{TV}(u) \leq (1 + L_\rho) (\mathbf{TV}(v) + \mathbf{TV}(\mathfrak{w})).$$

Moreover, by Proposition 3.4 we have that for any $t > 0$

$$\mathbf{TV}(v_n(t, \cdot)) + \mathbf{TV}(\mathfrak{w}_n(t, \cdot)) \leq \mathcal{T}_n(t) \leq C \mathcal{T}^0,$$

and therefore

$$\mathbf{TV}(u_n(t, \cdot)) \leq K \doteq (1 + L_\rho) C \mathcal{T}^0. \tag{21}$$

Proposition 3.7. *Fix $i \in \{0, 1, \dots, N\}$. For any $s, t \in (t_i, t_{i+1})$ we have*

$$\|u_n(t, \cdot) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L |t - s|, \tag{22}$$

with $L \doteq K \max\{V, R p'(R)\}$ which does not depend on i or n .

Proof. If no interaction occurs for times between s and t , then

$$\begin{aligned} \|u_n(t, \cdot) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} &\leq \sum_{i \in \mathbf{D}(t)} \left| (t-s) \dot{\delta}_n^i(t) \left(\rho_n(t, \delta_n^i(t)_-) - \rho_n(t, \delta_n^i(t)_+) \right) \right| \\ &\quad + \sum_{i \in \mathbf{D}(t)} \left| (t-s) \dot{\delta}_n^i(t) \left(v_n(t, \delta_n^i(t)_-) - v_n(t, \delta_n^i(t)_+) \right) \right| \leq L |t-s|, \end{aligned}$$

where $\delta_n^i(t) \in \mathbb{R}$, $i \in \mathbf{D}(t) \subset \mathbb{N}$, are the positions of the discontinuities of $u_n(t, \cdot)$. The case when one or more interactions take place for times between t and s is similar, because by the finite speed of propagation of the waves the map $(t_i, t_{i+1}) \ni t \mapsto u_n(t, \cdot)$ is $\mathbf{L}_{\text{loc}}^1$ -continuous across interaction times. \square

In general u_n is not Lipschitz continuous in time with respect to the \mathbf{L}^1 -norm in space, namely, (22) does not hold true for all $s, t > 0$. Indeed, the approximation at time t_i of $u_n^{i-1}(t_i, \cdot)$ with \bar{u}_n^i satisfies conditions listed in (17), which do not guarantee the continuity across t_i . This prevents an application of Helly's theorem directly to u_n . However, if we set $\Delta_n^i(x) \doteq u_n(t_i^+, x) - u_n(t_i^-, x)$, we observe that the functions u_n^{Lip} defined as

$$u_n^{\text{Lip}}(t, x) \doteq u_n(t, x) - \sum_{i=1}^N \Delta_n^i(x) \cdot \mathbb{1}_{[t_i, \infty)}(t),$$

are Lipschitz continuous in time. The most significant situation is with $t_{i+1} > t > t_i > s > t_{i-1}$ for some $i \in \{1, \dots, N\}$, and in this case

$$\begin{aligned} \|u_n^{\text{Lip}}(t, \cdot) - u_n^{\text{Lip}}(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} &= \|u_n(t, \cdot) - \Delta_n^i(\cdot) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \\ &\leq \|u_n(t, \cdot) - u_n(t_i^+, x)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} + \|u_n(t_i^-, x) - u_n(s, \cdot)\|_{\mathbf{L}^1(\mathbb{R}; \Omega)} \leq L |t-s|. \end{aligned}$$

All other cases are similar.

From the construction of the solutions u_n detailed in Section 3.4 and in particular from (17) (note that the constant $C(K)$ in (17) depends neither on n nor on F_i) we have that $u_n^{\text{Lip}} - u_n$ converges to the null function in $\mathbf{L}_{\text{loc}}^1$ as n goes to infinity. At the same time the sequence $\{u_n^{\text{Lip}}\}_n$ satisfies all the requirements of Helly's Theorem, so that it converges (up to a subsequence) in $\mathbf{L}_{\text{loc}}^1(\mathbb{R}_+ \times \mathbb{R}; \Omega)$ to a function $u \in \mathbf{L}^\infty(\mathbb{R}_+; \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{C}^0(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ and the limit satisfies the estimates listed in (14). Although the a.e. convergence is enough for the sake of the proof of Theorem 2.8, its extension sketched in Section 5 requires convergence of $\{u^n\}_n$ in the topology of $\mathbf{L}^\infty(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$. Observe that the above arguments do guarantee this convergence.

3.7 Characterization of the limit

We start by focusing on the renormalization property and the way it is used to handle entropy inequalities: these are the key arguments of the characterization of admissible discontinuities at the constraint. First, based on the fact that wave-front tracking solutions are piecewise constant weak solutions of the problem except at times $t = t_i$ and eventually at the constraint location $x = 0$, we readily assess the local renormalization property for approximate solutions.

Proposition 3.8. *The approximate solution u^n satisfies the renormalization property (7) with test functions supported in $\{(t, x) \mid t_i < t < t_{i+1}, x \neq 0\}$.*

Proof. Let $\phi \in \mathbf{C}_c^\infty((t_i, t_{i+1}); \mathbb{R}_*)$ be a test function. Due to the discrete nature of u^n , without loss of generality we can assume that its support intersects only one discontinuity curve $x = \delta(t)$ of u^n . We denote $u_+^n = u^n(t, \delta(t)_+)$ and so on, and for simplicity of notation we drop the subscript n . The Rankine-Hugoniot conditions (9), (10) along $x = \delta(t)$ give

$$\begin{aligned} &\begin{cases} \rho_+ \dot{\delta}(t) - f(u_+) = \rho_- \dot{\delta}(t) - f(u_-), \\ [\rho_+ \dot{\delta}(t) - f(u_+)] \mathbf{w}(u_+) = [\rho_- \dot{\delta}(t) - f(u_-)] \mathbf{w}(u_-), \end{cases} \\ \iff &\begin{cases} \rho_+ \dot{\delta}(t) = f(u_+), \\ \rho_- \dot{\delta}(t) = f(u_-), \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{w}(u_+) = \mathbf{w}(u_-), \\ \rho_+ \dot{\delta}(t) - f(u_+) = \rho_- \dot{\delta}(t) - f(u_-). \end{cases} \end{aligned}$$

In both the cases, for any continuous function $g: [w_-, w_+] \rightarrow \mathbb{R}$ we have

$$[\rho_+ \dot{\delta}(t) - f(u_+)] g(\mathbf{w}(u_+)) = [\rho_- \dot{\delta}(t) - f(u_-)] g(\mathbf{w}(u_-)).$$

As a consequence we have that u satisfies the renormalization property in $\mathbb{R}_-^2 \cup \mathbb{R}_+^2$. \square

Now, we can adapt to the present framework [4, Proposition 3.1].

Proposition 3.9. *For any test function $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\text{supp}(\phi) \subset (t_i, t_{i+1}) \times \mathbb{R}$, for an $i \in \{1, \dots, N\}$, we have that*

$$\int_0^\infty \mathbf{N}_{\mathbf{F}(t)}^k(u_n(t, 0_-)) \phi(t, 0) dt = \int_0^\infty \int_{-\infty}^0 \left(\rho_n \mathbf{N}_{\mathbf{F}(t)}^k(u_n) \psi_t \xi + f(u_n) \mathbf{N}_{\mathbf{F}(t)}^k(u_n) \psi \xi_x \right) dx dt$$

where $\mathbf{N}_{\mathbf{F}}^k(u)$ is defined by (4), $\psi(t) = \phi(t, 0)$ and ξ is an arbitrary $\mathbf{C}_c^\infty(\mathbb{R}; \mathbb{R})$ test function such that $\xi(0) = 1$.

Proof. By Proposition 3.8 with $g(W) \doteq \mathbf{n}_{\mathbf{F}}^k(W)$, see (4), we have that u_n satisfies the equation

$$\left(\rho_n \mathbf{n}_{\mathbf{F}}^k(\mathbf{w}(u_n)) \right)_t + \left(f(u_n) \mathbf{n}_{\mathbf{F}}^k(\mathbf{w}(u_n)) \right)_x = 0,$$

in the sense of distributions in $(t_i, t_{i+1}) \times (-\infty, 0)$. As a consequence by the Gauss-Green formula we have

$$\begin{aligned} \int_0^\infty \int_{-\infty}^0 \left(\rho_n \mathbf{n}_{\mathbf{F}}^k(\mathbf{w}(u_n)) \psi_t \xi + f(u_n) \mathbf{n}_{\mathbf{F}}^k(\mathbf{w}(u_n)) \psi \xi_x \right) dx dt &= \int_0^\infty f(u_n(t, 0_-)) \mathbf{n}_{\mathbf{F}}^k(\mathbf{w}(u_n(t, 0_-))) \psi dx dt \\ &= \int_0^\infty \mathbf{N}_{\mathbf{F}}^k(\mathbf{w}(u_n(t, 0_-))) \psi dx dt. \end{aligned}$$

□

Proposition 3.10. *Let $\bar{u} \in \mathbf{L}^1 \cap \mathbf{BV}(\mathbb{R}; \Omega)$ and $\mathbf{F} \in \mathbf{PC}((0, \infty); [0, f^+])$ satisfy **(H.1)** or **(H.2)**. If u is a limit of the sequence of approximate solutions $\{u_n\}_n$ constructed in Section 3.4, then u is a solution to constrained Cauchy problem (3) in the sense of Definition 2.1.*

Proof. We consider separately the conditions listed in Definition 2.1.

(S.1) Initial condition (3b) holds by (16), (22) and the $\mathbf{L}_{\text{loc}}^1$ -convergence of u_n to u .

(S.2) We prove now (6), that is for any test function $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ we have

$$\int_0^\infty \int_{\mathbb{R}} \left(\rho \phi_t + f(u) \phi_x \right) dx dt = 0.$$

Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. Since u_n is uniformly bounded and f is uniformly continuous on bounded sets, it is sufficient to prove that

$$\int_0^T \int_{\mathbb{R}} \left(\rho_n \phi_t + f(u_n) \phi_x \right) dx dt \rightarrow 0. \quad (23)$$

By the Gauss-Green formula the double integral above can be written as

$$\int_0^T \sum_{j \in \mathbf{D}(t)} \left(\delta_n^j(t) \Delta \rho_n^j(t) - \Delta f_n^j(t) \right) \phi(t, \delta_n^j(t)) dt + \sum_{i=1}^N \int_{\mathbb{R}} \left(\rho_n(t_i^-, x) - \rho_n(t_i^+, x) \right) \phi(t_i, x) dx,$$

where

$$\Delta \rho_n^j(t) \doteq \rho_n(t, \delta_n^j(t)_+) - \rho_n(t, \delta_n^j(t)_-), \quad \Delta f_n^j(t) \doteq f(u_n(t, \delta_n^j(t)_+)) - f(u_n(t, \delta_n^j(t)_-)).$$

By construction any discontinuity of $u_n(t, \cdot)$ satisfies the first Rankine-Hugoniot condition (9), therefore

$$\delta_n^j(t) \Delta \rho_n^j(t) - \Delta f_n^j(t) = 0, \quad j \in \mathbf{D}(t).$$

Moreover we have

$$\int_{\mathbb{R}} \sum_{i=1}^N \left(\rho_n(t_i^-, x) - \rho_n(t_i^+, x) \right) \phi(t_i, x) dx \leq \|\phi\|_{\mathbf{L}^\infty} L_\rho \sum_{i=1}^N \left(\|\bar{v}_n^i - v_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}_{\text{loc}}^1} + \|\bar{w}_n^i - w_n^{i-1}(t_i, \cdot)\|_{\mathbf{L}_{\text{loc}}^1} \right)$$

and (23) is trivial.

(S.3) Property (7) follows by Proposition 3.8, with the contribution of the restart times t_i controlled in the same way as in the above proof of property **(S.2)**.

(S.4) We prove now (8), namely that for any $k \in [0, V]$ and $\phi \in \mathbf{C}_c^\infty((0, \infty) \times \mathbb{R}; \mathbb{R})$ such that $\phi \geq 0$ we have

$$\int_0^\infty \left(\int_{\mathbb{R}} (\mathbf{E}^k(u) \phi_t + \mathbf{Q}^k(u) \phi_x) dx + \mathbf{N}_{\mathbb{F}(t)}^k(u(t, 0_-)) \phi(t, 0) \right) dt \geq 0,$$

where $\mathbf{N}_{\mathbb{F}}^k(u)$, \mathbf{E}^k and \mathbf{Q}^k are defined in (4) and (5). From Proposition 3.9 follows that for all $t \notin \{t_1, \dots, t_N\}$

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_0^\infty \mathbf{N}_{\mathbb{F}(t)}^k(u_n(t, 0_-)) \phi(t, 0) dt &= \int_0^\infty \mathbf{N}_{\mathbb{F}(t)}^k(u(t, 0_-)) \phi(t, 0) dt, \\ \lim_{n \rightarrow \infty} \int_0^\infty \mathbf{N}_{\mathbb{F}(t)}^k(u_n(t, 0_+)) \phi(t, 0) dt &= \int_0^\infty \mathbf{N}_{\mathbb{F}(t)}^k(u(t, 0_+)) \phi(t, 0) dt. \end{aligned}$$

Choose $T > 0$ such that $\phi(t, x) = 0$ whenever $t \geq T$. By the a.e. convergence of u_n to u and the uniform continuity of \mathbf{E}^k and \mathbf{Q}^k , it is sufficient to prove that

$$\liminf_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}} (\mathbf{E}^k(u_n) \phi_t + \mathbf{Q}^k(u_n) \phi_x) dx dt \geq 0. \quad (24)$$

By the Gauss-Green formula the double integral above can be written as

$$\begin{aligned} & \int_0^T \sum_{j \in \mathbf{D}(t)} \left(\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right) \phi(t, \delta_n^j(t)) dt + \sum_{i=1}^N \int_{\mathbb{R}} \left(\mathbf{E}^k(u_n(t_i^-, x)) - \mathbf{E}^k(u_n(t_i^+, x)) \right) \phi(t_i, x) dx \\ & \geq \int_0^T \sum_{j \in \mathbf{D}(t)} \left(\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right) \phi(t, \delta_n^j(t)) dt - \sum_{i=1}^N \int_{\mathbb{R}} \left| \mathbf{E}^k(u_n(t_i^-, x)) - \mathbf{E}^k(u_n(t_i^+, x)) \right| \phi(t_i, x) dx. \end{aligned}$$

Since the last sum above converges to zero as n goes to infinity, to prove (24) it is sufficient to show that

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{j \in \mathbf{D}(t)} \left(\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right) \phi(t, \delta_n^j(t)) dt \geq 0, \quad (25)$$

where

$$\Delta \mathbf{E}_n^{k,j}(t) \doteq \mathbf{E}^k(u_n(t, \delta_n^j(t)_+)) - \mathbf{E}^k(u_n(t, \delta_n^j(t)_-)), \quad \Delta \mathbf{Q}_n^{k,j}(t) \doteq \mathbf{Q}^k(u_n(t, \delta_n^j(t)_+)) - \mathbf{Q}^k(u_n(t, \delta_n^j(t)_-)).$$

To estimate the integral in (25) we have to distinguish the following cases.

- If the j -th discontinuity is a PT, then we let $x \doteq \delta_n^j(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) &< \min\{\rho_n(t, x_+), p^{-1}(w^- - k)\}, & \dot{\delta}_n^j(t) &= \Lambda(u_n(t, x_-), u_n(t, x_+)), \\ v_n(t, x_-) &= V > v_n(t, x_+), & \mathbf{W}(u_n(t, x_-)) &= w^- \leq \mathbf{w}(u_n(t, x_+)) = \mathbf{W}(u_n(t, x_+)), \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,j}(t) &= \begin{cases} \frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 & \text{if } v_n(t, x_+) < k \leq V, \\ 0 & \text{if } k \leq v_n(t, x_+), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,j}(t) &= \begin{cases} k - \frac{f(u_n(t, x_+))}{\rho_{n,+}^k} & \text{if } v_n(t, x_+) < k \leq V, \\ 0 & \text{if } k \leq v_n(t, x_+), \end{cases} \end{aligned}$$

where $\rho_{n,+}^k \doteq p^{-1}(\mathbf{w}(u_n(t, x_+)) - k)$. If $v_n(t, x_+) < k \leq V$, then

$$\begin{aligned} & \dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 \right] + k - \frac{f(u_n(t, x_+))}{\rho_{n,+}^k} \\ &= \underbrace{\left[\frac{\rho_n(t, x_+)}{\rho_{n,+}^k} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) - \Lambda((\rho_{n,+}^k, k), u_n(t, x_+)) \right]}_{>0} > 0. \end{aligned}$$

- If the j -th discontinuity is a CD, then we let $x \doteq \delta_n^j(t)$ and observe that $\dot{\delta}_n^j(t) = v_n(t, x_-) = v_n(t, x_+)$ implies that $\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) = 0$.
- If the j -th discontinuity is a S, then we let $x \doteq \delta_n^j(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) < \rho_n(t, x_+), & \quad f(u_n(t, x_-)) > f(u_n(t, x_+)), & \quad \dot{\delta}_n^j(t) = \Lambda(u_n(t, x_-), u_n(t, x_+)) < 0, \\ v_n(t, x_-) > v_n(t, x_+), & & \quad \mathbf{w}_\pm \doteq \mathbf{w}(u_n(t, x_-)) = \mathbf{w}(u_n(t, x_+)) \geq w^-, \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,j}(t) &= \begin{cases} \frac{\rho_n(t, x_+) - \rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_+) < v_n(t, x_-) < k, \\ \frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 & \text{if } v_n(t, x_+) < k \leq v_n(t, x_-), \\ 0 & \text{if } k \leq v_n(t, x_+) < v_n(t, x_-), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,j}(t) &= \begin{cases} \frac{f(u_n(t, x_-)) - f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_+) < v_n(t, x_-) < k, \\ k - \frac{f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_+) < k \leq v_n(t, x_-), \\ 0 & \text{if } k \leq v_n(t, x_+) < v_n(t, x_-). \end{cases} \end{aligned}$$

If $k > v_n(t, x_-)$ or $k \leq v_n(t, x_+)$, then it is immediate to see that $\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) = 0$. Furthermore, if $v_n(t, x_+) < k \leq v_n(t, x_-)$, then

$$\begin{aligned} & \dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right] + k - \frac{f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} \\ &= \underbrace{\left[\frac{\rho_n(t, x_+)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) - \Lambda\left(p^{-1}(\mathbf{w}_\pm - k), k, u_n(t, x_+)\right) \right]}_{>0} > 0. \end{aligned}$$

- If the j -th discontinuity is a RS, then we let $x \doteq \delta_n^j(t)$ and observe that

$$\begin{aligned} \rho_n(t, x_-) > \rho_n(t, x_+), & \quad f(u_n(t, x_-)) < f(u_n(t, x_+)), & \quad \dot{\delta}_n^j(t) = \Lambda(u_n(t, x_-), u_n(t, x_+)) < 0, \\ v_n(t, x_-) < v_n(t, x_+), & & \quad \mathbf{w}_\pm \doteq \mathbf{w}(u_n(t, x_-)) = \mathbf{w}(u_n(t, x_+)) \geq w^-, \end{aligned}$$

hence

$$\begin{aligned} \Delta \mathbf{E}_n^{k,j}(t) &= \begin{cases} \frac{\rho_n(t, x_+) - \rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_-) < v_n(t, x_+) < k, \\ \frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 & \text{if } v_n(t, x_-) < k \leq v_n(t, x_+), \\ 0 & \text{if } k \leq v_n(t, x_-) < v_n(t, x_+), \end{cases} \\ -\Delta \mathbf{Q}_n^{k,j}(t) &= \begin{cases} \frac{f(u_n(t, x_-)) - f(u_n(t, x_+))}{p^{-1}(\mathbf{w}_\pm - k)} & \text{if } v_n(t, x_-) < v_n(t, x_+) < k, \\ \frac{f(u_n(t, x_-))}{p^{-1}(\mathbf{w}_\pm - k)} - k & \text{if } v_n(t, x_-) < k \leq v_n(t, x_+), \\ 0 & \text{if } k \leq v_n(t, x_-) < v_n(t, x_+). \end{cases} \end{aligned}$$

If $k > v_n(t, x_+)$ or $k \leq v_n(t, x_-)$, then it is immediate to see that $\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) = 0$. Furthermore, if $v_n(t, x_-) < k \leq v_n(t, x_+)$, then

$$\begin{aligned} & \dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \\ &= \Lambda(u_n(t, x_-), u_n(t, x_+)) \left[\frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right] + \frac{f(u_n(t, x_-))}{p^{-1}(\mathbf{w}_\pm - k)} - k \end{aligned}$$

$$\begin{aligned}
&= \underbrace{\left[\frac{\rho_n(t, x_-)}{p^{-1}(\mathbf{w}_\pm - k)} - 1 \right]}_{>0} \underbrace{\left[\Lambda(u_n(t, x_-), u_n(t, x_+)) + \Lambda(u_n(t, x_-), (p^{-1}(\mathbf{w}_\pm - k), k)) \right]}_{<0} \\
&\geq -\frac{2}{\rho^-} p^{-1}(\mathbf{w}_\pm) p'(p^{-1}(\mathbf{w}_\pm)) [\rho_n(t, x_-) - \rho_n(t, x_+)]
\end{aligned}$$

because $\rho_n(t, x_-) > p^{-1}(\mathbf{w}_\pm - k) \geq \rho_n(t, x_+) \geq \rho^-$ and because by the concavity of $\mathfrak{L}_{\mathbf{w}_\pm}(\rho) = (\mathbf{w}_\pm - p(\rho)) \rho$ we have

$$\begin{aligned}
0 &> \Lambda(u_n(t, x_-), u_n(t, x_+)) > \Lambda(u_n(t, x_-), (p^{-1}(\mathbf{w}_\pm - k), k)) \\
&> \mathfrak{L}'_{\mathbf{w}_\pm}(\rho_n(t, x_-)) = \mathbf{w}_\pm - p(\rho_n(t, x_-)) - \rho_n(t, x_-) p'(\rho_n(t, x_-)) \\
&\geq \mathfrak{L}'_{\mathbf{w}_\pm}(p^{-1}(\mathbf{w}_\pm)) = -p^{-1}(\mathbf{w}_\pm) p'(p^{-1}(\mathbf{w}_\pm)).
\end{aligned}$$




Figure 7: Above $F \in (f^-, f^+)$, $v_0^\pm \doteq v_n(t, 0_\pm)$ and $v_{0,F}^\pm \doteq F/p^{-1}(\mathbf{w}(u_n(t, 0_\pm)) - k)$. With the first two pictures we show that if $v_0^- < k < v_0^+$, then $v_{0,F}^- < k$. In the last picture we consider the case $v_0^- < v_0^+ < k$ and show that $v_{0,F}^- < v_{0,F}^+ < k$.




Figure 8: Above $F \in (0, f^-)$, $v_0^\pm \doteq v_n(t, 0_\pm)$ and $v_{0,F}^\pm \doteq F/p^{-1}(\mathbf{w}(u_n(t, 0_\pm)) - k)$. With the first two pictures we show that if $v_0^- < k < v_0^+$, then $v_{0,F}^- < k$. In the last picture we consider the case $v_0^- < v_0^+ < k$ and show that $v_{0,F}^- < v_{0,F}^+ < k$.

- If the j -th discontinuity is a NS occurring at $x = 0$, then

$$\begin{aligned}
\delta_n^j(t) &= 0, & f(u_n(t, 0_\pm)) &= F, & v_F^- &\leq v_n(t, 0_-) < v_n(t, 0_+), \\
\dot{\delta}_n^j(t) &= 0, & \mathbf{w}(u_n(t, 0_-)) &= \mathbf{W}(u_n(t, 0_-)) \geq \mathbf{W}(u_n(t, 0_+)),
\end{aligned}$$

hence

$$-\Delta \mathbf{Q}_n^{k,j}(t) = \begin{cases} \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_-)) - k)} - \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_+)) - k)} & \text{if } v_n(t, 0_-) < v_n(t, 0_+) < k, \\ \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_-)) - k)} - k & \text{if } v_n(t, 0_-) < k \leq v_n(t, 0_+), \\ 0 & \text{if } k \leq v_n(t, 0_-) < v_n(t, 0_+), \end{cases}$$

$$\mathbf{N}_F^k(u_n(t, 0_-)) = \begin{cases} \left[k - \frac{F}{p^{-1}(\mathbf{w}(u_n(t, 0_-)) - k)} \right]_+ & \text{if } F \neq 0, \\ k & \text{if } F = 0. \end{cases}$$

Notice that if $F = 0$, then $u_n(t, 0_+) = (0, V)$ and $u_n(t, 0_-) \in [p^{-1}(w^-), R] \times \{0\}$. We observe, see Figures 7 and 8, that $-\Delta \mathbf{Q}_n^{k,j}(t) < 0$ and that $-\Delta \mathbf{Q}_n^{k,j}(t) + \mathbf{N}_F^k(u_n(t, 0_-)) \geq 0$ and therefore

$$\left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) + \mathbf{N}_F^k(u_n(t, 0_-)) \phi(t, 0) = \left[-\Delta \mathbf{Q}_n^{k,j}(t) + \mathbf{N}_F^k(u_n(t, 0_-)) \right] \phi(t, 0) \geq 0.$$

Thus, as in the cases above, we can conclude that (25) holds true. We underline that beside the NSs, the only possible stationary discontinuities at $x = 0$ are PTs and CDs, however in both of these cases we have $f(u_n(t, 0_-)) = 0$ and therefore $\mathbf{N}_F^k(u_n(t, 0_-)) = 0$.

The above case by case study shows that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}} \left[\mathbf{E}^k(u_n) \phi_t + \mathbf{Q}^k(u_n) \phi_x \right] dx dt \\ &= \liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \\ &\geq -\frac{2}{\rho^-} \max_{\rho \in [p^{-1}(w^-), R]} |\rho p'(\rho)| \liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\rho_n(t, \delta_n^j(t)_-) - \rho_n(t, \delta_n^j(t)_+) \right] \phi(t, \delta_n^j(t)) dt \\ &\geq -\frac{2T}{\rho^-} \|\phi\|_{\mathbf{L}^\infty} K \max_{\rho \in [p^-, R]} |\rho p'(\rho)| \doteq -M, \end{aligned}$$

where $\delta_n^j(t) \in \mathbb{R}$, $i \in \text{RS}_n(t) \subset \mathbb{N}$, are the positions of the RSs of $u_n(t, \cdot)$ and K is defined in (21).

We claim that for any fixed $h > 0$, there exists a dense set \mathcal{K}_h of values of k in $[0, V]$ such that

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \geq -\frac{1}{h}.$$

To prove it we fix $a, b \in [0, V]$ with $a < b$ and show that there exists $k \in (a, b)$ such that the above estimate is satisfied. Let $l \doteq \lceil 2(Mh + 1)/(b - a) \rceil$ and introduce the set

$$\mathcal{K}_h \doteq \frac{2\mathbb{N} + 1}{l} \cap (a, b).$$

Let $\mathcal{E}_n > 0$ be the maximal (v, w) -distance between two ‘‘consecutive’’ points in the grid $\mathcal{G}_{F,n}$ having the same w -coordinate, namely, with a slight abuse of notations, we let

$$\mathcal{E}_n \doteq \max_{\substack{(v^j, w), (v^{j+1}, w) \in \mathcal{G}_{F,n} \\ v^j \neq v^{j+1}}} (v^{j+1} - v^j).$$

Let $n_h \in \mathbb{N}$ be sufficiently large so that $\mathcal{E}_{n_h} < 2/l$. Take $n \geq n_h$. We claim that for any $j \in \text{RS}_n(t)$ we have

$$\mathcal{K}_h \cap \left(v_n(t, \delta_n^j(t)_-), v_n(t, \delta_n^j(t)_+) \right)$$

has at most one element. Indeed, if \mathcal{K}_h has more than one element then for any $i \in \text{RS}_n(t)$ we have

$$v_n(t, \delta_n^j(t)_+) - v_n(t, \delta_n^j(t)_-) \leq \mathcal{E}_n < \frac{2}{l} = \min_{\substack{k^1, k^2 \in \mathcal{K}_h \\ k^1 \neq k^2}} |k^1 - k^2|.$$

As a consequence the sum

$$\sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right]$$

has at most one nonzero element; moreover

$$-m \left(\rho_n(t, \delta_n^j(t)_-) - \rho_n(t, \delta_n^j(t)_+) \right) \leq \sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right],$$

where

$$m \doteq \frac{2}{\rho^-} \max_{\rho \in [\rho^-, R]} |\rho p'(\rho)| = \frac{M}{T C \|\phi\|_{\mathbf{L}^\infty}}.$$

Therefore we find

$$\sum_{i \in \text{RS}_n(t)} \sum_{k \in \mathcal{K}_h} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \geq -m K.$$

By exchanging the sums, multiplying by the test function and integrating in time we get

$$\sum_{k \in \mathcal{K}_h} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \geq -M.$$

Moreover, by construction we have that \mathcal{K}_h is a non-empty set with a finite number of elements (it has at most $h M$ elements), hence

$$h M \max_{k \in \mathcal{K}_h} \left[\int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \right] \geq -M.$$

In conclusion we proved that there exists $k \in \mathcal{K}_h \subseteq (a, b)$ such that the above estimate is satisfied for any $n \geq \mathfrak{n}_h$; therefore, since \mathcal{K}_h has a finite number of elements, we have

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \geq -\frac{1}{h}.$$

Since a and b are arbitrary, the above estimate holds true for a dense set of values of k in $[0, V]$.

Actually, the above estimate holds for any k in $[0, V]$ because the term in brackets in the above formula is continuous with respect to k . Finally, for the arbitrariness of h , we have that

$$\liminf_{n \rightarrow \infty} \int_0^T \sum_{i \in \text{RS}_n(t)} \left[\dot{\delta}_n^j(t) \Delta \mathbf{E}_n^{k,j}(t) - \Delta \mathbf{Q}_n^{k,j}(t) \right] \phi(t, \delta_n^j(t)) dt \geq 0$$

and this concludes the proof of (24).

(S.5) We prove now that (3c) holds for a.e. $t > 0$, namely

$$f(u(t, 0_\pm)) \leq F \quad \text{for a.e. } t > 0.$$

By construction $f(u_n(t, 0_\pm)) \leq F$ for any $t > 0$, namely the approximate solutions satisfy (3c). Since weak convergence preserves pointwise inequalities, it is sufficient to prove that $f(u_n(t, 0_\pm))$ weakly converges to $f(u(t, 0_\pm))$. If ϕ is a smooth test function of time with compact support in $(0, \infty)$ and φ is a smooth test function of space with compact support and such that $\varphi(0) = 1$, then

$$\int_0^\infty f(u_n(t, 0_-)) \phi(t) dt = \int_0^\infty \int_{-\infty}^0 \left[\rho_n(t, x) \dot{\phi}(t) \varphi(x) + f(u_n(t, x)) \phi(t) \dot{\varphi}(x) \right] dx dt.$$

The right-hand side passes to the limit, yielding the analogous expression with u_n replaced by u . By using again the Gauss-Green formula, one finally finds that

$$\lim_{n \rightarrow \infty} \int_0^\infty f(u_n(t, 0_-)) \phi(t) dt = \int_0^\infty f(u(t, 0_-)) \phi(t) dt.$$

As a consequence $f(u_n(t, 0_-))$ weakly converges to $f(u(t, 0_-))$, hence $f(u(t, 0_-)) \leq F$ for a.e. $t > 0$. At last, since we already proved that u satisfies the first Rankine-Hugoniot condition, we have $f(u(t, 0_-)) = f(u(t, 0_+))$, hence $f(u(t, 0_\pm)) \leq F$ for a.e. $t > 0$. \square

4 Proof of Proposition 2.4

We limit ourself to the proof of the last statement, which deals with non-classical discontinuities occurring at $x = 0$. Moreover, we consider only the case with $\mathbf{F} \equiv F$ constant belonging to $[0, f^+]$; the general case is analogous. The statement is obvious if $F = 0$, due to **(S.5)** and the fact that $f(u) \geq 0$. We can therefore assume that $F > 0$ and that $x \mapsto u(t_0, x)$ has a (stationary) non-classical shock (u_ℓ, u_r) , with $v_\ell < v_r$ and $f(u_\ell) = f(u_r) \doteq f \leq F$. We want to prove that $f = F$. Consider the test function

$$\phi(t, x) \doteq \left[\int_{|x|-\varepsilon}^{\infty} \varphi_\varepsilon(z) dz \right] \left[\int_{t-t_0+\varepsilon}^{t-t_0+2\varepsilon} \varphi_\varepsilon(z) dz \right],$$

where φ_ε is a smooth approximation of the Dirac mass centered at 0_+ , $\delta_{0_+}^D$, namely

$$\varphi_\varepsilon \in \mathbf{C}_c^\infty(\mathbb{R}; \mathbb{R}_+), \quad \varepsilon > 0, \quad \text{supp}(\delta_\varepsilon) \subseteq [0, \varepsilon], \quad \|\varphi_\varepsilon\|_{\mathbf{L}^1(\mathbb{R}; \mathbb{R})} = 1, \quad \varphi_\varepsilon \rightarrow \delta_{0_+}^D.$$

Observe that as ε goes to zero

$$\begin{aligned} \phi(t_0, x) &\equiv 0 \rightarrow 0, \\ \phi(t, 0) &= \int_{t-t_0+\varepsilon}^{t-t_0+2\varepsilon} \varphi_\varepsilon(z) dz \rightarrow \delta_{t_0-}^D(t), \\ \phi_t(t, x) &= \left[\int_{|x|-\varepsilon}^{\infty} \varphi_\varepsilon(z) dz \right] \left[\varphi_\varepsilon(t-t_0+2\varepsilon) - \varphi_\varepsilon(t-t_0+\varepsilon) \right] \rightarrow 0, \\ \chi_{\mathbb{R}_\pm}(x) \phi_x(t, x) &\rightarrow \mp \delta_{0_\pm}^D(x) \delta_{t_0-}^D(t). \end{aligned}$$

Then by (8) for all k belonging to the interval $(\check{v}(w_\ell, F), \check{v}(v_r, F))$ we have

$$\begin{aligned} &\mathbf{Q}^k(u_\ell) - \mathbf{Q}^k(u_r) + f \left[\frac{k}{F} - \frac{1}{p^{-1}(\mathbf{W}(u_\ell) - k)} \right]_+ \\ &= \left[\frac{f}{p^{-1}(\mathbf{W}(u_\ell) - k)} - k \right] + f \left[\frac{k}{F} - \frac{1}{p^{-1}(\mathbf{W}(u_\ell) - k)} \right] = \left[\frac{f}{F} - 1 \right] k \geq 0. \end{aligned}$$

Since $f \leq F$, the above estimate implies that $f = F$ and this concludes the proof of the constraint saturation claim.

It remains to prove that $\mathbf{W}(u_n(t, \cdot))$ may only have decreasing jumps at $x = 0$, in the precise sense (11). To do so, let us observe that whatever be the jump in u^n at time t across $x = 0$, there holds

$$f(u_n(t, 0_-)) = f(u_n(t, 0_+)) \geq 0 \quad \text{and} \quad \mathbf{W}(u_n(t, 0_-)) \geq \mathbf{W}(u_n(t, 0_+)). \quad (26)$$

It is possible to pass to the limit in (26) arguing in an indirect way. Indeed, we have the following weak form for comparing the fluxes of the generalized momentum at $x = 0^-$ and $x = 0^+$: for all $\phi \in \mathbf{C}_c^0(t_i, t_{i+1}); \mathbb{R}_+$

$$\int_0^T f(u_n(t, 0_-)) \mathbf{W}(u_n(t, 0_-)) \phi(t, 0) dt \geq \int_0^T f(u_n(t, 0_+)) \mathbf{W}(u_n(t, 0_+)) \phi(t, 0) dt. \quad (27)$$

Arguing as in Proposition 3.9 or in the proof of the property **(S.5)** of the main theorem, we use the Gauss-Green theorem to convert each of the integrals at $x = 0^\pm$ into volumic terms (with integrals over $(0, T) \times \mathbb{R}_\pm$), and then pass to the limit as $n \rightarrow \infty$ using the strong convergence of u^n . This argument shows that (27) is inherited at the limit where u^n is replaced by u . Localization with the test function ensures (11).

5 Time-discrete nonlocal constraints in the phase transition model

Assume that we are given a map $\mathcal{Q} : \mathcal{E} \mapsto \mathcal{F}$, with

$$\mathcal{E} \doteq \mathbf{L}^\infty((0, \infty); \mathbf{BV}(\mathbb{R}; \Omega)) \cap \mathbf{PC}(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega)), \quad \mathcal{F} \doteq \mathbf{PC}((0, \infty); [f^-, f^+]).$$

We supply \mathcal{E} with the $\mathbf{L}^\infty(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ topology, while for \mathcal{F} we consider the topology of pointwise a.e. convergence. We assume that \mathcal{Q} satisfies the following properties :

- (**Q.1**) The discontinuities of $\mathcal{Q}[u]$ can only occur at times $t_i = i\Delta$, for some fixed $\Delta > 0$ (the minimal switching time) ;
- (**Q.2**) The value of $\mathcal{Q}[u](t)$ only depends on $u|_{[0,t] \times \mathbb{R}}$;
- (**Q.3**) The operator \mathcal{Q} is continuous with respect to the above mentioned topologies on \mathcal{E} and \mathcal{F} .

Examples of such operators and the underlying modeling motivations are detailed in [2, Section 1.4.1]. Typically, they reproduce the adaptation of the constraint level made at discrete times $t_i = i\Delta$ in response to the upstream averaged density of agents measured by continuous or discrete in time observations.

Let us sketch the extension of the preceding theory to such nonlocally constrained problems; nontrivial details are discussed at each point.

- The definition of solution to such models is exactly the same as Definition 2.1 with the additional requirement that for all $t > 0$ there hold $\mathbf{F}(t) = \mathcal{Q}[u](t)$.
- The construction of solutions is fully analogous to the one of Section 3.4 with the only difference that the constraint level F_i^n on the interval $(t_i, t_{i+1}) = (i\Delta, (i+1)\Delta)$ (with the explicit dependence of the constraint level F_i on the approximation parameter n) is computed as $F_i^n \doteq \mathcal{Q}[u^n](t_i)$; note that (**Q.2**) makes this choice meaningful. This corresponds to the standard splitting procedure for approximation of coupled problems. At the practical level, the operator \mathcal{Q} can also be discretized (see [5] for typical examples and for the general view on consistency of such discretizations), however in principle the wave-front tracking procedure allows for the computation of \mathcal{Q} on u^n .
- The dependence of F_i^n on n is handled by the basic compactness argument, being understood that $F_i^n \in [f^-, f^+]$. Let us stress that the switching time Δ is independent of n . We denote by F_i the limit (along the suitable subsequence) of F_i^n . We can consider a finite number of switches, or even afford for $i \in \mathbb{N}$ upon using the diagonal extraction argument.
- The uniformity of the bounds on the total variation of u^n (at every fixed time horizon) is ensured by the uniformity of the Lipschitz constant in Lemma 2.6.
- In the passage to the limit, we have to care about the convergence $F_i^n \rightarrow F_i$. The constraint level \mathbf{F} appears explicitly at points (**S.4**), (**S.5**) of Definition 2.1. Handling the dependence of \mathbf{F}^n on n is easy due to the continuity of $N_F^k(u)$ with respect to $f \in [f^-, f^+]$, and to the obvious possibility to pass to the limit in (**S.5**).
- Finally, the link $F^n(t) = \mathcal{Q}[u^n](t)$ is preserved at the limit, for a.e. $t > 0$, due to (**Q.3**) and the convergence of (the suitable subsequence of) $(u^n)_n$ in the $\mathbf{L}^\infty(\mathbb{R}_+; \mathbf{L}_{\text{loc}}^1(\mathbb{R}; \Omega))$ topology, see the conclusion of §3.6.. In the case \mathcal{Q} is replaced by fully discrete approximations \mathcal{Q}^n , consistency properties are also required at this step of the argumentation. For the sake of conciseness, we do not pursue this line here.

To sum up, the result of Theorem 2.8 readily extends to $[f^-, f^+]$ -valued non-local constraints verifying the structural properties (**Q.1**), (**Q.2**), (**Q.3**).

While the two latter conditions are natural for the whole class of traffic models with non-local point constraint, the assumption (**Q.1**) and the restriction $F \geq f^-$ result from the technical limitations of our BV -based approach. Further work on this kind of models requires either smoother interaction potential terms (the Υ terms) in the Glimm-like functional \mathcal{T} , or less restrictive compactness tools such as compensated compactness.

Acknowledgements

The second author acknowledges the support of the Région Bourgogne Franche-Comté, projet 2017-2020 “Analyse mathématique et simulation numérique d’EDP issus de problèmes de contrôle et du trafic routier”. The last author is member of GNAMPA. He acknowledges the support of the National Science Centre, Poland, Project “Mathematics of multi-scale approaches in life and social sciences” No. 2017/25/B/ST1/00051, by the INdAM-GNAMPA Project 2020 “Dalla Buona Posizione alla Teoria dei Giochi nelle Leggi di Conservazione” and by University of Ferrara, FIR Project 2019 “Leggi di conservazione di tipo iperbolico: teoria ed applicazioni”. The publication has been prepared with the support of the “RUDN University Program 5-100”.

References

- [1] ANDREIANOV, B., DONADELLO, C., RAZAFISON, U., AND ROSINI, M. D. Qualitative behaviour and numerical approximation of solutions to conservation laws with non-local point constraints on the flux and modeling of crowd dynamics at the bottlenecks. *ESAIM: M2AN* 50, 5 (2016), 1269–1287.
- [2] ANDREIANOV, B., DONADELLO, C., RAZAFISON, U., AND ROSINI, M. D. Analysis and approximation of one-dimensional scalar conservation laws with general point constraints on the flux. *J. Math. Pures Appl. (9)* 116 (2018), 309–346.
- [3] ANDREIANOV, B., DONADELLO, C., AND ROSINI, M. D. Crowd dynamics and conservation laws with nonlocal constraints and capacity drop. *Mathematical Models and Methods in Applied Sciences* 24, 13 (2014), 2685–2722.
- [4] ANDREIANOV, B., DONADELLO, C., AND ROSINI, M. D. A second-order model for vehicular traffics with local point constraints on the flow. *Mathematical Models and Methods in Applied Sciences* 26, 04 (2016), 751–802.
- [5] ANDREIANOV, B., DONADELLO, C., AND ROSINI, M. D. A second-order model for vehicular traffics with local point constraints on the flow. *Mathematical Models and Methods in Applied Sciences* 26, 04 (2016), 751–802.
- [6] ANDREIANOV, B., GOATIN, P., AND SEGUIN, N. Finite volume schemes for locally constrained conservation laws. *Numerische Mathematik* 115, 4 (2010), 609–645.
- [7] ANDREIANOV, B., AND SYLLA, A. A macroscopic model to reproduce self-organization at bottlenecks. In *Finite volumes for complex applications IX*, R. Kloefkorn, Ed. Springer Proc. in Math. Stat., Cham, 2020, pp. 243–254.
- [8] AW, A., AND RASCLE, M. Resurrection of “second order” models of traffic flow. *SIAM J. Appl. Math.* 60, 3 (2000), 916–938 (electronic).
- [9] BENYAHIA, M., DONADELLO, C., DYMSKI, N., AND ROSINI, M. D. An existence result for a constrained two-phase transition model with metastable phase for vehicular traffic. *NoDEA Nonlinear Differential Equations Appl.* 25, 5 (2018), Art. 48, 42.
- [10] BENYAHIA, M., AND ROSINI, M. D. Entropy solutions for a traffic model with phase transitions. *Nonlinear Analysis: Theory, Methods & Applications* 141 (2016), 167 – 190.
- [11] BENYAHIA, M., AND ROSINI, M. D. A macroscopic traffic model with phase transitions and local point constraints on the flow. *Networks and Heterogeneous Media* 12, 2 (2017), 297–317.
- [12] BENYAHIA, M., AND ROSINI, M. D. Lack of bv bounds for approximate solutions to a twophase transition model arising from vehicular traffic. *Math. Methods Appl. Sci* (2020).
- [13] BLANDIN, S., WORK, D., GOATIN, P., PICCOLI, B., AND BAYEN, A. A general phase transition model for vehicular traffic. *SIAM J. Appl. Math.* 71, 1 (2011), 107–127.
- [14] BRESSAN, A. *Hyperbolic systems of conservation laws*, vol. 20 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2000.
- [15] COLOMBO, R. M. Hyperbolic phase transitions in traffic flow. *SIAM J. Appl. Math.* 63, 2 (2002), 708–721 (electronic).
- [16] COLOMBO, R. M., AND GOATIN, P. A well posed conservation law with a variable unilateral constraint. *J. Differential Equations* 234, 2 (2007), 654–675.
- [17] COLOMBO, R. M., GOATIN, P., AND PRIULI, F. S. Global well posedness of traffic flow models with phase transitions. *Nonlinear Anal.* 66, 11 (2007), 2413–2426.
- [18] DAL SANTO, E., ROSINI, M. D., DYMSKI, N., AND BENYAHIA, M. General phase transition models for vehicular traffic with point constraints on the flow. *Mathematical Methods in the Applied Sciences* (2017), 1–19.
- [19] DELLE MONACHE, M. L., AND GOATIN, P. Stability estimates for scalar conservation laws with moving flux constraints. *Netw. Heterog. Media* 12, 2 (2017), 245–258.
- [20] DYMSKI, N., GOATIN, P., AND ROSINI, M. D. Existence of BV solutions for a non-conservative constrained Aw-Rascle-Zhang model for vehicular traffic. working paper or preprint, Feb. 2018.
- [21] GARAVELLO, M., AND GOATIN, P. The Aw-Rascle traffic model with locally constrained flow. *Journal of Mathematical Analysis and Applications* 378, 2 (2011), 634 – 648.
- [22] GOATIN, P. The Aw–Rascle vehicular traffic flow model with phase transitions. *Mathematical and computer modelling* 44, 3 (2006), 287–303.
- [23] HOLDEN, H., AND RISEBRO, N. *Front Tracking for Hyperbolic Conservation Laws*. Applied Mathematical Sciences. Springer Berlin Heidelberg, 2013.
- [24] LIGHTHILL, M. J., AND WHITHAM, G. B. On kinematic waves. II. A theory of traffic flow on long crowded roads. *Proc. Roy. Soc. London. Ser. A.* 229 (1955), 317–345.
- [25] PANOV, E. Generalized solutions of the Cauchy problem for a transport equation with discontinuous coefficients. In *Instability in models connected with fluid flows. II*, vol. 7 of *Int. Math. Ser. (N. Y.)*. Springer, New York, 2008, pp. 23–84.
- [26] RICHARDS, P. I. Shock waves on the highway. *Operations Res.* 4 (1956), 42–51.
- [27] ZHANG, H. M. A non-equilibrium traffic model devoid of gas-like behavior. *Transportation Research Part B: Methodological* 36, 3 (2002), 275–290.

Boris Andreianov, INSTITUT DENIS POISSON (CNRS UMR7013), UNIVERSITÉ DE TOURS, UNIVERSITÉ D’ORLÉANS, PARC GRAND-MONT, 37200 TOURS, FRANCE
PEOPLES’ FRIENDSHIP UNIVERSITY OF RUSSIA (RUDN UNIVERSITY) 6 MIKLUKHO-MAKLAYA ST, MOSCOW, 117198, RUSSIAN FEDERATION
E - mail address: boris.andreianov@univ-tours.fr

Carlotta Donadello, LABORATOIRE DE MATHÉMATIQUES, CNRS UMR 6623, UNIVERSITÉ DE FRANCHE - COMTÉ, 16 ROUTE DE GRAY,
25030 BESANÇON, FRANCE
E - mail address: `carlotta.donadello@univ-fcomte.fr`

Massimiliano D. Rosini, (Corresponding author), DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI FERRARA, VIA
MACHIAVELLI 35, 44121 FERRARA, ITALY
INSTYTUT MATEMATYKI, UNIwersYTET MARIi CURIE-SKOŁODOWSKIEJ, PL. MARIi CURIE-SKOŁODOWSKIEJ 1, 20-031 LUBLIN, POLAND
E - mail address: `rsnmsm@unife.it`