



**HAL**  
open science

## **Regional trends in eutrophication across the Loire river basin during the 20th century based on multi-proxy paleolimnological reconstructions**

Anthony Foucher, Olivier Evrard, Sylvain Huon, Florence Curie, Irène Lefèvre, Véronique Vaury, Olivier Cerdan, Rosalie Vandromme, Sébastien Salvador-Blanes

### ► To cite this version:

Anthony Foucher, Olivier Evrard, Sylvain Huon, Florence Curie, Irène Lefèvre, et al.. Regional trends in eutrophication across the Loire river basin during the 20th century based on multi-proxy paleolimnological reconstructions. *Agriculture, Ecosystems & Environment*, 2020, 301, pp.107065. 10.1016/j.agee.2020.107065 . hal-02877523

**HAL Id: hal-02877523**

**<https://hal.science/hal-02877523>**

Submitted on 22 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manuscript Number: AGEE25394R2

Title: Regional trends in eutrophication across the Loire river basin during the 20th century based on multi-proxy paleolimnological reconstructions

Article Type: Research Paper

Keywords: terrigenous organic matter, autochthonous production, agricultural inputs, agriculture intensification, soil erosion, Anthropocene

Corresponding Author: Dr. Anthony Foucher,

Corresponding Author's Institution:

First Author: Anthony Foucher

Order of Authors: Anthony Foucher; Olivier Evrard; Sylvain Huon; Florence Curie; Irène Lefèvre; Véronique Vauray; Olivier Cerdan; Rosalie Vandromme; Sébastien Salvador-Blanes

Manuscript Region of Origin: FRANCE

Abstract: Excessive inputs of sediment and acceleration of primary production have been observed worldwide in a large number of water bodies. Human-environment interactions were recognized as one of the main drivers of this evolution during the 20th century with the occurrence of major landscape changes and a greater use of agricultural inputs. In this study, we used paleo-production proxies such as chlorophyll-a, organic matter properties (TOC and TN concentrations,  $\delta^{13}C$  and  $\delta^{15}N$ ) measured in sediment cores dated with fallout  $^{210}Pb_{ex}$  and  $^{137}Cs$  activities for reconstructing changes in accumulation rates and sources of organic matter during the recent period of agricultural intensification (1920-2020). In order to record these changes at the regional scale, sediment cores were collected at the outlet of several headwater catchments (n=9), covering a wide range of land covers / land uses across the Loire River basin (117,000 km<sup>2</sup>), France. The rates of sedimentary organic matter deposition in the studied water bodies accelerated from 1950 onwards (+48%). Between 1950 and 1970, the signature of sedimentary organic matter indicates a dominant contribution of soil-derived inputs. This period corresponds to major landscape modifications across the basin (land consolidation, stream re-design, implementation of tile drains) driving a general acceleration of erosion rates. Then, from 1960 onwards, chlorophyll-a and C/N proxies indicate an increase in primary production coupled with a decrease of terrigenous supply in agricultural catchments. These proxies were strongly correlated to the agricultural inputs during the 1955-1990 period (e.g.,  $r=0.9$  between chlorophyll-a content and N inputs), suggesting a progressive eutrophication of these reservoirs driven by increasing fertilizer use. During these 35 years, sedimentary organic matter deposition rates

1 **Regional trends in eutrophication across the Loire river basin during the**  
2 **20<sup>th</sup> century based on multi-proxy paleolimnological reconstructions**

3

4 Anthony Foucher (1), Olivier Evrard (1), Sylvain Huon (2), Florence Curie (3), Irène Lefèvre  
5 (1), Véronique Vaury (2), Olivier Cerdan (4), Rosalie Vandromme (4), Sébastien Salvador-  
6 Blanes (3)

7

8 (1) Laboratoire des Sciences du Climat et de l'Environnement (LSCE/IPSL), UMR 8212  
9 (CEA/CNRS/UVSQ), –Université Paris-Saclay, 91191 Gif-sur-Yvette Cedex, France

10 (2) Sorbonne Université, Institut d'Ecologie et des Sciences de l'Environnement de Paris (iEES), Case 237,  
11 4 place Jussieu. 75252 Paris cedex 05, France

12 (3) Laboratoire GéoHydrosystèmes Continentaux (GéHCO), E.A 6293, Université F. Rabelais de Tours,  
13 Faculté des Sciences et Techniques, Parc de Grandmont, 37200, Tours, France

14 (4) Département Risques et Prévention, Bureau de Recherches Géologiques et Minières (BRGM), 3 avenue  
15 Claude Guillemin, 45060, Orléans, France

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37 Abstract

38

39 Excessive inputs of sediment and acceleration of primary production have been observed  
40 worldwide in a large number of water bodies. Human-environment interactions were recognized as  
41 one of the main drivers of this evolution during the 20<sup>th</sup> century with the occurrence of major  
42 landscape changes and a greater use of agricultural inputs.

43 In this study, we used paleo-production proxies such as chlorophyll-a, organic matter  
44 properties (TOC and TN concentrations,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) measured in sediment cores dated with fallout  
45  $^{210}\text{Pb}_{\text{ex}}$  and  $^{137}\text{Cs}$  activities for reconstructing changes in accumulation rates and sources of organic  
46 matter during the recent period of agricultural intensification (1920-2020). In order to record these  
47 changes at the regional scale, sediment cores were collected at the outlet of several headwater  
48 catchments (n=9), covering a wide range of land covers / land uses across the Loire River basin  
49 (117,000 km<sup>2</sup>), France.

50 The rates of sedimentary organic matter deposition in the studied water bodies accelerated  
51 from 1950 onwards (+48%). Between 1950 and 1970, the signature of sedimentary organic matter  
52 indicates a dominant contribution of soil-derived inputs. This period corresponds to major landscape  
53 modifications across the basin (land consolidation, stream re-design, implementation of tile drains)  
54 driving a general acceleration of erosion rates. Then, from 1960 onwards, chlorophyll-a and C/N  
55 proxies indicate an increase in primary production coupled with a decrease of terrigenous supply in  
56 agricultural catchments. These proxies were strongly correlated to the agricultural inputs during the  
57 1955-1990 period (e.g.,  $r=0.9$  between chlorophyll-a content and N inputs), suggesting a progressive  
58 eutrophication of these reservoirs driven by increasing fertilizer use. During these 35 years,  
59 sedimentary organic matter deposition rates increased on average by 30%. During the 1990s, despite a  
60 slight reduction in fertilizer use (-13%), the paleo-production proxies (e.g. chlorophyll-a) still indicate  
61 a positive trend suggesting the contribution of another driving factor such as climate warming or again  
62 nutrient release from soils and sediments.

63 In the absence of long term geochemical and sediment input monitoring, paleolimnological  
64 reconstructions provide a powerful tool to reconstruct past agricultural pressures in rural

65 environments. This study illustrates the impact of intensive farming on water body siltation driven by  
66 varying sources of organic material during the 20<sup>th</sup> century. In addition, these results suggest that  
67 eutrophication processes of these reservoirs with contrasting land uses started during the 1960-1970  
68 period and are still ongoing nowadays.

69

70 Keywords: terrigenous organic matter, autochthonous production, agricultural inputs, agriculture  
71 intensification, soil erosion, Anthropocene

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

## 89 I. Introduction

90

91           Intensification of agricultural practices during the second half of the 20<sup>th</sup> century induced a  
92 significant acceleration of soil erosion and sediment delivery worldwide (Dearing & Jones, 2003;  
93 Cerdan *et al.*, 2010). This period of agriculture development is associated with a greater use of  
94 agricultural inputs including industrially produced fertilizers. Global fertilizer consumption was  
95 multiplied by 4.5 between 1960 and 2015 according to the FAO database (Tilman *et al.*, 2001; Lu &  
96 Tian, 2017). Both landscape management and the increase use of agricultural inputs have generated  
97 deleterious effects on terrestrial and aquatic ecosystems (e.g. Boardman & Poesen, 2006; Rowan *et al.*,  
98 2012). Among these adverse consequences, the excessive delivery of sediment may contribute to the  
99 siltation of water bodies including dams, ponds, lakes or river systems (e.g. Gellis *et al.*, 2006;  
100 Downing *et al.*, 2008; Ahn *et al.*, 2010). Two main sources of material can contribute to this sediment  
101 accumulation: sediment originating from terrestrial sources (e.g. soil erosion, river channel erosion)  
102 including mineral and organic material supply or organic matter associated with excessive primary  
103 production induced by eutrophication (e.g. Meyers & Ishiwatari, 1995; Gudas *et al.*, 2017). Causes  
104 and mechanisms of water body eutrophication were widely studied during the 1970s and 1980s in a  
105 large number of lakes in the Northern hemisphere (Glooschenko *et al.*, 1974; Hecky & Kilham, 1988).  
106 This research clearly outlined the key role of phosphorous (P) and nitrogen (N) point-based and  
107 diffuse source pollution in the bloom of algae in aquatic environments (Bennett *et al.*, 2001). Despite  
108 substantial management efforts implemented since this period to regulate agricultural N and P inputs  
109 and to improve wastewater treatment (EEC, 1991a; b), the excess of phytoplanktonic production  
110 remains problematic in coastal and continental waters (Le Moal *et al.*, 2019). According to the 2018  
111 European water assessment status, at least 38% of the continental water bodies have agricultural  
112 diffuse source pollution in Europe (EEA, 2018).

113           In the absence of long term geochemical and algae production monitoring in river systems  
114 (Minaudo *et al.*, 2015; Chen *et al.*, 2016), trajectories of eutrophication and paleoproduction are less

115 documented over the last century. Paleolimnological techniques based on the collection of sediment  
116 cores can provide a useful tool to reconstruct primary productivity over the last century (Brenner *et al.*,  
117 1999; Anderson *et al.*, 2014). To this end, a large number of methods was developed in order to  
118 reconstruct changes in paleo-productivity in sediment cores (e.g. Meyers, 1994; Lu *et al.*, 2010),  
119 including the analysis of pigment contents in lake sediments (Waters *et al.*, 2005; Florian *et al.*, 2015).  
120 Chlorophyll-a and its degradation products are the most abundant pigments in lake sediments (Leavitt  
121 & Hodgson, 2001). They were widely analyzed for reconstructing histories of eutrophication, and  
122 assessing the respective impacts of human activities and climate change (Korosi & Smol, 2012;  
123 Stewart *et al.*, 2015). Chlorophyll-a concentration can be quantified through a wide range of chromatic  
124 techniques (Millie *et al.*, 1993; Hodgson *et al.*, 1997). In recent years, the potential of the rapid and  
125 non-destructive Visual Reflectance Spectroscopy has been demonstrated to detect chlorophyll-a and  
126 the associated pheopigments in sedimentary sequences (Wolfe *et al.*, 2006). In this way, a variety of  
127 indices were used to infer chlorophyll-a and its degradation products abundances including its relative  
128 absorbance depth at 660/670 nm (Rein & Sirocko, 2002), sediment reflectance near 675 nm (Wolfe *et*  
129 *al.*, 2006), the absorbance peak between 650 and 700 nm (Michelutti *et al.*, 2010) or again the Q7/4  
130 ratio developed by Debret *et al.* (2011).

131 Stable C and N isotope signatures of sediment organic matter ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) provide one of the most  
132 widely used methods to identify changes in sources of organic material. This methodology was  
133 applied to palaeoenvironmental reconstructions based on the analysis of lake sediments (e.g. Meyers,  
134 1994; Huon *et al.*, 2018). The carbon isotopic composition of plants is variable and depends on the  
135 source of carbon assimilated. Terrestrial plant record  $\delta^{13}\text{C}$  values range from -13 to -30‰ (C3 plants  
136 are associated with values of -23 to -30‰ whereas C4 plants correspond to values between -9 and -  
137 17‰ with an average value of -13‰) - (Meyers, 1994; Bowsher *et al.*, 2008). Freshwater aquatic  
138 plants and plankton have  $\delta^{13}\text{C}$  values ranging between -25 and -30‰. Both terrestrial and aquatic  
139 organic matter may be distinguished through the comparison of their C/N ratio. Macrophyte residues  
140 have C/N ratios >20 but mineralization and aggregation in soils reduces C/N to *ca.* 10 (Meyers, 1994).  
141 In addition,  $\delta^{15}\text{N}$  can provide information on biogeochemical processes including N consumption by

142 organisms, varying sources of N, denitrification – nitrification and diagenetic processes (e.g. Jinglu *et*  
143 *al.*, 2007; Gälman *et al.*, 2009).

144 Although these proxies were widely used in sediment cores collected in estuarine, coastal  
145 areas or even in urban environments to investigate the relationship between nutrient inputs and a  
146 change in the water body trophic status (e.g. Voß & Struck, 1997; Rabalais *et al.*, 2007), many fewer  
147 studies were conducted in small reservoirs located in continental environments. In addition, to the best  
148 of our knowledge, the study of freshwater reservoirs located in contrasted agricultural environments at  
149 the large basin scale (<100,000 km<sup>2</sup>) has never been conducted especially considering the second part  
150 of the 20<sup>th</sup> century.

151 The current research investigates the potential of sedimentary sequences collected at the outlet  
152 of contrasted rural catchments (with land cover dominated by forest, grassland or arable land) to  
153 reconstruct regional paleoproductivity trends. The objectives of this work are to determine the rates of  
154 organic matter deposition in sediments and to identify the sources of organic matter accumulated  
155 during the 20<sup>th</sup> century in these reservoirs, through the discrimination of the terrigenous (soil-derived)  
156 *vs.* autochthonous (biological) inputs. Finally, these results will be compared to available agricultural  
157 N input data and to current knowledge on soil erosion in these catchments.

158

159

## 160 II. Site and methods

161

### 162 1. Study sites

163 The Loire river basin (117,000 km<sup>2</sup>, France) – (Fig. 1) is characterized by contrasted climate  
164 (annual rainfall varying between 600 and 1300 mm per year); landform (mean altitude = 300 m, 10%  
165 of the catchment >800 m); lithology (sedimentary; metamorphic and magmatic rocks) and land use  
166 conditions between its upper and lower parts (Fig.1). Characteristics of the studied sub-catchments  
167 were summarized in Table 1.

168 This large basin was impacted by major changes in agricultural practices during the second  
169 part of the 20<sup>th</sup> century. Some areas experienced an intensification of agricultural practices (mainly in



170 the middle and lower catchment sections) - (Foucher *et al.*, 2014) whereas other parts were impacted  
171 by land abandonment following rural depopulation (mainly in upper parts of the catchment) -  
172 (Foucher *et al.*, 2019b) – Table 2. After 1950, the spreading of “NPK” fertilizers has significantly  
173 increased in France in particular in the Loire River basin (Poisvert *et al.*, 2017). In response to these  
174 changes, the Loire river basin – as most European basins – has been impacted by eutrophication  
175 processes. The first monitoring surveys focusing on primary production have demonstrated the  
176 pronounced eutrophic conditions of the Loire river in the early 1980s (Meybeck *et al.*, 1988;  
177 Etcheber *et al.*, 2007). A recent study showed that the continuous decline then stabilization of  
178 phosphate and nitrate concentrations in the river since 1990 led to the significant reduction of  
179 phytoplankton biomass across the whole river system (Minaudo *et al.*, 2015). Nevertheless,  
180 eutrophication still represents a crucial environmental problem in this basin nowadays. According to  
181 a recent report of the regional water agency, eutrophication remains one of the main causes of  
182 degradation of the ecological status of water bodies (lakes, ponds, dams, etc.) with 29% of those in  
183 medium ecological status, 18% in low ecological status and even 26% in bad ecological status  
184 (AELB, 2013).

185         Nine reservoirs (lakes and ponds, with a surface ranging between 1.2 and 57 ha) located at the  
186 outlet of rural catchments (<100km<sup>2</sup>) were selected in various environments across the Loire river  
187 basin to reconstruct trends in rates of sedimentary organic material deposition and to identify the  
188 sources of sediment (Fig. 1). These reservoirs were selected among the 14,000 natural and artificial  
189 water bodies of the Loire basin because their catchment histories are well known. Furthermore, to the  
190 best of our knowledge, these reservoirs were never dredged or drained during prolonged periods.  
191 These water bodies are mainly artificial (n=8 artificial and one natural (Tazenat lake)) and they are  
192 connected by ephemeral river networks to their drainage area during heavy rainfall. These sites were  
193 selected in order to be representative of the diversity of landscape, climate, topography and  
194 agricultural practices found across the Loire basin. Although they are unevenly distributed in space  
195 across the basin, they cover all the characteristics of the agricultural landscapes found in this region.  
196 Among these catchments, four are mainly covered with grassland, three with arable land and two with  
197 forests.

198 The characteristics of these catchments are summarized in Table 2.

199

## 200 2. Materials and methods

201

202 Fieldwork was conducted between January 2017 and June 2018. Sediment cores (n=9) were  
203 collected in the nine water bodies using an Uwitec gravity corer (Uwitec®, Mondsee, Austria)  
204 equipped with a 90 mm PVC liner and a floating platform available at the Tours University (GeHCo  
205 laboratory, France). In each artificial reservoir, the sediment core was collected in the dam vicinity  
206 ( $\approx$ 50 to 100m from the dam) in the zone where the sedimentation rates are the highest. In the natural  
207 reservoir of Tazenat, the sediment core was retrieved in the center of the water body. Given the  
208 relatively small size of these reservoirs (<52 ha), it is assumed that the sediment properties recorded at  
209 these sampling sites are representative of those found across the entire water bodies with the exception  
210 of variations for the sedimentation rates.

211 Sampling locations and bathymetric data were recorded using a Garmin® Echomaps depth sounder  
212 (Garmin, Lenexa, USA).

213

### 214 2.1. Chronology

215 Sediment dating of the nine cores (selected to be representative of the entire water bodies)  
216 were established using Caesium-137 ( $^{137}\text{Cs}$ ) and excess Lead-210 ( $^{210}\text{Pb}_{\text{ex}}$ ) activities measured in 130  
217 samples of dried sediment material (*ca* 10g). These samples were regularly collected along the  
218 sediment cores with on average one sample taken every 6 cm. Gamma spectrometry measurements  
219 were obtained using coaxial N- and P- type HPGe detectors (Canberra/Ortec®) available at the  
220 Laboratoire des Sciences du Climat et de l'Environnement (Gif-sur-Yvette, France). The  $^{210}\text{Pb}$   
221 activities were determined at 46.5 keV.  $^{210}\text{Pb}_{\text{ex}}$  was calculated by subtracting the supported activity  
222 from the total  $^{210}\text{Pb}$  activity (measured at 46.5 keV) using two Radium-226 daughters, Lead-214  
223 (average count at 295.2 and 351.9 keV) and Bismuth-214 (609.3 keV). All measurements were

224 corrected for background level determined every two months as well as for detector and geometry  
225 efficiencies. Activities were also decay-corrected to the sampling date (Evrard *et al.*, 2016).

226 Age of the sediment sequence layers was determined using the Constant Rate of Supply model  
227 (CRS), (Appleby & Oldfield, 1978). This model assumes a constant rate of  $^{210}\text{Pb}_{\text{ex}}$  supply from  
228 atmospheric fallout, although sediment accumulation is allowed to vary throughout time. For  
229 improving the  $^{210}\text{Pb}_{\text{ex}}$  age model of some cores, the corrected CRS model described in Appleby (2001)  
230 was used. The  $^{210}\text{Pb}_{\text{ex}}$  age model validation was carried out through the identification of  $^{137}\text{Cs}$  peaks in  
231 the sedimentary sequences. This artificial radionuclide may originate from two sources in Western  
232 Europe: the Chernobyl accident (1986) and the thermonuclear weapons testing (maximal emissions in  
233 1963). Fukushima fallout was shown to be negligible in France (Evrard *et al.*, 2012). To distinguish  
234 between both potential  $^{137}\text{Cs}$  sources, Americium-241 ( $^{241}\text{Am}$ ), a daughter product of plutonium-241,  
235 was used to identify the  $^{137}\text{Cs}$  peak attributed to the maximum nuclear bomb fallout of 1963 (e.g  
236 Cambay *et al.*, 1989). Information related to sediment core dating was summarized in Table 3.

237

## 238 *2.2. Laboratory analyses*

239 Measurements of Visible Reflectance Spectroscopy (VRS) were obtained using a portable  
240 spectrophotometer (Konica® Minolta 700-D) equipped with a 3 mm target radius. Spectral reflectance  
241 was measured between 400 and 700 nm, with a 10 nm resolution. Before each set of measurements,  
242 the device was calibrated using a white and black calibration box. Measurements were continuously  
243 recorded on the fresh sediment surface with a 1-cm resolution, in each core.

244 The absorbance peak between 650 and 700 nm was extracted for a sequence of samples  
245 selected in the sedimentary sequences to cover 5-year periods on average as a spectral index to  
246 measure the summed concentration of all compounds associated with both primary and degraded  
247 chlorophyll-a (chlorophyll-a + all chlorophyll-a isomers + pheophytin-a + pheophorbide-a) -  
248 (Michelutti & Smol, 2016). The area under the absorbance peak from 650 to 700 nm was calculated  
249 following the procedure described in Das *et al* (2005). These values provide a relative information on  
250 chlorophyll-a occurrence index derived from the true area calculation. In order to reduce the noise of

251 the colorimetric time series, a LOESS smoothing model was applied using the free software PAST 3  
252 (Hammer *et al.*, 2001).

253

254 The occurrence of carbonate minerals was determined for 95 samples selected on the 9 master  
255 cores (on average one sample every 5 years) with a Bernard calcimeter (University of Tours, France)  
256 following the method described in NF ISO 10693 (1995). When detected, carbonates were  
257 systematically removed from the sample aliquots before performing analyses to determine the total  
258 organic carbon concentration (TOC) and stable C isotopes ratios of organic matter and clay-bound  
259 organic matter. To reduce possible leaching effects on sedimentary organic matter during carbonate  
260 removal, sediments were carefully treated with a 1N HCL solution during 96 hours under constant  
261 agitation and pH control in order not to drop below 2.5. Powder X-ray diffraction controls were  
262 conducted to ensure the full dissolution of all carbonate minerals in the sediment residue (calcite,  
263 dolomite, ankerite, ...). All the samples were rinsed six times in deionized water to remove dissolved  
264 salts. The solid residues were recovered by high velocity centrifugation (2500 rounds.min<sup>-1</sup>) for 15  
265 minutes each time (Huon *et al.*, 2002).

266 On these decarbonized sample aliquots, EA-IRMS analyses were conducted on dry sediment  
267 for elemental concentration (TOC and total nitrogen concentration TN) and stable C and N isotopes  
268 ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ ) measurements. These measurements were performed with a continuous flow Elementar®  
269 VarioPyro cube elemental analyzer (EA) coupled to a Micromass® Isoprime Isotope Ratios mass  
270 Spectrometer (IRMS) at the Institute of Ecology and Environmental Science of Paris. Oxygen for  
271 combustion was injected during 70 s (30 mL min<sup>-1</sup>) and temperatures were set at 850 °C and 1120 °C  
272 for the reduction and combustion furnaces, respectively (Agnihotri *et al.*, 2014). Analytical precision  
273 and repeatability were controlled with tyrosine samples calibrated against international standards  
274 (Coplen *et al.*, 1983). In the current research, the mean uncertainties were 0.2 ‰ for TN, 0.18 ‰ for  
275 TOC, 0.1 ‰ for  $\delta^{13}\text{C}$  and 0.2 ‰ for  $\delta^{15}\text{N}$ .

276

277 *2.3. Mass accumulation rate of organic material*

278 Organic matter Mass Accumulation Rates (MAR<sub>org</sub>, expressed in gC cm<sup>-2</sup> yr<sup>-1</sup>) were  
279 estimated for each individual core using the Sediment Accumulation Rate (SAR, cm yr<sup>-1</sup>, estimated  
280 with the <sup>210</sup>Pb<sub>ex</sub> decay), the Dry Bulk Density (DBD, expressed in g cm<sup>-3</sup> and calculated by measuring  
281 the amount of dry sediment in a known volume) and the total organic carbon concentration (TOC,  
282 expressed in gC g<sup>-1</sup>) following eq. 1. MAR<sub>org</sub> corresponds to the amount of organic material deposited  
283 at each individual coring site. This sediment may originate from terrestrial inputs (allochthonous  
284 inputs) or may be directly produced in the water body (autochthonous production).

285

$$286 \quad \text{MAR}_{\text{org}} [\text{g cm}^{-2} \text{ yr}^{-1}] = \text{DBD} [\text{g cm}^{-3}] \times \text{SAR} [\text{cm yr}^{-1}] \times \text{TOC} [\text{gC g}^{-1}] \text{ (eq. 1)}$$

287

#### 288 *2.4. Agricultural inputs*

289 Data from the CASSIS\_N model ([www.geosciences.univ-tours.fr/cassis](http://www.geosciences.univ-tours.fr/cassis)) - (Poisvert *et al.*,  
290 2017) was extracted for each individual catchment and for the entire Loire river basin. This model was  
291 used for estimating the mineral and organic nitrogen (N) consumption expressed in kg.ha<sup>-1</sup>. This  
292 model based on data from agricultural census is available between 1955 and 2015 with an annual  
293 resolution.

294

#### 295 *2.5. Statistical analyses*

296 The Mann-Kendall non-parametric test (MK-test) was used for detecting monotonic trends in  
297 temporal series (Warren & Gilbert, 1988). In this study, the temporal series correspond to the  
298 evolution of production proxies ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , TN, TOC, chlorophyll-a), MAR<sub>org</sub>, MAR<sub>min</sub> as well as  
299 agricultural inputs throughout time. This test was applied to confirm the occurrence of monotonic  
300 upward or downward trends of a given variable throughout time (P-value < 0.05). Trends can be  
301 positive, negative or non-null.

302 The non-parametric Buishand test (BU-test) was used for detecting the occurrence of changes in  
303 temporal series (Buishand, 1982). Buishand test with a P-value < 0.05 indicated a heterogeneous  
304 temporal trend between two periods.

305 In addition, the correlation coefficient (denoted  $r$ ) was used to quantify the direction (positive,  
306 negative) and strength of the linear association between two variables.

307

### 308 3. Results

309

#### 310 3.1. Chronology

311 According to the dating models, the deepest sediment layers in the cores were deposited before  
312 1925 (n=8) except for a single core where it is assumed to have deposited after 1950 (Table 3). The  
313 age controls provided by the detection of artificial radionuclides ( $^{137}\text{Cs}$  and  $^{241}\text{Am}$ ) were identified in  
314 four reservoirs for the Chernobyl fallout and in eight sequences for the 1963 fallout (Fig 4, 5 and 6).  
315 Errors between continuous  $^{210}\text{Pb}_{\text{ex}}$  age models and those obtained with the radiocesium were in  
316 average of 2 years for 1986 and 3 years for 1963. Chronology of two cores, respectively those of the  
317 Prugnolas pond and Malaguet lake (Fig. 1) was detailed in Foucher *et al.* (2019b; a). Chronology of  
318 the nine cores is detailed in Table 3.

319

#### 320 3.2. Trends in paleo-production proxies

321 Organic material production has significantly changed in seven of the nine study sites with a  
322 statistical increase of TOC concentrations in five of the reservoirs. For the two catchments dominated  
323 by forests, TOC contents varied between 94 and 180  $\text{gC kg}^{-1}$  (average value of  $130 \pm 24 \text{ gC kg}^{-1}$ ) –  
324 (Fig. 2). A non-significant trend in TOC content was recorded at both forested sites ( $P > 0.05$  MK-  
325 Test) – (Table 4). TOC content in catchments mainly occupied by grassland ranged between 18 and  
326 160  $\text{gC kg}^{-1}$  (average value of  $50 \pm 29 \text{ gC kg}^{-1}$ ) – (Fig. 3). Content in organic material increased by

327 0.04, 1.6, 2.6 and 7.8 % after 1950 (for Loroux, Goule, Beaurepaire and Tazenat reservoirs,  
328 respectively). The Loroux pond did not record any statistical change in TOC content. Nevertheless, a  
329 peak of organic material was identified in 1965 (TOC = 75 gC kg<sup>-1</sup>) – (Fig. 3). For the Malaguet lake,  
330 this peak was identified in 1956 (TOC = 159 gC kg<sup>-1</sup>) – (Fig. 2). During the same period, Tazenat lake  
331 recorded an acceleration of organic material accumulation, with TOC contents increasing from 2.6 to  
332 7.6 % (between 1950 and 1958). TOC concentrations measured in the three study sites dominated by  
333 arable land ranged between 28 and 157 gC kg<sup>-1</sup> (average value of 65 ± 28 gC kg<sup>-1</sup>) – (Fig. 4).  
334 Concentration in organic material decreased by 0.1 and 1.1% (for Brosse and Passavant ponds,  
335 respectively) although it increased by 1.3 % for the Boisvinet pond during the last century. Brosse and  
336 Passavant ponds recorded an acceleration of organic material accumulation in the middle of the 20<sup>th</sup>  
337 century (respectively between 1952 and 1960 (TOC = 157 gC kg<sup>-1</sup>) and between 1955 and 1966 (TOC  
338 = 95 gC kg<sup>-1</sup>) – (Fig. 4).

339 During the 20<sup>th</sup> century, δ<sup>13</sup>C results showed a significant positive trend in six of the studied  
340 reservoirs with a switch around 1973 ± 11 yr in grassland environments, in 1969 in catchments  
341 dominated by forest and in 1984 ± 1 yr in areas mainly occupied by arable land (Table 4). For the  
342 three remaining water bodies (Brosse, Boisvinet and Prugnolas pond), no statistical trend was  
343 observed (Table 4). In forested catchments, δ<sup>13</sup>C ranged between -27.4 ‰ and -29 ‰ (average value of  
344 -28 ± 0.4 ‰) – (Fig. 2). For the Malaguet lake, the δ<sup>13</sup>C decreased by 0.3 ‰ during the covered period  
345 while it decreased by 0.6 ‰ during the last century in the Prugnolas pond. For the pond located at the  
346 outlet of catchments dominated by grassland, the δ<sup>13</sup>C signature of organic matter ranged between -  
347 26.7 ‰ and -30.5 ‰ (average value of -28.6 ± 0.95 ‰) – (Fig. 3). For these four catchments, δ<sup>13</sup>C  
348 values decreased by 0.5, 0.6, 1.7 ‰ and increased by 0.54 ‰ during the 20<sup>th</sup> century (for the  
349 Beaurepaire, Goule, Tazenat and Loroux ponds, respectively). The δ<sup>13</sup>C of organic matter measured in  
350 sediment cores collected in reservoirs draining catchments under arable land ranged between -26.7 ‰  
351 and -32.3 ‰ (average value of -29.3 ± 1.4 ‰) – (Fig. 4). Values prior to 1950 were in average of -28.6  
352 ± 1.7 ‰ compared to -29.5 ± 1.1 ‰ afterwards. During this period, δ<sup>13</sup>C values decreased by 1.6 ‰  
353 and 0.32 ‰ (for Passavant and Boisvinet ponds, respectively) and increased by 0.2 ‰ in Brosse Pond.

354 The  $\delta^{15}\text{N}$  signature of organic matter displayed a significant positive trend in three reservoirs  
355 (Beaurepaire, Passavant and Boisvinet) and a negative trend in two other water bodies (Loroux and  
356 Prugnolas ponds) – (Table 4). Regarding the observed positive trend, the change in trajectory was  
357 recorded to occur around  $1977 \pm 11$  yr while it started in  $1976 \pm 26$  yr for negative trends. For the four  
358 remaining sites located under various land covers, no statistical trend was recorded (Table 4). For the  
359 forested catchments,  $\delta^{15}\text{N}$  decreased by 0.2 ‰ and 1.3 ‰ (for Malaguet and Prugnolas reservoirs,  
360 respectively) during the documented periods (Fig. 2). Under grassland environments, these values  
361 ranged between 4.1 ‰ and 8.8 ‰ (average  $7 \pm 1.4\%$ ).  $\delta^{15}\text{N}$  increased by 0.3‰ in the Beaurepaire  
362 pond and decreased by 0.4, 0.6 and 2.2 ‰ during the last century (for Goule, Loroux and Tazenat  
363 study sites, respectively) – (Fig. 3). In catchments dominated by arable land,  $\delta^{15}\text{N}$  ranged between 4.4  
364 ‰ and 8.1 ‰ ( $\pm 5.7$  ‰). When comparing values recorded prior to and after 1950,  $\delta^{15}\text{N}$  increased by  
365 0.5, 1 and 1.1 ‰ (for Brosse, Boisvinet and Passavant ponds, respectively) – (Fig. 4).

366 C/N ratios showed a significant negative trend in only two ponds (Loroux and Goule ponds)  
367 with a switch in this trend occurring around  $1969 \pm 6$  yr. For these two reservoirs, C/N ratio increased  
368 from 7.3 to 9.5 for the Goule pond and from  $8.6 \pm 0.2$  to  $8.9 \pm 0.6$  for the Loroux pond, in 1950 and  
369 2010, respectively (Fig. 3)

370 Statistical positive trends in chlorophyll-a were recorded in seven reservoirs. Only the  
371 Prugnolas pond and the Tazenat lake did not show any statistical trend (Table 4). In the forested  
372 catchment, this increase reached around  $160 \pm 205$  % (Fig. 2). In catchments dominated by grassland,  
373 chlorophyll-a values increased by  $136 \pm 87$  % after 1950 (Fig. 3). During this period, an increase of  
374  $190 \pm 160$  % was recorded in arable land environments (Fig. 4). Changes in these trends were recorded  
375 around  $1996 \pm 5$  yr  $1963 \pm 7$  yr and 1990 – for catchments dominated by arable land, grassland, and  
376 forest, respectively.

377

### 378 *3.3. Mass accumulation rate of organic material*

379 Accumulation rate of sedimentary organic material (originating from soil erosion and  
380 autochthonous production) has significantly changed during the last century. In the two catchments



381 occupied by forest, negative trends were recorded (decrease of  $160 \pm 100\%$  – Table 4). In these  
382 catchments, the deposition rates started to decrease around  $1963 \pm 23$  yr (Fig. 5). A significant positive  
383 trend in organic production was recorded in the four catchments occupied by grassland (MK test  $P$   
384  $< 0.05$ ) – (Table 4). In this context, organic matter deposition rate increased on average by  $58 \pm 23$  %  
385 between 1950 and 2015 (Fig. 5). In catchments dominated by arable land, a positive trend was only  
386 recorded in one catchment (Boisvinet catchment) with an acceleration of organic accumulation around  
387 1970. For the five catchments recording a positive trend, the average rate of deposition raised from  $67$   
388  $\pm 35$  gC m<sup>-2</sup> yr<sup>-1</sup> before 1950 to  $134 \pm 66$  gC m<sup>-2</sup> yr<sup>-1</sup> after 1950 (Fig. 5). In the two reservoirs where no  
389 statistical trend was observed (MK test  $P < 0.05$ ) - (Brosse and Passavant reservoirs), a significant  
390 increase of productivity was recorded between 1940 and 1975 (Fig. 5). On average, rates of organic  
391 deposits reached  $260 \pm 44$  gC m<sup>-2</sup> year<sup>-1</sup> before 1950,  $350 \pm 302$  gC m<sup>-2</sup> year<sup>-1</sup> between 1950 and 1975  
392 and  $318 \pm 126$  gC m<sup>-2</sup> year<sup>-1</sup> after 1975.

393

#### 394 *3.4. Agricultural inputs*

395 The analysis of N fertilizer inputs (mineral + organic) of the nine selected catchments shows  
396 the occurrence of three main trends. In the six catchments occupied by arable land and grassland, a  
397 significant increase in N fertilizer use was observed during the last century (MK test  $P < 0.05$ ) - (Table  
398 4) In these sites, a major acceleration occurred around  $1966 \pm 5$  yr (BU test) – (Fig. 6). On average,  
399 this consumption started to stabilize or to decrease in  $1994 \pm 5$  yr. For these catchments, consumption  
400 of N fertilizers increased on average by  $60 \pm 10\%$  for the 1955-2015 period. No statistical trend (MK  
401 test  $P > 0.05$ ) was observed for the Boisvinet catchment (site 5, Fig. 1) located in lowland  
402 environments dominated by arable land (Table 2).

403 Finally, the two catchments currently occupied by forest displayed a negative or the absence of any  
404 significant statistical trend in N fertilizer inputs (for the Malaguet and the Prugnolas catchment,  
405 respectively) – (Table 4). For the Malaguet site, a significant decrease in consumption was recorded  
406 around 1975. The use of N fertilizer inputs decreased by 20 % at this site (Fig. 6).

407

408

#### 409 4. Discussion

410

##### 411 *Local records*

412

413 The non-linear decrease and the independent variation of paleo-production proxies through  
414 time (chlorophyll-a, sedimentary bulk organic matter  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  and C/N concentration) does not  
415 suggest the post-depositional degradation of organic matter as no typical increasing logarithmic trend  
416 of TOC and TN concentrations was observed in the upper part of the sediment cores (Fig. 2, 3 and 4) -  
417 (Emerson *et al.*, 1985; Arndt *et al.*, 2013). Variation in individual proxies can therefore be attributed to  
418 a change in organic matter sources.

419 Results obtained on the nine sedimentary sequences collected at the outlet of contrasted  
420 catchments showed variable contributions of terrestrial versus autochthonous organic matter during  
421 the 20<sup>th</sup> century. These changes in organic matter sources were attributed to anthropogenic pressures  
422 observed in the landscape, with the intensification of anthropogenic activities and a greater use of  
423 agricultural inputs (e.g. Schindler Wildhaber *et al.*, 2012).

424 In the two forested catchments, the period prior to 1960 was characterized by the highest C/N  
425 ratios recorded in this study (ranging between 17 and 24), suggesting the dominant delivery of  
426 vascular debris (C/N >20 from vascular plants (Meyers, 1994)). These results were in agreement with  
427 our historical knowledge on the processes occurring in these catchments. During the 20<sup>th</sup> century, both  
428 catchments were affected by land abandonment, evolving from an open landscape, mostly occupied by  
429 grassland, to forested catchments. Between 1950 and 1960, major afforestation works were conducted  
430 in the Malaguet catchment inducing a significant acceleration of soil erosion (Foucher *et al.*, 2019b).  
431 The forest regrew spontaneously in the Prugnolas catchment and, as a consequence, soil erosion  
432 decreased progressively during the 20<sup>th</sup> century. The increase of values recorded by terrigenous  
433 organic matter proxies (C/N, TOC concentration) during the 1950-1960 period was recorded during  
434 this period of afforestation, soil disturbance and soil erosion. After this period, organic sediment  
435 sources have drastically changed as reflected by an increase of TOC contents in sediment (from 130 to  
436 170 gC kg<sup>-1</sup> and from 110 to 150 gC kg<sup>-1</sup>) as well as a decrease in the C/N ratios (from 24 to 13 and

437 from 16 to 11, for Prugnotas and Malaguet reservoirs, respectively) – (Fig. 2). In contrast, during this  
438 period, chlorophyll-a increased (450% and 160% for Prugnotas and Malaguet reservoirs, respectively).  
439 The strong negative relationship observed between chlorophyll-a and  $\delta^{13}\text{C}$  ( $r = 0.99$  and  $0.87$  for  
440 Prugnotas and Malaguet reservoirs, respectively) suggests a greater contribution of autochthonous  
441 production during the second half of the 20<sup>th</sup> century. During this period, agricultural inputs remained  
442 constant or even decreased (Fig. 6), and they were not correlated to paleo production proxies (e.g.  
443 correlation between chlorophyll-a vs N inputs  $r = 0.02$  and  $0.5$  for Prugnotas and Malaguet lakes). In  
444 the Prugnotas catchment, the increase of  $^{137}\text{Cs}$  activities recorded in sediment deposited after 1999 was  
445 attributed to a greater contribution of surface soil erosion during a landscape management period  
446 following the major 1999 windstorm (Foucher *et al.*, 2019a). During this period both TN  
447 concentration and  $\delta^{15}\text{N}$  increased as well as chlorophyll-a values ( $r = 0.98$  and  $0.99$ , respectively) –  
448 (Fig. 2). Acceleration of autochthonous production during the last 25 years can be therefore attributed  
449 to a release of soil nutrients.

450 In grassland areas (drained by the Beaurepaire, Tazenat and Loroux-Bottereau reservoirs),  
451 changes in sedimentary matter signatures were also recorded during the 20<sup>th</sup> century. Between 1950  
452 and 1970, a general increase in TOC contents (on average  $120 \pm 110$  %) associated with higher C/N  
453 organic ratios (average value of  $9.5 \pm 1$ ), an increase of MARorg rate ( $140 \pm 70\%$ ) and changes in  $\delta^{13}\text{C}$   
454 and  $\delta^{15}\text{N}$  proxies were observed (Fig. 3). A similar observation was made in catchments dominated by  
455 arable land (Passavant and Brosse basins). During this same period, TOC contents increased on  
456 average by  $63 \pm 1$  % and were associated with an acceleration of MARorg deposition ( $84 \pm 40$  %) and  
457 high C/N ( $11.5 \pm 0.5$ ). In addition, this period corresponds to an increase of  $^{137}\text{Cs}$  activity suggesting a  
458 mobilization of material originating from surface soil sources (Olley *et al.*, 2013) for le Loroux,  
459 Beaurepaire, Tazenat and Passavant catchments (Fig. 3 and 4).

460 In both contexts of arable land and grassland, this post Second World War (WWII) period corresponds  
461 to a phase of intensification of agricultural practices across the Loire river basin, with the  
462 implementation of major landscape modifications (e.g. land consolidation corresponding to the  
463 reorganization and the increase of plot size, stream redesign, tile drain creation) - (Vandromme *et al.*,  
464 2016; Chartin *et al.*, 2013; Grangeon *et al.*, 2017). The change in proxy values observed in the current

465 research together with this increase of accumulation of organic matter characterized by high C/N ratios  
466 very likely occurred in response to these landscape modifications (Mackie *et al.*, 2005).  
467 After this period of more intense management (after 1970) the MARorg, TOC and chlorophyll-a  
468 proxies increased following a statistically significant trend (Table 4). These results suggest a greater  
469 accumulation of autochthonous organic matter. MARorg values obtained in agricultural and  
470 abandoned catchments during the 2010-2017 period ( $255 \pm 170$  and  $78 \pm 22$  gC m<sup>-2</sup> yr<sup>-1</sup>, respectively)  
471 are in agreement with those obtained in previous studies (Fig. 5). For example, a median value of 350  
472 gC m<sup>-2</sup> yr<sup>-1</sup> and 56 gC m<sup>-2</sup> yr<sup>-1</sup> were respectively recorded in reservoirs draining cultivated and  
473 abandoned catchments - (Schlesinger & Bernhardt, 2013).

474

#### 475 *Regional trend*

476

477 The compilation of results obtained in the seven agricultural catchments demonstrates the  
478 continuous acceleration in organic material accumulation in ponds across the Loire River basin during  
479 the 20<sup>th</sup> century (MK-test P <0.05). Similar observations were made in agricultural catchments of the  
480 northern United-States (Heathcote *et al.*, 2013). In this area, the regional acceleration started around  
481 1950 and was accompanied by an increase of C/N ratios in sediment (Fig. 7). In previous  
482 paleolimnological studies, these increases were interpreted as historical periods characterized by the  
483 delivery of a high proportion of organic matter derived by soil erosion particularly during periods of  
484 greater human-induced environmental modifications (Guilizzoni *et al.*, 1996; Routh *et al.*, 2004). In  
485 addition to this change,  $\delta^{15}\text{N}$  signature recorded a general decrease suggesting a major change in  
486 organic matter sources. The beginning of acceleration of organic matter deposits (1950) recorded in  
487 these reservoirs occurred concomitantly with the major landscape changes that took place in the Loire  
488 river basin. These changes were associated with an acceleration of erosion processes and a higher  
489 sediment delivery to the reservoirs (Foucher *et al.*, 2014). This period of soil erosion was observed in  
490 various catchments across Europe. For example, a two to ten-fold acceleration of soil erosion rates was  
491 recorded in various catchments of the UK after 1950 in response to land-use change (Foster *et al.*,  
492 2011). In addition, the greater contribution of phytoplanktonic production was also observed from

493 1955 onwards: chlorophyll-a started to increase following the regional trends of N fertilizer inputs ( $r =$   
494  $0.9$  between chlorophyll-a contents and N inputs during the 1955-1990 period). From 1970 onwards,  
495 the terrigenous contribution decreased sharply following the trend of MARorg. After 1970, human  
496 pressure on landscape started to decrease in France (land consolidation programs mainly took place  
497 between 1955 and 1975, Andre & Polombo, 2013). After a period dominated by the terrigenous  
498 contribution to sediment deposited in the reservoirs, the MARorg showed a progressive acceleration  
499 from 1970 to 2015 (+40%). At the regional scale,  $\delta^{15}\text{N}$  is well correlated to N inputs ( $r = 0.76$ ), C/N  
500 ratio ( $r = 0.8$ ), to MARorg after 1970 ( $r = 0.72$ ) as well as to chlorophyll-a ( $r = 0.68$ ). The increase of  
501 organic matter deposition associated with a good correlation between phytoplanktonic proxies  
502 demonstrates a shift in the paleo productivity evolving from a system dominated by terrigenous inputs  
503 just after WWII to a period of dominant autochthonous production after 1970.

504 Between 1990 and 2015, the European legislation induced a slight reduction in fertilizer use in the  
505 Loire river basin (-13% on average) - (Poisvert *et al.*, 2017). Nevertheless, organic matter continues to  
506 accumulate in reservoirs (+17% in 15 years). The MARorg remained highly correlated to  
507 phytoplanktonic proxies during this period ( $r = 0.87$  between chlorophyll-a and MARorg). The  
508 negative correlation between N inputs and phytoplanktonic occurrence ( $r = 0.95$  between chlorophyll-a  
509 and N inputs and  $r = 0.97$  between MARorg and N inputs) highlights the effects of other driving  
510 factors and not only the trends in agricultural inputs. Past studies have demonstrated that in addition to  
511 human activities, climate warming or again secondary release of nutrients (Shayo & Limbu, 2018)  
512 may have a major impact on organic production and eutrophication phenomena.

513 Nowadays, although eutrophication represents a major threat for a large number of reservoirs  
514 across the world (Le *et al.*, 2010; Bhagowati & Ahamad, 2019), this process remains insufficiently  
515 studied in small headwater basins. The general positive trend of paleo production proxies observed in  
516 study could be improved by the collection of additional samples in various catchments of the Loire  
517 basin. Nevertheless, the results obtained in the current research emphasized the occurrence of a  
518 massive acceleration of organic matter deposition after 1950 in these reservoirs. Moreover, the  
519 sediment recorded in ponds draining contrasting land uses also showed a transition between a system  
520 dominated by terrigenous inputs to a system dominated by the autochthonous production during the

521 1950-2015 period. Although the paleo production proxies were highly correlated to the use of  
522 agricultural inputs until the 1990s, this relationship was no longer observed since then. Accordingly,  
523 questions remain to identify the factor(s) driving the recent autochthonous production in these  
524 reservoirs. Overall, soil erosion is classically considered to be the main factor inducing siltation of  
525 water bodies. As the excess of organic matter production can also lead to the filling of water bodies,  
526 the contribution of these processes and their respective contribution to reservoir siltation need to be  
527 further quantified. This will allow the design of effective control measures such as the systematic  
528 sowing of cover crops in winter to improve nutrient absorption by plants and better protect soils  
529 against erosion.

530

## 531 5. Conclusions

532

533 The collection and analysis of sediment accumulated in freshwater bodies provides a powerful  
534 technique for reconstructing changes in organic matter sources and for investigating their relationship  
535 with landscape management. Although these changes were studied during the 20<sup>th</sup> century in coastal  
536 areas, great lakes or during the last decades in riverine systems, there is currently a lack of records to  
537 support a regional analysis of these changes in continental areas over the last century. Results obtained  
538 in the current research underline an acceleration of organic matter deposition rates in cultivated areas  
539 during the second part of the 20<sup>th</sup> century. Between 1950 and 1965, organic matter that accumulated in  
540 water bodies draining agricultural catchments was mainly supplied by terrigenous sources. These  
541 deposits occurred during the implementation of major landscape management programs such as land  
542 consolidation, soil drainage or stream re-design. The increasing use of synthetic fertilizers which  
543 started in the 1970s induced a shift in organic matter accumulation with the acceleration of primary  
544 production in water bodies. This autochthonous production was highly correlated to the use of N  
545 fertilizers until the early 1990s. Nevertheless, the current research contributed to improve our  
546 understanding of long-term trends in eutrophication through the identification of a switch in  
547 production and the associated driving factors during the 1960-1970 period. Since then, although the

548 primary production continues to increase, this change cannot be exclusively explained by the enhanced  
549 use of fertilizers. The other factors driving this change should therefore be identified and quantified.  
550 This identification is of prime importance in order to reduce primary production in these water bodies  
551 and to improve the water quality of freshwater systems.

552

### 553 Acknowledgments

554 The authors are grateful to Anne Colmar, Xavier Bourrain and Jean-Noël Gautier for their  
555 technical support. This work was supported by a grant from the Loire-Brittany water agency  
556 (METEOR project). The authors would also like to thank Isabelle Pene-Galland, Jean-Paul Bakyono,  
557 Louis Maniere, Pierre Vanhooydonck, Thomas Grangeon, Anastasiia Bagaeva and Naresh Kumar for  
558 their help during the sediment core collection.

559

### 560 References

561

562 **AELB** (2013) Etat des lieux du bassin Loire-Bretagne établi en application de la directive  
563 cadre sur l'eau. 276 pp.

564 **Agnihotri, R., Kumar, R., Prasad, M.V.S.N., Sharma, C., Bhatia, S.K. and Arya, B.C.**  
565 (2014) Experimental Setup and Standardization of a Continuous Flow Stable Isotope  
566 Mass Spectrometer for Measuring Stable Isotopes of Carbon, Nitrogen and Sulfur in  
567 Environmental Samples. *Mapan - J Metrol Soc India*. doi: 10.1007/s12647-014-0099-8

568 **Ahn, Y.S., Nakamura, F. and Chun, K.W.** (2010) Recent history of sediment dynamics in  
569 Lake Toro and applicability of <sup>210</sup>Pb dating in a highly disturbed catchment in northern  
570 Japan. *Geomorphology*. doi: 10.1016/j.geomorph.2009.07.009

571 **Anderson, N.J., Bennion, H. and Lotter, A.F.** (2014) Lake eutrophication and its  
572 implications for organic carbon sequestration in Europe. *Glob Chang Biol*. doi:  
573 10.1111/gcb.12584

574 **Andre, M. and Polombo, N.** (2013) Soixante années de remembrement : essai de bilan  
575 critique de l'aménagement foncier en France.

576 **Appleby, P.G.** (2001) Tracking Environmental Change Using Lake Sediments. Volume 1:  
577 Basin Analysis, Coring, and Chronological Techniques. 171–201 pp.

578 **Appleby, P.G. and Oldfield, F.** (1978) The calculation of lead-210 dates assuming a constant  
579 rate of supply of unsupported <sup>210</sup>Pb to the sediment. *Catena*, **5**, 1–8.

- 580 **Arndt, S., Jørgensen, B.B., LaRowe, D.E., Middelburg, J.J., Pancost, R.D. and Regnier,**  
581 **P.** (2013) Quantifying the degradation of organic matter in marine sediments: A review  
582 and synthesis. *Earth-Science Rev.*, **123**, 53–86.
- 583 **Bennett, E.M., Carpenter, S.R. and Caraco, N.F.** (2001) Human Impact on Erodeable  
584 Phosphorus and Eutrophication: A Global Perspective Increasing accumulation of  
585 phosphorus in soil threatens rivers, lakes, and coastal oceans with. Source Biosci. doi:  
586 10.1641/0006-3568(2001)051[0227:hioepa]2.0.co;2
- 587 **Bhagowati, B. and Ahamad, K.U.** (2019) A review on lake eutrophication dynamics and  
588 recent developments in lake modeling. *Ecohydrol. Hydrobiol.*
- 589 **Boardman, J. and Poesen, J.** (2006) Soil Erosion in Europe.
- 590 **Bowsher, C., Steer, M. and Tobin, A.** (2008) Plant biochemistry, Garland Sc. New-York,  
591 446 pp.
- 592 **Brenner, M., Whitmore, T.J., Curtis, J.H., Hodell, D.A. and Schelske, C.L.** (1999) Stable  
593 isotope ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) signatures of sedimented organic matter as indicators of  
594 historic lake trophic state. *J Paleolimnol.* doi: 10.1023/A:1008078222806
- 595 **Buishand, T.A.** (1982) Some methods for testing the homogeneity of rainfall records. *J*  
596 *Hydrol.* doi: 10.1016/0022-1694(82)90066-X
- 597 **Cambray, R.S., Playford, K. and Carpenter, R.C.** (1989) Radioactive Fallout in Air and  
598 Rain: Results to the End of 1988.
- 599 **Cerdan, O., Govers, G., Le Bissonnais, Y., Van Oost, K., Poesen, J., Saby, N., Gobin, A.,**  
600 **Vacca, A., Quinton, J., Auerswald, K., Klik, A., Kwaad, F.J.P.M., Raclot, D., Ionita,**  
601 **I., Rejman, J., Rousseva, S., Muxart, T., Roxo, M.J. and Dostal, T.** (2010) Rates and  
602 spatial variations of soil erosion in Europe: A study based on erosion plot data.  
603 *Geomorphology.* doi: 10.1016/j.geomorph.2010.06.011
- 604 **Chartin, C., Evrard, O., Salvador-Blanes, S., Hirschberger, F., Van Oost, K., Lefèvre, I.,**  
605 **Daroussin, J. and Macaire, J.-J.** (2013) Quantifying and modelling the impact of land  
606 consolidation and field borders on soil redistribution in agricultural landscapes (1954–  
607 2009). *CATENA*, **110**, 184–195.
- 608 **Chen, D., Hu, M., Guo, Y. and Dahlgren, R.A.** (2016) Modeling forest/agricultural and  
609 residential nitrogen budgets and riverine export dynamics in catchments with contrasting  
610 anthropogenic impacts in eastern China between 1980–2010. *Agric. Ecosyst. Environ.*,  
611 **221**, 145–155.
- 612 **Coplen, T.B., Kendall, C. and Hopple, J.** (1983) Comparison of stable isotope reference  
613 samples. *Nature.* doi: 10.1038/302236a0
- 614 **Das, B., Vinebrooke, R.D., Sanchez-Azofeifa, A., Rivard, B. and Wolfe, A.P.** (2005)  
615 Inferring sedimentary chlorophyll concentrations with reflectance spectroscopy: a novel  
616 approach to reconstructing historical changes in the trophic status of mountain lakes. *Can*  
617 *J Fish Aquat Sci.* doi: 10.1139/f05-016
- 618 **Dearing, J.A. and Jones, R.T.** (2003) Coupling temporal and spatial dimensions of global  
619 sediment flux through lake and marine sediment records. *Glob. Planet. Change*, **39**, 147–  
620 168.
- 621 **Debret, M., Sebag, D., Desmet, M., Balsam, W., Copard, Y., Mourier, B., Susperrigui,**



- 622 **A.S., Arnaud, F., Bentaleb, I., Chapron, E., Lallier-Vergès, E. and Winiarski, T.**  
 623 (2011) Spectrocolorimetric interpretation of sedimentary dynamics: The new “Q7/4  
 624 diagram.” *Earth-Science Rev.*
- 625 **Directive, 91/271/EEC Council** (1991a) 91/271/EEC Council Directive 91/271/EEC of 21  
 626 May 1991 concerning urban waste-water treatment.
- 627 **Directive, 91/676/EEC Council** (1991b) 91/676/EEC Council Directive 91/676/EEC of 12  
 628 December 1991 concerning the protection of waters against pollution caused by nitrates  
 629 from agricultural sources.
- 630 **Downing, J.A., Cole, J.J., Middelburg, J.J., Striegl, R.G., Duarte, C.M., Kortelainen, P.,**  
 631 **Prairie, Y.T. and Laube, K.A.** (2008) Sediment organic carbon burial in agriculturally  
 632 eutrophic impoundments over the last century. *Global Biogeochem Cycles*. doi:  
 633 10.1029/2006GB002854
- 634 **Emerson, S., Fischer, K., Reimers, C. and Heggie, D.** (1985) Organic carbon dynamics and  
 635 preservation in deep-sea sediments. *Deep Sea Res. Part A. Oceanogr. Res. Pap.*, **32**, 1–  
 636 21.
- 637 **Etcheber, H., Taillez, A., Abril, G., Garnier, J., Servais, P., Moatar, F. and Commarieu,**  
 638 **M.V.** (2007) Particulate organic carbon in the estuarine turbidity maxima of the Gironde,  
 639 Loire and Seine estuaries: Origin and lability. In: *Hydrobiologia*,
- 640 **European Environment Agency** (2018) European waters: Assessment of status and  
 641 pressures 2018.
- 642 **Evrard, O., Lacey, J.P., Onda, Y., Wakiyama, Y., Jaegler, H. and Lefèvre, I.** (2016)  
 643 Quantifying the dilution of the radiocesium contamination in Fukushima coastal river  
 644 sediment (2011–2015). *Sci. Rep.*, **6**, 34828.
- 645 **Evrard, O., Van Beek, P., Gateuille, D., Pont, V., Lefèvre, I., Lansard, B. and Bonté, P.**  
 646 (2012) Evidence of the radioactive fallout in France due to the Fukushima nuclear  
 647 accident. *J Environ Radioact*. doi: 10.1016/j.jenvrad.2012.01.024
- 648 **Florian, C.R., Miller, G.H., Fogel, M.L., Wolfe, A.P., Vinebrooke, R.D. and Geirsdóttir,**  
 649 **Á.** (2015) Algal pigments in Arctic lake sediments record biogeochemical changes due to  
 650 Holocene climate variability and anthropogenic global change. *J Paleolimnol*. doi:  
 651 10.1007/s10933-015-9835-5
- 652 **Foster, I.D.L., Collins, A.L., Naden, P.S., Sear, D.A., Jones, J.I. and Zhang, Y.** (2011) The  
 653 potential for paleolimnology to determine historic sediment delivery to rivers. *J*  
 654 *Paleolimnol*. doi: 10.1007/s10933-011-9498-9
- 655 **Foucher, A., Evrard, O., Cerdan, O., Chabert, C., Lecompte, F., Lefèvre, I.,**  
 656 **Vandromme, R. and Salvador- Blanes, S.** (2019a) A quick and low- cost technique to  
 657 identify layers associated with heavy rainfall in sediment archives during the  
 658 Anthropocene. *Sedimentology*. doi: 10.1111/sed.12650
- 659 **Foucher, A., Evrard, O., Chabert, C., Cerdan, O., Lefèvre, I., Vandromme, R. and**  
 660 **Salvador-Blanes, S.** (2019b) Erosional response to land abandonment in rural areas of  
 661 Western Europe during the Anthropocene: A case study in the Massif-Central, France.  
 662 *Agric. Ecosyst. Environ.*, **284**, 106582.
- 663 **Foucher, A., Salvador-Blanes, S., Evrard, O., Simonneau, A., Chapron, E., Courp, T.,**  
 664 **Cerdan, O., Lefèvre, I., Adriaensen, H., Lecompte, F. and Desmet, M.** (2014)

- 665 Increase in soil erosion after agricultural intensification: Evidence from a lowland basin  
666 in France. *Anthropocene*, **7**, 30–41.
- 667 **Gälman, V., Rydberg, J. and Bigler, C.** (2009) Decadal diagenetic effects on  $\delta^{13}\text{C}$  and  
668  $\delta^{15}\text{N}$  studied in varved lake sediment. *Limnol Oceanogr.* doi: 10.4319/lo.2009.54.3.0917
- 669 **Gellis, A.C., Webb, R.M.T., McIntyre, S.C. and Wolfe, W.J.** (2006) Land-use effects on  
670 erosion, sediment yields, and reservoir sedimentation: A case study in the Lago Loíza  
671 Basin, Puerto Rico. *Phys Geogr.* doi: 10.2747/0272-3646.27.1.39
- 672 **Glooschenko, W.A., Moore, J.E. and Vollenweider, R.A.** (1974) Spatial and Temporal  
673 Distribution of Chlorophyll a and Pheopigments in Surface Waters of Lake Erie . *J Fish*  
674 *Res Board Canada.* doi: 10.1139/f74-046
- 675 **Grangeon, T., Manière, L., Foucher, A., Vandromme, R., Cerdan, O., Evrard, O., Pene-**  
676 **Galland, I. and Salvador-Blanes, S.** (2017) Hydro-sedimentary dynamics of a drained  
677 agricultural headwater catchment: A nested monitoring approach. *Vadose Zo J.* doi:  
678 10.2136/vzj2017.05.0113
- 679 **Gudasz, C., Ruppenthal, M., Kalbitz, K., Cerli, C., Fiedler, S., Oelmann, Y., Andersson,**  
680 **A. and Karlsson, J.** (2017) Contributions of terrestrial organic carbon to northern lake  
681 sediments. *Limnol Oceanogr Lett.* doi: 10.1002/lo2.10051
- 682 **Guilizzoni, P., Marchetto, A., Lami, A., Cameron, N.G., Appleby, P.G., Rose, N.L.,**  
683 **Schnell, A., Belis, C.A., Giorgis, A. and Guzzi, L.** (1996) The environmental history of  
684 a mountain lake (Lago Paione Superiore, Central Alps, Italy) for the last c. 100 years: A  
685 multidisciplinary, palaeolimnological study. *J Paleolimnol.* doi: 10.1007/BF00213044
- 686 **Hammer, Ø., Harper, D.A.T. and Ryan, P.D.** (2001) PAST: Paleontological Statistics  
687 Software Package for Education and Data Analysis. *Palaeontol. Electron.*, 4–9.
- 688 **Heathcote, A.J., Filstrup, C.T. and Downing, J.A.** (2013) Watershed Sediment Losses to  
689 Lakes Accelerating Despite Agricultural Soil Conservation Efforts. *PLoS One*, **8**,  
690 e53554.
- 691 **Hecky, R.E. and Kilham, P.** (1988) Nutrient limitation of phytoplankton in freshwater and  
692 marine environments: A review of recent evidence on the effects of enrichment. *Limnol*  
693 *Oceanogr.* doi: 10.4319/lo.1988.33.4part2.0796
- 694 **Hodgson, D.A., Wright, S.W. and Davies, N.** (1997) Mass Spectrometry and reverse phase  
695 HPLC techniques for the identification of degraded fossil pigments in lake sediments and  
696 their application in palaeolimnology. *J Paleolimnol.* doi: 10.1023/A:1007943119392
- 697 **Huon, S., Grousset, F.E., Burdloff, D., Bardoux, G. and Mariotti, A.** (2002) Sources of  
698 fine-sized organic matter in North Atlantic Heinrich Layers:  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  tracers.  
699 *Geochim Cosmochim Acta.* doi: 10.1016/S0016-7037(01)00776-1
- 700 **Huon, S., Hayashi, S., Lacey, J.P., Tsuji, H., Onda, Y. and Evrard, O.** (2018) Source  
701 dynamics of radiocesium-contaminated particulate matter deposited in an agricultural  
702 water reservoir after the Fukushima nuclear accident. *Sci. Total Environ.*, **612**, 1079–  
703 1090.
- 704 **Jinglu, W., Chengmin, H., Haihao, Z., Schleser, G.H. and Battarbee, R.** (2007)  
705 Sedimentary evidence for recent eutrophication in the northern basin of Lake Taihu,  
706 China: Human impacts on a large shallow lake. *J Paleolimnol.* doi: 10.1007/s10933-006-  
707 9058-x

- 708 **Korosi, J.B. and Smol, J.P.** (2012) Examining the effects of climate change, acidic  
709 deposition, and copper sulphate poisoning on long-term changes in cladoceran  
710 assemblages. *Aquat Sci.* doi: 10.1007/s00027-012-0261-8
- 711 **Le, C., Zha, Y., Li, Y., Sun, D., Lu, H. and Yin, B.** (2010) Eutrophication of lake waters in  
712 China: Cost, causes, and control. *Environ. Manage.*
- 713 **Le Moal, M., Gascuel-Oudou, C., Ménesguen, A., Souchon, Y., Étrillard, C., Levain, A.,  
714 Moatar, F., Pannard, A., Souchu, P., Lefebvre, A. and Pinay, G.** (2019)  
715 Eutrophication: A new wine in an old bottle? *Sci. Total Environ.*, **651**, 1–11.
- 716 **Leavitt, P. and Hodgson, D.A.** (2001) Tracking Environmental Change Using Lake  
717 Sediments. Volume 3: Terrestrial, Algal and Siliceous Indicators.
- 718 **Lu, C. and Tian, H.** (2017) Global nitrogen and phosphorus fertilizer use for agriculture  
719 production in the past half century: Shifted hot spots and nutrient imbalance. *Earth Syst*  
720 *Sci Data.* doi: 10.5194/essd-9-181-2017
- 721 **Lu, Y., Meyers, P.A., Johengen, T.H., Eadie, B.J., Robbins, J.A. and Han, H.** (2010)  $\delta^{15}$   
722 N values in Lake Erie sediments as indicators of nitrogen biogeochemical dynamics  
723 during cultural eutrophication. *Chem Geol.* doi: 10.1016/j.chemgeo.2010.02.002
- 724 **Meybeck, M., Cauwet, G., Dessery, S., Somville, M., Gouleau, D. and Billen, G.** (1988)  
725 Nutrients (organic C, P, N, Si) in the eutrophic River Loire (France) and its estuary.  
726 *Estuar Coast Shelf Sci.* doi: 10.1016/0272-7714(88)90071-6
- 727 **Meyers, P.A.** (1994) Preservation of elemental and isotopic source identification of  
728 sedimentary organic matter. *Chem Geol.* doi: 10.1016/0009-2541(94)90059-0
- 729 **Meyers, P.A. and Ishiwatari, R.** (1995) Organic Matter Accumulation Records in Lake  
730 Sediments. In: *Physics and Chemistry of Lakes*,
- 731 **Michelutti, N., Blais, J.M., Cumming, B.F., Paterson, A.M., Rühland, K., Wolfe, A.P.  
732 and Smol, J.P.** (2010) Do spectrally inferred determinations of chlorophyll a reflect  
733 trends in lake trophic status? *J Paleolimnol.* doi: 10.1007/s10933-009-9325-8
- 734 **Michelutti, N. and Smol, J.P.** (2016) Visible spectroscopy reliably tracks trends in paleo-  
735 production. *J Paleolimnol.* doi: 10.1007/s10933-016-9921-3
- 736 **Millie, D.F., Paerl, H.W. and Hurley, J.P.** (1993) Microalgal Pigment Assessments Using  
737 High-Performance Liquid Chromatography: A Synopsis of Organismal and Ecological  
738 Applications. *Can J Fish Aquat Sci.* doi: 10.1139/f93-275
- 739 **Minaudo, C., Meybeck, M., Moatar, F., Gassama, N. and Curie, F.** (2015) Eutrophication  
740 mitigation in rivers: 30 Years of trends in spatial and seasonal patterns of  
741 biogeochemistry of the Loire River (1980-2012). *Biogeosciences.* doi: 10.5194/bg-12-  
742 2549-2015
- 743 **NF ISO 10693** (1995) Qualité du sol - Détermination de la teneur en carbonate - Méthode  
744 volumétrique.
- 745 **Olley, J., Burton, J., Smolders, K., Pantus, F. and Pietsch, T.** (2013) The application of  
746 fallout radionuclides to determine the dominant erosion process in water supply  
747 catchments of subtropical South-east Queensland, Australia. *Hydrol Process.* doi:  
748 10.1002/hyp.9422
- 749 **Poisvert, C.** (2018) Analyse et modélisation des surplus azotés en France au cours du siècle

- 750           dernier: Application aux échelles départementales et communales. Université de Tours
- 751 **Poisvert, C., Curie, F. and Moatar, F.** (2017) Annual agricultural N surplus in France over a  
752           70-year period. *Nutr Cycl Agroecosystems*. doi: 10.1007/s10705-016-9814-x
- 753 **Rabalais, N.N., Turner, R.E., Gupta, B.K.S., Platon, E. and Parsons, M.L.** (2007)  
754           Sediments tell the history of eutrophication and hypoxia in the northern Gulf of Mexico.  
755           In: *Ecological Applications*,
- 756 **Rein, B. and Sirocko, F.** (2002) In-situ reflectance spectroscopy - Analysing techniques for  
757           high-resolution pigment logging in sediment cores. *Int J Earth Sci*. doi: 10.1007/s00531-  
758           002-0264-0
- 759 **Routh, J., Meyers, P.A., Gustafsson, Ö., Baskaran, M., Hallberg, R. and Schöldström, A.**  
760           (2004) Sedimentary geochemical record of human-induced environmental changes in the  
761           Lake Brunnsviken watershed, Sweden. *Limnol Oceanogr*. doi:  
762           10.4319/lo.2004.49.5.1560
- 763 **Rowan, J.S., Black, S. and Franks, S.W.** (2012) Sediment fingerprinting as an  
764           environmental forensics tool explaining cyanobacteria blooms in lakes. *Appl Geogr*. doi:  
765           10.1016/j.apgeog.2011.07.004
- 766 **Schindler Wildhaber, Y., Liechti, R. and Alewell, C.** (2012) Organic matter dynamics and  
767           stable isotope signature as tracers of the sources of suspended sediment. *Biogeosciences*.  
768           doi: 10.5194/bg-9-1985-2012
- 769 **Schlesinger, W.H. and Bernhardt, E.S.** (2013) *Biogeochemistry An Analysis : of Global*  
770           Change. *Elsevier*.
- 771 **Shayo, S. and Limbu, S.M.** (2018) Nutrient release from sediments and biological nitrogen  
772           fixation: Advancing our understanding of eutrophication sources in Lake Victoria,  
773           Tanzania. *Lakes Reserv Res Manag*. doi: 10.1111/lre.12242
- 774 **Stewart, E.M., Michelutti, N., Shenstone-Harris, S., Grooms, C., Weseloh, C., Kimpe,**  
775           **L.E., Blais, J.M. and Smol, J.P.** (2015) Tracking the history and ecological changes of  
776           rising double-crested cormorant populations using pond sediments from islands in  
777           Eastern Lake Ontario. *PLoS One*
- 778 **Tilman, D., Fargione, J., Wolff, B., D'Antonio, C., Dobson, A., Howarth, R., Schindler,**  
779           **D., Schlesinger, W.H., Simberloff, D. and Swackhamer, D.** (2001) Forecasting  
780           agriculturally driven global environmental change. *Science* (80- ). doi:  
781           10.1126/science.1057544
- 782 **Vandromme, R., Foucher, A., Cerdan, O. and Salvador-Blanes, S.** Quantification of bank  
783           erosion of artificial drainage networks using LIDAR data.
- 784 **Voß, M. and Struck, U.** (1997) Stable nitrogen and carbon isotopes as indicator of  
785           eutrophication of the Oder river (Baltic sea). *Mar. Chem.*, **59**, 35–49.
- 786 **Warren, J. and Gilbert, R.O.** (1988) *Statistical Methods for Environmental Pollution*  
787           Monitoring. *Technometrics*. doi: 10.2307/1270090
- 788 **Waters, M.N., Schelske, C.L., Kenney, W.F. and Chapman, A.D.** (2005) The use of  
789           sedimentary algal pigments to infer historic algal communities in Lake Apopka, Florida.  
790           *J Paleolimnol*. doi: 10.1007/s10933-004-1691-7
- 791 **Wolfe, A.P., Vinebrooke, R.D., Michelutti, N., Rivard, B. and Das, B.** (2006) Experimental

792 calibration of lake-sediment spectral reflectance to chlorophyll a concentrations:  
793 Methodology and paleolimnological validation. J Paleolimnol. doi: 10.1007/s10933-006-  
794 0006-6

795

796

797

798

799

800

801

802

803

804

805

806 Figures:

807

808 Fig.1 Localization of the nine studied catchments within the Loire river basin. (1) Le Loroux-  
809 Bottereau, (2) Beaurepaire, (3) Passavant-sur-Layon, (4) Brosse, (5) Boisvinet, (6) Prugnyolas,  
810 (7) Goule, (8) Tazenat, (9) Malaguet

811

812 Fig. 2 Evolution of sediment properties throughout time in reservoirs draining forested  
813 catchments: (6) Prugnyolas and (9) Malaguet. Grey lines correspond to the identification of  
814  $^{137}\text{Cs}$  peaks associated to the 1963 and 1986 fallouts.

815

816 Fig. 3 Evolution of sediment properties throughout time in reservoirs draining catchments  
817 dominated by grassland: (1) Le Loroux-Bottereau, (2) Beaurepaire, (7) Goule and (8) Tazenat.  
818 Grey lines correspond to the identification of  $^{137}\text{Cs}$  peaks associated to the 1963 and 1986  
819 fallouts.

820

821 Fig. 4 Evolution of sediment properties throughout time in reservoirs draining catchments  
822 dominated by arable land: (3) Passavant-sur-Layon, (4) Brosse and (5) Boisvinet. Grey line  
823 correspond to the identification of  $^{137}\text{Cs}$  peak associated to the 1963 fallouts.

824

825 Fig. 5 Evolution of Nitrogen consumption in the nine selected catchments under arable land,  
826 grassland or dominated by forest (1) Le Loroux-Bottereau, (2) Beaurepaire, (3) Passavant-sur-  
827 Layon, (4) Brosse, (5) Boisvinet, (6) Prugnolas, (7) Goule, (8) Tazenat, (9) Malaguet

828

829 Fig. 6 Evolution of Mass Accumulation Rates of organic material (MARorg) during the last  
830 century in reservoirs located at the outlet of the nine studied catchments: (1) Le Loroux-  
831 Bottereau, (2) Beaurepaire, (3) Passavant-sur-Layon, (4) Brosse, (5) Boisvinet, (6) Prugnolas,  
832 (7) Goule, (8) Tazenat, (9) Malaguet. Grey area correspond to a significant acceleration of  
833 MARorg.

834

835 Fig. 7 Evolution of sediment properties throughout time in reservoirs draining the seven  
836 cultivated catchments. The light grey part corresponds to the period dominated by terrigenous  
837 inputs whereas the dark grey colour corresponds to the period dominated by autochthonous  
838 primary productivity.

839

840

841 Tables:

842

843 Table 1. Characteristics of the Loire Basin sub-catchments

844

845 Table 2. Characteristics of the studied catchments

846

847 Table 3. Chronology of sediment cores and modern and historic Mass Accumulation Rates of  
848 organic matter (MAR<sub>org</sub> expressed in  $\text{gC m}^{-2} \text{yr}^{-1}$ : modern MAR<sub>org</sub> correspond to average  
849 value of MAR<sub>org</sub> for the 2017-1950 period and the historical MAR<sub>org</sub> to the average value  
850 before 1950). ND = non detected.

851

852 Table 4. Results of the Mann Kendal test with the significant  $p$  value  $<0.05$ . NT = No trend, +  
853 = positive trend and - = negative trend

854

855

	Sub catchment area (km <sup>2</sup> )	Average elevation (m)	Average slope (%)	Land use
Lower Loire	36,780	108	1.7	Arable land (60%), Grassland (24%), Forest (11%), Urban (5%)
Middle Loire	48,471	216	2.1	Arable land (48%), Forest (25%), Grassland (24%), Urban (4%)
Upper Loire	32,630	570	5.5	Grassland (43%), Forest (35%), Arable land (17%), Urban (4%)



Tables

[Click here to download Tables: Table 2 -review 2.docx](#)

Site ID	Reservoir name	Core location	Core length (cm)	Water depth (m)	Lake/Pond area (ha)	Catchment area (km <sup>2</sup> )	Average elevation (m)	Average slope (%)	Catchment location	Main land use	Land use	Landscape management
1	Loroux-Botterau	47.2379N, -1.3440E	71	1.2	1.7	4.5	68	2.7	Lower Loire	Grassland (65%)	Arable land (34%). urban area (1%)	Hedge removal during the 1965-1975 period
2	Beurepaire	47.0726N, -0.4737E	108	2.5	38	14.3	122	1.7	Lower Loire	Grassland (52%)	Arable land (45%). water bodies (3%). woodland (0.1%)	tile drain implementation and land consolidation in 1977
3	Passavant-sur-Layon	47.1013N, -0.3941E	116	1.4	14	108	121	2.4	Lower Loire	Arable land (70.5%)	Grassland (21%). woodland (6.36%). water bodies (0.9%). mining (0.6%). urban areas (0.3%)	Major land consolidation during the 1960's
4	Brosse	47.2761N, 1.0558E	127	2.4	11	24.5	105	2.3	Middle Loire	Arable land (83%)	Woodland (8%). grassland (7%). water bodies (1%). urban areas (1%)	Major land consolidation in 1959
5	Boisvinet	48.0733N, 0.8832E	97	2.1	39	9.5	189	2.5	Lower Loire	Arable land (63%)	Grassland (23%). woodland (9.5%). water bodies (4.3%)	
6	Prugnotas	45.8689N, 1.9032E	82	0.8	1.8	7.8	727	12	Middle Loire	Woodland (82%)	Grassland (16%). arable land 0.4%)	Afforestation after 1950, major windstorm in 1999 inducing damage on forest
7	Goule	46.7277N, 2.8004E	152	4.3	52	34	242	3	Middle Loire	Grassland (65.9%)	Woodland (18.4%). arable land (13.3%). water bodies (1.5%). urban areas (0.8%)	
8	Tazenat	45.9796N, 2.9900E	115	67	30	2.3	690	9.3	Upper Loire	Grassland (68%)	Woodland (16%). water bodies (15%)	Land consolidation in 1990
9	Malaguet	45.2512N, 3.7103E	69	3.1	20.5	3.7	1073	7.5	Upper Loire	Woodland (71%)	Grassland (13%). water bodies (7%). urban areas (6%). arable land (2%)	Afforestation works between 1950 and 1965

Site ID	Max $^{210}\text{Pb}_{\text{ex}}$ date / depth	1986 $^{137}\text{Cs}$ fallout	1963 $^{137}\text{Cs}$ fallout	Modern MARorg	Historic MARorg
1	1921 (71 cm)	ND	± 3 year (40 cm)	133 ± 32	46 ± 21
2	1925 (105 cm)	ND	± 2 year (70 cm)	219 ± 71	76 ± 12
3	<100 (74 cm)	ND	± 2 year (46 cm)	229 ± 50	133 ± 11
4	1922 (130 cm)	ND	± 2 year (92 cm)	460 ± 117	342 ± 126
5	1898 (90 cm)	ND	± 3 year (50 cm)	111 ± 28	76 ± 2
6	1918 (140 cm)	± 1 year (55 cm)	± 1 year (85 cm)	167 ± 12	114 ± 25
7	1901 (65 cm)	± 1 year (11.5 cm)	ND	141 ± 46	450 ± 330
8	1902 (44 cm)	± 1 year (11 cm)	± 3 year (19 cm)	41 ± 11	21 ± 12
9	1948 (76 cm)	± 4 year (16 cm)	ND	163 ± 73	-

Tables

[Click here to download Tables: Table 4 - review 2.docx](#)

Site ID	TOC		$\delta^{13}\text{C}$		$\delta^{15}\text{N}$		C/N		TN		chlorophyll-a		MARorg		N Inputs	
1	0.46	NT	<0.05	-	<0.05	-	<0.05	-	0.46	NT	<0.05	+	<0.05	+	<0.05	+
2	<0.05	+	<0.05	-	<0.05	+	0.3	NT	<0.05	+	<0.05	+	<0.05	+	<0.05	+
3	<0.05	+	<0.05	-	<0.05	+	0.27	NT	<0.05	-	<0.05	+	0.53	NT	<0.05	+
4	0.84	NT	0.82	NT	0.07	NT	0.11	NT	0.96	NT	<0.05	+	0.33	NT	<0.05	+
5	<0.05	+	0.6	NT	<0.05	+	0.46	NT	<0.05	+	<0.05	+	<0.05	+	0.43	NT
6	<0.05	+	<0.05	-	0.19	NT	<0.05	-	<0.05	+	<0.05	+	<0.05	+	<0.05	+
7	0.138	NT	0.408	NT	<0.05	-	0.5	NT	<0.05	+	0.06	NT	<0.05	-	0.38	NT
8	<0.05	+	<0.05	-	0.54	NT	0.46	NT	<0.05	+	0.38	NT	<0.05	+	<0.05	+
9	0.13	NT	<0.05	-	0.45	NT	0.23	NT	<0.05	+	<0.05	+	<0.05	-	<0.05	-

Figure 1  
[Click here to download high resolution image](#)




Figure 2  
[Click here to download high resolution image](#)




Figure 3  
[Click here to download high resolution image](#)




Figure 4  
[Click here to download high resolution image](#)




Figure 5  
[Click here to download high resolution image](#)






Figure 6  
[Click here to download high resolution image](#)




Figure 7  
[Click here to download high resolution image](#)

