



HAL
open science

Multidrug-Resistant Infections in U.S. Hospitals

Rémi Le Guern, Caroline Loiez, Frederic Wallet

► **To cite this version:**

Rémi Le Guern, Caroline Loiez, Frederic Wallet. Multidrug-Resistant Infections in U.S. Hospitals. New England Journal of Medicine, 2020, 383 (1), pp.93-94. 10.1056/NEJMc2014837 . hal-02888399

HAL Id: hal-02888399

<https://hal.science/hal-02888399>

Submitted on 3 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title:** Multidrug-Resistant Infections in U.S. Hospitals

2 **Author names and affiliations:** Rémi Le Guern, Caroline Loïez, Frédéric Wallet

3 Centre Hospitalier Régional Universitaire de Lille

4 Lille, France

5 **Corresponding author:** Rémi Le Guern

6 Email address: remi.leguern@chru-lille.fr

7

8 Rémi Le Guern, M.D., Ph.D.

9 CHU Lille, Lille, France

10 remi.leguern@chru-lille.fr

11

12 Caroline Loïez, M.D.

13 CHU Lille, Lille, France

14

15 Frédéric Wallet, M.D.

16 CHU Lille, Lille, France

17 Jernigan et al. reported national estimates of multidrug-resistant bacterial infections in the
18 U.S.¹, but extended-spectrum- β -lactamase-producing *Enterobacterales* (ESBL-E) were
19 underestimated. Indeed, only *Klebsiella spp.* (excepted *Klebsiella aerogenes*) and
20 *Escherichia coli* isolates were included in the pathogen phenotype definition (see Table
21 S2 in the Supplementary Appendix of the article, available at NEJM.org).

22 From January 2008 to March 2020, we conducted a retrospective study in our center on
23 ESBL-E carriage using rectal swabs culture. A total of 128,374 samples from 42,587
24 patients were included. Each ESBL-E isolate was counted only once per patient. Out of
25 9,945 isolates from 8,099 patients, *Klebsiella spp.* (excepted *K. aerogenes*) and *E. coli*
26 represented 78.11% of all the ESBL-E isolates, *Enterobacter* (and *K. aerogenes*) 16.38%,
27 *Citrobacter* 3.96%, and other *Enterobacterales* 1.55%.

28 Because *Enterobacter spp.*, *K. aerogenes*, and *C. freundii* naturally produce an AmpC
29 cephalosporinase², ESBL production cannot be inferred only from the antimicrobial
30 susceptibility results. Additional phenotypic confirmatory tests are needed³, which
31 explains the underestimation in this study based on antimicrobial susceptibility data.

32 **Bibliography**

- 33 1. Jernigan JA, Hatfield KM, Wolford H, et al. Multidrug-Resistant Bacterial Infections
34 in U.S. Hospitalized Patients, 2012-2017. *N Engl J Med* 2020;382:1309-19.
- 35 2. Bush K, Bradford PA. Epidemiology of beta-Lactamase-Producing Pathogens. *Clin*
36 *Microbiol Rev* 2020;33.
- 37 3. Poulou A, Grivakou E, Vrioni G, et al. Modified CLSI extended-spectrum beta-
38 lactamase (ESBL) confirmatory test for phenotypic detection of ESBLs among
39 Enterobacteriaceae producing various beta-lactamases. *J Clin Microbiol* 2014;52:1483-
40 9.