

The Role of Genetic Sex and Mitochondria in Response to COVID-19 Infection

Malgorzata Kloc, Rafik Mark Ghobrial, Jacek Z. Kubiak

▶ To cite this version:

Malgorzata Kloc, Rafik Mark Ghobrial, Jacek Z. Kubiak. The Role of Genetic Sex and Mitochondria in Response to COVID-19 Infection. International Archives of Allergy and Immunology, 2020, 181 (8), pp.629-634. 10.1159/000508560 . hal-02888603

HAL Id: hal-02888603 https://hal.science/hal-02888603

Submitted on 10 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mini review

The role of genetic sex and mitochondria in response to COVID-19 infection

Malgorzata Kloc*1,2,3, Rafik M. Ghobrial^{1,2}, Jacek Z Kubiak*4,5

¹The Houston Methodist Research Institute, Houston, Texas 77030, USA
²The Houston Methodist Hospital, Department of Surgery, Houston, Texas, USA
³The University of Texas, M.D. Anderson Cancer Center, Department of Genetics, Houston Texas, USA
⁴Laboratory of Regenerative Medicine and Cell Biology, Military Institute of Hygiene and Epidemiology (WIHE), Warsaw, Poland
⁵UnivRennes, UMR 6290, CNRS, Institute of Genetics and Development of Rennes, Cell Cycle Group,

Faculty of Medicine, Rennes, France

Corresponding authors:

Malgorzata Kloc, The Houston Methodist Research Institute, 6670 Bertner Ave, Houston, TX 77030, USA; email: mkloc@houstonmethodist.org

and

Jacek Z Kubiak, UMR 6290, CNRS/UR1, IGDR, Faculty of Medicine, 2 Ave. du Prof. Leon Bernard, 35043 Rennes cedex, France; e-mail: jacek.kubiak@univ-rennes1.fr

Keywords: COVID-19, SARS-CoV-2, macrophages, Y chromosome, X chromosome, immunity, melatonin

Abstract

The difference between the female and male immune response to COVID-19 infection, and infections in general, is multifactorial. The well-known determiners of the immune response such as X and Y chromosomes, sex hormones and microbiota, are functionally interconnected and influence each other in shaping the organism's immunity. We focus our commentary on the interplay between the genetic sex and mitochondria, and how this may affect a sex-dependent immune response in COVID-19 infection.

Realizing the existence of these interactions may help in designing novel methods or fine-tuning the existing and routine therapies to fight COVID-19 and other infections.

Introduction

Research and clinical studies indicate that the genetic sex of patients influences the immune response, and the course, and prognosis of many common noninfectious and infectious diseases. A similar correlation has also been observed recently in COVID-19 infected patients; the world-wide statistical analyses of COVID-19 cases indicate that the rate of infection and the direness of the outcome are much higher and sever in males than in females. The most popular explanation for this phenomenon is "the immunocompetence handicap model" proposing that in males, the testosterone-dependent secondary male sex traits develop at the expense of the immune response. Sex hormones are known to affect the innate and adaptive immunological response with the androgens being anti-proinflammatory, and the estrogens pro-inflammatory [1, 2]. However, the mechanisms of these sex-related differences are multifactorial and depend on very complex reciprocal interactions between sex chromosome-encoded and regulatory factors, hormones, and microbiota inhabiting the human body. In the last decade, it became clear that the microorganisms inhabiting the human gastrointestinal tract differ, in the type and abundance of species, between the sexes. This sex-dependent microbiome, called the microgenderome develops after puberty when the sex hormones kick in, and regulates local and systemic inflammation, and response to infection (Fig. 1); [1-3]. Recent studies also indicate that the sex-related difference in the immune response may depend on the sexually dimorphic populations of mitochondria. Below we give a short summary of how these different elements pitch in and shape the immune response to COVID-19 and other diseases.

X and Y chromosome regulation of immunity

Although some differences between the female and male immune response are directly related to the number of X chromosomes (two X in females versus one X in males), and the effect of sex hormones on the functions of immune cells, the studies of last decade indicate that both Y and X chromosomes regulate immune cell transcription and translation (transcriptome and proteome) profiles.

X chromosome

The Archive.ensabl.org 2017:

http://mar2017.archive.ensembl.org/Homo_sapiens/Location/Chromosome?r=X [4]

indicate that the human chromosome X consists of over 150 million DNA base pairs and contains more than 800 protein-coding genes, which include the highest number of the innate and adaptive immunity-related genes of the whole human genome, and several hundred non-coding genes. The immune response genes include fraktaline receptor CXCR3 (C–X–C motif chemokine receptor 3) that directs immune cells,

including macrophages, movement into inflamed tissues and organs; the gene for CD80 ligand that binds to the CD40 receptor on antigen-presenting cells, B cells, monocytes/macrophages and dendritic cells, and through interaction with CD154 transduces signals for T-dependent B-cell activation; and the Interleuk in-1 receptor-associated kinase 1 (IRAK1) gene. The X chromosome also contains 10 % of all genomic microRNAs (miRNAs) genes that regulate expression and degradation of autosomal gene products [5, 6]. Studies showed that the absence of, or extensive deletions in one of the X chromosomes lead to a variety of autoimmune diseases) (Fig. 2); [5, 6].

A higher rate of COVID-19 infection in males may be, in part, influenced by the X chromosome mosaicism in the female cells, due to the stochastic X chromosome inactivation [7]. For example, the angiotensin-converting enzyme 2, (ACE2), the receptor for β -type coronaviruses SARS-CoV-1 and SARS-CoV-2, is encoded by the ACE2 gene located on the X chromosome. It is possible that some alleles of this gene may code for the receptors with different efficiency of recognizing and binding the virus. Male cells always express a single ACE2 allele because all cells contain an identical X chromosome. In contrast, a female has a heterogenic ACE2 allele because of the mosaicism of X chromosomes, which are stochastically distributed between the cells. Thus, in the female, a potentially more efficient form of ACE2 receptor would be present in only half of all cells. This may limit infectability with SARS-CoV-1 and -2 viruses and, to some extent, give females a relative resistance to the infection (Fig. 2). It has also been debated if such ACE2 polymorphism may have an impact on hypertension and other diseases [8, 9].

Y chromosome

The human Y chromosome is much smaller than the X chromosome, consist of around 60 million DNA base pairs, and, in the majority of animals, its large portion is transcriptionally silent (heterochromatic). The Y chromosome is mainly composed of transposable elements, repeat sequences (comprising over 50% of its content), multicopy genes such as the polymorphic variants of the ribosomal genes, and a low number of protein-encoding genes, which mainly are male-specific and regulate sexual development, spermatogenesis, sex ratio, and fertility, and often evolved through the transposition/translocation from the autosomes. This male-specific region of the Y chromosome (MSY) does not recombine and thus remains unchanged between the consecutive male generations [10-13]. Studies showed that Y chromosome, can regulate the heterochromatin/euchromatin status of the autosomes and X chromosomes and thus affects silencing/expression of various genes, including the immune response genes, and regulate the tissue/cell-specific alternative gene splicing [10, 14-18]. Although the mechanism by which the Y chromosome influences heterochromatin/euchromatin content and thus the transcription of the other genes

is largely unknown, one of the hypotheses postulates that the Y chromosome sequesters heterochromatinization factors and another that it affects the architecture of cell nucleus, which makes specific genes inaccessible to the transcription factors [19-22]. Phylogenetic studies showed that the males with the haplogroup I, which is one of the most popular European lineages of the Y chromosome, have upregulated inflammatory response, downregulated adaptive immunity, and a higher risk of coronary diseases [23-25]. In the murine models, Y chromosomes influence the number of natural killer T cells, the gene expression pattern in CD4⁺ T cells, the immune response of macrophages, and mortality rate following the infection with coxsackievirus [26-28], which argues for the strong impact of the Y chromosome on a large variety of immune cells and immune processes. Studies of Case et al. [10] showed that the copy number of the Y chromosome gene *Sly* and RNA-binding motif gene *Rbmy* inversely regulates the susceptibility to autoimmune disease. Studies in mouse influenza model [29] showed that certain genetic variants of chromosome Y result in higher susceptibility to influenza A infection, activate proinflammatory cytokine expression in T cells, and increase pathogenic immune response in the lungs (Fig. 2).

Mitochondria

Studies of the past decades [30, 31] showed that mitochondria are not only the bioenergetic (they generate ATP) and biosynthetic (they generate components for the synthesis of various macromolecules) hubs but are also indispensable regulators of the innate and adaptive immune response, development and maintenance/survival and activation of the specific phenotypes of immune cells. Mitochondria can also affect various signaling pathways and transcription in immune cells by changing the ATP level, alternating their metabolic pathways, and releasing the reactive oxygen species (ROS) and mitochondrial DNA (mtDNA) signals. By changing the type of respiration from the broken TCA cycle to β -oxidation, the mitochondria can switch macrophage phenotype from the pro-inflammatory (M1) to the antiinflammatory (M2). In addition, the localization of mitochondria in the proximity of endoplasmic reticulum (ER) membranes of the immune cells directly affects their metabolism and immune-related functions [30, 31]. Also, the outer membrane of mitochondria contains the mitochondrial antiviral signaling (MAVS) protein that is activated by the viral RNA sensor, the retinoic acid-inducible gene I (RIG-I) that senses the presence of viral RNA. MAVS can also act as an antiviral defense mechanism. In the macrophages infected with RNA viruses MAVS interacts with the antiviral protein viperin, affecting the level of antiviral compound interferon (Fig. 3); [30,32]. Proper functioning of mitochondrial MAVS response can be especially relevant and significant for COVID-19 infection where the SARS-CoV-2 virus directly infects alveolar macrophages inducing them to switch on the cytokine storm in the lungs [33].

Moreover, because the survival and replication of the virus depend on energy produced by the host mitochondria, which in turn, are affected by the virus (Fig. 3), the novel antiviral strategies may include compounds that modulate mitochondrial bioenergetic functions [34].

All these data indicate that the healthy and properly functioning mitochondria are indispensable for the adequacy of the immune response. Thus, it is not surprising that one of the theories explaining the higher infection rate and severity of infections, such as COVID-19, in males, relates to the maternal transmission of mitochondria, and substandard quality of mitochondria in the males. In mammals, including humans, mitochondria are maternally transmitted in the egg cytoplasm. Following fertilization, during early embryogenesis, maternal mitochondria replicate and enter the prospective somatic and germ cells [20]. Although there are some, very rare, examples of paternal transmission of mitochondria in humans [35], the concept of maternal inheritance of mitochondria UNA in humans remains valid. During the formation and maturation of the mammalian egg, the mitochondria undergo the quality control [36] and those which are defective, or contain, harmful for the female, mutations, are eliminated. Because this mitochondrial culling is female-biased, the mitochondria remaining in the egg cytoplasm may contain mutations harmful for the male. This so-called "mother curse" [20] may also influence the quality and caliber of a male immune response.

Recently, the melatonin was suggested as a potential adjuvant for COVID-19 treatment [37]. Melatonin has positive effects on mitochondrial homeostasis by scavenging toxic oxygen species, nitrogen-based reactants, enhancing antioxidative enzymes, facilitating the electron transport chain, limiting electron leakage, free radical generation, and stimulating ATP synthesis [38]. In light of our hypothesis that the quality of mitochondria may influence COVID-19 infection, the melatonin treatment of COVID-19 patients may be more beneficial for men than women.

Although, at present, there is no proven effective remedy for the sex-biased outcome of COVID-19 and other microbial infections, the detailed knowledge of all mechanisms underlying the differences between female and male immune response should facilitate the development of new treatments and therapeutic approaches.

Acknowledgements

We acknowledge that some of the images used to make figures were from the Servier Medical ART: SMART, smart.servier.com.

We also acknowledge the support from the Polish Ministry of National Defense project "Kościuszko" # 5508/2017/DA to JZK.

Statement of Ethics

Not applicable

Disclosure Statement

Authors do not have conflict of interest and have nothing to disclose

Funding Sources

Support from the Polish Ministry of National Defense project "Kościuszko" # 5508/2017/DA to JZK.

Authors Contributions

Malgorzata Kloc- concept, writing, figures

Jacek Z Kubiak-concept, writing, editing, figures

Rafik Mark Ghobrial- writing, editing

References

- 1. Klein SL, Flanagan KL. Sex differences in immune responses. Nat Rev Immunol. 2016, 16:626–638
- 2. Vemuri R, Sylvia KE, Klein SL, Forster SC, Plebanski M, Eri R, Flanagan KL. The microgenderome revealed: sex differences in bidirectional interactions between the microbiota, hormones, immunity and disease susceptibilitySemin Immunopathol. 2019, 41: 265–275.
- 3. Steegenga WT, Mischke M, Lute C, Boekschoten MV, Pruis MG, Lendvai A, Verkade HJ, Boekhorst J, Timmerman HM, Plösch T, Müller M. Sexually dimorphic characteristics of the small intestine and colon of prepubescent C57BL/6 mice. Biol Sex Differ. 2014, 5:11
- 4. Archive.ensembl.org/Homo_sapiensChromosome X: 1-1 Chromosome summary Homo sapiens <u>http://mar2017.archive.ensembl.org/Homo_sapiens/Location/Chromosome?r=X</u>
- 5. Bianchi I, Lleo A, Gershwin ME, Invernizzi P. The X chromosome and immune associated genes. Journal of Autoimmunity. 2012), 38: J187-J192
- 6. Schurz H, Salie M, Tromp G, Hoal EG, Kinnear CJ, Möller M. The X chromosome and sexspecific effects in infectious disease susceptibility. Human Genomics, 2019, 13:2
- 7. Lyon MF. Gene action in the X-chromosome of the mouse (Mus musculus L.). Nature. 1961; 190: 372–3.
- 8. Benjafield AV1, Wang WY, Morris BJ. (2004). No association of angiotensin-converting enzyme 2 gene (ACE2) polymorphisms with essential hypertension. Am J Hypertens. 2004; 17:624-8
- 9. Pan Y, Wang T, Li Y, Guan T, Yanxian Lai Y, et al. Association of ACE2 polymorphisms with susceptibility to essential hypertension and dyslipidemia in Xinjiang, China. Lipids Health Dis. 2018; 17: 241.
- Case LK, Wall EH, Dragon JA, Saligrama N, Krementsov DN, Moussawi M, Zachary JF, Huber SA, Blankenhorn EP, Teuscher C. The Y chromosome as a regulatory element shaping immune cell transcriptomes and susceptibility to autoimmune disease. Genome Res. 2013, 23: 1474–1485. doi: 10.1101/gr.156703.113
- 11. Koerich LB, Wang X, Clark AG, Carvalho AB. Low conservation of gene content in the Drosophila Y chromosome. Nature, 2008, 456: 949–951.

- 12. Quintana-Murci L, Fellous M. The human Y chromosome: The biological role of a "Functional Wasteland". J Biomed Biotechnol. 2001, 1:18-24.
- Skaletsky H, Kuroda-Kawaguchi T, Minx PJ, Cordum HS, Hillier L, Brown LG, Repping S, Pyntikova T, Ali J, Bieri T, et al. The male-specific region of the human Y chromosome is a mosaic of discrete sequence classes. Nature, 2003, 423: 825–837.
- 14. Lemos B, Araripe LO, Hartl DL. Polymorphic Y chromosomes harbor cryptic variation with manifold functional consequences. Science, 2008, 319: 91–93.
- Lemos B, Branco AT, Hartl DL. Epigenetic effects of polymorphic Y chromosomes modulate chromatin components, immune response, and sexual conflict. Proc Natl Acad Sci, 2010, 107: 15826–15831.
- Paredes S, Branco AT, Hartl DL, Maggert KA, Lemos B. Ribosomal DNA deletions modulate genome-wide gene expression: "rDNA-sensitive" genes and natural variation. PLoS Genet. 2011, 7: e1001376.
- 17. Piergentili R. Multiple roles of the Y chromosome in the biology of Drosophila melanogaster. ScientificWorld Journal, 2010, 10: 1749–1767.
- Zhou J, Sackton TB, Martinsen L, Lemos B, Eickbush TH, Hartl DL. Y chromosome mediates ribosomal DNA silencing and modulates the chromatin state in Drosophila. Proc Natl Acad Sci. 2012, 109: 9941–9946.
- 19. Francisco F, Lemos B. How do Y-chromosomes modulate genome-wide epigenetic states: genome folding, chromatin sinks, and gene expression. J Genomics. 2014, 2: 94–103.
- 20. Silkaitis K, Lemos B. Sex-biased chromatin and regulatory cross-talk between sex chromosomes, autosomes, and mitochondria Biol Sex Differ. 2014, 5: 2.
- 21. Taddei A, Hediger F, Neumann FR, Gasser SM. The function of nuclear architecture: a genetic approach. Annu Rev Genet. 2004, 5:305–345.
- 22. Ye Q, Worman HJ. Interaction between an integral protein of the nuclear envelope inner membrane and human chromodomain proteins homologous to Drosophila HP1. J Biol Chem. 1996, 5:14653–14656.
- 23. Charchar FJ, Bloomer LD, Barnes TA, Cowley MJ, Nelson CP, Wang Y, Denniff M, Debiec R, Christofidou P, Nankervis S, et al. Inheritance of coronary artery disease in men: An analysis of the role of the Y chromosome. Lancet, 2012, 379: 915–922.
- 24. Khan SI, Andrews KL, Jennings GL, Sampson AK, Chin-Dusting JPF. Y Chromosome, Hypertension and Cardiovascular Disease: Is Inflammation the Answer? Int J Mol Sci. 2019, 20: 2892.
- 25. Maan AA, Eales J, Akbarov A, Rowland J, Xu X, Jobling MA, Charchar FJ, Tomaszewski M. The Y chromosome: a blueprint for men's health? European Journal of Human Genetics, 2017, 25:1181–1188
- 26. Case LK, Toussaint L, MoussawiM, Roberts B, Saligrama N, Brossay L, HuberSA, Teuscher C. Chromosome Y regulates survival following murine coxsackievirus B3 infection. G3-Genes Genom Genet. 2012, 2: 115–121.
- 27. Teuscher C, Noubade R, Spach K, McElvany B, Bunn JY, Fillmore PD, Zachary JF, Blankenhorn EP. Evidence that the Y chromosome influences autoimmune disease in male and female mice. Proc Natl Acad Sci. 2006, 103: 8024–8029.
- 28. Spach KM, Blake M, Bunn JY, McElvany B, Noubade R, Blankenhorn EP, Teuscher C. Cutting edge: The Y chromosome controls the age dependent experimental allergic encephalomyelitis sexual dimorphism in SJL/J mice. J Immunol. 2009, 182: 1789–1793
- 29. Krementsov DN, Case LK, Dienz O, Raza A, Fang Q, Ather JL, Poynter ME, Boyson JE, Bunn JY, Teuscher C. Genetic variation in chromosome Y regulates susceptibility to influenza A virus infection. PNAS. (2017). 114 : 3491–3496
- 30. Angajala A, Lim S, Phillips JB, Kim J-H, Yates C, You Z. Diverse Roles of Mitochondria in Immune Responses: Novel Insights Into Immuno-Metabolism. Front Immunol. 2018; 9:1605.

- 31. Weinberg SE, Sena LA, Chandel NS. Mitochondria in the regulation of innate and adaptive immunity. Immunity. Author manuscript; available in PMC Immunity. 2015, 42: 406–417.
- 32. Hee JS, Cresswell P. Viperin interaction with mitochondrial antiviral signaling protein (MAVS) limits viperin-mediated inhibition of the interferon response in macrophages. PLoS One, 2017, 12(2):e0172236.
- 33. Wang C, Xie J, Zhao L, Fei X, Zhang H, et al. Alveolar Macrophage Activation and Cytokine Storm in the Pathogenesis of Severe COVID-19. 2020, www.researchsquare.com doi:10.21203/rs.3.rs-19346/v1
- 34. El-Bacha T, Da Poian AT. Virus-induced changes in mitochondrial bioenergetics as potential targets for therapy The International Journal of Biochemistry & Cell Biology, 2013, 45, 41-46
- 35. Shiyu LC. Valencia A, Zhang J, Lee N-C, Slone J et al. Biparental Inheritance of Mitochondrial DNA in Humans. PNAS. 2018, 115: 13039-13044.
- 36. Lieber, T., Jeedigunta, S.P., Palozzi, J.M. et al. Mitochondrial fragmentation drives selective removal of deleterious mtDNA in the germline. Nature, 2019, 570, 380–384.
- 37. Zhang R, Wang X, Ni L, Di X, Ma B, Niu S, Liu C, Reiter RJ. COVID-19: Melatonin as a potential adjuvant treatment. Life Sci. 2020, 250:1175830
- 38. Leon J, Acuña-Castroviejo D, Sainz RM, Mayo JC, Tan DX, Reiter RJ. Melatonin and mitochondrial function. Life Sci. 2004 75:765-90.

Figure legends



Fig. 1. Microgenderome effect on the immune system

The microorganisms acquired during birth, and from the environment (air and food) colonize the human body, especially the digestive system. After puberty, the sex hormones influence the abundance and variety of species inhabiting the male and female body; less variety in males than in females. The compounds produced by the microgendorome reciprocally influence the function of innate and adaptive immunity, and brain (adapted from Vemuri et al. [2]).





Fig. 2. Effect of sex chromosomes on the immune-related functions

The X chromosome contains many immune response-related genes and 10% of all genomic miRNAs that regulate transcription, splicing, and degradation of many genes. It also, through the specification of female identity, directs maternal purging and transmission of mitochondria. Because in the female cells, the inactivation of one of the X chromosomes is random, the female body may contain different alleles of the same gene. For example, different alleles of viral receptor ACE2 may have a different affinity to the virus, which will affect the anti-viral immune response. The Y chromosome regulates many different immune-response functions, immune cell numbers, and immune cell phenotypes through the regulation of transcriptionally inert (heterochromatin) and active (euchromatin) status of X chromosome and autosomes.



Fig. 3. Effect of mitochondria on the anti-viral immune response.

The presence of the virus is sensed by the RIG-I that is a major sensor of viral RNA. This activates antiviral MAVS protein present in the mitochondrial membrane. Activated MAVS, through the interaction with the virus inhibitory protein, viperin, affects the level of interferon, which in turn regulates virus replication. In addition, viruses can directly affect mitochondrial functions such as aerobic or anaerobic respiration, which in turn affect immune cell phenotypes and responses.

10