



HAL
open science

GEOMETRIC CONDITIONS FOR THE EXACT CONTROLLABILITY OF FRACTIONAL FREE AND HARMONIC SCHRÖDINGER EQUATIONS

Jérémy Martin, Karel Pravda-Starov

► **To cite this version:**

Jérémy Martin, Karel Pravda-Starov. GEOMETRIC CONDITIONS FOR THE EXACT CONTROLLABILITY OF FRACTIONAL FREE AND HARMONIC SCHRÖDINGER EQUATIONS. *Journal of Evolution Equations*, 2021, 21 (1), pp.1059-1087. 10.1007/s00028-020-00618-6 . hal-02893885

HAL Id: hal-02893885

<https://hal.science/hal-02893885>

Submitted on 8 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GEOMETRIC CONDITIONS FOR THE EXACT CONTROLLABILITY OF FRACTIONAL FREE AND HARMONIC SCHRÖDINGER EQUATIONS

JÉRÉMY MARTIN, KAREL PRAVDA-STAROV

ABSTRACT. We provide necessary and sufficient geometric conditions for the exact controllability of the one-dimensional fractional free and fractional harmonic Schrödinger equations. The necessary and sufficient condition for the exact controllability of fractional free Schrödinger equations is derived from the Logvinenko-Sereda theorem and its quantitative version established by Kovrijkine, whereas the one for the exact controllability of fractional harmonic Schrödinger equations is deduced from an infinite dimensional version of the Hautus test for Hermite functions and the Plancherel-Rotach formula.

1. INTRODUCTION

The purpose of this article is to provide necessary and sufficient geometric conditions on control subsets to ensure exact controllability of fractional free Schrödinger equations and fractional harmonic Schrödinger equations. The notion of exact controllability is defined as follows:

Definition 1.1 (Exact controllability). *Let B be a closed operator on $L^2(\mathbb{R}^d)$ which is the infinitesimal generator of a strongly continuous semigroup $(e^{-tB})_{t \geq 0}$ on $L^2(\mathbb{R}^d)$, $T > 0$ and ω be a measurable subset of \mathbb{R}^d . The equation*

$$(1.1) \quad \begin{cases} (\partial_t + B)f(t, x) = u(t, x)\mathbb{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

is said to be exactly controllable from the control set ω in time $T > 0$ if, for any initial datum $f_0 \in L^2(\mathbb{R}^d)$ and any final datum $f_T \in L^2(\mathbb{R}^d)$, there exists $u \in L^2((0, T) \times \mathbb{R}^d)$, supported in $(0, T) \times \omega$, such that the mild (or semigroup) solution of (1.1) satisfies $f(T, \cdot) = f_T$. The equation (1.1) is said to be exactly controllable from the control set ω if there exists $T > 0$ such that this system is exactly controllable from ω in time $T > 0$.

By the Hilbert uniqueness method, see [3, 9], the exact controllability of the evolution equation (1.1) is equivalent to the exact observability of the adjoint system

$$(1.2) \quad \begin{cases} (\partial_t + B^*)g(t, x) = 0, & x \in \mathbb{R}^d, t > 0, \\ g|_{t=0} = g_0 \in L^2(\mathbb{R}^d), \end{cases}$$

where B^* denotes the $L^2(\mathbb{R}^d)$ -adjoint of the operator B . The notion of exact observability is defined as follows:

Definition 1.2 (Exact observability). *Let $T > 0$ and ω be a measurable subset of \mathbb{R}^d . The evolution equation (1.2) is said to be exactly observable from the control set ω in*

2010 *Mathematics Subject Classification.* 93B05.

Key words and phrases. Exact controllability, Schrödinger equation, harmonic oscillator.

time $T > 0$ if there exists a positive constant $C_T > 0$ such that, for any initial datum $g_0 \in L^2(\mathbb{R}^d)$, the mild (or semigroup) solution of (1.2) satisfies

$$\int_{\mathbb{R}^d} |g_0(x)|^2 dx \leq C_T \int_0^T \left(\int_{\omega} |g(t, x)|^2 dx \right) dt.$$

In this paper, we study evolution equations associated to skew-selfadjoint operators. More precisely, we consider evolution equations

$$(1.3) \quad \begin{cases} \partial_t f(t, x) + i(Af)(t, x) = 0, & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

where the selfadjoint operators A on $L^2(\mathbb{R}^d)$ generating a strongly continuous group $(e^{itA})_{t \in \mathbb{R}}$ on $L^2(\mathbb{R}^d)$ is specifically given by a fractional Laplacian $A = (-\Delta_x)^s$ or a fractional harmonic oscillator $A = (-\Delta_x + |x|^2)^s$, with $s > 0$.

Our starting point is the following result proved by Miller. This author shows in [11, Corollary 2.17] that the observability of the system (1.3) is equivalent to the following spectral estimates:

Proposition 1.3 (Miller [11, Corollary 2.17]). *Let A be a selfadjoint operator on $L^2(\mathbb{R}^d)$, which is the infinitesimal generator of a strongly continuous group $(e^{itA})_{t \in \mathbb{R}}$ on $L^2(\mathbb{R}^d)$. If the evolution equation (1.3) is exactly observable from a measurable subset $\omega \subset \mathbb{R}^d$ at time $T > 0$ then there exist some positive constants $k > 0$ and $D > 0$ such that*

$$(1.4) \quad \forall \lambda \in \mathbb{R}, \forall f \in \mathbb{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(L^2(\mathbb{R}^d)), \quad \|f\|_{L^2(\mathbb{R}^d)} \leq \sqrt{k} \|f\|_{L^2(\omega)}.$$

Conversely, when the spectral estimates (1.4) hold for some $k > 0$ and $D > 0$, then the system (1.3) is exactly observable from ω at any time

$$T > \pi \sqrt{\frac{1+k}{D}}.$$

The operators $\mathbb{1}_{\{|A-\lambda| \leq \sqrt{D}\}}$ appearing in (1.4) are defined by functional calculus. It is important to point out that the constants D and k do not depend on λ . We recall the proof given by Miller in Appendix (Section 5.2) since the minimal time obtained by Miller in [11, Corollary 2.17] is different to ours.

When the operator $A = (-\Delta_x)^s$, with $s > 0$, is a fractional Laplacian, the family $(\mathbb{1}_{\{|A-\lambda| \leq \sqrt{D}\}})_{\lambda \in \mathbb{R}}$ is a family of frequency cutoff operators on bounded frequency subsets and for all $D > 0$ and $\lambda \in \mathbb{R}$,

$$(1.5) \quad \mathbb{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(L^2(\mathbb{R}^d)) = \left\{ f \in L^2(\mathbb{R}^d) : \text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^d, ||\xi|^{2s} - \lambda| \leq \sqrt{D}\} \right\},$$

where $|\cdot|$ denotes the Euclidean norm on \mathbb{R}^d . The spectral estimates (1.4) are in this case related to the notion of annihilating pairs:

Definition 1.4 (Annihilating pairs). *Let S, Σ be two measurable subsets of \mathbb{R}^d .*

- *The pair (S, Σ) is said to be a weak annihilating pair if the only function $f \in L^2(\mathbb{R}^d)$ with $\text{supp } f \subset S$ and $\text{supp } \hat{f} \subset \Sigma$ is zero $f = 0$.*
- *The pair (S, Σ) is said to be a strong annihilating pair if there exists a positive constant $C = C(S, \Sigma) > 0$ such that for all $f \in L^2(\mathbb{R}^d)$,*

$$\int_{\mathbb{R}^d} |f(x)|^2 dx \leq C \left(\int_{\mathbb{R}^d \setminus S} |f(x)|^2 dx + \int_{\mathbb{R}^d \setminus \Sigma} |\hat{f}(\xi)|^2 d\xi \right).$$

It can be readily checked that a pair (S, Σ) is a strong annihilating one if and only if there exists a positive constant $\tilde{C} = \tilde{C}(S, \Sigma) > 0$ such that

$$(1.6) \quad \forall f \in L^2(\mathbb{R}^d), \text{ supp } \hat{f} \subset \Sigma, \quad \|f\|_{L^2(\mathbb{R}^d)} \leq \tilde{C} \|f\|_{L^2(\mathbb{R}^d \setminus S)}.$$

We deduce from the Hilbert uniqueness method, Proposition 1.3 and (1.5) that the exact controllability of the fractional free Schrödinger equation

$$\begin{cases} \partial_t f(t, x) + i(-\Delta_x)^s f(t, x) = u(t, x) \mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

with $s > 0$, from a measurable control set ω holds if and only if its complement $\mathbb{R}^d \setminus \omega$ forms a strong annihilating pair with the frequency set

$$\mathcal{C}_D(\lambda) = \{\xi \in \mathbb{R}^d : ||\xi|^{2s} - \lambda| \leq \sqrt{D}\},$$

for some $D > 0$ and all $\lambda \in \mathbb{R}$ with an uniform constant \tilde{C} with respect to λ in (1.6). An exhaustive description of annihilating pairs is for now out of reach. However, the Logvinenko-Sereda Theorem [10] gives a complete description of all support sets forming a strong annihilating pair with a given bounded frequency set:

Theorem 1.5 (Logvinenko-Sereda [10]). *Let $S, \Sigma \subset \mathbb{R}^d$ be measurable subsets with Σ bounded. Denoting $\tilde{S} = \mathbb{R}^d \setminus S$, the following assertions are equivalent:*

- *The pair (S, Σ) is a strong annihilating pair*
- *The subset \tilde{S} is thick, that is, there exists a cube $K \subset \mathbb{R}^d$ with sides parallel to coordinate axes and a positive constant $0 < \gamma \leq 1$ such that*

$$\forall x \in \mathbb{R}^d, \quad |(K + x) \cap \tilde{S}| \geq \gamma |K| > 0,$$

where $|A|$ denotes the Lebesgue measure of the measurable set A .

Notice that if the support set S forms a strong annihilating pair with some bounded frequency set Σ , then S forms a strong annihilating pair with any bounded frequency set. In order to be able to use the Logvinenko-Sereda theorem in control theory, it is essential to understand how the constant $\tilde{C}(S, \Sigma)$ appearing in (1.6) depends on the bounded frequency set Σ . In [8, Theorem 3], Kovrijkine addresses this question by establishing the following quantitative version of the Logvinenko-Sereda Theorem:

Theorem 1.6 (Kovrijkine [8, Theorem 3]). *Let $\omega \subset \mathbb{R}^d$ be a measurable subset γ -thick at scale $L > 0$, that is, satisfying*

$$(1.7) \quad \forall x \in \mathbb{R}^d, \quad |([0, L]^d + x) \cap \omega| \geq \gamma L^d > 0,$$

with $0 < \gamma \leq 1$. *There exists a universal positive constant $C > 0$ independent on the dimension $d \geq 1$ such that for all $f \in L^2(\mathbb{R}^d)$ satisfying $\text{supp } \hat{f} \subset J$, with J a cube with sides of length b parallel to coordinate axes,*

$$(1.8) \quad \|f\|_{L^2(\mathbb{R}^d)} \leq c(\gamma, d, L, b) \|f\|_{L^2(\omega)},$$

with

$$c(\gamma, d, L, b) = \left(\frac{C^d}{\gamma} \right)^{Cd(Lb+1)}.$$

This result is extended to the case of frequency supports covered by the union of finitely many parallelepipeds in [8, Theorem 4]. In this work, we only use this result in the one-dimensional setting:

Theorem 1.7 (Kovrijkine [8, Theorem 2]). *Let $\omega \subset \mathbb{R}$ be a measurable subset γ -thick at scale $L > 0$, that is, satisfying*

$$\forall x \in \mathbb{R}, \quad |([0, L] + x) \cap \omega| \geq \gamma L > 0,$$

with $0 < \gamma \leq 1$. There exists a universal positive constant $C > 0$ such that for all $f \in L^2(\mathbb{R})$ satisfying $\text{supp } \hat{f} \subset \bigcup_{k=1}^m J_k$, with $(J_k)_{1 \leq k \leq m}$ a finite family of intervals with length $|J_k| = b$ for all $1 \leq k \leq m$,

$$(1.9) \quad \|f\|_{L^2(\mathbb{R})} \leq c(\gamma, m, L, b) \|f\|_{L^2(\omega)},$$

with

$$c(\gamma, m, L, b) = \left(\frac{C}{\gamma}\right)^{Lb\left(\frac{C}{\gamma}\right)^m + m - \frac{1}{2}}.$$

Notice that the constant $c(\gamma, m, L, b)$ depends only on the number of intervals and their length but not on their locations. Thanks to this explicit dependence of the constant with respect to the length of the intervals in (1.9), Egidi and Veselić [4]; and Whang, Whang, Zhang and Zhang [13] have independently established that the heat equation

$$(1.10) \quad \begin{cases} (\partial_t - \Delta_x)f(t, x) = u(t, x)\mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

is null-controllable in any positive time $T > 0$ from a measurable control set $\omega \subset \mathbb{R}^d$ if and only if the control subset ω is thick in \mathbb{R}^d . In the recent work [2], Beauchard, Jaming and Pravda-Starov prove that this thickness condition is also sufficient for the null-controllability of the harmonic heat equation

$$\begin{cases} (\partial_t + (-\Delta_x + |x|^2))f(t, x) = u(t, x)\mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d). \end{cases}$$

More generally, the result of [2, Theorem 2.2] shows that the thickness condition is a sufficient condition for the null-controllability of a large class of hypoelliptic quadratic equations.

When the operator A appearing in the evolution equation (1.3) has a compact resolvent, the result of Proposition 1.3 is specified further as the following infinite dimensional version of the Hautus test:

Proposition 1.8 (Miller [11, Corollary 2.18]). *Let A be a selfadjoint operator on $L^2(\mathbb{R}^d)$, which is the infinitesimal generator of a strongly continuous group $(e^{itA})_{t \in \mathbb{R}}$ on $L^2(\mathbb{R}^d)$. When the operator A has a compact resolvent with a spectral gap $\gamma > 0$ in the following sense: $|\lambda - \mu| \geq \gamma$ for all distinct eigenvalues λ and μ , the evolution equation (1.3) is exactly observable at some time $T > 0$ from a measurable subset ω if and only if*

$$\exists \delta > 0, \quad \text{for all eigenvector } \psi \text{ of } A, \quad \|\psi\|_{L^2(\mathbb{R}^d)} \leq \delta \|\psi\|_{L^2(\omega)}.$$

When the spectral gap condition holds, the observability of the evolution equation (1.3) then depends only on some properties of the eigenvectors of A . Notice in particular that this spectral gap condition holds for the one-dimensional harmonic oscillator

$$\mathcal{H} = -\Delta_x + x^2,$$

as well as the fractional harmonic oscillators \mathcal{H}^s , when $s \geq 1$, since the spectra of these operators are given by

$$\sigma((-\Delta_x + x^2)^s) = \{(2n + 1)^s : n \in \mathbb{N}\}.$$

In this work, we aim at studying necessary and sufficient geometric conditions on the control subsets ω to ensure exact controllability of the free fractional Schrödinger equations

$$\begin{cases} \partial_t f(t, x) + i(-\Delta_x)^s f(t, x) = u(t, x)\mathbb{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

and the fractional harmonic Schrödinger equations

$$(1.11) \quad \begin{cases} \partial_t f(t, x) + i(-\Delta_x + |x|^2)^s f(t, x) = u(t, x)\mathbb{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

with $s > 0$. The first result contained in this paper (Corollary 2.7) shows that the thickness condition defined in Theorem 1.5 is a necessary and sufficient geometric condition for the exact controllability of the fractional free Schrödinger equations when the fractional parameter satisfies $s \geq \frac{1}{2}$. In the multidimensional setting, the thickness condition is always a necessary condition for exact controllability when the fractional parameter satisfies $s > 0$ (Theorem 2.5). A sufficient geometric condition to ensure the exact controllability from a neighborhood of the control subset is also given in the multidimensional case when $s \geq \frac{1}{2}$ (Proposition 2.9). These results are based on the Logvinenko-Sereda Theorem, its quantitative versions proved by Kovrijkine and new uncertainty principles recently established by Green, Jaye and Mitkovski [5].

Regarding the fractional harmonic Schrödinger equations, we prove that the following geometric condition

$$(1.12) \quad \forall A \in O(\mathbb{R}^d), \quad \liminf_{R_1 \rightarrow +\infty} \dots \liminf_{R_d \rightarrow +\infty} \frac{|A(\omega) \cap [-R_1, R_1] \times \dots \times [-R_d, R_d]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} > 0,$$

turns out to be necessary for exact controllability in any dimension $d \geq 1$ (Theorem 2.3) and sufficient in the one-dimensional setting (Theorem 2.2). The proof of this result is based on the infinite dimensional version of the Hautus test given by Proposition 1.8 applied with Hermite functions together with the Rotach-Plancherel formula for Hermite polynomials. This necessary condition enables us to derive a negative result of exact controllability from a cone in dimension 2. In the case of the one-dimensional harmonic Schrödinger equation, we establish that exact observability from a control subset satisfying (1.12) holds at time $T = \pi$.

2. STATEMENTS OF THE MAIN RESULTS

2.1. Hautus test and exact controllability of fractional harmonic Schrödinger equations. In this section, we state the results related to the exact controllability of fractional harmonic Schrödinger equations. We recall that the exact controllability of the evolution equations (1.11) is equivalent to the observability of the adjoint systems

$$(2.1) \quad \begin{cases} \partial_t f(t, x) - i(-\Delta_x + |x|^2)^s f(t, x) = 0, & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d). \end{cases}$$

The observability of (2.1) is deduced thanks to Proposition 1.8 by an infinite dimensional Hautus test on the Hermite functions seen as eigenvectors of the fractional harmonic oscillator.

Hermite functions are normalized $L^2(\mathbb{R})$ -functions defined as

$$(2.2) \quad \forall n \geq 0, \quad \psi_n(x) = \frac{1}{\pi^{\frac{1}{4}} 2^{\frac{n}{2}} \sqrt{n!}} e^{-\frac{x^2}{2}} H_n(x),$$

where H_n stands for the Hermite polynomial of degree $n \geq 0$, see e.g. [12], formula (5.5.1). The family $(\psi_n)_{n \geq 0}$ defines a Hilbert basis of $L^2(\mathbb{R})$. They satisfy the following infinite dimensional Hautus test:

Proposition 2.1. *Let ω be a measurable subset of \mathbb{R} and $(\psi_n)_{n \geq 0}$ be the Hermite basis on $L^2(\mathbb{R})$ defined in (2.2). The following assertions are equivalent:*

- (i) $\liminf_{R \rightarrow +\infty} \frac{|\omega \cap [-R, R]|}{|[-R, R]|} > 0,$
- (ii) $\exists c > 0, \forall n \geq 0, \quad 1 = \|\psi_n\|_{L^2(\mathbb{R})} \leq c \|\psi_n\|_{L^2(\omega)},$
- (iii) $\liminf_{n \rightarrow +\infty} \|\psi_n\|_{L^2(\omega)} > 0.$

The proof of Proposition 2.1 is given in Section 3 and is based on the Plancherel-Rotach formula. We readily deduce from Proposition 1.8 and 2.1 the following result of exact controllability in the one-dimensional case:

Theorem 2.2. *Let $s \geq 1$ and ω be a measurable set in \mathbb{R} . The two following assertions are equivalent:*

- (i) *The fractional harmonic Schrödinger equation*

$$\begin{cases} \partial_t f(t, x) + i(-\Delta_x + x^2)^s f(t, x) = u(t, x) \mathbb{1}_\omega(x), & x \in \mathbb{R}, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}), \end{cases}$$

is exactly controllable from the set ω for some positive time $T > 0$,

- (ii) *The control set obeys the following geometric condition*

$$\liminf_{R \rightarrow +\infty} \frac{|\omega \cap [-R, R]|}{|[-R, R]|} > 0.$$

If one of the two above conditions holds in the non fractional case $s = 1$, then the harmonic Schrödinger equation is exactly controllable from the control subset ω in time $T = \pi$.

The equivalence between (i) and (ii) directly follows from Propositions 1.8 and 2.1, since the Hermite basis $(\psi_n)_{n \geq 0}$ is a Hilbert basis of eigenvectors for the fractional harmonic oscillator \mathcal{H}^s associated to the eigenvalues $((2n+1)^s)_{n \geq 0}$ that satisfy the spectral gap condition when $s \geq 1$. Let us now check that when exact controllability holds for some positive time $T_0 > 0$ then it necessary holds in time $T = \pi$. To that end, we just notice that the function

$$t \in \mathbb{R} \mapsto \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2,$$

is π -periodic since

$$\begin{aligned} \forall t \in \mathbb{R}, \forall f_0 \in L^2(\mathbb{R}), \quad e^{i(t+\pi)(-\Delta_x + x^2)} f_0 &= \sum_{n \in \mathbb{N}} e^{i(t+\pi)(2n+1)} \langle f_0, \psi_n \rangle_{L^2(\mathbb{R})} \psi_n \\ &= -e^{it(-\Delta_x + x^2)} f_0. \end{aligned}$$

By recalling that exact controllability at time $T_0 > 0$ is equivalent to the observability of the evolution equation

$$\begin{cases} \partial_t f(t, x) - i(-\Delta_x + x^2) f(t, x) = 0, & x \in \mathbb{R}, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}), \end{cases}$$

at time $T_0 > 0$, that is,

$$\exists C_{T_0} > 0, \forall f_0 \in L^2(\mathbb{R}), \quad \|f_0\|_{L^2(\mathbb{R})}^2 \leq C_{T_0} \int_0^{T_0} \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2 dt.$$

The π -periodicity property readily implies observability and exact controllability at time $T = \pi$, since for all $f_0 \in L^2(\mathbb{R})$,

$$\begin{aligned} \|f_0\|_{L^2(\mathbb{R})}^2 &\leq C_{T_0} \int_0^{T_0} \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2 dt \leq C_{T_0} \int_0^{(\lfloor \frac{T_0}{\pi} \rfloor + 1)\pi} \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2 dt \\ &\leq C_{T_0} \sum_{k=0}^{\lfloor \frac{T_0}{\pi} \rfloor} \int_{k\pi}^{(k+1)\pi} \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2 dt \leq C_{T_0} \left(\lfloor \frac{T_0}{\pi} \rfloor + 1 \right) \int_0^\pi \|e^{it(-\Delta_x + x^2)} f_0\|_{L^2(\omega)}^2 dt, \end{aligned}$$

where $\lfloor \cdot \rfloor$ denotes the floor function.


Theorem 2.2 provides a necessary and sufficient geometric condition on the control subset ω to ensure exact controllability of fractional harmonic Schrödinger equations in the one-dimensional setting. In higher dimensions $d \geq 2$, it is natural to investigate if the geometric condition

$$(2.3) \quad \liminf_{R \rightarrow +\infty} \frac{|\omega \cap B_d(0, R)|}{|B_d(0, R)|} > 0,$$

where $B_d(0, R)$ denotes the Euclidean ball in \mathbb{R}^d centered at 0 with radius R , turns out to be sufficient to ensure exact controllability. This is actually not the case in general as the next result shows that condition (2.3) is not sufficient for the exact controllability of the harmonic Schrödinger equation posed on \mathbb{R}^2 . The stronger necessary geometric condition for exact controllability given by the following proposition proves in particular that the exact controllability of the harmonic Schrödinger equation posed on \mathbb{R}^2 cannot hold from the cone

$$(2.4) \quad \omega_\delta = \left\{ (r \cos \theta, r \sin \theta) : r \geq 0, |\theta| \leq \frac{\pi}{2} - \delta \right\} \subset \mathbb{R}^2,$$

with $0 < \delta \leq \frac{\pi}{2}$,



as the control subset ω_δ satisfies condition (2.3), but fails to satisfy the following necessary geometric condition

$$\liminf_{R_1 \rightarrow +\infty} \liminf_{R_2 \rightarrow +\infty} \frac{|\omega \cap [-R_1, R_1] \times [-R_2, R_2]|}{|[-R_1, R_1] \times [-R_2, R_2]|} > 0,$$

given by the following result:

Theorem 2.3. *Let $s \geq 1$ and ω be a measurable subset of \mathbb{R}^d with $d \geq 1$. If the fractional harmonic Schrödinger equation*

$$(2.5) \quad \begin{cases} \partial_t f(t, x) + i(-\Delta_x + |x|^2)^s f(t, x) = u(t, x) \mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

is exactly controllable from ω for some positive time $T > 0$, then the control subset ω does obey the following geometric condition

$$(2.6) \quad \forall A \in O(\mathbb{R}^d), \quad \liminf_{R_1 \rightarrow +\infty} \dots \liminf_{R_d \rightarrow +\infty} \frac{|A(\omega) \cap [-R_1, R_1] \times \dots \times [-R_d, R_d]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} > 0,$$

where $O(\mathbb{R}^d)$ stands for the orthogonal group in dimension d .

This result shows that the exact controllability of the fractional harmonic Schrödinger equations requires the control subset to be distributed in any space direction. The sufficiency of condition (2.6) for exact controllability is an open question in dimension $d \geq 2$. The proof of Theorem 2.3 is given in Section 4. An example of sufficient geometric condition is given by the following proposition, which is deduced from Theorem 2.2:

Example 2.4. Let $d \geq 1$, $s \geq 1$ and ω be a measurable subset of \mathbb{R} satisfying

$$\liminf_{R \rightarrow +\infty} \frac{|\omega \cap [-R, R]|}{|[-R, R]|} > 0.$$

The evolution equation

$$(2.7) \quad \begin{cases} \partial_t f(t, x) + i(-\Delta_x + |x|^2)^s f(t, x) = u(t, x) \mathbb{1}_{\omega \times \mathbb{R}^{d-1}}(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

is exactly controllable from $\omega \times \mathbb{R}^{d-1}$ for some positive time $T > 0$.

Proof. We first notice that the spectral gap condition in Proposition 1.8 is satisfied by the fractional harmonic operator $(-\Delta_x + |x|^2)^s$, since its spectrum is given by

$$\sigma((-\Delta_x + |x|^2)^s) = \{(2n + d)^s : n \in \mathbb{N}\}.$$

Let $N \in \mathbb{N}$ and f be an eigenvector associated to the eigenvalue $(2N + d)^s$. There exists a family of complex numbers $(C_\alpha)_{\alpha \in \mathbb{N}^d, |\alpha|=N}$ such that

$$f = \sum_{\substack{\alpha \in \mathbb{N}^d, \\ |\alpha|=N}} C_\alpha \Psi_\alpha,$$

with

$$\forall \alpha = (\alpha_1, \dots, \alpha_d) = (\alpha_1, \alpha') \in \mathbb{N}^d, \forall x = (x_1, \dots, x_d) = (x_1, x') \in \mathbb{R}^d, \quad \Psi_\alpha(x) = \prod_{i=1}^d \psi_{\alpha_i}(x_i).$$

We notice that

$$\begin{aligned} \|f\|_{L^2(\omega \times \mathbb{R}^{d-1})}^2 &= \int_\omega \left(\int_{\mathbb{R}^{d-1}} \left| \sum_{\substack{\alpha \in \mathbb{N}^d, \\ |\alpha|=N}} C_\alpha \Psi_\alpha(x_1, x') \right|^2 dx' \right) dx_1 \\ &= \int_\omega \left(\int_{\mathbb{R}^{d-1}} \left| \sum_{\substack{\alpha' \in \mathbb{N}^{d-1}, \\ |\alpha'| \leq N}} C_{N-|\alpha'|, \alpha'} \psi_{N-|\alpha'|}(x_1) \Psi_{\alpha'}(x') \right|^2 dx' \right) dx_1 \\ &= \int_\omega \sum_{\substack{\alpha' \in \mathbb{N}^{d-1}, \\ |\alpha'| \leq N}} |C_{N-|\alpha'|, \alpha'}|^2 |\psi_{N-|\alpha'|}(x_1)|^2 dx_1 = \sum_{\substack{\alpha' \in \mathbb{N}^{d-1}, \\ |\alpha'| \leq N}} |C_{N-|\alpha'|, \alpha'}|^2 \|\psi_{N-|\alpha'}\|_{L^2(\omega)}^2. \end{aligned}$$

By using Proposition 2.1,

$$\exists c > 0, \forall n \geq 0, \quad 1 = \|\psi_n\|_{L^2(\mathbb{R})} \leq c \|\psi_n\|_{L^2(\omega)},$$

we deduce that

$$(2.8) \quad \|f\|_{L^2(\omega \times \mathbb{R}^{d-1})}^2 \geq \frac{1}{c^2} \sum_{\substack{\alpha' \in \mathbb{N}^{d-1}, \\ |\alpha'| \leq N}} |C_{N-|\alpha'|, \alpha'}|^2 = \frac{1}{c^2} \sum_{\substack{\alpha \in \mathbb{N}^d, \\ |\alpha| = N}} |C_\alpha|^2 = \frac{1}{c^2} \|f\|_{L^2(\mathbb{R}^d)}^2.$$

Thanks to Proposition 1.8, we deduce from (2.8) the observability of the adjoint system

$$\begin{cases} \partial_t g(t, x) - i(-\Delta_x + |x|^2)^s g(t, x) = 0, & x \in \mathbb{R}^d, t > 0, \\ g|_{t=0} = g_0 \in L^2(\mathbb{R}^d), \end{cases}$$

and then the exact controllability of the evolution equation (2.7). \square

The above example ensures the exact controllability of fractional harmonic Schrödinger equations from the half space $\mathbb{R}_+ \times \mathbb{R}^{d-1}$. In particular, it appears that the cone (2.4) is a limit case for exact controllability.

2.2. Exact controllability of fractional free Schrödinger equations. In this section, we study the exact controllability of the fractional free Schrödinger equation

$$(2.9) \quad \begin{cases} (\partial_t + i(-\Delta_x)^s) f(t, x) = u(t, x) \mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

with $s > 0$, from a measurable control subset $\omega \subset \mathbb{R}^d$. By the Hilbert uniqueness method, the exact controllability of the evolution equation (2.9) is equivalent to the observability of the adjoint system

$$(2.10) \quad \begin{cases} (\partial_t - i(-\Delta_x)^s) g(t, x) = 0, & x \in \mathbb{R}^d, t > 0, \\ g|_{t=0} = g_0 \in L^2(\mathbb{R}^d). \end{cases}$$

We first notice that the Logvinenko-Sereda Theorem recalled in Theorem 1.5 shows that the thickness condition is a necessary geometric condition for the exact controllability of the fractional Schrödinger equation (2.9) in any dimension $d \geq 1$:

Theorem 2.5. *Let $s > 0$, ω be a measurable subset of \mathbb{R}^d and $T > 0$. If the fractional free Schrödinger equation (2.9) is exactly controllable from ω in some time $T > 0$, then the control subset ω is a thick set.*

Proof. The assumptions of Theorem 2.5 imply that the system (2.10) is exactly observable from ω for the time $T > 0$. We deduce from Proposition 1.3 that there exist some positive constants $k > 0$ and $D > 0$ such that for all $\lambda \in \mathbb{R}$ and $f \in L^2(\mathbb{R}^d)$,

$$(2.11) \quad \text{supp } \widehat{f} \subset \{\xi \in \mathbb{R}^d : \|\xi\|^{2s} - \lambda \leq \sqrt{D}\} \implies \|f\|_{L^2(\mathbb{R}^d)} \leq \sqrt{k} \|f\|_{L^2(\omega)}.$$

While taking $\lambda = 0$, the assertion (2.11) implies in particular that the sets $\mathbb{R}^d \setminus \omega$ and the Euclidean ball $B_d(0, D^{\frac{1}{4s}})$ in \mathbb{R}^d make a strong annihilating pair. Since $B_d(0, D^{\frac{1}{4s}})$ is a bounded set, the Logvinenko-Sereda Theorem (Theorem 1.5) ensures that the control subset ω must be thick. \square

In the one-dimensional setting, the quantitative version of the Logvinenko-Sereda Theorem established by Kovrijkine (Theorem 1.6) together with Proposition 1.3 show that the thickness condition is also sufficient for the exact controllability of the fractional free Schrödinger equations (2.9) when $s \geq \frac{1}{2}$:

Theorem 2.6. *If $s \geq \frac{1}{2}$ and ω is a thick measurable subset of \mathbb{R} , then there exists a positive constant $T_0 > 0$ such that for any time $T > T_0$, the fractional free Schrödinger equation (2.9) is exactly controllable from ω in time T .*

Proof. Let $\omega \subset \mathbb{R}$ be a measurable subset γ -thick at scale $L > 0$. Let $D > 0$, $\lambda \in \mathbb{R}$ and $f \in L^2(\mathbb{R})$ such that

$$\text{supp } \hat{f} \subset \{\xi \in \mathbb{R} : \|\xi\|^{2s} - \lambda \leq \sqrt{D}\}.$$

If $|\lambda| > \sqrt{D} + 1$, the Fourier transform \hat{f} is supported in the union of two intervals

$$\text{supp } \hat{f} \subset \left[-(\sqrt{D} + |\lambda|)^{\frac{1}{2s}}, -(-\sqrt{D} + |\lambda|)^{\frac{1}{2s}} \right] \cup \left[(-\sqrt{D} + |\lambda|)^{\frac{1}{2s}}, (\sqrt{D} + |\lambda|)^{\frac{1}{2s}} \right]$$

and

$$\left| \left[-(\sqrt{D} + |\lambda|)^{\frac{1}{2s}}, -(-\sqrt{D} + |\lambda|)^{\frac{1}{2s}} \right] \right| = \left| \left[(-\sqrt{D} + |\lambda|)^{\frac{1}{2s}}, (\sqrt{D} + |\lambda|)^{\frac{1}{2s}} \right] \right| \leq \frac{\sqrt{D}}{s},$$

since $s \geq \frac{1}{2}$ and $|\lambda| \geq \sqrt{D} + 1$. It follows from Theorem 1.7 that there exists a universal positive constant $C' > 0$ such that

$$(2.12) \quad \|f\|_{L^2(\mathbb{R})} \leq \left(\frac{C'}{\gamma}\right)^{L\frac{\sqrt{D}}{s}(\frac{C'}{\gamma})^2 + \frac{3}{2}} \|f\|_{L^2(\omega)}.$$

On the other hand, we notice that $\text{supp } \hat{f} \subset [0, (2\sqrt{D} + 1)^{\frac{1}{2s}}]$ when $|\lambda| \leq \sqrt{D} + 1$. We deduce from Theorem 1.6 that there exists a universal positive constant $C > 0$ such that

$$(2.13) \quad \|f\|_{L^2(\mathbb{R})} \leq \left(\frac{C}{\gamma}\right)^{C(1+L(2\sqrt{D}+1)^{\frac{1}{2s}})} \|f\|_{L^2(\omega)}.$$

By setting

$$d = \max \left(\left(\frac{C'}{\gamma}\right)^{L\frac{\sqrt{D}}{s}(\frac{C'}{\gamma})^2 + \frac{3}{2}}, \left(\frac{C}{\gamma}\right)^{C(1+L(2\sqrt{D}+1)^{\frac{1}{2s}})} \right)^2,$$

it follows from (2.12) and (2.13) that for all $\lambda \in \mathbb{R}$ and $f \in L^2(\mathbb{R})$ with $\text{supp } \hat{f} \subset \{\xi \in \mathbb{R} : \|\xi\|^{2s} - \lambda \leq \sqrt{D}\}$,

$$\|f\|_{L^2(\mathbb{R})} \leq \sqrt{d} \|f\|_{L^2(\omega)}.$$

Theorem 2.6 is then a direct consequence of the Hilbert uniqueness method and Proposition 1.3. \square

We deduce the following necessary and sufficient geometric condition for the exact controllability of the fractional free Schrödinger equations:

Corollary 2.7. *Let $s \geq \frac{1}{2}$ and ω be a measurable subset of \mathbb{R} . The one-dimensional fractional free Schrödinger equation*

$$\begin{cases} (\partial_t + i(-\Delta_x)^s)f(t, x) = u(t, x)\mathbb{1}_\omega(x), & x \in \mathbb{R}, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}), \end{cases}$$

is exactly controllable from ω if and only if ω is thick.

It is interesting to point out that the sharp geometric condition for the exact controllability of the free Schrödinger equation is actually the same as the one for the null-controllability of the free heat equation (1.10), even if the behavior of these two equations are very different. This thickness condition is also a necessary and sufficient condition for the null-controllability of the fractional heat equations as showed by Alphonse and Bernier [1, Remark 1.13], when the fractional parameter satisfies $s > \frac{1}{2}$.

In a recent work [5], Green, Jaye and Mitkovski prove new uncertainty principles. These results can be seen as a generalization of the Logvinenko-Sereda Theorem and its quantitative versions given by Kovrijkine. They establish in particular the following uncertainty principle:

Theorem 2.8 (Green, Jaye & Mitkovski [5, Corollary 3]). *Let ω be a measurable subset of \mathbb{R}^d satisfying the following one-dimensional geometric control condition: there exist some positive constants $L > 0$ and $0 < \gamma \leq 1$ such that for all straight lines $\mathcal{D} \subset \mathbb{R}^d$ and for all line segment $S \subset \mathcal{D}$ of length L ,*

$$\mathcal{H}(S \cap \omega) \geq \gamma L,$$

where \mathcal{H} denotes the one-dimensional Lebesgue measure of \mathcal{D} . Then, for all $\delta > 0$, $\beta > 0$, there exists $C > 0$ such that for all $R > 0$ and $f \in L^2(\mathbb{R}^d)$,

$$\text{supp } \hat{f} \subset \{\xi \in \mathbb{R}^d : R - \beta \leq |\xi| \leq R + \beta\} \implies \|f\|_{L^2(\mathbb{R}^d)} \leq C \|f\|_{L^2(\mathcal{U}_\delta(\omega))},$$

where $\mathcal{U}_\delta(\omega)$ denotes the δ -neighborhood of ω defined by $\mathcal{U}_\delta(\omega) = \{x \in \mathbb{R}^d : d(x, \omega) < \delta\}$.

This result together with Proposition 1.3 and the Hilbert uniqueness method allow to derive the following sufficient geometric condition on the control subset ω to ensure the exact controllability of the fractional free Schrödinger equation (2.9):

Proposition 2.9. *Let $d \geq 1$, $s \geq \frac{1}{2}$ and ω be a measurable subset of \mathbb{R}^d satisfying that there exist some positive constants $L > 0$ and $0 < \gamma \leq 1$ such that for all straight lines $\mathcal{D} \subset \mathbb{R}^d$ and for all line segment $S \subset \mathcal{D}$ of length L ,*

$$(2.14) \quad \mathcal{H}(S \cap \omega) \geq \gamma L,$$

where \mathcal{H} denotes the one-dimensional Lebesgue measure of \mathcal{D} . Then, for all $\delta > 0$, the fractional free Schrödinger equation

$$\begin{cases} (\partial_t + i(-\Delta_x)^s)f(t, x) = u(t, x)\mathbf{1}_\omega(x), & x \in \mathbb{R}^d, t > 0, \\ f|_{t=0} = f_0 \in L^2(\mathbb{R}^d), \end{cases}$$

is exactly controllable from the control subset $\mathcal{U}_\delta(\omega)$.

Notice that any measurable subset $\omega \subset \mathbb{R}^d$ satisfying the condition (2.14) is a thick subset in the sense of (1.7). The condition (2.14) is therefore stronger than the thickness condition.

Addendum. After the completion of this work, we find out the recent preprint [7], which points out the same necessary and sufficient geometric condition on control subsets for the exact controllability of the one-dimensional free and harmonic Schrödinger equations. The authors of [7] also derive a necessary and sufficient geometric condition for the exact controllability of the Schrödinger equation associated to the one-dimensional operator $P = -\Delta_x + V(x)$, with $V(x) = x^{2m}$, when $m \geq 2$, which is not discussed in this work. On the other hand, the fractional case is not discussed in [7] and the multidimensional results contained in this work are also not covered by the results of [7]. The proofs given in the two papers are also different. Regarding the free Schrödinger equation, the proof of [7] given only in the non-fractional case $s = 1$ makes the use of explicit computations of the Fourier transform of Gaussians which cannot be directly used in the fractional case. This proof does not make explicit the condition $s \geq \frac{1}{2}$ found out in Corollary 2.7 to obtain the necessary and sufficient geometric condition for exact controllability. Regarding the harmonic Schrödinger equation, the proof of [7] is much more involved since it is given in the general case when $m \geq 1$ with no explicit simplification in the harmonic case $m = 1$. The use of the Plancherel-Rotach formula in this work allows to obtain more directly the result in the harmonic case and accounts for the fact that the present paper is much shorter than the preprint [7].

3. PROOF OF PROPOSITION 2.1

This section is devoted to the proof of Proposition 2.1. The assertions (ii) and (iii) are clearly equivalent. Indeed, (ii) implies (iii) by passing to the limit inferior

$$\liminf_{n \rightarrow +\infty} \|\psi_n\|_{L^2(\omega)} \geq \frac{1}{c} > 0.$$

On the other hand, we readily deduce from formula (2.2) that for any measurable set $\omega \subset \mathbb{R}$ of positive measure $|\omega| > 0$,

$$(3.1) \quad \forall n \geq 0, \quad \|\psi_n\|_{L^2(\omega)} > 0,$$

since H_n is a polynomial of degree n . Then, assertion (ii) directly follows from (iii) and (3.1). It is therefore sufficient to prove the equivalence of assertions (i) and (iii).

As a preliminary step, we begin by establishing that

$$(3.2) \quad \forall 0 < \varepsilon < \frac{\pi}{2}, \quad \lim_{n \rightarrow +\infty} \|\psi_n\|_{L^2(I_{\varepsilon,n})} = \sqrt{1 - \frac{2\varepsilon}{\pi}},$$

where $I_{\varepsilon,n}$ stands for the open interval $(-\sqrt{2n+1} \cos \varepsilon, \sqrt{2n+1} \cos \varepsilon)$. Let $0 < \varepsilon < \frac{\pi}{2}$. The Plancherel-Rotach formula for Hermite polynomials [12] (Theorem 8.22.9) provides that for all $\varepsilon \leq \theta \leq \pi - \varepsilon$,

$$(3.3) \quad e^{-\frac{x^2}{2}} H_n(x) = \frac{2^{\frac{n}{2} + \frac{1}{4}} \sqrt{n!}}{(\pi n)^{\frac{1}{4}} \sqrt{\sin \theta}} \left[\sin \left(\left(\frac{n}{2} + \frac{1}{4} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{4} \right) + R_n(x) \right],$$

with $x = \sqrt{2n+1} \cos \theta$, where the remainder term satisfies

$$(3.4) \quad \exists C > 0, \forall n \geq 0, \forall y \in [-\sqrt{2n+1} \cos \varepsilon, \sqrt{2n+1} \cos \varepsilon], \quad |R_n(y)| \leq \frac{C}{n+1}.$$

It follows from (2.2) and (3.3) that the Hermite function

$$\psi_n(x) = \frac{1}{\pi^{\frac{1}{4}} 2^{\frac{n}{2}} \sqrt{n!}} e^{-\frac{x^2}{2}} H_n(x), \quad \|\psi_n\|_{L^2(\mathbb{R})} = 1, \quad n \geq 0,$$

satisfies that for all $\varepsilon \leq \theta \leq \pi - \varepsilon$ and $n \geq 1$,

$$(3.5) \quad \psi_n(x) = \frac{2^{\frac{1}{4}}}{\sqrt{\pi n^{\frac{1}{4}} \sqrt{\sin \theta}}} \left[\sin \left(\left(\frac{n}{2} + \frac{1}{4} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{4} \right) + R_n(x) \right],$$

with $x = \sqrt{2n+1} \cos \theta$. We obtain from (3.5) that for all $n \geq 1$,

$$(3.6) \quad \|F_n\|_{L^2(I_{\varepsilon,n})} - \|G_n\|_{L^2(I_{\varepsilon,n})} \leq \|\psi_n\|_{L^2(I_{\varepsilon,n})} \leq \|F_n\|_{L^2(I_{\varepsilon,n})} + \|G_n\|_{L^2(I_{\varepsilon,n})},$$

where the two functions F_n and G_n are defined for all $\varepsilon < \theta < \pi - \varepsilon$,

$$(3.7) \quad F_n(x) = \frac{2^{\frac{1}{4}}}{\sqrt{\pi n^{\frac{1}{4}} \sqrt{\sin \theta}}} \sin \left(\left(\frac{n}{2} + \frac{1}{4} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{4} \right)$$

and

$$(3.8) \quad G_n(x) = \frac{2^{\frac{1}{4}}}{\sqrt{\pi n^{\frac{1}{4}} \sqrt{\sin \theta}}} R_n(x),$$

with $x = \sqrt{2n+1} \cos \theta$. We deduce from (3.4) that for all $n \geq 1$,

$$\|G_n\|_{L^2(I_{\varepsilon,n})}^2 = \frac{\sqrt{2(2n+1)}}{\pi \sqrt{n}} \int_{\varepsilon}^{\pi - \varepsilon} |R_n(\sqrt{2n+1} \cos \theta)|^2 d\theta \leq \frac{C^2 \sqrt{2(2n+1)} (\pi - 2\varepsilon)}{\pi \sqrt{n} (n+1)^2},$$

implying that

$$(3.9) \quad \lim_{n \rightarrow +\infty} \|G_n\|_{L^2(I_{\varepsilon,n})} = 0.$$

On the other hand, we deduce from (3.7) that for all $n \geq 1$,

$$(3.10) \quad |F_n(x)|^2 = \frac{1}{\sqrt{2n\pi} \sin \theta} \left[1 - \cos \left(\left(n + \frac{1}{2} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{2} \right) \right].$$

It follows from (3.10) that for all $n \geq 1$,

$$(3.11) \quad \int_{I_{\varepsilon,n}} |F_n(x)|^2 dx = \int_{I_{\varepsilon,n}} \frac{\sqrt{2n+1}}{\sqrt{2n\pi} \sqrt{(2n+1) - x^2}} dx \\ - \frac{\sqrt{2n+1}}{\sqrt{2n\pi}} \int_{\varepsilon}^{\pi-\varepsilon} \cos \left(\left(n + \frac{1}{2} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{2} \right) d\theta,$$

since $\sin^2 \theta = 1 - \frac{x^2}{2n+1}$, when $x = \sqrt{2n+1} \cos \theta$. We have for all $n \geq 1$,

$$(3.12) \quad \int_{I_{\varepsilon,n}} \frac{\sqrt{2n+1}}{\sqrt{2n\pi} \sqrt{(2n+1) - x^2}} dx = \frac{1}{\pi} \sqrt{\frac{2n+1}{2n}} \int_{-\cos \varepsilon}^{\cos \varepsilon} \frac{dy}{\sqrt{1-y^2}} \\ = \frac{2}{\pi} \sqrt{\frac{2n+1}{2n}} \arcsin(\cos \varepsilon) = \frac{2}{\pi} \sqrt{\frac{2n+1}{2n}} \left(\frac{\pi}{2} - \arccos(\cos \varepsilon) \right) = \sqrt{\frac{2n+1}{2n}} \left(1 - \frac{2\varepsilon}{\pi} \right),$$

since

$$\forall -1 \leq x \leq 1, \quad \arccos x + \arcsin x = \frac{\pi}{2}.$$

It follows from (3.12) that

$$(3.13) \quad \lim_{n \rightarrow +\infty} \int_{I_{\varepsilon,n}} \frac{\sqrt{2n+1}}{\sqrt{2n\pi} \sqrt{(2n+1) - x^2}} dx = 1 - \frac{2\varepsilon}{\pi}.$$

On the other hand, we have for all $n \geq 1$,

$$(3.14) \quad \frac{\sqrt{2n+1}}{\sqrt{2n\pi}} \int_{\varepsilon}^{\pi-\varepsilon} \cos \left(\left(n + \frac{1}{2} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{2} \right) d\theta \\ = \frac{\sqrt{2n+1}}{\sqrt{2n\pi}} \int_{\varphi(\varepsilon)}^{\varphi(\pi-\varepsilon)} \frac{1}{\varphi'(\varphi^{-1}(t))} \cos \left(\left(n + \frac{1}{2} \right) t - \frac{3\pi}{2} \right) dt,$$

where $\varphi(\theta) = 2\theta - \sin(2\theta)$ is an increasing C^∞ -diffeomorphism from $[\varepsilon, \pi - \varepsilon]$ to

$$[\varphi(\varepsilon), \varphi(\pi - \varepsilon)] = [2\varepsilon - \sin(2\varepsilon), 2\pi - 2\varepsilon + \sin(2\varepsilon)].$$

An integration by parts shows that for all $n \geq 1$,

$$\int_{\varphi(\varepsilon)}^{\varphi(\pi-\varepsilon)} \frac{1}{\varphi'(\varphi^{-1}(t))} \cos \left(\left(n + \frac{1}{2} \right) t - \frac{3\pi}{2} \right) dt = \left[\frac{\sin \left(\left(n + \frac{1}{2} \right) t - \frac{3\pi}{2} \right)}{\varphi'(\varphi^{-1}(t)) \left(n + \frac{1}{2} \right)} \right]_{\varphi(\varepsilon)}^{\varphi(\pi-\varepsilon)} \\ - \frac{1}{n + \frac{1}{2}} \int_{\varphi(\varepsilon)}^{\varphi(\pi-\varepsilon)} \frac{d}{dt} \left(\frac{1}{\varphi'(\varphi^{-1}(t))} \right) \sin \left(\left(n + \frac{1}{2} \right) t - \frac{3\pi}{2} \right) dt,$$

implying that

$$(3.15) \quad \left| \int_{\varphi(\varepsilon)}^{\varphi(\pi-\varepsilon)} \frac{1}{\varphi'(\varphi^{-1}(t))} \cos \left(\left(n + \frac{1}{2} \right) t - \frac{3\pi}{2} \right) dt \right| \leq \frac{C_0}{n + \frac{1}{2}} (2 + 2\pi - 4\varepsilon + 2 \sin(2\varepsilon)),$$

with

$$0 < C_0 = \sup_{t \in [2\varepsilon - \sin(2\varepsilon), 2\pi - 2\varepsilon + \sin(2\varepsilon)]} \left\{ \left| \frac{1}{\varphi'(\varphi^{-1}(t))} \right| + \left| \frac{d}{dt} \left(\frac{1}{\varphi'(\varphi^{-1}(t))} \right) \right| \right\} < +\infty.$$

We deduce from (3.14) and (3.15) that

$$(3.16) \quad \lim_{n \rightarrow +\infty} \frac{\sqrt{2n+1}}{\sqrt{2n\pi}} \int_{\varepsilon}^{\pi-\varepsilon} \cos \left(\left(n + \frac{1}{2} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{2} \right) d\theta = 0.$$

It follows from (3.11), (3.13) and (3.16) that

$$(3.17) \quad \lim_{n \rightarrow +\infty} \|F_n\|_{L^2(I_{\varepsilon,n})} = \sqrt{1 - \frac{2\varepsilon}{\pi}}.$$

Thanks to (3.6), (3.9) and (3.17), we obtain that (3.2) holds.

Let ω be a measurable subset of \mathbb{R} verifying

$$(3.18) \quad \delta = \liminf_{n \rightarrow +\infty} \|\psi_n\|_{L^2(\omega)} > 0.$$

Let $0 < \varepsilon_0 < \frac{\pi}{2}$ satisfying

$$0 < \varepsilon_0 < \frac{\pi}{2} \delta^2.$$

By using that $\|\psi_n\|_{L^2(\mathbb{R})} = 1$, we observe that for all $0 < \varepsilon < \frac{\pi}{2}$ and $n \geq 0$,

$$(3.19) \quad \begin{aligned} \|\psi_n\|_{L^2(\omega)}^2 &= \|\psi_n\|_{L^2(\omega \cap I_{\varepsilon,n})}^2 + \|\psi_n\|_{L^2(\omega \cap (\mathbb{R} \setminus I_{\varepsilon,n}))}^2 \\ &\leq \|\psi_n\|_{L^2(\omega \cap I_{\varepsilon,n})}^2 + \|\psi_n\|_{L^2(\mathbb{R} \setminus I_{\varepsilon,n})}^2 = \|\psi_n\|_{L^2(\omega \cap I_{\varepsilon,n})}^2 + 1 - \|\psi_n\|_{L^2(I_{\varepsilon,n})}^2. \end{aligned}$$

It follows from (3.2), (3.18) and (3.19) that for all $0 < \varepsilon \leq \varepsilon_0$,

$$(3.20) \quad \liminf_{n \rightarrow +\infty} \|\psi_n\|_{L^2(\omega \cap I_{\varepsilon,n})} > 0.$$

On the other hand, we deduce from (3.5), (3.7) and (3.8) that for all $n \geq 1$ and $0 < \varepsilon < \frac{\pi}{2}$,

$$(3.21) \quad \|F_n\|_{L^2(\omega_{\varepsilon,n})} - \|G_n\|_{L^2(\omega_{\varepsilon,n})} \leq \|\psi_n\|_{L^2(\omega_{\varepsilon,n})} \leq \|F_n\|_{L^2(\omega_{\varepsilon,n})} + \|G_n\|_{L^2(\omega_{\varepsilon,n})},$$

with

$$(3.22) \quad \omega_{\varepsilon,n} = \omega \cap I_{\varepsilon,n}.$$

By using that $\|G_n\|_{L^2(\omega_{\varepsilon,n})} \leq \|G_n\|_{L^2(I_{\varepsilon,n})}$, it follows from (3.9) that

$$(3.23) \quad \forall 0 < \varepsilon < \frac{\pi}{2}, \quad \lim_{n \rightarrow +\infty} \|G_n\|_{L^2(\omega_{\varepsilon,n})} = 0.$$

We deduce from (3.10) and (3.22) that for all $n \geq 1$,

$$(3.24) \quad \begin{aligned} \int_{\omega_{\varepsilon_0,n}} |F_n(x)|^2 dx &\leq \int_{\omega_{\varepsilon_0,n}} \frac{2\sqrt{2n+1}}{\sqrt{2n\pi}\sqrt{(2n+1)-x^2}} dx \\ &\leq \int_{\omega_{\varepsilon_0,n}} \frac{\sqrt{2}}{\sqrt{n\pi}\sqrt{1-\cos^2\varepsilon_0}} dx \leq \frac{\sqrt{2}}{\pi \sin \varepsilon_0} \frac{|\omega \cap (-\sqrt{2n+1}, \sqrt{2n+1})|}{\sqrt{n}}. \end{aligned}$$

By using (3.20), (3.21), (3.22), (3.23) and (3.24), it follows that

$$(3.25) \quad \begin{aligned} \liminf_{n \rightarrow +\infty} \frac{|\omega \cap (-\sqrt{2n+1}, \sqrt{2n+1})|}{\sqrt{n}} &\geq \liminf_{n \rightarrow +\infty} \frac{\pi \sin \varepsilon_0}{\sqrt{2}} \int_{\omega_{\varepsilon_0,n}} |F_n(x)|^2 dx \\ &= \liminf_{n \rightarrow +\infty} \frac{\pi \sin \varepsilon_0}{\sqrt{2}} \int_{\omega_{\varepsilon_0,n}} |\psi_n(x)|^2 dx > 0. \end{aligned}$$

We readily obtain from (3.25) that

$$(3.26) \quad \liminf_{n \rightarrow +\infty} \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{\sqrt{n}} > 0.$$

Since the floor function satisfies

$$\forall R > 0, \quad [R] \leq R < [R] + 1,$$

it follows that for all $R \geq 2$,

$$(3.27) \quad \sqrt{2n+1} \leq R < \sqrt{2n+3},$$

with $n = n(R) = \lfloor \frac{R^2-1}{2} \rfloor \in \mathbb{N}$. We obtain that for all $R \geq 2$,

$$(3.28) \quad \frac{|\omega \cap [-R, R]|}{|[-R, R]|} \geq \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{2R} = \frac{\sqrt{n}}{2R} \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{\sqrt{n}}.$$

Notice that (3.27) implies that

$$(3.29) \quad \lim_{R \rightarrow +\infty} \frac{\sqrt{n}}{2R} = \frac{1}{2\sqrt{2}}.$$

It follows from (3.26), (3.28) and (3.29) that

$$\liminf_{R \rightarrow +\infty} \frac{|\omega \cap [-R, R]|}{|[-R, R]|} \geq \frac{1}{2\sqrt{2}} \liminf_{n \rightarrow +\infty} \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{\sqrt{n}} > 0.$$

It proves that assertion (ii) implies assertion (i).

Conversely, let ω be a measurable subset of \mathbb{R} verifying

$$\liminf_{R \rightarrow +\infty} \frac{|\omega \cap [-R, R]|}{|[-R, R]|} > 0$$

and define

$$\tilde{\delta} = \liminf_{n \rightarrow +\infty} \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{\sqrt{n}} > 0.$$

In order to prove (iii), it follows from (3.22) that it is actually sufficient to show that

$$\exists 0 < \varepsilon < \frac{\pi}{2}, \quad \liminf_{n \rightarrow +\infty} \|\psi_n\|_{L^2(\omega_{\varepsilon, n})} > 0.$$

According to (3.21) and (3.23), it is then sufficient to show that

$$(3.30) \quad \exists 0 < \varepsilon < \frac{\pi}{2}, \quad \liminf_{n \rightarrow +\infty} \|F_n\|_{L^2(\omega_{\varepsilon, n})} > 0.$$

While using the substitution rule with $x = \sqrt{2n+1} \cos \theta$, we deduce from (3.7) and (3.22) that for all $0 < \varepsilon < \frac{\pi}{2}$ and $n \geq 1$,

$$(3.31) \quad \int_{\omega_{\varepsilon, n}} |F_n(x)|^2 dx = \frac{1}{\pi} \sqrt{\frac{4n+2}{n}} \int_{\tilde{\omega}_{\varepsilon, n}} \sin^2 \left(\left(\frac{n}{2} + \frac{1}{4} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{4} \right) d\theta,$$

with

$$(3.32) \quad \tilde{\omega}_{\varepsilon, n} = \arccos \left(\frac{\omega_{\varepsilon, n}}{\sqrt{2n+1}} \right) \subset (\varepsilon, \pi - \varepsilon).$$

According to (3.30) and (3.31), it is then sufficient to show that

$$(3.33) \quad \exists 0 < \varepsilon < \frac{\pi}{2}, \quad \liminf_{n \rightarrow +\infty} \int_{\tilde{\omega}_{\varepsilon, n}} \sin^2 \left(\left(\frac{n}{2} + \frac{1}{4} \right) (\sin(2\theta) - 2\theta) + \frac{3\pi}{4} \right) d\theta > 0.$$

Let $0 < \varepsilon_0 < \frac{\pi}{2}$ verifying

$$0 < 2\sqrt{2}(1 - \cos \varepsilon_0) < \tilde{\delta}.$$

By noticing from (3.22) that for all $0 < \varepsilon < \frac{\pi}{2}$ and $n \geq 1$,

$$\frac{|\omega_{\varepsilon, n}|}{\sqrt{n}} \geq \frac{|\omega \cap [-\sqrt{2n+1}, \sqrt{2n+1}]|}{\sqrt{n}} - \frac{2\sqrt{2n+1}}{\sqrt{n}}(1 - \cos \varepsilon),$$

we deduce that

$$\liminf_{n \rightarrow +\infty} \frac{|\omega_{\varepsilon_0, n}|}{\sqrt{n}} \geq \tilde{\delta} - 2\sqrt{2}(1 - \cos \varepsilon_0) > 0.$$

It follows that there exist some positive constants $c_0 > 0$ and $n_0 \geq 1$ such that

$$(3.34) \quad \forall n \geq n_0, \quad |\omega_{\varepsilon_0, n}| \geq c_0 \sqrt{n}.$$

On the other hand, we notice from (3.32) and (3.34) that for all $n \geq n_0$,

$$(3.35) \quad |\tilde{\omega}_{\varepsilon_0, n}| = \int_{\omega_{\varepsilon_0, n}} |\kappa'_n(x)| dx = \int_{\omega_{\varepsilon_0, n}} \frac{dx}{\sqrt{(2n+1) - x^2}} \geq \frac{|\omega_{\varepsilon_0, n}|}{\sqrt{2n+1}} \geq c_0 \sqrt{\frac{n}{2n+1}},$$

with $\kappa_n(x) = \arccos\left(\frac{x}{\sqrt{2n+1}}\right)$. It follows from (3.35) that there exists a positive constant $c_1 > 0$ such that

$$(3.36) \quad \forall n \geq n_0, \quad |\tilde{\omega}_{\varepsilon_0, n}| \geq c_1 > 0.$$

By using anew the substitution rule with $t = 2\theta - \sin(2\theta)$, we observe that

$$(3.37) \quad \int_{\tilde{\omega}_{\varepsilon_0, n}} \sin^2\left(\left(\frac{n}{2} + \frac{1}{4}\right)(\sin(2\theta) - 2\theta) + \frac{3\pi}{4}\right) d\theta \\ = \int_{\varphi(\tilde{\omega}_{\varepsilon_0, n})} \frac{1}{\varphi'(\varphi^{-1}(t))} \sin^2\left(\left(\frac{n}{2} + \frac{1}{4}\right)t - \frac{3\pi}{4}\right) dt \geq c_2 \int_{\varphi(\tilde{\omega}_{\varepsilon_0, n})} \sin^2\left(\left(\frac{n}{2} + \frac{1}{4}\right)t - \frac{3\pi}{4}\right) dt,$$

with

$$0 < c_2 = \inf_{t \in [2\varepsilon_0 - \sin(2\varepsilon_0), 2\pi - 2\varepsilon_0 + \sin(2\varepsilon_0)]} \frac{1}{\varphi'(\varphi^{-1}(t))} < +\infty,$$

where $\varphi(\theta) = 2\theta - \sin(2\theta)$ is the increasing C^∞ -diffeomorphism from $[\varepsilon_0, \pi - \varepsilon_0]$ to $[2\varepsilon_0 - \sin(2\varepsilon_0), 2\pi - 2\varepsilon_0 + \sin(2\varepsilon_0)]$ already used above. According to (3.33) and (3.37), it therefore sufficient to prove that

$$(3.38) \quad \liminf_{n \rightarrow +\infty} \int_{\varphi(\tilde{\omega}_{\varepsilon_0, n})} \sin^2\left(\left(\frac{n}{2} + \frac{1}{4}\right)t - \frac{3\pi}{4}\right) dt > 0,$$

where according to (3.36), the subset $\varphi(\tilde{\omega}_{\varepsilon_0, n}) \subset (2\varepsilon_0 - \sin(2\varepsilon_0), 2\pi - 2\varepsilon_0 + \sin(2\varepsilon_0))$ satisfies for all $n \geq n_0$,

$$(3.39) \quad |\varphi(\tilde{\omega}_{\varepsilon_0, n})| = \int_{\tilde{\omega}_{\varepsilon_0, n}} |\varphi'(x)| dx \geq \left(\inf_{[\varepsilon_0, \pi - \varepsilon_0]} |\varphi'| \right) |\tilde{\omega}_{\varepsilon_0, n}| \geq c_3,$$

with

$$0 < c_3 = c_1 \inf_{[\varepsilon_0, \pi - \varepsilon_0]} |\varphi'| < +\infty.$$

By using anew the substitution rule with $x = \left(\frac{n}{2} + \frac{1}{4}\right)t - \frac{3\pi}{4}$, we observe that

$$(3.40) \quad \int_{\varphi(\tilde{\omega}_{\varepsilon_0, n})} \sin^2\left(\left(\frac{n}{2} + \frac{1}{4}\right)t - \frac{3\pi}{4}\right) dt = \frac{4}{2n+1} \int_{\Omega_{\varepsilon_0, n}} \sin^2 x \, dx,$$

where according to (3.32) and (3.39), the subset

$$\Omega_{\varepsilon_0, n} = \left(\frac{n}{2} + \frac{1}{4}\right)\varphi(\tilde{\omega}_{\varepsilon_0, n}) - \frac{3\pi}{4} \subset \left(\left(\frac{n}{2} + \frac{1}{4}\right)\varphi(\varepsilon_0) - \frac{3\pi}{4}, \left(\frac{n}{2} + \frac{1}{4}\right)\varphi(\pi - \varepsilon_0) - \frac{3\pi}{4}\right),$$

satisfies

$$(3.41) \quad \exists c_4 > 0, \forall n \geq n_0, \quad |\Omega_{\varepsilon_0, n}| \geq c_4 n.$$

According to (3.38) and (3.40), it is therefore sufficient to check that

$$(3.42) \quad \liminf_{n \rightarrow +\infty} \frac{1}{n} \int_{\Omega_{\varepsilon_0, n}} \sin^2 x \, dx > 0.$$

Let $k \geq 1$ be an integer depending only on ε_0 such that

$$(3.43) \quad \forall n \geq n_0, \quad \Omega_{\varepsilon_0, n} \subset \left(-kn\pi + \frac{\pi}{2}, kn\pi + \frac{\pi}{2}\right).$$

We denote $N_n = \#I_n$ the cardinality of the following set

$$(3.44) \quad I_n = \left\{ i \in [-kn, kn - 1] \cap \mathbb{Z} : \left| \Omega_{\varepsilon_0, n} \cap \left[i\pi + \frac{\pi}{2}, \pi(i+1) + \frac{\pi}{2} \right] \right| \geq \frac{c_4}{2k(\pi+1)} \right\},$$

with $n \geq n_0$. It follows from (3.43) and (3.44) that for all $n \geq n_0$,

$$\begin{aligned}
 (3.45) \quad |\Omega_{\varepsilon_0, n}| &= \sum_{i=-kn}^{kn-1} \left| \Omega_{\varepsilon_0, n} \cap \left[i\pi + \frac{\pi}{2}, (i+1)\pi + \frac{\pi}{2} \right] \right| \\
 &= \sum_{i \in I_n} \left| \Omega_{\varepsilon_0, n} \cap \left[i\pi + \frac{\pi}{2}, (i+1)\pi + \frac{\pi}{2} \right] \right| + \sum_{\substack{i \notin I_n, \\ i \in [-kn, kn-1] \cap \mathbb{Z}}} \left| \Omega_{\varepsilon_0, n} \cap \left[i\pi + \frac{\pi}{2}, (i+1)\pi + \frac{\pi}{2} \right] \right| \\
 &\leq \pi N_n + 2kn \frac{c_4}{2k(\pi+1)} = \pi N_n + \frac{nc_4}{\pi+1}.
 \end{aligned}$$

We deduce from (3.41) and (3.45) that

$$(3.46) \quad \forall n \geq n_0, \quad N_n \geq \frac{nc_4}{\pi+1}.$$

Let us now prove that the estimate (3.42) holds. By using the π -periodicity of the function $x \mapsto \sin^2 x$ and (3.43), we obtain that for all $n \geq n_0$,

$$\begin{aligned}
 (3.47) \quad \frac{1}{n} \int_{\Omega_{\varepsilon_0, n}} \sin^2 x \, dx &= \frac{1}{n} \sum_{i=-kn}^{kn-1} \int_{\Omega_{\varepsilon_0, n} \cap [i\pi + \frac{\pi}{2}, (i+1)\pi + \frac{\pi}{2}]} \sin^2 x \, dx \\
 &\geq \frac{1}{n} \sum_{i \in I_n} \int_{(\Omega_{\varepsilon_0, n} - (i+1)\pi) \cap [-\frac{\pi}{2}, \frac{\pi}{2}]} \sin^2 x \, dx.
 \end{aligned}$$

An application of Lemma 5.1 in appendix together with (3.44), (3.46) and (3.47) provide that for all $n \geq n_0$,

$$\begin{aligned}
 (3.48) \quad \frac{1}{n} \int_{\Omega_{\varepsilon_0, n}} \sin^2 x \, dx &\geq \frac{1}{n} \sum_{i \in I_n} \int_{-\frac{m_{i,n}}{2}}^{\frac{m_{i,n}}{2}} \sin^2 x \, dx \geq \frac{N_n}{n} \int_{-\frac{c_4}{4k(\pi+1)}}^{\frac{c_4}{4k(\pi+1)}} \sin^2 x \, dx \\
 &\geq \frac{c_4}{\pi+1} \int_{-\frac{c_4}{4k(\pi+1)}}^{\frac{c_4}{4k(\pi+1)}} \sin^2 x \, dx > 0,
 \end{aligned}$$

with

$$m_{i,n} = \left| (\Omega_{\varepsilon_0, n} - (i+1)\pi) \cap \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \right| = \left| \Omega_{\varepsilon_0, n} \cap \left[i\pi + \frac{\pi}{2}, (i+1)\pi + \frac{\pi}{2} \right] \right|,$$

for all $n \geq n_0$ and $i \in [-kn, kn-1] \cap \mathbb{Z}$. This establishes (3.42) and concludes that assertion (i) implies assertion (iii). This ends the proof of Proposition 2.1.

4. PROOF OF THEOREM 2.3

This section is devoted to the proof of Theorem 2.3. For any $\alpha = (\alpha_1, \dots, \alpha_d) \in (\mathbb{N} \setminus \{0\})^d$ and $\varepsilon \geq 0$, we consider the parallelepipeds

$$\mathcal{C}_{\alpha, \varepsilon} = \prod_{j=1}^d [-\sqrt{2\alpha_j + 1} \cos \varepsilon, \sqrt{2\alpha_j + 1} \cos \varepsilon].$$

Let $0 < \varepsilon < \frac{\pi}{2}$. We deduce from the one-dimensional Plancherel-Rotach formula (3.5) that for all $\alpha \in (\mathbb{N} \setminus \{0\})^d$, $0 < \varepsilon < \frac{\pi}{2}$ and $x \in \mathcal{C}_{\alpha, \varepsilon}$,

$$(4.1) \quad \Psi_{\alpha}(x) = F_{\alpha}(x) + G_{\alpha}(x),$$

with

$$(4.2) \quad F_{\alpha}(x) = \frac{2^{\frac{d}{4}}}{\pi^{\frac{d}{2}} \alpha^{\frac{1}{4}}} \prod_{i=1}^d \frac{\sin \left(\left(\frac{\alpha_i}{2} + \frac{1}{4} \right) (\sin(2\theta_i) - 2\theta_i) + \frac{3\pi}{4} \right)}{\sqrt{\sin \theta_i}}$$

and

$$(4.3) \quad G_\alpha(x) = \frac{2^{\frac{d}{4}}}{\pi^{\frac{d}{2}} \alpha^{\frac{1}{4}} \prod_{i=1}^d \sqrt{\sin \theta_i}} R_\alpha(x),$$

with $x = (\sqrt{2\alpha_1 + 1} \cos \theta_1, \dots, \sqrt{2\alpha_d + 1} \cos \theta_d) \in \mathcal{C}_{\alpha, \varepsilon}$. The remainder terms R_α satisfy

$$(4.4) \quad \exists C_\varepsilon > 0, \forall \alpha \in (\mathbb{N} \setminus \{0\})^d, \forall x \in \mathcal{C}_{\alpha, \varepsilon}, \quad |R_\alpha(x)| \leq C_\varepsilon \sum_{i=1}^d \frac{1}{\alpha_i + 1}.$$

By using the preliminary step in the proof of Proposition 2.1, we obtain from (3.2) that for all $0 < \varepsilon < \frac{\pi}{2}$,

$$(4.5) \quad \begin{aligned} & \lim_{\alpha_1 \rightarrow +\infty} \dots \lim_{\alpha_d \rightarrow +\infty} \|\Psi_\alpha\|_{L^2(\mathcal{C}_{\alpha, \varepsilon})}^2 \\ &= \lim_{\alpha_1 \rightarrow +\infty} \dots \lim_{\alpha_d \rightarrow +\infty} \prod_{j=1}^d \|\psi_{\alpha_j}\|_{L^2([- \sqrt{2\alpha_j + 1} \cos \varepsilon, \sqrt{2\alpha_j + 1} \cos \varepsilon])}^2 \\ &= \prod_{j=1}^d \lim_{\alpha_j \rightarrow +\infty} \|\psi_{\alpha_j}\|_{L^2([- \sqrt{2\alpha_j + 1} \cos \varepsilon, \sqrt{2\alpha_j + 1} \cos \varepsilon])}^2 = \left(1 - \frac{2\varepsilon}{\pi}\right)^{\frac{d}{2}}. \end{aligned}$$

Let us assume that the fractional harmonic Schrödinger equation (2.5) with $s \geq 1$ is exactly controllable from a measurable subset $\omega \subset \mathbb{R}^d$ for some time $T > 0$. The spectral gap condition in Proposition 1.8 is satisfied by the fractional harmonic oscillator $(-\Delta_x + |x|^2)^s$ since its spectrum is given by

$$\sigma((-\Delta_x + |x|^2)^s) = \{(2n + d)^s : n \in \mathbb{N}\}.$$

We can deduce from the infinite dimensional Hautus test [11, Corollary 2.18] recalled in Proposition 1.8 that there exists $\delta > 0$ such that

$$(4.6) \quad \forall \alpha \in \mathbb{N}^d, \quad \delta = \delta \|\Psi_\alpha\|_{L^2(\mathbb{R}^d)} \leq \|\Psi_\alpha\|_{L^2(\omega)}.$$

Let $0 < \varepsilon_0 < \frac{\pi}{2}$ such that

$$\delta - \left(1 - \left(1 - \frac{2\varepsilon_0}{\pi}\right)^{\frac{d}{2}}\right) > 0.$$

By writing for all $\alpha \in \mathbb{N}^d$,

$$\|\Psi_\alpha\|_{L^2(\omega)}^2 = \|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})}^2 + \|\Psi_\alpha\|_{L^2(\omega \setminus \mathcal{C}_{\alpha, \varepsilon_0})}^2 \leq \|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})}^2 + 1 - \|\Psi_\alpha\|_{L^2(\mathcal{C}_{\alpha, \varepsilon_0})}^2,$$

we deduce from (4.5) and (4.6) that

$$\liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} \|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})} \geq \sqrt{\delta^2 - \left(1 - \left(1 - \frac{2\varepsilon_0}{\pi}\right)^{\frac{d}{2}}\right)} > 0.$$

It follows from (4.3) and (4.4) that

$$\forall \alpha \in (\mathbb{N} \setminus \{0\})^d, \quad \|G_\alpha\|_{L^2(\mathcal{C}_{\alpha, \varepsilon_0})}^2 \leq C_{\varepsilon_0}^2 \frac{2^{\frac{d}{2}}}{\alpha^{\frac{1}{2}}} \left(\sum_{i=1}^d \frac{1}{\alpha_i + 1}\right)^2 \left(\prod_{i=1}^d \sqrt{2\alpha_i + 1}\right) \left(1 - \frac{2\varepsilon_0}{\pi}\right)^d$$

and

$$(4.7) \quad \limsup_{\alpha_1 \rightarrow +\infty} \dots \limsup_{\alpha_d \rightarrow +\infty} \|G_\alpha\|_{L^2(\mathcal{C}_{\alpha, \varepsilon_0})} = 0.$$

We can therefore deduce from (4.1) and (4.7) that

$$\begin{aligned}
 (4.8) \quad & \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} \|F_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})} \\
 & \geq \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} [\|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})} - \|G_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})}] \\
 & \geq \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} [\|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})} - \|G_\alpha\|_{L^2(\mathcal{C}_{\alpha, \varepsilon_0})}] \\
 & \geq \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} \|\Psi_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})} > 0.
 \end{aligned}$$

On the other hand, we notice from (4.2) that for all $\alpha \in (\mathbb{N} \setminus \{0\})^d$,

$$\forall x \in \mathcal{C}_{\alpha, \varepsilon_0}, \quad |F_\alpha(x)|^2 \leq \frac{2^{\frac{d}{2}}}{\pi^d \alpha^{\frac{1}{2}}} \prod_{i=1}^d \frac{\sqrt{2\alpha_i + 1}}{\sqrt{2\alpha_i + 1 - x_i^2}}$$

and this implies that for all $\alpha \in (\mathbb{N} \setminus \{0\})^d$,

$$\begin{aligned}
 (4.9) \quad \|F_\alpha\|_{L^2(\omega \cap \mathcal{C}_{\alpha, \varepsilon_0})}^2 & \leq \frac{2^{\frac{d}{2}}}{\pi^d \alpha^{\frac{1}{2}}} \int_{\omega \cap \mathcal{C}_{\alpha, \varepsilon_0}} \prod_{i=1}^d \frac{\sqrt{2\alpha_i + 1}}{\sqrt{2\alpha_i + 1 - x_i^2}} dx \\
 & \leq \frac{2^{\frac{d}{2}}}{\pi^d \sin^d \varepsilon_0} \frac{|\omega \cap [-\sqrt{2\alpha_1 + 1}, \sqrt{2\alpha_1 + 1}] \times \dots \times [-\sqrt{2\alpha_d + 1}, \sqrt{2\alpha_d + 1}]|}{\alpha^{\frac{1}{2}}}.
 \end{aligned}$$

We deduce from (4.8) and (4.9) that

$$(4.10) \quad \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} \frac{|\omega \cap [-\sqrt{2\alpha_1 + 1}, \sqrt{2\alpha_1 + 1}] \times \dots \times [-\sqrt{2\alpha_d + 1}, \sqrt{2\alpha_d + 1}]|}{|[-\sqrt{2\alpha_1 + 1}, \sqrt{2\alpha_1 + 1}] \times \dots \times [-\sqrt{2\alpha_d + 1}, \sqrt{2\alpha_d + 1}]|} > 0.$$

It follows from (3.27) that for all $(R_1, \dots, R_d) \in [2, +\infty)^d$,

$$(4.11) \quad \forall 1 \leq i \leq d, \quad \sqrt{2n_i + 1} \leq R_i < \sqrt{2n_i + 3},$$

where $n_i = n_i(R_i) = \lfloor \frac{R_i^2 - 1}{2} \rfloor \in \mathbb{N}$ for all $1 \leq i \leq d$. We obtain that

$$\begin{aligned}
 (4.12) \quad & \frac{|\omega \cap [-R_1, R_1] \times \dots \times [-R_d, R_d]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} \\
 & \geq \frac{|\omega \cap [-\sqrt{2n_1 + 1}, \sqrt{2n_1 + 1}] \times \dots \times [-\sqrt{2n_d + 1}, \sqrt{2n_d + 1}]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} \\
 & = \left(\prod_{i=1}^d \frac{\sqrt{2n_i + 1}}{R_i} \right) \frac{|\omega \cap [-\sqrt{2n_1 + 1}, \sqrt{2n_1 + 1}] \times \dots \times [-\sqrt{2n_d + 1}, \sqrt{2n_d + 1}]|}{|[-\sqrt{2n_1 + 1}, \sqrt{2n_1 + 1}] \times \dots \times [-\sqrt{2n_d + 1}, \sqrt{2n_d + 1}]|}.
 \end{aligned}$$

We observe that (4.11) implies that

$$(4.13) \quad \forall 1 \leq i \leq d, \quad \lim_{R_i \rightarrow +\infty} \frac{\sqrt{2n_i + 1}}{R_i} = 1.$$

It follows from (4.10), (4.12) and (4.13) that

$$\begin{aligned}
 & \liminf_{R_1 \rightarrow +\infty} \dots \liminf_{R_d \rightarrow +\infty} \frac{|\omega \cap [-R_1, R_1] \times \dots \times [-R_d, R_d]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} \\
 & \geq \liminf_{\alpha_1 \rightarrow +\infty} \dots \liminf_{\alpha_d \rightarrow +\infty} \frac{|\omega \cap [-\sqrt{2\alpha_1 + 1}, \sqrt{2\alpha_1 + 1}] \times \dots \times [-\sqrt{2\alpha_d + 1}, \sqrt{2\alpha_d + 1}]|}{|[-\sqrt{2\alpha_1 + 1}, \sqrt{2\alpha_1 + 1}] \times \dots \times [-\sqrt{2\alpha_d + 1}, \sqrt{2\alpha_d + 1}]|} > 0.
 \end{aligned}$$

It establishes (2.6) with $A = I_d$. Let $A \in O(\mathbb{R}^d)$. By using the invariance of the Weyl quantization under affine symplectic transformations (see e.g. [6, Theorem 18.5.9]), we notice that

$$(4.14) \quad R_A \circ \mathcal{H} = \mathcal{H} \circ R_A,$$

with $\mathcal{H} = -\Delta_x + |x|^2$, where R_A is the unitary operator on $L^2(\mathbb{R}^d)$ defined by $R_A f = f(A \cdot)$ for $f \in L^2(\mathbb{R}^d)$. The identity (4.14) implies in particular that $(R_A \Psi_\alpha)_{\alpha \in \mathbb{N}^d}$ is a Hilbert basis of $L^2(\mathbb{R}^d)$ satisfying

$$\forall \alpha \in \mathbb{N}^d, \quad \mathcal{H}(R_A \Psi_\alpha) = (2|\alpha| + d)R_A \Psi_\alpha,$$

when $A \in O(\mathbb{R}^d)$. By using that the operator

$$R_A : \text{Span}((\Psi_\alpha)_{\alpha \in \mathbb{N}^d, |\alpha|=N}) \rightarrow \text{Span}((\Psi_\alpha)_{\alpha \in \mathbb{N}^d, |\alpha|=N}),$$

is a unitary operator for any $N \in \mathbb{N}$, we notice that for all $s \geq 1$ and $f \in \mathcal{D}(\mathcal{H}^s)$,

$$\begin{aligned} (R_A \circ \mathcal{H}^s)f &= (R_A \circ \mathcal{H}^s) \left(\sum_{\alpha \in \mathbb{N}^d} \langle f, R_{A^{-1}} \Psi_\alpha \rangle_{L^2(\mathbb{R}^d)} R_{A^{-1}} \Psi_\alpha \right) \\ &= R_A \left(\sum_{\alpha \in \mathbb{N}^d} (2|\alpha| + d)^s \langle f, R_{A^{-1}} \Psi_\alpha \rangle_{L^2(\mathbb{R}^d)} R_{A^{-1}} \Psi_\alpha \right) \\ &= \sum_{\alpha \in \mathbb{N}^d} (2|\alpha| + d)^s \langle R_A f, \Psi_\alpha \rangle_{L^2(\mathbb{R}^d)} \Psi_\alpha = (\mathcal{H}^s \circ R_A)f, \end{aligned}$$

that is

$$(4.15) \quad \forall A \in O(\mathbb{R}^d), \forall s \geq 1, \quad R_A \circ \mathcal{H}^s = \mathcal{H}^s \circ R_A.$$

Let $f_0, f_T \in L^2(\mathbb{R}^d)$. By assumption, there exists a control function $u \in L^2([0, T] \times \mathbb{R}^d)$ supported in $\mathbb{R}_+ \times \omega$ such that the semigroup solution $g : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ of the system (2.5) associated to the initial condition $f_0(A \cdot)$ satisfies $g(T, \cdot) = f_T(A \cdot)$. We deduce from (4.15) that the function f defined on $\mathbb{R}_+ \times \mathbb{R}^d$ by

$$\forall t \in \mathbb{R}_+, \forall x \in \mathbb{R}^d, \quad f(t, x) = g(t, A^{-1}x),$$

is the semigroup solution of (2.5) associated to the initial condition f_0 and the control function $v(t, x) = u(t, A^{-1}x) \in L^2([0, T] \times \mathbb{R}^d)$ supported in $\mathbb{R}_+ \times A(\omega)$; and satisfies $f(T, \cdot) = f_T(\cdot)$. It implies that the harmonic Schrödinger equation (2.5) is exactly controllable from the control set $A(\omega)$. By using the first part of this proof with ω replaced by $A(\omega)$, we obtain that

$$\liminf_{R_1 \rightarrow +\infty} \dots \liminf_{R_d \rightarrow +\infty} \frac{|A(\omega) \cap [-R_1, R_1] \times \dots \times [-R_d, R_d]|}{|[-R_1, R_1] \times \dots \times [-R_d, R_d]|} > 0.$$

This ends the proof of Theorem 2.3.

5. APPENDIX

This appendix contains the proof of two instrumental results. The first part is devoted to the proof of a technical lemma used in the proof of Proposition 2.1. The purpose of the second part is to recall the proof of Miller's result stated in Proposition 1.3.

5.1. A technical lemma. The following technical result is used in the proof of Proposition 2.1:

Lemma 5.1. *If A is a measurable subset of $[-\frac{\pi}{2}, \frac{\pi}{2}]$ and $\delta = |A|$ denotes its Lebesgue measure then*

$$\int_A \sin^2 x \, dx \geq \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \sin^2 x \, dx.$$

Proof. By using that the function $x \mapsto \sin^2 x$ is even and increasing on $[0, \frac{\pi}{2}]$, we observe that

$$\begin{aligned}
 (5.1) \quad & \int_A \sin^2 x \, dx = \int_{A \cap [-\frac{\delta}{2}, \frac{\delta}{2}]} \sin^2 x \, dx + \int_{A \cap ([-\frac{\pi}{2}, \frac{\pi}{2}] \setminus [-\frac{\delta}{2}, \frac{\delta}{2}])} \sin^2 x \, dx \\
 & \geq \int_{A \cap [-\frac{\delta}{2}, \frac{\delta}{2}]} \sin^2 x \, dx + \left| A \cap \left(\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \setminus \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \right) \right| \sin^2 \left(\frac{\delta}{2} \right) \\
 & \geq \int_{A \cap [-\frac{\delta}{2}, \frac{\delta}{2}]} \sin^2 x \, dx + \frac{|A \cap ([-\frac{\pi}{2}, \frac{\pi}{2}] \setminus [-\frac{\delta}{2}, \frac{\delta}{2}])|}{|[-\frac{\delta}{2}, \frac{\delta}{2}] \cap ([-\frac{\pi}{2}, \frac{\pi}{2}] \setminus A)|} \int_{[-\frac{\delta}{2}, \frac{\delta}{2}] \cap ([-\frac{\pi}{2}, \frac{\pi}{2}] \setminus A)} \sin^2 x \, dx.
 \end{aligned}$$

By using that $\delta = |A| = |[-\frac{\delta}{2}, \frac{\delta}{2}]|$, we notice that

$$\begin{aligned}
 (5.2) \quad & \left| A \cap \left(\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \setminus \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \right) \right| = |A| - \left| A \cap \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \right| \\
 & = \left| \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \right| - \left| A \cap \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \right| = \left| \left[-\frac{\delta}{2}, \frac{\delta}{2} \right] \cap \left(\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \setminus A \right) \right|.
 \end{aligned}$$

It follows from (5.1) and (5.2) that

$$\int_A \sin^2 x \, dx \geq \int_{A \cap [-\frac{\delta}{2}, \frac{\delta}{2}]} \sin^2 x \, dx + \int_{[-\frac{\delta}{2}, \frac{\delta}{2}] \cap ([-\frac{\pi}{2}, \frac{\pi}{2}] \setminus A)} \sin^2 x \, dx = \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \sin^2 x \, dx.$$

This ends the proof of Lemma 5.1. \square

5.2. Spectral inequalities and exact controllability. This section is devoted to recall the proof of Miller's result [11, Corollary 2.17] stated in Proposition 1.3 which provides necessary and sufficient spectral estimates for the observability of system (1.3) to hold. The proof of Proposition 1.3 is based on another Miller's result [11, Theorem 2.4] characterizing the observability with resolvent conditions which is recalled below and whose proof is omitted:

Proposition 5.2 (Miller [11, Theorem 2.4]). *Let $(A, \mathcal{D}(A))$ be a selfadjoint operator on $L^2(\mathbb{R}^d)$, which is the infinitesimal generator of a strongly continuous group $(e^{itA})_{t \in \mathbb{R}}$ on $L^2(\mathbb{R}^d)$. The system (1.3) is exactly observable from a measurable subset $\omega \subset \mathbb{R}^d$ if and only if there exist some positive constants $M > 0$ and $m > 0$ such that*

$$(5.3) \quad \forall f \in \mathcal{D}(A), \forall \lambda \in \mathbb{R}, \quad \|f\|_{L^2(\mathbb{R}^d)}^2 \leq M \|(A - \lambda)f\|_{L^2(\mathbb{R}^d)}^2 + m \|f\|_{L^2(\omega)}^2.$$

When condition (5.3) is satisfied, exact observability holds in any time $T > \pi\sqrt{M}$.

We consider the system (1.3). If this system is exactly observable from a measurable subset $\omega \subset \mathbb{R}^d$ at time $T > 0$, Proposition 5.2 proves that there exist some positive constants $M > 0$ and $m > 0$ such that the resolvent estimate (5.3) holds. Let $\lambda \in \mathbb{R}$, $0 < D < \frac{1}{M}$ and $f \in \mathbf{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(\mathcal{D}(A))$. The functional calculus shows that

$$(5.4) \quad \|(A - \lambda)f\|_{L^2(\mathbb{R}^d)} \leq \sqrt{D} \|f\|_{L^2(\mathbb{R}^d)}.$$

It follows from (5.3) and (5.4) that

$$(5.5) \quad \forall \lambda \in \mathbb{R}, \forall f \in \mathbf{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(\mathcal{D}(A)), \quad \|f\|_{L^2(\mathbb{R}^d)}^2 \leq MD \|f\|_{L^2(\mathbb{R}^d)}^2 + m \|f\|_{L^2(\omega)}^2,$$

that is

$$\forall \lambda \in \mathbb{R}, \forall f \in \mathbf{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(\mathcal{D}(A)), \quad \|f\|_{L^2(\mathbb{R}^d)}^2 \leq \frac{m}{1 - MD} \|f\|_{L^2(\omega)}^2.$$

It establishes the spectral estimates (1.4) for all $f \in \mathbf{1}_{\{|A-\lambda| \leq \sqrt{D}\}}(L^2(\mathbb{R}^d))$ since the domain $\mathcal{D}(A)$ is dense in $L^2(\mathbb{R}^d)$.

Conversely, let us assume that there exist some positive constants $D > 0$ and $k > 0$ such that the spectral estimates (1.4) holds. Let $\lambda \in \mathbb{R}$ and $f \in \mathcal{D}(A)$. With $f_\lambda = \mathbb{1}_{\{|A-\lambda| \leq \sqrt{D}\}} f$ and $f_\perp = f - f_\lambda$, we deduce from (1.4) and the functional calculus that for all $\lambda \in \mathbb{R}$ and $f \in \mathcal{D}(A)$,

$$(5.6) \quad \|f\|_{L^2(\mathbb{R}^d)}^2 = \|f_\lambda\|_{L^2(\mathbb{R}^d)}^2 + \|f_\perp\|_{L^2(\mathbb{R}^d)}^2 \leq k\|f_\lambda\|_{L^2(\omega)}^2 + \frac{1}{D}\|(A-\lambda)f_\perp\|_{L^2(\mathbb{R}^d)}^2 \\ \leq k\|f_\lambda\|_{L^2(\omega)}^2 + \frac{1}{D}\|(A-\lambda)f\|_{L^2(\mathbb{R}^d)}^2.$$

Let $T > \pi\sqrt{\frac{1+k}{D}}$ and $\varepsilon > 0$ such that

$$T^2 > \pi^2 \frac{1 + (1 + \varepsilon^2)k}{D}.$$

The functional calculus shows that for all $\lambda \in \mathbb{R}$ and $f \in \mathcal{D}(A)$,

$$(5.7) \quad \|f_\lambda\|_{L^2(\omega)}^2 = \|f - f_\perp\|_{L^2(\omega)}^2 \leq (1 + \varepsilon^{-2})\|f\|_{L^2(\omega)}^2 + (1 + \varepsilon^2)\|f_\perp\|_{L^2(\omega)}^2 \\ \leq (1 + \varepsilon^{-2})\|f\|_{L^2(\omega)}^2 + (1 + \varepsilon^2)\|f_\perp\|_{L^2(\mathbb{R}^d)}^2 \leq (1 + \varepsilon^{-2})\|f\|_{L^2(\omega)}^2 + \frac{1 + \varepsilon^2}{D}\|(A-\lambda)f_\perp\|_{L^2(\mathbb{R}^d)}^2 \\ \leq (1 + \varepsilon^{-2})\|f\|_{L^2(\omega)}^2 + \frac{1 + \varepsilon^2}{D}\|(A-\lambda)f\|_{L^2(\mathbb{R}^d)}^2.$$

It follows from (5.6) and (5.7) that

$$\forall \lambda \in \mathbb{R}, \forall f \in \mathcal{D}(A), \quad \|f\|_{L^2(\mathbb{R}^d)}^2 \leq k(1 + \varepsilon^{-2})\|f\|_{L^2(\omega)}^2 + \frac{1 + k(1 + \varepsilon^2)}{D}\|(A-\lambda)f\|_{L^2(\mathbb{R}^d)}^2.$$

We can deduce from Proposition 5.2 that the system (1.3) is exactly observable from ω at time T .

REFERENCES

- [1] P. Alphonse, J. Bernier, *Smoothing properties of fractional Ornstein-Uhlenbeck semigroups and null-controllability*, preprint (2018), arXiv:1810.02629
- [2] K. Beauchard, P. Jaming, K. Pravda-Starov, *Spectral inequality for finite combinations of Hermite functions and null-controllability of hypoelliptic quadratic equations*, preprint (2018), arXiv:1804.04895
- [3] J.-M. Coron, *Control and nonlinearity*, Mathematical Surveys and Monographs 136, AMS, Providence, RI (2007)
- [4] M. Egidi, I. Veselić, *Sharp geometric condition for null-controllability of the heat equation on \mathbb{R}^d and consistent estimates on the control cost*, Arch. Math. (Basel) 111 (2018), no. 1, 85-99
- [5] W. Green, B. Jaye, M. Mitkovski, *Uncertainty principles associated to sets satisfying the geometric control condition*, preprint (2019), arXiv:1912.05077
- [6] L. Hörmander, *The analysis of linear partial differential Operators. III. Pseudo-differential operators*, Reprint of the 1994 edition. Classics in Mathematics. Springer, Berlin (2007)
- [7] S. Huang, G. Wang, M. Wang, *Observable sets, potentials and Schrödinger equations*, preprint (2020), arXiv:2003.11263
- [8] O. Kovrijkine, *Some results related to the Logvinenko-Sereda Theorem*, Proc. Amer. Math. Soc. 129, (2001), no. 10, 3037-3047
- [9] J.-L. Lions, *Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués*, Tome 1 et 2, vol. 8, Recherches en Mathématiques Appliquées, Masson, Paris (1988)
- [10] V.N. Logvinenko, J.F. Sereda, *Equivalent norms in spaces of entire functions of exponential type*, Teor. Funkcii Funkcional. Anal. i Priložen. Vyp. 20 (1974), 102-111, 175
- [11] L. Miller, *Resolvent conditions for the control of unitary groups and their approximations*, J. Spectr. Theory 2 (2012), no. 1, 1-55
- [12] G. Szegő, *Orthogonal polynomials*, Fourth edition, American Mathematical Society, Colloquium Publications, Vol. XXIII. American Mathematical Society, Providence, R.I. (1975)
- [13] G. Wang, M. Wang, C. Zhang, Y. Zhang, *Observable set, observability, interpolation inequality and spectral inequality for the heat equation in \mathbb{R}^d* , J. Math. Pures Appl. 126 (2019), 144-194

JÉRÉMY MARTIN, UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE
E-mail address: `jeremy.martin@univ-rennes1.fr`

KAREL PRAVDA-STAROV, UNIV RENNES, CNRS, IRMAR - UMR 6625, F-35000 RENNES, FRANCE
E-mail address: `karel.pravda-starov@univ-rennes1.fr`