



HAL
open science

Entropy solutions in BV^s for a class of triangular systems involving a transport equation

Christian Bourdarias, Anupam Pal Choudhury, Billel Guelmame, Stéphane Junca

► **To cite this version:**

Christian Bourdarias, Anupam Pal Choudhury, Billel Guelmame, Stéphane Junca. Entropy solutions in BV^s for a class of triangular systems involving a transport equation. *SIAM Journal on Mathematical Analysis*, 2022, 54 (1), pp.791-817. 10.1137/20M1351783 . hal-02895603v3

HAL Id: hal-02895603

<https://hal.science/hal-02895603v3>

Submitted on 21 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Entropy solutions in BV^s for a class of triangular systems involving a transport equation

Christian Bourdarias ^{*}, Anupam Pal Choudhury [†],
Billel Guelmame [‡], Stéphane Junca [§]

ABSTRACT. – In this article, we consider a class of strictly hyperbolic triangular systems involving a transport equation. Such systems are known to create measure solutions for the initial value problem. Adding a stronger transversality assumption on the fields, we are able to obtain solutions in L^∞ under optimal fractional BV regularity of the initial data. Our results show that the critical fractional regularity is $s = 1/3$. We also construct an initial data that is not in $BV^{1/3}$ but for which a blow-up in L^∞ occurs, proving the optimality of our results.

AMS Classification: 35L65, 35L67, 35Q35, 26A45, 76N10.

Key words: conservation laws; triangular systems; entropy solution; fractional BV spaces; linear transport; chromatography; Cauchy problem; non convex flux; regularity.

Contents

1	Introduction	2
1.1	The hyperbolic triangular system	3
1.2	Weak and entropy solutions	4
1.3	BV^s functions	5
2	Main results	6
3	The uniformly strictly hyperbolic system	8
3.1	Riemann invariants	8
3.2	The Riemann problem	9
4	The Lax curves	12
4.1	The Lax cubic estimate on the Rankine-Hugoniot curve	13
4.2	Cubic estimates for the Riemann problem	15

^{*}Université Savoie-Mont Blanc, LAMA, Chambéry, France, bourdarias@univ-savoie.fr

[†]School of Mathematical Sciences, National Institute of Science Education and Research, Bhubaneswar 752050, India, and Homi Bhabha National Institute, Training School Complex, Anushaktinagar, Mumbai 400094, India, anupampcmath@gmail.com

[‡]Université Côte d’Azur, Inria & CNRS, LJAD, Nice, France, Billel.Guelmame@univ-cotedazur.fr

[§]Université Côte d’Azur, Inria & CNRS, LJAD, Nice, France, Stephane.Junca@univ-cotedazur.fr

5	Existence in BV^s	16
5.1	The Wave Front Tracking algorithm	17
5.2	Uniform estimates	20
5.3	Passage to the limit in the weak formulation	24
6	Blow-up of an entropy solution with $u_0 \notin BV^{1/3}$	25
A	On Lipschitz test functions	28
B	Uniqueness of Riemann problem in L^∞	28

1 Introduction

We consider triangular systems of the form, $t > 0$, $x \in \mathbb{R}$,

$$\partial_t u + \partial_x f(u) = 0, \tag{1.1}$$

$$\partial_t v + \partial_x (a(u)v) = 0. \tag{1.2}$$

Here f is the scalar flux function (which we shall henceforth refer to as the flux) for the equation (1.1) and the function a denotes the velocity of the linear equation with respect to v (1.2). The system has a decoupled nonlinear conservation law and a coupled “linear” transport equation with a discontinuous velocity. The above system is complemented by a set of initial data, $x \in \mathbb{R}$,

$$u(x, 0) = u_0(x), \tag{1.3}$$

$$v(x, 0) = v_0(x). \tag{1.4}$$

The Pressure Swing Adsorption process (PSA) in chemistry [9] has such a triangular structure after a change of variables from Euler to Lagrange [12, 43]. Such systems are already of mathematical interest due to the coupling of the theory of scalar conservation laws with the theory of transport equations. This system was studied in [35] in a non hyperbolic setting, $f' = a$, with measure solutions for v . That even a strictly hyperbolic setting is not enough to avoid measure solutions was shown in [21, 25]. Here, strengthening the hyperbolicity of the system by a stronger transversality condition, global weak bounded entropy solutions are provided with an optimal fractional BV regularity for the initial data u_0 .

At the first sight, such a system seems easy to solve in the “triangular” manner, that is solving the first equation (1.1) to get u and then solving the linear equation (1.2) keeping u fixed. This method works well for smooth solutions [23], but, when a shock wave appears in u the velocity $a(u)$ becomes discontinuous. The theory of linear transport equations with discontinuous velocity is a delicate topic yielding measure solutions and a loss of uniqueness [5, 6, 44].

In this paper, we propose a different approach to obtain global weak solutions in L^∞ . A main idea is to consider the system (1.1), (1.2) not as a triangular system, but, as a 2×2 hyperbolic system as in [3]. If f is nonlinear, one field is nonlinear and the other one is linearly degenerate. When f is uniformly convex it is a particular case of 2×2

systems with one genuinely nonlinear field and a linearly degenerate one [26]. An example of such a system when the flux f is piecewise convex or concave arises from gas-liquid chromatography [12]. The presence of linearly degenerate fields can simplify the study of solutions [42, 43], but can also produce blow-up behaviours [12, 40].

We show that the behaviour of the proposed entropy solutions is linked to the fractional BV regularity of u , indeed, $u_0 \in BV^s$, $0 < s \leq 1$. The BV^s framework is optimal to study the regularity of entropy solutions of scalar conservation laws [16, 37]. This framework is recalled in Section 1.3. For the scalar case, the theory works well for all $s > 0$. For the triangular system (1.1), (1.2), we prove that the regularity $s = 1/3$ of u_0 is critical for the existence of L^∞ entropy solutions. The exponent $s = 1/3$ is directly linked to the cubic estimate on the Lax curves [33]. For nonconvex fields, it is known that the Lax curves are less regular [4, 2, 36]. However, for the triangular system (1.1), (1.2) with the flux $f \in C^4$, the velocity $a \in C^3$, and satisfying a uniformly strict hyperbolicity assumption, we prove that the exponent $s = 1/3$ is optimal for the existence theory of bounded entropy solutions. Our proof is based on a new cubic estimate that holds even when the flux f is non-convex. Our estimate generalizes the well known cubic estimates for genuinely nonlinear (convex) systems by Lax [33].

The paper is organized as follows. The hyperbolicity of the triangular system and the key assumptions are given in Section 1.1. The definitions of weak and entropy solutions are stated in Section 1.2. The BV^s framework is recalled in Section 1.3. In Section 2, the two main results are stated: existence for $s \geq 1/3$ and blow-up for $s < 1/3$. In Section 3, the Riemann invariants and the Riemann problem are studied. The Lax curves and the key cubic estimates are studied in Section 4. The proof of existence is discussed in Section 5 and the blow-up is studied in Section 6. In the appendices A and B, we discuss on the uniqueness of the Riemann problem in the class of bounded weak entropy solutions.

1.1 The hyperbolic triangular system

The system (1.1), (1.2) of conservation laws is hyperbolic when $f' \neq a$ and can be rewritten, using the vectorial flux \mathbf{F} , as $\partial_t \mathbf{U} + \partial_x \mathbf{F}(\mathbf{U}) = 0$, $\mathbf{U} = (u, v)^\top$. The matrix of the linearized system has a triangular structure,

$$D\mathbf{F}(\mathbf{U}) = \begin{pmatrix} f'(u) & 0 \\ a'(u)v & a(u) \end{pmatrix}.$$

This system has the unbounded invariant region $[-M, M]_u \times \mathbb{R}_v$, where $M = \|u_0\|_\infty$. In this paper, the system is assumed to be strictly hyperbolic through the condition

$$\forall u \in [-M, M], \quad f'(u) > a(u). \quad (1.5)$$

Of course, the symmetric assumption: $\forall u \in [-M, M]$, $f'(u) < a(u)$, yields a similar study. For large data, the strict hyperbolicity condition (1.5) has to be strengthened on the set $[-M, M]$, by the following uniformly strict hyperbolicity (USH) condition:

$$\inf_{|u| \leq M} f'(u) > \sup_{|u| \leq M} a(u). \quad (1.6)$$

An interesting case is already when f is convex, $f'' > 0$ everywhere, or concave $f'' < 0$. This case occurs for a chromatography system with a convex isotherm written in appropriate Lagrangian coordinates [12].

In this paper, the flux $f(\cdot)$ belongs to C^4 and the velocity $a(\cdot)$ belongs to C^3 . Few times in the paper, to present the nonconvex case, the flux is assumed to have a finite number of inflection points,

$$\mathcal{Z} = \{u, f''(u) = 0, |u| \leq M\} \text{ is finite.} \quad (1.7)$$

Thus, Riemann problems are solved by a finite number of elementary waves, the so called composite wave [27]. The finiteness of the number of inflection points is not mandatory for the existence Theorem 2.1 below, because the proof uses approximate piecewise linear flux f_ν . The important point is that f_ν is also locally piecewise convex or concave so the solution of the Riemann problem has only a finite number of wavefronts [19].

1.2 Weak and entropy solutions

A weak solution of the system (1.1)-(1.2) satisfying the initial conditions (1.3)-(1.4) is defined as follows.

Definition 1.1 (weak solution). *The pair (u, v) is a weak solution of the system (1.1)-(1.2) with initial data (1.3)-(1.4) if for all compactly supported test functions $\varphi, \psi \in C_c^1(\mathbb{R} \times [0, +\infty[, \mathbb{R})$, the following integral identities hold:*

$$\int_0^{+\infty} \int_{\mathbb{R}} (u \partial_t \varphi + f(u) \partial_x \varphi) dx dt + \int_{\mathbb{R}} u_0(x) \varphi(x, 0) dx = 0, \quad (1.8)$$

$$\int_0^{+\infty} \int_{\mathbb{R}} (v \partial_t \psi + a(u) v \partial_x \psi) dx dt + \int_{\mathbb{R}} v_0(x) \psi(x, 0) dx = 0. \quad (1.9)$$

The following regularity is usually required for (u, v) : $u \in L_{loc}^\infty$, $v \in L_{loc}^1$ or v is a measure. In the case when v is a measure, there are some issues in defining the product $a(u) \times v$ [5]. But our main focus in this paper is on bounded weak solutions $u, v \in L^\infty$.

We propose below a notion of entropy solutions for the system (1.1)-(1.2). As in [8, 30] the entropy condition is only tested on the nonlinear component u . Contact discontinuity waves linked to a linearly degenerate field are well known not to affect the entropy inequality which, therefore, remains an equality [20].

Definition 1.2 (entropy solution). *The pair (u, v) is an entropy solution of the system (1.1)-(1.2) with initial data (1.3)-(1.4) if it is a weak solution and for all convex function η and all non-negative test functions $\varphi \in C_c^1(\mathbb{R} \times]0, +\infty[, \mathbb{R})$, with $q' = \eta' f'$, u satisfies the following inequality:*

$$\int_0^{+\infty} \int_{\mathbb{R}} (\eta(u) \partial_t \varphi + q(u) \partial_x \varphi) dx dt + \int_{\mathbb{R}} \eta(u_0(x)) \varphi(x, 0) dx \geq 0. \quad (1.10)$$

Thus, it suffices only to have a weak solution of the system and the entropy solution of (1.1).

Uniqueness of entropy solutions with a fixed initial data (u_0, v_0) is a delicate matter [14]. It is due to the lack of uniqueness of weak solutions for the linear transport equation with a discontinuous velocity ([5, 6]).

For triangular systems as discussed in this paper, one can have measure solutions. A prototypical example of such triangular systems [25] is given by

$$\begin{aligned} \partial_t u + \partial_x \left[\frac{u^2}{2} \right] &= 0, \\ \partial_t v + \partial_x [(u-1)v] &= 0, \end{aligned} \tag{1.11}$$

with $f(u) = \frac{u^2}{2}$ and $a(u) = u - 1$. Clearly this satisfies the strict hyperbolicity condition (1.5) but not the more restrictive uniform strict hyperbolicity condition (1.6) for too large data. Similar phenomenon is well known for transport equations [5] and is also widely observed in the case of non-strictly hyperbolic systems [6].

The main approach in solving the system (1.11) in [25] was to observe that the first equation in u can be solved independently. The second equation can then be considered as a transport equation in v with a discontinuous coefficient u (see also [5]), but this idea of looking at the two equations separately might not be a good one. Our aim in this article is to prove the existence of solutions to systems of type (1.11) in the class of fractional BV functions and hence one need not appeal to δ -shock wave type solutions. Moreover, to obtain L^∞ solution with a wave front algorithm, we suppose that the system (1.1), (1.2) is uniformly strictly hyperbolic (1.6). Note that we already need the uniform strict hyperbolicity condition (1.6) to solve the Riemann problem appropriately.

1.3 BV^s functions

For one dimensional scalar conservation laws, the spaces BV^s are known to give optimal results for weak entropy solutions [11, 16, 37, 38], first on the fractional Sobolev regularity, and second on the structure of such solutions. Such results on the maximal regularity have been extended for some systems and the multidimensional case [17, 23].

In this section, basic facts on BV^s functions are recalled.

Definition 1.3. [39] *A function u is said to be in $BV^s(\mathbb{R})$ with $0 < s \leq 1$ if $TV^s u < +\infty$, where*

$$TV^s u := \sup_{n \in \mathbb{N}, x_1 < \dots < x_n} \sum_{i=1}^n |u(x_{i+1}) - u(x_i)|^{1/s}.$$

The BV^s semi-norm is defined by

$$|u|_{BV^s} := (TV^s u)^s,$$

and a norm on this space is defined by

$$\|u\|_{BV^s} := \|u\|_{L^\infty} + |u|_{BV^s}.$$

Fractional BV functions have traces like BV functions. This is a fundamental property to define the Rankine-Hugoniot condition for shock waves. Moreover, this property is not true for the Sobolev functions in $W^{s,1/s}$, the Sobolev space nearest to BV^s [11].

Theorem 1.1. [39] *For all $s \in]0, 1[$, BV^s functions are regulated functions.*

The fractional total variation only depends on the local extrema of the function and the order of this extrema.

Lemma 1.1. [11, 26] *If u is a piecewise monotonic function and if its local extrema are located in the increasing sequence $(x_i)_{i \in I}$, then $TV^s u$ only depends on the sequence $(u(x_i))_{i \in I}$. Moreover, there exists an ordered subset J of I such that*

$$TV^s u = \sum_{j \in J, j \neq \max J} |u(x_{\text{suc}(j)}) - u(x_j)|^{1/s},$$

where $\text{suc}(j)$ denotes the successor of j in J , $\text{suc}(j) = \min\{k \in J, k > j\}$.

Moreover, it can be dangerous to refine the mesh to compute the fractional total variation [11, example 2.1], [15]. Consider $u(x) \equiv x$ on $[0, R]$, $p = 1/s > 1$. Then $TV^s u[0, R] = R^p$ but, when $n \rightarrow +\infty$,

$$\sum_{i=1}^n |u(iR/n) - u((i-1)R/n)|^p = n(R/n)^p = R^p/n^{p-1} \rightarrow 0.$$

This property, which is not true in BV , is used later to prove the existence of weak solutions in $BV^{1/3}$ for the triangular system (1.1), (1.2) when the flux f is convex.

2 Main results

The main results are stated in the BV^s framework. The basic facts on this setting were recalled in Section 1.3. The critical space for the existence theory is $BV^{1/3}$.

Theorem 2.1 (Existence in L^∞ with $(u_0, v_0) \in BV^{1/3} \times L^\infty$). *Suppose that the flux $f(\cdot)$ and the transport velocity $a(\cdot)$ satisfy the following assumptions on $[-M, M]$ where $M = \|u_0\|_\infty$:*

1. $f'(\cdot)$ and $a(\cdot)$ belong to $C^3([-M, M], \mathbb{R})$,
2. $f'(u) > a(u)$ and satisfy the uniform strict hyperbolicity assumption (USH) (1.6),

Then, if (u_0, v_0) belongs to $BV^s \times L^\infty$ and $s \geq 1/3$, there exists an entropy solution (u, v) of the system (1.1), (1.2), $u \in L^\infty([0, +\infty)_t, BV^s(\mathbb{R}_x, \mathbb{R}))$, $v \in L^\infty((0, +\infty)_t \times \mathbb{R}_x, \mathbb{R})$. In addition, the positivity of the initial data is preserved, that is, if $\inf v_0 > 0$ then $\inf v > 0$.

Two mandatory conditions are required to avoid singular phenomena at $t = 0+$. The strong transversality condition (USH) (1.6) is the key assumption to avoid instantaneous δ -shock. The BV^s regularity of u_0 is the other key condition to avoid instantaneous blow-up in L^∞ .

The positivity of v is the key tool to get the uniform strict hyperbolicity of the (PSA) system [8]. The complete proof of the positivity of v for the (PSA) system is obtained in [13] without using the triangular structure of the (PSA) system [12].

Notice that the sign of v is preserved a.e. by Theorem 2.1. Of course, by linearity of the equation (1.2) with respect to v , the negativity of the initial data is also preserved, that is, if $\sup v_0 < 0$ then $\sup v < 0$.

The triangular approach, that means solving first (1.1) with the unique Krushkov entropy solutions [32] and then (1.2), usually yields non-unique measure solutions v [5, 44]. It is the reason why the uniqueness of entropy solutions for the whole triangular system is an open problem. Of course, we have uniqueness for u but the problem of uniqueness remains for v . For a weakly coupled system with some linearly degenerate fields, the entropy condition only on the nonlinear fields is enough to ensure uniqueness in [30]. For the triangular system, the coupling by the transport velocity is too nonlinear to achieve uniqueness in the same way.

For the existence result, a Wave Front Tracking algorithm (WFT) is proposed as in [3] for the whole system. As a matter of fact the approximate solution u of the scalar equation satisfies the BV^s uniform bounds as in [11, 28]. The difficult point is to bound v in L^∞ . Since the system is linear with respect to v , an L^∞ bound for v is enough to get an existence theorem as in [12].

Theorem (2.1) is optimal and, in general, we cannot reduce the $BV^{1/3}$ regularity of u_0 , else a blow-up can occur. For this purpose, we build a sharp example in a space very near to $BV^{1/3}$, namely, $BV^{1/3-0}$

$$BV^{1/3} \subsetneq BV^{1/3-0} := \bigcap_{s < 1/3} BV^s.$$

The simple example (1.11) from [25] with a Burgers' flux f and a linear velocity a is enough to provide a blow-up.

Theorem 2.2 (Blow-up in L^∞ at $t = 0+$ for $u_0 \in BV^{1/3-0}$). *For the system (1.11) with the Burgers' flux $f(u) = u^2/2$ and the velocity $a(u) = u - 1$, there exists $u_0 \in BV^{1/3-0}$, $v_0 \in L^\infty$ such that*

- *the system (1.1), (1.2) is uniformly strictly hyperbolic (USH) (1.6), and*
- *there doesn't exist any bounded entropy solution of the system (1.1), (1.2) with the initial data u_0, v_0 .*

Of course $u_0 \notin BV^{1/3}$, otherwise the existence theorem 2.1 gives a bounded entropy solution. There is no blow-up for u since the entropy solution u of the scalar conservation

law satisfies the maximum principle. Indeed, only the function v has a blow-up at time $t = 0^+$. Notice also that the blow-up depends only on the regularity of u_0 and not of v_0 .

The example presented below follows the construction of particular explicit solutions [1] which provide optimal example in BV^s for convex scalar conservation laws [23]. On any compact set avoiding the blow-up point, the entropy solution u is locally Lipschitz except on a finite number of lines. For initial data u_0 leading to such an entropy solution u , we show that a uniqueness result for the entropy solution (u, v) holds at least upto the time of first interaction of waves in u . Thus there is no way to avoid the blow-up. In general, a possible dense jump set in plane (t, x) is possible for the entropy solution u of a scalar conservation law [24].

Another consequence of this blow-up example is that for general nonlinear 2×2 strictly hyperbolic systems with a genuinely nonlinear eigenvalue and a linearly degenerate one, the existence result proven in [26] is optimal.

3 The uniformly strictly hyperbolic system

In this section, we study the Riemann invariants and the solution of the Riemann problem for the triangular system (1.1)-(1.2).

3.1 Riemann invariants

A 2×2 strictly hyperbolic system admits, at least locally, a set of coordinates which diagonalizes the hyperbolic system for smooth solutions [46]. The knowledge of this coordinate system, given by the Riemann invariants, is often useful in understanding the structure of the system. Next we study the Riemann invariants for the system (1.1)-(1.2).

The eigenvalues of the system (1.1)-(1.2) are

$$\lambda_1 = f'(u) > a(u) = \lambda_2. \quad (3.1)$$

Notice that the eigenvalues are functions of u only. Let r_1 and r_2 denote the corresponding right eigenvectors.

Clearly, u is a 2-Riemann invariant associated to the right eigenvector $r_2 = (0, 1)^\top$ and satisfies

$$\partial_t u + f'(u) \partial_x u = 0.$$

A 1-Riemann invariant, which we denote as $z(u, v)$, corresponding to the right eigenvector r_1 , can be computed in the following manner. We note that a right eigenvector of the matrix $D\mathbf{F}(u)$ corresponding to the eigenvalue $f'(u)$ is given by

$$r_1 = \begin{pmatrix} 1 \\ \frac{a'(u)v}{f'(u) - a(u)} \end{pmatrix}.$$

Then z satisfies

$$\frac{1}{v} \partial_u z = \frac{a'(u)}{a(u) - f'(u)} \partial_v z.$$

This can be solved using a separation of variables.

$$\partial_u z = \frac{a'(u)}{a(u) - f'(u)}, \quad \partial_v z = \frac{1}{v}.$$

For instance, a Riemann invariant is given by

$$z(u, v) = A(u) + \ln(|v|), \quad A'(u) = \frac{a'(u)}{a(u) - f'(u)}.$$

To avoid the singularity at $v = 0$, it suffices to take the exponential

$$Z = \exp(z) = v \exp(A(u)), \quad (3.2)$$

which satisfies the transport equation

$$\partial_t Z + a(u) \partial_x Z = 0. \quad (3.3)$$

3.2 The Riemann problem

We study the Riemann problem for the system (1.1)-(1.2) with initial data:

$$u_0(x) = u_{\pm}, \pm x > 0, \quad (3.4)$$

$$v_0(x) = v_{\pm}, \pm x > 0. \quad (3.5)$$

A direct and somewhat naive approach is to solve the conservation law (1.1) first and then the second equation (1.2) using the solution u . In such an approach, one faces the difficulty of solving the linear transport equation with a discontinuous coefficient.

Instead, we consider the two equations together as a system. This is a key point as in [3]. The solution of the Riemann problem consists of two waves separated by an intermediary state (u_m, v_m) where $u_m = u_-$ and v_m is unknown. In accordance with the labeling of the eigenvalues (3.1), a wave associated to u is called a 1-wave and a wave associated to Z is called a 2-wave.

1. A wave associated to the eigenvalue $\lambda_1 = f'$ is a shock wave or a rarefaction wave if f is convex or concave. For a non-convex flux f , this wave is a composite wave.
2. A linearly degenerate wave associated to the eigenvalue $\lambda_2 = a$ is called a contact discontinuity. The speed of this 2-wave is $a(u_-)$.

The intermediate value v_m has to be computed through the 1-wave. Now, the various 1-waves that can occur are detailed. For this purpose, we consider Riemann problems yielding only a 1-wave.

Shock waves

Let us denote $U = (u, v)$ and $\tilde{U} = (u, Z)$.

The Rankine-Hugoniot condition gives

$$s[u] = [f(u)], \quad s[v] = [a(u)v], \quad (3.6)$$

where s denotes the speed of the discontinuity (or the slope of the jump).

Thus, the slope of the jump is determined by u_{\pm} and the flux f ,

$$s = \frac{[f(u)]}{[u]}. \quad (3.7)$$

Since entropy solutions are considered, the Oleinik-entropy condition [14] for (1.1) enforces the Lax-entropy conditions,

$$f'(u_-) \geq s \geq f'(u_+), \quad (3.8)$$

whence

$$s \geq f'(u_+) > a(u_+). \quad (3.9)$$

Using the second equation of (3.6) yields

$$(s - a(u_+))v_+ = (s - a(u_-))v_-, \quad (3.10)$$

$$v_+ = v_- \frac{s - a(u_-)}{s - a(u_+)} := \mathcal{S}_-(u_+; U_-). \quad (3.11)$$

The equation (3.11) can be interpreted in terms of the Lax shock curve. For a fixed $U_- = (u_-, v_-)$, the right hand side of (3.11) is only a function of u_+ as s is given by (3.7) as a function of u_+ . In the plane $U = (u, v)$, (3.11) describes the set of U_+ such that the Riemann problem with initial data U_{\pm} is solved by a shock. On this curve, only U_+ satisfying the Oleinik condition (3.8) are considered to allow an entropic shock.

Conversely, if U_+ is fixed, the Lax shock curve is parametrized by u_- and reads

$$v_- = v_+ \frac{s - a(u_+)}{s - a(u_-)} := \mathcal{S}_+(u_-; U_+). \quad (3.12)$$

Proceeding similarly with $Z = v \exp(A(u))$, $\tilde{U}_- = (u_-, Z_-)$ and keeping the notation \mathcal{S} in coordinates (u, Z) yields,

$$Z_+ = Z_- \frac{s - a(u_-)}{s - a(u_+)} \exp[A(u_+) - A(u_-)] := \mathcal{S}_-(u_+; \tilde{U}_-),$$

and

$$Z_- = Z_+ \frac{s - a(u_+)}{s - a(u_-)} \exp[A(u_-) - A(u_+)] := \mathcal{S}_+(u_-; \tilde{U}_+).$$

1-Contact discontinuity

This is a limiting case of the preceding one, when f' is constant on the interval $[u_-, u_+]$. The same formula for the shock wave follows.

$$s = f'(u_{\pm}) = f'(u) > a(u_{\pm}), \quad u \in [u_-, u_+], \quad (3.13)$$

$$v_+ = v_- \frac{f'(u_-) - a(u_-)}{f'(u_+) - a(u_+)} = \mathcal{S}_-(u_+; U_-), \quad (3.14)$$

$$Z_+ = Z_- \frac{f'(u_-) - a(u_-)}{f'(u_+) - a(u_+)} \exp[A(u_+) - A(u_-)] := \mathcal{S}_-(u_+; \tilde{U}_-). \quad (3.15)$$

Such waves arise in the wave-front tracking (WFT) method when the flux is approximated by piecewise linear functions.

Rarefaction waves

Since Z is a 1-Riemann invariant, the Lax rarefaction curve $\mathcal{R}_-(\tilde{U}_-)$ is simply,

$$Z_+ = Z_-. \quad (3.16)$$

Using the definition of Z , this implies $v_+ \exp(A(u_+)) = v_- \exp(A(u_-))$ and thus the Lax rarefaction curve can be written explicitly,

$$v_+ = \mathcal{R}_-(u_+; U_-) = v_- \exp(A(u_-) - A(u_+)), \quad (3.17)$$

$$v_- = \mathcal{R}_+(u_-; U_+) = v_+ \exp(A(u_+) - A(u_-)). \quad (3.18)$$

Notice that v_+ and v_- have the same signs. In particular, if $v_- = 0$, $v = 0$ is constant through the rarefaction wave.

1-Composite waves

The composite waves only occur for the waves associated to the first eigenvalue $f'(u)$. In general, the entropy solution u is a composite wave [27]. If f has a finite number N of inflection points, then there are at most N contact-shock waves [20, 36]. The Lax curves associated to such waves are studied below in Section 4.

2-Contact discontinuity

u is a 2-Riemann invariant and hence is constant along the 2-contact discontinuity. Thus the Lax curve is simply a vertical line in the plane $U = (u, v)$ or the plane $\tilde{U} = (u, Z)$,

$$u_- = u_+. \quad (3.19)$$

4 The Lax curves

A fundamental theorem due to Lax [33] states that the shock curve and the rarefaction curve emanating from a constant state U_- in the plane (u, Z) or (u, v) have a contact of the second order for a genuinely nonlinear eigenvalue. This means that the shock curve can be replaced by the rarefaction curve up to an error of order $[u]^3$ where $[u] = u_+ - u_-$ [14, 20, 45]. For the triangular system, a genuinely nonlinear eigenvalue means $f'' > 0$ (or $f'' < 0$) everywhere. For nonconvex cases, typically f'' locally has a finite number of roots where f'' changes its sign and the Lax curves are less regular due to the occurrence of contact-shocks. Under a concave-convex assumption, which means here that f''' does not vanish, the regularity of the Lax curves is only piecewise C^2 [36], see also [2]. As a consequence, the error becomes of order $[u]^2$ for the variation of Z through a contact-shock [36]. The situation is worse in general, the Lax curves are only Lipschitz [4]. However, we prove that cubic estimates are still valid for the triangular system (1.1), (1.2). It is mainly due to the existence of Riemann invariant coordinates.

In this section, cubic estimates on the Lax curves for the triangular system in the plane (u, Z) are proved.

Let (u_-, v_-) , (u_+, v_+) be two constant states connected by a rarefaction or a shock wave. It is more convenient to use the Riemann invariant coordinates (u, Z) . The rarefaction curves in the plane (u, Z) are simply,

$$Z_- = Z_+, \quad (4.1)$$

which means that in the (u, v) plane

$$v_- \exp(A(u_-)) = v_+ \exp(A(u_+)). \quad (4.2)$$

For the shock curve, the Rankine-Hugoniot condition is written in the conservative variables (u, v) ,

$$s = \frac{[f(u)]}{[u]} = \frac{f(u_+) - f(u_-)}{u_+ - u_-}, \quad (4.3)$$

$$v_-(s - a(u_-)) = v_+(s - a(u_+)). \quad (4.4)$$

Since $f \in C^4$, s is a C^3 function of its arguments. Moreover, fixing (u_+, v_+) and considering u_- as a variable, the Lax shock curve is C^3 with respect to u_- ,

$$v_- = v_+ \frac{s - a(u_+)}{s - a(u_-)}. \quad (4.5)$$

Indeed, the denominator never vanishes due to the uniformly strict hyperbolicity assumption (USH) (1.6). The same regularity of the shock curve holds in the variables (u, Z)

$$Z_- = Z_+ \frac{s - a(u_+)}{s - a(u_-)} \exp(A(u_-) - A(u_+)). \quad (4.6)$$

Of course, the Lax rarefaction curve and the Lax shock curve has to be restricted on the subset satisfying entropy conditions. Nevertheless, we use these curves for all range of u_- in \mathbb{R} (at least for $-M \leq u_- \leq M$) to obtain a generalized Lax cubic estimate for the nonconvex case.

4.1 The Lax cubic estimate on the Rankine-Hugoniot curve

The Lax cubic estimate [33] can be written as follows for a shock wave connecting (u_-, Z_-) to (u_+, Z_+) for the triangular system (1.1),(1.2), as soon as Z is bounded,

$$[Z] = \mathcal{O}([u])^3, \quad [Z] = Z_+ - Z_-, \quad [u] = u_+ - u_-. \quad (4.7)$$

The Riemann invariant Z is constant along the rarefaction curves. The Lax cubic estimate means that the shock curve and the rarefaction curve have a contact of order 2. The Lax cubic estimate was written in a genuinely nonlinear framework [33]. This means that f'' does not vanish. Indeed, the Lax computations are still valid without this convex assumption and without only considering the entropic part of the Rankine-Hugoniot curve. Here, we also use the non-entropic part of the Rankine-Hugoniot curve. It is a useful tool used many times in this paper, first in the next section 4.2 to get a cubic estimate for the entropy solution of the Riemann problem for the triangular system with a non-convex flux f .

For the triangular system, the Rankine-Hugoniot curve is global, explicit and well defined thanks to the uniformly strict hyperbolicity assumption (1.6). For 2×2 systems, such a global curve does not always exist [31].

Now, to prove cubic estimates, we have to write the Rankine-Hugoniot curve. Here, we choose to write Z_- as a function of u_- when (u_+, Z_+) are fixed for the following reasons.

1. To solve the Riemann problem and compute the intermediary state v_m which corresponds to Z_m and here Z_- .
2. To obtain the cubic estimate on the global Rankine-Hugoniot curve below.
3. To obtain the existence result, bounding Z along the 2-characteristics.
4. To build a blow-up, again computing Z from the right to the left on the 2-characteristics.

The Rankine-Hugoniot curve \mathcal{RH}_+ when $\tilde{U}_+ = (u_+, Z_+)$ is fixed and Z_- is a function of u_- , is given by

$$Z_- = Z_+ \frac{s - a(u_+)}{s - a(u_-)} \exp [A(u_-) - A(u_+)] = Z_+ r(u_-, u_+) := \mathcal{RH}_+(u_-; u_+, Z_+). \quad (4.8)$$

Note that the Rankine-Hugoniot curve \mathcal{RH}_+ is a property of (u, v) , which means this curve is always computed in the conservative variables (u, v) [20]. For our convenience, here we write it in terms of (u, Z) due to the fundamental role of the line $Z = \text{constant}$ in the Lax cubic estimate [33]. This classic cubic estimate on the shock curve is generalized on the global Rankine-Hugoniot curve.

Lemma 4.1 (Cubic flatness of the global Rankine-Hugoniot curve). *If f' and a belong to*

$C^3(\mathbb{R}, \mathbb{R})$ and satisfy the uniform strict hyperbolicity condition (1.6), then

$$s = s(u_-, u_+) = \frac{[f]}{[u]} = \frac{f(u_+) - f(u_-)}{u_+ - u_-} \in C^3([-M, M]^2, \mathbb{R}),$$

$$r = r(u_-, u_+) = \frac{s - a(u_+)}{s - a(u_-)} \exp[A(u_-) - A(u_+)] \in C^3([-M, M]^2, \mathbb{R}),$$

$$r = 1 + \mathcal{O}(1)[u]^3 > 0, \quad \forall (u_-, u_+) \in [-M, M]^2, \quad (4.9)$$

$$Z_- = Z_+(1 + \mathcal{O}(1)[u]^3), \quad \forall (u_-, u_+, Z_+) \in [-M, M]^2 \times \mathbb{R}, \quad (4.10)$$

where the constant $\mathcal{O}(1)$ depends only on the derivatives of f' and a on $[-M, M]$. Moreover, Z_- has the same sign as Z_+ , more precisely,

$$Z_+ = 0 \Rightarrow Z_- = 0,$$

$$Z_+ \neq 0 \Rightarrow Z_- Z_+ > 0.$$

The cubic flatness of the Rankine-Hugoniot curve \mathcal{RH}_+ is expressed in (4.10). It does not depend on the convexity of the fields, although in the classical textbooks [14, 20, 27, 36, 45, 46] some nonlinearity assumptions on the fields are given. The reason of these nonlinear assumptions is usually to introduce the rarefaction curve and the shock curve. Here, the global Rankine-Hugoniot curve defined for all $u_- \in [-M, M]$ is the main subject without looking at the entropic parts of this curve. A careful reading of the classical proof of the cubic estimate in textbooks shows that it is a geometric property of the Rankine-Hugoniot curve itself, without referring at the entropic or nonlinearity assumptions. This geometric property is a consequence of the symmetry of the Rankine-Hugoniot condition with respect to U_- and U_+ as explained in [45]. It is very important in this paper to prove the cubic estimate for the non-entropic part of the Rankine-Hugoniot curve for two reasons.

1. The cubic estimates on the Lax curve (which is not piecewise C^3 for nonconvex f [36]) uses the global Rankine-Hugoniot curve, as discussed in the next section 4.2.
2. Rarefaction wave fans are replaced by weak non-entropic jumps in the wave front tracking algorithm [14]. The error in the weak formulation is controlled by the cubic estimate to pass to the limit and get a weak solution of (1.1), (1.2).

The proof appears as a direct consequence of (4.9). An elementary and self-contained proof using only Taylor's expansions is proposed. A more tedious computation can give the more precise result

$$r(u_-, u_+) = 1 + E[u]^3 + \mathcal{O}([u]^3), \quad (4.11)$$

where E depends in a quite complicated way on the derivatives of f' and a at $u = u_+$. The computation of E is quite intricate and not useful here, except in the last section 6 on the blow-up where a direct computation of E at $u_+ = 0$ is given when f is quadratic and a is linear.

Now, Lemma 4.1 is proven.

Proof. The positivity on r is a consequence of the assumption (1.6). This positivity implies that Z_- has the same sign as Z_+ .

To obtain (4.9), we use Taylor expansion. Here u_+ is fixed and u_- is the variable near u_+ . The notations $a_- = a(u_-)$, $a_+ = a(u_+)$ and so on are used to shorten the expressions.

$$\begin{aligned} a_- &= a_+ - a'_+[u] + (a''_+/2)[u]^2 + \mathcal{O}([u]^3), \\ s &= \frac{[f]}{[u]} = \frac{f_- - f_+}{-[u]} = f'_+ - (f''_+/2)[u] + (f'''_+/6)[u]^2 + \mathcal{O}([u]^3). \end{aligned}$$

The hyperbolic quantity h is used,

$$h := f' - a > 0. \quad (4.12)$$

The sign of h is the consequence of the strict hyperbolicity assumption (1.6). We start with the fraction part of r and $h_+ = f'_+ - a_+ \neq 0$.

$$\frac{s - a_+}{s - a_-} = \frac{(f'_+ - a_+) - (f''_+/2)[u] + (f'''_+/6)[u]^2}{(f'_+ - a_+) - (f''_+/2 - a'_+)[u] + (f'''_+/6 - a''_+/2)[u]^2} + \mathcal{O}([u]^3) \quad (4.13)$$

$$= 1 - \frac{a'_+[u]}{h_+} + \frac{h_+ a''_+ - a'_+ f''_+ + 2a_+^2}{2h_+^2} [u]^2 + \mathcal{O}([u]^3). \quad (4.14)$$

For the term $\exp(-[A])$ in r , the Taylor expansion of A'_- is used at the first order.

$$\begin{aligned} -A'_- &= \frac{a'_-}{h_-} = \frac{a'_+ - a''_+[u]}{h_+ - h'_+[u]} + \mathcal{O}([u]^2) \\ &= \frac{a'_+}{h_+} + \frac{a'_+ h'_+ - a''_+ h_+}{h_+^2} [u] + \mathcal{O}([u]^2). \end{aligned}$$

Integrating with respect to u_- yields,

$$A_- - A_+ = \frac{a'_+[u]}{h_+} + \frac{a'_+ h'_+ - a''_+ h_+}{2h_+^2} [u]^2 + \mathcal{O}([u]^3).$$

Since $\exp(x) = 1 + x + x^2/2 + \mathcal{O}(x)^3$, it yields

$$\exp[A_- - A_+] = 1 + \frac{a'_+[u]}{h_+} + \frac{a'_+ h'_+ - a''_+ h_+ + a_+^2}{2h_+^2} [u]^2 + \mathcal{O}([u]^3). \quad (4.15)$$

Now, multiplying (4.14) and (4.15) yields $r = 1 + \mathcal{O}([u]^3)$. \square

4.2 Cubic estimates for the Riemann problem

Now, the intermediary state Z_m of a Riemann problem has to be estimated. For this purpose, the variation of Z along a composite wave is studied. When the flux is convex, Lax proved that the variation of Z is a cubic order of the variation of u [33]. For a non-convex flux, it is well known that the Lax curve is less regular, piecewise C^2 [36] or only Lipschitz [4]. However, we prove that for our triangular system we are able to keep a cubic

order. This is mainly due to the existence of Riemann coordinates for 2×2 systems and the cubic estimate for the global Rankine-Hugoniot locus, Lemma 4.1. As a consequence, we can prove a similar estimate for the variation of Z over a composite 1-wave. This improves the well known square root estimate for concave-convex eigenvalues [36], which correspond to cubic degeneracies for f . That means that for the triangular system (1.1), (1.2), the estimate is as precise as for the convex case [33].

Proposition 4.1 (Variation of Z through a composite wave).

Assume that the flux has N_{infl} points.

Let the states $\tilde{U}_i = (u_i, Z_i)$, $i = 1, \dots, m$, where $u_0 < u_1 < \dots < u_m$ ($m \leq N_{infl} + 1$) comprise a composite 1-wave, and $Z_- = Z_0$ and $Z_+ = Z_m$. Then the total variation of Z through a 1-wave is,

$$\|Z\|_\infty \leq |Z_+| \exp(\mathcal{O}(|u_+ - u_-|^3)), \quad (4.16)$$

$$TVZ \leq \mathcal{O}(1)|Z_+||u_+ - u_-|^3. \quad (4.17)$$

Later, as the consequence of the existence proof, the assumption of the finite number of inflection points can be removed and the same inequalities (4.16), (4.17) hold.

Proof. We note that Z is a 1-Riemann invariant and therefore it remains constant over a rarefaction wave. Moreover, if the states \tilde{U}_i and \tilde{U}_{i+1} are joined by a jump, by Lemma 4.1 and the classical inequality $1 + x \leq \exp(x)$, we have the estimate

$$|Z_i| = |Z_{i+1}r(u_i, u_{i+1})| = |Z_{i+1}|(1 + \mathcal{O}(|u_i - u_{i+1}|^3)) \leq |Z_{i+1}| \exp(\mathcal{O}(|u_i - u_{i+1}|^3)).$$

Summing up and noting that u_0, u_1, \dots, u_m are ordered, the estimate (4.16) follows,

$$\max_i |Z_i| \leq |Z_+| \exp(\mathcal{O}(|u_+ - u_-|^3)).$$

Now, the BV bound for Z through the composite wave is computed.

$$\begin{aligned} Z_i - Z_{i+1} &= Z_{i+1}(r(u_i, u_{i+1}) - 1) = Z_{i+1}\mathcal{O}(|u_i - u_{i+1}|^3), \\ TVZ &\leq \max_i |Z_i| \sum_i \mathcal{O}(|u_i - u_{i+1}|^3) \leq |Z_+|\mathcal{O}(|u_+ - u_-|^3) \exp(\mathcal{O}(|u_+ - u_-|^3)). \end{aligned}$$

Since u satisfies the maximum principle, we further have $\exp(\mathcal{O}(|u_+ - u_-|^3)) = \mathcal{O}(1)$ which only depends on the L^∞ bound of the initial data u_0 . Hence the inequality (4.17) follows. \square

5 Existence in BV^s

In this section, Theorem 2.1 is proved using a simplified Wave Front Tracking (WFT) [14, 27] algorithm for such a triangular system (1.1), (1.2). The BV^s estimates for u are a consequence of such estimates for scalar conservation laws [11, 29, 28]. The L^∞ bound for v and the proof of existence of a weak solution for the triangular system are based on an approach using the $BV^{1/3}$ regularity for u .

5.1 The Wave Front Tracking algorithm

The WFT depends on an integer parameter $\nu > 0$. The approximate solutions will be denoted by u_ν, v_ν, Z_ν . We shall mostly use the Riemann invariant coordinates (u_ν, Z_ν) except when passing to the limit in the weak formulation.

This algorithm is explained in many books [14, 20, 27] on hyperbolic systems. Taking advantage of the structure of the triangular system (1.1), (1.2), we will mix the WFT for the scalar case [19] and for systems [14, 27]. The main principle is to work with piecewise constant approximations.

As in the scalar case [14, 27, 29, 28], the values of u_ν are taken on a uniform grid parametrized by the integer ν , that is,

$$u_\nu \in \nu^{-1}\mathbb{Z}. \quad (5.1)$$

On the other hand, v_ν , or equivalently Z_ν , is not required to stay on the uniform grid, that is,

$$v_\nu, Z_\nu \in \mathbb{R}, \quad (5.2)$$

as we solve the Riemann problem with the exact flux f to compute v_ν , or equivalently Z_ν . In this way, we can use the cubic estimate on the global Rankine-Hugoniot curves of Lemma 4.1. A similar approach for a convex case was used in [26].

The initial data are approximated as follows. Let $N_0 + 1$ be the number of constant states for the approximate initial data $u_{0,\nu}, v_{0,\nu}$ and the corresponding $Z_{0,\nu}$. Here $u_{0,\nu}$ takes values on the grid and N_0 is a function of ν .

To ensure that $(u_{0,\nu}, v_{0,\nu})$ converges towards the initial data (u_0, v_0) , we need the condition that N_0 goes to infinity as ν tends to infinity.

The approximate initial data can be chosen to satisfy the following uniform estimates with respect to ν [29, 28]:

$$\|u_{0,\nu}\|_\infty \leq \|u_0\|_\infty, \quad (5.3)$$

$$TV^s u_{0,\nu} \leq TV^s u_0, \quad (5.4)$$

$$\|v_{0,\nu}\|_\infty \leq \|v_0\|_\infty. \quad (5.5)$$

Moreover, $(u_{0,\nu}, v_{0,\nu})$ converges pointwise almost everywhere to (u_0, v_0) when $\nu \rightarrow +\infty$ and therefore, the previous inequalities become equalities at the limit $\nu \rightarrow +\infty$.

The flux f is replaced by a piecewise linear continuous flux f_ν [19] that coincides with f on the uniform grid [14, Ch. 6], [27, p. 70],

$$f_\nu(k/\nu) = f(k/\nu), \quad \forall k \in \mathbb{Z}. \quad (5.6)$$

At $t = 0+$, N_0 Riemann problem are solved. The approximate solution u_ν is the weak entropy solution of

$$\partial_t u_\nu + \partial_x f_\nu(u_\nu) = 0, \quad u_\nu(x, 0) = u_{0,\nu}(x). \quad (5.7)$$

With strong compactness on u_ν , the Kruzkov entropy solution is recovered [14, 27].

For (u_ν, v_ν) the approximate vectorial flux $F_\nu(u, v) = (f_\nu(u), a(u)v)$ is used, as detailed below. The reason for this choice is twofold. First, to have a simpler wave front tracking for the scalar equation for u with a non-convex flux following the seminal paper [19] (this case is also very well detailed in the textbook [27]). This wave front tracking algorithm does not lead to any consistency error in the weak formulation [19]. Secondly, to use the new generalized cubic estimates for Z_ν which also hold for the solution (u_ν, v_ν) of the system with the approximate flux F_ν , as $f_\nu = f$ on the grid $\{(u, v) \in (\nu^{-1}\mathbb{Z}) \times \mathbb{R}\}$.

After some time, some of the Riemann problems interact and hence, new Riemann problems have to be solved. The non-linear interactions for the triangular system are described in detail below. The process continues until the second time of interactions and so on.

This process can be continued for all time because there is only a finite number of interactions for all time. For u_ν this follows from the existing results for the scalar case. An explicit bound on the number of interactions is given in [27, p. 71-72]. The 2-waves, associated to the linearly degenerate eigenvalue $a(u_\nu)$, never interact together since they are contact discontinuities. Due to the transversality assumption (USH), a 2-wave can interact only once with a 1-wave and creates a new 2-wave (and does not modify the 1-wave). Thus, the number of such interactions and of 2-waves is finite. This proves that the WFT is well defined for all time.

Now, the approximate Riemann solver and the nonlinear interactions of the waves are detailed.

Approximate Riemann solver

In this short section, the approximate Riemann solver is detailed. Let (u_-, Z_-) , (u_+, Z_+) denote the left and right states in the initial data. For u_ν the solution is a series of entropic jumps $u_- = u_0 < u_1 < \dots < u_m = u_+$ for the piecewise linear flux f_ν . For Z_ν , there are many possibilities. We want to approximate the exact solution of the Riemann problem and keep the cubic estimates (Proposition 4.1) which generalize the Lax cubic estimates for genuinely nonlinear waves. So, we use the Rankine-Hugoniot curve to determine Z_{m-1}, \dots, Z_1 . The states Z_i , $i = m, m-1, \dots, 2$ are built as follows:

- If there is a jump between u_{i-1} and u_i corresponding to a shock or a contact discontinuity for f_ν , then Z_{i-1} is given by the Rankine-Hugoniot curve (4.8),

$$Z_{i-1} = \mathcal{RH}_+(u_{i-1}; u_i, Z_i). \quad (5.8)$$

Note that $\mathcal{RH}_+^\nu = \mathcal{RH}_+$ on the grid $\{(u, v) \in (\nu^{-1}\mathbb{Z}) \times \mathbb{R}\}$, where \mathcal{RH}_+^ν denotes the Rankine-Hugoniot curve corresponding to the approximate flux F_ν . Thus, the jump between (u_{i-1}, Z_{i-1}) and (u_i, Z_i) satisfies the Rankine-Hugoniot condition and there is no error in the weak formulation of the linear equation satisfied by v and hence, no error in the weak formulation of the approximate triangular system (5.7), (1.2), where the exact flux f is replaced by the piecewise-linear flux f_ν in (1.1).

- If there is a 2-contact discontinuity (along which u is constant and there is a jump between v_{\pm}), then the contact discontinuity is solved with the exact speed $a(u)$. Thus, again, there is no error of consistency for the approximate system (5.7), (1.2).

Nonlinear wave interactions

We briefly describe the different possible nonlinear wave interactions and the details of the interactions are given in the (u, Z) plane. Here Z denotes the 1-Riemann invariant with respect to the exact flux F .

Lemma 5.1. *The cubic estimates for Z proved in Proposition 4.1 also hold true across 1-waves (shock and 1-contact discontinuity) corresponding to the approximate flux F_{ν} .*

Proof. This immediately follows from (5.8), the fact that $\mathcal{RH}_+^{\nu} = \mathcal{RH}_+$ on the grid and the cubic estimates for Z across 1-waves corresponding to f (Proposition 4.1). \square

In all the cases, we consider three states $\tilde{U}_-, \tilde{U}_0, \tilde{U}_+$ before the interaction. Here \tilde{U}_0 denotes the intermediary state which disappears after the interaction. The states \tilde{U}_-, \tilde{U}_0 are connected by an elementary wave and similarly the states \tilde{U}_0, \tilde{U}_+ are connected by an elementary wave. An elementary wave is either a 1-wave: shock or a small jump, or a 2-wave: a contact discontinuity. At a time of interaction $t_{interact}$, the Riemann problem is solved with a new intermediary state \tilde{U}_m (see Figure 1).




Figure 1: Nonlinear interaction of two waves. The 1-wave on the left crosses the interaction point with the same speed and the same value u_- on the left of the 1-wave ($u_m = u_-$) and u_+ on the right. On the other hand, the second wave is affected by the interaction: the speed of the 2-wave and a new value Z_m appears.

We shall use the following notations.

- S or S_1 stands for a shock wave which is always a 1-wave.

- C_1 or C_2 stands for a contact discontinuity associated to λ_1 or λ_2 , a 1-wave or a 2-wave. C_1 is considered as a degenerate shock S_1 .

The key point here is to understand the effect of the L^∞ norm of Z_ν after the interaction. First of all, we note that there is no self interaction for the second family since there are only C_2 waves. Also the case of interactions between the 1- waves of the first family have already been well-studied [14], [20],[27] for the component u_ν and are not presented here. The new feature is the effect of this interaction on the second component Z_ν . We have seen in lemma 5.1 that the change in L^∞ norm of Z_ν is of the order of the cube of the change in u (or u_ν).


Finally, we consider the interaction of a 1-wave with a 2-wave.

1. In the case of an interaction of the form $S_1 - C_2$ (which means that a 1-shock interacts with a 2-contact discontinuity), the outgoing wave is of the form $C_2 - S_1$. The shock continues with the same slope and the same value u_-, u_+ and still satisfies the entropy condition (3.8). Thus, for u there is no change before and after the interaction. Roughly speaking, the interaction of a 1-shock with a 2-contact discontinuity is transparent for u . On the contrary, there is a change for Z_ν and following [26], it can be shown that the change in L^∞ norm of Z_ν is of the order of cube of the change in u_ν .
2. The interaction $C_1 - C_2$ (that is, when a 1-contact discontinuity interacts with a 2-contact discontinuity) generates outgoing waves of the form $C_2 - C_1$. This case can be dealt in a similar manner as in the case of $S_1 - C_2$ and it follows that the change in L^∞ norm of Z_ν is of the order of cube of the change in u_ν .

The reader interested by all the cases $S_1 - C_1$, $C_1 - S_1$, $S_1 - C_2$, $C_1 - C_2$ can consult [26] (and different notations where the 1 and 2 waves have to be exchanged) in the plane of Riemann invariant, a technique used in [46] for the isentropic Euler system. Here we present the last two cases for the triangular system, the case $S_1 - C_2$ in figure 2 and the case $C_1 - C_2$ in figure 3.


5.2 Uniform estimates

The BV^s estimates for u_ν are already known since u is the entropy solution of the scalar conservation law (1.1). These estimates are recalled briefly in the first paragraph. The only difficulty in this section is to obtain the L^∞ estimates for v . For this purpose, we generalize the approach first used in [12] and recently in [26]. The approach consists of bounding v_ν , indeed Z_ν , along the 2-characteristics. For the chromatography system [12], the 2-characteristics are simply straight lines. In general, here, the 2-characteristics are piecewise linear. They are uniquely defined since the second eigenvalue $a(u)$ is linearly degenerate and its integral curves are transverse to the discontinuity lines of the first field (assumption (USH)). The precise definition of such characteristics is given in the second paragraph. Then the estimate on Z_ν along 2-characteristics, as in [12, 26], is given in the last paragraph using our generalized estimates on the Lax curves.



Interaction Shock & Contact Discontinuity

Figure 2: Interaction of a shock with a 2-contact discontinuity. The interacting waves are represented by dotted lines, a 1-wave in black followed by a 2-wave in red. The full lines represent the resulting waves.



Interaction 1 and 2 Contact Discontinuities

Figure 3: Interaction between a 1-contact discontinuity and a 2-contact discontinuity. The interacting waves are represented by dotted lines, a 1-wave in black (horizontal line) followed by a 2-wave in red (vertical line). The full lines represent the resulting waves.

BV^s estimates for u

The TV^s decay is known for the Glimm scheme, the Godunov scheme and the Wave Front Tracking algorithm [11, 12]. Another argument is that u_ν is also the exact entropy solution of the scalar conservation law (5.7) with the piecewise-constant initial data $u_{0,\nu}$, and the decay of $TV^s u$ gives the uniform estimates with respect to ν ,

$$TV^s u_\nu(\cdot, t) \leq TV^s u_{0,\nu} \leq TV^s u_0. \quad (5.9)$$

The approximate 2-characteristics

Essentially, for the WFT, an approximate i -characteristic is a continuous curve which is piecewise linear following the velocity λ_i , $i = 1, 2$. Since the eigenvalues depend only on u , there is a problem to define an i -characteristic where u is not defined. For $i = 2$, there is no problem of uniqueness, since a 2-characteristic is always transverse to the discontinuity lines of u . Thus, the 2-characteristic crosses the u discontinuity with a kink.

Let $\gamma_\nu(x_0, t)$ be the forward generalized 2-characteristic starting at the point x_0 , that is $\gamma_\nu(x_0, t)$ is a solution of the differential inclusion

$$\frac{d}{dt}\gamma_\nu(x_0, t) \in [a(u_\nu(\gamma_\nu(x_0, t) - 0, t)), a(u_\nu(\gamma_\nu(x_0, t) + 0, t))], \quad \gamma_\nu(x_0, 0) = x_0.$$

For the wave front tracking, these 2-characteristics are uniquely determined and are piecewise linear continuous curves (thanks to the transversality assumption (USH)) and satisfy the differential equation

$$\frac{d}{dt}\gamma_\nu(x_0, t) = a(u_\nu(\gamma_\nu(x_0, t), t)), \quad \gamma_\nu(x_0, 0) = x_0, \quad (5.10)$$

except for a finite number of times which correspond to a jump of the piecewise constant function u_ν .

L^∞ estimates for v

The L^∞ estimate on v_ν is first obtained on the approximate Riemann invariant Z_ν . The key point is then to bound Z_ν in L^∞ . Z_ν is not constant through a 1-wave: a shock wave or a 1-contact discontinuity. But, we know that the variation of Z_ν is of order of the cube of the variation of u_ν , by lemma 5.1. When there is no 1-wave, the simple curve to bound Z_ν is the 2-characteristic. In this case, $Z_\nu(\gamma_\nu(x_0), t) = Z_\nu(x_0, 0)$.

Note that the 2-characteristics starting at time $t = 0$ span the whole (t, x) half-plane but Z_ν is only defined outside a finite number of 2-characteristics.

For example, Z_ν is not well-defined on a 2-characteristic that contains a 2-wave (contact discontinuity for the second family). But these are only finite in number as the number of 2-wave fronts is finite.

Also Z_ν is not well-defined on 2-characteristics passing through the points of intersections of waves corresponding to the first family. Again there are only finite number of such curves.

Z_ν is also not well-defined on a finite number of 2-characteristics originating from the points on $t = 0$ where the Riemann problems are solved initially.




Figure 4: Wave front tracking algorithm. For the picture, $f'(u) > 0 > a(u)$. Thus the 1-waves in blue go to the right and the 2-waves in black to the left. Notice that these 1-waves are not affected by the interaction with the 2-waves, but the 2-waves are affected by the interaction.

But since there are only finite number of such 2-characteristics, this is enough to estimate $\|Z_\nu\|_\infty$.

Now, due to the transversality conditions, a 2-characteristic meets many 1-waves. Thanks to lemma 5.1, we have a cubic estimate for the L^∞ norm and the total variation of Z_ν through a 1-wave. More precisely, if t_1 is a time just before the 2-characteristic meets a 1-wave and t_2 is the time just after the 2-characteristic crosses the 1-wave, we have the following estimate,

$$\begin{aligned} \|Z_\nu\|_{\infty, (\gamma_\nu(x_0, \cdot), \cdot)[t_1, t_2]} &\leq |Z_\nu(t_1)| \exp(\mathcal{O}(TV^{1/3}u_\nu(\gamma_\nu(x_0, t_2), t_2))), \\ TV Z_\nu(\gamma_\nu(x_0, \cdot), \cdot)[t_1, t_2] &\leq \|Z_\nu\|_{\infty, (\gamma_\nu(x_0, \cdot), \cdot)[t_1, t_2]} (\mathcal{O}(TV^{1/3}u_\nu(\gamma_\nu(x_0, \cdot), \cdot)[t_1, t_2])). \end{aligned}$$

That means that the L^∞ norm and the total variation of Z_ν along the piece of curve $\{(\gamma_\nu(x_0, t), t), t \in [t_1, t_2]\}$ is controlled by $TV^{1/3}u_\nu$ along the same piece of curve. Notice also that the sign of Z_ν is constant along a 2-characteristic by lemma 4.1.

First, the L^∞ norm of Z is bounded on the 2-characteristic and, second, the estimate on $TV Z_\nu$ follows. The total variation is additive and the fractional total variation is sub-additive [15]. So adding all these estimates, on the whole 2-characteristic $\Gamma(x_0) =$

$\{(\gamma_\nu(x_0, t), t), t > 0\}$ starting at $x = x_0$, we have the estimates

$$\begin{aligned} \|Z\|_{\infty, \Gamma(x_0)} &\leq |Z_{0,\nu}(x_0)| \exp(\mathcal{O}(TV^{1/3}u_\nu[\Gamma(x_0)])), \\ &\leq \|Z_0\|_{\infty} \exp(\mathcal{O}(TV^{1/3}u_\nu[\Gamma(x_0)])), \\ TVZ_\nu[\Gamma(x_0)] &\leq \|Z_0\|_{\infty} \Psi(\mathcal{O}(TV^{1/3}u_\nu[\Gamma(x_0)])), \\ \Psi(x) &= x \exp(x). \end{aligned}$$

Now note that $\Gamma(x_0)$ is a space like curve for the 1-characteristics of (5.7) ([20]). Hence

$$TV^{1/3}u_\nu[\Gamma(x_0)] \leq TV^{1/3}u_{0,\nu} \leq TV^{1/3}u_0.$$

This follows from the fact the wave-fans corresponding to u_ν are monotone with respect to the left and right end states. Therefore no extremal values are added when we measure the fractional variation along $\Gamma(x_0)$.

Together these yield L^∞ and BV bounds for Z_ν (uniform with respect to ν) along the 2-characteristics,

$$\|Z_\nu\|_{\infty} \leq \|Z_0\|_{\infty} \exp(\mathcal{O}(1)TV^{1/3}u_0), \quad (5.11)$$

$$TVZ_\nu[\Gamma(x_0)] \leq \|Z_0\|_{\infty} \Psi(\mathcal{O}(TV^{1/3}u_0)). \quad (5.12)$$

Notice also that the positivity is preserved. If $\inf v_0 > 0$, then $\inf Z_0 > 0$ and by similar arguments presented above, we have $\inf Z_\nu > \inf Z_0 \exp(\mathcal{O}(1)TV^{1/3}u_0) > 0$, where the constant $\mathcal{O}(1)$ is negative. Finally, a L^∞ bound (respectively a positivity) for Z_ν yields a L^∞ bound (respectively a positivity) for v_ν .

For the triangular system, the L^∞ bound of Z_ν and hence of v_ν is enough to pass to the weak limit in (1.2) since the left hand side is linear with respect to v_ν . The uniform BV bound on Z_ν along the 2-characteristics can be used to recover a strong trace at $t = 0$, like in [8] (at $x = 0$).

5.3 Passage to the limit in the weak formulation

Passing to the strong-weak limit in the equation (1.2) which is linear with respect to v allows us to get a weak solution. But to do so, we need to understand the error of consistency of the scheme. There is no error of consistency in the independent scalar equation for u_ν (5.7) [19]. By construction, our approximate solver does not produce error for the linear equation (1.2). That means that the approximate solution (u_ν, v_ν) is an exact solution of the approximate system (5.7),(1.2):

$$\begin{aligned} \partial_t u_\nu + \partial_x f_\nu(u_\nu) &= 0, \\ \partial_t v_\nu + \partial_x (a(u_\nu)v_\nu) &= 0. \end{aligned}$$

Therefore as in [19], one can pass to the limit and obtain a weak solution of the exact system (1.1),(1.2).

6 Blow-up of an entropy solution with $u_0 \notin BV^{1/3}$

In this section, we provide a proof of Theorem 2.2. For this purpose, we construct initial data u_0, Z_0 that satisfy

$$\begin{cases} u_0 \in BV^{1/3-0}(\mathbb{R}), \\ Z_0 \in L^\infty(\mathbb{R}), \end{cases}$$

and exhibit blow-up in L^∞ norm of Z at $t = 0+$ for the system (1.1)-(1.2).

Recall that the notation $u_0 \in BV^{1/3-0}(\mathbb{R})$ means that u_0 is in BV^s for all $s \in [0, 1/3)$. Our idea of construction is motivated by similar examples studied in [1, 17, 22, 23].

We would like to emphasize that the solution proposed below is not based on the Wave-Front Tracking algorithm discussed earlier, rather it is an exact solution.

Notice that when $v_0 \equiv 0$, i.e. $Z_0 \equiv 0$, no blow-up occurs since $(u, v) \equiv (u, 0)$ gives a global entropy solution where u the entropy solution associated to the L^∞ initial data u_0 . Thus, in the following construction, we have to avoid the value 0 for Z .

Let us consider the system (1.1)-(1.2) with flux $f(u) = \frac{u^2}{2}$, $a(u) = u - 1$ and initial data u_0 as described below and $Z_0 \equiv 1$.

Let $x_0 = 0$ and $x_n = 1 - \frac{1}{2^n}$, $n \geq 1$. Let B_n be chosen such that

$$x_n = x_{n-1} + 2B_n,$$

that is, $B_n = \frac{1}{2^{n+1}}$. Let $b_n = \frac{1}{(n+26)^{\frac{1}{3}}}$ and we define

$$U_0(x, b, B) = b\mathbb{1}_{[0, B)}(x) - b\mathbb{1}_{[B, 2B)}(x).$$

Using this we define the initial data u_0 as

$$u_0(x) = \sum_{n \geq 1} U_0(x - x_{n-1}, b_n, B_n).$$

Note that $\|u_0\|_\infty \leq \frac{1}{3}$ and therefore, the condition of uniform strict hyperbolicity is satisfied.

The first interaction times T_n for the Riemann problems for the equation

$$\partial_t u + \partial_x f(u) = 0$$

with initial data u_0 is given by

$$T_n = \frac{B_n}{b_n} = \frac{(n+26)^{\frac{1}{3}}}{2^{n+1}}.$$

Note that the first interaction time T_n satisfies the relation

$$T_n > 1 - x_n. \tag{6.1}$$




Figure 5: A typical building-block

The initial data u_0 , described above, clearly does not belong to $BV^{\frac{1}{3}}(\mathbb{R})$, but, as in [12, 17] we can conclude that $u_0 \in BV^{1/3-0}(\mathbb{R})$.

We use the forward generalized characteristic for Z . Since $a(u) = u - 1$ and the first interaction times of the Riemann problems for u satisfy (6.1), it follows that the forward generalized characteristic for Z starting at the point $x_\infty := 1$ crosses infinitely many shocks before the first interaction of the waves in u .

As we have already seen from the nonlinear wave interactions, the L^∞ norm of Z does not change when it interacts with a 1-rarefaction wave.

Now, let us consider a left state u_- and a right state u_+ connected by a 1-shock wave. In our example, we have $u_- = b > 0$ and $u_+ = -b < 0$. Hence, the speed of the shock $s = 0$. Also by construction $0 < b_n < \frac{1}{2} < 1$ and hence we assume that the prototype b satisfies the same.

Now

$$\begin{aligned}
\frac{Z_-}{Z_+} &= \frac{s - a(u_+)}{s - a(u_-)} \exp [A(u_-) - A(u_+)] \\
&= \frac{a(u_+)}{a(u_-)} \exp [A(u_-) - A(u_+)] \\
&= \frac{u_+ - 1}{u_- - 1} \exp [A(u_-) - A(u_+)] \\
&= \frac{1 + b}{1 - b} \exp [A(u_-) - A(u_+)].
\end{aligned} \tag{6.2}$$

Also

$$A'(u) = \frac{a'(u)}{a(u) - f'(u)} = \frac{1}{(u - 1) - u} = -1 < 0,$$

and hence

$$A(u_-) - A(u_+) = \int_{u_+}^{u_-} A'(u) du = -(u_- - u_+) = -2b.$$

Therefore from (6.2), we have

$$\frac{Z_-}{Z_+} = \frac{1+b}{1-b} e^{-2b}. \quad (6.3)$$

We show that for b positive and small enough,

$$\frac{1+b}{1-b} e^{-2b} > 1, \quad (6.4)$$

and therefore $Z_- > Z_+$. Thus, Z increases in strength as the forward 2-generalized characteristic crosses a shock (from right to left).

Notice that we simply need that Z increases when $b \sim 0$. It is for very small oscillations that Z blows-up. Inequality (6.4) follows from a Taylor expansion up to the third order:

$$\begin{aligned} (1+b)\exp(-2b) &= (1+b)(1 - 2b + 1/2 \cdot (2b)^2 - 1/6 \cdot (2b)^3 + \mathcal{O}(b)^4) \\ &= 1 + (1-2)b + (-2+2)b^2 + (2-4/3)b^3 + \mathcal{O}(b)^4 \\ &= 1 - b + 2/3b^3 + \mathcal{O}(b)^4 \\ &> 1 - b, \end{aligned}$$

for b sufficiently small and hence $Z_- > Z_+ > 0$.

Thus, due to an interaction with a 1-shock wave, there is a change of order $[u]^3$ in the L^∞ norm of Z . Since

$$\sum_{n \geq 1} |[u]^3| = \sum_{n \geq 1} \left(\frac{2}{(n+26)^{\frac{1}{3}}} \right)^3 = 8 \sum_{n \geq 1} \frac{1}{(n+26)} = +\infty,$$

we find that the L^∞ norm for Z blows up.

Remark 6.1. Let $\Gamma(x_0) = \{(\gamma(x_0, t), t), t \geq 0\}$ be the 2-characteristic issued from $x = x_0$ at $t = 0$. The solution is well defined under $\Gamma(1)$ that is on the set $\{(x, t), x < \gamma(1, t), t \geq 0\}$. But, over $\Gamma(1)$, $\{(x, t), 1 > x > \gamma(1, t), t > 0\}$, Z and v blow up, $v = +\infty$.

Remark 6.2. This example does not contradict the Lax-Oleinik smoothing effect ([33]) as the blow-up for Z occurs only at $t = 0$, that is, there is an immediate blow-up. Such a blow-up is not possible for a time $t_0 > 0$ due to the BV smoothing of u .

Now suppose that there exists a bounded weak solution for the system (1.1)-(1.2) with the above initial data. Then since the solution of the Riemann problem in the class of bounded weak solutions is unique upto the time of first interaction of the waves (see appendix B), any bounded weak solution will satisfy the construction discussed above. This, in turn, implies that the bounded weak solution exhibits a blow-up in L^∞ norm, which contradicts its definition.

Thus, there cannot exist a bounded weak solution for the system (1.1)-(1.2) with the above initial data, which proves Theorem 2.2. This also shows that the existence Theorem 2.1 is optimal in the class of bounded weak solutions.

Acknowledgments

A part of this work was done when the last author was visiting NISER, Bhubaneswar, India. We also thank the IFCAM project “ BV^s , control & interfaces” which allowed the interaction between the Indian hyperbolic community with the French one. We also thank Boris Haspot for fruitful discussions.

A On Lipschitz test functions

Here we briefly discuss the use of Lipschitz test functions instead of C^1 test functions for bounded weak entropy solutions for triangular systems (1.1)-(1.2). We show that if $u, v \in L^\infty$ is a weak bounded solution (and therefore the solution constructed in Section 5) for the triangular system, then the C^1 test functions can be enlarged to Lipschitz functions. This can be proved using the following observations. Note that (u, v) is a weak solution and hence satisfies the integral identities (1.8)-(1.9) for compactly supported smooth test functions. Now if we consider a compactly supported Lipschitz continuous test function ψ , we can construct (using a standard mollifier) a sequence of compactly supported smooth test functions $\{\phi_n\}$ that converges pointwise almost everywhere to ψ . An application of dominated convergence theorem shows that $\partial_t \phi_n$ and $\partial_x \phi_n$ converge to $\partial_t \psi$ and $\partial_x \psi$ respectively at points of continuity (and therefore almost everywhere) of ψ . Since the integral identities (1.8)-(1.9) are satisfied for each ϕ_n , we can use the above facts to pass to the limit in the identities to prove the integral identities for the Lipschitz continuous test function ψ .

B Uniqueness of Riemann problem in L^∞

We next show that in the case of strictly convex fluxes f , the solution of the Riemann problem is unique in the class of bounded entropy solutions at least upto the time of first interaction of waves. In this direction, we first note the following result.

Lemma B.1. *If (u, v) is a bounded entropy solution to the system (1.1)-(1.2) with initial data $(u_0, v_0) \in L^\infty(\mathbb{R}) \times L^\infty(\mathbb{R})$, then $Z := v \exp(A(u))$ satisfies*

$$\partial_t Z + a(u) \partial_x Z \stackrel{E}{=} 0, \tag{B.1}$$

in open sets where u is Lipschitz continuous.

Here $E := Lip'_c(\mathbb{R} \times [0, +\infty[, \mathbb{R})$ denotes the dual of the space of compactly supported Lipschitz continuous functions.

Proof. Since u is Lipschitz continuous and $A \in C^1(\mathbb{R})$, we can write any $\psi \in Lip_c$ in the form $\psi = \phi \exp(-A(u))$ for a suitable $\phi \in Lip_c$. Therefore, for $\psi \in Lip_c(\mathbb{R} \times [0, +\infty[, \mathbb{R})$,

we can write,

$$\begin{aligned}
& - \int_0^\infty \int_{\mathbb{R}} [Z \partial_t \psi + Z \partial_x (a(u) \psi)] \, dx dt + \int_{\mathbb{R}} Z_0(x) \psi(x, 0) \, dx \\
& = - \int_0^\infty \int_{\mathbb{R}} [Z \partial_t (\phi \exp(-A(u))) + Z \partial_x (a(u) \phi \exp(-A(u)))] \, dx dt \\
& \quad + \int_{\mathbb{R}} Z_0(x) \phi(x, 0) \exp(-A(u_0)) \, dx \\
& = - \int_0^\infty \int_{\mathbb{R}} [Z \exp(-A(u)) \partial_t \phi + Z a(u) \exp(-A(u)) \partial_x \phi] \, dx dt \\
& \quad + \int_{\mathbb{R}} Z_0(x) \phi(x, 0) \exp(-A(u_0)) \, dx \\
& \quad + \int_0^\infty \int_{\mathbb{R}} [Z A'(u) \partial_t u \exp(-A(u)) \phi + Z A'(u) \partial_x u \phi a(u) \exp(-A(u))] \, dx dt \\
& \quad - \int_0^\infty \int_{\mathbb{R}} Z \exp(-A(u)) \phi a'(u) \partial_x u \, dx dt \\
& = \int_0^\infty \int_{\mathbb{R}} Z \phi \exp(-A(u)) (A'(u) \partial_t u + A'(u) a(u) \partial_x u - a'(u) \partial_x u) \, dx dt,
\end{aligned} \tag{B.2}$$

where we use the fact that v satisfies the integral identity (1.9) for any $\phi \in Lip_c(\mathbb{R} \times [0, +\infty[, \mathbb{R})$.

Now $\partial_t u$ and $\partial_x f(u)$ belong to L^∞ and hence in L^1_{loc} , and also satisfy the integral identity (1.8) for all compactly supported smooth functions. Therefore $\partial_t u = -\partial_x f(u)$ almost everywhere. Also using the fact that $f \in C^1$ and u is continuous, using Volpert's formula [47], it follows that $-\partial_x f(u) = f'(u) \partial_x u$ almost everywhere.

Using these facts and the definition of A in (B.2), it follows that

$$- \int_0^\infty \int_{\mathbb{R}} [Z \partial_t \psi + Z \partial_x (a(u) \psi)] \, dx dt + \int_{\mathbb{R}} Z_0(x) \psi(x, 0) \, dx = 0,$$

for $\psi \in Lip_c(\mathbb{R} \times [0, +\infty[, \mathbb{R})$. □

Now suppose that f is strictly convex and consider the Riemann problem with left and right states given by (u_-, v_-) and (u_+, v_+) respectively. Let the corresponding states in (u, Z) coordinates be given by (u_-, Z_-) and (u_+, Z_+) respectively. Without loss of generality, we assume that the Riemann problem is based at the point $(0, 0)$.

First we consider the case when there is a rarefaction wave in u . Note that in this case u is a Lipschitz continuous function away from a ball $B(0, r)$ of radius r around $(0, 0)$, for any $r > 0$.

From the previous lemma, we conclude that away from the ball $B(0, r)$, Z (corresponding to any bounded weak solution (u, v)) satisfies the identity (B.1) and by unicity of u and the unicity of weak solutions of a transport equation with Lipschitz continuous velocity field, Z is uniquely determined.

Since this is true for any $r > 0$, it follows that in the class of bounded weak solutions, (u, Z) (and therefore (u, v)) is unique away from the set $\{x = a(u_-)t\}$ of measure zero.

Next let us consider the case when there is a shock wave in u . In this case, using the previous lemma, we see that Z (corresponding to any bounded weak solution (u, v)) satisfies the transport equation (B.1) with velocity $a(u_-)$ in the region

$$\left\{ (x, t) : -\infty < \frac{x}{t} < s := \frac{[f(u)]}{[u]} \right\},$$

and therefore is uniquely determined in this region and has a trace on the line $x = st$. Similarly, Z satisfies the transport equation (B.1) with velocity $a(u_+)$ in the region

$$\left\{ (x, t) : s < \frac{x}{t} < +\infty \right\},$$

and therefore is uniquely determined in this region as well.

But any bounded weak solution (u, v) satisfies the same Rankine-Hugoniot condition across the shock curve, and therefore, the left trace of Z on the line $x = st$ is uniquely determined by the states u_-, u_+, v_+ , which implies the unicity of Z as well.

Therefore, for a strictly convex flux f , at least upto the time of first interaction of waves, the solution of the Riemann problem is unique in the class of bounded entropy solutions.

Note that the above proof holds for initial data u_0 which gives rise to entropy solutions u that are locally Lipschitz except for a finite number of lines, and hence applies to the initial data that we construct in Section 6. Uniqueness for the slightly smaller class of piecewise C^1 entropy solutions are well known [34] and has been already obtained for the (PSA) system [7].

Notice also that for the Kruzkov solution u a continuity in time with respect to L^1_{loc} in space is required [32], but for a nonlinear flux this property is automatically fulfilled [18]. Notice that since u_0 belongs to BV^s , $s > 0$, the solution is continuous in time with values in L^1_{loc} in space [11, 29]. Thus, the initial data is recovered.

References

- [1] Adimurthi, S.S. Ghoshal, and G.D.V. Gowda. Structure of entropy solutions to scalar conservation laws with strictly convex flux, *J. Hyperbolic Differ. Equ.* 4 (2012),571–611.
- [2] F. Ancona and A. Marson. Well-posedness for general 2×2 systems of conservation laws. *Mem. Amer. Math. Soc.* 169 (2004), no. 801, x+170 pp.
- [3] P. Baiti and H.K. Jenssen. Well-posedness for a class of 2×2 conservation laws with L^∞ data, *J. Differential Equations* 140 (1997), no. 1, 161–185.
- [4] S. Bianchini. On the Riemann problem for nonconservative hyperbolic systems. Preprint I.A.C, Roma, 2002
- [5] F. Bouchut and F. James. One-dimensional transport equations with discontinuous coefficients. *Nonlinear Analysis TMA*, 32 (1998), 891-933.
- [6] F. Bouchut and F. James. Duality solutions for pressureless gases, monotone scalar conservations laws and uniqueness. *Comm. Partial Diff. Eq.* 24 (1999), no. 11-12, 2173-2189.

- [7] C. Bourdarias, M. Gisclon, and S. Junca. Some mathematical results on a system of transport equations with an algebraic constraint describing fixed-bed adsorption of gases. *J. Math. Anal. Appl.* 313(2), 551-571, (2006).
- [8] C. Bourdarias, M. Gisclon, and S. Junca. Existence of entropy solutions for gas chromatography with one or two active species and non convex isotherms. *Commun. Math. Sci.* 5(1), 67-84, (2007).
- [9] C. Bourdarias, M. Gisclon, and S. Junca. Hyperbolic models in gas-solid chromatography. *Bol. Soc. Esp. Mat. Apl.* 43, 29-57, (2008).
- [10] C. Bourdarias, M. Gisclon, and S. Junca. Blow up at the hyperbolic boundary for a 2×2 chemical system. *J. Hyperbolic Differ. Equ.* 7 (2010), no.2, 297-316.
- [11] C. Bourdarias, M. Gisclon, and S. Junca. Fractional BV spaces and first applications to conservation laws. *J. Hyperbolic Differ. Equ.* 11 (2014), no. 4, 655-677.
- [12] C. Bourdarias, M. Gisclon, S. Junca, and Y.-Y. Peng. Eulerian and Lagrangian formulations in BV^s for gas-solid chromatography. *Commun. Math. Sci.* 14 (2016), no. 6, 1665-1685.
- [13] C. Bourdarias, M. Gisclon, and S. Junca. Kinetic formulation of a 2×2 hyperbolic system arising in gas chromatography. *Kinet. Relat. Models* 13 (2020), no. 5, 869–888.
- [14] A. Bressan. Hyperbolic Systems of Conservation Laws, The One-Dimensional Cauchy Problem. Oxford lecture series in mathematics and its applications. Oxford University Press, 20, 2000.
- [15] M. Bruneau. La variation totale d'une fonction. (French). Lecture Notes in Mathematics. 413. Springer-Verlag. XIV, 332 p., 1974.
- [16] P. Castelli and S. Junca. Oscillating waves and the maximal smoothing effect for one dimensional nonlinear conservation laws. *AIMS on Applied Mathematics* 8 (2014), 709-716.
- [17] P. Castelli and S. Junca. On the maximal smoothing effect for multidimensional scalar conservation laws. *Nonlinear Anal.* 155 (2017), 207–218.
- [18] G.Q. Chen and M. Rasle. Initial layers and uniqueness of weak entropy solutions to hyperbolic conservation laws. *Arch. Ration. Mech. Anal.* 153 (2000), no. 3, 205–220.
- [19] C.-M. Dafermos. Polygonal approximations of solutions of the initial value problem for a conservation law. *J. Math. Anal. Appl.* 38 (1972),33-41.
- [20] C.-M. Dafermos. Hyperbolic Conservation Laws in Continuum Physics. Springer Verlag, Berlin-Heidelberg, 2016. Fourth edition.
- [21] V. G. Danilov, D. Mitrovic. Delta shock wave formation in the case of triangular hyperbolic system of conservation laws. *J. Differential Equations* 245 (2008), no. 12, 3704–3734.

- [22] C. De Lellis and M. Westdickenberg, On the optimality of velocity averaging lemmas, *Ann. Inst. H. Poincaré, Anal. Nonlinéaire* 20 (6), (2003), 1075–1085.
- [23] S. Ghoshal, B. Guélame, A. Jana, and S. Junca. Optimal BV^s regularity for all time for entropy solutions of conservation laws. *NoDEA Nonlinear Differential Equations Appl.* 27 (2020), no. 5, Paper No. 46, 30 pp.
- [24] S.S. Ghoshal and A. Jana. Optimal jump set in hyperbolic conservation laws. *J. Hyperbolic Differ. Equ.* 17 (2020), no. 4, 765–784.
- [25] B.T. Hayes and P.G. Lefloch. Measure solutions to a strictly hyperbolic system of conservation laws. *Nonlinearity* 9 (1996), no. 6, 1547–1563.
- [26] B. Haspot and S. Junca. BV^s solutions for 2×2 systems of conservation laws with a linearly degenerate field. preprint 2020, hal-02532444
- [27] H. Holden and N.H. Risebro. Front tracking for hyperbolic conservation laws. Second edition. Applied Mathematical Sciences, 152. Springer, Heidelberg, 2015. xiv+515 pp.
- [28] H.K. Jenssen and J. Ridder. One-dimensional scalar conservation laws with regulated data. *SIAM J. Math. Anal.* 52 (2020), no. 3, 3114–3130.
- [29] H.K. Jenssen and J. Ridder. On Φ -variation for 1-d scalar conservation laws. *J. Hyperbolic Differ. Equ.* 17 (2020), no. 4, 843–861.
- [30] S. Junca and B. Lombard. Analysis of a Sugimoto’s model of nonlinear acoustics in an array of Helmholtz resonators. *SIAM J. Appl. Math.* 80 (2020), no. 4, 1704–1722.
- [31] B. Keyfitz and H. C. Kranzer. A strictly hyperbolic system of conservation laws admitting singular shocks. *Nonlinear Evolution Equations That Change Type (IMA Vol. Math. Appl.)* 27 ed B Keyfitz and M Shearer, 1990.
- [32] S. N. Kruzkov. First order quasilinear equations with several independent variables. (Russian) *Mat. Sb. (N.S.)* 81 (123) 1970, 228–255.
- [33] P.D. Lax. Hyperbolic systems of conservation laws II. *Comm. Pure Appl. Math.* 10 (1957), 537-566.
- [34] P.D. Lax. Hyperbolic partial differential equations. With an appendix by Cathleen S. Morawetz. Courant Lecture Notes in Mathematics, 14. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2006. viii+217 pp.
- [35] P. G. LeFloch. An existence and uniqueness result for two nonstrictly hyperbolic systems. *Nonlinear evolution equations that change type* 126–138, IMA Vol. Math. Appl., 27, Springer, New York, 1990.
- [36] P. G. LeFloch. Hyperbolic Systems of Conservation Laws: the theory of classical and nonclassical shock waves. Lectures in Mathematics, ETH Zürich, Birkhäuser, 2002.

- [37] E. Marconi. Regularity estimates for scalar conservation laws in one space dimension. *J. Hyperbolic Differ. Eq.* 15 (2018), no. 4, 623-691.
- [38] E. Marconi. Structure and regularity of solutions to 1D scalar conservation laws. *AIMS Series on Applied Mathematics* 10 (2020), 546-553.
- [39] J. Musielak and W. Orlicz. On generalized variations. *I. Studia Math.* 18 (1959), 11-41.
- [40] W. Neves and D. Serre. Ill-posedness of the Cauchy problem for totally degenerate system of conservation laws. *Electron. J. Differ. Equ.* 2005, Paper No. 124, 25 p.
- [41] O.-A. Oleinik. Discontinuous solutions of nonlinear differential equations, (in russian 1957), *Transl. Amer. Math. Soc. Ser. 2* (1963), no. 26, 95-172.
- [42] Y.-J. Peng. Explicit solutions for 2×2 linearly degenerate systems. *Appl. Math. Lett.* 11 (1998), no. 5, 75-78.
- [43] Y.-J. Peng. Euler-Lagrange change of variables in conservation laws. *Nonlinearity* 20 (2007), no. 8, 1927-1953.
- [44] F. Poupaud and M. Rascle. Measure solutions to the linear multi-dimensional transport equation with non-smooth coefficients. *Comm. Partial Differential Equations* 22 (1997), no. 1-2, 337-358.
- [45] D. Serre. Systems of conservation laws. I: Hyperbolicity, entropy, shock waves. (Systèmes de lois de conservation. I: Hyperbolicité, entropies, ondes de choc.) (French) Fondations. Paris: Diderot Editeur. xii, 298 p. (1996).
- [46] J. Smoller. Shock waves and reaction-diffusion equations. Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 258. Springer-Verlag, New York, 1994. xxiv+632 pp.
- [47] A.I. Volpert, Spaces BV and quasilinear equations. *Mat. Sb. (N.S.)* 73 (1967), no. 115, 255–302.

August 26, 2021