

Properties of linear spectral statistics of frequency-smoothed estimated spectral coherence matrix of high-dimensional Gaussian time series

Philippe Loubaton, Alexis Rosuel

▶ To cite this version:

Philippe Loubaton, Alexis Rosuel. Properties of linear spectral statistics of frequency-smoothed estimated spectral coherence matrix of high-dimensional Gaussian time series. Electronic Journal of Statistics , 2021, 15 (2), pp.5380-5454. hal-02900814v3

HAL Id: hal-02900814 https://hal.science/hal-02900814v3

Submitted on 4 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Properties of linear spectral statistics of frequency-smoothed estimated spectral coherence matrix of high-dimensional Gaussian time series

Philippe Loubaton^{*} and Alexis Rosuel^{*}

Laboratoire d'Informatique Gaspard Monge, UMR 8049, Université Paris-Est Marne la Vallée, France e-mail: alexis.rosuel@univ-eiffel.fr; philippe.loubaton@univ-eiffel.fr

Abstract: The asymptotic behaviour of Linear Spectral Statistics (LSS) of the smoothed periodogram estimator of the spectral coherency matrix of a complex Gaussian high-dimensional time series $(\mathbf{y}_n)_{n\in\mathbb{Z}}$ with independent components is studied under the asymptotic regime where the sample size N converges towards $+\infty$ while the dimension M of \mathbf{y} and the smoothing span of the estimator grow to infinity at the same rate in such a way that $\frac{M}{N} \to 0$. It is established that, at each frequency, the estimated spectral coherency matrix is close from the sample covariance matrix of an independent identically $\mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_M)$ distributed sequence, and that its empirical eigenvalue distribution converges towards the Marcenko-Pastur distribution. This allows to conclude that each LSS has a deterministic behaviour that can be evaluated explicitly. Using concentration inequalities, it is shown that the order of magnitude of the supremum over the frequencies of the deviation of each LSS from its deterministic approximation is of the order of $\frac{1}{M} + \frac{\sqrt{M}}{N} + (\frac{M}{N})^3$ where N is the sample size. Numerical simulations supports our results.

MSC2020 subject classifications: Primary 60B20, 62H15; secondary 62M15.

Keywords and phrases: Random Matrices, Spectral Analysis, High Dimensional Statistics, Time Series, Independence Test.

Contents

1	Intro	$\operatorname{pduction} \ldots $	3	
	1.1	The addressed problem and the results	3	
	1.2	Motivation	5	
	1.3	On the literature	3	
	1.4	General approach 8	3	
	1.5	Assumptions and general notations	L	
	1.6	Overview of the paper 14	1	
2	Useful technical tools			
	2.1	Stochastic domination	5	

*This work is funded by ANR Project HIDITSA, reference ANR-17-CE40-0003.

	<i>P. L</i>	oubaton and A. Rosuel/Coherency matrices of high-dimensional time series	2
	2.2	Properties of the eigenvalues and of the resolvent of	
		large Wishart matrices	16
		2.2.1 Concentration of the largest and the smallest	
		eigenvalues	16
		2.2.2 Asymptotic behaviour of the resolvent of $\frac{X_N X_N^*}{D}$	17
	2.3	Concentration of functionals of Gaussian entries	18
	2.4	Hanson-Wright inequality	19
	2.5	Helffer-Sjöstrand formula	19
3	Stoc	hastic representations of $\tilde{\mathbf{C}}(\nu)$ and $\hat{\mathbf{C}}(\nu)$	21
	3.1	Step 1: Stochastic representation of \tilde{C}	21
	3.2	Step 2: Estimates for $\hat{s}_m(\nu)$	29
	3.3	Step 3: Stochastic representation of \hat{C}	29
4	Stoc	hastic domination of the family $\psi_N(f,\nu), N \ge 1, \nu \in [0,1]$	30
	4.1	Step 1: Evaluation of $\mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\frac{\mathbf{X}_{N}(\nu)\mathbf{X}_{N}^{*}(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^{+}} f \mathrm{d}\mu_{MP}^{(c_{N})}$	31
	4.2	Step 2: Evaluation of $\frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right) - \mathbb{E} \left[\frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right) \right]$	32
	4.3	Step 3: Evaluation of $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right)$.	33
		4.3.1 Reduction to the study of ζ	36
		4.3.2 Proof of Lemma 4.2 and Lemma 4.4 \ldots	42
	4.4	Step 4: evaluation of $\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\mathbf{C}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{K}(\nu)\mathbf{K}(\nu)}{B+1})\right)\right]$	49
	4.5	Estimation of $r_N(\nu)$	53
5	\mathbf{Use}	of Lipschitz properties of functions $\nu \rightarrow \psi_N(f,\nu)$ and	
	$\nu \rightarrow c$	$\hat{\psi}_N(f, u)$	54
	5.1	Lipschitz properties	55
		5.1.1 Lipschitz constant of $\nu \to \frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}(\nu))$	55
		5.1.2 Lipschitz constant of $\nu \rightarrow \hat{r}_N(\nu)$	61
	5.2	Stochastic domination of $\sup_{\nu \in [0,1]} \psi_N(f,\nu) $ and $\sup_{\nu \in [0,1]} \hat{\psi}_N(f,\nu) $	$ _{N}(f,\nu) = 62$
6	Num	nerical simulations	63
\mathbf{A}	App	endix	64
	A.1	Proof of Lemma A.1	64
	A.2	Proof of Lemma 3.1	68
	A.3	Proof of Lemma 3.3	70
	A.4	Proof of Lemma A.5	70
	A.5	Proof of Lemma 4.5	71
	A.6	Proof of Lemma 4.6	72
Re	feren	ces	73

1. Introduction

1.1. The addressed problem and the results

We consider a M-variate zero-mean complex Gaussian stationary time series ¹ $(\mathbf{y}_n)_{n\in\mathbb{Z}}$ and assume that the samples $\mathbf{y}_1, \ldots, \mathbf{y}_N$ are available. We introduce the traditional frequency smoothed periodogram estimate $\hat{\mathbf{S}}(\nu)$ of the spectral density of \mathbf{y} at frequency ν defined by

$$\hat{\mathbf{S}}(\nu) = \frac{1}{B+1} \sum_{b=-B/2}^{B/2} \boldsymbol{\xi}_{\mathbf{y}} \left(\nu + \frac{b}{N}\right) \, \boldsymbol{\xi}_{\mathbf{y}} \left(\nu + \frac{b}{N}\right)^* \tag{1.1}$$

where B is an even integer, which represents the smoothing span, and

$$\boldsymbol{\xi}_{\mathbf{y}}(\nu) = \frac{1}{\sqrt{N}} \sum_{n=1}^{N} \mathbf{y}_n e^{-2i\pi(n-1)\nu}$$
(1.2)

is the renormalized Fourier transform of $(\mathbf{y}_n)_{n=1...,N}$. The corresponding estimated spectral coherency matrix is defined as:

$$\hat{\mathbf{C}}(\nu) = \operatorname{diag}\left(\hat{\mathbf{S}}(\nu)\right)^{-\frac{1}{2}} \hat{\mathbf{S}}(\nu) \operatorname{diag}\left(\hat{\mathbf{S}}(\nu)\right)^{-\frac{1}{2}}$$
(1.3)

where diag($\hat{\mathbf{S}}(\nu)$) = $\hat{\mathbf{S}}(\nu) \odot \mathbf{I}_M$, with \odot denoting the Hadamard product (ie. entrywise product) and \mathbf{I}_M is the *M*-dimensional identity matrix. Under the hypothesis \mathcal{H}_0 that the *M* components $(y_{1,n})_{n \in \mathbb{Z}}, \ldots, (y_{M,n})_{n \in \mathbb{Z}}$ of \mathbf{y} are mutually uncorrelated, we evaluate the behaviour of certain Linear Spectral Statistics (LSS) of the eigenvalues of $\hat{\mathbf{C}}(\nu)$ in asymptotic regimes where $N \to +\infty$ and both M = M(N) and B = B(N) converge towards $+\infty$ in such a way that $M(N) = \mathcal{O}(N^{\alpha})$ for $\alpha \in (1/2, 1)$ and $c_N = \frac{M(N)}{B(N)} \to c$ where $c \in (0, 1)$. We denote by $\mu_{MP}^{(c)}$ the Marcenko-Pastur distribution with parameter c < 1 defined by

$$d\mu_{MP}^{(c)}(\lambda) = \frac{\sqrt{(\lambda_{+} - \lambda)(\lambda - \lambda_{-})}}{2\pi c \lambda} \mathbb{1}_{\lambda \in [\lambda_{-};\lambda_{+}]}(\lambda) \,\mathrm{d}\lambda, \quad \lambda_{\pm} = (1 \pm \sqrt{c})^{2}$$

and define the sequences $(u_N)_{N\geq 1}$ and $(v_N)_{N\geq 1}$ by

$$u_N = \frac{1}{B} + \frac{\sqrt{B}}{N} + \left(\frac{B}{N}\right)^3 \tag{1.4}$$

and

$$v_N = \frac{1}{B+1} \sum_{b=-B/2}^{B/2} \left(\frac{b}{N}\right)^2.$$
 (1.5)

¹any finite linear combination x of the components of $(\mathbf{y}_n)_{n \in \mathbb{Z}}$ is a complex Gaussian random variable, i.e. $\operatorname{Re}(x)$ and $\operatorname{Im}(x)$ are independent zero-mean Gaussian random variables having the same variance

We notice that

$$u_N = \mathcal{O}\left(\frac{1}{B}\right) \mathbf{1}_{\frac{1}{2} \le \alpha \le \frac{2}{3}} + \mathcal{O}\left(\frac{\sqrt{B}}{N}\right) \mathbf{1}_{\frac{2}{3} \le \alpha \le \frac{4}{5}} + \mathcal{O}\left(\frac{B}{N}\right)^3 \mathbf{1}_{\alpha \ge \frac{4}{5}}$$
(1.6)

and $v_N = \mathcal{O}\left((\frac{B}{N})^2\right)$, as well as $\frac{u_N}{v_N} \to 0$ if $\alpha > 2/3$ and $\frac{u_N}{v_N} \to +\infty$ if $\alpha < 2/3$. Then, if $(s_m)_{m=1,...,M}$ represent the spectral densities of the scalar time series $((y_{m,n})_{n\in\mathbb{Z}})_{m=1,...,M}$, for each function f defined on \mathbb{R}^+ and \mathcal{C}^{∞} in a neighbourhood of the support $[\lambda_-; \lambda_+]$ of $\mu_{MP}^{(c)}$, it holds that for each $\epsilon > 0$, there exist a $\gamma(\epsilon) := \gamma > 0$ such that for each N large enough:

$$\mathbb{P}\left[\sup_{\nu\in[0,1]}\left|\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \int_{\mathbb{R}^{+}} f\,\mathrm{d}\mu_{MP}^{(c_{N})} - r_{N}(\nu)\,\phi_{N}(f)\,v_{N}\,\mathbf{1}_{\alpha>2/3}\right| > N^{\epsilon}u_{N}] \leq \exp{-N^{\gamma}} \quad (1.7)$$

where $r_N(\nu)$ is defined by

$$r_N(\nu) = \left(\frac{1}{M} \sum_{m=1}^M \frac{s'_m(\nu)}{s_m(\nu)}\right)^2$$
(1.8)

4

and where $\phi_N(f)$ is a deterministic $\mathcal{O}(1)$ term which coincides with the action of function f on a certain compactly supported distribution D_N (to be precised later) depending on the Marcenko-Pastur distribution $\mu_{MP}^{(c_N)}$. In other words, under \mathcal{H}_0 , uniformly w.r.t. the frequency ν , $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right)$ behaves as $\int_{\mathbb{R}} f \, d\mu_{MP}^{(c_N)}$. If $\alpha \leq 2/3$, with high probability, the order of magnitude of the corresponding error is not larger than $u_N = \frac{1}{B} = \mathcal{O}(\frac{1}{N^{\alpha}})$. If $\alpha > 2/3$, $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \int_{\mathbb{R}} f \, d\mu_{MP}^{(c_N)}$ behaves as the deterministic $\mathcal{O}(\frac{B}{N})^2$ term $r_N(\nu) \, \phi_N(f) \, v_N$, and the rate of convergence towards 0 of the corrected statistics $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \int_{\mathbb{R}^+} f \, d\mu_{MP}^{(c_N)} - r_N(\nu) \, \phi_N(f) \, v_N \, \mathbf{1}_{\alpha>2/3}$ appears to be u_N which verifies $\frac{u_N}{v_N} \to 0$.

Our approach is based on the observation that in the above asymptotic regime, $\hat{\mathbf{S}}(\nu)$ can be interpreted as the sample covariance matrix of the large vectors $(\boldsymbol{\xi}_{\mathbf{y}}(\nu + \frac{b}{N}))_{b=-B/2,...,B/2}$. Classical time series analysis results suggest that the vectors $(\boldsymbol{\xi}_{\mathbf{y}}(\nu + \frac{b}{N}))_{b=-B/2,...,B/2}$ appear as "nearly" i.i.d. zero mean complex random vectors with covariance matrix $\mathbf{S}(\nu)$ where $\mathbf{S}(\nu) =$ diag $(s_1(\nu), \ldots, s_M(\nu))$. $\hat{\mathbf{C}}(\nu)$ can be interpreted as the sample autocorrelation matrix of the above vectors. As it is well-known that the empirical eigenvalue distribution of the sample autocorrelation matrix of i.i.d. large random vectors converges towards the Marcenko-Pastur distribution (see e.g. [21]), it is not surprising that $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right)$ behaves as $\int_{\mathbb{R}^+} f \, d\mu_{MP}^{(c_N)}$. Our main results are thus obtained using tools borrowed from large random matrix theory (see e.g. [30], [1]) and from frequency domain time series analysis techniques (see e.g. [4]).

5

1.2. Motivation

This paper is motivated by the problem of testing whether the components of \mathbf{y} are uncorrelated or not when the dimension M of \mathbf{y} is large and the number of observations N is significantly larger than M. For this, a possible way would be to estimate the spectral coherency matrix, equal to \mathbf{I}_M at each frequency ν under \mathcal{H}_0 , by the standard estimate $\hat{\mathbf{C}}(\nu)$ defined by (1.3) for a relevant choice of B, and to compare, for example, the supremum over ν of the spectral norm $\|\hat{\mathbf{C}}(\nu) - \mathbf{I}_M\|$ to a threshold. To understand the conditions under which such an approach should provide satisfying results, we mention that under some mild extra assumptions, it can be shown that

$$\sup_{\nu} \|\hat{\mathbf{S}}(\nu) - \mathbf{S}(\nu)\| \xrightarrow[N \to +\infty]{a.s.} 0$$

as well as

$$\sup_{\nu} \|\hat{\mathbf{C}}(\nu) - \mathbf{I}_M\| \xrightarrow[N \to +\infty]{a.s.} 0$$

in asymptotic regimes where N, B, M converge towards $+\infty$ in such a way that $\frac{B}{N} \to 0$ and $\frac{M}{B} \to 0$. Therefore, $\hat{\mathbf{C}}(\nu)$ is likely to be close from \mathbf{I}_M for each ν if both $\frac{B}{N}$ and $\frac{M}{B}$ are small enough. However, if M is large and that the number of available samples N is not arbitrarily large w.r.t. M, it may be impossible to choose the smoothing span B in such a way that $\frac{B}{N} \ll 1$ and $\frac{M}{B} \ll 1$. In such a context, the predictions provided by the asymptotic regime $\frac{B}{N} \to 0$ and $\frac{M}{B} \to 0$ will not be accurate, and any test comparing $\hat{\mathbf{C}}(\nu)$ to \mathbf{I}_M for each ν will provide poor results. To solve this issue, we propose to choose B of the same order of magnitude as M. In this case, $\hat{\mathbf{C}}(\nu)$ has of course no reason to be close from \mathbf{I}_M for each ν . If $\frac{M}{N}$, or equivalently if $\frac{B}{N}$ is small enough, the asymptotic regime where both M and B converge towards $+\infty$ at the same rate appears relevant to understand the behaviour of $\hat{\mathbf{C}}(\nu)$. We mention in particular that the condition $\alpha > 1/2$ implies that the rate of convergence of $\frac{M}{N}$ towards 0 is moderate, which is in accordance with practical situations in which the sample size is not arbitrarily large. Our asymptotic results thus suggest that if $\frac{M}{N}$ is small enough and if B is chosen of the same order of magnitude as M, then it seems reasonable to test that the components of y are uncorrelated by comparing

$$\frac{1}{u_N} \sup_{\nu \in [0,1]} \left| \frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) - \int_{\mathbb{R}^+} f \, \mathrm{d}\mu_{MP}^{(c_N)} - \hat{r}_N(\nu) \, \phi_N(f) \, v_N \, \mathbf{1}_{\alpha > 2/3} \right| \right|$$

to a well chosen threshold, where $\hat{r}_N(\nu)$ represents an estimate of $r_N(\nu)$ accurate enough to keep equal to u_N the convergence rate towards 0 of the modified statistics. We notice that our results just characterize the order of magnitude of the above statistics under \mathcal{H}_0 , and that we do not provide asymptotic approximation of its distribution. While the derivation of such an approximation would be quite useful to design a well defined statistical test and to study and compare its performance with existing approaches, our results represent a first necessary

step that has its own interest. We notice that we consider the supremum on the whole frequency interval [0, 1] because, compared to a solution where the maximum is over a low number of fixed frequencies, this allows to increase the power of the test in contexts of alternatives for which, under \mathcal{H}_1 ,

$$\nu \to \left| \frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \int_{\mathbb{R}^+} f \, \mathrm{d}\mu_{MP}^{(c_N)} - \hat{r}_N(\nu) \, \phi_N(f) \, v_N \, \mathbf{1}_{\alpha > 2/3} \right|$$
(1.9)

exhibits narrow peaks that would not be visible on a low density frequency grid. We also mention that other statistics could also be considered, e.g. the integral on the frequency domain of function (1.9) or of this function to the square.

We finally remark that the most usual asymptotic regime considered in the context of large random matrices is $M \to +\infty, N \to +\infty$ in such a way that $\frac{M}{N}$ converges towards a non zero constant. In this regime, it is still possible to develop large random matrix-based approaches testing that the components of \mathbf{y} are uncorrelated or not, see e.g. the contribution [29] to be presented below which, under the extra assumption that the components of \mathbf{y} share the same spectral density, is based on a Gaussian approximation of linear spectral statistics of the empirical covariance matrix $\hat{\mathbf{R}}_N$ defined by

$$\hat{\mathbf{R}}_N = \frac{1}{N} \sum_{n=1}^N \mathbf{y}_n \mathbf{y}_n^* \tag{1.10}$$

6

under \mathcal{H}_0 . However, when the ratio $\frac{M}{N}$ is small enough, the asymptotic regime considered in the present paper seems more relevant than the standard large random matrix regime $M \to +\infty, N \to +\infty$, and test statistics that depend on the estimated spectral coherency matrix $\hat{\mathbf{C}}(\nu)$ should provide better performance than functionals of the matrix $\hat{\mathbf{R}}_N$.

1.3. On the literature

The problem of testing whether various jointly stationary and jointly Gaussian time series are uncorrelated is an important problem that was extensively addressed in the past. Apart from a few works that will be discussed later, almost all the previous contributions addressed the case where the number M of available time series remains finite when the sample size increases. Two classes of methods were mainly studied. The first class uses lag domain approaches based on the observation that M jointly stationary time series $(y_{1,n})_{n\in\mathbb{Z}}, \ldots, (y_{M,n})_{n\in\mathbb{Z}}$ are mutually uncorrelated if and only if for each integer L, the covariance matrix of the ML dimensional vector $\mathbf{y}_n^{(L)}$ defined by

$$\mathbf{y}_{n}^{(L)} = (y_{1,n}, \dots, y_{1,n+L-1}, \dots, y_{M,n}, \dots, y_{M,n+L-1})^{T}$$

is block diagonal. The lag domain approach was in particular used in [17] for M = 2, and extended and developed in [24], [25], [19], [20], [8] and [12].

 $\overline{7}$

The second approach is based on the observation that the M jointly stationary time series $(y_{1,n})_{n\in\mathbb{Z}},\ldots,(y_{M,n})_{n\in\mathbb{Z}}$ are uncorrelated if and only the spectral density matrix $\mathbf{S}(\nu)$ of $\mathbf{y}_n = (y_{1,n},\ldots,y_{M,n})^T$ is diagonal for each frequency ν , or equivalently, if its spectral coherence matrix $\mathbf{C}(\nu)$ is reduced to \mathbf{I}_M for each ν . [35] is one of the first related contribution. This work was followed by [10], [33], as well as [11].

We now review the existing works devoted to the case where the number M of time series converges towards $+\infty$. The particular context where the observations $\mathbf{y}_1, \ldots, \mathbf{y}_N$ are i.i.d. and where the ratio $\frac{M}{N}$ converges towards a constant $d \in (0,1)$ is the most popular. In contrast with the asymptotic regime considered in the present paper, M and N are of the same order of magnitude. This is because, in this context, the time series are mutually uncorrelated if and only the covariance matrix $\mathbb{E}[\mathbf{y}_n \mathbf{y}_n^*]$ is diagonal. Therefore, it is reasonable to consider test statistics that are functionals of the sample covariance matrix \mathbf{R}_N defined by (1.10). In particular, when the observations are Gaussian random vectors, the generalized likelihood ratio test (GLRT) consists in comparing the test statistics $\log \det(\hat{\mathbf{C}}_N)$ to a threshold, where $\hat{\mathbf{C}}_N$ represents the sample autocorrelation matrix. [21] proved that under \mathcal{H}_0 , the empirical eigenvalue distribution of \mathbf{C}_N converges almost surely towards the Marcenko-Pastur distribution $\mu_{MP}^{(d)}$ and therefore, that $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}_N) \right)$ converges towards $\int f d\mu_{MP}^{(d)}$ for each bounded continuous function f. In the Gaussian case, [23] also established a central limit theorem (CLT) for $\log \det(\hat{\mathbf{C}}_N)$ under \mathcal{H}_0 using the moment method. In the real Gaussian case, [7] remarked that $\left(\det \hat{\mathbf{C}}_N\right)^{\tilde{N/2}}$ is the product of independent beta distributed random variables. Therefore, log det $(\hat{\mathbf{C}}_N)$ appears as the sum of independent random variables, thus deducing the CLT. More recently, in [28] is established a CLT on LSS of $\hat{\mathbf{C}}_N$ in the Gaussian case using large random matrix techniques when the covariance matrix $\mathbb{E}[\mathbf{y}_n \mathbf{y}_n^*]$ is not necessarily diagonal. This allows studying the asymptotic performance of the GLRT under a certain class of alternatives. We also mention that [22] studied the behaviour of $\max_{i,j} |(\hat{\mathbf{C}}_N)_{i,j}|$ under \mathcal{H}_0 , and established that $\max_{i,j} |(\hat{\mathbf{C}}_N)_{i,j}|$, after recentering and appropriate normalization, converges in distribution towards a Gumble distribution, which, of course, allows to test the hypothesis \mathcal{H}_0 . This first contribution was extended later in several works, in particular in [6] who considered the case where the samples $\mathbf{y}_1, \ldots, \mathbf{y}_N$ have some specific correlation pattern. Still, in the asymptotic regime $\frac{M}{N} \to d$, [29] proposed to test hypothesis \mathcal{H}_0 when the components of y share the same spectral density. In this case, the rows of the $M \times N$ matrix $(\mathbf{y}_1, \ldots, \mathbf{y}_N)$ are independent and identically distributed under \mathcal{H}_0 . [29] established a central limit theorem for linear spectral statistics of the empirical covariance matrix $\hat{\mathbf{R}}_N$ defined by (1.10), and deduced from this test statistics to check whether \mathcal{H}_0 holds or not. We notice that the results of [29] are valid in the non-Gaussian case.

In our knowledge, no existing work studied the behaviour of linear spectral

8

statistics of matrix $\hat{\mathbf{C}}(\nu)$ in the asymptotic regime defined in the present paper. However, we mention that this regime was considered in [3] in order to solve a completely different problem, i.e. the use of shrinkage in the frequency domain in order to enhance the performance of the spectral density estimate (1.1) when the components of \mathbf{y} are not uncorrelated. We notice that $\frac{B^{3/2}}{N}$ is supposed to converge towards 0 in [3]. When $B = \mathcal{O}(N^{\alpha})$, this condition is equivalent to $\alpha <$ 2/3, while we rather study situations where $\alpha > 1/2$. We finally mention that our works [27] and [31] also consider the present asymptotic regime and study respectively the behaviour of $\sup_{i < j, \nu \in \mathcal{G}_N} |\hat{\mathbf{C}}_{i,j}(\nu)|$ (\mathcal{G}_N is the set $\{k \frac{B+1}{N}, k =$ $0, \ldots, \frac{N}{B+1}\}$) and the largest eigenvalues of $\hat{\mathbf{C}}(\nu)$ in the presence of an extra signal, independent from \mathbf{y} , and having a low-rank spectral density matrix.

1.4. General approach

In order to simplify the notations, we denote by $\psi_N(f,\nu)$ the statistics defined by

$$\psi_N(f,\nu) = \frac{1}{M} \operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \int_{\mathbb{R}^+} f \,\mathrm{d}\mu_{MP}^{(c_N)} - r_N(\nu) \,\phi_N(f) \,v_N \,\mathbf{1}_{\alpha>2/3}.$$
 (1.11)

In order to be able to study the behaviour of $\sup_{\nu} |\psi_N(f,\nu)|$, we establish exponential concentration inequalities that allow to evaluate $\mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N)$ for each ν as well as $\mathbb{P}(\sup_{\nu \in \mathcal{V}_N} |\psi_N(f,\nu)| > N^{\epsilon}u_N)$ for some relevant finite discrete grid \mathcal{V}_N of the interval [0, 1]. (1.7) is then obtained by using Lipschitz properties of function $\nu \to \psi_N(f,\nu)$.

In order to evaluate $\mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N)$ for each ν , we use the following approach:

• We first study the behaviour of the modified sample spectral coherency matrix $\tilde{\mathbf{C}}(\nu)$ defined by

$$\tilde{\mathbf{C}}(\nu) = \operatorname{diag}\left(\mathbf{S}(\nu)\right)^{-\frac{1}{2}} \hat{\mathbf{S}}(\nu) \operatorname{diag}\left(\mathbf{S}(\nu)\right)^{-\frac{1}{2}}.$$
(1.12)

We notice that $\hat{\mathbf{C}}(\nu)$ is obtained from $\hat{\mathbf{C}}(\nu)$ by replacing the estimated diagonal matrix diag $(\hat{\mathbf{S}}(\nu))$ by its true value diag $(\mathbf{S}(\nu))$. Using classical results of [4], we establish that for each ν , $\tilde{\mathbf{C}}(\nu)$ can be represented as

$$\tilde{\mathbf{C}}(\nu) = \frac{\mathbf{X}(\nu)\mathbf{X}^*(\nu)}{B+1} + \tilde{\mathbf{\Delta}}(\nu)$$
(1.13)

where $\mathbf{X}(\nu)$ is a $M \times (B+1)$ random matrix with $\mathcal{N}_{\mathbb{C}}(0,1)$ i.i.d. entries, and $\tilde{\mathbf{\Delta}}(\nu)$ is another matrix such that, for any $\epsilon > 0$, there exists $\gamma > 0$, independent from ν , such that for each large enough $N \in \mathbb{N}$:

$$\mathbb{P}\left[\|\tilde{\mathbf{\Delta}}(\nu)\| > N^{\epsilon} \frac{B}{N}\right] \le \exp{-N^{\gamma}}.$$

We deduce from (1.13) that $\hat{\mathbf{C}}(\nu)$ can be written as

$$\hat{\mathbf{C}}(\nu) = \frac{\mathbf{X}(\nu)\mathbf{X}^*(\nu)}{B+1} + \mathbf{\Delta}(\nu)$$
(1.14)

where $\Delta(\nu)$ verifies the concentration inequality

$$\mathbb{P}\left[\|\boldsymbol{\Delta}(\boldsymbol{\nu})\| > N^{\epsilon} \left(\frac{1}{\sqrt{B}} + \frac{B}{N}\right)\right] \le \exp{-N^{\gamma}}$$

for each $\epsilon > 0$, where γ does not depend on ν . Using (1.13) and (1.14), we establish that the eigenvalues of $\tilde{\mathbf{C}}(\nu)$ and $\hat{\mathbf{C}}(\nu)$ are localized with high probability in a neighbourhood of the support of the Marcenko-Pastur distribution $\mu_{MP}^{(c)}$.

 $\tilde{\mathbf{C}}(\nu)$ appears as a useful intermediate matrix because the study of $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \int_{\mathbb{R}^+} f \, \mathrm{d} \mu_{MP}^{(c_N)}$ is based on the evaluation of each term of the following decomposition:

$$\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \int_{\mathbb{R}^{+}} f \, \mathrm{d}\mu_{MP}^{(c_{N})} = \frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) + \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right] + \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right] + \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] + \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^{+}} f \, \mathrm{d}\mu_{MP}^{(c_{N})}. \tag{1.15}$$

Using the above-mentioned results related to the localization of the eigenvalues of $\tilde{\mathbf{C}}(\nu)$ and $\hat{\mathbf{C}}(\nu)$, we also argue that it is sufficient to do so when f is compactly supported.

• The term $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right)$ is studied using the Helffer-Sjöstrand formula which allows, in a certain sense, to be back to the study of $\frac{1}{M} \operatorname{Tr} \left(\hat{\mathbf{Q}}(z) - \tilde{\mathbf{Q}}(z) \right)$ for $z \in \mathbb{C}^+$, where $\hat{\mathbf{Q}}(z)$ and $\tilde{\mathbf{Q}}(z)$ represent the resolvents of matrices $\hat{\mathbf{C}}(\nu)$ and $\tilde{\mathbf{C}}(\nu)$ (see below for a formal definition). Using (1.13) and (1.14), we express $\frac{1}{M} \operatorname{Tr} \left(\hat{\mathbf{Q}}(z) - \tilde{\mathbf{Q}}(z) \right)$ in terms of the resolvent $\mathbf{Q}(z)$ of matrix $\frac{\mathbf{X}(\nu)\mathbf{X}^*(\nu)}{B+1}$. As matrix $\mathbf{X}(\nu)$ is Gaussian, it is possible to use standard Gaussian tools (Poincaré-Nash inequality and the integration by part formula) to have a good understanding of the behaviour of $\mathbf{Q}(z)$, and to prove that for each $\epsilon > 0$, it exists γ independent from ν such that

$$\mathbb{P}\left(\left|\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s_m''(\nu)}{s_m(\nu)}\right)\,\tilde{\phi}_N(f)\,v_N\,\mathbf{1}_{\alpha>2/3}\right| > N^{\epsilon}u_N\right) \le \exp{-N^{\gamma}}$$

where $\tilde{\phi}_N(f)$ is a deterministic term defined as the action of f on a compactly supported distribution \tilde{D}_N depending on $\mu_{MP}^{(c_N)}$.

• Using a standard Gaussian concentration inequality as well as the structure of matrix $\tilde{\mathbf{C}}(\nu)$, we obtain that for each $\epsilon > 0$, it exists γ independent from ν such that

$$\mathbb{P}\left[\left|\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right]\right| > N^{\epsilon}\frac{1}{B}\right] \le \exp{-N^{\gamma}}$$
(1.16)

for each N large enough.

• We still analyse the deterministic term $\mathbb{E}\left[\frac{1}{M}\mathrm{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\mathrm{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right]$ using the Helffer-Sjöstrand formula. We first show that for each $z \in \mathbb{C}^{+}$, $\mathbb{E}\left[\frac{1}{M}\mathrm{Tr}(\tilde{\mathbf{Q}}(z) - \mathbf{Q}(z))\right]$ is a $\mathcal{O}(\frac{B}{N})^{2}$ term, a non obvious result because the evaluation (1.13) just leads to the conclusion that the above term is $\mathcal{O}(\frac{B}{N})$. Moreover, using long and very tedious Gaussian calculations, we obtain that if $\alpha > \frac{2}{3}$, it holds that

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}(\tilde{\mathbf{Q}}(z) - \mathbf{Q}(z))\right] = -\left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s_m''(\nu)}{s_m(\nu)}\right) \tilde{p}_N(z) v_N + \left(\frac{1}{M}\sum_{m=1}^{M}\frac{s_m'(\nu)}{s_m(\nu)}\right)^2 p_N(z) v_N + \mathcal{O}\left(\frac{B}{N}\right)^3$$

where p_N and \tilde{p}_N are the Stieltjes transforms of the compactly supported distributions D_N and \tilde{D}_N introduced previously. This immediately implies that if $\alpha \leq \frac{2}{3}$, then

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right]$$
$$= \mathcal{O}\left(\frac{B}{N}\right)^{2} = o\left(\frac{1}{B}\right) = o(u_{N})$$

while if $\alpha > \frac{2}{3}$, then,

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] = -\left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s_{m}'(\nu)}{s_{m}(\nu)}\right) \tilde{\phi}_{N}(f) v_{N} + \left(\frac{1}{M}\sum_{m=1}^{M}\frac{s_{m}'(\nu)}{s_{m}(\nu)}\right)^{2} \phi_{N}(f) v_{N} + \mathcal{O}(u_{N})$$

because $(\frac{B}{N})^3 \ll u_N$ if $2/3 < \alpha \le 4/5$ and $(\frac{B}{N})^3$ is equivalent to u_N if $\alpha > 4/5$.

• Finally, classical results imply that

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^{+}} f \,\mathrm{d}\mu_{MP}^{(c_{N})} = \mathcal{O}\left(\frac{1}{B^{2}}\right) = o(u_{N})$$

Gathering the above evaluations and using the Lipschitz properties of function $\nu \to \psi(f, \nu)$, we eventually obtain (1.7).

We also indicate how the use of lag window estimators of the spectral densities $(s_m)_{m=1,...,M}$ allows to design an estimator $\hat{r}_N(\nu)$ of $r_N(\nu)$ defined by (1.8) for which the rate of convergence towards 0 of the statistics $\hat{\psi}_N(f,\nu)$ obtained by replacing $r_N(\nu)$ by $\hat{r}_N(\nu)$ in Eq. (1.11) is still u_N . In particular, we establish that for each $\epsilon > 0$, $\mathbb{P}\left(\sup_{\nu} |\hat{\psi}_N(f,\nu)| > N^{\epsilon}u_N\right)$ converges towards 0 exponentially.

1.5. Assumptions and general notations

Assumption 1.1. For each $m \ge 1$, $(y_{m,n})_{n \in \mathbb{Z}}$ is a zero mean stationary complex Gaussian time series, ie.

- 1. $\mathbb{E}[y_{m,n}] = 0$ for any $m \ge 1$ and any $n \in \mathbb{Z}$
- 2. every finite linear combination x of the random variables $(y_{m,n})_{n\in\mathbb{Z}}$ is a $\mathcal{N}_{\mathbb{C}}(0,\sigma^2)$ distributed random variable for some σ^2 , i.e. $\operatorname{Re}(x)$ and $\operatorname{Im}(x)$ are independent and $\mathcal{N}(0,\sigma^2/2)$ distributed.

Assumption 1.2. If $m_1 \neq m_2$, then the scalar time series $(y_{m_1,n})_{n \in \mathbb{Z}}$ and $(y_{m_2,n})_{n \in \mathbb{Z}}$ are independent.

We now formulate the following assumptions on the growth rate of the quantities N, M, B:

Assumption 1.3.

$$B, M = \mathcal{O}(N^{\alpha}) \text{ where } \frac{1}{2} < \alpha < 1, \quad \frac{M}{B+1} = c_N, \quad c_N \xrightarrow[N \to +\infty]{} c \in (0, 1).$$

As M = M(N) converges towards $+\infty$, we assume that an infinite sequence $(y_{1,n})_{n \in \mathbb{Z}}, (y_{2,n})_{n \in \mathbb{Z}}, \ldots, (y_{k,n})_{n \in \mathbb{Z}}, \ldots$ of mutually independent zero mean complex Gaussian time series is given.

We denote by $(s_m)_{m\geq 1}$ the corresponding sequence of spectral densities (i.e. s_m coincides with the spectral density of the times series $(y_{m,n})_{n\in\mathbb{Z}}$). For each $m\geq 1$, we denote by $r_m = (r_{m,u})_{u\in\mathbb{Z}}$ the autocovariance sequence of $(y_{m,n})_{n\in\mathbb{Z}}$, i.e. $r_{m,u} = \mathbb{E}[y_{m,n+u}y_{m,n}^*]$. We formulate the following assumptions on $(s_m)_{m\geq 1}$ and $(r_m)_{m\geq 1}$:

Assumption 1.4. The time series $((y_{m,n})_{n \in \mathbb{Z}})_{m \geq 1}$ are such that:

$$\inf_{m \ge 1} \inf_{\nu \in [0,1]} |s_m(\nu)| > 0 \tag{1.17}$$

and

$$\sup_{m \ge 1} \sum_{u \in \mathbb{Z}} (1+|u|)^{\gamma_0} |r_{m,u}| < +\infty$$
(1.18)

where $\gamma_0 \geq 3$. Assumption (1.18) of course implies that the spectral densities $(s_m)_{m\geq 1}$ are \mathcal{C}^3 and that

$$\sup_{m \ge 1} \sup_{\nu \in [0,1]} |s_m^{(i)}(\nu)| < +\infty$$
(1.19)

for i = 0, 1, 2, 3 ($s_m^{(i)}$ represents the derivative of order i of s_m). We notice that (1.18) holds as soon as we have

$$\sup_{m \ge 1} |r_{m,u}| \le \frac{C}{|u|^{1+\gamma_0+\delta}}$$

for each $u \neq 0$ as well as $\sup_{m\geq 1} |r_{m,0}| < \infty$ (C > 0 and $\delta > 0$ represent constants). If z represents the backward shift operator, a simple example of time series verifying Assumption 1.4 is to consider ARMA time series generated as

$$y_{m,n} = [h_m(z)]\epsilon_{m,n}$$

where $((\epsilon_{m,n})_{n\in\mathbb{Z}})_{m\geq 1}$ are mutually independent i.i.d. $\mathcal{N}_{\mathbb{C}}(0,1)$ sequences, and where $h_m(z) = \frac{b_m(z)}{a_m(z)}$, a_m and b_m being 2 polynomials having no pole nor zero in the closed unit disk $\overline{\mathbb{D}}$. Moreover, $\sup_{m\geq 1} \max(\deg(a_m), \deg(b_m)) < +\infty$, and if $(z_{k,m})_{k=1,\dots,\deg(b_m)}$ and $(p_{k,m})_{k=1,\dots,\deg(a_m)}$ are the zeros of b_m and a_m , then we should have

$$\inf_{m \ge 1} \inf_{k} \operatorname{dist}(z_{k,m}, \overline{\mathbb{D}}) > 0, \ \inf_{m \ge 1} \inf_{k} \operatorname{dist}(p_{k,m}, \overline{\mathbb{D}}) > 0 \\ \sup_{m \ge 1} \sup_{k} |z_{k,m}| < +\infty, \ \sup_{m \ge 1} \sup_{k} |p_{k,m}| < +\infty.$$

It is easy to check that (1.18) holds for each $\gamma_0 > 0$, and that (1.17) is verified as well.

Notations. A zero mean complex valued random vector \mathbf{y} is said to be $\mathcal{N}_{\mathbb{C}}(0, \mathbf{\Sigma})$ distributed if $\mathbb{E}(\mathbf{y}\mathbf{y}^*) = \mathbf{\Sigma}$ and if each linear combination x of the entries of \mathbf{y} is a complex Gaussian random variable, i.e. $\operatorname{Re}(x)$ and $\operatorname{Im}(x)$ are

independent Gaussian random variables sharing the same variance. If x is a random variable, we denote by x° the random variable defined by

$$x^{\circ} = x - \mathbb{E}[x]. \tag{1.20}$$

If **A** is a $P \times Q$ matrix, $\|\mathbf{A}\|$ and $\|\mathbf{A}\|_F$ denote its spectral norm and Frobenius norm respectively. If P = Q and **A** is Hermitian, $\lambda_1(\mathbf{A}) \geq \ldots \geq \lambda_P(\mathbf{A})$ are the eigenvalues of **A**. The spectrum of **A**, which is here the set of its eigenvalues $(\lambda_k(\mathbf{A}))_{k=1,\ldots,P}$, is denoted by $\sigma(\mathbf{A})$. For **A** and **B** square Hermitian matrices, if all the eigenvalues of $\mathbf{A} - \mathbf{B}$ are non negative, we write $\mathbf{A} \geq \mathbf{B}$. We define Re $\mathbf{A} = (\mathbf{A} + \mathbf{A}^*)/2$ and Im $\mathbf{A} = (\mathbf{A} - \mathbf{A}^*)/2$ where \mathbf{A}^* is the conjugate transpose of matrix **A**.

 \mathcal{C}^p represents the set of all real-valued functions defined on \mathbb{R} whose p first derivatives exist and are continuous, and \mathcal{C}^p_c is the set of all compactly supported functions of \mathcal{C}^p .

We recall that $\mathbf{S}(\nu)$ represents the $M \times M$ diagonal matrix $\mathbf{S}(\nu) = \text{diag}(s_1(\nu), \ldots, s_M(\nu))$. We notice that \mathbf{S} depends on M, thus on N (through M := M(N)), but we often omit to mention the corresponding dependency in order to simplify the notations. In the following, we will denote by \mathbf{y}_m the N-dimensional vector $\mathbf{y}_m = (y_{m,1}, \ldots, y_{m,N})^T$.

A nice constant is a positive a constant that does not depend on the frequency ν , the time series index m, the complex variable z of the various resolvents and Stieltjes transforms used throughout the paper, as well as on the dimensions B, M and N. A nice polynomial is a polynomial whose degree and coefficients are nice constants. If $z \in \mathbb{C}^+$ and if P_1 and P_2 are two nice polynomials, terms such as $P_1(z)P_2(\frac{1}{\text{Im}z})$ play an important role in the following. C and C(z) will represent a generic notation for respectively a nice constant and a term $P_1(z)P_2(\frac{1}{\text{Im}z})$, and the values of C and C(z) may change from one line to the other.

If $(a_N)_{N\geq 1}$ and $(b_N)_{N\geq 1}$ are two sequences of positive real numbers, we write $a_N \ll b_N$ if $\frac{a_N}{b_N} \to 0$ when $N \to +\infty$.

We also recall how a function can be applied to Hermitian matrices. For a $M \times M$ Hermitian matrix **A** with spectral decomposition \mathbf{UAU}^* where $\mathbf{A} = \text{diag}(\lambda_m, m = 1, \dots, M)$ and the $(\lambda_m)_{m=1,\dots,M}$ are the real eigenvalues of **A**, then for any function f defined on \mathbb{R} , we define $f(\mathbf{A})$ as:

$$f(\mathbf{A}) = \mathbf{U} \begin{pmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_M) \end{pmatrix} \mathbf{U}^*$$

 \mathbb{C}^+ is the upper half-plane of $\mathbb{C},$ i.e. the set of all complex numbers z for

which $\operatorname{Im} z > 0$.

For μ a probability measure, its Stieltjes transform s_{μ} is the function defined on $\mathbb{C} \setminus \text{Supp } \mu$ as

$$s_{\mu}(z) = \int \frac{\mathrm{d}\mu(\lambda)}{\lambda - z}.$$
(1.21)

We recall that

$$|s_{\mu}(z)| \le \frac{1}{\operatorname{Im} z} \tag{1.22}$$

for each $z \in \mathbb{C}^+$. Moreover, if μ is carried by \mathbb{R}^+ , then for any a > 0, function $-\frac{1}{z(1+as_{\mu}(z))}$ is also the Stieljes transform of a probability distribution carried by \mathbb{R}^+ , a property which implies that

$$\left|\frac{1}{1+as_{\mu}(z)}\right| \le \frac{|z|}{\mathrm{Im}z} \tag{1.23}$$

for each $z \in \mathbb{C}^+$ (see [15], Proposition 5-1, item 4).

If $\lambda_1, \ldots, \lambda_M$ denote the eigenvalues of an Hermitian matrix **A** and if $\mu := \frac{1}{M} \sum_{i=1}^{M} \delta_{\lambda_i}$ denotes the empirical eigenvalue distribution of **A**, then we have the following relation:

$$s_{\mu}(z) = \frac{1}{M} \operatorname{tr} \mathbf{Q}_{\mathbf{A}}(z)$$

where $\mathbf{Q}_{\mathbf{A}}(z)$ represents the resolvent of \mathbf{A} defined by

$$\mathbf{Q}_{\mathbf{A}}(z) = (\mathbf{A} - z\mathbf{I}_M)^{-1}.$$
 (1.24)

We finally mention the following useful control for the norm $\mathbf{Q}_{\mathbf{A}}$. For each $z \in \mathbb{C}^+$, we have

$$\|\mathbf{Q}_{\mathbf{A}}\| \le \frac{1}{\operatorname{Im} z}.\tag{1.25}$$

1.6. Overview of the paper

We first recall in Section 2 useful technical tools: in Paragraph 2.1, the concept of stochastic domination adapted from [13] which allows to considerably simplify the exposition of the following results, in Paragraph 2.2 some useful properties of the extreme eigenvalues and of the resolvent of large Wishart matrices, two well-known Gaussian concentration inequalities expressed using the stochastic domination framework in Paragraphs 2.3 and 2.4, and the Helffer-Sjöstrand formula in Paragraph 2.5. We establish in Section 3 the stochastic representations (1.13) and (1.14) of $\tilde{\mathbf{C}}(\nu)$ and $\hat{\mathbf{C}}(\nu)$. In Section 4, we prove for each ν the concentration of $|\psi_N(f,\nu)|$ defined by (1.11), and indicate how it is possible to estimate the term $r_N(\nu)$ in order to keep equal to u_N the rate of convergence of the statistics $\hat{\psi}_N(f,\nu)$ obtained by replacing $r_N(\nu)$ by $\hat{r}_N(\nu)$ in (1.11). In Section 5, we establish Lipschitz properties for the functions $\nu \to \psi_N(f,\nu)$ and $\nu \to \hat{\psi}_N(f,\nu)$ that allow to establish the concentration of $\sup_{\nu} |\psi_N(f,\nu)|$ and $\sup_{\nu} |\hat{\psi}_N(f,\nu)|$. We finally provide in Section 6 some numerical simulations that support our results.

2. Useful technical tools

2.1. Stochastic domination

We now present the concept of stochastic domination introduced in [13]. A nice introduction to this tool can also be found in the lecture notes [2].

Definition 2.1. Stochastic Domination. Let

$$X = (X^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)}), \quad Y = (Y^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)})$$

be two families of nonnegative random variables, where $U^{(N)}$ is a set that may possibly depend on N. We say that X is stochastically dominated by Y if for all (small) $\epsilon > 0$, there exists some $\gamma > 0$ (which of course depends on ϵ) such that:

$$\mathbb{P}\left[X^{(N)}(u) > N^{\epsilon}Y^{(N)}(u)\right] \le \exp{-N^{\gamma}}$$

for each $u \in U^{(N)}$ and for each large enough $N > N_0(\epsilon)$, where $N_0(\epsilon)$ is independent of u, or equivalently

$$\sup_{u \in U^{(N)}} \mathbb{P}\left[X^{(N)}(u) > N^{\epsilon} Y^{(N)}(u)\right] \le \exp{-N^{\gamma}}.$$
(2.1)

for each large enough $N > N_0(\epsilon)$. If X is stochastically dominated by Y we use the notation $X^{(N)}(u) \prec Y^{(N)}(u)$. To simplify the notations, we will very often denote $X^{(N)} \prec Y^{(N)}$ or $X \prec Y$ when the context will be clear enough. Moreover, if for some complex valued family X we have $|X| \prec Y$ we also write $X = \mathcal{O}_{\prec}(Y)$.

Finally, we say that a family of events $\Xi = \Xi^{(N)}(u)$ holds with exponentially high (small) probability if there exist N_0 and $\gamma > 0$ such that for $N \ge N_0$, $\mathbb{P}[\Xi_N(u)] > 1 - \exp{-N^{\gamma}}$ ($\mathbb{P}[\Xi_N(u)] < \exp{-N^{\gamma}}$) for each $u \in U^{(N)}$.

Remark 2.1. Suppose $(X_N)_{N \in \mathbb{N}}$ is a sequence of positive random variables, satisfying $X_N \prec a_N N^{\epsilon}$ for any $\epsilon > 0$ for some positive real numbers sequence $(a_N)_{N \in \mathbb{N}}$. It turns out that this precisely means that $X_N \prec a_N$. Indeed, consider an arbitrary $\epsilon' > 0$. By the stochastic domination property of X_N , one can take ϵ such that $0 < \epsilon < \epsilon'$ and write

$$\mathbb{P}\left[X_N > a_N \times N^{\epsilon'}\right] \le \mathbb{P}\left[X_N > a_N \times N^{\epsilon} \times \underbrace{N^{\epsilon'-\epsilon}}_{\gg 1}\right] \le \mathbb{P}\left[X_N > a_N \times N^{\epsilon}\right]$$

which goes to zero exponentially since $X_N \prec a_N N^{\epsilon}$ for the ϵ chosen. This argument will be used in the proof of Lemma 4.2.

Lemma 2.1. Take four families of non negative random variables X_1, X_2, Y_1 and Y_2 defined as in Definition 2.1. Then the following holds:

$$X_1 \prec Y_1 \text{ and } X_2 \prec Y_2 \implies X_1 + X_2 \prec Y_1 + Y_2 \text{ and } X_1 X_2 \prec Y_1 Y_2.$$

We omit the proof of this lemma.

Remark 2.2. Note that Definition 2.1 is slightly different from the original one [13] which states that the left hand side of (2.1) should be bounded by a quantity of order N^{-D} for any finite D > 0. In the present paper, all the random variables are Gaussian, and exponential concentration rates can be achieved.

2.2. Properties of the eigenvalues and of the resolvent of large Wishart matrices

In this paper we will at multiple occasion use properties of the eigenvalues of matrices $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$ where \mathbf{X}_N is a $M \times (B+1)$ complex Gaussian matrix with i.i.d. $\mathcal{N}_{\mathbb{C}}(0,1)$ entries when M = M(N) and B = B(N) follow Assumption 1.3.

2.2.1. Concentration of the largest and the smallest eigenvalues

We first recall concentration results of the largest and smallest eigenvalue of $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$ due to [14]. We have for any $\epsilon > 0$

$$\mathbb{P}\left[\lambda_M\left(\frac{\mathbf{X}_N\mathbf{X}_N^*}{B+1}\right) < (1-\sqrt{c})^2 - \epsilon\right] \leq (B+1)\exp{-C(B+1)\epsilon^2} \quad (2.2)$$
$$\mathbb{P}\left[\lambda_1\left(\frac{\mathbf{X}_N\mathbf{X}_N^*}{B+1}\right) > (1+\sqrt{c})^2 + \epsilon\right] \leq (B+1)\exp{-C(B+1)\epsilon^2} \quad (2.3)$$

for some nice constant C.

Consider for $\epsilon > 0$, the ϵ -expansion of the support of the Marchenko-Pastur distribution $\mu_{MP}^{(c)}$:

$$\operatorname{Supp} \mu_{MP}^{(c)} + \epsilon := \left[(1 - \sqrt{c})^2 - \epsilon, (1 + \sqrt{c})^2 + \epsilon \right]$$

and the event:

$$\Lambda_{N,\epsilon} = \left\{ \sigma \left(\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1} \right) \subset \operatorname{Supp} \mu_{MP}^{(c)} + \epsilon \right\}.$$
(2.4)

It is clear that using (2.2) and (2.3), $\Lambda_{N,\epsilon}$ holds with exponentially high probability for any $\epsilon > 0$. This will be of high importance in the following since it will enable us to work on events of exponentially high probability where the norm of $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$ and the norm of its inverse are bounded.

Eventually, the following (weaker) statement is a simple consequence of the equations (2.2) and (2.3), which will sometimes be enough in the following:

$$\lambda_1 \left(\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1} \right) + \frac{1}{\lambda_M \left(\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1} \right)} \prec 1.$$
 (2.5)

We finally notice that if we consider a family $\mathbf{X}_N(u) \in \mathbb{C}^{M \times (B+1)}$ with i.i.d. $\mathcal{N}_{\mathbb{C}}(0,1)$ entries, $u \in U^{(N)}$, where $U^{(N)}$ is a certain set possibly depending on N, then (2.2) and (2.3) hold for each $u \in U^{(N)}$ because the constant C in (2.2) and (2.3) is universal. This implies that the stochastic domination (2.5) is still verified by the family $\mathbf{X}_N(u)$, $u \in U^{(N)}$. Moreover, the family of events $\Lambda_{N,\epsilon}(u)$ defined by (2.4) when \mathbf{X}_N is replaced by $\mathbf{X}_N(u)$ still holds with exponentially high probability.

2.2.2. Asymptotic behaviour of the resolvent of $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$

We next review known results related to the asymptotic behaviour of the resolvent $\mathbf{Q}_N(z)$ of matrix $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$ that can be deduced from standard Gaussian tools. The Poincaré-Nash inequality (see e.g. [30, Proposition 2.1.6] in the Gaussian real case and Eq. (18) in [16] in the complex Gaussian case) implies immediately that the following Lemma holds.

Lemma 2.2. Consider deterministic $M \times M$ and $(B+1) \times (B+1)$ matrices **A** and $\tilde{\mathbf{A}}$. Then, it holds that

$$\operatorname{Var} \frac{1}{M} \operatorname{tr} \mathbf{A} \mathbf{Q}_{N}^{i}(z) \leq \frac{C(z)}{M^{2}} \frac{1}{M} \operatorname{tr} \mathbf{A} \mathbf{A}^{*}$$
(2.6)

$$\operatorname{Var} \frac{1}{M} \operatorname{tr} \left(\frac{\mathbf{X} \tilde{\mathbf{A}} \mathbf{X}^*}{B+1} \mathbf{Q}_N^i(z) \right) \leq \frac{C(z)}{M^2} \frac{1}{B+1} \operatorname{tr} \tilde{\mathbf{A}} \tilde{\mathbf{A}}^*$$
(2.7)

for i = 1, 2

We recall that C(z) represents a generic notation for $P_1(z)P_2(\frac{1}{\text{Im}z})$ where P_1 and P_2 are nice polynomials.

The integration by parts formula states that if $h(\mathbf{X}, \mathbf{X}^*)$ is a \mathcal{C}^1 function of the entries of \mathbf{X} and \mathbf{X}^* with polynomially bounded first derivatives, then, it holds that

$$\mathbb{E}(X_{ij}h(\mathbf{X}, \mathbf{X}^*)) = \mathbb{E}|X_{ij}|^2 \mathbb{E}\left[\frac{\partial h}{\partial \overline{X}_{ij}}(\mathbf{X}, \mathbf{X}^*)\right].$$
(2.8)

(2.8), in conjunction with the Poincaré-Nash inequality, allows to evaluate easily the asymptotic behaviour of the entries of $\mathbb{E}(\mathbf{Q}_N(z))$ (see e.g. [30]). We first notice that properties of the distribution of matrix \mathbf{X}_N immediately imply that $\mathbb{E}(\mathbf{Q}_N(z))$ is reduced to $\beta_N(z)\mathbf{I}_M$ where $\beta_N(z)$ coincides with $\mathbb{E}(\mathbf{Q}_{m,m}(z))$ for each *m*. Then, it holds that

$$\beta_N(z) = t_N(z) + \epsilon_N(z) \tag{2.9}$$

where the error term $\epsilon_N(z)$ verifies $|\epsilon_N(z)| \leq \frac{C(z)}{M^2}$ and where $t_N(z)$ is the Stieltjes transform of the Marcenko-Pastur distribution $\mu_{MP}^{(c_N)}$. In other words, $t_N(z)$ is the unique Stieltjes transform satisfying the equation

$$t_N(z) = \frac{1}{-z + \frac{1}{1 + c_N t_N(z)}}.$$
(2.10)

It is also convenient to define $\tilde{t}_N(z)$ by

$$\tilde{t}_N(z) = -\frac{1}{z(1 + c_N t_N(z))}$$
(2.11)

so that $t_N(z)$ is also given by

$$t_N(z) = -\frac{1}{z(1+\tilde{t}_N(z))}.$$
(2.12)

It is well-known that $\tilde{t}_N(z)$ is the Stieltjes transform of the probability distribution $c_N \mu_{MP}^{(c_N)} + (1 - c_N) \delta_0$.

We finally mention that $\mathbb{E}(\mathbf{Q}'_N(z)) = \mathbb{E}(\mathbf{Q}^2_N(z)) = \beta'_N(z)\mathbf{I}_M$ (where ' stands for the derivative w.r.t. z), and that $\epsilon'_N(z) = \beta'_N(z) - t'_N(z)$ still satisfies

$$|\epsilon'_N(z)| \le \frac{C(z)}{M^2}.$$
 (2.13)

2.3. Concentration of functionals of Gaussian entries

It is well-known (see e.g. [34, Th. 2.1.12]) that for any 1-Lipschitz real valued function f defined on \mathbb{R}^N and any N-dimensional random variable $\mathbf{X} \sim \mathcal{N}(0, \mathbf{I}_N)$, there exists a universal constant C such that:

$$\mathbb{P}\left[|f(\mathbf{X}) - \mathbb{E}f(\mathbf{X})| > t\right] \le C \exp{-Ct^2}.$$
(2.14)

This inequality is still valid when $\mathbf{X} \sim \mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_N)$: in this context, $f(\mathbf{X})$ is replaced by a real-valued function $f(\mathbf{X}, \mathbf{X}^*)$ depending on the entries of \mathbf{X} and \mathbf{X}^* . $f(\mathbf{X}, \mathbf{X}^*)$ can of course be written as $f(\mathbf{X}, \mathbf{X}^*) = \tilde{f}(\sqrt{2}\operatorname{Re}(\mathbf{X}), \sqrt{2}\operatorname{Im}(\mathbf{X}))$ for some function \tilde{f} defined on \mathbb{R}^{2N} . As $(\sqrt{2}\operatorname{Re}(\mathbf{X}), \sqrt{2}\operatorname{Im}(\mathbf{X}))$ is $\mathcal{N}(0, \mathbf{I}_{2N})$ distributed, the concentration inequality is still valid for $f(\mathbf{X}, \mathbf{X}^*) = \tilde{f}(\sqrt{2}\operatorname{Re}(\mathbf{X}), \sqrt{2}\operatorname{Im}(\mathbf{X}))$. We just finally mention that f, considered as a function of $(\mathbf{X}, \mathbf{X}^*)$, and \tilde{f} have Lipschitz constants that are of the same order of magnitude. More precisely, if we define the differential operators $\frac{\partial}{\partial z}$ and $\frac{\partial}{\partial \overline{z}}$ by

$$\frac{\partial}{\partial z} = \frac{\partial}{\partial x} - i\frac{\partial}{\partial y}, \quad \frac{\partial}{\partial \bar{z}} = \frac{\partial}{\partial x} + i\frac{\partial}{\partial y}$$

we can verify immediately that

$$\sum_{i=1}^{N} \left(\left| \frac{\partial f}{\partial X_i} \right|^2 + \left| \frac{\partial f}{\partial X_i^*} \right|^2 \right) = \| \left(\nabla f \right)_{(\mathbf{X}, \mathbf{X}^*)} \|^2 = 4 \| \left(\nabla \tilde{f} \right)_{(\sqrt{2} \operatorname{Re}(\mathbf{X}), \sqrt{2} \operatorname{Im}(\mathbf{X}))} \|^2.$$

Within the stochastic domination framework, the concentration inequality (2.14) implies that for a family $\mathbf{X}_N(u) \sim \mathcal{N}(0, \mathbf{I}_N)$ for $u \in U^{(N)}$:

$$|f(\mathbf{X}_N(u)) - \mathbb{E}f(\mathbf{X}_N(u))| \prec 1$$

The proof is immediate: consider $\epsilon > 0$ and obtain that

$$\mathbb{P}[|f(\mathbf{X}_N(u)) - \mathbb{E}f(\mathbf{X}_N(u))| > N^{\epsilon}] \le C \exp - CN^{2\epsilon}$$

for each u as expected. This result can easily be extended in the complex case, i.e. when $\mathbf{X}_N(u) \sim \mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_N)$.

2.4. Hanson-Wright inequality

The Hanson-Wright inequality [32] is useful to control deviations of a quadratic form from its expectation. While it is proved in the real case in [32], it can easily be understood that it can be extended in the complex case as follows: let $\mathbf{X} \sim \mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_N)$ and $\mathbf{A} \in \mathbb{C}^{N \times N}$. Then

$$\mathbb{P}[|\mathbf{X}^*\mathbf{A}\mathbf{X} - \mathbb{E}\mathbf{X}^*\mathbf{A}\mathbf{X}| > t] \le 2\exp{-C\min\left(\frac{t^2}{\|\mathbf{A}\|_F^2}, \frac{t}{\|\mathbf{A}\|}\right)}.$$
 (2.15)

We now write (2.15) in the stochastic domination framework. Consider a family of independent $\mathcal{N}_{\mathbb{C}}(0,1)$ random variables $(X_n(u))_{n=1,...,N}$ where $u \in U^{(N)}$ and a sequence of $N \times N$ matrices $\mathbf{A}_N(u)$ that possibly depend on u. Take $\epsilon > 0$ and $t = N^{\epsilon} \|\mathbf{A}_N(u)\|_F$. Since $\|\mathbf{A}_N(u)\| > 0$, $\|\mathbf{A}_N(u)\|_F > 0$, and $\|\mathbf{A}_N(u)\|_F$:

$$\min\left(\frac{t}{\|\mathbf{A}_N(u)\|}, \frac{t^2}{\|\mathbf{A}_N(u)\|_F^2}\right) = \min\left(N^{\epsilon} \frac{\|\mathbf{A}_N(u)\|_F}{\|\mathbf{A}_N(u)\|}, N^{2\epsilon} \frac{\|\mathbf{A}_N(u)\|_F^2}{\|\mathbf{A}_N(u)\|_F^2}\right)$$
$$\geq \min(N^{\epsilon}, N^{2\epsilon}) = N^{\epsilon}.$$

Denote $\mathbf{X}_N(u) = (X_1(u), \dots, X_N(u))^T$. For any $u \in U^{(N)}$, it holds that:

$$\mathbb{P}\left[|\mathbf{X}_{N}^{*}(u)\mathbf{A}_{N}(u)\mathbf{X}_{N}(u) - \mathbb{E}\mathbf{X}_{N}^{*}(u)\mathbf{A}_{N}(u)\mathbf{X}_{N}(u)| > N^{\epsilon}\|\mathbf{A}_{N}(u)\|_{F}\right] \leq 2\exp{-CN^{\epsilon}}.$$
 (2.16)

We can therefore rewrite (2.16) as the following stochastic domination:

$$\|\mathbf{X}_{N}^{*}(u)\mathbf{A}_{N}(u)\mathbf{X}_{N}(u) - \mathbb{E}\mathbf{X}_{N}^{*}(u)\mathbf{A}_{N}(u)\mathbf{X}_{N}(u)\| \prec \|\mathbf{A}_{N}(u)\|_{F}.$$
 (2.17)

2.5. Helffer-Sjöstrand formula

If μ is a probability measure, the Helffer-Sjöstrand formula can be seen as an alternative to the Stieltjes inversion formula that allows to express $\int f d\mu$ in terms of the Stieltjes transform $s_{\mu}(z)$ of μ (see (1.21)) when f is a regular enough

compactly supported function. In order to introduce this tool, we consider a class \mathcal{C}^{k+1} compactly supported function f for a certain integer k, and denote by $\Phi_k(f): \mathbb{C} \to \mathbb{C}$ the function defined on \mathbb{C} by

$$\Phi_k(f)(x+iy) = \sum_{l=0}^k \frac{(iy)^l}{l!} f^{(l)}(x)\rho(y)$$

where $\rho : \mathbb{R} \to \mathbb{R}^+$ is smooth, compactly supported, with value 1 in a neighbourhood of 0. Function $\Phi_k(f)$ coincides with f on the real line and extends it to the complex plane. Let $\bar{\partial} = \partial_x + i\partial_y$. It is well-known that

$$\bar{\partial}\Phi_k(f)(x+iy) = \frac{(iy)^k}{k!} f^{(k+1)}(x)$$
(2.18)

(a proof of this result can be found in [9] or [18]) if y belongs to the neighbourhood of 0 in which ρ is equal to 1. The Helffer-Sjöstrand formula can be written as

$$\int f \,\mathrm{d}\mu = \frac{1}{\pi} \operatorname{Re} \int_{\mathbb{C}^+} \bar{\partial} \Phi_k(f)(z) s_\mu(z) \,\mathrm{d}x \,\mathrm{d}y.$$
(2.19)

In order to understand why the integral at the right hand side of (2.19) is well defined, we take, to fix the ideas, $\rho \in C^{\infty}$ such that $\rho(y) = 1$ for $|y| \leq 1$ and $\rho(y) = 0$ for |y| > 2, and denote by $[a_1, a_2]$ an interval containing the support of f. Then, it appears that the integral on \mathbb{C}^+ is in fact over the compact set $\mathcal{D} = \{x + iy : x \in [a_1, a_2], y \in [0, 2]\}$. Moreover, as $|\mathbf{s}_{\mu}(z)| \leq \frac{1}{y}$ if $z \in \mathcal{D}$ (see (1.22)), (2.18) for k = 1 leads to the conclusion that

$$|\bar{\partial}\Phi_k(f)(z)s_\mu(z)| \le C$$

for $z \in \{x + iy \in D, y \leq 1\}$. Therefore, the right hand side of (2.19) is well defined.

We finally mention that the Helffer-Sjöstrand formula remains still valid for any compactly supported distribution D (see e.g. [26], section 9). The Stieltjes transform of D, denoted by $s_D(z)$, is defined for each $z \in \mathbb{C}^+$ as the action of function $\lambda \to \frac{1}{\lambda - z}$ on D, i.e. $s_D(z) = \langle D, \frac{1}{\lambda - z} \rangle$, and verifies

$$|s_D(z)| \le C\left(1 + \frac{1}{(\operatorname{Im} z)^{n_0}}\right)$$

for each $z \in \mathbb{C}^+$ where n_0 is related to the order of the distribution. We refer the reader to [5] (Theorem 4.3) and the references therein for more details on Stieltjes transforms of distributions. Then, if f is a \mathcal{C}^{∞} function supported by $[a_1, a_2], < D, f >$ is given by

$$\langle D, f \rangle = \frac{1}{\pi} \operatorname{Re} \int_{\mathcal{D}} \bar{\partial} \Phi_k(f)(z) s_D(z) \,\mathrm{d}x \,\mathrm{d}y$$
 (2.20)

for $k \ge n_0$. We also recall that an alternative expression for $\langle D, f \rangle$ is given by the Stieltjes inversion formula, also valid for distributions, i.e.

$$\langle D, f \rangle = \frac{1}{\pi} \lim_{y \to 0} \int_{a_1}^{a_2} f(\lambda) \operatorname{Im} s_D(\lambda + iy) d\lambda.$$
 (2.21)

3. Stochastic representations of $\tilde{C}(\nu)$ and $\hat{C}(\nu)$

The first step is to show that $\tilde{\mathbf{C}}(\nu)$ and $\hat{\mathbf{C}}(\nu)$ can be approximated by the sample covariance matrix of a sequence of i.i.d. Gaussian random vectors, and to control the order of magnitude of the corresponding errors. This is the objective of the following result.

Theorem 3.1. Under Assumptions 1.1, 1.2, 1.3 and 1.4, for any $\nu \in [0, 1]$, it exists a $M \times (B+1)$ random matrix $\mathbf{X}_N(\nu)$ with $\mathcal{N}_{\mathbb{C}}(0, 1)$ i.i.d. entries, and two matrices $(\tilde{\mathbf{\Delta}}_N(\nu), \mathbf{\Delta}_N(\nu))$ such that:

$$\tilde{\mathbf{C}}_{N}(\nu) = \frac{\mathbf{X}_{N}(\nu)\mathbf{X}_{N}^{*}(\nu)}{B+1} + \tilde{\mathbf{\Delta}}_{N}(\nu), \quad \|\tilde{\mathbf{\Delta}}_{N}(\nu)\| \prec \frac{B}{N}$$
(3.1)

$$\hat{\mathbf{C}}_{N}(\nu) = \frac{\mathbf{X}_{N}(\nu)\mathbf{X}_{N}^{*}(\nu)}{B+1} + \mathbf{\Delta}_{N}(\nu), \quad \|\mathbf{\Delta}_{N}(\nu)\| \prec \frac{1}{\sqrt{B}} + \frac{B}{N}.$$
 (3.2)

Remark 3.1. Therefore, up to small additive perturbations, $\tilde{\mathbf{C}}_N(\nu)$ and $\hat{\mathbf{C}}_N(\nu)$ appear as empirical covariance matrices of *i.i.d.* $\mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_M)$ random vectors. We thus expect that $\tilde{\mathbf{C}}_N(\nu)$ and $\hat{\mathbf{C}}_N(\nu)$ will satisfy a number of useful properties of empirical covariance matrices of *i.i.d.* $\mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_M)$ random vectors.

In particular, Theorem 3.1 allows to precise the location of the eigenvalues of $\tilde{\mathbf{C}}_N(\nu)$ and $\hat{\mathbf{C}}_N(\nu)$. In order to formulate the corresponding result, we define some notations. We introduce the events $\Lambda_{N,\epsilon}^{\tilde{\mathbf{C}}}(\nu)$ and $\Lambda_{N,\epsilon}^{\hat{\mathbf{C}}}(\nu)$ defined by

$$\Lambda_{N,\epsilon}^{\tilde{\mathbf{C}}}(\nu) = \{\sigma(\tilde{\mathbf{C}}_N(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} + \epsilon\}$$
(3.3)

$$\Lambda_{N,\epsilon}^{\mathbf{C}}(\nu) = \{\sigma(\hat{\mathbf{C}}_N(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} + \epsilon\}.$$
(3.4)

Then, we establish in the following the Corollary:

Corollary 3.1. For each $\epsilon > 0$, the family of events $\Lambda_{N,\epsilon}^{\tilde{\mathbf{C}}}(\nu), N \ge 1, \nu \in [0,1]$ and $\Lambda_{N,\epsilon}^{\hat{\mathbf{C}}}(\nu), N \ge 1, \nu \in [0,1]$ hold with exponential high probability.

Remark 3.2. In the following, we will often omit to mention that the various matrices under consideration depend on N and ν . Matrices $\hat{\mathbf{C}}_N(\nu), \tilde{\mathbf{C}}_N(\nu), \mathbf{X}_N(\nu), \mathbf{\Delta}_N(\nu), \ldots$ will therefore be denoted by $\hat{\mathbf{C}}(\nu), \tilde{\mathbf{C}}(\nu), \mathbf{X}(\nu), \mathbf{\Delta}(\nu), \ldots$ or $\hat{\mathbf{C}}, \tilde{\mathbf{C}}, \mathbf{X}, \mathbf{\Delta}, \ldots$ We will also denote $\Lambda_{N,\epsilon}^{\tilde{\mathbf{C}}}(\nu)$ and $\Lambda_{N,\epsilon}^{\hat{\mathbf{C}}}(\nu)$ by $\Lambda_{\epsilon}^{\tilde{\mathbf{C}}}(\nu)$ or $\Lambda_{\epsilon}^{\tilde{\mathbf{C}}}$ and $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}(\nu)$ or $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$.

The proof of Theorem 3.1 will proceed in three steps: first we provide the result for matrix $\tilde{\mathbf{C}}(\nu)$, then control the deviations between diag $(\mathbf{S}(\nu))^{-\frac{1}{2}}$ and diag $(\hat{\mathbf{S}}(\nu))^{-\frac{1}{2}}$, and eventually extend the stochastic representation of $\tilde{\mathbf{C}}(\nu)$ to $\hat{\mathbf{C}}(\nu)$.

3.1. Step 1: Stochastic representation of \tilde{C}

In order to establish (3.1), we prove the following Proposition.

Proposition 3.1. Under Assumptions 1.1, 1.2, 1.3 and 1.4, for any $\nu \in [0, 1]$, there exist a $M \times (B + 1)$ random matrix $\mathbf{X}_N(\nu)$ with $\mathcal{N}_{\mathbb{C}}(0, 1)$ i.i.d. entries, and another matrix $\Gamma_N(\nu)$ such that:

$$\tilde{\mathbf{C}}_{N}(\nu) = \frac{(\mathbf{X}_{N}(\nu) + \mathbf{\Gamma}_{N}(\nu))(\mathbf{X}_{N}(\nu) + \mathbf{\Gamma}_{N}(\nu))^{*}}{B+1}$$
(3.5)

where the family of random variables $\frac{\|\Gamma_N(\nu)\|^2}{B+1}$, $\nu \in [0,1]$ verifies

$$\frac{\|\mathbf{\Gamma}_N(\nu)\|^2}{B+1} \prec \frac{B^2}{N^2}.$$
(3.6)

Proof. Denote by Σ the $M \times (B+1)$ random matrix defined by

$$\boldsymbol{\Sigma} = \left(\boldsymbol{\xi}_{\mathbf{y}}(\nu - \frac{B}{2N}), \dots, \boldsymbol{\xi}_{\mathbf{y}}(\nu + \frac{B}{2N})\right)$$
(3.7)

where we recall that the normalized Fourier transform $\boldsymbol{\xi}_{\mathbf{y}}$ is defined in (1.2), so that $\hat{\mathbf{S}}$ defined in (1.1) is equal to $\boldsymbol{\Sigma}\boldsymbol{\Sigma}^*/(B+1)$. Denote by $\boldsymbol{\omega}_m$ the *m*-th row of $\boldsymbol{\Sigma}$. In other words, $\boldsymbol{\omega}_m$ coincides with the (B+1)-dimensional Gaussian complex row vector defined by:

$$\boldsymbol{\omega}_m = \left(\xi_{y_m}(\nu - \frac{B}{2N}), \dots, \xi_{y_m}(\nu + \frac{B}{2N})\right).$$

The covariance matrix $\mathbb{E}[\boldsymbol{\omega}_m^*\boldsymbol{\omega}_m]$ of $\boldsymbol{\omega}$ is given by:

$$\mathbb{E}[\boldsymbol{\omega}_m^*\boldsymbol{\omega}_m] = \mathbb{E}\left[\left\{\xi_{y_m}(\nu + \frac{b_1}{N})^*\xi_{y_m}(\nu + \frac{b_2}{N})\right\}_{b_1, b_2 = -B/2}^{B/2}\right]$$

By Lemma A.1 in Appendix, we have for b and $b_1 \neq b_2$:

~ 7

$$\mathbb{E}\left[\left|\xi_{y_m}\left(\nu+\frac{b}{N}\right)\right|^2\right] = s_m\left(\nu+\frac{b}{N}\right) + \mathcal{O}\left(\frac{1}{N}\right)$$
$$\mathbb{E}\left[\xi_{y_m}(\nu+\frac{b_1}{N})^*\xi_{y_m}(\nu+\frac{b_2}{N})\right] = \mathcal{O}\left(\frac{1}{N}\right)$$

where the error is uniform over $m \geq 1$ and $\nu \in [0, 1]$. Therefore one can write that there exists some Hermitian matrix $\Upsilon_m(\nu)$ and some nice constant C such that:

$$\mathbb{E}[\boldsymbol{\omega}_m^*\boldsymbol{\omega}_m] = \operatorname{diag}\left(s_m\left(\nu + \frac{b}{N}\right) : b = -B/2, \dots, B/2\right) + \boldsymbol{\Upsilon}_m$$

where $\boldsymbol{\Upsilon}_m$ satisfies

$$\sup_{m\geq 1,b_1,b_2} \left| (\boldsymbol{\Upsilon}_m)_{b_1,b_2} \right| \leq \frac{C}{N}.$$

Moreover, the regularity of the applications $\nu \mapsto s_m(\nu)$ specified in Assumption 1.4 implies that there exists quantities ϵ_m such that:

$$s_m(\nu + \frac{b}{N}) = s_m(\nu) + s'_m(\nu) \frac{b}{N} + \frac{1}{2} s''_m(\nu) \left(\frac{b}{N}\right)^2 + \epsilon_m(\nu + \frac{b}{N})$$

where:

$$\sup_{m \ge 1} \sup_{-B/2 \le b \le B/2} |\epsilon_m(\nu + \frac{b}{N})| \le C \left(\frac{B}{N}\right)^3$$

for some nice constant C. Therefore, it holds that

$$\operatorname{diag}\left(s_m(\nu + \frac{b}{N}) : b = -B/2, \dots, B/2\right)$$
$$= s_m(\nu)\mathbf{I}_{B+1} + s'_m(\nu)\operatorname{diag}\left(\frac{b}{N} : b = -B/2, \dots, B/2\right) + \frac{1}{2}s''_m(\nu)\operatorname{diag}\left((\frac{b}{N})^2 : b = -B/2, \dots, B/2\right) + \operatorname{diag}\left(\epsilon_m(\nu + \frac{b}{N}) : b = -B/2, \dots, B/2\right)$$

If we define matrix Φ_m as:

$$\boldsymbol{\Phi}_m = \frac{1}{s_m} \left[\boldsymbol{\Upsilon}_m + \operatorname{diag} \left(\mathbf{s}_m(\nu + \frac{b}{N}) - s_m(\nu) : b = -B/2, \dots, B/2 \right) \right]$$

then $\mathbb{E}[\boldsymbol{\omega}_m^*\boldsymbol{\omega}_m] = s_m \left(\mathbf{I}_{B+1} + \boldsymbol{\Phi}_m\right)$ with

$$\sup_{m \ge 1, b_1 \ne b_2} |(\mathbf{\Phi}_m)_{b_1, b_2}| \le \frac{C}{N}, \quad \sup_{m \ge 1, b} |(\mathbf{\Phi}_m)_{b, b}| \le \frac{CB}{N}$$
(3.8)

as well as

$$\frac{1}{B+1} \operatorname{tr} \mathbf{\Phi}_m = \frac{1}{2} \frac{s_m''(\nu)}{s_m(\nu)} v_N + \mathcal{O}\left(\left(\frac{B}{N}\right)^3 + \frac{1}{N}\right)$$
(3.9)

where we recall that v_N is defined by (1.5). The spectral norm of Φ_m can be roughly bounded by the following inequality:

$$\sup_{m \ge 1} \|\Phi_m\| \le \sup_{m \ge 1 - B/2 \le b_1 \le B/2} \sum_{b_2 = -B/2}^{B/2} |(\Phi_m)_{b_1, b_2}| \le C \frac{B}{N}.$$

Moreover, it is easily checked that the Frobenius norm of $\frac{\Phi_m}{B+1}$ verifies

$$\left\|\frac{\mathbf{\Phi}_m}{B+1}\right\|_F \le C \, \frac{\sqrt{B}}{N} = \mathcal{O}(u_N). \tag{3.10}$$

Using the Gaussianity of vector $\boldsymbol{\omega}_m$ and the expression (3.8), we obtain that $\boldsymbol{\omega}_m$ can be represented as

$$\boldsymbol{\omega}_m = \sqrt{s_m} \, \mathbf{x}_m \, \left(\mathbf{I} + \boldsymbol{\Phi}_m \right)^{1/2}, \quad \mathbf{x}_m \sim \mathcal{N}_{\mathbb{C}}(0, I_{B+1}) \tag{3.11}$$

where \mathbf{x}_{m_1} and \mathbf{x}_{m_2} are independent for $m_1 \neq m_2$. This comes from the mutual independence of the time series $((y_{m,n})_{n \in \mathbb{Z}})_{m=1,\dots,M}$. It is clear that $(\mathbf{I} + \boldsymbol{\Phi}_m)^{1/2}$ can be written as

$$\left(\mathbf{I} + \boldsymbol{\Phi}_m\right)^{1/2} = \mathbf{I} + \boldsymbol{\Psi}_m \tag{3.12}$$

where matrix Ψ_m verifies

$$\sup_{m} \|\Psi_m\| \le C \frac{B}{N} \tag{3.13}$$

Therefore, it holds that:

$$\boldsymbol{\omega}_{m} = \sqrt{s_{m}} \mathbf{x}_{m} \left(\mathbf{I} + \boldsymbol{\Psi}_{m} \right) = \sqrt{s_{m}} \left(\mathbf{x}_{m} + \mathbf{x}_{m} \boldsymbol{\Psi}_{m} \right)$$

We denote by **X** and Γ the $M \times (B+1)$ matrices with rows $(\mathbf{x}_m)_{m=1,...,M}$, and $(\mathbf{x}_m \Psi_m)_{m=1,...,M}$ respectively. Then, it holds that

$$\boldsymbol{\Sigma} = \operatorname{diag}\left(\sqrt{s_m}, m = 1, \dots, M\right) \left(\mathbf{X} + \boldsymbol{\Gamma}\right)$$
(3.14)

where we recall that Σ is defined by (3.7). We recall the definition of matrix $\tilde{\mathbf{C}}$ given by

$$\tilde{\mathbf{C}} = \operatorname{diag}(\sqrt{s_m}, m = 1, \dots, M)^{-1/2} \, \hat{\mathbf{S}} \, \operatorname{diag}(\sqrt{s_m}, m = 1, \dots, M)^{-1/2} \, (3.15) \\ = \operatorname{diag}(\sqrt{s_m}, m = 1, \dots, M)^{-1/2} \frac{\mathbf{\Sigma} \mathbf{\Sigma}^*}{B+1} \operatorname{diag}(\sqrt{s_m}, m = 1, \dots, M)^{-1/2}.$$

The representation (3.14) implies that $\tilde{\mathbf{C}}$ can also be written as

$$\tilde{\mathbf{C}} = \frac{(\mathbf{X} + \mathbf{\Gamma})(\mathbf{X} + \mathbf{\Gamma})^*}{B+1}.$$

Equivalently, for each m_1, m_2 , the entry $(\tilde{\mathbf{C}})_{m_1, m_2}$ is given by

$$(\tilde{\mathbf{C}})_{m_1,m_2} = \frac{1}{B+1} \mathbf{x}_{m_1} (\mathbf{I} + \mathbf{\Phi}_{m_1})^{1/2} (\mathbf{I} + \mathbf{\Phi}_{m_2})^{1/2} \mathbf{x}_{m_2}^*.$$
 (3.16)

This completes the proof of (3.5). It remains to show (3.6). We denote by **Z** the $M \times M$ matrix $\mathbf{Z} = \frac{1}{B+1} \Gamma \Gamma^*$. As $\|\mathbf{Z}\|$ verifies

$$\|\mathbf{Z}\| \le \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| + \|\mathbb{E}\mathbf{Z}\|$$

it is enough to prove the two following facts:

$$\|\mathbb{E}\mathbf{Z}\| \le C \,\frac{B^2}{N^2} \tag{3.17}$$

$$\|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \prec \frac{B^2}{N^2}.$$
 (3.18)

We start with (3.17). The definition of Γ leads to

$$\mathbb{E}[\mathbf{Z}_{i,j}] = \frac{1}{B+1} \mathbb{E}[\mathbf{\Gamma}\mathbf{\Gamma}^*]_{i,j} = \frac{1}{B+1} \mathbb{E}[\mathbf{x}_i \boldsymbol{\Psi}_i \boldsymbol{\Psi}_j^* \mathbf{x}_j^*] = \delta_{ij} \frac{1}{B+1} \operatorname{tr} \, \boldsymbol{\Psi}_i \boldsymbol{\Psi}_j^*$$

so that it is clear that $\mathbb{E}[\mathbf{Z}]$ is the diagonal matrix with diagonal entries $(\frac{1}{B+1} \operatorname{tr} \Psi_m \Psi_m^*)_{m=1,\ldots,M}$. By the estimation in equation (3.13), we easily have (3.17).

It remains to prove (3.18). We use the observation that $\|\mathbf{Z} - \mathbb{E}[\mathbf{Z}]\| = \max_{\|\mathbf{h}\|=1} |\mathbf{h}^*(\mathbf{Z} - \mathbb{E}[\mathbf{Z}])\mathbf{h}|$, and use a classical ϵ -net argument that allows to deduce the behaviour of $\|\mathbf{Z} - \mathbb{E}[\mathbf{Z}]\|$ from the behaviour of any recentered quadratic form $\mathbf{g}^*\mathbf{Z}\mathbf{g} - \mathbb{E}\mathbf{g}^*\mathbf{Z}\mathbf{g}$ where $\mathbf{g} \in \mathbb{C}^M$ is a deterministic unit norm vector. We thus first concentrate $\mathbf{g}^*\mathbf{Z}\mathbf{g} - \mathbb{E}\mathbf{g}^*\mathbf{Z}\mathbf{g}$ using the Hanson-Wright inequality (2.17). For this, we need to express $\mathbf{g}^*\mathbf{Z}\mathbf{g}$ as a quadratic form of a certain complex Gaussian random vector with i.i.d. entries. We denote by \mathbf{z} the *M*-dimensional random vector $\mathbf{z} = \frac{\mathbf{\Gamma}^*(\nu)\mathbf{g}}{\sqrt{B+1}}$. Its covariance matrix $\mathbf{G} = \mathbf{G}(\nu)$ is equal to

$$\mathbf{G}(\nu) = \mathbb{E}[\mathbf{z}\mathbf{z}^*] = \frac{1}{B+1} \sum_{m=1}^M |\mathbf{g}_m|^2 (\boldsymbol{\Psi}_m(\nu))^* \boldsymbol{\Psi}_m(\nu).$$

Therefore, \mathbf{z} can be written $\mathbf{z} = \mathbf{G}^{1/2}\mathbf{w}$ for some $\mathbf{w} \sim \mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_M)$ random vector. As a consequence, the quadratic form $\mathbf{g}^*\mathbf{Z}\mathbf{g} - \mathbb{E}\mathbf{g}^*\mathbf{Z}\mathbf{g}$ can be written as

$$\mathbf{g}^*\mathbf{Z}\mathbf{g} - \mathbb{E}\mathbf{g}^*\mathbf{Z}\mathbf{g} = \mathbf{w}^*\mathbf{G}\mathbf{w} - \mathbb{E}\mathbf{w}^*\mathbf{G}\mathbf{w}.$$

The Hanson-Wright inequality (2.17) can now be applied:

$$|\mathbf{w}^* \mathbf{G} \mathbf{w} - \mathbb{E} \mathbf{w}^* \mathbf{G} \mathbf{w}| \prec ||\mathbf{G}||_F.$$
(3.19)

Since $\sum_{m=1}^{M} |\mathbf{g}_m|^2 = 1$, it is clear that $\|\mathbf{G}\| \leq \frac{1}{B+1} \sup_{m=1,\dots,M} \|\Psi_m(\nu)\|^2$. Therefore, (3.13) and the rough evaluation $\|\mathbf{G}\|_F^2 \leq (B+1) \|\mathbf{G}\|^2$ leads to

$$\|\mathbf{G}\| \le C \frac{1}{B+1} \left(\frac{B}{N}\right)^2, \quad \|\mathbf{G}\|_F^2 \le C \frac{1}{B+1} \left(\frac{B}{N}\right)^4 \tag{3.20}$$

The substitution of (3.20) in equation (3.19) gives the following control of $\mathbf{g}^*\mathbf{Z}\mathbf{g} - \mathbb{E}\mathbf{g}^*\mathbf{Z}\mathbf{g}$:

$$|\mathbf{g}^* \mathbf{Z} \mathbf{g} - \mathbb{E} \mathbf{g}^* \mathbf{Z} \mathbf{g}| \prec \frac{1}{\sqrt{B}} \left(\frac{B}{N}\right)^2$$
 (3.21)

Consider $\epsilon > 0$, and an ϵ -net N_{ϵ} of \mathbb{C}^{M} , that is a set of \mathbb{C}^{M} unit norm vectors $\{\mathbf{h}_{k} : k = 1, \ldots, \mathcal{K}\}$ such that for each unit norm vector $\mathbf{u} \in \mathbb{C}^{M}$, it exists a vector $\mathbf{h} \in N_{\epsilon}$ for which $\|\mathbf{u} - \mathbf{h}\| \leq \epsilon$. It is well known that the cardinal of N_{ϵ} is bounded by $C_{0} \left(\frac{1}{\epsilon}\right)^{2M}$ where C_{0} is a universal constant. Then, denote \mathbf{g}_{s} a (random) unit norm vector such that $|\mathbf{g}_{s}^{*}\mathbf{Z}\mathbf{g}_{s} - \mathbb{E}\mathbf{g}_{s}^{*}\mathbf{Z}\mathbf{g}_{s}| = \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\|$, and define $\mathbf{h}_{s} \in \mathbf{N}_{\epsilon}$ as the closest vector from \mathbf{g}_{s} . Therefore, we have

$$\begin{split} \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| &= |\mathbf{g}_s^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{g}_s| \\ &= |(\mathbf{g}_s^* - \mathbf{h}_s^* + \mathbf{h}_s^*)(\mathbf{Z} - \mathbb{E}\mathbf{Z})(\mathbf{g}_s - \mathbf{h}_s + \mathbf{h}_s)| \\ &\leq |(\mathbf{g}_s^* - \mathbf{h}_s^*)(\mathbf{Z} - \mathbb{E}\mathbf{Z})(\mathbf{g}_s - \mathbf{h}_s)| + |(\mathbf{g}_s^* - \mathbf{h}_s^*)(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}_s| \\ &+ |\mathbf{h}_s^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})(\mathbf{g}_s - \mathbf{h}_s)| + |\mathbf{h}_s^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}_s|. \end{split}$$

It is clear that:

$$\begin{aligned} |(\mathbf{g}_s^* - \mathbf{h}_s^*)(\mathbf{Z} - \mathbb{E}\mathbf{Z})(\mathbf{g}_s - \mathbf{h}_s)| &\leq \epsilon^2 \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \\ |(\mathbf{g}_s^* - \mathbf{h}_s^*)(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}_s| &\leq \epsilon \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \end{aligned}$$

and

$$\|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \le |\mathbf{h}_s^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}_s| + \epsilon^2 \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| + 2\epsilon \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\|$$

which leads to

$$(1 - 2\epsilon - \epsilon^2) \|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| \le |\mathbf{h}_s^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}_s|$$

This implies that for each t > 0,

$$\{ \| \mathbf{Z} - \mathbb{E}\mathbf{Z} \| > t \} \subset \cup_{h \in N_{\epsilon}} \{ |\mathbf{h}^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}| > C_1 t \}$$

where $C_1 = (1 - 2\epsilon - \epsilon^2)$. Using the union bound, we obtain that

$$\mathbb{P}\left[\|\mathbf{Z} - \mathbb{E}\mathbf{Z}\| > t\right] \le \sum_{h \in N_{\epsilon}} \mathbb{P}\left[|\mathbf{h}^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}| > C_1 t\right].$$
(3.22)

Here, we would like to use equation (3.21) to conclude. By the definition of \prec , (3.21) is valid uniformly on any set of vector with cardinality polynomial in N. Here, the cardinal of the set N_{ϵ} is a $\mathcal{O}(\epsilon^{-2M})$ term and therefore exponential in M. As a consequence, we have to accept to lose some speed when going from the stochastic domination of $|\mathbf{g}^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{g}|$ for a fixed \mathbf{g} to the same stochastic domination but uniformly over N_{ϵ} .

More precisely, write again (3.21) but here without the notation \prec in order to understand precisely how a change in speed affects the probability. Take t_N a sequence of positive number such that $t_N \geq B^2/N^2$. Using the estimates (3.20) of $\|\mathbf{G}\|$ and $\|\mathbf{G}\|_F^2$, and the fact that $\min(a_1, a_2) > \min(b_1, b_2)$ when $a_1 > b_1$ and $a_2 > b_2$, we obtain that there exist some nice constant C > 0 such that:

$$\min\left(\frac{t_N}{\|\mathbf{G}\|}, \frac{t_N^2}{\|\mathbf{G}\|_F^2}\right) \ge CB\min\left(t_N\left(\frac{N}{B}\right)^2, \left(t_N\left(\frac{N}{B}\right)^2\right)^2\right)$$
$$= CBt_N\left(\frac{N}{B}\right)^2.$$

The Hanson-Wright inequality (2.15) provides:

$$\mathbb{P}\left[\left|\mathbf{g}^{*}\left[\mathbf{Z}-\mathbb{E}\mathbf{Z}\right)\right]\mathbf{g}\right| > C_{1}t_{N}\right] \leq 2 \exp\left\{-CB\frac{t_{N}}{(B/N)^{2}}\right\}$$

for some nice constant C that depends on C_1 . Eventually, the union bound on N_{ϵ} gives:

$$\mathbb{P}\left[\|\mathbf{Z} - \mathbb{E}(\mathbf{Z})\| > t_N\right] \le \sum_{h \in N_{\epsilon}} \mathbb{P}\left[|\mathbf{h}^*(\mathbf{Z} - \mathbb{E}\mathbf{Z})\mathbf{h}| > C_1 t_N\right]$$
$$\le 2 C_0 \exp\left\{-CB \frac{t_N}{(B/N)^2} + 2M \log\frac{1}{\epsilon}\right\}. \quad (3.23)$$

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 27

If we take $t_N = N^{\epsilon'}(B^2/N^2)$, then, it exists $\gamma > 0$ such that

$$\exp\left\{-CB\frac{t_N}{(B/N)^2} + 2CM\log\frac{1}{\epsilon}\right\} \le \exp{-N^{\gamma}}$$

holds for each N large enough. (3.22) thus implies (3.18). This completes the proof of (3.5). \Box

Corollary 3.2 is a rewriting of Proposition 3.1 in a more concise way. Define:

$$\tilde{\Delta} = \frac{\mathbf{X}\Gamma^* + \Gamma\mathbf{X}^* + \Gamma\Gamma^*}{B+1}.$$
(3.24)

Corollary 3.2. For any $\nu \in [0,1]$, $\tilde{\mathbf{C}}(\nu)$ can be written as

$$\tilde{\mathbf{C}}(\nu) = \frac{\mathbf{X}(\nu)\mathbf{X}^*(\nu)}{B+1} + \tilde{\boldsymbol{\Delta}}(\nu)$$
(3.25)

where the family of random variable $\|\tilde{\Delta}(\nu)\|, \nu \in [0, 1]$ verifies

$$\|\tilde{\mathbf{\Delta}}\| \prec \frac{B}{N}.\tag{3.26}$$

Proof. Let $\nu \in [0, 1]$. By equation (3.6) from Theorem 3.1 and equation (2.5) from Paragraph 2.2, we have the two following estimates:

$$\frac{\|\mathbf{\Gamma}\|}{\sqrt{B+1}} \prec \frac{B}{N}, \quad \frac{\|\mathbf{X}\|}{\sqrt{B+1}} \prec 1$$

The result is immediate using decomposition $\tilde{\Delta}$ from (3.24):

We now take benefit of Corollary 3.2 to establish the first part of Corollary 3.1 and to analyse the location of the eigenvalues of matrices $\hat{\mathbf{S}}$. We denote by \mathbf{D} and $\hat{\mathbf{D}}$ the matrices $\mathbf{D} = \mathbf{D}(\nu) := \text{diag}(\mathbf{S}(\nu))^{\frac{1}{2}}$ and $\hat{\mathbf{D}} = \hat{\mathbf{D}}(\nu) := \text{diag}(\hat{\mathbf{S}}(\nu))^{\frac{1}{2}}$. Denote by \bar{s} and \underline{s} the quantities such that:

$$\underline{s} := \inf_{m \ge 1} \inf_{\nu \in [0,1]} s_m(\nu), \quad \overline{s} := \sup_{m \ge 1} \sup_{\nu \in [0,1]} s_m(\nu)$$

which are by Assumption 1.4 in $(0, +\infty)$. We consider the event:

$$\Lambda_{\epsilon}^{\hat{\mathbf{S}}}(\nu) = \left\{ \sigma(\hat{\mathbf{S}}(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}] + \epsilon \right\}$$
(3.27)

where the notation $\operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}]$ stands for $[(1 - \sqrt{c})^2 \underline{s}, (1 + \sqrt{c})^2 \overline{s}]$. Note that in our settings, $c \in (0, 1)$ so $\operatorname{Supp} \mu_{MP}^{(c)}$ is bounded and away from zero. In conjunction with Assumption 1.4, the same holds for $\operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}]$. We also note that $\Lambda_{\epsilon}^{\hat{\mathbf{s}}}(\nu)$ of course depends on N.

Corollary 3.3. For any $\epsilon > 0$, the families of events $\Lambda_{\epsilon}^{\tilde{\mathbf{C}}}(\nu)$, $\nu \in [0,1]$ and $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}(\nu)$, $\nu \in [0,1]$ hold with exponentially high probability.

Proof. Equation (3.25) implies that

$$\frac{\mathbf{X}\mathbf{X}^*}{B+1} - \|\tilde{\mathbf{\Delta}}\|\mathbf{I}_M \le \tilde{\mathbf{C}} \le \frac{\mathbf{X}\mathbf{X}^*}{B+1} + \|\tilde{\mathbf{\Delta}}\|\mathbf{I}_M.$$

Therefore, the event $\{\lambda_1(\tilde{\mathbf{C}}) > (1+\sqrt{c})^2 + \epsilon\}$ is included in $\{\lambda_1(\frac{\mathbf{X}\mathbf{X}^*}{B+1}) + \|\tilde{\mathbf{\Delta}}\| > (1+\sqrt{c})^2 + \epsilon\}$, which is itself included into

$$\left\{\lambda_1(\frac{\mathbf{X}\mathbf{X}^*}{B+1}) > (1+\sqrt{c})^2 + \epsilon/2\right\} \cup \left\{\|\tilde{\mathbf{\Delta}}\| > \epsilon/2\right\}.$$

Therefore,

$$\mathbb{P}\left[\lambda_1(\tilde{\mathbf{C}}) > (1+\sqrt{c})^2 + \epsilon\right] \le \mathbb{P}\left[\lambda_1(\frac{\mathbf{X}\mathbf{X}^*}{B+1}) > (1+\sqrt{c})^2 + \epsilon/2\right] \\ + \mathbb{P}\left[\|\tilde{\mathbf{\Delta}}\| > \epsilon/2\right].$$

Equations (2.3) and (3.26) imply that $\mathbb{P}\left[\lambda_1(\tilde{\mathbf{C}}) > (1+\sqrt{c})^2 + \epsilon\right]$ converges towards 0 exponentially. A similar evaluation of $\mathbb{P}\left[\lambda_M(\tilde{\mathbf{C}}) < (1-\sqrt{c})^2 - \epsilon\right]$ leads to the same conclusion. This, in turn, establishes that $\Lambda_{\epsilon}^{\tilde{\mathbf{C}}}(\nu), \nu \in [0,1]$ holds with exponential high probability.

In order to establish that the same property holds for $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}(\nu), \nu \in [0, 1]$, we just need to write (1.12) as $\hat{\mathbf{S}} = \mathbf{D}^{1/2} \tilde{\mathbf{C}} \mathbf{D}^{1/2}$. Therefore, for each $k = 1, \ldots, M$, the eigenvalues of $\hat{\mathbf{S}}$ verify

$$\underline{s} \lambda_M(\tilde{\mathbf{C}}) \le \lambda_k(\hat{\mathbf{S}}) \le \overline{s} \lambda_1(\tilde{\mathbf{C}}).$$

This, of course, implies that $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}(\nu), \nu \in [0, 1]$ holds with exponential high probability (indeed, one can change ϵ to $\tilde{\epsilon}$ such that $(\operatorname{Supp} \mu_{MP}^{(c)} + \tilde{\epsilon}) \times [\underline{s}, \overline{s}] \subset \operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}] + \epsilon$.

Remark 3.3. Corollary 3.3 implies the following weaker property, which will be useful:

$$\|\mathbf{S}(\nu)\| \prec 1. \tag{3.28}$$

Before ending the section and proving Theorem 3.1, we need some stochastic control on the diagonal elements of $\hat{\mathbf{S}}$ in order to evaluate $\boldsymbol{\Theta}$ defined by

$$\boldsymbol{\Theta} := \hat{\mathbf{C}} - \hat{\mathbf{C}}.\tag{3.29}$$

Using the definition of $\hat{\mathbf{C}}$ from (1.3) and $\tilde{\mathbf{C}}$ from (1.12), $\boldsymbol{\Theta}$ can be written as

$$\boldsymbol{\Theta} = (\hat{\mathbf{D}}^{-1/2} - \mathbf{D}^{-1/2})\hat{\mathbf{S}}\hat{\mathbf{D}}^{-1/2} + \mathbf{D}^{-1/2}\hat{\mathbf{S}}(\hat{\mathbf{D}}^{-1/2} - \mathbf{D}^{-1/2}).$$
(3.30)

Since we proved that $\|\hat{\mathbf{S}}\| \prec 1$, it remains to show that $\|\hat{\mathbf{D}}^{-1/2}\|$ and $\|\hat{\mathbf{D}}^{-1/2} - \mathbf{D}^{-1/2}\|$ can also be stochastically dominated by some relevant quantity in order to control $\|\mathbf{\Theta}\|$. Define

$$\hat{s}_m(\nu) := \hat{\mathbf{S}}_{m,m}(\nu) \tag{3.31}$$

the diagonal elements of $\hat{\mathbf{S}}(\nu)$ spectral density estimator (note that they coincide with the traditional smoothed periodogram estimator of the spectral density s_m). The aim of the following Paragraph 3.2 is to establish stochastic domination results for \hat{s}_m , $\|\hat{\mathbf{D}}^{-1/2}\|$ and $\|\hat{\mathbf{D}}^{-1/2} - \mathbf{D}^{-1/2}\|$.

3.2. Step 2: Estimates for $\hat{s}_m(\nu)$

We write $s_m(\nu) := s_m$, $\mathbf{D}(\nu) := \mathbf{D}$, in order to simplify the notations. Define as in (3.27) the following quantity

$$\Lambda_{\epsilon}^{\mathbf{D}}(\nu) = \{ \sigma(\hat{\mathbf{D}}(\nu)) \subset [\underline{s}, \overline{s}] + \epsilon \}.$$
(3.32)

Lemma 3.1. Let $\epsilon > 0$. The family of events $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu), \nu \in [0,1]$ holds with exponentially high probability.

Proof. See Appendix A.2.

Roughly speaking, this ensures that with exponentially high probability, \hat{s}_m stays bounded and away from zero. This result implies the following (weaker) statement, but will still be enough for some proofs and reduces the complexity of the arguments.

Lemma 3.2. The family of random variables $(|\hat{s}_m(\nu)| + \frac{1}{|\hat{s}_m(\nu)|})_{m=1,...,M}, \nu \in [0,1]$, verifies

$$\left(|\hat{s}_m| + \frac{1}{|\hat{s}_m|}\right) \prec 1.$$

Proof. Immediate from Lemma 3.1.

Lemma 3.3. The set of random variable $(|\hat{s}_m(\nu)^{-1/2} - s_m(\nu)^{-1/2}|)_{m=1,...,M}$ and $(|\sqrt{\frac{s_m(\nu)}{\hat{s}_m(\nu)}} - 1|)_{m=1,...,M}, \nu \in [0, 1]$, verifies

$$\left|\hat{s}_{m}^{-1/2} - s_{m}^{-1/2}\right| \prec \frac{1}{\sqrt{B}} + \frac{B^{2}}{N^{2}}, \quad \left|\sqrt{\frac{s_{m}}{\hat{s}_{m}}} - 1\right| \prec \frac{1}{\sqrt{B}} + \frac{B^{2}}{N^{2}}.$$
 (3.33)

Proof. See Appendix A.3

3.3. Step 3: Stochastic representation of \hat{C}

We are now in position to prove the result concerning $\hat{\mathbf{C}}$ of Theorem 3.1 and of Corollary 3.1.

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 30

Proof. We have first to control the operator norm of:

$$\boldsymbol{\Delta} = \hat{\mathbf{C}} - \frac{\mathbf{X}\mathbf{X}^*}{B+1} = \hat{\mathbf{C}} - \tilde{\mathbf{C}} + \tilde{\mathbf{C}} - \frac{\mathbf{X}\mathbf{X}^*}{B+1} = \boldsymbol{\Theta} + \tilde{\boldsymbol{\Delta}}.$$
 (3.34)

The operator norm of $\|\tilde{\boldsymbol{\Delta}}\|$ has already been proved in Corollary 3.2 to satisfy $\|\tilde{\boldsymbol{\Delta}}\| \prec (\frac{B}{N})$. Moreover, recall that $\boldsymbol{\Theta}$ can be written as a function of $\hat{\mathbf{D}}^{-1/2} - \mathbf{D}^{-1/2}$ in (3.30), so that one can use Lemma 3.2 and Lemma 3.3 to dominate each term and get:

$$\|\boldsymbol{\Theta}\| \prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}.$$
(3.35)

Summing the estimate of Θ and the one of $\hat{\Delta}$, one get:

$$\|\mathbf{\Delta}\| \prec \frac{1}{\sqrt{B}} + \frac{B}{N}$$

which is the desired result.

As a consequence, we state here Corollary 3.4 about the localization of the eigenvalues of $\hat{\mathbf{C}}(\nu)$.

Corollary 3.4. For each $\epsilon > 0$, we define $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}(\nu)$ as the event

$$\Lambda_{\epsilon}^{\hat{\mathbf{C}}}(\nu) = \left\{ \sigma(\hat{\mathbf{C}}(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} + \epsilon \right\}.$$
(3.36)

Then, the family of events $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}(\nu), \nu \in [0, 1]$ holds with exponentially high probability.

Proof. We simply write:

$$\frac{\mathbf{X}\mathbf{X}^*}{B+1} - \|\mathbf{\Delta}\|\mathbf{I}_M \leq \hat{\mathbf{C}} \leq \frac{\mathbf{X}\mathbf{X}^*}{B+1} + \|\mathbf{\Delta}\|\mathbf{I}_M$$

and use the same arguments as in the proof of Corollary 3.3.

4. Stochastic domination of the family $\psi_N(f,\nu), N \ge 1, \nu \in [0,1]$

We have first to define the distribution D_N introduced in the definition (1.11) of $\psi_N(f,\nu)$. For this, we consider the function $p_N(z)$ defined by

$$p_N(z) = -\frac{c_N \left(z \, t_N(z) \, \tilde{t}_N(z)\right)^3}{1 - c(z \, t_N(z) \, \tilde{t}_N(z))^2} \tag{4.1}$$

where we recall that t_N and \tilde{t}_N are defined by (2.10) and (2.11). Then (see Lemma 9.2 in [26]), p_N is the Stieltjes transform of a distribution whose support is contained into the support Supp $\mu_{MP}^{(c_N)} = [(1 - \sqrt{c_N})^2, (1 + \sqrt{c_N})^2]$ of the Marcenko-Pastur distribution $\mu_{MP}^{(c_N)}$. This distribution is D_N introduced in (1.11). In the following, we consider LSS for function f satisfying the following assumptions.

Assumption 4.1. *f* is defined on \mathbb{R}_+ and there exist some $\epsilon > 0$ such that its restriction on $\operatorname{Supp}_{MP}^{(c)} + \epsilon$ is \mathcal{C}^{∞} .

We now state the main result of this section.

Theorem 4.1. Let f be an application satisfying the conditions of Assumption 4.1. Then, under Assumptions 1.1, 1.2, 1.3 and 1.4, the family $|\psi_N(f,\nu)|, N \ge 1, \nu \in [0,1]$ verifies

$$|\psi_N(f,\nu)| \prec u_N. \tag{4.2}$$

Before starting the proof of Theorem 4.1, we first mention that it is sufficient to establish (4.2) when f is compactly supported by a neighbourhood of $\operatorname{Supp} \mu_{MP}^{(c)}$. To justify this claim, we consider $\kappa > 0$ and define $\chi : \mathbb{R} \to \mathbb{R}$ as a \mathbb{C}^{∞} function such that:

$$\chi(\lambda) = \begin{cases} 1 & \text{if } \lambda \in \operatorname{Supp} \mu_{MP}^{(c)} + \kappa \\ 0 & \text{if } \lambda \notin \operatorname{Supp} \mu_{MP}^{(c)} + 2\kappa. \end{cases}$$
(4.3)

We consider function \bar{f} given by $\bar{f} = f \times \chi$. Then, as $c_N \to c$, for N large enough, Supp $\mu_{MP}^{(c_N)}$ is contained in Supp $\mu_{MP}^{(c)} + \kappa$. Therefore, $f = \bar{f}$ on Supp $\mu_{MP}^{(c_N)}$ for N large enough, and it holds that $\langle D_N, f \rangle = \langle D_N, \bar{f} \rangle$ and $\int f d\mu_{MP}^{(c_N)} = \int \bar{f} d\mu_{MP}^{(c_N)}$. For each $\epsilon > 0$, we express $\mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N)$ as

$$\begin{aligned} \mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N) \\ &= \mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N, \Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu)) + \mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N, (\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu))^c) \\ &\leq \mathbb{P}(|\psi_N(f,\nu)| > N^{\epsilon}u_N, \Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu)) + \mathbb{P}\left((\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu))^c\right) \\ &\leq \mathbb{P}(|\psi_N(\bar{f},\nu)| > N^{\epsilon}u_N) + \mathbb{P}\left((\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu))^c\right) \end{aligned}$$

where the last inequality follows from the observation that $\frac{1}{M}$ tr $f(\hat{\mathbf{C}}) = \frac{1}{M}$ tr $\bar{f}(\hat{\mathbf{C}})$ on $\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu)$. Moreover, the family of events $\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu)$ holds with exponential high probability, which implies that $\mathbb{P}\left((\Lambda_{\kappa}^{\hat{\mathbf{C}}}(\nu))^{c}\right)$ converges towards 0 exponentially. Therefore, $|\psi_{N}(\bar{f},\nu)| \prec u_{N}$ implies (4.2) as expected. From now on, we thus assume that function f is supported by $\operatorname{Supp} \mu_{MP}^{(c)} + 2\kappa$

In order to establish (4.2), we evaluate the four terms of the righhandside of (1.15).

4.1. Step 1: Evaluation of $\mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\frac{X_N(\nu)X_N^*(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^+} f \,\mathrm{d}\mu_{MP}^{(c_N)}$

We evaluate this term using the Helffer-Sjöstrand formula. We keep the notations of paragraphs 2.5 and 2.2: we assume that the support of f is included into $[a_1, a_2]$ with $a_1 = (1 - \sqrt{c_N})^2 - 2\kappa$ and $a_2 = (1 + \sqrt{c_N})^2 + 2\kappa$. Moreover, the resolvent of matrix $\frac{\mathbf{X}_N \mathbf{X}_N^*}{B+1}$ is denoted $\mathbf{Q}_N(z)$ (we omit to mention that the matrices depend on ν), and $\beta_N(z)$ represents $\mathbb{E}((\mathbf{Q}_N(z))_{mm})$ for each m. We also denote by $\epsilon_N(z)$ the error term defined by (2.9) which satisfies $|\epsilon_N(z)| \leq \frac{1}{M^2} P_1(|z|) P_2(\frac{1}{\text{Im}z})$ on \mathbb{C}^+ for some nice polynomials P_1 and P_2 . Then, for $k \geq \text{deg}(P_2)$, it holds that

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^{+}} f \,\mathrm{d}\mu_{MP}^{(c_{N})}$$
$$= \frac{1}{\pi}\operatorname{Re}\int_{\mathcal{D}} \bar{\partial}\Phi_{k}(f)(z)(\beta_{N}(z) - t_{N}(z)) \,\mathrm{d}x \,\mathrm{d}y$$

where \mathcal{D} is defined as in paragraph 2.5. $\int_{\mathcal{D}} \left| \bar{\partial} \Phi_k(f)(z) \right| P_1(|z|) P_2(\frac{1}{\mathrm{Im}z}) \,\mathrm{d}x \,\mathrm{d}y$ is finite, and by (2.9), the following bound holds:

$$\begin{aligned} \left| \mathbb{E} \left[\frac{1}{M} \operatorname{Tr} \left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^*(\nu)}{B+1}) \right) \right] - \int_{\mathbb{R}^+} f \, \mathrm{d}\mu_{MP}^{(c_N)} \right| \leq \\ \frac{1}{M^2} \int_{\mathcal{D}} \left| \bar{\partial} \Phi_k(f)(z) \right| P_1(|z|) P_2(\frac{1}{\mathrm{Im}z}) \, \mathrm{d}x \, \mathrm{d}y \leq \frac{C}{B^2} \end{aligned}$$

for some nice constant C. We have therefore established the following result.

Lemma 4.1. It exists a nice constant C such that, for each ν ,

$$\left| \mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1}) \right) \right] - \int_{\mathbb{R}^{+}} f \, \mathrm{d}\mu_{MP}^{(c_{N})} \right| \leq \frac{C}{B^{2}}.$$
 (4.4)

4.2. Step 2: Evaluation of
$$\frac{1}{M} \operatorname{Tr} \left(f(\tilde{C}(\nu)) \right) - \mathbb{E} \left[\frac{1}{M} \operatorname{Tr} \left(f(\tilde{C}(\nu)) \right) \right]$$

In order to evaluate the above term, we use the Gaussian concentration inequality introduced in Paragraph 2.3. We recall that $\tilde{\mathbf{C}}$ can be interpreted as a function of $(\mathbf{X}, \mathbf{X}^*)$ (see (3.16))). Therefore, $\frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right)$ can be written as $g(\mathbf{X}, \mathbf{X}^*)$ for some real valued function g. We establish in the following that g is $\mathcal{O}(\frac{1}{B})$ -Lipschitz, which in turn, will imply that

$$\left|\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right]\right| \prec \frac{1}{B}.$$
(4.5)

For this, we evaluate

$$\|\nabla g(\mathbf{X}, \mathbf{X}^*)\|^2 = \sum_{i,j} \left|\frac{\partial g}{\partial X_{i,j}}\right|^2 + \left|\frac{\partial g}{\partial \overline{X_{i,j}}}\right|^2 = 2\sum_{i,j} \left|\frac{\partial g}{\partial X_{i,j}}\right|^2.$$
(4.6)

Using classic identities for the derivation of Hermitian matrices, we obtain that

$$\frac{1}{M} \frac{\partial \operatorname{tr} f(\tilde{\mathbf{C}})}{\partial X_{ij}} = \frac{1}{M} \operatorname{tr} \left(f'(\tilde{\mathbf{C}}) \frac{\partial \tilde{\mathbf{C}}}{\partial X_{ij}} \right)$$

Straightforward calculations lead to

$$\sum_{i,j} \left| \frac{\partial g}{\partial X_{i,j}} \right|^2 = \frac{1}{M^2 (B+1)^2} \sum_{i=1}^M \left(f'(\tilde{\mathbf{C}}) (\mathbf{X} + \mathbf{\Gamma}) (\mathbf{I} + \mathbf{\Phi}_i) (\mathbf{X} + \mathbf{\Gamma})^* f'(\tilde{\mathbf{C}}) \right)_{ii}.$$

Using $\sup_i \|\mathbf{I} + \mathbf{\Phi}_i\| \leq C$ for some nice constant C as well as $\tilde{\mathbf{C}} = \frac{1}{B+1}(\mathbf{X} + \mathbf{\Gamma})(\mathbf{X} + \mathbf{\Gamma})^*$, we obtain immediately that

$$\sum_{i,j} \left| \frac{\partial g}{\partial X_{i,j}} \right|^2 \le \frac{C}{B^2} \frac{1}{M} \operatorname{tr} \left(f^{\prime 2}(\tilde{\mathbf{C}}) \tilde{\mathbf{C}} \right).$$

As $f \in C^{\infty}$ and is compactly supported, the function $\lambda \to \lambda f^{\prime 2}(\lambda)$ is bounded by some constant, and there exists a nice constant C such that

$$\|\nabla g(\mathbf{X}, \mathbf{X}^*)\|^2 \le \frac{C}{B^2}$$

This proves that g is $\mathcal{O}(\frac{1}{B})$ -Lipschitz. Paragraph 2.3 thus leads to (4.5).

4.3. Step 3: Evaluation of $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right)$

The goal of this paragraph is to establish the following Proposition.

Proposition 4.1. Let \tilde{D}_N the distribution supported by $\operatorname{Supp}(\mu_{MP}^{(c_N)})$ with Stieltjes transform

$$\tilde{p}_N(z) = (z t_N(z))' = \frac{(z t_N(z) \tilde{t}_N(z))^2}{1 - c(z t_N(z) \tilde{t}_N(z))^2}.$$
(4.7)

Then, if we denote $< \tilde{D}_N, f > by \ \tilde{\phi}_N(f)$, we have

$$\left|\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s_m''(\nu)}{s_m(\nu)}\right) \tilde{\phi}_N(f) v_N \mathbf{1}_{\alpha>2/3}\right| \prec u_N. \quad (4.8)$$

Remark 4.1. (4.8) implies that $\left|\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right| \prec \frac{1}{B}$ if $\alpha \leq 2/3$. If $\alpha > 2/3$, the dominant term of $\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)$ is the deterministic $\mathcal{O}\left(\frac{B}{N}\right)^2$ term $\left(\frac{1}{M}\sum_{m=1}^M \frac{s'_m(\nu)}{s_m(\nu)}\right) \tilde{\phi}_N(f) v_N$, and its substraction from $\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right)$ allows to retrieve a term stochastically dominated by u_N .

Remark 4.2. We notice that (3.35) leads immediately to

$$\left|\frac{1}{M}\operatorname{tr} f(\hat{\mathbf{C}})(\nu) - \frac{1}{M}\operatorname{tr} f(\tilde{\mathbf{C}})(\nu)\right| \prec \frac{B}{N} + \frac{1}{\sqrt{B}}$$
(4.9)

an evaluation which is considerably more pessimistic that (4.8). As seen below, the derivation of (4.8) is rather demanding, and is based on subtle effects. In order to understand why (4.9) can be improved, we consider the simple case $f(\lambda) = \log \lambda$. We thus have

$$\frac{1}{M}\operatorname{tr} f(\hat{\mathbf{C}})(\nu) - \frac{1}{M}\operatorname{tr} f(\tilde{\mathbf{C}})(\nu) = \frac{1}{M}\sum_{m=1}^{M} \left(\log s_m(\nu) - \log \hat{s}_m(\nu)\right)$$

which depends only on the estimators $(\hat{s}_m(\nu))_{m=1,...,M}$. We just provide a brief analysis of the above term. For this, we first remark that is possible to study $\frac{1}{M}\sum_{m=1}^{M} (\log s_m(\nu) - \log \hat{s}_m(\nu))$ on the event $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$ defined by (3.32). For each m, we expand around s_m the logarithm up to the second order, and obtain that

$$\frac{1}{M} \sum_{m=1}^{M} \left(\log s_m(\nu) - \log \hat{s}_m(\nu) \right)$$
$$= -\frac{1}{M} \sum_{m=1}^{M} (\hat{s}_m - s_m) \frac{1}{s_m} + \frac{1}{M} \sum_{m=1}^{M} \frac{1}{2} \left(\frac{\hat{s}_m - s_m}{\theta_m} \right)^2 \quad (4.10)$$

where for each m, θ_m is located between s_m and \hat{s}_m . Lemma A.5 allows to conclude that the second term of the right hand side of (4.10) is dominated by $\frac{1}{B} + \left(\frac{B}{N}\right)^4 = \mathcal{O}(u_N)$ term. In order to evaluate the first term of the r.h.s. of (4.10), we remark that (A.9) leads to

$$\frac{1}{M}\sum_{m=1}^{M} (\mathbb{E}(\hat{s}_m - s_m)) \frac{1}{s_m} = \frac{1}{2M}\sum_{m=1}^{M} \frac{s''_m}{s_m} v_N + \mathcal{O}\left(\left(\frac{B}{N}\right)^3 + \frac{1}{N}\right)$$

As $\frac{\hat{s}_m}{s_m} = \frac{\mathbf{x}_m(\mathbf{I} + \mathbf{\Phi}_m)\mathbf{x}_m^*}{B+1}$ we finally remark that

$$\frac{1}{M}\sum_{m=1}^{M}\frac{\hat{s}_m - \mathbb{E}(\hat{s}_m)}{s_m}$$

can be interpreted as a recentered quadratic form of the M(B+1)-dimensional vector $\mathbf{x} = (\mathbf{x}_1^T, \dots, \mathbf{x}_M^T)^T$. The stochastic domination relation

$$\left|\frac{1}{M}\sum_{m=1}^{M}\frac{\hat{s}_m - \mathbb{E}(\hat{s}_m)}{s_m}\right| \prec \frac{1}{B}$$

then follows from the Hanson-Wright inequality. Putting all the pieces together, and using that $\frac{1}{B} + \mathcal{O}\left(\left(\frac{B}{N}\right)^3 + \frac{1}{N}\right) = \mathcal{O}(u_N)$ and that $v_N = o(u_N)$ if $\alpha < 2/3$,

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 35

 $we \ obtain \ that$

$$\left|\frac{1}{M}\sum_{m=1}^{M}\left(\log s_{m}(\nu) - \log \hat{s}_{m}(\nu)\right) + \frac{1}{2M}\sum_{m=1}^{M}\frac{s_{m}'(\nu)}{s_{m}(\nu)}v_{N}\,\mathbf{1}_{\alpha>2/3}\right| \prec u_{N}$$

Comparing this result with (4.8), we deduce that $\langle \tilde{D}_N, f \rangle = -1$. We just check this formula directly. For this, we notice that function $z \to \log z$ is holomorphic inside a neighbourhood of the interval $[a_1, a_2]$. We consider the expression (2.21) of $\langle \tilde{D}_N, f \rangle$ and remark that if $(\partial \mathcal{R}_{\epsilon})_{-}$ denotes the negatively oriented contour

$$(\partial \mathcal{R}_{\epsilon})_{-} = \{\lambda \pm i\epsilon, \lambda \in [a_1, a_2]\} \cup \{a_1 + iy, y \in [-\epsilon, \epsilon]\} \cup \{a_2 + iy, y \in [\epsilon, -\epsilon]\}$$

then, by (2.21), $\langle \tilde{D}_N, f \rangle$ can also be written as the contour integral

$$< \tilde{D}_N, f > = \lim_{\epsilon \to 0} \frac{1}{2i\pi} \int_{(\partial \mathcal{R}_\epsilon)_-} \log z \, \tilde{p}_N(z) \, dz.$$

But, the above contour integral does not depend on ϵ , so that for each ϵ , we have

$$< \tilde{D}_N, f >= \frac{1}{2i\pi} \int_{(\partial \mathcal{R}_{\epsilon})_-} \log z \, \tilde{p}_N(z) \, dz.$$

Using the expression of $\tilde{p}_N(z)$ and the integration by part trick, we get that

$$< \tilde{D}_N, f > = -\frac{1}{2i\pi} \int_{(\partial \mathcal{R}_\epsilon)_-} t_N(z) \, dz$$

Taking the limit $\epsilon \to 0$, and using the Stieltjes inversion formula for the Marcenko-Pastur distribution $\mu_{MP}^{(c_N)}$, we eventually obtain that

$$< \tilde{D}_N, f> = -\frac{1}{\pi} \lim_{\epsilon \to 0} \int_{a_1}^{a_2} \operatorname{Im}(t_N(\lambda + i\epsilon)) d\lambda = -\mu_{MP}^{(c_N)}([a_1, a_2]) = -1$$

which is the expected result.

Proof. We now establish (4.8). In order to simplify the notations, we put

$$\tilde{r}_N(\nu) = \frac{1}{2M} \sum_{m=1}^M \frac{s_m''(\nu)}{s_m(\nu)}.$$
(4.11)

The Helffer-Sjöstrand formula implies that

$$\frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}) - \frac{1}{M} \operatorname{tr} f(\tilde{\mathbf{C}}) - \tilde{r}_N(\nu) \,\tilde{\phi}_N(f) \, v_N \, \mathbf{1}_{\alpha > 2/3} = \\ \frac{1}{\pi} \operatorname{Re} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\bar{\partial} \Phi_k(f)(z) \left[\frac{1}{M} (\operatorname{tr} \,\hat{\mathbf{Q}}(z) - \operatorname{tr} \,\tilde{\mathbf{Q}}(z)) - \tilde{r}_N(\nu) \,\tilde{p}_N(z) \, v_N \, \mathbf{1}_{\alpha > 2/3} \right].$$
4.3.1. Reduction to the study of ζ

We define

$$\zeta = \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial}\Phi_k(f)(z) \frac{1}{M} \sum_{m=1}^M (z\mathbf{Q})'_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right) \tag{4.12}$$

where we recall that the row vectors $(\mathbf{x}_m)_{m=1,...,M}$ are the rows of the i.i.d. matrix **X**. We establish in this paragraph that

$$\left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \, \bar{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \mathrm{tr} \left\{ \hat{\mathbf{Q}} - \tilde{\mathbf{Q}} \right\} - \tilde{r}_N(\nu) \, \tilde{p}_N(z) \, v_N \, \mathbf{1}_{\alpha > 2/3} \right) - \zeta \right| \prec u_N.$$
(4.13)

It turns out that by Lemma 4.2 and Lemma 4.4 in Paragraph 4.3.2 below, $\dot{\zeta}$ verifies the key properties :

$$|\zeta| \le |\zeta - \mathbb{E}\zeta| + |\mathbb{E}\zeta| \prec \frac{1}{B}.$$

(4.8) will then follow directly from (4.13).

Plugging in the integral expression of ζ , and using the expression (4.7), we get:

$$\begin{split} \left| \int_{\mathcal{D}} \bar{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \operatorname{tr} \left\{ \hat{\mathbf{Q}} - \tilde{\mathbf{Q}} \right\} - \tilde{r}_N(\nu) \, \tilde{p}_N(z) \, v_N \, \mathbf{1}_{\alpha > 2/3} \right) \mathrm{d}x \, \mathrm{d}y - \zeta \right| \\ &= \left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \overline{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \operatorname{tr} \left\{ \hat{\mathbf{Q}} - \tilde{\mathbf{Q}} \right\} - \tilde{r}_N \, (zt_N(z))' \, v_N \, \mathbf{1}_{\alpha > 2/3} \right. \\ &\left. - \frac{1}{M} \sum_{m=1}^M (z \mathbf{Q})'_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right) \right|. \end{split}$$

We recall the definition of $\Theta := \hat{\mathbf{C}} - \tilde{\mathbf{C}}$ from (3.29). We will proceed in three steps, which, in turn, will imply (4.13):

$$\left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \mathrm{tr} \left\{ \hat{\mathbf{Q}} - \tilde{\mathbf{Q}} \right\} + \frac{1}{M} \mathrm{tr} \left\{ \mathbf{Q}^2 \boldsymbol{\Theta} \right\} \right) \right| \prec u_N \quad (4.14)$$

2.

1.

$$\left| \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \operatorname{tr} \{ \mathbf{Q}^2 \boldsymbol{\Theta} \} -2 \frac{1}{M} \operatorname{tr} \frac{\mathbf{X} \mathbf{X}^*}{B+1} \mathbf{Q}^2 (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \right) \right| \prec u_N \quad (4.15)$$

3.

$$\left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \times \left(2 \frac{1}{M} \mathrm{tr} \, \frac{\mathbf{X} \mathbf{X}^*}{B+1} \mathbf{Q}^2 (\mathbf{I} - \hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2}) - \tilde{r}_N \left(z t_N(z) \right)' v_N \, \mathbf{1}_{\alpha > 2/3} - \frac{1}{M} \sum_{m=1}^M (z \mathbf{Q})'_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right) \right| \prec u_N \quad (4.16)$$

Step 1. Using the well-known identity $\mathbf{A}^{-1} - \mathbf{B}^{-1} = \mathbf{B}^{-1}(\mathbf{B} - \mathbf{A})\mathbf{A}^{-1}$, we express $\hat{\mathbf{Q}} - \tilde{\mathbf{Q}}$ as:

$$\hat{\mathbf{Q}} - \tilde{\mathbf{Q}} = -\tilde{\mathbf{Q}}\boldsymbol{\Theta}\hat{\mathbf{Q}}.$$
(4.17)

We claim that it is possible to approximate tr $\tilde{\mathbf{Q}}\Theta\hat{\mathbf{Q}}$ by tr $\mathbf{Q}\Theta\mathbf{Q}.$ Indeed, we have

$$\begin{aligned} \operatorname{tr} \, \mathbf{Q} \Theta \mathbf{Q} - \operatorname{tr} \, \mathbf{Q} \Theta \mathbf{Q} | \\ &= |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \hat{\mathbf{Q}} - \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} + \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} - \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \mathbf{Q} - \operatorname{tr} \, \mathbf{Q} \Theta \mathbf{Q} | \\ &\leq |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \hat{\mathbf{Q}} - \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} | + |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} - \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \mathbf{Q} | + |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \mathbf{Q} - \operatorname{tr} \, \mathbf{Q} \Theta \mathbf{Q} | \\ &:= T_1 + T_2 + T_3. \end{aligned}$$

The following rough bounds are enough to control T_1 (we used (1.25) to control the norm of the resolvents):

$$T_1 = |\mathrm{tr} \; \tilde{\mathbf{Q}} \Theta(\hat{\mathbf{Q}} - \tilde{\mathbf{Q}})| = |\mathrm{tr} \; \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} \Theta \hat{\mathbf{Q}}| \le M \|\tilde{\mathbf{Q}}\|^2 \|\hat{\mathbf{Q}}\| \|\|\Theta\|^2 \le \frac{1}{\mathrm{Im}^3 z} M \|\Theta\|^2.$$

Concerning T_2 and T_3 , we write similarly that $\tilde{\mathbf{Q}} - \mathbf{Q} = -\tilde{\mathbf{Q}}\tilde{\mathbf{\Delta}}\mathbf{Q}$, and obtain that

$$T_{2} = |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \tilde{\mathbf{Q}} - \operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \mathbf{Q}| \le M \|\tilde{\mathbf{Q}}\|^{2} \|\mathbf{Q}\| \|\tilde{\mathbf{\Delta}}\| \Theta\| \le \frac{1}{\operatorname{Im}^{3} z} M \|\tilde{\mathbf{\Delta}}\| \|\Theta\|$$
$$T_{3} = |\operatorname{tr} \, \tilde{\mathbf{Q}} \Theta \mathbf{Q} - \operatorname{tr} \, \mathbf{Q} \Theta \mathbf{Q}| \le M \|\tilde{\mathbf{Q}}\| \|\mathbf{Q}\|^{2} \|\tilde{\mathbf{\Delta}}\| \Theta\| \le \frac{1}{\operatorname{Im}^{3} z} M \|\tilde{\mathbf{\Delta}}\| \|\Theta\|.$$

Plugging these estimations into the left hand side of (4.14), we obtain that

$$\begin{split} \left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \left(\frac{1}{M} \mathrm{tr} \left\{ \hat{\mathbf{Q}} - \tilde{\mathbf{Q}} \right\} - \frac{1}{M} \mathrm{tr} \left\{ \mathbf{Q}^2 \boldsymbol{\Theta} \right\} \right) \right| \\ & \leq \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y |\overline{\partial} \Phi_k(f)(z)| \frac{1}{M} \left(T_1 + T_2 + T_3 \right) \\ & \leq C(\|\boldsymbol{\Theta}\|^2 + 2\|\tilde{\boldsymbol{\Delta}}\|\boldsymbol{\Theta}\|). \end{split}$$

Moreover, the concentration results (3.35) for $\|\Theta\|$ and (3.26) for $\|\dot{\Delta}\|$ from Proposition 3.1, imply that

$$\|\mathbf{\Theta}\|^2 + 2\|\mathbf{\Theta}\|\|\tilde{\mathbf{\Delta}}\| \prec \frac{1}{B} + \frac{1}{\sqrt{B}}\frac{B}{N} + \left(\frac{B}{N}\right)^3 = u_N.$$

This eventually establishes (4.14).

Step 2. We claim that:

$$\left\| \boldsymbol{\Theta} - \left((\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^*}{B+1} + \frac{\mathbf{X} \mathbf{X}^*}{B+1} (\mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \right) \right\| \prec u_N.$$
(4.18)

We recall that $\hat{\mathbf{S}}$ can be written using the definition (1.12) of $\tilde{\mathbf{C}}$, and use the decomposition (3.25) of $\tilde{\mathbf{C}}$ from Corollary 3.2. Using these results, we get that

$$\hat{\mathbf{S}} = \mathbf{D}^{1/2} \, \tilde{\mathbf{C}} \, \mathbf{D}^{1/2} = \mathbf{D}^{1/2} \left(\frac{\mathbf{X} \mathbf{X}^*}{B+1} + \tilde{\mathbf{\Delta}} \right) \mathbf{D}^{1/2}.$$

Plugging this expression of $\hat{\mathbf{S}}$ into (3.30), we obtain easily that

$$\begin{split} \boldsymbol{\Theta} = & (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \left(\frac{\mathbf{X} \mathbf{X}^*}{B+1} + \tilde{\mathbf{\Delta}} \right) \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} \\ & + \left(\frac{\mathbf{X} \mathbf{X}^*}{B+1} + \tilde{\mathbf{\Delta}} \right) (\mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \\ & := \boldsymbol{\Theta}_1 + \boldsymbol{\Theta}_2. \end{split}$$

As $\tilde{\Delta}$ is a negligible quantity, one should expect that the leading quantity in Θ_1 and Θ_2 is respectively $(\hat{\mathbf{D}}^{-1/2}\mathbf{D}^{1/2} - \mathbf{I})\frac{\mathbf{X}\mathbf{X}^*}{B+1}\mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2}$ and $\frac{\mathbf{X}\mathbf{X}^*}{B+1}(\mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2} - \mathbf{I})$. To prove it, write:

$$\begin{aligned} \left\| \boldsymbol{\Theta}_{1} - (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} \right\| \\ &= \left\| (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \tilde{\boldsymbol{\Delta}} \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} \right\| \\ &\leq \| \hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I} \| \| \tilde{\boldsymbol{\Delta}} \| \| \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} \|. \end{aligned}$$
(4.19)

 $\tilde{\Delta}$ is controlled by (3.26) from Corollary 3.2, and $\hat{\mathbf{D}}^{-1/2}\mathbf{D}^{1/2}-\mathbf{I}$ is controlled by (3.33) from Lemma 3.3 (it is a diagonal matrix which elements are stochastically dominated by Lemma 3.3). Moreover, from Lemma 3.3, it holds that $\|\mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2}\| \prec 1$. Combining these estimations into (4.19), one get:

$$\left\| \boldsymbol{\Theta}_{1} - (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} \right\| \prec \left(\frac{1}{\sqrt{B}} + \frac{B^{2}}{N^{2}} \right) \frac{B}{N} = \mathcal{O}(u_{N}).$$

$$(4.20)$$

Using that $\|\hat{\mathbf{D}}^{-1/2}\mathbf{D}^{1/2} - \mathbf{I}\| \prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}$ as well as (2.5) from Paragraph 2.2 to control the norm of $\mathbf{XX}^*/(B+1)$, one can further approximate $(\hat{\mathbf{D}}^{-1/2}\mathbf{D}^{1/2} - \mathbf{I})\frac{\mathbf{XX}^*}{B+1}\mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2}$ by $(\hat{\mathbf{D}}^{-1/2}\mathbf{D}^{1/2} - \mathbf{I})\frac{\mathbf{XX}^*}{B+1}$. In particular, it is easy to check that

$$\left\| \boldsymbol{\Theta}_1 - (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^*}{B+1} \right\| \prec u_N.$$
(4.21)

Similarly for Θ_2 , one would obtain:

$$\left\| \boldsymbol{\Theta}_2 - \frac{\mathbf{X}\mathbf{X}^*}{B+1} (\mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \right\| \prec u_N.$$
(4.22)

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 39

Combining (4.21) and (4.22), we obtain (4.18). To finish the proof of Step 2, it remains to consider tr $\mathbf{Q}^2 \Theta$ and prove (4.15). Remark that $\mathbf{X}\mathbf{X}^*/(B+1)$ and its resolvent \mathbf{Q} commutes.

tr
$$\mathbf{Q}^{2} \left((\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} + \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} (\mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \right)$$

= 2 tr $\frac{\mathbf{X} \mathbf{X}^{*}}{B+1} \mathbf{Q}^{2} (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I})$ (4.23)

Therefore, using (4.23):

$$\left\| \frac{1}{M} \operatorname{tr} \mathbf{Q}^{2} \mathbf{\Theta} - 2 \frac{1}{M} \operatorname{tr} \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} \mathbf{Q}^{2} (\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \right\|$$

$$\leq \|\mathbf{Q}\|^{2} \left\| \mathbf{\Theta} - \left((\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} + \frac{\mathbf{X} \mathbf{X}^{*}}{B+1} (\mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \right) \right\| \quad (4.24)$$

so that the left hand side of (4.18) is recognised in the right hand side of (4.24). We can eventually prove (4.15) by following the same idea as in Step 1:

$$\begin{aligned} \left\| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \, \frac{1}{M} \left(\mathrm{tr} \left\{ \mathbf{Q}^2 \mathbf{\Theta} \right\} - 2\mathrm{tr} \, \frac{\mathbf{X} \mathbf{X}^*}{B+1} \mathbf{Q}^2(\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \right) \right\| \\ &\leq \left\| \mathbf{\Theta} - \left((\hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2} - \mathbf{I}) \frac{\mathbf{X} \mathbf{X}^*}{B+1} + \frac{\mathbf{X} \mathbf{X}^*}{B+1} (\mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2} - \mathbf{I}) \right) \right\| \\ &\qquad \times \underbrace{\int_{\mathcal{D}} |\overline{\partial} \Phi_k(f)(z)| \frac{1}{\mathrm{Im}^2 z} \, \mathrm{d}x \, \mathrm{d}y}_{<+\infty} .\end{aligned}$$

This proves (4.15) and ends Step 2.

Step 3. By definition of the resolvent, the following identity holds $\left(\frac{\mathbf{X}\mathbf{X}^*}{B+1} - z\mathbf{I}_M\right)\mathbf{Q}(z) = \mathbf{I}_M$, which leads to the so-called resolvent identity:

$$\frac{\mathbf{X}\mathbf{X}^*}{B+1}\mathbf{Q} = \mathbf{I}_M + z\mathbf{Q}.$$
(4.25)

Using (4.25) as well the identity $\mathbf{Q}'(z) = \mathbf{Q}^2(z)$ one can write:

$$\frac{1}{M} \operatorname{tr} \frac{\mathbf{X}\mathbf{X}^*}{B+1} \mathbf{Q}^2 (\mathbf{I} - \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2}) = \frac{1}{M} \operatorname{tr} (\mathbf{I} + z\mathbf{Q}) \mathbf{Q} (\mathbf{I} - \mathbf{D}^{1/2} \hat{\mathbf{D}}^{-1/2}) = \frac{1}{M} \sum_{m=1}^M (z\mathbf{Q})'_{mm} \left(1 - \sqrt{\frac{s_m}{\hat{s}_m}} \right). \quad (4.26)$$

To handle $1 - \sqrt{\frac{s_m}{\hat{s}_m}}$ we use the following Taylor expansion: define the application h by $h(u) = 1 - \frac{1}{\sqrt{u}}$, with $h'(u) = \frac{1}{2} \frac{1}{u^{3/2}}$ and $h''(u) = -\frac{3}{4} \frac{1}{u^{5/2}}$. A Taylor

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 40

expansion to the second order of h around 1 provides:

$$h\left(\frac{\hat{s}_m}{s_m}\right) = h(1) + \left(\frac{\hat{s}_m}{s_m} - 1\right)h'(1) + \frac{1}{2}\left(\frac{\hat{s}_m}{s_m} - 1\right)^2 h''(\theta_m)$$
$$= \frac{1}{2s_m}(\hat{s}_m - s_m) + \frac{1}{2}\frac{h''(\theta_m)}{s_m^2}(\hat{s}_m - s_m)^2$$

where θ_m is some random quantity between \hat{s}_m and s_m . Therefore (4.26) becomes

$$\frac{1}{M} \operatorname{tr} \left((z\mathbf{Q})'(\mathbf{I} - \mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2}) \right)$$

= $\frac{1}{M} \operatorname{tr} \left((z\mathbf{Q})' \operatorname{diag} \left(\frac{\hat{s}_m - s_m}{2s_m} + \frac{1}{2} \frac{h''(\theta_m)(\hat{s}_m - s_m)^2}{s_m^2} : m \in \{1, \dots, M\} \right) \right).$

Lemma 3.1 implies that the set $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$ defined by (3.32) holds with exponentially high probability. Therefore, it is sufficient to study the term $\frac{1}{M}$ tr $(z\mathbf{Q})'(\mathbf{I} - \mathbf{D}^{1/2}\hat{\mathbf{D}}^{-1/2})$ on the event $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$. If $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$ holds, θ_m belongs to $[\underline{s}, \overline{s}] + \epsilon$ for each $m \in \{1, \ldots, M\}$, and $\sup_{m \ge 1} |h''(\theta_m)|$ is bounded by a nice constant. Moreover, as $\inf_{\nu} \inf_{m \ge 1} s_m(\nu)$ is bounded away from zero, there exists a nice constant C for which the inequality

$$\begin{aligned} \left| \frac{1}{M} \text{ tr } \left((z\mathbf{Q})' \text{diag} \left(\frac{1}{2} \frac{h''(\theta_m)(\hat{s}_m - s_m)^2}{s_m^2} : m \in \{1, \dots, M\} \right) \right) \right| \\ & \leq C(\|\mathbf{Q}\| + z \|\mathbf{Q}\|^2) \frac{1}{M} \sum_{m=1}^M (\hat{s}_m - s_m)^2 \leq C(z) \frac{1}{M} \sum_{m=1}^M (\hat{s}_m - s_m)^2 \end{aligned}$$

holds on $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$, where we recall that C(z) can be written as $P_1(|z|)P_2(\frac{1}{\mathrm{Im}z})$ for some nice polynomials P_1 and P_2 . Following again the same argument as in Step 1, we obtain that

$$\begin{split} \left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \overline{\partial} \Phi_k(f)(z) \left\{ \frac{1}{M} \operatorname{tr} (z\mathbf{Q})' (\mathbf{I} - \hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2}) \right. \\ \left. - \frac{1}{M} \operatorname{tr} (z\mathbf{Q})' \operatorname{diag} \left(\frac{\hat{s}_m - s_m}{2s_m} : m \in \{1, \dots, M\} \right) \right\} \right| \\ \leq C \frac{1}{M} \sum_{m=1}^M (\hat{s}_m - s_m)^2 \end{split}$$

on $\Lambda_\epsilon^{\hat{\mathbf{D}}}(\nu)$ provided $k\geq \mathrm{Deg}(P_2)$. Lemma A.5 in Appendix implies that

$$\frac{1}{M} \sum_{m=1}^{M} (\hat{s}_m - s_m)^2 \prec \frac{1}{B} + \frac{B^4}{N^4} = \mathcal{O}(u_N).$$

We have thus shown that

$$\left| \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \overline{\partial} \Phi_k(f)(z) \left\{ \frac{1}{M} \mathrm{tr} \, (z\mathbf{Q})' (\mathbf{I} - \hat{\mathbf{D}}^{-1/2} \mathbf{D}^{1/2}) - \frac{1}{M} \, \mathrm{tr} \, (z\mathbf{Q})' \mathrm{diag} \left(\frac{\hat{s}_m - s_m}{2s_m} : m \in \{1, \dots, M\} \right) \right\} \right|$$

 $\prec u_N.$

We denote by $\eta_N(z)$ the term defined by

$$\eta_N(z) = \frac{1}{M} \sum_{m=1}^M \left(z \mathbf{Q} \right)'_{mm} \left(\frac{\hat{s}_m - s_m}{s_m} - \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right) - \tilde{r}_N(z) (z t_N(z))' v_N \mathbf{1}_{\alpha > 2/3} \quad (4.27)$$

and define δ_N as

$$\delta_N = \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \,\eta_N(z).$$

In order to establish (4.16), it is sufficient to prove that $|\delta_N| \prec u_N$. For this, we first remark that $\hat{s}_m = s_m \frac{\mathbf{x}_m(\mathbf{I} + \boldsymbol{\Phi}_m)\mathbf{x}_m^*}{B+1}$, so that $\eta_N(z)$ can also be written as

$$\eta_N(z) = \frac{1}{M} \sum_{m=1}^M (z\mathbf{Q})'_{mm} \frac{\mathbf{x}_m \Phi_m \mathbf{x}_m^*}{B+1} - \tilde{r}_N(zt_N(z))' v_N \mathbf{1}_{\alpha > 2/3}.$$
 (4.28)

We express $\eta_N(z)$ as $\eta_N(z) = \eta_{1,N}(z) + \eta_{2,N}(z) + \eta_{3,N}(z)$ where $(\eta_{i,N})_{i=1,2,3}$ are defined by

$$\eta_{1,N}(z) = \frac{1}{M} \sum_{m=1}^{M} (z\mathbf{Q})'_{mm} \left(\frac{\mathbf{x}_m \mathbf{\Phi}_m \mathbf{x}_m^*}{B+1} - \frac{1}{B+1} \text{tr } \mathbf{\Phi}_m \right)$$
$$\eta_{2,N}(z) = \frac{1}{M} \sum_{m=1}^{M} \mathbb{E}[(z\mathbf{Q})'_{mm}] \frac{1}{B+1} \text{tr } \mathbf{\Phi}_m - \tilde{r}_N(zt_N(z))' v_N \mathbf{1}_{\alpha > 2/3}$$
$$\eta_{3,N}(z) = \frac{1}{M} \sum_{m=1}^{M} ((z\mathbf{Q})'_{mm})^\circ \frac{1}{B+1} \text{tr } \mathbf{\Phi}_m$$

and denote by $(\delta_{i,N})_{i=1,2,3}$ the contributions of $(\eta_{i,N})_{i=1,2,3}$ to δ_N . We recall the definition (1.20) of $((z\mathbf{Q})'_{mm})^\circ$. In order to evaluate $\delta_{1,N}$, we remark that $|(z\mathbf{Q})'_{mm}| = |Q_{mm} + z\mathbf{Q}^2_{mm}| \leq C(z)$ and that

$$|\eta_{1,N}(z)| \le C(z) \sup_{m=1,\dots,M} \left| \frac{\mathbf{x}_m \boldsymbol{\Phi}_m \mathbf{x}_m^*}{B+1} - \frac{1}{B+1} \operatorname{tr} \boldsymbol{\Phi}_m \right|.$$

Therefore, for k large enough, $\delta_{1,N}$ verifies $|\delta_{1,N}| \leq C \sup_{m=1,\dots,M} \left| \frac{\mathbf{x}_m \mathbf{\Phi}_m \mathbf{x}_m^*}{B+1} - \frac{1}{B+1} \operatorname{tr} \mathbf{\Phi}_m \right|$. The Hanson-Wright inequality as well

as the evaluation (3.10) of the Frobenius norm of Φ_m imply that $|\delta_{1,N}| \prec u_N$. We now evaluate $\delta_{2,N}$. For this, we notice that the results reviewed in Paragraph 2.2.2 imply that $\mathbb{E}(z\mathbf{Q})'_{mm} = (z\beta_N(z))' = (zt_N(z))' + (z\epsilon_N(z))'$ where $|(z\epsilon_N(z))'| \leq \frac{C(z)}{M^2}$. Therefore, using (3.9), we obtain that

$$\eta_{2,N}(z)) = \left(\frac{1}{M} \sum_{m=1}^{2M} \frac{s_m''}{s_m}\right) (zt_N(z))' v_N + \epsilon_{1,N}(z) - \tilde{r}_N (zt_N(z))' v_N \mathbf{1}_{\alpha > 2/3} = \tilde{r}_N (zt_N(z))' v_N \mathbf{1}_{\alpha \le 2/3} + \epsilon_{1,N}(z)$$

where $\epsilon_{1,N}(z)$ verifies $|\epsilon_{1,N}(z)| \leq C(z) u_N$. We then deduce that $|\eta_{2,N}(z)| \leq C(z)u_N$ because if $\alpha \leq 2/3$, $v_N \leq u_N$. This implies that $|\delta_{2,N}| = \mathcal{O}(u_N)$. In order to address $\delta_{3,N}$, we interpret $\delta_{3,N}$ as a function g of $(\mathbf{X}, \mathbf{X}^*)$, and use the Gaussian concentration inequality presented in Paragraph 2.3. In particular, we verify that

$$\|\nabla g\| \le C \frac{1}{\sqrt{B}} \left(\frac{B}{N}\right)^2 = o(u_N).$$

As $\mathbb{E}(\delta_{3,N}) = 0$, this leads immediately to $|\delta_{3,N}| \prec u_N$. We just check that

$$\sum_{i,j} \left| \frac{\partial g}{\partial X_{ij}} \right|^2 \le C \frac{1}{B} \left(\frac{B}{N} \right)^4.$$
(4.29)

For this, we express $(z\mathbf{Q})'_{mm}$ as $(z\mathbf{Q})'_{mm} = Q_{mm} + z\mathbf{Q}^2_{mm}$ and notice that

$$\frac{\partial Q_{mm}}{\partial X_{ij}} = -Q_{mi} \left(\frac{\mathbf{X}^*}{B+1}\mathbf{Q}\right)_{jm}$$
$$\frac{\partial \mathbf{Q}^2_{mm}}{\partial X_{ij}} = -(\mathbf{Q}^2)_{mi} \left(\frac{\mathbf{X}^*}{B+1}\mathbf{Q}\right)_{jm} - Q_{mi} \left(\frac{\mathbf{X}^*}{B+1}\mathbf{Q}^2\right)_{jm}$$

Using the Jensen inequality, we obtain that

$$\left|\frac{\partial g}{\partial X_{ij}}\right|^2 \leq \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y |\overline{\partial}\Phi_k(f)(z)|^2 \frac{1}{M} \sum_{m=1}^M \left|\frac{\partial (z\mathbf{Q})'_{mm}}{\partial X_{ij}}\right|^2 \left(\frac{1}{B+1} \mathrm{tr} \, \Phi_m\right)^2.$$

Summing over i, j leads to the expected evaluation of (4.29) and to $|\delta_{3,N}| \prec u_N$. This, in turn, completes the proof of (4.16) and of (4.13).

Up to the Lemma 4.2 and Lemma 4.4, Theorem 4.1 is proved. $\hfill \Box$

4.3.2. Proof of Lemma 4.2 and Lemma 4.4

We now establish Lemma 4.2 and Lemma 4.4.

Lemma 4.2. The family of random variables $\zeta(\nu) - \mathbb{E}\zeta(\nu)$, $\nu \in [0, 1]$ verifies the following property:

$$|\zeta(\nu) - \mathbb{E}\zeta(\nu)| \prec \frac{1}{B}.$$
(4.30)

Proof. ζ defined by (4.12) can be written as

$$\zeta = \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \frac{1}{M} \sum_{m=1}^M Q_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right) + \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \frac{1}{M} \sum_{m=1}^M z(\mathbf{Q}^2)_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right)$$

 $:= \zeta_1 + \zeta_2.$

In the following, we omit to evaluate $|\zeta_1(\nu) - \mathbb{E}(\zeta_1(\nu))|$, and just establish that $|\zeta_2(\nu) - \mathbb{E}(\zeta_2(\nu))| \prec \frac{1}{B}$ using the Gaussian concentration inequality from Paragraph 2.3.

Recall that $\|\mathbf{x}_m\|_2^2$ is a $\chi^2_{2(B+1)}$ random variable. Therefore it is clear that:

$$\left|\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right| \prec \frac{1}{\sqrt{B}}.$$

Knowing this, the idea is to show that, conditioned on the event where the random variables $\left(\frac{\|\mathbf{x}_m\|_2^2}{B+1}-1\right)_{m=1,\ldots,M}$ are localized, which holds with exponentially high probability, ζ_2 is a $\mathcal{O}(\frac{1}{B^{1-\epsilon}})$ -Lipschitz function of the entries of matrix \mathbf{X} for any $\epsilon > 0$. Let $0 < \epsilon < \frac{1}{2}$, and define the family of events $A_{m,\epsilon}(\nu)$, $m = 1, \ldots, M, \nu \in [0, 1]$ given by

$$A_{m,\epsilon}(\nu) = \left\{ \frac{\|\mathbf{x}_m(\nu)\|_2^2}{B+1} \in \left[1 - \frac{B^{\epsilon}}{\sqrt{B}}, 1 + \frac{B^{\epsilon}}{\sqrt{B}}\right] \right\}$$
(4.31)

as well as $A_{\epsilon}(\nu) = \bigcap_{m=1}^{M} A_{m,\epsilon}(\nu)$. It is clear that the family of events $A_{m,\epsilon}(\nu)$, $m = 1, \ldots, M, \nu \in [0, 1]$ holds with exponentially high probability, and that the same property holds for the family $A_{\epsilon}(\nu), \nu \in [0, 1]$. We claim that it exists a family of \mathcal{C}^{∞} functions $(g_{B,\epsilon})_{B\geq 1}$ verifying

$$g_{B,\epsilon}(t) = \begin{cases} t-1 & \text{if } t \in \left[1 - \frac{B^{\epsilon}}{\sqrt{B}}, 1 + \frac{B^{\epsilon}}{\sqrt{B}}\right] \\ 0 & \text{if } t \notin \left[1 - 2\frac{B^{\epsilon}}{\sqrt{B}}, 1 + 2\frac{B^{\epsilon}}{\sqrt{B}}\right] \end{cases}$$

and

$$\sup_{t} |g_{B,\epsilon}(t)| \le C \frac{B^{\epsilon}}{\sqrt{B}}, \quad \sup_{t} |g'_{B,\epsilon}(t)| \le C$$
(4.32)

for each B, where C is a nice constant. Indeed consider $h \in C^{\infty}$ such that it verifies $|h(t)| \le 2|t|$ for each t and

$$h(t) = \begin{cases} t & \text{if } t \in [-1,1] \\ 0 & \text{if } t \notin [-2,2]. \end{cases}$$

Then, it is easy to check that the family $(g_{B,\epsilon})_{B\geq 1}$ defined by

$$g_{B,\epsilon}(t) = \frac{B^{\epsilon}}{\sqrt{B}} h\left(\frac{\sqrt{B}}{B^{\epsilon}} (t-1)\right)$$

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 44

satisfies the requirements (4.32).

We define $\tilde{\zeta}_{2,\epsilon}$ by

$$\tilde{\zeta}_{2,\epsilon} = \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \overline{\partial} \Phi_k(f)(z) \frac{1}{M} \sum_{m=1}^M (z\mathbf{Q}^2)_{mm} \,g_{B,\epsilon}\left(\frac{\|\mathbf{x}_m\|_2^2}{B+1}\right)$$

and notice that ζ_2 and $\tilde{\zeta}_{2,\epsilon}$ coincide on the exponentially high probability event $A_{\epsilon}(\nu)$. We claim that if $|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| \prec \frac{1}{B^{1-\epsilon}}$, then $|\zeta_2 - \mathbb{E}(\zeta_2)| \prec \frac{1}{B^{1-\epsilon}}$. Since ϵ is arbitrary and $B^{\epsilon} = \mathcal{O}(N^{\alpha\epsilon})$, Remark 2.1 will imply that $|\zeta_2 - \mathbb{E}(\zeta_2)| \prec \frac{1}{B}$. To justify this, we evaluate $\mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{1}{B^{1-\epsilon}}N^{\delta}\right)$ for each $\delta > 0$. It holds that

$$\mathbb{P}\left(|\zeta_2 - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha \epsilon + \delta}}{B}\right) \le \mathbb{P}\left(|\zeta_2 - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha \epsilon + \delta}}{B}, A_{\epsilon}\right) + \mathbb{P}(A_{\epsilon}^c).$$

As $\mathbb{P}(A^c_{\epsilon})$ converges towards zero exponentially, we have just to consider

$$\mathbb{P}\left(|\zeta_2 - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha \epsilon + \delta}}{B}, A_{\epsilon}\right)$$

and write, since ζ_2 and $\zeta_{2,\epsilon}$ coincide on A_{ϵ} ,

$$\mathbb{P}\left(|\zeta_2 - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha\epsilon + \delta}}{B}, A_\epsilon\right) = \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha\epsilon + \delta}}{B}, A_\epsilon\right)$$
$$\leq \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{N^{\alpha\epsilon + \delta}}{B} - |E(\zeta_2 - \tilde{\zeta}_{2,\epsilon})|, A_\epsilon\right).$$

We now prove that $|\mathbb{E}(\zeta_2 - \tilde{\zeta}_{2,\epsilon})|$ converges towards 0 exponentially. For this, we notice that as ζ_2 and $\tilde{\zeta}_{2,\epsilon}$ coincide on A_{ϵ} , then

$$\left|\mathbb{E}(\zeta_{2}-\tilde{\zeta}_{2,\epsilon})\right| = \left|\mathbb{E}((\zeta_{2}-\tilde{\zeta}_{2,\epsilon})\mathbb{I}_{A_{\epsilon}^{c}})\right| \leq \left(\mathbb{E}\left|\zeta_{2}-\tilde{\zeta}_{2,\epsilon}\right|^{2}\right)^{1/2} \left(\mathbb{P}(A_{\epsilon}^{c})\right)^{1/2}.$$

A rough evaluation of $\left(\mathbb{E}\left|\zeta_{2}-\tilde{\zeta}_{2,\epsilon}\right|^{2}\right)^{1/2}$ leads to $\left(\mathbb{E}\left|\zeta_{2}-\tilde{\zeta}_{2,\epsilon}\right|^{2}\right)^{1/2} \leq C$ for some nice constant C. Therefore, $\left(\mathbb{E}\left|\zeta_{2}-\tilde{\zeta}_{2,\epsilon}\right|^{2}\right)^{1/2}$ $(\mathbb{P}(A_{\epsilon}^{c}))^{1/2}$, and thus $|\mathbb{E}(\zeta_{2}-\tilde{\zeta}_{2,\epsilon})|$, converge towards 0 exponentially. For each N large enough, we thus have

$$\begin{split} \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{N^{\alpha\epsilon+\delta}}{B} \ -|\mathbb{E}(\zeta_2 - \tilde{\zeta}_{2,\epsilon})|, A_\epsilon\right) \\ & \leq \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{N^{\alpha\epsilon+\delta/2}}{B}, A_\epsilon\right) \\ & \leq \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{N^{\alpha\epsilon+\delta/2}}{B}\right). \end{split}$$

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 45

We have therefore established that

$$\mathbb{P}\left(|\zeta_2 - \mathbb{E}(\zeta_2)| > \frac{N^{\alpha \epsilon + \delta}}{B}, A_\epsilon\right) \le \mathbb{P}\left(|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| > \frac{N^{\alpha \epsilon + \delta/2}}{B}\right)$$

which eventually justifies that if $|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| \prec \frac{B^{\epsilon}}{B}$, then $|\zeta_2 - \mathbb{E}(\zeta_2)| \prec \frac{B^{\epsilon}}{B}$.

Therefore, it remains to prove that $|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| \prec \frac{B^{\epsilon}}{B}$. This is true by Lemma 4.3 below. The stochastic domination relation $|\zeta_1 - \mathbb{E}\zeta_1| \prec \frac{B^{\epsilon}}{B}$ is proved similarly. The completes the proof of Lemma 4.2.

Lemma 4.3.

$$|\tilde{\zeta}_{2,\epsilon} - \mathbb{E}(\tilde{\zeta}_{2,\epsilon})| \prec \frac{B^{\epsilon}}{B}$$

Proof. In the following, we evaluate the norm square of the gradient of $\tilde{\zeta}_{2,\epsilon}$ w.r.t. the variables $X_{i,j}, X_{i,j}^*$ and just compute $\sum_{i,j} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} \right|^2$ because $\sum_{i,j} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}^*} \right|^2$ is of the same order of magnitude.

We recall that

$$\frac{\partial (\mathbf{Q}^2)_{mm}}{\partial X_{ij}} = -\left(\frac{(\mathbf{Q}^2)_{mi}(\mathbf{X}^*\mathbf{Q})_{jm}}{B+1} + \frac{Q_{mi}(\mathbf{X}^*\mathbf{Q}^2)_{jm}}{B+1}\right).$$
(4.33)

Moreover it is clear that

$$\frac{\partial}{\partial X_{ij}} \left(g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \right) = \delta_{im} \frac{\overline{X_{m,j}}}{B+1} g'_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right).$$
(4.34)

Collecting the derivatives (4.33) and (4.34) we get after some algebra that

$$\frac{\partial}{\partial X_{ij}} \left(\sum_{m=1}^{M} (\mathbf{Q}^2)_{mm} g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \right) = \frac{\overline{X_{i,j}}}{B+1} g'_{B,\epsilon} \left(\frac{\|\mathbf{x}_i\|_2^2}{B+1} \right) (\mathbf{Q}^2)_{ii} - \sum_{m=1}^{M} g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \left(\frac{(\mathbf{Q}^2)_{mi} (\mathbf{X}^* \mathbf{Q})_{jm}}{B+1} + \frac{Q_{mi} (\mathbf{X}^* \mathbf{Q}^2)_{jm}}{B+1} \right). \quad (4.35)$$

It remains to control $\sum_{i,j} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} \right|^2$. From the integral representation of $\tilde{\zeta}_{2,\epsilon}$, the derivative with respect to X_{ij} is applied only on the integrand as follows:

$$\frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} = \frac{1}{M} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \frac{\partial}{\partial X_{ij}} \left(\sum_{m=1}^M z(\mathbf{Q}^2)_{mm} g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \right).$$

Plugging in the derivative computed in (4.35) we get:

$$\frac{\partial \bar{\zeta}_{2,\epsilon}}{\partial X_{ij}} = \frac{1}{M} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \, \overline{\partial} \Phi_k(f)(z) \, z \left\{ \frac{\overline{X_{i,j}}}{B+1} g'_{B,\epsilon} \left(\frac{\|\mathbf{x}_i\|_2^2}{B+1} \right) (\mathbf{Q}^2)_{ii} - \sum_{m=1}^M g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \left(\frac{(\mathbf{Q}^2)_{mi} (\mathbf{X}^* \mathbf{Q})_{jm}}{B+1} + \frac{Q_{mi} (\mathbf{X}^* \mathbf{Q}^2)_{jm}}{B+1} \right) \right\}.$$

Using the bounds of $g_{B,\epsilon}$ and $g'_{B,\epsilon}$ from inequalities (4.32), the observation that $g'_{B,\epsilon}(t) = 0$ if $|t-1| \ge \frac{2B^{\epsilon}}{\sqrt{B}}$, and that |z| is bounded on \mathcal{D} , one can write:

$$\begin{split} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} \right|^2 &\leq \frac{C}{M^2} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \left| \frac{\overline{X_{i,j}}}{B+1} g'_{B,\epsilon} \left(\frac{\|\mathbf{x}_i\|_2^2}{B+1} \right) (\mathbf{Q}^2)_{ii} \right. \\ &\quad \left. - \sum_{m=1}^M g_{B,\epsilon} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} \right) \left(\frac{(\mathbf{Q}^2)_{mi} (\mathbf{X}^* \mathbf{Q})_{jm}}{B+1} + \frac{Q_{mi} (\mathbf{X}^* \mathbf{Q}^2)_{jm}}{B+1} \right) \right|^2 \\ &\leq \frac{C}{M^2} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \left| \frac{\overline{X_{i,j}}}{B+1} (\mathbf{Q}^2)_{ii} \right|^2 \, \mathbbm{1} \left(\left| \frac{\|\mathbf{x}_i\|_2^2}{B+1} - 1 \right| \leq \frac{2B^\epsilon}{\sqrt{B}} \right) \\ &\quad + \frac{C}{M^2} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \left(\frac{B^\epsilon}{\sqrt{B}} \right)^2 \left| \sum_{m=1}^M \frac{|(\mathbf{Q}^2)_{mi}| \left| (\mathbf{X}^* \mathbf{Q})_{jm} \right|}{B+1} \right|^2 \\ &\quad + \frac{C}{M^2} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \left(\frac{B^\epsilon}{\sqrt{B}} \right)^2 \left| \sum_{m=1}^M \frac{|Q_{mi}| \left| (\mathbf{X}^* \mathbf{Q}^2)_{jm} \right|}{B+1} \right|^2 \\ &\quad = \frac{C}{M^2} \left(T_{ij}^{(1)} + T_{ij}^{(2)} + T_{ij}^{(3)} \right). \end{split}$$

It remains to sum over i, j.

$$\begin{split} &\sum_{i,j=1}^{M} T_{ij}^{(1)} \\ &= \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_{k}(f)(z) \right|^{2} \sum_{i,j=1}^{M} \left| \frac{\overline{X_{i,j}}}{B+1} (\mathbf{Q}^{2})_{ii} \right|^{2} \mathbb{1} \left(\left| \frac{\|\mathbf{x}_{i}\|_{2}^{2}}{B+1} - 1 \right| \leq \frac{2B^{\epsilon}}{\sqrt{B}} \right) \\ &\leq \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_{k}(f)(z) \right|^{2} \sum_{i=1}^{M} |(\mathbf{Q}^{2})_{ii}|^{2} \mathbb{1} \left(\left| \frac{\|\mathbf{x}_{i}\|_{2}^{2}}{B+1} - 1 \right| \leq \frac{2B^{\epsilon}}{\sqrt{B}} \right) \sum_{j=1}^{M} \left| \frac{\overline{X_{i,j}}}{B+1} \right|^{2} \\ &= \frac{C}{B+1} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_{k}(f)(z) \right|^{2} \sum_{i=1}^{M} |(\mathbf{Q}^{2})_{ii}|^{2} \frac{\|\mathbf{x}_{i}\|_{2}^{2}}{B+1} \mathbb{1} \left(\left| \frac{\|\mathbf{x}_{i}\|_{2}^{2}}{B+1} - 1 \right| \leq \frac{2B^{\epsilon}}{\sqrt{B}} \right) \\ &\leq \frac{C}{B+1} (1 + \frac{2B^{\epsilon}}{\sqrt{B}}) \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_{k}(f)(z) \right|^{2} \sum_{i=1}^{M} |(\mathbf{Q}^{2})_{ii}|^{2}. \end{split}$$

Since

$$\sum_{i=1}^{M} |(\mathbf{Q}^2)_{ii}|^2 \le M ||\mathbf{Q}||^4$$

it can be written that:

$$\sum_{i,j=1}^{M} T_{ij}^{(1)} \le C \frac{M}{B+1} \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \| \mathbf{Q} \|^4.$$

P. Loubaton and A. Rosuel/Coherency matrices of high-dimensional time series 47

Inspecting $T_{ij}^{(2)}$, one can see that by Jensen's inequality

$$\left|\sum_{m=1}^{M} \left| (\mathbf{Q}^{2})_{mi} \right| \left| (\mathbf{X}^{*} \mathbf{Q})_{jm} \right| \right|^{2} \le M \sum_{m=1}^{M} \left| (\mathbf{Q}^{2})_{mi} \right|^{2} \left| (\mathbf{X}^{*} \mathbf{Q})_{jm} \right|^{2}$$

so summing over i and j provides:

$$\sum_{i,j=1}^{M} T_{ij}^{(2)} \leq \frac{B^{2\epsilon} M}{(B+1)^3} \times \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \sum_{m=1}^{M} \left(\sum_{i=1}^{M} \left| (\mathbf{Q}^2)_{mi} \right|^2 \right) \left(\sum_{j=1}^{M} \left| (\mathbf{X}^* \mathbf{Q})_{jm} \right|^2 \right).$$

Notice that since $\sum_{i=1}^{M} |(\mathbf{Q}^2)_{mi}|^2$ is the square euclidean norm of line m of \mathbf{Q}^2 :

$$\sum_{i=1}^{M} \left| (\mathbf{Q}^2)_{mi} \right|^2 \le \|\mathbf{Q}^2\|^2 \le \|\mathbf{Q}\|^4.$$

Moreover,

$$\sum_{m=1}^{M} \left(\sum_{j=1}^{M} \left| (\mathbf{X}^* \mathbf{Q})_{jm} \right|^2 \right) = \text{tr } \mathbf{X}^* \mathbf{Q} \mathbf{Q}^* \mathbf{X}$$
$$= (B+1) \text{tr } ((\mathbf{I} + z \mathbf{Q}) \mathbf{Q}^*) \le M(B+1) (\|\mathbf{Q}\| + |z| \|\mathbf{Q}\|^2)$$

therefore

$$\sum_{i,j=1}^{M} T_{ij}^{(2)} \le B^{2\epsilon} \left(\frac{M}{B+1}\right)^2 \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \left|\overline{\partial}\Phi_k(f)(z)\right|^2 \|\mathbf{Q}^4\| (\|\mathbf{Q}\| + |z|\|\mathbf{Q}\|^2)$$

and similarly for $T_{ij}^{(3)}$ one get:

$$\sum_{i,j=1}^{M} T_{ij}^{(3)} \le B^{2\epsilon} \left(\frac{M}{B+1}\right)^2 \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \left|\overline{\partial}\Phi_k(f)(z)\right|^2 \|\mathbf{Q}^2\| (\|\mathbf{Q}\|^3 + |z|\|\mathbf{Q}\|^4).$$

Collecting the terms in $T_{ij}^{(1)}, T_{ij}^{(2)}$ and $T_{ij}^{(3)}$, and since $M/(B+1) = \mathcal{O}(1)$ by Assumption 1.3, we can write:

$$\sum_{i,j} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} \right|^2 \le \frac{C}{M^2} B^{2\epsilon} \int_{\mathcal{D}} \mathrm{d}x \, \mathrm{d}y \left| \overline{\partial} \Phi_k(f)(z) \right|^2 \left(\|\mathbf{Q}\|^4 + \|\mathbf{Q}^5\| + |z| \|\mathbf{Q}^6\| \right)$$

As $\|\mathbf{Q}\|^4 + \|\mathbf{Q}^5\| + |z|\|\mathbf{Q}^6\| \le C(z)$, we obtain that for k large enough,

$$\sum_{i,j} \left| \frac{\partial \tilde{\zeta}_{2,\epsilon}}{\partial X_{ij}} \right|^2 = \mathcal{O}\left(\frac{B^{2\epsilon}}{B^2} \right)$$

as expected.

It remains to study $\mathbb{E}[\zeta]$, and establish the following Lemma.

Lemma 4.4.

$$\mathbb{E}\zeta| = \mathcal{O}\left(\frac{1}{B}\right).$$

Proof. As in the proof of Lemma 4.2, we only consider

$$\mathbb{E}[\zeta_2] = \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\overline{\partial} \Phi_k(f)(z) \frac{1}{M} \sum_{m=1}^M z \mathbb{E}\left[(\mathbf{Q}^2)_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right]$$

as $\mathbb{E}[\zeta_1]$ is shown to be also $\mathcal{O}(\frac{1}{B})$ with the same argument. As $\mathbb{E}\left[\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right] = 0$, we have

$$\mathbb{E}\left[(\mathbf{Q}^2)_{mm} \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right] = \mathbb{E}\left[\left((\mathbf{Q}^2)_{mm} - \mathbb{E}[(\mathbf{Q}^2)_{mm}] \right) \left(\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right) \right].$$

Apply now the Cauchy-Schwartz inequality:

$$|\mathbb{E}[\zeta_2]| \leq \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,|\overline{\partial}\Phi_k(f)(z)| \frac{1}{M} \sum_{m=1}^M |z| \sqrt{\mathrm{Var}(\mathbf{Q}^2)_{mm}} \sqrt{\mathbb{E}\left|\frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1\right|^2}.$$
(4.36)

As it is clear that $\mathbb{E} \left| \frac{\|\mathbf{x}_m\|_2^2}{B+1} - 1 \right|^2 = \mathcal{O} \left(\frac{1}{B} \right)$, it remains to control $\operatorname{Var}(\mathbf{Q}^2)_{mm} = \operatorname{Var}(\operatorname{tr} \mathbf{Q}^2 \mathbf{e}_m \mathbf{e}_m^T)$ where $(\mathbf{e}_m)_{m=1,\ldots,M}$ is the canonical basis of \mathbb{C}^M . A direct application of (2.6) for i = 2 leads immediately to

$$\operatorname{Var}(\mathbf{Q}^2)_{mm} \le \frac{C(z)}{B} \tag{4.37}$$

for some nice constant C. Using (4.37) in (4.36), we get that for k large enough:

$$\begin{aligned} |\mathbb{E}\zeta_2| &\leq \frac{1}{B} \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,|\overline{\partial}\Phi_k(f)(z)| \sqrt{C(z)} \\ &\leq \frac{1}{B} \int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,|\overline{\partial}\Phi_k(f)(z)| (1+C(z)) \leq C \frac{1}{B}. \end{aligned}$$

This completes the proof of Lemma 4.4.

Remark 4.3. We notice that, instead of using (1.15), an alternative approach to study $\frac{1}{M}$ tr $f(\hat{\mathbf{C}}(\nu)) - \int f d\mu_{MP}^{(c_N)}$ could have been based on the decomposition

$$\frac{1}{M} \operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \int_{\mathbb{R}^{+}} f \, \mathrm{d}\mu_{MP}^{(c_{N})} = \frac{1}{M} \operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right)\right] + \\
\mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right)\right] - \mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right)\right] + \\
\mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M} \operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] + \\
\mathbb{E}\left[\frac{1}{M} \operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] - \int_{\mathbb{R}^{+}} f \, \mathrm{d}\mu_{MP}^{(c_{N})}. \tag{4.38}$$

The first term of the r.h.s. of (4.38) can be addressed using the Gaussian concentration inequality. However, the calculations are more complicated than the evaluation of $\frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right) - \mathbb{E} \left[\frac{1}{M} \operatorname{Tr} \left(f(\tilde{\mathbf{C}}(\nu)) \right) \right]$ because, considered as a function of $(\mathbf{X}, \mathbf{X}^*)$, $\frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right)$ is not a Lipschitz function. Using technics similar to those developed to evaluate $\zeta - \mathbb{E}(\zeta)$ (see Lemma 4.2), it could however be shown that

$$\left|\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right) - \mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\hat{\mathbf{C}}(\nu))\right)\right]\right| \prec \frac{1}{B}.$$
(4.39)

In order to evaluate the second term of the r.h.s. of (4.38), one should prove that

$$\mathbb{E}\left[\int_{\mathcal{D}} \mathrm{d}x \,\mathrm{d}y \,\left(\bar{\partial}\Phi_k(f)(z)\frac{1}{M} \mathrm{tr}\,\left\{\hat{\mathbf{Q}}-\tilde{\mathbf{Q}}\right\}-\tilde{r}_N(\nu)\,\tilde{p}_N(z)\,v_N\,\mathbf{1}_{\alpha>2/3}\right)-\zeta\right]\\=\mathcal{O}(u_N) \quad (4.40)$$

and $\mathbb{E}(\zeta) = \mathcal{O}(\frac{1}{B})$. The proof of (4.40) does not appear simpler than the proof of (4.13): the 3 steps that allowed to establish (4.13) should still be used, except that the stochastic domination properties should be replaced by properties of the mathematical expectation of the various terms. However, proving stochastic domination appears simpler than showing the desired properties of the above mathematical expectations. In sum, while the use of decomposition (4.38) allows to avoid Lemma 4.2, the justification of (4.39) needs to develop tools that are similar to those of Lemma 4.2, and the proof of (4.40) tends to be more complicated than the proof of (4.13). This explains why we have chosen to use decomposition (1.15) rather than (4.38).

4.4. Step 4: evaluation of
$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{C}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{X(\nu)X^{*}(\nu)}{B+1})\right)\right]$$

The Helffer-Sjöstrand formula implies that

$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right]$$
$$= \frac{1}{\pi}\operatorname{Re}\int_{\mathcal{D}} \mathrm{d}x\,\mathrm{d}y\,\overline{\partial}\Phi_{k}(f)(z)\mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z) - \mathbf{Q}_{N}(z)\right)\right].$$

Therefore, we are back to evaluate $\mathbb{E}\left[\frac{1}{M} \operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z) - \mathbf{Q}_{N}(z)\right)\right]$.

In order to simplify the exposition of the results of this paragraph, we introduce the following notation. If $(h_N(z))_{N\geq 1}$ is a sequence of complex-valued functions defined on \mathbb{C}^+ and if $(w_N)_{N\geq 1}$ is a sequence of positive real numbers, the notation $h_N(z) = \mathcal{O}_z(w_N)$ means that it exists two nice polynomials P_1 and P_2 such that $|h_N(z)| \leq w_N P_1(|z|) P_2(\frac{1}{\operatorname{Im} z})$ for each $z \in \mathbb{C}^+$.

In this paragraph, we establish the following Proposition.

Proposition 4.2. $\mathbb{E}\left[\frac{1}{M} \operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z) - \mathbf{Q}_{N}(z)\right)\right]$ can be written as

$$\mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z)-\mathbf{Q}_{N}(z)\right)\right] = \left(\frac{1}{M}\sum_{m=1}^{M}\frac{s_{m}'}{s_{m}}\right)^{2}p_{N}(z)v_{N} - \left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s_{m}''}{s_{m}}\right)\tilde{p}_{N}(z)v_{N} + \mathcal{O}_{z}\left(\left(\frac{B}{N}\right)^{3}+\frac{1}{N}\right).$$
 (4.41)

The Helffer-Sjöstrand formula thus leads to the following Corollary: $\begin{bmatrix} x & y \\ y \end{bmatrix}$

Corollary 4.1.
$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\mathbf{C}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] \text{ is given by}$$
$$\mathbb{E}\left[\frac{1}{M}\operatorname{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\operatorname{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right] = \left(\frac{1}{M}\sum_{m=1}^{M}\frac{s'_{m}}{s_{m}}\right)^{2}\phi_{N}(f) v_{N} - \left(\frac{1}{2M}\sum_{m=1}^{M}\frac{s'_{m}}{s_{m}}\right)\tilde{\phi}_{N}(f) v_{N} + \mathcal{O}\left(\left(\frac{B}{N}\right)^{3} + \frac{1}{N}\right).$$
(4.42)

Corollary 4.1 first implies that $\mathbb{E}\left[\frac{1}{M}\mathrm{Tr}\left(f(\tilde{\mathbf{C}}(\nu))\right) - \frac{1}{M}\mathrm{Tr}\left(f(\frac{\mathbf{X}(\nu)\mathbf{X}^{*}(\nu)}{B+1})\right)\right]$ is a $\mathcal{O}\left(\frac{B}{N}\right)^{2}$, a result which is not a priori obvious. In particular, the stochastic representation (3.1) of matrix $\tilde{\mathbf{C}}$ can be shown to provide the more pessimistic $\mathcal{O}(\frac{B}{N})$ rate of convergence. The comparison of (4.42) with (4.8) also leads to the conclusion that if $\alpha > 2/3$, the dominant $\mathcal{O}\left(\frac{B}{N}\right)^{2}$ deterministic term of $\frac{1}{M}\mathrm{tr}\left(f(\hat{\mathbf{C}}(\nu)) - f(\tilde{\mathbf{C}}(\nu)\right)$ is cancelled by the second term of the righthandside of (4.42), thus explaining the structure of the $\mathcal{O}\left(\frac{B}{N}\right)^{2}$ deterministic correction of $\frac{1}{M}\mathrm{tr}\left(f(\hat{\mathbf{C}}(\nu)) - \int f d\mu_{MP}^{(c_{N})}$. In particular, establishing (4.41) (and thus (4.42)) will complete the proof of Theorem 4.1.

Proof. The proof of (4.41) is based on the Gaussian tools reviewed in Paragraph 2.2.2, and needs long and very tedious calculations. Therefore, we just provide a sketch of proof. In particular, we justify that $\mathbb{E}\left[\frac{1}{M} \operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z) - \mathbf{Q}_{N}(z)\right)\right]$ is a $\mathcal{O}_{z}\left(\frac{B}{N}\right)^{2}$ term, but do not establish its expression (4.41).

The starting point of the proof is to express $\tilde{\mathbf{Q}} - \mathbf{Q}$ as

$$\dot{\mathbf{Q}} - \mathbf{Q} = -\dot{\mathbf{Q}}\dot{\Delta}\mathbf{Q} = -\mathbf{Q}\dot{\Delta}\mathbf{Q} + \mathbf{Q}\dot{\Delta}\mathbf{Q}\dot{\Delta}\mathbf{Q} - \dot{\mathbf{Q}}\,\,\dot{\Delta}\mathbf{Q}\dot{\Delta}\mathbf{Q}\dot{\Delta}\mathbf{Q}.$$

Therefore, $\mathbb{E}\left[\frac{1}{M} \text{tr} \left(\tilde{\mathbf{Q}}_N(z) - \mathbf{Q}_N(z)\right)\right]$ can be written as

$$\mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\tilde{\mathbf{Q}}-\mathbf{Q}\right)\right] = -\mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\mathbf{Q}^{2}\tilde{\boldsymbol{\Delta}}\right)\right] + \mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\mathbf{Q}^{2}\tilde{\boldsymbol{\Delta}}\mathbf{Q}\tilde{\boldsymbol{\Delta}}\right)\right] \\ - \mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\tilde{\mathbf{Q}}\tilde{\boldsymbol{\Delta}}\mathbf{Q}\tilde{\boldsymbol{\Delta}}\mathbf{Q}\tilde{\boldsymbol{\Delta}}\right)\right]. \quad (4.43)$$

It is clear that the moduli of the second and third terms of the right hand side of (4.43) are controlled by $C(z)\mathbb{E}(\|\tilde{\Delta}\|^2)$ and $C(z)\mathbb{E}(\|\tilde{\Delta}\|^3)$ respectively. We now state the following useful Lemma, proved in the Appendix, which implies that these terms are $\mathcal{O}_z\left(\frac{B}{N}\right)^2$ and $\mathcal{O}_z\left(\frac{B}{N}\right)^3$ respectively.

Lemma 4.5. For each $k \ge 1$, it exist a nice constant C depending on k such that $\mathbb{E}\left(\|\tilde{\mathbf{\Delta}}\|^k\right) \le C\left(\frac{B}{N}\right)^k$

In order to prove that $\mathbb{E}\left[\frac{1}{M}\operatorname{tr}\left(\tilde{\mathbf{Q}}_{N}(z)-\mathbf{Q}_{N}(z)\right)\right]=\mathcal{O}_{z}\left(\frac{B}{N}\right)^{2}$, we thus have to check that

$$\mathbb{E}\left[\frac{1}{M} \operatorname{tr} \mathbf{Q}^{2} \tilde{\boldsymbol{\Delta}}\right] = \mathcal{O}_{z} \left(\frac{B}{N}\right)^{2}.$$
(4.44)

For this, we first express $\mathbb{E}\left[\frac{1}{M} \operatorname{tr} \mathbf{Q}^2 \tilde{\mathbf{\Delta}}\right]$ as

$$\mathbb{E}\left[\frac{1}{M}\mathrm{tr} \ \mathbf{Q}^{2}\tilde{\boldsymbol{\Delta}}\right] = \mathbb{E}\left(\frac{1}{M}\mathrm{tr} \ \mathbf{Q}^{2}\frac{\boldsymbol{\Gamma}\mathbf{X}^{*}}{B+1}\right) + \mathbb{E}\left(\frac{1}{M}\mathrm{tr} \ \mathbf{Q}^{2}\frac{\mathbf{X}\boldsymbol{\Gamma}^{*}}{B+1}\right) \\ + \mathbb{E}\left(\frac{1}{M}\mathrm{tr} \ \mathbf{Q}^{2}\frac{\boldsymbol{\Gamma}\boldsymbol{\Gamma}^{*}}{B+1}\right).$$

The third term of the right hand side is clearly $\mathcal{O}_z((\frac{B}{N})^2)$. We thus need to check that the first two terms are also $\mathcal{O}_z((\frac{B}{N})^2)$. We just verify this property for the first term. For this, we evaluate $\mathbb{E}\left(\frac{1}{M} \operatorname{tr} \mathbf{Q} \frac{\mathbf{\Gamma} \mathbf{X}^*}{B+1}\right)$ using the Gaussian tools, and take the derivative w.r.t. z to obtain the expression of $\mathbb{E}\left(\frac{1}{M} \operatorname{tr} \mathbf{Q}^2 \frac{\mathbf{\Gamma} \mathbf{X}^*}{B+1}\right)$.

In order to simplify the notations, we denote by \mathbf{W} the matrix $\mathbf{W} = \frac{\mathbf{X}}{\sqrt{B+1}}$, and denote by $\mathbf{w}_1 = \frac{\mathbf{x}_1}{\sqrt{B+1}}, \dots, \mathbf{w}_M = \frac{\mathbf{x}_M}{\sqrt{B+1}}$ its M rows. In particular, the row m of matrix $\frac{\mathbf{\Gamma}}{\sqrt{B+1}}$ coincides with $\mathbf{w}_m \Psi_m$ where we recall that matrix Ψ_m is defined by (3.12). If $(\mathbf{e}_1, \dots, \mathbf{e}_m)$ represents the canonical basis of \mathbb{C}^M , $\mathbb{E}\left(\frac{1}{M} \operatorname{tr} \mathbf{Q} \frac{\mathbf{\Gamma} \mathbf{X}^*}{B+1}\right)$ can be written as

$$\mathbb{E}\left(\frac{1}{M} \operatorname{tr} \, \mathbf{Q} \frac{\mathbf{\Gamma} \mathbf{X}^*}{B+1}\right) = \frac{1}{M} \sum_{m=1}^M \mathbb{E}\left(\mathbf{w}_m \boldsymbol{\Psi}_m \mathbf{W}^* \mathbf{Q} \mathbf{e}_m\right).$$

We now state the following Lemma whose proof is given in Appendix. We recall that $\beta_N(z) = \mathbb{E}((\mathbf{Q}_N(z))_{mm} \text{ for each } m.$

Lemma 4.6. If **A** represents a $(B+1) \times (B+1)$ matrix, the following equality holds

$$\mathbb{E}\left(\mathbf{w}_{m}\mathbf{A}\mathbf{W}^{*}\mathbf{Q}\mathbf{e}_{m}\right) = \frac{\beta}{1+\beta c}\frac{1}{B+1}\operatorname{tr}\mathbf{A} - \mathbb{E}\left[\left(\frac{1}{B+1}\operatorname{tr}\mathbf{W}\mathbf{A}\mathbf{W}^{*}\mathbf{Q}\right)^{\circ}\mathbf{Q}_{m,m}^{\circ}\right] + \frac{\beta c}{1+\beta c}\mathbb{E}\left[\left(\frac{1}{B+1}\operatorname{tr}\mathbf{W}\mathbf{A}\mathbf{W}^{*}\mathbf{Q}\right)^{\circ}\frac{1}{B+1}\operatorname{tr}\mathbf{Q}^{\circ}\right].$$
 (4.45)

Using (1.23) in the case $\mathbf{s}_{\mu}(z) = \beta(z)$ as well as (2.9), we easily obtain that $\frac{\beta}{1+\beta c} = \frac{t}{1+ct} + \epsilon_1(z) = -zt(z)\tilde{t}(z) + \epsilon_1(z)$ where $\epsilon_1(z) = \mathcal{O}_z(\frac{1}{B^2})$. Moreover, it follows from (2.13) that $\epsilon'_1(z)$ is also a $\mathcal{O}_z(\frac{1}{B^2})$. We now use (4.45) for $\mathbf{A} = \Psi_m$, and differentiate (4.45) for $\mathbf{A} = \Psi_m$ w.r.t. z. Using the Schwartz inequality the inequalities (2.6) and (2.7), and (3.13), we obtain immediately that

$$\mathbb{E}\left(\mathbf{w}_{m}\boldsymbol{\Psi}_{m}\mathbf{W}^{*}\mathbf{Q}\mathbf{e}_{m}\right) = -\left(zt(z)\tilde{t}(z)\right)'\frac{1}{B+1}\operatorname{tr}\boldsymbol{\Psi}_{m} + \mathcal{O}_{z}\left(\frac{B}{N}\right)^{3} + \mathcal{O}_{z}\left(\frac{1}{\sqrt{B}N}\right)$$

and that

$$\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{Q}^{2}\frac{\mathbf{\Gamma}\mathbf{X}^{*}}{B+1}\right) = -\left(zt(z)\tilde{t}(z)\right)'\frac{1}{B+1}\operatorname{tr}\left(\frac{1}{M}\sum_{m=1}^{M}\boldsymbol{\Psi}_{m}\right) + \mathcal{O}_{z}\left(\frac{B}{N}\right)^{3} + \mathcal{O}_{z}\left(\frac{1}{\sqrt{B}N}\right).$$

It is easily checked that

$$\Psi_m = (\mathbf{I} + \Phi_m)^{1/2} - \mathbf{I} = \frac{1}{2} \Phi_m - \frac{1}{8} \Phi_m^2 + \boldsymbol{\Xi}_m$$

where $\|\mathbf{\Xi}_m\| \leq C\left(\frac{B}{N}\right)^3$. It is easily seen that $\frac{1}{B+1} \operatorname{tr} \mathbf{\Phi}_m^2 = \left(\frac{s'_m}{s_m}\right)^2 v_N + \mathcal{O}((\frac{B}{N})^3 + \frac{1}{N})$. Using (3.9), we thus obtain that

$$\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{Q}^{2}\frac{\mathbf{\Gamma}\mathbf{X}^{*}}{B+1}\right) = -(zt_{N}(z)\tilde{t}_{N}(z))'\left(\frac{1}{M}\sum_{m=1}^{M}\left(\frac{s_{m}''}{2s_{m}} - \frac{(s_{m}')^{2}}{8(s_{m})^{2}}\right)\right)v_{N} + \mathcal{O}_{z}(u_{N})$$

because

$$\mathcal{O}_z\left(\frac{B}{N}\right)^3 + \mathcal{O}_z\left(\frac{1}{\sqrt{BN}}\right) + \mathcal{O}_z\left(\frac{1}{N}\right) = \mathcal{O}_z(u_N).$$

We have thus established that $\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{Q}^{2}\frac{\mathbf{\Gamma}\mathbf{X}^{*}}{B+1}\right)$ is a $\mathcal{O}_{z}\left(\frac{B}{N}\right)^{2}$ term, and have evaluated the corresponding principal term. Using similar calculations, we can obtain easily the expression of the $\mathcal{O}_{z}\left(\frac{B}{N}\right)^{2}$ term of $\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{Q}^{2}\tilde{\boldsymbol{\Delta}}\right)$. In order to establish (4.42), it is necessary to evaluate the $\mathcal{O}_{z}\left(\frac{B}{N}\right)^{2}$ term of $\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{Q}\tilde{\boldsymbol{\Delta}}\mathbf{Q}\tilde{\boldsymbol{\Delta}}\mathbf{Q}\right)$. This step needs very long calculations that are omitted.

4.5. Estimation of $r_N(\nu)$

The term $\sup_{\nu} |\psi_N(f,\nu)|$ depends on the unknown true spectral densities $(s_m)_{m=1,...,M}$ through the term $r_N(\nu)$ defined by (1.8). In order to be able to use Theorem 4.1 in practice, it appears necessary to estimate $r_N(\nu)$ by an accurate enough estimate $\hat{r}_N(\nu)$, and to replace $\psi_N(f,\nu)$ by $\hat{\psi}_N(f,\nu)$ defined by

$$\hat{\psi}_N(f,\nu) = \frac{1}{M} \operatorname{Tr} \left(f(\hat{\mathbf{C}}(\nu)) \right) - \int_{\mathbb{R}^+} f \, \mathrm{d}\mu_{MP}^{(c_N)} - \hat{r}_N(\nu) \, \phi_N(f) \, v_N \, \mathbf{1}_{\alpha > 2/3} \quad (4.46)$$

 $\hat{r}_N(\nu)$ has to be chosen in such a way that $|\hat{\psi}_N(f,\nu)| \prec u_N$, a condition that will be verified if $|\hat{r}_N(\nu) - r_N(\nu)| \prec \frac{u_N}{v_N}$ if $\alpha > \frac{2}{3}$. A natural choice for $\hat{r}_N(\nu)$ would be to replace the true spectral densities $(s_m)_{m=1,\dots,M}$ by their frequency smoothed estimates $(\hat{s}_m)_{m=1,\dots,M}$ defined by (3.31), and the derivatives $(s'_m)_{m=1,\dots,M}$ by $(\hat{s}'_m)_{m=1,\dots,M}$. However, \hat{s}'_m is not an accurate estimate of s'_m so that the corresponding estimate of $r_N(\nu)$ does not verify $|\hat{r}_N(\nu) - r_N(\nu)| \prec \frac{u_N}{v_N}$ if $\alpha > \frac{2}{3}$. If L < N is an integer, we introduce the lag window estimator $\hat{s}_{m,L}$ of s_m defined by

$$\hat{s}_{m,L}(\nu) = \int_0^1 |\xi_{\mathbf{y}_m}(\mu)|^2 w_L(\nu - \mu) d\mu = \sum_{l=-L}^L \hat{r}_{m,l} e^{-2i\pi l\nu}$$
(4.47)

where $w_L(\nu) = \sum_{l=-L}^{L} e^{-2i\pi l\nu}$ is the Fourier transform of the rectangular window and $\hat{r}_{m,l}$ represents the biased estimate of the autocovariance coefficient $r_{m,l}$ of y_m at lag l defined by

$$\hat{r}_{m,l} = \frac{1}{N} \sum_{n=1}^{N-l} y_{m,n+l} y_{m,n}^*$$
(4.48)

and $\hat{r}_{m,-l} = \hat{r}_{m,l}^*$ for $l \ge 0$. Then, the following result holds.

Proposition 4.3. Assume that $L = L(N) = \mathcal{O}(N^{\frac{1}{2\gamma_0+1}})$, where $\gamma_0 \geq 3$ is defined by (1.18). Then, the estimate $\hat{r}_N(\nu)$ defined by

$$\hat{r}_N(\nu) = \left(\frac{1}{M} \sum_{m=1}^M \frac{\hat{s}'_{m,L}(\nu)}{\hat{s}_{m,L}(\nu)}\right)^2$$
(4.49)

verifies

$$|\hat{r}_N(\nu) - r_N(\nu)| \prec \frac{1}{N^{(\gamma_0 - 1)/(2\gamma_0 + 1)}}$$
(4.50)

$$|\hat{r}_N(\nu) - r_N(\nu)| \quad \prec \quad \frac{u_N}{v_N} \text{ if } \alpha > \frac{2}{3}$$

$$(4.51)$$

 $as \ well \ as$

$$|\hat{\psi}_N(f,\nu)| \prec u_N. \tag{4.52}$$

Proof. We denote by $\mathbf{d}_N(\nu)$ the *N*-dimensional vector defined by $\mathbf{d}_N(\nu) = (1, e^{-2i\pi\nu}, \ldots, e^{-2i\pi(N-1)\nu})^T$. We recall that \mathbf{y}_m is the *N*-dimensional vector $\mathbf{y}_m = (y_{m,1}, \ldots, y_{m,N})^T$. \mathbf{y}_m can be written as $\mathbf{y}_m = \mathbf{R}_m^{1/2} \mathbf{z}_m$ where $\mathbf{R}_m = \mathbb{E}(\mathbf{y}_m \mathbf{y}_m^*)$ and \mathbf{z}_m is $\mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_N)$ distributed. It is clear that $\hat{s}_{m,L}(\nu)$ can be written as

$$\hat{s}_{m,L}(\nu) = \mathbf{z}_m^* \mathbf{R}_m^{1/2} \mathbf{\Omega}(\nu) \mathbf{R}_m^{1/2} \mathbf{z}_m$$

with

$$\mathbf{\Omega}(\nu) = \frac{1}{N} \int \mathbf{d}_N(\mu) \mathbf{d}_N(\mu)^* w_L(\nu - \mu) d\mu$$

while $\hat{s}'_{m,L}(\nu)$ is equal to

$$\hat{s}_{m,L}'(\nu) = \mathbf{z}_m^* \mathbf{R}_m^{1/2} \mathbf{\Omega}'(\nu) \mathbf{R}_m^{1/2} \mathbf{z}_m$$

with

$$\mathbf{\Omega}'(\nu) = \frac{-2i\pi}{N} \int \mathbf{d}_N(\mu) \mathbf{d}_N(\mu)^* \left(\sum_{l=-L}^L l \, e^{-2i\pi l(\nu-\mu)}\right) \, d\mu$$

It is easy to check that $\|\mathbf{\Omega}'(\nu)\|_F = \mathcal{O}(\frac{L^{3/2}}{N^{1/2}})$ and therefore that $\|\mathbf{R}_m^{1/2}\mathbf{\Omega}'(\nu)\mathbf{R}_m^{1/2}\|_F = \mathcal{O}(\frac{L^{3/2}}{N^{1/2}})$. The Hanson-Wright inequality leads immediately to $|\hat{s}'_{m,L}(\nu) - \mathbb{E}(\hat{s}'_{m,L}(\nu))| \prec \frac{L^{3/2}}{N^{1/2}}$. Moreover, it is easy to check that (1.18) implies that

$$|\mathbb{E}(\hat{s}'_{m,L}(\nu)) - s'_m(\nu)| \le \frac{C}{L^{\gamma_0 - 1}}$$

where C is a nice constant. For L = L(N) in such a way that $\mathcal{O}(\frac{L^{3/2}}{N^{1/2}}) = \frac{1}{L^{\gamma_0-1}}$, i.e. $L = \mathcal{O}(N^{\frac{1}{2\gamma_0+1}})$, we obtain that

$$|\hat{s}'_{m,L}(\nu) - s'_m(\nu)| \prec \frac{1}{N^{(\gamma_0 - 1)/(2\gamma_0 + 1)}}.$$

Moreover, a similar analysis leads to

$$|\hat{s}_{m,L}(\nu) - s_m(\nu)| \prec \frac{1}{N^{\gamma_0/(2\gamma_0+1)}}$$

from which we deduce that the estimate $\hat{r}_N(\nu)$ defined by (4.49) verifies (4.50). It is then easily checked that if $\gamma_0 \geq 3$, then (4.51) holds, which implies that $|\hat{\psi}_N(f,\nu)| \prec u_N$ holds.

5. Use of Lipschitz properties of functions $\nu \to \psi_N(f,\nu)$ and $\nu \to \hat{\psi}_N(f,\nu)$

In this section, we establish Lipschitz properties of $\nu \to \psi_N(f,\nu)$ and $\nu \to \hat{\psi}_N(f,\nu)$, and deduce that the stochastic domination properties (4.2) and (4.52) are still valid for $\sup_{\nu \in [0,1]} |\psi_N(f,\nu)|$ and $\sup_{\nu \in [0,1]} |\hat{\psi}_N(f,\nu)|$ where $\hat{\psi}_N(f,\nu)$ is defined by (4.46, 4.49).

5.1. Lipschitz properties

The goal of this paragraph is to prove the following Proposition.

Proposition 5.1. Functions $\nu \to \psi_N(f,\nu)$ and $\nu \to \hat{\psi}_N(f,\nu)$ verify

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{\|\psi_N(f,\nu) - \psi_N(f,\nu+\delta)\|}{|\delta|} \prec M N^{3/2}$$
(5.1)

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{\|\hat{\psi}_N(f,\nu) - \hat{\psi}_N(f,\nu+\delta)\|}{|\delta|} \prec MN^{3/2}$$
(5.2)

In the following, we just establish (5.2). For this, we evaluate separately the Lipschitz constants of $\nu \to \frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu))$ and of $\nu \to \hat{r}_N(\nu)$.

5.1.1. Lipschitz constant of $\nu \to \frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu))$

To show that $\nu \to \frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu))$ is $MN^{3/2}$ -Lipschitz with overwhelming probability, we need to establish a number of intermediate properties.

Proposition 5.2. It holds that

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{\|\widehat{\mathbf{S}}(\nu) - \widehat{\mathbf{S}}(\nu + \delta)\|}{|\delta|} \prec MN^{3/2}.$$
(5.3)

Proof. Let $\delta \in \mathbb{R}$ and $\nu \in [0, 1]$. As the random variables $(y_{m,n})_{m=1,\ldots,M,n=1,\ldots,N}$ are complex Gaussian and that $\sup_{m\geq 1} \mathbb{E}|y_{m,n}|^2 < +\infty$, the family $(y_{m,n})_{m=1,\ldots,N,n=1,\ldots,N}$ verifies $|y_{m,n}| \prec 1$. Therefore, it holds that

$$\frac{1}{\sqrt{N}}\sum_{n=1}^{N}|y_{m,n}| \prec \sqrt{N}.$$
(5.4)

For the same reasons, the family $\xi_{y_m}(\nu), m = 1, \dots, M, \nu \in [0, 1]$ satisfies.

$$\xi_{y_m}(\nu)| \prec 1. \tag{5.5}$$

We also claim that

$$\sup_{\nu \in [0,1]} |\xi_{y_m}(\nu)| \prec 1.$$
(5.6)

In order to verify (5.6), we first observe that for any $n \ge 1$, we have the following control:

 $|e^{-2i\pi n\nu} - e^{-2i\pi n(\nu+\delta)}| \le 2|\sin \pi n\delta| \le 2\pi n|\delta|.$

(5.4) implies that

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \left| \frac{\xi_{y_m}(\nu) - \xi_{y_m}(\nu + \delta)}{\delta} \right|$$

$$= \sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{1}{\sqrt{N}} \left| \sum_{n=1}^{N} y_{m,n} \frac{e^{-2i\pi n\nu} - e^{-2i\pi n(\nu + \delta)}}{\delta} \right|$$

$$\leq 2\pi N \frac{1}{\sqrt{N}} \sum_{n=1}^{N} |y_{m,n}|$$

$$\prec N^{3/2}. \tag{5.7}$$

We consider a frequency $\nu_* \in [0, 1]$ (depending on *m*) where $|\xi_{y_m}(\nu)|$ is maximum, and have thus to establish that for each $\epsilon > 0$, then it exists $\gamma > 0$ depending only on ϵ such that

$$\mathbb{P}(|\xi_{y_m}(\nu_*)| > N^{\epsilon}) \le \exp{-N^{\gamma}}$$

for each N larger than a certain integer $N_0(\epsilon)$. We introduce the discrete the set

$$\mathcal{V}_{N}^{p} = \left\{ \frac{k}{N^{p}} : k \in \{0, \dots, N^{p} - 1\} \right\}$$
(5.8)

which cardinal is $|\mathcal{V}_N^p| = N^p$. We notice that (5.5) in conjunction with the union bound implies that $\sup_{\nu_p \in \mathcal{V}_N^p} |\xi_{y_m}(\nu_p)| \prec 1$. We denote by $\nu_{*,p}$ the element of \mathcal{V}_N^p for which $|\nu_* - \nu_p|$ is minimum, and notice that $|\nu_* - \nu_{*,p}| \leq \frac{1}{N^p}$. Then, we have the following inequality

$$\mathbb{P}(|\xi_{y_m}(\nu_*)| > N^{\epsilon}) \\
\leq \mathbb{P}\left(|\xi_{y_m}(\nu_*) - \xi_{y_m}(\nu_{*,p})| > \frac{N^{\epsilon}}{2}\right) + \mathbb{P}\left(|\xi_{y_m}(\nu_{*,p})| > \frac{N^{\epsilon}}{2}\right) \\
\leq \mathbb{P}\left(|\xi_{y_m}(\nu_*) - \xi_{y_m}(\nu_{*,p})| > \frac{N^{\epsilon}}{2}\right) + \mathbb{P}\left(\sup_{\nu_p \in \mathcal{V}_N^p} |\xi_{y_m}(\nu_p)| > \frac{N^{\epsilon}}{2}\right).$$
(5.9)

As $\sup_{\nu_p \in \mathcal{V}_N^p} |\xi_{y_m}(\nu_p)| \prec 1$, the second term of the right hand side of (5.9) converges exponentially towards 0. In order to evaluate the first term of the r.h.s. of (5.9), we use (5.7), and obtain that

$$\mathbb{P}\left(\left|\xi_{y_m}(\nu_*) - \xi_{y_m}(\nu_{*,p})\right| > \frac{N^{\epsilon}}{2}\right) \le \mathbb{P}\left(N\frac{1}{\sqrt{N}}\sum_{n=1}^{N}|y_{m,n}| \ge \frac{\pi}{2|\nu_* - \nu_{*,p}|}N^{\epsilon}\right)$$
$$\le \mathbb{P}\left(\frac{1}{\sqrt{N}}\sum_{n=1}^{N}|y_{m,n}| \ge \frac{\pi}{2}N^{p+\epsilon-1}\right).$$

We choose p so that p-1 > 3/2, and use (5.4) to conclude that $\mathbb{P}\left(|\xi_{y_m}(\nu_*) - \xi_{y_m}(\nu_{*,p})| > \frac{N^{\epsilon}}{2}\right)$ converges towards 0 exponentially. This

establishes (5.6).

In order to complete the proof of Proposition 5.2, we consider an individual entry $\hat{s}_{ij}(\nu)$ of $\hat{\mathbf{S}}(\nu)$ for $i, j \leq M$, and write that

$$\begin{aligned} |\hat{s}_{ij}(\nu) - \hat{s}_{ij}(\nu + \delta)| \\ &= \frac{1}{B+1} \left| \sum_{b=-B/2}^{B/2} \xi_i \left(\nu + \frac{b}{N}\right) \xi_j \left(\nu + \frac{b}{N}\right)^* \right| \\ &- \xi_i \left(\nu + \delta + \frac{b}{N}\right) \xi_j \left(\nu + \delta + \frac{b}{N}\right)^* \right| \\ &\leq \frac{1}{B+1} \sum_{b=-B/2}^{B/2} \left| \xi_i \left(\nu + \frac{b}{N}\right) \left(\xi_j \left(\nu + \frac{b}{N}\right)^* - \xi_j \left(\nu + \delta + \frac{b}{N}\right)^* \right) \right| \\ &+ \left| \left(\xi_i \left(\nu + \frac{b}{N}\right) - \xi_i \left(\nu + \delta + \frac{b}{N}\right) \right) \xi_j \left(\nu + \delta + \frac{b}{N}\right)^* \right|. \end{aligned}$$

Using the estimations (5.6) and (5.7), we get:

$$\sup_{i,j} \sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \left| \frac{\hat{s}_{ij}(\nu) - \hat{s}_{ij}(\nu + \delta)}{\delta} \right| \prec N^{3/2}$$
(5.10)

and deduce (5.3) from the rough bound

$$\sup_{\nu \in [0,1]} \| \hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu + \delta) \| \leq \sup_{\nu \in [0,1]} \sup_{i} \sum_{j} |\hat{s}_{ij}(\nu) - \hat{s}_{ij}(\nu + \delta)|$$
$$\leq M \sup_{\nu \in [0,1]} \sup_{i,j} |\hat{s}_{ij}(\nu) - \hat{s}_{ij}(\nu + \delta)|.$$

Combining the eigenvalue localisation result from Corollary 3.3 and the Lipschitz behaviour of $\hat{\mathbf{S}}$ from Proposition 5.2, the following statement holds.

Corollary 5.1. (ν uniform version of Corollary 3.3.) Denote for $\epsilon > 0$:

$$\begin{split} \Lambda^{\hat{\mathbf{S}}}_{\epsilon} &= \left\{ \forall \nu \in [0,1] : \sigma(\hat{\mathbf{S}}(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}] + \epsilon \right\} \\ \Lambda^{\hat{\mathbf{D}}}_{\epsilon} &= \left\{ \forall \nu \in [0,1] : \sigma(\hat{\mathbf{D}}(\nu)) \subset [\underline{s}, \overline{s}] + \epsilon \right\}. \end{split}$$

Then, $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}$ and $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}$ hold with exponentially high probability.

Proof. As the proof for $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}$ is strictly similar to the one of $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}$, we will only write the arguments for $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}$. For any fixed $\nu \in [0, 1]$, Corollary 3.3 ensures that $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}(\nu)$ holds with exponentially high probability. For $p \geq 1$, we still consider the set \mathcal{V}_{N}^{p} defined by (5.8) and denote by $\Lambda_{\epsilon,p}^{\hat{\mathbf{S}}}$ the event defined by

$$\Lambda_{\epsilon,p}^{\hat{\mathbf{S}}} = \left\{ \forall \nu_p \in \mathcal{V}_N^p : \sigma(\hat{\mathbf{S}}(\nu_p)) \subset \operatorname{Supp} \mu_{MP}^{(c)} \times [\underline{s}, \overline{s}] + \epsilon \right\}$$

which is $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}$ but where ν runs only on the finite grid \mathcal{V}_{N}^{p} . It is immediate (by the union bound) that $\Lambda_{\epsilon,p}^{\hat{\mathbf{S}}}$ holds with exponentially high probability for any fixed $p \in \mathbb{N}$. Moreover, it is clear from the definitions of $\Lambda_{\epsilon}^{\hat{\mathbf{S}}}$ and $\Lambda_{\epsilon,p}^{\hat{\mathbf{S}}}$ that $\Lambda_{\epsilon}^{\hat{\mathbf{S}}} \subset \Lambda_{\epsilon,p}^{\hat{\mathbf{S}}}$. We now show the following inclusion:

$$\left(\Lambda_{\epsilon}^{\hat{\mathbf{S}}}\right)^{c} \subset \left(\Lambda_{\epsilon/2,p}^{\hat{\mathbf{S}}}\right)^{c} \\ \cup \left\{ \exists \nu \in [0,1] : \|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu_{p}^{*})\| > \epsilon/2 \text{ where } \nu_{p}^{*} \in \operatorname*{argmin}_{\nu_{p} \in \mathcal{V}_{N}^{p}} |\nu - \nu_{p}| \right\}.$$

$$(5.11)$$

Suppose that $(\Lambda_{\epsilon}^{\hat{\mathbf{S}}})^c$ is realized, and denote by $\nu^* \in [0,1]$ a frequency such that $\sigma(\hat{\mathbf{S}})(\nu^*) \not\subset \operatorname{Supp}_{MP}^{(c)} \times [\underline{s}, \overline{s}] + \epsilon$. Denote also $\nu_p^* \in \operatorname{argmin}_{\nu_p \in \mathcal{V}_N^p} |\nu_p - \nu^*|$. We just consider the case where $\lambda_1(\hat{\mathbf{S}}(\nu^*)) > \overline{s}(1 + \sqrt{c})^2 + \epsilon$, since in the case where $\lambda_M(\hat{\mathbf{S}}(\nu^*)) < \underline{s}(1 - \sqrt{c})^2 - \epsilon$, the proof is similar. Then, either:

1. $\|\hat{\mathbf{S}}(\nu_p^*) - \hat{\mathbf{S}}(\nu^*)\| \leq \epsilon/2$, which implies the following estimation for the location of $\lambda_1(\hat{\mathbf{S}}(\nu_p^*))$:

$$\lambda_1(\hat{\mathbf{S}}(\nu^*)) - \frac{\epsilon}{2} \le \lambda_1(\hat{\mathbf{S}}(\nu_p^*)) \le \lambda_1(\hat{\mathbf{S}}(\nu^*)) + \frac{\epsilon}{2}$$

and in particular, $\lambda_1(\hat{\mathbf{S}}(\nu_p^*)) \geq \bar{s}(1+\sqrt{c})^2 + \epsilon/2$. This means that $\left(\Lambda_{\epsilon/2,p}^{\hat{\mathbf{S}}}\right)^c$ holds.

2. $\|\hat{\mathbf{S}}(\nu_p^*) - \hat{\mathbf{S}}(\nu^*)\| > \epsilon/2$, which exactly means that $\left\{ \exists \nu \in [0,1] : \|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu_p^*)\| > \epsilon/2$ where $\nu_p^* \in \operatorname{argmin}_{\nu_p \in \mathcal{V}_N^p} |\nu - \nu_p| \right\}$ is realized

(5.11) is now proved.

We already showed that $\left(\Lambda_{\epsilon/2,p}^{\hat{\mathbf{S}}}\right)^c$ holds with exponentially small probability, and establish now that the set

$$\left\{ \exists \nu \in [0,1] : \| \hat{\mathbf{S}}(\nu_p^*) - \hat{\mathbf{S}}(\nu) \| > \epsilon/2 \text{ where } \nu_p^* \in \operatorname*{argmin}_{\nu_p \in \mathcal{V}_N^p} |\nu - \nu_p| \right\}$$

has the same property. To justify this claim, we remark that Proposition 5.2 implies that for each $\kappa > 0$, the probability

$$\mathbb{P}\left[\left\{\exists \nu, \nu' \in [0,1], \ \frac{\|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu')\|}{|\nu - \nu'|} > N^{\kappa} M N^{3/2}\right\}\right]$$

converges to 0 exponentially fast. As the following inclusion

$$\left\{ \exists \nu \in [0,1], \ \frac{\|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu_p^*)\|}{|\nu - \nu_p^*|}, > N^{\kappa} M N^{3/2}, \text{ where } \nu_p^* \in \underset{\nu_p \in \mathcal{V}_N^p}{\operatorname{argmin}} |\nu - \nu_p| \right\} \\ \subset \left\{ \exists \nu, \nu' \in [0,1], \ \frac{\|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu')\|}{|\nu - \nu'|} > N^{\kappa} M N^{3/2} \right\}$$

holds, we get that

$$\mathbb{P}\left[\left\{\exists \nu \in [0,1], \|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu_p^*)\| > |\nu - \nu_p^*| N^{\kappa} M N^{3/2}\right\}\right] \to 0$$

exponentially fast. Moreover, as for each ν , $|\nu - \nu_p^*| \leq \frac{1}{N^p}$, we obtain that

$$\mathbb{P}\left[\left\{\exists \nu \in [0,1], \|\hat{\mathbf{S}}(\nu) - \hat{\mathbf{S}}(\nu_p^*)\| > \frac{1}{N^p} N^{\kappa} M N^{3/2}\right\}\right] \to 0$$

exponentially fast as well. For p large enough, $N^{\kappa} \frac{1}{N^p} M N^{3/2}$ will eventually become smaller than $\epsilon/2$. This proves that

$$\left\{ \exists \nu \in [0,1], \ \| \hat{\mathbf{S}}(\nu_p^*) - \hat{\mathbf{S}}(\nu) \| > \epsilon/2 \text{ where } \nu_p^* \in \operatorname*{argmin}_{\nu_p \in \mathcal{V}_N^p} |\nu - \nu_p| \right\}$$

holds with exponentially small probability.

The same argument can be used in order to control $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}$. This completes the proof of Corollary 5.1.

We deduce immediately from Corollary 5.1 the following result that can be seen as a refinement of (3.28) and of Lemma 3.1.

Corollary 5.2. It holds that

$$\sup_{\nu \in [0,1]} \| \hat{\mathbf{D}}(\nu)^{-1/2} \| \prec 1, \quad \sup_{\nu \in [0,1]} \| \hat{\mathbf{S}}(\nu) \| \prec 1.$$

A useful consequence of this is the following Corollary, which states that the Lipschitz result holds for $\hat{\mathbf{C}}(\nu)$.

Corollary 5.3. It holds that

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \left\| \frac{\hat{\mathbf{C}}(\nu) - \hat{\mathbf{C}}(\nu + \delta)}{\delta} \right\| \prec MN^{3/2}$$
(5.12)

Proof. For more clarity in the following argument, denote $\nu_1 = \nu$ and $\nu_2 = \nu + \delta$. Recall that $\hat{\mathbf{D}} = \text{diag}\hat{\mathbf{S}}$. Using the definition of $\hat{\mathbf{C}}$ from equation (1.3), we write:

$$\begin{split} \hat{\mathbf{C}}(\nu_2) - \hat{\mathbf{C}}(\nu_1) &= \hat{\mathbf{D}}^{-1/2}(\nu_2)\hat{\mathbf{S}}(\nu_2)\hat{\mathbf{D}}^{-1/2}(\nu_2) - \hat{\mathbf{D}}^{-1/2}(\nu_1)\hat{\mathbf{S}}(\nu_1)\hat{\mathbf{D}}^{-1/2}(\nu_1) \\ &= (\hat{\mathbf{D}}^{-1/2}(\nu_2) - \hat{\mathbf{D}}^{-1/2}(\nu_1))\hat{\mathbf{S}}(\nu_2)\hat{\mathbf{D}}^{-1/2}(\nu_2) \\ &+ \hat{\mathbf{D}}^{-1/2}(\nu_1)(\hat{\mathbf{S}}(\nu_2)\hat{\mathbf{D}}^{-1/2}(\nu_2) - \hat{\mathbf{S}}(\nu_1)\hat{\mathbf{D}}^{-1/2}(\nu_1)) \end{split}$$

Moreover, we write that

$$\begin{split} \hat{\mathbf{S}}(\nu_2) \hat{\mathbf{D}}^{-1/2}(\nu_2) &- \hat{\mathbf{S}}(\nu_1) \hat{\mathbf{D}}^{-1/2}(\nu_1) \\ &= (\hat{\mathbf{S}}(\nu_2) - \hat{\mathbf{S}}(\nu_1)) \hat{\mathbf{D}}^{-1/2}(\nu_2) + \hat{\mathbf{S}}(\nu_1) (\hat{\mathbf{D}}^{-1/2}(\nu_2) - \hat{\mathbf{D}}^{-1/2}(\nu_1)). \end{split}$$

Therefore, applying the operator norm, we get by the triangle inequality:

$$\begin{split} \|\hat{\mathbf{C}}(\nu_{2}) - \hat{\mathbf{C}}(\nu_{1})\| &\leq \|\hat{\mathbf{D}}^{-1/2}(\nu_{2}) - \hat{\mathbf{D}}^{-1/2}(\nu_{1})\| \|\hat{\mathbf{S}}(\nu_{2})\| \|\hat{\mathbf{D}}^{-1/2}(\nu_{2})\| \\ &+ \|\hat{\mathbf{D}}^{-1/2}(\nu_{1})\| \|\hat{\mathbf{S}}(\nu_{2}) - \hat{\mathbf{S}}(\nu_{1})\| \|\hat{\mathbf{D}}^{-1/2}(\nu_{2})\| \\ &+ \|\hat{\mathbf{D}}^{-1/2}(\nu_{1})\| \|\hat{\mathbf{S}}(\nu_{1})\| \|\hat{\mathbf{D}}^{-1/2}(\nu_{2}) - \hat{\mathbf{D}}^{-1/2}(\nu_{1})\| \end{split}$$

It is easy to check that

$$\sup_{\delta \neq 0} \sup_{|\nu_2 - \nu_1| = \delta} \left\| \frac{\hat{\mathbf{D}}^{-1/2}(\nu_2) - \hat{\mathbf{D}}^{-1/2}(\nu_1)}{\delta} \right\| \prec N^{3/2}$$

holds. Therefore, Proposition 5.2 and Corollary 5.2 immediately imply (5.12). $\hfill \Box$

Finally, we can write for the spectrum of $\hat{\mathbf{C}}$ the same kind of result as in Corollary 5.1.

Corollary 5.4. For each $\epsilon > 0$, we define $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$ as the event

$$\Lambda_{\epsilon}^{\hat{\mathbf{C}}} = \left\{ \forall \nu \in [0,1] : \sigma(\hat{\mathbf{C}}(\nu)) \subset \operatorname{Supp} \mu_{MP}^{(c)} + \epsilon \right\}$$

Then, $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$ holds with exponentially high probability.

Proof. The proof is similar to the proof of Corollary 5.1 and is thus omitted. \Box

We finally use the above results to prove that $\nu \to \frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu)) - \int f \, d\mu_{MP}^{(c_N)}$ is $MN^{3/2}$ -Lipschitz with overwhelming probability. For this, we establish the following Proposition.

Proposition 5.3. It holds that

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{1}{|\delta|} \left| \frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}(\nu)) - \frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}(\nu+\delta)) \right| \prec MN^{3/2}.$$
(5.13)

Proof. By Corollary 5.4, the event $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$ holds with exponentially high probability. Therefore, it is sufficient to establish that

$$\mathbf{1}_{\Lambda_{\epsilon}^{\mathbf{C}}} \sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{1}{|\delta|} \left| \frac{1}{M} \mathrm{tr} \ f(\hat{\mathbf{C}}(\nu + \delta)) - \frac{1}{M} \mathrm{tr} \ f(\hat{\mathbf{C}}(\nu)) \right| \prec M N^{3/2}$$

We express $\frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu + \delta)) - \frac{1}{M} \text{tr } f(\hat{\mathbf{C}}(\nu))$ as

$$\frac{1}{M}\operatorname{tr} f(\hat{\mathbf{C}}(\nu+\delta)) - \frac{1}{M}\operatorname{tr} f(\hat{\mathbf{C}}(\nu)) = \frac{1}{M}\sum_{m=1}^{M} f(\lambda_m(\hat{\mathbf{C}}(\nu+\delta))) - f(\lambda_m(\hat{\mathbf{C}}(\nu))).$$

As f is \mathcal{C}^{∞} on a neighborhood of $\operatorname{Supp}_{MP}^{(c)}$, on the set $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$, there exist some random quantities $(\tilde{\lambda}_m)_{1 \leq m \leq M}$ between $\lambda_m(\hat{\mathbf{C}}(\nu))$ and $\lambda_m(\hat{\mathbf{C}}(\nu+\delta))$ such that

$$\begin{split} \frac{1}{M} \sum_{m=1}^{M} f(\lambda_m(\hat{\mathbf{C}}(\nu+\delta))) &- f(\lambda_m(\hat{\mathbf{C}}(\nu))) \\ &= \frac{1}{M} \sum_{m=1}^{M} \left(\lambda_m(\hat{\mathbf{C}}(\nu+\delta)) - \lambda_m(\hat{\mathbf{C}}(\nu)) \right) \, f'(\tilde{\lambda}_m). \end{split}$$

Using the following eigenvalue inequality for Hermitian matrices:

$$\left|\lambda_m(\hat{\mathbf{C}}(\nu+\delta)) - \lambda_m(\hat{\mathbf{C}}(\nu))\right| \le \|\hat{\mathbf{C}}(\nu+\delta) - \hat{\mathbf{C}}(\nu)\|$$

in conjunction with the fact that $\sup_{1 \leq m \leq M} |f'(\tilde{\lambda}_m)|$ is bounded by some nice constant C on the event $\Lambda_{\epsilon}^{\hat{\mathbf{C}}}$, we obtain that

$$\mathbb{P}\left[\sup_{\delta\neq 0}\sup_{\nu\in[0,1]}\left|\frac{1}{M}\sum_{m=1}^{M}f'(\tilde{\lambda}_{m})(\lambda_{m}(\hat{\mathbf{C}}(\nu+\delta))-\lambda_{m}(\hat{\mathbf{C}}(\nu))\right|\right.\\\left.\left.\left.\left.\left|\delta\right|N^{\kappa}MN^{3/2},\Lambda_{\epsilon}^{\hat{\mathbf{C}}}\right]\right]\right.\\\left.\leq\mathbb{P}\left[\sup_{\delta\neq 0}\sup_{\nu\in[0,1]}C\|\hat{\mathbf{C}}(\nu+\delta)-\hat{\mathbf{C}}(\nu)\|>|\delta|N^{\kappa}MN^{3/2},\Lambda_{\epsilon}^{\hat{\mathbf{C}}}\right]\right]$$

(5.12) eventually leads to (5.13).

	-	-	-	

5.1.2. Lipschitz constant of $\nu \to \hat{r}_N(\nu)$.

The function $\nu \to \hat{r}_N(\nu)$ verifies the following property:

Proposition 5.4.

$$\sup_{\delta \neq 0} \sup_{\nu \in [0,1]} \frac{1}{|\delta|} |\hat{r}_N(\nu + \delta) - \hat{r}_N(\nu)| \prec N^{3/(2\gamma_0 + 1)}.$$
(5.14)

We just provide the main steps the proof, and left the details to the reader. We first prove that $\sup_{\nu \in [0,1]} \sum_{m=1}^{M} \frac{1}{\hat{s}_{m,L}(\nu)} \prec 1$ by verifying that the set $\{\forall \nu \in [0,1], \forall m = 1, \ldots, M, \hat{s}_{m,L}(\nu) \in [\underline{s}, \overline{s}] + \epsilon\}$ holds with exponentially high probability. Then, we establish that $\nu \to \hat{s}_{m,L}(\nu)$ and $\nu \to \hat{s}'_{m,L}(\nu)$ are $N^{2/(2\gamma_0+1)}$ Lipschitz and $N^{3/(2\gamma_0+1)}$ Lipschitz with overwhelming probability. This leads immediately to (5.14).

As $v_N N^{3/(2\gamma_0+1)} \ll M N^{3/2}$, Propositions 5.3 and 5.4 lead to (5.2). This completes the proof of Proposition 5.1.

5.2. Stochastic domination of $\sup_{\nu \in [0,1]} |\psi_N(f,\nu)|$ and $\sup_{\nu \in [0,1]} |\hat{\psi}_N(f,\nu)|$

We are now in position to establish the main result of this paper.

Theorem 5.1. $\sup_{\nu \in [0,1]} |\psi_N(f,\nu)|$ and $\sup_{\nu \in [0,1]} |\hat{\psi}_N(f,\nu)|$ verify the following stochastic domination property:

$$\sup_{\nu \in [0,1]} |\psi_N(f,\nu)| \quad \prec \quad u_N \tag{5.15}$$

$$\sup_{\nu \in [0,1]} |\hat{\psi}_N(f,\nu)| \quad \prec \quad u_N. \tag{5.16}$$

Proof. We just establish (5.16). We consider $\epsilon > 0$ and evaluate

$$\mathbb{P}\left[\sup_{\nu\in[0,1]}\left|\hat{\psi}_N(f,\nu)\right|>N^{\epsilon}u_N\right].$$

We denote by $\nu^* \in [0,1]$ an element where the supremum is achieved, and consider ν_p^* the closest element of \mathcal{V}_N^p from ν^* , where we recall that \mathcal{V}_N^p is defined by (5.8). Therefore, one can write:

$$\mathbb{P}\left[\sup_{\nu\in[0,1]}\left|\hat{\psi}_{N}(f,\nu)\right| > N^{\epsilon}u_{N}\right] \leq \mathbb{P}\left[\left|\hat{\psi}_{N}(f,\nu^{*}) - \hat{\psi}_{N}(f,\nu_{p}^{*})\right| > \frac{1}{2}N^{\epsilon}u_{N}\right] + \mathbb{P}\left[\left|\hat{\psi}_{N}(f,\nu_{p}^{*})\right| > \frac{1}{2}N^{\epsilon}u_{N}\right].$$

(4.52) implies that $\mathbb{P}\left[\left|\hat{\psi}_{N}(f,\nu_{p}^{*})\right| > \frac{1}{2}N^{\epsilon}u_{N}\right]$ converges exponentially towards 0. It thus remains to study $\mathbb{P}\left[\left|\hat{\psi}_{N}(f,\nu^{*})-\hat{\psi}_{N}(f,\nu_{p}^{*})\right|\right| > \frac{1}{2}N^{\epsilon}u_{N}\right]$. For this, we of course use (5.2), Corollary 5.3, and write that

$$\mathbb{P}\left[\left| \hat{\psi}_{N}(f,\nu^{*}) - \hat{\psi}_{N}(f,\nu_{p}^{*}) \right| > \frac{1}{2}N^{\epsilon}u_{N} \right]$$

$$= \mathbb{P}\left[\left| \frac{\hat{\psi}_{N}(f,\nu^{*}) - \hat{\psi}_{N}(f,\nu_{p}^{*}))}{\nu^{*} - \nu_{p}^{*}} \right| > \frac{1}{2|\nu^{*} - \nu_{p}^{*}|}N^{\epsilon}u_{N} \right]$$

$$\leq \mathbb{P}\left[\left| \frac{\hat{\psi}_{N}(f,\nu^{*}) - \hat{\psi}_{N}(f,\nu_{p}^{*}))}{\nu^{*} - \nu_{p}^{*}} \right| > \frac{1}{2}N^{p}N^{\epsilon}u_{N} \right].$$

If we choose p large enough, $MN^{3/2}$ verifies $MN^{3/2} \ll N^p u_N$, and $\mathbb{P}\left[\left|\frac{\hat{\psi}_N(f,\nu^*)-\hat{\psi}_N(f,\nu_p^*))}{\nu^*-\nu_p^*}\right| > \frac{1}{2}N^pN^\epsilon u_N\right]$ converges towards 0 exponentially as expected. This completes the proof of (5.16).

6. Numerical simulations

In this section we examine the impact of the correction quantity $r_N(\nu)\phi_N(f)v_N$ when $\alpha > \frac{2}{3}$ and see how it improves the estimation of the LSS $\frac{1}{M}$ tr $f(\hat{\mathbf{C}}(\nu))$. More precisely, we start by examining the behaviour of the LSS

$$\left|\frac{1}{M} \operatorname{tr} \, f(\hat{\mathbf{C}}(\nu)) - \int_{\mathbb{R}} f \, \mathrm{d}\mu_{MP}^{(c_N)}\right|$$

and the impact of the corrective term

$$\left(\frac{1}{M}\sum_{m=1}^{M}\frac{s'_{m}(\nu)}{s_{m}(\nu)}\right)^{2}\phi_{N}(f)\left(\frac{1}{B+1}\sum_{b=-B/2}^{B/2}\left(\frac{b}{N}\right)\right)^{2} = r_{N}(\nu)\phi_{N}(f)v_{N}$$
$$\left(\frac{1}{M}\sum_{m=1}^{M}\frac{\hat{s}'_{m}(\nu)}{\hat{s}_{m}(\nu)}\right)^{2}\phi_{N}(f)\left(\frac{1}{B+1}\sum_{b=-B/2}^{B/2}\left(\frac{b}{N}\right)\right)^{2} = \hat{r}_{N}(\nu)\phi_{N}(f)v_{N}.$$

under \mathcal{H}_0 . We recall that $\phi_N(f)$ is the deterministic term defined as the action of f on the compactly supported distribution D_N , whose Stieltjes transform is:

$$p_N(z) = -\frac{c_N(zt_N(z)\tilde{t}_N(z))^3}{1 - c_N(zt_N(z)\tilde{t}_N(z))^2}$$

Motivated by [28], we consider $f(\lambda) = (\lambda - 1)^2$ where it can be verified with a bit of algebra and residue calculus that

$$\int_{\mathbb{R}} f(\lambda) \, \mathrm{d}\mu_{MP}^{(c_N)}(\lambda) = c_N$$

and $\phi_N(f) = c_N$. Take \mathbf{y}_n generated by the following simple model:

$$\mathbf{y}_{n+1} = \mathbf{A}\mathbf{y}_n + \boldsymbol{\epsilon}_n \tag{6.1}$$

where $(\epsilon_n)_{n\in\mathbb{Z}}$ is an independent sequence of $\mathcal{N}_{\mathbb{C}}(0, \mathbf{I}_M)$ distributed random vectors, and where **A** is the diagonal matrix defined by $\mathbf{A} = \theta \mathbf{I}_M$ for $\theta \in \mathbb{C}$ such that $|\theta| < 1$. Under (6.1), each time series are independent AR(1) processes. In Figure 1 is represented on the left the values of the LSS associated to $f(\lambda) = (\lambda - 1)^2$ for each $\nu \in (0, 1)$ when (N, B, M, L) = (10119, 1600, 800, 21) (so $\alpha = 0.8$ and c = 1/2) and $\theta = 0.4$, where we recall that L represent the lag window size in the estimation of $\hat{r}_N(\nu)$. We see that the correction term captures the majority of the deviation of the LSS from zero. Moreover, the correction where the spectral densities s_m and s'_m are estimated still provide a good approximation of the $\mathcal{O}(\frac{B}{N})^2$ term. On the right side is represented the LSS against $\psi_N(f, \nu)$ and $\hat{\psi}_N(f, \nu)$. We again observe that the majority of the deviation from zero of the LSS is corrected by the $\mathcal{O}(\frac{B}{N})^2$ terms. Around $\nu = \pm 0.1$, the corrections precision seems degraded. This can be understood since $\nu = \pm 0.1$ corresponds



FIG 1. Linear Spectral Statistics vs the correction term. $f(\lambda) = (\lambda - 1)^2$, (N, B, M, L) = (10119, 1600, 800, 21), and $\theta = 0.4$.

to peaks in s'_m , which leads to greater estimation errors for \hat{s}'_m at this frequency than for the other ones.

We now check the derived speed of convergence towards zero in Theorem 5.1, and more precisely that the following estimations hold true:

$$\begin{split} \sup_{\nu \in [0,1]} \left| \frac{1}{M} \mathrm{tr} \ f(\hat{\mathbf{C}}(\nu)) - \int_{\mathbb{R}} f \, \mathrm{d}\mu_{MP}^{(c_N)} \right| &= \mathcal{O}\left(\frac{1}{B}\right) \mathbbm{1}_{1/2 \le \alpha \le 2/3} + \mathcal{O}\left(\frac{B}{N}\right)^2 \mathbbm{1}_{\alpha \ge 2/3} \\ \sup_{\nu \in [0,1]} |\psi(f,\nu)| &= \mathcal{O}(u_N) \end{split}$$

and we abbreviate $\frac{1}{M}$ tr $f(\hat{\mathbf{C}}(\nu)) - \int_{\mathbb{R}} f d\mu_{MP}^{(c_N)}$ by $LSS(f,\nu)$. In the following we take $c = \frac{M}{B+1} = \frac{1}{2}$ and $\alpha = 4/5$. In this case we recall that $u_N = \mathcal{O}(\frac{B}{N})^3$. On the left of Figure 2 is represented for $M \in \{20, 30, \dots, 1500\}$ the value of $\sup_{\nu \in [0,1]} |LSS(f,\nu)|$ against $\sup_{\nu \in [0,1]} |\psi(f,\nu)|$ and $\sup_{\nu \in [0,1]} |\hat{\psi}(f,\nu)|$. On the right of Figure 2 we rescale all quantities by $(\frac{N}{B})^2$ and observe, in accordance with Theorem 4.1 that $LSS(f,\nu)$ remains $\mathcal{O}(1)$ while the corrected quantities are o(1). Eventually, on Figure 3 are represented $\sup_{\nu \in [0,1]} |\psi(f,\nu)|$ and $\sup_{\nu \in [0,1]} |\hat{\psi}(f,\nu)|$ rescaled by $(\frac{N}{B})^3$, and observe that these quantities are now $\mathcal{O}(1)$, again in accordance with Theorem 4.1.

Eventually on Figure 4 is represented 20000 realisations of the LSS $\sup_{\nu \in \mathcal{F}_N} \left| \frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}(\nu)) - \int f \, d\mu_{MP}^{(c_N)} \right|$ against its improved estimations $\sup_{\nu \in \mathcal{F}_N} |\psi_N(f,\nu)|$ and $\sup_{\nu \in \mathcal{F}_N} |\hat{\psi}_N(f,\nu)|$. We see that the oracle corrected statistics $\psi(f,\nu)$ is more concentrated around 0, and that its estimated counterpart $\hat{\psi}(f,\nu)$ is close to $\psi(f,\nu)$ but exhibits more spread due to the additional estimation step of the $\hat{s}_m(\nu)$.

Appendix A: Appendix

A.1. Proof of Lemma A.1

Lemma A.1 is a slight variation of Theorem 4.3.2 [4].



FIG 2. $\sup_{\nu \in \mathcal{F}_N} \left| \frac{1}{M} \operatorname{tr} f(\hat{\mathbf{C}}(\nu)) - \int f \, \mathrm{d}\mu_{MP}^{(c_N)} \right|$ against $\sup_{\nu \in \mathcal{F}_N} \psi_N(f,\nu)$ and $\sup_{\nu \in \mathcal{F}_N} \hat{\psi}_N(f,\nu)$ as functions of M. On the right the quantities are rescaled by $(\frac{N}{B})^2$. $\alpha = 0.8, c = 1/2, \theta = 0.4$



FIG 3. $\sup_{\nu \in \mathcal{F}_N} \psi_N(f,\nu)$ and $\sup_{\nu \in \mathcal{F}_N} \hat{\psi}_N(f,\nu)$ rescaled by $(\frac{N}{B})^3$ as functions of M. $\alpha = 0.8$, c = 1/2, $\theta = 0.4$.

Lemma A.1. For any ν_1 and ν_2 in [0,1], such that there exists $k \in \{0,1,\ldots,N-1\}$ satisfying $\nu_2 - \nu_1 = k/N$, the following bound holds:

$$\sup_{m \ge 1} |\mathbb{E}\left[\xi_{y_m}(\nu_1)\xi_{y_m}(\nu_2)^*\right] - s_m(\nu_1)\delta_{\nu_1 = \nu_2}| = \mathcal{O}\left(\frac{1}{N}\right).$$
(A.1)



 $\begin{array}{lll} \mbox{Fig} & 4. & \sup_{\nu \in \mathcal{F}_N} |\frac{1}{M} \mbox{tr} & f(\hat{\mathbf{C}}(\nu)) & - \int f \, \mathrm{d} \mu_{MP}^{(c_N)} | & against & \sup_{\nu \in \mathcal{F}_N} |\psi_N(f,\nu)| & and \\ \sup_{\nu \in \mathcal{F}_N} |\hat{\psi}_N(f,\nu)|. & (N,B,M,L) = (4254,800,400,16), \ \theta = 0.4. \end{array}$

Proof.

$$\mathbb{E}\left[\xi_{y_m}(\nu_1)\xi_{y_m}(\nu_2)^*\right]$$

$$= \frac{1}{N}\sum_{n_1,n_2=1}^{N}\mathbb{E}[y_{m,n_1}y_{m,n_2}^*]e^{-2i\pi(n_1-1)\nu_1}e^{2i\pi(n_2-1)\nu_2}$$

$$= \frac{1}{N}\sum_{n_1,n_2=1}^{N}r_{m,n_1-n_2}e^{-2i\pi(n_1-1)\nu_1+2i\pi(n_2-1)\nu_2}$$

$$= \frac{1}{N}\sum_{u=-(N-1),n_1,n_2\in 0,\dots,N-1}^{(N-1)}r_{m,u}\sum_{n_1-n_2=u}e^{-2i\pi n_1\nu_1+2i\pi n_2\nu_2}$$

Splitting this expression for u = 0, u > 0 and u < 0 provides

$$\mathbb{E}[\xi_{y_m}(\nu_1)\xi_{y_m}(\nu_2)^*] = \frac{1}{N}r_{m,0}\sum_{n_1=0}^{N-1} e^{-2i\pi n_1(\nu_2-\nu_1)} + \frac{1}{N}\sum_{u=1}^{(N-1)} r_{m,u}\sum_{n_2=0}^{N-1-u} e^{-2i\pi(u+n_2)\nu_1} e^{2i\pi n_2\nu_2} + \frac{1}{N}\sum_{u=-(N-1)}^{-1} r_{m,u}\sum_{n_2=-u}^{N-1} e^{-2i\pi(u+n_2)\nu_1} e^{2i\pi n_2\nu_2}.$$
 (A.2)

The first term of the right hand side of (A.2) can be computed in the case

 $\nu_1 = \nu_2$:

:

$$\frac{1}{N}r_{m,0}\sum_{n_1=0}^{N-1}e^{-2i\pi n_1(\nu_2-\nu_1)}=r_{m,0}$$

and in the case $\nu_1 \neq \nu_2$,

$$\frac{1}{N}r_{m,0}\sum_{n_1=0}^{N-1}e^{-2i\pi n_1\frac{k}{N}}=0.$$

Therefore, the first term of the right hand side of (A.2) is equal to $r_{m,0}\delta_{\nu_1=\nu_2}$. Consider now the second term of (A.2) (where u > 0):

$$\frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} \sum_{n_2=0}^{N-1-u} e^{-2i\pi(u+n_2)\nu_1} e^{2i\pi n_2\nu_2}$$
$$= \frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)}.$$
(A.3)

The right hand side of (A.3) can also be explicitly written in the case $\nu_1 = \nu_2$

$$\frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)}$$
$$= \frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} (N-u)$$
$$= \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \frac{N-u}{N}$$
$$= \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} - \frac{1}{N} \sum_{u=1}^{N-1} u r_{m,u} e^{2i\pi u\nu_1}$$

By Assumption 1.4, $\sup_{m\geq 1} \sum_{u\in\mathbb{Z}} |u| |r_{m,u}| < +\infty$, so we have:

$$\sup_{m\geq 1} \frac{1}{N} \left| \sum_{u=1}^{N-1} u r_{m,u} e^{2i\pi u\nu_1} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$

Therefore:

$$\sup_{m\geq 1} \left| \frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)} - \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$
(A.4)

In the case where $\nu_1 \neq \nu_2$, note that $\nu_1 - \nu_2 = k/N$ with $k \neq 0$, therefore:

$$\sum_{n_2=0}^{N-1} e^{-2i\pi n_2(\nu_2 - \nu_1)} = \sum_{n_2=0}^{N-1} e^{-2i\pi n_2 \frac{k}{N}} = 0.$$
(A.5)

Using (A.5), one can rewrite the right hand side of (A.3) as

$$\begin{aligned} \left| \frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)} \right| \\ &= \left| -\frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=N-u}^{N} e^{-2i\pi n_2(\nu_2-\nu_1)} \right| \\ &\leq \frac{1}{N} \sum_{u=1}^{N-1} |u| |r_{m,u}| \end{aligned}$$

which, again by Assumption 1.4, provides the bound:

$$\sup_{m\geq 1} \left| \frac{1}{N} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)} \right| = \mathcal{O}\left(\frac{1}{N}\right).$$
(A.6)

Combining (A.4) and (A.6), the second term of the right hand side of (A.2)can be estimated as follow:

$$\sup_{m\geq 1} \left| \frac{1}{N} \sum_{u=1}^{(N-1)} r_{m,u} e^{-2i\pi u\nu_1} \sum_{n_2=0}^{N-1-u} e^{-2i\pi n_2(\nu_2-\nu_1)} - \delta_{\nu_1=\nu_2} \sum_{u=1}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \right|$$
$$= \mathcal{O}\left(\frac{1}{N}\right).$$

The term for u < 0 in equation (A.2) is similar. Gathering the three terms of equation (A.2) leads to

$$\sup_{m\geq 1} \left| \mathbb{E}[\xi_{y_m}(\nu_1)\xi_{y_m}(\nu_2)^*] - \delta_{\nu_1=\nu_2} \left(\sum_{u=-(N-1)}^{N-1} r_{m,u} e^{-2i\pi u\nu_1} \right) \right| = \mathcal{O}\left(\frac{1}{N}\right).$$
(A.7)

Eventually, using again Assumption 1.4 we have:

$$\left|\sum_{|u|>N} r_m(u) e^{-2i\pi u\nu_1}\right| \leq \frac{1}{N} \sum_{|u|>N} |u| |r_m(u)| = \mathcal{O}\left(\frac{1}{N}\right).$$

Inserting this into equation (A.7), we obtain equation (A.1)

ī

A.2. Proof of Lemma 3.1

Proof. Consider the complement of the event $\Lambda_{\epsilon}^{\hat{\mathbf{D}}}(\nu)$ and notice that:

$$\Lambda^{\hat{\mathbf{D}}}_{\epsilon}(\nu)^{c} \subset \{\exists m \in \{1, \dots, M\} : \hat{s}_{m} > \bar{s} + \epsilon\} \cup \{\exists m \in \{1, \dots, M\} : \hat{s}_{m} < \underline{s} - \epsilon\}.$$
(A.8)

We start by proving that the first set of the right hand side of (A.8) holds with is exponentially small probability, ie. for any $\epsilon > 0$, there exist $\gamma > 0$ such that:

$$\mathbb{P}\left[\exists m \in \{1, \dots, M\} : \hat{s}_m > \bar{s} + \epsilon\right] \le \exp{-N^{\gamma}}.$$

By Lemma A.2 (see below), $|\mathbb{E}\hat{s}_m - s_m| = \mathcal{O}(B^2/N^2)$ so for N large enough, this bias term will be smaller than $\epsilon/2$. Moreover, for any $m \in \{1, \ldots, M\}$, $s_m - \bar{s} \leq 0$. Therefore, one can write for large enough N:

$$\mathbb{P}\left[\exists m \in \{1, \dots, M\} : \hat{s}_m > \bar{s} + \epsilon\right]$$

$$= \mathbb{P}\left[\sup_{m \in \{1, \dots, M\}} (\hat{s}_m - \mathbb{E}\hat{s}_m + \mathbb{E}\hat{s}_m - s_m + s_m - \bar{s}) > \epsilon\right]$$

$$\leq \mathbb{P}\left[\sup_{m \in \{1, \dots, M\}} |\hat{s}_m - \mathbb{E}\hat{s}_m| > \epsilon/2\right]$$

which holds with exponentially high probability by Lemma A.3 (see below). The proof for the lower bound is similar. $\hfill \Box$

It remains to prove Lemma A.2 and Lemma A.3. Concerning the proof of Lemma A.2, we follow the same approach as the one used in Theorem 5.4.2 in [4].

Lemma A.2. For any $\nu \in [0, 1]$, the following results hold:

$$\mathbb{E}(\hat{s}_m(\nu)) - s_m(\nu) = \frac{s_m''(\nu)}{2} v_N + \mathcal{O}\left(\left(\frac{B}{N}\right)^3 + \frac{1}{N}\right)$$
(A.9)

and

$$\sup_{=1,\dots,M} |\mathbb{E}\hat{s}_m(\nu) - s_m(\nu)| = \mathcal{O}\left(\frac{B}{N}\right)^2.$$
(A.10)

Proof. It is clear that $\hat{s}_m(\nu) = \hat{\mathbf{S}}_{m,m}(\nu) = s_m(\nu)\tilde{\mathbf{C}}_{m,m}(\nu)$ can be written as

$$\hat{s}_m(\nu) = s_m(\nu) \frac{x_m (\mathbf{I} + \mathbf{\Phi}_m) x_m^*}{B+1}.$$
 (A.11)

Therefore, $\mathbb{E}(\hat{s}_m(\nu)) = s_m(\nu)(1 + \frac{1}{B+1} \text{tr } \Phi_m)$. (A.9) thus follows immediately from (3.9). (A.10) is an immediate consequence of (A.9).

Lemma A.3. The family of random variables $\sup_{m=1,...,M} |\hat{s}_m(\nu) - \mathbb{E}[\hat{s}_m(\nu)]|, \nu \in [0, 1]$ verifies

$$\sup_{n=1,\dots,M} |\hat{s}_m - \mathbb{E}[\hat{s}_m]| \prec \frac{1}{\sqrt{B}}.$$
 (A.12)

Proof. (A.11) implies that $\hat{s}_m - \mathbb{E}[\hat{s}_m]$ can be written as $\hat{s}_m - \mathbb{E}[\hat{s}_m] = s_m \left(\frac{x_m(\mathbf{I}+\boldsymbol{\Phi}_m)x_m^*}{B+1} - \frac{1}{B+1}\mathrm{Tr}(\mathbf{I}+\boldsymbol{\Phi}_m)\right)$. It is clear that $\sup_m \left\|\frac{(\mathbf{I}+\boldsymbol{\Phi}_m)}{B+1}\right\|_F \leq \frac{C}{B}$ for some nice constant C. Therefore, (A.12) leads immediately to (2.17).

A.3. Proof of Lemma 3.3

Proof. These estimates can be proved in a compact way by using the calculus rules available in the stochastic domination framework introduced in Definition 2.1 and proved in Lemma 2.1. Using Lemma 3.2 and Lemma A.4 (see below):

$$\begin{aligned} \frac{1}{\sqrt{\hat{s}_m}} - \frac{1}{\sqrt{s_m}} \bigg| &= \bigg| \frac{\sqrt{s_m} - \sqrt{\hat{s}_m}}{\sqrt{s_m}\sqrt{\hat{s}_m}} \bigg| \\ &\leq \underbrace{\left| \sqrt{s_m} - \sqrt{\hat{s}_m} \right|}_{\mathcal{O}_{\prec}(\frac{1}{\sqrt{B}} + \frac{B^2}{N^2})} \times \underbrace{\left| \sqrt{\frac{1}{s_m}} \right|}_{\mathcal{O}_{\prec}(1)} \times \underbrace{\left| \sqrt{\frac{1}{\hat{s}_m}} \right|}_{\mathcal{O}_{\prec}(1)} \\ &\prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}. \end{aligned}$$

The second inequality is similar to prove:

$$\begin{split} \left| \sqrt{\frac{s_m}{\hat{s}_m}} - 1 \right| &= \left| \frac{\sqrt{s_m} - \sqrt{\hat{s}_m}}{\sqrt{\hat{s}_m}} \right| \\ &\leq \underbrace{|s_m - \hat{s}_m|}_{\mathcal{O}_{\prec}(\frac{1}{\sqrt{B}} + \frac{B^2}{N^2})} \times \underbrace{\frac{1}{\hat{s}_m(\sqrt{s_m + \hat{s}_m})}}_{\mathcal{O}_{\prec}(1)} \\ &\prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}. \end{split}$$

Lemma A.4. The family of random variables $(\sup_{m=1,...,M} |\hat{s}_m(\nu) - s_m(\nu)|), \nu \in [0,1]$ verifies

$$\sup_{m=1,...,M} |\hat{s}_m - s_m| \prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}$$

Proof. It is sufficient to check that the family of random variables $(|\hat{s}_m - s_m|)_{m=1,...,M}, \nu \in [0,1]$ verifies $|\hat{s}_m - s_m| \prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}$. Using Lemma A.2 and Lemma A.3, we obtain as expected that

$$|\hat{s}_m - s_m| = |s_m - \mathbb{E}\hat{s}_m + \mathbb{E}\hat{s}_m - \hat{s}_m| \le \underbrace{|s_m - \mathbb{E}\hat{s}_m|}_{\mathcal{O}(\frac{B^2}{N^2})} + \underbrace{|\mathbb{E}\hat{s}_m - \hat{s}_m|}_{\mathcal{O}_{\prec}(\frac{1}{\sqrt{B}})} \prec \frac{1}{\sqrt{B}} + \frac{B^2}{N^2}.$$

A.4. Proof of Lemma A.5

Lemma A.5. The set of random variable $(\sum_{m=1}^{M} |\hat{s}_m(\nu) - s_m(\nu)|^2), \nu \in [0, 1]$ verifies

$$\sum_{m=1}^{M} |\hat{s}_m - s_m|^2 \prec 1 + \frac{B^5}{N^4}.$$

Proof. Using Lemma A.4, we have

$$|\hat{s}_m - s_m|^2 \prec \frac{1}{B} + \frac{B^4}{N^4}$$

and summing over $m = 1 \dots M$, one immediately get:

$$\sum_{m=1}^{M} |\hat{s}_m - s_m|^2 \prec 1 + \frac{B^5}{N^4}$$

A.5. Proof of Lemma 4.5

. We express $\tilde{\Delta}$ as $\tilde{\Delta} = \frac{\mathbf{X}\Gamma^*}{B+1} + \frac{\Gamma\mathbf{X}^*}{B+1} + \frac{\Gamma\Gamma^*}{B+1}$. Therefore, we have

$$\|\tilde{\mathbf{\Delta}}\|^k \le C\left(\left\|\frac{\mathbf{X}}{\sqrt{B+1}}\right\|^k \left\|\frac{\mathbf{\Gamma}}{\sqrt{B+1}}\right\|^k + \left\|\frac{\mathbf{\Gamma}\mathbf{\Gamma}^*}{B+1}\right\|^k\right).$$

Using the Schwartz inequality, we obtain that

$$\mathbb{E}\|\tilde{\mathbf{\Delta}}\|^{k} \leq C\left(\left(\mathbb{E}\left\|\frac{\mathbf{X}\mathbf{X}^{*}}{B+1}\right\|^{k}\right)^{1/2} \left(\mathbb{E}\left\|\frac{\mathbf{\Gamma}\mathbf{\Gamma}^{*}}{B+1}\right\|^{k}\right)^{1/2} + \mathbb{E}\left\|\frac{\mathbf{\Gamma}\mathbf{\Gamma}^{*}}{B+1}\right\|^{k}\right).$$

It is well-known that $\mathbb{E}\left(\left\|\frac{\mathbf{X}\mathbf{X}^*}{B+1}\right\|^k\right) \leq C$ for some nice constant depending on k. Therefore, we establish that

$$\mathbb{E} \left\| \frac{\mathbf{\Gamma} \mathbf{\Gamma}^*}{B+1} \right\|^k \le C \left(\frac{B}{N} \right)^{2k}$$

a property which will imply that $\mathbb{E}\|\tilde{\boldsymbol{\Delta}}\|^k \leq C\left(\frac{B}{N}\right)^k$. For this, we put $\mathbf{Z} = \frac{\Gamma\Gamma^*}{B+1}$. As (3.17) holds, it remains to verify that $\mathbb{E}(\|\mathbf{Z}-\mathbb{E}(\mathbf{Z})\|^k) = \mathcal{O}\left(\frac{B}{N}\right)^{2k}$. For this, we use the concentration inequality (3.23). We choose $t_N = w^{1/k} (\frac{B}{N})^2$, and obtain that

$$\mathbb{P}\left[\frac{\|\mathbf{Z} - \mathbb{E}(\mathbf{Z})\|}{(\frac{B}{N})^2} > w^{1/k}\right] \le 2C_0 \exp{-CB(w^{1/k} - w_0^{1/k})}$$
(A.13)

for some $w_0 > 0$. If we denote by z_N the random variable $z_N = \left(\frac{\|\mathbf{Z} - \mathbb{E}(\mathbf{Z})\|}{(\frac{B}{N})^2}\right)^k$, we have to establish that $\mathbb{E}(z_N) = \mathcal{O}(1)$. For this, we express $\mathbb{E}(z_N)$ as

$$\mathbb{E}(z_N) = \int_0^{+\infty} \mathbb{P}(z_N > w) \, dw = \int_0^{w_0} \mathbb{P}(z_N > w) \, dw + \int_{w_0}^{+\infty} \mathbb{P}(z_N > w) \, dw.$$

As $\mathbb{P}(z_N > w) = \mathbb{P}(z_N^{1/k} > w^{1/k})$, (A.13) immediately implies that $\mathbb{E}(z_N) = \mathcal{O}(1)$.
A.6. Proof of Lemma 4.6

We denote by η_m the term of interest, i.e. $\eta_m = \mathbb{E}(\mathbf{w}_m \mathbf{A} \mathbf{W}^* \mathbf{Q} \mathbf{e}_m)$. η_m can be written as

$$\eta_m = \sum_{n_1} \left(\sum_{n_2, m'} \mathbb{E}(\mathbf{W}_{m, n_2} \overline{\mathbf{W}}_{m', n_1} \mathbf{Q}_{m', m}) \mathbf{A}_{n_2, n_1} \right).$$

The integration by part formula (2.8) leads to

$$\mathbb{E}(\mathbf{W}_{m,n_2}\overline{\mathbf{W}}_{m',n_1}\mathbf{Q}_{m',m}) = \delta_{m-m'}\delta_{n_1-n_2}\frac{1}{B+1}\mathbb{E}(\mathbf{Q}_{m,m}) - \frac{1}{B+1}\mathbb{E}\left[\overline{\mathbf{W}}_{m',n_1}(\mathbf{Q}\mathbf{W})_{m',n_2}\mathbf{Q}_{m,m}\right].$$

Therefore, we obtain that

$$\sum_{n_2,m'} \mathbb{E}(\mathbf{W}_{m,n_2} \overline{\mathbf{W}}_{m',n_1} \mathbf{Q}_{m',m}) \mathbf{A}_{n_2,n_1}$$
$$= \frac{1}{B+1} \mathbb{E}(\mathbf{Q}_{m,m}) \mathbf{A}_{n_1,n_1} - \frac{1}{B+1} \mathbb{E}\left[(\mathbf{W}^* \mathbf{Q} \mathbf{W} \mathbf{A})_{n_1,n_1} \mathbf{Q}_{m,m}\right]$$

and that

$$\eta_{m} = \mathbb{E}[\mathbf{Q}_{m,m}] \frac{1}{B+1} \operatorname{tr} \mathbf{A} - \mathbb{E}\left[\left(\frac{1}{B+1} \operatorname{tr} \mathbf{W}^{*} \mathbf{Q} \mathbf{W} \mathbf{A}\right) \mathbf{Q}_{m,m}\right]$$
$$= \beta \frac{1}{B+1} \operatorname{tr} \mathbf{A} - \beta \mathbb{E}\left(\frac{1}{B+1} \operatorname{tr} \mathbf{W}^{*} \mathbf{Q} \mathbf{W} \mathbf{A}\right)$$
$$- \mathbb{E}\left[\left(\frac{1}{B+1} \operatorname{tr} \mathbf{W}^{*} \mathbf{Q} \mathbf{W} \mathbf{A}\right)^{\circ} \mathbf{Q}_{m,m}^{\circ}\right].$$
(A.14)

In order to evaluate $\mathbb{E}\left(\frac{1}{B+1} \operatorname{tr} \mathbf{W}^* \mathbf{QWA}\right)$, we remark that

$$\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{W}^{*}\mathbf{QWA}\right) = \frac{1}{M}\sum_{m=1}^{M}\eta_{m}$$
$$= \beta \frac{1}{B+1}\operatorname{tr} \mathbf{A} - \beta c \mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{W}^{*}\mathbf{QWA}\right) - \mathbb{E}\left[\left(\frac{1}{B+1}\operatorname{tr} \mathbf{W}^{*}\mathbf{QWA}\right)^{\circ}\frac{1}{M}\operatorname{tr} \mathbf{Q}^{\circ}\right]$$

from which we deduce that

$$\mathbb{E}\left(\frac{1}{M}\operatorname{tr} \mathbf{W}^{*}\mathbf{Q}\mathbf{W}\mathbf{A}\right) = \frac{\beta}{1+\beta c} \frac{1}{B+1}\operatorname{tr} \mathbf{A} - \frac{1}{1+\beta c} \mathbb{E}\left[\left(\frac{1}{B+1}\operatorname{tr} \mathbf{W}^{*}\mathbf{Q}\mathbf{W}\mathbf{A}\right)^{\circ} \frac{1}{M}\operatorname{tr} \mathbf{Q}^{\circ}\right].$$

Plugging this relation into (A.14) leads immediately to (4.45).

References

- [1] BAI, Z. and SILVERSTEIN, J. W. (2010). Spectral analysis of large dimensional random matrices **20**. Springer.
- [2] BENAYCH-GEORGES, F. and KNOWLES, A. (2016). Lectures on the local semicircle law for Wigner matrices. arXiv preprint arXiv:1601.04055.
- [3] BÖHM, H. and VON SACHS, R. (2009). Shrinkage estimation in the frequency domain of multivariate time series. *Journal of Multivariate Analysis* 100 913–935.
- [4] BRILLINGER, D. R. (1981). Time series: data analysis and theory 36. Siam.
- [5] CAPITAINE, M., DONATI-MARTIN, C. and FÉRAL, D. (2009). The largest eigenvalue of finite rank deformation of large Wigner matrices. *Ann. Probab.* 37 1-47.
- [6] CHEN, X. and LIU, W. (2018). Testing independence with highdimensional correlated samples. Ann. Stat. 46(2) 866–895.
- [7] DETTE, H. and DÖRNEMANN, N. (2020). Likelihood ratio tests for many groups in high dimensions. *Journal of Multivariate Analysis* 104605.
- [8] DUCHESNE, P. and ROY, R. (2003). Robust tests for independence of two time series. *Statistica Sinica* 827–852.
- [9] DYN'KIN, E. M. (1972). An operator calculus based on the Cauchy–Green formula. Zapiski Nauchnykh Seminarov POMI 30 33–39.
- [10] EICHLER, M. (2007). A frequency-domain based test for non-correlation between stationary time series. *Metrika* 65 133–157.
- [11] EICHLER, M. (2008). Testing nonparametric and semiparametric hypotheses in vector stationary processes. *Journal of Multivariate Analysis* 99 968– 1009.
- [12] EL HIMDI, K., ROY, R. and DUCHESNE, P. (2003). Tests for nonnorrelation of two multivariate time series: A nonparametric approach. *Lecture Notes-Monograph Series, vol. 42* 397–416.
- [13] ERDOS, L., KNOWLES, A. and YAU, H.-T. (2013). Averaging fluctuations in resolvents of random band matrices. In *Annales Henri Poincaré* 14 1837– 1926. Springer.
- [14] HAAGERUP, U. and THORBJORNSEN, S. (2003). Random matrices with complex Gaussian entries. *Expositiones Mathematicae* **21** 293–337.
- [15] HACHEM, W., LOUBATON, P. and NAJIM, J. (2007). Deterministic equivalents for certain functionals of large random matrices. *The Annals of Applied Probability* 17 875–930.
- [16] HACHEM, W., KOHRUNZHIY, O., LOUBATON, P., NAJIM, J. and PAS-TUR, L. (2008). A New Approach for Capacity Analysis of Large Dimensional Multi-Antenna Channels. *IEEE Transactions on Information Theory* 54 3987–4003.
- [17] HAUGH, L. D. (1976). Checking the independence of two covariancestationary time series: a univariate residual cross-correlation approach. *Journal of the American Statistical Association* **71** 378–385.
- [18] HELFFER, B. and SJOSTRAND, J. (1989). Equation de Schrödinger avec

champ magnétique et équation de Harper. In *Schrödinger operators* 118–197. Springer.

- [19] HIMDI, K. E. and ROY, R. (1997). Tests for noncorrelation of two multivariate ARMA time series. *Canadian Journal of Statistics* **25** 233–256.
- [20] HONG, Y. (1996). Testing for independence between two covariance stationary time series. *Biometrika* 83 615–625.
- [21] JIANG, T. (2004a). The limiting distributions of eigenvalues of sample correlation matrices. Sankhya: The Indian Journal of Statistics 66 35–48.
- [22] JIANG, T. (2004b). The asymptotic distributions of the largest entries of sample correlation matrices. Ann. Appl. Probab. 14(2) 865–880.
- [23] JIANG, T. and YANG, F. (2013). Central limit theorems for classical likelihood ratio tests for high-dimensional distributions. Ann. of Stats. 41 2029– 2074.
- [24] KOCH, P. D. and YANG, S.-S. (1986). A method for testing the independence of two time series that accounts for a potential pattern in the cross-correlation function. *Journal of the American Statistical Association* 81 533–544.
- [25] LI, W. and HUI, Y. (1994). Robust residual cross correlation tests for lagged relations in time series. *Journal of Statistical Computation and Simulation* **49** 103–109.
- [26] LOUBATON, P. (2016). On the almost sure location of the singular values of certain Gaussian block-Hankel large random matrices. J. Theor. Prob. 4 1339-1443.
- [27] LOUBATON, P., ROSUEL, A. and VALLET, P. (2021). On the asymptotic distribution of the maximum sample spectral coherence of Gaussian time series in the high dimensional regime. Preprint arXiv:2107.02891.
- [28] MESTRE, X. and VALLET, P. (2017). Correlation tests and linear spectral statistics of the sample correlation matrix. *IEEE Transactions on Informa*tion Theory 63 4585–4618.
- [29] PAN, G., JITI, G. and YANRONG, Y. (2014). Testing independence among a large numbers of high-dimensional random vectors. J. Amer. Stat. Assoc. 109 600-612.
- [30] PASTUR, L. and SHCHERBINA, M. (2011). *Eigenvalue Distribution of Large Random Matrices*. Mathematical Surveys and Monographs, AMS.
- [31] ROSUEL, A., VALLET, P., LOUBATON, P. and MESTRE, X. (2021). On the detection of low-rank signal in the presence of spatially uncorrelated noise: a frequency domain approach. Preprint . arXiv:2106.12815.
- [32] RUDELSON, M., VERSHYNIN, R. et al. (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability* 18.
- [33] TANIGUCHI, M., PURI, M. L. and KONDO, M. (1996). Nonparametric approach for non-Gaussian vector stationary processes. *Journal of Multi*variate Analysis 56 259–283.
- [34] TAO, T. (2011). Topics in random matrix theory. Graduate Studies in Mathematics 132.
- [35] WAHBA, G. (1971). Some tests of independence for stationary multivariate

time series. Journal of the Royal Statistical Society: Series B (Methodological) **33** 153–166.