



HAL
open science

Changes in nutrient concentrations of leaves and roots in response to global change factors

Jordi Sardans, Oriol Grau, Han Chen, Ivan Janssens, Philippe Ciais, Shilong Piao, Josep Penuelas

► **To cite this version:**

Jordi Sardans, Oriol Grau, Han Chen, Ivan Janssens, Philippe Ciais, et al.. Changes in nutrient concentrations of leaves and roots in response to global change factors. *Global Change Biology*, 2017, 23 (9), pp.3849-3856. 10.1111/gcb.13721 . hal-02903769

HAL Id: hal-02903769

<https://hal.science/hal-02903769>

Submitted on 7 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC0 - Public Domain Dedication 4.0 International License

1 **Changes in nutrient concentrations of leaves and roots in**
2 **response to Global Change factors.**

3

4

5

6 JORDI SARDANS², ORIOL GRAU^{1,2}, HAN Y. H. CHEN³, IVAN A. JANSSENS⁴, PHILIPPE
7 CIAIS⁵, SHILONG PIAO⁶, JOSEP PEÑUELAS^{1,2}

8

9

10 ¹ CSIC, Global Ecology Unit CREAM-CEAB-UAB, Cerdanyola del Vallès, 08193 Catalonia, Spain.

11 ² CREAM, Cerdanyola del Valles, 08193 Catalonia, Spain.

12 ³Faculty of Natural Resources Management, Lakehead University, 955 Oliver Road, Thunder Bay,
13 Ontario, P7G 1A6, Canada.

14 ⁴Research Group of Plant and Vegetation Ecology (PLECO), Department of Biology, University of
15 Antwerp, B-2610 Wilrijk, Belgium.

16 ⁵ Laboratoire des Sciences du Climat et de l'Environnement, IPSL, 91191 Gif-sur-Yvette, France.

17 ⁶Sino-French Institute for Earth System Science, College of Urban and Environmental Sciences,
18 Peking University, Beijing 100871, China.

19

20 Running head: Foliar and root nutrients under global change

21

22

23

24

25

26

27

28

29

30

31

This is the accepted version of the following article: Sardans, J. et al. "Changes in nutrient concentrations of leaves and roots in response to global change factors" in *Global change biology*, vol. 23, issue 9 (Sep. 2017), p. 3849-3856, which has been published in final form at DOI 10.1111/gcb.13721. **This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.**

32 **Abstract**

33 Global Change impacts on biogeochemical cycles have been widely studied, but our
34 understanding of whether the responses of plant elemental composition to Global Change
35 drivers differ between above- and belowground plant organs remains incomplete. We
36 conducted a meta-analysis of 201 reports including 1,687 observations of studies that
37 have analyzed simultaneously N and P concentrations changes in leaves and roots in the
38 same plants in response to drought, elevated [CO₂], and N and P fertilization around the
39 world, and contrasted the results within those obtained with a general database (838
40 reports and 14,772 observations) that analyzed the changes in N and P concentrations in
41 leaves and/or roots of plants submitted to the commented global change drivers. At global
42 level, elevated [CO₂] decreased N concentrations in leaves and roots and decreased N:P
43 ratio in roots but no in leaves, but was not related with P concentration changes. However,
44 the response differed among vegetation types. In temperate forests elevated [CO₂] was
45 related with lower N concentrations in leaves but not in roots, whereas in crops the
46 contrary patterns were observed. Elevated [CO₂] decreased N concentrations in leaves
47 and roots in tundra plants, whereas not clear relationships were observed in temperate
48 grasslands. However, when elevated [CO₂] and N fertilization coincided, leaves had lower
49 N concentrations whereas root had higher N concentrations suggesting that more nutrients
50 will be allocated to roots to improve uptake of the soil resources not directly provided by
51 the global change drivers. N fertilization and drought increased foliar and root N
52 concentrations while the effects on P concentrations were less clear. The changes in N
53 and P allocation to leaves and root, especially those occurring in opposite direction
54 between them have the capacity to differentially affect above- and below-ground
55 ecosystem functions, such as litter mineralization and above- and belowground food webs.

56 **Keywords.** CO₂; ecological stoichiometry; drought, N deposition; nitrogen; N:P;
57 phosphorus.

58

59

60

61

62 **Introduction**

63 Global Change is characterized by simultaneous shifts in multiple factors that affect
64 ecosystem functioning, such as elevated [CO₂], N deposition, warming and drought,
65 species invasion, and many others (Elser *et al.*, 2010; Sardans *et al.*, 2012; Peñuelas *et*
66 *al.*, 2013). Several of these Global Change drivers are known to affect nutrient cycling and
67 availability. Several interactions among drivers of Global Change, such as elevated CO₂
68 concentrations, N and P fertilization, drought or species invasion, with ecosystemic
69 nutrient availability and cycles have been reported (Sardans & Peñuelas, 2012). A recent
70 meta-analysis by Yuan and Chen (2015) reported different shifts in whole-plant N and P
71 concentrations and in C:P, C:N and N:P ratios in response to elevated [CO₂], N and P
72 fertilization, drought and warming, consistent with previous studies (Sardans *et al.*, 2008;
73 Duval *et al.*, 2012). Another recent meta-analysis (Deng *et al.*, 2015) reported decreases
74 in plant N and P concentrations and N:P ratios under elevated [CO₂]. Foliar nutrient
75 concentrations correlate with overall shifts in plant and ecosystemic functioning (Kerkhoff
76 *et al.*, 2005; Kerkhoff & Enquist, 2006), so the results of these meta-analyses have
77 motivated modelers to include N, and more recently P controls on photosynthesis and
78 plant growth for projecting the future evolution of carbon sinks (Peñuelas *et al.*, 2013; Van
79 der Velde *et al.*, 2014).

80 Changes in nutrient use and allocation in response to changes in environmental
81 conditions are likely to vary among plant organs as a consequence of their particular
82 functions. Recent studies by Gargallo-Garriga *et al.* (2014, 2015) have observed different
83 metabolic and nutrient responses in photosynthetic tissues and roots in different grasses
84 submitted to distinct levels of drought and warming. Root metabolism under drought
85 shifted toward the up-regulation of primary metabolism linked with growth and energy

86 production, whereas photosynthetic tissues in the same plants shifted toward the up-
87 regulation of secondary metabolism associated with anti-stress and conservative functions
88 and toward the down-regulation of primary metabolism (growth and energy production).
89 These asymmetrical effects on metabolic function between photosynthetic organs and
90 roots of the same plants were also correlated with asymmetrical allocations of N and P to
91 photosynthetic organs and roots, with more nutrients to roots to allow that up-regulation
92 and less to leaves related to the observed down-regulation (Gargallo-Garriga *et al.*, 2014,
93 2015).. Interactions between different drivers such as CO₂ and nutrient fertilization
94 introduce further complexity, because their effects can be synergistic or antagonistic, and
95 not simply additive (Churkina *et al.*, 2009; Leuzinger *et al.*, 2011; Dieleman *et al.*, 2012;
96 Reich & Hobbie, 2012; Thomas *et al.*, 2013). Meyerholt & Zaehle (2015) reported that a
97 model of forest production consistently produced the best results when changes in the C:N
98 ratio were decoupled among different plant organs. Thus, the possible different
99 relationships of global change drivers with distinct plant organs warrant investigation.
100 Despite previous studies have demonstrated general effects of global change drivers such
101 as elevated [CO₂], drought, warming, N and P fertilization and species invasion on plant N
102 and P concentrations and N:P ratios (Yuan and Chen, 2015; Deng *et al.*, 2015; Sardans *et*
103 *al.*, 2017), these studies have not distinguished among plant organs and have mainly
104 focused on aboveground data, so a clear understanding of whether the effects of global
105 change drivers are different in distinct plant organs, issues that can only be determined by
106 using studies that provided simultaneous data for the different plant organs of the same
107 plants submitted to different environmental conditions.

108 All these empirical data illustrate the need to account separately for nutrient
109 adjustments in different plant organs to better understand carbon and nutrient cycles.
110 Several experiments have observed different responses to elevated CO₂ of both N (Li *et*

111 *al.*, 2013; Liu *et al.*, 2013a) and P (Duval *et al.*, 2012; Li *et al.*, 2013; Liu *et al.*, 2013b)
112 concentrations in different plant organs. *In situ* climate manipulation experiments have also
113 revealed asymmetrical relationships of warming and/or drought, with contrasting changes
114 in N (Sardans *et al.*, 2008a; Akmal *et al.*, 2010) and P (Sardans *et al.*, 2008b)
115 concentrations among different plant organs. These asymmetrical relationships between
116 leaves and roots have been observed not only for N and P but also for other elements
117 (Gargallo-Garriga *et al.*, 2014, 2015). The response of the elemental compositions of
118 leaves and roots to environmental changes, hereafter elevated [CO₂], N and P fertilization
119 and drought, however, has not yet been analyzed globally and for different vegetation
120 types. Such global analysis of plant leaves and roots is crucial to project global change
121 impacts on the functional composition of plant communities, plant-microbe-soil interactions
122 and possible mismatches between aboveground versus belowground processes.

123 We hypothesized that the impacts of Global-Change drivers on plant elemental
124 composition differ between leaves and roots due to their different functions. We
125 hypothesize that more nutrients will be allocated to roots to improve uptake of the
126 resources not provided by the global change drivers. For example, more nutrients should
127 be allocated to roots under drought, to enhance root primary metabolism and growth for
128 uptake of soil water. To test this hypothesis, we conducted a global meta-analysis of
129 published data on the response of nutrient concentrations in foliar and root tissues of
130 different vegetation types elevated [CO₂], N and P fertilization and drought and their
131 interactions.

132 **Material and Methods**

133 *Data collection*

134 We searched the ISI Web of Science using combinations of the following keywords:
135 availability, carbon, concentration, content, C:N, C:P, deposition, foliar, CO₂, leaf, needle,

136 nitrogen, N:K, N:P, phosphorus, plant, potassium, P:K, ratio, root, soil, solution,
137 stoichiometric and stoichiometry. We obtained 838 reports with 14,772 observations from
138 around the world (see Figures 1-3, S1, S3-S5 ; Table S1, Supplementary Material).

139

140 *Data analyses*

141 We examined the effects of Global-Change drivers on the differences of foliar and root
142 elemental compositions and stoichiometries between control and treated plants by
143 calculating the response ratios from each study, as described by Hedges *et al.* (Hedges et
144 al., 1999). The natural-log response ratio ($\ln RR$) was calculated as $\ln (X_i/X_n) = \ln X_i - \ln X_n$,
145 where X_i and X_n are the values of each observation in the treated and control plants,
146 respectively. The sampling variance for each $\ln RR$ was calculated as $\ln[(1/n_i) \times (S_i/X_i)^2 +$
147 $(1/n_n) \times (S_n/X_n)^2]$ using the R package metafor 1.9–2 (Viechtbauer, 2012), where n_i and n_n ,
148 S_i and S_n and X_i and X_n are the treated and control sample sizes, standard deviations and
149 mean response values, respectively. The natural-log response ratios were determined by
150 specifying studies as random factors using the *rma* model in metafor. The difference in the
151 N and P concentrations and N:P ratios between the leaves and roots of treated and control
152 plants were considered significant if the 95% confidence interval of $\ln RR$ did not overlap
153 zero. All statistical analyses were performed in RStudio 3.1.2 (R Core team, 2015). We
154 analyzed only the variables with more than 30 observations available at the Global scale.
155 We thereafter examined the sensitivities of plant nutrient concentrations and their
156 stoichiometries to the Global-change drivers using REML estimation in the *rma.uni* model
157 for metafor. We defined sensitivity as the change of the variable (N and P concentrations,
158 in mg g^{-1} , and N:P ratios in roots and leaves) as a function of the unit of the Global-Change
159 driver, i.e. ppm CO_2 for elevated $[\text{CO}_2]$, kg ha^{-1} for N and P fertilization and l m^{-2} of rainfall
160 for drought. To avoid the possible bias we have focused on the meta-analyses of the
161 studies that simultaneously analyzed leaves and roots of the same plants. We have also

162 analyzed data from the studies that analyzed leaves and roots of the same plants or
163 different plants to only confirm the results (results provided in supplementary material).
164 The number of reports and observations used for the statistical analysis of each foliar and
165 root variable studied are shown in Figures 1-3. The equivalent meta-analysis on global
166 data set with all reports with information on leaf and/or roots was also shown (Figures S1-
167 S6, Supplementary Material). We also analyzed the natural-log response ratio of different
168 biomes when possible (more than 15 different observations).

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191 **Results**192 *General stoichiometric responses of leaves and roots*193 Studies providing data for leaves and roots of the same plants showed that elevated [CO₂]

194 is associated with decreases in both foliar and root N concentrations (Figure 1; Table 1).

195 The natural-log response ratios (from here onward simplified as response ratios) of N

196 concentrations was thus negative in leaves and roots associated to elevated [CO₂],

197 whereas no effect was observed for foliar and root P concentrations (Figure 1) With

198 drought, both leaf and root N concentrations increased, root P increased marginally

199 ($P < 0.1$), whereas leaf P did not change (Figure 1). N fertilization increased N

200 concentrations in both leaves and roots and decreased in P concentrations in roots (Figure

201 1). These changes translated into increases in the leaf and root N:P ratio under drought

202 (Figure 1).

203 The combination of elevated [CO₂] and N fertilization was related with an increase

204 of root N and P concentrations and also a decrease in leaf N concentrations (Figure 1).

205 Fertilization with both N and P increased both foliar and root N and P concentrations

206 (Figure 1). Table 1 summarizes these results from studies where leaves and roots were

207 simultaneously measured.

208 Studies providing all available data, including also root-only or foliage-only

209 responses from different experiments, reported similar results (Figure S1). There were

210 similar responses of N and P concentrations to elevated [CO₂] (Figures S1). Leaves had211 lower N:P ratio under elevated [CO₂], whereas no effects were observed in roots (Figure

212 S1). Drought was associated with higher N concentrations in leaves and roots, but with a

213 more pronounced increase in roots than in leaves. Drought was related with higher root P
214 concentrations and lower foliar P concentrations (Figure S1). The increases in the
215 response ratio of root P concentrations were more accentuated for severe droughts
216 (Figure S2). N fertilization was associated with higher leaf and even higher root N
217 concentrations (Figure S1). Higher leaf P concentrations and higher root N:P ratio were
218 also associated to N fertilization (Figure S1). Similarly, P fertilization was related with
219 increases in N and P concentrations in leaves and roots (Figure S1). The combination of
220 elevated CO₂ and N fertilization was related with an increase of root N concentration
221 (Figure S1).

222 The response ratios of N and P concentrations in leaves and roots in studies
223 providing data for leaves and roots of the same plants thus show a negative sensitivity of
224 leaf and root N concentration to elevated [CO₂] and positive N sensitivities of N
225 concentrations in leaves and roots to N fertilization whereas no differences were observed
226 in the corresponding sensitivities of natural log response ratios of P (Figure 2).

227 *Differential stoichiometry responses among vegetation types*

228 Elevated [CO₂] differentially altered nutrient concentrations in different vegetation types.
229 Decreases in leaf N concentrations but not in root N concentrations were observed in
230 grasslands, whereas the opposite pattern was observed in croplands exposed to elevated
231 [CO₂] (Figure 3). Decreases in foliar N concentrations and, on the contrary, increases in
232 root N concentrations were observed in temperate forests (Figure 3). As also observed
233 when analyzing all vegetation types together, leaf and root N concentrations increased in
234 temperate forests under N fertilization (Figure 3). However, we had not sufficient data to
235 analyze boreal and tropical ecosystems separately.

236 Studies providing data for leaves and/or roots but not of the same plants also
237 reported that elevated $[\text{CO}_2]$ and drought were associated with different changes
238 depending on vegetation type. Under elevated $[\text{CO}_2]$, decreases in the leaf N
239 concentrations were observed in the three vegetation types studied (temperate forests,
240 temperate grasslands and crops) (Figure S3), whereas decreases in root N concentrations
241 were observed only in temperate forests (Figure S3).

242 Drought was associated with increases in foliar N concentrations in temperate
243 grasslands and decreases of foliar P concentration in temperate forest, and with increases
244 in root N and P concentrations in crops (Figure S3). All studied vegetation types showed
245 increases in foliar and root N concentrations associated with N fertilization (except root N
246 concentrations in tropical forests) (Figure S3). N fertilization was also associated to an
247 increase in response ratio of P leaves concentrations in temperate grasslands (Figure S3).

248 A sensitivity analysis of these heterogeneous data confirmed the positive
249 relationships between drought and N and P concentrations in roots and the negative
250 relationships between drought with N and P concentrations in leaves (Fig. S4). As a result,
251 the N:P ratios in roots presented positive responses to elevated $[\text{CO}_2]$, whereas the N:P
252 ratio in leaves presented negative responses (Fig S4).

253

254

255 **Discussion**

256 The changes in N and P concentrations and N:P ratios in response to the drivers studied
257 were mostly similar in leaves and roots, but in some cases and vegetation types were of
258 different magnitudes and even of different directions. Our results confirmed previous
259 studies (Deng *et al.*, 2015; Peñuelas & Matamala, 1990) that reported a dilution effect
260 under elevated [CO₂], with lower leaf and root N concentrations. Moreover, elevated [CO₂]
261 was associated with changes in N but not in P concentrations. The decreases in leaf and
262 root N concentrations were not accompanied by significant decreases in P leaf and root
263 concentrations in response to elevated [CO₂] observing a decrease in roots N:P ratio.
264 Deng *et al.* (2015) also showed that CO₂ enhancement had no effect on P concentrations
265 in plant tissues, when analyzing data coming only from FACE experiments. These results
266 suggest that increased plant biomass under elevated [CO₂] can be achieved with a more
267 efficient use of N (consistent with a faster turnover of N-rich Rubisco enzymes to fix more
268 carbon), but not of P, resulting in lower tissue N:P ratios. These lower N:P ratios,
269 especially when both nutrients are not limiting, are thus associated with the higher growth
270 rates observed under elevated [CO₂], consistent with the growth rate hypothesis (Stern
271 Elser, 2002). Faster growth would be associated with lower N:P ratios due to the need for
272 more P-rich RNA to sustain rapid protein synthesis (Stern & Elser, 2002).

273 Moreover, elevated [CO₂] in temperate forests decreased foliar N concentration but
274 not root N concentration. In contrast, in temperate grasslands, tundra and crops N
275 concentration changes under elevated [CO₂] were not different between leaves and roots.
276 This shows that the impacts of global change drivers are different depending on vegetation
277 types and biomes. In this case, the results suggested that the root nutrient concentrations
278 in trees are more responsive than those of grasslands or crops. More N is thus allocated to

279 roots of trees under elevated [CO₂] suggesting a tree investment in the uptake of more soil
280 resources.

281 Higher N inputs were associated with a general increase in N concentrations and
282 N:P ratios in leaves and roots. Higher N inputs may thus increase the uptake of N but can
283 decrease P concentration in roots but not in leaves. Leaf N concentration increased in all
284 studied vegetation types by N fertilization, whereas leaf P concentrations increased in
285 response to N fertilization depending on the vegetation types. Higher foliar N and P
286 concentrations may be associated with larger C-assimilation and growth-rate capacities
287 (Sterner & Elser 2002; Bandau *et al.*, 2015). Increases in N availability at short to medium
288 terms (months to a few years, depending on input rate) enhance the capacity of plants to
289 take up other nutrients such as P (Coblentz *et al.*, 2004; Zhang *et al.*, 2004; Silva *et al.*,
290 2015). Long-term N inputs, however, may eventually decrease growth by causing soil-
291 plant nutrient imbalances (Jonard *et al.*, 2015; Kou *et al.*, 2015), which can in turn
292 decrease foliar P concentrations (Wang *et al.*, 2015). However, when increasing
293 atmospheric [CO₂] coincides with N fertilization higher allocation of N and P to roots is
294 observed suggesting that the lack of P drives the allocation of N mainly to roots to increase
295 the uptake capacity of soil resources. Thus, at the root level, there was a compensatory
296 effect on the response ratio of N and P concentrations when increasing atmospheric [CO₂]
297 were combined with higher N availability. The N-dilution effect frequently observed under
298 increasing atmospheric [CO₂] is thus logically counteracted by the higher nutrient
299 availability in those combined experiments. However, leaves responded differently than
300 roots. Despite the increased N availability, concentrations of N increased only in roots and
301 still declined in the leaves (Fig. 1).

302 The observed differences between N concentrations in leaves and in roots under N
303 fertilization and increasing atmospheric [CO₂] should be embedded into models of N and P

304 cycles for projecting future scenarios of increasing atmospheric [CO₂], climate change and
305 N deposition and they are likely to affect above- and belowground food webs differently.

306 P enrichment is less widespread than N enrichment in natural terrestrial
307 ecosystems (Peñuelas *et al.*, 2013; Wang *et al.*, 2015), but P enrichment does
308 nonetheless occur in specific ecosystems, such as wetlands (Osborne *et al.*, 2014),
309 especially near intensively fertilized agricultural lands or near urban areas (Chen *et al.*,
310 2014; Yan *et al.*, 2016). P fertilization experiments showed increased foliar and root P
311 concentrations, but also higher foliar and root N concentrations. These results are
312 consistent with the fact that most studies have observed increased N-uptake capacity due
313 to increases in P availability (Scheiner *et al.*, 2002; Graciano *et al.*, 2006), despite some
314 studies that did not observe this pattern (Graciano *et al.*, 2006). P fertilization can improve
315 N uptake in different ways. Firstly, P fertilization can increase soil N availability (Reed *et al.*,
316 2007), by increasing the activity of free living N₂-fixing soil microbes (Reed *et al.*, 2007;
317 Alguacil *et al.*, 2010) and epiphytic lichens (Abbasi *et al.*, 2008) and increasing plant
318 nodulation in N₂-fixing plants (Alguacil *et al.*, 2010; Djumaeva *et al.*, 2013; Benner &
319 Vitousek, 2007). The increase in P availability, though, can also increase plant N content
320 by enhancing N-uptake capacity from increases in the density (Sheng *et al.*, 2012) and
321 length (Sheng *et al.*, 2012; Fageria *et al.*, 2014) of fine roots or by improving the efficiency
322 of N resorption (Wienend & Stock, 1995; Ruiz & Romero, 2000). The increases in N
323 concentrations associated with P fertilization can have great consequences in nutrient
324 limited ecosystems where plants have a low investment in reproduction that is strongly
325 related to plant nutrient concentrations (Fujita *et al.*, 2010).

326 Under drought P concentrations increased in roots but not in leaves. This result is
327 consistent with previous experiments showing that plants under drought reallocate more
328 resources from growth and energy metabolism, including the allocation of N and P and

329 other nutrients towards roots for increasing water uptake (Gargallo-Garriga *et al.*, 2014,
330 2015). Changes in P concentrations were thus observed in roots but not in leaves when
331 plants suffered from drought. Increases in root N concentration and decreases in foliar N
332 concentration occurred when increasing atmospheric [CO₂] coincided with N fertilization
333 All these different responses in roots than in leaves will likely have different ecological
334 consequences on above- and belowground processes. For example, they could imply a
335 different impact on root-based relative to leaf-based food webs. Changes in plant-host
336 stoichiometry affect the stoichiometry (Kay *et al.*, 2003; Abbas *et al.*, 2004) and growth and
337 survival (Audusseau *et al.*, 2015) of herbivorous insects. Different changes in foliar and
338 root N and P concentrations can also affect the respective resorption processes differently
339 and thus above- versus belowground food webs (Sterner & Elser, 2002). Similarly, other
340 important ecosystemic processes such as the decomposition of organic matter, food fluxes
341 and generally all nutrient and C cycles can also be affected by changes in N and P
342 concentrations and N:P ratios (Schade *et al.*, 2003; Grover, 2003; Cleveland, 2006; Jonas
343 & Joern, 2008; Sardans *et al.*, 2012b). Some studies of terrestrial ecosystems show that
344 higher N availability increases body N:P ratios and reduces the species diversity of
345 communities, whereas increases in P availability have opposite effects on food resources,
346 with lower N:P ratios increasing the transfer of energy to higher food levels, further
347 favoring larger food webs (Sardans *et al.*, 2012b). All these shifts in the elemental
348 composition of plant organs can thus have important flow-on effects on food webs and
349 ecosystemic structure that could be asymmetrical between above- and belowground
350 ecosystemic compartments in some cases and some ecosystem types by the influence
351 that plant above- and below-ground elemental composition exert on the own food web but
352 also with the food web of the other respective plant compartment (Peñuelas & Matamala
353 1990, Sterner & Elser, 2002). Johnson *et al.* (2013) observed that the relationships
354 between belowground herbivores and root plant nutrient contents also affect aboveground

355 herbivores and ultimately populations of parasites and predators of aboveground
356 herbivores. A global data synthesis showed that the C:N:P ratios of leaves and roots and
357 the respective above- and below-ground food webs are interconnected (Mulder *et al.*,
358 2013).

359 We conclude that increasing atmospheric [CO₂], N and P fertilization and drought
360 typically affect the concentrations of N and P in leaves and roots similarly, but in some
361 cases they affect them differently or even oppositely, such as when increasing
362 atmospheric [CO₂] is combined with N fertilization. The overall analyses of the changes in
363 N and P concentrations in leaves and roots in response to the studied global change
364 drivers and their combinations suggested that an equilibrium of the concentration effect (in
365 the case of drought or N and P fertilization) and the dilution effect (in the case of elevated
366 [CO₂]) with the increasing N and P allocation to roots to enhance the uptake of other soil
367 resources when N and/or P availability increased. Then the results suggest that when
368 several different resources are provided at once enhancing the potential for plant
369 production capacity, more sources are allocated to roots to improve the uptake of
370 complementary resources to those enhanced by environmental changes. This occurs for N
371 and P under drought, for N under N fertilization, for P under P fertilization, for N and P
372 under N and P fertilization and for N when N and CO₂ increase. This is not observed under
373 elevated CO₂. The dilution effect together with the necessity to maintain N in
374 photosynthetic tissues to take profit of higher CO₂ could explain it. The results also
375 suggest that above- and belowground processes could sometimes be affected differently
376 and even oppositely by the drivers of Global Change. Such differential shifts in above-
377 versus below-ground elemental composition can change ecosystem function, for example
378 by affecting the corresponding food webs differently. Furthermore, several models of
379 carbon cycles in terrestrial ecosystems have been developed to determine the

380 mechanisms behind the shoot/root allocation of carbon (Wilson, 1988; Cannell & Dewar,
381 1994). The results suggest the need to also incorporate flexible allocations of N and P
382 concentrations to different plant organs in the distinct biomes into models projecting effects
383 under different scenarios of climate and fertility.

384

385 **Acknowledgements**

386 The authors would like to acknowledge the financial support from the European Research Council
387 Synergy grant ERC-SyG-2013-610028 IMBALANCE-P, the Spanish Government grant CGL2013-
388 48074-P and the Catalan Government grant SGR 2014-274.

389

390 **References**

- 391 Abbas M, Klein AM, Ebeling A, Oelmann Y, Ptanick R, Weisserg WW, Hillebrand H (2014)
392 Plant diversity effects on pollinating and herbivorous insects can be linked to plant
393 stoichiometry. *Basic Applied Ecology*, **15**, 169-178.
- 394 Abbasi MK, Majeed A, Sadiq A, Khan SR (2008) Application of *Bradyrhizobium japonicum*
395 and phosphorus fertilization improved growth, yield and nodulation of soybean in the
396 sub-humid hilly region of Azad Jammu and Kashmir, Pakistan. *Plant Production*
397 *Science*, **11**, 368-376.
- 398 Akmal M, Schelberg J, Khattak RA (2010) Biomass allocation and nitrogen distribution in
399 ryegrass under water and nitrogen supplies. *Journal of Plant Nutrition*, **35**, 1777-
400 1788.
- 401 Alguacil MD, Lozano Z, Campoy MJ, Roldan (2010) A Phosphorus fertilization
402 management modifies the biodiversity of AM fungi in a tropical savanna forage
403 system. *Soil Biology and Biochemistry*, **42**, 1114-1122.
- 404 Audusseau H, Kolb G, Janz N (2015) Plant fertilization interacts with life history: variation
405 in stoichiometry and performance in Nettle-Feeding butterflies. *Plos One*, **10**,
406 e0124616.
- 407 Bandau F, Decker VHG, Gundale MJ, Albrechtsen BR (2015) Genotypic tannin levels in
408 *Populus tremula* impact the way nitrogen enrichment affects growth and allocation
409 responses for some traits and not for others. *PlosOne*, **10**, e0140971.
- 410 Benner JW, Vitousek PM (2007) Development of a diverse epiphyte community in
411 response to phosphorus fertilization. *Ecology Letters*, **10**, 628-636.
- 412 Chen FS, Yavitt J, Hu XF (2014) Phosphorus enrichment helps increase soil carbon
413 mineralization in vegetation along an urban-to-rural gradient, Nanchang, China.
414 *Applied Soil Ecology*, **75**, 181-188.
- 415 Churkina G, Brovkin V, von Bloh W, Trusilova K, Jung M, Dentener F (2009) Synergy of
416 rising nitrogen depositions and atmospheric CO₂ on land carbon uptake moderately
417 offsets global warming. *Global Biogeochemical Cycles*, **23**, GB4027.
- 418 Cleveland CC, Reed SC, Townsend AR (2006) Nutrient regulation of organic matter
419 decomposition in a tropical rain forest. *Ecology*, **87**, 492-503.

- 420 Coblenz WK, Daniels MB, Gunsaulis JL *et al.* (2004) Effects of nitrogen fertilization on
 421 phosphorus uptake in Bermudagrass forage grown on high soil-test phosphorus
 422 sites. *Profesional Animal Science*, **20**, 146-154.
- 423 Crous JW, Morris AR, Scholes MC. (2011) Changes in topsoil, standing litter and tree
 424 nutrient content of a *Pinus patula* plantation after phosphorus and potassium
 425 fertilization. *European Journal of Forest Research*, **130**, 277-292.
- 426 Deng Q, Hui D, Luo Y, Elser JJ, Wang YP, Loladze I, Zhang Q, Dennis S (2015) Down-
 427 regulation of tissue N:P ratios in terrestrial plants by elevated CO₂. *Ecology*, **96**,
 428 3354-3362.
- 429 Dieleman WIJ, Vicca S, Dijkstra FA *et al.* (2012) Simple additive effects are rare: a
 430 quantitative review of plant biomass and soil process responses to combined
 431 manipulations of CO₂ and temperature. *Global Change Biology*, **18**, 2681-2693.
- 432 Djumaeva D, Lamers JPA, Khamzina A, Vlek PL (2013) The benefits of phosphorus
 433 fertilization of trees grown on salinized croplands in the lower reaches of Amu Darya,
 434 Uzbekistan. *Agroforestry Systems*, **87**, 555-569.
- 435 Duval BD, Blankinship Dijkstra P, Hungate BA (2012) CO₂ effects on plant nutrient
 436 concentration depend on plant functional group and available nitrogen: a meta-
 437 analysis. *Plant Ecology*, **213**, 505-521.
- 438 Fageria NK, Moreira A, Moraes LAC, Moraes MF (2014) Root growth, Nutrient uptake, and
 439 Nutrient-use-efficiency by roots of tropical legume cover crops as influenced by
 440 phosphorus fertilization. *Commun. Soil Science and Plant Analysis*, **45**, 555-569.
- 441 Fujita Y, Olde Venterink H, van Bodegom PM *et al.* (2010) Low investment in sexual
 442 reproduction threatens plants adapted to phosphorus limitation. *Nature*, **505**, 82-86.
- 443 Gargallo-Garriga A, Sardans J, Pérez-Trujillo M *et al.*, (2014) Opposite metabolic
 444 responses of shoots and roots to drought. *Scientific Reports*, **4**, 6829.
- 445 Gargallo-Garriga A, Sardans J, Ovaric M *et al.* (2015) Warming differently influences the
 446 effects of drought on Stoichiometry and metabolomics in shoots and roots. *New*
 447 *Phytologist*, **207**, 591-603.
- 448 Graciano C, Goya JF, Frangi JL, Guiamet JJ (2006) Fertilization with phosphorus
 449 increases soil nitrogen absorption in Young plants of *Eucalyptus grandis*. *Forest*
 450 *Ecology and Management*, **236**, 202-210.
- 451 Grover P (2003). The impact of variable stoichiometry on predator-prey interactions: a
 452 multinutrient approach. *American Naturalist* **162**, 29-43
- 453 Hedges LV, Gurevitch J. Curtis PS (1999) The meta-analysis of response ratios in
 454 experimental ecology. *Ecology*, **80**, 1150-1156.
- 455 Jonard M, Fuerst A, Verstraeten A *et al.* (2015) Tree mineral nutrition is deteriorating in
 456 Europe. *Global Change Biology*, **21**, 418-430
- 457 Jonas JL, Joern A (2008) Host-plant quality alters grass/forb consumption by a mixed-
 458 feeding insect herbivore, *Melanoplus bivittatus* (Orthoptera: Acrididae). *Ecological*
 459 *Entomology*, **33**, 546-554.
- 460 Johnson SN, Mitchell C, McNicol JW, Thompson J, Karley AJ (2013) Downstairs drovers-
 461 root herbivores shape communities of above-ground herbivores and natural enemies
 462 via changes in plant nutrients. *Journal of Animal Ecology*, **82**, 1021-1030.
- 463 Kay A, Schade J, Novotny A, Hernández D, Ogdahl M, Sterner RW, Hobbie, SE (2003)
 464 Body size and N:P stoichiometry of insect herbivores vary with host plant chemistry.
 465 *Ecological Society American Annual Meeting Abstracts*, **88**, 176.
- 466 Kerkhoff AJ, Enquist BJ, Elser JJ, Fagan WF (2005) Plant allometry, stoichiometry and the
 467 temperature-dependence of primary productivity. *Global Ecology and Biogeography*,
 468 **14**, 585-598.
- 469 Kerkhoff AJ, Enquist BJ (2006) Ecosystem allometry: the scaling of nutrient stocks and
 470 primary productivity across plant communities. *Ecology Letters*, **9**, 419-427.

- 471 Kou L, Guo DL, Yang H, Gao WL, Li SG (2015) Growth, morphological traits and
472 mycorrhizal colonization of fine roots respond differently to nitrogen addition in a
473 slash pine plantation in subtropical China. *Plant Soil*, **391**, 207-218.
- 474 Leuzinger S, Luo Y, Beier C, Dieleman W, Vicca V, Körner CD (2011) global change
475 experiments overestimate impacts on terrestrial ecosystems? *Trends in Ecology and*
476 *Evolution*, **26**, 236–241.
- 477 Li J, Zhou JM, Duan ZQ, Du CW, Wang HY (2007) Effect of CO₂ enrichment on the growth
478 and nutrient uptake of tomato seedlings. *Pedosphere*, **17**, 343-351.
- 479 Liu XJ, Zhang Y, Han WX, Tang AH, Shen JL, Cui ZL, Vitousek P, Erisman JW, Goulding
480 K, *et al.* (2013) Enhanced nitrogen deposition over China. *Nature*, **494**, 459-462.
- 481 Liu J, Huang W, Zhou G, Zhang D, Liu S, Li Y (2013) Nitrogen to phosphorus ratios of
482 three species response to elevated carbon dioxide and nitrogen addition in
483 subtropical forests. *Global Change Biology*, **19**, 208-216.
- 484 Meyerholds J, Zaehle S (2015) The role of stoichiometry flexibility in modelling forest
485 ecosystem responses to nitrogen fertilization. *New Phytologist*, **208**, 1042-1055.
- 486 Mulder C, Ahrestani FS, Bahn M *et al.* (2013) Connecting the green and brown worlds:
487 allometric and stoichiometric predictability of above- and below-ground networks.
488 *Advances in Ecology Research*, **49**, 69-175.
- 489 Osborne TZ, Reddy KR, Ellis LR, Aumen NG, Surratt DD, Zimmerman MS, Sadle J (2014)
490 Evidence of recent phosphorus enrichment in surface soils of Taylor slough and
491 Northeast everglades National Park. *Wetlands*, **34**, S37-S45.
- 492 Peñuelas J, Poulter B, Sardans J *et al.*, (2013) Human-induced nitrogen-phosphorus
493 imbalances alter natural and managed ecosystems across the globe. *Nature*
494 *Communications*, **42934**, 1-10
- 495 Peñuelas J, Matamala R (1990) Changes in N and S leaf content, stomatal density and
496 specific leaf area of 14 plant species during the last three centuries of CO₂ increase.
497 *Journal of Experimental Botany*, **41**, 1119-1124.
- 498 R Core Team (2015) R: A language and environment for statistical computing. R
499 Foundation for Statistical Computing, Vienna, Austria. URL [https://www.R-](https://www.R-project.org/)
500 [project.org/](https://www.R-project.org/).
- 501 Reed SC, Seastedt TR, Mann CM, Suding KN, Townsend AR, Cherwin KL (2007)
502 Phosphorus fertilization stimulates nitrogen fixation and increases inorganic nitrogen
503 concentrations in a restored prairie. *Applied Soil Ecology*, **36**, 238-242.
- 504 Reich PB, Hobbie SE (2012) Decade-long soil nitrogen constraint on the CO₂ fertilization
505 of plant biomass. *Nature Climate Change*, **3**, 278-282.
- 506 Ruiz JM, Romero L (2000) Nitrogen metabolism and yield response of cucumber (*Cucumis*
507 *sativus* L cv Brunex) plants to phosphorus fertilization. *Journal of Science Food*
508 *Agriculture*, **80**, 2069-2073.
- 509 Sardans J, Peñuelas J (2012) The role of plants in the effects of global change on Nutrient
510 availability and nutrient stoichiometry in the Plant-Soil system. *Plant Physiology*, **160**,
511 1741-1761.
- 512 Sardans J, Peñuelas J, Ogaya R (2008) Drought induced changes in C and N
513 stoichiometry in a *Quercus ilex* Mediterranean Forest. *Forest Science*, **54**, 513-522.
- 514 Sardans J, Peñuelas J, Prieto P, Estiarte M (2008) Drought and warming induced changes
515 in P and K concentration and accumulation in plant biomass and soil in a
516 Mediterranean shrubland. *Plant Soil*, **306**, 261-271.
- 517 Sardans J, Rivas-Ubach A, Peñuelas J (2012b) The C:N:P Stoichiometry of organisms
518 and Ecosystems in a changing world: A review and perspectives. *Perspectives in*
519 *Plant Ecology, Evolution and Systematics*, **14**, 33-47.

- 520 Sardans J, Bartrons M, Margalef O. *et al.* (2017) Plant Invasion is associated with higher
521 plant-soil nutrient concentrations in nutrient-poor environments. *Global Change*
522 *Biology* **23**, 1282-1291.
- 523 Schade JD, Kyle M, Hobbie SE, Fagan WF, Elser JJ (2003) Stoichiometric tracking of soil
524 nutrients by a desert insect herbivore. *Ecological Letters*, **6**, 96-101.
- 525 Scheiner JD, Gutierrez-Boem FH, Lavado RS (2002) Sunflower nitrogen requirement and
526 ¹⁵N fertilizer recovery in western Pampas, Argentina. *European Journal of*
527 *Agronomy*, **17**, 73-79.
- 528 Sheng M, Lalande R, Hamel C, Ziadi N, Shi YC (2012) Growth of corn roots and
529 associated arbuscular mycorrhizae are affected by long-term tillage and phosphorus
530 fertilization. *Agronomy Journal*, **104**, 1672-1678.
- 531 Silva LCR, Gomez-Guerrero A, Doane T, Howarth WR (2015) Isotopic and nutritional
532 evidence for species- and site-specific responses to N deposition and elevated CO₂
533 in temperate forests. *Journal of Geophysical Research Biogeosciences*, **120**, 1110-
534 1123.
- 535 Sterner RW, Elser JJ (2002) Ecological Stoichiometry: The Biology of Elements from
536 Molecules to the Biosphere. Princeton University Press.
- 537 Thomas RQ, Bonan GB, Goodale CL (2013) Insights into mechanisms governing forest
538 carbon response to nitrogen deposition: a model-data comparison using observed
539 responses to nitrogen addition. *Biogeosciences*, **10**, 3869-3887.
- 540 Van der Velde M, Folberth C, Balkovic J *et al.*, (2014) African crop yield reductions due to
541 increasingly unbalanced nitrogen and phosphorus consumption. *Global Change*
542 *Biology*, **20**, 1278-1288.
- 543 Viechtbauer WR (2012) package: *Metafor*. Available at website [http://](http://www.metafor-project.org/doku.php)
544 www.metafor-project.org/doku.php.
- 545 Wang R, Balkanski Y, Boucher O, Ciais P, Peñuelas J, Tao S (2015) Significant
546 contribution of combustion-related emissions to the atmospheric phosphorus budget.
547 *Nature Geosciences*, **8**, 48-54.
- 548 Wienend KT, Stock WD (1995) Long-term phosphorus fertilization effects on the litter
549 dynamics of an age sequence of *Pinus-elliottii* plantations in the Southern Cape of
550 South-Africa. *Forest Ecology Management*, **75**, 135-146
- 551 Wilson JB (1988) Shoot Competition and Root Competition. *J. Appl. Ecol.* **25**, 279–296.
- 552 Cannell MGR, Dewar RC (1994) Carbon Allocation in Trees: a Review of Concepts for
553 Modelling. *Advances in Ecological Research*, **25**, 59–104.
- 554 Yan Z, Han W, Peñuelas J, Sardans J, Elser JJ, Du E, Reich PB, Fang J (2016)
555 Phosphorus accumulates faster than nitrogen globally in freshwater ecosystems
556 under anthropogenic impacts. *Ecology Letters*, **19**, 1237-1246.
- 557 Yuan ZY, Chen HYH (2015) Decoupling of nitrogen and phosphorus in terrestrial plants
558 associated with global changes. *Nature Climate Change*, **5**, 465-469.
- 559 Zhang F, Kang S, Zhang J, Zhang R, Li F (2004) Nitrogen fertilization on uptake of soil
560 inorganic phosphorus fractions in the wheat root zone. *Soil Science Society*
561 *American Journal*, **68**, 1890-1895.

562

563

564

565

566 **Additional information**

567 Supplementary information for only on-line publication is added

568

569 **Competing interests**

570 The authors declare no competing financial interests.

571

572 Table 1. Summary of the results of the meta-analyses for the concentrations of N and P
 573 and the corresponding ratios in response to the Global-Change drivers.

Global-Change driver	[N]		[P]	
	Leaves	Roots	Leaves	Roots
Elevated [CO ₂]	↓	↓	=	=
N fertilization	↑	↑	=	↓
P fertilization	NA	NA	↑	↑
Drought	↑	↑	=	↑
[CO ₂] + N fertilization	↓ ^b	↑ ^a	=	↑
N + P fertilization	↑	↑	↑	↑

574 ↑, significant ($P < 0.05$) increases in meta-analysis; ↓, significant decreases ($P < 0.05$) in
 575 meta-analysis; =, no change in meta-analysis; NA, not available. Different letters indicate
 576 significant differences ($P < 0.05$) in response ratios between leaves and roots. Different
 577 letters indicate significant differences ($P < 0.05$) in response ratios between photosynthetic
 578 tissues and roots only when overall data was analyzed including studies with data of leaf
 579 and/or roots.

580

581

582

583

584

585

586

587

588

589

590

591 Figure Captions

592 Figure 1. Natural log response ratios of N and P concentrations and N:P ratios in leaves
593 (blue) and roots (grey) due to elevated [CO₂], drought, N fertilization, P fertilization,
594 elevated [CO₂] + N enrichment, N + P enrichment and elevated [CO₂] + drought. The
595 meta-analyses were conducted only with studies that provided data of leaves and roots
596 measured simultaneously in the same plants. Error bars indicate the mean response ratio
597 with 95% confidence intervals. The numbers between brackets indicate the number of
598 reports and observations, respectively, used in the meta-analysis of each variable. (***),
599 (**), and (*) indicate significant differences at ($P < 0.001$), ($P < 0.01$) and ($P < 0.05$)
600 respectively, between treated plants compared with the control plants. Positive response
601 ratios mean that the corresponding global change drivers increased the value of the
602 variable with respect to the control conditions.

603 Figure 2. Sensitivity of the natural log response ratios of N and P concentrations (by ppm
604 of elevated [CO₂]) in leaves and roots to elevated [CO₂] and sensitivity of the natural log
605 response ratios of N and P concentrations (by kg ha⁻¹ of N enhancement) in leaves and
606 roots to N fertilization obtained in studies that provided data of leaves and roots of the
607 same plants. Error bars indicate the 95% confidence intervals for the mean response ratio.
608 The numbers between brackets indicate the number of reports and observations,
609 respectively, used in the meta-analysis of each variable. (***), (**), and (*) indicate
610 significant differences, at ($P < 0.001$), ($P < 0.01$) and ($P < 0.05$) respectively, between treated
611 plants compared with the control plants. Positive response ratios mean that the
612 corresponding global change drivers increased the value of the variable with respect to the
613 control conditions.

614 Figure 3. Natural log response ratios due to elevated [CO₂] and to N fertilization on N
615 concentrations in leaves and roots in different vegetation types obtained in studies that
616 provided data of leaves and roots of the same plants. Error bars indicate the 95%
617 confidence intervals for the mean response ratio. The numbers between brackets indicate
618 the number of reports and observations, respectively, used in the meta-analysis of each
619 variable. (***), (**), and (*) indicate significant differences, at ($P < 0.001$), ($P < 0.01$) and
620 ($P < 0.05$) respectively, between treated plants compared with the control plants. Positive
621 response ratios mean that the corresponding global change drivers increased the value of
622 the variable with respect to the control conditions.

623

624

625

626

627




Figure 1. Natural log response ratios of N and P concentrations and N:P ratios in leaves (blue) and roots (grey) due to elevated [CO₂], drought, N fertilization, P fertilization, elevated [CO₂] + N enrichment, N + P enrichment and elevated [CO₂] + drought. The meta-analyses were conducted only with studies that provided data of leaves and roots measured simultaneously in the same plants. Error bars indicate the mean response ratio with 95% confidence intervals. The numbers between brackets indicate the number of reports and observations, respectively, used in the meta-analysis of each variable. (***), (**), and (*) indicate significant differences at (P<0.001), (P<0.01) and (P<0.05) respectively, between treated plants compared with the control plants. Positive response ratios mean that the corresponding global change drivers increased the value of the variable with respect to the control conditions.




Figure 2. Sensitivity of the natural log response ratios of N and P concentrations (by ppm of elevated [CO₂]) in leaves and roots to elevated [CO₂] and sensitivity of the natural log response ratios of N and P concentrations (by kg ha⁻¹ of N enhancement) in leaves and roots to N fertilization obtained in studies that provided data of leaves and roots of the same plants. Error bars indicate the 95% confidence intervals for the mean response ratio. The numbers between brackets indicate the number of reports and observations, respectively, used in the meta-analysis of each variable. (***), (**), and (*) indicate significant differences, at (P<0.001), (P<0.01) and (P<0.05) respectively, between treated plants compared with the control plants. Positive response ratios mean that the corresponding global change drivers increased the value of the variable with respect to the control conditions.

338x451mm (96 x 96 DPI)




Figure 3. Natural log response ratios due to elevated [CO₂] and to N fertilization on N concentrations in leaves and roots in different vegetation types obtained in studies that provided data of leaves and roots of the same plants. Error bars indicate the 95% confidence intervals for the mean response ratio. The numbers between brackets indicate the number of reports and observations, respectively, used in the meta-analysis of each variable. (***), (**), and (*) indicate significant differences, at (P < 0.001), (P < 0.01) and (P < 0.05) respectively, between treated plants compared with the control plants. Positive response ratios mean that the corresponding global change drivers increased the value of the variable with respect to the control conditions.