

The Sensitivity of Southeast Pacific Heat Distribution to Local and Remote Changes in Ocean Properties

Daniel C. Jones, Emma Boland, Andrew J. S. Meijers, Gael Forget, Simon Josey, Jean-Baptiste Sallée, Emily Shuckburgh

To cite this version:

Daniel C. Jones, Emma Boland, Andrew J. S. Meijers, Gael Forget, Simon Josey, et al.. The Sensitivity of Southeast Pacific Heat Distribution to Local and Remote Changes in Ocean Properties. Journal of Physical Oceanography, 2020, 50 (3), pp.773-790. $10.1175/jpo-d-19-0155.1$. hal-02904199

HAL Id: hal-02904199 <https://hal.science/hal-02904199>

Submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

¹³ *Corresponding author address: D.C. Jones, British Antarctic Survey, NERC, UKRI, Cambridge,

UK. 14

¹⁵ E-mail: dannes@bas.ac.uk

ABSTRACT

The Southern Ocean features ventilation pathways that transport surface waters into the subsurface thermocline on timescales of decades to centuries, sequestering anomalies of heat and carbon away from the atmosphere and thereby regulating the rate of surface warming. Despite its importance for climate sensitivity, the factors that control the distribution of heat along these pathways are not well understood. In this study, we use an observationallyconstrained, physically consistent global ocean model to examine the sensitivity of heat distribution in the recently ventilated subsurface Pacific (RVP) sector of the Southern Ocean to changes in ocean temperature and salinity. First, we define the RVP using numerical passive tracer release experiments that highlight the ventilation pathways. Next, we use an ensemble of adjoint sensitivity experiments to quantify the sensitivity of the RVP heat content to changes in ocean temperature and salinity. In terms of sensitivities to surface ocean properties, we find that RVP heat content is most sensitive to anomalies along the Antarctic Circumpolar Current (ACC), upstream of the subduction hotspots. In terms of sensitivities to subsurface ocean properties, we find that RVP heat content is most sensitive to basin-scale changes in the subtropical Pacific Ocean, around the same latitudes as the RVP. Despite the localized nature of mode water subduction hotspots, changes in basin-scale density gradients are an important controlling factor on heat distribution in the Southeast Pacific. 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36

37 1. Introduction

³⁸ The Southern Ocean (SO) is an important region for the uptake, storage, and transport of heat and ³⁹ carbon; SO upwelling brings up water that has not seen the surface for centuries or even millennia ⁴⁰ and is thus far out of equilibrium with surface temperature and atmospheric carbon (Sabine et al. ⁴¹ 2004; Sarmiento et al. 2004; Lenton and Matear 2007; Ito et al. 2010; Talley 2013; Roemmich et al. ⁴² 2015; Armour et al. 2016). Over the period 1861-2005, the SO, defined as the ocean south of 30[°]S, ⁴³ is estimated to account for $43\% \pm 3\%$ of anthropogenic carbon dioxide uptake and $75\% \pm 22\%$ of 44 heat uptake, despite only occupying 30% of global surface ocean area (Frölicher et al. 2015). The ⁴⁵ SO's ability to absorb and transport this disproportionately high fraction of heat and carbon comes ⁴⁶ in part from a balance between powerful overlying westerly winds, strong buoyancy fluxes, and 47 internal dynamics that maintains steeply tilted surfaces of constant density and localized pools ⁴⁸ of weakly stratified water (Russell et al. 2006). These pools of weakly stratified water, referred ⁴⁹ to collectively as Subantarctic Mode Water (SAMW), are refreshed by localized wintertime deep ⁵⁰ convection that occurs just north of the Antarctic Circumpolar Current (ACC) (Speer et al. 2000; $_{51}$ Hanawa and Talley 2001; Lumpkin and Speer 2007; Talley 2008; Herraiz-Borreguero and Rintoul ⁵² 2011; Speer and Forget 2013). The processes of lateral induction, eddy-induced transport, and ₅₃ advection via the mean flow subduct the SAMW into the subsurface ocean at a rate that varies ⁵⁴ across interannual and decadal timescales (Karsten and Marshall 2002; Sallée et al. 2010b; Sallée 55 and Rintoul 2011; Sallée et al. 2012; Liu and Huang 2012). Together with the denser and relatively ⁵⁶ fresher Antarctic Intermediate Water (AAIW), SAMW ventilates the subtropical thermocline on σ timescales of decades to centuries (Iudicone et al. 2007; Sallée et al. 2010a; Cerovecki et al. 2013; ⁵⁸ Jones et al. 2016). Here, ventilation refers to the broad set of processes by which atmospheric ⁵⁹ and surface ocean properties are able to affect subsurface ocean properties. Ventilation is driven

⁶⁰ in part by the formation, subduction, and destruction of water masses and can be considered a ⁶¹ consequence of the ocean's global overturning circulation (Marshall and Speer 2012; Cerovecki 62 and Mazloff 2015).

⁶³ SAMW and AAIW both display complex patterns of decadal variability. In the Western Pacific, ⁶⁴ decadal-scale surface warming and/or freshening at high-latitude SAMW and AAIW subduction ⁶⁵ sites can lead to cooling and freshening along isopycnals in the subsurface (Johnson and Orsi ⁶⁶ 1997). Decadal trends in SAMW temperature and salinity display complex spatial variability; in ⁶⁷ recent decades, both the SAMW and AAIW have become warmer and saltier in the South Atlantic ⁶⁸ but cooler and fresher in the South Atlantic (Katsumata and Fukasawa 2011). The AAIW core (its ⁶⁹ salinity minimum) has warmed and shoaled overall, with small salinity trends whose sign varies ⁷⁰ by region (Schmidtko and Johnson 2012). Decadal trends of SAMW and AAIW properties are τ_1 consistent with an amplification of the global hydrological cycle and broad-scale surface warming π (Helm et al. 2010; Durack and Wijffels 2010). The spatial and temporal complexity of these 73 SAMW and AAIW trends suggests that we will need a full four-dimensional description of the 74 sensitivity of ocean structure in order to make robust projections of potential future ocean states.

 75 Regional differences in subduction, ventilation, and the properties of the ventilated subsurface π can impact (i.) the sequestration of anthropogenic heat and carbon into the subsurface thermocline π and (ii.) the supply of nutrients to low latitudes via the overturning circulation (Sarmiento et al. ⁷⁸ 2004; Sabine et al. 2004; Khatiwala et al. 2009; Ito et al. 2010; Roemmich et al. 2015). The ⁷⁹ regionally specific nature of subduction and ventilation, and how those localized characteristics ⁸⁰ affect the distribution of heat, carbon, and nutrients in the ocean, is an active area of oceanographic 81 study (Cerovecki et al. 2013; Jones et al. 2016). Despite these efforts, we still have relatively little ⁸² knowledge on how regional variations in ocean state variables (e.g. temperature, salinity) and ⁸³ surface forcing can ultimately impact the properties of the ventilated subsurface. Advancements in ⁸⁴ this area may be especially helpful for improving projections of future ocean states, as changes in ⁸⁵ the Southern Ocean forcing-subduction-ventilation mechanism are expected to have a considerable ⁸⁶ impact on future climate (Cessi and Otheguy 2003; Downes et al. 2009; Lovenduski and Ito 2009; Morrison et al. 2011; Sallée et al. 2012).

 In order to quantify how regional variations in ocean state variables may affect the heat distribu- tion in the ventilated subsurface ocean, we perform a set of adjoint sensitivity experiments using an observationally-constrained, physically consistent state estimate. First, we identify the recently ventilated subsurface ocean using a combination of physical state variables (e.g. potential vor- ticity, density, mixed layer depth) and numerical passive tracer distributions that track ventilation pathways from the surface into the subsurface. Our method for identifying the recently ventilated subsurface treats ventilation as a general process that may affect many different water masses, in contrast to approaches that consider one particular water mass (e.g. SAMW). We focus our atten- tion on the Eastern Pacific because it features an especially efficient export pathway of water from the surface ocean into the subsurface thermocline, as measured by numerical passive tracer exper- iments (Jones et al. 2016). For convenience, we refer to the recently ventilated subsurface Eastern Pacific as the RVP and the heat content of the RVP as RVPh. Although there is overlap between the two, we note that the RVP is more general than the region occupied by SAMW or any other particular water mass. Once we have identified the RVP, we perform a set of adjoint sensitivity experiments using, for consistency, the same numerical model setup that we used for the passive tracer experiments. Our adjoint model produces linear, time-evolving sensitivity maps, allowing us to identify locations and timescales where hypothetical anomalies could potentially impact the heat distribution in the ventilated subsurface, as revealed by the heat content of the fixed-volume RVP. Although we do not use this framework here, the sensitivity fields can roughly be interpreted as optimal linear perturbations, i.e. perturbations with the largest possible impact on RVPh, within the linear framework of the adjoint model (see a related but not identical example in Sevellec and ´ Fedorov (2015)).

 This paper is structured as follows: in section 2, we describe the ECCOv4 model setup, the design of our numerical tracer release experiments, and the design of our adjoint sensitivity ex- periments. In section 3, we examine the results of the tracer experiments in order to define the RVP, which is the control volume for our adjoint sensitivity experiments. In section 4, we examine the time-evolving sensitivities of RVPh to changes in temperature and salinity, decomposed into changes at constant density (i.e. kinematic sensitivities) and changes with varying density (i.e. dynamic sensitivities). In section 5, we discuss factors that should be considered when comparing our results with other studies. Finally, in section 6, we summarize our results and conclusions.

2. Methods

¹¹⁹ In this section, we describe the observationally-constrained global ocean model setup (subsec- tion a), the design of our numerical passive tracer experiments used to define the RVP (subsection b), and the design of our adjoint sensitivity experiments (subsection c).

a. Global ocean model setup

¹²³ We use the modelling setup associated with ECCOv4 (release 2, hereafter ECCOv4-r2 or just ECCOv4). ECCOv4 is a *state estimate*, meaning that it has been adjusted to minimize the mis- fits between the model state and a suite of observations from various sources over the time period 1992-2011 (e.g. Argo float temperature and salinity profiles, ship hydrography, satellite altimetry). ¹²⁷ The model's initial conditions, surface forcing fields, and mixing parameters were iteratively ad-128 justed in order to reduce model-data misfit. Because no artificial sources or sinks of heat were used in the ocean subsurface, ECCOv4 features closed budgets of heat, salt, and volume. The model

 setup is available for download on GitHub.com (https://github.com/gaelforget/ECCOv4) as an instance of the open source MIT general circulation model (MITgcm, http://mitgcm.org/, ¹³² also available on GitHub). We briefly describe the relevant features of the ECCOv4 setup below; ¹³³ a more thorough description is available in Forget et al. (2015a) and references therein.

 $ECCOv4$ is a global ocean model that uses a Lat-Lon-Cap (LLC) grid. Its horizontal grid spac- ing ranges from around 40-50 km at high latitudes up to roughly 110 km at the equator. It features parameterized diffusion, including both diapycnal and isopycnal components, simple convective ¹³⁷ adjustment, and the Gaspar-Gregoris-Lefevre (GGL) mixed layer turbulence closure scheme (Gas- par et al. 1990). To represent the along-isopycnal effect of unresolved eddies, Forget et al. (2015a) used a bolus transport parameterization (Gent and Mcwilliams 1990, hereafter GM). Although the horizontal resolution of ECCOv4 is relatively coarse (roughly 1°), its mixing properties are in ¹⁴¹ good agreement with observations, as indicated by a comparison of simulated ECCOv4 oxygen distribution and World Ocean Atlas 2013 (Forget et al. 2015b, Figure 4). This agreement is thanks in part to the use of optimized, spatially-varying turbulent transport coefficients. ECCOv4 features fully interactive dynamic sea ice, so that buoyancy and mass fluxes at the sea surface are calculated based on the thermodynamic balance of Losch et al. (2010). Open ocean rain, evaporation, and runoff simply carry (advect through the free surface) the local SST and a salinity value of zero, and runoff is provided by a monthly climatology (Fekete et al. 2002). ECCOv4 calculates buoyancy, radiative, and mass fluxes using the bulk formulae of Large and Yeager (2009) with 6-hourly ERA- Interim re-analysis fields (Dee et al. 2011) as a "first guess" for the forcing fields. Specifically, we use wind stress, 2 m air temperature, 2 m specific humidity, wind speed, downward longwave radiation, and downward shortwave radiation as model inputs. These atmospheric state fields have been iteratively adjusted by the state estimation process in order to minimize model-data misfits. The ECCOv4-r2 setup that we use here does *not* use surface salinity restoring. For additional val-

 idation information, see Forget et al. (2015a). For calculations relevant to mode water formation 155 and distribution in ECCOv4, see Speer and Forget (2013).

b. Design of the numerical passive tracer release experiments

 Generally, on a selected timescale (e.g. 1 year, 10 years), we can consider the volume of the ocean that has been affected by near-surface properties (e.g. the temperature and salinity charac- teristics of the mode water formation regions) as having been ventilated via advection, diffusion, and mixing. In terms of an operational definition, the ventilated subsurface ocean can be identified using physical state variables, tracer distributions, or a combination of the two. Luyten et al. (1983) showed that, in a simple, inviscid, analytical model, pathways of potential vorticity (PV) can be used to separate the ventilated thermocline from the unventilated thermocline and also to separate regions of weak stratification from regions of strong stratification. This result helps justify the use of potential vorticity as a "dynamical tracer" that indicates thermocline ventilation pathways. Alternatively, one can derive ventilation pathways using the observed distributions of tracers such as tritium, helium-3 and chlorofluorocarbons, which are carried into the subsurface by both advec- tion along ventilation pathways and by isopycnal and diapycnal mixing (Musgrave 1990; Speer and Tziperman 1992; Primeau and Holzer 2006). In the subsurface, diapycnal mixing contributes to the erosion of water masses by homogenizing their properties across density surfaces (Tross- man et al. 2012). In combination with PV and tracer considerations, the ventilated subsurface is generally considered to be located below the mixed layer, isolated from immediate contact with the surface. In this study, we use a combination of physical state variables (e.g. PV, mixed layer depth, density) and numerical passive tracer release experiments to identify the recently ventilated subsurface ocean in ECCOv4-r2, described in detail below.

 First, we use stratification (PV) and mixed layer depth (MLD) to identify the Southern Ocean regions that tend to ventilate the subsurface thermocline (Figure 1). We calculate $PV = f\rho_0^{-1}$ ¹⁷⁷ gions that tend to ventilate the subsurface thermocline (Figure 1). We calculate $PV = f\rho_0^{-1} d\rho/dz$, ¹⁷⁸ where $f = 2\Omega \sin(\phi)$ is the Coriolis parameter and ρ_0 is the reference density, and we calculate mixed layer depth using a criteria based on the density change associated with a temperature vari-¹⁸⁰ ation of $\Delta T = 0.8$ °C (Kara et al. 2000). For each year between 1996 and 2001, we calculate the ¹⁸¹ June-July-August (JJA) mean PV and both the JJA mean and annual maximum MLD, and we use these values with a set of criteria to construct a "mixed layer mask" for that year. Specifically, for a grid cell to be included in the mixed layer mask for a given year, it must:

1. be within the annual maximum mixed layer and

2. be at a latitude and longitude where one or both of the following conditions are satisfied:

(a) the JJA mean $|PV|$ at the base of the JJA mean mixed layer must be less than $10^{-10.5}$ $1/\text{ms}$, or

(b) the JJA mean mixed layer must be greater than 300 m.

 These relatively weak conditions ensure that we target likely ventilation pathways without being overly prescriptive.

¹⁹¹ We perform six different tracer release experiments (one for each starting year between 1996 and 2001). For each experiment, we initialize passive tracer within the corresponding year's mixed layer mask on 1 January; our method of initializing tracer above the maximum mixed layer for the release year ensures that variation associated with seasonal release timings are minimal. We ¹⁹⁵ integrate the tracer equations in "online" mode for 10 years, i.e. we explicitly solve the momentum, buoyancy, and tracer equations at the same time. Note that although the bulk of the tracer used to define the RVP originates in the Pacific, a smaller fraction also comes from the Indian sector. The 10-year timescale of the experiments allows a large fraction of weakly stratified water to subduct

¹⁹⁹ into the subsurface (Jones et al. 2016), and using a 10-year timescale also allows us to carry out a ²⁰⁰ suite of passive tracer and adjoint sensitivity runs within the 20-year observationally-constrained ²⁰¹ time period of ECCOv4. We then use the annual mean tracer distribution after 10 years of model ²⁰² integration to map out the ventilation pathways. See section 3 for more details and results.

²⁰³ *c. Design of the adjoint sensitivity experiments*

²⁰⁴ An adjoint model is a tool for sensitivity analysis; all other applications derive from there (Errico 1997). Adjoint methods can be used to construct state estimates (e.g. the ECCOv4 state estimate used in this paper) via the iterative optimization of model initial conditions, model parameters, and surface forcing (collectively called the "controls"), guided by *gradients* or *sensitivities* that are calculated by an adjoint model. In such applications, the sensitivities indicate how one needs to adjust the controls in order to reduce the *cost function*, which is a measure of model-data misfit over the duration of the model run. This method enables objective, physically consistent, and ²¹¹ efficient model optimization that would otherwise require considerable ad-hoc parameter tuning ₂₁₂ and/or the introduction of artificial sources or sinks of heat and salt into the subsurface.

213 More generally, adjoint methods also enable us to carry out efficient and comprehensive sensi-²¹⁴ tivity studies, such as the one presented in this manuscript. In a traditional "forward" perturbation ²¹⁵ experiment, the experimenter introduces a perturbation into the model (e.g. a change in the initial $_{216}$ temperature pattern) and examines the effect on some quantity of interest (e.g. RVPh). Using this ₂₁₇ method, the experimenter cannot feasibly determine if the perturbation they selected is the ideal 218 or optimal one (i.e. the one with the largest effect on RVPh) without performing a very large num-²¹⁹ ber of numerical experiments. By contrast, in an adjoint sensitivity experiment, the experimenter ²²⁰ selects a quantity of interest (e.g. annual mean RVPh) and uses an adjoint approach to calculate ²²¹ *gradients* or *sensitivities*. The sensitivity fields indicate the perturbations with the largest possible

²²² linear impact on RVPh. For examples of adjoint sensitivity experiments, see Verdy et al. (2014); $_{223}$ Sévellec and Fedorov (2015); Jones et al. (2018), and many others. We refer the reader to these ²²⁴ works for a more thorough and general description to adjoint modeling (Thacker and Long 1988; ²²⁵ Marotzke et al. 1999b; Heimbach 2008; Mazloff et al. 2010; Griewank and Walther 2012; Verdy $_{226}$ et al. 2014; Forget et al. 2015a; Fukumori et al. 2015; Jones et al. 2018, for example).

²²⁷ We perform a set of such adjoint sensitivity experiments in order to examine the sensitivity of ²²⁸ RVPh to subsurface ocean properties. The objective function is defined as the annual- and volume-²²⁹ mean RVPh:

$$
\tilde{J} = \frac{1}{V\Delta t} \int_{V} \int_{\Delta t} \rho_0 c_p \theta(\mathbf{r}, t) dt dV,
$$
\n(1)

²³⁰ where ρ_0 is the reference density, c_p is the heat capacity of seawater, θ is the potential temperature, 231 **r** is the position vector, *t* is the time, *V* is the control volume, and Δt is the time period of the 232 integration. For convenience, we scale \tilde{J} by the constant $\rho_0 c_p$, so RVPh= $J = \tilde{J}/\rho_0 c_p$, meaning that RVPh has units of $°C$.

²³⁴ We compute an ensemble of six adjoint sensitivity experiments, with the objective function ²³⁵ defined over the last year of each run, i.e. from 1 January to 31 December, with objective function ²³⁶ years ranging from 2006 through 2011. Each experiment spans a minimum of 15 years, up to a ₂₃₇ maximum of 20 years. We allow the RVP to vary by ensemble member based on the results of ₂₃₈ the passive tracer experiments. Our ECCOv4 adjoint model calculates the sensitivities of these 239 objective functions to a set of independent variables, including temperature $\left(\frac{\partial J}{\partial T}\right)$ and salinity ²⁴⁰ (∂ *J*/∂*S*); the sensitivity fields vary with space (latitude, longitude, depth) and time. We use 14-day ²⁴¹ averaged sensitivity fields throughout.

242 3. Identifying the recently ventilated subsurface ocean

²⁴³ We use a six-member ensemble of passive tracer experiments with release years ranging from $_{244}$ 1996 to 2001 to identify the RVP in ECCOv4. The ensemble mean, annual mean tracer distribution ²⁴⁵ after 10 years of integration (Figure 2) highlights the ventilation pathways from the Southern ²⁴⁶ Ocean mixed layer into the subsurface thermocline.

²⁴⁷ The normalized tracer distribution features several cores, i.e. local maxima with some degree ²⁴⁸ of spatial coherence across latitudes and longitudes (Supplemental Animation S1). In the Atlantic ²⁴⁹ sector, we find a core below the 1992-2011 maximum MLD in the subtropical latitudes (roughly ₂₅₀ between 30°S-10°S) that drifts southwards before merging with a broad pattern that stretches $_{251}$ from the Weddell Sea, across the ACC, and into the subtropics (Figure 2(a)). The approximately ²⁵² zonal Atlantic distribution bears the imprint of the convergence of the Brazil current and the ACC. ²⁵³ In the Indian sector, the tracer distribution is influenced by the Agulhas current and the broader ²⁵⁴ circulation of the South Indian Gyre that is characterized by relatively shallow overturning in ²⁵⁵ density space; here, we find the core of the distribution at roughly 500 m and densities lighter than ²⁵⁶ the $\sigma_0 = 26.9$ kg m^{−3} density surface (Figure 2(b)). In the Pacific sector, we find relatively large ²⁵⁷ values throughout the entire South Pacific Gyre (SPG), with two distinct cores. The core of the 258 distribution in the western SPG is found at densities lighter than $\sigma_0 = 26.9$ kg m⁻³ and is affected ²⁵⁹ by tracer that subducts in the Indian sector, whereas the core of the distribution in the eastern SPG ²⁸⁰ largely straddles the $\sigma_0 = 26.9$ kg m^{−3} line and is relatively less affected by tracer that subducts $_{261}$ in the Indian sector (Figure 2(c,d)). High tracer concentrations in the eastern region of the SPG ²⁶² highlights the relatively efficient Eastern Pacific export pathway that was previously identified in ²⁶³ high-resolution simulations (Jones et al. 2016, Figure 8d).

₂₆₄ In order to define the RVP, we use the annual mean tracer concentrations after 10 years of ²⁶⁵ transport and some additional physical and geographical criteria, selecting grid cells that satisfy ²⁶⁶ the following four conditions:

 \bullet be located below the maximum mixed layer over the entire ECCOv4-r2 period (i.e. 1992- 268 2011)

²⁶⁹ • tracer concentration is at least 10% of the global ocean maximum value

• be located in the Southeast Pacific, between 170° W-60°W and 60° S-20°S

• potential density is greater than or equal to $\sigma_0 = 26.9 \text{ kg m}^{-3}$ 271

²⁷² The $\sigma_0 = 26.9 \text{ kg m}^{-3}$ surface delineates lighter waters from waters that are at least as dense ²⁷³ as the SAMW. The resulting RVP for any given release year is a three-dimensional volume with $_{274}$ a lateral imprint that varies with depth (Figure 3 and 4). Between roughly 300-500m (Figure 275 3(a)-(c)), the areal extent of the RVP changes with depth and sits largely in the Southeast Pacific, ²⁷⁶ north of the regions with deep mixed layers. The structure of the RVP at these depths reflects the ²⁷⁷ subduction and ventilation pattern associated with the Eastern Pacific export pathway (Jones et al. $_{278}$ 2016). Between roughly 500-700 m (Figure 3(d)-(e)), the RVP reaches its maximum areal extent, ²⁷⁹ covering a large fraction of the SPG.

280 1) THE RVP IS MORE GENERAL THAN SAMW

²⁸¹ Although the RVP has considerable overlap with the density ranges associated with SAMW, it ²⁸² is more general than SAMW alone. Our passive tracers ventilate a wide range of water masses, ²⁸⁸ including SAMW, AAIW, and the lightest part of the Circumpolar Deep Water (CDW). In this ²⁸⁴ way, our RVP approach is broader than an approach focused on a particular density range, in that ²⁸⁵ it allows us to consider a wider range of ventilated water mass types than SAMW alone. Because

 RVPh is the heat content of a fixed volume, RVPh is a measure of the heat distribution in the ²⁸⁷ target region, as RVPh may be affected by both isopycnal heave and by changes in properties along isopycnals. RVPh may have different sensitivities than the SAMW heat content, which may be affected by changes in the volume of SAMW and/or changes in the properties of water in the SAMW density range. We further explore this distinction in section 5.

 Having defined the RVP in ECCOv4, we can now perform a set of adjoint sensitivity experiments using the adjoint capability of the ECCOv4 model setup.

4. Sensitivities of subsurface heat content to local and remote anomalies

²⁹⁴ In this section, we examine the sensitivity of RVPh to changes in temperature and salinity, de- composed into changes at constant density (i.e. kinematic sensitivities) and changes with varying density (i.e. dynamic sensitivities). In subsection a, we show how sensitivities to temperature and ²⁹⁷ salinity perturbations can be cast in terms of kinematic and dynamic sensitivities. In subsection b, we investigate the spatial and temporal structure of the kinematic sensitivities, focusing on both the subsurface and the surface. Finally, in subsection c, we examine the spatial and temporal structure of the dynamic sensitivities, including regional patterns in the subsurface and at the surface. In ³⁰¹ this analysis, we use the time-variable -0.25 m ECCOv4 sea surface height contour as a proxy for ³⁰² the Subantarctic Front (SAF) that roughly divides the Southern Ocean into two regions - one dom-³⁰³ inated by gyre-like circulation and another region dominated by the circumpolar flow associated 304 with the ACC (Kim and Orsi 2014).

a. Defining kinematic and dynamic sensitivities

³⁰⁶ Following Marotzke et al. (1999b) and Jones et al. (2018), we decompose the adjoint sensitivity ³⁰⁷ fields into sensitivities to changes that propagate along isopycnals (i.e. kinematic) and sensitivities ³⁰⁸ to changes in density (i.e. dynamic). Writing the objective function in terms of density and 309 temperature $J = J[\rho(T, S), T]$ allows us to write the sensitivities as follows:

$$
\left(\frac{\partial J}{\partial T}\right)_S = \left(\frac{\partial J}{\partial \rho}\right)_T \left(\frac{\partial \rho}{\partial T}\right)_S + \left(\frac{\partial J}{\partial T}\right)_{\rho}.
$$
\n(2)

310 The first term on the right-hand side of equation 2 is the "dynamic" component of the sensitivity 311 (i.e. sensitivity to changes in density), and the second term on the right-hand side is the "kine-312 matic" component (i.e. dynamically-inactive sensitivities to temperature anomalies). Using the 313 coefficient of thermal expansion α and coefficient of haline contraction β , defined as

$$
\alpha \equiv -\frac{1}{\rho} \left(\frac{\partial \rho}{\partial T} \right)_S \text{ and } \beta \equiv \frac{1}{\rho} \left(\frac{\partial \rho}{\partial S} \right)_T,
$$
\n(3)

314 we can write

$$
\left(\frac{\partial J}{\partial S}\right)_T = \left(\frac{\partial J}{\partial \rho}\right)_T \left(\frac{\partial \rho}{\partial S}\right)_T = \beta \rho \left(\frac{\partial J}{\partial \rho}\right)_T,\tag{4}
$$

315 and the dynamic sensitivity becomes:

$$
F_{dyn} = \left(\frac{\partial J}{\partial \rho}\right)_T \left(\frac{\partial \rho}{\partial T}\right)_S = \frac{1}{\beta \rho} \left(\frac{\partial J}{\partial S}\right)_T \left(\frac{\partial \rho}{\partial T}\right)_S = -\frac{\alpha}{\beta} \left(\frac{\partial J}{\partial S}\right)_T.
$$
(5)

316 We can write the kinematic sensitivity as:

$$
F_{kin} = \left(\frac{\partial J}{\partial T}\right)_S + \frac{\alpha}{\beta} \left(\frac{\partial J}{\partial S}\right)_T.
$$
\n(6)

317 Physically, kinematic sensitivities indicate the linear response of the objective function to the 318 simultaneous application of a small temperature change and a compensating salinity change of ³¹⁹ the form $\Delta S = \Delta T \alpha / \beta$, such that density remains constant following the linearized equation of 320 state for seawater, $\rho = \rho_0(1 - \alpha\Delta T + \beta\Delta S)$. We can refer to these perturbations as "density-³²¹ compensated temperature anomalies". In contrast, dynamic sensitivities indicate the linear re-³²² sponse of the objective function to the application of a temperature change ∆*T* or a density-³²³ equivalent salinity change of $\Delta S = -\Delta T \alpha/\beta$ in the equation of state. The linear response is

³²⁴ given by $\Delta J = F_{dyn} \Delta T = F_{dyn} (-\Delta S\beta/\alpha)$. We use 14-day averaged, three-dimensional α/β fields ³²⁵ derived from ECCOv4-r2 potential temperatures and salinities using the TEOS-10 toolbox (Mc-³²⁶ Dougall and Barker 2011). Next, we examine the kinematic and dynamic sensitivities of RVPh. ³²⁷ In this paper, we follow the naming and conceptual conventions established in the adjoint mod-³²⁸ eling literature, specifically the use of kinematic and dynamic sensitivities (Marotzke et al. 1999a; ³²⁹ Jones et al. 2018, e.g.). Kinematic sensitivities are conceptually similar to sensitivities to "spice" ³³⁰ anomalies (Flament 2002), and dynamic sensitivities are conceptually similar to sensitivities to ³³¹ "heave" anomalies (Bindoff and Mcdougall 1994). However, despite their resemblance, it would ³³² be misleading to conflate kinematic sensitivities with sensitivities to spice anomalies, and it would ³³³ be misleading to conflate dynamic sensitivities with sensitivities to heave anomalies. Their deriva-³³⁴ tions and resulting definitions are not rigorously identical. In addition, there is a difference in how ³³⁵ these quantities are normally used; spice and heave are typically used to decompose local potential ³³⁶ temperature anomalies into these two components, whereas kinematic and dynamic adjoint sensi-³³⁷ tivities are more generally used to show impacts on an objective function over different spatial and ³³⁸ temporal lags.

³³⁹ At present, adjoint sensitivity fields in ECCOv4 do *not* include sensitivity to changes in mixing ³⁴⁰ as parameterized by the GGL mixed layer turbulence closure scheme. Also, the sensitivity fields ³⁴¹ do *not* include the sensitivity to changes in sea ice. The adjoint sensitivity calculations essentially ³⁴² treat GGL mixing and sea ice as fixed processes. We apply a mask of the form (1− *f*), where *f* is ³⁴³ the local sea ice concentration, to all surface sensitivity fields to account for the lack of sensitivity ³⁴⁴ to sea ice change.

³⁴⁵ *b. Sensitivity to changes at constant density*

³⁴⁶ Kinematic sensitivity fields approximately indicate potential source water pathways. More ³⁴⁷ specifically, one can think of the kinematic sensitivity fields as indicating the eventual response of annual mean RVP heat content to a spatially uniform, density-compensated potential temperature perturbation that is sustained for some time (in our case, two weeks) and allowed to propagate. Regions with zero sensitivity cannot affect RVPh; such regions are not connected to the RVP via advection and mixing on the associated timescales. The kinematic sensitivity fields are positive or zero nearly everywhere; an increase/decrease in temperature in regions with non-zero sensitivities eventually increases/decreases RVP heat content on the indicated lag timescale (Figure 5).

₃₅₄ The near-surface kinematic sensitivity fields highlight the potential propagation pathways of ³⁵⁵ anomalies into the SAMW formation regions, where they can be subducted below the mixed layer ³⁵⁶ and exported into the thermocline (Figure 5, left column, and Supplemental Animation S2). In the ³⁵⁷ Pacific basin, the largest near-surface sensitivity fields tend to be found just north of the SAF proxy, 358 within or upstream of regions of deep mixed layers and mostly south of the RVP areal extent. In ³⁵⁹ the Indian basin, the largest near-surface sensitivity fields are found along the SAF proxy or just ³⁶⁰ south of it. The fields cross over the SAF around Campbell Plateau and South Australian Basin, 361 suggesting cross-front exchange in this region.

³⁶² Density-compensated anomalies cannot typically propagate from directly above the RVP into ³⁶³ the RVP itself on the timescales considered here; they are not typically advected into the regions ³⁶⁴ of deep mixing. On timescales longer than 10 years, the near-surface sensitivities extend across the ³⁶⁵ entire Southern Ocean, putting a lower bound on the temperature anomaly propagation timescale ³⁶⁶ from the Atlantic to the RVP (Figure 5, left column). On timescales longer than roughly 5-6 ³⁶⁷ years, we find sensitivities south of the SAF proxy in the Indian basin; here, wind stress may

³⁶⁸ advect anomalies across the SAF via Ekman transport. On shorter timescales, the sensitivity fields ₃₆₉ are more localized around the southeast Pacific mixed layer region and are typically smaller in ₃₇₀ magnitude relative to the deeper levels. Temperature anomalies can only affect RVPh if they ³⁷¹ have sufficient time to propagate into the mixed layer, subduct, and get exported; only very local, ³⁷² targeted anomalies can affect RVPh on timescales shorter than 1-2 years. The persistent local 373 maximum in the southeast Pacific indicates a relatively rapid route from the surface into the RVP, ³⁷⁴ which is consistent with a well-documented subduction and export pathway in that location (Sallée) 375 et al. 2010b).

₃₇₆ The subsurface sensitivity fields highlight the long residence time of local hypothetical pertur-³⁷⁷ bations in the subsurface of the gyre, as well as the pathways of subsurface ocean advection into ³⁷⁸ the RVP (Figure 5, right column, and Supplemental Animation S3). On timescales longer than 10 379 years, we find sensitivities extending along the East Australian Current and, in contrast with the ³⁸⁰ surface sensitivities, sensitivities along the Agulhas Current and associated retroflection as well. ³⁸¹ Surface anomalies in the Agulhas Current cannot affect RVPh on the timescales considered here, ³⁸² but anomalies in the subsurface (here, roughly 550 m) can eventually alter RVPh. We also find ³⁸³ some sensitivity at this depth extending along the Brazil current. Notably, the Agulhas current is ³⁸⁴ the only major Southern Hemispheric western boundary current with zero kinematic sensitivity in ³⁸⁵ the near-surface.

 386 The global mean RMS kinematic sensitivity decreases with timescale (Figure 6(a)). It is reason-³⁸⁷ ably well described ($R^2 > 0.99$) by an exponential function with a decay rate of about 10 years. ³⁸⁸ The decay partly reflects the fact that as we consider longer timescales, RVPh becomes sensitive ³⁸⁹ to a larger fraction of the ocean, so the global RMS sensitivity decreases. The decay also partly ₃₉₀ reflects the tendency of air-sea exchange, represented in ECCOv4 as bulk formulae with a pre-³⁹¹ scribed set of atmospheric variables, to dampen potential temperature anomalies at the surface

₃₉₂ and thereby reducing the sensitivities. The linear change in sensitivity between lag 0 and lag 1 yr ³⁹³ reflects the one-year integration period over which the objective function is calculated. There is no sensitivity to anomalies after year 1, after the integration has finished. The ensemble standard deviation is small, indicating a relatively low level of interannual variability in sensitivity across our six-member ensemble.

³⁹⁷ We find non-zero sensitivities to hypothetical anomalies within the local RVP throughout the entire 16-year adjoint sensitivity experiment, indicating that RVPh is sensitive to local density- compensated temperature anomalies on timescales of at least 16 years (Figure 6(b)). The ensemble ⁴⁰⁰ mean RMS local sensitivity time series is well-described ($R^2 > 0.99$) by a two-term exponential with decay rates of 4.4 years and 10.9 years. The non-local RMS sensitivities outside of the RVP exceed the local sensitivities on timescales longer than roughly 5 years (Figure 6(b)). RMS sensitivities outside of the MLD mask are larger than those within the MLD mask on timescales shorter than about 8 years (Figure 6(c)). This is largely due to the fact that sensitivities exist below the surface, e.g. in the RVP itself, where the overlying mixed layers are typically shallow.

⁴⁰⁶ Outside of the RVP, we mostly find non-zero sensitivities just north of the SAF proxy in all three ocean basins, in what may be considered the northern extent of the ACC (Figure 6(d)). The near- surface sensitivity fields display a clear seasonal cycle, with the RMS mean peaking in August in the Pacific and Indian basins, during the strong mixing phase that occurs as the mixed layer deepens before reaching its maximum depth in mid-September (Figure 6e). The sensitivity fields ⁴¹¹ spread westward with lead time, reflecting the advective timescale between basins (Figure $6(e,f)$). ⁴¹² The RMS sensitivity in the West Pacific exceeds the East Pacific value on timescales longer than roughly 4.4 years.

c. Sensitivity to changes in density

⁴¹⁵ The dynamic sensitivity fields indicate patterns of density change that can alter RVPh, poten- tially by inducing changes in isopycnal tilt and thereby altering the associated transport and heat convergence into the RVP. More specifically, one can think of the dynamic sensitivity fields as indicating the eventual response of annual mean RVP heat content to a spatially uniform potential temperature perturbation, or a density-equivalent salinity perturbation, that is sustained for some time (two weeks, as with the kinematic sensitivity case) and allowed to affect the oceanic density ⁴²¹ structure. Regions of zero sensitivity cannot affect RVPh; such regions are not dynamically con- nected to the RVP. The dynamic sensitivity fields are both positive and negative, with the dipoles highlighting regions where a change in isopycnal tilt will change the associated circulation and 424 ultimately RVPh (Figure 7).

⁴²⁵ The near-surface dynamic sensitivity fields are mostly negative across all lags, with local ex- ceptions south of Australia and in the eastern tropical Pacific (Figure 7, left column). Negative values indicate regions where a hypothetical temperature increase, which will decrease density, will ultimately decrease RVP heat content. Similarly, in these regions a hypothetical temperature decrease, which will increase density, will ultimately increase RVP heat content. It is important to note that, due to technical limitations, our adjoint sensitivity fields *do not* represent non-linear changes. They *do* reflect changes in circulation and diffusion. So the negative near-surface values ⁴³² may represent the fact that, even under fixed mixing conditions, relatively lighter water may not reach the depths of the RVP as readily as relatively denser water, where "relative" refers to the background ECCOv4 state around which the adjoint sensitivities are calculated. Alternatively, the negative near-surface values may represent a hypothetical decrease in RVP heat convergence in-duced by a surface decrease in density, or the values may represent a hypothetical increase in RVP heat convergence induced by a surface increase in density. As with the kinematic sensitivity fields, on shorter timescales the dynamic sensitivities become increasingly localized to the southeastern 439 edge of the RVP, near the southern tip of South America.

 In contrast with the near-surface kinematic sensitivity fields, the near-surface dynamic sensitiv- ity fields feature a persistent negative local minimum along the western coast of South America (Figure 7, left column, and Supplemental Animation S4). Considered together with the small positive values in the eastern tropical Pacific, this dipole reflects a sensitivity to the across-shelf pressure gradient. Variations in the across-shelf pressure gradients are associated with changes in basin-scale pressure gradients and the associated basin-scale circulation, which can ultimately change heat convergence and thus heat content within a selected ocean volume (Fukumori et al. 2015; Jones et al. 2018; Hughes et al. 2018).

⁴⁴⁸ The subsurface dynamic sensitivity fields highlight the sensitivity of annual mean RVP heat con- tent to basin-wide density contrasts, equivalently expressed as tilted basin-scale isopycnal surfaces (Figure 7, right column, and Supplemental Animation S5). This basin-scale structure is associated ⁴⁵¹ with the circulation of the SPG, which is in part maintained by pressure gradients induced by wind stress curl. We find a persistent sensitivity dipole stretched across the RVP, with positive values to the east and negative values to the west. The positive values reach into the tropics and extend westward with timescale, across both the Pacific and Indian basins. To illustrate the structure of these fields, consider a hypothetical potential temperature anomaly characterized by warming to ⁴⁵⁶ the east of the RVP and cooling to the west of the RVP, imposed on the model vertical level cen- tered at 553 m (Figure 7(d)). The warming to the east would decrease density at 553 m and, based on the positive sign of the sensitivity field, ultimately contribute to an increase RVPh. The cooling to the west would increase density at 553 m and, based on the negative sign of the sensitivity field, it would also increase RVPh. The change in RVPh induced by any anomaly is the product of the

461 sensitivity field and the imposed anomaly, i.e. $\Delta J = F_{dyn} \Delta T$. In this scenario, we have imposed ⁴⁶² isopycnal deepening to the east and isopycnal shoaling to the west. In the RVP, isopycnal surfaces at roughly 500-600 m tend to shoal from west to east, so our imposed perturbation alters isopycnal surfaces locally, ultimately changing transport and increasing heat convergence in the RVP (Jones et al. 2018).

 The subsurface sensitivity fields also feature a persistent sensitivity along the western coast of South America (Figure 7, right column). In contrast to the near-surface fields, which are fairly consistently negative along the shelf, the along-shelf subsurface fields are positive to the north and negative to the south. The north-south sign contrast indicates a sensitivity to barotropic structure, whereas the surface-subsurface sign contrast highlights an overall sensitivity to baroclinic den-⁴⁷¹ sity structure above the continental shelf. In addition, the subsurface fields display characteristic 472 fingerprints of baroclinic Rossby waves, as seen by westward propagating positive anomalies at ⁴⁷³ depth (Supplemental Animation S5). As with the surface fields, on shorter timescales the subsur-⁴⁷⁴ face dynamic sensitivities become increasingly localized, albeit with a different structure than the surface fields. The subsurface fields are localized around the edge of the RVP, indicating a sensi- tivity to density contrasts across the RVP that can alter fluxes across the boundary of the RVP, and ⁴⁷⁷ the subsurface sensitivities are also localized along the western coast of South America, indicating a short-timescale sensitivity to density changes along the continental shelf.

⁴⁷⁹ The globally averaged RMS dynamic sensitivity increases with timescale, reaching a maximum at roughly 8.7 years (Figure 8(a)). Much of this sensitivity comes from outside the RVP, partic- ularly near its boundary (Figure 8(b)). RMS sensitivity is similar inside and outside of the MLD mask, indicating that when considering RVPh, we need to study dynamic changes both inside and far outside of the regions of deep mixing (Figure 8(c)). The RMS sensitivity is persistently larger north of the SAF proxy, although much of this is located just north of the SAF, and the RMS

485 sensitivity south of the SAF increases roughly linearly with timescale (Figure 8(d)). The near- surface fields also display a seasonal cycle, with similar timings as the kinematic fields (Figure $8(e)$). Notably, the RMS dynamic sensitivity is persistently higher in the West Pacific than in the East Pacific, indicating the importance of dynamic basin-scale changes (Figure 8(h)). The RMS sensitivity in the Indian and Atlantic basins increases with timescale, although the values are neg- $_{490}$ ligibly small (less than 3% of the maximum) for short timescales (about 2.5 years for the Indian basin and 5 years for the Atlantic basin).

492 5. Discussion

⁴⁹³ In the canonical view of Southern Ocean meridional overturning circulation (MOC), Circum-⁴⁹⁴ polar Deep Water upwells south of the ACC, where it is brought to the surface and exported northwards; this water is modified by air-sea fluxes and is ultimately subducted as SAMW and AAIW (Marshall and Speer 2012). As a result of this overturning circulation, the properties of SAMW and AAIW are potentially sensitive to anomalies in near-Antarctic waters. For example, the variability of the Amundsen Sea low (ASL) can affect the interannual and interdecadal vari- ability of SAMW in the Southeast Pacific via cross-frontal Ekman transport of Antarctic surface waters (Garabato et al. 2009; Close et al. 2013). In 2008-2010, a deep ASL enhanced the merid-₅₀₁ ional wind-driven sea ice export from the eastern Ross Sea; the sea ice melt introduced a cold ₅₀₂ surface freshwater anomaly that was advected across the PF and SAF by Ekman transport and then transported into the SAMW formation region by the ACC (Cerovecki et al. 2019). In this ₅₀₄ context, our adjoint sensitivity results may appear surprising, in that they show relatively weak sensitivities to anomalies in the Ross Sea. Our results are not inconsistent with those studies, but there are several factors to consider when comparing our results with those obtained using other approaches, as discussed below.

a. RVPh is a fixed-volume quantity

 Our control volume is fixed relative to the model grid. Although RVP captures the mean position ₅₁₀ of the ventilated water, it does not move with isopycnal heave. Because it overlaps with both $_{511}$ SAMW and AAIW, it is more general than either water mass, but we must be careful not to ₅₁₂ conflate SAMW/AAIW with the RVP. Isopycnal heave in the subtropical gyre can affect RVPh without necessarily changing the heat content of the recently subducted SAMW. However, RVPh can still tell us something about subduction. The majority of recent heat uptake by SAMW is driven by a thickening of the associated isopycnal layers, driven by changes in wind forcing and subduction in the deep mixed layer region Gao et al. (2018). Considering SAMW in the Pacific basin, its local mean temperature is correlated with remote forcing (e.g. air-sea heat flux, wind stress) with a lag that is attributed to transport, whereas its total heat content is controlled largely 519 by layer thickness, which is driven by local wintertime forcing (Meijers et al. 2019). These results are consistent with the our adjoint sensitivity fields, which largely capture mechanisms that can affect mean temperature within the SAMW as opposed to those that can affect total heat content via changing layer thickness. It would be useful to examine the sensitivities in an adjoint model that uses isopycnal surfaces as its vertical coordinate, but this is beyond the scope of this paper.

b. Zonal flow dominates meridional overturning

₅₂₅ The largely zonal volume transport of the ACC is at least an order of magnitude larger than the volume transport associated with the MOC (Talley 2013; Donohue et al. 2016). As such, we might ₅₂₇ expect our adjoint sensitivity fields to reflect the dominance of ACC transport in the Southeast Pacific. According to these fields, sensitivity to high-latitude processes is small but nonzero. To examine the impacts of actual temperature and salinity anomalies on RVPh, one could examine ₅₃₀ the combination of sensitivity fields together with temperature and/or salinity anomaly fields. In

⁵³¹ this work, we focus on the sensitivity fields themselves, as these could potentially be combined ⁵³² with a large variety of anomaly fields in order to address different questions (e.g. how does the ⁵³³ SAM forcing pattern affect RVPh across different timescales?)

⁵³⁴ *c. Cross-front exchange rates differ between data products*

₅₃₅ Cross-front exchange in the SO is highly localized, occurring in relatively narrow longitude ₅₃₆ bands in the lee of bathymetric features (Thompson and Sallée 2012). The propagation of anoma- 537 lies from the Ross Gyre to the subtropics occurs largely via two distinct exchange windows in the ⁵³⁸ Southeast Pacific and the central Pacific; on timescales of roughly 1-3 years, the central Pacific ₅₃₉ pathway enables transport from Antarctic surface waters to the deep mixed layer region (Cerovecki ⁵⁴⁰ et al. 2019). The subsurface exchange pathway in the central Pacific (around the Eltanin fracture $_{541}$ zone, between roughly 110-160°W) is driven by enhanced diapycnal mixing near the relatively ₅₄₂ rough topography and temporal circulation variability on timescales longer than 90 days (McAu-₅₄₃ field 2019). Because of its relatively coarse resolution, it is possible that ECCOv4-r2 may have ⁵⁴⁴ a lower cross-frontal exchange rate than that of higher-resolution data products, but a lack of ob-⁵⁴⁵ servational data constraints makes it challenging to discriminate amongst estimates. A thorough, ⁵⁴⁶ quantitative comparison of exchange rates between ECCOv4-r2 and other data products would be ⁵⁴⁷ a useful addition to this work.

⁵⁴⁸ *d. Mixed layer depth*

⁵⁴⁹ In this work, we calculate mixed layer depth using a criteria based on the density change associ- σ ated with a temperature variation of $\Delta T = 0.8$ °C (Kara et al. 2000). Compared with other methods, ⁵⁵¹ this threshold approach tends to be biased towards deeper values (Forget et al. 2015c, supporting ₅₅₂ information). Because we discard all tracer above the maximum mixed layer, this implies that our

₅₅₃ estimate of the subducted volume may be somewhat conservative. It is possible that our results would be different using another mixed layer scheme. That being said, the stratification and mixed layer properties in ECCOv4 compare well spatially with in-situ profiles (Forget et al. 2015c, Fig. 6).

e. Adjoint sensitivities use fixed vertical mixing and sea ice

 At present, adjoint sensitivity fields in ECCOv4 do *not* include sensitivities to changes in mix- ing as parameterized by the GGL mixed layer turbulence closure scheme. Also, the sensitivity fields do *not* include the sensitivity to changes in sea ice. Although the forward and adjoint runs do feature dynamic sea ice and vertical mixing as parameterized by GLL, the adjoint sensitivity ₅₆₂ calculations essentially treat them as fixed processes. Any processes that affect RVPh primar- ily through *changing* sea ice concentration and/or vertical mixing will not be represented in our sensitivity fields.

6. Summary and conclusions

 We used numerical passive tracer experiments and physical state variables (i.e. mixed layer depth, potential density, potential vorticity) in an observationally-constrained state estimate to ₅₆₈ identify the recently ventilated Eastern Pacific sector of the Southern Ocean (RVP), which is a relatively efficient export pathway from the surface ocean into the subsurface thermocline. The RVP, defined on a 10-year timescale, is located north of the -0.25 m sea surface height contour (i.e. the SAF proxy) and north of the deep mixed layer regions, highlighting the ventilation path- way from the tip of South America, along the eastern Pacific boundary and out across the South Pacific Gyre (SPG) in the subsurface. The RVP varies with depth, reaching maximum areal extent between 500-600 m, and it features some variation with tracer release year, mostly in its western

₅₇₅ extent. The overall spatial pattern of the subducted tracer is consistent with estimates from an eddy-permitting model (Jones et al. 2016).

₅₇₇ We used the adjoint capabilities of MITgcm to calculate the sensitivities of annual mean heat ₅₇₈ content within the fixed volume RVP (i.e. RVPh) to anomalies in potential temperature and salin-₅₇₉ ity. These sensitivities are four-dimensional (i.e. latitude, longitude, depth, and time) and indicate the anomaly patterns that can have the largest potential impact on RVPh after the appropriate lag time has elapsed (Figure 9). In terms of sensitivities to surface anomalies, RVPh is most sensitive to anomalies along the SAF proxy, upstream of the subduction hotspot in the Southeast Pacific. The sensitivity signal crosses the SAF proxy around Campbell Plateau and just south of Australia, ₅₈₄ indicating possible wind-driven cross-frontal exchange in that region. This result is consistent ₅₈₅ with estimates of the spatial pattern of meridional Ekman transport of atmospheric carbon in an eddy-permitting state estimate (Ito et al. 2010). In terms of sensitivities to subsurface anomalies, RVPh is most sensitive to basin-scale changes in density, as indicated by the persistent dipole in subsurface dynamic sensitivity (Figure 7). The sensitivity dipole is centered around the same lati- tudes as the RVP, extending upstream along the ACC and northwards into the tropical Pacific with ₅₉₀ increasing timescale. This persistent dipole implies that property gradients between the SPG and 591 the ACC are important for interannual variability in RVPh.

 By the end of the 21st century, Southern Ocean wind stress is expected to strengthen in magni- tude and move poleward, resulting in a poleward shift of the gyres, and mixed layers are expected to shoal (Meijers 2013). Our results suggest that the resulting subtropical, basin-scale temperature and salinity changes induced by the expected shifts in wind stress and heat flux can have a consid- erable impact on the heat distribution of the ventilated Southeast Pacific, across the density ranges ₅₉₇ of the base of the SAMW, AAIW, and the lightest CDW. In a companion manuscript, we examine the sensitivities of RVP heat content to changes in surface forcing (Jones et al. 2019).

₅₉₉ *Acknowledgments.* This study is supported by grants from the Natural Environment Research Council (NERC), including [1] The North Atlantic Climate System Integrated Study (ACSIS) 601 (grant NE/N018028/1, authors DJ, ES), [2] Securing Multidisciplinary UndeRstanding and Pre-602 diction of Hiatus and Surge events (SMURPHS) (grant NE/N006038/1, author EB), and [3] Ocean Regulation of Climate by Heat and Carbon Sequestration and Transports (ORCHESTRA) (grant NE/N018095/1, authors EB, AM, SJ). GF is supported by NASA award #6937342 and Simons ⁶⁰⁵ Foundation award #549931. Numerical model runs were carried out on ARCHER, the UK national HPC facility [http://archer.ac.uk/]. Adjoint code was generated using the TAF software tool, created and maintained by FastOpt GmbH [http://www.fastopt.com/]. We used the gcmfaces package for plotting and analysis (https://github.com/gaelforget/gcmfaces). The authors ⁶⁰⁹ wish to thank two anonymous reviewers whose comments greatly improved the quality of this paper.

 Data availability statement: The model used here is an instance of the ECCOv4-r2 state es- timate (https://github.com/gaelforget/ECCOv4) as an instance of the MIT general circula- tion model (MITgcm, http://mitgcm.org/). Specific model configuration files, input data, and sample output available via Zenodo (Jones 2020).

References

616 Armour, K. C., J. Marshall, J. R. Scott, A. Donohoe, and E. R. Newsom, 2016: Southern Ocean warming delayed by circumpolar upwelling and equatorward transport. *Nature Geoscience*, 9 (7), 549–554, doi:10.1038/ngeo2731, URL http://www.nature.com/articles/ngeo2731.

619 Bindoff, N. L., and T. J. Mcdougall, 1994: Diagnosing Climate Change and Ocean Ven-tilation Using Hydrographic Data. *Journal of Physical Oceanography*, 24 (6), 1137–

 1152, doi:10.1175/1520-0485(1994)024 $\langle 1137:DCCAOV \rangle$ 2.0.CO;2, URL https://doi.org/10. 1175/1520-0485(1994)024 \langle 1137:DCCAOV \rangle 2.0.CO;2.

 ϵ_{023} Cerovečki, I., and M. Mazloff, 2015: The spatiotemporal structure of diabatic processes governing the evolution of Subantarctic Mode Water in the Southern Ocean. *Journal of Physical Oceanog-raphy*, 46, 683–710, doi:10.1175/JPO-D-14-0243.1.

- 626 Cerovecki, I., A. J. S. Meijers, M. R. Mazloff, S. T. Gille, V. M. Tamsitt, P. R. Holland, and V. M. Tamsitt, 2019: The effects of enhanced sea ice export from the Ross Sea on recent cooling and freshening of the Southeast Pacific. *Journal of Climate*, JCLI–D–18–0205.1, doi:10.1175/ $_{629}$ JCLI-D-18-0205.1.
- 630 Cerovecki, I., L. D. Talley, M. R. Mazloff, and G. Maze, 2013: Subantarctic Mode Water Forma- tion, Destruction, and Export in the Eddy-Permitting Southern Ocean State Estimate. *Journal of Physical Oceanography*, 43 (7), 1485–1511, doi:10.1175/jpo-d-12-0121.1.
- Cessi, P., and P. Otheguy, 2003: Oceanic teleconnections: Remote response to decadal wind forc-ing. *Journal of Physical Oceanography*, 33 (8), 1604–1617.

Close, S. E., A. C. Naveira Garabato, E. L. McDonagh, B. A. King, M. Biuw, and L. Boehme,

⁶³⁶ 2013: Control of Mode and Intermediate Water Mass Properties in Drake Passage by the

Amundsen Sea Low. *Journal of Climate*, 26 (14), 5102–5123, doi:10.1175/JCLI-D-12-00346.1,

⁶³⁸ URL http://journals.ametsoc.org/doi/10.1175/JCLI-D-12-00346.1.

639 Dee, D. P., and Coauthors, 2011: The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137 (656),

 $_{641}$ 553–597, doi:10.1002/qj.828.

662 version 4: an integrated framework for non-linear inverse modeling and global ocean state esti-

mation. *Geoscientific Model Development*, 8 (10), 3071–3104, doi:10.5194/gmd-8-3071-2015.

- Forget, G., D. Ferreira, and X. Liang, 2015b: On the observability of turbulent transport rates by Argo: supporting evidence from an inversion experiment. *Ocean Science Discussions*, 12 (3), 1107–1143, doi:10.5194/osd-12-1107-2015.
- ₆₆₇ Forget, G., D. Ferreira, and X. Liang, 2015c: On the observability of turbulent transport rates by Argo: supporting evidence from an inversion experiment. *Ocean Science*, 11 (5), 839–853, doi:10.5194/os-11-839-2015.
- 670 Frölicher, T. L., J. L. Sarmiento, D. J. Paynter, J. P. Dunne, J. P. Krasting, and M. Winton, 2015: ⁶⁷¹ Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Mod-els. *Journal of Climate*, 28 (2), 862–886, doi:10.1175/jcli-d-14-00117.1.
- ⁶⁷³ Fukumori, I., O. Wang, W. Llovel, I. Fenty, and G. Forget, 2015: A near-uniform fluctuation of ⁶⁷⁴ ocean bottom pressure and sea level across the deep ocean basins of the Arctic Ocean and the Nordic Seas. *Progress in Oceanography*, 134 (C), 152–172, doi:10.1016/j.pocean.2015.01.013.
- Gao, L., S. R. Rintoul, and W. Yu, 2018: Recent wind-driven change in Subantarctic Mode Water and its impact on ocean heat storage. *Nature Publishing Group*, 1–7, doi:10.1038/ $\frac{678}{1541558-017-0022-8}$.
- Garabato, A. C. N., L. Jullion, D. P. Stevens, K. J. Heywood, and B. A. King, 2009: *Journal of Climate*, 22 (13), 3661–3688, doi:10.1175/2009jcli2621.1.
- ⁶⁸¹ Gaspar, P., Y. Grégoris, and J. M. Lefevre, 1990: A simple eddy kinetic energy model for ⁶⁸² simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site. *Journal of Geophysical Research: Atmospheres*, 95 (C9), 16 179–16 193, doi: 10.1029/JC095iC09p16179.
- Gent, P. R., and J. C. Mcwilliams, 1990: Isopycnal Mixing in Ocean Circulation Models. *Journal* ₆₈₆ *of Physical Oceanography*, **20** (1), 150–155, doi:10.1175/1520-0485(1990)020 $\langle 0150$:imiocm \rangle 2.0.co; 2.
- Griewank, A., and A. Walther, 2012: *Evaluating Derivatives*. 2nd ed., Principles and Techniques of Algorithmic Differentiation, Second Edition, Society for Industrial and Applied Mathemat-
- Hanawa, K., and L. Talley, 2001: Mode Waters. *Ocean Circulation and Climate*, G. Siedler, and J. Church, Eds., International Geophysics Series, 373–386.
- Heimbach, P., 2008: *The MITgcm/ECCO adjoint modelling infrastructure*. Clivar Exchanges.
- 694 Helm, K. P., N. L. Bindoff, and J. A. Church, 2010: Changes in the global hydrological-cycle in- ferred from ocean salinity. *Geophysical Research Letters*, 37 (18), doi:10.1029/2010GL044222, URL https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010GL044222.
- 697 Herraiz-Borreguero, L., and S. R. Rintoul, 2011: Subantarctic mode water: distribution and circu-lation. *Ocean Dynamics*, 61 (1), 103–126, doi:10.1007/s10236-010-0352-9.
- Hughes, C. W., J. Williams, A. Blaker, A. Coward, and V. Stepanov, 2018: A window on the deep ocean: The special value of ocean bottom pressure for monitoring the large-scale, deep-ocean circulation. *Progress in Oceanography*, 161, 19–46, doi:10.1016/j.pocean.2018.01.011.
- Ito, T., M. Woloszyn, and M. Mazloff, 2010: Anthropogenic carbon dioxide transport in the South-ern Ocean driven by Ekman flow. *Nature*, 463, 80, doi:10.1038/nature08687.
- Iudicone, D., K. Rodgers, R. Schopp, and G. Madec, 2007: An exchange window for the injection
- of Antarctic Intermediate Water into the South Pacific. *Journal of Physical Oceanography*, 37,
- 31–49, doi:http://dx.doi.org/10.1175/JPO2985.1.

690 ics, doi:10.1137/1.9780898717761.

- Johnson, G. C., and A. H. Orsi, 1997: Southwest Pacific Ocean Water-Mass Changes between 1968/69 and 1990/91*,. *JOURNAL OF CLIMATE*, 10, 11.
- Jones, D., 2020: DanJonesOcean/ecco v4 r2 config: Initial release. Zenodo, URL https://doi.org/ 10.5281/zenodo.3602900, doi:10.5281/zenodo.3602900.
- Jones, D. C., E. Boland, A. J. Meijers, G. Forget, S. A. Josey, J.-B. Sallee, and E. Shuckburgh, 2019: Heat distribution in the Southeast Pacific is only weakly sensitive to high-latitude heat flux and wind stress. *Journal of Geophysical Research: Oceans*, 124, doi:10.1029/2019JC015460, URL https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JC015460.
- Jones, D. C., G. Forget, B. Sinha, S. A. Josey, E. J. D. Boland, A. J. S. Meijers, and E. Shuckburgh, 2018: Local and Remote Influences on the Heat Content of the Labrador Sea: An Adjoint Sensitivity Study. *Journal of Geophysical Research - Oceans*, 105 (2-3), 182, doi:10.1002/ ⁷¹⁸ 2018JC013774.
- Jones, D. C., A. J. S. Meijers, E. Shuckburgh, J.-B. Sallee, P. Haynes, E. K. McAufield, and ´ M. R. Mazloff, 2016: How does Subantarctic Mode Water ventilate the Southern Hemi- sphere subtropics? *Journal of Geophysical Research - Oceans*, 121 (9), 6558–6582, doi: 10.1002/2016jc011680.
- Kara, A. B., P. A. Rochford, and H. E. Hurlburt, 2000: An optimal definition for ocean mixed layer depth. *Journal of Geophysical Research - Oceans*, 105 (C7), 16 803–16 821, doi:10.1029/ 2000JC900072.
- Karsten, R., and J. Marshall, 2002: Testing theories of the vertical stratification of the ACC against observations. *Dynamics of Atmospheres and Oceans*, 36, 233–246.

 Lumpkin, R., and K. Speer, 2007: Global ocean meridional overturning. *Journal of Physical Oceanography*, 37, 2550–2562, doi:10.1175/JPO3130.1.

- Luyten, J., J. Pedlosky, and H. Stommel, 1983: The ventilated thermocline. *Journal of Physical Oceanography*, 13 (2), 292–309.
- Marotzke, J., R. Giering, K. Zhang, D. Stammer, C. Hill, and T. Lee, 1999a: Construction of the Adjoint MIT Ocean General Circulation Model and Application to Atlantic Heat Transport Sensitivity. *MIT report*, 1–50.
- Marotzke, J., R. Giering, K. Q. Zhang, D. Stammer, C. Hill, and T. Lee, 1999b: Construction of the adjoint MIT ocean general circulation model and application to Atlantic heat transport sensitivity. *Journal of Geophysical Research: Atmospheres*, 104 (C12), 29 529–29 547, doi: 10.1029/1999JC900236.
- Marshall, J., and K. Speer, 2012: Closure of the meridional overturning circulation through South-ern Ocean upwelling. *Nature Geoscience*, 5, 171–180, doi:10.1038/ngeo1391.
- Mazloff, M., P. Heimbach, and C. Wunsch, 2010: An Eddy-Permitting Southern Ocean State Estimate. *Journal of Physical Oceanography*, 40, 880–899.
- McAufield, E. K., 2019: Lagrangian study of the Southern Ocean circulation. Ph.D. thesis, Uni-versity of Cambridge, URL https://doi.org/10.17863/CAM.36004.
- McDougall, T., and P. Barker, 2011: Getting started with TEOS-10 and the Gibbs Seawater (GSW) Oceanographic Toolbox. *SCOR/IAPSO WG127*.
- Meijers, A. J. S., 2013: The Southern Ocean in CMIP5. *Proceedings of the Royal Society A*, 1–21.
- Meijers, A. J. S., I. Cerovecki, B. A. King, and V. Tamsitt, 2019: A See-Saw in Pacific Sub- ˇ
- antarctic Mode Water Formation Driven by Atmospheric Modes. *Geophysical Research Letters*,
- 2019GL085280, doi:10.1029/2019GL085280, URL https://onlinelibrary.wiley.com/doi/abs/10.
- 1029/2019GL085280.
- Morrison, A., A. Hogg, and M. L. Ward, 2011: Sensitivity of the Southern Ocean overturning circulation to surface buoyancy forcing. *Geophysical Research Letters*, 38 (L14602), doi:10. 1029/2011GL048031.
- Musgrave, D., 1990: Numerical studies of tritium and helium-3 in the thermocline. *Journal of Physical Oceanography*, 20, 344–373.
- Primeau, F., and M. Holzer, 2006: The Ocean's Memory of the Atmosphere: Residence-Time and Ventilation-Rate Distributions of Water Masses. *Journal of Physical Oceanography*, 36, 1439–1456.
- Roemmich, D., J. Church, J. Gilson, D. Monselesan, P. Sutton, and S. Wijffels, 2015: Unabated planetary warming and its ocean structure since 2006. *Nature Publishing Group*, 5 (3), 240, doi:10.1038/nclimate2513.
- Russell, J., K. Dixon, A. Gnanadesikan, R. Stouffer, and J. Toggweiler, 2006: The Southern Hemi- sphere westerlies in a warming world: Propping open the door to the deep ocean. *Journal of Climate*, 19, 6382–6390.
- Sabine, C., and Coauthors, 2004: The oceanic sink for anthropogenic CO2. *Science*, 305 (367), doi:10.1126/science.1097403.
- Sallee, J., and S. Rintoul, 2011: Parameterization of eddy-induced subduction in the Southern ´ Ocean surface-layer. *Ocean Modelling*, 39, 146–153.
- Sallee, J., K. Speer, S. Rintoul, and S. Wijffels, 2010a: Southern Ocean thermocline ventilation. ´
- *Journal of Physical Oceanography*, 40, 509–529.

- Sallee, J.-B., R. J. Matear, S. R. Rintoul, and A. Lenton, 2012: Localized subduction of anthro- ´ pogenic carbon dioxide in the Southern Hemisphere oceans. *Nature Geoscience*, 5 (8), 579–584, doi:10.1038/ngeo1523.
- Sallee, J.-B., K. Speer, S. Rintoul, and S. Wijffels, 2010b: Southern Ocean Thermocline Ventila- ´ tion. *Journal of Physical Oceanography*, 40 (3), 509–529, doi:10.1175/2009jpo4291.1.
- Sarmiento, J. L., N. Gruber, M. A. Brzezinski, and J. P. Dunne, 2004: High-latitude controls of thermocline nutrients and low latitude biological productivity. *Nature*, 427 (6969), 56–60, doi:10.1038/nature02127.
- 800 Schmidtko, S., and G. C. Johnson, 2012: Multidecadal Warming and Shoaling of Antarctic Inter- mediate Water*. *Journal of Climate*, 25 (1), 207–221, doi:10.1175/JCLI-D-11-00021.1, URL 802 http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-11-00021.1.
- 803 Sévellec, F., and A. V. Fedorov, 2015: Optimal excitation of AMOC decadal variability: Links to the subpolar ocean. *Progress in Oceanography*, 132, 287–304, doi:10.1016/j.pocean.2014.02. 805 006.
- Speer, K., and G. Forget, 2013: Global Distribution and Formation of Mode Waters. *Ocean Circulation and Climate - A 21st Century Perspective*, Elsevier, 211–226, doi:10.1016/ 808 B978-0-12-391851-2.00009-X.
- Speer, K., S. Rintoul, and B. Sloyan, 2000: The Diabatic Deacon Cell. *Journal of Physical Oceanography*, 30 (12), 3212–3222.
- 811 Speer, K., and E. Tziperman, 1992: Rates of water mass formation in the North Atlantic Ocean. *Journal of Physical Oceanography*, 22, 93–104.

813 Talley, L., 2008: Freshwater transport estimates and the global overturning circulation: Shallow, ⁸¹⁴ deep and throughflow components. *Progress in Oceanography*, **78** (3), 257–303, doi:10.1016/j. 815 pocean.2008.05.001.

816 Talley, L., 2013: Closure of the Global Overturning Circulation Through the Indian, Pa-⁸¹⁷ cific, and Southern Oceans: Schematics and Transports. *Oceanography*, **26** (1), 80–97, doi: 818 10.5670/oceanog.2013.07.

- 819 Thacker, W. C., and R. B. Long, 1988: Fitting dynamics to data. *Journal of Geophysical Research*, 820 93 (C2), 1227–1240, doi:10.1029/JC093iC02p01227.
- 821 Thompson, A. F., and J.-B. Sallée, 2012: Jets and Topography: Jet Transitions and the Impact ⁸²² on Transport in the Antarctic Circumpolar Current. *Journal of Physical Oceanography*, 42 (6), 823 956–972, doi:10.1175/JPO-D-11-0135.1.
- 824 Trossman, D. S., L. Thompson, S. Mecking, and M. J. Warner, 2012: On the formation, ventilation,
- ⁸²⁵ and erosion of mode waters in the North Atlantic and Southern Oceans. *Journal of Geophysical*
- ⁸²⁶ *Research Oceans*, 117, doi:10.1029/2012JC008090.
- 827 Verdy, A., M. R. Mazloff, B. D. Cornuelle, and S. Y. Kim, 2014: Wind-Driven Sea Level Variabil-
- ⁸²⁸ ity on the California Coast: An Adjoint Sensitivity Analysis. *Journal of Physical Oceanography*,
- 829 $\,44$ (1), 297–318, doi:10.1175/JPO-D-13-018.1.

830 LIST OF FIGURES

831 Fig. 1. Southern Ocean potential vorticity and mixed layer depth. Shown are median values of annual minimum $\log_{10}(|PV|)$ across the years 1992-2011 as (a) a section through z=300 m. 833 The yellow contour indicates a section through the median mixed layer mask at z=300 m 834 depth, and the yellow dot-dash line indicates the location of the section in panel (b). (b) ss Section of median values of annual minimum $log_{10}(|PV|)$ across the years 1992-2011 at $100.5^\circ W$, with ensemble median values for the annual mean (solid white line) and ASO

FIG. 1. Southern Ocean potential vorticity and mixed layer depth. Shown are median values of annual minimum $log_{10}(|PV|)$ across the years 1992-2011 as (a) a section through $z=300$ m. The yellow contour indicates a section through the median mixed layer mask at z=300 m depth, and the yellow dot-dash line indicates the location of the section in panel (b). (b) Section of median values of annual minimum $log_{10}(|PV|)$ across the years 1992-2011 at 100.5◦*W*, with ensemble median values for the annual mean (solid white line) and ASO mean (dashed white line) mixed layer depths. PV has units of 1/ms. 894 895 896 897 898 899

FIG. 2. Defining the recently ventilated subsurface. Shading indicates the ensemble mean, annual mean tracer after 10 years of transport. Values have been normalized by the maximum value. Also shown is the mean mixed layer depth (dashed lines) and maximum mixed layer depth (dash-dot lines) over the length of the 10 year simulation. Solid lines indicate potential density surfaces ($\sigma_0 = 26.9$ kg m⁻³ to $\sigma_0 = 27.2$ kg m⁻³, spaced every 0.1 kg m⁻³). Tracer above the maximum mixed layer has been discarded. Although we discard the tracer lighter than $\sigma_0 = 26.9 \text{ kg m}^{-3}$ when defining the RVP, we kept this tracer in the figure for visualization purposes. Inset figures are column-integrated tracer histogram, which is normalized by the maximum column-integrated value, after discarding the tracer above the maximum mixed layer, along with a section through the ensemble median mixed layer mask at z=300 depth (yellow dashed contours) and a section through the ensemble median ventilated water mask at z=553 m (white solid contour). Sections are shown for (a) 33.5◦W, (b) 78.5◦E, (c) 131.5◦W, and (d) 80.5◦W. 900 901 902 903 904 905 906 907 908 909 910

FIG. 3. Vertical structure of the ensemble median RVP. Yellow regions are included in the RVP. A section through the ensemble median mixed layer mask at roughly 300 m is shown for reference (white solid contours). 911 912

FIG. 4. (a)-(f) Interannual variability in the RVP expressed as column-integrated volume in units of 10^{12} m³. The years listed in the titles indicate the dates of the tracer transport (i.e. released 1 Jan 1996 and transported until 31 Dec 2006). Percentages in the top left of each panel indicate volume changes relative to the 1996-2006 experiment. (g) Comparison of ventilated volume by depth level across release years. We used these tracer distributions to define the RVP in a self-consistent way; in this work, we do not examine the causes for the variation in RVP. 913 914 915 916 917 918

FIG. 5. Relative ensemble mean sensitivities of RVPh to density-compensated temperature anomalies (i.e. kinematic sensitivities). Also shown is a section through the RVP at roughly 553 m depth (black solid line), a section through the mixed layer mask at roughly 300 m depth (green dashed line), a depth contour (red solid line; 95 m for the left column, 552 m for the right column), and the SAF proxy (magenta solid line). All fields have been scaled by the grid cell thickness ∆*z* to allow for comparison between different depth levels. The plots have been further scaled by the global maximum sensitivity for the entire adjoint run to allow for comparison between different lags. 919 920 921 922 923 924 925

FIG. 6. Kinematic RMS sensitivities averaged over different volumes, indicating propagation timescales and relative magnitudes. Timescale increases from right to left, as indicated by more negative lags (i.e. longer lead times). Lines indicate ensemble means, and shading indicates one standard deviation across the ensemble. All values have been scaled by the same maximum of 0.0017 [°]*C*/[°]*C* and can be compared relative to each other. We show (a) the global mean sensitivity, (b) within the control volume (RVP) or not, (c) inside or outside of the 300 m MLD mask, (d), north or south of the SAF proxy, (e) in the upper 100 m in different basins, and (e) below 100 m in different basins. Note that the axes in panel (e) differ from the others. To define the basins, we used the standard ECCOv4 masks that are part of the gcmfaces package (see acknowledgements), and we chose 120[°]W as the dividing longitude between the East and West Pacific basins. South of 30◦S, the basins are roughly demarcated by longitude: Atlantic (70◦W-20◦E), Indian (20◦E-150◦E), Western Pacific (150◦E-120◦W), and Eastern Pacific (120◦W-70◦W). 926 927 928 929 930 931 932 933 934 935 936

FIG. 7. Relative ensemble mean sensitivities to changes in density (i.e. dynamic sensitivities). Scaling and units are the same as in Figure 5. 937 938

FIG. 8. RMS dynamic sensitivity time series. The panels and scaling are identical to those in Figure 6. Timescale increases from right to left. 939 940

FIG. 9. Schematic illustrating the main kinematic and dynamic sensitivities up to approximately 10 years lag. Chalk lines with arrows indicate kinematic sensitivities, with the vertical structure indicated by color. Weaker pathways by dashed lines. The circles connected by lines indicate where dynamic sensitivities resemble dipoles. Hot-spots are defined as regions that are either perennially sensitive or regions that feature a large seasonal amplitude throughout the 10-year timescale of the schematic. The reference contours include the RVP at 553 m (thick solid black line), the mixed layer mask at roughly 300 m depth (green dashed line), the SAF proxy (magenta solid line), and the 3000 m depth contour (thin gray lines). 941 942 943 944 945 946 947