



HAL
open science

Wave equation on certain noncompact symmetric spaces

Hong-Wei Zhang

► **To cite this version:**

| Hong-Wei Zhang. Wave equation on certain noncompact symmetric spaces. 2020. hal-02905019v2

HAL Id: hal-02905019

<https://hal.science/hal-02905019v2>

Preprint submitted on 24 Jul 2020 (v2), last revised 22 Jul 2021 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

WAVE EQUATION ON CERTAIN NONCOMPACT SYMMETRIC SPACES

HONG-WEI ZHANG

ABSTRACT. In this paper, we prove sharp pointwise kernel estimates and dispersive properties for the linear wave equation on noncompact Riemannian symmetric spaces G/K of any rank with G complex. As a consequence, we deduce Strichartz inequalities for a large family of admissible pairs and prove global well-posedness results for the corresponding semilinear equation with low regularity data as on hyperbolic spaces.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Pointwise estimates of the wave kernel	5
4. Dispersive estimates	16
4.1. Small time dispersive estimate	17
4.2. Large time dispersive estimate	18
5. Strichartz inequality and applications	19
5.1. Strichartz inequality	20
5.2. Global well-posedness in $L^p(\mathbb{R}, L^q(X))$	21
6. Further results on locally symmetric spaces	22
References	23

1. INTRODUCTION

This paper is devoted to study the dispersive properties of the linear wave equation on noncompact symmetric space G/K of any rank with G complex, and their applications to nonlinear Cauchy problems. This theory is well established for the wave equation on \mathbb{R}^d with $d \geq 3$:

$$\begin{cases} \partial_t^2 u(t, x) - \Delta_{\mathbb{R}^d} u(t, x) = F(t, x), \\ u(0, x) = f(x), \quad \partial_t|_{t=0} u(t, x) = g(x), \end{cases}$$

where the solutions u satisfy the Strichartz inequality: ¹

$$\|\nabla_{\mathbb{R} \times \mathbb{R}^d} u\|_{L^p(I; H^{-\sigma, q}(\mathbb{R}^d))} \lesssim \|f\|_{H^1(\mathbb{R}^d)} + \|g\|_{L^2(\mathbb{R}^d)} + \|F\|_{L^{\tilde{p}'}(I; H^{\tilde{\sigma}, \tilde{q}'}(\mathbb{R}^d))},$$

1991 *Mathematics Subject Classification.* 22E30, 35L05, 43A85, 47J35, 43A90.

Key words and phrases. Symmetric space, Wave equation, Dispersive estimate.

¹The symbol \lesssim , let us recall, means precisely that there exists a constant $0 < C < +\infty$ such that $\|\nabla_{\mathbb{R} \times \mathbb{R}^d} u\|_{L^p(I; H^{-\sigma, q}(\mathbb{R}^d))} \leq C(\|f\|_{H^1(\mathbb{R}^d)} + \|g\|_{L^2(\mathbb{R}^d)} + \|F\|_{L^{\tilde{p}'}(I; H^{\tilde{\sigma}, \tilde{q}'}(\mathbb{R}^d))})$, where \tilde{p}' is the dual exponent of \tilde{p} , defined by the formula $\frac{1}{\tilde{p}} + \frac{1}{\tilde{p}'} = 1$, and similarly for \tilde{q}' .

on any (possibly unbounded) interval $I \subseteq \mathbb{R}$ under the assumptions that

$$\sigma = \frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{q} \right), \quad \tilde{\sigma} = \frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{\tilde{q}} \right),$$

and the couples $(p, q), (\tilde{p}, \tilde{q}) \in (2, +\infty) \times [2, 2\frac{d-1}{d-3})$ fulfill the admissibility conditions:

$$\frac{1}{p} = \frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{q} \right), \quad \frac{1}{\tilde{p}} = \frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{\tilde{q}} \right).$$

Notice that this inequality also holds at the endpoint $(2, 2\frac{d-1}{d-3})$ when $d \geq 4$, but fails without additional assumptions when $d = 3$ (see [GV95] and [KT98] for more details). These estimates serve as a tool in order to find minimal regularity conditions on the initial data ensuring well-posedness for corresponding semilinear wave equations, which is addressed in [Kap94], and almost fully answered in [DGK01, GLS97, KT98, LS95].

Given the rich Euclidean theory, several attempts have been made in order to establish Strichartz inequality for dispersive equations in other settings. We are interested in Riemannian symmetric spaces of noncompact type where relevant questions are now well answered in the rank one case, see for instance [AP14, APV12, Fon97, MT11, MT12, Tat01] on hyperbolic spaces, and [APV15] on Damek-Ricci spaces. A first study of the wave equation on general symmetric spaces of higher rank was carried out in [Has11], where some non optimal estimates were obtained under the strong smoothness assumption.

In this paper, we prove sharp estimates for the non-shifted wave equation on noncompact symmetric spaces G/K of any rank with G complex. In this case, the Harish-Chandra \mathbf{c} -function and the spherical function have elementary expressions, which allow us to analyze accurately the wave kernel. For lack of such expressions in general, our present approach is limited to the class of symmetric spaces G/K with G complex, and maybe to a few other cases.

Consider the operator $W_t^\sigma = \tilde{D}^{-\sigma} e^{it\sqrt{-\Delta}}$ defined on the symmetric space $X = G/K$, for suitable exponents $\sigma \in \mathbb{C}$, where ρ denotes the half sum of positive roots, and $\tilde{D} = \sqrt{-\Delta - |\rho|^2 + \tilde{\rho}^2}$ is the differential operator with a fixed constant $\tilde{\rho} \geq |\rho|$. To avoid possible singularities, we may consider the analytic family of operators

$$\tilde{W}_t^\sigma = \frac{e^{\sigma^2}}{\Gamma(\frac{d+1}{2} - \sigma)} \tilde{D}^{-\sigma} e^{it\sqrt{-\Delta}},$$

in the vertical strip $0 \leq \operatorname{Re} \sigma \leq \frac{d+1}{2}$, see (3.2). We denote by $\tilde{\omega}_t^\sigma$ its K -bi-invariant convolution kernel. Our first and main result is the following pointwise kernel estimate, which summarizes Theorem 3.1 and Theorem 3.3 proved in Sect. 3.

Theorem 1.1 (Pointwise kernel estimates). *For all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$, there exists $N \in \mathbb{N}$ such that the following estimates hold for all $x \in X$:*

$$|\tilde{\omega}_t^\sigma(x)| \lesssim (1 + |H_x|)^N e^{-\langle \rho, H_x \rangle} \begin{cases} |t|^{-\frac{d-1}{2}} & \text{if } 0 < |t| < 1, \\ |t|^{-\frac{d}{2}} & \text{if } |t| \geq 1, \end{cases}$$

where $H_x \in \overline{\mathfrak{a}^+}$ denotes the middle component according to the Cartan decomposition of x .

Remark 1.2. These kernel estimates are sharp in time and similar results hold obviously in the easier case where $\operatorname{Re} \sigma > \frac{d+1}{2}$. The value of N will be specified in [Theorem 3.1](#) and [Theorem 3.3](#). However, the polynomial $(1 + |H_x|)^N$ is not crucial for further computations since there is an exponential decay following.

By interpolation arguments, we deduce our second result.

Theorem 1.3 (Dispersion property). *Let $2 < q, \tilde{q} < +\infty$ and $\sigma \geq (d+1) \max(\frac{1}{2} - \frac{1}{q}, \frac{1}{2} - \frac{1}{\tilde{q}})$. Then there exists a constant $C > 0$ such that the following dispersive estimates hold:*

$$\|\widetilde{W}_t^\sigma\|_{L^{\tilde{q}'}(X) \rightarrow L^q(X)} \leq C \begin{cases} |t|^{-(d-1) \max(\frac{1}{2} - \frac{1}{q}, \frac{1}{2} - \frac{1}{\tilde{q}})} & \text{if } 0 < |t| < 1, \\ |t|^{-\frac{d}{2}} & \text{if } |t| \geq 1. \end{cases}$$

Remark 1.4. At the endpoint $q = \tilde{q} = 2$, $t \mapsto e^{it\sqrt{-\Delta}}$ is a one-parameter group of unitary operators on $L^2(X)$.

Remark 1.5. These estimates, which are proved on real hyperbolic spaces in [\[AP14\]](#), extend straightforwardly to all the noncompact symmetric spaces of rank 1, and more generally to all Damek-Ricci spaces. In these cases, the large time decay is $|t|^{-\frac{3}{2}}$, where "3" corresponds to the so-called dimension at infinity of X . In higher rank, the dimension at infinity coincides with the manifold dimension d of G/K if and only if G is complex.

Remark 1.6. In [\[CGM02\]](#), M. Cowling, S. Giulini and S. Meda have described the L^p - L^q boundedness of a semi-group of operators related to the shifted Laplacian in constant time $t = 1$ on general noncompact symmetric spaces. However, sharp dispersive properties with a general $t \in \mathbb{R}^*$ are crucial for proving Strichartz type inequalities and for studying related PDE problems. Along the lines in [\[CGM02\]](#), we may derive a sharp dispersive inequality in small time $|t| \leq 1$. But as mentioned by these authors, their method does not yield good estimates in large time.

This paper is organized as follows. After recalling some basic notations and reviewing harmonic analysis on noncompact symmetric spaces in [Sect. 2](#), we derive pointwise kernel estimates in [Sect. 3](#). We prove the dispersive estimates by interpolation arguments in [Sect. 4](#). As a consequence, we deduce in [Sect. 5](#), Strichartz inequalities for a large family of admissible pairs and obtain well-posedness results for the associated semilinear wave equation. In [Sect. 6](#), we discuss similar results on a class of locally symmetric spaces.

The present paper is part of the author's Ph.D. thesis, supervised by J.-Ph. Anker and N. Burq. The author would like to thank them for sharing their knowledge and experience with him.

2. PRELIMINARIES

We review in this section some elementary notations and facts about noncompact symmetric spaces. We refer to [\[Hel62, Hel00\]](#) for more details.

Let G be a complex semisimple Lie group, connected, noncompact, with finite center, and K be a maximal compact subgroup of G . The homogeneous space

$X = G/K$ is a Riemannian symmetric space of noncompact type and dimension $d \geq 3$. Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition of the Lie algebra of G . The Killing form of \mathfrak{g} induces a K -invariant inner product on \mathfrak{p} , hence a G -invariant Riemannian metric on X , whose tangent space at the origin is identified with \mathfrak{p} .

Fix a maximal abelian subspace \mathfrak{a} in \mathfrak{p} . The rank of X is the dimension ℓ of \mathfrak{a} . Let $\Sigma \subset \mathfrak{a}$ be the root system of $(\mathfrak{g}, \mathfrak{a})$ which is reduced, and denote by W the Weyl group associated to Σ . Choose a set Σ^+ of positive roots, let $\mathfrak{a}^+ \subset \mathfrak{a}$ be the corresponding positive Weyl chamber and $\overline{\mathfrak{a}^+}$ be its closure. Notice that $d = \ell + 2|\Sigma^+|$ in our case. As usual, $\rho \in \mathfrak{a}^+$ denotes the half sum of positive roots, counted with their multiplicities, which is given in our case by $\rho = \sum_{\alpha \in \Sigma^+} \alpha$. It is well known that the spectrum of the negative Laplace-Beltrami operator $-\Delta$ on $L^2(X)$ is the half-line $[|\rho|^2, +\infty)$.

Denote by \mathfrak{n} the nilpotent Lie subalgebra of \mathfrak{g} associated with Σ^+ , and by N the corresponding Lie subgroup of G . Then we have the following two decompositions of G :

$$\begin{cases} G = N(\exp \mathfrak{a})K & (Iwasawa), \\ G = K(\exp \overline{\mathfrak{a}^+})K & (Cartan). \end{cases}$$

In the Cartan decomposition, the Haar measure on G writes

$$(2.1) \quad \int_G f(g)dg = \text{const.} \int_K dk_1 \int_{\mathfrak{a}^+} \delta(H)dH \int_K f(k_1(\exp H)k_2)dk_2,$$

where ²

$$\delta(H) = \prod_{\alpha \in \Sigma^+} (\sinh \langle \alpha, H \rangle)^2 \asymp \left\{ \prod_{\alpha \in \Sigma^+} \frac{\langle \alpha, H \rangle}{1 + \langle \alpha, H \rangle} \right\}^2 e^{\langle 2\rho, H \rangle} \quad \forall H \in \overline{\mathfrak{a}^+}.$$

Denote by $\mathcal{S}(K \backslash G/K)$ the Schwartz space of K -bi-invariant functions on G . The spherical Fourier transform \mathcal{H} is defined by

$$\mathcal{H}f(\lambda) = \int_G f(x)\varphi_\lambda(x)dx \quad \forall \lambda \in \mathfrak{a}, \forall f \in \mathcal{S}(K \backslash G/K).$$

Here $\varphi_\lambda \in \mathcal{C}^\infty(K \backslash G/K)$ denotes the spherical function of index $\lambda \in \mathfrak{a}_\mathbb{C}$, which is a radial eigenfunction of the Laplace-Beltrami operator Δ satisfying

$$(2.2) \quad \begin{cases} -\Delta\varphi_\lambda(x) = (|\lambda|^2 + |\rho|^2)\varphi_\lambda(x), \\ \varphi_\lambda(0) = 1. \end{cases}$$

In the noncompact case, the spherical functions are given by

$$(2.3) \quad \varphi_\lambda(x) = \int_K dk e^{\langle i\lambda + \rho, A(kx) \rangle} \quad \forall \lambda \in \mathfrak{a}_\mathbb{C},$$

where $A(kx)$ is the unique \mathfrak{a} -component in the Iwasawa decomposition of kx . It satisfies the basic estimate

$$(2.4) \quad |\varphi_\lambda(x)| \leq \varphi_0(x) \quad \forall \lambda \in \mathfrak{a}, \forall x \in X,$$

²The symbol $f \asymp g$ between two nonnegative expressions means that there exist two constants $0 < c_1 \leq c_2 < +\infty$ such that $c_1g \leq f \leq c_2g$.

where

$$(2.5) \quad \varphi_0(\exp H) = \prod_{\alpha \in \Sigma^+} \frac{\langle \alpha, H \rangle}{\sinh \langle \alpha, H \rangle} \asymp \left\{ \prod_{\alpha \in \Sigma^+} (1 + \langle \alpha, H \rangle) \right\} e^{-\langle \rho, H \rangle} \quad \forall H \in \overline{\mathfrak{a}^+}.$$

If G is complex, we have also

$$(2.6) \quad \varphi_\lambda(x) = \varphi_0(x) \int_K dk e^{i\langle (\text{Ad } k) \cdot \lambda, x \rangle}.$$

Denote by $\mathcal{S}(\mathfrak{a})^W$ the subspace of W -invariant functions in the Schwartz space $\mathcal{S}(\mathfrak{a})$. Then \mathcal{H} is an isomorphism between $\mathcal{S}(K \backslash G / K)$ and $\mathcal{S}(\mathfrak{a})^W$. The inverse spherical Fourier transform is defined by

$$(2.7) \quad f(x) = \text{const.} \int_{\mathfrak{a}} d\lambda \mathcal{H}f(\lambda) \varphi_\lambda(x) |\mathbf{c}(\lambda)|^{-2} \quad \forall x \in G, \forall f \in \mathcal{S}(\mathfrak{a})^W,$$

where $\mathbf{c}(\lambda)$ is the Harish-Chandra \mathbf{c} -function. If G is complex, the Plancherel density reads

$$(2.8) \quad |\mathbf{c}(\lambda)|^{-2} = \pi(\lambda)^2 = \prod_{\alpha \in \Sigma^+} \langle \alpha, \lambda \rangle^2,$$

and in particular, the inverse spherical Fourier transform (2.7) becomes

$$f(x) = \text{const.} \varphi_0(x) \int_{\mathfrak{p}} d\lambda \mathcal{H}f(\lambda) e^{-i\langle (\text{Ad } k) \cdot \lambda, x \rangle}.$$

Recall at last the asymptotic expansion of the Bessel function of the first kind:

$$(2.9) \quad J_m(r) \sim r^{-1/2} e^{ir} \sum_{j=0}^{\infty} a_j(m) r^{-j} + r^{-1/2} e^{-ir} \sum_{j=0}^{\infty} b_j(m) r^{-j} \quad \text{as } r \rightarrow +\infty,$$

where $a_j(m)$ and $b_j(m)$ are suitable coefficients.

3. POINTWISE ESTIMATES OF THE WAVE KERNEL

In this section, we derive pointwise estimates for the K -bi-invariant convolution kernel ω_t^σ of the operator $W_t^\sigma = \tilde{D}^{-\sigma} e^{it\sqrt{-\Delta}}$ on the symmetric space X :

$$W_t^\sigma f(x) = f * \omega_t^\sigma(x) = \int_G dy \omega_t^\sigma(y^{-1}x) f(y),$$

for suitable exponents $\sigma \in \mathbb{C}$. Via the spherical Fourier transform and (2.2), the negative Laplace-Beltrami operator $-\Delta$ corresponds to $|\lambda|^2 + |\rho|^2$, hence the operator \tilde{D} to $\sqrt{|\lambda|^2 + \tilde{\rho}^2}$. The inverse spherical Fourier transform implies that

$$(3.1) \quad \omega_t^\sigma(x) = \text{const.} \int_{\mathfrak{a}} d\lambda |\mathbf{c}(\lambda)|^{-2} (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} \varphi_\lambda(x).$$

As already observed for hyperbolic spaces (see for instance [APV12]), the classical Euclidean rescaling method fails since this kernel has different behaviors depending whether t is small or large. We will prove sharp time pointwise estimates for this kernel along the lines of [AP14].

Consider smooth even cut-off functions χ_0 and χ_∞ on \mathbb{R} such that

$$\chi_0(r) + \chi_\infty(r) = 1 \quad \text{and} \quad \begin{cases} \chi_0(r) = 1 & \text{if } |r| \leq 1, \\ \chi_\infty(r) = 1 & \text{if } |r| \geq 2. \end{cases}$$

Let us split up

$$\begin{aligned} \omega_t^\sigma(x) &= \omega_t^{\sigma,0}(x) + \omega_t^{\sigma,\infty}(x) \\ &= \text{const.} \int_{\mathfrak{a}} d\lambda \chi_0^\rho(\lambda) |\mathbf{c}(\lambda)|^{-2} (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} \varphi_\lambda(x) \\ &\quad + \text{const.} \int_{\mathfrak{a}} d\lambda \chi_\infty^\rho(\lambda) |\mathbf{c}(\lambda)|^{-2} (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} \varphi_\lambda(x) \end{aligned}$$

where $\chi_0^\rho(\lambda) = \chi_0(|\lambda|/|\rho|)$ and $\chi_\infty^\rho(\lambda) = \chi_\infty(|\lambda|/|\rho|)$ are radial cut-off functions defined on \mathfrak{a} . We shall see later that the kernel $\omega_t^{\sigma,\infty}(x)$ has a logarithmic singularity on the sphere $|x| = t$ when $\sigma = \frac{d+1}{2}$. To get around this problem, we consider the analytic family of operators

$$(3.2) \quad \widetilde{W}_t^{\sigma,\infty} = \frac{e^{\sigma^2}}{\Gamma(\frac{d+1}{2} - \sigma)} \chi_\infty^\rho(\sqrt{-\Delta - |\rho|^2}) \widetilde{D}^{-\sigma} e^{it\sqrt{-\Delta}},$$

in the vertical strip $0 \leq \text{Re } \sigma \leq \frac{d+1}{2}$ and the corresponding kernels

$$\widetilde{\omega}_t^{\sigma,\infty}(x) = \frac{e^{\sigma^2}}{\Gamma(\frac{d+1}{2} - \sigma)} \int_{\mathfrak{a}} d\lambda \chi_\infty^\rho(\lambda) |\mathbf{c}(\lambda)|^{-2} (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} \varphi_\lambda(x).$$

Notice that the Gamma function will allow us to deal with the boundary point $\sigma = \frac{d+1}{2}$, while the exponential function ensures boundedness at infinity in the vertical strip.

Theorem 3.1. *Let $\sigma \in \mathbb{C}$. The following pointwise estimates hold for the kernel $\omega_t^{\sigma,0}$:*

(i) *For all $t \in \mathbb{R}^*$, we have*

$$|\omega_t^{\sigma,0}(x)| \lesssim \varphi_0(x).$$

(ii) *Assume that $|t| \geq 1$.*

(a) *If $\frac{|x|}{|t|} \geq \frac{1}{2}$, then*

$$|\omega_t^{\sigma,0}(x)| \lesssim |t|^{-N} (1 + |x|)^N \varphi_0(x),$$

for every $N \in \mathbb{N}$.

(b) *If $\frac{|x|}{|t|} \leq \frac{1}{2}$, then*

$$|\omega_t^{\sigma,0}(x)| \lesssim |t|^{-\frac{d}{2}} (1 + |x|)^{\frac{d-\ell}{2}} \varphi_0(x).$$

Proof. By symmetry we may assume that $t > 0$. Recall that

$$(3.3) \quad \omega_t^{\sigma,0}(x) = \text{const.} \int_{\mathfrak{a}} d\lambda \chi_0^\rho(\lambda) |\mathbf{c}(\lambda)|^{-2} (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} \varphi_\lambda(x).$$

(i) follows from the representation (2.8) and the estimate (2.4) that

$$|\omega_t^{\sigma,0}(x)| \lesssim \int_{|\lambda| < 2|\rho|} d\lambda |\lambda|^{d-\ell} |\varphi_\lambda(x)| \lesssim \varphi_0(x).$$

Moreover (ii).(a) is straightforward since $\frac{|x|}{t} \geq \frac{1}{2}$.

We prove (ii).(b) by substituting in (3.3) the integral representation (2.3) of φ_λ and the expression (2.8) of $|\mathbf{c}(\lambda)|^{-2}$. Specifically,

$$\omega_t^{\sigma,0}(x) = \text{const.} \int_K dk e^{\langle \rho, A(kx) \rangle} \int_{\mathfrak{a}} d\lambda a(\lambda) e^{it\sqrt{|\lambda|^2 + |\rho|^2} + i\langle \lambda, A(kx) \rangle},$$

where $a(\lambda) = \chi_0^\rho(\lambda)\pi(\lambda)^2(|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}}$. Since

$$\int_K dk e^{\langle \rho, A(kx) \rangle} = \varphi_0(x),$$

it remains to estimate the oscillatory integral

$$I_0(t, x) = \int_{\mathfrak{a}} d\lambda a(\lambda) e^{it\psi(\lambda)}$$

with amplitude $a(\lambda)$ and phase

$$\psi(\lambda) = \sqrt{|\lambda|^2 + |\rho|^2} + \left\langle \frac{A}{t}, \lambda \right\rangle,$$

where $A = A(kx)$ is a vector in \mathfrak{a} such that $|A(kx)| \leq |x|$. This is achieved by carrying out in our case, the stationary phase analysis described in [Ste93, VIII.2.3]. The critical point λ_0 of the phase ψ is given by

$$(|\lambda_0|^2 + |\rho|^2)^{-\frac{1}{2}} \lambda_0 = -\frac{A}{t}.$$

Hence

$$(3.4) \quad |\lambda_0| = |\rho| \frac{|A|}{t} \left(1 - \frac{|A|^2}{t^2}\right)^{-\frac{1}{2}} \leq |\rho| \frac{|x|}{t} \left(1 - \frac{|x|^2}{t^2}\right)^{-\frac{1}{2}}.$$

Denote by $B(\lambda_0, \eta)$ the ball in \mathfrak{a} centered at λ_0 :

$$B(\lambda_0, \eta) = \{\lambda \in \mathfrak{a} \mid |\lambda - \lambda_0| \leq \eta\},$$

where the radius η will be chosen later. Notice that if $\frac{|x|}{t} \leq \frac{1}{2}$, then $|\lambda_0| < \frac{|\rho|}{\sqrt{3}}$ and $|\lambda| < |\rho| + \eta$ for all $\lambda \in B(\lambda_0, \eta)$. Let P_λ be the projection onto the vector spanned by $\frac{\lambda}{|\lambda|}$. Then $|\lambda|^2 P_\lambda = \lambda \otimes \lambda$ and the Hessian matrix of ψ is given by

$$\begin{aligned} \text{Hess } \psi(\lambda) &= (|\lambda|^2 + |\rho|^2)^{-\frac{1}{2}} I_\ell - (|\lambda|^2 + |\rho|^2)^{-\frac{3}{2}} \lambda \otimes \lambda \\ &= (|\lambda|^2 + |\rho|^2)^{-\frac{3}{2}} \{ |\rho|^2 P_\lambda + (|\lambda|^2 + |\rho|^2)(I_\ell - P_\lambda) \} \\ &= (|\lambda|^2 + |\rho|^2)^{-\frac{3}{2}} \left(\begin{array}{c|c} |\rho|^2 & 0 \\ \hline 0 & (|\lambda|^2 + |\rho|^2) I_{\ell-1} \end{array} \right), \end{aligned}$$

which is a positive definite symmetric matrix. Hence λ_0 is a nondegenerate critical point. Since $\nabla\psi(\lambda_0) = 0$, we have

$$\psi(\lambda) - \psi(\lambda_0) = (\lambda - \lambda_0)^T \underbrace{\left\{ \int_0^1 ds (1-s) \text{Hess } \psi(\lambda_0 + s(\lambda - \lambda_0)) \right\}}_{=M(\lambda)} (\lambda - \lambda_0),$$

where $M(\lambda)$ belongs, for every $\lambda \in B(\lambda_0, \eta)$, to a compact subset of the set of positive definite symmetric matrices. We introduce a new variable $\mu = M(\lambda)^{\frac{1}{2}}(\lambda -$

λ_0), then $|\mu|^2 = \psi(\lambda) - \psi(\lambda_0)$ and $\mu = 0$ if and only if $\lambda = \lambda_0$. Notice that for every $k \in \mathbb{N}$, there exists $C_k > 0$ such that

$$(3.5) \quad |\nabla^k M(\lambda)^{\frac{1}{2}}| \leq C_k \quad \forall \lambda \in B(\lambda_0, \eta).$$

Denote by $J(\lambda)$ the Jacobian matrix such that $d\mu = \det[J(\lambda)] d\lambda$, then we can choose $\eta > 0$ small enough such that

$$(3.6) \quad \det[J(\lambda)] > \frac{1}{2} \det\left[M(\lambda)^{\frac{1}{2}}\right] \quad \forall \lambda \in B(\lambda_0, \eta).$$

Now let us divide the study of $I_0(t, x)$ into two parts, corresponding to the decomposition

$$I_0(t, x) = \underbrace{\int_{\mathbf{a}} d\lambda \chi_{\eta}^{-}(\lambda) a(\lambda) e^{it\psi(\lambda)}}_{=I_0^{-}(t, x)} + \underbrace{\int_{\mathbf{a}} d\lambda \chi_{\eta}^{+}(\lambda) a(\lambda) e^{it\psi(\lambda)}}_{=I_0^{+}(t, x)},$$

associated with the smooth cut-off functions

$$\chi_{\eta}^{-}(\lambda) = \chi_0\left(\frac{|\lambda - \lambda_0|}{2\eta}\right) \quad \text{and} \quad \chi_{\eta}^{+}(\lambda) = \chi_{\infty}\left(\frac{|\lambda - \lambda_0|}{2\eta}\right).$$

Estimate of $I_0^{-}(t, x)$. Notice that $\text{supp } \chi_{\eta}^{-} \subset B(\lambda_0, \eta)$. There exist $0 < \tilde{\eta}_1 \leq \tilde{\eta}_2$ such that $\mu \in B(0, \tilde{\eta}_1)$ implies $\lambda \in B(\lambda_0, \eta)$, and $\lambda \in B(\lambda_0, \eta)$ implies $\mu \in B(0, \tilde{\eta}_2)$. By substituting $\psi(\lambda) = |\mu|^2 + \psi(\lambda_0)$, we get

$$I_0^{-}(t, x) = e^{it\psi(\tilde{\lambda})} \int_{\mathbf{a}} d\mu \tilde{a}(\lambda(\mu)) e^{it|\mu|^2}$$

where the amplitude

$$(3.7) \quad \tilde{a}(\lambda(\mu)) = \chi_{\eta}^{-}(\lambda(\mu)) \chi_0^{\rho}(\lambda(\mu)) \pi(\lambda(\mu))^2 (|\lambda(\mu)|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} \det[J(\lambda(\mu))]^{-1}$$

is smooth and compactly supported in $B(0, \tilde{\eta}_2)$. We deduce, from (3.5) and (3.6) that $\tilde{a}(\lambda(\mu))$ is bounded, together with all its derivatives. Let $\tilde{\chi} \in \mathcal{C}_c^{\infty}(\mathbf{a})$ be a bump function such that $\tilde{\chi} = 1$ on $B(0, \tilde{\eta}_2)$. Let $M \geq \frac{d}{2}$ be an integer. Then

$$I_0^{-}(t, x) = e^{it\psi(\lambda_0)} \int_{\mathbf{a}} d\mu e^{it|\mu|^2} e^{-|\mu|^2} [e^{|\mu|^2} \tilde{a}(\lambda(\mu))] \tilde{\chi}(\mu),$$

where the coefficients of the Taylor expansion of $e^{|\mu|^2} \tilde{a}(\lambda(\mu))$ at the origin:

$$e^{|\mu|^2} \tilde{a}(\lambda(\mu)) = \sum_{|k| \leq 2M} c_k \mu^k + R_{2M}(\mu),$$

satisfy

$$(3.8) \quad |c_k| \lesssim \left(\frac{|x|}{t}\right)^{d-\ell},$$

and the remainder

$$(3.9) \quad |\nabla^k R_{2M}(\mu)| \lesssim |\mu|^{2M+1-k} \quad \forall 0 \leq k \leq 2M+1,$$

according to (3.4) and (3.7). By substituting this expansion in the above integral, $I_0^-(t, x)$ is equal to the sum of following three terms:

$$I_1 = \sum_{|k| \leq 2M} c_k \int_{\mathfrak{a}} d\mu e^{it|\mu|^2} e^{-|\mu|^2} \mu^k,$$

$$I_2 = \int_{\mathfrak{a}} d\mu e^{it|\mu|^2} R_{2M}(\mu) e^{-|\mu|^2} \tilde{\chi}(\mu),$$

and

$$I_3 = \int_{\mathfrak{a}} d\mu e^{it|\mu|^2} \left(\sum_{|k| \leq 2M} c_k \mu^k \right) e^{-|\mu|^2} (\tilde{\chi}(\mu) - 1).$$

In order to estimate I_1 , we write the integral as a product

$$I_1 = \sum_{|k| \leq 2M} c_k \prod_{j=1}^{\ell} \int_{-\infty}^{+\infty} d\mu_j e^{it\mu_j^2} e^{-\mu_j^2} \mu_j^{k_j},$$

where

$$\int_{-\infty}^{+\infty} d\mu_j e^{-(1-it)\mu_j^2} \mu_j^{k_j} = 0$$

if k_j is odd, while

$$\int_{-\infty}^{+\infty} d\mu_j e^{-(1-it)\mu_j^2} \mu_j^{k_j} = 2(1-it)^{-\frac{k_j+1}{2}} \int_0^{+\infty} dz_j e^{-z_j^2} z_j^{k_j}$$

by a change of contour if k_j is even. From (3.8), we obtain

$$|I_1| \lesssim \left(\frac{|x|}{t}\right)^{d-\ell} t^{-\frac{\ell}{2}} = t^{-\frac{d}{2}} \left(\frac{|x|}{\sqrt{t}}\right)^{d-\ell} \lesssim t^{-\frac{d}{2}} |x|^{-\frac{d-\ell}{2}},$$

since $\frac{|x|}{t} < \frac{1}{2}$. Next, we perform M integrations by parts based on

$$(3.10) \quad e^{it|\mu|^2} = -\frac{i}{2t} \sum_{j=1}^{\ell} \frac{\mu_j}{|\mu|^2} \frac{\partial}{\partial \mu_j} e^{it|\mu|^2}$$

and obtain

$$I_2 = O(t^{-M}),$$

according to (3.9). Finally, as $\mu \mapsto \left(\sum_{|k| \leq 2M} c_k \mu^k\right) e^{-|\mu|^2} (\tilde{\chi}(\mu) - 1)$ is exponentially decreasing and vanishes near the origin, we perform $N \geq \frac{d}{2}$ integrations by parts based on (3.10) again and obtain

$$I_3 = O(t^{-N}).$$

By summing up the estimates of I_1 , I_2 and I_3 , we deduce that

$$(3.11) \quad |I_0^-(t, x)| \lesssim t^{-\frac{d}{2}} (1 + |x|)^{\frac{d-\ell}{2}}.$$

Estimate of $I_0^+(t, x)$. Since the phase ψ has a unique critical point λ_0 , then for all $\lambda \in \text{supp } \chi_\eta^+$, $\nabla\psi(\lambda) \neq 0$. In order to get a large time decay, we estimate the oscillatory integral $I_0^+(t, x)$ by using several integrations by parts based on

$$(3.12) \quad e^{it\psi(\lambda)} = \frac{1}{it} (\tilde{\psi}(\lambda))^{-1} \sum_{j=1}^{\ell} \lambda_j \frac{\partial}{\partial \lambda_j} e^{it\psi(\lambda)},$$

where

$$\tilde{\psi}(\lambda) = |\lambda|^2 (|\lambda|^2 + |\rho|^2)^{-\frac{1}{2}} + \langle \frac{A}{t}, \lambda \rangle$$

is a nonzero symbol of order zero for all $\lambda \in \text{supp } \chi_\eta^+$. After performing N such integrations by parts, $I_0^+(t, x)$ becomes

$$I_0^+(t, x) = \text{const.} (it)^{-N} \underbrace{\int_{\mathfrak{a}} d\lambda e^{it\psi(\lambda)} \left\{ - \sum_{j=1}^{\ell} \frac{\partial}{\partial \lambda_j} \circ \frac{\lambda_j}{\tilde{\psi}(\lambda)} \right\}^N \left\{ \chi_\eta^+(\lambda) a(\lambda) \right\}}_{<+\infty},$$

for every $N \in \mathbb{N}$. The last integral is finite since the amplitude a is supported in $B(0, 2|\rho|)$ thanks to the cut-off χ_0^ρ . Hence

$$(3.13) \quad |I_0^+(t, x)| \lesssim t^{-N},$$

for every integer $N \geq 0$. According to (3.11) and (3.13), we conclude that

$$|\omega_t^{\sigma, 0}(x)| \lesssim |I_0(t, x)| \int_K dk e^{-\langle \rho, A(xk) \rangle} \lesssim t^{-\frac{d}{2}} |x|^{\frac{d-\ell}{2}} \varphi_0(x).$$

□

Remark 3.2. In the proof of [Theorem 3.1](#), we have replaced the spherical function $\varphi_\lambda(x)$ by its general integral expression (2.3). Moreover, this proof also works without using the explicit expression (2.8) of the density $|\mathbf{c}(\lambda)|^{-2}$. Thus we can derive similar estimates for any symmetric space G/K by the same approach.

Theorem 3.3. *Let $\sigma \in \mathbb{C}$ with $\text{Re } \sigma = \frac{d+1}{2}$. The following pointwise estimates hold for the kernel $\tilde{\omega}_\infty^\sigma(t, x)$, for all $t \in \mathbb{R}^*$ and $x \in \mathfrak{a}$:*

$$|\tilde{\omega}_t^{\sigma, \infty}(x)| \lesssim \begin{cases} |t|^{-\frac{d-1}{2}} \varphi_0(x) & \text{if } |t| < 1, \\ |t|^{-N} (1 + |x|)^N \varphi_0(x) & \text{if } |t| \geq 1, \end{cases}$$

for every $N \in \mathbb{N}$.

Remark 3.4. Similar estimates hold in the easier case $\text{Re } \sigma > \frac{d+1}{2}$. By symmetry we assume again that $t > 0$. We divide the proof into two parts, depending whether $\frac{|x|}{t} \geq \frac{1}{2}$ or not.

Proof for $\frac{|x|}{t} \geq \frac{1}{2}$. Substituting the integral representation (2.6) of φ_λ allows us to write the kernel as

$$\tilde{\omega}_t^{\sigma, \infty}(x) = \text{const.} \frac{e^{\sigma^2}}{\Gamma((d+1)/2 - \sigma)} \varphi_0(x) \underbrace{\int_{\mathfrak{p}} d\lambda \chi_\infty^\rho(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} e^{-i\langle x, \lambda \rangle}}_{=I_\infty(t, x)}.$$

We split up $I_\infty(t, x)$ into two parts

$$I_\infty(t, x) = \underbrace{\int_{\mathfrak{p}} d\lambda \chi_0^\rho(t\lambda) \chi_\infty^\rho(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} e^{-i\langle x, \lambda \rangle}}_{I_\infty^-(t, x)} + \underbrace{\int_{\mathfrak{p}} d\lambda \chi_\infty^\rho(t\lambda) \chi_\infty^\rho(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{|\lambda|^2 + |\rho|^2}} e^{-i\langle x, \lambda \rangle}}_{I_\infty^+(t, x)},$$

by using the smooth radial cut-off functions $\chi_0^\rho(t\lambda)$ and $\chi_\infty^\rho(t\lambda)$. The first integral $I_\infty^-(t, x)$ vanishes if $t \geq 2$ and is easily estimated as follows, if $0 < t < 2$:

$$\begin{aligned} |I_\infty^-(t, x)| &\lesssim \int_{\mathfrak{p}} d\lambda \chi_0^\rho(t\lambda) \chi_\infty^\rho(\lambda) |\lambda|^{-\text{Re } \sigma} \\ &\leq t^{-\frac{d-1}{2}} \int_{\mathfrak{p}} d\lambda \chi_0^\rho(\lambda) |\lambda|^{-\frac{d+1}{2}} \lesssim t^{-\frac{d-1}{2}} \end{aligned}$$

for all $\sigma \in \mathbb{C}$ with $\text{Re } \sigma = \frac{d+1}{2}$. In order to study the second integral $I_\infty^+(t, x)$, we introduce polar coordinates and rewrite

$$(3.14) \quad I_\infty^+(t, x) = \int_0^{+\infty} dr \chi_\infty^\rho(tr) \chi_\infty^\rho(r) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\sqrt{r^2 + |\rho|^2}} \int_{|\lambda|=r} d\sigma(\lambda) e^{-i\langle x, \lambda \rangle}.$$

Here the inner integral is a modified Bessel function. Specifically

$$(3.15) \quad \int_{|\lambda|=r} d\sigma(\lambda) e^{-i\langle x, \lambda \rangle} = r^{\frac{d}{2}} |x|^{\frac{2-d}{2}} J_{\frac{d-2}{2}}(r|x|),$$

where $J_{(d-2)/2}$ denotes the classical Bessel function of the first kind. Since $I_\infty^+(t, x)$ vanishes if $tr < |\rho|$, we may restrict ourselves to $tr \geq |\rho|$, hence $r|x| \geq \frac{rt}{2} \geq \frac{|\rho|}{2}$. Then from (2.9), there exist constants C_d^+ and C_d^- such that

$$(3.16) \quad J_{\frac{d-2}{2}}(r|x|) = C_d^+(r|x|)^{-1/2} e^{ir|x|} + C_d^-(r|x|)^{-1/2} e^{-ir|x|} + R(r|x|),$$

where the remainder $R(r|x|)$ is $O((r|x|)^{-\frac{3}{2}})$. Substituting (3.15) and (3.16) in (3.14) implies

$$\begin{aligned} I_\infty^+(t, x) &= I_\infty^{+,1}(t, x) + I_\infty^{+,2}(t, x) + I_\infty^{+,3}(t, x) \\ &= C_d^+ |x|^{-\frac{d-1}{2}} \int_0^{+\infty} dr \chi_\infty^\rho(tr) \chi_\infty^\rho(r) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} r^{\frac{d-1}{2}} e^{it\sqrt{r^2+|\rho|^2}} e^{i|x|r} \\ &\quad + C_d^- |x|^{-\frac{d-1}{2}} \int_0^{+\infty} dr \chi_\infty^\rho(tr) \chi_\infty^\rho(r) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} r^{\frac{d-1}{2}} e^{it\sqrt{r^2+|\rho|^2}} e^{-i|x|r} \\ &\quad + |x|^{-\frac{d-2}{2}} \int_0^{+\infty} dr \chi_\infty^\rho(tr) \chi_\infty^\rho(r) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} r^{\frac{d}{2}} e^{it\sqrt{r^2+|\rho|^2}} R(r|x|). \end{aligned}$$

On the one hand, the last integral is estimated as follows:

$$|I_\infty^{+,3}(t, x)| \lesssim |x|^{-\frac{d+1}{2}} \int_0^{+\infty} dr \chi_\infty^\rho(tr) \chi_\infty^\rho(r) r^{-2},$$

for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$. Hence

$$(3.17) \quad |I_\infty^{+,3}(t, x)| \lesssim |x|^{-\frac{d+1}{2}} \int_{|\rho|}^{+\infty} dr r^{-2} \lesssim |x|^{-\frac{d+1}{2}}$$

for large time and

$$(3.18) \quad |I_\infty^{+,3}(t, x)| \lesssim |x|^{-\frac{d+1}{2}} \int_{|\rho|t^{-1}}^{+\infty} dr r^{-2} \lesssim t|x|^{-\frac{d+1}{2}} \lesssim |x|^{-\frac{d-1}{2}}$$

for small time.

On the other hand, in order to estimate $I_\infty^{+,1}(t, x)$ and $I_\infty^{+,2}(t, x)$, it is sufficient to study the integral

$$\tilde{I}_\infty^\pm(t, x) = \int_0^{+\infty} dr \chi_\infty^\rho(r) \chi_\infty^\rho(tr) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} r^{\frac{d-1}{2}} e^{it\sqrt{r^2+|\rho|^2}} e^{\pm i|x|r}.$$

Notice that

$$(3.19) \quad \chi_\infty^\rho(r) \chi_\infty^\rho(tr) = \begin{cases} \chi_\infty^\rho(r) & \text{if } t \geq 2, \\ \chi_\infty^\rho(tr) & \text{if } t \leq \frac{1}{2}. \end{cases}$$

We split up $\tilde{I}_\infty^\pm(t, x)$ into three parts, corresponding to the decomposition

$$\tilde{I}_\infty^\pm(t, x) = \tilde{I}_\infty^{\pm,1}(t, x) + \tilde{I}_\infty^{\pm,2}(t, x) + \tilde{I}_\infty^{\pm,3}(t, x)$$

where

$$\tilde{I}_\infty^{\pm,1}(t, x) = \int_0^{+\infty} dr \chi_\infty^\rho(r) \chi_\infty^\rho(tr) (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} r^{\frac{d-1}{2}} \{e^{it\sqrt{r^2+|\rho|^2}} - e^{itr}\} e^{\pm i|x|r},$$

$$\tilde{I}_\infty^{\pm,2}(t, x) = \int_0^{+\infty} dr \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{\frac{d-1}{2}} \{(r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} - r^{-\sigma}\} e^{i(t \pm |x|)r},$$

and

$$\tilde{I}_\infty^{\pm,3}(t, x) = \int_0^{+\infty} dr \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i\operatorname{Im} \sigma} e^{i(t \pm |x|)r},$$

for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$.

Estimates of $\tilde{I}_\infty^{+,1}$ and $\tilde{I}_\infty^{+,2}$. The first two parts are easily estimated. By using

$$e^{it\sqrt{r^2+|\rho|^2}} - e^{itr} = O\left(\frac{t}{r}\right) \quad \text{and} \quad (r^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} - r^{-\sigma} = O(r^{-\operatorname{Re}\sigma-2})$$

for all $r \in \operatorname{supp} \chi_\infty^\rho$, we obtain

$$(3.20) \quad \tilde{I}_\infty^{+,1}(t, x) = O(t) \quad \text{and} \quad \tilde{I}_\infty^{+,2}(t, x) = O(1)$$

for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$.

Estimate of $\tilde{I}_\infty^{+,3}$. We claim that

$$(3.21) \quad |\tilde{I}_\infty^{+,3}(t, x)| \lesssim \underbrace{|\operatorname{Im} \sigma| + \frac{1}{|\operatorname{Im} \sigma|}}_{C_\sigma}.$$

Denoting by $\xi = t \pm |x|$, we divide our estimate into two parts, depending whether $|\xi| \geq \frac{1}{2|\rho|}$ or not.

- If $|\xi| \geq \frac{1}{2|\rho|}$, we perform an integration by parts based on $e^{i\xi r} = \frac{1}{i\xi} \frac{\partial}{\partial r} e^{i\xi r}$. Then

$$(3.22) \quad \tilde{I}_\infty^{+,3}(t, x) = \frac{i}{\xi} \int_0^{+\infty} dr e^{i\xi r} \frac{\partial}{\partial r} \{ \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i\operatorname{Im} \sigma} \}.$$

If the derivative hits either $\chi_\infty^\rho(r)$ or $r^{-1-i\operatorname{Im} \sigma}$, then the corresponding contributions to (3.22) are bounded by

$$C(1 + |\operatorname{Im} \sigma|).$$

If the derivative hits $\chi_\infty^\rho(tr)$, which matters only in the case $t < 2$, according to (3.19), then the corresponding contribution to (3.22) is bounded by

$$Ct \int_{|\rho|t^{-1}}^{2|\rho|t^{-1}} dr r^{-1} < 2(\ln 2)C.$$

Hence $\tilde{I}_\infty^{+,3}(t, x) = O(1 + |\operatorname{Im} \sigma|)$ for all $t > 0$ and $x \in \mathfrak{a}$ satisfying $\frac{|x|}{t} \geq \frac{1}{2}$ and $|t \pm |x|| \geq \frac{1}{2|\rho|}$.

- If $|\xi| < \frac{1}{2|\rho|}$, we split up

$$\tilde{I}_\infty^{+,3}(t, x) = \left(\int_0^{\frac{1}{|\xi|}} + \int_{\frac{1}{|\xi|}}^{+\infty} \right) dr \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i\operatorname{Im} \sigma} e^{i\xi r},$$

We estimate both terms by performing different integration by parts. On the one hand,

$$\begin{aligned}
& \int_0^{\frac{1}{|\xi|}} dr r^{-1-i \operatorname{Im} \sigma} \chi_\infty^\rho(r) \chi_\infty^\rho(tr) e^{i\xi r} \\
&= \frac{i}{\operatorname{Im} \sigma} \underbrace{r^{-i \operatorname{Im} \sigma} \chi_\infty^\rho(r) \chi_\infty^\rho(tr) e^{i\xi r} \Big|_{r=0}^{r=\frac{1}{|\xi|}}}_{= O(1)} \\
&\quad - \frac{i}{\operatorname{Im} \sigma} \underbrace{\int_0^{\frac{1}{|\xi|}} dr r^{-i \operatorname{Im} \sigma} \frac{\partial}{\partial r} \{ \chi_\infty^\rho(r) \chi_\infty^\rho(tr) e^{i\xi r} \}}_{= O(1) \quad (*)} \\
&= O\left(\frac{1}{|\operatorname{Im} \sigma|}\right).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \int_{\frac{1}{|\xi|}}^{+\infty} dr e^{i\xi r} \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i \operatorname{Im} \sigma} \\
&= \frac{1}{i\xi} \underbrace{e^{i\xi r} \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i \operatorname{Im} \sigma} \Big|_{r=\frac{1}{|\xi|}}^{r=+\infty}}_{= O(1)} \\
&\quad + \frac{i}{\xi} \underbrace{\int_{\frac{1}{|\xi|}}^{+\infty} dr e^{i\xi r} \frac{\partial}{\partial r} \{ \chi_\infty^\rho(r) \chi_\infty^\rho(tr) r^{-1-i \operatorname{Im} \sigma} \}}_{= O(1+|\operatorname{Im} \sigma|) \quad (**)} \\
&= O(1+|\operatorname{Im} \sigma|).
\end{aligned}$$

Let us explain the estimates (*) and (**). In both cases, the derivative $\frac{\partial}{\partial r}$ produces 3 or 2 terms, depending whether $t \in (\frac{1}{2}, 2)$ or not. The contributions of $\frac{\partial}{\partial r} e^{i\xi r} = i\xi e^{i\xi r}$ to (*) and of $\frac{\partial}{\partial r} r^{-1-i \operatorname{Im} \sigma} = -(1+i \operatorname{Im} \sigma) r^{-2-i \operatorname{Im} \sigma}$ to (**) are easily dealt with. The contribution of $\frac{\partial}{\partial r} \chi_\infty^\rho(r)$, whose support is contained in the interval $[\rho, 2|\rho|]$, to (*) is obviously bounded, while its contribution to (**) vanishes, as $\frac{1}{|\xi|} > 2|\rho|$.

Consider finally the contributions of $\frac{\partial}{\partial r} \chi_\infty^\rho(tr) = t(\chi_\infty^\rho)'(tr)$ when t is small. Regarding (*), it is easily estimated by $t \int_{|\rho|t^{-1}}^{2|\rho|t^{-1}} dr = |\rho|$. Regarding (**), notice that it vanishes unless $|\xi| \geq \frac{t}{2|\rho|}$. In this case, it is estimated by

$$\frac{t}{|\xi|} \int_{|\rho|t^{-1}}^{2|\rho|t^{-1}} dr r^{-1} \leq 2|\rho| \ln 2.$$

This concludes the estimate of (3.21).

We deduce from (3.20) and (3.21) that

$$\tilde{I}_\infty^+(t, x) = O(t + C_\sigma).$$

By combining this estimate with (3.17) and (3.18), we get next

$$|I_\infty^+| \lesssim C_\sigma |x|^{-\frac{d-1}{2}} \lesssim C_\sigma t^{-\frac{d-1}{2}},$$

if t is small and

$$|I_\infty^+| \lesssim C_\sigma (t|x|)^{-\frac{d-1}{2}} + |x|^{-\frac{d+1}{2}} \lesssim C_\sigma t^{-N} (1+|x|)^{N-\frac{d-3}{2}},$$

if t is large, for any $N \in \mathbb{N}$. Together with the estimate of I_∞^- , this allows us to conclude the proof of [Theorem 3.3](#) in the case $\frac{|x|}{t} \geq \frac{1}{2}$. \square

Remark 3.5. This proof relies on the expression (2.6) of spherical functions, which is available if G is complex but not in general.

Proof for $\frac{|x|}{t} \leq \frac{1}{2}$. Substituting the expressions (2.3) of φ_λ and (2.8) of $|\mathbf{c}(\lambda)|^{-2}$ implies

$$\begin{aligned} \tilde{\omega}_t^{\sigma, \infty}(x) = \text{const.} \frac{e^{\sigma^2}}{\Gamma((d+1)/2 - \sigma)} \int_K dk e^{\langle \rho, A(xk) \rangle} \\ \underbrace{\int_{\mathfrak{a}} d\lambda \chi_\infty^\rho(\lambda) \pi^2(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\psi(\lambda)}}_{=\tilde{II}_\infty(t, x)}. \end{aligned}$$

Since $\frac{|A(xk)|}{t} \leq \frac{|x|}{t} \leq \frac{1}{2}$, we know from (3.4) that the critical point λ_0 of ψ satisfies $|\lambda_0| < \frac{|\rho|}{\sqrt{3}}$. Hence ψ has no critical point in the support of χ_∞^ρ , and $\tilde{II}_\infty(t, x)$ makes sense after several integrations by parts based on (3.12).

Consider first the large time case where $t \geq 1$. After performing N integrations by parts based on (3.12), $\tilde{II}_\infty(t, x)$ becomes

$$\begin{aligned} \tilde{II}_\infty(t, x) = \text{const.} (it)^{-N} \\ \int_{\mathfrak{a}} d\lambda e^{it\psi(\lambda)} \left\{ - \sum_{j=1}^{\ell} \frac{\partial}{\partial \lambda_j} \circ \frac{\lambda_j}{\tilde{\psi}(\lambda)} \right\}^N \left\{ \chi_\infty^\rho(\lambda) \pi^2(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} \right\}. \end{aligned}$$

If some derivatives hit $\chi_\infty^\rho(\lambda)$, then the range of integration reduces to a spherical shell where $|\lambda| \asymp |\rho|$ and we obtain the bound $O(t^{-N})$. Otherwise, since (in the support of χ_∞^ρ) $\lambda_j (\tilde{\psi}(\lambda))^{-1}$, $\pi^2(\lambda)$ and $(|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}}$ are inhomogeneous symbols of order 0, $2|\Sigma^+$ and $-\text{Re } \sigma = -\frac{d+1}{2}$ respectively, we have

$$|\tilde{II}_\infty(t, x)| \lesssim t^{-N} \int_{\mathfrak{a}} d\lambda \chi_\infty(\lambda) |\lambda|^{2|\Sigma^+| - \text{Re } \sigma - N} \lesssim t^{-N}$$

provided that $N > \text{Re } \sigma = \frac{d+1}{2}$. Hence, for all $t \geq 1$ such that $\frac{|x|}{t} \leq \frac{1}{2}$,

$$|\tilde{\omega}_t^{\sigma, \infty}(x)| \lesssim t^{-N} \int_K dk e^{\langle \rho, A(xk) \rangle} = t^{-N} \varphi_0(x),$$

for all $\sigma \in \mathbb{C}$ with $\text{Re } \sigma = \frac{d+1}{2}$, and for every integer $N > \frac{d+1}{2}$.

Notice that the same argument implies similar estimates in small time with $t^{-\frac{d+1}{2}}$, where the negative power is too large. In order to reduce it to $t^{-\frac{d-1}{2}}$, we

split up $\widetilde{II}_\infty(t, x)$ into two parts

$$\begin{aligned}\widetilde{II}_\infty(t, x) &= \widetilde{II}_\infty^-(t, x) + \widetilde{II}_\infty^+(t, x) \\ &= \int_{\mathfrak{a}} d\lambda \chi_0^\rho(t\lambda) \chi_\infty^\rho(\lambda) \pi^2(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\psi(\lambda)} \\ &\quad + \int_{\mathfrak{a}} d\lambda \chi_\infty^\rho(t\lambda) \chi_\infty^\rho(\lambda) \pi^2(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} e^{it\psi(\lambda)},\end{aligned}$$

where $\widetilde{II}_\infty^-(t, x)$ is easily estimated:

$$(3.23) \quad |\widetilde{II}_\infty^-(t, x)| \lesssim \int_{|\rho| \leq |\lambda| \leq 2|\rho|t^{-1}} d\lambda |\lambda|^{2|\Sigma^+| - \operatorname{Re} \sigma} \lesssim t^{-2|\Sigma^+| + \operatorname{Re} \sigma - \ell} = t^{-\frac{d-1}{2}},$$

for all $0 < t < 1$ and for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$. After M integrations by parts based on (3.12), $\widetilde{II}_\infty^+(t, x)$ becomes

$$\begin{aligned}\widetilde{II}_\infty^+(t, x) &= \text{const.}(it)^{-M} \\ &\int_{\mathfrak{a}} d\lambda e^{it\psi(\lambda)} \left\{ -\sum_{j=1}^{\ell} \frac{\partial}{\partial \lambda_j} \circ \frac{\lambda_j}{\widetilde{\psi}(\lambda)} \right\}^M \left\{ \chi_\infty^\rho(t\lambda) \chi_\infty^\rho(\lambda) \pi^2(\lambda) (|\lambda|^2 + \tilde{\rho}^2)^{-\frac{\sigma}{2}} \right\}.\end{aligned}$$

Hence

$$\begin{aligned}|\widetilde{II}_\infty^+(t, x)| &\lesssim t^{-M} \int_{|\lambda| \geq |\rho|t^{-1}} d\lambda |\lambda|^{2|\Sigma^+| - \operatorname{Re} \sigma - M} \\ &\lesssim t^{-M} \int_{r \geq |\rho|t^{-1}} \frac{dr}{r} r^{2|\Sigma^+| - \operatorname{Re} \sigma - M + \ell}.\end{aligned}$$

Therefore, for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$, we have

$$(3.24) \quad |\widetilde{II}_\infty^+(t, x)| \lesssim t^{-M} t^{-d + \operatorname{Re} \sigma + M} = t^{-\frac{d-1}{2}}$$

provided that $M > \frac{d-1}{2}$. From (3.23) and (3.24), we deduce that

$$|\widetilde{\omega}_t^{\sigma, \infty}(x)| \lesssim t^{-\frac{d-1}{2}} \int_K dk e^{\langle \rho, A(xk) \rangle} = t^{-\frac{d-1}{2}} \varphi_0(x),$$

for all $0 < t < 1$ such that $\frac{|x|}{t} \leq \frac{1}{2}$ and for all $\sigma \in \mathbb{C}$ with $\operatorname{Re} \sigma = \frac{d+1}{2}$. Thus we have proved the last case, and the proof of [Theorem 3.3](#) is complete. \square

Remark 3.6. In the last proof, we have used the integral expression (2.3) of the spherical function $\varphi_\lambda(x)$, which holds in general, and the particular expression (2.8) of the Plancherel density $|\mathbf{c}(\lambda)|^{-2}$ when G is complex. In general, $|\mathbf{c}(\lambda)|^{-2}$ is not a polynomial, nor even an inhomogeneous symbol of order $d - \ell$, which is known to be a major difficulty in higher rank analysis.

4. DISPERSIVE ESTIMATES

In this section, we prove our main result about the $L^q \rightarrow L^q$ estimates for the operator $W_t^\sigma = \widetilde{D}^{-\sigma} e^{it\sqrt{-\Delta}}$. We introduce the following criterion based on the Kunze-Stein phenomenon, which is a straightforward generalization of [\[APV11, Theorem 4.2\]](#) and which is crucial for our dispersive estimates.

Lemma 4.1. *Let κ be a reasonable K -bi-invariant function on G . Then*

$$\|\cdot * \kappa\|_{L^{q'}(X) \rightarrow L^q(X)} \leq \left\{ \int_G dx \varphi_0(x) |\kappa(x)|^{\frac{q}{2}} \right\}^{\frac{2}{q}}$$

for any $q \in [2, +\infty]$.

Proof. For $s \in [2, +\infty)$, we define \mathcal{A}_s as the space of all K -bi-invariant functions on G such that

$$\int_G dx \varphi_0(x) |\kappa(x)|^{s/2} < \infty.$$

Given κ in \mathcal{A}_s , we set

$$\|\kappa\|_{\mathcal{A}_s} = \left(\int_G dx \varphi_0(x) |\kappa(x)|^{s/2} \right)^{2/s}.$$

For $s = +\infty$, we denote by \mathcal{A}_∞ the space of $L^\infty(G, dx)$ K -bi-invariant functions on G and by $\|\cdot\|_{\mathcal{A}_\infty}$ the L^∞ -norm. We show in the following that, for any $q \in [2, +\infty]$, we have

$$L^{q'}(G) * \mathcal{A}_q \subset L^q(G),$$

which yields our lemma.

The case where $q = 2$ follows by Herz's criterion (see [Cow97]):

$$\|\cdot * \kappa\|_{L^2(G) \rightarrow L^2(G)} \leq \int_G dx \varphi_0(x) |\kappa(x)| = \|\kappa\|_{\mathcal{A}_2}$$

for every κ in \mathcal{A}_2 . When $q = +\infty$, taking f in $L^1(G)$ and κ in \mathcal{A}_∞ , we have that for every x in G ,

$$|f * \kappa(x)| \leq \int_G dy |f(y)| |\kappa(y^{-1}x)| \leq \|\kappa\|_{\mathcal{A}_\infty} \|f\|_{L^1(G)}.$$

By interpolation between the cases $q = 2$ and $q = \infty$, we obtain that

$$[L^2(G), L^1(G)]_\theta * [\mathcal{A}_2, \mathcal{A}_\infty]_\theta \subset [L^2(G), L^\infty(G)]_\theta$$

with $\theta = 2/q$. On the one hand,

$$[L^2(G), L^1(G)]_\theta = L^{q'}(G) \quad \text{and} \quad [L^2(G), L^\infty(G)]_\theta = L^q(G).$$

On the other hand, since

$$[L^1(G, \varphi_0 dx), L^\infty(G, \varphi_0 dx)]_\theta = L^{q/2}(G, \varphi_0 dx),$$

we have $[\mathcal{A}_2, \mathcal{A}_\infty]_\theta = \mathcal{A}_q$. This concludes the proof of Lemma 4.1. \square

4.1. Small time dispersive estimate.

Theorem 4.2. *Assume that $0 < |t| < 1$, $2 < q < +\infty$ and $\sigma = (d+1)(\frac{1}{2} - \frac{1}{q})$. Then*

$$\|\widetilde{D}^{-\sigma} e^{it\sqrt{-\Delta}}\|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim |t|^{(d-1)(\frac{1}{2} - \frac{1}{q})}.$$

Proof. We divide the proof into two parts, corresponding to the kernel decomposition $\omega_t^\sigma = \omega_t^{\sigma,0} + \omega_t^{\sigma,\infty}$. According to Lemma 4.1, we have

$$\|\cdot * \omega_t^{\sigma,0}\|_{L^{q'}(X) \rightarrow L^q(X)} \leq \left\{ \int_G dx \varphi_0(x) |\omega_t^{\sigma,0}(x)|^{\frac{q}{2}} \right\}^{\frac{2}{q}}.$$

By using the Cartan decomposition, together with the fact that $\delta(H) \lesssim e^{2\langle \rho, H \rangle}$, we obtain

$$\int_G dx \varphi_0(x) |\omega_t^{\sigma,0}(x)|^{\frac{q}{2}} \lesssim \int_{\mathfrak{a}^+} dH \varphi_0(H) |\omega_t^{\sigma,0}(H)|^{\frac{q}{2}} e^{2\langle \rho, H \rangle}.$$

According to [Theorem 3.1](#) and to the estimate (2.4) of the ground spherical function φ_0 , we obtain next

$$\int_{\mathfrak{a}^+} dH \varphi_0(H) |\omega_t^{\sigma,0}(H)|^{\frac{q}{2}} e^{2\langle \rho, H \rangle} \lesssim \int_{\mathfrak{a}^+} dH (1 + |H|)^{\frac{q}{2}+1} e^{-(\frac{q}{2}-1)\langle \rho, H \rangle} < +\infty$$

for any $q \in (2, +\infty)$. Hence

$$\| \cdot * \omega_t^{\sigma,0} \|_{L^{q'}(X) \rightarrow L^q(X)} < +\infty.$$

For the second part $\omega_t^{\sigma,\infty}$, we use an analytic interpolation between $L^2 \rightarrow L^2$ and $L^1 \rightarrow L^\infty$ estimates for the family of operators $\widetilde{W}_t^{\sigma,\infty}$ defined by (3.2) in the vertical strip $0 \leq \operatorname{Re} \sigma \leq \frac{d+1}{2}$. When $\operatorname{Re} \sigma = 0$, the spectral theorem yields

$$\| \widetilde{W}_t^{\sigma,\infty} \|_{L^2(X) \rightarrow L^2(X)} \lesssim \| e^{it\sqrt{-\Delta}} \|_{L^2(X) \rightarrow L^2(X)} = 1.$$

for all $t \in \mathbb{R}^*$. When $\operatorname{Re} \sigma = \frac{d+1}{2}$, [Theorem 3.3](#) gives

$$\| \widetilde{W}_t^{\sigma,\infty} \|_{L^1(X) \rightarrow L^\infty(X)} \lesssim \| \widetilde{\omega}_t^{\sigma,\infty} \|_{L^\infty(X)} \lesssim t^{-\frac{d-1}{2}}.$$

By applying Stein's interpolation theorem for an analytic family of operators, we obtain

$$\| \widetilde{W}_t^{\frac{d+1}{2}(1-\theta),\infty} \|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim t^{-\frac{d-1}{2}(1-\theta)},$$

with $\theta = \frac{2}{q}$. In conclusion,

$$\| W_t^\sigma \|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim |t|^{-(d-1)(\frac{1}{2}-\frac{1}{q})},$$

for $0 < |t| < 1$, $2 < q < +\infty$ and $\sigma = (d+1)(\frac{1}{2} - \frac{1}{q})$. \square

4.2. Large time dispersive estimate.

Theorem 4.3. *Assume that $|t| \geq 1$, $2 < q < +\infty$ and $\sigma = (d+1)(\frac{1}{2} - \frac{1}{q})$. Then*

$$\| \widetilde{D}^{-\sigma} e^{it\sqrt{-\Delta}} \|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim |t|^{-\frac{d}{2}}.$$

Proof. We divide the proof into three parts, corresponding to the kernel decomposition

$$\omega_t^\sigma = \mathbf{1}_{B(0, \frac{|t|}{2})} \omega_t^{\sigma,0} + \mathbf{1}_{X \setminus B(0, \frac{|t|}{2})} \omega_t^{\sigma,0} + \omega_t^{\sigma,\infty}.$$

By using [Lemma 4.1](#) and [Theorem 3.1](#), we have

$$\begin{aligned} & \| \cdot * \{ \mathbf{1}_{B(0, \frac{|t|}{2})} \omega_t^{\sigma,0} \} \|_{L^{q'}(X) \rightarrow L^q(X)} \\ & \lesssim \left\{ \int_{\{H \in \mathfrak{a}^+ : 0 < |H| < \frac{|t|}{2}\}} dH \varphi_0(H) |\omega_t^{\sigma,0}(H)|^{\frac{q}{2}} e^{2\langle \rho, H \rangle} \right\}^{\frac{2}{q}} \\ & \lesssim t^{-\frac{d}{2}} \underbrace{\left\{ \int_{|H| < \frac{|t|}{2}} dH (1 + |H|)^{\frac{q}{2}(d+1)} e^{-(\frac{q}{2}-1)\langle \rho, H \rangle} \right\}^{\frac{2}{q}}}_{< +\infty} \end{aligned}$$

and

$$\begin{aligned} & \|\cdot * \{\mathbf{1}_{X \setminus B(0, \frac{|t|}{2})} \omega_t^{\sigma, 0}\}\|_{L^{q'}(X) \rightarrow L^q(X)} \\ & \lesssim \left\{ \int_{|H| \geq \frac{|t|}{2}} dH (1 + |H|)^{\frac{d}{2}(d+1)} e^{-(\frac{d}{2}-1)\langle \rho, H \rangle} \right\}^{\frac{2}{q}}, \end{aligned}$$

which is $O(|t|^{-N})$, for any $N \in \mathbb{N}$. Instead of $\omega_t^{\sigma, \infty}$, we consider again the kernel $\tilde{\omega}_t^{\sigma, \infty}$. The associated operators satisfy

$$\|\tilde{W}_t^{\sigma, \infty}\|_{L^1(X) \rightarrow L^\infty(X)} \lesssim t^{-N} \quad \forall N \in \mathbb{N},$$

when $\operatorname{Re} \sigma = \frac{d+1}{2}$, according to [Theorem 3.3](#). By applying Stein's interpolation theorem for an analytic family of operators and by summing up these estimates, we obtain

$$\|W_t^\sigma\|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim |t|^{-\frac{d}{2}},$$

for $|t| \geq 1$, $2 < q < +\infty$ and $\sigma = (d+1)(\frac{1}{2} - \frac{1}{q})$. \square

Remark 4.4. The standard TT^* method used to prove the Strichartz inequality breaks down in the critical case. The dyadic decomposition method carried out in [\[KT98\]](#) takes care of the endpoints, but it requires a stronger dispersive property than [Theorem 4.2](#) in small time, namely our main theorem.

Proof of Theorem 1.3. It follows from [Theorem 4.2](#), [Theorem 4.3](#) and the L^q kernel estimate

$$\|\omega_t^\sigma\|_{L^q(X)} \leq \|\omega_t^{\sigma, 0}\|_{L^q(X)} + \|\tilde{\omega}_t^{\sigma, \infty}\|_{L^q(X)} \lesssim \begin{cases} |t|^{-\frac{d-1}{2}}, & \text{if } 0 < |t| < 1, \\ |t|^{-\frac{d}{2}}, & \text{if } |t| \geq 1, \end{cases}$$

for $2 < q < +\infty$ and $\sigma = \frac{d+1}{2}(\frac{1}{2} - \frac{1}{q})$. Indeed, by using standard interpolation arguments between

$$\begin{cases} \|W_t^\sigma\|_{L^1(X) \rightarrow L^q(X)} \lesssim \|\omega_t^\sigma\|_{L^q(X)} \lesssim t^{-\frac{d-1}{2}}, \\ \|W_t^\sigma\|_{L^{q'}(X) \rightarrow L^\infty(X)} \lesssim \|\omega_t^\sigma\|_{L^q(X)} \lesssim t^{-\frac{d-1}{2}}, \\ \|W_t^\sigma\|_{L^2(X) \rightarrow L^2(X)} = 1, \end{cases}$$

for small time and

$$\begin{cases} \|W_t^\sigma\|_{L^1(X) \rightarrow L^q(X)} \lesssim \|\omega_t^\sigma\|_{L^q(X)} \lesssim t^{-\frac{d}{2}}, \\ \|W_t^\sigma\|_{L^{q'}(X) \rightarrow L^\infty(X)} \lesssim \|\omega_t^\sigma\|_{L^q(X)} \lesssim t^{-\frac{d}{2}}, \\ \|W_t^\sigma\|_{L^{q'}(X) \rightarrow L^q(X)} \lesssim |t|^{-\frac{d}{2}}, \end{cases}$$

for large time, we conclude. \square

5. STRICHARTZ INEQUALITY AND APPLICATIONS

Let $\sigma \in \mathbb{R}$ and $1 < q < \infty$. Recall that the Sobolev space $H^{\sigma, q}(X)$ is the image of $L^q(X)$ under the operator $(-\Delta)^{-\frac{\sigma}{2}}$, equipped with the norm

$$\|f\|_{H^{\sigma, q}(X)} = \|(-\Delta)^{\frac{\sigma}{2}} f\|_{L^q(X)}.$$

If $\sigma = N$ is a nonnegative integer, then $H^{\sigma, q}(X)$ coincides with the classical Sobolev space

$$W^{N, q}(X) = \{f \in L^q(X) \mid \nabla^j f \in L^q(X), \forall 1 \leq j \leq N\},$$

defined by means of covariant derivatives. The following Sobolev embedding theorem is used in next subsection.

Theorem 5.1. *Let $1 < q_1, q_2 < \infty$ and $\sigma_1, \sigma_2 \in \mathbb{R}$ such that $\sigma_1 - \sigma_2 \geq \frac{d}{q_1} - \frac{d}{q_2} \geq 0$. Then*

$$(5.1) \quad H^{\sigma_1, q_1}(X) \subset H^{\sigma_2, q_2}(X).$$

We refer to [Tri92] for more details about function spaces on Riemannian manifolds. Let us state next the Strichartz inequality and some applications. The proofs are adapted straightforwardly from [AP14, APV12] and are therefore omitted.

5.1. Strichartz inequality. Recall the linear inhomogenous Klein-Gordon equation on X :

$$(5.2) \quad \begin{cases} \partial_t^2 u(t, x) - \Delta u(t, x) = F(t, x), \\ u(0, x) = f(x), \quad \partial_t|_{t=0} u(t, x) = g(x). \end{cases}$$

whose solution is given by Duhamel's formula:

$$u(t, x) = (\cos t\sqrt{-\Delta})f(x) + \frac{\sin t\sqrt{-\Delta}}{\sqrt{-\Delta}}g(x) + \int_0^t \frac{\sin(t-s)\sqrt{-\Delta}}{\sqrt{-\Delta}}F(s, x)ds.$$

We consider first the case $d \geq 4$ and discuss the 3-dimensional case in the final remark. Recall that a couple (p, q) is called admissible if $(\frac{1}{p}, \frac{1}{q})$ belongs to the triangle

$$\left\{ \left(\frac{1}{p}, \frac{1}{q} \right) \in \left(0, \frac{1}{2} \right] \times \left(0, \frac{1}{2} \right) \mid \frac{1}{p} \geq \frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{q} \right) \right\} \cup \left\{ \left(0, \frac{1}{2} \right) \right\}.$$




FIGURE 1. Admissibility in dimension $d \geq 4$.

Theorem 5.2. *Let (p, q) and (\tilde{p}, \tilde{q}) be two admissible couples, and let*

$$\sigma \geq \frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{q} \right) \quad \text{and} \quad \tilde{\sigma} \geq \frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{\tilde{q}} \right).$$

Then all solutions u to the Cauchy problem (5.2) satisfy the following Strichartz inequality:

$$(5.3) \quad \|\nabla_{\mathbb{R} \times X} u\|_{L^p(I; H^{-\sigma, q}(X))} \lesssim \|f\|_{H^1(X)} + \|g\|_{L^2(X)} + \|F\|_{L^{\tilde{p}'}(I; H^{\tilde{\sigma}, \tilde{q}'}(X))}.$$

The admissible range in (5.3) can be widened by using the Sobolev embedding theorem.


Corollary 5.3. *Assume that (p, q) and (\tilde{p}, \tilde{q}) are two couples corresponding to the square*

$$\left[0, \frac{1}{2}\right] \times \left(0, \frac{1}{2}\right) \cup \left\{\left(0, \frac{1}{2}\right)\right\},$$

see Fig.(A) in the following. Let $\sigma, \tilde{\sigma} \in \mathbb{R}$ such that $\sigma \geq \sigma(p, q)$, where

$$\sigma(p, q) = \frac{d+1}{2} \left(\frac{1}{2} - \frac{1}{q}\right) + \max \left\{0, \frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{q}\right) - \frac{1}{p}\right\},$$

and similarly $\tilde{\sigma} \geq \sigma(\tilde{p}, \tilde{q})$. Then the Strichartz inequality (5.3) holds for all solutions to the Cauchy problem (5.2).



(A) Case $d \geq 4$.

(B) Case $d = 3$.

Remark 5.4. *Theorem 5.2 and Corollary 5.3 still hold true in lower dimension $d = 3$ with similar proofs. But the endpoint $(\frac{1}{2}, \frac{1}{2} - \frac{1}{d-1})$ is excluded from the admissible triangle in this case, see Fig.(B).*

5.2. Global well-posedness in $L^p(\mathbb{R}, L^q(X))$. We refer to [AP14, APV12] for more detailed proofs of the following well-posedness results. By using the classical fixed point scheme with the previous Strichartz inequality, one obtains the global well-posedness for the semilinear equation

$$(5.4) \quad \begin{cases} \partial_t^2 u(t, x) - \Delta u(t, x) = F(u(t, x)), \\ u(0, x) = f(x), \quad \partial_t|_{t=0} u(t, x) = g(x). \end{cases}$$

on X with power-like nonlinearities F satisfying

$$|F(u)| \leq C|u|^\gamma, \quad |F(u) - F(v)| \leq C(|u|^{\gamma-1} + |v|^{\gamma-1})|u - v|, \quad \gamma > 1.$$

and small initial datas f and g . For every $d \geq 3$, consider the following powers

$$\begin{aligned} \gamma_1 &= 1 + \frac{3}{d}, \quad \gamma_2 = 1 + \frac{2}{\frac{d-1}{2} + \frac{2}{d-1}}, \quad \gamma_c = 1 + \frac{4}{d-1}, \\ \gamma_3 &= \begin{cases} \frac{\frac{d+6}{2} + \frac{2}{d-1} + \sqrt{4d + \left(\frac{6-d}{2} + \frac{2}{d-1}\right)^2}}{d} & \text{if } d \leq 5, \\ 1 + \frac{2}{\frac{d-1}{2} - \frac{1}{d-1}} & \text{if } d \geq 6, \end{cases} \\ \gamma_4 &= \begin{cases} 1 + \frac{4}{d-2} & \text{if } d \leq 5, \\ \frac{d-1}{2} + \frac{3}{d+1} - \sqrt{\left(\frac{d-3}{2} + \frac{3}{d+1}\right)^2 - 4\frac{d-1}{d+1}} & \text{if } d \geq 6, \end{cases} \end{aligned}$$

and the following curves

$$\begin{aligned} \sigma_1(\gamma) &= \frac{d+1}{4} - \frac{(d+1)(d+5)}{8d} \frac{1}{\gamma - \frac{d+1}{2d}}, \\ \sigma_2(\gamma) &= \frac{d+1}{4} - \frac{1}{\gamma-1}, \quad \sigma_3(\gamma) = \frac{d}{2} - \frac{2}{\gamma-1}. \end{aligned}$$

Denote by 0^+ any small positive constant. In dimension $d \geq 3$, the equation (5.4) is globally well-posed for small initial data in $H^{\sigma,2}(X) \times H^{\sigma-1,2}(X)$ provided that

$$\begin{cases} \sigma = 0^+, & \text{if } 1 < \gamma \leq \gamma_1, \\ \sigma = \sigma_1(\gamma), & \text{if } \gamma_1 < \gamma \leq \gamma_2, \\ \sigma = \sigma_2(\gamma), & \text{if } \gamma_2 < \gamma \leq \gamma_c, \\ \sigma = \sigma_3(\gamma), & \text{if } \gamma_c < \gamma \leq \gamma_4, \end{cases}$$

Observe that one obtains the same global well-posedness results on noncompact symmetric spaces of arbitrary rank with G complex as on real hyperbolic spaces, without further assumptions.

6. FURTHER RESULTS ON LOCALLY SYMMETRIC SPACES

Let Γ be a discrete torsion-free subgroup of G . The locally symmetric space $\Gamma \backslash X$, equipped with the Riemannian structure inherited from X becomes a Riemannian manifold. Consider the Poincaré series

$$P(s; x, y) = \sum_{\gamma \in \Gamma} e^{-sd(x, \gamma y)}, \quad s > 0, \quad x, y \in X,$$

and denote by $\delta(\Gamma) = \inf\{s > 0 \mid P(s; x, y) < +\infty\}$ its critical exponent. In [Zha19], the author has studied the wave equation and has obtained similar Strichartz inequality and global well-posedness results as in Sect. 5, in the case where $\Gamma \backslash X$ is a rank one locally symmetric space such that Γ is convex cocompact and $\delta(\Gamma) < |\rho|$.

Recall that, in the rank one setting, Γ is called convex cocompact if the quotient group $\Gamma \backslash \text{CH}(\Lambda_\Gamma)$ is compact, where $\text{CH}(\Lambda_\Gamma)$ is the convex hull of the limit set Λ_Γ of Γ . While this notion yields many interesting examples in rank one, it is known to yield a rather limited class in higher rank (see [KL06] and [Qui05] for more details).

However, thanks to our wave kernel estimates Theorem 3.1 and Theorem 3.3, we can study along the lines of [Zha19] the wave equation on higher rank noncompact locally symmetric spaces, under slightly different assumptions:

- (1) G is complex,
- (2) $\delta(\Gamma) < |\rho|$,
- (3) there exists $C > 0$ such that for all $x, y \in X$, $P(s; x, y) \leq CP(s; \mathbf{0}, \mathbf{0})$, where $\mathbf{0} = eK$ denotes the origin of X .

The first assumption ensures sharp wave kernel estimates on X , from which we can deduce wave kernel estimates on $\Gamma \backslash X$. Notice that such information is still lacking for G real. The second assumption plays the same role as in rank one. On the one hand, it ensures that the wave kernel on $\Gamma \backslash X$ is well defined. On the other hand, there is a L^2 Kunze-Stein phenomenon under this assumption. In order to get the desired dispersive estimates on $\Gamma \backslash X$, a uniform upper bound of the Poincaré series is required. Notice that we could deduce the last condition (3) from the convex cocompactness of Γ in rank one. We refer to [Zha19] and the references therein for more details about wave type equations on locally symmetric spaces.

REFERENCES

- [AP14] J.-Ph. Anker and V. Pierfelice. Wave and Klein-Gordon equations on hyperbolic spaces. *Anal. PDE*, 7(4):953–995, 2014.
- [APV11] J.-Ph. Anker, V. Pierfelice, and M. Vallarino. Schrödinger equations on Damek-Ricci spaces. *Comm. Partial Differential Equations*, 36(6):976–997, 2011.
- [APV12] J.-Ph. Anker, V. Pierfelice, and M. Vallarino. The wave equation on hyperbolic spaces. *J. Differential Equations*, 252(10):5613–5661, 2012.
- [APV15] J.-Ph. Anker, V. Pierfelice, and M. Vallarino. The wave equation on Damek-Ricci spaces. *Ann. Mat. Pura Appl. (4)*, 194(3):731–758, 2015.
- [CGM02] M. Cowling, S. Giulini, and S. Meda. Oscillatory multipliers related to the wave equation on noncompact symmetric spaces. *J. London Math. Soc. (2)*, 66(3):691–709, 2002.
- [Cow97] M. Cowling. Herz’s “principe de majoration” and the Kunze-Stein phenomenon. In *Harmonic analysis and number theory (Montreal, PQ, 1996)*, volume 21 of *CMS Conf. Proc.*, pages 73–88. Amer. Math. Soc., Providence, RI, 1997.
- [DGK01] P. D’Ancona, V. Georgiev, and H. Kubo. Weighted decay estimates for the wave equation. *J. Differential Equations*, 177(1):146–208, 2001.
- [Fon97] J. Fontaine. A semilinear wave equation on hyperbolic spaces. *Comm. Partial Differential Equations*, 22(3-4):633–659, 1997.
- [GLS97] V. Georgiev, H. Lindblad, and C. Sogge. Weighted Strichartz estimates and global existence for semilinear wave equations. *Amer. J. Math.*, 119(6):1291–1319, 1997.
- [GV95] J. Ginibre and G. Velo. Generalized Strichartz inequalities for the wave equation. *J. Funct. Anal.*, 133(1):50–68, 1995.
- [Has11] A. Hassani. Wave equation on riemannian symmetric spaces. *J. Math. Phys.*, 52(4):043514, 2011.
- [Hel62] S. Helgason. *Differential geometry and symmetric spaces*. Pure and Applied Mathematics, Vol. XII. Academic Press, New York-London, 1962.
- [Hel00] S. Helgason. *Groups and geometric analysis: integral geometry, invariant differential operators, and spherical functions (Corrected reprint of the 1984 original)*, volume 83 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2000.
- [Kap94] L. Kapitanski. Weak and yet weaker solutions of semilinear wave equations. *Comm. Partial Differential Equations*, 19(9-10):1629–1676, 1994.
- [KL06] B. Kleiner and B. Leeb. Rigidity of invariant convex sets in symmetric spaces. *Invent. Math.*, 163(3):657–676, 2006.
- [KT98] M. Keel and T. Tao. Endpoint Strichartz estimates. *Amer. J. Math.*, 120(5):955–980, 1998.
- [LS95] H. Lindblad and Christopher D. Sogge. On existence and scattering with minimal regularity for semilinear wave equations. *J. Funct. Anal.*, 130(2):357–426, 1995.
- [MT11] J. Metcalfe and M. Taylor. Nonlinear waves on 3D hyperbolic space. *Trans. Amer. Math. Soc.*, 363(7):3489–3529, 2011.

- [MT12] J. Metcalfe and M. Taylor. Dispersive wave estimates on 3D hyperbolic space. *Proc. Amer. Math. Soc.*, 140(11):3861–3866, 2012.
- [Qui05] J.-F. Quint. Groupes convexes cocompacts en rang supérieur. *Geom. Dedicata*, 113:1–19, 2005.
- [Ste93] E.M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993.
- [Tat01] D. Tataru. Strichartz estimates in the hyperbolic space and global existence for the semilinear wave equation. *Trans. Amer. Math. Soc.*, 353(2):795–807, 2001.
- [Tri92] H. Triebel. *Theory of function spaces. II*, volume 84 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1992.
- [Zha19] H.-W. Zhang. Wave and Klein–Gordon equations on certain locally symmetric spaces. *To appear in J. Geom. Anal.*, *Doi: 10.1007/s12220-019-00246-8*, 2019.

INSTITUT DENIS POISSON (UMP 7013), UNIVERSITÉ D'ORLÉANS, UNIVERSITÉ DE TOURS
& CNRS, BÂTIMENT DE MATHÉMATIQUES - RUE DE CHARTRES B.P. 6759 - 45067 ORLÉANS
CEDEX 2 - FRANCE

Email address: hong-wei.zhang@univ-orleans.fr