

# Inside the shell: body composition of free-ranging tortoises (Testudo hermanni)

Ljiljana Tomović, Dragan Arsovski, Ana Golubović, Xavier Bonnet

## ▶ To cite this version:

Ljiljana Tomović, Dragan Arsovski, Ana Golubović, Xavier Bonnet. Inside the shell: body composition of free-ranging tortoises (Testudo hermanni). Zoology, 2020, 142, pp.125821. 10.1016/j.zool.2020.125821. hal-02905179

## HAL Id: hal-02905179 https://hal.science/hal-02905179v1

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0944200620300805 Manuscript\_1b818b0ad5d293aa42b7df2ac230ff0e

1	
2	Inside the shell: body composition of free-ranging
3	tortoises (Testudo hermanni)
4	
5	
6	
7	
8	Ljiljana Tomović <sup>1</sup> , Dragan Arsovski <sup>2</sup> , Ana Golubović <sup>1</sup> and Xavier Bonnet <sup>3*</sup>
9	
10	
11	
12	
13	<sup>1</sup> University of Belgrade, Faculty of Biology, Institute of Zoology, Studentski trg 16, 11000
14	Belgrade, Serbia.
15	<sup>2</sup> Macedonian Ecological Society, Arhimedova 5, 1000 Skopje, North Macedonia.
16	<sup>3</sup> Centre d'Etudes Biologiques de Chizé, UMR 7372, Centre National de la Recherche
17	Scientifique Université de la Rochelle, 79360 Villiers-en-Bois, France.
18	
19	
20	* Corresponding author: Xavier Bonnet, e-mail: bonnet@cebc.cnrs.fr
21	
22	All authors declare a lack of conflict of interest.
23	
24	

#### 25 Abstract

26 Body condition indices (BCI - mass scaled by size) are widely used in ecological 27 studies. They presumably reflect variations of endogenous fat reserves in freeranging animals. In the field, however, accurately quantifying internal body reserves 28 29 is a difficult task. This is especially true in armoured animals where convenient clues 30 that may guide BCI assessment (e.g. visible subcutaneous fat deposits) remain 31 inaccessible. Alternatively, inclusive dissections may provide anatomical abacuses to 32 estimate body reserves in living individuals. Sacrificing animals for this purpose is 33 not acceptable. We opportunistically tested the ability of BCI to estimate body 34 reserves in 13 free-ranging Hermann's tortoises (Gmelin, 1789) dissected soon after 35 they died from natural causes. On average, BCI values were lower in dissected 36 tortoises relative to living individuals (N>10,000 measurements), but they remained within the range of variation of the studied populations. Shell mass relative to body 37 mass was high and showed considerable inter-individual variation (33.5% to 52.3%). 38 39 Stomach and digestive tract content represented another important and variable part 40 of total body mass (4.4% to 14.5%). The contribution of fat bodies was negligible 41 (0.0% to 0.5%). Overall, in the studied tortoises, variations of body condition are weakly determined by variations of fat stores. Other endogenous (e.g. muscles, 42 visceral tissues, liver) and "exogenous" (e.g. digestive tract content, clutch) elements 43 44 should be considered to better understand age and sex specific life-history trade-offs faced by chelonians. 45

46

47

49

<sup>48</sup> Key words: Body condition index; fat reserves; dissection; Chelonians

#### 50 **1. Introduction**

51 Direct access to the internal elements that compose an animal, for example 52 reproductive organs, body reserves or digestive tract content, is essential to examine a wide range of questions in ecology, evolutionary ecology and toxicology (Schmidt-53 Nielsen, 1975; Sakai et al., 2000; Zera and Harshman, 2001; Polilov, 2015). But in 54 55 Vertebrates, most studies have focused on a single segment. Typically, diet has been 56 assessed by scrutinizing stomach contents while other organs were overlooked 57 (Evans and Lampo, 1996; Paltridge et al., 1997). Similarly, during inspection of 58 ovarian structures, liver or muscles were generally ignored (Sica et al., 2001; Pizzatto 59 et al., 2008; Gherissi et al., 2018). Whole-body composition based on dissection has 60 been examined in only a handful of studies (Sakai et al., 2000; Landgraf et al., 2006).

61 Thus, for the vast majority of living Vertebrates, we cannot quantify the respective 62 input of each main body components to overall body mass (e.g. relative mass of the 63 skin or locomotor muscles). This limits the possibility to address ecological issues where body condition indices (BCI, mass scaled by size) are used to evaluate lipidic 64 reserves (Stephens et al., 2009). For example, calculation and interpretation of BCI 65 can be misleading since total body mass includes unknown shares of fat stores, 66 67 muscles, partly digested food, and ovulated follicles. In most cases, measuring all 68 main internal body components is not feasible in the field, even when using 69 sophisticated techniques (Speakman, 2001; Cojean et al., 2018; Hofmeyr et al., 2020). Currently, only comprehensive dissections allow isolating and measuring organs 70 71 (mass, dimensions), extracting and weighing whole content of the stomach or of the 72 intestine, to exactly assess body reserves, etc., and thus to eventually obtain a 73 comprehensive picture of the elements that constitute an animal as a whole.

But performing inclusive dissections relies on the availability of dead animals and on the willingness to devote a considerable amount of time to the partitioning and measuring of internal elements (Sakai et al., 2000). Although sacrificing free ranging animals should be avoided, accidentally or naturally dead animals can be collected without raising major ethical concern. Picking and fully dissecting randomly encountered cadavers is not in the culture of most field ecologists.

3

80 The lack of convenient technique to assess an individual's body composition 81 might prove particularly taxing when dealing with animals protected by an external 82 armour (e.g. shell, large scales, spines or very thick skin). First, the mass of external 83 protection may well be substantial (e.g. ~30% of total body mass in pangolins; 84 Wilson and Mittermeier 2011). Second, the mass of all internal components that do not belong to the tissues of the individual sensu stricto, like digestive tract content 85 (e.g. faeces) or developing embryos for example, is inaccessible via palpation 86 (ultrasound and radiography enable the visualization of follicles and eggs, however 87 - Hofmeyr et al., 2020). Third, convenient clues to assess BCI, like visible fat 88 deposits, cannot be used (e.g. camel hump; Gherissi et al., 2018). Overall, the 89 relationship between body condition and actual endogenous body reserves is 90 obscured using mass/size relationships, since exogenous matter and reproductive 91 items also contribute to overall body mass. These complications are almost 92 93 systematically ignored in the ecological literature where body condition is considered as a reliable estimator of fat reserves (Schulte-Hostedde et al., 2005; Peig 94 95 and Green, 2009). Yet, the paucity of information also implies that we cannot gauge 96 the severity and relevance of this issue.

97 In this study we dissected fresh, naturally dead tortoises in the course of capture-98 mark-recaptures studies. This offered the opportunity to examine to what extent the use of a classical BCI may have tarnished our ecological interpretations during 99 100 previous studies (Bonnet et al., 2001; Djordjević et al., 2011; Lecq et al., 2014). Finally, 101 we expected that a better knowledge of tortoise body composition may improve 102 further ecological studies. To the best of our knowledge, this study provides the first 103 precise data on the whole-body composition of terrestrial Chelonians, and provides 104 exemplary points of prudence when interpreting BCI.

105

#### 106 **2. Material and methods**

107 *Study species and study sites* 

108 The Hermann's tortoise (*Testudo hermanni*) can grow up to 25 cm, attain 50 years, and 109 displays delayed maturity and low fecundity (Bertolero et al., 2011). Estimates for body size at maturity range between 13 cm and 15 cm of straight carapace length
(SCL); we classified individuals larger than 13 cm as adults (Willemsen and Hailey,
112 1999).

From 2008 until 2019, we studied the ecology and morphology of tortoises in two dense populations (> 20 adults / ha) of *T. hermanni* in two nearby sites (~4.5 km away from each other) of the Prespa Lake region in North Macedonia: Golem Grad Island (GG, ~18 ha; N 40°52'; E 20°59'), and Konjsko village (KV, ~20 ha, N 40°54'; E 20°99') (Djordjević et al., 2011; Arsovski et al., 2018a; Golubović et al., 2013, 2017, 2018).

119

## 120 Data collection

121 Each individual was sexed when possible (small individuals cannot always be sexed - Arsovski et al., 2018b; Golubović et al., 2018) and marked using a notch-code on 122 the marginal-scutes (Stubbs et al., 1984). Overall, we encountered tortoises more than 123 18,500 times in the field (2,798 captures plus 15,789 recaptures). Straight carapace 124 125 length (SCL) and body mass (BM) were recorded, and individuals were rapidly 126 released at the exact place of capture. To limit unnecessary handling during frequent 127 recaptures, SCL and BM were not systematically recorded. We obtained 9,680 and 1,432 measurements of both SCL and BM from Golem Grad and Konjsko 128 129 populations, respectively.

During field work, we examined all dead (or dying) tortoises. In total, we found 130 315 dead tortoises on Golem Grad (223 males, 41 females and 51 immatures; this 131 proportion of dead males reflects the highly biased sex-ratio in this site; Bonnet et al. 132 2016) and 35 in Konjsko (20 males, 14 females and 1 immature). Most of them were 133 partly rotten or fully skeletonised (i.e. empty shells) and could not be used for 134 135 dissection. We retained only 13 very fresh individuals for dissection (12 from Golem 136 Grad and one from Konjsko; Table 1). Most of them had big cracks on the shell, 137 revealing that they fell from cliffs, or serious injuries on the cloaca suggesting they 138 had died from sexual harassment (Golubović et al., 2013, 2018). Animals in poor 139 condition (e.g. severely wounded) but still alive were kept in outdoor enclosures,

wounds cleaned, and provided with water and food until they recovered or not. One
recently wounded female could not be cured in the field (very deep hole in the back),
thus she was euthanized.

143 Prior to dissection, we recorded SCL and BM. Then we opened the shell ventrally 144 using a hacksaw to remove the plastron. The main organs of the tortoise were separated (using scissors and pliers), carefully examined and weighed using 145 146 electronic scales. The mass of the shell was recorded to the nearest 0.1 g. Most components could be actually separated and weighed with precision of 0.1 g: heart, 147 liver, lungs, stomach, intestine, gonads, growing follicles, eggs or digestive tract 148 149 content for example. Several parts of the body could not be easily dissected. For 150 example, we did not tease apart the skin, muscles and bones for the legs, head, neck 151 and tail; thus, we only recorded the total mass of the abovementioned elements. Fluids and blood lost during dissection, along with parts which were not easily 152 153 identified or assigned, like mesenteric tissues and blood vessels for example, were 154 classified as miscellaneous material for conciseness.

155

#### 156 Analyses

Body condition index (BCI) was calculated using residuals from a linear regression 157 158 (living and dissected individuals pooled, dead but not-dissected tortoises discarded) of In-BM against In-SCL (Bonnet et al., 2001). This allowed us to evaluate to what 159 extent the BCI of freshly dissected tortoises was representative of the values 160 measured in the studied populations. The small proportion of dissected tortoises 161 (0.1%) limited their contribution in the calculation of the parameters of the 162 163 regression. BCI values were normally distributed (Shapiro-Wilk test, p > 0.05). Many 164 living individuals were represented more than once (recaptured over time), therefore, individual identity was included as a random factor. Not-dissected dead 165 166 tortoises (e.g. partly rotten, empty shells) were not included in the body condition 167 analyses, but they were plotted on Figure 1 for comparison. Generalized linear 168 mixed models were built in R Version 1.2.5001 (R Core Team, 2019) using the package nlme (Pinheiro et al., 2019). Competitive models were ranked using theAkaike information criterion (AIC, Burnham and Anderson, 2002).

The masses of certain organs were highly correlated (e.g. left versus right legs). 171 172 Furthermore, for the purpose of the current study, considering morpho-functional 173 components instead of each separate element improved the conciseness of analyses. Therefore, we assigned all dissected elements into 11 simple units (henceforth 174 designated as components, Table 2) that were used in most analyses. The 175 176 relationship between the total body mass and the respective mass of these 177 components was examined with a backward stepwise regression. We first verified 178 that the mass of these components, used as the dependent variables, was correlated 179 (0.57 < r < 0.98; all p < 0.05) with the body mass (the independent variable). Because 180 the sum of the components represented total body mass, this stepwise analysis 181 enabled us to identify potential major contributors of BCI.

Sex difference in body composition was tested using ANCOVA with the main 182 component as the dependent variable, sex as the factor and body mass as the 183 covariable. The strong correlation between the mass of each body component and 184 185 body mass facilitated this analysis. However, due to the small sample size of 186 dissected tortoises, we also used non-parametric Kruskal-Wallis ANOVA to compare the body composition (e.g. shell, somatic and visceral organs mass relative 187 188 to the body mass, ratios were arcsine transformed prior to the analyses) of females (N = 4) relative to males (N = 9). 189

190

#### 191 **3. Results**

## 192 Body condition: effect of sex, age and population

The data obtained enabled us to encompass the range of variations of body mass as a function of body size in both populations (Figure 1). Model selection revealed that interaction of locality (Golem Grad Island or Konjsko village), state (living *vs.* dissected) and sex (male, female or immature) best explained variation of BCI (Table 3). Tortoises from Konjsko exhibited higher body condition in general, regardless of 198 state and sex (t = 9.88, p < 0.001, Figure 1), whereas males exhibited lower body 199 condition than females in both localities (t = -3.33, p < 0.001).

200

#### 201 Body condition of living and dissected individuals

Dissected tortoises exhibited lower body condition compared to living individuals (t = 4.0, p < 0.001), but Figure 1 shows that the body mass relative to body size of the dissected tortoises remained within the range of variation exhibited by living ones, especially in Golem Grad where most dissected tortoises were found (Figure 1A). But considering the Konjsko population, most dissected tortoises could be classified as lean (Figure 1B).

208

## 209 Body composition of dissected tortoises

Body composition was highly variable among dissected individuals (Table 1). For example, focusing on adult males, the relative mass of the shell ranged from 33.5% to 51.8%. This strong variability was also observed in other body components (see Supplementary Material – Table I).

214 The final step of the backward regression (adjusted  $R^2 = 0.999$ ,  $F_{9,3} = 4894.2$ , p < 215 0.0001) retained nine components among eleven, that significantly contributed to 216 total body mass (Table 4). The masses of somatic organs (legs, head, neck and tail) 217 and shell mass were the most important contributors (standardized coefficients > 218 0.30). Miscellaneous material and digestive tract (net) were also substantial 219 contributors (standardized coefficients > 0.10). Finally, the masses of kidneys, 220 digestive tract content (i.e. partly digested material), and fat bodies, were retained in 221 the model, but the standardized coefficients were small (~0.05). The masses of the 222 heart and respiratory system provided almost null standardized coefficients (~-0.08). 223 The masses of the liver and reproductive organs were not retained.

Regarding sexual dimorphism, ANCOVA and Kruskal-Wallis ANOVA provided similar trends (although some non-parametric tests failed to reach statistical significance). We found sex differences for the most important components: females exhibited relatively heavier shells, while males showed higher relative somatic organs mass - (Table 5; Figure 2). A closer inspection of the somatic organs showed
that males had relatively heavier limbs and heads (Table 5). They also had higher
relative values of some visceral organs, e.g. hearts and livers compared to females
(Table 5). We found no other significant sex differences.

232

## 233 4. Discussion

Dissections of free-ranging tortoises provided new insight on the significance of body mass, and by extension of body condition in terrestrial chelonians. Several results support previous studies (e.g. about sexual dimorphism), but others were unexpected (e.g. regarding the relative contribution of digestive tract content *versus* fat bodies).

239

## 240 Dissected versus living tortoises

On average, dissected individuals were in the lower part of the range of variation of BCI exhibited by living tortoises (Figure 1). Dead tortoises may dehydrate rapidly in the field, especially those with large cracks in the shell. Some tortoises may have lost significant amounts of blood. Rapid loss of fluids may reduce body mass and thus body condition. One female harassed by nine males exhibited low body condition and may have died from exhaustion. On the other side of the spectrum, our sample also included one large female with a high BCI.

Overall, our dissection data encompass most of the variation in adult body size, but are not fully illustrative of the whole range of body condition observed in the populations monitored (e.g. heaviest males and small individuals were lacking). However, comparisons with other studies provide means to assess to what extent our dissection results are representative of what is currently known in chelonians.

253

#### 254 Shell mass

The protective role of the armour is crucial in Chelonians (Polly et al., 2016), but it also imposes major morpho-functional and energetic constraints. The shell can obstruct movements in dense vegetation and limits tortoise agility (Bonnet et al., 2001; Golubović et al., 2014, 2015; Chiari et al., 2017). Abundant resources and time are needed to build an effective shell; this shapes the growth trajectory during the first decade of life in the Hermann's tortoise (Arsovski et al., 2018a, b). Likely, crucial structural materials of the shell are not easily mobilised; we thus speculate that they play no (or limited) role as body reserves.

The fresh mass of the shell represents the main component of a tortoise, averaging 263 41.5% of total body mass (39.6% when removing two extreme values in very lean 264 individuals). Dissections of captive tortoises indicate that the mean dry mass of the 265 shell relative to the dry mass of the carcass represents 31%, and 35% in adults (Table 266 267 18 in Kopsch, 2006). Despite differences in methodologies (fresh vs. dry masses, relative to total body vs. carcass mass), previous and current studies show that the 268 269 mass of the shell is considerable in tortoises (suggested by Iverson 1984). This proportion is approximately 30% (27% in adults) in terrapins (Miller and Birchard, 270 271 2005; Kopsch, 2006), and is markedly lower (17%) in adult sea turtles (Sakai et al., 2000). 272

We observed very strong variations among individuals in the relative fresh mass of the shell (34% to 52%). Dry masses measured in different species support our results (e.g. 23% to 80%; Kopsch, 2006). The mass of the shell correlates with body size, yet not perfectly ( $r^2 = 0.81$ , N = 13) due to variations in shell shape. Part of the variation in total body mass (~19%) is caused by interindividual variations in relative shell mass. Thus, BCI includes an unknown albeit substantial shellcomponent independent of body reserves.

280

#### 281 Body reserves

Fat bodies were virtually absent in all dissected individuals, representing only 0.0% to 0.5% of the total body mass. Kopsch (2006) did not provide information regarding fat bodies in the dissections of 92 captive individuals (3 *Testudo*, 2 terrapins, and *Geochelone pardalis*); suggesting that adipose tissues were negligible. Sporadic cases of obesity have been reported in overfed captive tortoises, with pads protruding from the openings of the shell (Burchfield et al., 1980; Lamberski, 2013). But this 288 situation may not occur in natural conditions: we never found an obese individual in 289 our large datasets (> 5,000 individuals) collected in different tortoise species (Bonnet 290 et al., 2001; El Mouden et al., 2006; Djordjević et al., 2011; Lecq et al., 2014). 291 Noticeably, in the current study, dissected individuals exhibited average to low 292 body condition (except one very large female). Thus, we cannot ascertain that fat bodies are always negligible, especially in heaviest individuals. It would be 293 informative to assess if fat bodies develop above a BCI threshold (this was not the 294 case in the high-BCI dissected female, 0.3% fat). Abundant fat deposits, however, 295 296 have been observed in sea turtles (Kwan, 1994), reaching 12% of the total body mass 297 in female loggerhead turtles (calculated from Table 1 in Sakai et al., 2000). Lipidic 298 reserves may well be more developed in aquatic than in terrestrial chelonians.

The liver is an important source of endogenous reserves (e.g. lipids and proteins) mobilized during reproduction (Telford, 1970; Bonnet et al., 1994; Heck et al., 1997; Finkler, 2013). This organ also stores reserves beyond reproduction (Collins and Anderson, 1995). Dissections showed that the liver represents an important proportion of body mass in Chelonians, 2.7% to 7.2% in the current study, 2.6% to 3.5% in female sea turtles (Sakai et al., 2001) and 1.5% to 5.0% in tortoises and terrapins (Table 32 in Kopsch, 2006).

Fat bodies, the most "classical" proxy for BCI, did not contribute significantly to variation of body mass in tortoises. Somatic organs (e.g. locomotor muscles) and the digestive tract were better contributors. Tortoises may first allocate resources in their locomotor and digestive structures, and build up fat bodies only later. Future investigations focusing on the relationships between body condition and dynamic aspects of mass variation of the main components are needed.

312

313 Sexual dimorphism in body plan

In our study, the relative mass of the shell was greater in females (38.7% – 52.3% *vs.* 315 33.5% – 51.8% in males; Supplementary Material, Table I). This supports the notion 316 of a more developed shell in females to maximize protective functions and to 317 enhance abdominal volume to hold the clutch, compared to a relatively lighter shell with larger openings to favor mobility in males (Bonnet et al., 2001; Djordjević et al.,
2011).The relative mass of organs involved in locomotor performance (e. g. limbs,
heart) was greater in males compared to females (21.5% - 30.0% *versus* 15.8% - 24.3%
of the total body mass; Supplementary Material, Table I). These result fit well with
previous studies on sexual dimorphism of body plan in chelonians (Bonnet et al.,
2010), and more generally regarding sexual dimorphism of body composition in
snakes and fishes (Bonnet et al., 1998; Casselman and Schulte-Hostedde, 2004).

325

## 326 Exogenous components inside the body

In all dissected individuals, dissections revealed substantial amounts (4.4% – 10.2% of the total body mass in adults) of exogenous and partly digested materials in the digestive tract. Incidentally, the highest proportion (14.5%) was observed in the immature male; perhaps small and rapidly growing individuals need to feed a lot. Regardless, partly processed food contributed to total body mass more than fat bodies.

The resources invested into large follicles and ovulated eggs cannot be considered as endogenous reserves (Bonnet et al. 1994). One female carried abundant and welldeveloped follicles and eggs (Supplementary Material), but in the others, the ovaries and oviducts were almost empty, only small follicles being visible. This considerable variation (0.1% – 10.0% of the total body mass) explains the significant contribution of the reproductive organs + follicles and eggs in total body mass variations.

339

## 340 Body condition versus body composition

Body condition integrates the mass of endogenous body reserves (fat and protein stores) along with exogenous material. Fluctuations of the relative mass of the shell cannot be factored out. Further, hydration status influences body mass (Packard and Packard, 2001), as suggested by the importance of the fluids lost during dissections. Thus, the meaning of body condition should be considered with caution and certainly not reduced to lipidic stores. Yet, BCI is a useful index to assess health status in tortoises; for example, it strongly correlates with the mass of crucial 348 locomotor structures. Additionally, in females it may well be a good indicator of 349 reproductive status. The mass of the reproductive items of the dissected female with 350 well-developed eggs and follicles was 5 to 100 times greater than in non-351 reproductive females (Supplementary Material – Table I).

352

353 *Consequences for ecological studies* 

354 The two populations we studied differ considerably in terms of body condition; this contrast requires explanation(s). Only one individual from Konjsko was dissected, 355 356 precluding direct comparisons of body composition between sites however. In 357 Golem Grad fat bodies were negligible, but this might not be true for the heaviest 358 Konjsko tortoises. Yet, using BCI as a mere indicator of fat stores may lead to invalid 359 conclusions. For example, on Golem Grad, tortoises with a high BCI might be 360 incorrectly considered as fat compared to those with a low BCI - a classical shortcut in ecological studies. Instead, our results suggest that somatic organs were more 361 362 important. This information may help to address specific issues. For instance, we may hypothesize that Golem Grad tortoises are locally adapted to extremely uneven 363 topography where lightness and agility are advantageous (Golubović et al., 2017). 364 365 Alternatively, does low food availability explain why Golem Grad tortoises are lean 366 compared to Konjsko? The ecological and evolutionary processes that underpin 367 these questions differ considerably. Well-informed BCI values would be then crucial. In conclusion, field dissections provided support to previous studies on sexual 368 body shape dimorphism. More importantly, they enabled us to broaden the meaning 369 of body condition in free-ranging tortoises, notably to shift away from a fat store 370 371 focus toward a more complex vision that includes various body components. This 372 might be useful for future studies of sexual dimorphism, feeding ecology, 373 reproductive investment, demography, and other life-history traits.

374

#### 375 Acknowledgements

Our research was funded by Ministries of Education, Sciences and TechnologicalDevelopment (project No. 173043). Permissions were issued by the Ministry of

379 Park "Galičica" (11-2817/2, 11-4093/5, and 03-246). Many thanks to our colleagues
380 Rastko Ajtić, Bogoljub Sterijovski, Sonja Nikolić and Ana Ivanović, as well as to
381 students Uroš Ilić and Stefan Avramović who considerably helped us in dissections.
382
383 References
384 Arsovski, D., Olivier, A., Bonnet, X., Drilholle, S., Tomović, L., Béchet, A., Golubović,
385 A., Besnard, A. 2018. Covariates streamline age-specific early life survival

Environment and Spatial Planning of Macedonia and the authorities of the National

estimates of two chelonian species. J. Zool. 306, 223-234.

387 Arsovski, D., Tomović, L., Golubović, A., Nikolić, S., Sterijovski, B., Ajtić, R.,

Ballouard, J.M., Bonnet, X. 2018. When carapace governs size: variation among

389 age classes and individuals in a free-ranging ectotherm with delayed maturity.

390 Oecologia 186, 953-963.

378

391 Bertolero, A., Cheylan, M., Hailey, A., Livoreil, B., Willemsen, R.E. 2011. Testudo

392 *hermanni* Gmelin 1789 – Hermann's tortoise. Conservation biology of freshwater

393 turtles and tortoises: a compilation project of the IUCN/SSC Tortoise and

394 Freshwater Turtle Specialist Group. Chelonian Res. Monogr. 5, 059-051.

Bonnet, X., Naulleau, G., Mauget, R. 1994. The influence of body condition on 17-β

396 estradiol levels in relation to vitellogenesis in female *Vipera aspis* Reptilia,

397 Viperidae. Gen. Comp. Endocr. 93, 424-437.

Bonnet, X., Shine, R., Naulleau, G., Vacher-Vallas, M. 1998. Sexual dimorphism in
snakes: different reproductive roles favour different body plans. Proc. Royal. Soc.
B 265, 179-183.

401 Bonnet, X., Lagarde, F., Henen, B.T., Corbin, J., Nagy, K.A., Naulleau, G., Balhoul, K.,

402 Chastel, O. Legrand, A., Cambag, R. 2001. Sexual dimorphism in steppe tortoises

403 *Testudo horsfieldii*: influence of the environment and sexual selection on body

404 shape and mobility. Biol. J. Linn. Soc. 72, 357-372.

405 Bonnet, X., Delmas, V., El-Mouden, H., Slimani, T., Sterijovski, B., Kuchling, G. 2010.

406 Is sexual body shape dimorphism consistent in aquatic and terrestrial chelonians?

407 Zoology 113, 213-220.

408	Bonnet, X., Golubović, A., Arsovski, D., Đorđević, S., Ballouard, JM., Sterijovski, B.,
409	Ajtić, R., Barbraud, C. Tomović, L. 2016. A prison effect in a wild population: a
410	scarcity of females induces homosexual behaviors in males. Behav. Ecol. 27, 1206-
411	1215.
412	Burchfield, P.M., Doucette, C.S., Beimler, T.F. 1980. Captive management of the
413	Radiated Tortoise Geochelone radiata at Gladys Porter Zoo, Brownsville. Int. Zoo
414	Yearb. 20, 1-6.
415	Burnham, K.P., Anderson, D.R. 2002. Model Selection and Multimodel Inference: A
416	Practical Information-Theoretic Approach. Springer-Verlag, Berlin.
417	Casselman, S.J., Schulte-Hostedde, A.I. 2004. Reproductive roles predict sexual
418	dimorphism in internal and external morphology of lake whitefish, Coregonus
419	<i>clupeaformis</i> . Ecol. Freshw. Fish 13, 217-222.
420	Chiari, Y., Van Der Meijden, A., Caccone, A., Claude, J., Gilles, B. 2017. Self-righting
421	potential and the evolution of shell shape in Galápagos tortoises. Sci. Rep. 7,
422	15828.
423	Cojean, O., Vergneau-Grosset, C., Masseau, I. 2018. Ultrasonographic anatomy of
424	reproductive female leopard geckos Eublepharis macularius. Vet. Radiol.
425	Ultrasound 59, 333-344.
426	Collins, A.L., Anderson, T.A. 1995. The regulation of endogeneous energy stores
427	during starvation and refeeding in the somatic tissues of the golden perch. J. Fish
428	Biol. 47, 1004-1015.
429	Djordjević, S., Djurakić, M., Golubović, A., Ajtić, R., Tomović, L., Bonnet, X. 2011.
430	Sexual body size and body shape dimorphism of Testudo hermanni in central and
431	eastern Serbia. Amphibia-Reptilia 32, 445-458.
432	El Mouden, E.H., Slimani, T., Kaddour, K.B., Lagarde, F., Ouhammou, A., Bonnet, X.
433	2006. Testudo graeca graeca feeding ecology in an arid and overgrazed zone in
434	Morocco. J. Arid Environ. 64, 422-435.
435	Evans, M., Lampo, M. 1996. Diet of <i>Bufo marinus</i> in Venezuela. J. Herpetol. 30, 73-76.
	15

- 436 Finkler, M.S. 2013. Sexual dimorphism in visceral organ mass and hematology in
- 437 spotted salamanders, *Ambystoma maculatum*, before and after breeding. Copeia,
  438 2013, 338-345.
- 439 Gherissi, D.E., Afri-Bouzebda, F., Bouzebda, Z., Bonnet, X. 2018. Are female camels
- capital breeders? Influence of seasons, age, and body condition on reproduction inan extremely arid region. Mamm. Biol. 93, 124-134.
- Golubović, A., Arsovski, D., Ajtić, R., Tomović, L., Bonnet, X. 2013. Moving in the
  real world: tortoises take the plunge to cross steep steps. Biol. J. Linn. Soc. 108,
  719-726.
- 445 Golubović, A., Andjelković, M., Arsovski, D., Vujović, A., Iković, V., Djordjević, S.,
- 446 Tomović, L. 2014. Skills or strength how tortoises cope with dense vegetation?
  447 Acta Ethol. 17, 141-147.
- Golubović, A., Tomović, L., Ivanović, A. 2015. Geometry of self righting: the case of
  Hermann's tortoises. Zool. Anz. 254, 99-105.
- 450 Golubović, A., Andjelković, M., Arsovski, D., Bonnet, X., Tomović, L. 2017.
- 451 Locomotor performances reflect habitat constraints in an armoured species.
- 452 Behav. Ecol. Sociobiol. 71, 93.
- 453 Golubović, A., Arsovski, D., Tomović, L., Bonnet, X. 2018. Is sexual brutality
- 454 maladaptive under high population density? Biol. J. Linn. Soc. 124, 394-402.
- 455 Heck, J., MacKenzie, D.S., Rostal, D., Medler, K., Owens, D. 1997. Estrogen induction
- 456 of plasma vitellogenin in the Kemp's ridley sea turtle *Lepidochelys kempi*. Gen.
- 457 Comp. Endocrinol. 107, 280-288.
- 458 Hofmeyr, M.D., Henen, B.T., Loehr, V.J.T. 2020. Reproductive investments of a small,
- 459 arid zone tortoise *Chersobius signatus*: Follicle and egg development. Acta Zool.
  460 101, 39-50.
- 461 Iverson, J.B. 1984. Proportional skeletal mass in turtles. Fla. Scientist 47, 1-11.
- 462 Kopsch, G. 2006. Untersuchungen zur Körperzusammensetzung von Schildkröten.
- 463 PhD Thesis, Faculty of Veterinary Medicine, Imu München.
- 464 Kwan, D. 1994. Fat reserves and reproduction in the green turtle, *Chelonia mydas*.
- 465 Wildl. Res. 21, 257-265.

- 466 Lamberski, N. 2013. Body condition scores for desert tortoises. Retrieved from
- 467 www.fws.gov/nevada/desert\_tortoise/documents/reports/2013/assess/Desert468 Tortoise-BCS-2013-lamberskipo.pdf
- 469 Landgraf, S., Susenbeth, A., Knap, P., Looft, H., Plastow, G., Kalm, E., Roehe, R.

2006. Developments of carcass cuts, organs, body tissues and chemical body
composition during growth of pigs. Anim. Sci. 82, 889-899.

- 472 Lecq, S., Ballouard, J.-M., Caron, S., Livoreil, B., Seynaeve, V., Matthieu, L.-A.,
- Bonnet, X. 2014. Body condition and habitat use by Hermann's tortoises in burnt
  and intact habitats. Conserv. Physiol. 2, cou019.
- 475 Miller, K., Birchard, G.F. 2005. Influence of body size on shell mass in the ornate box
  476 turtle, *Terrapene ornata*. J. Herpetol. 39, 158-162.
- 477 Packard, G., Packard, M. 2001. Environmentally induced variation in size, energy
- 478 reserves and hydration of hatchling painted turtles, *Chrysemys picta*. Funct. Ecol.
  479 15, 481-489.
- Paltridge, R., Gibson, D., Edwards, G. 1997. Diet of the feral cat *Felis catus* in central
  Australia. Wildl. Res. 24, 67-76.
- Peig, J., Green, A.J. 2009. New perspectives for estimating body condition from
  mass/length data: the scaled mass index as an alternative method. Oikos 118,
  1883-1891.
- 485 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Team, R.C. 2019. Nlme: Linear and
- 486 Nonlinear Mixed Effects Models. R package version 3.1-131. Retrieved from
  487 https://CRAN.R-project.org/package=nlme
- 488 Pizzatto, L., Jordão, R.S., Marques, O.A. 2008. Overview of reproductive strategies in
- 489 Xenodontini Serpentes: Colubridae: Xenodontinae with new data for *Xenodon*490 *neuwiedii* and *Waglerophis merremii*. J. Herpetol. 42, 153-163.
- 491 Polilov, A.A. 2015. Small is beautiful: features of the smallest insects and limits to
  492 miniaturization. Annu. Rev. Entomol. 60, 103-121.
- 493 Polly, P.D., Stayton, C.T., Dumont, E.R., Pierce, S.E., Rayfield, E.J., Angielczyk, K.D.
- 494 2016. Combining geometric morphometrics and finite element analysis with
- 495 evolutionary modeling: towards a synthesis. J. Vertebr. Paleontol. 36, e1111225.

- 496 R Core Team. 2019. R: A Language and Environment for Statistical Computing.
- 497 Vienna, Austria: R Foundation for Statistical Computing.
- 498 https://www.gbif.org/tool/81287/r-a-language-and-environment-for-statistical499 computing.
- 500 Sakai, H., Saeki, K., Ichihashi, H., Suganuma, H., Tanabe, S., Tatsukawa, R. 2000.
- 501 Species-specific distribution of heavy metals in tissues and organs of loggerhead
- 502 turtle *Caretta caretta* and green turtle *Chelonia mydas* from Japanese coastal waters.
- 503 Mar. Pollut. Bull. 40, 701-709.
- Schmidt-Nielsen, K. 1997. Animal Physiology: Adaptation and Environment.
  Cambridge University Press.
- 506 Schulte-Hostedde, A.I., Zinner, B., Millar, J.S., Hickling, G.J. 2005. Restitution of
- 507 mass-size residuals: validating body condition indices. Ecology 86, 155-163.
- 508 Sica, S., Fierro, D., Iodice, C., Muoio, R., Filosa, S., Motta, C. 2001. Control of oocyte
- recruitment: regulative role of follicle cells through the release of a diffusiblefactor. Mol. Reprod. Dev. 58, 444-450.
- 511 Speakman, J.R. 2001. Body Composition Analysis of Animals: A Handbook of Non512 destructive Methods. Cambridge University Press, Cambridge.
- 513 Stephens, P.A., Boyd, I.L., McNamara, J.M., Houston, A.I. 2009. Capital breeding and
- 514 income breeding: their meaning, measurement, and worth. Ecology 90, 2057-2067.
- 515 Stubbs, D., Hailey, A., Pulford, E., Tyler, W. 1984. Population ecology of European
- 516 tortoises: review of field techniques. Amphibia-Reptilia 5, 57-68.
- 517 Telford, Jr, S.R. 1970. Seasonal fluctuations in liver and fat body weights of the
- 518 Japanese lacertid *Takydromus tachydromoides* Schlegel. Copeia, 1970 681-688.
- Willemsen, R.E., Hailey, A. 1999. Variation of adult body size of the tortoise *Testudo hermanni* in Greece: proximate and ultimate causes. J. Zool. 248, 379-396.
- 521 Wilson, D.E., Mittermeier, R.A. eds. 2011. Handbook of the Mammals of the World.
- 522 Vol. 2. Hoofed Mammals. Lynx Edicions, Barcelona.
- 523 Zera, A.J., Harshman, L.G. 2001. The physiology of life history trade-offs in animals.
- 524 Annu. Rev. of Ecol. Syst. 32, 95-126.
- 525

**Table 1.** Summary data of dissected tortoises from Golem Grad Island (GG) and Konjsko village (KV). M (I) – immature male, M – adult male, F – adult female. SCL stands for straight carapace length (mm), BM1 and BM2 respectively for body mass measured when the animal was found (dead or alive, see text) or just before dissection. BCI means Body Condition Index (mass scaled by size, residuals). The most important components of total body mass are provided (shell, soma and viscera). Soma includes front and hind legs, head, neck and tail; Viscera includes liver, heart, thymus, spleen, respiratory organs, kidneys, reproductive organs, digestive system (stomach + intestine + food + faeces), fat bodies, miscellaneous material (blood. etc.).

Sex, ID and	Biometry				Mass at dissection (g)				
location	SCL	BM1	BCI	BM2	Shell	Soma	Viscera		
M (I) 994 (GG)	95.9	211.0	0.08945	203.0	81.0 (39.9)	56.7 (27.9)	65.3 (32.2)		
M 843 (GG)	182.2	1199.0	0.00065	1016.5	397.7 (39.1)	387.0 (38.1)	231.8 (22.8)		
M 234 (GG)	188.0	1113.0	-0.16300	1062.0	393.9 (37.1)	418.3 (39.4)	249.8 (23.5)		
M 722 (GG)	148.0	426.0	-0.44304	423.0	219.0 (51.8)	133.3 (31.5)	70.7 (16.7)		
M 303 (GG)	139.2	556.0	-0.00222	555.0	224.0 (40.4)	179.2 (32.3)	151.8 (27.3)		
M 1296 (GG)	187.0	1151.0	-0.11485	1078.0	473.0 (43.9)	360.5 (33.4)	244.5 (22.7)		
M 1206 (GG)	174.0	968.0	-0.08291	964.0	323.0 (33.5)	381.3 (39.6)	259.7 (26.9)		
M 924 (GG)	186.0	1286.0	0.01132	1220.0	450.0 (36.9)	453.6 (37.2)	316.4 (25.9)		
M 338 (GG)	178.0	1076.0	-0.04183	1063.6	395.0 (37.1)	361.6 (34.0)	306.4 (28.9)		
F 1776 (GG)	152.0	672.0	-0.06313	651.0	278.0 (42.7)	187.7 (28.8)	185.3 (28.5)		
F 1058 (GG)	178.0	911.0	-0.20830	914.0	424.0 (46.4)	276.6 (30.3)	213.4 (23.3)		
F 1712 (GG)	163.0	766.0	-0.13108	771.0	403.0 (52.3)	161.5 (20.9)	206.5 (26.8)		
F NEW (KV)	225.0	2348.0	0.07152	2348.0	908.0 (38.7)	629.0 (26.8)	811.0 (34.5)		

**Table 2.** List of the main components used to describe the body composition of the dissected tortoises. They were implemented in the stepwise backward regression analysis. Element pooled explain which organs or elements were pooled in certain components.

Component (g)	Elements pooled				
Shell	-				
Somatic organs	See Supplementary Material – Table I				
Liver	-				
Heart	-				
<b>Respiratory system</b>	-				
Kidneys	-				
Reproductive organs	Both gonads, epididymis, oviducts plus follicles and				
	ovulated eggs				
Digestive tract	Gullet, stomach and intestine (net)				
Fat bodies	-				
Digestive tract content	Stomach and intestine content				
Miscellaneous material	Blood, lymph, several mesenteric elements and blood				
	vessels				

**Table 3.** Model ranking of generalized linear mixed models of Hermann's tortoise body condition variation conditional to univariate, additive (+) and interactive (\*) effects of sex and state (immature, male or female) and locality (Golem Grad Island and Konjsko village). One model built to evaluate whether dissected tortoises' body condition differed from that of living tortoises (dissected *vs.* alive) had a low AIC (likely due to small sample size of dissected tortoises). Models were ranked according to their respective Akaike Information Criteria (AIC) values,  $\Delta$ AIC and AIC weights.

Madal			AIC
Model	AIC	ΔΑΙ	weight
Dissected <i>vs.</i> alive	-28.734,8	479.6	0.0
Sex and state	-28.905,9	308.5	0.0
Locality	-29.204,2	10.2	0.0
Sex + Locality	-29.214,2	0.2	0.5
Sex * Locality	-29.214,4	0.0	0.5
Sex * Locality	-29.214,4	0.0	0.5

**Table 4.** Final selection from the stepwise backward regression analysis (following 7 steps). Nine components among 11 (see Table 2) were retained. Standardized coefficient and coefficient are expressed  $\pm$  SE; t-test t with degree of freedom in brackets.

Component	Standardized	Coofficient	+ (6)	р	
Component	coefficient	Coefficient	ι (0)		
Somatic organs	$0.448 \pm 0.027$	$1.458 \pm 0.087$	16.781	< 0.001	
Shell	$0.314 \pm 0.020$	$0.836 \pm 0.052$	15.969	< 0.001	
Miscellaneous material	$0.208 \pm 0.014$	1.975±0.129	15.223	< 0.001	
Digestive tract	$0.124 \pm 0.014$	2.609±0.282	9.280	< 0.003	
Kidneys	$0.060 \pm 0.014$	9.362±2.196	4.265	0.025	
Digestive tract content	$0.058 \pm 0.013$	1.014±0.219	4.631	0.019	
Fat bodies	$0.047 \pm 0.013$	11.218±3.233	3.470	0.040	
Heart	-0.083±0.018	-15.591±3.360	-4.641	0.018	
Respiratory system	-0.076±0.018	-5.499±1.315	-4.183	0.025	

**Table 5.** Results from GLM (ANCOVA, columns 2 to 4) and Kruskal-Wallis ANOVA (columns 6 and 7) to compare the relative mass of different components or organs between the sexes (4 females and 9 males). SD (column 5) indicates the direction of sexual dimorphism. Components where only non-significant results were found are not reported.

Component	SC	F	Р	SD	Н	Р
Shell	9089.3	7.466	0.021	F	2.381	0.122
Somatic organs	21476.3	12.192	0.006	Μ	6.095	0.013
Legs	9732.0	12.472	0.005	Μ		
Head	424.8	8.385	0.016	Μ		
Heart	29.3	12.510	0.005	Μ		
Liver	1508.9	10.633	0.009	Μ	5.357	0.020

## **Figure captions:**

**Figure 1.** Relationship between body mass (g) and body size (straight carapace length, mm) of live tortoises from Golem Grad Island (Top panel A) and Konjsko village (Lower panel B). Dissected (black circles), and not dissected but dead (grey crosses) tortoises are displayed on both panels for population comparison (scales identical). One large dissected female was in very good body condition and one male was particularly lean (grey arrows). This large female was found in Konjsko, all the other dissected tortoises were found in Golem Grad.

**Figure 2.** Comparison of the body composition of male and female Hermann's tortoises. Mean values (± SD) provide the proportions (% relative to body mass) of the main body components: shell, somatic organs, visceral organs, fat bodies and faeces (see table 2 and text for details and statistics).

Figure 1.



Figure 2.

