



HAL
open science

Treatment-induced shrinking of tumour aggregates: A nonlinear volume-filling chemotactic approach

Luis Almeida, Gissell Estrada-Rodriguez, Lisa Oliver, Diane Peurichard,
Alexandre Poulain, François M. Vallette

► **To cite this version:**

Luis Almeida, Gissell Estrada-Rodriguez, Lisa Oliver, Diane Peurichard, Alexandre Poulain, et al..
Treatment-induced shrinking of tumour aggregates: A nonlinear volume-filling chemotactic approach.
Journal of Mathematical Biology, 2021, 10.1007/s00285-021-01642-x . hal-02906240v2

HAL Id: hal-02906240

<https://hal.science/hal-02906240v2>

Submitted on 27 Jan 2021 (v2), last revised 17 Sep 2021 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Treatment-induced shrinking of tumour aggregates: A nonlinear volume-filling chemotactic approach

Luis Almeida* Gissell Estrada-Rodriguez †‡ Lisa Oliver§
Diane Peurichard¶ Alexandre Poulain || Francois Vallette**

Abstract

Motivated by experimental observations in 3D/organoid cultures derived from glioblastoma, we develop a mathematical model where tumour aggregate formation is obtained as the result of nutrient-limited cell proliferation coupled with chemotaxis-based cell movement. The introduction of a chemotherapeutic treatment induces mechanical changes at the cell level, with cells undergoing a transition from rigid bodies to semi-elastic entities. We analyse the influence of these individual mechanical changes on the properties of the aggregates obtained at the population level by introducing a nonlinear volume-filling chemotactic system of partial differential equations. The elastic properties of the cells are taken into account through the so-called *squeezing probability*, which allows us to change the packing capacity of the aggregates, depending on the concentration of the treatment in the extracellular microenvironment. We explore two scenarios for the effect of the treatment: firstly, the treatment acts only on the mechanical properties of the cells and, secondly, we assume it also prevents cell proliferation. A linear stability analysis enables us to study the ability of the system to create patterns. We provide numerical simulations in 1D and 2D that illustrate the shrinking of the aggregates due to the presence of the treatment.

1 Introduction

Cell migration in the extracellular microenvironment (ECM) and the organisation of cells in response to chemical and mechanical cues are successfully studied using continuum descriptions based on differential equations [1, 42]. In a continuous setting, the chemotactic behaviour of cells (*i.e.* the ability to move along a chemical gradient) is often modelled using a Keller-Segel system of equations [27]. This model was originally proposed for pattern formation in bacterial populations but turned out to be pertinent to describe a wide variety of self-organisation behaviours [11, 6, 59, 28, 56].

*Sorbonne Université, Inria, CNRS, Laboratoire Jacques-Louis Lions, F-75005 Paris. email: luis.almeida@sorbonne-universite.fr

†Sorbonne Université, Laboratoire Jacques-Louis Lions, F-75005 Paris. email: estradarodriguez@ljl.math.upmc.fr

‡Corresponding author

§Centre de Recherche en Cancérologie et Immunologie Nantes-Angers, UMR 1232, Université de Nantes. email: Lisa.Oliver@univ-nantes.fr

¶Sorbonne Université, Inria, CNRS, Laboratoire Jacques-Louis Lions, F-75005 Paris. email: diane.a.peurichard@inria.fr

||Sorbonne Université, Laboratoire Jacques-Louis Lions, F-75005 Paris. email: poulain@ljl.math.upmc.fr

**Centre de Recherche en Cancérologie et Immunologie Nantes-Angers, UMR 1232, Université de Nantes. email: Francois.Vallette@univ-nantes.fr

Different variations of the Keller-Segel model have been adopted in order to better understand the way cells aggregate [8, 4, 40, 46, 16].

Cancer cells have been shown to respond to chemical and mechanical signals from components of the tumour micro-environment (TME) and *vice versa*, cells in the TME acquire a more tumourigenic phenotype, which would sustain tumour growth. The interaction of tumour cells with the TME has been the subject of recent biological surveys [22, 18]. Many *in vitro* (and *ex vivo*) experiments have shown that cells that are cultured on ECM often have a tendency to form aggregate patterns that are dependent on the particular cell lines and physical properties of the media [20]. The exact consequences of the dynamic interplay between heterogeneous cellular entities and their response to alterations in the TME have not yet been elucidated. In particular, little is known about the role of mechanics in the spatial organization of the tumour spheroids. Biological evidence presented in [33, 29, 12] suggests that the formation of aggregates in glioma cells can be explained through a chemotaxis process, rather than, *e.g.*, cell-cell adhesion. In this paper, we follow the chemotactic approach to explain the formation of glioma aggregates. For the case of breast cancer cells, a recent report [9] has proposed a chemotaxis-based explanation for spheroid formation based on theoretical analysis and numerical simulations of the Turing instabilities of such systems.

Inspired by experimental observations in 3D/organoid cultures derived from freshly operated Glioblastoma (GBM), which reproduce *in vivo* behaviours as described in [37] (see Section 2 for more details), in this paper we explore a simple setting where GBM aggregate formation is the result of nutrient-limited cell proliferation coupled with a chemotaxis-based cell movement. A chemotherapeutic treatment introduced after the formation of the aggregates induces mechanical changes at the cell level. We study the influence of these individual mechanical changes on the characteristics of the aggregates obtained at the population level.

GBM are solid tumours characterised by intra- and inter-tumoural heterogeneity and resistance to conventional treatments that result in a poor prognosis [31]. They are the most common and aggressive primary brain tumour in adults. Standard treatments include surgical resection (when possible), combined with radiotherapy and chemotherapy using the DNA alkylating agent Temozolomide (TMZ) [48]. In fact, the overall survival of treated patients is about 15 months versus 3 months without treatment, with fewer than 5% of patients surviving longer than 5 years [47].

One reason behind this relative therapeutic failure is the poor response of GBM tumours to this chemotherapeutic treatment due to their plasticity. Several studies have looked at the genetic compounds of TMZ-resistant cells focusing on the genes responsible for DNA mismatch repair protein [50], while other studies focused on spatial and temporal variations in signalling pathways, which lead to functional and phenotypic changes in GBM [38]. The communication between the tumour cells and the TME as well as the properties of the ECM have a large impact on tumour evolution and invasion, as shown in recent studies [15, 7, 58]. From a biological and medical perspective, it is difficult to investigate the connections between clinically observable glioma behaviour and the underlying molecular and cellular processes. The challenge is to integrate the theoretical and empirical acquired knowledge to better understand the mechanisms and factors that contribute to GBM resistance to treatment. In this context, mathematical models provide useful tools towards identifying dependencies between different phenomena and how they are affected by the different therapeutic strategies.

Much effort has been dedicated to the modelling of GBM formation and invasion of the surrounding tissue, as well as to improving diagnosis and treatment. The exhaustive review [2] discusses different modelling approaches as well as some of the main mechanisms that are observed in GBM formation and invasion.

In this work, we explore two scenarios: the case where the treatment only acts on the mechanical properties of the cells, and the case where it also prevents cell proliferation [30]. We adopt a macro-

scopic approach where cells are represented by their macroscopic density and are supposed to move in the environment via chemotaxis, *i.e.* towards zones of high concentrations of a chemoattractant that is produced by the tumour cells. Moreover, cell proliferation is assumed to depend on the local concentration of nutrients available. Finally, we suppose that when the treatment - represented by its continuous concentration - is introduced, it diffuses in the environment and is naturally consumed by the cells.

Under these hypotheses, which are motivated by the experimental results discussed in Section 2, we obtain a nonlinear volume-filling Keller-Segel model for the cell density, coupled with reaction diffusion equations for the chemoattractant and treatment concentrations. Moreover, we provide a linear stability analysis that enables us to study the ability of the system to generate patterns, and we provide numerical simulations in 1D and 2D.

The paper is organised as follows. In Section 2 we describe the *in vitro* experiments and the main experimental observations that motivated our model. Section 3 is devoted to the description of the model for the first part of the experiments (without the treatment, Section 3.1) and the second part, when the treatment is introduced (Section 3.2). In Section 4, we present the stability analysis for each of these two parts, and Section 5 is devoted to numerical simulations. Section 5.1 presents the numerical scheme, Section 5.3 presents the results in 1D including a discussion on the comparison between numerical experiments and theoretical predictions of the stability analysis. Section 5.4 shows the 2D simulations. Finally, we discuss the results and give some perspectives in Section 6.

2 Description of the experiments

To address the question of the response of GBM cells to TMZ treatment, we took advantage of recently developed 3D biosphere experiments, using GBM patient-derived cultures in a simple 3D scaffold composed of alginate and gelatin [39]. GBMG5 cells were cultured at a concentration of 4×10^6 cells/ml for 14 days until the formation of cell aggregates could be observed, corresponding to the first part of the experiments (**P1**). After that, the cultures were treated with 100 μ M TMZ for two hours once every week (second part of the experiments, **P2**). Over the 30 days, the proliferation was determined counting 3 representative samples, and the cell number was determined as follows. The biospheres were dissociated by incubation for 3 min in 100 mM Na-Citrate and the cell number and cell viability were determined using the Countess optics and image automated cell counter (Life Technologies). In addition the aggregates were photographed to analyse their morphology, and the diameter of the cell aggregates were measured from pictographs using FIJI. To determine the diameters of these cell structures, more than 200 cell aggregates were measured.

We show in Figure 1 (I) the mean length (in μ m) of the cell aggregates computed from the microscopic images without TMZ treatment (round markers), and with TMZ weekly administered (squared markers). Figures 1 A and B show typical microscopic images of the spheroids at day 24, without and with weekly TMZ treatment respectively. In Figure 1 (II), we show the evolution of the cell number as a function of time, where we do not observe big changes in cell number with and without TMZ, once the carrying capacity of the system is reached.

Using clinically relevant concentrations [44] the total number of cells in the biospheres does not seem to be significantly impacted by the TMZ treatment. However, the mean size of the GBMG5 cell aggregates decreases when TMZ is introduced weekly as compared to control cultures (Figure 1 (I)), suggesting that in the presence of the treatment, GBMG5 cells tend to self-organize into smaller and more compact cell clusters.

Another observation supporting a tendency for GBM cells to form more compact structures with higher intracellular adhesion under this type of treatment is the increased expression of claudin, a




Figure 1: Biological experiments of GBMG5 cells cultured in 3D scaffolds with or without 100µM TMZ. (I) Evolution of the mean cell aggregates diameter determined from the microscopic images as function of time, without treatment (circle markers) and with a weekly TMZ (square markers). (A) Typical microscopic image on day 24 of control cell aggregates without treatment, (B) microscopic image at day 24 with 100µM of TMZ administrated weekly for 2 hours. (II) Evolution of the total cell number in the biospheres as function of time, without treatment (circle markers) and with weekly TMZ (square markers).




Figure 2: Claudin expression marking the tight junctions between the cells.

marker of tight junctions formed between cells (Figure 2). In Figure 2 we have GBMG5 cells that were cultured for 14 days until the formation of cell aggregates could be observed, and the cultures were treated with different doses of TMZ until day 23. Furthermore, cell aggregates were fixed then labeled with an anti-claudin antibody. The degree of staining was determined in a double-blind experiment. As one can observe from the images of Figure 2 (left), the cell aggregates seem to be smaller and more compact when TMZ is present compared to the control group (top left figure), and these cultures are associated with higher levels of claudin (see Figure 2 (right)).

Inspired by these observations and results presented in [33, 29, 12], we propose a general model for the chemotaxis-driven formation of cell clusters and shrinking of the aggregates via the action of a non-cytotoxic drug whose effect would be characterised by a change in the mechanical properties of cells that would just modify their packing properties. Moreover, we also consider a second scenario where the treatment would also affect cell proliferation and compare the results obtained.

3 Mathematical model

Motivated by the experiments described in Section 2, we assume that glioma cells have a chemotactic behavior, *i.e.* they move in response to some signaling chemical (chemoattractant), which is secreted by themselves and diffuses in the environment. The chemotactic movement of cell populations plays a fundamental role for example in gastrulation [17], during embryonic development; it directs the movement of immune cells to sites of infection and it is crucial to understand tumour cell invasion [14] and cancer development [59]. Motivated by these applications, chemotaxis and related phenomena have received significant attention in the theoretical community, see the reviews [23, 24].

We suppose that in normal conditions (first part of the experiments described in Section 2, where there is no treatment), tumour cells proliferate and move via chemotaxis as described before. We suppose that cell proliferation is limited by the nutrients available in the environment, *i.e.* cell proliferation is only active as long as the local density does not exceed a given threshold corresponding to the carrying capacity of the environment. Moreover, in order to take into account the finite size of cells and volume limitations, cell motion is only allowed in locations where the local cell density is much smaller than another threshold value corresponding to the tight packing state. In normal conditions, cells are supposed to behave as rigid bodies. In stressed conditions however, (second part of the experiments described in Section 2) we suppose that cells respond to the chemotherapeutic stress (induced by the presence of the treatment) by changing their mechanical properties and behaving as a semi-elastic material.

These hypotheses are modelled via a system of partial differential equations (PDE) which corresponds to a volume-filling chemotaxis equation [43] for the first part (to describe the self-organization of cells into aggregates), and a “semi-elastic” volume-filling chemotaxis approach [55] for the second part, when the treatment is introduced.

For convenience we denote the density of the population of cancer cells in the first part of the experiments by $u(\mathbf{x}, t)$ and that of the second part by $w(\mathbf{x}, t)$. Here $\mathbf{x} \in \Omega \subset \mathbb{R}^2$ where Ω is a bounded domain. The time $t \in [0, T]$ where $T = T_1 + T_2$ represents the total time corresponding to the first and second parts. The main difference between these two populations is the change in the elastic properties of the cells due to the presence of the treatment. If the concentration of the treatment is zero, $u(\mathbf{x}, t) = w(\mathbf{x}, t)$. Cells follow a biased random walk according to the distribution of the chemoattractant of concentration $c(\mathbf{x}, t)$ that is secreted by the cells. We start by detailing the different components of the mathematical models corresponding to the two parts of the experiments described in Section 2.

Logistic growth model for cell proliferation As previously described, in order to take into account the nutrient-limited population growth, cell proliferation is modelled by a logistic growth process. At the population level, we consider a source term $f(u)$ in the PDE for the evolution of the cell density which depends nonlinearly on the local cell density u and reads:

$$f(u) = ru \left(1 - \frac{u}{u_{\max}} \right). \quad (1)$$

Here, $r > 0$ is the proliferation rate and u_{\max} corresponds to the maximum density of the population, also referred to as the carrying capacity of the environment. Alternative cell kinetics could be considered. For example, we can assume that the proliferation is also mediated by the chemoattractant concentration such that $f(u, c) = ruc(1 - u/u_{\max})$ [43]. In this paper we focus on the case given by (1).

Chemoattractant dynamics As described before, we suppose that cell aggregates spontaneously emerge as the result of a self-organization phenomenon of chemotaxis type. To this aim, we suppose that the cells themselves produce the signaling chemical (chemoattractant) that drives their motion. The chemical secreted is therefore supposed to be continuously produced by the cells at rate $\alpha > 0$ and diffuses in the surrounding environment with diffusion coefficient $d_2 > 0$. It is further assumed that the chemical has a finite lifetime and degrades at rate $\beta > 0$. The evolution of the chemical concentration $c(\mathbf{x}, t)$ is therefore given by the following reaction-diffusion equation

$$\partial_t c = d_2 \Delta c + \alpha u - \beta c, \quad (2)$$

where u is the cell density.

Treatment dynamics We suppose that the treatment is introduced at the beginning of **P2**. This treatment is supposed to diffuse in the environment with diffusion coefficient d_4 , and to be consumed by the cells at rate δ . This is modelled by a reaction-diffusion equation for the drug concentration $M(\mathbf{x}, t)$:

$$\partial_t M = d_4 \Delta M - \delta w,$$

where $w(\mathbf{x}, t)$ represents the cell density corresponding to the second part of the experiments. We consider different initial conditions for the drug: either uniformly distributed in the simulation domain, or introduced in the center as a very steep Gaussian function (see Section 5).

3.1 Volume-filling approach for chemotaxis: first part P1

The classical Keller-Segel system of equations [27] describes how cells move along the gradient to local maxima of the chemoattractant. At the same time, this chemical, which is produced by the cells, promotes aggregation leading to the so-called “overcrowding scenarios”, and eventually, the cell density may blow up in finite time (see the comprehensive reviews [23, 49] and references therein).

In this paper, in order to take into account volume limitations and the finite cell size, we consider a modified version of the Keller-Segel model called the *volume-filling* approach for cell motion. This approach was introduced in [43], where the authors provided a detailed derivation in one dimension as well as a comprehensive numerical study of the model.




Figure 3: Squeezing probability for different values of γ_M for $\bar{u} = 1$.

Moreover, we introduce the so-called *squeezing probability*, which describes the probability that a cell finds an empty space at a neighbouring location before moving (see [55]). It takes the form

$$q(u(\mathbf{x}, t)) = \begin{cases} 1 - \left(\frac{u}{\bar{u}}\right)^{\gamma_M}, & \text{for } 0 \leq u \leq \bar{u}, \\ 0, & \text{otherwise,} \end{cases} \quad (3)$$

where $u(\mathbf{x}, t)$ is the cell density, $\gamma_M \geq 1$ is the *squeezing parameter*, $M \equiv M(\mathbf{x}, t) \geq 0$ denotes the concentration of the treatment, and \bar{u} is the *crowding capacity* which corresponds to the tight packing state of the cells. The function $q(u)$ satisfies the following properties,

$$q(\bar{u}) = 0, \quad 0 < q(u) \leq 1, \quad \text{and } q'(u) \leq 0 \text{ for } 0 \leq u < \bar{u}. \quad (4)$$

Moreover, $|q'(u)|$ is bounded and $q''(u) \leq 0$.

The exponent γ_M is chosen to be

$$\gamma_M = (\bar{\gamma} - 1)M + 1, \quad (5)$$

where $\bar{\gamma}$ is a positive constant, $M = 0$ when there is no drug in the environment (part **P1** of the experiments), and $M \equiv M(\mathbf{x}, t)$ when the drug is introduced (part **P2**, described below). Such choice of γ_M enables to take into account different forms of squeezing probability, corresponding to different mechanical behaviors of the cells. In Figure 3, we plot the squeezing probabilities as a function of the cell density, corresponding to different values of γ_M . We see that when there is no treatment present in the environment ($M = 0$, $\gamma_M = 1$, blue curve of Figure 3), the squeezing probability decreases linearly with the local cell density, corresponding to cells modelled as solid particles.

However for larger values of γ_M (when the treatment is present, $M > 0$ and $\gamma_M > 1$, red and yellow curves in Figure 3), the squeezing probability becomes a nonlinear function of the cell density, modelling the fact that in the presence of a drug, cells change their mechanical state to behave as semi-elastic entities that can squeeze into empty spaces. We refer to [54] for more details on the link between the cells elastic properties and the squeezing probability.

For convenience of notation, we distinguish the case without treatment ($M = 0$) for **P1** and define

$$q_1(u(\mathbf{x}, t)) = 1 - \frac{u}{\bar{u}}. \quad (6)$$

The Keller-Segel model built with this specific squeezing probability (6) has been widely studied in the literature, from modelling [43, 54], analytic [57, 21, 32, 16] and numerical [26] perspectives.

Complete PDE system for the first part P1 Taking into account all the previous ingredients, the complete PDE system for part **P1** (when no treatment is present in the environment) reads:

$$\begin{aligned}\partial_t u &= \nabla \cdot (d_1 D_1(u) \nabla u - \chi_u \phi_1(u) \nabla c) + f(u) , \\ \partial_t c &= d_2 \Delta c + \alpha u - \beta c ,\end{aligned}\tag{7}$$

where the first equation describes the evolution of the cell density u and the second is the reaction-diffusion equation for the chemoattractant, previously introduced. The equation for u describes the volume-filling chemotactic motion associated with the squeezing probability q_1 . This equation has been obtained as the hydrodynamic limit of the continuous space-time biased random walk model that corresponds to the squeezing probability q_1 . In the hydrodynamic limit, the cell density evolves according to a nonlinear transport diffusion equation with source term, which corresponds to a volume filling Keller-Segel model with logistic growth. The density-dependent diffusion coefficient $D_1(u)$ and the chemotactic sensitivity $\phi_1(u)$ relate to q_1 via

$$D_1(u) = q_1 - q_1' u , \quad \phi_1(u) = q_1 u .$$

For this part **P1**, where q_1 is given by (6), these coefficients take the values $D_1 = 1$ and $\phi_1(u) = u(1 - u/\bar{u})$. In equation (7), $d_1, \chi_u, \alpha, \beta$ are all positive parameters, and $f(u)$ is the logistic growth source term previously mentioned and defined by (1).

The PDE system is supplemented with the following zero-flux boundary conditions

$$(d_1 D_1(u) \nabla u - \chi_u \phi_1(u) \nabla c) \cdot \eta = 0 , \quad d_2 \nabla c \cdot \eta = 0 ,\tag{8}$$

where η is the outer unit normal at $\partial\Omega$. The initial conditions are given by

$$u(\mathbf{x}, 0) = u_0 , \quad c(\mathbf{x}, 0) = c_0 .\tag{9}$$

3.2 PDE system including the treatment: Part P2

We now describe the dynamics of the cell population when the drug is introduced (part **P2** of the experiments described in Section 2). As previously mentioned in Section 2 and motivated by the observations in [39], where the treatment TMZ does not seem to induce cell death, we suppose that the drug only affects the elastic properties of the cells. For a cell density $w(\mathbf{x}, t)$ the squeezing probability of part **P2** is

$$q_2(w(\mathbf{x}, t), M) = 1 - \left(\frac{w}{\bar{u}}\right)^{\gamma_M} .\tag{10}$$

Note that we have supposed that the crowding capacity \bar{u} which corresponds to the tight packing state remains unchanged from **P1** to **P2**. This corresponds to the hypothesis that the treatment does not modify the volume of the cells but only change their elastic properties.

The complete PDE system corresponding to the second part of the experiments therefore reads:

$$\begin{aligned}\partial_t w &= \nabla \cdot (d_3 D_2(w, M) \nabla w - \chi_w \phi_2(w, M) \nabla c) + f(w) , \\ \partial_t c &= d_2 \Delta c + \alpha w - \beta c , \\ \partial_t M &= d_4 \Delta M - \delta w ,\end{aligned}\tag{11}$$

where $f(w)$ is again the logistic growth source term given by (1), and the first equation has been derived using the squeezing probability q_2 defined by (10). Note that the carrying capacity u_{\max} remains unchanged in the two parts of the experiments: we have supposed here that the drug does not interact with the nutrients. Analogous to (7), the movement of the cells is described by a chemotactic system where, in this case, the diffusion and chemosensitive coefficients depend also on the concentration of the treatment. These modified coefficients will lead to changes in the size of the aggregates as shown in the numerical experiments in Section 5. The evolution of the concentration c is the same as in (7) where, in this case, the chemoattractant is produced by the new population w . Including proliferation also in this second part allows us to assess the effect of the treatment in the population at earlier times, while the population of cells is still growing and aggregates are still forming.

Different initial conditions for the cell density and concentration of the treatment will be considered as described in Section 5. Each initial condition for **P2** will correspond to a density profile solution of the system **P1** at a given time (solution of (16)), *i.e.* $w(\mathbf{x}, 0) = u(\mathbf{x}, T_1)$ for some given T_1 . We will consider cases where the treatment is introduced on already formed and stable aggregates (steady state of (16)), but also cases where it is introduced at earlier times (during the formation of the aggregates, see Section 5). The initial condition for the concentration of the drug is considered to be either homogeneously distributed in the simulation domain or introduced in the center.

Remark 3.1. *In both parts of the experiments, we assume that the crowding capacity \bar{u} is larger than the carrying capacity u_{\max} . The carrying capacity is defined as the maximum population density beyond which there are not enough nutrients to support growth, while the crowding capacity describes the maximum density in an aggregate depending on the space available.*

4 Linear stability analysis and pattern formation

The system for part **P1** (7) without source term is well known in the literature as the volume-filling Keller-Segel model, for which emergence of patterns has been characterised and is now well documented. Pattern formation refers to the phenomenon by which, after varying a bifurcation parameter, the spatially homogeneous steady state loses stability and inhomogeneous solutions appear. In the following, we investigate in which parameter region we can expect instability of homogeneous solutions, corresponding to the formation of patterns. The linear stability analysis followed here is classical and follows the lines of [43, 55, 35].

We first recall the two systems associated with the dynamics described in parts **P1** and **P2** of the model. Using the fact that in the first part of the experiments, the squeezing probability is chosen to be $q_1(u) = 1 - \frac{u}{\bar{u}}$, we have

$$\begin{aligned}\partial_t u &= \nabla \cdot (d_1 \nabla u - \chi_u \phi_1(u) \nabla c) + ru \left(1 - \frac{u}{u_{\max}}\right), \\ \partial_t c &= d_2 \Delta c + \alpha u - \beta c,\end{aligned}\tag{12}$$

where

$$D_1 = 1, \quad \text{and} \quad \phi_1(u) = u \left(1 - \frac{u}{\bar{u}}\right).\tag{13}$$

For part **P2**, when the squeezing probability function is given by (10), the system writes

$$\begin{aligned}\partial_t w &= \nabla \cdot (d_3 D_2(w, M) \nabla w - \chi_w \phi_2(w, M) \nabla c) + \tilde{r} w \left(1 - \frac{w}{u_{\max}}\right), \\ \partial_t c &= d_2 \Delta c + \alpha w - \beta c, \\ \partial_t M &= d_4 \Delta M - \delta w,\end{aligned}\tag{14}$$

with diffusion and chemotactic coefficients given by

$$D_2(w, M) = 1 + (\gamma_M - 1) \left(\frac{w}{\bar{u}} \right)^{\gamma_M} \quad \text{and} \quad \phi_2(w, M) = w \left(1 - \left(\frac{w}{\bar{u}} \right)^{\gamma_M} \right). \quad (15)$$

4.1 Dimensionless model

To get a deeper insight to the behaviour of the system we introduce the characteristic values of the physical quantities appearing in the models. Denoting by X and T the macroscopic units of space and time, respectively, such that $\bar{\mathbf{x}} = \frac{\mathbf{x}}{X}$, $\bar{t} = \frac{t}{T}$, then we choose

$$(\bar{\mathbf{x}}, \bar{t}) = \left(\sqrt{\frac{\beta}{d_2}} \mathbf{x}, \frac{\beta d_1}{d_2} t \right).$$

Using these new variables, the dimensionless systems write for part **P1**

$$\begin{aligned} \partial_t u &= \nabla \cdot (\nabla u - A \phi_1(u) \nabla c) + r_0 u \left(1 - \frac{u}{u_{\max}} \right), \\ \zeta \partial_t c &= \Delta c + u - c. \end{aligned} \quad (16)$$

Similarly, we obtain for **P2**

$$\begin{aligned} \theta \partial_t w &= \nabla \cdot (D_2(w, M) \nabla w - B \phi_2(w, M) \nabla c) + \tilde{r}_0 n \left(1 - \frac{w}{u_{\max}} \right), \\ \zeta \partial_t c &= \Delta c + n - c, \\ m \partial_t M &= \Delta M - \delta_0 w, \end{aligned} \quad (17)$$

where

$$A = \frac{\chi u}{d_1}, \quad r_0 = \frac{d_2 r}{d_1 \beta}, \quad \zeta = \frac{d_1}{d_2}, \quad \theta = \frac{d_1}{d_3}, \quad B = \frac{\chi w}{d_3}, \quad \tilde{r}_0 = \frac{d_2 \tilde{r}}{d_3 \beta}, \quad m = \frac{d_3}{d_4}, \quad \delta_0 = \frac{\delta}{d_1}. \quad (18)$$

The parameters ζ and m are assumed to be small since the chemoattractant and the chemotherapeutic treatment diffuse faster than the cells. On the other hand, $\theta \simeq 1$ since both population densities u and w are assumed to be diffusing at similar rates. In the following, we state the linear stability for both systems in separate sections.

4.2 First part: Formation of the aggregates

We first consider the system (16), which can be re-written in a more general form as

$$\begin{aligned} \partial_t u &= \nabla \cdot (\nabla u - A \phi_1(u) \nabla c) + f(u), \\ \zeta \partial_t c &= \Delta c + g(u, c), \end{aligned} \quad (19)$$

where $\phi_1(u)$ is given in (13), $f(u) = r_0 u \left(1 - \frac{u}{u_{\max}} \right)$ and $g(u, c) = u - c$. This system is subject to uniformly distributed initial conditions and zero-flux boundary conditions as in (8).

The main result in this section is the following theorem, which gives the conditions for pattern formation for the system (19).

Theorem 4.1. Consider (u^*, c^*) a spatially homogeneous steady state. Then pattern formation for the system (19) with zero flux boundary conditions (8) and initial data (9) is observed if the following conditions are satisfied,

$$\begin{aligned} f_u^* + \zeta^{-1}g_c^* < 0, \quad f_u^*g_c^* > 0, \quad \zeta^{-1}g_c^* + f_u^* - \zeta^{-1}g_u^*A\phi_1(u^*) > 0, \\ g_c^* + f_u^* + g_u^*A\phi_1(u^*) > 2\sqrt{f_u^*g_c^*}. \end{aligned} \quad (20)$$

The critical chemosensitivity is given by

$$A^c = \frac{2\sqrt{r_0} + 1 + r_0}{u_{\max} \left(1 - \frac{u_{\max}}{u}\right)}, \quad (21)$$

and for $A > A^c$ patterns can be expected. The wavenodes k^2 are in the interval defined by

$$k_1^2 = \frac{-m - \sqrt{m^2 - 4f_u^*g_c^*}}{2} < k^2 < k_2^2 = \frac{-m + \sqrt{m^2 - 4f_u^*g_c^*}}{2}, \quad (22)$$

where $m = -(g_c^* + f_u^* + g_u^*A\phi_1(u^*))$.

Proof. See Appendix A for the proof of this theorem. \square

4.3 Second part: Treatment

We now consider the system (17) which corresponds to the second part of the experiments, when the treatment is introduced. The parameter range where patterns are observed is summarised in the following theorem.

Theorem 4.2. Consider (w^*, c^*, M^*) a spatially homogeneous steady state. Also, consider (17) with zero flux boundary conditions given by

$$(d_3D_2(w, M)\nabla w - \chi_w\phi_2(w, M)\nabla c) \cdot \eta = 0, \quad d_2\nabla c \cdot \eta = 0, \quad d_4\nabla M \cdot \eta = 0,$$

and initial conditions

$$\begin{aligned} w(\mathbf{x}, 0) &= u(\mathbf{x}, T_1), \\ c(\mathbf{x}, 0) \text{ of } \mathbf{P2} &\text{ is equal to } c(\mathbf{x}, T_1) \text{ from } \mathbf{P1}, \\ M(\mathbf{x}, 0) &= \begin{cases} \text{constant in } \Omega, & \text{or} \\ C e^{-\frac{(\mathbf{x}-\mathbf{x}_0)^2}{2\sigma^2}}, & \end{cases} \end{aligned} \quad (23)$$

where C is the amplitude, \mathbf{x}_0 is the centre of the Gaussian and σ is the width. Then, the critical chemosensitivity is given by

$$B^c = \frac{2\sqrt{r_0}D_2(u_{\max}, M_s) + D_2(u_{\max}, M_s) + \tilde{r}_0}{u_{\max} \left(1 - \left(\frac{u_{\max}}{u}\right)^{\gamma_{M_s}}\right)},$$

where

$$D_2(u_{\max}, M_s) = 1 + (\gamma_{M_s} - 1) \left(\frac{u_{\max}}{u}\right)^{\gamma_{M_s}}.$$

Patterns can be expected if $B > B^c$ and the wavenodes k^2 are in the interval defined by

$$k_1^2 = \frac{-\bar{m} - \sqrt{\bar{m}^2 - 4D_2(u_{\max}, M_s)(f_w^*g_c^*)}}{2D_2(u_{\max}, M_s)} < k^2 < k_2^2 = \frac{-\bar{m} + \sqrt{\bar{m}^2 - 4D_2(u_{\max}, M_s)(f_w^*g_c^*)}}{2D_2(u_{\max}, M_s)},$$

for $\bar{m} = -(D_2(u_{\max}, M_s)g_c^* + g_w^*B\phi_2(u_{\max}, M_s) + f_w^*)$.

Proof. The proof of this theorem can also be found in Appendix A. \square

Remark 4.1. For the case of 2 dimensions, we can rewrite the systems (16) and (17) using polar coordinates (ρ, θ) where we use the transformation $x = \rho \sin \theta$, $y = \rho \cos \theta$ and the Laplace operator is now given by $\Delta_p = \frac{1}{R} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial}{\partial \rho} \right) + \rho^2 \frac{\partial^2}{\partial \theta^2}$ where R is the radius of the domain. The eigenvalue problem (35) is now written as $-\Delta_p \psi_k = k^2 \psi_k$ with boundary conditions $\partial \psi_k / \partial \rho = 0$ at $\rho = R$.

The eigenfunctions are obtained by separation of variables and are given by $\psi_k(x, y) = \mathcal{R}(\rho)\Theta(\theta)$. Here $\Theta(\theta) = e^{is\theta} = A \cos(s\theta) + B \sin(s\theta)$ for some $s \in \mathbb{Z}$. The radial part $\mathcal{R}(\rho)$ is given in terms of Bessel functions $\mathcal{R}(\rho) = \mathcal{J}_s(k\rho)$ (see [45]) where $k = \frac{c_{s,p}}{R}$ and $c_{s,p}$ denotes the p th zero derivative of \mathcal{J}_s , which is a first kind Bessel function of order m . Finally we can write $\psi_k^{s,p}(\rho, \theta) = \mathcal{J} \left(\frac{c_{s,p}}{R} \rho \right) (A \cos(s\theta) + B \sin(s\theta))$.

The stability analysis reveals that several competing effects control the system's ability to create patterns (aggregates). The criteria obtained both in **P1** or **P2** show that the chemotactic sensitivity must be large enough to compensate the smoothing effect of the diffusion term and of the logistic growth. On the other hand, one can observe from the bifurcation formulae that the ratio $\frac{u_{\max}}{\bar{u}}$ (carrying capacity *vs* density of the tight packing state) plays an important role in the emergence of patterns: larger values lead to more aggregated states. These results show that the logistic growth term has an intrinsic smoothing property, *i.e.*, it tends to force the density to equate the carrying capacity, while the chemotactic term acts as a attractive force and creates zones of higher density (recall that $u_{\max} < \bar{u}$). The aggregates are an expression of a balance in between these two competing effects, which are completely characterised by the stability criterion.




Figure 4: Wavenumbers for different values of γ_M when (a) $r_0 = 0.1$, $u_{\max} = u_0 = 0.5$; and for (b) $r_0 = 0.05$, $u_{\max} = u_0 = 0.1$. Circles indicate the values of k_c .

In order to give more insights about the size of the emerging patterns, we show in Figure 4 the values of the inverse of the maximal wavenumber as function of the chemotactic sensitivity (denoted by A for part **P1** and B for part **P2**), for different values of the exponent γ_M . We recall that $\gamma_M = 1$ in **P1** (blue curve) where cells act as rigid bodies and $\gamma_M > 1$ in the presence of the treatment, where cells behave as semi-elastic entities (**P2**). Figure 4a shows the results for growth rate $r_0 = 0.1$ and carrying capacity $u_{\max} = 0.5$, Figure 4b for $r_0 = 0.05$ (slower growth) and $u_{\max} = 0.1$ (lower carrying capacity). In both figures, the black circles indicate the critical values for the chemotactic sensitivity above which the system is unstable. Here, the tight packing density is set to $\bar{u} = 1$. The maximal wavenumber corresponds to the most unstable mode, *i.e.* the perturbed wave that will

grow the fastest. Therefore, the inverse of this maximal wavenumber is directly related to the size of the emerging patterns. As one can observe, an increase in the chemotactic sensitivity parameter correlates with a decrease in the observed pattern size, suggesting that the aggregates are smaller: larger chemotactic sensitivity leads to more aggregated clusters. Moreover, the aggregate size also decreases as cells pass from rigid bodies to semi-elastic entities (when γ_M increases). This is due to the fact that for larger values of γ_M , cells are more easily deformed and can aggregate more efficiently than when they behave as rigid spheres.

When we increase the ratio $\frac{u_{\max}}{\bar{u}}$ (compare Figure 4a and 4b), we observe that the critical value of the chemosensitivity above which patterns are generated is larger than for smaller ratios $\frac{u_{\max}}{\bar{u}}$. These results highlight once again the smoothing effect of the logistic growth: When the cell tight packing density is unchanged, decreasing the carrying capacity of the environment enhances cell death in the aggregates formed by chemotaxis, where cells try to reach the tight packing state. In this case, the critical chemosensitivity value must be large enough to compensate for the cell death induced by the logistic growth. Moreover, we observe that larger ratios $\frac{u_{\max}}{\bar{u}}$ induce less influence of the parameter γ_M . The cell aggregation abilities are mainly driven by the chemosensitivity intensity and not so much by the cell mechanical properties for large values of $\frac{u_{\max}}{\bar{u}}$.

5 Numerical simulations

In addition to the analytic results obtained in Section 4, we present numerical simulations for the two problems (16) and (17). This allows us to investigate the behavior of the solution of the models for different scenarios and range of parameters. It is well-known that a standard discretization of the Keller-Segel models can lead to nonphysical solutions due to the convective term. Here, we focus on a numerical method that preserves the non-negativity of the cell density using the upwind finite element method described in [10] for the simulation of the Cahn-Hilliard equation.

The calculation of the chemotactic coefficient follows the lines of [3]. Indeed, the finite volume scheme proposed in [3] is identical to the numerical method presented in this paper in dimension one. However, in higher dimensions, and since we also use a finite element method, the numerical scheme presented in [3] differs from the one in this section as detailed in Appendix B.

5.1 Description of the scheme

The domain Ω is discretized by a mesh \mathcal{T}^h constituted of N_{el} piecewise linear finite elements. We denote the total number of nodes by N_h and the standard \mathbb{P}^1 finite element space by S^h . The standard L^2 inner product is indicated by (\cdot, \cdot) and the lumped scalar product defined in Appendix B by $(\cdot, \cdot)^h$. Given $N_T \in \mathbb{N}^*$, let $\Delta t := T/N_T$ be the time-step where T is the time corresponding to the end of the simulation. Let $t_n := n\Delta t$, $n = 0, \dots, N_T - 1$ be the temporal mesh. We approximate the continuous time derivative by $\frac{\partial u_h}{\partial t} \approx \frac{u_h^{n+1} - u_h^n}{\Delta t}$. We define

$$u_h^n(x) := \sum_{j=1}^{N_h} u_j^n \chi_j(x), \quad \text{and} \quad c_h^n(x) := \sum_{j=1}^{N_h} c_j^n \chi_j(x),$$

the finite element approximations of the cell density u and the concentration of chemoattractant c where $\{u_j^n\}_{j=1, \dots, N_h}$ and $\{c_j^n\}_{j=1, \dots, N_h}$ are unknowns and $\{\chi_j\}_{j=1, \dots, N_h}$ is the finite element basis. Then, the finite element problem associated with the system (16) reads as follows.

For each $n = 0, \dots, N_T - 1$, find $\{u_h^{n+1}, c_h^{n+1}\}$ in $S^h \times S^h$, such that for all $\chi \in S^h$

$$\begin{aligned} \left(\frac{u_h^{n+1} - u_h^n}{\Delta t}, \chi \right)^h + (D(u_h^n) \nabla u_h^{n+1}, \nabla \chi) = \\ (A(\phi^{\text{upw}}(u_h^n)) \nabla c_h^n, \nabla \chi) + r_0 \left(u_h^n \left(1 - \frac{u_h^n}{u_{\max}} \right), \chi \right)^h, \end{aligned} \quad (24)$$

$$\zeta \left(\frac{c_h^{n+1} - c_h^n}{\Delta t}, \chi \right)^h = -(\nabla c_h^{n+1}, \nabla \chi) + (u_h^{n+1} - c_h^{n+1}, \chi)^h. \quad (25)$$

The function ϕ^{upw} is defined by (66) in Appendix B. The finite element scheme associated with the system (17) including the effect of the treatment is the following

$$\begin{aligned} \theta \left(\frac{w_h^{n+1} - w_h^n}{\Delta t}, \chi \right)^h + (\bar{D}(w_h^n, M_h^n) \nabla w_h^{n+1}, \nabla \chi) \\ = \left(B(\bar{\phi}_2^{\text{upw}}(w_h^n, M_h^n)) \nabla c_h^n, \nabla \chi \right) + \tilde{r} \left(w_h^n \left(1 - \frac{w_h^n}{u_{\max}} \right), \chi \right)^h, \end{aligned} \quad (26)$$

$$\zeta \left(\frac{c_h^{n+1} - c_h^n}{\Delta t}, \chi \right)^h = -(\nabla c_h^{n+1}, \nabla \chi) + (w_h^{n+1} - c_h^{n+1}, \chi)^h, \quad (27)$$

$$m \left(\frac{M_h^{n+1} - M_h^n}{\Delta t}, \chi \right)^h = -(\nabla M_h^{n+1}, \nabla \chi) - \delta (M_h^{n+1}, \chi)^h. \quad (28)$$

The method to compute the coefficient $\bar{\phi}^{\text{upw}}$ and the diffusion coefficient $\bar{D}(w_h^n, M_h^n)$ is the same as for the problem (24)–(25) (see Appendix B).

5.2 Biological relevance of the model parameters

Here, we comment on the choice of the model parameters that we will use for the numerical simulations and how they relate to experimental known data. In [19, 13], the proliferation rate for well oxygenated glioma cells in vitro r was shown to lie between 0.5 and 1 day⁻¹. As the proliferation rate relies significantly on the nutrient, also smaller value seems to be biologically admissible in real conditions and following [13] we choose $r = 0.4$ day⁻¹ and $r = 0.8$ day⁻¹ (corresponding to the non-dimensionalised parameter $r_0 = 0.05$ and $r_0 = 0.1$). As hypoxia-inducible factors (HIF) are supposed to be responsible for the chemotaxis motion of GBM cells, we suppose that the diffusion coefficient d_2 and consumption rate β for the chemoattractant are linked to biological measurements of the oxygen diffusion in human brain which were estimated in [52, 19] to $d_2 = 86.4$ mm² day⁻¹ and $\beta = 8640$ day⁻¹. However, we chose to consider slightly lower values $d_2 = 8.64$ mm² day⁻¹ and $\beta = 864$ day⁻¹ owing to the large variability of this parameter according to the type of tissue (see [51]). For such values and using the scaling of Section 4.1, one unit of time of our model corresponds to approximately 0.1 day and one unit of space is 0.1 mm. As we found no experimental data on the chemotactic coefficient χ_u of glioma cells in response to chemoattractant concentration, the choice of this parameter is driven by the stability analysis and we find that the interesting regimes are obtained for a dimensionless chemosensitivity in between 7 and 70, corresponding to a chemotactic coefficient $\chi_u \in [0.6, 6]$ mm² day⁻¹. Moreover, as no measurements for glioma cells diffusion coefficient are available in the literature, the parameter d_1 is arbitrarily chosen to be 100 times smaller than the chemoattractant diffusion speed and we choose $d_1 = d_3 = 0.086$ mm² day⁻¹, i.e the non-dimensionalised parameter $\zeta = \frac{d_1}{d_2} = 0.01$.

5.3 Numerical results for a one dimensional case

For all numerical computations we choose the packing capacity $\bar{u} = 1$. We consider different proliferation rates $r_0 = 0.1, 0.05$ and different initial conditions and carrying capacities $u_{\max} = u_0 = 0.1, 0.5$. The nondimensional parameters given in (18) are $\zeta = m = 0.01$ and $\theta = 1$ since we assume that the chemoattractant c and the treatment diffuse much faster than the cells, while the motility of the cells is not affected by the treatment, so $d_1 \approx d_3$. The initial condition for the cell density u_0 is assumed to be randomly distributed in space. Similarly, we can also define the initial chemoattractant concentration c_0 .

In this section we start by solving the systems (16) and (17) on the interval $[0, L]$ with homogeneous non-flux boundary conditions using the method described in Section 5.1. In Appendix C we investigate the effect of the size of the domain as well as the effect of the parameters A and B on the formation and evolution of patterns. Moreover, using (17) we study the effect of the treatment using the solution of (16) at the final time T_1 as initial condition. We explore the case when we introduce the treatment at earlier stages of the formation of the aggregates.

We consider two different scenarios for the evolution of the concentration of the treatment. First, we assume that the treatment diffuses very fast in the whole domain so that the concentration is homogeneous from time $T_2 = 0$. The other case we consider, which is closer to real experiments, starts with a high concentration of the drug in the centre of the domain and this concentration diffuses over time according to the third equation in (17).

Comparison with the linear stability analysis In order to quantify the aggregate sizes and compare it to the ones predicted by the stability analysis, we use the Fourier transform of the numerical solution and extract the frequency that corresponds to the maximal Fourier mode. For the sake of this analysis, periodic boundary conditions are therefore considered. To this aim, we compute the discrete Fourier transform $\mathcal{F}[u](\mathbf{x}, t) = \hat{u}(\lambda, t)$ and define

$$k_{\max} = \arg \max_{\lambda} (|\hat{u}(\lambda)|) = \arg \max_{\lambda} \left(\sqrt{\operatorname{Re}(\hat{u}(\lambda))^2 + \operatorname{Im}(\hat{u}(\lambda))^2} \right),$$

which corresponds to the frequency of the largest Fourier mode. The inverse $(k_{\max})^{-1}$ of this maximal frequency relates to the pattern size. This maximal frequency of the Fourier transform of the solution is expected to correspond to the maximal wavenumber predicted by the stability analysis. We show in Figure 5 the values of $(k_{\max})^{-1}$ computed from the numerical solution (blue dotted line) compared to the predictions of the stability analysis (red curve), as function of the chemosensitivity, for $\gamma_M = 1$ (left figure) and $\gamma_M = 5$ (right figure). As one can observe, we obtain a very good agreement between the numerical values and the predictions of the stability analysis, and we recover the critical value of the chemosensitivity parameter above which the system generates patterns.

Introduction of the treatment on already-formed aggregates In this part, we aim to study the influence of the treatment on already formed aggregates. For this, we let the system run in **P1** (without treatment, $M = 0, \gamma_M = 1$) until time $T_1 = 200$ and introduce the treatment uniformly in the domain ($M = 1, \gamma_M = 5$).

In Figure 6 left and middle panels, we choose values of the chemosensitivity very close to the critical values corresponding to k_c , where the wavenumbers are very different for the cases $\gamma_M = 1$ and $\gamma_M = 5$ as we see in Figure 4a. In Figure 6, the blue curves describe the formation of aggregates for a time $T_1 = 200$ without the treatment, while the cells are proliferating with rate $r_0 = 0.1$. We




Figure 5: Comparison of the wavelength obtained analytically, using (45) and (46), and numerically, using the Fourier transform of the solution for (a) $\gamma_M = 1$ and (b) $\gamma_M = 5$.

consider two different scenarios when introducing the treatment: either cells stop proliferating (red curves), or they continue with the same rate as before $r_0 = 0.1$ (yellow curves).

When we introduce the treatment for values of A and B close to the instability threshold ($A = B = 7$, left figure), we observe that the aggregates become steeper and the density in each aggregate reaches the packing capacity $\bar{u} = 1$. This clearly leads to more compact aggregates as a result of the nonlinearity introduced in (11) by the parameter γ_M . The main physical difference between changing the parameter γ_M and changing the chemosensitivity coefficients A or B is the following. By changing A or B depending on the concentration of the treatment, we are enhancing aggregation over diffusion, essentially we are changing the motility of cells. By changing γ_M the motility, as well as the elastic properties of the cells in the aggregates are affected. When we introduce the treatment while cells keep proliferating, aggregates tend to merge together since the density is growing, as we see in Figure 6 middle panel.

It is noteworthy that for large values of the chemosensitivity parameter ($A = B = 70$, right panel in Figure 6), the treatment does not impact the aggregate dynamics. In this case, cell aggregation is mainly driven by the chemotactic term and the cell mechanical properties have little influence. These observations are in agreement with the stability analysis, which shows that the parameter γ_M has more influence when the chemotactic sensitivity is close to the instability threshold.

Finally, by comparing the red and yellow curves of Figure 6, it is clear that cell proliferation has a major impact on the size of the aggregates. If the treatment has the double effect to stop proliferation as well as modifying the cell mechanical properties, cell aggregates will become very dense and well-separated, while merging aggregates are still observed if the treatment has the sole effect to change the cell mechanical properties. Here, we chose to introduce the treatment at time $T_1 = 200$, we now aim to study the effect of the treatment introduced at different times in the aggregation process.

Introduction of the treatment at different times Here, we consider the case when the treatment is introduced at different times in the aggregation process. As before, we let the system run in **P1** (without drug, $M = 0, \gamma_M = 1$) until time $t = T_1$ and introduce the treatment uniformly in the domain ($M = 1, \gamma_M = 5$). We consider the cases when the treatment has the ability to stop proliferation, and when the treatment only acts on the cell mechanical properties. In Figure 7, we show the results at time $T = T_1 + 200$ (red curve), when the treatment is introduced at times $T_1 = 50$ (left plots), $T_1 = 100$ (middle plots) and $T_1 = 300$ (right plots). Figures 7a show the results




Figure 6: Aggregation pattern when $A = B$ for $A = 7$ (left), $A = 12$ (middle) and $A = 70$ (right). Blue curve: solution at $T_1 = 200$ without treatment, red curves: solution at $T = T_1 + 200$ when the treatment is introduced uniformly with $\gamma_M = 5$ and proliferation is stopped, yellow curves: solution with the same parameters as the red curves when proliferation goes on with $r_0 = 0.1$.

when the treatment stops proliferation, Figures 7b when the treatment only changes the mechanical properties of the cells. For each, the blue curves are the density profiles before introducing the treatment. As one can observe, in Figures 7a and 7b, introducing the treatment at different times of the aggregation process have a major impact on the size of the aggregated patterns formed at a latter time. Introducing the treatment at an earlier time ($T_1 = 50$, left figures) enables to obtain smaller aggregates compared to when the treatment is introduced on already formed aggregates ($T_1 = 300$, right figures). This effect is more visible when the treatment has the double effect of blocking cell proliferation and changing the elasticity (compare red curves in Figures 7a and 7b). In this case, the earlier the treatment is introduced, the smaller the aggregated patterns. When the treatment stops proliferation as well as the cell mechanical properties and is introduced at later times (right panel of Figure 7a) we recover the observation of the real systems, where the treatment induces a shrinking of the aggregate and favors the formation of more compact cell structures. This effect is not observed when proliferation is active with the treatment, (right panel of Figure 7b) where aggregates are merging and they are larger than before the treatment introduction. This suggests indeed that the treatment has the double effect of blocking cell proliferation as well as changing the cell mechanical properties. The model suggests that introducing the treatment at earlier times of the tumour development could enable us to control the size and separation of the tumour aggregates.

Introduction of the treatment in the middle of the domain Finally, we aim to study the case when the treatment is introduced in the center of the domain and diffuses in the environment. Here we assume that the treatment is not consumed or escapes the domain, therefore $\delta_0 = 0$ in (17). In Figure 8, we show the density profiles of the solution before introducing the treatment (blue curves), and when the treatment is present (red curves), at times $T_2 = 0$ (left), $T_2 = 5$ (center) and $T_2 = 30$ (right). The distribution of the treatment follows a Gaussian of the form $M(\mathbf{x}, 0) = Ce^{-\frac{(\mathbf{x}-\mathbf{x}_0)^2}{2\sigma^2}}$, where $C = 40$ is the amplitude, \mathbf{x}_0 is the center of the Gaussian and $\sigma = 0.5$ describes the spread. As one can observe, the large concentration of the treatment in the middle immediately sharpens the interface between already-formed aggregates, and favors the separation of the cell clusters. As the treatment diffuses (middle figure), the clusters interfaces which sense a high concentration of the treatment sharpen, creating denser and well-separated cell clusters.




Figure 7: Introduction of the the treatment at different times $T_1 = 50, 100, 300$ when $u_0 = 0.5$ and $A = B = 12$. The blue curves gives the initial condition for the part **P2** of the experiment with the treatment, represented by the red curve. (a) Without proliferation. (b) With proliferation, $r_0 = 0.1$. The red curves are at $T = T_1 + 200$.




Figure 8: Evolution of the pattern when the treatment is introduced in the center of the domain with amplitude 40. The blue curves correspond to the solution without treatment at $T_1 = 200$ and the red curves are the solution with the treatment at $T = T_1 + 5$ and $T = T_1 + 30$.

5.4 Numerical results for a two dimensional case

For the 2D simulations we consider that Ω is a disk of radius R which can be defined as $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < R^2\}$ where the boundary is given by $\partial\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = R^2\}$. The proliferation rate is chosen to be $r_0 = 0.05$ and the initial homogeneous density as well as the carrying capacity are set to $u_{\max} = u_0 = 0.1$ or $u_{\max} = u_0 = 0.5$. The other parameters can be found at the beginning of Section 5.3.

In Figure 9 we show the formation of aggregates for different values of A without the treatment, for $u_0 = 0.1$ and $r_0 = 0.05$. We observe that for $A = 10$ (Figure 9a) we do not have patterns, in agreement with the analytic results obtained in Figure 7b since this value of A is less than $A^c \approx 16.7$. As we increase the chemosensitivity parameter, the aggregates become more compact. From Figure 9b we observe the phenomena of two aggregates merging together, analogous to the one dimensional results in Figure 15. As expected, by changing the carrying capacity and the initial density of cells to $u_0 = u_{\max} = 0.5$, the patterns change shape. We observe a transition from spot-like patterns in Figure 9 to maze-like structures in Figure 10. This behaviour has been widely studied experimentally [41], numerically [34, 36] and more recently, also including a volume-filling approach [43].




Figure 9: Formation of aggregates at $T_1 = 200$ when $u_0 = 0.1$, $r_0 = 0.05$ and (a) $A = 10$, (b) $A = 20$ and (c) $A = 70$.




Figure 10: Formation of aggregates at $T_1 = 200$ without the treatment and when $u_{\max} = u_0 = 0.5$, $r_0 = 0.05$ and (a) $A = 7$, (b) $A = 12$ and (c) $A = 50$.

Analogous to the one dimensional case, we consider two different initial conditions for the treatment: (i) we first include the treatment uniformly in the domain with $M = 5$, and (ii) we introduce the treatment in the middle of the domain and let it diffuse in space. In these simulations, the treatment is supposed to block proliferation as well as changing the cell mechanical properties.

In order to compare the change in size of the aggregates before and after the treatment, we compute the difference between the solution of the first part of the experiments $u(\mathbf{x}, T_1)$ coming

from (16), when the aggregates are formed (at $T_1 = 200$), and the solution $w(\mathbf{x}, t)$ of (17), once the treatment has been inserted (at time $T = 200 + T_2$). In Figure 11, we show the results for different values of the chemosensitivity parameter $A = B$, when the aggregates have been exposed to a uniform distribution of the treatment for a time $T_2 = 30$. We notice that the aggregates without the treatment are bigger and therefore the treatment induces a shrinking in the size of the pattern, as described in Section 3. When we compare the results in Figure 11a and 11b for different values of A and B we observe that the effect of the treatment is stronger when the value of the chemosensitivity parameter B is closer to its critical value (see Figure 4b). This is in accordance with the results of the stability analysis. The cell mechanical properties (controlled by the parameter γ_M) have less influence on the cell cluster sizes when the chemosensitivity parameter $A = B$ is increased (see Figure 4b).




Figure 11: Difference between the solutions $u(\mathbf{x}, 200) - w(\mathbf{x}, 230)$ when (a) $A = B = 20$ and (b) $A = B = 70$. Here $\gamma_M = 11$, $r_0 = 0.05$ and $u_{\max} = u_0 = 0.1$.




Figure 12: Difference between the solutions when the initial concentration of the treatment is a Gaussian function centered in the domain. (a) $T = 210$, (b) $T = 230$ (c) $T = 300$ for $T_1 = 200$, $r_0 = 0.05$ and $u_0 = 0.1$.

We now study the case when the treatment is introduced in the middle of the domain and diffuses in the environment. To this aim, the initial concentration of the treatment is supposed to be a Gaussian function with width 5 centered in the middle of the domain. We assume that the treatment is not consumed by the cells in the time scales we are interested in, and choose $\delta_0 = 0$. In Figure 12 we show the evolution of the difference between the two solutions $u(\mathbf{x}, T_1) - w(\mathbf{x}, t)$, where $T_1 = 200$ is the time at which the treatment is introduced and T is the duration of the treatment. We explore different times $T_2 = 10, 30, 100$. For short times, the effect of the treatment is only noticed by the aggregates at the center of the domain and therefore the difference between the two

solutions close to the boundaries is zero. As time increases, the concentration of the treatment reaches the whole domain as is observed in Figure 12c.




Figure 13: Effect of the treatment at different times while the aggregates are forming. Top row: Cell aggregates for $A = 20$ without treatment at (a) $T_1 = 30$, (b) $T_1 = 50$ and (c) $T_1 = 100$. Bottom row: Cell aggregates at time $T = T_1 + 200$, when the treatment has been introduced at times (d) $T_1 = 30$, (e) $T_1 = 50$ and (f) $T_1 = 100$.

Finally, we also study the effect of the treatment at earlier stages of the formation of the aggregates. Figure 13 shows the different patterns obtained during the formation of the aggregates at different times, top row, and the corresponding effect of the treatment, bottom row. For example, introducing the treatment at $T_1 = 50$ leads to a significant reduction of the size of the pattern with a reasonably low concentration of the treatment. Identifying this specific time in real patients could make the treatment much more effective and reduce the spread of the cancer cells.

6 Discussion of results and perspectives

In this paper we propose a mechanism for the effect of certain treatments on tumours formed by a chemotaxis type self-organization phenomenon. Inspired by the experiments concerning the action of TMZ on Glioblastoma cells mentioned in Section 2, we considered the particular case of treatments that do not act as cytotoxic agents but rather induce stress in the environment, which may induce changes in the mechanical properties of individual tumour cells by making them pass from rigid bodies to semi-elastic entities. We explored two scenarios: a first one where only cell's plasticity is impacted by the treatment, and a second one where the treatment has a double effect of preventing cell proliferation as well as changing cell mechanics.

Under these hypotheses, we obtained a modified version of the Keller-Segel model, known as the nonlinear volume-filling approach for cell motion, where the cell mechanical properties are taken into account in the form of the so-called squeezing probability. In the nonlinear volume-filling Keller-Segel

model, this squeezing probability function could be related to the amplitude of the transport term towards zones of high chemoattractant density (chemosensitivity), as well as with the (nonlinear) diffusion coefficient.

By performing a linear stability analysis, we are able to detect and characterise the parameter ranges for which the homogeneous distribution is unstable, *i.e.* the ranges for which patterns appear as the result of the dynamics. We show that the emergence of patterns without treatment (*i.e.* when cells act as rigid bodies) is driven by a fine interplay between the chemotactic sensitivity, which tends to aggregate the cells, and the diffusion, together with the logistic growth, which tend to smoothen the solution. We are able to compute the critical chemosensitivity value above which the system self-organizes into aggregates, and characterise the size of the aggregates as a function of the model parameters.

Under treatment, *i.e.* when cells behave as semi-elastic entities, we show that the critical value of the chemosensitivity above which patterns emerge is smaller than that in control cultures, showing that as cells become more elastic, they tend to aggregate more easily than when they behave as rigid entities.

We are able to completely characterise the size of the patterns and show that semi-elastic cells create smaller aggregates than rigid entities for the same value of the chemosensitivity. These results suggest that the mechanical properties of individual cells have a huge impact on the shape and size of the aggregated patterns at the population level.

Moreover, we show that the ratio between the tight packing cell density and the carrying capacity of the TME plays a major role in the size and shape of the obtained patterns. For large values of this ratio, the aggregation abilities of the system are essentially driven by the chemotactic transport term while the individual cell mechanical behaviour has little impact on the shape and size of the patterns. However, for smaller values of this ratio, *i.e.* when the tight packing density is closer to the carrying capacity of the environment, cell mechanics has a huge influence on the behavior of the population.

These results are confirmed by numerical experiments in 1D and 2D for which, given an initial perturbation of the homogeneous cell distribution, we observe the emergence of cell aggregates and we recover the critical values of the chemosensitivity predicted by the stability analysis. We obtain a very good correspondence between the simulations and the theoretical predictions, for the appearance of patterns as well as their size.

By performing simulations of the whole system, we recover the experimental observations: introducing the treatment (TMZ in the experiments mentioned in Section 2) on already-formed aggregates leads to the quick formation, of more compact patterns. As the treatment diffuses in the domain (changing locally the cell mechanical properties as it goes), it sharpens the border of the cell aggregates and leads to denser and well-separated clusters.

While the border sharpening of the clusters is independent on whether proliferation is activated or not during treatment, the shrinking of the clusters is more clear when the treatment has the double effect of changing the cell mechanical properties as well as blocking cell proliferation. Indeed, when proliferation is still active in the presence of the treatment, we observe the merging of existing clusters and this results in aggregates being larger than before treatment. These results suggest a possible mechanism for the shrinking of the aggregates observed under the experimental conditions described in Section 2: TMZ might not only stop cell proliferation, but might also generate a stress in the environment to which cells respond to by changing their mechanical state.

While alterations of mechanical properties, around or inside the tumour, are common in solid tumours including GBM, the question of the nature and the regulation of cancer cells through mechano-sensitive pathways are largely unknown. Recently, in a *Drosophila* model, glioma progression has been associated to a regulatory loop mediated by the mechano-sensitive ion channel

Piezol and tissue stiffness [12]. A direct perspective of these works consists in verifying the potential mechano-sensing effect of TMZ proposed in the present model, through direct measures in real systems by studying the mechanical properties of individual GBM cells which have been exposed to TMZ treatment. The targeting of mechano-sensitive pathways after TMZ treatment may provide new therapeutic angles in GBM and in more general settings.

Moreover, it would be interesting to identify other clinical settings where the effects of the treatment are similar to those of TMZ in GBM, and to check if the effects are due to changes in the tumor cell properties corresponding to the general hypothesis of the model constructed in this work.

In the future, better quantitative comparison with experiments will allow for systematic choice of parameters and validation of the mechanisms we propose here. From a biological point of view, a natural sequel of this work consists in studying the coupled effect of TMZ and irradiation. Indeed, even if TMZ alone seems not to suffice to decrease the tumour mass, the coupling of TMZ treatment with irradiation has been shown to have more efficient effects than irradiation alone [47, 5, 53]. It would also be interesting to introduce a second treatment with cytotoxic effects in this model, to study whether the mechanical changes of individual cells induced by TMZ could explain the better response of the system to irradiation treatments.

A Stability analysis

First part of the experiments We first observe that the homogeneous distributions $u(\mathbf{x}, t) = u^*$ and $c(\mathbf{x}, t) = c^*$ are steady-states solutions of system (16) for u^* and c^* such that $f(u^*) = 0$ and $g(u^*, c^*) = 0$. In order to study their stability, we consider the system without spatial variations

$$\partial_t u = f(u) , \quad \zeta \partial_t c = g(u, c) , \quad (29)$$

and linearize the solution at (u^*, c^*) . We obtain

$$\partial_t \sigma = G \sigma , \quad \text{where } \sigma = \begin{pmatrix} u - u^* \\ c - c^* \end{pmatrix} \quad \text{and } G = \begin{pmatrix} f_u^* & 0 \\ \zeta^{-1} g_u^* & \zeta^{-1} g_c^* \end{pmatrix} , \quad (30)$$

where the quantities f_u^* , g_u^* and g_c^* are the linearization slopes of f and g : $f_u^* = f'(u^*)$, $g_u^* = \partial_u g(u^*, c^*)$, $g_c^* = \partial_c g(u^*, c^*)$. The steady state is linearly stable if $\text{tr}(G) < 0$ and $\det(G) > 0$, which imposes the following constraints on the kinetic functions $f(u)$ and $g(u, c)$,

$$f_u^* + \zeta^{-1} g_c^* < 0 \quad \text{and} \quad f_u^* g_c^* > 0 . \quad (31)$$

Note that in our case, $f_u^* = -r_0$, $g_u^* = 1$, $g_c^* = -1$ so the conditions are satisfied.

We now go back to the full chemotactic system (19). In order to investigate the stability of the homogeneous steady-state, *i.e.* the ability of the system to create patterns, we introduce a small parameter $\varepsilon \ll 1$ and write

$$u = u^* + \varepsilon \tilde{u}(\mathbf{x}, t) , \quad c = c^* + \varepsilon \tilde{c}(\mathbf{x}, t) . \quad (32)$$

We substitute (32) into (19) and, computing the first order terms with respect to ε and neglecting higher order terms, the linearized system reads

$$\begin{aligned} \partial_t u &= \Delta u - A \phi_1(u^*) \Delta c + u f_u^* + c f_c^* , \\ \zeta \partial_t c &= \Delta c + u g_u^* + c g_c^* , \end{aligned} \quad (33)$$

where $\phi_1(u^*) = u^* q_1(u^*)$. We now look for perturbations of the form

$$u(\mathbf{x}, t) = \sum_k a_k(t) \psi_k(\mathbf{x}) \quad \text{and} \quad c(\mathbf{x}, t) = \sum_k b_k(t) \psi_k(\mathbf{x}) , \quad (34)$$

where $(\psi_k)_{k \geq 1}$ is an orthonormal basis of $L^2(\Omega)$ and satisfies the following spatial eigenvalue problem

$$-\Delta \psi_k = k^2 \psi_k , \quad \frac{\partial \psi_k}{\partial \eta} = 0 . \quad (35)$$

Then, the linearized system (33) can be written as

$$\begin{aligned} \partial_t (a_k) &= -a_k k^2 + A \phi_1(u^*) b_k k^2 + a_k f_u^* + b_k f_c^* , \\ \zeta \partial_t (b_k) &= -b_k k^2 + a_k g_u^* + b_k g_c^* , \end{aligned} \quad (36)$$

where k is the spatial eigenfunction, also called the wavenumber and $1/k$ is proportional to the wavelength ω . In matrix form we can write (36) as $\partial_t X_k(t) = P_k(t) X_k(t)$ where

$$X_k = \begin{pmatrix} a_k \\ b_k \end{pmatrix} , \quad P_k = \begin{pmatrix} -k^2 + f_u^* & A \phi_1(u^*) k^2 + f_c^* \\ \zeta^{-1} g_u^* & \zeta^{-1} (-k^2 + g_c^*) \end{pmatrix} . \quad (37)$$

Remark A.1. Since the solutions of the eigenvalue problem (35) are simply sines and cosines, the “size” of various spatial patterns is measured by the wavelength of the trigonometric functions. For example, in one dimension when $0 < x < L$, $\psi \propto \cos(n\pi x/L)$ and the wavelength is $\omega = 1/k = L/n\pi$, where $n \in \mathbb{Z}$.

If the matrix P_k has eigenvalues with positive real part, then the homogeneous steady state (u^*, c^*) is unstable, resulting in pattern formation. The characteristic polynomial related to (37) is given by $\ell^2 + a(k^2)\ell + b(k^2) = 0$ where

$$a(k^2) = (1 + \zeta^{-1})k^2 - (f_u^* + \zeta^{-1}g_c^*) , \quad (38)$$

$$b(k^2) = \zeta^{-1}k^4 - \zeta^{-1}(g_c^* + f_u^* + g_u^*A\phi_1(u^*))k^2 + \zeta^{-1}f_u^*g_c^* . \quad (39)$$

The eigenvalues ℓ determine the temporal growth of the eigenmodes, and we require $\mathcal{Re}(\ell(k^2)) > 0$ for the homogeneous steady state to be unstable. Note that we only look for the eigenmodes $k \neq 0$ since we already guaranteed that the steady state is stable in the absence of spatial perturbations, i.e. $\mathcal{Re}(\ell(k^2 = 0)) < 0$ in (31).

From the conditions (31), we know that $a(k^2) > 0$, hence the instability can only occur if $b(k^2) < 0$ for some k so that the characteristic polynomial has one positive and one negative root. This implies

$$k^4 - (g_c^* + f_u^* + g_u^*A\phi_1(u^*))k^2 + f_u^*g_c^* < 0 . \quad (40)$$

We also know from (31) that $f_u^*g_c^* > 0$, then a necessary (but not sufficient) condition for $b(k^2) < 0$ is

$$g_c^* + f_u^* + g_u^*A\phi_1(u^*) > 0 .$$

Remark A.2. The bifurcation between spatially stable and unstable modes occurs when the critical expression $b_{\min}(k_{\min}^2) = 0$ is satisfied.

Moreover, to satisfy (40) the minimum b_{\min} must be negative [35]. Differentiation with respect to k^2 in (39) leads to

$$b_{\min}(k_{\min}^2) = -\frac{(g_c^* + f_u^* + g_u^*A\phi_1(u^*))^2}{4} + f_u^*g_c^* . \quad (41)$$

Hence, the condition $b_{\min} < 0$ implies

$$g_c^* + f_u^* + g_u^*A\phi_1(u^*) > 2\sqrt{f_u^*g_c^*} . \quad (42)$$

To summarise, we have obtained the following conditions for the generation of spatial patterns for the chemotaxis system (19),

$$\begin{aligned} f_u^* + \zeta^{-1}g_c^* < 0 , \quad f_u^*g_c^* > 0 , \quad \zeta^{-1}g_c^* + f_u^* - \zeta^{-1}g_u^*A\phi_1(u^*) > 0 , \\ g_c^* + f_u^* + g_u^*A\phi_1(u^*) > 2\sqrt{f_u^*g_c^*} . \end{aligned} \quad (43)$$

From the analysis in this section, and using the particular forms of $\phi_1(u)$, f and g as in (19), it is easy to see that the spatially homogeneous steady states are $(0, 0)$ and (u_{\max}, u_{\max}) . We can check that $(0, 0)$ is an unstable steady state, therefore we only work with (u_{\max}, u_{\max}) which, on the contrary, is stable. The first and second properties in (43) are immediately satisfied, i.e., $-(r_0 + \zeta^{-1}) < 0$ and $r_0/\zeta > 0$, respectively. Finally, we have to check that the third and fourth conditions are satisfied as well. We have that

$$-1 - r_0 + A\phi_1(u^*) > 2\sqrt{r_0} . \quad (44)$$

Therefore, (44) is a necessary condition for pattern formation for the original system (12). Considering A as a bifurcation parameter, we can obtain a critical value A^c , so that we observe pattern formation if $A > A^c$. From (44) we get

$$A^c = \frac{2\sqrt{r_0} + 1 + r_0}{u_{\max} \left(1 - \frac{u_{\max}}{\bar{u}}\right)}. \quad (45)$$

The corresponding critical wavenumber k_c^2 is obtained from (41) using (44) as follows,

$$k_c^2 = \frac{g_c^* + f_u^* + g_u^* A^c \phi_1(u^*)}{2} = \sqrt{f_u^* g_c^*} = \sqrt{r_0}. \quad (46)$$

This means that, within the unstable range, $\mathcal{R}e(\ell(k^2)) > 0$ has a maximum wavenumber given by k_c^2 . The range of linearly unstable modes $k_1^2 < k^2 < k_2^2$ is obtained from $b(k^2) = 0$,

$$k_1^2 = \frac{-m - \sqrt{m^2 - 4f_u^* g_c^*}}{2} < k^2 < k_2^2 = \frac{-m + \sqrt{m^2 - 4f_u^* g_c^*}}{2}, \quad (47)$$

where $m = -(g_c^* + f_u^* + g_u^* A \phi_1(u^*))$.

Second part of the experiments Following the same steps as before we linearise the system (17) to get

$$\begin{aligned} \partial_t w &= D_2(w^*, M^*) \Delta w - B \phi_2(w^*, M^*) \Delta c + w f_w^*, \\ \zeta \partial_t c &= \Delta c + w g_w^* + c g_c^*, \\ m \partial_t M &= \Delta M - \delta_0 w, \end{aligned} \quad (48)$$

where

$$D_2(w^*, M^*) = 1 + (\gamma_M^* - 1) \left(\frac{w^*}{\bar{u}}\right)^{\gamma_M^*}, \quad \phi_2(w^*, M^*) = w \left(1 - \left(\frac{w^*}{\bar{u}}\right)^{\gamma_M^*}\right),$$

and γ_M^* is given by (5) evaluated at M^* . As in (34), we let

$$w(\mathbf{x}, t) = \sum_k a_k(t) \psi_k(\mathbf{x}), \quad c(\mathbf{x}, t) = \sum_k b_k(t) \psi_k(\mathbf{x}), \quad M(\mathbf{x}, t) = \sum_k c_k(t) \psi_k(\mathbf{x}), \quad (49)$$

where $\psi_k(\mathbf{x})$ satisfies (35) and we obtain a system $\partial_t X_k(t) = P_k(t) X_k(t)$ where

$$X_k = \begin{pmatrix} a_k \\ b_k \\ c_k \end{pmatrix}, \quad P_k = \begin{pmatrix} -D_2(w^*, M^*) k^2 + f_w^* & B \phi_2(w^*, M^*) k^2 & 0 \\ \frac{1}{\zeta} g_w^* & \frac{1}{\zeta} (-k^2 + g_c^*) & 0 \\ -\frac{\delta_0}{m} & 0 & -\frac{1}{m} k^2 \end{pmatrix}. \quad (50)$$

Similar to the previous section, the characteristic polynomial is given by $a(k^2)\ell^3 + b(k^2)\ell^2 + c(k^2)\ell + d(k^2) = 0$ where $a(k^2) = -1$ and

$$b(k^2) = -\left(D_2 + \frac{1}{\zeta} + \frac{1}{m}\right) k^2 + f_w^* + \frac{g_c^*}{\zeta}, \quad (51)$$

$$\begin{aligned} c(k^2) &= -\left(\frac{D_2}{\zeta} + \frac{D_2}{m} + \frac{1}{m\zeta}\right) k^4 + \left(\frac{g_c^* D_2}{\zeta} + \frac{g_c^*}{\zeta m} + \frac{f_w^*}{\zeta} + \frac{f_w^*}{m} + \frac{g_w^* B \phi_2}{\zeta}\right) k^2 \\ &\quad - \frac{f_w^* g_c^*}{\zeta}, \end{aligned} \quad (52)$$

$$d(k^2) = -\frac{D_2}{m\zeta} k^6 + \frac{1}{m\zeta} (D_2 g_c^* + g_w^* B \phi_2 + f_w^*) k^4 + \frac{1}{m\zeta} (-f_w^* g_c^*) k^2. \quad (53)$$

In general, the stability analysis for this cubic polynomial will require the Ruth–Hurwitz stability criterion [25] which states that the steady state is unstable if the coefficients of $a(k^2)\ell^3 + b(k^2)\ell^2 + c(k^2)\ell + d(k^2) = 0$ satisfy the condition

$$\frac{1}{(a(k^2))^2} (b(k^2)c(k^2) - a(k^2)d(k^2)) < 0 .$$

However, from (50) we observe that one of the eigenvalues of the matrix P_k is given by $\ell_1 = \frac{-k^2}{m} < 0$. The remaining two eigenvalues can be computed from the upper-left matrix

$$\begin{pmatrix} -D_2(w^*, M^*)k^2 + f_w^* & B\phi_2(w^*, M^*)k^2 \\ \frac{1}{\zeta}g_w^* & \frac{1}{\zeta}(-k^2 + g_c^*) \end{pmatrix} , \quad (54)$$

following the same analysis as for the case without TMZ.

The characteristic polynomial $\ell^2 + \bar{a}(k^2)\ell + \bar{b}(k^2) = 0$ related to (54) has coefficients

$$\bar{a}(k^2) = \left(D_2(w^*, M^*) + \frac{1}{\zeta} \right) k^2 - f_w^* - \frac{g_c^*}{\zeta} , \quad (55)$$

$$\begin{aligned} \bar{b}(k^2) &= \frac{D_2(w^*, M^*)}{\zeta} k^4 - \left(\frac{D_2(w^*, M^*)g_c^*}{\zeta} + \frac{B\phi_2(w^*, M^*)g_n^*}{\zeta} + \frac{f_w^*}{\zeta} \right) k^2 \\ &\quad + \frac{f_w^*g_c^*}{\zeta} . \end{aligned} \quad (56)$$

For the steady state to be unstable we require, as before, that $\mathcal{R}e(\ell(k^2)) > 0$. Since $\bar{a}(k^2) > 0$ the instability can only occur if $\bar{b}(k^2) < 0$. Computing $\frac{d\bar{b}(k^2)}{dk^2} = 0$ from (56) we obtain

$$k_{\min}^2 = \frac{D_2(w^*, M^*)g_c^* + g_w^*B\phi_2(w^*, M^*) + f_w^*}{2D_2(w^*, M^*)} . \quad (57)$$

Hence from the condition $\bar{b}_{\min}(k_{\min}^2) < 0$ we get

$$D_2(w^*, M^*)g_c^* + g_w^*B\phi_2(w^*, M^*) + f_w^* > \sqrt{4D_2(w^*, M^*)f_w^*g_c^*} . \quad (58)$$

The spatially homogeneous steady state is $(w^*, c^*, M^*) = (u_{\max}, u_{\max}, M_s)$, where $M_s = |\Omega|^{-1} \int_{\Omega} M(\mathbf{x}, 0) d\mathbf{x}$. Therefore, from (58) we obtain a critical constant B^c so that for any $B > B^c$ we observe pattern formation. This critical constant is given by

$$B^c = \frac{2\sqrt{\tilde{r}_0}D_2(u_{\max}, M_s) + D_2(u_{\max}, M_s) + \tilde{r}_0}{u_{\max} \left(1 - \left(\frac{u_{\max}}{\bar{u}} \right)^{\gamma_{M_s}} \right)} , \quad (59)$$

where

$$D_2(u_{\max}, M_s) = 1 + (\gamma_{M_s} - 1) \left(\frac{u_{\max}}{\bar{u}} \right)^{\gamma_{M_s}} . \quad (60)$$

The corresponding critical wavemode is given by

$$k_c^2 = \frac{D_2(u_{\max}, M_s)g_c^* + g_w^*B^c\phi_2(u_{\max}, M_s) + f_w^*}{2D_2(u_{\max}, M_s)} = \frac{\sqrt{D_2(u_{\max}, M_s)(f_w^*g_c^*)}}{D_2(u_{\max}, M_s)} . \quad (61)$$

Finally, the unstable modes are $k^2 < k_c^2$, where from $\bar{b}(k^2) = 0$ we get

$$\begin{aligned} k_1^2 &= \frac{-\bar{m} - \sqrt{\bar{m}^2 - 4D_2(u_{\max}, M_s)(f_w^* g_c^*)}}{2D_2(u_{\max}, M_s)} < k^2 < k_2^2 \\ &= \frac{-\bar{m} + \sqrt{\bar{m}^2 - 4D_2(u_{\max}, M_s)(f_w^* g_c^*)}}{2D_2(u_{\max}, M_s)}, \end{aligned} \quad (62)$$

for $\bar{m} = -(D_2(u_{\max}, M_s)g_c^* + g_w^* B\phi_2(u_{\max}, M_s) + f_w^*)$.

B Description of the numerics

We denote $H^1(\Omega) = W^{1,2}(\Omega)$ the usual Sobolev space and the standard L^2 inner product is denoted by (\cdot, \cdot) . Let \mathcal{T}^h , $h > 0$, be a quasi-uniform mesh of the domain Ω consisting of N disjoint piecewise linear mesh elements K such that the discretized domain $\Omega_h = \bigcup_{K \in \mathcal{T}^h} K$. Let $h_K := \text{diam}(K)$ and $h = \max_K h_K$ and for $d = 2$, we choose linear triangular elements. In addition, we assume that the mesh is acute *i.e.* for $d = 2$, each angle of the triangles can not exceed $\frac{\pi}{2}$. We must stress that for $d = 2$ since the domain Ω is circular, a small error of approximation is committed using Ω_h . We consider the standard finite element space associated with \mathcal{T}^h

$$S^h := \{\chi \in C(\bar{\Omega}) : \chi|_K \in \mathbb{P}^1(K), \quad \forall K \in \mathcal{T}^h\} \subset H^1(\Omega), \quad (63)$$

where $\mathbb{P}^1(K)$ denotes the space of first-order polynomials on K . Let N_h be the total number of nodes of \mathcal{T}^h , J_h the set of nodes and $\{x_j\}_{j=1, \dots, N_h}$ their coordinates. We call $\{\chi_j\}_{j=1, \dots, N_h}$ the standard Lagrangian basis functions associated with the spatial mesh. We denote by $\pi^h : C(\bar{\Omega}) \rightarrow S^h$ the standard Lagrangian interpolation operator. We also need the lumped scalar product to define the problem

$$(\eta_1, \eta_2)^h = \int_{\Omega} \pi^h(\eta_1(x)\eta_2(x)) \, dx \equiv \sum_{x_j \in J_h} (1, \chi_j)\eta_1(x_j)\eta_2(x_j), \quad \eta_1, \eta_2 \in C(\bar{\Omega}).$$

We define the standard mass and stiffness finite element matrices as G and K , where

$$\begin{aligned} G_{ij} &= \int_{\Omega} \chi_i \chi_j \, dx, \quad \text{for } i, j = 1, \dots, N_h, \\ K_{ij} &= \int_{\Omega} \nabla \chi_i \nabla \chi_j \, dx, \quad \text{for } i, j = 1, \dots, N_h. \end{aligned}$$

In the following finite element approximation of the Keller-Segel problem, the mass matrix is lumped, *i.e.* the matrix becomes diagonal with each term being the row-sum of the corresponding row of the standard mass matrix,

$$G_{l,ii} := \sum_{j=1}^{N_h} G_{ij}, \quad \text{for } i = 1, \dots, N_h.$$

In order to describe how the chemotactic coefficient ϕ^{uwp} is computed, let us rewrite the discrete equation (24) into its matrix form

$$(G_L + \Delta t K_D) \underline{u}^{n+1} = G_L \underline{u}^n + \Delta t K_{\phi} \underline{c}^n + \Delta t G_L \underline{g}^n,$$

where \underline{u}^n and \underline{c}^n are the vectors of coefficients which are the unknowns of the problem and \underline{g}^n is a vector defined by

$$[\underline{g}^n]_i = \left(u_h^n \left(1 - \frac{u_h^n}{u_{\max}} \right) \right) (x_i), \quad \text{for } i = 1, \dots, N_h.$$

We define the finite element matrices associated with the diffusion K_D and the advection K_ϕ

$$K_{D,ij} = \int_{\Omega} D(u_h^n) \nabla \chi_i \nabla \chi_j \, dx \quad \text{for } i, j = 1, \dots, N_h, \quad (64)$$

$$K_{\phi,ij} = \int_{\Omega} \phi^{\text{upw}}(u_h^n(x_i), u_h^n(x_j)) \nabla \chi_i \nabla \chi_j \, dx \quad \text{for } i, j = 1, \dots, N_h. \quad (65)$$

In (64), the integral is computed using Gauss quadrature to deal with a potential choice of nonlinear functional for $D(u_h^n)$. The exactness of the quadrature is obtained using the adequate number of Gauss points since $D(u_h^n)$ is a polynomial of order $\gamma_M + 1$.

The chemotactic coefficient is computed using an upwing approach. For each element and depending on the direction of the gradient of the chemoattractant we have

$$\phi^{\text{upw}}(u_h^n(x_i), u_h^n(x_j)) = \begin{cases} u_h^n(x_j) \left(1 - \frac{u_h^n(x_i)}{u} \right), & \text{if } c_h^n(x_j) - c_h^n(x_i) < 0, \\ u_h^n(x_i) \left(1 - \frac{u_h^n(x_j)}{u} \right), & \text{otherwise.} \end{cases} \quad (66)$$

Therefore, the chemotactic coefficient is chosen as function of the sign of the difference of chemoattractant between nodes connected by an edge. The property of non-negativity of the cell density satisfied by our numerical scheme can be proved using similar arguments as in [10].

C One dimensional numerical results

Influence of the domain size The unstable wavenumbers are discrete values, $k = n\pi/L$, that satisfy the relation (47) from Section 4.2. The wavemode n determines the number of aggregates depending on the length of the domain. For $A = 7$ and the parameters specified at the beginning of this section we have $0.25 < n\pi/L < 0.4$. As shown in Figure 14, as we increase the length of the domain, the number of aggregates also increases. When the domain is large, as in Figure 14c, we observe that some aggregates are merging together while others are emerging, *i.e.*, they are formed from a zone of low cell density. This process is called coarsening [43] and is not observed in a small domain such as in Figure 14a.




Figure 14: Relationship between the wavenumber k and the length of the domain L for $A = 7$.

For the remaining simulations in this section we fix the length of the domain to $L = 40$, and all simulations are performed with carrying capacity $u_{\max} = 0.5$.

Comparison of the results with the stability analysis predictions Here, we study the influence of the chemosensitivity parameter A on the pattern dynamics and size of the aggregates, in the presence or absence of TMZ. In order to compare the solutions to the predictions of the stability analysis, the initial condition is a small perturbation around the homogeneous distribution $u_0 = 0.5$. Results of this section are obtained with a proliferation rate $r_0 = 0.1$. For such parameters, using the results of Section 4, the critical value of the chemosensitivity parameter without the treatment (in **P1**, when $M = 0$ and therefore $\gamma_M = 1$) is $A^c \approx 6.92$ and with the treatment uniformly distributed (for $\gamma_M = 5$) is $B^c \approx 3.9$.

In the first part of the experiment, *i.e.* without any treatment, we show in Figure 15 the formation of patterns at different times, for $A = 7$ (close to the instability threshold, Figure 15a), $A = 50$ (Figure 15b) and $A = 150$ (Figure 15c). We observe here the process of merging and emerging patterns through time.




Figure 15: Formation of the pattern without the treatment for (a) $A = 7$, (b) $A = 50$ and (c) $A = 150$ when $r_0 = 0.1$ and $u_{\max} = u_0 = 0.5$.

As predicted by the stability analysis, larger values of the chemotactic sensitivity A favor the emergence of smaller aggregates (compare Figures 15a and 15c for $A = 7$ and $A = 150$, respectively). This is due to the fact that for larger chemotactic sensitivity, cells are more attracted to zones of high concentration of chemoattractant. The creation of patterns instead of the expansion of a homogeneous cell distribution is due to an instability which results from a positive feedback loop between the production of the chemical by the cells on one hand, and their attraction to high density zones of this chemical on the other. The chemotactic sensitivity A must be large enough to trigger this instability, in order to compensate the competing effects of diffusion and of the logistic growth term, which on the contrary, tends to regulate the local cell density to the carrying capacity of the environment u_{\max} , and therefore induces cell death inside the aggregated patterns for which the density is above u_{\max} .

When the drug is introduced uniformly in the domain starting from a homogeneous distribution of cells (*i.e.* for $M = 1$, $\gamma_M = 5$) at time $t = 0$, we show in Figure 16 the formation of patterns at different times, for chemosensitivity $B = 5$ (close to the instability threshold, Figure 16a), $B = 30$ (Figure 16b) and $B = 150$ (Figure 16c). Note that in this case we also let cells to proliferate with rate $r_0 = 0.1$.

As one can see in Figure 16, we first observe again that increasing the chemosensitivity parameter B results in the formation of smaller cell aggregates (compare Figures 16a and 16b). Very close to the instability threshold (Figure 16a), the system converges quickly to one aggregate, while for larger values of B (Figure 16c) a large number of well-separated small aggregates arises. These clusters merge in time to form bigger clusters as for the case without the treatment. Moreover, comparing Figures. 15c and 16c, we clearly observe that varying the mechanical state of cells (*i.e.* passing from $\gamma_M = 1$ to $\gamma_M = 5$), leads to a change in the cell's aggregate size. When cells are more elastic, they tend to create smaller aggregates than when they behave as rigid spheres.

Acknowledgments

The authors would like to thank K. J. Painter and H. Gimpelrein for fruitful discussions and helpful suggestions. This research was funded by the Plan Cancer/ITMO HTE (tumor heterogeneity and ecosystem) program (MoGlimaging).

References

- [1] A. Agosti, S. Marchesi, G. Scita, and P. Ciarletta. Modelling cancer cell budding in-vitro as a self-organised, non-equilibrium growth process. *Journal of Theoretical Biology*, 492:110203, 2020.
- [2] J. Alfonso, K. Talkenberger, M. Seifert, B. Klink, A. Hawkins-Daarud, K. Swanson, H. Hatzikirou, and A. Deutsch. The biology and mathematical modelling of glioma invasion: a review. *Journal of the Royal Society Interface*, 14(136):20170490, 2017.
- [3] L. Almeida, F. Bubba, B. Perthame, and C. Pouchol. Energy and implicit discretization of the Fokker-Planck and Keller-Segel type equations. *Network and Heterogeneous Media*, 14(1):23–41, 2019.
- [4] W. Alt. Biased random walk models for chemotaxis and related diffusion approximations. *Journal of mathematical biology*, 9(2):147–177, 1980.




Figure 16: Formation of the pattern with treatment ($M = 1, \gamma_M = 5$), included at $t = 0$ when $u_0 = 0.5, r_0 = 0.1$ for (a) $B = 5$, (b) $B = 30$ and (c) $B = 150$.

- [5] L. Barazzuol, R. Jena, N. G. Burnet, J. C. Jaynes, M. J. Merchant, K. J. Kirkby, and N. F. Kirkby. In vitro evaluation of combined temozolomide and radiotherapy using x rays and high-linear energy transfer radiation for glioblastoma. *Radiation research*, 177(5):651–662, 2012.
- [6] E. Ben-Jacob, I. Cohen, and H. Levine. Cooperative self-organization of microorganisms. *Advances in Physics*, 49(4):395–554, 2000.
- [7] M. Brandao, T. Simon, G. Critchley, and G. Giamas. Astrocytes, the rising stars of the glioblastoma microenvironment. *Glia*, 67(5):779–790, 2019.
- [8] F. Bubba, T. Lorenzi, and F. R. Macfarlane. From a discrete model of chemotaxis with volume-filling to a generalized Patlak–Keller–Segel model. *Proceedings of the Royal Society A*, 476(2237):20190871, 2020.
- [9] F. Bubba, C. Pouchol, N. Ferrand, G. Vidal, L. Almeida, B. Perthame, and M. Sabbah. A chemotaxis-based explanation of spheroid formation in 3d cultures of breast cancer cells. *Journal of Theoretical Biology*, 479:73–80, 2019.
- [10] F. Bubba and A. Poulain. A positivity-preserving finite element scheme for the relaxed cahn-hilliard equation with single-well potential and degenerate mobility. *ArXiv*, page 20, 2019.
- [11] M. A. Chaplain and G. Lolas. Mathematical modelling of cancer invasion of tissue: dynamic heterogeneity. *Networks & Heterogeneous Media*, 1(3):399, 2006.
- [12] X. Chen, S. Wanggou, A. Botalia, M. Zhu, W. Dong, J. J. Fan, W. C. Yin, H.-K. Min, M. Hu, D. Draghici, et al. A feedforward mechanism mediated by mechanosensitive ion channel piezo1 and tissue mechanics promotes glioma aggression. *Neuron*, 100(4):799–815, 2018.
- [13] M. C. Colombo, C. Giverso, E. Faggiano, C. Boffano, F. Acerbi, and P. Ciarletta. Towards the personalized treatment of glioblastoma: integrating patient-specific clinical data in a continuous mechanical model. *PLoS One*, 10(7):e0132887, 2015.
- [14] J. Condeelis, R. H. Singer, and J. E. Segall. The great escape: when cancer cells hijack the genes for chemotaxis and motility. *Annual Review of Cell and Developmental Biology*, 21:695–718, 2005.
- [15] S. L. Di Jia, D. Li, H. Xue, D. Yang, and Y. Liu. Mining tcga database for genes of prognostic value in glioblastoma microenvironment. *Aging (Albany NY)*, 10(4):592, 2018.
- [16] Y. Dolak and C. Schmeiser. The Keller–Segel model with logistic sensitivity function and small diffusivity. *SIAM Journal on Applied Mathematics*, 66(1):286–308, 2005.
- [17] D. Dormann and C. J. Weijer. Chemotactic cell movement during dictyostelium development and gastrulation. *Current Opinion in Genetics & Development*, 16(4):367–373, 2006.
- [18] Y. A. Fouad and C. Aanei. Revisiting the hallmarks of cancer. *American Journal of Cancer Research*, 7(5):1016, 2017.
- [19] H. B. Frieboes, J. S. Lowengrub, S. Wise, X. Zheng, P. Macklin, E. L. Bearer, and V. Cristini. Computer simulation of glioma growth and morphology. *Neuroimage*, 37:S59–S70, 2007.
- [20] P. Friedl. Prespecification and plasticity: shifting mechanisms of cell migration. *Current opinion in cell biology*, 16(1):14–23, 2004.

- [21] Y. Han, Z. Li, J. Tao, and M. Ma. Pattern formation for a volume-filling chemotaxis model with logistic growth. *Journal of Mathematical Analysis and Applications*, 448(2):885–907, 2017.
- [22] D. Hanahan and R. A. Weinberg. Hallmarks of cancer: the next generation. *Cell*, 144(5):646–674, 2011.
- [23] T. Hillen and K. J. Painter. A user’s guide to pde models for chemotaxis. *Journal of Mathematical Biology*, 58(1-2):183, 2009.
- [24] D. Horstmann. The Keller-Segel model in chemotaxis and its consequences. *I, Jahresber. Deutsch. Math.-Verein*, pages 103–165, 2004.
- [25] A. Hurwitz et al. On the conditions under which an equation has only roots with negative real parts. *Selected Papers on Mathematical Trends in Control Theory*, 65:273–284, 1964.
- [26] M. Ibrahim and M. Saad. On the efficacy of a control volume finite element method for the capture of patterns for a volume-filling chemotaxis model. *Computers & Mathematics with Applications*, 68(9):1032–1051, 2014.
- [27] E. F. Keller and L. A. Segel. Model for chemotaxis. *Journal of Theoretical Biology*, 30(2):225–234, 1971.
- [28] J. S. Kennedy and D. Marsh. Pheromone-regulated anemotaxis in flying moths. *Science*, 184(4140):999–1001, 1974.
- [29] Y. Kim and S. Kumar. Cd44-mediated adhesion to hyaluronic acid contributes to mechanosensing and invasive motility. *Molecular Cancer Research*, 12(10):1416–1429, 2014.
- [30] S. Y. Lee. Temozolomide resistance in glioblastoma multiforme. *Genes & Diseases*, 3(3):198–210, 2016.
- [31] D. N. Louis, A. Perry, G. Reifenberger, A. Von Deimling, D. Figarella-Branger, W. K. Cavenee, H. Ohgaki, O. D. Wiestler, P. Kleihues, and D. W. Ellison. The 2016 world health organization classification of tumors of the central nervous system: a summary. *Acta Neuropathologica*, 131(6):803–820, 2016.
- [32] M. Ma, C. Ou, and Z.-A. Wang. Stationary solutions of a volume-filling chemotaxis model with logistic growth and their stability. *SIAM Journal on Applied Mathematics*, 72(3):740–766, 2012.
- [33] G. D. Maurer, D. P. Brucker, and J. P. Steinbach. Loss of cell-matrix contact increases hypoxia-inducible factor-dependent transcriptional activity in glioma cells. *Biochemical and biophysical research communications*, 515(1):77–84, 2019.
- [34] H. Meinhardt. Models for positional signalling with application to the dorsoventral patterning of insects and segregation into different cell types. *Development*, 107(Supplement):169–180, 1989.
- [35] J. Murray. *Mathematical biology II: spatial models and biomedical applications*. Springer New York, 2001.
- [36] B. Nagorcka and J. Mooney. From stripes to spots: prepatterns which can be produced in the skin by a reaction-diffusion system. *Mathematical Medicine and Biology: A Journal of the IMA*, 9(4):249–267, 1992.

- [37] K. Oizel, C. Chauvin, L. Oliver, C. Gratas, F. Geraldo, U. Jarry, E. Scotet, M. Rabe, M.-C. Alves-Guerra, R. Teusan, et al. Efficient mitochondrial glutamine targeting prevails over glioblastoma metabolic plasticity. *Clinical Cancer Research*, 23(20):6292–6304, 2017.
- [38] L. Oliver, L. Lalier, C. Salaud, D. Heymann, P. F. Carton, and F. Vallette. Drug resistance in glioblastoma: Are persisters the key to therapy? *Under review*, 2020.
- [39] L. Oliver, A. Álvarez Arenas, C. Salaud, et al. A simple 3d cell culture method for studying the interactions between primary glioblastoma cells and tumor-activated stromal cells. *Under review*, 2020.
- [40] H. G. Othmer and T. Hillen. The diffusion limit of transport equations ii: Chemotaxis equations. *SIAM Journal on Applied Mathematics*, 62(4):1222–1250, 2002.
- [41] Q. Ouyang and H. L. Swinney. Transition from a uniform state to hexagonal and striped turing patterns. *Nature*, 352(6336):610–612, 1991.
- [42] K. Painter. Modelling cell migration strategies in the extracellular matrix. *Journal of Mathematical Biology*, 58(4-5):511, 2009.
- [43] K. J. Painter and T. Hillen. Volume-filling and quorum-sensing in models for chemosensitive movement. *Canadian Applied Mathematics Quarterly*, 10(4):501–543, 2002.
- [44] W. Roos, L. Batista, S. Naumann, W. Wick, M. Weller, C. Menck, and B. Kaina. Apoptosis in malignant glioma cells triggered by the temozolomide-induced dna lesion o 6-methylguanine. *Oncogene*, 26(2):186–197, 2007.
- [45] W. Sarfaraz and A. Madzvamuse. Domain-dependent stability analysis of a reaction–diffusion model on compact circular geometries. *International Journal of Bifurcation and Chaos*, 28(08):1830024, 2018.
- [46] A. Stevens. The derivation of chemotaxis equations as limit dynamics of moderately interacting stochastic many-particle systems. *SIAM Journal on Applied Mathematics*, 61(1):183–212, 2000.
- [47] R. Stupp, M. E. Hegi, W. P. Mason, M. J. Van Den Bent, M. J. Taphoorn, R. C. Janzer, S. K. Ludwin, A. Allgeier, B. Fisher, K. Belanger, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase iii study: 5-year analysis of the eortc-ncic trial. *The Lancet Oncology*, 10(5):459–466, 2009.
- [48] R. Stupp, W. P. Mason, M. J. Van Den Bent, M. Weller, B. Fisher, M. J. Taphoorn, K. Belanger, A. A. Brandes, C. Marosi, U. Bogdahn, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *New England Journal of Medicine*, 352(10):987–996, 2005.
- [49] T. Suzuki. *Chemotaxis, reaction, network*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2018. Mathematics for self-organization.
- [50] I. Talhaoui, B. T. Matkarimov, T. Tchenio, D. O. Zharkov, and M. K. Saparbaev. Aberrant base excision repair pathway of oxidatively damaged dna: implications for degenerative diseases. *Free Radical Biology and Medicine*, 107:266–277, 2017.
- [51] L. Tang, A. L. Van De Ven, D. Guo, V. Andasari, V. Cristini, K. C. Li, and X. Zhou. Computational modeling of 3d tumor growth and angiogenesis for chemotherapy evaluation. *PloS one*, 9(1):e83962, 2014.

- [52] A. Toma, L. R. C. Castillo, T. A Schuetz, S. Becker, A. Mang, A. Régnier-Vigouroux, and T. M Buzug. A validated mathematical model of tumour-immune interactions for glioblastoma. *Current Medical Imaging*, 9(2):145–153, 2013.
- [53] J. van Rijn, J. J. Heimans, J. van den Berg, P. van der Valk, and B. J. Slotman. Survival of human glioma cells treated with various combination of temozolomide and x-rays. *International Journal of Radiation Oncology* Biology* Physics*, 47(3):779–784, 2000.
- [54] Z. Wang. On chemotaxis models with cell population interactions. *Mathematical Modelling of Natural Phenomena*, 5(3):173–190, 2010.
- [55] Z. Wang and T. Hillen. Classical solutions and pattern formation for a volume filling chemotaxis model. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 17(3):037108, 2007.
- [56] S. Ward. Chemotaxis by the nematode caenorhabditis elegans: identification of attractants and analysis of the response by use of mutants. *Proceedings of the National Academy of Sciences*, 70(3):817–821, 1973.
- [57] D. Wrzosek. Volume filling effect in modelling chemotaxis. *Mathematical Modelling of Natural Phenomena*, 5(1):123–147, 2010.
- [58] R. Würth, A. Bajetto, J. K. Harrison, F. Barbieri, and T. Florio. Cxcl12 modulation of cxcr4 and cxcr7 activity in human glioblastoma stem-like cells and regulation of the tumor microenvironment. *Frontiers in Cellular Neuroscience*, 8:144, 2014.
- [59] H. Yamaguchi, J. Wyckoff, and J. Condeelis. Cell migration in tumors. *Current Opinion in Cell Biology*, 17(5):559–564, 2005.