



HAL
open science

Combining planar and central chirality in ferrocene thiophosphine-sulfoxides

Raluca Malacea, Jean-Claude Daran, Rinaldo Poli, Eric Manoury

► **To cite this version:**

Raluca Malacea, Jean-Claude Daran, Rinaldo Poli, Eric Manoury. Combining planar and central chirality in ferrocene thiophosphine-sulfoxides. *Tetrahedron: Asymmetry*, 2013, 24 (9-10), pp.612-620. 10.1016/j.tetasy.2013.04.002 . hal-02908226

HAL Id: hal-02908226

<https://hal.science/hal-02908226>

Submitted on 2 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Combining planar and central chirality in ferrocene thiophosphine-sulfoxides

Raluca Malacea,^{a,b} Jean-Claude Daran,^{a,b} Rinaldo Poli,^{a,b,c} Eric Manoury^{a,b} *

^a CNRS, LCC (Laboratoire de Chimie de Coordination), 205 route de Narbonne, BP 44099, F-31077

Toulouse Cedex 4, France; Fax: (+) 33-561553003; E-mail: manoury@lcc-toulouse.fr


^b Université de Toulouse, UPS, INPT, F-31077 Toulouse Cedex 4, France

^c Institut Universitaire de France, 103, bd Saint-Michel, 75005 Paris, France

Abstract

New chiral ferrocenylthiophosphine-sulfoxide and phosphine-sulfoxide derivatives possessing planar chirality for the ferrocene moiety and central chirality at the sulfur atom have been synthesized. These ligands could be obtained as pure diastereoisomers in both racemic and enantiomerically pure forms. Complete characterization by XRD analysis has allowed the absolute configuration assignment in each case. Preliminary coordination studies of the phosphine-sulfoxide ligand on platinum are also reported. They show chelating complexation by the phosphorous and sulfur atoms.

Image for the graphical abstracts:



R = Ph, Et, *t*Bu

1. Introduction

Various chiral bidentate ligands were developed over the past decades, possessing the chiral information on the carbon backbone or on the donor atom. Ligands with various combinations of donor atoms (in particular, P–P, P–N, or N–N ligands) have been extensively studied and a few P-chirogenic compounds such as QuinoxP¹ or dipamp² are very effective in a variety of asymmetric catalytic processes. Recently, very efficient systems for asymmetric catalysis using P–S ligands with various backbones, have been reported.^{3,4} Furthermore, several reports have described the preparation and the coordination chemistry of chiral phosphine sulfoxide ligands possessing the chirality on the sulfur atom, for example α -phosphinosulfoxides (PCSO⁵ and PNSO⁶) or β -phosphinosulfoxides⁷ (Figure 1). Riera and Verdager⁸ described ligands with an extra chiral center in the central carbon atom between P and SO and also camphor derivatives combining a chiral carbon backbone with sulfur chirality.⁹ Kagan's methodology based on diastereoselective *ortho*-lithiation of ferrocenesulfoxide derivatives was used in order to synthesize phosphine-sulfoxide ligands with additional planar chirality.¹⁰ We have recently developed syntheses of new chiral ferrocenyl γ -P,S ligands in both racemic and enantiomerically pure form (R_{Fc} or S_{Fc} configuration), and reported on their coordination chemistry and applications to asymmetric catalysis, namely in asymmetric allylic substitution, alkene methoxycarbonylation and ketone hydrogenation.¹¹ Here we present the synthesis of new γ -ferrocenyl phosphine sulfoxide ligands (Figure 1) and preliminary studies on their coordination chemistry with platinum.





Figure 1 α , β and γ phosphine sulfoxides

2. Results and Discussion

Starting from the chiral phosphine-thioether ligands already synthesized in our group, we have developed a new class of ferrocenyl phosphine-sulfoxides ligands based on the selective oxidation of the thioether functionality to sulfoxide. Thus, a central chirality on the sulfur atom (R_S or S_S) is added to the planar chirality of ferrocene (R_{Fc} or S_{Fc}). The two diastereoisomers that can *a priori* be formed must therefore be subsequently separated.


A first potential issue in the synthetic pathway to the thiophosphine-sulfoxides **2** from thiophosphine-thioethers **1** (see scheme 1) is the possible over-oxidation to the corresponding sulfones as well as oxidation of the iron atom to ferrocenium or conversion of

thiophosphine to phosphine oxide, as already observed in the oxidation of related (2-diphenylthiophosphinoferrocenyl)methanol to the corresponding aldehyde.¹² Therefore, the mild method described by Liang Xu¹³ using an aqueous solution of H₂O₂ in the presence of phenol was selected. Under these conditions, the racemic compounds **1a-c** (R = Ph, Et, *t*Bu) were transformed into the corresponding thiophosphine-sulfoxides **2a-c** with total conversion and chemoselectivity (no sulfone or other by-products were observed). For each thioether, two diastereoisomers (each one as a pair of enantiomers) were obtained in a 1:1 ratio. These reactions are fast (2-3 minutes) and strongly exothermic. For safety reasons, we therefore limited the thioether amount (typically 250 mg) transformed in each run. Each diastereomer could be isolated in good yields after separation by column chromatography (Scheme 1).



Scheme 1

After diastereomer separation, the sulfur atom configurations were established by single crystal X-ray diffraction analyses. Single crystals of **2a-Dia1**, **2a-Dia2**, **2b-Dia1**, **2c-Dia1** and **2c-Dia2** were obtained by slow diffusion of pentane vapors into a dichloromethane solution. These products were obtained from the racemic thiophosphine-thioethers **1a-c** (*R/S*_{Fc}) and crystallize as racemates, except for **2c-Dia2** which affords a twin by inversion, therefore also containing both enantiomers (70:30) in the crystal.






2a-Dia1 (R_{Fc} , S_S)/(*)**2a-Dia2** (S_{Fc} , S_S)/(*)**2b-Dia1** (R_{Fc} , R_S)/(*)**2c-Dia1** (S_{Fc} , R_S)/(*)**2c-Dia2** (S_{Fc} , S_S)/(*)

Figure 2. ORTEP views of **2a-Dia1**, **2a-Dia2**, **2b-Dia1**, **2c-Dia1** and **2c-Dia2** (* the other enantiomer is present in the crystal structure but not represented in the picture) with the atom labeling scheme. Ellipsoids are drawn at the 50% probability level. H atoms have been omitted for the sake of clarity.

To the best of our knowledge, these compounds are the first structurally characterized ferrocenylthiophosphine-sulfoxides. Selected geometrical parameters are reported in Table 1. The five compounds have similar structures with the protecting $S(1)$ *endo* with respect to the Cp ring whereas the sulfido $S(2)$ is *exo* (Table 1). The distances and angles within the sulfoxidemoiety are identical within experimental error. The two Cp rings are either nearly

eclipsed for **2a-Dia2**, **2c-Dia1** and **2c-Dia2B** and in between eclipsed-staggered for **2a-Dia1, 2b-Dia1** and **2c-Dia2A** (τ values in Table 1). In compound **2c-Dia2**, there are two molecules within the asymmetric unit. These molecules have similar configuration as indicated by the molecular fitting shown in Figure 3. The only marked difference is the twist angles observed between the two Cp rings, $12.7(3)^\circ$ for molecule A and $4.5(3)^\circ$ for B.





Figure 3. Molecular fitting of the two molecules **2c-Dia2A** and **2c-Dia2B**.

Table 1. Selected geometrical parameters for the structures of compounds **2**, **3** and **4**.

	2a-Dia1	2a-Dia2	2b-Dia1	2c-Dia1	2c-Dia2A	2c-Dia2B	3a-Dia2^a	4a^b
Distances (Å)								
Fe(1) - Cp(1)	1.6440(11)	1.642(4)	1.635(1)	1.640(1)	1.642(2)	1.644(2)	1.639(1)	1.646(2)
Fe(1) - Cp(2)	1.6582(14)	1.654(6)	1.652(2)	1.655(1)	1.655(2)	1.659(2)	1.653(1)	1.661(3)
C(1) - P(1)	1.788(2)	1.788(8)	1.802(3)	1.792(2)	1.788(4)	1.798(4)	1.816(2)	1.796(5)
P(1) - S(1)	1.9611(8)	1.958(3)	1.957(1)	1.9584(8)	1.958(2)	1.956(2)		
C(2) - C(21)	1.491(3)	1.502(11)	1.502(4)	1.500(3)	1.496(5)	1.495(5)	1.487(3)	1.481(8)
C(21) - S(2)	1.817(2)	1.828(8)	1.825(3)	1.811(3)	1.820(4)	1.817(4)	1.823(2)	1.810(5)
S(2) - O(2)	1.499(2)	1.485(6)	1.508(2)	1.494(2)	1.495(4)	1.480(4)	1.470(2)	1.459(3)
S(2) - C(22)	1.782(3)	1.787(9)	1.803(3)	1.841(3)	1.826(5)	1.840(4)	1.790(2)	1.782(6)
Angles (°)								
Cp(1) - Fe(1) - Cp(2)	178.53(7)	177.7(2)	177.29(8)	177.42(7)	178.5(1)	178.2(1)	177.90(5)	176.21(13)
C(2) - C(21) - S(2)	115.90(17)	111.0(5)	113.1(2)	110.5(2)	108.1(3)	107.5(3)	110.3(1)	116.2(3)
C(21) - S(2) - O(2)	103.28(13)	105.5(4)	105.5(1)	105.0(1)	103.8(2)	105.8(2)	105.6(1)	108.1(2)
C(21) - S(2) - C(22)	100.84(12)	94.4(4)	97.8(2)	101.0(1)	100.4(2)	100.6(2)	95.65(9)	97.5(2)
O(2) - S(2) - C(22)	107.50(13)	106.8(4)	106.9(2)	108.0(1)	108.0(2)	108.0(2)	107.5(1)	109.1(2)
Distances from plane (Å)								
S(1)---Cp(1)	-0.859(1)	-1.06(2)	-0.921(6)	-1.063(5)	-0.992(8)	-1.154(9)		
S(2)---Cp(1)	1.500(1)	1.63(1)	1.647(5)	0.457(5)	1.751(7)	1.757(7)	1.619(4)	0.970(9)
O(2)---Cp(1)	1.228(2)	2.49(1)	1.619(8)	1.945(5)	1.654(9)	1.895(8)	2.236(4)	2.186(9)
Twist angle for the Cp rings (°)								
τ°	12.5(2)	4.0(9)	13.3(2)	1.6(3)	12.7(3)	4.5(3)	4.7(2)	14.5(3)


^a: **3a dia2**: (*S*_{Fc}, *S*_S)/(*R*_{Fc}, *R*_S)-**3a**^b: **4a**: (*R*_{Fc}, *R*_S)/(*S*_{Fc}, *S*_S)-**4a**

As stated above, no diastereoselectivity was observed under these oxidation conditions. Other achiral oxidants such as meta-chloroperbenzoic acid¹⁴, *t*BuOOH/MoO₂(acac)₂,¹⁵ and also chiral ones such as *t*BuOOH/diethyltartrate/Ti(O*i*Pr)₄¹⁶ and *t*BuOOH/diethyltartrate/MoO₂(acac)₂ were then employed. Unfortunately, all of them gave lower conversions and no significant diastereoselectivity improvement relative to Xu's method. Therefore, our investigations were continued using the H₂O₂/phenol system. The next step was the oxidation of enantiomerically pure thiophosphine-thioethers (*R*_{Fc}) **1a-c**, **11a**, **12** which gave the corresponding thiophosphine-sulfoxides **2a-c** as a mixture of two diastereomers in enantiomerically pure form (Scheme 2). The results obtained working with optically pure thioethers are quite similar to those obtained using the compounds in racemic form. Thus, we have access to the desired ferrocenylthiophosphine-sulfoxides in diastereoisomerically and enantiomerically pure form.



Scheme 2

As the thioether oxidations were performed on ferrocene derivatives with thiophosphine group in order to prevent phosphorus oxidation, the next step was the phosphorus deprotection in order to liberate the free phosphine group and generate ferrocenylphosphine sulfoxides that can be used as chiral bidentate P-S ligands. This reaction was carried out using tris(dimethylamino)phosphine in refluxing toluene for 3 hours (Scheme 3).^{11a} Longer reaction time gave partial deoxygenation of the sulfoxide group. Phosphine-sulfoxide compounds **3a-c** being air sensitive, their purification was accomplished by column chromatography under argon with yields of isolated products from 65 to 93%. Single crystals of (*S*_{Fc}, *S*_S)/(*R*_{Fc}, *R*_S)-**3a** were obtained by slow diffusion of pentane vapors into a dichloromethane solution, and the XRD structure is presented in Figure 4. The conformation of this unprotected ligand does not show significant differences relative to the protected one (*S*_{Fc}, *S*_S)/(*R*_{Fc}, *R*_S)-**3a** (Table 1).



Scheme 3




Figure 4. ORTEP view of the phosphine sulfoxide **3a-Dia2** (S_{Fc} , S_S) with the atom labeling scheme. Ellipsoids are drawn at the 50% probability level. H atoms have been omitted for the sake of clarity.

Given that the P,S,O ligands could adopt different coordination modes (κ^2 -PS, κ^2 -PO, or κ^1 -P),¹⁷ we carried out preliminary coordination chemistry investigations with platinum, by reacting the phosphine sulfoxide ligand **3a** (as the racemic diastereoisomeric mixture, Dia1/Dia2 = 1/1) with the platinum precursor $PtCl_2(CH_3CN)_2$. The $^{31}P\{^1H\}$ NMR analysis of the final mixture shows the presence of two signals corresponding to two similar platinum complexes ($\delta_1 = 0.72$ ppm, $J_{P-Pt} = 3338$ Hz and $\delta_2 = 0.47$ ppm, $J_{P-Pt} = 3444$ Hz) in a 1:1 ratio. The $^1J_{PP}$ value is characteristic of square planar complexes with *cis* chlorine atoms.¹⁸ Crystals of one complex were obtained by diffusion of pentane into a dichloromethane solution, and thus the coordination mode of the sulfoxide group as κ^2 -PS was determined from the XRD structure (See figure 5).





Figure 5. ORTEP view of platinum complex (R_{Fc}, R_S)-**4a** with the atom labeling scheme. The other enantiomer (not shown) is also present in the crystal. Ellipsoids are drawn at the 50% probability level. H atoms have been omitted for the sake of clarity. Selected bond lengths (Å): Pt-P: 2.2350(12); Pt-S: 2.2291(12); Pt-Cl1(trans S): 2.2973(12); Pt-Cl2(trans P): 2.3312(13); S-O: 1.459.

The Pt atom is surrounded by the S2, P1, Cl1 and Cl2 donor atoms in square planar coordination geometry. It deviates from the least squares plane defined by these four atoms by only $-0.0181(7)$ Å. The C1, C2, C21, S2, Pt1, P1 metallacycle displays a boat conformation as indicated by the puckering parameters $\varphi = 302.6(3)^\circ$ and $\theta = 87.9(3)^\circ$.¹⁹ The S2 and O2 atoms are *endo* with respect to the Cp1 ring by $-0.970(9)$ Å and $-2.186(9)$ Å respectively. The coordination has induced a slight twist of the two Cp rings ($\tau = 14.5(4)^\circ$) when compared to the uncoordinated ligand which was roughly eclipsed with a twist angle of $4.7(2)^\circ$.

The crystallized compound corresponds to the racemic diastereomer with the (R_{Fc}, R_S)/(S_{Fc}, S_S) configuration. Given the similarity of the NMR signals, we attribute to the second compound the structure **4a'** with the (S_{Fc}, R_S)/(R_{Fc}, S_S) configuration (Scheme 4).



Scheme 4

The presence of two complexes in a 1:1 ratio in the product suggests that the two diastereomeric ferrocenyl phosphine-sulfoxide ligands do not have significantly different reactivity with the platinum precursor. This might be surprising given that in the case of the corresponding ferrocenyl phosphine thioethers, we observed a total discrimination between the two electron pairs for coordination with platinum, palladium²⁰ or iridium.²¹ Our attempts to obtain complex **4a** by oxidation of the sulfur atom in the coordination sphere of the corresponding phosphine-thioether platinum precursor (Ph₂P-Fc-CH₂SPh)PtCl₂²⁰ were unsuccessful (conditions : mCPBA, THF, reflux, 20h). The oxidation reaction did not occur and the starting material was recovered quantitatively.

3. Conclusion

In conclusion, we have described the synthesis of new ferrocenyl derivatives thiophosphine-sulfoxide **2a-c** and phosphine-sulfoxide **3a-c** having aromatic or aliphatic substituents on the sulfur atom. These ligands possess planar chirality for the ferrocene moiety and central chirality at the sulfur atom and could be obtained as pure diastereoisomers in both racemic and enantiomerically pure form. Most of these compounds were fully characterized by XRD analysis allowing the absolute configuration assignment for each stereoisomer. Preliminary coordination studies of these ligands were carried out with platinum, demonstrating bidentate κ^2 -PS complexation in this case. Tests of enantiomerically pure ligands in various catalytic asymmetric reactions are in progress in our laboratories and will be presented in due course.

4. Experimental

4.1 General

All reactions were carried out under argon using standard Schlenk line techniques. Solvents were carefully dried by conventional methods and distilled under argon before use. The NMR spectra were recorded on a Bruker DRX-500 Avance instrument with degassed solvents. Elemental analyses were recorded on Perkin Elmer 2400 II. The mass spectra were recorded on a Nermag R-10-10H spectrometer.

4.2 2-thiodiphenylphosphino-(phenyl-sulfoxymethyl)ferrocene, **2a**.

In a Schlenk tube under argon were introduced 240 mg (0.45 mmol) of racemic thioether (*R/S*_{Fc})-**1a** and 500 mg (5.4 mmol) of phenol. The mixture was heated to 35°C, then 0.1 ml of an aqueous solution of H₂O₂ (30%) were slowly added to the reaction medium. The reaction is highly exothermic and within minutes the solution turned black. After 5 minutes, 10 ml of ethyl acetate, then 10 ml of a saturated solution of Na₂SO₃, and finally 20 ml of an aqueous solution of NaOH (10%) were added. The phases were separated and the aqueous phase was

extracted with three portions (10ml) of ethyl acetate. The combined organic layers were washed with water, dried over MgSO₄ and then the solvent was removed by evaporation under reduced pressure. Compound **2a** was recovered as an orange solid (yield: 226 mg, 92%). The two diastereomers were separated by chromatography on silica using a mixture of ether and dichloromethane (1/1) as eluent to give 100 mg of **2a-Dia1**, (**R**_{Fe},**S**_S)/(**S**_{Fe},**R**_S)-**2a** (yield: 41%) and 108 mg of **2a-Dia2**, (**R**_{Fe},**R**_S)/(**S**_{Fe},**S**_S)-**2a** (yield: 44%). MS (DCI, NH₃): m/z = 541 (M + 1, 20 %); 558 (M + NH₄, 100%).

(R_{Fe},S_S)/(S_{Fe},R_S)-2a: ¹H NMR (δ, 500 MHz, CDCl₃): 7.86 (2H, m, Ph); 7.65-7.40 (13H, m, Ph); 4.48 (1H, broad s, substCp); 4.44 (5H, s, Cp); 4.35 (1H, broad s, substCp); 4.32 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 4.30 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 3.87 (1H, broad s, substCp). ¹³C{¹H} NMR (δ, 125.8 MHz, CDCl₃): 144.5 (s, quatSPh); 134.8 (d, J_{CP} = 87 Hz, quat PPh₂); 133.1 (d, J_{CP} = 86 Hz, quat PPh₂); 132.5 (d, J_{CP} = 11 Hz, PPh₂); 132.3 (d, J_{CP} = 11 Hz, PPh₂); 131.96 (s, PPh₂); 131.9 (s, PPh₂); 131.3 (s, SPh); 129.4 (s, SPh); 128.9 (d, J_{CP} = 13 Hz, PPh₂); 128.6 (d, J_{CP} = 13 Hz, PPh₂); 124.8 (s, SPh); 83.1 (d, J_{CP} = 12 Hz, quatCp); 75.8 (d, J_{CP} = 95 Hz, quatCp); 75.4 (d, J_{CP} = 9 Hz, substCp); 74.6 (d, J_{CP} = 12 Hz, substCp); 71.6 (s, Cp); 70.4 (d, J_{CP} = 10 Hz, substCp); 58.0 (s, CH₂Cp). ³¹P{¹H} NMR (δ, 202.5 MHz, CDCl₃): 43.5. Elemental analysis: C₂₉H₂₅FePS₂O; %C (calcd 64.44, found 62.84), % H (calcd 4.63, found 5.03).

(R_{Fe},R_S)/(S_{Fe},S_S)-2a: ¹H NMR (δ, 500 MHz, CDCl₃): 7.89 (2H, m, Ph); 7.87 (2H, m, Ph); 7.6-7.40 (11H, m, Ph); 5.34 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 4.85 (1H, m, substCp); 4.49 (1H, m, substCp); 4.30 (5H, s, Cp); 3.87 (1H, m, substCp); 3.61 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp). ¹³C{¹H} NMR (δ, 125.8 MHz, CDCl₃): 134.9 (d, J_{CP} = 87 Hz, quat PPh₂); 133.2 (d, J_{CP} = 87 Hz, quat PPh₂); 132.7 (d, J_{CP} = 11 Hz, PPh₂); 132.4 (d, J_{CP} = 11 Hz, PPh₂); 132.1 (d, J_{CP} = 3 Hz, PPh₂); 132.0 (d, J_{CP} = 3 Hz, PPh₂); 130.9 (s, SPh); 129.5 (s, SPh); 129.4 (s, quatSPh); 128.8 (d, J_{CP} = 13 Hz, PPh₂); 128.6 (d, J_{CP} = 13 Hz, PPh₂); 123.9 (s, SPh); 83.0 (d, J_{CP} = 12 Hz, quatCp); 76.2 (d, J_{CP} = 9 Hz, substCp); 75.8 (d, J_{CP} = 12 Hz, substCp); 74.8 (d, J_{CP} = 95 Hz, quatCp); 71.4 (s, Cp); 71.1 (d, J_{CP} = 10 Hz, substCp); 61.1 (s, CH₂Cp). ³¹P{¹H} NMR (δ, 202.5 MHz, CDCl₃): 43.9; Elemental analysis: C₂₉H₂₅FePS₂O; %C (calcd 64.44, found 64.12), % H (calcd 4.63, found 4.32).

4.3 2-thiodiphenylphosphino-(ethyl-sulfoxymethyl)-ferrocene **2b**

Following the same procedure described above for the synthesis of **2a**, the reaction was realized starting from 250 mg (0.52 mmol) of racemic thioether (**R/S**)_{Fe}-**1b**, 500 mg of phenol

and 0.10 ml of an aqueous solution of H₂O₂ (30%). **2b** was recovered as an orange solid (yield: 241 mg, 93%). The two diastereomers were separated by chromatography on silica using a mixture of ether and dichloromethane (1/1) as eluent to give 114 mg of **2b-Dia1**, (**R_{FC},R_S**)/(**S_{FC},S_S**)-**2b** (yield: 44%) and 109 mg of **2b-Dia2**, (**R_{FC},S_S**)/(**S_{FC},R_S**)-**2b** (yield: 42%). MS (DCI, NH₃): m/z = 493 (M + 1, 45 %); 510 (M + NH₄, 100%).

(R_{FC},R_S)/(S_{FC},S_S**)-**2b****: ¹H NMR (δ, 500 MHz, CDCl₃): 7.85-7.81 (2H, m, PPh₂); 7.66-7.47 (6H, m, PPh₂); 7.43 (2H, broad s, PPh₂); 4.90 (1H, s, subst Cp); 4.47 (1H, s, subst Cp); 4.36 (5H, s, Cp); 4.45-4.28 (2H, m, CH₂Cp); 3.87 (1H, s, subst Cp); 2.35 (2H, broad s, CH₂CH₃); 1.1 (3H, broad s, CH₂CH₃). ¹³C{¹H} NMR (δ, 125.8 MHz, CDCl₃): 133.9 (d, J_{CP} = 87 Hz, quat PPh₂); 132.1 (d, J_{CP} = 86 Hz, quat PPh₂); 132.0 (d, J_{CP} = 11 Hz, PPh₂); 131.9 (s, PPh₂); 131.8 (broad s, PPh₂); 128.7 (d, J_{CP} = 12 Hz, PPh₂); 128.5 (d, J_{CP} = 12 Hz, PPh₂); 82.0 (d, J_{CP} = 12 Hz, quat Cp); 74.8 (s, subst Cp); 74.7 (d, J_{CP} = 12 Hz, subst Cp); 74.0 (d, J_{CP} = 96 Hz, quat Cp); 71.3 (s, Cp); 70.8 (d, J_{CP} = 10 Hz, subst Cp); 50.6 (s, CH₂Cp); 44.7 (s, CH₂CH₃); 7.6 (s, CH₂CH₃). ³¹P{¹H} NMR (δ, 202.5 MHz, CDCl₃): 41.06. Elemental analysis: C₂₅H₂₅FePS₂O; % C (calcd 60.98, found 61.08), % H (calcd 5.08, found 5.17).

(R_{FC},S_S)/(S_{FC},R_S**)-**2b****: ¹H NMR (δ, 500 MHz, CDCl₃): 7.83-7.79 (2H, m, PPh₂); 7.59-7.48 (6H, m, PPh₂); 7.41 (2H, broad s, PPh₂); 5.26 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 4.81 (1H, s, subst Cp); 4.47 (1H, s, subst Cp); 4.32 (5H, s, Cp); 3.82 (1H, s, subst Cp); 3.53 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 2.65 (2H, q, J_{HH} = 7 Hz, CH₂CH₃); 1.5 (3H, broad s, CH₂CH₃). ¹³C{¹H} NMR (δ, 125.8 MHz, CDCl₃): 134.1 (d, J_{CP} = 88 Hz, quat PPh₂); 132.3 (d, J_{CP} = 88 Hz, quat PPh₂); 132.0 (d, J_{CP} = 11 Hz, PPh₂); 131.9 (broad s, PPh₂); 128.6 (d, J_{CP} = 12 Hz, PPh₂); 128.5 (d, J_{CP} = 13 Hz, PPh₂); 82.5 (d, J_{CP} = 10 Hz, quat Cp); 75.8 (d, J_{CP} = 9 Hz, subst Cp); 75.5 (d, J_{CP} = 12 Hz, subst Cp); 73.1 (d, J_{CP} = 95 Hz, quat Cp); 71.2 (s, Cp); 71.1 (s, subst Cp); 53.4 (s, CH₂Cp); 46.7 (s, CH₂CH₃); 7.1 (s, CH₂CH₃). ³¹P{¹H} NMR (δ, 202.5 MHz, CDCl₃): 41.6. Elemental analysis: C₂₅H₂₅FePS₂O; % C (calcd 60.98, found 60.98), % H (calcd 5.08, found 4.83).

4.4 2-thiodiphenylphosphino-(*tert*-butyl-sulfoxymethyl)-ferrocene**2c**

Following the same procedure described above for the synthesis of **2a**, the reaction was carried out starting from 250 mg (0.496 mmol) of racemic thioether (**R/S**)_{FC}-**1c**, 500 mg of phenol and 0.10 ml of an aqueous solution of H₂O₂ (30%). **2c** was recovered as an orange solid (yield: 237 mg, 92%). The two diastereomers were separated by chromatography on silica using a mixture of ether and acetone (5%) as eluent to give 111 mg of **2c-Dia1**, (**R_{FC},S_S**)/(**S_{FC},R_S**)-**2c** (yield:

43%) and 103 mg of **2c-Dia2**, (**R_{Fe},R_S**)/(**S_{Fe},S_S**)-**2c** (yield: 40%).MS (DCI, NH₃): m / z = 521 (M + 1, 100 %); 538 (M + NH₄, 80%).

(**R_{Fe},S_S**)/(**S_{Fe},R_S**)-**2c**: ¹H NMR (δ,500 MHz, CDCl₃):7.8 (2H, m, PPh₂); 7.58-7.47 (6H, m, PPh₂); 7.40 (2H, m, PPh₂); 5.07 (1H, broad s, subst Cp); 4.47 (1H, broad s, CH₂Cp); 4.41(5H, s, Cp) ; 4.41(1H, s, subst Cp); 3.78 (1H, s, subst Cp); 3.05 (1H, d(AB), J_{HH} = 14 Hz, CH₂Cp); 1.32 (9H, s, C(CH₃)₃). ¹³C{¹H} NMR (δ,125.8 MHz, CDCl₃): 134.6 (d, J_{CP} = 87 Hz, quat PPh₂); 132.4 (d, J_{CP} = 86 Hz, quat PPh₂); 131.9 (d, J_{CP} = 11 Hz, PPh₂); 131.7 (d, J_{CP} = 11 Hz, PPh₂); 131.6 (d, J_{CP} = 3 Hz, PPh₂); 131.5 (d, J_{CP} = 3 Hz, PPh₂); 128.6 (d, J_{CP} = 13 Hz, PPh₂); 128.2 (d, J_{CP} = 13 Hz, PPh₂); 85.2 (d, J_{CP} = 12 Hz, quat Cp); 74.9 (d, J_{CP} = 96 Hz, quat Cp); 73.8 (d, J_{CP} = 12 Hz, subst Cp); 73.6 (d, J_{CP} = 19 Hz, subst Cp); 71.1 (s, Cp); 70.0 (d, J_{CP} = 9 Hz, subst Cp); 54.2 (broad s, C(CH₃)₃); 44.9 (s, CH₂Cp); 23.1 (s, C(CH₃)₃). ³¹P{¹H} NMR (δ,202.5 MHz, CDCl₃): 41.8. Elemental analysis: C₂₇H₂₉FePS₂O; %C (calcd 62.31, found 62.16), % H (calcd 5.58, found 5.54).

(**R_{Fe},R_S**)/(**S_{Fe},S_S**)-**2c**: ¹H NMR (δ,500 MHz, CDCl₃): 7.83 (1H, dd, J_{HH} = 2 Hz, J_{HP} = 14 Hz, m C₆H₅); 7.82 (1H, dd, J_{HH} = 2 Hz, J_{HP} = 15Hz, m C₆H₅); 7.66 (1H, dd, J_{HH} = 2 Hz, J_{HP} = 15Hz, m C₆H₅); 7.65 (1H, dd, J_{HH} = 2 Hz, J_{HP} = 15Hz, m C₆H₅); 7.58-7.53 (1H, m, p C₆H₅); 7.52-7.45 (3H, m, p+2o C₆H₅); 7.42-7.37 (2H, m, 2o C₆H₅); 5.09 (1H, d(AB). J_{HH} = 13 Hz, CH₂Cp); 4.89 (1H, m, subst Cp); 4.43 (1H, m, subst Cp); 4.37 (5H, s, Cp); 3.75 (1H, m, subst Cp); 3.28 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 1.04 (9H, s, C(CH₃)₃). ¹³C{¹H} NMR (δ,125.8 MHz. CDCl₃):134.02 (d, J_{CP} = 87 Hz, quat PPh₂); 132.87 (d, J_{CP} = 82 Hz, quat PPh₂); 132.53 (d, J_{CP} = 10 Hz, m PPh₂); 131.97 (d, J_{CP} = 10 Hz, m PPh₂); 131.50 (d, J_{CP} = 3 Hz, pPPh₂); 131.45 (d, J_{CP} = 3 Hz, p PPh₂); 128.23 (d, J_{CP} = 4 Hz, o PPh₂); 128.13 (d, J_{CP} = 4 Hz, o PPh₂); 83.50 (d, J_{CP} = 12 Hz, quat Cp); 75.20 (d, J_{CP} = 9 Hz, subst Cp); 75.07 (d, J_{CP} = 13 Hz, subst Cp); 74.47 (d, J_{CP} = 95Hz, quat Cp); 71.04 (s, Cp); 70.41 (d, J_{CP} = 10 Hz, subst Cp); 53.23 (s, C(CH₃)₃); 47.84 (s, CH₂Cp); 22.75 (s, C(CH₃)₃). ³¹P{¹H} NMR (δ,202.5 MHz. CDCl₃):41.4.Elemental analysis:C₂₇H₂₇FePS₂O; %C (calcd 62.31, found 59.81), % H (calcd 5.58, found 5.58).

4.5 Synthesis of enantiomerically pure (**R_{Fe}**) 2-thiodiphenylphosphino-(phenylsulfoxymethyl) ferrocene, (**R_{Fe}**)-**2a**.

The reaction was carried out using enantiomerically pure (**R_{Fe}**)-**1a** following the same procedure described above for the synthesis of racemic **2a**. **Dia1**, (**R_{Fe},S_S**)-**2a** : [α]_D = -100.2; (CHCl₃. c=0.49); **Dia2**, (**R_{Fe},R_S**)-**2a** : [α]_D = +14.2; (CHCl₃. c=0.49).

4.6 (R_{Fc}) 2-thiodiphenylphosphino-(ethylsulfoxymethyl) ferrocene, (R_{Fc})-2b.

The reaction was carried out using enantiomerically pure (R_{Fc})-1b following the same procedure described above for the synthesis of racemic 2b.Dia1, (R_{Fc},R_S)-2b: $[\alpha]_D = -110.2$; ($CHCl_3$, $c=0.5$); Dia2, (R_{Fc},S_S)-2b: $[\alpha]_D = +32.1$; ($CHCl_3$, $c=0.505$).

4.7 (R_{Fc}) 2-thiodiphenylphosphino-(*tert*-butylsulfoxymethyl) ferrocene, (R_{Fc})-2c.

The reaction was carried out using enantiomerically pure (R_{Fc})-1c following the same procedure described above for the synthesis of racemic 2c.Dia1, (R_{Fc},S_S)-2c: $[\alpha]_D = -152.2$; ($CHCl_3$, $c=0.5$); Dia2, (R_{Fc},R_S)-2c: $[\alpha]_D = +5.0$; ($CHCl_3$, $c=0.515$).

4.8 (R_{Fc},S_S)/(S_{Fc},R_S)-2-diphenylphosphino-(phenyl-sulfoxymethyl)-ferrocene, 3a-Dia1.

In a Schlenktube under argon were introduced 81 mg (0.15 mmol) of racemic sulfoxide (R_{Fc},S_S)/(S_{Fc},R_S)-2a-Dia1 in 5 ml of toluene and 0.2 ml (0.18 g, 1.1 mmol) of tris(dimethylamino)phosphine. After reflux for 3 h, the solvent was removed by evaporation and the crude sulfoxide phosphine (R_{Fc},S_S)/(S_{Fc},R_S)-3a-Dia1 was obtained as an orange liquid (yield: 71 mg, 93%). 1H NMR (δ , 500 MHz, $CDCl_3$): 7.64-7.49 (4H, m, Ph); 7.49-7.32 (6H, m, Ph); 7.32-7.12 (5H, m, Ph); 4.43 (1H, dd(ABX), $J_{HH} = 13$ Hz, $J_{HP} = 2$ Hz, CH_2Cp); 4.42 (1H, broad s, substCp); 4.22 (1H, t, $J_{HH} = 2$ Hz, substCp); 4.01 (5H, s, Cp); 3.90 (1H, broad s, substCp); 3.74 (1H, d(AB), $J_{HH} = 13$ Hz, CH_2Cp). $^{31}P\{^1H\}$ NMR (δ , 202.5 MHz, $CDCl_3$): -22.5.

4.9 (R_{Fc},R_S)/(S_{Fc},S_S)-2-diphenylphosphino-(phenyl-sulfoxymethyl)-ferrocene, 3a-Dia2.

Following the same procedure described above for the desulfurisation of (R_{Fc},S_S)/(S_{Fc},R_S)-2a-Dia1, the reaction was realized from 92 mg (0.17 mmol) of racemic sulfoxide (R_{Fc},R_S)/(S_{Fc},S_S)-2a-Dia2 and 0.2 ml (0.18 g, 1.1 mmol) of tris(dimethylamino)phosphine. The sulfoxide phosphine product (R_{Fc},R_S)/(S_{Fc},S_S)-3a-Dia2 was obtained as an orange solid (yield: 73 mg, 85%). 1H NMR (δ , 500 MHz, $CDCl_3$): 7.65-7.5 (3H, m, Ph); 7.5-7.35 (6H, m, Ph); 7.35-7.2 (6H, m, Ph); 4.77 (1H, d, $J_{HH} = 1$ Hz, substCp); 4.43 (1H, t, $J_{HH} = 3$ Hz, substCp); 4.17 (1H, dd(ABX), $J_{HH} = 13$ Hz, $J_{HP} = 3$ Hz, CH_2Cp); 3.99 (5H, s, Cp); 3.85 (1H, t, $J_{HH} = 1$ Hz, substCp); 3.77 (1H, d(AB), $J_{HH} = 13$ Hz, CH_2Cp). $^{31}P\{^1H\}$ NMR (δ , 202.5 MHz, $CDCl_3$): -20.9.

4.10 (R_{Fc},R_S)/(S_{Fc},S_S)-2-diphenylphosphino-(ethylsulfoxymethyl) ferrocene, 3b-Dia1.

In a Schlenktube under argon were introduced 77 mg (0.157mmol) of the racemic sulfoxide(**R_{FC},R_S**)/(**S_{FC},S_S**)-**2b-Dia1** in 5 ml of toluene and 0.2 ml (0.18 g, 1.1 mmol) of tris(dimethylamino)phosphine. After reflux for 2 h, the solvent was evaporated under reduced pressure and the crude reaction product was purified by column chromatography on silica using degassed ether/dichloromethane mixture (1/0 - 1/1) as eluent. The sulfoxide phosphine product (**R_{FC},R_S**)/(**S_{FC},S_S**)-**3b-Dia1** was obtained as an orange solid (yield: 48mg, 66%). ¹H NMR (δ, 500 MHz, CDCl₃): 7.62-7.50 (3H, m, Ph); 7.44-7.35 (2H, m, Ph); 7.28-7.08 (5H, m, Ph); 4.62 (1H, q, J_{HH} = 1 Hz, substCp); 4.39 (1H, t, J_{HH} = 3 Hz, substCp); 4.15 (1H, dd(ABX), J_{HH} = 14 Hz, J_{HP} = 2 Hz, CH₂Cp); 4.03 (5H, s, Cp); 3.93 (1H, d, J_{HH} = 1 Hz substCp); 3.76 (1H, d(AB), J_{HH} = 14 Hz, CH₂Cp); 2.37 (1H, dq, J_{HH} = 7.7 Hz, J_{HP} = 2.7 Hz, CH₂CH₃); 1.12 (1H, t, J_{HH} = 7.7 Hz, CH₂CH₃). ³¹P{¹H} NMR (δ, 202.5 MHz, CDCl₃): -22.5. MS (DCI, NH₃): m/z = 461 (M + 1, 100 %).

4.11 (**R_{FC},S_S**)/(**S_{FC},R_S**)-2-diphenylphosphino-(ethylsulfoxymethyl) ferrocene, **3b-Dia2**.

Following the same procedure described above for the desulfurisation of (**R_{FC},R_S**)/(**S_{FC},S_S**)-**2b-Dia1**, the reaction was realized from 55mg (0.112 mmol) of racemic sulfoxide(**R_{FC},S_S**)/(**S_{FC},R_S**)-**2b-Dia2** and 0.2 ml (0.18 g, 1.1 mmol) of tris(dimethylamino)phosphine. The crude reaction product was purified by column chromatography on silica using degassed ether/methanol mixture (1/0 - 95/5) as eluent. The sulfoxide phosphine product (**R_{FC},S_S**)/(**S_{FC},R_S**)-**3b-Dia2** was obtained as an orange solid (yield: 33 mg, 65%). ¹H NMR (δ, 500 MHz, CDCl₃): 7.62-7.48 (3H, m, Ph); 7.43-7.35 (3H, m, Ph); 7.30-7.13 (4H, m, Ph); 4.72 (1H, d, J_{HH} = 1.5 Hz, substCp); 4.41 (1H, t, J_{HH} = 1 Hz, substCp); 4.10 (1H, dd(ABX), J_{HH} = 13 Hz, J_{HP} = 3 Hz, CH₂Cp); 4.01 (5H, s, Cp); 3.87 (1H, t, J_{HH} = 1 Hz, substCp); 3.72 (1H, d(AB), J_{HH} = 13 Hz, CH₂Cp); 2.45 (1H, dq, J_{HH} = 7.6 Hz, J_{HP} = 3 Hz, CH₂CH₃); 1.10 (1H, t, J_{HH} = 7.6 Hz, CH₂CH₃). ³¹P{¹H} NMR (δ, 202.8 MHz, CDCl₃): -21.5.

4.12 (**R_{FC},S_S**)/(**S_{FC},R_S**)-2-diphenylphosphino-(*tert*-butylsulfoxymethyl) ferrocene, **3c-Dia1**.

In a Schlenktube under argon were introduced 50 mg (0.096mmol) of the sulfoxide compound (**R_{FC},S_S**)/(**S_{FC},R_S**)-**2c-Dia1** in 5 ml of toluene and 0.1 ml (90 g, 0.55 mmol) of tris(dimethylamino)phosphine. After reflux for 2 h, the solvent was removed and the crude reaction product was purified by column chromatography on silica using degassed ether/dichloromethane mixture (1/1) as eluent. The sulfoxide phosphine product (**R_{FC},S_S**)/(**S_{FC},R_S**)-**3c-Dia1** was obtained as an orange liquid (yield: 36mg, 76%). ¹H NMR (δ, 500 MHz, CDCl₃): 7.58-7.42 (2H, m, Ph); 7.43-7.32 (3H, m, Ph); 7.30-7.18 (3H, m, Ph);

7.18-7.05 (2H, m, Ph); 4.79 (1H, s broad, substCp); 4.34 (1H, t, $J_{\text{HH}} = 1$ Hz, substCp); 4.08 (5H, s, Cp); 3.82 (1H, t, $J_{\text{HH}} = 1$ Hz, substCp); 3.73 (1H, dd(ABX), $J_{\text{HH}} = 12$ Hz, $J_{\text{HP}} = 2$ Hz, CH_2Cp); 3.30 (1H, d(AB), $J_{\text{HH}} = 12$ Hz, CH_2Cp); 1.2 (9H, s, $\text{C}(\text{CH}_3)_3$). $^{31}\text{P}\{^1\text{H}\}$ NMR (δ , 202.8 MHz, CDCl_3): -20.6.

4.13 ($R_{\text{Fc}}, R_{\text{S}}$)/($S_{\text{Fc}}, S_{\text{S}}$)-2-diphenylphosphino-(*tert*-butylsulfoxymethyl) ferrocene, 3c-Dia2.

Following the same procedure described above for the desulfurisation of ($R_{\text{Fc}}, S_{\text{S}}$)/($S_{\text{Fc}}, R_{\text{S}}$)-2c-Dia1, the reaction was realized from 70 mg (0.135 mmol) of the sulfoxide compound ($R_{\text{Fc}}, R_{\text{S}}$)/($S_{\text{Fc}}, S_{\text{S}}$)-2c-Dia2 and 0.2 ml (0.18 g, 1.1 mmol) of tris(dimethylamino)phosphine. The sulfoxide phosphine product ($R_{\text{Fc}}, R_{\text{S}}$)/($S_{\text{Fc}}, S_{\text{S}}$)-3c-Dia2 was obtained as an orange solid (yield: 57 mg, 86%). ^1H NMR (δ , 500 MHz, CDCl_3): 7.57-7.46 (3H, m, Ph); 7.42-7.34 (2H, m, Ph); 7.28-7.12 (5H, m, Ph); 4.78 (1H, d, $J_{\text{HH}} = 2$ Hz, substCp); 4.38 (1H, t, $J_{\text{HH}} = 1$ Hz, substCp); 4.06 (1H, dd(ABX), $J_{\text{HH}} = 13$ Hz, $J_{\text{HP}} = 2$ Hz, CH_2Cp); 4.04 (5H, s, Cp); 3.75 (1H, d, $J_{\text{HH}} = 1$ Hz, substCp); 3.36 (1H, d(AB), $J_{\text{HH}} = 13$ Hz, CH_2Cp); 1.05 (9H, s, $\text{C}(\text{CH}_3)_3$). $^{31}\text{P}\{^1\text{H}\}$ NMR (δ , 202.8 MHz, CDCl_3): -20.9.

X-ray analyses

A single crystal of each compound was mounted under inert perfluoropolyether at the tip of a glass fibre and cooled in the cryostream of either an Oxford-Diffraction XCALIBUR CCD diffractometer for 2a-Dia1, 2a-Dia2, 2b-Dia1 and 3a-Dia2 or a Stoe IPDS diffractometer for 2c-Dia1, 2c-Dia2 and 4a. Data were collected using the monochromatic $\text{MoK}\alpha$ radiation ($\lambda = 0.71073$).

The structures were solved by direct methods (SIR97²¹) and refined by least-squares procedures on F^2 using SHELXL-97.²² All H atoms attached to carbon were introduced in calculation in idealised positions and treated as riding models. The drawing of the molecules was realised with the help of ORTEP32.²⁴ Crystal data and refinement parameters are shown in Table 2.

Table 2. Crystal data and structure refinement parameters.

Identification code	2a-Dia1	2a-Dia2	2b-Dia1
Empirical formula	C ₂₉ H ₂₅ FeOPS ₂	C ₂₉ H ₂₅ FeOPS ₂	C ₂₅ H ₂₅ FeOPS ₂
Formula weight	540.43	540.43	492.39
Temperature, K	180(2)	180(2)	180(2)
Wavelength, Å	0.71073	0.71073	0.71073
Crystal system	Monoclinic	Monoclinic	Orthorhombic
Space group	Cc	P 2 ₁ /a	Pcab
a, Å	14.5928(10)	12.812(2)	8.1197(6)
b, Å	18.4079(11)	9.4172(16)	16.9489(11)
c, Å	9.4244(6)	21.896(4)	33.823(2)
α, °	90.0	90.0	90.0
β, °	93.858(5)	104.832(16)	90.0
γ, °	90.0	90.0	90.0
Volume, Å ³	2525.9(3)	2553.8(8)	4654.7(5)
Z	4	4	8
Density (calcd), Mg/m ³	1.421	1.322	1.405
Absorption coefficient, mm ⁻¹	0.847	0.754	0.911
F(000)	1120	1056	2048
Crystal size, mm ³	0.45x0.42x0.2	0.16 x 0.10 x 0.05	0.52x 0.39 x 0.12
Theta range, °	3.56 to 27.81	2.71 to 26.03	3.32 to 26.37
Reflections collected	9974	18130	33297
Independent reflections [R(int)]	4487 (0.0242)	5055 (0.1527)	4751 (0.0634)
Completeness, %	99.7	99.7	99.8
Absorption correction	Multiscan	Multi-scan	Multi-scan
Max. , min. transmission	0.8415 and 0.6881	1.0500 and 0.8078	0.8230 and 0.6750
Refinement method	F ²	F ²	F ²
Data / restraints / parameters	4487 / 2 / 308	5055 / 0 / 307	4751 / 0 / 272
Goodness-of-fit on F ²	1.055	0.892	1.081
Final R indices [I>2sigma(I)]	0.0278, 0.0642	0.0662, 0.1564	0.0514, 0.1035
R indices (all data)	0.0305, 0.0656	0.1713, 0.2309	0.0704, 0.1118
Absolute structure parameter	0.019(14)		
Largest diff. peak and hole, e.Å ⁻³	0.326 and -0.219	0.711 and -0.865	0.417 and -0.367

Table 1. Crystal data and structure refinement(Contd.).

Identification code	2c-Dia1	2c-Dia2	3a-Dia2
Empirical formula	C ₂₇ H ₂₉ FeOPS ₂	C ₂₇ H ₂₉ FeOPS ₂	C ₂₉ H ₂₅ FeOPS
Formula weight	520.44	520.44	508.37
Temperature, K	180(2)	180(2)	180(2)
Wavelength, Å	0.71073	0.71073	0.70930
Crystal system	Monoclinic	Monoclinic	Triclinic
Space group	Cc	P2 ₁	P -1
a, Å	11.8130(12)	14.7673(14)	7.8035(8)
b, Å	14.8503(12)	10.1912(7)	9.3075(9)
c, Å	13.9653(16)	17.0533(18)	17.5040(15)
α, °	90.0	90.0	79.331(7)
β, °	91.394(13)	98.030(12)	87.727(7)
γ, °	90.0	90.0	77.299(8)
Volume, Å ³	2449.2(4)	2541.3(4)	1218.8(2)
Z	4	4	2
Density (calcd), Mg/m ³	1.411	1.360	1.385
Absorption coefficient, mm ⁻¹	0.870	0.838	0.790
F(000)	1088	1088	528
Crystal size, mm ³	0.76 x 0.22 x 0.2	0.56 x 0.52 x 0.02	0.35 x 0.32 x 0.2
Theta range, °	2.62 to 26.04	2.33 to 26.08	3.47 to 28.22
Reflections collected	11892	21668	10784
Independent reflections [R(int)]	4769 (0.0442)	9539 (0.0730)	6009 (0.0237)
Completeness, %	98.8	97.7	99.1
Absorption correction	Multi-scan	Multi-scan	Multi-scan
Max. , min. transmission	0.943 and 0.610	0.8058 and 0.7391	0.7521 and 0.6862
Refinement method	F ²	F ²	F ²
Data / restraints / parameters	4769 / 2 / 292	9539 / 1 / 584	6009 / 0 / 298
Goodness-of-fit on F ²	1.025	0.843	0.991
Final R indices [I>2sigma(I)]	0.0302, 0.0778	0.0379, 0.0611	0.0389, 0.1012
R indices (all data)	0.0314, 0.0785	0.0716, 0.0685	0.0507, 0.1067
Absolute structure parameter	-0.004(11)	0.284(14)	
Largest diff. peak and hole, e.Å ⁻³	0.318 and -0.384	0.328 and -0.425	1.418 and -0.385

Table 1. Crystal data and structure refinement(Contd.).

Identification code	4a
Empirical formula	C ₂₉ H ₂₅ Cl ₂ FeOPPt S
Formula weight	774.36
Temperature, K	180(2)
Wavelength, Å	0.71073
Crystal system	Monoclinic
Space group	P2 ₁ /n
a, Å	18.6057(14)
b, Å	7.5780(7)
c, Å	19.3027(14)
α, °	90.0
β, °	107.659(8)
γ, °	90.0
Volume, Å ³	2593.3(4)
Z	4
Density (calcd), Mg/m ³	1.983
Absorption coefficient, mm ⁻¹	6.319
F(000)	1504
Crystal size, mm ³	0.18 x 0.08 x 0.08
Theta range, °	2.21 to 26.06
Reflections collected	24800
Independent reflections [R(int)]	5081 (0.0562)
Completeness, %	99.0
Absorption correction	Multi-scan
Max. , min. transmission	0.5281 and 0.4506
Refinement method	F ²
Data / restraints / parameters	5081 / 0 / 325
Goodness-of-fit on F ²	0.981
Final R indices [I>2σ(I)]	0.0289, 0.0585
R indices (all data)	0.0445, 0.0624
Absolute structure parameter	
Largest diff. peak and hole, e.Å ⁻³	0.907 and -0.927

- ¹Imamoto, T.; Sugita, K.; Yoshida, K. *J. Am. Chem. Soc.* **2005**, *127*, 11934.
- ²Knowles, W. S.; Sabacky, M. J.; Vineyard, B. D.; Weinkauff, D. J. *J. Am. Chem. Soc.* **1975**, *97*, 2567.
- ³For recent reviews, see: (a) Masdeu-Bulto, A. M.; Dieguez, M.; Martin, E.; Gomez, M. *Coord. Chem. Rev.* **2003**, *159*; (b) Pellissier, H. *Tetrahedron* **2007**, *63*, 1297; (c) Mellah, M.; Voituriez, A.; Schulz, E. *Chem. Rev.* **2007**, *107*, 5133; (d) Malacea, R.; Manoury, E. in « *Phosphorus-Sulfur Ligands* » from “*Phosphorus Ligands in Asymmetric Catalysis*” edited by Armin Börner (Wiley-VCH, Weinheim, Allemagne) **2005**, 749; (e) Lam, F. L.; Kwong, F. Y.; Chan, A. S. C. *Chem. Commun.* **2010**, *46*, 4649.
- ⁴ Selected references (a) Albinati, A.; Pregosin, P. S.; Wick, K. *Organometallics* **1996**, *15*, 2419; (b) Spencer, J.; Gramlich, V.; Hausel, R.; Togni, R. *Tetrahedron: Asymmetry* **1996**, *7*, 41; (c) Enders, D.; Peters, R.; Lochtmann, R.; Raabe, G.; Runsink, J.; Bats, J. W. *Eur. J. Org. Chem.* **2000**, 3399; (d) Verdaguer, X.; Moyano, A.; Pericas, M. A.; Riera, A.; Maestro, M. A.; Mahia, J. *J. Am. Chem. Soc.* **2000**, *122*, 10242; (e) Evans, D. A.; Michael, F. E.; Tedrow, J. S.; Campos, K. R. *J. Am. Chem. Soc.* **2003**, *125*, 3534; (f) Nakano, H.; Yokohama, J.; Okuyama, Y.; Fujita, R.; Hongo, H. *Tetrahedron: Asymmetry* **2003**, *14*, 2361; (g) Tu, T.; Zhou, Y.-G.; Hou, X.-L.; Dai, L.-X.; Dong, X.-C.; Yu, Y.-H.; Sun, J. *Organometallics* **2003**, *22*, 1255; (h) Priego, J.; Mancheño, O. G.; Cabrera, S.; Arrayás, R.; Llamas, T.; Carretero, J. C. *Chem. Commun.* **2002**, 2512; (i) Mancheño, O. G.; Priego, J.; Cabrera, S.; Arrayas, R. G.; Llamas, T.; Carretero, J. C. *J. Org. Chem.* **2003**, *68*, 3679. (l) Zeng, W.; Zhou, Y. G. *Tet. Lett.* **2007**, *48*, 4619.
- ⁵ (a) Alcock, N. X.; Brown, J. M.; Evans, P. L. *J. Organomet. Chem.* **1988**, *356*, 233; (b) Quek, G. H.; Leung, P. H.; Mok, K. F. *Inorg. Chim. Acta* **1995**, *239*, 185
- ⁶ (a) Sola, J.; Reves, M.; Riera, A.; Verdaguer, X. *Angew. Chem. Int. Ed.* **2007**, *46*, 5020; (b) Reves, M.; Achard, T.; Sola, J.; Riera, A.; Verdaguer, X. *J. Org. Chem.* **2008**, *73*, 7080; (c) Ji, Y.; Riera, A.; Verdaguer, X. *Org. Lett.* **2009**, *11*, 19 4346.
- ⁷ (a) Chooi, S. Y. M.; Leung, P. H.; Mok, K. F. *Inorg. Chim. Acta* **1993**, *205*, 245–247; (b) Hiroi, K.; Suzuki, Y.; Kawagishi, R. *Tet. Lett.* **1999**, *40*, 715; (c) Xing, J.; Cao, P.; Liao, J. *Tetrahedron: Asymmetry* **2012**, *23*, 527–535.
- ⁸ Ferrer, C.; Riera, A.; Verdaguer, X. *Organometallics* **2009**, *28*, 4571.
- ⁹ Solà, J.; Riera, A.; Verdaguer, X. *J. Organomet. Chem.* **2010**, *695*, 2377.
- ¹⁰ (a) Riant, O.; Argouarch, G.; Guillaneux, D.; Samuel, O.; Kagan, H. B. *J. Org. Chem.* **1998**, *63*, 3511; (b) Cheemala, M. N.; Knochel, P. *Org. Lett.* **2007**, *9*, 3089; (c) Zhang, H. L.; Hou, X. L.; Dai, L. X.; Luo, Z. B. *Tetrahedron: Asymmetry* **2007**, *18*, 224.
- ¹¹ (a) Routaboul, L.; Vincendeau, S.; Daran, J. C.; Manoury, E. *Tetrahedron: Asymmetry* **2005**, *16*, 2685; (b) Diab, L.; Gouygou, M.; Manoury, E.; Kalck, P.; Urrutigoity, M. *Tet. Lett.* **2008**, *49*, 5186. (c) Le Roux, E.; Malacea, R.; Manoury, E.; Poli, R.; Gonsalvi, L.; Peruzzini, M. *Adv. Synth. Catal.* **2007**, *349*, 309; (d) Manoury, E.; Poli, R. “*Phosphine-Containing Planar Chiral ferrocenes: Synthesis, Coordination Chemistry and Applications to Asymmetric Catalysis*.” in the Series: *Catalysis by Metal Complexes (CMCO)*, Volume 36 (*Phosphorus Chemistry: Catalysis and Material Science Applications*), edited by Maurizio Peruzzini and Luca Gonsalvi (Springer Verlag, Germany) **2011**, p121.
- ¹² Mateus, N.; Routaboul, L.; Daran, J.-C.; Manoury, E. *J. Organomet. Chem.* **2006**, *691*, 2297.
- ¹³ Xu, W. L.; Zheng Li, Y.; Shan Zhang, Q.; Sun Zhu, H. *Synthesis* **2004**, *2*, 227.¹⁴ Weigand, W.; Bosl, G. *Zeitschrift fuer Naturforschung, Chemical Sciences* **1992**, *47*, 1165.
- ¹⁵ Di Furia, F.; Modena, G.; Seraglia, R. *Synthesis*, **1984**, 325.
- ¹⁶ Conte, V.; Di Furia, F.; Lucini, G.; Modena, G. *Tet. Lett.*, **1989**, *30*, 4859.
- ¹⁷ (a) Calligaris, M. *Coord. Chem. Rev.* **2004**, *248*, 351; (b) Kagan, H. B.; Ronan, B., *Rev. Heteroatom Chem. Rev.* **1992**, *248*, 92.
- ¹⁸ Anderson, G. K.; Kumar, R. *Inorg. Chem.* **1984**, *23*, 4064.
- ¹⁹ Cremer, D.; Pople, J. A. *J. Am. Chem. Soc.* **1975**, *97*, 1354.
- ²⁰ Malacea, R.; Manoury, E.; Daran, J. C.; Poli, R. *J. Molec. Str.* **2008**, *890*, 249.
- ²¹ (a) Malacea, R.; Manoury, E.; Routaboul, L.; Daran, J. C.; Poli, R.; Dunne, J. P.; Withood, A. C.; Godard, C.; Duckett, S. B. *Eur. J. Inorg. Chem.* **2006**, *9*, 1803; (b) Malacea, R.; Daran, J.-C.; Duckett, S. B.; Dunne, J. P.; Manoury, E.; Poli, R.; Withood, A. C. *Dalton Trans.* **2006**, 3350.
- ¹⁸ Altomare, A.; Burla, M. C.; Camalli, M.; Cascarano, G. L.; Giacovazzo, C.; Guagliardi, A.; Moliterni, A. G. G.; Polidori, G.; Spagna, R. *J. Appl. Cryst.* **1999**, *32*, 115.
- ¹⁹ Sheldrick, G. M. *Acta Cryst.* **2008**, *A64*, 112.
- ²⁰ Farrugia, L. J. *J. Appl. Cryst.* **1999**, *32*, 837.
- ²¹ Farrugia, L. J. *J. Appl. Cryst.* **1997**, *30*, 565.
- ²² Spek, A. L. *J. Appl. Cryst.* **2003**, *36*, 7.