



HAL
open science

Adaptive reconstruction of imperfectly-observed monotone functions, with applications to uncertainty quantification

Luc Bonnet, Jean-Luc Akian, Eric Savin, Tim J. Sullivan

► **To cite this version:**

Luc Bonnet, Jean-Luc Akian, Eric Savin, Tim J. Sullivan. Adaptive reconstruction of imperfectly-observed monotone functions, with applications to uncertainty quantification. *Algorithms*, 2020, 13 (8), pp.196. 10.3390/a13080196 . hal-02908957v2

HAL Id: hal-02908957

<https://hal.science/hal-02908957v2>

Submitted on 17 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Adaptive reconstruction of imperfectly-observed monotone functions, with applications to uncertainty quantification

L. Bonnet^{1,2} J.-L. Akian¹ É. Savin¹ T. J. Sullivan^{3,4}

August 11, 2020

Abstract: Motivated by the desire to numerically calculate rigorous upper and lower bounds on deviation probabilities over large classes of probability distributions, we present an adaptive algorithm for the reconstruction of increasing real-valued functions. While this problem is similar to the classical statistical problem of isotonic regression, the optimisation setting alters several characteristics of the problem and opens natural algorithmic possibilities. We present our algorithm, establish sufficient conditions for convergence of the reconstruction to the ground truth, and apply the method to synthetic test cases and a real-world example of uncertainty quantification for aerodynamic design.

Keywords: adaptive approximation • isotonic regression • optimisation under uncertainty • uncertainty quantification • aerodynamic design

2010 Mathematics Subject Classification: 49M30 • 62G08 • 68T37

1 Introduction

This paper considers the problem of adaptively reconstructing a monotonically increasing function F^\dagger from imperfect pointwise observations of this function. In the statistical literature, the problem of estimating a monotone function is commonly known as *isotonic regression*, and it is assumed that the observed data consist of noisy pointwise evaluations of F^\dagger . However, we consider this problem under assumptions that differ from the standard formulation, and these differences motivate our algorithmic approach to the problem. To be concrete, our two motivating examples are that

$$F^\dagger(x) := \mathbb{P}_{\Xi \sim \mu}[g(\Xi) \leq x] \quad (1.1)$$

is the cumulative distribution function (CDF) of a known real-valued function g of a random variable Ξ with known distribution μ , or that

$$F^\dagger(x) := \sup_{(g,\mu) \in \mathcal{A}} \mathbb{P}_{\Xi \sim \mu}[g(\Xi) \leq x] \quad (1.2)$$

is the supremum of a family of such CDFs over some class \mathcal{A} . We assume that we have access to a numerical optimisation routine that can, for each x and some given numerical parameters q (e.g. the number of iterations or other convergence tolerance parameters), produce a *numerical estimate* or

¹ ONERA, 29 Avenue de la Division Leclerc, 92320 Châtillon, France (luc.bonnet@onera.fr, jean-luc.akian@onera.fr, eric.savin@onera.fr)

² Laboratoire MSSMat - UMR CNRS 8579, CentraleSupélec, 8–10 rue Joliot Curie, 91190 Gif sur Yvette, France (luc.bonnet@ens-paris-saclay.fr)

³ Mathematics Institute and School of Engineering, University of Warwick, Coventry CV4 7AL, United Kingdom (t.j.sullivan@warwick.ac.uk)

⁴ Zuse Institute Berlin, Takustraße 7, 14195 Berlin, Germany (sullivan@zib.de)

observation $G(x, q)$ of $F^\dagger(x)$; furthermore, we assume that $G(x, q) \leq F^\dagger(x)$ is always true, i.e. the numerical optimisation routine always under-estimates the true optimum value, and that the positive error $F^\dagger(x) - G(x, q)$ can be controlled to some extent through the choice of the optimisation parameters q , but remains essentially influenced by randomness in the optimisation algorithm for each x . The assumption $G(x, q) \leq F^\dagger(x)$ is for example coherent with either [Equation \(1.1\)](#), which may be approached by increasing the number of samples (say q) in a Monte Carlo simulation, or [Equation \(1.2\)](#), which is a supremum over a set that may be explored only partially by the algorithm.

A single observation $G(x, q)$ yields some limited information about $F^\dagger(x)$; a key limitation is that one may not even know a priori how accurate $G(x, q)$ is. Naturally, one may repeatedly evaluate G at x , perhaps with different values of the optimisation parameters q , in order to more accurately estimate $F^\dagger(x)$. However, a key observation is that a *suite* of observations $G(x_i, q_i)$, $i = 1, \dots, I$, contains much more information than simply estimates of $F^\dagger(x_i)$, $i = 1, \dots, I$, and this information can and must be used. For example, suppose that the values $(G(x_i, q_i))_{i=1}^I$ are not increasing, e.g. because

$$G(x_i, q_i) > G(x_{i'}, q_{i'}) \quad \text{and} \quad x_i < x_{i'}.$$

Such a suite of observations would be inconsistent with the axiomatic requirement that F^\dagger is an increasing function. In particular, while the observation at x_i may be relatively good or bad on its own merits, the observation $G(x_{i'}, q_{i'})$ at $x_{i'}$, which violates monotonicity, is in some sense “useless” as it gives no better lower bound on $F^\dagger(x_{i'})$ than the observation at x_i does. The observation at $x_{i'}$ is thus a good candidate for repetition with more stringent optimisation parameters q — and this is not something that could have been known without comparing it to the rest of the data set.

The purpose of this article is to leverage this and similar observations to define an algorithm for the reconstruction of the function F^\dagger , repeating old observations of insufficient quality and introducing new ones as necessary. The principal parameter in the algorithm is an “exchange rate” \mathcal{E} that quantifies the degree to which the algorithm prefers to have a few high-quality evaluations versus many poor-quality evaluations. Our approach is slightly different from classical isotonic (or monotonic) regression, which is understood as the least-squares fitting of an increasing function to a set of points in the plane. The latter problem is uniquely solvable and its solution can be constructed by the pool adjacent violators algorithm (PAVA) extensively studied in [Barlow et al. \(1972\)](#). This algorithm consists of exploring the data set from left to right until the monotonicity condition is violated, and replacing the corresponding observations by their average while back-averaging to the left if needed to maintain monotonicity. Extensions to the PAVA have been developed by [de Leeuw et al. \(2009\)](#) to consider non least-squares loss functions and repeated observations, by [Tibshirani et al. \(2011\)](#) to consider “nearly-isotonic” or “nearly-convex” fits, and by [Jordan et al. \(2019\)](#) to consider general loss functions and partially ordered data sets. Useful references on isotonic regression also include [Robertson et al. \(1988\)](#) and [Groeneboom and Jongbloed \(2014\)](#).


The remainder of this paper is structured as follows. [Section 2](#) presents the problem description and notation, after which the proposed adaptive algorithm for the reconstruction of F^\dagger is presented in [Section 3](#). We demonstrate the convergence properties of the algorithm in [Section 3.2](#) and study its performance on several analytically tractable test cases in [Section 4](#). [Section 5](#) details the application of the algorithm to a challenging problem of the form [Equation \(1.2\)](#) drawn from aerodynamic design. Some closing remarks are given in [Section 6](#).

2 Notation and problem description

In the following, the “ground truth” response function that we wish to reconstruct is denoted $F^\dagger: [a, b] \rightarrow \mathbb{R}$ and has inputs $x \in [a, b] \subset \mathbb{R}$. It is assumed that F^\dagger is monotonically increasing and non-constant on $[a, b]$. In contrast, $G: [a, b] \times \mathbb{R}_+ \rightarrow \mathbb{R}$ denotes the numerical process used to obtain an imperfect pointwise observation y of $F^\dagger(x)$ at some point $x \in [a, b]$ for some numerical parameter $q \in \mathbb{R}_+$. Here, on a heuristic level, $q > 0$ stands for the “quality” of the noisy evaluation $G(x, q)$.

The main aim of this paper is to show the effectiveness of the proposed algorithm for the adaptive reconstruction of F^\dagger , which could be continuous or not, from imperfect pointwise observations $G(x_i, q_i)$ of F^\dagger , where we are free to choose x_{i+1} and q_{i+1} adaptively based upon x_j , q_j , and $G(x_j, q_j)$ for $j \leq i$.

First, we associate with I imperfect pointwise observations $\{x_i, y_i := G(x_i, q_i)\}_{i=1}^I \subset [a, b] \times \mathbb{R}$, positive numbers $\{q_i\}_{i=1}^I \subset \mathbb{R}_+$ which we will call *qualities*. The quality q_i quantifies the confidence we have in



(a) Possible ground truth functions between two consecutive points x_1 and x_2 . The ground truth function must lie in the area formed by these two points.

(b) Right-continuous piecewise constant interpolation function.

Figure 2.1: Possible ground truth functions between two consecutive points x_1 and x_2 , and our choice of piecewise constant interpolant.

the pointwise observation y_i of $F^\dagger(x_i)$ using the numerical process $G(x_i, q_i)$. The higher this value, the greater the confidence. We divide this quality as the product of two different numbers c_i and r_i , $q_i = c_i \times r_i$, with the following definitions:

- *Consistency* $c_i \in \{0, 1\}$: This describes the fact that two successive points must be monotonically consistent with respect to each other. That is, when one takes two input values $x_2 > x_1$, one should have $y_2 \geq y_1$ as y must be monotonically increasing. There is no consistency associated with the very first data point as it does not have any predecessor.
- *Reliability* $r_i \in \mathbb{R}_+$: This describes how confident we are about the numerical value. Typically, it will be related to some error estimator if one is available, or the choice of optimisation parameters. It is expected that the higher the reliability, the closer the pointwise observation is to the true value, in average.

Typically, if the observation $y_{i+1} = G(x_{i+1}, q_{i+1})$ is consistent with regard to the observation $y_i = G(x_i, q_i)$ where $x_{i+1} > x_i$, the quality q_{i+1} associated with y_{i+1} will be equal to $q_{i+1} = r_{i+1} \in \mathbb{R}_+^*$ since $c_{i+1} = 1$ in this case. If the value is not consistent, we have $q_{i+1} = r_{i+1} \times c_{i+1} = 0$. Finally, if $x = a$ there is no notion of consistency as there is no point preceding it. Thereby, the quality associated with this point is only equal to its reliability.

Moreover, we associate to these pointwise observations a notion of area, illustrated in Figure 2.1 and defined as follows. Consider two consecutive points x_i and x_{i+1} with their respective observations y_i and y_{i+1} , the area a_i for these two points is

$$a_i = (x_{i+1} - x_i) \times (y_{i+1} - y_i). \quad (2.1)$$

Thus, we can define a vector $\mathbf{a} = \{a_i\}_{i=1}^{I-1}$ which contains all the computed areas for the whole dataset. In addition, we can assure that if we take two points x_1 and $x_2 > x_1$ with $y_1 = F^\dagger(x_1)$ and $y_2 = F^\dagger(x_2)$ —namely, the error at these point is equal to zero, the graph of ground truth function F^\dagger must lie in the rectangular area spanned by the two points $(x_1, F^\dagger(x_1))$ and $(x_2, F^\dagger(x_2))$.

To adopt a conservative point of view, we choose as the approximating function F of F^\dagger a piecewise constant interpolation function, say:

$$F(x) = \sum_{i=1}^{I-1} y_i \mathbb{1}_{[x_i, x_{i+1})}(x), \quad (2.2)$$

where $\mathbb{1}_{\mathcal{I}}$ denotes the indicator function of the interval \mathcal{I} . We do not want this interpolation function

to overestimate the true function F^\dagger as one knows that the numerical estimate in our case always underestimates the ground truth function $F^\dagger(x)$. See [Figure 2.1](#) for an illustration of this choice, which can be viewed as a worst-case approach. Indeed, this chosen interpolation function is the worst possible function underestimating F^\dagger given two points x_1 and x_2 and their respective observations y_1 and y_2 .

3 Reconstruction algorithms

The reconstruction algorithm that we propose, [Algorithm 1](#), is driven to produce a sequences of reconstructions that converges to F^\dagger by following a principle of *area minimisation*: we associate to the discrete data set $\{x_i, y_i\}_{i=1}^I \subset [a, b] \times \mathbb{R}$ a natural notion of area [\(2.1\)](#) as explained above, and seek to drive this area towards zero. The motivation behind this objective is in [Proposition 3.6](#) which states that the area converges to 0 as more points are added to the data set. However, the objective of minimising the area is complicated by the fact that evaluations of F^\dagger are imperfect. Therefore, a key user-defined parameter in the algorithm is $\mathcal{E} \in (0, \infty)$, which can be thought of as an “exchange rate” that quantifies to what extent the algorithm prefers to redo poor-quality evaluations of the target function versus driving the area measure to zero.

3.1 Algorithm

The main algorithm is organized as follows, starting from $I^{(0)} \geq 2$ points and a dataset that is assumed to be consistent at the initial step $n = 0$. It goes through N iterations, where N is either fixed a priori, or obtained a posteriori once a stopping criterion is met. Note that q_{new} stands for the quality of a newly generated observation y_{new} for any new point x_{new} introduced by the algorithm. The latter is driven by the user-defined “exchange rate” \mathcal{E} as explained just above. At each step n , the algorithm computes the weighted area $\text{WA}^{(n)}$ as the minimum of the quality times the sum of the areas of the data points:

$$\text{WA}^{(n)} = q_-^{(n)} \times \text{A}^{(n)}, \quad (3.1)$$

where

$$q_-^{(n)} = \min_{1 \leq i \leq I^{(n)}} \{q_i^{(n)}\}, \quad \text{A}^{(n)} = \sum_{i=1}^{I^{(n)}-1} a_i^{(n)}, \quad (3.2)$$

$a_i^{(n)}$ is the area computed by [Equation \(2.1\)](#) at step n (see also [Equation \(3.5\)](#)), and $I^{(n)}$ is the number of data points. Then it is divided into two parts according to the value of $\text{WA}^{(n)}$ compared to \mathcal{E} .

- If $\text{WA}^{(n)} < \mathcal{E}$, then the algorithm aims at increasing the quality $q_-^{(n)}$ of the worst data point (the one with the lowest quality) with index $i_-^{(n)} = \arg \min_{1 \leq i \leq I^{(n)}} \{q_i^{(n)}\}$ at step n . It stores the corresponding old value y_{old} , searches for a new value y_{new} by improving successively the quality of this very point, and stops when $y_{\text{new}} > y_{\text{old}}$.
- If $\text{WA}^{(n)} \geq \mathcal{E}$, then the algorithm aims at driving the total area $\text{A}^{(n)}$ to zero. In that respect, it identifies the biggest rectangle

$$a_+^{(n)} = \max_{1 \leq i \leq I^{(n)}-1} \{a_i^{(n)}\} \quad (3.3)$$

and its index

$$i_+^{(n)} = \arg \max_{1 \leq i \leq I^{(n)}-1} \{a_i^{(n)}\} \quad (3.4)$$

and adds a new point x_{new} at the middle of this biggest rectangle. Then, it computes a new data value $y_{\text{new}} = G(x_{\text{new}}, q_{\text{new}})$ with a new quality q_{new} .

In both cases, the numerical parameters q_{new} (for example a number of iterations, or the size of a sampling set or a population) are arbitrary and any value can be chosen in practice each time a new point x_{new} is added to the dataset. They can be increased arbitrarily as well each time such a new point has to be improved. Indeed, the numerical parameters q of the optimisation routine we have access to can be increased as much as desired, and increasing them will improve the estimates $G(x, q)$ of the true function $F^\dagger(x)$ uniformly in x ; see [Assumption 3.1](#). The algorithm then verifies the consistency of the dataset by checking the quality of each point. If there is any inconsistent point, the algorithm computes a new value until obtaining consistency by improving successively the corresponding reliability. This is achieved in a

finite number of steps starting from an inconsistent point and exploring the dataset from the left to the right.

Finally, the algorithm updates the quality vector $\{q_i^{(n+1)}\}_{i=1}^{I^{(n+1)}}$, the area vector $\{a_i^{(n+1)}\}_{i=1}^{I^{(n+1)}}$, the worst quality $q_-^{(n+1)}$ and the index $i_-^{(n+1)}$ of the corresponding point, the biggest rectangle $a_+^{(n+1)}$ and its index $i_+^{(n+1)}$, and then the new weighted area $WA^{(n+1)}$.

Algorithm 1: Adaptive algorithm to reconstruct a monotonically increasing function F^\dagger

Input: $I^{(0)} \geq 2$, $\{x_i^{(0)}, y_i^{(0)}, q_i^{(0)}\}_{i=1}^{I^{(0)}}$ and \mathcal{E} .

Output: $\{x_i^{(N)}, y_i^{(N)}, q_i^{(N)}\}_{i=1}^{I^{(N)}}$ with $I^{(N)} \geq I^{(0)}$.

Initialization:

Get the worst quality point and its index:

- $q_-^{(0)} = \min_{1 \leq i \leq I^{(0)}} \{q_i^{(0)}\}$;
- $i_-^{(0)} = \arg \min_{1 \leq i \leq I^{(0)}} \{q_i^{(0)}\}$.

Compute the area of each pair of data points: $a_i^{(0)} = (x_{i+1}^{(0)} - x_i^{(0)}) \times (y_{i+1}^{(0)} - y_i^{(0)})$.

Get the biggest rectangle and its index:

- $a_+^{(0)} = \max_{1 \leq i \leq I^{(0)}-1} \{a_i^{(0)}\}$;
- $i_+^{(0)} = \arg \max_{1 \leq i \leq I^{(0)}-1} \{a_i^{(0)}\}$.

Define the weighted area at step $n = 0$ as $WA^{(0)} = q_-^{(0)} \times \sum_{i=1}^{I^{(0)}-1} a_i^{(0)}$.

while $n \leq N$ **do**

if $WA^{(n)} < \mathcal{E}$ **then**

Data points are unchanged: $I^{(n+1)} = I^{(n)}$ and $\{x_i^{(n+1)}\}_{i=1}^{I^{(n+1)}} = \{x_i^{(n)}\}_{i=1}^{I^{(n)}}$;

Store the old value $y_{\text{old}} = y_{i_-^{(n)}}$;

while $y_{\text{new}} \leq y_{\text{old}}$ **do**

Compute a new value $y_{\text{new}} = G(x_{i_-^{(n)}}, q_{\text{new}})$;

end

else

Introduce a new point at the middle of the biggest rectangle: $I^{(n+1)} = I^{(n)} + 1$,

$x_{\text{new}} = \frac{1}{2}(x_{i_+^{(n)}}^{(n)} + x_{i_+^{(n)}+1}^{(n)})$, and

$(x_1^{(n+1)}, \dots, x_{i_+^{(n)}}^{(n+1)}, x_{i_+^{(n)}+1}^{(n+1)}, x_{i_+^{(n)}+2}^{(n+1)}, \dots, x_{I^{(n+1)}}^{(n+1)}) = (x_1^{(n)}, \dots, x_{i_+^{(n)}}^{(n)}, x_{\text{new}}, x_{i_+^{(n)}+1}^{(n)}, \dots, x_{I^{(n)}}^{(n)})$;

Compute the new value $y_{\text{new}} = G(x_{\text{new}}, q_{\text{new}})$;

end

Verify consistency of the pointwise observations $\{y_i^{(n+1)}\}_{i=1}^{I^{(n+1)}}$ by checking their quality. If there are not consistent, recompute them until obtaining consistency and then update the quality vector;

Compute the new quality vector $\{q_i^{(n+1)}\}_{i=1}^{I^{(n+1)}}$ and area vector $\{a_i^{(n+1)}\}_{i=1}^{I^{(n+1)}}$;

Update $q_-^{(n+1)}$, $i_-^{(n+1)}$, $a_+^{(n+1)}$ and $i_+^{(n+1)}$;

Compute $WA^{(n+1)} = q_-^{(n+1)} \times \sum_{i=1}^{I^{(n+1)}-1} a_i^{(n+1)}$;

$n = n + 1$;

end

3.2 Proof of convergence

We denote by $I^{(n)}$ the number of data points, and $\{x_i^{(n)}, y_i^{(n)}, q_i^{(n)}\}_{i=1}^{I^{(n)}}$ the positions of the data points, the observations given by the optimization algorithm at these positions, and the qualities associated with the optimization algorithm at the step n of [Algorithm 1](#). For each $i = 1, \dots, I^{(n)} - 1$, we define

$s_i^{(n)} = [x_i^{(n)}, x_{i+1}^{(n)}[\subset [a, b]$ and the vector containing all rectangle areas $\{a_i^{(n)}\}_{i=1}^{I^{(n)}-1}$ by:

$$a_i^{(n)} = (x_{i+1}^{(n)} - x_i^{(n)}) \times (y_{i+1}^{(n)} - y_i^{(n)}). \quad (3.5)$$

The pointwise observation $y_i^{(n)} = G(x_i^{(n)}, q_i^{(n)})$ is thus associated to the quality $q_i^{(n)} \in \mathbb{R}_+$, which quantifies the confidence we have in this observation as outlined in the problem description in [Section 2](#). This number can represent the inverse error achieved by the optimization algorithm, for example, or the number of iterations, or the number of individuals in a population, or any other numerical parameter pertaining to this optimization process. The higher it is, the closer the observation is to the true target value. Therefore we consider the following assumption on the numerical process G .

Assumption 3.1. $G(x, q)$ converges to $F^\dagger(x)$ as $q \rightarrow +\infty$ uniformly in $x \in [a, b]$, that is:

$$\forall \epsilon > 0, \exists Q > 0 \text{ such that } \forall q \geq Q, \forall x \in [a, b], |G(x, q) - F^\dagger(x)| \leq \epsilon.$$

Moreover, we can guarantee that:

$$\forall x \in [a, b], \forall q \in \mathbb{R}_+, G(x, q) \leq F^\dagger(x). \quad (3.6)$$

That is, the optimisation algorithm will always underestimate the true value $F^\dagger(x)$. In this way, one can model the relationship between the numerical estimate G and the true value F^\dagger as:

$$\forall x \in [a, b], \forall q \in \mathbb{R}_+, G(x, q) = F^\dagger(x) - \epsilon(x, q), \quad (3.7)$$

where ϵ is a positive random variable. These assumptions imply some robustness and stability of the algorithm we use.

In the following, we will assume that $I^{(0)} \geq 2$. That is, we have at least two data points at the beginning of the reconstruction algorithm. Also among these points, we have one point at $x = a$ and another one at $x = b$. Moreover, we will assume that the initial dataset is consistent. Since [Algorithm 1](#) recomputes the inconsistent points at all steps, we can also consider in the following that any new numerical observation is actually consistent. Also, we need to guarantee that the weighted area $\text{WA}^{(n)}$ will permanently oscillate about \mathcal{E} as the iteration step n is increasing; this is the purpose of [Assumption 3.3](#) below as shown in the subsequent [Proposition 3.5](#). From these properties it will then be shown that [Algorithm 1](#) is convergent, as stated in [Theorem 3.8](#).

Assumption 3.2. Any new numerical value obtained by [Algorithm 1](#) is consistent.

Assumption 3.3. $q_-^{(n)} \rightarrow +\infty$ as $n \rightarrow \infty$.

Within [Assumption 3.2](#) all points have a consistency of 1, and therefore $q = r > 0$ the reliability. Besides, one has $G(x_i^{(n)}, q_i^{(n)}) \leq G(x_{i+1}^{(n)}, q_{i+1}^{(n)})$, that is, $y_i^{(n)} \leq y_{i+1}^{(n)}$ for all points i and steps n . We finally define the sequence of piecewise constant reconstruction functions $F^{(n)}$ as follows.

Definition 3.4. For each $x \in [a, b]$, we define the reconstructing function $F^{(n)}$ at step n as:

$$F^{(n)}(x) = \sum_{i=1}^{I^{(n)}-1} y_i^{(n)} \mathbb{1}_{s_i^{(n)}}(x),$$

and $F^{(n)}(x_{I^{(n)}}^{(n)}) = F^{(n)}(b) = y_{I^{(n)}}^{(n)}$.

Now let

$$E^+ := \{n \in \mathbb{N}; \text{WA}^{(n)} \geq \mathcal{E}\}, \quad E^- := \{n \in \mathbb{N}; \text{WA}^{(n)} < \mathcal{E}\}, \quad (3.8)$$

which are such that $E^+ \cup E^- = \mathbb{N}$ and $E^+ \cap E^- = \emptyset$. In order to prove the convergence (in a sense to be given) of [Algorithm 1](#), we first need to establish the following intermediate results, [Proposition 3.5](#), [Proposition 3.6](#), and [Proposition 3.7](#). They clarify the behaviour of the sequence $\text{WA}^{(n)}$ when points are added to the dataset and the largest area $a_+^{(n)}$ is divided into four parts at each iteration step n ; see [Figure 3.1](#).

Proposition 3.5. E^+ is infinite.

Proof. Let us assume that E^+ is finite: $\exists N$ such that $\forall n \geq N, n \in E^-$. Therefore we are in the situation $\text{WA}^{(n)} < \mathcal{E}$, the minimum quality $q_{-}^{(n)}$ of the data goes to infinity, and the total area $A^{(n)}$ is modified although the evaluation points $\{x_i^{(n)}\}_{i=1}^{I^{(n)}}$ and their number $I^{(n)}$ are unchanged; thus they are independent of n . Repeating this step yields

$$\lim_{n \rightarrow \infty} A^{(n)} = \sum_{i=1}^{I-1} (x_{i+1} - x_i)(F^\dagger(x_{i+1}) - F^\dagger(x_i)) = A > 0$$

since F^\dagger is monotonically increasing and non-constant on $[a, b]$, and [Assumption 3.1](#) is used. Consequently $\text{WA}^{(n)} \rightarrow +\infty$ as $n \rightarrow \infty$, that is $\text{WA}^{(n)} \geq \mathcal{E} \forall n \geq N_1$ for some N_1 , which is a contradiction. \blacksquare

The set E^+ is therefore of the form

$$E^+ = \bigcup_{k \geq 1} \llbracket m_k, n_k \rrbracket,$$

where

$$\llbracket m_k, n_k \rrbracket := \{n \in \mathbb{N}; m_k \leq n \leq n_k\}.$$

Let us introduce the strictly increasing application $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ such that $\varphi(p)$ is the p^{th} element of E^+ (in increasing order), and $\llbracket m_k, n_k \rrbracket = \varphi(\llbracket p_k + 1, p_{k+1} \rrbracket)$. p is the counter of the elements of E^+ , and n is the corresponding iteration number.

Proposition 3.6. Let $I^{(\varphi(p))} = I^{(\varphi(0))} + p$. Then

$$A^{(\varphi(p))} = \sum_{i=1}^{I^{(\varphi(p))}-1} a_i^{(\varphi(p))} = O\left(\frac{1}{\sqrt{p}}\right)$$

as $p \rightarrow \infty$, and $A^{(n)} \rightarrow 0$ as $n \rightarrow 0$.

Proof. Let $k \geq 1$ and $n = \varphi(p) \in \llbracket m_k, n_k \rrbracket$, where $p \in \llbracket p_k + 1, p_{k+1} \rrbracket$. Let $A^{(n)}$ be given by [Equation \(3.2\)](#), $a_+^{(n)}$ be given by [Equation \(3.3\)](#), and $i_+^{(n)}$ be given by [Equation \(3.4\)](#). At iteration $n + 1$ one has:

$$x_i^{(n+1)} = \begin{cases} x_i^{(n)} & \text{for } 1 \leq i \leq i_+^{(n)}, \\ \frac{1}{2} \left(x_{i_+^{(n)}}^{(n)} + x_{i_+^{(n)}+1}^{(n)} \right) & \text{for } i = i_+^{(n)} + 1, \\ x_{i-1}^{(n)} & \text{for } i_+^{(n)} + 2 \leq i \leq I^{(n+1)}. \end{cases}$$

Also $y_i^{(n+1)} \leq y_{i+1}^{(n+1)}$ for $1 \leq i \leq I^{(n+1)} - 1$. One may check that $a_+^{(n)} = 2a_{i_+^{(n)}}^{(n+1)} + 2a_{i_+^{(n)}+1}^{(n+1)}$ (see [Figure 3.1](#)) and therefore:

$$A^{(n+1)} = A^{(n)} - a_+^{(n)} + a_{i_+^{(n)}}^{(n+1)} + a_{i_+^{(n)}+1}^{(n+1)} = A^{(n)} - \frac{1}{2}a_+^{(n)}. \quad (3.9)$$

Besides $A^{(n)} \leq (I^{(n)} - 1)a_+^{(n)}$ so that one has:

$$\begin{aligned} A^{(n+1)} &\leq A^{(n)} - \frac{A^{(n)}}{2(I^{(n)} - 1)} \\ &\leq A^{(n)} \left(\frac{2(I^{(n)} - 1) - 1}{2(I^{(n)} - 1)} \right), \end{aligned}$$

or:

$$A^{(\varphi(p)+1)} \leq A^{(\varphi(p))} \left(\frac{2(I^{(\varphi(p))} - 1) - 1}{2(I^{(\varphi(p))} - 1)} \right). \quad (3.10)$$

At this stage two situations arise:




Figure 3.1: New area when one adds a point at the middle of the biggest rectangle.

- either $p \in \llbracket p_k + 1, p_{k+1} - 1 \rrbracket$, in which case $\varphi(p) + 1 = \varphi(p + 1)$;
- or $p = p_{k+1}$, in which case by our algorithm $A^{(n)}$ is kept constant from $n = n_k + 1$ to $n = m_{k+1}$; that is $A^{(n_k+1)} = A^{(m_{k+1})}$, or:

$$A^{(\varphi(p_{k+1})+1)} = A^{(\varphi(p_{k+1}+1))}.$$

The choice of k being arbitrary, one concludes that Equation (3.10) also reads $\forall p \in \mathbb{N}$:

$$\begin{aligned} A^{(\varphi(p+1))} &\leq A^{(\varphi(p))} \left(\frac{2(I^{(\varphi(p))}) - 1}{2(I^{(\varphi(p))}) - 1} \right) \\ &\leq A^{(\varphi(p))} \left(\frac{2(I^{(\varphi(0))}) + p - 1}{2(I^{(\varphi(0))}) + p - 1} \right). \end{aligned}$$

Thus:

$$\begin{aligned} A^{(\varphi(p))} &\leq A^{(\varphi(1))} \prod_{i=1}^{p-1} \left(\frac{2(I^{(\varphi(0))}) + i - 1}{2(I^{(\varphi(0))}) + i - 1} \right) \\ &\leq A^{(\varphi(1))} \prod_{i=1}^{p-1} \left(1 + \frac{\alpha}{i} \right), \end{aligned}$$

letting $\alpha = I^{(\varphi(0))} - \frac{3}{2}$ and $\beta = I^{(\varphi(0))} - 1$. However,

$$\sum_{i=1}^p \log \left(1 + \frac{\alpha}{i} \right) = \alpha \sum_{i=1}^p \frac{1}{i} + C_p''$$

where $\lim_{p \rightarrow \infty} C_p'' = C''$, and

$$\sum_{i=1}^p \frac{1}{i} = \log p + \gamma + \epsilon_p',$$

where γ is the Euler constant and $\lim_{p \rightarrow \infty} \epsilon_p' = 0$. Consequently:

$$\begin{aligned} \sum_{i=1}^{p-1} \log \left(1 + \frac{\alpha}{i} \right) - \sum_{i=1}^{p-1} \log \left(1 + \frac{\beta}{i} \right) &= (\alpha - \beta) \log(p-1) + C_p' \\ &= (\alpha - \beta) \left[\log p + \log \left(1 - \frac{1}{p} \right) \right] + C_p' \\ &= \log \left(\frac{1}{\sqrt{p}} \right) + C_p, \end{aligned}$$

since $\alpha - \beta = -\frac{1}{2}$; again C_p and C'_p are sequences with constant limits $\lim_{p \rightarrow \infty} C_p = C$ and $\lim_{p \rightarrow \infty} C'_p = C'$. Therefore,

$$\prod_{i=1}^{p-1} \left(\frac{1 + \frac{\alpha}{i}}{1 + \frac{\beta}{i}} \right) = \frac{C}{\sqrt{p}} (1 + \epsilon_p)$$

where C is a constant, and $\lim_{p \rightarrow \infty} \epsilon_p = 0$. One also concludes that $A^{(n)}$, which is either kept constant or equal to $A^{(\varphi(p))}$, converges to 0 as $n \rightarrow \infty$. Hence the claimed results hold. \blacksquare

Proposition 3.7. E^- is infinite.

Proof. Let us assume that E^- is finite: $\exists N$ such that $\forall n \geq N, n \in E^+$. Therefore we are in the situation $WA^{(n)} \geq \mathcal{E} > 0$, and $\varphi(n)$ has the form $\varphi(n) = n - n_0, n \geq N$ for some $n_0 \in \mathbb{N}$. From [Proposition 3.6](#):

$$A^{(n-n_0)} = O\left(\frac{1}{\sqrt{n}}\right),$$

thus $A^{(n)} \rightarrow 0$ and $WA^{(n)} \rightarrow 0$ as $n \rightarrow \infty$ since $q_-^{(n)}$ is kept unchanged, which is a contradiction. \blacksquare

We now provide three results on the convergence of [Algorithm 1](#). As is to be expected, the algorithm can only be shown to converge uniformly when the target response function F^\dagger is sufficiently smooth; otherwise, the convergence is at best pointwise or in mean.

Theorem 3.8 (Algorithm convergence). *Assume that F^\dagger is strictly increasing. Then, for any choice of $\mathcal{E} > 0$, [Algorithm 1](#) is convergent in the following senses:*

- If F^\dagger is piecewise continuous on $[a, b]$, then $\lim_{n \rightarrow \infty} F^{(n)}(x) = F^\dagger(x)$ at all points $x \in [a, b]$ where F^\dagger is continuous;
- If F^\dagger is continuous on $[a, b]$, then convergence holds uniformly: $\|F^{(n)} - F^\dagger\|_\infty \xrightarrow[n \rightarrow \infty]{} 0$.

Proof. Let $\mathcal{E} > 0$. We know from [Propositions 3.5](#) and [3.7](#) that $WA^{(n)}$ will oscillate about \mathcal{E} in the iterating process as $n \rightarrow \infty$, while $\lim_{n \rightarrow \infty} q_-^{(n)} = +\infty$ from [Assumption 3.3](#). Furthermore, let

$$\Delta^{(n)} := \sup_{1 \leq i \leq I^{(n)} - 1} \left| x_{i+1}^{(n)} - x_i^{(n)} \right|.$$

Assuming for example that for some j , $s_j^{(n)} = [x_j^{(n)}, x_{j+1}^{(n)})$ is never divided in two in the iteration process and is thus independent of n , it turns out that $a_j^{(n)} \rightarrow (x_{j+1} - x_j)(F^\dagger(x_{j+1}) - F^\dagger(x_j)) > 0$ as $n \rightarrow \infty$, which is impossible because $A^{(n)}$ goes to 0 as $n \rightarrow \infty$ from [Proposition 3.6](#). Therefore there exists some $m \in \mathbb{N}^*$ (depending on n) such that $\Delta^{(n+m)} \leq \frac{1}{2} \Delta^{(n)}$; also the sequence $\Delta^{(n)}$ is decreasing, hence $\Delta^{(n)} \rightarrow 0$ as $n \rightarrow \infty$. Now let $x \in [x_i^{(n)}, x_{i+1}^{(n)})$. Then:

$$\begin{aligned} \left| F^{(n)}(x) - F^\dagger(x) \right| &= \left| G(x_i^{(n)}, q_i^{(n)}) - F^\dagger(x) \right| \\ &\leq \left| G(x_i^{(n)}, q_i^{(n)}) - F^\dagger(x_i^{(n)}) \right| + \left| F^\dagger(x_i^{(n)}) - F^\dagger(x) \right|. \end{aligned}$$

But $x_i^{(n)} \rightarrow x$ as $n \rightarrow \infty$ because $\Delta^{(n)} \rightarrow 0$; thus if F^\dagger is continuous at x , the second term on the right hand side above goes to 0 as $n \rightarrow \infty$. However, if F^\dagger is continuous everywhere on $[a, b]$, it is in addition uniformly continuous on $[a, b]$ by Heine's theorem, and the second term goes to 0 as $n \rightarrow \infty$ uniformly on $[a, b]$. Finally, invoking [Assumption 3.1](#), the first term on the right hand side above also tends to 0 as $n \rightarrow \infty$. This completes the proof. \blacksquare

Proposition 3.9 (Convergence in mean). *Let $F^\dagger: [a, b] \rightarrow \mathbb{R}$ be piecewise continuous. Then [Algorithm 1](#) is convergent in mean in the sense that*

$$\|F^{(n)} - F^\dagger\|_1 \xrightarrow[n \rightarrow \infty]{} 0.$$

Proof. We can check that the sequence $F^{(n)}$ is monotone. Indeed, if $WA^{(n)} < \mathcal{E}$, then by construction we have

$$F^{(n+1)}(x) - F^{(n)}(x) \geq \left(y_{i_-}^{(n+1)} - y_{i_-}^{(n)} \right) \mathbb{1}_{s_-^{(n)}}(x) \geq 0$$

where $s_-^{(n)} = [x_{i_-^{(n)}}^{(n)}, x_{i_-^{(n)}+1}^{(n)}]$. However, if $\text{WA}^{(n)} > \mathcal{E}$, then consistency implies that

$$F^{(n+1)}(x) - F^{(n)}(x) \geq \left(y_{i_+^{(n)}+1}^{(n+1)} - y_{i_+^{(n)}}^{(n)} \right) \mathbb{1}_{s_+^{(n+1)}}(x) \geq 0$$

where $s_+^{(n+1)} = [x_{i_+^{(n)}+1}^{(n+1)}, x_{i_+^{(n)}+2}^{(n+1)}]$. The claim now follows from the monotone convergence theorem and the fact that $F^{(0)}$ is integrable. \blacksquare

4 Test cases

To show the effectiveness of [Algorithm 1](#), we try it on two cases, in which F^\dagger is a continuous function and a discontinuous function respectively. For both cases, the error between the numerical estimate and the ground truth function is modelled as a random variable following a Log-normal distribution. That is,

$$\forall x \in [a, b], \epsilon(x) \sim \text{Log}\mathcal{N}(\mu(x), \sigma^2), \quad (4.1)$$

with $\sigma^2 = 1$ and $\mu(x)$ is chosen as $\mathbb{P}[0 \leq \epsilon(x) \leq 0.1 \cdot F^\dagger(x)] = 0.9$. Thus, the mean μ is different for each $x \in [a, b]$.

As we have access to the ground truth function and for validation purpose, the quality value associated to a numerical point is the inverse of the relative error. Moreover, we assume that the initial points are consistent.

For illustrative purposes, we set the parameter $\mathcal{E} = 15$ for the examples considered below.

4.1 F^\dagger is a continuous function

First, consider the function $F^\dagger \in C^0([1, 2], [1, 2])$ defined as follows:

$$F^\dagger(x) = \begin{cases} F_1^\dagger(x) & \text{if } x \in [1, \frac{3}{2}], \\ F_2^\dagger(x) & \text{if } x \in [\frac{3}{2}, 2], \end{cases}$$

with

$$\begin{aligned} F_1^\dagger(x) &= a_1 \exp(x^3) + b_1, \\ F_2^\dagger(x) &= a_2 \exp((3-x)^3) + b_2, \end{aligned} \quad (4.2)$$

where:

$$a_1 = -\frac{1}{2(\exp(1) - \exp(27/8))}, \quad b_1 = \frac{3 - 2\exp(19/8)}{2(1 - \exp(19/8))}, \quad a_2 = -a_1, \quad b_2 = 2a_1 \exp(27/8) + b_1.$$

The target function F^\dagger and the reconstructions $F^{(n)}$ obtained through the algorithm for several values of the step n are shown on [Figure 4.1](#). For each n , the reconstruction $F^{(n)}$ is increasing and the initial points are consistent. The ∞ -norm and 1-norm of the error appear to converge to zero with approximate rates -0.512 and -0.534 respectively.

4.2 F^\dagger is a discontinuous function

Now, consider the discontinuous function F^\dagger defined as follows:

$$F^\dagger(x) = \begin{cases} F_1^\dagger & \text{if } x \in [1, \frac{3}{2}], \\ F_2^\dagger & \text{if } x \in (\frac{3}{2}, 2], \end{cases}$$

where F_1^\dagger and F_2^\dagger are given by [\(4.2\)](#), and:

$$\begin{aligned} a_1 &= -\frac{1}{2(\exp(1) - \exp(27/8))}, & b_1 &= \frac{3 - 2\exp(19/8)}{2(1 - \exp(19/8))}, \\ a_2 &= \frac{2}{5(\exp(8) - \exp(27/8))}, & b_2 &= \frac{10 - 8\exp(37/8)}{5(1 - \exp(37/8))}. \end{aligned}$$




Figure 4.1: Evolution of $F^{(n)}$ and the ∞ - and 1-norms of the error $F^\dagger - F^{(n)}$ as functions of the iteration count, n , for a smooth ground truth F^\dagger .

Here, F^\dagger is piecewise continuous on $[1, \frac{3}{2}]$ and $]\frac{3}{2}, 2]$. In this case, one can apply [Proposition 3.9](#). The target function F^\dagger and the reconstructions $F^{(n)}$ obtained through the algorithm for several values of the step n are shown on [Figure 4.2](#). Observe that the approximation quality, as measured by the ∞ -norm of the error $F^\dagger - F^{(n)}$, quite rapidly saturates and does not converge to zero. This is to be expected for this discontinuous target F^\dagger , since closeness of two functions in the supremum norm mandates that they have approximately the same discontinuities in exactly the same places. The 1-norm error, in contrast, appears to converge at the rate -0.561 .

Regarding computational cost, the number of calls to the numerical model is lower when F^\dagger is continuous than when it is discontinuous. For both examples above and for the same number of data points, the number of evaluations of the numerical model (analytical formula in the present case) in the discontinuous case is about six times higher than the number of evaluations in the continuous case. This is because the algorithm typically adds more points near discontinuities and the effort of making them consistent increases the number of calls to the model.

4.3 Influence of the user-defined parameter \mathcal{E}

We consider the case in which F^\dagger is discontinuous, as in [Section 4.2](#). We will show the influence of the choice of the parameter \mathcal{E} on the reconstruction function $F^{(n)}$.

4.3.1 Case $\mathcal{E} \ll 1$

Let us consider the case $\mathcal{E} = 10^{-4} \ll 1$. This choice corresponds to the case where one wishes to split over and over the worst quality point. This can be seen on [Figure 4.3](#) where the worst quality is almost constant over 100 steps while the sum of areas strongly decreases; see [Figure 4.3\(e\)](#) and [Figure 4.3\(f\)](#) respectively. At each step, the algorithm is adding a new point by splitting the biggest rectangle. One can note on [Figure 4.3\(f\)](#) that the minimum of the quality is not constant. It means that when the




Figure 4.2: Evolution of $F^{(n)}$ and the ∞ - and 1-norms of the error $F^\dagger - F^{(n)}$ as functions of the iteration count, n , for a discontinuous ground truth F^\dagger .

algorithm added a new data point, the point with the worst quality was not consistent any more and had to be recomputed. In summary, in this case, we obtain more points but with lower quality values.

4.3.2 Case $\mathcal{E} \gg 1$

We now consider the case $\mathcal{E} = 10^4 \gg 1$. This choice corresponds to the case where one wishes to redo the worst quality point over split. This can be seen on [Figure 4.4](#) where the sum of areas stays more or less the same over 100 steps while the minimum of the quality surges; see [Figure 4.4\(f\)](#) and [Figure 4.4\(e\)](#) respectively. There is no new point. The algorithm is only redoing the worst quality point to improve it. To sum up, we obtain fewer points with higher quality values.

5 Application to optimal uncertainty quantification

5.1 Optimal uncertainty quantification

In the *optimal uncertainty quantification* paradigm proposed by [Owhadi et al. \(2013\)](#) and further developed by, e.g., [Sullivan et al. \(2013\)](#) and [Han et al. \(2015\)](#), upper and lower bounds on the performance of an incompletely-specified system are calculated via optimisation problems. More concretely, one is interested in the probability that a system, whose output is a function $g^\dagger: \mathcal{X} \rightarrow \mathbb{R}$ of inputs Ξ distributed according to a probability measure μ^\dagger on an input space \mathcal{X} , satisfies $g^\dagger(\Xi) \leq x$, where x is a specified performance threshold value. We emphasise that although we focus on a scalar performance measure, the input Ξ may be a multivariate random variable.

In practice, μ^\dagger and g^\dagger are not known exactly; rather, it is known only that $(\mu^\dagger, g^\dagger) \in \mathcal{A}$ for some admissible subset \mathcal{A} of the product space of all probability measures on \mathcal{X} with the set of all real-valued




Figure 4.3: Evolution of $F^{(n)}$ and the minimum of the quality and the total area as functions of the iteration count, n , for a discontinuous ground truth F^\dagger with $\mathcal{E} = 10^{-4}$.

functions on \mathcal{X} . Thus, one is interested in

$$\underline{P}_{\mathcal{A}}(x) := \inf_{(\mu, g) \in \mathcal{A}} \mathbb{P}_{\Xi \sim \mu} [g(\Xi) \leq x] \quad \text{and} \quad \overline{P}_{\mathcal{A}}(x) := \sup_{(\mu, g) \in \mathcal{A}} \mathbb{P}_{\Xi \sim \mu} [g(\Xi) \leq x].$$

The inequality

$$0 \leq \underline{P}_{\mathcal{A}}(x) \leq \mathbb{P}_{\Xi \sim \mu^\dagger} [g^\dagger(\Xi) \leq x] \leq \overline{P}_{\mathcal{A}}(x) \leq 1$$

is, by definition, the tightest possible bound on the quantity of interest $\mathbb{P}_{\Xi \sim \mu^\dagger} [g^\dagger(\Xi) \leq x]$ that is compatible with the information used to specify \mathcal{A} . Thus, the optimal UQ perspective enriches the principles of worst- and best-case design to account for distributional and functional uncertainty. We concentrate our attention hereafter, without loss of generality, on the least upper bound $\overline{P}_{\mathcal{A}}(x)$.

Remark 5.1. The main focus of this paper is the dependency of $\overline{P}_{\mathcal{A}}(x)$ on x . In practice, an underlying task is, for any individual x , reducing the calculation of $\overline{P}_{\mathcal{A}}(x)$ to a tractable finite-dimensional optimisation problem. Central enabling results here are the *reduction theorems* of [Owhadi et al. \(2013, Section 4\)](#), which, loosely speaking, say that if, for each g , $\{\mu \mid (\mu, g) \in \mathcal{A}\}$ is specified by a system of m equality or inequality constraints on expected values of arbitrary test functions under μ , then for the determination of $\overline{P}_{\mathcal{A}}(x)$ it is sufficient to consider only distributions μ that are convex combinations of at most $m + 1$ point masses; the optimisation variables are then the m independent weights and $m + 1$ locations in \mathcal{X} of these point masses. If μ factors as a product of distributions (i.e. Ξ is a vector with independent components), then this reduction theorem applies componentwise.

As a function of the performance threshold x , $\overline{P}_{\mathcal{A}}(x)$ is an increasing function, and so it is potentially advantageous to determine $\overline{P}_{\mathcal{A}}(x)$ jointly for a wide range of x values using the algorithm developed above. Indeed, determining $\overline{P}_{\mathcal{A}}(x)$ for many values of x , rather than just one value, is desirable for multiple reasons:




Figure 4.4: Evolution of $F^{(n)}$ and the minimum of the quality and the total area as functions of the iteration count, n , for a discontinuous ground truth F^\dagger with $\mathcal{E} = 10^4$.

1. Since numerical optimisation to determine $\bar{P}_{\mathcal{A}}(x)$ may be affected by errors, computing several values of $\bar{P}_{\mathcal{A}}(x)$ could lead to validate their consistency as the function $x \mapsto \bar{P}_{\mathcal{A}}(x)$ must be increasing;
2. The function $\bar{P}_{\mathcal{A}}(x)$ can be discontinuous. Thus, by computing several values of $\bar{P}_{\mathcal{A}}(x)$, one can highlight potential discontinuities and can identify key threshold values of $x \mapsto \bar{P}_{\mathcal{A}}(x)$.

5.2 Test case

For the application of [Algorithm 1](#) to OUQ, we study the robust shape optimization of the two-dimensional RAE2822 airfoil ([Cook et al., 1979](#), Appendix A6) using ONERA's CFD software *elsA* ([Cambier et al., 2013](#)). The following example is taken from [Dumont et al. \(2019\)](#). The shape of the original RAE2822 is altered using four bumps located at four different locations: 5%, 20%, 40%, and 60% of the way along the chord c (see [Figure 5.1](#)). These bumps are characterised by B-splines functions.

The lift-to-drag ratio $\frac{C_l}{C_d}$ of the RAE2822 wing profile (see [Figure 5.2](#)) at Reynolds Number $Re = 6.5 \cdot 10^6$, Mach number $M_\infty = 0.729$ and angle of attack $\alpha = 2.31^\circ$ is chosen as the performance function g^\dagger with inputs $\Xi = (\Xi_1, \Xi_2, \Xi_3, \Xi_4)$, where $(\Xi_i)_{i=1\dots 4}$ is the amplitude of each bump. They will be considered as random variables over their respective range given in [Table 5.1](#).

The corresponding flow values are the ones described in test case #6 together with the wall interferences corrections formulas given in [Garner et al. \(1966, Chapter 6\)](#) and in [Haase et al. \(1993, Section 5.1\)](#). Moreover, we will assume that $(\Xi_i)_{i=1\dots 4}$ are mutually independent. An ordinary Kriging procedure has been chosen to build a metamodel (or response surface) of g^\dagger , which is identified with the actual response function g^\dagger in the subsequent analysis. A tensorised grid of 9 equidistributed abscissas for each parameter is used. The model is then based on $N = 9^4 = 6561$ observations. In that respect, a Gaussian




Figure 5.1: Black lines: Maximum and minimum deformation of the RAE2822 profile. Red: Maximum deformation of the third bump alone. Blue: Minimum deformation of the third bump alone. This image is taken from [Dumont et al. \(2019\)](#).




Figure 5.2: Picture depicting the lift C_l and the drag C_d of an airfoil.

kernel

$$K(\Xi, \Xi') = \exp\left(-\frac{1}{2} \sum_{i=1}^4 \frac{(\Xi_i - \Xi'_i)^2}{\gamma_i^2}\right)$$

has been chosen, where $\Xi = (\Xi_1, \Xi_2, \Xi_3, \Xi_4)$ and $\Xi' = (\Xi'_1, \Xi'_2, \Xi'_3, \Xi'_4)$ are inputs of the function g^\dagger , and where $\gamma = (\gamma_1, \gamma_2, \gamma_3, \gamma_4)$ are the parameters of the kernel. These parameters are chosen to minimize the variance between the ground truth data defined by the N observations and their Kriging metamodel g^\dagger . The response surfaces in the (Ξ_1, Ξ_3) plan for two values of (Ξ_2, Ξ_4) are shown on [Figure 5.3](#).


One seeks to determine $\bar{P}_{\mathcal{A}}(x) := \sup_{\mu \in \mathcal{A}} \mathbb{P}_{\Xi \sim \mu}[g^\dagger(\Xi) \leq x]$, where the admissible set \mathcal{A} is defined as follows:

$$\mathcal{A} = \left\{ (g, \mu) \left| \begin{array}{l} \Xi \in \mathcal{X} = \mathcal{X}_1 \times \mathcal{X}_2 \times \mathcal{X}_3 \times \mathcal{X}_4 \\ g : \mathcal{X} \mapsto \mathcal{Y} \text{ is known equal to } g^\dagger \\ \mu = \mu_1 \otimes \mu_2 \otimes \mu_3 \otimes \mu_4 \\ \mathbb{E}_{\Xi \sim \mu}[g(\Xi)] = \text{LD} \end{array} \right. \right\}. \quad (5.1)$$

A priori, finding $\bar{P}_{\mathcal{A}}(x)$ is not computationally tractable because it requires a search over a infinite-dimensional space of probability measures defined by \mathcal{A} . Nevertheless, as described briefly in [Remark 5.1](#), it has been shown in [Owhadi et al. \(2013\)](#) that this optimisation problem can be reduced to a finite-dimensional one, where now the probability measures are products of finite convex combinations of Dirac masses.

	Range	Law
Bump 1: Ξ_1	$[-0.0025c; +0.0025c]$	μ_1^\dagger : Beta law with $\alpha = 6, \beta = 6$
Bump 2: Ξ_2	$[-0.0025c; +0.0025c]$	μ_2^\dagger : Beta law with $\alpha = 2, \beta = 2$
Bump 3: Ξ_3	$[-0.0025c; +0.0025c]$	μ_3^\dagger : Beta law with $\alpha = 2, \beta = 2$
Bump 4: Ξ_4	$[-0.0025c; +0.0025c]$	μ_4^\dagger : Beta law with $\alpha = 2, \beta = 2$

Table 5.1: Range of each input parameter.


Figure 5.3: Response surface in the (Ξ_1, Ξ_3) plane with $(\Xi_2 = -0.0025, \Xi_4 = 0)$ (a) and $(\Xi_2 = 0.0025, \Xi_4 = 0)$ (b). These images are taken from [Dumont et al. \(2019\)](#).


Remark 5.2. The ground truth law μ^\dagger of each input variable given in [Table 5.1](#) is only used to compute the expected value $\mathbb{E}_{\Xi \sim \mu}[g(\Xi)] = \text{LD}$. This expected value is computed with 10^4 samples.

Remark 5.3. The admissible set \mathcal{A} from (5.1) can be understood as follows:


- One knows the range of each input parameter $(\Xi_i)_{i=1, \dots, 4}$;
- g is exactly known as $g = g^\dagger$;
- $(\Xi_i)_{i=1, \dots, 4}$ are independent;
- One only knows the expected value of g : $\mathbb{E}_{\Xi \sim \mu}[g(\Xi)]$.

The optimisation problem of determining $\overline{P}_{\mathcal{A}}(x)$ for each chosen x was solved using the Differential Evolution algorithm of [Storn and Price \(1997\)](#) within the *mystic* optimisation framework ([McKerns et al., 2011](#)). Ten iterations of [Algorithm 1](#) have been performed using $\mathcal{E} = 1 \times 10^4$. The evolution of $\overline{P}_{\mathcal{A}}(x)$ as function of the iteration count, n , is shown on [Figure 5.4](#). At $n = 0$ — see [Figure 5.4\(a\)](#) — two consistent points are present at $x = 57.51$ and $x = 67.51$. At this step, $\text{WA}^{(0)} = 35289$. As $\text{WA}^{(0)} \geq \mathcal{E}$, at next step $n = 1$, the algorithm adds a new point at the middle of the biggest rectangle — see [Figure 5.4\(b\)](#) and [Figure 5.5\(b\)](#). After $n = 10$ steps, eight points are now present in total with a minimum quality increasing from 5000 to 11667 and with a total area decreasing from 7.05 to 0.84; see [Figure 5.5\(a\)](#) and [Figure 5.5\(b\)](#) respectively.


The number of iterations in this complex numerical experiment has been limited to 10 because obtaining new or improved data points consistent throughout the optimization algorithm may take up to two days (wall-clock time on a personal computer equipped with an Intel Core i5-6300HQ processor with 4 cores and 6 MB cache memory) for one single point. This running time is increased further for data points of higher quality. Nevertheless, this experiment shows that the proposed algorithm can be used for real-world examples in an industrial context.




(a) $n = 0$



(b) $n = 1$




(c) $n = 5$




(d) $n = 10$

Figure 5.4: Evolution of $\bar{P}_{\mathcal{A}}(x)$ as function of the iteration count, n .



(a) Evolution of the minimum of the quality.



(b) Evolution of the total area.

Figure 5.5: Evolution of the minimum of the quality and the total area as function of the iteration count, n .

6 Concluding remarks

In this paper we have developed an algorithm to reconstruct a monotonically increasing function such as the cumulative distribution function of a real-valued random variable, or the least upper bound of the performance criterion of a system as a function of its performance threshold. In particular, this latter setting has relevance to the optimal uncertainty quantification (OUQ) framework of [Owhadi et al. \(2013\)](#) we have in mind for applications to real-world incompletely specified systems. The algorithm uses imperfect pointwise evaluations of the target function, subject to partially controllable one-sided errors, to direct further evaluations either at new sites in the function’s domain or to improve the quality of evaluations at already-evaluated sites. It allows for some flexibility at targeting either strategy through a user-defined “exchange rate” parameter, yielding an approximation of the target function with few high-quality points or alternatively more lower-quality points. We have studied its convergence properties and have applied it to several examples: known target functions that are either continuous and discontinuous, and a performance function for aerodynamic design of a well-documented standard profile in the OUQ setting.

[Algorithm 1](#) is reminiscent of the classical PAVA approach to isotonic regression that applies to statistical inference with order restrictions. Examples of its use can be found in shape constrained or parametric density problems as illustrated in e.g. [Groeneboom and Jongbloed \(2014\)](#). Possible improvements and extensions of our algorithm include weighting the areas $a_i^{(n)}$ as they are summed up to form the total weighted area $WA^{(n)}$ driving the iterative process, in order to optimally enforce both the addition of “steps” $s_i^{(n)}$ in the reconstruction function $F^{(n)}$ of [Definition 3.4](#), and the improvement of their “heights” $y_i^{(n)}$. This could be achieved considering for example the following alternative definition $i_+^{(n)} = \arg \max_i \{(I^{(n)} - i - 1)a_i^{(n)}\}$ in [Algorithm 1](#), which results in both adding a step to the $i_+^{(n)}$ -th current one and possibly improving all subsequent evaluations $y_i^{(n+1)}$, $i > i_+^{(n)}$. We may further envisage to adapt the ideas elaborated in this research to the reconstruction of convex functions by extending the notion of consistency. These perspectives shall be considered in future works.

Acknowledgements

The work of J.L.A. and É.S. has been partially supported by ONERA within the Laboratoire de Mathématiques Appliquées pour l’Aéronautique et Spatial (LMA²S). L.B. is supported by a CDSN grant from the French Ministry of Higher Education (MESRI) and a grant from the German Academic Exchange Service (DAAD), Program #57442045. T.J.S. has been partially supported by the Freie Universität Berlin within the Excellence Strategy of the DFG, including project TrU-2 of the Excellence Cluster “MATH+ The Berlin Mathematics Research Center” (EXC-2046/1, project 390685689) and DFG project 415980428.

References

- R. E. Barlow, D. J. Bartholomew, J. M. Bremner, and H. D. Brunk. *Statistical Interference under Order Restrictions. The Theory and Application of Isotonic Regression*. John Wiley & Sons, London-New York-Sydney, 1972.
- L. Cambier, S. Heib, and S. Plot. The ONERA elsA CFD software: input from research and feedback from industry. *Mechanics & Industry*, 14(3):159–174, 2013. doi:10.1051/meca/2013056.
- P. H. Cook, M. A. McDonald, and M. C. P. Firmin. Aerofoil RAE 2822 pressure distributions, and boundary layer and wake measurements. In *Experimental data base for computer program assessment. AGARD Advisory Report No. 138*. NATO, 1979. URL http://eda-ltd.com.tr/caeeda_doc/AGARD-AR-138.pdf.
- J. de Leeuw, K. Hornik, and P. Mair. Isotone optimization in R: Pool-Adjacent-Violators Algorithm (PAVA) and active set methods. *J. Stat. Software*, 32(5):1–24, 2009. doi:10.18637/jss.v032.i05.
- A. Dumont, J.-L. Hantrais-Gervois, P.-Y. Passaggia, J. Peter, I. Salah el Din, and É. Savin. Ordinary kriging surrogates in aerodynamics. In C. Hirsch, D. Wunsch, J. Szumbariski, L. Łaniewski-Wołk, and J. Pons-Prats, editors, *Uncertainty Management for Robust Industrial Design in Aeronautics: Findings and Best Practice Collected During UMRIDA, a Collaborative Research Project (2013–2016)*

- Funded by the European Union*, pages 229–245. Springer International Publishing, Cham, 2019. doi:[10.1007/978-3-319-77767-2_14](https://doi.org/10.1007/978-3-319-77767-2_14).
- H. C. Garner, E. W. E. Rogers, W. E. A. Acum, and E. C. Maskell. Subsonic wind tunnel jwall corrections. AGARDo-graph 109, NATO, 1966.
- P. Groeneboom and G. Jongbloed. *Nonparametric Estimation under Shape Constraints: Estimators. Algorithms and Asymptotics*. Cambridge University Press, Cambridge, 2014. doi:[10.1017/CBO9781139020893](https://doi.org/10.1017/CBO9781139020893).
- W. Haase, F. Bradsma, E. Elsholz, M. Leschziner, and D. Schwamborn. *EUROVAL - An European Initiative on Validation of CFD Codes*. Vieweg Verlag, Wiesbaden, 1993. doi:[10.1007/978-3-663-14131-0](https://doi.org/10.1007/978-3-663-14131-0).
- S. Han, M. Tao, U. Topcu, H. Owhadi, and R. M. Murray. Convex Optimal Uncertainty Quantification. *SIAM J. Optim.*, 25(3):1368–1387, 2015. doi:[10.1137/13094712X](https://doi.org/10.1137/13094712X).
- A. I. Jordan, A. Mhlemann, and J. F. Ziegel. Optimal solutions to the isotonic regression problem, 2019. URL <https://arxiv.org/abs/1904.04761>.
- M. M. McKerns, L. Strand, T. J. Sullivan, A. Fang, and M. A. G. Aivazis. Building a framework for predictive science. In S. van der Walt and J. Millman, editors, *Proceedings of the 10th Python in Science Conference (SciPy 2011), June 2011*, pages 67–78, 2011. doi:[10.25080/Majora-ebaa42b7-00d](https://doi.org/10.25080/Majora-ebaa42b7-00d).
- H. Owhadi, C. Scovel, T. J. Sullivan, M. McKerns, and M. Ortiz. Optimal Uncertainty Quantification. *SIAM Rev.*, 55(2):271–345, 2013. doi:[10.1137/10080782X](https://doi.org/10.1137/10080782X).
- T. Robertson, F. T. Wright, and R. L. Dykstra. *Order Restricted Statistical Inference*. John Wiley & Sons, London-New York-Sydney, 1988.
- R. Storn and K. Price. Differential evolution – a simple and efficient heuristic for global optimization over continuous spaces. *Journal of Global Optimization*, 11:341–359, 1997. doi:[10.1023/A:1008202821328](https://doi.org/10.1023/A:1008202821328).
- T. J. Sullivan, M. McKerns, D. Meyer, F. Theil, H. Owhadi, and M. Ortiz. Optimal Uncertainty Quantification for legacy data observations of Lipschitz functions. *ESAIM Math. Model. Numer. Anal.*, 47(6):1657–1689, 2013. doi:[10.1051/m2an/2013083](https://doi.org/10.1051/m2an/2013083).
- R. J. Tibshirani, H. Hoefling, and R. Tibshirani. Nearly-isotonic regression. *Technometrics*, 53(1):54–61, 2011. doi:[10.1198/TECH.2010.10111](https://doi.org/10.1198/TECH.2010.10111).