



HAL
open science

Boundary singular solutions of a class of equations with mixed absorption-reaction

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron

► **To cite this version:**

Marie-Françoise Bidaut-Veron, Marta Garcia-Huidobro, Laurent Veron. Boundary singular solutions of a class of equations with mixed absorption-reaction. *Calculus of Variations and Partial Differential*, 2022, 61 (113), pp.1-46. 10.1007/s00526-022-02200-z . hal-02909840v4

HAL Id: hal-02909840

<https://hal.science/hal-02909840v4>

Submitted on 23 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Boundary singular solutions of a class of equations with mixed absorption-reaction

Marie-Françoise Bidaut-Véron*
Marta Garcia-Huidobro †
Laurent Véron ‡

Abstract

We study properties of positive functions satisfying (E) $-\Delta u + u^p - M|\nabla u|^q = 0$ in a domain Ω or in \mathbb{R}_+^N when $p > 1$ and $1 < q < \min\{p, 2\}$. We concentrate our research on the solutions of (E) vanishing on the boundary except at one point. This analysis depends on the existence of separable solutions in \mathbb{R}_+^N . We construct various types of positive solutions with an isolated singularity on the boundary. We also study conditions for the removability of compact boundary sets and the Dirichlet problem associated to (E) with a measure as boundary data.

2010 Mathematics Subject Classification: 35J62-35J66-35J75-31C15

Keywords: Elliptic equations, boundary singularities, Bessel capacities, measures, supersolutions, subsolutions.

Contents

1	Introduction	1
2	Singular boundary value problems	8
2.1	A priori estimates	8
2.2	Removable singularities	13
2.3	Proof of Theorem 1.2	17
2.4	Measure boundary data	19

*Laboratoire de Mathématiques et Physique Théorique, Université de Tours, 37200 Tours, France. E-mail: veronmf@univ-tours.fr

†Departamento de Matemáticas, Pontificia Universidad Católica de Chile Casilla 307, Correo 2, Santiago de Chile. E-mail: mgarcia@mat.puc.cl

‡Laboratoire de Mathématiques et Physique Théorique, Université de Tours, 37200 Tours, France. E-mail: veronl@univ-tours.fr

3	Separable solutions	24
3.1	Existence of singular solutions	24
3.2	Existence or non-existence of separable solutions	28
3.2.1	Proof of Theorem 1.9	28
3.2.2	Non-existence	30
4	Solutions with an isolated boundary singularity	32
4.1	Uniqueness of singular solutions	32
4.2	Construction of fundamental solutions	34
4.3	Solutions with a strong singularity	40
4.3.1	The case $1 < q \leq \frac{2p}{p+1}$	40
4.3.2	The case $\frac{2p}{p+1} < q < p$	43
4.3.3	Open problems	47

1 Introduction

The aim of this article is to study some properties of solutions of the following equation

$$\mathcal{L}_{q,M}u := -\Delta u + |u|^{p-1}u - M|\nabla u|^q = 0 \tag{1.1}$$

in a bounded domain Ω of \mathbb{R}^N or in the half-space \mathbb{R}_+^N , where $M > 0$ and $p > q > 1$. We are particularly interested in the analysis of boundary singularities of such solutions. If $M = 0$ the boundary singularities problem has been investigated since thirty years, starting with the work of Gmira and Véron [15] who obtained an almost complete description of the solutions with isolated boundary singularities. When $M > 0$ there is a balance between the absorption term $|u|^{p-1}u$ and the source term $M|\nabla u|^q$, a confrontation which can create very new effects. Furthermore, the scale of the two opposed reaction terms depends upon the position of q with respect to $\frac{2p}{p+1}$. This is due to the fact that (1.1) is equivariant with respect to the scaling transformation T_ℓ defined for $\ell > 0$ by $T_\ell[u](x) = \ell^{\frac{2}{p-1}}u(\ell x)$.

If $q < \frac{2p}{p+1}$, the absorption term is dominant and the behaviour of the singular solutions is modelled by the equation studied in [15]

$$-\Delta u + |u|^{p-1}u = 0. \tag{1.2}$$

If $q > \frac{2p}{p+1}$, the source term is dominant and the behaviour of the singular solutions is modelled by positive separable solutions of the equation without diffusion

$$u^p - M|\nabla u|^q = 0. \tag{1.3}$$

Another associated equation which plays an important role in the construction of singular solutions since its positive solutions are supersolutions of (1.1) is

$$-\Delta u - M|\nabla u|^q = 0. \tag{1.4}$$

Note that in (1.3) and (1.4), M can be fixed to be 1 by replacing u by ℓu .

If $q = \frac{2p}{p+1}$, the coefficient $M > 0$ plays a fundamental role in the properties of the set of solutions, in particular for the existence of singular solutions and removable singularities. This situation is similar in some sense to what happens for equation

$$-\Delta u = |u|^{p-1}u + M|\nabla u|^q \quad (1.5)$$

which is studied thoroughly in [5], [6] in the case $M > 0$ and in [26] in the case $M < 0$. In this last article the opposition of a forcing term $|u|^{p-1}u$ and an absorption term $M|\nabla u|^q$ creates a very rich configuration of unexpected phenomena and new effects.

In the present paper we will consider the case where $1 < q < 2$, with a special emphasis on the case $q = \frac{2p}{p+1}$ which allows to put into light the role of the value of M . We first analyze the following problem: given a smooth bounded domain $\Omega \subset \mathbb{R}^N$ such that $0 \in \partial\Omega$, under what conditions involving p , q and M is the point 0 a removable singularity for a solution of (1.1) continuous in $\overline{\Omega} \setminus \{0\}$ and vanishing on $\partial\Omega \setminus \{0\}$? In the sequel we denote $\rho(x) = \text{dist}(x, \partial\Omega)$ and for $1 \leq s < \infty$, $L_\rho^s(\Omega) := L^s(\Omega; \rho dx)$ and the space of test functions in Ω is defined by

$$\mathbb{X}(\Omega) = \{\zeta \in C^1(\overline{\Omega}) : \zeta = 0 \text{ on } \partial\Omega, \Delta\zeta \in L^\infty(\Omega)\}. \quad (1.6)$$

If Ω is replaced by \mathbb{R}_+^N , then

$$\mathbb{X}(\mathbb{R}_+^N) = \{\zeta \in C^1(\overline{\mathbb{R}_+^N}) \text{ with compact support in } \overline{\mathbb{R}_+^N}, \Delta\zeta \in L^\infty(\mathbb{R}_+^N)\}. \quad (1.7)$$

Our first result is the following:

Theorem 1.1 *Assume $p \geq \frac{N+1}{N-1}$, $M > 0$ and*

(i) *either $p = \frac{N+1}{N-1}$ and $1 < q < 1 + \frac{1}{N}$.*

(ii) *or $p > \frac{N+1}{N-1}$ and $1 < q \leq \frac{2p}{p+1}$.*

Then any nonnegative solution $u \in C^2(\Omega) \cap C^1(\overline{\Omega} \setminus \{0\})$ of

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= 0 && \text{in } \partial\Omega \setminus \{0\} \end{aligned} \quad (1.8)$$

verifies $\nabla u \in L_\rho^q(\Omega)$, $u \in L_\rho^p(\Omega)$ and is a weak solution of

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= 0 && \text{in } \partial\Omega, \end{aligned} \quad (1.9)$$

in the sense that

$$\int_{\Omega} (-u\Delta\zeta + (|u|^{p-1}u - M|\nabla u|^q)\zeta) dx = 0 \quad \text{for all } \zeta \in \mathbb{X}(\Omega). \quad (1.10)$$

Furthermore, if we assume either (i), or

(iii) $p > \frac{N+1}{N-1}$ and $1 < q < \frac{2p}{p+1}$ or

(iv) $p > \frac{N+1}{N-1}$, $q = \frac{2p}{p+1}$ and

$$M < m^{**} := (p+1) \left(\frac{(N-1)p - (N+1)}{2p} \right)^{\frac{p}{p+1}}, \quad (1.11)$$

then $u = 0$.

This result is optimal in the case $p = \frac{N+1}{N-1}$, $q = \frac{2p}{p+1}$ as we will see in Section 4. Combining the method used in proving Theorem 1.1 with the result of [18] we prove the removability of compact boundary sets on $\partial\Omega$, provided they satisfy some zero Bessel capacity property.

Theorem 1.2 *Assume $p > \frac{N+1}{N-1}$ and $\frac{N+1}{N-1} < r < p$. If one of the following conditions is satisfied:*

(i) *either $q = \frac{2p}{p+1}$ and*

$$M < m_r^{**} := (p+1) \left(\frac{p-r}{p(r-1)} \right)^{\frac{p}{p+1}}, \quad (1.12)$$

(ii) *or $1 < q < \frac{2p}{p+1}$, $r \leq 3$ and M is arbitrary.*

Then if $K \subset \partial\Omega$ is a compact set such that $\text{cap}_{\frac{2}{r}, r}^{\partial\Omega}(K) = 0$, any solution u of

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega \setminus K, \end{aligned} \quad (1.13)$$

is identically 0.

Note that $m_{\frac{N+1}{N-1}}^{**} = m^{**}$. The capacity framework allows to consider the Dirichlet problem for (1.1)

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= \mu && \text{in } \partial\Omega, \end{aligned} \quad (1.14)$$

where μ is a Radon measure on $\partial\Omega$. By a weak solution of (1.14) we understand a function $u \in L^1(\Omega) \cap L^p_\rho(\Omega)$ such that $|\nabla u| \in L^q_\rho(\Omega)$, which satisfies

$$\int_{\Omega} (-u\Delta\zeta + (|u|^{p-1}u - M|\nabla u|^q)\zeta) dx = - \int_{\partial\Omega} \frac{\partial\zeta}{\partial\mathbf{n}} d\mu \quad \text{for all } \zeta \in \mathbb{X}(\Omega). \quad (1.15)$$

When the two exponents are super-critical with respect to the equations (1.2) and (1.4), the admissibility condition on the measure for (1.1) requires the introduction of two different Bessel capacities defined on Borel subsets of $\partial\Omega$.

Theorem 1.3 *Let $p > 1$, $1 < q < 2$ and μ be a nonnegative Radon measure on $\partial\Omega$ which satisfies*

$$\mu(E) \leq C \min \left\{ \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(E), \text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(E) \right\} \quad \text{for all Borel set } E \subset \partial\Omega, \quad (1.16)$$

for some $C > 0$. Then there exists $c_0 > 0$ such that for any $0 < c \leq c_0$ there exists a nonnegative weak solution of (1.14) with boundary data $c\mu$. Furthermore the boundary trace of u is the measure $c\mu$.

The proof is based upon a non-standard application of the sub and supersolutions technique since it relies of the dynamical (and more natural) aspect of the boundary trace as it is exposed in [20]. Another surprising fact is the use of the equation

$$\begin{aligned} -\Delta u &= u^p && \text{in } \Omega \\ u &= \mu && \text{in } \partial\Omega, \end{aligned}$$

which yields key estimates for our construction. The theorem admits several corollaries the proof of which is based on properties of Bessel capacities as exposed in [1]. It is noticeable that the results therein cover the full range $(p, q) \in (1, \infty) \times (1, 2)$.

Corollary 1.4 *Assume $p \geq \frac{N+1}{N-1}$ and $\frac{2p}{p+1} \leq q < 2$. If μ is a nonnegative Radon measure on $\partial\Omega$ which satisfies, for some $C > 0$,*

$$\mu(E) \leq C \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(E) \quad \text{for all Borel set } E \subset \partial\Omega, \quad (1.17)$$

there the conclusions of Theorem 1.3 hold.

The condition on the measure is also fulfilled under the following conditions.

Corollary 1.5 *Assume $\frac{N+1}{N} \leq q < \frac{2p}{p+1}$. If μ is a nonnegative Radon measure on $\partial\Omega$ such that for some constant $C > 0$, there holds for any Borel set $E \subset \partial\Omega$,*

$$\mu(E) \leq C \text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(E), \quad (1.18)$$

then the conclusions of Theorem 1.3 hold.

Since the exponents p and q can be separately super or sub-critical, or even both sub-critical, we have the following result in different configurations of exponents.

Corollary 1.6 *Let $p > 1$, $1 < q < 2$ and $\mu \in \mathfrak{M}_+(\partial\Omega)$. There exists a function $u \in L^1(\Omega) \cap L^p_\rho(\Omega)$ such that $\nabla u \in L^q_\rho(\Omega)$ which is a weak solution to (1.14) in the following cases:*

- (i) *When $p < \frac{N+1}{N-1}$, $q < \frac{N+1}{N}$ and there exists some $c_1 > 0$ such that $\|\mu\|_{\mathfrak{M}} \leq c_1$.*
- (ii) *When $p < \frac{N+1}{N-1}$, $q \geq \frac{N+1}{N}$ and μ satisfies (1.17); in that case μ has to be replaced by $c\mu$ with $0 < c \leq c_2$, for some $c_2 > 0$, in problem (1.14).*
- (iii) *When $p \geq \frac{N+1}{N-1}$, $q < \frac{N+1}{N}$, and μ satisfies $\|\mu\|_{\mathfrak{M}} \leq c_3$ for some $c_3 > 0$ and*

$$\mu(E) = 0 \quad \text{for all Borel set } E \subset \partial\Omega \text{ such that } \text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(E) = 0. \quad (1.19)$$

In [7] the same authors study the problem

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= \mu && \text{in } \Omega \\ u &= 0 && \text{in } \partial\Omega, \end{aligned} \quad (1.20)$$

where μ is a bounded Borel measure in Ω . There too sufficient conditions for solving the problem involves Bessel capacities, but since the boundary trace argument is no longer valid, an intensive utilization of potential theory with various kernels has to be used.

In the sub-critical case (i) and when μ is a Dirac mass at 0 on the boundary we have no restriction on its weight.

Theorem 1.7 *Assume $1 < p < \frac{N+1}{N-1}$ and $1 < q < \frac{N+1}{N}$. Then for any $k \geq 0$ there exists a minimal positive solution u_k of*

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \mathbb{R}_+^N \\ u &= 0 && \text{in } \partial\mathbb{R}_+^N \setminus \{0\}, \end{aligned} \quad (1.21)$$

satisfying

$$\lim_{x \rightarrow 0} \frac{u_k(x)}{P_N(x)} = k \quad (1.22)$$

where $P_N(x) = c_N x_N |x|^{-N}$ is the Poisson kernel in \mathbb{R}_+^N . Furthermore this solution is unique among the positive solutions of (1.21)-(1.22) if $q \leq \frac{2p}{p+1}$. This function satisfies $u_k \in L_{loc}^1(\overline{\mathbb{R}_+^N}) \cap L_{loc}^p(\overline{\mathbb{R}_+^N}; x_N dx)$, $\nabla u_k \in L_{loc}^q(\overline{\mathbb{R}_+^N}; x_N dx)$ and

$$\int_{\mathbb{R}_+^N} (-u_k \Delta \zeta + (u_k^p - M|\nabla u_k|^q) \zeta) dx = k \frac{\partial \zeta}{\partial x_N}(0) \quad \text{for all } \zeta \in \mathbb{X}(\mathbb{R}_+^N). \quad (1.23)$$

The proof is completely different from the ones of Theorem 1.3 and Corollary 1.6 and is based upon a delicate construction of supersolutions and subsolutions. A similar result holds if \mathbb{R}_+^N is replaced by a bounded smooth domain $\Omega \subset \mathbb{R}_+^N$ such that $0 \in \partial\Omega$.

Theorem 1.8 *Assume $1 < p < \frac{N+1}{N-1}$ and $0 < q < \frac{N+1}{N}$. Then for any $M > 0$ and $k > 0$ there exists a minimal solution $u_k \in C^1(\overline{\Omega} \setminus \{0\})$ of (1.8) satisfying*

$$\lim_{x \rightarrow 0} \frac{u_k(x)}{P_\Omega(x)} = k, \quad (1.24)$$

where P_Ω is the Poisson kernel in Ω . Furthermore $u_k \in L^1(\Omega) \cap L_\rho^p(\Omega)$, $\nabla u_k \in L_\rho^q(\Omega)$, and

$$\int_{\Omega} (-u_k \Delta \zeta + (u_k^p - M|\nabla u_k|^q) \zeta) dx = -k \frac{\partial \zeta}{\partial \mathbf{n}}(0) \quad \text{for all } \zeta \in \mathbb{X}(\Omega). \quad (1.25)$$

In order to study the behaviour of these solutions u_k when $k \rightarrow \infty$ we have to introduce separable solutions of (1.1) in the model case \mathbb{R}_+^N . They are solutions of

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^{\frac{2p}{p+1}} &= 0 && \text{in } \mathbb{R}_+^N \\ u &= 0 && \text{on } \partial\mathbb{R}_+^N \setminus \{0\}, \end{aligned} \quad (1.26)$$

which have the following expression in spherical coordinates

$$u(r, \sigma) = r^{-\frac{2}{p-1}}\omega(\sigma) \quad \text{for all } (r, \sigma) \in (0, \infty) \times S_+^{N-1}.$$

Put

$$\alpha = \frac{2}{p-1}, \quad (1.27)$$

and denote by Δ' and ∇' the Laplace-Beltrami operator and the spherical gradient, then ω satisfies

$$\begin{aligned} -\Delta'\omega + \alpha(N-2-\alpha)\omega + |\omega|^{p-1}\omega - M(\alpha^2\omega^2 + |\nabla'\omega|^2)^{\frac{p}{p+1}} &= 0 && \text{in } S_+^{N-1} \\ \omega &= 0 && \text{in } \partial S_+^{N-1}. \end{aligned} \quad (1.28)$$

Theorem 1.9 *There exists a positive solution ω to problem (1.28) if one of the following conditions is satisfied:*

- (i) either $1 < p < \frac{N+1}{N-1}$ and $M \geq 0$,
- (ii) or $p = \frac{N+1}{N-1}$ and $M > 0$,
- (iii) or $1 < p < 3$ or $p > \frac{N+1}{N-1}$, and $M \geq M_{N,p}$ for some explicit value $M_{N,p} > 0$.

The positive solutions of (1.28) allow to characterize the limit u_∞ of the solutions u_k constructed in Theorem 1.7.

Theorem 1.10 *Let $1 < p < \frac{N+1}{N-1}$, $1 < q < \frac{N+1}{N}$ and $M > 0$, then*

$$\lim_{x \rightarrow 0} \frac{u_\infty(x)}{P_N(x)} = \infty. \quad (1.29)$$

Furthermore

- (i) If $1 < q < \frac{2p}{p+1}$

$$\lim_{r \rightarrow 0} r^\alpha u_\infty(r, \cdot) = \psi \quad \text{uniformly on } S_+^{N-1}, \quad (1.30)$$

where ψ is the unique positive solution of

$$\begin{aligned} -\Delta'\psi + \alpha(N-2-\alpha)\psi + |\psi|^{p-1}\psi &= 0 && \text{in } S_+^{N-1} \\ \psi &= 0 && \text{in } \partial S_+^{N-1}, \end{aligned} \quad (1.31)$$

and u_∞ is the unique positive function solution of (1.21) and satisfying (1.29).

- (ii) If $q = \frac{2p}{p+1}$

$$\lim_{r \rightarrow 0} r^\alpha u_\infty(r, \cdot) = \omega \quad \text{uniformly on } S_+^{N-1}, \quad (1.32)$$

where ω is the minimal positive solution of (1.28).

A similar result holds if \mathbb{R}_+^N is replaced by a bounded smooth domain $\Omega \subset \mathbb{R}_+^N$, which boundary contains 0. In that case we assume that $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$ (i.e. $\partial\mathbb{R}_+^N$ is the tangent hyperplane to $\partial\Omega$ at 0 in order to use the spherical coordinates (r, σ) as above. Finally, if $(p, q) = \left(\frac{N+1}{N-1}, \frac{N+1}{N}\right)$ and $\partial\Omega$ is "very flat" near 0 in the sense that $\text{dist}(x, T_{\partial\Omega}(0)) \leq c|x|^N$ for all $x \in \partial\Omega$ close to 0, we prove that the function u_∞ defined in the previous theorem still satisfies (1.32). Note that the above flatness condition is always satisfied if $N = 2$ since $\partial\Omega$ is locally the graph of a C^k real valued function ($k \geq 2$) defined on $T_{\partial\Omega}(0) \cap B_\delta$ and degenerate at 0.

When $\frac{2p}{p+1} < q < \min\{2, p\}$, the situation is completely changed and the solutions with strong boundary blow-up are modeled by equation (1.3). If $1 < q < 2$ we set

$$\beta = \frac{2-q}{q-1}, \quad (1.33)$$

and if $1 < q < p$

$$\gamma = \frac{q}{p-q}. \quad (1.34)$$

We prove the following result in the statement of which ϕ_1 denotes the first eigenfunction of $-\Delta'$ in $W_0^{1,2}(S_+^{N-1})$.

Theorem 1.11 *Assume $M > 0$ and $\frac{2p}{p+1} < q < \min\{2, p\}$. Then there exists a positive solution u of (1.1) in \mathbb{R}_+^N , which vanishes on $\partial\mathbb{R}_+^N \setminus \{0\}$ such that*

$$m\phi_1(\sigma)r^{-\gamma} \leq u(r, \sigma) \leq c_4 \max\left\{r^{-\alpha}, M^{\frac{1}{p-q}}r^{-\gamma}\right\} \quad \text{for all } (r, \sigma) \in (0, r^*) \times \overline{S_+^{N-1}}. \quad (1.35)$$

for some $m > 0$, $r^* \in (0, \infty]$ and where $c_4 = c_4(N, p, q) > 0$. If $Nq \geq (N-1)p$, $r^* = \infty$.

Note that our construction which is made by mean of supersolutions and subsolutions does not imply that in the case $\frac{2p}{p+1} < q < \frac{N+1}{N}$, the solution u_∞ obtained in Theorem 1.10 satisfies (1.35). A similar result holds if \mathbb{R}_+^N is replaced by a bounded smooth domain $\Omega \subset \mathbb{R}_+^N$, such that $0 \in \partial\Omega$, under the flatness condition $\text{dist}(x, T_{\partial\Omega}(0)) \leq c|x|^{\gamma+1}$ for $x \in \partial\Omega$ near 0.

In the sequel $C > 0$ denotes a constant the value of which can change from one occurrence to another and c_j ($j = 0, 1, 2, \dots$) a more specific positive constant the value of which depends of more precise elements such as p, q, N or other previous constants c_i .

In a forthcoming article [8] we study the isolated singularities of positive solution in a domains. Due to the number of parameters even the radial solutions present an amazing rich complexity.

Acknowledgements The authors are grateful to the anonymous referee for his careful checking of the work and his pertinent observations. This article has been prepared with the support FONDECYT grants 1210241 and 1190102 for the three authors.

2 Singular boundary value problems

2.1 A priori estimates

We give two series of estimates for solutions of (1.1) with a boundary singularity according to the sign of M .

Theorem 2.1 *Let Ω be a domain such that $0 \in \partial\Omega$, $M \in \mathbb{R}$ and $1 < q < \min\{p, 2\}$. If $u \in C^1(\overline{\Omega} \setminus \{0\})$ is a solution of (1.1) vanishing on $\partial\Omega \setminus \{0\}$, there holds*

1- *If $M > 0$, there exists $c_5(N, p, q) > 0$ such that*

$$u_+(x) \leq c_5 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{q}{p-q}}, |x|^{-\frac{2}{p-1}} \right\} \quad \text{for all } x \in \Omega. \quad (2.1)$$

2- *If $M \leq 0$, there exist $c_6 = c_6(N, q) > 0$ and $c_7 = c_7(N, p) > 0$ such that*

$$u_+(x) \leq \min \left\{ c_6 |M|^{-\frac{1}{q-1}} |x|^{-\frac{2-q}{q-1}}, c_7 |x|^{-\frac{2}{p-1}} \right\} \quad \text{for all } x \in \Omega. \quad (2.2)$$

Proof. We first assume that $\overline{\Omega} \subset B_{R_0}$ for some $R_0 > 0$. Let $\epsilon > 0$, we set

$$j_\epsilon(r) = \begin{cases} 0 & \text{if } r \leq 0 \\ \frac{r^2}{2\epsilon} & \text{if } 0 \leq r \leq \epsilon \\ r - \frac{\epsilon}{2} & \text{if } r \geq \epsilon. \end{cases}$$

If we extend u by 0 in $\overline{\Omega}^c \cap B_{2R_0}$ and set $v_\epsilon = j_\epsilon(u)$ we have

$$\begin{aligned} -\Delta v_\epsilon + v_\epsilon^p - M|\nabla v_\epsilon|^q &= -j'_\epsilon(u)\Delta u - j''_\epsilon(u)|\nabla u|^2 + (j_\epsilon(u))^p - M(j'_\epsilon(u))^q |\nabla u|^q \\ &\leq M j'_\epsilon(u) (1 - (j'_\epsilon(u))^{q-1}) |\nabla u|^q + (j_\epsilon(u))^p - j'_\epsilon(u) u_+^p \\ &\leq M \frac{u}{\epsilon} \left(1 - \frac{u^{q-1}}{\epsilon^{q-1}} \right) |\nabla v_\epsilon|^q \chi_{\{0 < u < \epsilon\}}. \end{aligned}$$

Letting $\epsilon \rightarrow 0$, we deduce from the dominated convergence theorem that $v_0 = \lim_{\epsilon \rightarrow 0} v_\epsilon$ is nonnegative (actually it is the extension of u^+ by 0 outside $\overline{\Omega} \setminus \{0\}$) and satisfies

$$L v_0 := -\Delta v_0 + v_0^p - M|\nabla v_0|^q \leq 0 \quad \text{in } \mathcal{D}'(B_{2R_0} \setminus \{0\}). \quad (2.3)$$

The case $M > 0$. Following the method of Keller [16] and Osserman [23], we fix $a \in B_{R_0} \setminus \{0\}$, and introduce $U(x-a) = \lambda(|a|^2 - |x-a|^2)^{-b}$ for some $b > 0$. Then putting $r = |x-a|$ and $\tilde{U}(r) = U(x-a)$, we have

$$\begin{aligned} L\tilde{U} &= -\tilde{U}'' - \frac{N-1}{r}\tilde{U}' - M|\tilde{U}'|^q + \tilde{U}^p \\ &= \lambda(|a|^2 - r^2)^{-2-b} [\lambda^{p-1}(|a|^2 - r^2)^{2-b(p-1)} + 2b(N-2(b+1))r^2 - 2Nb|a|^2 \\ &\quad - M2^q b^q \lambda^{q-1} r^q (|a|^2 - r^2)^{2+b-q(b+1)}]. \end{aligned}$$

If $M > 0$, the two necessary conditions on b to be fulfilled in order \tilde{U} be a supersolution in $B_{|a|}(a)$ are

- (i) $2 - b(p - 1) \leq 0 \iff b(p - 1) \geq 2$,
- (ii) $2 + b - q(b + 1) \geq 2 - b(p - 1) \iff b(p - q) \geq q$.

The above inequalities are satisfied if

$$b = \max \left\{ \frac{2}{p-1}, \frac{q}{p-q} \right\} = \max \{ \alpha, \gamma \}. \quad (2.4)$$

If $q > \frac{2p}{p+1}$ then $b = \frac{q}{p-q}$ and

$$L\tilde{U} \geq \lambda (|a|^2 - r^2)^{-\frac{2p-q}{p-q}} \left[\lambda^{q-1} (\lambda^{p-q} - M2^q b^q r^q) (|a|^2 - r^2)^{\frac{2p-q(p+1)}{p-q}} - (3b+1)N|a|^2 \right].$$

There exists $c_5 > 0$ depending on N , p and q such that if we choose

$$\lambda = c_5 \max \left\{ M^{\frac{1}{p-q}} |a|^{\frac{q}{p-q}}, |a|^{\frac{2p(q-1)}{(p-1)(p-q)}} \right\},$$

there holds

$$L\tilde{U} \geq 0. \quad (2.5)$$

Since $\tilde{U}(x) \rightarrow \infty$ when $|x| \rightarrow |a|$, we obtain by the maximum principle (see [24] for a) that $v_0 \leq \tilde{U}$ in $B_{|a|}(a)$. In particular

$$u_+(a) = v_0(a) \leq \tilde{U}(a) = \lambda |a|^{-\frac{2q}{p-q}} = c_5 \max \left\{ M^{\frac{1}{p-q}} |a|^{-\frac{q}{p-q}}, |a|^{-\frac{2}{p-1}} \right\}. \quad (2.6)$$

If $q \leq \frac{2p}{p+1}$ then $b = \frac{2}{p-1}$ and

$$\begin{aligned} L\tilde{U} &\geq \lambda (|a|^2 - r^2)^{-\frac{2p}{p-1}} \left[\lambda^{p-1} + \frac{2}{p-1} \left(N - \frac{2(p+1)}{p-1} \right) r^2 - \frac{2N}{p-1} |a|^2 \right. \\ &\quad \left. - M2^q \left(\frac{2}{p-1} \right)^q \lambda^{q-1} r^q (|a|^2 - r^2)^{\frac{2p-q(p+1)}{p-1}} \right] \\ &\geq \lambda (|a|^2 - r^2)^{-\frac{2p}{p-1}} \left[\lambda^{p-1} - C|a|^2 - C' \lambda^{q-1} M |a|^{\frac{4p-q(p+3)}{p-1}} \right]. \end{aligned}$$

Hence, if $q = \frac{2p}{p+1}$, (2.5) holds if for some $c_5 = c_5(N, p, q) > 0$,

$$\lambda = c_5 \max \left\{ M^{\frac{p+1}{p(p-1)}}, 1 \right\} |a|^{\frac{2}{p-1}},$$

which yields

$$u_+(a) = v_0(a) \leq \tilde{U}(a) = \lambda |a|^{-\frac{4}{p-1}} = c_5 \max \left\{ M^{\frac{p+1}{p(p-1)}}, 1 \right\} |a|^{-\frac{2}{p-1}}. \quad (2.7)$$

While if $q < \frac{2p}{p+1}$, we choose

$$\lambda = c_5 \max \left\{ M^{\frac{1}{p-q}} |a|^{\frac{4p-q(p+3)}{(p-1)(p-q)}}, |a|^{\frac{2}{p-1}} \right\},$$

where $c_5 > 0 = c_5(N, p, q)$, which yields

$$u_+(a) = v_0(a) \leq \tilde{U}(a) = \lambda |a|^{-\frac{4}{p-1}} = c_5 \max \left\{ M^{\frac{1}{p-q}} |a|^{-\frac{q}{p-q}}, |a|^{-\frac{2}{p-1}} \right\}. \quad (2.8)$$

The case $M \leq 0$. We first assume that $M < 0$. By [22, Lemma 3.3] v_0 satisfies

$$-\Delta v_0 + |M| |\nabla v_0|^q \leq 0 \quad \text{in } \mathcal{D}'(B_{2R_0} \setminus \{0\}). \quad (2.9)$$

Therefore, since $1 < q < 2$,

$$u_+(a) = v_0(a) \leq c_6 |M|^{-\frac{1}{q-1}} |a|^{-\frac{2-q}{q-1}}. \quad (2.10)$$

If $M \leq 0$ there also holds

$$-\Delta v_0 + v_0^p \leq 0 \quad \text{in } \mathcal{D}'(B_{2R_0} \setminus \{0\}). \quad (2.11)$$

Hence

$$u_+(a) = v_0(a) \leq c_7 |a|^{-\frac{2}{p-1}}. \quad (2.12)$$

In the above inequalities $c_6 = c_6(q, N) > 0$ and $c_7 = c_7(p, N) > 0$. Combining these estimates we derive

$$u_+(a) \leq \min \left\{ c_7 |a|^{-\frac{2}{p-1}}, c_6 |M|^{-\frac{1}{q-1}} |a|^{-\frac{2-q}{q-1}} \right\}. \quad (2.13)$$

Since the estimate is independent of R_0 , the assumption that $\bar{\Omega} \subset B_{R_0}$ is easily ruled out. This ends the proof. \square

Remark. If $M = 0$, estimate (2.1) is just

$$u_+(x) \leq c_7 |x|^{-\frac{2}{p-1}}. \quad (2.14)$$

If $M < 0$, (2.14) is valid what ever is the value of q . Furthermore there also holds

$$u_+(x) \leq c_6 |M| |x|^{-\frac{2-q}{q-1}}, \quad (2.15)$$

whatever is the value of p , provided $1 < q < 2$.

The equation is not invariant by $u \mapsto -u$, hence the lower and upper estimates are not symmetric.

Corollary 2.2 *Under the assumptions of Theorem 2.1, there holds*

1- *If $M > 0$*

$$\begin{aligned} -\min \left\{ c_6 |M|^{-\frac{1}{q-1}} |x|^{-\frac{2-q}{q-1}}, c_7 |x|^{-\frac{2}{p-1}} \right\} &\leq u(x) \\ &\leq c_5 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{q}{p-q}}, |x|^{-\frac{2}{p-1}} \right\} \quad \text{for all } x \in \Omega. \end{aligned} \quad (2.16)$$

2- *If $M \leq 0$, there exist $c_6 = c_6(N, q) > 0$ and $c_7 = c_7(N, p) > 0$ such that*

$$\begin{aligned} -c_5 \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{q}{p-q}}, |x|^{-\frac{2}{p-1}} \right\} &\leq u(x) \\ &\leq \min \left\{ c_6 |M|^{-\frac{1}{q-1}} |x|^{-\frac{2-q}{q-1}}, c_7 |x|^{-\frac{2}{p-1}} \right\} \quad \text{for all } x \in \Omega. \end{aligned} \quad (2.17)$$

We infer from Theorem 2.1 an estimate of the gradient of u near 0.

Theorem 2.3 *Let Ω be a smooth bounded domain such that $0 \in \partial\Omega$ and $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$, $M > 0$, $p > 1$ and $1 < q < \min\{2, p\}$. If $u \in C^1(\bar{\Omega} \setminus \{0\})$ is a nonnegative solution of (1.1) vanishing on $\partial\Omega \setminus \{0\}$, for any $r_0 > 0$ there holds there exists $c_8 = c_8(N, p, q, \Omega, r_0, M) > 0$ such that*

$$|\nabla u(x)| \leq c_8 \max \left\{ |x|^{-\frac{p}{p-q}}, |x|^{-\frac{p+1}{p-1}} \right\} \quad \text{for all } x \in \Omega \cap B_{r_0}. \quad (2.18)$$

The restriction that $|x| \leq 1$ is not needed if $q = \frac{2p}{p+1}$.

Proof. We assume first that $B_2^+ \subset \Omega$.

Case 1: $1 < q \leq \frac{2p}{p+1}$. For $0 < r < 1$ we set

$$u(x) = r^{-\frac{2}{p-1}} u_r\left(\frac{x}{r}\right) = r^{-\frac{2}{p-1}} u_r(y) \quad \text{with } y = \frac{x}{r}.$$

If $\frac{r}{2} < |x| < 2r$, then $\frac{1}{2} < |y| < 2$ and $u_r > 0$ satisfies

$$-\Delta u_r + u_r^p - M r^{\frac{2p-q(p+1)}{p-1}} |\nabla u_r|^q = 0 \quad \text{in } B_2^+ \setminus B_{\frac{1}{2}}^+,$$

and vanishes on $\partial(B_2^+ \setminus B_{\frac{1}{2}}^+)$. Since $0 < M r^{\frac{2p-q(p+1)}{p-1}} \leq M$ as $2p - q(p+1) \geq 0$, by the standard regularity theory we have the estimate

$$\max \left\{ |\nabla u_r(z)| : \frac{2}{3} < |z| < \frac{3}{2} \right\} \leq c_9 \max \left\{ |u_r(z)| : \frac{1}{2} < |z| < 2 \right\}, \quad (2.19)$$

where c_9 depends on N, p, q and M . Now it follows that

$$\max \left\{ |u_r(z)| : \frac{1}{2} < |z| < 2 \right\} \leq 2^{\frac{2}{p-1}} c_5 \max \left\{ M^{\frac{1}{p-q}} r^{\frac{2p-q(p+1)}{(p-1)(p-q)}}, 1 \right\},$$

by (2.1). Therefore

$$\begin{aligned} \max \left\{ |\nabla u(y)| : \frac{r}{2} < |y| < 2r \right\} &\leq 2^{\frac{2}{p-1}} c_5 c_9 r^{-\frac{p+1}{p-1}} \max \left\{ M^{\frac{1}{p-q}} r^{\frac{2p-q(p+1)}{(p-1)(p-q)}}, 1 \right\} \\ &\leq c_8 \max \left\{ |x|^{-\frac{p}{p-q}}, |x|^{-\frac{p+1}{p-1}} \right\}, \end{aligned} \quad (2.20)$$

which is (2.18).

Case 2: $\frac{2p}{p+1} < q < 2$. For $0 < r < 1$ we set

$$u(x) = r^{-\frac{2-q}{q-1}} u_r\left(\frac{x}{r}\right) = r^{-\frac{2-q}{q-1}} u_r(y) \quad \text{with } y = \frac{x}{r}.$$

If $\frac{r}{2} < |x| < 2r$, then $\frac{1}{2} < |y| < 2$ and $u_r > 0$ satisfies

$$-\Delta u_r + r^{\frac{q(p+1)-2p}{q-1}} u_r^p - M |\nabla u_r|^q = 0 \quad \text{in } B_2^+ \setminus B_{\frac{1}{2}}^+,$$

We notice that $q(p+1) - 2p > 0$. Then inequality (2.19) holds. Now

$$\max \left\{ |u_r(z)| : \frac{1}{2} < |z| < 2 \right\} \leq c'_9 r^{\frac{2-q}{q-1}} \max \left\{ r^{-\frac{2}{p-1}}, r^{-\frac{q}{p-q}} \right\},$$

thus

$$\max \left\{ |\nabla u_r(z)| : \frac{2}{3} < |z| < \frac{3}{2} \right\} \leq c''_9 r^{\frac{2-q}{q-1}-1} \max \left\{ r^{-\frac{2}{p-1}}, r^{-\frac{q}{p-q}} \right\}, \quad (2.21)$$

which implies

$$\max \left\{ |\nabla u(x)| : \frac{2r}{3} < |x| < \frac{3r}{2} \right\} \leq c_8 \max \left\{ r^{-\frac{p+1}{p-1}}, r^{-\frac{p}{p-q}} \right\}. \quad (2.22)$$

The general case; If $\partial\Omega$ is not flat near 0 we proceed as in the proof of [22, Lemma 3.4], using the same scaling as in the flat case which transform the domain $B_2^+ \setminus B_1^+$ into $(B_2 \setminus B_1) \cap \frac{1}{r}\Omega$, the curvature of which is bounded when $0 < r < 1$. The same estimates holds, up to the value of the constant c_8 and we derive (2.18). \square

As a consequence we have the following.

Corollary 2.4 *Under the assumptions of Theorem 2.3 the function u satisfies*

$$u(x) \leq c_8 \rho(x) \max \left\{ M^{\frac{1}{p-q}} |x|^{-\frac{p}{p-q}}, |x|^{-\frac{p+1}{p-1}} \right\} \quad \text{for all } x \in \Omega \cap B_1. \quad (2.23)$$

The restriction that $|x| \leq 1$ is not needed if $q = \frac{2p}{p+1}$.

2.2 Removable singularities

Proof of Theorem 1.1. If $M \leq 0$, u is a nonnegative subsolution of $-\Delta u + v^p = 0$ which vanishes on $\partial\Omega \setminus \{0\}$, hence it is identically zero by [15].

Step 1. We assume $M > 0$. It is straightforward to verify from estimates (2.18) that under conditions (i) or (ii), $|\nabla u(x)| \leq c_8 |x|^{-a}$ with $a \leq N$. Since these conditions imply $q < \frac{N+1}{N}$, it follows that $|\nabla u|^q \in L^1_\rho(\Omega)$.

For any $\epsilon > 0$ we denote by w_ϵ the solution of

$$\begin{aligned} -\Delta w + w^p &= M|\nabla u|^q && \text{in } \Omega_\epsilon := \Omega \cap \overline{B}_\epsilon^c \\ w &= 0 && \text{in } \partial\Omega \cap \overline{B}_\epsilon^c \\ \lim_{|x| \rightarrow \epsilon} w(x) &= \infty && \text{on } \partial B_\epsilon \cap \Omega, \end{aligned} \quad (2.24)$$

which exists since $|\nabla u|^q \in L_\rho^1(\Omega)$, see [19]. Then $u \leq w_\epsilon$ in Ω_ϵ . Let z_ϵ be the solution of

$$\begin{aligned} -\Delta z + z^p &= 0 && \text{in } \Omega_\epsilon \\ z &= 0 && \text{in } \partial\Omega \cap \overline{B}_\epsilon^c \\ \lim_{|x| \rightarrow \epsilon} z(x) &= \infty && \text{on } \partial B_\epsilon \cap \Omega. \end{aligned} \quad (2.25)$$

Denote by $\mathbb{G}_\Omega[\cdot]$ the Green operator in Ω . Since $z_\epsilon + M\mathbb{G}_\Omega[|\nabla u|^q]_{\Omega_\epsilon}$ is a supersolution of (2.24) in Ω_ϵ we deduce

$$u \leq z_\epsilon + M\mathbb{G}_\Omega[|\nabla u|^q]_{\Omega_\epsilon} \quad \text{in } \Omega_\epsilon. \quad (2.26)$$

When $\epsilon \rightarrow 0$, z_ϵ decreases to z_0 which satisfies

$$\begin{aligned} -\Delta w + w^p &= 0 && \text{in } \Omega \\ w &= 0 && \text{in } \partial\Omega \setminus \{0\}. \end{aligned} \quad (2.27)$$

Since $p \geq \frac{N+1}{N-1}$ it is proved in [15] that any solution of (2.27) extends as a continuous solution in Ω with boundary value 0, hence $z_0 = 0$ by the maximum principle. Therefore $u \leq M\mathbb{G}_\Omega[|\nabla u|^q]$ in Ω and the boundary trace $Tr_{\partial\Omega}[u]$ of u is zero. By [20] the fact that $|\nabla u|^q \in L_\rho^1(\Omega)$ jointly with $Tr_{\partial\Omega}[u] = 0$ implies in turn that $u^p \in L_\rho^1(\Omega)$ and u is a weak solution of

$$\begin{aligned} -\Delta u + u^p &= M|\nabla u|^q && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega, \end{aligned} \quad (2.28)$$

in the sense that there holds

$$\int_{\Omega} (-u\Delta\zeta + u^p\zeta - M|\nabla u|^q\zeta) dx = 0 \quad \forall \zeta \in W^{2,\infty}(\Omega) \cap C_c^1(\overline{\Omega}). \quad (2.29)$$

Step 2. Let us assume that $p > \frac{N+1}{N-1}$. If u is nonnegative and not identically zero, then by the maximum principle it is positive in Ω . We set $u = v^b$ with $0 < b \leq 1$. Then

$$-\Delta v - (b-1)\frac{|\nabla v|^2}{v} + \frac{1}{b}v^{(p-1)b+1} - Mb^{q-1}v^{(b-1)(q-1)}|\nabla v|^q = 0. \quad (2.30)$$

For $\epsilon > 0$,

$$v^{(b-1)(q-1)}|\nabla v|^q \leq \frac{q\epsilon^{\frac{2}{q}}}{2} \frac{|\nabla v|^2}{v} + \frac{2-q}{2\epsilon^{\frac{2}{2-q}}} v^{\frac{(2b-1)q-2(b-1)}{2-q}}.$$

Therefore

$$-\Delta v + \left(1 - b - M \frac{qb^{q-1}\epsilon^{\frac{2}{q}}}{2}\right) \frac{|\nabla v|^2}{v} + \frac{1}{b} v^{(p-1)b+1} - Mb^{q-1} \frac{2-q}{2\epsilon^{\frac{2}{2-q}}} v^{\frac{(2b-1)q-2(b-1)}{2-q}} = 0. \quad (2.31)$$

We notice that the following relation is independent of b

$$\frac{(2b-1)q-2(b-1)}{2-q} \leq (p-1)b+1 \iff q \leq \frac{2p}{p+1},$$

with simultaneous equality. We take

$$(p-1)b+1 = \frac{N+1}{N-1} \iff b = \frac{2}{(N-1)(p-1)}, \quad (2.32)$$

hence $p > \frac{N+1}{N-1}$ if and only if $0 < b < 1$.

We first assume that $0 < q < \frac{2p}{p+1}$ and choose $\epsilon > 0$ such that

$$1 - b - M \frac{qb^{q-1}\epsilon^{\frac{2}{q}}}{2} = 0 \iff \epsilon = \left(\frac{2(1-b)}{Mqb^{q-1}}\right)^{\frac{q}{2}} = \left(\frac{2((N-1)p-N-1)}{Mqb^{q-1}(N-1)(p-1)}\right)^{\frac{q}{2}}. \quad (2.33)$$

This transforms (2.31) into

$$-\Delta v + \frac{(N-1)(p-1)}{2} v^{\frac{N+1}{N-1}} - \frac{(2-q)b^{\frac{2(q-1)}{2-q}}}{2} \left(\frac{q}{2(1-b)}\right)^{\frac{q}{2-q}} M^{\frac{2}{2-q}} v^{\frac{(2b-1)q-2(b-1)}{2-q}} \leq 0. \quad (2.34)$$

Then, as

$$\frac{(2b-1)q-2(b-1)}{2-q} < \frac{N+1}{N-1},$$

there exists $A > 0$, depending on M and b , such that

$$-\Delta v + \frac{(N-1)(p-1)}{4} v^{\frac{N+1}{N-1}} \leq A. \quad (2.35)$$

Since v vanishes on $\partial\Omega \setminus \{0\}$, $\tilde{v} = (v - c_{10}A^{\frac{N-1}{N+1}})^{\frac{N+1}{N-1}}_+$ with $c_{10} = \left(\frac{4}{(N-1)(p-1)}\right)^{\frac{N+1}{N-1}}$ satisfies

$$-\Delta \tilde{v} + \frac{(N-1)(p-1)}{4} \tilde{v}^{\frac{N+1}{N-1}} \leq 0. \quad (2.36)$$

By [15], $\tilde{v} = 0$ which implies $v \leq c_{10}A^{\frac{N-1}{N+1}}$ and therefore $u(x) \leq c_{11}A^{\frac{2}{(N+1)(p-1)}}$ in Ω . Since u vanishes on $\partial\Omega \setminus \{0\}$ we extend it in a neighborhood of 0 by odd reflection trough $\partial\Omega$ and denote by \tilde{u} the new function defined in B_α where it satisfies

$$-\operatorname{div}A(x, \nabla \tilde{u}) + \tilde{u}^p + B(x, \nabla \tilde{u}) = 0 \quad \text{in } B_\alpha \setminus \{0\}. \quad (2.37)$$

In this expression the operator $A : (x; \xi) \in B_\alpha \times \mathbb{R}^N \mapsto A(x, \xi) \in \mathbb{R}^N$ is smooth in x in B_α and linear in ξ (see e.g. [4, Lemma 2.5] in a more general setting), it and satisfies for all $(x; \xi) \in B_\alpha \times \mathbb{R}^N$,

$$A(x, \xi) \cdot \xi \geq 2|\xi|^2 \quad \text{and} \quad |A(x, \xi)| \leq 4|\xi| \quad \text{for all } (x; \xi) \in B_\alpha \times \mathbb{R}^N.$$

Since we can write $|B(\cdot, \nabla \tilde{u})| \leq 2|\nabla \tilde{u}|^q = 2|\nabla \tilde{u}|^{q-1}|\nabla \tilde{u}| = C(x)|\nabla \tilde{u}|$ in B_α , then $B : (x, \xi) \in B_\alpha \times \mathbb{R}^N \mapsto B(x, \xi) \in \mathbb{R}$ verifies

$$|B(x, \xi)| \leq C(x)|\xi|,$$

and $C(x) \leq 2c_8|x|^{-\frac{(p+1)(q-1)}{p-1}}$ by Theorem 2.3. Since $q < \frac{2p}{p+1}$, $\frac{(p+1)(q-1)}{p-1} < 1$. Hence $C \in L^{N+\tau}$ for some $\tau > 0$. By Serrin's theorem [25, Theorem 10] the singularity at 0 is removable and \tilde{u} can be extended as a regular solution of (2.37) in B_α . Hence $\tilde{u} \in C^1(B_{\frac{\alpha}{2}})$, and as a consequence $u \in C^1(\bar{\Omega})$. If u is not zero, it is positive in Ω and achieves its maximum at some $x_0 \in \Omega$ where $\Delta u(x_0) \leq 0$ and $\nabla u(x_0) = 0$. Contradiction.

Next we assume that $q = \frac{2p}{p+1}$. By the choice of b in (2.32), inequality (2.31) becomes

$$-\Delta v + \left(1 - b - \frac{Mpb^{\frac{p-1}{p+1}}\epsilon^{\frac{p+1}{p}}}{p+1}\right) \frac{|\nabla v|^2}{v} + \left(\frac{1}{b} - \frac{Mb^{\frac{p-1}{p+1}}}{(p+1)\epsilon^{p+1}}\right) v^{(p-1)b+1} \leq 0. \quad (2.38)$$

We need to make both coefficients positive so that we obtain

$$\begin{aligned} -\Delta v + \tau v^{\frac{N+1}{N-1}} &\leq 0 && \text{in } \Omega \\ v &= 0 && \text{on } \partial\Omega \setminus \{0\}. \end{aligned} \quad (2.39)$$

We first choose

$$\epsilon^{\frac{p+1}{p}} > \left(\frac{M}{p+1}\right)^{\frac{1}{p}} b^{\frac{2}{p+1}},$$

say

$$\epsilon^{\frac{p+1}{p}} = \left(\frac{M}{p+1}\right)^{\frac{1}{p}} b^{\frac{2}{p+1}} + \tilde{\epsilon}, \quad (2.40)$$

with $\tilde{\epsilon} > 0$ so that the coefficient of $v^{\frac{N+1}{N-1}}$ is positive, and we can choose $\tilde{\epsilon}$ thanks to the assumption $m^{**} > M$: we have

$$\begin{aligned} 1 - b - \frac{Mpb^{\frac{p-1}{p+1}}}{p+1} \left(\left(\frac{M}{p+1}\right)^{\frac{1}{p}} b^{\frac{2}{p+1}} + \tilde{\epsilon} \right) &= 1 - b - \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} pb - \frac{Mpb^{\frac{p+1}{p-1}}}{p+1} \tilde{\epsilon} \\ &= b \left(\frac{1-b}{b} - \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} p \right) - \frac{Mpb^{\frac{p+1}{p-1}}}{p+1} \tilde{\epsilon} \\ &= pb \left(\frac{(N-1)p - (N+1)}{2p} - \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} \right) - \frac{Mpb^{\frac{p+1}{p-1}}}{p+1} \tilde{\epsilon} \\ &= pb \left(\left(\frac{m^{**}}{p+1}\right)^{\frac{p+1}{p}} - \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} \right) - \frac{Mpb^{\frac{p+1}{p-1}}}{p+1} \tilde{\epsilon} \end{aligned} \quad (2.41)$$

and the right-hand side is positive if $\tilde{\epsilon}$ small enough. Hence we obtain (2.39). By [15], $v = 0$ and the same holds for u . This ends the case $p > \frac{N+1}{N-1}$.

Step 3. Finally we assume $p = \frac{N+1}{N-1}$ and $1 < q < \frac{2p}{p+1} = \frac{N+1}{N}$, then

$$M|\nabla u(x)|^q \leq c_{12}|x|^{-q\frac{p+1}{p-1}} = c_{12}|x|^{-qN} := c_{13}Q(x).$$

Hence $u \leq u_1 := c_{13}\mathbb{G}_\Omega[Q]$. At this point we need the following intermediate result:

Claim. Assume $w_\alpha = \mathbb{G}_\Omega[Q_\alpha]$ where $Q_\alpha(x) = |x|^{-\alpha}$ with $\alpha < N + 1$, then

$$w_\alpha(x) \leq c_\alpha|x|^{2-\alpha} \quad \text{for all } x \in \Omega. \quad (2.42)$$

If this holds true, then $u(x) \leq c_{13}c_{qN}|x|^{2-qN}$. By the scaling method of Theorem 2.3, we obtain

$$|\nabla u(x)| \leq c_8c_{13}c_{qN}|x|^{1-qN} \implies |\nabla u(x)|^q \leq c_{14}|x|^{q(1-qN)} := c_{14}Q_{q(Nq-1)}(x), \quad (2.43)$$

and thus

$$w_{q(Nq-1)}(x) = c_{14}\mathbb{G}_\Omega[Q_{q(Nq-1)}](x) \leq c_{14}c_{q(Nq-1)}|x|^{2-q(Nq-1)} \quad \text{for all } x \in \Omega. \quad (2.44)$$

Since $q < 1 + \frac{1}{N}$, $q(Nq - 1) - 2 < Nq - 2$. Iterating this process, we finally obtain that u is bounded and we end the proof as in Step 2. \square

Remark. It is noticeable that the equation exhibits a phenomenon which is characteristic of Emden-Fowler type equations

$$\Delta u = u^p \quad \text{in } B_1 \setminus \{0\}. \quad (2.45)$$

If u is nonnegative then there exists $a \geq 0$ such that

$$\Delta u = u^p + a\delta_0 \quad \text{in } \mathcal{D}'(B_1). \quad (2.46)$$

If $1 < p < \frac{N}{N-2}$ then a can be positive, but if $p \geq \frac{N}{N-2}$, then $a = 0$. This means that the singularity cannot be seen in the sense of distributions, however there truly exist singular solutions, e.g. if $p > \frac{N}{N-2}$,

$$u_s(x) = c_{N,p}|x|^{-\frac{2}{p-1}}. \quad (2.47)$$

A similar phenomenon exists for solutions of

$$\begin{aligned} -\Delta u &= u^p & \text{in } B_1^+ \\ u &= 0 & \text{in } \partial B_1^+ \setminus \{0\}. \end{aligned} \quad (2.48)$$

In such a case the critical value is $\frac{N+1}{N-1}$ since for $p \geq \frac{N+1}{N-1}$ the boundary value is achieved in the sense of distributions in ∂B_1^+ .

2.3 Proof of Theorem 1.2

As in Theorem 1.1, the proof differs according to whether $0 < q < \frac{2p}{p+1}$ or $q = \frac{2p}{p+1}$, and we first assume that $u > 0$. We perform the same change of unknown as in the previous theorem putting $u = v^b$, but now we choose b as follows

$$(p-1)b + 1 = r \iff b = \frac{r-1}{p-1}, \quad (2.49)$$

and we first assume that

$$1 - b - M \frac{qb^{q-1}\epsilon^{\frac{2}{q}}}{2} = 0 \iff \epsilon = \left(\frac{2(1-b)}{Mqb^{q-1}} \right)^{\frac{q}{2}} = \left(\frac{2(p-r)}{Mq(p-1)b^{q-1}} \right)^{\frac{q}{2}}. \quad (2.50)$$

Hence (2.34) becomes

$$-\Delta v + \frac{p-1}{r-1}v^r - \frac{(2-q)b^{q-1}}{2} \left(\frac{q}{2(1-b)} \right)^{\frac{q}{2-q}} M^{\frac{2}{2-q}} v^{\frac{(2r-p-1)q+2(p-r)}{(p-1)(2-q)}} \leq 0. \quad (2.51)$$

The condition $r \geq \frac{(2r-p-1)q+2(p-r)}{(p-1)(2-q)}$ is equivalent to $2p - q(p+1) \leq r(2p - q(p+1))$ since $1 < r < p$.

Assuming first that $q < \frac{2p}{p+1}$, we obtain from (2.51)

$$-\Delta v + \frac{p-1}{2(r-1)}v^r \leq A. \quad (2.52)$$

for some constant $A \geq 0$. Since $cap_{\frac{2}{r}, r'}^{\partial\Omega}(K) = 0$ and v vanishes on $\partial\Omega \setminus K$, it follows from [18] that $v \leq cA^{\frac{1}{r}}$ for some $c > 0$, hence u is also uniformly bounded above in Ω by some constant a . Next we have to show that $\nabla u \in L^2(\Omega)$. We also denote by Φ_1 the first eigenfunction of $-\Delta$ in $W_0^{1,2}(\Omega)$ normalized by $\sup \Phi_1 = 1$ and by λ_1 the corresponding eigenvalue. Since $\frac{N+1}{N-1} < r \leq 3$ we infer from [1, Theorem 5.5.1], that

$$\left(cap_{\frac{1}{2}, 2}^{\partial\Omega}(K) \right)^{\frac{1}{N-2}} \leq B \left(cap_{\frac{2}{r}, r'}^{\partial\Omega}(K) \right)^{\frac{1}{N-1-\frac{2}{r-1}}}.$$

Therefore $cap_{\frac{2}{r}, r'}^{\partial\Omega}(K) = 0$ implies $cap_{\frac{1}{2}, 2}^{\partial\Omega}(K) = 0$ and there exists a decreasing sequence $\{\zeta_n\} \subset C_0^2(\partial\Omega)$ such that $\zeta_n = 1$ in a neighborhood of K , $0 \leq \zeta_n \leq 1$ and $\|\zeta_n\|_{W^{1,2}(\partial\Omega)} \rightarrow 0$ when $n \rightarrow \infty$, furthermore $\zeta_n \rightarrow 0$ quasi everywhere. Let $\mathbb{P}_\Omega : C^2(\partial\Omega) \mapsto C^2(\overline{\Omega})$ be the Poisson operator. It is an admissible lifting in the sense of [18, Section 1] in the sense that

$$\mathbb{P}_\Omega[\eta]|_{\partial\Omega} = \eta \quad \text{and} \quad \eta \geq 0 \implies \mathbb{P}_\Omega[\eta] \geq 0.$$

Put $\eta_n = 1 - \zeta_n$. Then, multiplying equation (1.13) by $u(\mathbb{P}_\Omega[\eta_n])^2$ and integrating, we obtain

$$\begin{aligned} & \int_{\Omega} |\nabla u|^2 (\mathbb{P}_\Omega[\eta_n])^2 dx + 2 \int_{\Omega} u \mathbb{P}_\Omega[\eta_n] \nabla u \cdot \nabla \mathbb{P}_\Omega[\eta_n] dx \\ & + \int_{\Omega} u^{p+1} (\mathbb{P}_\Omega[\eta_n])^2 dx - M \int_{\Omega} |\nabla u|^q u (\mathbb{P}_\Omega[\eta_n])^2 dx = 0, \end{aligned}$$

which implies

$$\begin{aligned} \int_{\Omega} |\nabla u|^2 (\mathbb{P}_{\Omega}[\eta_n])^2 dx - 2 \left(\int_{\Omega} |\nabla u|^2 (\mathbb{P}_{\Omega}[\eta_n])^2 dx \right)^{\frac{1}{2}} \left(\int_{\Omega} |\nabla \mathbb{P}_{\Omega}[\eta_n]|^2 u^2 dx \right)^{\frac{1}{2}} \\ + \int_{\Omega} u^{p+1} (\mathbb{P}_{\Omega}[\eta_n])^2 dx - M \int_{\Omega} |\nabla u|^q u (\mathbb{P}_{\Omega}[\eta_n])^2 dx \leq 0. \end{aligned}$$

It is standard that

$$\int_{\Omega} |\nabla \mathbb{P}_{\Omega}[\eta_n]|^2 dx \leq c_{12} \|\eta_n\|_{W^{\frac{1}{2},2}(\partial\Omega)}^2 = A_n.$$

Set $X_n = \|\mathbb{P}_{\Omega}[\eta_n]|\nabla u\|_{L^2}$, then

$$X_n^2 - 2A_n X_n - Ma|\Omega|^{\frac{2-q}{2}} X_n^q \leq 0.$$

Hence there exist two positive real numbers a_1 and a_2 depending only on q , $|\Omega|$ and $a = \|u\|_{L^\infty}$ such that

$$X_n \leq a_1 A_n^{\frac{1}{q-1}} + a_2 M^{\frac{1}{2-q}}. \quad (2.53)$$

Now $A_n \rightarrow 0$ and $X_n \rightarrow \|\nabla u\|_{L^2}^2$, therefore by Fatou's Lemma

$$|\Omega|^{1-\frac{2}{q}} \|\nabla u\|_{L^q}^2 \leq \|\nabla u\|_{L^2}^2 \leq a_2 M^{\frac{1}{2-q}} < \infty.$$

Let $\zeta \in C_0^1(\overline{\Omega})$ and η_n as above. Since η_n vanishes in a neighborhood of K and ζ vanishes on $\partial\Omega$,

$$\int_{\Omega} \mathbb{P}_{\Omega}[\eta_n] \nabla u \cdot \nabla \zeta dx + \int_{\Omega} \zeta \nabla u \cdot \nabla \mathbb{P}_{\Omega}[\eta_n] dx + \int_{\Omega} u^p \zeta \mathbb{P}_{\Omega}[\eta_n] dx = M \int_{\Omega} |\nabla u|^q \zeta \mathbb{P}_{\Omega}[\eta_n] dx.$$

Letting n to infity and using the fact that $\nabla u \in L^2(\Omega)$ and $\nabla \mathbb{P}_{\Omega}[\eta_n] \rightarrow 0$ in $L^2(\Omega)$, we derive

$$\int_{\Omega} \nabla u \cdot \nabla \zeta dx + \int_{\Omega} u^p \zeta dx = M \int_{\Omega} |\nabla u|^q \zeta dx.$$

Hence u is a nonnegative bounded weak solution of

$$\begin{aligned} -\Delta u + |u|^{p-1}u - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega. \end{aligned} \quad (2.54)$$

It is therefore C^2 . Again, by the maximum principle we see that u cannot achieve a positive maximum in Ω , this yields a contradiction.

Next we assume $q = \frac{2p}{p+1}$. We choose $b = \frac{r-1}{p-1}$ and (2.38) becomes

$$-\Delta v + \left(1 - b - \frac{Mpb^{\frac{p-1}{p+1}} \epsilon^{\frac{p+1}{p}}}{p+1} \right) \frac{|\nabla v|^2}{v} + \left(\frac{1}{b} - \frac{Mb^{\frac{p-1}{p+1}}}{(p+1)\epsilon^{p+1}} \right) v^r \leq 0. \quad (2.55)$$

From there the argument is similar to the one of Step 2-Case $q = \frac{2p}{p+1}$ in the proof of Theorem 1.1: we claim that for some suitable choices the function v satisfies

$$\begin{aligned} -\Delta v + \tau v^r &\leq 0 && \text{in } \Omega \\ v &= 0 && \text{in } \partial\Omega \setminus K. \end{aligned}$$

We first choose $\epsilon > 0$ so that (2.40) holds, hence the coefficient of v , say τ is positive. Then the expression

$$1 - b - \frac{Mpb^{\frac{p-1}{p+1}}\epsilon^{\frac{p+1}{p}}}{p+1} = \frac{p(r-1)}{p-1} \left(\left(\frac{m_r^{**}}{p+1} \right)^{\frac{p+1}{p}} - \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \right) - \frac{Mpb^{\frac{p+1}{p-1}}}{p+1} \tilde{\epsilon} \quad (2.56)$$

is positive provided $\tilde{\epsilon} > 0$ is small enough. Since $\text{cap}_{\frac{2}{r}, r'}^{\partial\Omega}(K) = 0$ it follows from [18] that $v = 0$. Hence $u = 0$, which ends the proof. \square

2.4 Measure boundary data

Let μ be a nonnegative Radon measure on $\partial\Omega$. The results concerning the following two types of equations

$$\begin{aligned} -\Delta v + v^p &= 0 && \text{in } \Omega \\ v &= \mu && \text{in } \partial\Omega, \end{aligned} \quad (2.57)$$

and

$$\begin{aligned} -\Delta w &= M|\nabla w|^q && \text{in } \Omega \\ w &= c\mu && \text{in } \partial\Omega, \end{aligned} \quad (2.58)$$

allow us to consider the measure boundary data for equation (1.1). We recall the results concerning (2.57) and (2.58).

1- Assume $p > 1$. If μ satisfies

$$\text{For all Borel set } E \subset \partial\Omega, \text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(E) = 0 \implies \mu(E) = 0, \quad (2.59)$$

then problem (2.57) admits a necessarily unique weak solution $v := v_\mu$, see [18], i.e. $v_\mu \in L^1(\Omega) \cap L_\rho^p(\Omega)$ and for any function $\zeta \in \mathbb{X}(\Omega) := \{\eta \in C_0^1(\overline{\Omega}) \text{ s.t. } \Delta\eta \in L^\infty(\Omega)\}$, there holds

$$\int_{\Omega} (-v\Delta\zeta + v^p\zeta) dx = - \int_{\Omega} \frac{\partial\zeta}{\partial\mathbf{n}} d\mu. \quad (2.60)$$

Notice that there is no condition on μ if $1 < p < \frac{N+1}{N-1}$.

2- Assume $1 < q < 2$. If there exists $C > 0$ such that μ satisfies

$$\text{For all Borel set } E \subset \partial\Omega, \mu(E) \leq C \text{cap}_{\frac{2-q}{2}, q'}^{\partial\Omega}(E), \quad (2.61)$$

then problem (2.58) admits at least a positive solution w for $c > 0$ small enough, see [9, Theorem 1.3], in the sense that $w \in L^1(\Omega)$, $\nabla w \in L_\rho^q(\Omega)$ and for any $\zeta \in \mathbb{X}(\Omega)$, there holds

$$\int_{\Omega} (-w\Delta\zeta - M|\nabla w|^q\zeta) dx = - \int_{\Omega} \frac{\partial\zeta}{\partial\mathbf{n}} d\mu. \quad (2.62)$$

Notice that if $1 < q < \frac{N+1}{N}$ there is no capacity condition on μ .

We use also the following result.

Lemma 2.5 *Let $p > \frac{N+1}{N-1}$ and $\mu \in \mathfrak{M}_+(\partial\Omega)$. If $\mu \in W^{-\frac{2}{p},p}(\partial\Omega)$, then there exists $C > 0$ such that*

$$\mu(E) \leq C \left(\text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(E) \right)^{\frac{1}{p'}} \quad \text{for all Borel set } E \subset \partial\Omega. \quad (2.63)$$

Conversely, if μ satisfies

$$\mu(E) \leq C \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(E) \quad \text{for all Borel set } E \subset \partial\Omega, \quad (2.64)$$

for some $C > 0$, then $\mu \in W^{-\frac{2}{p},p}(\partial\Omega)$.

Proof. Assume $\mu \in W^{-\frac{2}{p},p}(\partial\Omega) \cap \mathfrak{M}_+(\partial\Omega)$. If E is a compact subset of $\partial\Omega$ and $\zeta \in C^2(\partial\Omega)$ with $0 \leq \zeta \leq 1$, with $\zeta = 1$ on E , then

$$\mu(E) \leq \int_{\partial\Omega} \zeta d\mu = \langle \mu, \zeta \rangle \leq \|\mu\|_{W^{-\frac{2}{p},p}} \|\zeta\|_{W^{\frac{2}{p},p}}.$$

Therefore, by the definition of the capacity,

$$\mu(E) \leq \|\mu\|_{W^{-\frac{2}{p},p}} \left(\text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(E) \right)^{\frac{1}{p'}}.$$

Conversely, if (2.64) holds, then there exists c_{16} such that for any $0 < c \leq c_{16}$ there exists a $z_{c\mu}$ to

$$\begin{aligned} -\Delta z &= z^p & \text{in } \Omega \\ z &= c\mu & \text{in } \partial\Omega, \end{aligned} \quad (2.65)$$

(see [9, Theorem 1.5]) in the sense that $z_{c\mu} \in L^1(\Omega) \cap L^p_\rho(\Omega)$ and $c\mathbb{P}_\Omega[\mu] \leq z_{c\mu}$. Hence $\mathbb{P}_\Omega[\mu] \in L^p_\rho(\Omega)$, which implies $\mu \in W^{-\frac{2}{p},p}(\Omega)$ by [18]. \square

Those weak solutions are characterized by their boundary trace. Let $\Sigma_\epsilon = \{x \in \Omega : \rho(x) = \epsilon > 0\}$ and $\Sigma_0 = \partial\Omega$. For $0 < \epsilon \leq \epsilon_0$ the hypersurfaces Σ_δ defines a foliation of the set $\Omega_{\epsilon_0} = \{x \in \Omega : 0 < \rho(x) \leq \epsilon_0\}$. Let $\pi(x)$ be the orthogonal projection of $x \in \Omega_{\epsilon_0}$ on $\partial\Omega$. Then $|x - \pi(x)| = \rho(x)$ and $\mathbf{n}_x = (\rho(x))^{-1}(\pi(x) - x)$. The mapping

$$x \mapsto \Pi(x) = (\rho(x), \pi(x)),$$

from Ω_{ϵ_0} onto $(0, \epsilon_0] \times \Sigma_0$ is a C^2 diffeomorphism and the restriction Π_ϵ of Π to Σ_ϵ is a C^2 diffeomorphism from Σ_ϵ onto Σ_0 . Let dS_ϵ be the surface measure on Σ_ϵ , then a continuous function u defined in Ω has boundary trace the Radon measure μ on $\partial\Omega$ if

$$\lim_{\epsilon \rightarrow 0} \int_{\Sigma_\epsilon} u Z dS_\epsilon = \int_{\Sigma} Z d\mu \quad \text{for all } Z \in C(\overline{\Omega}). \quad (2.66)$$

Equivalently, if $\zeta \in C(\partial\Omega)$ and $\zeta_\epsilon = \zeta \circ \Pi_\epsilon^{-1} \in C(\Sigma_\epsilon)$, then

$$\lim_{\epsilon \rightarrow 0} \int_{\Sigma_\epsilon} u \zeta_\epsilon dS_\epsilon = \int_{\Sigma} \zeta d\mu \quad \text{for all } \zeta \in C(\partial\Omega). \quad (2.67)$$

The functions v_μ solution of (2.57) and w solution of (2.58) admit for respective boundary trace μ and $c\mu$. Furthermore, for the equations in (2.57) and (2.58), the existence of a boundary trace of a positive solution is equivalent to the fact that $v_\mu \in L^1(\Omega) \cap L_\rho^p(\Omega)$ and $w \in L^1(\Omega)$ with $\nabla w \in L_\rho^q(\Omega)$ respectively.

Proof of Theorem 1.3. If we assume that (1.16) holds, the measure μ is Lipschitz continuous with respect to $cap_{\frac{2}{p}, p'}^{\partial\Omega}$ and $cap_{\frac{2-q}{q}, q'}^{\partial\Omega}$. By [9, Theorem 1.3] there exists $c_{17} > 0$ such that for any $0 < c \leq c_{17}$ there exists a weak solution $w = w_{c\mu}$ to (2.58) and there holds for some positive constant c_{18} depending on q and Ω

$$w_{c\mu} \leq c_{18} c \mathbb{P}_\Omega[\mu]. \quad (2.68)$$

By [18] there exists a unique solution $v_{c\mu}$ to (2.57) with μ replaced by $c\mu$. The functions $w_{c\mu}$ and $v_{c\mu}$ are respectively supersolution and subsolution of (2.57) with boundary data $c\mu$ and there holds,

$$v_{c\mu} \leq c \mathbb{P}_\Omega[\mu] \leq w_{c\mu} \quad (2.69)$$

Hence there exists a nonnegative function u satisfying (1.1) and such that

$$0 \leq v_{c\mu} \leq u \leq w_{c\mu} \leq c_{18} c \mathbb{P}_\Omega[\mu]. \quad (2.70)$$

Moreover $v_{c\mu} \in L_\rho^p(\Omega)$ and $\nabla w_{c\mu} \in L_\rho^q(\Omega)$. Because $v_{c\mu}$ and $w_{c\mu}$ have boundary trace $c\mu$ in the sense of (2.66) and (2.67), the function u has the same property and we denote it by $u_{c\mu}$. Assuming that $c \leq \min\{c_{16}, c_{17}\}$, there exists also $z_{c\mu}$ solution of (2.65) which satisfies $z_{c\mu} \in L_\rho^p(\Omega)$ and $c \mathbb{P}_\Omega[\mu] \leq z_{c\mu}$ by the maximum principle. Therefore $w_{c\mu} \in L_\rho^p(\Omega)$ and finally $u_{c\mu} \in L_\rho^p(\Omega)$.

Let $\phi = \mathbb{G}_\Omega[u_{c\mu}]$, then $\phi \geq 0$ and

$$-\Delta(u_{c\mu} + \phi) = |\nabla u_{c\mu}|^q.$$

The function $u_{c\mu} + \phi$ is a nonnegative superharmonic function in Ω . By Doob's theorem [12, Chapter II], $-\Delta(u_{c\mu} + \phi) \in L_\rho^1(\Omega)$. Hence $|\nabla u_{c\mu}| \in L_\rho^q(\Omega)$. This implies that $u_{c\mu}$ is a weak solution of (1.15). \square

Proof of Corollary 1.4. We use [1, Theorem 5.5.1], with the same cases (a), (b), (c) and (d), and we denote by K is any compact subset of $\partial\Omega$ and by A a positive constant the value of which may vary from one case to another.

(a) If $q > \frac{2p}{p+1}$ and $p > \frac{N+1}{N-1}$, equivalently $\frac{2-q}{q-1} < \frac{2}{p-1} < N-1$, then

$$cap_{\frac{2-q}{q}, q'}^{\partial\Omega}(K) \leq A \left(cap_{\frac{2}{p}, p'}^{\partial\Omega}(K) \right)^{\frac{N-1-\frac{2-q}{q-1}}{N-1-\frac{2}{p-1}}}. \quad (2.71)$$

Since $\frac{N-1-\frac{2-q}{2}}{N-1-\frac{2}{p-1}} > 1$, thus

$$\text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) \leq c_{21} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K). \quad (2.72)$$

If (1.17) holds, then

$$\begin{aligned} \mu(E) &\leq C \text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) = C \min \left\{ \text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K), c_{21} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K) \right\} \\ &\leq C \max\{1, c_{21}\} \min \left\{ \text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K), \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K) \right\} \end{aligned}$$

and the proof follows by Theorem 1.3.

(b) If $q = \frac{2p}{p-1}$ and $p > \frac{N+1}{N-1}$, then $p > q$, thus

$$\text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) \leq c_{22} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K). \quad (2.73)$$

The proof follows as in (a).

(c) If $p = \frac{N+1}{N-1}$ and $q > \frac{2p}{p+1}$, then for some $A > \text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(\partial\Omega)$,

$$\left(\ln \frac{A}{\text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K)} \right)^{-1} \leq A \left(\text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K) \right)^{\frac{2}{N-1}}. \quad (2.74)$$

Since for any $r \geq 1$

$$(\ln r)^{-1} > \frac{2}{N-1} r^{-\frac{2}{N-1}},$$

we deduce

$$\text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) \leq \left(\frac{N-1}{2} \right)^{\frac{N-1}{2}} A^{\frac{N+1}{2}} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K) := c_{23} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K). \quad (2.75)$$

The proof follows as in (a).

(d) If $p = \frac{N+1}{N-1}$ and $q = \frac{2p}{p+1} = \frac{N+1}{N}$, then as above

$$\left(\text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) \right)^{q-1} \leq A \left(\text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K) \right)^{p-1} \implies \text{cap}_{\frac{2-q}{q},q'}^{\partial\Omega}(K) \leq c_{24} \text{cap}_{\frac{2}{p},p'}^{\partial\Omega}(K). \quad (2.76)$$

The proof follows. □

Proof of Corollary 1.5. We adapt again the formulation of [1, Theorem 5.5.1] to our framework permuting the two capacities and only statement (a) and (c) therein apply.

(a) If $\frac{N+1}{N} < q < \frac{2p}{p+1}$ there exists a constant $A > 0$ such that if $K \subset \partial\Omega$ is a compact set, then

$$\text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(K) \leq A \left(\text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(K) \right)^{\frac{N-1-\frac{2}{p-1}}{N-1-\frac{2-q}{q-1}}}. \quad (2.77)$$

Since $\frac{N+1}{N} < q < \frac{2p}{p+1}$ is equivalent to $N-1-\frac{2}{p-1} > N-1-\frac{2-q}{q-1} > 0$, we deduce

$$\text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(K) \leq c_{25} \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(K). \quad (2.78)$$

We end the proof as in the proof of Corollary 1.4(a).

(c) If $\frac{N+1}{N} = q < \frac{2p}{p+1}$, then for some $A > \text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(\partial\Omega)$,

$$\left(\ln \frac{A}{\text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(K)} \right)^{-1} \leq A \left(\text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(K) \right)^{\frac{1}{N}}. \quad (2.79)$$

Since for $r > 1$,

$$(\ln r)^{-1} > \frac{1}{N} r^{-\frac{1}{N}},$$

we infer

$$\text{cap}_{\frac{2}{p}, p'}^{\partial\Omega}(K) \leq N^N A^{N+1} \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(K) := c_{26} \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(K), \quad (2.80)$$

and the proof follows. \square

The proof in the partially sub-critical case is simpler.

Proof of Corollary 1.6. If $1 < p < \frac{N+1}{N-1}$ for any $\mu \in \mathfrak{M}_+(\partial\Omega)$ problem (2.57) admits a unique solution v_μ (see [15]). If $1 < q < \frac{N+1}{N}$, then there exists $a_0 > 0$ such that for any non-empty Borel set $E \subset \partial\Omega$, $\text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(E) \geq a_0$. Therefore

$$\mu(E) \leq \|\mu\|_{\mathfrak{M}} \leq \frac{\|\mu\|_{\mathfrak{M}}}{a_0} \text{cap}_{\frac{2-q}{q}, q'}^{\partial\Omega}(E).$$

It follows from [9, Theorem 1.3] that problem (2.58) admits a solution w_μ whenever $\|\mu\|_{\mathfrak{M}}$ is small enough. By [10] problem (2.65) admits a solution z_μ with $c\mu$ replaced by μ provided $\|\mu\|_{\mathfrak{M}}$ is small enough. Furthermore

$$w_\mu \leq \mathbb{P}_\Omega[\mu] \leq z_\mu. \quad (2.81)$$

Since $z_\mu \in L_\rho^p(\Omega)$, $w_\mu \in L_\rho^p(\Omega)$. Hence by the same argument as in Theorem 1.3, there exists a solution u_μ of (1.1) which satisfies $v_\mu \leq u_\mu \leq w_\mu$. Hence $u_\mu \in L_\rho^p(\Omega)$ and by the previous argument $\nabla u_\mu \in L_\rho^q(\Omega)$. This implies again that u_μ is a weak solution of (1.15).

If $1 < p < \frac{N+1}{N-1}$ and $\frac{N+1}{N} \leq q < 2$, then problem (2.57) is uniquely solvable for any $\mu \in \mathfrak{M}_+(\partial\Omega)$, while problem (2.65) admits a solution z_μ with $c\mu$ replaced by

μ provided $\|\mu\|_{\mathfrak{M}}$ is small enough and since (2.61) holds, problem (2.58) admits a weak solution provided $0 < c \leq c_0$. Since (2.81) holds with $z_\mu \in L_\rho^p(\Omega)$, the result follows as above.

If $p \geq \frac{N+1}{N-1}$, $1 < q < \frac{N+1}{N}$ and $\mu \in \mathfrak{M}_+(\partial\Omega)$ absolutely continuous with respect to $\text{cap}_{\frac{2}{p}, p}^{\partial\Omega}$, there exists u_μ solution of (2.57) and w_μ solution of (2.58) provided $c\|\mu\|_{\mathfrak{M}}$ is small enough. Since $|\nabla w_\mu|^q \in L_\rho^1(\Omega)$ the function w_μ belongs to the Marcinkiewicz space $M_\rho^{\frac{N+1}{N-1}}(\Omega)$ (see eg. [31]). Since $M_\rho^{\frac{N+1}{N-1}}(\Omega) \subset L_\rho^p(\Omega)$ as $1 < p < \frac{N+1}{N-1}$, it implies that w_μ and therefore u_μ , belongs to $L_\rho^p(\Omega)$. The end of the proof is as before. \square

3 Separable solutions

Separable solutions of (1.1) in $\mathbb{R}^N \setminus \{0\}$ are solutions which have the form

$$u(x) = u(r, \sigma) = r^{-\kappa} \omega(\sigma) \quad \text{for } (r, \sigma) \in \mathbb{R}_+ \times S^{N-1}.$$

This forces q to be equal to $\frac{2p}{p+1}$, κ to $\frac{2}{p-1}$ (recall that this defines α) and ω satisfies

$$-\Delta' \omega + \alpha(N-2-\alpha)\omega + |\omega|^{p-1}\omega - M(\alpha^2\omega^2 + |\nabla' \omega|^2)^{\frac{p}{p+1}} = 0 \quad \text{in } S^{N-1}. \quad (3.1)$$

Constant positive solutions are solutions of

$$X^{p-1} - M\alpha^{\frac{2p}{p+1}} X^{\frac{p-1}{p+1}} + \alpha(N-2-\alpha) = 0. \quad (3.2)$$

This existence of solutions to (3.2) and their stability properties will be detailed in a forthcoming article [8]. The understanding of boundary singularities of solutions of (1.1) is conditioned by the knowledge of separable solutions in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}^N \setminus \{0\}$. Then ω is a solution of

$$\begin{aligned} -\Delta' \omega + \alpha(N-2-\alpha)\omega + |\omega|^{p-1}\omega - M(\alpha^2\omega^2 + |\nabla' \omega|^2)^{\frac{p}{p+1}} &= 0 & \text{in } S_+^{N-1} \\ \omega &= 0 & \text{in } \partial S_+^{N-1}. \end{aligned} \quad (3.3)$$

3.1 Existence of singular solutions

We recall the following result proved in [32, Corollary 1.4.5] is a variant of Boccardo-Murat-Puel's result [11, Theorem 2.1] dealing with the quasilinear equation in a domain $G \subset \mathbb{R}^N$.

$$\mathcal{Q}(u) := -\Delta u + B(\cdot, u, \nabla u) = 0 \quad \text{in } \mathcal{D}'(G), \quad (3.4)$$

where $B \in C(G \times \mathbb{R} \times \mathbb{R}^N)$ satisfies, for some continuous increasing function Γ from \mathbb{R}^+ to \mathbb{R}^+ ,

$$|B(x, r, \xi)| \leq \Gamma(|r|)(1 + |\xi|^2) \quad \text{for all } (x, r, \xi) \in G \times \mathbb{R} \times \mathbb{R}^N. \quad (3.5)$$

Theorem 3.1 *Let G be a bounded domain in \mathbb{R}^N . If there exist a supersolution ϕ and a subsolution ψ of the equation $\mathcal{Q}v = 0$ belonging to $W^{1,\infty}(G)$ and such that $\psi \leq \phi$, then for any $\chi \in W^{1,\infty}(G)$ satisfying $\psi \leq \chi \leq \phi$ there exists a function $u \in W^{1,2}(G)$ solution of $\mathcal{Q}u = 0$ such that $\psi \leq u \leq \phi$ and $u - \chi \in W_0^{1,2}(G)$.*

Remark. *Mutatis mutandi*, the same result holds if \mathbb{R}^N is replaced by a Riemannian manifold.

Their result is actually more general since the Laplacian can be replaced by a quasilinear p -Laplacian-type operator and B by a perturbation with the *natural* p -growth. This theorem has direct applications in the construction of solutions on S_+^{N-1} , but also for the construction of singular solutions in several configurations.

Proposition 3.2 *Let Ω be a bounded smooth domain containing 0, $p > 1$, $1 \leq q \leq 2$ and $M \in \mathbb{R}$. Assume that equation*

$$-\Delta u + u^p - M|\nabla u|^q = 0, \quad (3.6)$$

admits a radial positive and decreasing solution v in $\mathbb{R}^N \setminus \{0\}$ satisfying

$$\lim_{|x| \rightarrow 0} v(x) = \infty. \quad (3.7)$$

Then there exists a positive function u satisfying (3.6) in $\Omega \setminus \{0\}$, vanishing on $\partial\Omega$ and such that

$$(v(x) - \max\{v(z) : |z| = \delta_0\})_+ \leq u(x) \leq v(x) \quad \text{for all } x \in \Omega \setminus \{0\}. \quad (3.8)$$

where $\delta_0 = \text{dist}(0, \partial\Omega)$.

Proof. Put $m = \max\{v(z) : |z| = \delta_0\} = v(\delta_0)$. The function $v_m = (v - m)_+$ is a radial subsolution of (3.6) in Ω , positive in $B_{\delta_0} \setminus \{0\}$ and vanishing in $\Omega \setminus B_{\delta_0}$. For $\epsilon > 0$ set $\Omega_\epsilon = \Omega \setminus \overline{B_\epsilon}$. Since v_m is dominated by the supersolution v , there exists a solution u_ϵ of (3.6) in Ω_ϵ such that $v_m \leq u_\epsilon \leq v$ and $u_\epsilon - v_m \in H_0^1(\Omega_\epsilon)$. By standard regularity estimates, u_ϵ is C^2 , hence it solves

$$\begin{aligned} -\Delta u_\epsilon + u_\epsilon^p - M|\nabla u_\epsilon|^q &= 0 && \text{in } \Omega_\epsilon \\ u_\epsilon &= v_m && \text{on } \partial B_\epsilon \\ u_\epsilon &= 0 && \text{on } \partial\Omega. \end{aligned} \quad (3.9)$$

Notice that u_ϵ is unique by the comparison principle. If $0 < \epsilon' < \epsilon$ the function $u_{\epsilon'}$ solution of (3.9) in $\Omega_{\epsilon'}$ with the corresponding boundary data is larger than v_m and in particular $u_{\epsilon'}|_{\partial B_{\epsilon'}} \geq v_m|_{\partial B_{\epsilon'}} = u_\epsilon|_{\partial B_{\epsilon'}}$. Hence $u_{\epsilon'} \geq u_\epsilon$ in Ω_ϵ . When $\epsilon \downarrow 0$, u_ϵ increase and converges in the $C_{loc}^{1,\theta}(\overline{\Omega} \setminus \{0\})$ -topology toward some function u which satisfies (3.6) in $\Omega \setminus \{0\}$, is larger than v_m and smaller than v , vanishes on $\partial\Omega$ and such that (3.9) holds. \square

The previous result can be adapted to the study of solutions with a boundary singularity in bounded domains which are flat enough near the singular point or in \mathbb{R}_+^N .

Proposition 3.3 *Let $p > 1$, $1 \leq q \leq 2$ and $M \in \mathbb{R}$. Assume that the equation (3.6) admits a positive solution w in \mathbb{R}_+^N belonging to $C(\overline{\mathbb{R}_+^N} \setminus \{0\})$, radially decreasing in \mathbb{R}_+^N and satisfying*

$$\lim_{t \rightarrow 0} w(t\sigma) = \infty \quad \text{uniformly on compact sets } K \subset S_+^{N-1}. \quad (3.10)$$

Assume also

(i) either $w|_{\partial\mathbb{R}_+^N \setminus \{0\}}$ is bounded,

(ii) or $\Omega \subset \mathbb{R}_+^N$ is a bounded smooth domain such that $0 \in \partial\Omega$ starshaped with respect to 0 and such that $w|_{\partial\Omega \setminus \{0\}}$ is bounded.

Then there exists a positive function u satisfying (3.6) in \mathbb{R}_+^N in case (i), or Ω in case (ii), vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ in case (i), or $\partial\Omega \setminus \{0\}$ in case (ii), and such that

$$(w(x) - \sup \{w(z) : z \in \partial\mathbb{R}_+^N \setminus \{0\}\})_+ \leq u(x) \leq w(x) \quad \text{for all } x \in \mathbb{R}_+^N, \quad (3.11)$$

where $K = \sup \left\{ \limsup_{|z| \rightarrow \infty} w(z), \sup \{w(z) : z \in \partial\mathbb{R}_+^N \setminus \{0\}\} \right\}$ in case (i) or

$$(w(x) - \sup \{w(z) : z \in \partial\Omega \setminus \{0\}\})_+ \leq u(x) \leq w(x) \quad \text{for all } x \in \Omega. \quad (3.12)$$

in case (ii).

Proof. The proof is a variant of the preceding one, only the geometry of the domains is changed.

In case (ii) set $m = \sup \{w(z) : z \in \partial\Omega \setminus \{0\}\}$. Then the function $z \mapsto w_m := (w(z) - m)_+$ is a subsolution of (3.6) in Ω . It vanishes on $\partial\Omega \setminus \{0\}$ and is dominated by w . For $\epsilon < \delta_0$, let Ω_ϵ denote $\Omega \cap \overline{B_\epsilon^c}$. We consider the problem of finding u_ϵ solution of

$$\begin{aligned} -\Delta u_\epsilon + u_\epsilon^p - M|\nabla u_\epsilon|^q &= 0 && \text{in } \Omega_\epsilon \\ u_\epsilon &= w_m && \text{on } \partial B_\epsilon \cap \Omega \\ u_\epsilon &= 0 && \text{on } B_\epsilon^c \cap \partial\Omega. \end{aligned} \quad (3.13)$$

Again since $u_\epsilon - w_m \in H_0^1(\Omega_\epsilon)$ and since w_m is smaller than $w|_{\Omega_\epsilon}$, the solution u_ϵ exists and it satisfies $w_m \leq u_\epsilon \leq w$ in Ω_ϵ . If $0 < \epsilon' < \epsilon$, $u_{\epsilon'}|_{\partial\Omega_\epsilon} \geq u_\epsilon|_{\partial\Omega_\epsilon} = w_m$. Hence $u_{\epsilon'} \geq \epsilon$ in Ω_ϵ . As in the proof of Proposition 3.2 the sequence $\{u_\epsilon\}$ is relatively compact in the $C_{loc}^{1,\theta}(\overline{\Omega} \setminus \{0\})$ -topology, which ends the proof.

In case (i), for $n > 0$ set $K_n = \sup \{w(z) : z \in \partial B_n^+ \setminus \{0\}\}$ where, we recall it, $B_n^+ = B_n \cap \mathbb{R}_+^N$. The function $w_{K_n} = (w - K_n)_+$ is a subsolution of (3.6) in B_n^+ which vanishes on $\partial B_n^+ \setminus \{0\}$ and is smaller than w . For $0 < \epsilon < n$ we denote by $u_{\epsilon,n}$ the unique function satisfying

$$\begin{aligned} -\Delta u_{\epsilon,n} + u_{\epsilon,n}^p - M|\nabla u_{\epsilon,n}|^q &= 0 && \text{in } \Gamma_{\epsilon,n} := B_n^+ \setminus \overline{B_\epsilon^+} \\ u_{\epsilon,n} &= w_m && \text{on } \partial B_\epsilon \cap \mathbb{R}_+^N \\ u_{\epsilon,n} &= 0 && \text{on } (\partial B_n^+ \cap \mathbb{R}_+^N) \cup (\overline{\Gamma_{\epsilon,n}} \cap \partial\mathbb{R}_+^N). \end{aligned} \quad (3.14)$$

For $\epsilon' \leq \epsilon < n \leq n'$ there holds $w_{K_n} \leq u_{\epsilon,n} \leq u_{\epsilon',n'} \leq w$ in $\Gamma_{\epsilon,n}$. Letting $n \rightarrow \infty$ and $\epsilon \rightarrow 0$ there exists a subsequence still denoted by $\{u_{\epsilon,n}\}$ which converges to a solution of u of (3.6) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and satisfying (3.11). \square

Remark. The assumption that $w|_{\partial\Omega \setminus \{0\}}$ is bounded is restrictive. For example if $w(t\sigma) = t^{-\alpha}\omega(\sigma)$ the flatness assumption means that $\text{dist}(x, \mathbb{R}_+^N) = O(|x|^{\alpha+1})$ for all $x \in \partial\Omega$ near 0 (remember that $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$). This assumption is always satisfied if $p \geq 3$ since $\alpha \leq 1$, and it can be avoided if there exists a subsolution.

Proposition 3.4 *Let $p > 1$, $1 \leq q \leq 2$ and $M \in \mathbb{R}$. Assume that the equation (3.6) admits a positive supersolution w in \mathbb{R}_+^N belonging to $C(\overline{\mathbb{R}_+^N} \setminus \{0\})$ satisfying (3.10). Assume also*

(i) *either there exists a positive subsolution $Z \in C(\overline{\mathbb{R}_+^N} \setminus \{0\})$ vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$, smaller than w and satisfying (3.10),*

(ii) *or $\Omega \subset \mathbb{R}_+^N$ is a bounded smooth domain such that $0 \in \partial\Omega$ and there exists a positive subsolution $Z \in C(\overline{\Omega} \setminus \{0\})$, vanishing on $\partial\Omega \setminus \{0\}$ such that $Z \leq w|_{\Omega}$ and satisfying (3.10).*

Then there exists a positive function u satisfying (3.6) in \mathbb{R}_+^N (resp. Ω), vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ (resp. $\partial\Omega \setminus \{0\}$) and such that

$$Z(x) \leq u(x) \leq w(x) \quad \text{for all } x \in \mathbb{R}_+^N + \quad (\text{resp. } x \in \Omega). \quad (3.15)$$

Example. If $1 < p < \frac{N+1}{N-1}$ it is proved in [15] that if $\Omega \subset \mathbb{R}_+^N$ is a smooth bounded domain such that $0 \in \partial\Omega$, there exists a nonnegative function $Z_\infty \in C(\overline{\Omega} \setminus \{0\}) \cap C^2(\Omega)$ satisfying the equation

$$\begin{aligned} -\Delta Z + Z^p &= 0 & \text{in } \Omega \\ Z &= 0 & \text{on } \partial\Omega \setminus \{0\}, \end{aligned} \quad (3.16)$$

and such that $t^{\frac{2}{p-1}}Z_\infty(t\sigma) \rightarrow \psi(\sigma)$ uniformly on compact sets $K \subset S_+^{N-1}$ as $t \rightarrow 0$ where ψ is the unique a positive solution of

$$\begin{aligned} -\Delta' \psi + \alpha(N-2-\alpha)\psi + \psi^p &= 0 & \text{in } S_+^{N-1} \\ \psi &= 0 & \text{on } \partial S_+^{N-1}. \end{aligned} \quad (3.17)$$

Furthermore, for any $k > 0$ there exists a nonnegative function $Z_k \in C(\overline{\Omega} \setminus \{0\}) \cap C^2(\Omega)$ satisfying (3.16) and such that $t^{N-1}Z_k(t\sigma) \rightarrow k\phi_1(\sigma)$ where ϕ_1 has been introduced in Theorem 1.11, uniformly on compact subsets of S_+^{N-1} . Furthermore $Z_k \uparrow Z_\infty$ when $k \rightarrow \infty$. If the equation (3.6) admits a positive supersolution w in \mathbb{R}_+^N belonging to $C(\overline{\mathbb{R}_+^N} \setminus \{0\})$ and such that $Z_k \leq w$ in Ω for some $0 < k \leq \infty$, then there exists a positive function u satisfying (3.6) in Ω , vanishing on $\partial\Omega \setminus \{0\}$ and such that

$$Z_k(x) \leq u(x) \leq w(x) \quad \text{for all } x \in \Omega. \quad (3.18)$$

The same result holds if Ω is replaced by \mathbb{R}_+^N .

3.2 Existence or non-existence of separable solutions

Since any large enough constant is a supersolution of (3.1), it follows by Theorem 3.1 that if there exists a nonnegative subsolution $z \in W_0^{1,\infty}(S_+^{N-1})$, there exists a solution in between.

3.2.1 Proof of Theorem 1.9

We recall that ϕ_1 is the first eigenfunction of $-\Delta'$ in $W_0^{1,2}(S_+^{N-1})$ with corresponding eigenvalue $\lambda_1 = N - 1$. Put

$$H(\omega) = -\Delta'\omega + \alpha(N - 2 - \alpha)\omega + |\phi|^{p-1}\omega - M(\alpha^2\omega^2 + |\nabla'\omega|^2)^{\frac{p}{p+1}},$$

then

$$H(\phi_1) = (N - 1 + \alpha(N - 2 - \alpha))\phi_1 + \phi_1^p - M(\alpha^2\phi_1^2 + |\nabla'\phi_1|^2)^{\frac{p}{p+1}}.$$

If ϕ_1 is small enough, there holds $\phi_1^p - M(\alpha^2\phi_1^2 + |\nabla'\phi_1|^2)^{\frac{p}{p+1}} < 0$, hence ϕ_1 is a subsolution. However the condition $N - 1 + \alpha(N - 2 - \alpha) \leq 0$ is too stringent. We can use the fact that, up to a good choice of coordinates, $\phi_1 = \phi_1(\sigma) = \cos \sigma$ with $\sigma \in [0, \frac{\pi}{2}]$. Furthermore the statement " ϕ_1 is small enough " can be achieved by $\phi_1 = \delta \cos \sigma$ with $\delta > 0$ small enough. Then

$$\begin{aligned} & \delta^{-1}H(\delta^{\frac{p+1}{p-1}} \cos \sigma) \\ &= (N - 1 + \alpha(N - 2 - \alpha)) \cos \sigma + \delta^{p+1} \cos^p \sigma - M\delta(\alpha^2 \cos^2 \sigma + \sin^2 \sigma)^{\frac{p}{p+1}}. \end{aligned}$$

The problem is to find $\delta > 0$ such that for all $\sigma \in [0, \frac{\pi}{2}]$ we have $H(\delta^{\frac{p+1}{p-1}} \cos \sigma) \leq 0$. Put $Z = \cos \sigma$ and $\delta^{-1}H(\delta^{\frac{p+1}{p-1}} \cos \sigma) = \delta^{-1}H(\delta^{\frac{p+1}{p-1}} Z) = K_\delta(Z)$, then

$$K_\delta(Z) = (N - 1 + \alpha(N - 2 - \alpha))Z + \delta^{p+1}Z^p - M\delta((\alpha^2 - 1)Z^2 + 1)^{\frac{p}{p+1}},$$

where $0 \leq Z \leq 1$. We use the fact that

$$\alpha^2 \cos^2 \sigma + \sin^2 \sigma \geq \min\{\alpha^2, 1\}(\cos^2 \sigma + \sin^2 \sigma) := \kappa^2 > 0,$$

hence

$$\delta^{-1}H(\delta^{\frac{p+1}{p-1}} \cos \sigma) \leq (N - 1 + \alpha(N - 2 - \alpha)) \cos \sigma + \delta^{p+1} \cos^p \sigma - M\delta\kappa^{\frac{2p}{p+1}}.$$

Then

$$K_\delta(Z) \leq \tilde{K}_\delta(Z) := (N - 1 + \alpha(N - 2 - \alpha))Z + \delta^{p+1}Z^p - M\delta\kappa^{\frac{2p}{p+1}}, \quad (3.19)$$

and

$$\tilde{K}'_\delta(Z) = N - 1 + \alpha(N - 2 - \alpha) + p\delta^{p+1}Z^{p-1}. \quad (3.20)$$

If $N - 1 + \alpha(N - 2 - \alpha) \geq 0$, equivalently $p \geq \frac{N+1}{N-1}$, then $\tilde{K}'_\delta \geq 0$ on $[0, 1]$, hence

$$\tilde{K}_\delta(Z) \leq \tilde{K}_\delta(1) = N - 1 + \alpha(N - 2 - \alpha) + \delta^{p+1} - M\delta\kappa^{\frac{2p}{p+1}}.$$

The function $\delta \mapsto \tilde{K}_\delta(1)$ achieves its minimum for $\delta = \delta_0 := \beta^{\frac{2}{p+1}} \left(\frac{M}{p+1}\right)^{\frac{1}{p}}$ and

$$\tilde{K}_{\delta_0}(1) = N - 1 + \alpha(N - 2 - \alpha) - p\kappa^2 \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}}.$$

Therefore, when $p \geq \frac{N+1}{N-1}$, $K_{\delta_0} \leq 0$ on $[0, 1]$ if

$$\begin{aligned} \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} &\geq \left(\frac{M_{N,p}}{p+1}\right)^{\frac{p+1}{p}} := \frac{N - 1 + \alpha(N - 2 - \alpha)}{p \min\{1, \alpha^2\}} \\ &= \frac{(p+1)(p(N-1) - (N+1))}{p \min\{(p-1)^2, 4\}}. \end{aligned} \quad (3.21)$$

If $N - 1 + \alpha(N - 2 - \alpha) \leq 0$, equivalently $p \leq \frac{N+1}{N-1}$, it is clear from (3.19) that $\tilde{K}_\delta(Z) \leq 0$ for any $Z \in [0, 1]$ as soon as $\delta \leq \kappa^{\frac{1}{p+1}} M^{\frac{1}{p}}$.

Improvement in the case $\alpha > 1$, equivalently $1 < p < 3$. We set

$$F(Z) = \frac{(\alpha^2 - 1)Z^2 + 1}{Z^{\frac{p+1}{p}}}.$$

Then

$$\frac{F'(Z)}{F(Z)} = \frac{(p-1)(\alpha^2 - 1)Z^2 - (p+1)}{p((\alpha^2 - 1)Z^2 + 1)Z}.$$

Since

$$K_\delta(Z) \leq 0 \iff (N - 1 + \alpha(N - 2 - \alpha)) + \delta^{p+1}Z^{p-1} \leq M\delta F^{\frac{p}{p+1}}(Z) \quad (3.22)$$

for all $Z \in (0, 1]$, it is sufficient to prove

$$(N - 1 + \alpha(N - 2 - \alpha)) + \delta^{p+1} \leq M\delta \min_{Z \in (0,1]} F^{\frac{p}{p+1}}(Z) \quad (3.23)$$

The function F is minimal on $(0, 1]$ at $Z = Z_0 = \sqrt{\alpha^2 - 1}$ (remember that $\alpha = \frac{2}{p-1}$) and $F(Z_0) = (\alpha + 2)(\alpha - 1)^{\frac{\alpha+1}{\alpha+2}}$.

If $Z_0 \leq 1$, equivalently $\alpha \geq 2$, inequality (3.23) is satisfied if one find δ such that

$$(N - 1 + \alpha(N - 2 - \alpha)) + \delta^{p+1} \leq M\delta F^{\frac{p}{p+1}}(Z_0),$$

and a sufficient condition is

$$p \left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} \geq p \left(\frac{M_{N,p}}{p+1}\right)^{\frac{p+1}{p}} := \frac{N - 1 + \alpha(N - 2 - \alpha)}{F(Z_0)} \quad (3.24)$$

If $Z_0 > 1$, equivalently $1 < \alpha < 2$, the minimum of F on $(0, 1]$ is achieved at $Z = 1$ with value $F(1) = \alpha^2$, hence a sufficient condition is

$$(N - 1 + \alpha(N - 2 - \alpha)) + \delta^{p+1} \leq M\delta\alpha^{\frac{2p}{p+1}},$$

and we obtain the desired inequality as soon as

$$p \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \geq p \left(\frac{M_{N,p}}{p+1} \right)^{\frac{p+1}{p}} := \frac{N - 1 + \alpha(N - 2 - \alpha)}{\alpha^2}. \quad (3.25)$$

This ends the proof. \square

Remark. Introducing m^{**} defined in (1.11), inequality (3.21) takes the form

$$M \geq \left(\frac{2(p+1)}{\min\{(p-1)^2, 4\}} \right)^{\frac{p}{p+1}} m^{**}, \quad (3.26)$$

in the general case and a more complicated expression in the case $\alpha > 1$.

3.2.2 Non-existence

Theorem 3.5 *Let $p > \frac{N+1}{N-1}$ and $M \leq m^{**}$, defined by (1.11). Then equation (3.1) admits no positive solution.*

Proof. If ω is a positive solution of (3.1) the function η defined by $\omega = \eta^b$ for some $b > 0$ satisfies

$$\begin{aligned} -\Delta' \eta + (1-b) \frac{|\nabla' \eta|^2}{\eta} + \frac{\alpha(N-2-\alpha)}{b} \eta + \frac{1}{b} \eta^{1+(p-1)b} \\ - \frac{M \eta^{\frac{(b-1)(p-1)}{p+1}}}{b} (\alpha^2 \eta^2 + b^2 |\nabla' \eta|^2)^{\frac{p}{p+1}} = 0. \end{aligned}$$

Since for any $\epsilon > 0$ we have by Hölder's inequality,

$$\begin{aligned} \int_{S_+^{N-1}} \eta^{1+\frac{(b-1)(p-1)}{p+1}} (\alpha^2 \eta^2 + b^2 |\nabla' \eta|^2)^{\frac{p}{p+1}} dS \\ \leq \frac{\epsilon^{\frac{p+1}{p}}}{p+1} \int_{S_+^{N-1}} (\alpha^2 \eta^2 + b^2 |\nabla' \eta|^2) + \frac{1}{(p+1)\epsilon^{p+1}} \int_{S_+^{N-1}} \eta^{2+(p-1)b} dS, \end{aligned}$$

it follows that

$$\begin{aligned} \left(2 - b - M \frac{\epsilon^{\frac{p+1}{p}}}{p+1} pb \right) \int_{S_+^{N-1}} |\nabla' \eta|^2 dS + \frac{\alpha}{b} \left(N - 2 - \alpha - M \frac{\epsilon^{\frac{p+1}{p}}}{p+1} \alpha p \right) \int_{S_+^{N-1}} \eta^2 dS \\ + \frac{1}{b} \left(1 - \frac{M}{(p+1)\epsilon^{p+1}} \right) \int_{S_+^{N-1}} \eta^{2+(p-1)b} dS \leq 0. \end{aligned} \quad (3.27)$$

If $b \in (0, 2)$, $\epsilon > 0$ and $M > 0$ are linked by the relation

$$2 - b - M \frac{\epsilon^{\frac{p+1}{p}} pb}{p+1} \geq 0 \iff M \epsilon^{\frac{p+1}{p}} \leq \frac{(2-b)(p+1)}{bp}, \quad (3.28)$$

inequality (3.27) turns into

$$\begin{aligned} & \left((2-b)(N-1) + \frac{\alpha(N-2-\alpha)}{b} - \frac{M \epsilon^{\frac{p+1}{p}} p}{p+1} \left((N-1)b + \frac{\alpha^2}{b} \right) \right) \int_{S_+^{N-1}} \eta^2 dS \\ & + \frac{1}{b} \left(1 - \frac{M}{(p+1)\epsilon^{p+1}} \right) \int_{S_+^{N-1}} \eta^{2+(p-1)b} dS \leq 0. \end{aligned} \quad (3.29)$$

Next we choose

$$\epsilon^{p+1} = \frac{M}{p+1}, \quad (3.30)$$

and we define the function $b \mapsto L(b)$ by

$$L(b) := (2-b)(N-1) + \frac{\alpha(N-2-\alpha)}{b} - p \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \left((N-1)b + \frac{\alpha^2}{b} \right). \quad (3.31)$$

Because $N-1$ is the first eigenvalue of $-\Delta'$ in $W_0^{1,2}(S^{N-1})$, (3.29) combined with (3.30) yields

$$L(b) \int_{S_+^{N-1}} \eta^2 dS \leq 0. \quad (3.32)$$

Furthermore, if inequality (3.28) is strict, and since η is not a first eigenfunction, inequality (3.32) is also strict. Then $L(b) \geq 0$ if

$$p \left(\frac{M}{p+1} \right)^{\frac{p+1}{p}} \leq f(b) := \frac{b(2-b)(N-1) + \alpha(N-2-\alpha)}{(N-1)b^2 + \alpha^2}. \quad (3.33)$$

Now

$$f'(b) = \frac{-2(N-1)^2}{((N-1)b^2 + \alpha^2)^2} (b + \alpha) \left(b - \frac{\alpha}{N-1} \right).$$

Notice that

$$\frac{\alpha}{N-1} \leq 1 \iff p \geq \frac{N+1}{N-1}.$$

If $1 < p \leq \frac{N+1}{N-1}$, then $f' \geq 0$ and in such a case the maximum of f over $(0, 1]$ is achieved at $b = 1$ and for such a value $L(b) \leq 0$.

If $p > \frac{N+1}{N-1}$, then f is increasing on $[0, \frac{\alpha}{N-1})$ and decreasing on $(\frac{\alpha}{N-1}, 1]$, hence the maximum is achieved at $b = \frac{\alpha}{N-1}$, which gives

$$f\left(\frac{\alpha}{N-1}\right) = \frac{N-1-\alpha}{\alpha} = \frac{(N-1)p - (N+1)}{2}. \quad (3.34)$$

Therefore there exists no solution if $p \geq \frac{N+1}{N-1}$ and

$$\left(\frac{M}{p+1}\right)^{\frac{p+1}{p}} \leq \left(\frac{m^{**}}{p+1}\right)^{\frac{p+1}{p}} := \frac{(N-1)p - (N+1)}{2p}. \quad (3.35)$$

□

Remark. Using Theorem 1.1 we can prove the previous result in the case $M < m^{**}$. Indeed, if ω is a positive solution of (3.1), $u_\omega(r, \cdot) = r^{-\frac{2}{p-1}}\omega(\cdot)$ is a positive solution of (1.1) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$. Let $\Omega \subset \mathbb{R}_+^N$ be any smooth domain such that $0 \in \partial\Omega$ and $\partial\Omega$ is flat near 0. Then $u_\omega \leq K$ on $\partial\Omega$ for some $K > 0$. Put $v = (u_\omega - K)_+$, then it is a nonnegative subsolution of (3.1). For any $\epsilon > 0$ small enough there exists a solution u_ϵ of

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^{\frac{2p}{p+1}} &= 0 & \text{in } \Omega_\epsilon := \Omega \cap \overline{B}_\epsilon^c \\ u &= v & \text{on } \partial B_\epsilon \cap \Omega \\ u &= 0 & \text{on } B_\epsilon^c \cap \partial\Omega. \end{aligned} \quad (3.36)$$

Then $v \leq u_\epsilon \leq u_\omega$. Furthermore, for $0 < \epsilon' < \epsilon$, $u_\epsilon \leq u_{\epsilon'}$ in Ω_ϵ . Hence $\{u_\epsilon\}$ converges, when $\epsilon \rightarrow 0$ to a solution u_0 of (1.9), which satisfies $v \leq u_0 \leq u_\omega$ and therefore vanishes on $\partial\Omega \setminus \{0\}$, which is a contradiction.

4 Solutions with an isolated boundary singularity

4.1 Uniqueness of singular solutions

In this section we use scaling transformations to prove uniqueness of singular solutions.

Theorem 4.1 *Assume $N \geq 2$, $p > 1$, $1 < q < 2$ and $M > 0$. Let a such that*

$$\begin{aligned} (i) \quad & 0 \leq a < \beta & \text{if } q \leq \frac{2p}{p+1} \\ (ii) \quad & \beta < a & \text{if } q > \frac{2p}{p+1}. \end{aligned} \quad (4.37)$$

Let $\theta \in C^1(\overline{S_+^{N-1}})$ be a nonnegative function, positive in S_+^{N-1} , vanishing on ∂S_+^{N-1} , and \tilde{a} be a real number. Then there exists at most one nonnegative solution of (1.21) satisfying

$$u(r, \sigma) = r^{-a} |\ln r|^{\tilde{a}} \theta(\sigma) (1 + o(1)) \quad \text{as } r \rightarrow 0. \quad (4.38)$$

Proof. The proof is an adaptation to the configuration where the singularity lies on the boundary of [8, Theorem 1.15]. If u is a solution of (1.1) in \mathbb{R}_+^N , $\ell > 0$ and $b > 0$, define $u_{\ell,b}$ by

$$u_{\ell,b}(x) = \ell^b u(\ell x).$$

Then

$$-\Delta u_{\ell,b} + \ell^{2-b(p-1)} u_{\ell,b}^p - M \ell^{2-q-b(q-1)} |\nabla u_{\ell,b}|^q = 0 \quad \text{in } \mathbb{R}_+^N.$$

If $\ell > 1$, $u_{\ell,b}$ is a supersolution of (1.1) in \mathbb{R}_+^N if and only if

$$\alpha \leq b \leq \beta.$$

These conditions are compatible if and only if $1 < q \leq \frac{2p}{p+1}$. Then we take $b \in (a, \infty) \cap [\alpha, \beta]$, then

$$u_{\ell,b}(r, \sigma) = \ell^{b-a} r^{-a} |\ln r|^{\tilde{a}} \theta(\sigma) (1 + o(1)) \quad \text{as } r \rightarrow 0.$$

By (2.1) all $u(x)$ tends to 0 when $|x| \rightarrow \infty$. Hence, for any $\epsilon > 0$ the super solution $u_{\ell,b} + \epsilon$ which is larger than u for $|x|$ small enough and large enough is larger than another solution \tilde{u} in \mathbb{R}_+^N . Letting $\ell \downarrow 1$ and $\epsilon \rightarrow 0$ yields $u \geq \tilde{u}$. In the same way $\tilde{u} \geq u$.

If $\ell < 1$, $u_{\ell,b}$ is a supersolution of (1.1) in $\frac{1}{\ell}G$ if and only if

$$\beta \leq b \leq \alpha,$$

and these conditions are compatible if and only if $\frac{2p}{p+1} \leq q < 2$. If $\alpha > \beta$ we choose $b \in (0, a) \cap [\beta, \alpha]$ and we conclude as in the first case. \square

Remark. In the case $a = \beta$ a more precise expansion of the singular solution u at $x = 0$ yields uniqueness as it is proved in [8] in the case of an internal singularity. Since the proof of the next result is based upon a easy adaptation of the ideas in [8, Theorem 4.4], we omit it.

Theorem 4.2 *Assume $N \geq 2$, $p > 1$, $1 < q \leq \frac{2p}{p+1}$, $M > 0$ and $a \in [0, \beta]$. Assume θ and $\tilde{\theta}$ are $C^1(\overline{S_+^{N-1}})$ functions positive in S_+^{N-1} and vanishing on ∂S_+^{N-1} and \tilde{a} is a real smaller than a . Then there exists at most one nonnegative solution of (1.21) satisfying*

$$u(r, \sigma) = r^{-a} \theta(\sigma) + r^{-\tilde{a}} \tilde{\theta}(\sigma) (1 + o(1)) \quad \text{as } r \rightarrow 0. \quad (4.39)$$

When problem (1.21) is replaced by (1.8) the scaling method becomes much more delicate to apply. However we give below an easy extension when $\partial\Omega$ is flat near $x = 0$.

Theorem 4.3 *Assume $N \geq 2$, $p > 1$, $1 < q < 2$, $M > 0$ and Ω is a bounded smooth domain such that $0 \in \partial\Omega$ and there exists $\delta > 0$ such that $\partial\Omega \cap B_\delta = T_{\partial\Omega}(0) \cap B_\delta$. Let a such that*

$$\begin{aligned} (i) \quad & 0 \leq a < \beta & \text{if } q \leq \frac{2p}{p+1} \\ (ii) \quad & \beta < a & \text{if } q > \frac{2p}{p+1}. \end{aligned} \quad (4.40)$$

Let $\theta \in C^1(\overline{S_+^{N-1}})$ be a nonnegative function, positive in S_+^{N-1} , vanishing on ∂S_+^{N-1} , and \tilde{a} be a real number. If Ω is starshaped with respect to 0, then there exists at most one nonnegative solution of (1.21) satisfying

$$u(r, \sigma) = r^{-a} |\ln r|^{\tilde{a}} \theta(\sigma) (1 + o(1)) \quad \text{as } r \rightarrow 0. \quad (4.41)$$

Proof. We use the same change of scale as in Theorem 4.1. In case (i) with $\ell > 1$ and $b \in (a, \infty) \cap [\alpha, \beta]$, $u_{\ell,b}$ is a supersolution in $\Omega_\ell = \frac{1}{\ell}\Omega \subset \Omega$ and $\partial\Omega_\ell \cap B_{\delta/\ell} = T_{\partial\Omega}(0) \cap B_{\delta/\ell}$. If \tilde{u} is another solution, $\tilde{u}|_{\partial\Omega_\ell} = h_\ell$ and $h_\ell \rightarrow 0$ uniformly as $\ell \downarrow 1$ since $u \in C^1(\overline{\Omega} \cap B_\epsilon^c)$ for any $\epsilon > 0$. The function $u_{\ell,b} + \max_{\partial\Omega_\ell} h_\ell$ is a supersolution of 1.1 in Ω_ℓ larger than \tilde{u} on $\partial\Omega_\ell \setminus \{0\}$ and near $x = 0$, hence it is larger than \tilde{u} in Ω_ℓ . Letting $\ell \downarrow 1$ yields $u \geq \tilde{u}$.

In case (ii), for $\ell < 1$, $\Omega \subset \Omega_\ell$ and with $b \in (0, a) \cap [\beta, \alpha]$, $u_{\ell,b}$ is a supersolution in Ω larger than \tilde{u} on $\partial\Omega \setminus \{0\}$ and near $x = 0$ it is larger than \tilde{u} in Ω . We conclude as in case (i). \square

4.2 Construction of fundamental solutions

Let Ω be either \mathbb{R}_+^N or a bounded domain with $0 \in \partial\Omega$. A function u satisfying (1.8) is a *fundamental solution* if it has a singularity of potential type, that is

$$\lim_{x \rightarrow 0} \frac{|x|^N u(x)}{\rho(x)} = c_N k, \quad (4.42)$$

for some $k > 0$. The function u can also be looked for as a solution of

$$\begin{aligned} -\Delta u + u^p - M|\nabla u|^q &= 0 && \text{in } \Omega \\ u &= k\delta_0 && \text{in } \partial\Omega, \end{aligned} \quad (4.43)$$

in the sense that $u \in L^p_\rho(\Omega \cap B_r)$, $\nabla u \in L^q_{\rho,loc}(\Omega \cap B_r)$ for any $r > 0$, and for any $\zeta \in C^1_c(\overline{\Omega}) \cap W^{2,\infty}(\Omega)$ there holds

$$\int_{\Omega} (-u\Delta\zeta + u^p\zeta - M|\nabla u|^q\zeta) dx = -k \frac{\partial\zeta}{\partial\mathbf{n}}(0). \quad (4.44)$$

We first consider the problem in \mathbb{R}_+^N .

Proof of Theorem 1.7. The scheme of the proof is surprising since we first show that, in the case $q = \frac{2p}{p+1}$, there exists $M_1 > 0$ such that for any $k > 0$ and any $0 < M < M_1$ there exists a solution. Using this result we prove that if $1 < q < \frac{2p}{p+1}$, then for any $M > 0$ and $k > 0$ there exists a solution. Then we return to the case $q = \frac{2p}{p+1}$ and using the result in the previous case, we prove that when $q = \frac{2p}{p+1}$ we can get rid of the restriction on $M > 0$ and $k > 0$ for the existence of solutions.

I- The case $q = \frac{2p}{p+1}$ and M upper bounded.

For $\ell > 0$ the transformation T_ℓ defined by

$$T_\ell[u(x)] = \ell^{\frac{2}{p-1}} u(\ell x), \quad (4.45)$$

leaves the operator $\mathcal{L}_{\frac{2p}{p+1}, M}$ invariant. We can therefore write

$$T_\ell[u_k] = u_{k\ell^{\frac{2}{p-1}+1-N}},$$

in the sense that if u_k satisfies (4.42) then $T_\ell[u_k]$ satisfies the same limit with k replaced by $k\ell^{\frac{2}{p-1}+1-N}$. However this identity to hold needs some uniqueness for the solutions under consideration satisfying (4.42). This is achieved if u_k is the minimal solution satisfying (4.42) in which case $T_\ell[u_k]$ is the minimal solution satisfying (4.42) with k replaced by $k\ell^{\frac{2}{p-1}+1-N}$. Therefore if there exists a solution to (1.8) in \mathbb{R}_+^N , vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ satisfying (4.47) for some $k > 0$, then there exists such a solution for any $k > 0$.

Step 1- Construction of a subsolution. For $k > 0$ we denote by v_k the solution of

$$\begin{aligned} -\Delta v + v^p &= 0 & \text{in } \mathbb{R}_+^N \\ v &= k\delta_0 & \text{on } \partial\mathbb{R}_+^N \setminus \{0\}. \end{aligned} \quad (4.46)$$

Such a solution exists thanks to [15] if \mathbb{R}_+^N is replaced by a bounded domain Ω . If case of a half-space the problem is first solved in B_n^+ and by letting $n \rightarrow \infty$, we obtain the solution in \mathbb{R}_+^N . Clearly v_k is a subsolution of problem (1.8), and it satisfies

$$\lim_{x \rightarrow 0} \frac{u(x)}{P_N(x)} = k, \quad (4.47)$$

for some $c'_N > 0$, where $P_N(x) = c_N \frac{x^N}{|x|^N}$ is the Poisson kernel in \mathbb{R}_+^N .

Step 2- Construction of a supersolution. It is known that

$$|\nabla P_N(x)|^2 = |x|^{-2N} c^2(x), \quad (4.48)$$

where $c(\cdot)$ is smooth and verifies

$$0 < \tilde{c}_1 \leq c(x) \leq \tilde{c}_2 \quad \text{for some } \tilde{c}_1, \tilde{c}_2 > 0.$$

We construct w_k in \mathbb{R}_+^N of the form

$$w_k = kP_N + w, \quad (4.49)$$

where w satisfies

$$\begin{aligned} -\Delta w + w^p &= a\gamma_2 |x|^{-\frac{2Np}{p+1}} & \text{in } \mathbb{R}_+^N \\ w &= 0 & \text{on } \partial\mathbb{R}_+^N, \end{aligned} \quad (4.50)$$

for some $a > 0$ to be chosen later on. Then

$$\begin{aligned} \mathcal{L}_{\frac{2p}{p+1}, M} w_k &= -\Delta w + (kP_N + w)^p - M (|k\nabla P_N + \nabla w|^2)^{\frac{p}{p+1}} \\ &= (kP_N + w)^p - w^p + a\gamma_2 |x|^{-\frac{2Np}{p+1}} - M (|k\nabla P_N + \nabla w|^2)^{\frac{p}{p+1}} \\ &\geq pkP_N w^{p-1} + a\gamma_2 |x|^{-\frac{2Np}{p+1}} - 2M \left(k^{\frac{2p}{p+1}} \gamma_2^{\frac{2p}{p+1}} |x|^{-\frac{2Np}{p+1}} + |\nabla w|^{\frac{2p}{p+1}} \right). \end{aligned}$$

Now it is easy to check using Osserman's type construction as in [29, Lemma 2.1] and scaling techniques that

$$w(x) \leq \gamma_3 \min \left\{ a^{\frac{1}{p}} |x|^{-\frac{2N}{p+1}}, a |x|^{2(1-\frac{Np}{p+1})} \right\},$$

and

$$|\nabla w(x)| \leq \gamma_4 \min \left\{ a^{\frac{1}{p}} |x|^{-\frac{2N}{p+1}-1}, a |x|^{1-\frac{2Np}{p+1}} \right\}$$

$$\implies |\nabla w(x)|^{\frac{2p}{p+1}} \leq \gamma_5 \min \left\{ a^{\frac{2}{p+1}} |x|^{-\frac{2p(2N+p+1)}{(p+1)^2}}, a^{\frac{2p}{p+1}} |x|^{\frac{2p(p+1-2Np)}{(p+1)^2}} \right\}.$$

Therefore, if we put

$$\tau = \frac{p^2 - 1}{2p(N + 1 - p(N - 1))},$$

then $\tau > 0$ since $N + 1 > p(N - 1)$ and

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1}, M} w_k \geq \gamma_2 \left(a - 2Mk^{\frac{2p}{p+1}} \gamma_2^{\frac{p-1}{p+1}} \right) - 2M\gamma_5 a^{\frac{2p}{p+1}} |x|^{\frac{2p(N+1-p(N-1))}{(p+1)^2}}$$

$$\geq \gamma_2 \left(a - 2Mk^{\frac{2p}{p+1}} \gamma_2^{\frac{p-1}{p+1}} \right) - 2M\gamma_5 a \quad \text{in } B_{a^\tau}^+, \quad (4.51)$$

and similarly,

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1}, M} w_k \geq \gamma_2 \left(a - 2Mk^{\frac{2p}{p+1}} \gamma_2^{\frac{p-1}{p+1}} \right) - 2M\gamma_5 a \quad \text{in } (B_{a^\tau}^+)^c. \quad (4.52)$$

Replacing τ by its value, we obtain a very simple expression from (4.51) and (4.52), valid both in $B_{a^\tau}^+$ and $(B_{a^\tau}^+)^c$, namely

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1}, M} w_k \geq \gamma_2 \left(a - 2Mk^{\frac{2p}{p+1}} \gamma_2^{\frac{p-1}{p+1}} \right) - 2M\gamma_5 a \quad \text{in } \mathbb{R}_+^N. \quad (4.53)$$

When

$$M < M_1 := \frac{\gamma_2}{2\gamma_5}, \quad (4.54)$$

then for fixed k , if we take

$$a > \frac{2M_1 \gamma_2^{\frac{2p}{p+1}} k^{\frac{2p}{p+1}}}{\gamma_2 - 2M\gamma_5},$$

we infer that the right-hand side of (4.53) is nonnegative, hence w_k is a supersolution.

Step 3-Existence. For $0 < k \leq k_0$ w_k is a supersolution which dominates the subsolution v_k . Hence, by [32, Theorem 1-4-6] there exists a solution u_k to (1.8) in \mathbb{R}_+^N , vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and such that $v_k \leq u_k \leq w_k$. Since

$$\lim_{x \rightarrow 0} \frac{v_k(x)}{P_N(x)} = \lim_{x \rightarrow 0} \frac{w_k(x)}{P_N(x)} = k,$$

it follows that u_k inherits the same asymptotic behaviour. Since $k < k_0$ can be replaced by any $k > 0$, the existence of a solution follows.

II- The case $1 < q < \frac{2p}{p+1}$. Assume $M < M_1$, $k > 0$ and \tilde{u}_k is the minimal solution of (1.8) in \mathbb{R}_+^N with $q = \frac{2p}{p+1}$, vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and such that (4.47). Since $|\nabla\phi|^{\frac{2p}{p+1}} \geq |\nabla\phi|^q - 1$, there holds

$$-\Delta\tilde{u}_k + \tilde{u}_k^p + M - M|\nabla\tilde{u}_k|^q \geq 0.$$

Hence $\tilde{u}_k^* = \tilde{u}_k + M^{\frac{1}{p}}$ is a supersolution (1.8) in \mathbb{R}_+^N and it dominates v_k defined in (4.46). By [32, Theorem 1-4-6] there exists a solution u_k of (1.8), vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and satisfying (4.47) under the following weaker form

$$\lim_{t \rightarrow 0} \frac{u_k(tx)}{P_N(tx)} = k \quad \text{uniformly on compact subsets of } \mathbb{R}_+^N. \quad (4.55)$$

Since $|x|^{N-1}u_k(x)$ is uniformly bounded and vanishes on $\partial\mathbb{R}_+^N \setminus \{0\}$, it is bounded in the $C_{loc}^1(\overline{\mathbb{R}_+^N})$ -topology. Hence (4.47) holds. This proves the result when $M < M_1$.

Next let $M > 0$ arbitrary and $k > 0$. In order to find a solution $u := u_k$ to (1.8), we set $u(x) = \ell^{-\frac{2}{p-1}}U_\ell(\frac{x}{\ell})$. Then $\mathcal{L}_{q,M_\ell}u = 0$ is equivalent to

$$\mathcal{L}_{q,M_\ell}U_\ell := -\Delta U_\ell + U_\ell^p - M_\ell|\nabla U_\ell|^q = 0 \quad \text{with } M_\ell = M\ell^{\frac{2p-q(p+1)}{p-1}},$$

and (4.47) is equivalent to

$$\lim_{x \rightarrow 0} \frac{U_\ell(x)}{P_N(x)} = \ell^{\frac{2}{p-1}+1-N}k.$$

Since $2p - q(p+1) > 0$ it is enough to choose $\ell > 0$ such that $M\ell^{\frac{2p-q(p+1)}{p-1}} < M_1$, and we end the proof using the result when $M < M_1$.

III- The case $q = \frac{2p}{p+1}$ revisited. Let $p < \tilde{p} < \frac{N+1}{N-1}$. Then $\frac{2p}{p+1} < \frac{2\tilde{p}}{\tilde{p}+1}$. This implies that for any $M > 0$ and $k > 0$ there exists a positive solution \tilde{u}_k to

$$-\Delta\tilde{u}_k + \tilde{u}_k^{\tilde{p}} - M|\nabla\tilde{u}_k|^{\frac{2p}{p+1}} = 0 \quad \text{in } \mathbb{R}_+^N,$$

vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and such that

$$\lim_{x \rightarrow 0} \frac{\tilde{u}_k(x)}{P_N(x)} = k.$$

Since $\tilde{p} > p$ we have $\tilde{u}_k^{\tilde{p}} > \tilde{u}_k^p - 1$ and therefore

$$-\Delta\tilde{u}_k + \tilde{u}_k^p - M|\nabla\tilde{u}_k|^{\frac{2p}{p+1}} \geq 1 > 0 \quad \text{in } \mathbb{R}_+^N. \quad (4.56)$$

The function \tilde{v}_k solution of

$$\begin{aligned} -\Delta v + v^p &= 0 & \text{in } \mathbb{R}_+^N \\ v &= k\delta_0 & \text{on } \partial\mathbb{R}_+^N \setminus \{0\}, \end{aligned} \quad (4.57)$$

is a subsolution of (4.56), hence there exists a solution u_k of such that $\tilde{v}_k < u_k < \tilde{u}_k$ of (1.8) in \mathbb{R}_+^N , vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and such that (4.42) holds.

IV- The case $\frac{2p}{p+1} < q < \frac{1+N}{N}$. We follow the ideas of Case I. We look for a supersolution w_k of the form (4.49) where w_k satisfies

$$\begin{aligned} -\Delta w + w^p &= a\gamma_2|x|^{-Nq} && \text{in } \mathbb{R}_+^N \\ w &= 0 && \text{on } \partial\mathbb{R}_+^N, \end{aligned} \quad (4.58)$$

for some $a > 0$. Then

$$\begin{aligned} \mathcal{L}_{q,M} w_k &= -\Delta w + (kP_N + w)^p - M(|k\nabla P_N + \nabla w|^2)^{\frac{q}{2}} \\ &= (kP_N + w)^p - w^p + a\gamma_2|x|^{-Nq} - M(|k\nabla P_N + \nabla w|^2)^{\frac{q}{2}} \\ &\geq pkP_N w^{p-1} + a\gamma_2|x|^{-Nq} - 2M(k^q\gamma_2^q|x|^{-Nq} + |\nabla w|^q). \end{aligned}$$

As in Case I, by scaling techniques,

$$w(x) \leq \gamma_3 \min \left\{ a^{\frac{1}{p}}|x|^{-\frac{Nq}{p}}, a|x|^{2-Nq} \right\}$$

and

$$|\nabla w(x)| \leq \gamma_4 \min \left\{ a^{\frac{1}{p}}|x|^{-\frac{Nq}{p}-1}, a|x|^{1-Nq} \right\}.$$

Hence

$$|\nabla w(x)|^q \leq \gamma_5 \min \left\{ a^{\frac{q}{p}}|x|^{-\frac{Nq^2}{p}-q}, a^q|x|^{q(1-Nq)} \right\}.$$

We set

$$\tau = -\frac{1}{2p' - Nq} = -\frac{p-1}{2p - Nq(p-1)}.$$

Then, by the definition of τ ,

$$\begin{aligned} |x|^{Nq} \mathcal{L}_{q,M} w_k &\geq \gamma_2 \left(a - 2Mk^q\gamma_2^{q-1} \right) - 2M\gamma_5 a^q |x|^{q(N+1-Nq)} \\ &\geq \gamma_2 \left(a - 2Mk^q\gamma_2^{q-1} \right) - 2M\gamma_5 a^{\frac{1+N-p(N-1)}{q} - N(p-1)} \quad \text{in } B_{a^\tau}^+, \end{aligned} \quad (4.59)$$

and

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1},M} w_k \geq \gamma_2 \left(a - 2Mk^q\gamma_2^{q-1} \right) - 2M\gamma_5 a^{\frac{1+N-p(N-1)}{q} - N(p-1)} \quad \text{in } (B_{a^\tau}^+)^c. \quad (4.60)$$

We obtain a very simple expression from (4.59) and (4.60), valid both in $B_{a^\tau}^+$ and $(B_{a^\tau}^+)^c$, hence

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1},M} w_k \geq \gamma_2 \left(a - 2Mk^q\gamma_2^{q-1} \right) - 2M\gamma_5 a^{\frac{1+N-p(N-1)}{q} - N(p-1)} \quad \text{in } \mathbb{R}_+^N. \quad (4.61)$$

Using the scaling transformation T_ℓ defined in (4.45), the problem of finding u_k solution of (4.43) is equivalent to looking for a solution of

$$\begin{aligned} -\Delta u + u^p - M\ell^{\frac{2p-q(p+1)}{p-1}} |\nabla u|^q &= 0 && \text{in } \mathbb{R}_+^N \\ u &= k\ell^{\frac{p+1}{p-1}-N} \delta_0 && \text{in } \partial\mathbb{R}_+^N. \end{aligned} \quad (4.62)$$

If we replace M by $M_\ell := M\ell^{\frac{2p-q(p+1)}{p-1}}$ and k by $k_\ell := k\ell^{\frac{p+1}{p-1}-N}$, the inequality (4.53) turns into

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1}, M} w_{k,\ell} \geq \gamma_2 \left(a - 2M_\ell k_\ell^q \gamma_2^{q-1} \right) - 2M_\ell \gamma_5 a^{\frac{1+N-p(N-1)}{\frac{2p}{q}-N(p-1)}} \quad \text{in } \mathbb{R}_+^N, \quad (4.63)$$

where $w_{k,\ell} = w + k_\ell R$ instead of (4.49). Notice that $M_\ell k_\ell^q = M\ell^{\frac{2p}{p-1}-Nq} k^q$. We choose $\ell > 0$ such that $M_\ell k_\ell^q \gamma_2^{q-1} = \frac{a}{4}$, hence

$$|x|^{\frac{2Np}{p+1}} \mathcal{L}_{\frac{2p}{p+1}, M} w_{k,\ell} \geq \frac{a\gamma_2}{2} \left(1 - \gamma_5 \gamma_2^{-q} k^{-q} a^{\frac{1+N-p(N-1)}{\frac{2p}{q}-N(p-1)}} \right) \quad \text{in } \mathbb{R}_+^N. \quad (4.64)$$

It is now sufficient to choose $a > 0$ such that the right-hand side of (4.64) is nonnegative and thus $w_{k,\ell}$ is a supersolution. Since $\tilde{v}_{k,\ell}$ is a subsolution smaller than $w_{k,\ell}$, we end the proof as in Case I.

V- Uniqueness or existence of a minimal solution. If $1 < q \leq \frac{2p}{p+1}$, uniqueness follows from Theorem 4.1 applied with $a = N - 1 < \beta = \frac{2-q}{q-1}$. If $\frac{2p}{p+1} < q < \frac{N+1}{N}$ and if $u_{k,1}$ and $u_{k,2}$ are solutions, they are larger than v_k and the function $u_{k,1,2} = \inf\{u_{k,1}, u_{k,2}\}$ is a supersolution larger than v_k . Hence there exists a solution \tilde{u}_k such that

$$v_k \leq \tilde{u}_k \leq u_{k,1,2}.$$

Let \mathcal{E}_k be the set of nonnegative solutions of (1.8) in \mathbb{R}_+^N , vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and such that (4.42) and put

$$u_k = \inf\{v : v \in \mathcal{E}_k\}.$$

Then there exists a decreasing sequence $\{v_j\}$ such that v_j converges to u_k on a countable dense subset of \mathbb{R}_+^N . By standard elliptic equation regularity theory, v_j converges to u_k on any compact subset of $\overline{\mathbb{R}_+^N} \setminus \{0\}$. Hence u_k is a solution of (1.8) in \mathbb{R}_+^N , it vanishes on $\partial\mathbb{R}_+^N \setminus \{0\}$ and (4.47) since $u_k \geq v_k$. Hence u_k is the minimal solution. \square

Next of we consider the same problem in a bounded domain Ω .

Proof of Theorem 1.8. We give first proof when $\Omega \subset \mathbb{R}_+^N$. We adapt the proof of Theorem 1.7. The solution v_k of

$$\begin{aligned} -\Delta v + v^p &= 0 && \text{in } \Omega \\ v &= k\delta_0 && \text{on } \partial\Omega, \end{aligned} \quad (4.65)$$

is a subsolution for (1.8) in Ω and satisfies (1.24). The solution u_k of (1.8) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and satisfying (4.42) is larger than v_k in Ω . Hence the result follows by Proposition 3.4.

When Ω is not included in \mathbb{R}_+^N , estimates (4.48) is valid with the same type of bounds on c . We also consider separately the cases $q = \frac{2p}{p+1}$ and M upper bounded, $q < \frac{2p}{p+1}$ and $M > 0$ arbitrary and $q = \frac{2p}{p+1}$ and $M > 0$ arbitrary and finally $\frac{2p}{p+1} < q < \frac{N+1}{N}$. As supersolution we consider the function $w_k := kP^\Omega + w$ where w satisfies

$$\begin{aligned} -\Delta w + w^p &= a\gamma_2|x|^{-\frac{2Np}{p+1}} && \text{in } \Omega \\ w &= 0 && \text{on } \partial\Omega, \end{aligned} \quad (4.66)$$

for some $a > 0$. The estimates on w endow the form

$$w(x) \leq \gamma_3 a^{\frac{1}{p}} |x|^{2(1-\frac{Np}{p+1})},$$

and

$$|\nabla w(x)| \leq \gamma_4 a^{\frac{1}{p}} |x|^{1-\frac{2Np}{p+1}},$$

where γ_3 and γ_4 depend on Ω . Hence (4.53) holds in Ω instead of \mathbb{R}_+^N , and we have existence for $M < M_1$, where M_1 is defined by (4.54). Then we prove existence for any $M > 0$ and $k > 0$ when $q < \frac{2p}{p+1}$ then for any $M > 0$ when $q = \frac{2p}{p+1}$ and finally when $\frac{2p}{p+1} < q < \frac{N+1}{N}$ as in Theorem 1.7. \square

4.3 Solutions with a strong singularity

4.3.1 The case $1 < q \leq \frac{2p}{p+1}$

If $p = \frac{N+1}{N-1}$ and $1 < q < \frac{N+1}{N}$ and if $p > \frac{N+1}{N-1}$ and either $1 < q < \frac{2p}{p+1}$ and $M > 0$ or $q = \frac{2p}{p+1}$ and $M > m^{**}$ defined in (1.11), the singularity is removable by Theorem 1.1. Thus the ranges of exponents that we consider are the following,

$$\begin{aligned} (i) \quad & 1 < q \leq \frac{2p}{p+1} \quad \text{and} \quad 1 < p < \frac{N+1}{N-1}, \\ (ii) \quad & (p, q) = \left(\frac{N+1}{N-1}, \frac{N+1}{N} \right). \end{aligned} \quad (4.67)$$

If (4.67)-(i) holds, $q < \frac{N+1}{N}$, and in this range the limit of the fundamental solutions u_k when $k \rightarrow \infty$ is a solution with a strong singularity with an explicit blow-up rate. In the case of a bounded domain our construction requires a geometric *flatness* condition of $\partial\Omega$ near 0. We consider first the case $\Omega = \mathbb{R}_+^N$.

Theorem 4.4 *Assume (4.67)-(i) holds, then for any $M \geq 0$ there exists a positive solution u of (1.1) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ such that*

$$\lim_{x \rightarrow 0} \frac{u(x)}{P_N(x)} = \infty. \quad (4.68)$$

Furthermore,

(i) If $1 < q < \frac{2p}{p+1}$,

$$\lim_{r \rightarrow 0} r^{\frac{2}{p-1}} u(r, \cdot) = \psi \quad \text{uniformly in } S_+^{N-1}, \quad (4.69)$$

where ψ is the unique positive solution of (3.17).

(ii) If $q = \frac{2p}{p+1}$,

$$\lim_{r \rightarrow 0} r^{\frac{2}{p-1}} u(r, \cdot) = \omega \quad \text{uniformly in } S_+^{N-1}, \quad (4.70)$$

where ω is the minimal positive solution of (1.28).

Proof. If $k > 0$, we denote by $u = u_{k,M}$ the solution of

$$\begin{aligned} -\Delta u + u^p &= M|\nabla u|^q && \text{in } \mathbb{R}_+^N \\ u &= k\delta_0 && \text{in } \partial\mathbb{R}_+^N. \end{aligned} \quad (4.71)$$

The mapping $k \mapsto u_k$ is increasing. We set $T_\ell[u] = u_\ell$, where T_ℓ is defined in (4.45).

Since $1 < q \leq \frac{2p}{p+1}$,

$$T_\ell[u_{k,M}] = u_{k\ell^{\frac{2}{p-1}+1-N}, M\ell^{\frac{2p-q(p+1)}{p-1}}}.$$

It follows from Theorem 2.1 and Theorem 2.3 that the sequences $\{u_{k,M}\}$ and $\{\nabla u_{k,M}\}$ converge locally uniformly in \mathbb{R}_+^N , when $k \rightarrow \infty$, to a function $u_{\infty,M}$ which satisfies (1.1) in \mathbb{R}_+^N . Furthermore

$$T_\ell[u_{\infty,M}] = u_{\infty, M\ell^{\frac{2p-q(p+1)}{p-1}}} \quad \text{for all } \ell > 0. \quad (4.72)$$

In the case $q = \frac{2p}{p+1}$ the function $u_{\infty,M}$ is self-similar, hence

$$u_{\infty,M}(r, \sigma) = r^{-\alpha} \tilde{\omega}(\sigma),$$

where $\tilde{\omega}$ is a nonnegative solution of (1.28). Inasmuch $u_{k,M} \geq u_{k,0} = v_k$ (already defined by (4.46)), it follows that

$$u_{\infty,M}(r, \sigma) \geq u_{\infty,0}(r, \sigma) = r^{-\alpha} \psi(\sigma) \implies \tilde{\omega} \geq \psi \quad \text{in } S_+^{N-1}. \quad (4.73)$$

Since $u_{k,M}$ is dominated by any self-similar solution of (1.1), it implies that $\tilde{\omega}$ is the minimal positive solution of (1.28) that we denote by ω hereafter. Up to a subsequence, $\{T_{\ell_n}[u_{\infty,M}]\}$ converges locally uniformly in $\overline{\mathbb{R}_+^N} \setminus \{0\}$ to $u_{\infty,M}$. Consequently

$$\lim_{\ell_n \rightarrow 0} \ell_n^\alpha u_{\infty,M}(\ell_n, \sigma) = \omega(\sigma) \quad \text{uniformly in } S_+^{N-1}.$$

Because of uniqueness, the whole sequence converges, which implies (4.70).

In the case $q < \frac{2p}{p+1}$, using the a priori estimates from Theorem 2.1 and Theorem 2.3, we obtain that $T_{\ell_n}[u_{\infty,M}](1, \sigma) = \ell_n^\alpha u_{\infty,M}(\ell_n, \sigma)$ converges locally uniformly in $\overline{S_+^{N-1}}$ to $u_{\infty,0}(1, \sigma)$. Since $u_{\infty,0}(1, \cdot) \geq \psi$, it follows that

$$\lim_{\ell_n \rightarrow 0} \ell_n^\alpha u_{\infty,M}(\ell_n, \sigma) = \psi(\sigma) \quad \text{uniformly in } S_+^{N-1}.$$

Hence (4.70) follows by uniqueness of the function ψ .

Uniqueness of positive solution of (1.21) satisfying (4.68) follows from Theorem 4.1 applied with $a = \alpha = \frac{2}{p-1} < \frac{2-q}{q-1} = \beta$. □

As a consequence of Theorem 1.9-(ii) we have

Theorem 4.5 *Assume (4.67)-(ii) holds, then for any $M > 0$ there exists a positive separable solution u of (1.1) in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$*

When \mathbb{R}_+^N is replaced by a bounded domain there holds.

Theorem 4.6 *Assume $\Omega \subset \mathbb{R}_+^N$ is a bounded smooth domain such that $0 \in \partial\Omega$ and $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$, and (p, q) satisfies (4.67)-(i). Then for any $M \geq 0$ there exists a positive solution u of (1.1) in Ω vanishing on $\partial\Omega \setminus \{0\}$ such that*

$$\lim_{x \rightarrow 0} \frac{u(x)}{P_\Omega(x)} = \infty, \quad (4.74)$$

where P_Ω is the Poisson kernel in Ω . Furthermore

(i) *If $1 < q < \frac{2p}{p+1}$, then*

$$\lim_{r \rightarrow 0} r^\alpha u(r, \cdot) = \psi \quad \text{locally uniformly in } S_+^{N-1}, \quad (4.75)$$

where ψ is the unique positive solution of

$$\begin{aligned} -\Delta' \psi + \alpha(N-2-\alpha)\psi + \psi^p &= 0 && \text{in } S_+^{N-1} \\ \psi &= 0 && \text{in } \partial S_+^{N-1}. \end{aligned}$$

(ii) *If $q = \frac{2p}{p+1}$, then*

$$\psi \leq \liminf_{r \rightarrow 0} r^\alpha u(r, \cdot) \leq \limsup_{r \rightarrow 0} r^\alpha u(r, \cdot) \leq \omega \quad \text{locally uniformly in } S_+^{N-1}. \quad (4.76)$$

Proof. As in the proof of Theorem 4.4, the sequence $\{u_k\}$ of the solution of (1.8) which satisfy (4.42) is increasing. Since it is bounded from above by the restriction to Ω of the solutions of the same equation in \mathbb{R}_+^N , vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ and satisfying (4.68), it admits a limit u_∞ which is a solution of 1.8 which vanishes on $\partial\Omega \setminus \{0\}$ and satisfies (4.74). In order to have an estimate of the blow-up rate, we recall that the solution v_k of (4.65) is a subsolution of (1.1) and $u_k \geq v_k$. Furthermore $\{v_k\}$ converges to $\{v_\infty\}$ which is a positive solution of (1.1) in Ω , vanishing on $\partial\Omega \setminus \{0\}$ and such that

$$\lim_{r \rightarrow 0} r^\alpha v_\infty(r, \sigma) = \psi(\sigma) \quad \text{locally uniformly in } S_+^{N-1}. \quad (4.77)$$

Combined with (4.69) and (4.70) it implies (4.75) and (4.76) since the solution u_k in Ω is bounded from above by the solution in \mathbb{R}_+^N .

$$\liminf_{r \rightarrow 0} r^\alpha u_\infty(r, \sigma) \geq \psi(\sigma) \quad \text{locally uniformly in } S_+^{N-1}. \quad (4.78)$$

□

Theorem 4.7 Assume $\Omega \subset \mathbb{R}_+^N$ is a bounded smooth domain such that $0 \in \partial\Omega$ and $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$, $p = \frac{N+1}{N-1}$ and $q = \frac{2p}{p+1} = \frac{N+1}{N}$. If

$$\text{dist}(x, \mathbb{R}_+^N) \leq c_{27}|x|^N \quad \text{for all } x \in \partial\Omega \cap B_\delta, \quad (4.79)$$

for some constants $\delta, c_{27} > 0$, then there exists a positive solution u of (1.1) in Ω , vanishing on $\partial\Omega \setminus \{0\}$ such that

$$\lim_{r \rightarrow 0} r^\alpha u(r, \sigma) = \omega(\sigma) \quad \text{locally uniformly in } S_+^{N-1}. \quad (4.80)$$

Proof. The function $u_\omega(r, \cdot) = r^{1-N}\omega$ satisfies (1.1) in \mathbb{R}_+^N and vanishes on $\partial\mathbb{R}_+^N \setminus \{0\}$. Since $\nabla\omega$ is bounded, it satisfies

$$u(x) \leq c_{19} \quad \text{for all } x \in \partial\Omega \setminus \{0\},$$

for some constant $c_{19} > 0$. Then the result follows from Proposition 3.3. \square

4.3.2 The case $\frac{2p}{p+1} < q < p$

If

$$1 < p < \frac{N+1}{N-1} \quad \text{and} \quad \frac{2p}{p+1} < q < \frac{N+1}{N}, \quad (4.81)$$

there exists fundamental solutions u_k in \mathbb{R}_+^N by Theorem 1.7, or in Ω by Theorem 1.8. Since the mapping $k \mapsto u_k$ is increasing and u_k is bounded from above the function $u_\infty = \lim_{k \rightarrow \infty} u_k$ is a solution of (1.1) in \mathbb{R}_+^N (resp. Ω) vanishing on $\mathbb{R}_+^N \setminus \{0\}$ (resp. $\Omega \setminus \{0\}$) which satisfies (4.68) (resp. (4.74)). However the blow-up rate of u_∞ is not easy to obtain from scaling methods since the transformation T_ℓ transform (1.1) into (4.62) where M is replaced by $M\ell^{\frac{2p-q(p+1)}{p-1}}$ which is not bounded when $\ell \rightarrow 0$. When $q > \frac{2p}{p+1}$, the natural exponent is γ defined by (1.34) The transformation S_ℓ defined for $\ell > 0$ by

$$S_\ell[u](x) = \ell^\gamma u(\ell x), \quad (4.82)$$

transforms (1.1) into

$$-\ell^{\frac{q(p+1)-2p}{p-q}} \Delta u + |u|^{p-1}u - M|\nabla u|^q = 0. \quad (4.83)$$

When $\ell \rightarrow 0$, the limit equation is an eikonal equation (up to change of unknown),

$$|u|^{p-1}u - M|\nabla u|^q = 0. \quad (4.84)$$

Separable solutions of (1.3) in \mathbb{R}_+^N are under the form $u_\eta(r, \cdot) = r^{-\gamma}\eta$ and η satisfies

$$|\eta|^{p-1}\eta - M(\gamma^2\eta^2 + |\nabla'\eta|^2)^{\frac{q}{2}} = 0 \quad \text{in } S_+^{N-1}. \quad (4.85)$$

Clearly this equation admits no C^1 solution but for the constant ones. As limit of solutions with vanishing viscosity, the solutions that we obtain are viscosity solutions

outside the origin. We will look for solutions having a strong singularity by the method of sub and supersolutions. Note that (1.3) admits an explicit radial singular solution, namely

$$U(x) = \omega_0 |x|^{-\gamma} := \gamma^\gamma M^{\frac{1}{p-q}} |x|^{-\gamma}. \quad (4.86)$$

Proof of Theorem 1.11. For $n > 0$ set $U_n(r) = nr^{-\gamma}$. As

$$\gamma(p-1) + 2 = -q(p+1) + \gamma + 2 = \frac{2p - q(p+1)}{p-q},$$

we have

$$n^{-1} r^{-2-\gamma} \mathcal{L}_{q,M} U_n = -\gamma(\gamma+2-N) + n^{q-1} (n^{p-q} - \gamma^q M) r^{2-(p-1)\gamma}.$$

Since $\gamma+2-N > 0$ because $q > \frac{2p}{p+1}$ and $p < \frac{N+1}{N-1}$, for any $n > \omega_0$ there exists $r_n > 0$ such that

$$n^{q-1} (n^{p-q} - \gamma^q M) r_n^{2-(p-1)\gamma} = \gamma(\gamma+2-N).$$

It implies that U_n is a super solution of (1.1) in $B_{r_n} \setminus \{0\}$. Furthermore

$$r_n = \left(\frac{n^{p-1}}{\gamma(\gamma+2-N)} \right)^{\frac{1}{(p-1)\gamma-2}} (1 + o(1)) \quad \text{when } n \rightarrow \infty. \quad (4.87)$$

For a subsolution we set

$$W_m(r, \sigma) = mr^{-\gamma} \phi_1(\sigma), \quad (4.88)$$

where $m > 0$. Then

$$\begin{aligned} r^{p\gamma} \mathcal{L}_{q,M} W_m &= -mr^{\frac{q(p+1)-2p}{p-q}} (\gamma^2 - (N-2)\gamma + 1 - N) \phi_1 \\ &\quad + m^q \left(m^{p-q} \phi_1^p - M (\gamma^2 \phi_1^2 + |\nabla' \phi_1|^2)^{\frac{q}{2}} \right), \end{aligned} \quad (4.89)$$

and this expression is negative for $m > 0$ small enough. Set

$$P(X) = X^2 - (N-2)X + 1 - N = (X+1)(X+1-N).$$

Then

$$P(\gamma) = \frac{p(Nq - (N-1)p)}{(p-q)^2}.$$

We first give the proof when $Nq \geq (N-1)p$. In such case $P(\gamma) \geq 0$. Hence there exists $m_0 > 0$ such that for any $0 < m \leq m_0$, W_m is a subsolution in \mathbb{R}_+^N , smaller than U_n and it is bounded on $\partial B_{r_n}^+ \setminus \{0\}$. When $m \leq m_0$, the function W_m defined in (4.88) is a subsolution of (1.1) in \mathbb{R}_+^N . Since W_m is bounded on $\partial B_{r_n}^+ \setminus \{0\}$ there exists a nonnegative solution u_n of (1.1) in $B_{r_n}^+$ which vanishes on $B_{r_n}^+ \setminus \{0\}$ and there holds

$$(W_m(x) - mr_n^{-\gamma})_+ \leq u_n(x) \leq U_n(x) \quad \text{for all } x \in B_{r_n}^+. \quad (4.90)$$

The fact that $B_{r_n}^+$ is just a Lipschitz domain is easily bypassed by smoothing it in a neighborhood of $\partial B_{r_n} \cap \mathbb{R}_+^N$. Furthermore, by (2.1) and (2.18),

$$u_n(x) \leq c_5 \max \left\{ |x|^{-\alpha}, M^{\frac{1}{p-q}} |x|^{-\gamma} \right\}. \quad (4.91)$$

and for any $r_0 > 0$, there exists $c_8 > 0$ depending on r_0 such that

$$|\nabla u_n(x)| \leq c_8 \max \left\{ |x|^{-\alpha-1}, M^{\frac{1}{p-q}} |x|^{-\gamma-1} \right\}. \quad (4.92)$$

By standard local regularity theory, there exists a subsequence $\{u_{n_j}\}$ which converges in the $C^1(K)$ -topology for any compact set $K \subset \overline{\mathbb{R}_+^N} \setminus \{0\}$ to a positive solution u of (1.1) in \mathbb{R}_+^N which vanishes on $\partial \mathbb{R}_+^N \setminus \{0\}$ and satisfies (1.35).

Next we assume $Nq < (N-1)p$. Observe that $\gamma^2 \phi_1^2 + |\nabla' \phi_1|^2 \geq \delta^2 > 0$, then

$$m^p \phi_1^p - Mm^q (\gamma^2 \phi_1^2 + |\nabla' \phi_1|^2)^{\frac{q}{2}} \leq m^p - Mm^q \delta^q.$$

Thus, from (4.89) we obtain

$$r^{p\gamma} \mathcal{L}_{q,M} W_m \leq -mr^{\frac{q(p+1)-2p}{p-q}} P(\gamma) + m^p - Mm^q \delta^q, \quad (4.93)$$

and $P(\gamma) < 0$. If we choose

$$m = \delta^\gamma \left(\frac{M}{2} \right)^{\frac{1}{p-q}},$$

then

$$m^p - Mm^q \delta^q = -\frac{Mm^q \delta^q}{2}.$$

Therefore $\mathcal{L}_{q,M} W_m \leq 0$ on $B_{r_*}^+$ where

$$r_* = \left(\frac{Mm^{q-1} \delta^q}{-2P(\gamma)} \right)^{\frac{p-q}{q(p+1)-2p}}.$$

If $a = mr_*^{-\gamma}$, then $W_m \leq a$ in $\partial B_{r_*}^+$, thus $W_{m,a} = (W_m - a)_+$ is nonnegative in $B_{r_*}^+$ and it is a subsolution of (1.1) in $B_{r_*}^+$ which vanishes on $\partial B_{r_*}^+ \setminus \{0\}$. If we extend it by 0 in \mathbb{R}_+^N , the new function is a subsolution of (1.1) which belongs to $W_{loc}^{1,\infty}(\overline{\mathbb{R}_+^N} \setminus \{0\})$. We end the proof using Proposition 3.4 as in the previous case. \square

If \mathbb{R}_+^N is replaced by a bounded domain we have the following result.

Theorem 4.8 *Let $M > 0$ and $\frac{2p}{p+1} < q < p$. If $\Omega \subset \mathbb{R}_+^N$ is a bounded smooth domain such that $0 \in \partial\Omega$ and $T_{\partial\Omega}(0) = \partial\mathbb{R}_+^N$. If*

$$\text{dist}(x, \partial\mathbb{R}_+^N) \leq c_{20} |x|^{\frac{p}{p-q}} \quad \text{for all } x \in \partial\Omega \text{ near } 0, \quad (4.94)$$

for some constant $c_{23} > 0$. Then there exists a positive solution u of (1.1) in Ω vanishing on $\partial\Omega \setminus \{0\}$ satisfying, for some $m > 0$,

$$m\phi_1(\sigma) \leq \liminf_{r \rightarrow 0} r^\gamma u(r, \sigma) \leq \limsup_{r \rightarrow 0} r^\gamma u(r, \sigma) \leq \omega_0, \quad (4.95)$$

uniformly on any compact set $K \subset S_+^{N-1}$.

Proof. We recall that ϕ_1 is the first eigenfunction of $-\Delta'$ in $W_0^{1,2}(S^{N-1})$. Let $R > 0$ and $B := B_R(a) \subset \Omega$ be an open ball tangent to $\partial\Omega$ at 0. Up to rescaling and since the result does not depend on the value of M we can assume that $R = 1$. We set $w_m(x) = m|x|^{-\theta}P_B(x)$ where $\theta = \gamma + 1 - N$ and P_B is the Poisson kernel in B expressed by

$$P_B(x) = \frac{1 - |x - a|^2}{\sigma_N |x|^N},$$

where σ_N is the volume of the unit sphere in \mathbb{R}^N . Then

$$\begin{aligned} m^{-1}\mathcal{L}_{q,M}w_m &= -(\theta^2 + (2 - N)\theta)|x|^{-\theta-2}P_B(x) + 2\theta|x|^{-\theta-1}\langle \nabla P_B(x), \frac{x}{|x|} \rangle + m^{p-1}|x|^{-p\theta}P_B^p(x) \\ &\quad - Mm^{q-1}\left(\theta^2|x|^{-2(\theta-1)}P_B^2(x) + |x|^{-2\theta}|\nabla P_B(x)|^2 - 2\theta|x|^{-2\theta-1}\langle \nabla P_B(x), \frac{x}{|x|} \rangle\right)^{\frac{q}{2}}. \end{aligned} \quad (4.96)$$

Since

$$\nabla P_B(x) = -\frac{1}{\sigma_N} \left(\frac{N(1 - |x - a|^2)}{|x|^{N+1}} \frac{x}{|x|} + \frac{2(x - a)}{|x|^N} \right),$$

then

$$\begin{aligned} \langle \nabla P_B(x), \frac{x}{|x|} \rangle &= -\frac{1}{\sigma_N |x|^{N+1}} ((N - 1)(1 - |x - a|^2) + |x|^2) \\ &= -\frac{N - 1}{|x|} P_B(x) - \frac{1}{\sigma_N |x|^{N-1}}, \end{aligned}$$

which implies in particular

$$|\nabla P_B(x)| \geq \frac{N - 1}{|x|} P_B(x) + \frac{1}{\sigma_N |x|^{N-1}}.$$

If $q \geq \frac{N-1}{N}p$, equivalently $\theta \geq 0$, we have

$$\begin{aligned} |\nabla w_m|^2 &= \theta^2|x|^{-2(\theta+1)}P_B^2(x) + |x|^{-2\theta}|\nabla P_B(x)|^2 - 2\theta|x|^{-2\theta-1}\langle \nabla P_B(x), \frac{x}{|x|} \rangle \\ &\geq \theta^2|x|^{-2(\theta+1)}P_B^2(x) + |x|^{-2\theta} \left(\frac{N - 1}{|x|} P_B(x) + \frac{1}{\sigma_N |x|^{N-1}} \right)^2 \\ &\quad + 2\theta|x|^{-2\theta-1} \left(\frac{N - 1}{|x|} P_B(x) + \frac{1}{\sigma_N |x|^{N-1}} \right) \\ &\geq (\theta^2 + (N - 1)^2)|x|^{-2(\theta+1)}P_B^2(x). \end{aligned} \quad (4.97)$$

Hence

$$\begin{aligned} m^{-1}\mathcal{L}_{q,M}w_m &\leq -(\theta^2 + N\theta)|x|^{-\theta-2}P_B(x) + m^{p-1}|x|^{-p\theta}P_B^p(x) \\ &\quad - m^{q-1}M(\theta^2 + (N - 1)^2)^{\frac{q}{2}}|x|^{-q(\theta+1)}P_B^q(x) \\ &\leq m^{q-1}|x|^{-p\theta}P_B^q(x) \left(m^{p-q}P_B^{p-q}(x) - M(\theta^2 + (N - 1)^2)|x|^{(p-q)\theta-q} \right). \end{aligned} \quad (4.98)$$

Now

$$P_B(x) \leq \frac{2}{\sigma_N |x|^{N-1}} \implies P_B^{p-q}(x) \leq \left(\frac{2}{\sigma_N}\right)^{p-q} |x|^{(1-N)(p-q)}.$$

Since $(1-N)(p-q) = (p-q)\theta - q$, we obtain finally that,

$$m^{-1} \mathcal{L}_{q,M} w_m \leq m^{q-1} |x|^{-q\theta} P_B^q(x) \left(m^{p-q} \left(\frac{2}{\sigma_N}\right)^{p-q} - M(\theta^2 + (N-1)^2) \right).$$

Choosing m small enough we deduce that w_m is a subsolution in B . If we extend it by 0 in $\Omega \setminus B$, the new function denoted by \tilde{w} is a nonnegative subsolution of (1.1) in Ω which vanishes on $\partial\Omega \setminus \{0\}$ and satisfies (4.95). The proof follows from Proposition 3.4.

If $q < \frac{N-1}{N}p$, then $\theta < 0$. Since $\langle \nabla P_B(x), \frac{x}{|x|} \rangle \leq 0$, (4.97) is replaced by

$$\begin{aligned} |\nabla w_m|^2 &= \theta^2 |x|^{-2(\theta+1)} P_B^2(x) + |x|^{-2\theta} |\nabla P_B(x)|^2 - 2\theta |x|^{-2\theta-1} \langle \nabla P_B(x), \frac{x}{|x|} \rangle \\ &\geq \theta^2 |x|^{-2(\theta+1)} P_B^2(x) + |x|^{-2\theta} \left(\frac{N-1}{|x|} P_B(x) + \frac{1}{\sigma_N |x|^{N-1}} \right)^2 \\ &\quad + 2\theta |x|^{-2\theta-1} \left(\frac{N-1}{|x|} P_B(x) + \frac{1}{\sigma_N |x|^{N-1}} \right) \\ &\geq (\theta^2 + (N-1)^2) |x|^{-2(\theta+1)} P_B^2(x) + \left(\frac{1}{\sigma_N^2 |x|^{2(N+\theta-1)}} + \frac{2\theta}{\sigma_N |x|^{N+2\theta}} \right) \\ &\quad + 2(N-1) \left(\frac{1}{\sigma_N |x|^{N+2\theta}} + \frac{\theta}{|x|^{2\theta+2}} \right) P_B(x). \end{aligned} \tag{4.99}$$

Set

$$\tilde{r} = \min \left\{ 2, \left(\frac{1}{2\sigma_N |\theta|} \right)^{\frac{1}{N-2}} \right\}. \tag{4.100}$$

If $x \in B \cap B_{\tilde{r}}(0)$, the two last terms in (4.99) are nonnegative, hence

$$|\nabla w|^2 \geq (\theta^2 + (N-1)^2) |x|^{-2(\theta+1)} P_B^2(x) \quad \text{for all } x \in B \cap B_{\tilde{r}}(0). \tag{4.101}$$

Note that $B \cap B_{\tilde{r}}(0) = B$ if $\tilde{r} = 2$. Choosing $m > 0$ small enough we infer that w_m is a subsolution of (1.1) in $B \cap B_{\tilde{r}}(0)$. Denoting by \hat{m} the maximum of w_m on $\partial(B \cap B_{\tilde{r}}(0)) \setminus \{0\}$, then $(w_m - \hat{m})_+$ is a subsolution in Ω . Since the restriction to Ω of the solution constructed in Theorem 1.11 dominates $(w_m - \hat{m})_+$, the proof follows as in the first case. \square

4.3.3 Open problems

Problem 1. Under what conditions are the positive solutions of problem (1.28) unique? If instead of separable solutions in \mathbb{R}_+^N vanishing on $\partial\mathbb{R}_+^N \setminus \{0\}$ one looks

for separable radial solutions of (1.1) in $\mathbb{R}^N \setminus \{0\}$ (with $q = \frac{2p}{p+1}$), then they are under the form

$$U(x) = A|x|^{-\alpha} \quad (4.102)$$

and A is a positive root of the polynomial

$$P(X) = X^{p-1} - M\alpha^{\frac{2p}{p+1}}X^{\frac{p-1}{p+1}} + \alpha(N-2-\alpha). \quad (4.103)$$

A complete study of the radial solutions of (1.1) is provided in [7], however it is straightforward to check that if $1 < p < \frac{N}{N-2}$, there exists a unique positive root, hence a unique positive separable solution, while if $p > \frac{N}{N-2}$, there exists a unique positive root (resp. two positive roots) if

$$M = (p+1) \left(\frac{p(N-2) - N}{2p} \right)^{\frac{p}{p+1}} := m^*, \quad (4.104)$$

(resp. $M > m^*$). Uniqueness of solution plays a fundamental role in the description and classification of all the positive solutions with an isolated singularity at 0.

Problem 2. It is proved in [7] that if $\max\{\frac{N}{N-1}, \frac{2p}{p+1}\} < q < \min\{2, p\}$ and $M > 0$, there exist infinitely many local radial solutions of (1.1) in $\mathbb{R}^N \setminus \{0\}$ which satisfies

$$u(r) = \xi_M r^{-\beta}(1 + o(1)) \quad \text{as } r \rightarrow 0 \quad (4.105)$$

where

$$\beta = \frac{2-q}{q-1} \quad \text{and} \quad \xi_M = \frac{1}{\beta} \left(\frac{(N-1)q - N}{M(p-1)} \right)^{\frac{1}{p-1}}. \quad (4.106)$$

These solutions present the property that their blow-up is smaller than the one of the explicit radial separable solution. It would be interesting to construct such solutions of (1.1) in \mathbb{R}_+^N (or more likely B_R^+), vanishing on $\partial\mathbb{R}^N \setminus \{0\}$.

Problem 3. Is it possible to define a boundary trace for any positive solution of (1.1) in \mathbb{R}_+^N , noting the fact such a result holds separately for positive solutions of (1.2) and (1.4)? A related problem would be to define an initial trace for any positive solution of the parabolic equation

$$\partial_t u - \Delta u + u^p - M|\nabla u|^q = 0, \quad (4.107)$$

in $(0, T) \times \mathbb{R}^N$. Initial trace of semilinear parabolic equations ($M = 0$ in (4.107)) are studied in [17], [14].

Problem 4. Are the positive solutions of 1.8 satisfying (1.24) or (4.75) unique without the flatness and the starshapedness assumptions of Theorem 4.2. More generally, are the weak solutions of the Dirichlet problem with measure boundary data (1.14) unique?

References

- [1] D. Adams, L. Hedberg. *Function Spaces and Potential Theory*, Theory, Springer-Verlag, London-Berlin-Heidelberg-New York (1996).
- [2] D. R. Adams, M. Pierre. Capacitary strong type estimates in semilinear problems, *Ann. Inst. Fourier (Grenoble)* **41**, 117-135 (1991).
- [3] P. Baras, M. Pierre. Singularités éliminables pour des équations semi-linéaires, *Ann. Inst. Fourier (Grenoble)* **34**, 185-206 (1986).
- [4] M.F. Bidaut-Véron, M. Garcia-Huidobro, L. Véron. Boundary singularities of positive solutions of quasilinear Hamilton-Jacobi equations, *Cal. Var. Part. Diff. Equ.* **54**, 3471-3515 (2015).
- [5] M.F. Bidaut-Véron, M. Garcia-Huidobro, L. Véron. A priori estimates for elliptic equations with reaction terms involving the function and its gradient, *Math. Annalen* **378**, 13-56 (2020).
- [6] M.F. Bidaut-Véron, M. Garcia-Huidobro, L. Véron. Radial solutions of scaling invariant nonlinear elliptic equations with mixed reaction terms. *Discrete Cont. Dyn. Systems* **40**, 933-982 (2020).
- [7] M.F. Bidaut-Véron, M. Garcia-Huidobro, L. Véron. Measure Data Problems for a Class of Elliptic Equations with Mixed Absorption-Reaction, *Adv Nonlinear Studies* **21**, 261-280 (2020).
- [8] M.F. Bidaut-Véron, M. Garcia-Huidobro, L. Véron. Singular solutions of some elliptic equations involving mixed absorption-reaction, *submitted*, arXiv:2107.13399v2.
- [9] M.F. Bidaut-Véron, G. Hoang, Q. H. Nguyen, L. Véron. An elliptic semilinear equation with source term and boundary measure data: the supercritical case, *J. Funct. Anal.* **269**, 1995-2017 (2015).
- [10] M.F. Bidaut-Véron, L. Vivier. An elliptic semilinear equation with source term involving boundary measures: the subcritical case, *Rev. Mat. Iberoamericana* **16**, 477-513 (2000).
- [11] L. Boccardo, F. Murat, J. P. Puel. Résultats d'existence pour certains problèmes elliptiques quasilineaires, *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze* **Ser. 4, T. 11**, 213-235 (1984).
- [12] J. Doob. *Classical Potential Theory and Its Probabilistic Counterpart*, Springer-Verlag, London-Berlin-Heidelberg-New York (1984).
- [13] D. Gilbarg, N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*, reprint of the 1998 edition, Classics in Math. Springer-Verlag, New York (2001).
- [14] K. Gkikas, L. Véron. Complete Classification of the Positive Solutions of Heat Equation with Super Critical Absorption, *Advanced Nonlinear Studies* **14**, 47-113 (2014).

- [15] A. Gmira, L. Véron. Boundary singularities of solutions of nonlinear elliptic equations, *Duke Mathematical J.* **64**, 271-324 (1991).
- [16] J. B. Keller. On the solutions of $\Delta u = f(u)$, *Comm. Pure Appl. Math.* **10**, 503-510 (1957).
- [17] M. Marcus, L. Véron. Initial trace of positive solutions of some nonlinear parabolic equations, *Comm. part. Diff. Equ.* **24**, 1445-1499 (1999).
- [18] M. Marcus, L. Véron. Removable singularities and boundary traces, *J. Math. Pures Appl.* **80**, 879-900 (2001).
- [19] M. Marcus, L. Véron. Maximal solutions of semilinear elliptic equations with locally integrable forcing term, *Israel J. Math.* **152**, 333-348 (2006)
- [20] M. Marcus, L. Véron. *Nonlinear Elliptic Equations involving Measures*, De Gruyter Series in Nonlinear Analysis and Applications. **21** xiii+pp. 1-262 (2013).
- [21] V. H. Nguyen. Sharp weighted Sobolev and Gagliardo-Nirenberg inequalities on half space via mass transport and consequences, *Proc. London Math. Soc.* **111**, 127-148 (2015).
- [22] P. T. Nguyen, L. Véron. Boundary singularities of solutions to elliptic viscous Hamilton-Jacobi equations, *J. Funct. Anal.* **263**, 1487-1538 (2012).
- [23] R. Osserman. On the inequality $\Delta u \geq f(u)$, *Pacific J. Math.* **7**, 1641-1647 (1957).
- [24] P. Pucci, J. Serrin. *The maximum principle*, Progress in Nonlinear Differential Equations and Their Applications, Birkhäuser Verlag, Basel (2007).
- [25] J. Serrin. Isolated singularities of solutions of quasilinear equations, *Acta Mathematica* **34**, 247-302 (1964).
- [26] J. Serrin, H. Zou. Existence and nonexistence results for ground states of quasilinear elliptic equations., *Arch. Rat. Mech. Anal.* **121**, 101-130 (1992).
- [27] E. M. Stein *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, New Jersey (1970).
- [28] I. E. Verbitsky, R. L. Wheeden. Weighted norm inequalities for integral operators, *Trans. Amer. Math. Soc.* **350**, 3371-3391 (1998).
- [29] L. Véron. Singular solutions of some nonlinear elliptic equations, *Ann. Fac. Sci. Toulouse* **6**, 1-31 (1984).
- [30] L. Véron. Singular solutions of some nonlinear elliptic equations, *Nonlinear Anal. Theory, Methods Appl.* **5**, 225-242 (1981).
- [31] L. Véron. *Elliptic Equations Involving Measures*, Handbook of Differential Equations, Vol. 1, Chapter 8, pp 593-712. Stationary Partial Differential Equations (2004).
- [32] L. Véron. *Local and Global Aspects of Quasilinear Degenerate Elliptic Equations*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ (2017).