



HAL
open science

Production of perdeuterated fucose from glyco-engineered bacteria

Lukas Gajdos, V Trevor Trevor Forsyth, Matthew P. Blakeley, Michael Haertlein, Anne Imberty, Eric Samain, Juliette M Devos

► **To cite this version:**

Lukas Gajdos, V Trevor Trevor Forsyth, Matthew P. Blakeley, Michael Haertlein, Anne Imberty, et al.. Production of perdeuterated fucose from glyco-engineered bacteria. *Glycobiology*, 2021, 31, pp.151-158. 10.1093/glycob/cwaa059 . hal-02911649

HAL Id: hal-02911649

<https://hal.science/hal-02911649v1>

Submitted on 4 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Production of perdeuterated fucose from glyco-engineered bacteria

Lukas Gajdos^{a,b,c}, V. Trevor Forsyth^{a,b,d}, Matthew P. Blakeley^e, Michael Haertlein^{a,b}, Anne Imberty^{c*}, Eric Samain^{c*} & Juliette M. Devos^{a,b*}

^a Life Sciences Group, Institut Laue-Langevin, 71 Avenue des Martyrs, 38000 Grenoble, France

^b Partnership for Structural Biology (PSB), 71 Avenue des Martyrs, 38000 Grenoble, France

^c Univ. Grenoble Alpes, CNRS, CERMAV, 38000 Grenoble, France

^d Faculty of Natural Sciences, Keele University, ST5 5BG Staffordshire, UK

^e Large Scale Structures Group, Institut Laue-Langevin, 71 Avenue des Martyrs, 38000 Grenoble, France.

* Authors for correspondence: Anne Imberty (anne.imberty@cermav.cnrs.fr), Eric Samain (eric.samain@cermav.cnrs.fr), Juliette Devos (devosj@ill.fr).

Keywords: deuteration / engineering / *Escherichia coli* / fucose / neutron scattering

Supplementary data: Figure S1, Figure S2

Abstract

L-Fucose and L-fucose-containing polysaccharides, glycoproteins or glycolipids play an important role in a variety of biological processes. L-Fucose-containing glycoconjugates have been implicated in many diseases including cancer and rheumatoid arthritis. Interest in fucose and its derivatives is growing in cancer research, glyco-immunology, and the study of host-pathogen interactions. L-Fucose can be extracted from bacterial and algal polysaccharides, or produced (bio)synthetically. While deuterated glucose and galactose are available, and are of high interest for metabolic studies and biophysical studies, deuterated fucose is not easily available.

Here, we describe the production of perdeuterated L-fucose, using glyco-engineered *Escherichia coli* in a bioreactor with the use of a deuterium oxide-based growth medium and a deuterated carbon source. The final yield was 0.2 g L⁻¹ of deuterated sugar, which was fully characterized by mass spectrometry and nuclear magnetic resonance spectroscopy. We anticipate that the perdeuterated fucose produced in this way will have numerous applications in structural biology where techniques such as NMR, solution neutron scattering and neutron crystallography are widely used. In the case of neutron macromolecular crystallography, the availability of perdeuterated fucose can be exploited in identifying the details of its interaction with protein receptors and notably the hydrogen bonding network around the carbohydrate binding site.

Introduction

The deuteration of biomolecules by stable isotope labelling is of high interest for nuclear magnetic resonance (NMR) spectroscopy and neutron scattering studies of the structure and dynamics of biological macromolecules and is now widely used (Haertlein et al. 2016, Blakeley and Podjarny 2018). Tailor-designed deuteration can be used very effectively in studies of multi-component systems by small-angle neutron solution scattering (Laux et al. 2008, Cuypers et al. 2013b, Dunne et al. 2017, Josts et al. 2018, Maric et al. 2019) and neutron reflection (Grage et al. 2011, Hellstrand et al. 2013, Waldie et al. 2018). In Laue neutron crystallography, the use of perdeuterated protein imparts major benefits in terms of data quality and interpretation (Haupt et al. 2014, Dajnowicz et al. 2017, Yee et al. 2019, Kwon et al. 2020) and the same is true in the case of monochromatic neutron crystallography (Cuypers et al. 2013a, Cuypers et al. 2016). Deuteration of small biomolecules has wide application in analytical methods, with growing development for the study of metabolism (Shimba et al. 1990) and in living cells imaging techniques such as Raman microscopy (Wei et al. 2013). MeV ion beam analysis techniques such as ^3He Nuclear Reaction Analysis (NRA), where high sensitivity to deuterium was utilized to monitor the uptake of deuterated sugars in bacteria (Lowery et al. 2015). More recently, it has been demonstrated that deuteration of key hydrogen atoms at positions where drugs are enzymatically metabolised for elimination has direct effects on pharmacological properties of these substances by improving their stability, and therefore their half-life, and by decreasing their toxicities (Schmidt 2017). The basis of the increased stability of these “heavy hydrogen drugs” is related to the stronger deuterium-carbon bond making the drug more resistant to enzymatic modification and elimination; deuteration of known drugs, referred to as the “deuterium switch”, resulted in a boost of patenting of labelled

drugs, with several of them now in clinical trials (Timmins 2017). The first approved deuterated drug for therapeutic use was deutetrabenazine, and it has been used for the treatment of chorea in Huntington's disease (Dean and Sung 2018).

Proteins and lipids are generally obtained in their perdeuterated form through the use of recombinant bacteria or yeasts (Meilleur et al. 2009, Haertlein et al. 2016, Moulin et al. 2018). The availability of perdeuterated, and well characterized glycans is more limited. For selective deuteration, chemical synthesis of perdeuterated monosaccharides has previously been described. Through the use of appropriate protecting groups, direct H–D exchange reactions in the presence of an activated carbon-supported platinum group catalyst were used to produce a variety of deuterated monosaccharides (Koch and Stuart 1978, Sawama et al. 2012). Using biotechnology approaches, it has been demonstrated more than 50 years ago that certain algae can be adapted to grow in 99.6% D_2O , resulting in the large scale production of perdeuterated glucose and mannose (Crespi et al. 1959, Chorney et al. 1960). It is only recently that cyanidin 3-*O*-glucoside was produced with a perdeuterated glucose moiety using recombinant *Escherichia coli* (Gupta et al. 2018). Polysaccharides have been produced in bacteria, in the form of perdeuterated cellulose (O'Neill et al. 2015) and perdeuterated heparin (Cress et al. 2019).

The current work focuses on L-fucose (6-deoxy-L-galactose), which is a non-classical monosaccharide with an L-configuration correlated to unusual ring conformation, and a methyl group at C6. L-fucose (Fuc) is present in all living kingdoms, from giant viruses (De Castro et al. 2013) to bacterial polysaccharides and plant cell walls. In mammals, it is a common component of glycolipids and glycoproteins (Vanhooren and Vandamme 1999, Schneider et al. 2017). Terminal fucosylation of oligosaccharides create bioactive epitopes, such as the histo-blood group oligosaccharides (ABO and Lewis epitopes) (Painter et al. 1965). Human

milk oligosaccharides are also rich in terminal fucose, which serve as receptor analogues for pathogens, and modulator of immune responses (Bode 2015). Fucose is therefore of interest as a biomarker for some cancers as well as a target for pathogen receptors (Heggelund et al. 2017).

The production of perdeuterated fucose, L-fucose-d₁₂ (Fuc-d₁₂) is therefore of high interest. The chemical approach has been described but is complex and time-consuming (Koch and Stuart 1978, Sawama et al. 2012). An alternative approach is based on the production of “recombinant” oligosaccharides in microorganisms, by engineering of their metabolic pathways. This area of research was initiated at the end of the last century with the production of chito-oligosaccharides in *Escherichia coli* (Samain et al. 1997). During the last decades, a large number of oligosaccharides have been produced using this approach, including human milk oligosaccharides, that are of high commercial interest for improving infant formula milk (Priem et al. 2002). For example, α -2'-fucosyllactose, a major trisaccharide of human milk, has been obtained by introducing the FUT2 gene from *Helicobacter pylori* into *E. coli* (Albermann et al. 2001, Drouillard et al. 2006, Yu et al. 2018). This trisaccharide can serve as a substrate for a fucosidase. With appropriate engineering of microorganisms, fucose can be recombinantly produced, as demonstrated in *Saccharomyces cerevisiae* (Liu et al. 2018) and then in *E. coli* (Liu et al. 2019).

In this work, the production of perdeuterated Fuc-d₁₂ (per-C-deuterated Fuc-d₈ when dissolved in H₂O) is described. Based on the previous construct for production of α -2'-fucosyllactose (Drouillard et al. 2006), the metabolic pathways of the previously described *E. coli* strain were further engineered with the addition of the appropriate hydrolase and removal of sugar transporter. After adaptation of the strains for cultivation in an appropriate deuterated medium, a large amount of perdeuterated fucose was produced and fully characterized.

Results

The fucose-producing strain

A schematic diagram of the production of L-fucose using the engineered FUC5 strain is shown in Figure 1. FUC5 is a derivative of the *E. coli* K12 strain DH1, with genetic modifications for overexpression of genes involved in GDP-fucose production, introduction of 2'fucosyltransferase gene from *Helicobacter pylori* and fucosidase gene from *Bifidobacterium bifidum* strain JCM1254, and deletion of genes involved in the metabolism of lactose. In addition, several genes have been knocked out to optimize the production of fucose: the *fucI* gene (coding for fucose isomerase) to prevent isomerisation to fuculose and the *fucP* gene (fucose permease) to prevent internalization of extracellular fucose. A detailed list of the plasmids, genes and *E. coli* strains described in this work can be found in Table I.




Figure 1. Schematic representation of the metabolic network for fucose production using the engineered *E. coli* FUC5 strain. Inserted genes are shown in italics and knockout genes are crossed out.

Table I. Genes, plasmids and *E. coli* strains used in this study

Description	Reference or source	
Genes		
<i>futC</i>	α -1,2-fucosyltransferase from <i>Helicobacter pylori</i> strain 26695	GenBank KY499613
<i>afcA</i>	α -1,2-fucosidase from <i>Bifidobacterium bifidum</i> strain JCM1254	GenBank AY303700
<i>gmd</i> , <i>wcaG</i> , <i>manC</i> , <i>manB</i>	<i>E. coli</i> genes coding GDP-Man dehydratase, fucose synthase, GDP-Man pyrophosphorylase and phosphomannomutase respectively	GenBank U38473
Plasmids		
pEXT20-futC	pEXT20 carrying <i>futC</i>	(Drouillard et al. 2006)
pBP-futC	Derived from pEX20-futC	This study
pBBRGAB	pBBR1MCS-3 derivative carrying <i>gmd</i> , <i>wcaG</i> , <i>manC</i> and <i>manB</i>	(Dumon et al. 2006)
pSU-fucase	pSU2718 derivative carrying active α -1,2-fucosidase domain of <i>afcA</i> from <i>Bifidobacterium bifidum</i>	This study
Strains		
DC	DH1 <i>lacZ lacA</i>	(Dumon et al. 2006)
FUC	DC <i>fucI fucP</i>	This study
FUC5	FUC (pBP-futC, pBBRGAB, pSU-fucase)	This study

Adaptation to D₂O and batch fermentation

In order to compare fucose production in different conditions, the FUC5 strain was grown in both unlabelled and deuterated minimal medium supplemented with hydrogenated glycerol and deuterated glycerol-d₈, respectively. Its growth was stepwise adapted to deuterated media as previously described (Haertlein et al. 2016). Both hydrogenated and deuterated fucose

were produced in simple batch-fermentations. The hydrogenated culture showed a higher growth rate in the exponential phase, reaching a maximum OD₆₀₀ of 26, 73 hours after inoculation. The deuterated culture showed a similar lag-phase profile but slower growth in the exponential phase with the highest OD₆₀₀ being 17, 112 hours after the start of the fermentation (Figure 2A).




Figure 2: (A) Growth curves of hydrogenated and deuterated *E. coli* FUC5 in high cell-density cultures. The optical density was measured at 600 nm using a spectrophotometer and was plotted against time (hours). (B, C) Production of hydrogenated (B) and deuterated (C) fucose using the genetically modified *E. coli* FUC5. The concentrations of extracellular and intracellular fucose were measured spectrophotometrically using the K-Fucose kit (Megazyme). Symbols represent a mean value of two independent measurements. Error bars indicate the standard deviation and some are smaller than the symbol size used in the panels.

The bacterial growth and the deuterated fucose concentration showed a cell-density-associated production pattern (Figure 2). A maximum Fuc-d₁₂ concentration of 0.20 g L⁻¹ was obtained by the end of the fermentation process. The yield of fucose produced per gram of carbon source in the growth medium was 10 fold lower for the deuterated culture (0.2 vs. 2.1 g L⁻¹, see Figure 2B and Figure 2C) when supplied with the same quantity of carbon source, *ie* 30 g of glycerol used per liter of growth medium as indicated in the experimental section. Analysis by thin layer chromatography (TLC) showed gradually increasing concentrations of intracellular lactose transported from the medium to the cells by lactose permease. The TLC also revealed the increase in concentration of

extracellular fucose over time (Figures S1 and S2).

Optimisation of purification method

Deuterated fucose was purified directly from the crude extracellular fraction. After the first ion exchange chromatography purification step, TLC analysis showed the presence of lactose and isopropyl- β -D-thiogalactopyranoside (IPTG) that were added to the minimal medium during high cell-density culture. Further fractionation of these compounds was achieved by size-exclusion chromatography (SEC) using a TOYOPEARL HW-40 column made of hydroxylated methacrylic polymer (Figure 3 bottom). This technique was used as a

polishing step to efficiently separate lactose, fucose and IPTG, three molecules of similar sizes that could be eluted as separate single peaks. After purification, 220 mg of

lyophilized Fuc-d₁₂ were obtained from a culture using 1.5 L of D₂O and 45 g of deuterated glycerol.




Figure 3: (Top) Mass spectrometry and (Bottom) chromatography analyses of per-C-deuterated fucose produced. Analyses were carried out in H₂O and thus correspond to Fuc-d₈. Bottom: Size-exclusion chromatography (SEC) profiles of the saccharides present in the extracellular fraction of the fucose-producing strain after batch-fermentation in deuterated media: (a) lactose, (b) deuterated fucose, and (c) isopropyl-β-D-thiogalactopyranoside (IPTG) in the black line, superimposed with a SEC profile of standard solutions in the grey line. Top: Only the positive electrospray ionization (ESI) mass spectrum of the deuterated fucose (Fuc-d₈) is displayed with sodium and potassium adduct ions at m/z 196 and 211 respectively.

Deuterated fucose characterization

The metabolic incorporation of deuterium in the fucose molecule was examined and demonstrated by electrospray ionization (ESI) mass spectrometry, and quantified by ¹H NMR. The results (Figure 3 top) showed

that the purified product dissolved in a water/methanol solution generated two major peaks corresponding to the exact masses of sodium adduct ions [M+Na]⁺ and potassium adduct ions [M+K]⁺ at m/z values of 195 and 211, respectively, confirming the presence of per-C-deuterated L-fucose (Fuc-d₈).




Figure 4: ^1H (Top and Middle) and ^2H (Bottom) NMR spectra comparing the signal intensity for hydrogenated L-fucose and deuterated L-fucose. Peaks in the dashed boxes correspond to the hydrogen (^1H spectra) and deuterium (^2H spectrum) atoms of the fucose molecule. A signal of sodium succinate is observed at 2.4 ppm for the ^1H spectra.

The extent of deuteration of the fucose ring and the methyl carbon was also verified by ^1H and ^2H NMR. Proton NMR spectra were measured on samples of commercially available hydrogenated L-fucose and on the deuterated L-fucose produced in this work, both solubilized in D_2O . The deuterium NMR spectrum was measured on the deuterated fucose solubilized in water. The ^1H spectrum of the purified compound exhibited the loss of intensity of all proton signals compared to the non-labelled standard (Figure 4). Moreover, the deuterium signals observed in the ^2H spectrum of the perdeuterated product demonstrated that deuterium replacement has taken place successfully on all of the carbon atoms of the fucose molecule.

Discussion

In this study, glyco-engineered *E. coli* has been used to produce deuterated fucose from

deuterated glycerol and D_2O via biosynthetic pathways. For the production of deuterated fucose- d_{12} , deuterated glycerol- d_8 was used as the sole carbon source. Even though glucose is a preferred carbon source for *E. coli*, several studies showed that engineered *E. coli* strains producing fucosyllactose and fucose give higher titers of fucose when grown on glycerol (Jung et al. 2019, Liu et al. 2019). Compared to the hydrogenated culture, the perdeuterated high cell-density culture grew slower in the exponential phase and gave lower titers of fucose. Similar behaviour has been observed in deuteration of organic molecules using engineered organisms (Meilleur et al. 2009, Moulin et al. 2018).

The engineered FUC5 strain produced 0.2 g L^{-1} of deuterated fucose from a limited amount of glycerol in a simple batch-fermentation. The fully deuterated fucose was characterized by mass spectrometry,

proton and deuterium NMR, confirming full deuteration of the final product. It should be noted that the final yield of both the hydrogenated and the perdeuterated monosaccharides in this study was strongly dependent on the amount of glycerol used during the fermentation process. Due to the cost of the deuterated carbon source, this amount was limited to 45 g in both the H- and the D-study. Under optimal hydrogenated conditions where the quantity of carbon source is not limited, the highest titer of fucose obtained was 20 g L⁻¹ (data not shown).

Perdeuteration using engineered *E. coli* is therefore a convenient and effective method for a high-quantity production of fully deuterated simple and more complex sugars for use in neutron scattering experiments as well as in pharmacokinetic studies. In a first application, Fuc-d₁₂ will be used as a ligand for fucose-specific lectins in single crystal neutron diffraction experiments.

Due to the larger and positive coherent scattering of deuterium compared to hydrogen in neutron scattering experiments, there is a better visibility of deuterium atoms versus hydrogen. In this study, the ¹H atoms changed for deuterium (²H) atoms in the fucose molecule will allow clearer visualisation of crucial interactions between the sugar and bacterial lectins, as well as an optimisation of the overall quality of the neutron data through the reduction of hydrogen incoherent scattering.

Materials and methods

Strains and plasmid construction

The fucose-producing strain FUC5 was obtained by transforming the host strain FUC with the three following plasmids: pBP-futC which contains the α -1,2-fucosyltransferase gene *futC* from *H. pylori* strain 26695; pBBRGAB which contains the four *E. coli* *gmd*, *wcaG*, *manC*, and *manB* genes coding GDP-Man dehydratase, fucose synthase, GDP-Man pyrophosphorylase and phosphomannomutase respectively; pSU-fucase which contains the sequence of the

active α -1,2-fucosidase domain of the *afcA* gene from *Bifidobacterium bifidum* strain JCM1254 (Katayama et al. 2004).

The host strain FUC was designed from strain DC (Dumon et al. 2006) by knocking out the genes *fucI* and *fucP* encoding fucose isomerase and fucose permease respectively. Strain DC was a *lacA lacZ* null mutant derived from the *Escherichia coli* K12 strain DH1 (DSM 4235). In order to knockout the *fucPI* genes, a 1.435 kb segment located between nucleotides 913 of *fucP* and 998 of *fucI* was deleted and replaced by the 5'AAGCTT sequence as follows: two DNA segments flanking the deleted sequence were amplified by PCR. The upstream 0.873 kb segment was amplified with primers 5'GGATCCGTAGATAAAGATGCAGG GCAAAGCAGAAG and 5'AAGCTTGGTTCCGGTTAAATAGT TAGCGGCAAAG and the downstream 0.806 kb segment was amplified with primers 5'AAGCTTTCGTGGCGACCGAAAACG ACAG and 5'CTCGAGACCGGGCATCACATCAG GGAG. The two amplified fragments were ligated at their terminal *HindIII* restriction site and cloned together into the *BamHI Sall* sites of the suicide vector pKO3. The deletion was then carried out according to the pKO3 gene replacement protocol (Link et al. 1997).

The plasmid pBP-futC was designed from plasmid pEXT20-futC (Drouillard et al. 2006) by removing the ampicillin gene using *DraI* digestion and by inserting via blunt-end ligation a kanamycin cassette, obtained by *PstI* digestion of the pUC4K vector.

The design of pSU-fucase plasmid was carried out as follows: a 2.728 kb DNA fragment containing the sequence of the active α -1,2-fucosidase domain of the *afcA* gene was amplified by PCR using genomic DNA of *Bifidobacterium bifidum* strain JCM1254 as a template and the following primers: 5'CTCGAGTAAGGAGGTAATAT AATGGTCATCGCCAGTGTCTGAGGACG

and 5' AAGCTTAGGCGCTCGCCTTCTTC GTGATCGTGAC. The amplified fragment was first cloned into pCR4Blunt-TOPO vector (Invitrogen) and then sub-cloned into the *Xho*I and *Hind*III sites of the pSU2718 expression vector to form pSU-fucase (Martinez et al. 1988).

Construction of pBBRGAB plasmid was previously described (Dumon et al. 2006).

Production of deuterated fucose in batch-fermentation

Precultures of the fucose-producing strain were first grown in LB medium. All culture media were supplemented with 15 $\mu\text{g mL}^{-1}$ tetracycline, 20 $\mu\text{g mL}^{-1}$ chloramphenicol, 50 $\mu\text{g mL}^{-1}$ kanamycin and 4 $\mu\text{g mL}^{-1}$ thiamine. All cultures were grown at 28 °C with shaking at 160 rpm. The strain was adapted to deuterated minimal medium with the following composition: 5 g L^{-1} $\text{NH}_4\text{H}_2\text{PO}_4$, 5 g L^{-1} KH_2PO_4 , 0.5 g L^{-1} $\text{C}_6\text{H}_8\text{O}_7$ (citric acid), 1.65 g L^{-1} KOH, 0.65 g L^{-1} NaOH, 7.5 mL trace mineral solution [13 g L^{-1} $\text{N}(\text{CH}_2\text{CO}_2\text{H})_3$ (nitrilotriacetic acid), 7 g L^{-1} KOH, 7.5 g L^{-1} $\text{C}_6\text{H}_5\text{FeO}_7$ (ferric citrate), 1.3 g L^{-1} $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 1.2 g L^{-1} $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.25 g L^{-1} H_3BO_3 , 0.15 g L^{-1} $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 0.21 g L^{-1} $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.13 g L^{-1} $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$], 45 g glycerol- d_8 (Eurisotop). A single colony of *E. coli* FUC5 cells containing pBP-futC, pBBRGAB and pSU-fucase plasmids grown overnight on LB agar plates supplemented with the 3 antibiotics was used to inoculate 15 mL of minimal medium and was grown overnight at 28 °C. The culture was then used to inoculate 15 mL of 100% D_2O minimal medium (with deuterated glycerol- d_8) at OD_{600} of 0.1 and was grown overnight. This step was repeated five times until the doubling time for *E. coli* reached values similar to those for hydrogenated cultures. The last deuterated preculture of 200 ml was used to inoculate the 1.5 L of deuterated minimal medium in a 3 L bioreactor used for batch-fermentation. The pD of the culture medium was regulated at 7.2 by addition of 4% NaOD. The temperature was maintained at 28 °C. IPTG inducer and lactose acceptor were added to a

final concentration of 0.2 mM and 0.5 g L^{-1} respectively at the inoculation time. The fermentation was stopped after consumption of the deuterated glycerol- d_8 from the culture medium.

Analytical methods

Cell growth was monitored by measuring turbidity at 600 nm (OD_{600}) using a spectrophotometer. Culture aliquots (1 mL) were taken at different time points throughout the high cell-density culture. Samples were centrifuged and the supernatants were treated as the extracellular fraction. The pellets were resuspended in 1 mL of Milli-Q water and boiled for 20 min at 100 °C. After centrifugation, the supernatants were treated as the intracellular fraction. Samples were analysed using thin layer chromatography on silica gels with an eluent composed of 1-propanol: acetic acid: water in 2:1:1 ratio. After dipping the plate in orcinol-sulphuric acid reagent, the spots were visualized by heating at 100 °C.

Purification of deuterated fucose

At the end of the fermentation, the bacterial cells were recovered by centrifugation (8000 g for 1 h at 8 °C). The supernatant containing the deuterated fucose is referred to as *extracellular fraction*. The purification was performed in two steps. The first step was carried out using ion-exchange chromatography. The crude extracellular fraction was treated with a strongly acidic cation-exchanger resin (Amberlite IR120 hydrogen form, Sigma-Aldrich). The resin was added progressively until the pH dropped below 3. After decantation, precipitated proteins were eliminated by centrifugation (16 000 g for 1 h at 8 °C). The supernatant was filtered and loaded onto a column with a strongly acidic cation-exchanger resin (DOWEX 50WX4 hydrogen form, Sigma-Aldrich) and immediately neutralized by passing through a weakly basic anion-exchanger resin (DOWEX 66 free base, Sigma-Aldrich). The final eluate was colourless with neutral pH. The purity

and amount of fucose were analyzed by thin layer chromatography. Fractions containing fucose were pooled together, sterile filtered and lyophilized. The resulting mix was further fractionated by size-exclusion chromatography using a semi-preparative TOYOPEARL[®] HW-40 column (Tosoh Bioscience GmbH) equipped with a refractive index detector (Knauer). The chromatography was performed at ambient temperature with water as a mobile phase and a flow rate of 120 mL h⁻¹. Fractions containing pure product were pooled together and submitted to three cycles of resuspension in D₂O and freeze-drying.

Quantification of hydrogenated and deuterated L-fucose

Quantification of the intracellular and extracellular fucose was carried out on the cell paste and crude supernatant samples, respectively, using an enzymatic L-fucose assay kit (K-FUCOSE, Megazyme). The assay is based on the oxidation of L-fucose to L-fucono-1,5-lactone by L-fucose dehydrogenase in the presence of nicotinamide adenine dinucleotide phosphate (NADP⁺). The amount of L-fucose is stoichiometric to the production of reduced nicotinamide adenine dinucleotide phosphate (NADPH), measured by an increase of absorbance at 340 nm. The final amount of the deuterated fucose obtained was quantified by weighing the lyophilized powder after purification.

Structural analysis of deuterated fucose

Positive and negative-ion ESI mass spectrometry data were recorded on the Bruker amaZon speed ion-trap mass spectrometer. A sample of deuterated fucose was solubilized in D₂O to a final concentration of 50 µM.

The proton and deuterium NMR spectra were recorded with a Bruker Avance[™] III NMR spectrometer operating at a frequency of 400.13 MHz for ¹H and 61.42 MHz for ²H. For the ¹H NMR experiments, samples of hydrogenated and deuterated fucose were solubilized in D₂O with 1.82 g L⁻¹ sodium

succinate as a standard. Residual signal of the solvent was used as the internal standard: HOD at 4.8 ppm at 298 K. The ¹H spectra were recorded at 298 K, with a 4006 Hz spectral width, 32 768 data points, 4.096 s acquisition times, 10 s relaxation delays and 16 scans.

The ²H NMR spectrum was measured on a sample of 7 mg of deuterated fucose dissolved in H₂O at pH 6.7. The ²H spectrum was acquired at 298 K, without broadband ²H-decoupling, using a 90° pulse, 10 015 Hz sweep width, 3030 number of data points, 1 s acquisition time and 1024 number of scans.

Acknowledgements

We thank the ILL for the provision of studentship funding to LG, and for the award of neutron beamtime. We also thank the ILL for access to the Deuteration Laboratory platform in ILL's Life Sciences Group (<https://www.ill.eu/lsg>). We would like to thank the CERMAV Common Chromatography Service (SC3) platform, especially Laurine Buon for help with the fucose purification, and the Chimie Nanobio Joint Technology Platform of the Institute of Molecular Chemistry of Grenoble (PCN-ICMG) for mass spectrometry and NMR measurements. VTF wishes to acknowledge the UK Engineering and Physical Sciences Research Council (EPSRC) for grants GR/R99393/01 and EP/C015452/1 that funded the creation of the Deuteration Laboratory within ILL's Life Sciences Group. AI wishes to acknowledge the support of French ANR through Glyco@Alps (ANR-15-IDEX-02) and Labex ARCANÉ / CBH-EUR-GS (ANR-17-EURE-0003).

Abbreviations ESI, electrospray ionization; Fuc, fucose; FUT2, fucosyltransferase 2; IPTG, isopropyl-β-D-thiogalactopyranoside; LB, Luria Bertani medium; NMR, nuclear magnetic resonance; PCR, polymerase chain reaction; SEC, size-exclusion chromatography; TLC, thin layer chromatography.

References

- Albermann C, Piepersberg W, Wehmeier UF. 2001. Synthesis of the milk oligosaccharide 2'-fucosyllactose using recombinant bacterial enzymes. *Carbohydr Res.* 334:97-103.
- Blakeley MP, Podjarny AD. 2018. Neutron macromolecular crystallography. *Emerging Topics in Life Sciences.* 2:39-55.
- Bode L. 2015. The functional biology of human milk oligosaccharides. *Early Hum Dev.* 91:619-622.
- Chorney W, Scully NJ, Crespi HL, Katz JJ. 1960. The growth of algae in deuterium oxide. *Biochim Biophys Acta.* 37:280-287.
- Crespi HL, Archer SM, Katz JJ. 1959. Culture of algae and other micro-organisms in deuterium oxide. *Nature.* 184(Suppl 10):729-730.
- Cress BF, Bhaskar U, Vaidyanathan D, Williams A, Cai C, Liu X, Fu L, V MC, Zhang F, Mousa SA, *et al.* 2019. Heavy Heparin: A Stable Isotope-Enriched, Chemoenzymatically-Synthesized, Poly-Component Drug. *Angew Chem Int Ed Engl.* 58:5962-5966.
- Cuypers MG, Mason SA, Blakeley MP, Mitchell EP, Haertlein M, Forsyth VT. 2013a. Near-atomic resolution neutron crystallography on perdeuterated *Pyrococcus furiosus* rubredoxin: implication of hydronium ions and protonation state equilibria in redox changes. *Angew Chem Int Ed Engl.* 52:1022-1025.
- Cuypers MG, Mason SA, Mossou E, Haertlein M, Forsyth VT, Mitchell EP. 2016. Macromolecular structure phasing by neutron anomalous diffraction. *Sci Rep.* 6:31487.
- Cuypers MG, Trubitsyna M, Callow P, Forsyth VT, Richardson JM. 2013b. Solution conformations of early intermediates in Mos1 transposition. *Nucleic Acids Res.* 41:2020-2033.
- Dajnowicz S, Johnston RC, Parks JM, Blakeley MP, Keen DA, Weiss KL, Gerlits O, Kovalevsky A, Mueser TC. 2017. Direct visualization of critical hydrogen atoms in a pyridoxal 5'-phosphate enzyme. *Nat Commun.* 8:955.
- De Castro C, Molinaro A, Piacente F, Gurnon JR, Sturiale L, Palmigiano A, Lanzetta R, Parrilli M, Garozzo D, Tonetti MG, *et al.* 2013. Structure of N-linked oligosaccharides attached to chlorovirus PBCV-1 major capsid protein reveals unusual class of complex N-glycans. *Proc Natl Acad Sci U S A.* 110:13956-13960.
- Dean M, Sung VW. 2018. Review of deutetrabenazine: a novel treatment for chorea associated with Huntington's disease. *Drug Des Devel Ther.* 12:313-319.
- Drouillard S, Driguez H, Samain E. 2006. Large-scale synthesis of H-antigen oligosaccharides by expressing *Helicobacter pylori* alpha1,2-fucosyltransferase in metabolically engineered *Escherichia coli* cells. *Angew Chem Int Ed Engl.* 45:1778-1780.
- Dumon C, Bosso C, Utile JP, Heyraud A, Samain E. 2006. Production of Lewis x tetrasaccharides by metabolically engineered *Escherichia coli*. *Chembiochem.* 7:359-365.
- Dunne O, Weidenhaupt M, Callow P, Martel A, Moulin M, Perkins SJ, Haertlein M, Forsyth VT. 2017. Matchout deuterium labelling of proteins for small-angle neutron scattering studies using prokaryotic and eukaryotic expression systems and high cell-density cultures. *Eur Biophys J.* 46:425-432.
- Grage SL, Keleshian AM, Turzeladze T, Battle AR, Tay WC, May RP, Holt SA, Contera SA, Haertlein M, Moulin M, *et al.* 2011. Bilayer-mediated clustering and functional interaction of MscL channels. *Biophys J.* 100:1252-1260.
- Gupta M, Zha J, Zhang X, Jung GY, Linhardt RJ, Koffas MAG. 2018. Production of Deuterated Cyanidin 3-O-Glucoside from Recombinant *Escherichia coli*. *ACS Omega.* 3:11643-11648.

- Haertlein M, Moulin M, Devos JM, Laux V, Dunne O, Forsyth VT. 2016. Biomolecular Deuteration for Neutron Structural Biology and Dynamics. *Methods Enzymol.* 566:113-157.
- Haupt M, Blakeley MP, Fisher SJ, Mason SA, Cooper JB, Mitchell EP, Forsyth VT. 2014. Binding site asymmetry in human transthyretin: insights from a joint neutron and X-ray crystallographic analysis using perdeuterated protein. *IUCrJ.* 1:429-438.
- Heggelund JE, Varrot A, Imberty A, Krengel U. 2017. Histo-blood group antigens as mediators of infections. *Curr Opin Struct Biol.* 44:190-200.
- Hellstrand E, Grey M, Ainalem ML, Ankner J, Forsyth VT, Fragneto G, Haertlein M, Dauvergne MT, Nilsson H, Brundin P, *et al.* 2013. Adsorption of alpha-synuclein to supported lipid bilayers: positioning and role of electrostatics. *ACS Chem Neurosci.* 4:1339-1351.
- Josts I, Nitsche J, Maric S, Mertens HD, Moulin M, Haertlein M, Prevost S, Svergun DI, Busch S, Forsyth VT, *et al.* 2018. Conformational States of ABC Transporter MsbA in a Lipid Environment Investigated by Small-Angle Scattering Using Stealth Carrier Nanodiscs. *Structure.* 26:1072-1079 e1074.
- Jung SM, Park YC, Seo JH. 2019. Production of 3-Fucosyllactose in Engineered *Escherichia coli* with alpha-1,3-Fucosyltransferase from *Helicobacter pylori*. *Biotechnol J.* 14:e1800498.
- Katayama T, Sakuma A, Kimura T, Makimura Y, Hiratake J, Sakata K, Yamanoi T, Kumagai H, Yamamoto K. 2004. Molecular cloning and characterization of *Bifidobacterium bifidum* 1,2-alpha-L-fucosidase (AfcA), a novel inverting glycosidase (glycoside hydrolase family 95). *J Bacteriol.* 186:4885-4893.
- Koch HJ, Stuart RS. 1978. The synthesis of per-C-deuterated D-glucose. *Carbohydrate Research.* 64:127-134.
- Kwon H, Basran J, Devos JM, Suardiaz R, van der Kamp MW, Mulholland AJ, Schrader TE, Ostermann A, Blakeley MP, Moody PCE, *et al.* 2020. Visualizing the protons in a metalloenzyme electron proton transfer pathway. *Proc Natl Acad Sci U S A.* 117:6484-6490.
- Laux V, Callow P, Svergun DI, Timmins PA, Forsyth VT, Haertlein M. 2008. Selective deuteration of tryptophan and methionine residues in maltose binding protein: a model system for neutron scattering. *Eur Biophys J.* 37:815-822.
- Link AJ, Phillips D, Church GM. 1997. Methods for generating precise deletions and insertions in the genome of wild-type *Escherichia coli*: application to open reading frame characterization. *J Bacteriol.* 179:6228-6237.
- Liu JJ, Kwak S, Pathanibul P, Lee JW, Yu S, Yun EJ, Lim H, Kim KH, Jin YS. 2018. Biosynthesis of a Functional Human Milk Oligosaccharide, 2'-Fucosyllactose, and 1-Fucose Using Engineered *Saccharomyces cerevisiae*. *ACS Synth Biol.* 7:2529-2536.
- Liu JJ, Lee JW, Yun EJ, Jung SM, Seo JH, Jin YS. 2019. L-Fucose production by engineered *Escherichia coli*. *Biotechnol Bioeng.* 116:904-911.
- Lowery R, Gibson MI, Thompson RL, Fullam E. 2015. Deuterated carbohydrate probes as 'label-free' substrates for probing nutrient uptake in mycobacteria by nuclear reaction analysis. *Chem Commun (Camb).* 51:4838-4841.
- Maric S, Lind TK, Raida MR, Bengtsson E, Fredrikson GN, Rogers S, Moulin M, Haertlein M, Forsyth VT, Wenk MR, *et al.* 2019. Time-resolved small-angle neutron scattering as a probe for the dynamics of lipid exchange between human lipoproteins and naturally derived membranes. *Sci Rep.* 9:7591.
- Martinez E, Bartolome B, de la Cruz F. 1988. pACYC184-derived cloning vectors containing the multiple cloning site and lacZ alpha reporter gene of pUC8/9 and pUC18/19 plasmids. *Gene.* 68:159-162.

- Meilleur F, Weiss KL, Myles DA. 2009. Deuterium labeling for neutron structure-function-dynamics analysis. *Methods Mol Biol.* 544:281-292.
- Moulin M, Strohmeier GA, Hirz M, Thompson KC, Rennie AR, Campbell RA, Pichler H, Maric S, Forsyth VT, Haertlein M. 2018. Perdeuteration of cholesterol for neutron scattering applications using recombinant *Pichia pastoris*. *Chem Phys Lipids.* 212:80-87.
- O'Neill H, Shah R, Evans BR, He J, Pingali SV, Chundawat SP, Jones AD, Langan P, Davison BH, Urban V. 2015. Production of bacterial cellulose with controlled deuterium-hydrogen substitution for neutron scattering studies. *Methods Enzymol.* 565:123-146.
- Painter TJ, Watkins WM, Morgan WT. 1965. Serologically active fucose-containing oligosaccharides isolated from human blood-group A and B substances. *Nature.* 206:594-597.
- Priem B, Gilbert M, Wakarchuk WW, Heyraud A, Samain E. 2002. A new fermentation process allows large-scale production of human milk oligosaccharides by metabolically engineered bacteria. *Glycobiology.* 12:235-240.
- Samain E, Drouillard S, Heyraud A, Driguez H, Geremia RA. 1997. Gram-scale synthesis of recombinant chitoooligosaccharides in *Escherichia coli*. *Carbohydr Res.* 302:35-42.
- Sawama Y, Yabe Y, Iwata H, Fujiwara Y, Monguchi Y, Sajiki H. 2012. Stereo- and regioselective direct multi-deuterium-labeling methods for sugars. *Chemistry.* 18:16436-16442.
- Schmidt C. 2017. First deuterated drug approved. *Nat Biotechnol.* 35:493-494.
- Schneider M, Al-Shareffi E, Haltiwanger RS. 2017. Biological functions of fucose in mammals. *Glycobiology.* 27:601-618.
- Shimba S, Unno K, Okada S. 1990. Study on the biodistribution of deuterated biomolecules in mice aiming at new diagnostic radio-imaging agents. *Chem Pharm Bull (Tokyo).* 38:2610-2613.
- Timmins GS. 2017. Deuterated drugs; updates and obviousness analysis. *Expert Opin Ther Pat.* 27:1353-1361.
- Vanhooren PT, Vandamme EJ. 1999. L-Fucose: occurrence, physiological role, chemical, enzymatic and microbial synthesis. *Journal of Chemical Technology & Biotechnology.* 74:479-497.
- Waldie S, Lind TK, Browning K, Moulin M, Haertlein M, Forsyth VT, Luchini A, Strohmeier GA, Pichler H, Maric S, *et al.* 2018. Localization of Cholesterol within Supported Lipid Bilayers Made of a Natural Extract of Tailor-Deuterated Phosphatidylcholine. *Langmuir.* 34:472-479.
- Wei L, Yu Y, Shen Y, Wang MC, Min W. 2013. Vibrational imaging of newly synthesized proteins in live cells by stimulated Raman scattering microscopy. *Proc Natl Acad Sci U S A.* 110:11226-11231.
- Yee AW, Aldeghi M, Blakeley MP, Ostermann A, Mas PJ, Moulin M, de Sanctis D, Bowler MW, Mueller-Dieckmann C, Mitchell EP, *et al.* 2019. A molecular mechanism for transthyretin amyloidogenesis. *Nat Commun.* 10:925.
- Yu S, Liu JJ, Yun EJ, Kwak S, Kim KH, Jin YS. 2018. Production of a human milk oligosaccharide 2'-fucosyllactose by metabolically engineered *Saccharomyces cerevisiae*. *Microb Cell Fact.* 17:101.