



HAL
open science

New Insights into Asthma Inflammation: Focus on iNKT, MAIT, and $\gamma\delta$ T Cells

Jefferson Russo Victor, Guillaume Lezmi, Maria Leite-De-Moraes

► **To cite this version:**

Jefferson Russo Victor, Guillaume Lezmi, Maria Leite-De-Moraes. New Insights into Asthma Inflammation: Focus on iNKT, MAIT, and $\gamma\delta$ T Cells. *Clinical Reviews in Allergy and Immunology*, 2020, 10.1007/s12016-020-08784-8 . hal-02911836

HAL Id: hal-02911836

<https://hal.science/hal-02911836v1>

Submitted on 6 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **New insights into asthma inflammation: focus on iNKT, MAIT and $\gamma\delta$ T cells**

2
3 **Short title: Innate-like T cells and asthma**

4
5 Jefferson Russo Victor^{a, b} PhD, Guillaume Lezmi^{c, d} MD, PhD and
6 Maria Leite-de-Moraes^c PhD
7

8 **Affiliations:**

9 ^a-Laboratory of Medical Investigation LIM 56, Division of Clinical Dermatology, Medical
10 School, University of Sao Paulo, Sao Paulo, Brazil

11 ^b-Division of Environmental Health, FMU, Laureate International Universities, Sao Paulo,
12 Brazil

13 ^c-Laboratory of Immunoregulation and Immunopathology, INEM (Institut Necker-
14 Enfants Malades), CNRS UMR8253, INSERM UMR1151, and Université Paris Descartes,
15 Paris, France

16 ^d AP-HP, Hôpital Necker-Enfants Malades, Service de Pneumologie et d'Allergologie
17 Pédiatriques, Paris, France;
18
19

20 **Corresponding author:**

21 Maria Leite-de-Moraes, Laboratory of Immunoregulation and Immunopathology,
22 Institut Necker-Enfants Malades, CNRS UMR 8253, INSERM UMR 1151 and Université
23 Paris Descartes Sorbonne Paris Cité, 75015, Paris, France.

24 E-mail: maria.leite-de-moraes@parisdecartes.fr
25
26

27 **Word count: 3475**
28
29

30 **Abstract**

31

32 Asthma is a chronic immunological disease affecting all age groups, but often starting in
33 childhood. Although it has long been ascribed to a single pathology, recent studies have
34 highlighted its heterogeneity due to the potential involvement of various pathogenic
35 mechanisms. Here, we present our current understanding of the role of innate-like T
36 (ILT) cells in asthma pathogenesis. These cells constitute a specific family mainly
37 comprising $\gamma\delta$ T, invariant Natural Killer (iNKT) and Mucosal-Associated Invariant (MAIT)
38 T cells. They all share the ability to massively secrete a wide range of cytokines in a T cell
39 receptor (TCR)-dependent or -independent manner. ILT cells are prevalent in mucosal
40 tissues, including airways, where their innate and adaptive immune functions consist
41 primarily in protecting tissue integrity. However, ILT cells may also have detrimental
42 effects leading to asthma symptoms. The immune mechanisms through which this
43 pathogenic effect occurs will be discussed in this overview.

44

45

46 **Key Words:** Asthma, $\gamma\delta$ T cells, iNKT cells, MAIT cells, children, patients, CD1d, MR1

47

48 INTRODUCTION

49

50 Asthma is now considered an umbrella term encompassing various conditions
51 characterized by wheeze, cough, shortness of breath, chest tightness and variable
52 degrees of airflow limitation. These symptoms are associated with different patterns of
53 inflammation [1-4]. For the sake of clarity, an allergic and a non-allergic type of asthma
54 are commonly distinguished. In the first case, inflammation is primarily caused by type
55 2 immune responses mediated through the Th2 cytokines IL-4, IL-5, and IL-13 and
56 associated with increased Th2 cells and eosinophils in the airways [5]. By contrast, non-
57 allergic asthma is mainly triggered by an inflammatory response to viral infections with
58 a major neutrophilic component [6]. There is mounting evidence that neutrophilic forms
59 of murine and human asthma are associated with IL-17A (hereafter referred to as IL-17)
60 [7-9]. This cytokine is involved in the induction and persistence of neutrophilic
61 inflammation[10] and can contribute to airway smooth muscle remodeling in
62 association with TGF- β [11].

63

64 Asthma can adopt diverse phenotypes associated with a number of endotypes[12,2,4].
65 Previous reports have implicated Th2 and Th17 cells in mechanisms favoring asthmatic
66 immune endotypes [13]. However, these two populations alone cannot account for the
67 great diversity of symptoms, mainly because other cell types present in the airways can
68 also contribute to the production of Th2- and Th17-type cytokines. The best potential
69 candidates for this purpose are “unconventional” T cells also currently named “Innate-
70 Like T (ILT) lymphocytes” that recognize non-peptide antigens presented by specialized
71 MHC class I-like molecules [14-16].

72

73 Innate-like T (ILT) cells are unique unconventional T lymphocytes sharing features of
74 both innate and adaptive immunity. This population recognizes non-peptide antigens
75 presented by specialized MHC class I-like molecules. It includes invariant natural killer T
76 (iNKT), mucosal-associated invariant T (MAIT) and gamma-delta T ($\gamma\delta$ T) cells, all
77 characterized by a conserved and restricted T cell receptor (TCR) [14]. iNKT cells express
78 an antigen-specific semi-invariant TCR, V α 14-J α 18 (TRAV11-TRAJ18) and V α 24-J α 18
79 (TRAV10-TRAJ18) in mice and humans, respectively [17,16,14]. Similarly, MAIT cells
80 display V α 19-J α 33 (TRAV1-2-TRAJ33) in mice and V α 7.2-J α 33 (TRAV1-2-TRAJ33) in
81 humans [18-20]. The $\gamma\delta$ T cell population is not homogeneous, even though the V δ 2⁺
82 subset is predominant in human peripheral blood [21]. These cells can produce large
83 amounts of cytokines shortly after stimulation and play an important role in first-line
84 defense against microbial infections [22]. $\gamma\delta$ T, iNKT and MAIT cells are currently divided
85 into distinct subsets according to their cytokine profile (Table 1). They have all been
86 associated with a growing number of inflammatory diseases including asthma.

87

88 The next paragraphs will be focused on the development and functional properties of
89 ILT cells and their impact on the immune mechanisms leading to asthma.

90

91

92 **INNATE-LIKE T CELLS**

93

94 $\gamma\delta$ T cells

95

96 $\gamma\delta$ T cells constitute a distinct population of T lymphocytes in mice and humans, which
97 upon activation can promptly generate a number of cytokines and chemokines with
98 modulatory activities in innate and adaptive immunity [23-25]. Compared to other
99 unconventional T cells, they can readily perform effector functions in response to tissue
100 stress [22,23]. Furthermore, in epithelial tissues including lungs, $\gamma\delta$ T cells make up the
101 major resident T cell population and their functions, although still controversial, can be
102 related to tissue integrity and maintenance of host homeostasis [22].

103

104 $\gamma\delta$ T cells can be classified by the expression of their $\gamma\delta$ TCR chains. In humans, the
105 differential expression of δ (δ 1-3) and γ genes (γ 2-5, γ 8, γ 9, and γ 11) has been used to
106 this end, whereas in mice $\gamma\delta$ T cell subsets are defined by the expression of distinct γ
107 chains[26] (Table 1). Contrasting with conventional T cells, which recognize peptides
108 presented by MHC molecules, $\gamma\delta$ T cells can distinguish uncommon antigens, such as
109 stress molecules (MICA and MICB), heat-shock proteins or synthetic phosphoantigens
110 (pAg) [27,28]. It is clear by now that $\gamma\delta$ T cells do not interact directly with pAg but
111 require the presence of butyrophilin BTN3A as an intermediate [29]. This indicates the
112 usage of a fundamentally different mechanism of antigen detection whose molecular
113 basis remains to be elucidated.

114

115 Although $\gamma\delta$ T cells can produce several cytokines, two major murine subsets defined by
116 their capacity to produce IL-17 or IFN- γ have been described in recent years [30] (Table
117 1). Compared to conventional TCR $\alpha\beta$ ⁺ CD4 and CD8 T cells, $\gamma\delta$ T cells are more prompt in
118 producing significant amounts of IFN- γ [31], thus providing an early source of this factor
119 that could be protective in tumor immunity [32-34]. However, in some cases, such as
120 fulminant viral hepatitis in a murine model, the production of IFN- γ by $\gamma\delta$ T cells can
121 become deleterious [35]. $\gamma\delta$ T cells that produce substantial amounts of IL-17 are
122 frequently termed $\gamma\delta$ T17 and display a CD27⁻NK1.1⁺CCR6⁺ phenotype [30,36]. They can
123 take part in tissue surveillance and repair and mediate effective immune responses
124 against bacterial and fungal infections [37-40]. Conversely, they are potential promoters
125 of inflammation in type 1 autoimmune diseases, such as arthritis or colitis [41,42].
126 According to recent reports, $\gamma\delta$ T17 cells could also promote tumor progression in
127 experimental models [43].

$\gamma\delta$ T cells in experimental models of asthma

The role of $\gamma\delta$ T cells in lung inflammation has been debated in the literature for more than 30 years. In 1998, one of the first publications supporting the requirement of $\gamma\delta$ T cells for promoting IgE production and Th2 airway inflammation in an IL-4-dependent manner came from experiments with $\gamma\delta$ T cell-deficient mice, which failed to develop the pathology [44]. One year later, another team reported that airway reactivity was increased in the absence of $\gamma\delta$ T cells, suggesting a protective role of these cells [45]. These controversial data could be explained by the distinct genetic backgrounds, namely BALB/c [44] and C57BL/6 [45] in these $\gamma\delta$ T cell-deficient mice. Indeed, these strains respond differently in the asthma protocol since BALB/c display more severe AHR than C57BL/6 mice, while airway eosinophilia is more pronounced in the C57BL/6 strain [46].

Further studies using blocking anti- $\gamma\delta$ TCR mAb administered during the resolution phase of OVA-induced allergic inflammation, provided evidence for an exacerbation of lung injury [47]. However, it turned out that GL3 mAb used in this study did not deplete $\gamma\delta$ T cells in vivo, but induced the downmodulation of their $\gamma\delta$ TCR [48].

The complexity of $\gamma\delta$ T cell regulation of asthmatic inflammation may be explained by the intervention of distinct subsets. In this line of evidence, it has been shown that the effect of $\gamma\delta$ T cells on IgE production in an OVA-induced allergic asthma model changed according to the phenotype of the $\gamma\delta$ T cell population involved [49]. Indeed V γ 1 $\gamma\delta$ T cells promoted an enhanced IgE response, while their V γ 4 counterpart suppressed this activity [50]. These non-overlapping functions of V γ 4 and V γ 1 $\gamma\delta$ T cell subsets were also observed in terms of AHR [51]. Consistent with this observation, it has been shown more recently that IL-4-producing V γ 1⁺ $\gamma\delta$ T cells promoted airway eosinophilia and AHR in experimental allergic asthma induced by the dust mite *Blomia tropicalis* [52].

$\gamma\delta$ T cells also produce IFN- γ and IL-17. It has been proposed that the IFN- γ thus generated might favor allergic inhibition by reducing IgE production [53], while IL-17-producing $\gamma\delta$ T cells could ameliorate AHR and accelerate the resolution of eosinophilic airway inflammation [47]. Furthermore, a relationship between reduced thymic maturation of IL-17-producing $\gamma\delta$ T cells and decreased lung inflammation has been described in a murine protocol of OVA-induced allergy [54].

$\gamma\delta$ T cells in asthmatic patients

In humans, peripheral $\gamma\delta$ T cells were reported to be decreased in difficult-to-control asthmatic patients compared to controls [55]. Further, a higher frequency of IL-4-producing versus a lower incidence of IFN- γ -producing $\gamma\delta$ T cells was described in patients with allergic asthma, as compared with healthy controls [56].

169

170 In another study $\gamma\delta$ T cells were found to be more frequent at the site of inflammation
171 in bronchoalveolar lavage fluid (BALF) of untreated asthmatic patients, as compared
172 with control volunteers [57]. Among the scarce data available on this issue, according to
173 another group, $\gamma\delta$ T cells occurred at similar levels in BALF or lung biopsies, whether the
174 donors were suffering from asthma or not [58], while another research confirmed that
175 the frequency of $\gamma\delta$ T cells in BALF from stable asthmatic subjects was not significantly
176 different from controls [59]. However, after *in vivo* allergen challenge, BALF $\gamma\delta$ T cells
177 increased their capacity to produce Th2-type cytokines, including IL-5 and IL-13, and
178 reduced their secretion of Th1-type cytokines, such as IFN- γ [59]. These findings suggest
179 that the $\gamma\delta$ T cytokine profile is flexible in BALF, ready to be skewed toward the Th2-type
180 in asthmatic patients [59].

181

182 Overall, the studies cited here reveal that $\gamma\delta$ T cells exert a highly complex control in lung
183 inflammation, possibly because of the contribution of various subsets whose
184 composition could differ according to the clinical status of asthmatic patients (Fig.1).

185

186 **invariant Natural Killer T cells**

187

188 Differentiation and maturation of iNKT cells occur in the thymus, where they appear
189 slightly later than MHC-restricted conventional T cells [60-63]. Thymic iNKT cell
190 differentiation requires a particular rearrangement between V α and J α gene segments
191 in the TCR α locus. This is believed to be a random process that occurs at very low
192 frequency leading to the characteristic iNKT cell invariant TCR α chain: V α 14J α 18
193 (TRAV11-TRAJ18) in mice and V α 24J α 18 (TRAV10-TRAJ18) in humans[64,65]. The TCR α
194 chain associates with a limited set of TCR β chains [66-68] (Table 1).

195

196 Once they have completed a tightly controlled thymic differentiation pathway, iNKT cells
197 migrate to peripheral organs as CD4⁺CD8⁻ and double-negative (CD4⁻CD8⁻) cells. In
198 humans, the CD4⁻CD8⁺ subset does likewise occur [69]. iNKT cells are mostly resident in
199 tissues, where they are believed to act as sentinels of tissue integrity by “patrolling” to
200 identify threats to the body [70]. They are rare in most peripheral sites (0.1-1% among
201 T lymphocytes) but numerous in the liver [16]. The invariant TCR α chains in mice and
202 humans are conserved, enabling iNKT cells to recognize the same glycolipids in both
203 species, as is the case for the glycolipid α -galactosylceramide (α -GalCer) [71,72].

204

205 iNKT cells are restricted by lipids and glycolipids loaded onto CD1d molecules present
206 on the surface of APCs [73]. Moreover, their cytokine production along with their
207 cognate interactions, enable them to modulate the activities of different immune cells
208 playing a pivotal role in the orchestration of the immune response [74].

209

iNKT cells in experimental models of asthma

210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250

In the nineties, several studies demonstrated that iNKT cells promptly and massively secreted IL-4 [75-78]. Consequently, it was proposed that their major role was to promote Th2 differentiation via this cytokine [79]. This has led a number of groups to test IL-4-producing iNKT cells in specific experimental models requiring Th2 cells, including allergic asthma models.

Initial studies using iNKT cell-deficient mice, namely β_2 microglobulin (β_2m)^{-/-} and CD1d^{-/-} strains, revealed no major differences in the severity of allergic airway inflammation [80-82]. More recently, other investigators using these same mutants concluded that NKT cells were dispensable for T cell-dependent allergic airway inflammation [83,84]. However, it turned out that both β_2m ^{-/-} and CD1d^{-/-} mice are also deficient for other T cell subsets (CD8⁺ T and CD1d-dependent T lymphocytes, respectively), thus masking a potential effect of iNKT cells. To solve this issue, two groups used more specific iNKT cell-deficient ($J\alpha 18$ ^{-/-}) mice, for which they reported attenuated asthma symptoms, including AHR, airway eosinophilia, Th2 inflammation and OVA-specific anti-IgE production relative to wild-type controls [85,86]. The adoptive transfer of wild type iNKT cells was sufficient to restore asthma severity in $J\alpha 18$ ^{-/-} mice. Of note, adoptive transfers of iNKT cells from IL-4^{-/-} or IL-13^{-/-} mice were much less effective and iNKT cells from double knockout IL-4^{-/-}IL-13^{-/-} mice had no effect at all [86]. This finding demonstrated that iNKT cells enhanced allergic asthma symptoms through the production of IL-4 and IL-13 [85,86].

Recently, a new mouse strain with the deletion of *Traj18* was created on a C57BL/6 genetic background [90]. These *Traj18*-deficient mice lack iNKT cells but have a normal $J\alpha$ -region repertoire. They display significantly less eosinophils and AHR in an airway hyper-reactivity model [90], thereby confirming the implication of iNKT cells in asthmatic airway inflammation. In further support of this conclusion, *Ezh2* (Enhancer of zeste homolog 2)-deficient mice spontaneously developed an asthma-like disease [91] associated with abnormally high iNKT cell counts, increased IL-4 and IL-13 production, AHR, lung inflammation and IgE levels, all consistent with a contribution of iNKT cells to asthma pathogenesis [91].

Even though iNKT cells do not recognize OVA as an antigen, their ability to promote lung inflammation is TCR-dependent since the injection of blocking anti-CD1d antibodies reduced the effect, indicating that endogenous lipidic antigens stimulated the iNKT cells [85]. Indeed, in this setting we cannot exclude that commensal microbial antigens present in the lung could take part in the activation of iNKT cells. In the same line of evidence, an elegant study by Blumberg's team showed that iNKT cells accumulated in the lung and in the colonic lamina propria in germ-free (GF) mice, rendering these

251 animals more susceptible to OVA-induced asthma and oxazolone-induced colitis [87].
252 The colonization of neonatal GF mice with a normal flora or *Bacteroides fragilis*
253 decreased the number of iNKT cells and protected the mice against these diseases,
254 clearly establishing a link between iNKT cells and the microbiota [87,88]. Knowing that
255 a bacterial community is present in the airways and that this microbiota could be altered
256 in the case of asthma [89], we postulate that some bacterial antigens might activate
257 iNKT cells in the lung, thus promoting inflammation. Further studies are required to test
258 this hypothesis.

259

260 For asthma that is not associated with pro-Th2 allergic immune responses, it has been
261 reported that i.n. administration of α -GalCer activates IL-17-secreting iNKT cells, which,
262 in turn, promote airway neutrophilia [92]. IL-17-secreting iNKT cells are also required for
263 the pathogenic mechanisms leading to AHR after exposure to ozone, a major component
264 of air pollution [93]. Hence, iNKT cells may promote asthma inflammation in different
265 ways, depending on their cytokine profile (Fig. 2).

266

267 **iNKT cells in asthmatic patients**

268

269 Analysis of the incidence of iNKT cells in the peripheral blood of adult asthmatic patients,
270 revealed no significant differences when compared to control subjects [94]. Further
271 studies suggested however that IL-4-producing iNKT cells were more frequent in these
272 cases and that they were associated with lung functions [95]. In children with asthma,
273 the percentage of peripheral blood iNKT cells did not differ significantly between
274 asthmatic children classified as exacerbators (1 or more severe exacerbations in the last
275 12 months) and non-exacerbators [96]. Similarly, no relationship was observed between
276 the frequency of circulating iNKT cells and IL-4- or IFN- γ -producing iNKT cells in one-
277 year-old children and asthma-related clinical outcomes at the age of seven years [97].

278

279 In addition to peripheral blood analysis, a number of studies have evaluated iNKT cells
280 in BALF or bronchial biopsies of asthmatic patients. The first data in adult asthmatic
281 patients revealed that about 60% of the pulmonary CD4⁺CD3⁺ T cells in the group with
282 moderate-to-severe persistent asthma were iNKT cells [98]. These results were not
283 reproduced by others [99,100], who found that up to 2% of the T cells recovered from
284 airway biopsies, BALF and sputum induction from subjects with mild or moderately
285 severe asthma were iNKT cells. Umetsu and collaborators reported later that only a
286 small fraction of T cells in the lung of adult asthmatic patients were iNKT cells [101],
287 which accounted for less than 1% of T cells in BALF from children with severe asthma
288 [102]. The discrepancies between these studies could be explained by the limited
289 number of samples, the heterogeneity of the cohorts, or by non-specific staining of cells
290 in BALF, as suggested by the study of Thomas *et al* [100].

291

292 Taken together, these findings show that large adult and children cohorts composed of
293 asthmatic patients with well-defined symptoms and grades of severity (controlled, non-
294 controlled and severe asthma) associated with relevant controls are required to
295 definitely assess whether iNKT cells are involved in asthma pathology or not.

296

297 Mucosal-Associated Invariant T cells

298

299 MAIT cells are restricted by the MHC class I-related molecule MR1 and recognize
300 microbial-derived vitamin B2 (riboflavin) metabolites [103] (Table 1). Riboflavin
301 metabolites are not present in mammalian cells but occur in several bacteria and yeast
302 [104], which suggests that MAIT cells can sense antigens displayed by a limited number
303 of microbes.

304

305 MAIT cells develop in the thymus, where they become functionally competent and able
306 to promptly produce Th1 (IFN- γ , TNF- α) and Th17 (IL-17 and IL-22) cytokines in response
307 to stimulation [105]. In contrast, MAIT cells produce only low to moderate levels of Th2
308 (IL-4 and IL-13) cytokines [106-109]. Similarly to iNKT cells, MAIT precursors undergo a
309 positive selection of MR1⁺ double-positive (CD4⁺CD8⁺) thymocytes [110,111]. Previous
310 reports have stressed the key role of the microbiota in MAIT cell expansion since these
311 cells are nearly absent in germ-free mice [20]. Recently, Lantz's team made a major
312 breakthrough in this field by demonstrating that microbial metabolites control the
313 thymic development of MAIT cells [112]. This finding paved the way for revisiting the
314 concept of exogenous and self-antigens.

315

316 Human MAIT cells express high levels of CD161 and IL-18R α and are mainly CD8⁺ [113].
317 A smaller proportion among these is double-negative (CD4⁻CD8⁻) and some CD4⁺ cells
318 have been described in the literature [114]. MAIT cells can be activated in a TCR-
319 dependent and -independent manner. They account for up to 10% of peripheral blood
320 T cells and are numerous in the gut, lung, and liver [115,113,20]. Their abundance in
321 mucosal tissues supports their involvement in antimicrobial responses [20].

322

323 Because of the scarce pieces of evidence for a role of MAIT cells in experimental models
324 of asthma, we will not discuss this issue, but pass on directly to studies with asthmatic
325 patients.

326

327 **MAIT cells in asthmatic patients**

328

329 As indicated by their name, MAIT cells are preferentially associated with mucosal tissues
330 including lungs, where they promote antimicrobial defenses through the secretion of
331 IFN- γ , TNF- α , IL-17, perforin and granzyme B [116-120]. Evidence for MAIT cell expansion
332 in the lungs of MAIT cell-enriched V α 19iC α ^{-/-}MR1^{+/+} mice experimentally infected with

333 *Mycobacterium tuberculosis* supports their role during pulmonary infection [121]. This
334 proliferative response was rapid and depended on the MR1 presentation of riboflavin
335 biosynthesis-derived bacterial ligands [121]. In line with these experimental data it has
336 been reported that adult patients suffering from active pulmonary tuberculosis (TB) had
337 more MAIT cells in BALF and lung parenchyma than uninfected controls [122]. However,
338 in children with similar symptoms, reduced MAIT cell counts were found both in
339 peripheral blood and BALF [123]. This discrepancy might be due to clinical differences
340 between pediatric and adult active TB, calling for further studies to ascertain the role of
341 MAIT cells in in this infectious disease.

342

343 The data gained from lung infections prompted several groups to address the
344 involvement of MAIT cells in other lung pathologies like asthma. Pioneer studies by
345 Hinks et al. reported that the frequency of MAIT cells was significantly lower in
346 peripheral blood, sputum and bronchial biopsy samples of asthmatic patients, relative
347 to control subjects [124]. This deficiency was correlated with clinical severity and the
348 use of oral corticosteroids [124].

349

350 A recent study provided evidence for a relationship between a high frequency of
351 circulating MAIT cells in one-year-old children and a lower risk of developing asthma by
352 the age of seven years [97]. Furthermore, these high MAIT cell levels were correlated
353 with higher frequency of IFN- γ -producing CD4⁺ T cells, suggesting a possible protective
354 effect of MAIT cells as children grow older [97]. IL-17 production by MAIT cells did not
355 correlate with asthma in this study. In children with mild asthma, the frequency of
356 circulating MAIT cells did not differ between exacerbators and non-exacerbators [96].
357 However, the proportion of MAIT cells producing IL-17 correlated positively with the
358 number of severe exacerbations and negatively with the asthma control test score in
359 these children, which was not the case for their IFN- γ -producing counterpart [96]. More
360 recently, the same team showed that MAIT cells, including their IL-17-producing subset,
361 were present in BALF from children with severe asthma [125]. It is noteworthy that when
362 this group was divided into two according to the frequency of their IL-17-producing
363 MAIT cells in BALF into MAIT17-low and MAIT17-high, the latter presented more severe
364 exacerbations than their MAIT17-low counterpart [125].

365

366 Overall, the results of these studies suggest that IL-17-producing MAIT cells may be
367 associated with asthma symptoms, whereas the pro-Th1 subset might provide
368 protection [97,96,125] (Fig. 3).

369

370

371

372 **CONCLUDING REMARKS**

373

374 Much progress has been achieved in understanding the physiological and pathological
375 role of $\gamma\delta$ T, iNKT and MAIT cells, which have by now acquired a lasting position among
376 the complex immune mechanisms leading to distinct asthma phenotypes and
377 endotypes. Nonetheless, major uncertainties remain. For instance, asthma often begins
378 early in childhood, at a time when the amount and functional properties of lung $\gamma\delta$ T,
379 iNKT and MAIT cells are about to be established. However, very few studies have
380 addressed the evolution of these populations in the airways with age. The cross-talk
381 between lung $\gamma\delta$ T, iNKT and MAIT cells needs likewise to be investigated in more detail,
382 as well as the influence of microorganisms present in the lung. In sum, studies of
383 experimental models and pediatric asthma will be crucial for the comprehension of
384 these complex interactions and the development of new therapeutic approaches.

385

386

387 **Funding**

388 This work was supported by the grant ANR-18-CE14-0011-01 SevAsthma-children, Paris,
389 France.

390

391 **Compliance with Ethical Standards**

392 **Conflict of Interest.** The authors declare that they have no conflict of interest.

393 **Ethical Approval and Informed Consent**

394 No approvals or informed consents were obtained, as this manuscript does not contain
395 primary research data.

396

397 **REFERENCES**

- 398 1. Reddel HK, FitzGerald JM, Bateman ED, Bacharier LB, Becker A, Brusselle G, Buhl R,
399 Cruz AA, Fleming L, Inoue H, Ko FW, Krishnan JA, Levy ML, Lin J, Pedersen SE, Sheikh
400 A, Yorgancioglu A, Boulet LP (2019) GINA 2019: a fundamental change in asthma
401 management: Treatment of asthma with short-acting bronchodilators alone is no
402 longer recommended for adults and adolescents. *Eur Respir J* 53 (6).
403 doi:10.1183/13993003.01046-2019
- 404 2. Lötvall J, Akdis CA, Bacharier LB, Bjermer L, Casale TB, Custovic A, Lemanske RF,
405 Wardlaw AJ, Wenzel SE, Greenberger PA (2011) Asthma endotypes: a new approach
406 to classification of disease entities within the asthma syndrome. *J Allergy Clin
407 Immunol* 127 (2):355-360. doi:10.1016/j.jaci.2010.11.037
- 408 3. Kaur R, Chupp G (2019) Phenotypes and endotypes of adult asthma: Moving toward
409 precision medicine. *J Allergy Clin Immunol* 144 (1):1-12.
410 doi:10.1016/j.jaci.2019.05.031
- 411 4. Pavord ID, Beasley R, Agusti A, Anderson GP, Bel E, Brusselle G, Cullinan P, Custovic A,
412 Ducharme FM, Fahy JV, Frey U, Gibson P, Heaney LG, Holt PG, Humbert M, Lloyd
413 CM, Marks G, Martinez FD, Sly PD, von Mutius E, Wenzel S, Zar HJ, Bush A (2018)
414 After asthma: redefining airways diseases. *Lancet* 391 (10118):350-400.
415 doi:10.1016/S0140-6736(17)30879-6
- 416 5. Wenzel SE (2006) Asthma: defining of the persistent adult phenotypes. *Lancet* 368
417 (9537):804-813. doi:10.1016/S0140-6736(06)69290-8
- 418 6. Wenzel SE, Schwartz LB, Langmack EL, Halliday JL, Trudeau JB, Gibbs RL, Chu HW
419 (1999) Evidence that severe asthma can be divided pathologically into two
420 inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J
421 Respir Crit Care Med* 160 (3):1001-1008. doi:10.1164/ajrccm.160.3.9812110
- 422 7. Wilson RH, Whitehead GS, Nakano H, Free ME, Kolls JK, Cook DN (2009) Allergic
423 sensitization through the airway primes Th17-dependent neutrophilia and airway

- 424 hyperresponsiveness. *Am J Respir Crit Care Med* 180 (8):720-730.
425 doi:10.1164/rccm.200904-0573OC
- 426 8. Bullens DM, Truyen E, Coteur L, Dilissen E, Hellings PW, Dupont LJ, Ceuppens JL (2006)
427 IL-17 mRNA in sputum of asthmatic patients: linking T cell driven inflammation and
428 granulocytic influx? *Respir Res* 7:135. doi:10.1186/1465-9921-7-135
- 429 9. Chesné J, Braza F, Chadeuf G, Mahay G, Cheminant MA, Loy J, Brouard S, Sauzeau V,
430 Loirand G, Magnan A (2015) Prime role of IL-17A in neutrophilia and airway smooth
431 muscle contraction in a house dust mite-induced allergic asthma model. *J Allergy
432 Clin Immunol* 135 (6):1643-1643.e1643. doi:10.1016/j.jaci.2014.12.1872
- 433 10. Lindén A (2001) Role of interleukin-17 and the neutrophil in asthma. *Int Arch Allergy
434 Immunol* 126 (3):179-184. doi:10.1159/000049511
- 435 11. Evasovic JM, Singer CA (2019) Regulation of IL-17A and implications for TGF- β 1
436 comodulation of airway smooth muscle remodeling in severe asthma. *Am J Physiol
437 Lung Cell Mol Physiol* 316 (5):L843-L868. doi:10.1152/ajplung.00416.2018
- 438 12. Agache I (2019) Severe asthma phenotypes and endotypes. *Semin Immunol*:101301.
439 doi:10.1016/j.smim.2019.101301
- 440 13. Choy DF, Hart KM, Borthwick LA, Shikotra A, Nagarkar DR, Siddiqui S, Jia G, Ohri CM,
441 Doran E, Vannella KM, Butler CA, Hargadon B, Sciurba JC, Gieseck RL, Thompson
442 RW, White S, Abbas AR, Jackman J, Wu LC, Egen JG, Heaney LG, Ramalingam TR,
443 Arron JR, Wynn TA, Bradding P (2015) TH2 and TH17 inflammatory pathways are
444 reciprocally regulated in asthma. *Sci Transl Med* 7 (301):301ra129.
445 doi:10.1126/scitranslmed.aab3142
- 446 14. Godfrey DI, Uldrich AP, McCluskey J, Rossjohn J, Moody DB (2015) The burgeoning
447 family of unconventional T cells. *Nat Immunol* 16 (11):1114-1123.
448 doi:10.1038/ni.3298
- 449 15. Salio M, Silk JD, Jones EY, Cerundolo V (2014) Biology of CD1- and MR1-restricted T
450 cells. *Annu Rev Immunol* 32:323-366. doi:10.1146/annurev-immunol-032713-
451 120243
- 452 16. Crosby CM, Kronenberg M (2018) Tissue-specific functions of invariant natural killer
453 T cells. *Nat Rev Immunol* 18 (9):559-574. doi:10.1038/s41577-018-0034-2
- 454 17. Lantz O, Bendelac A (1994) An invariant T cell receptor alpha chain is used by a
455 unique subset of major histocompatibility complex class I-specific CD4+ and CD4-8-
456 T cells in mice and humans. *J Exp Med* 180 (3):1097-1106
- 457 18. Porcelli S, Yockey CE, Brenner MB, Balk SP (1993) Analysis of T cell antigen receptor
458 (TCR) expression by human peripheral blood CD4-8- alpha/beta T cells
459 demonstrates preferential use of several V beta genes and an invariant TCR alpha
460 chain. *J Exp Med* 178 (1):1-16. doi:10.1084/jem.178.1.1
- 461 19. Tilloy F, Treiner E, Park SH, Garcia C, Lemonnier F, de la Salle H, Bendelac A,
462 Bonneville M, Lantz O (1999) An invariant T cell receptor alpha chain defines a novel
463 TAP-independent major histocompatibility complex class Ib-restricted alpha/beta T
464 cell subpopulation in mammals. *J Exp Med* 189 (12):1907-1921.
465 doi:10.1084/jem.189.12.1907
- 466 20. Treiner E, Duban L, Bahram S, Radosavljevic M, Wanner V, Tilloy F, Affaticati P,
467 Gilfillan S, Lantz O (2003) Selection of evolutionarily conserved mucosal-associated
468 invariant T cells by MR1. *Nature* 422 (6928):164-169. doi:10.1038/nature01433

- 469 21. Kalyan S, Kabelitz D (2013) Defining the nature of human $\gamma\delta$ T cells: a biographical
470 sketch of the highly empathetic. *Cell Mol Immunol* 10 (1):21-29.
471 doi:10.1038/cmi.2012.44
- 472 22. Nielsen MM, Witherden DA, Havran WL (2017) $\gamma\delta$ T cells in homeostasis and host
473 defence of epithelial barrier tissues. *Nat Rev Immunol* 17 (12):733-745.
474 doi:10.1038/nri.2017.101
- 475 23. Hayday AC (2009) Gammadelta T cells and the lymphoid stress-surveillance
476 response. *Immunity* 31 (2):184-196. doi:10.1016/j.immuni.2009.08.006
- 477 24. Hayday A, Tigelaar R (2003) Immunoregulation in the tissues by gammadelta T cells.
478 *Nat Rev Immunol* 3 (3):233-242. doi:10.1038/nri1030
- 479 25. Bonneville M, O'Brien RL, Born WK (2010) Gammadelta T cell effector functions: a
480 blend of innate programming and acquired plasticity. *Nat Rev Immunol* 10 (7):467-
481 478. doi:10.1038/nri2781
- 482 26. Adams EJ, Gu S, Luoma AM (2015) Human gamma delta T cells: Evolution and ligand
483 recognition. *Cell Immunol* 296 (1):31-40. doi:10.1016/j.cellimm.2015.04.008
- 484 27. Wu YL, Ding YP, Tanaka Y, Shen LW, Wei CH, Minato N, Zhang W (2014) $\gamma\delta$ T cells
485 and their potential for immunotherapy. *Int J Biol Sci* 10 (2):119-135.
486 doi:10.7150/ijbs.7823
- 487 28. Gu S, Nawrocka W, Adams EJ (2014) Sensing of Pyrophosphate Metabolites by
488 $V\gamma 9V\delta 2$ T Cells. *Front Immunol* 5:688. doi:10.3389/fimmu.2014.00688
- 489 29. Vantourout P, Laing A, Woodward MJ, Zlatareva I, Apolonia L, Jones AW, Snijders AP,
490 Malim MH, Hayday AC (2018) Heteromeric interactions regulate butyrophilin (BTN)
491 and BTN-like molecules governing $\gamma\delta$ T cell biology. *Proc Natl Acad Sci U S A* 115
492 (5):1039-1044. doi:10.1073/pnas.1701237115
- 493 30. Ribot JC, deBarros A, Pang DJ, Neves JF, Peperzak V, Roberts SJ, Girardi M, Borst J,
494 Hayday AC, Pennington DJ, Silva-Santos B (2009) CD27 is a thymic determinant of
495 the balance between interferon-gamma- and interleukin 17-producing gammadelta
496 T cell subsets. *Nat Immunol* 10 (4):427-436. doi:10.1038/ni.1717
- 497 31. Chen L, He W, Kim ST, Tao J, Gao Y, Chi H, Intlekofer AM, Harvey B, Reiner SL, Yin Z,
498 Flavell RA, Craft J (2007) Epigenetic and transcriptional programs lead to default
499 IFN-gamma production by gammadelta T cells. *J Immunol* 178 (5):2730-2736
- 500 32. He W, Hao J, Dong S, Gao Y, Tao J, Chi H, Flavell R, O'Brien RL, Born WK, Craft J, Han
501 J, Wang P, Zhao L, Wu J, Yin Z (2010) Naturally activated V gamma 4 gamma delta T
502 cells play a protective role in tumor immunity through expression of eomesodermin.
503 *J Immunol* 185 (1):126-133. doi:10.4049/jimmunol.0903767
- 504 33. Gao Y, Yang W, Pan M, Scully E, Girardi M, Augenlicht LH, Craft J, Yin Z (2003) Gamma
505 delta T cells provide an early source of interferon gamma in tumor immunity. *J Exp*
506 *Med* 198 (3):433-442. doi:10.1084/jem.20030584
- 507 34. Girardi M, Oppenheim DE, Steele CR, Lewis JM, Glusac E, Filler R, Hobby P, Sutton B,
508 Tigelaar RE, Hayday AC (2001) Regulation of cutaneous malignancy by gammadelta
509 T cells. *Science* 294 (5542):605-609. doi:10.1126/science.1063916
- 510 35. Wu D, Yan WM, Wang HW, Huang D, Luo XP, Ning Q (2018) $\gamma\delta$ T Cells Contribute to
511 the Outcome of Murine Fulminant Viral Hepatitis via Effector Cytokines TNF- α and
512 IFN- γ . *Curr Med Sci* 38 (4):648-655. doi:10.1007/s11596-018-1926-x
- 513 36. Haas JD, González FH, Schmitz S, Chennupati V, Föhse L, Kremmer E, Förster R, Prinz
514 I (2009) CCR6 and NK1.1 distinguish between IL-17A and IFN-gamma-producing

- 515 gammadelta effector T cells. *Eur J Immunol* 39 (12):3488-3497.
516 doi:10.1002/eji.200939922
- 517 37. Puel A, Cypowyj S, Bustamante J, Wright JF, Liu L, Lim HK, Migaud M, Israel L,
518 Chrabieh M, Audry M, Gumbleton M, Toulon A, Bodemer C, El-Baghdadi J, Whitters
519 M, Paradis T, Brooks J, Collins M, Wolfman NM, Al-Muhsen S, Galicchio M, Abel L,
520 Picard C, Casanova JL (2011) Chronic mucocutaneous candidiasis in humans with
521 inborn errors of interleukin-17 immunity. *Science* 332 (6025):65-68.
522 doi:10.1126/science.1200439
- 523 38. Conti HR, Peterson AC, Brane L, Huppler AR, Hernández-Santos N, Whibley N, Garg
524 AV, Simpson-Abelson MR, Gibson GA, Mamo AJ, Osborne LC, Bishu S, Ghilardi N,
525 Siebenlist U, Watkins SC, Artis D, McGeachy MJ, Gaffen SL (2014) Oral-resident
526 natural Th17 cells and $\gamma\delta$ T cells control opportunistic *Candida albicans* infections. *J*
527 *Exp Med* 211 (10):2075-2084. doi:10.1084/jem.20130877
- 528 39. Cho JS, Pietras EM, Garcia NC, Ramos RI, Farzam DM, Monroe HR, Magorien JE,
529 Blauvelt A, Kolls JK, Cheung AL, Cheng G, Modlin RL, Miller LS (2010) IL-17 is essential
530 for host defense against cutaneous *Staphylococcus aureus* infection in mice. *J Clin*
531 *Invest* 120 (5):1762-1773. doi:10.1172/JCI40891
- 532 40. Sumaria N, Roediger B, Ng LG, Qin J, Pinto R, Cavanagh LL, Shklovskaya E, Fazekas de
533 St Groth B, Triccas JA, Weninger W (2011) Cutaneous immunosurveillance by self-
534 renewing dermal gammadelta T cells. *J Exp Med* 208 (3):505-518.
535 doi:10.1084/jem.20101824
- 536 41. Papotto PH, Ribot JC, Silva-Santos B (2017) IL-17. *Nat Immunol* 18 (6):604-611.
537 doi:10.1038/ni.3726
- 538 42. Papotto PH, Reinhardt A, Prinz I, Silva-Santos B (2018) Innately versatile: $\gamma\delta$ 17 T cells
539 in inflammatory and autoimmune diseases. *J Autoimmun* 87:26-37.
540 doi:10.1016/j.jaut.2017.11.006
- 541 43. Rei M, Gonçalves-Sousa N, Lança T, Thompson RG, Mensurado S, Balkwill FR, Kulbe
542 H, Pennington DJ, Silva-Santos B (2014) Murine CD27(-) V γ 6(+) $\gamma\delta$ T cells producing
543 IL-17A promote ovarian cancer growth via mobilization of protumor small
544 peritoneal macrophages. *Proc Natl Acad Sci U S A* 111 (34):E3562-3570.
545 doi:10.1073/pnas.1403424111
- 546 44. Zuany-Amorim C, Ruffié C, Hailé S, Vargaftig BB, Pereira P, Pretolani M (1998)
547 Requirement for gammadelta T cells in allergic airway inflammation. *Science* 280
548 (5367):1265-1267
- 549 45. Lahn M, Kanehiro A, Takeda K, Joetham A, Schwarze J, Köhler G, O'Brien R, Gelfand
550 EW, Born W, Kanehio A (1999) Negative regulation of airway responsiveness that is
551 dependent on gammadelta T cells and independent of alphabeta T cells. *Nat Med* 5
552 (10):1150-1156. doi:10.1038/13476
- 553 46. Gueders MM, Paulissen G, Crahay C, Quesada-Calvo F, Hacha J, Van Hove C, Tournoy
554 K, Louis R, Foidart JM, Noël A, Cataldo DD (2009) Mouse models of asthma: a
555 comparison between C57BL/6 and BALB/c strains regarding bronchial
556 responsiveness, inflammation, and cytokine production. *Inflamm Res* 58 (12):845-
557 854. doi:10.1007/s00011-009-0054-2
- 558 47. Murdoch JR, Lloyd CM (2010) Resolution of allergic airway inflammation and airway
559 hyperreactivity is mediated by IL-17-producing $\gamma\delta$ T cells. *Am J Respir*
560 *Crit Care Med* 182 (4):464-476. doi:10.1164/rccm.200911-1775OC

- 561 48. Koenecke C, Chennupati V, Schmitz S, Malissen B, Förster R, Prinz I (2009) In vivo
562 application of mAb directed against the gammadelta TCR does not deplete but
563 generates "invisible" gammadelta T cells. *Eur J Immunol* 39 (2):372-379.
564 doi:10.1002/eji.200838741
- 565 49. Schramm CM, Puddington L, Yiamouyiannis CA, Lingenheld EG, Whiteley HE,
566 Wolyniec WW, Noonan TC, Thrall RS (2000) Proinflammatory roles of T-cell receptor
567 (TCR)gammadelta and TCRalphabeta lymphocytes in a murine model of asthma. *Am*
568 *J Respir Cell Mol Biol* 22 (2):218-225. doi:10.1165/ajrcmb.22.2.3620
- 569 50. Huang Y, Jin N, Roark CL, Aydintug MK, Wands JM, Huang H, O'Brien RL, Born WK
570 (2009) The influence of IgE-enhancing and IgE-suppressive gammadelta T cells
571 changes with exposure to inhaled ovalbumin. *J Immunol* 183 (2):849-855.
572 doi:10.4049/jimmunol.0804104
- 573 51. Hahn YS, Taube C, Jin N, Sharp L, Wands JM, Aydintug MK, Lahn M, Huber SA, O'Brien
574 RL, Gelfand EW, Born WK (2004) Different potentials of gamma delta T cell subsets
575 in regulating airway responsiveness: V gamma 1+ cells, but not V gamma 4+ cells,
576 promote airway hyperreactivity, Th2 cytokines, and airway inflammation. *J*
577 *Immunol* 172 (5):2894-2902
- 578 52. Belkadi A, Dietrich C, Machavoine F, Victor JR, Leite-de-Moraes M (2018) $\gamma\delta$ T cells
579 amplify *Blomia tropicalis*-induced allergic airway disease. *Allergy*.
580 doi:10.1111/all.13618
- 581 53. Huang Y, Aydintug MK, Loomis J, Macleod MK, McKee AS, Kirchenbaum G, Jakubzick
582 CV, Kedl RM, Sun D, Jacobelli J, O'Brien RL, Born WK (2013) Antigen-specific
583 regulation of IgE antibodies by non-antigen-specific $\gamma\delta$ T cells. *J Immunol* 190
584 (3):913-921. doi:10.4049/jimmunol.1202230
- 585 54. de Oliveira MG, de Lima Lira AA, da Ressureição Sgnotto F, Inoue AHS, Santos LS,
586 Nakamatsu BY, Duarte AJDS, Leite-de-Moraes M, Victor JR (2019) Maternal IgG
587 impairs the maturation of offspring intrathymic IL-17-producing $\gamma\delta$ T cells:
588 implications for murine and human allergies. *Clin Exp Allergy*.
589 doi:10.1111/cea.13393
- 590 55. Krejsek J, Král B, Vokurková D, Derner V, Tousková M, Paráková Z, Kopecký O (1998)
591 Decreased peripheral blood gamma delta T cells in patients with bronchial asthma.
592 *Allergy* 53 (1):73-77. doi:10.1111/j.1398-9995.1998.tb03776.x
- 593 56. Zhao Y, Yang J, Gao YD (2011) Altered expressions of helper T cell (Th)1, Th2, and
594 Th17 cytokines in CD8(+) and $\gamma\delta$ T cells in patients with allergic asthma. *J Asthma* 48
595 (5):429-436. doi:10.3109/02770903.2011.570403
- 596 57. Spinozzi F, Agea E, Bistoni O, Forenza N, Monaco A, Bassotti G, Nicoletti I, Riccardi C,
597 Grignani F, Bertotto A (1996) Increased allergen-specific, steroid-sensitive gamma
598 delta T cells in bronchoalveolar lavage fluid from patients with asthma. *Ann Intern*
599 *Med* 124 (2):223-227
- 600 58. Urboniene D, Babusyte A, Lötvall J, Sakalauskas R, Sitkauskiene B (2013) Distribution
601 of $\gamma\delta$ and other T-lymphocyte subsets in patients with chronic obstructive
602 pulmonary disease and asthma. *Respir Med* 107 (3):413-423.
603 doi:10.1016/j.rmed.2012.11.012
- 604 59. Krug N, Erpenbeck VJ, Balke K, Petschallies J, Tschernig T, Hohlfeld JM, Fabel H (2001)
605 Cytokine profile of bronchoalveolar lavage-derived CD4(+), CD8(+), and
606 gammadelta T cells in people with asthma after segmental allergen challenge. *Am J*
607 *Respir Cell Mol Biol* 25 (1):125-131. doi:10.1165/ajrcmb.25.1.4194

- 608 60. Pellicci DG, Hammond KJ, Uldrich AP, Baxter AG, Smyth MJ, Godfrey DI (2002) A
609 natural killer T (NKT) cell developmental pathway involving a thymus-dependent
610 NK1.1(-)CD4(+) CD1d-dependent precursor stage. *J Exp Med* 195 (7):835-844
611 61. Benlagha K, Kyin T, Beavis A, Teyton L, Bendelac A (2002) A thymic precursor to the
612 NK T cell lineage. *Science* 296 (5567):553-555
613 62. Benlagha K, Wei DG, Veiga J, Teyton L, Bendelac A (2005) Characterization of the
614 early stages of thymic NKT cell development. *J Exp Med* 202 (4):485-492.
615 doi:10.1084/jem.20050456
616 63. Gapin L, Matsuda JL, Surh CD, Kronenberg M (2001) NKT cells derive from double-
617 positive thymocytes that are positively selected by CD1d. *Nat Immunol* 2 (10):971-
618 978. doi:10.1038/ni710
619 64. Shimamura M, Ohteki T, Beutner U, MacDonald HR (1997) Lack of directed V alpha
620 14-J alpha 281 rearrangements in NK1+ T cells. *Eur J Immunol* 27 (6):1576-1579.
621 doi:10.1002/eji.1830270638
622 65. Egawa T, Eberl G, Taniuchi I, Benlagha K, Geissmann F, Hennighausen L, Bendelac A,
623 Littman DR (2005) Genetic evidence supporting selection of the Valpha14i NKT cell
624 lineage from double-positive thymocyte precursors. *Immunity* 22 (6):705-716.
625 doi:10.1016/j.immuni.2005.03.011
626 66. Kronenberg M (2014) When less is more: T lymphocyte populations with restricted
627 antigen receptor diversity. *J Immunol* 193 (3):975-976.
628 doi:10.4049/jimmunol.1401491
629 67. Godfrey DI, Stankovic S, Baxter AG (2010) Raising the NKT cell family. *Nat Immunol*
630 11 (3):197-206. doi:10.1038/ni.1841
631 68. Ohteki T, MacDonald HR (1996) Stringent V beta requirement for the development
632 of NK1.1+ T cell receptor-alpha/beta+ cells in mouse liver. *J Exp Med* 183 (3):1277-
633 1282. doi:10.1084/jem.183.3.1277
634 69. Gumperz JE, Miyake S, Yamamura T, Brenner MB (2002) Functionally distinct subsets
635 of CD1d-restricted natural killer T cells revealed by CD1d tetramer staining. *J Exp*
636 *Med* 195 (5):625-636
637 70. Geissmann F, Cameron TO, Sidobre S, Manlongat N, Kronenberg M, Briskin MJ,
638 Dustin ML, Littman DR (2005) Intravascular immune surveillance by CXCR6+ NKT
639 cells patrolling liver sinusoids. *PLoS Biol* 3 (4):e113.
640 doi:10.1371/journal.pbio.0030113
641 71. Kawano T, Cui J, Koezuka Y, Taura I, Kaneko Y, Motoki K, Ueno H, Nakagawa R, Sato
642 H, Kondo E, Koseki H, Taniguchi M (1997) CD1d-restricted and TCR-mediated
643 activation of valpha14 NKT cells by glycosylceramides. *Science* 278 (5343):1626-
644 1629. doi:10.1126/science.278.5343.1626
645 72. Matsuda JL, Naidenko OV, Gapin L, Nakayama T, Taniguchi M, Wang CR, Koezuka Y,
646 Kronenberg M (2000) Tracking the response of natural killer T cells to a glycolipid
647 antigen using CD1d tetramers. *J Exp Med* 192 (5):741-754.
648 doi:10.1084/jem.192.5.741
649 73. Rossjohn J, Pellicci DG, Patel O, Gapin L, Godfrey DI (2012) Recognition of CD1d-
650 restricted antigens by natural killer T cells. *Nat Rev Immunol* 12 (12):845-857.
651 doi:10.1038/nri3328
652 74. Brennan PJ, Brigl M, Brenner MB (2013) Invariant natural killer T cells: an innate
653 activation scheme linked to diverse effector functions. *Nat Rev Immunol* 13 (2):101-
654 117. doi:10.1038/nri3369

- 655 75. Leite-de-Moraes MC, Dy M (1997) Natural killer T cells: a potent cytokine-producing
656 cell population. *Eur Cytokine Netw* 8 (3):229-237
- 657 76. Yoshimoto T, Paul WE (1994) CD4pos, NK1.1pos T cells promptly produce interleukin
658 4 in response to in vivo challenge with anti-CD3. *J Exp Med* 179 (4):1285-1295.
659 doi:10.1084/jem.179.4.1285
- 660 77. Bendelac A, Matzinger P, Seder RA, Paul WE, Schwartz RH (1992) Activation events
661 during thymic selection. *J Exp Med* 175 (3):731-742. doi:10.1084/jem.175.3.731
- 662 78. Arase H, Arase N, Nakagawa K, Good RA, Onoé K (1993) NK1.1+ CD4+ CD8-
663 thymocytes with specific lymphokine secretion. *Eur J Immunol* 23 (1):307-310.
664 doi:10.1002/eji.1830230151
- 665 79. Singh N, Hong S, Scherer DC, Serizawa I, Burdin N, Kronenberg M, Koezuka Y, Van
666 Kaer L (1999) Cutting edge: activation of NK T cells by CD1d and alpha-
667 galactosylceramide directs conventional T cells to the acquisition of a Th2
668 phenotype. *J Immunol* 163 (5):2373-2377
- 669 80. Brown DR, Fowell DJ, Corry DB, Wynn TA, Moskowitz NH, Cheever AW, Locksley RM,
670 Reiner SL (1996) Beta 2-microglobulin-dependent NK1.1+ T cells are not essential
671 for T helper cell 2 immune responses. *J Exp Med* 184 (4):1295-1304
- 672 81. Korsgren M, Persson CG, Sundler F, Bjerke T, Hansson T, Chambers BJ, Hong S, Van
673 Kaer L, Ljunggren HG, Korsgren O (1999) Natural killer cells determine development
674 of allergen-induced eosinophilic airway inflammation in mice. *J Exp Med* 189
675 (3):553-562
- 676 82. Zhang Y, Rogers KH, Lewis DB (1996) Beta 2-microglobulin-dependent T cells are
677 dispensable for allergen-induced T helper 2 responses. *J Exp Med* 184 (4):1507-1512
- 678 83. Das J, Eynott P, Jupp R, Bothwell A, Van Kaer L, Shi Y, Das G (2006) Natural killer T
679 cells and CD8+ T cells are dispensable for T cell-dependent allergic airway
680 inflammation. *Nat Med* 12 (12):1345-1346; author reply 1347.
681 doi:10.1038/nm1206-1345
- 682 84. McKnight CG, Morris SC, Perkins C, Zhu Z, Hildeman DA, Bendelac A, Finkelman FD
683 (2017) NKT cells contribute to basal IL-4 production but are not required to induce
684 experimental asthma. *PLoS One* 12 (11):e0188221.
685 doi:10.1371/journal.pone.0188221
- 686 85. Lisbonne M, Diem S, de Castro Keller A, Lefort J, Araujo LM, Hachem P, Fourneau JM,
687 Sidobre S, Kronenberg M, Taniguchi M, Van Endert P, Dy M, Askenase P, Russo M,
688 Vargaftig BB, Herbelin A, Leite-de-Moraes MC (2003) Cutting edge: invariant V alpha
689 14 NKT cells are required for allergen-induced airway inflammation and
690 hyperreactivity in an experimental asthma model. *J Immunol* 171 (4):1637-1641
- 691 86. Akbari O, Stock P, Meyer E, Kronenberg M, Sidobre S, Nakayama T, Taniguchi M,
692 Grusby MJ, DeKruyff RH, Umetsu DT (2003) Essential role of NKT cells producing IL-
693 4 and IL-13 in the development of allergen-induced airway hyperreactivity. *Nat Med*
694 9 (5):582-588. doi:10.1038/nm851
- 695 87. Olszak T, An D, Zeissig S, Vera MP, Richter J, Franke A, Glickman JN, Siebert R, Baron
696 RM, Kasper DL, Blumberg RS (2012) Microbial exposure during early life has
697 persistent effects on natural killer T cell function. *Science* 336 (6080):489-493.
698 doi:10.1126/science.1219328
- 699 88. An D, Oh SF, Olszak T, Neves JF, Avci FY, Erturk-Hasdemir D, Lu X, Zeissig S, Blumberg
700 RS, Kasper DL (2014) Sphingolipids from a symbiotic microbe regulate homeostasis

- 701 of host intestinal natural killer T cells. *Cell* 156 (1-2):123-133.
702 doi:10.1016/j.cell.2013.11.042
- 703 89. Remot A, Descamps D, Noordine ML, Boukadiri A, Mathieu E, Robert V, Riffault S,
704 Lambrecht B, Langella P, Hammad H, Thomas M (2017) Bacteria isolated from lung
705 modulate asthma susceptibility in mice. *ISME J* 11 (5):1061-1074.
706 doi:10.1038/ismej.2016.181
- 707 90. Chandra S, Zhao M, Budelsky A, de Mingo Pulido A, Day J, Fu Z, Siegel L, Smith D,
708 Kronenberg M (2015) A new mouse strain for the analysis of invariant NKT cell
709 function. *Nat Immunol* 16 (8):799-800. doi:10.1038/ni.3203
- 710 91. Tumes D, Hirahara K, Papadopoulos M, Shinoda K, Onodera A, Kumagai J, Yip KH,
711 Pant H, Kokubo K, Kiuchi M, Aoki A, Obata-Ninomiya K, Tokoyoda K, Endo Y, Kimura
712 MY, Nakayama T (2019) Ezh2 controls development of natural killer T cells, which
713 cause spontaneous asthma-like pathology. *J Allergy Clin Immunol* 144 (2):549-
714 560.e510. doi:10.1016/j.jaci.2019.02.024
- 715 92. Michel ML, Keller AC, Paget C, Fujio M, Trottein F, Savage PB, Wong CH, Schneider E,
716 Dy M, Leite-de-Moraes MC (2007) Identification of an IL-17-producing NK1.1(neg)
717 iNKT cell population involved in airway neutrophilia. *J Exp Med* 204 (5):995-1001.
718 doi:10.1084/jem.20061551
- 719 93. Pichavant M, Goya S, Meyer EH, Johnston RA, Kim HY, Matangkasombut P, Zhu M,
720 Iwakura Y, Savage PB, DeKruyff RH, Shore SA, Umetsu DT (2008) Ozone exposure in
721 a mouse model induces airway hyperreactivity that requires the presence of natural
722 killer T cells and IL-17. *J Exp Med* 205 (2):385-393. doi:10.1084/jem.20071507
- 723 94. Koh YI, Shim JU (2010) Association between sputum natural killer T cells and
724 eosinophilic airway inflammation in human asthma. *Int Arch Allergy Immunol* 153
725 (3):239-248. doi:10.1159/000314364
- 726 95. Shim JU, Koh YI (2014) Increased Th2-like Invariant Natural Killer T cells in Peripheral
727 Blood From Patients With Asthma. *Allergy Asthma Immunol Res* 6 (5):444-448.
728 doi:10.4168/aair.2014.6.5.444
- 729 96. Lezmi G, Abou Taam R, Dietrich C, Chatenoud L, de Blic J, Leite-de-Moraes M (2018)
730 Circulating IL-17-producing mucosal-associated invariant T cells (MAIT) are
731 associated with symptoms in children with asthma. *Clin Immunol* 188:7-11.
732 doi:10.1016/j.clim.2017.11.009
- 733 97. Chandra S, Wingender G, Greenbaum JA, Khurana A, Gholami AM, Ganesan AP,
734 Rosenbach M, Jaffee K, Gern JE, Wood R, O'Connor G, Sandel M, Kattan M, Bacharier
735 L, Togias A, Horner AA, Kronenberg M (2018) Development of Asthma in Inner-City
736 Children: Possible Roles of MAIT Cells and Variation in the Home Environment. *J*
737 *Immunol* 200 (6):1995-2003. doi:10.4049/jimmunol.1701525
- 738 98. Akbari O, Faul JL, Hoyte EG, Berry GJ, Wahlström J, Kronenberg M, DeKruyff RH,
739 Umetsu DT (2006) CD4+ invariant T-cell-receptor+ natural killer T cells in bronchial
740 asthma. *N Engl J Med* 354 (11):1117-1129. doi:10.1056/NEJMoa053614
- 741 99. Vijayanand P, Seumois G, Pickard C, Powell RM, Angco G, Sammut D, Gadola SD,
742 Friedmann PS, Djukanovic R (2007) Invariant natural killer T cells in asthma and
743 chronic obstructive pulmonary disease. *N Engl J Med* 356 (14):1410-1422.
744 doi:10.1056/NEJMoa064691
- 745 100. Thomas SY, Lilly CM, Luster AD (2006) Invariant natural killer T cells in bronchial
746 asthma. *N Engl J Med* 354 (24):2613-2616; author reply 2613-2616.
747 doi:10.1056/NEJMc066189

- 748 101. Matangkasombut P, Marigowda G, Ervine A, Idris L, Pichavant M, Kim HY, Yasumi
749 T, Wilson SB, DeKruyff RH, Faul JL, Israel E, Akbari O, Umetsu DT (2009) Natural killer
750 T cells in the lungs of patients with asthma. *J Allergy Clin Immunol* 123 (5):1181-
751 1185. doi:10.1016/j.jaci.2009.02.013
- 752 102. Pham-Thi N, de Blic J, Le Bourgeois M, Dy M, Scheinmann P, Leite-de-Moraes MC
753 (2006) Enhanced frequency of immunoregulatory invariant natural killer T cells in
754 the airways of children with asthma. *J Allergy Clin Immunol* 117 (1):217-218.
755 doi:10.1016/j.jaci.2005.09.052
- 756 103. Kjer-Nielsen L, Patel O, Corbett AJ, Le Nours J, Meehan B, Liu L, Bhati M, Chen Z,
757 Kostenko L, Reantragoon R, Williamson NA, Purcell AW, Dudek NL, McConville MJ,
758 O'Hair RA, Khairallah GN, Godfrey DI, Fairlie DP, Rossjohn J, McCluskey J (2012) MR1
759 presents microbial vitamin B metabolites to MAIT cells. *Nature* 491 (7426):717-723.
760 doi:10.1038/nature11605
- 761 104. Kjer-Nielsen L, Corbett AJ, Chen Z, Liu L, Mak JY, Godfrey DI, Rossjohn J, Fairlie DP,
762 McCluskey J, Eckle SB (2018) An overview on the identification of MAIT cell antigens.
763 *Immunol Cell Biol* 96 (6):573-587. doi:10.1111/imcb.12057
- 764 105. Koay HF, Gherardin NA, Enders A, Loh L, Mackay LK, Almeida CF, Russ BE, Nold-
765 Petry CA, Nold MF, Bedoui S, Chen Z, Corbett AJ, Eckle SB, Meehan B, d'Udekem Y,
766 Konstantinov IE, Lappas M, Liu L, Goodnow CC, Fairlie DP, Rossjohn J, Chong MM,
767 Kedzierska K, Berzins SP, Belz GT, McCluskey J, Uldrich AP, Godfrey DI, Pellicci DG
768 (2016) A three-stage intrathymic development pathway for the mucosal-associated
769 invariant T cell lineage. *Nat Immunol* 17 (11):1300-1311. doi:10.1038/ni.3565
- 770 106. Lee OJ, Cho YN, Kee SJ, Kim MJ, Jin HM, Lee SJ, Park KJ, Kim TJ, Lee SS, Kwon YS, Kim
771 N, Shin MG, Shin JH, Suh SP, Ryang DW, Park YW (2014) Circulating mucosal-
772 associated invariant T cell levels and their cytokine levels in healthy adults. *Exp*
773 *Gerontol* 49:47-54. doi:10.1016/j.exger.2013.11.003
- 774 107. Lepore M, Kalinichenko A, Colone A, Paleja B, Singhal A, Tschumi A, Lee B, Poidinger
775 M, Zolezzi F, Quagliata L, Sander P, Newell E, Bertolotti A, Terracciano L, De Libero
776 G, Mori L (2014) Parallel T-cell cloning and deep sequencing of human MAIT cells
777 reveal stable oligoclonal TCRbeta repertoire. *Nat Commun* 5:3866.
778 doi:10.1038/ncomms4866
- 779 108. Kurioka A, Jahun AS, Hannaway RF, Walker LJ, Fergusson JR, Sverremark-Ekstrom
780 E, Corbett AJ, Ussher JE, Willberg CB, Klenerman P (2017) Shared and Distinct
781 Phenotypes and Functions of Human CD161⁺⁺ Valpha7.2⁺ T Cell Subsets. *Front*
782 *Immunol* 8:1031. doi:10.3389/fimmu.2017.01031
- 783 109. Rahimpour A, Koay HF, Enders A, Clanchy R, Eckle SB, Meehan B, Chen Z, Whittle B,
784 Liu L, Fairlie DP, Goodnow CC, McCluskey J, Rossjohn J, Uldrich AP, Pellicci DG,
785 Godfrey DI (2015) Identification of phenotypically and functionally heterogeneous
786 mouse mucosal-associated invariant T cells using MR1 tetramers. *J Exp Med* 212
787 (7):1095-1108. doi:10.1084/jem.20142110
- 788 110. Lantz O, Legoux F (2019) MAIT cells: programmed in the thymus to mediate
789 immunity within tissues. *Curr Opin Immunol* 58:75-82.
790 doi:10.1016/j.coi.2019.04.016
- 791 111. Legoux F, Gilet J, Procopio E, Echasserieau K, Bernardeau K, Lantz O (2019)
792 Molecular mechanisms of lineage decisions in metabolite-specific T cells. *Nat*
793 *Immunol* 20 (9):1244-1255. doi:10.1038/s41590-019-0465-3

- 794 112. Legoux F, Bellet D, Daviaud C, El Morr Y, Darbois A, Niort K, Procopio E, Salou M,
795 Gilet J, Ryffel B, Balvay A, Foussier A, Sarkis M, El Marjou A, Schmidt F, Rabot S, Lantz
796 O (2019) Microbial metabolites control the thymic development of mucosal-
797 associated invariant T cells. *Science*. doi:10.1126/science.aaw2719
- 798 113. Martin E, Treiner E, Duban L, Guerri L, Laude H, Toly C, Premel V, Devys A, Moura
799 IC, Tilloy F, Cherif S, Vera G, Latour S, Soudais C, Lantz O (2009) Stepwise
800 development of MAIT cells in mouse and human. *PLoS Biol* 7 (3):e54.
801 doi:10.1371/journal.pbio.1000054
- 802 114. Li C, Lu Z, Bi K, Wang K, Xu Y, Guo P, Chen Y, Zhou P, Wei Z, Jiang H, Cao Y (2019)
803 CD4. *Reprod Biol Endocrinol* 17 (1):78. doi:10.1186/s12958-019-0524-5
- 804 115. Dusseaux M, Martin E, Serriari N, Peguillet I, Premel V, Louis D, Milder M, Le Bourhis
805 L, Soudais C, Treiner E, Lantz O (2011) Human MAIT cells are xenobiotic-resistant,
806 tissue-targeted, CD161hi IL-17-secreting T cells. *Blood* 117 (4):1250-1259.
807 doi:10.1182/blood-2010-08-303339
- 808 116. Chen Z, Wang H, D'Souza C, Sun S, Kostenko L, Eckle SB, Meehan BS, Jackson DC,
809 Strugnell RA, Cao H, Wang N, Fairlie DP, Liu L, Godfrey DI, Rossjohn J, McCluskey J,
810 Corbett AJ (2017) Mucosal-associated invariant T-cell activation and accumulation
811 after in vivo infection depends on microbial riboflavin synthesis and co-stimulatory
812 signals. *Mucosal Immunol* 10 (1):58-68. doi:10.1038/mi.2016.39
- 813 117. Ussher JE, Bilton M, Attwod E, Shadwell J, Richardson R, de Lara C, Mettke E,
814 Kurioka A, Hansen TH, Klenerman P, Willberg CB (2014) CD161⁺⁺ CD8⁺ T cells,
815 including the MAIT cell subset, are specifically activated by IL-12+IL-18 in a TCR-
816 independent manner. *Eur J Immunol* 44 (1):195-203. doi:10.1002/eji.201343509
- 817 118. van Wilgenburg B, Scherwitzl I, Hutchinson EC, Leng T, Kurioka A, Kulicke C, de Lara
818 C, Cole S, Vasanawathana S, Limpitikul W, Malasit P, Young D, Denney L, Moore MD,
819 Fabris P, Giordani MT, Oo YH, Laidlaw SM, Dustin LB, Ho LP, Thompson FM,
820 Ramamurthy N, Mongkolsapaya J, Willberg CB, Screaton GR, Klenerman P,
821 consortium S-H (2016) MAIT cells are activated during human viral infections. *Nat*
822 *Commun* 7:11653. doi:10.1038/ncomms11653
- 823 119. Sattler A, Dang-Heine C, Reinke P, Babel N (2015) IL-15 dependent induction of IL-
824 18 secretion as a feedback mechanism controlling human MAIT-cell effector
825 functions. *Eur J Immunol* 45 (8):2286-2298. doi:10.1002/eji.201445313
- 826 120. Salou M, Franciszkiewicz K, Lantz O (2017) MAIT cells in infectious diseases. *Curr*
827 *Opin Immunol* 48:7-14. doi:10.1016/j.coi.2017.07.009
- 828 121. Sakala IG, Kjer-Nielsen L, Eickhoff CS, Wang X, Blazevic A, Liu L, Fairlie DP, Rossjohn
829 J, McCluskey J, Fremont DH, Hansen TH, Hoft DF (2015) Functional Heterogeneity
830 and Antimycobacterial Effects of Mouse Mucosal-Associated Invariant T Cells
831 Specific for Riboflavin Metabolites. *J Immunol* 195 (2):587-601.
832 doi:10.4049/jimmunol.1402545
- 833 122. Wong EB, Gold MC, Meermeier EW, Xulu BZ, Khuzwayo S, Sullivan ZA, Mahyari E,
834 Rogers Z, Kløverpris H, Sharma PK, Worley AH, Lalloo U, Baijnath P, Ambaram A,
835 Naidoo L, Suleman M, Madansein R, McLaren JE, Ladell K, Miners KL, Price DA, Behar
836 SM, Nielsen M, Kasprovicz VO, Leslie A, Bishai WR, Ndung'u T, Lewinsohn DM
837 (2019) TRAV1-2 + CD8 + T-cells including oligoclonal expansions of MAIT cells are
838 enriched in the airways in human tuberculosis. *Commun Biol* 2:203.
839 doi:10.1038/s42003-019-0442-2

- 840 123. Malka-Ruimy C, Ben Youssef G, Lambert M, Turret M, Ghazarian L, Faye A, Caillat-
841 Zucman S, Houdouin V (2019) Mucosal-Associated Invariant T Cell Levels Are
842 Reduced in the Peripheral Blood and Lungs of Children With Active Pulmonary
843 Tuberculosis. *Front Immunol* 10:206. doi:10.3389/fimmu.2019.00206
- 844 124. Hinks TS, Zhou X, Staples KJ, Dimitrov BD, Manta A, Petrossian T, Lum PY, Smith CG,
845 Ward JA, Howarth PH, Walls AF, Gadola SD, Djukanovic R (2015) Innate and adaptive
846 T cells in asthmatic patients: Relationship to severity and disease mechanisms. *J*
847 *Allergy Clin Immunol* 136 (2):323-333. doi:10.1016/j.jaci.2015.01.014
- 848 125. Lezmi G, Abou-Taam R, Garcelon N, Dietrich C, Machavoine F, Delacourt C, Adel-
849 Patient K, Leite-de-Moraes M (2019) Evidence for a MAIT-17-high phenotype in
850 children with severe asthma. *J Allergy Clin Immunol*. doi:10.1016/j.jaci.2019.08.003
851
852

853 **Figure Captations**

854

855 **Figure 1: $\gamma\delta$ T cell immune response in asthma.** In experimental asthma models (left) IL-
856 4-producing $\gamma\delta$ T cells could enhance Th2 inflammation, IgE production, airway
857 eosinophilia and airway hyperreactivity (AHR), while IL-17-producing $\gamma\delta$ T17 cells could
858 have an opposite effect. In BALF (bronchoalveolar lavage fluid) from asthmatic patients
859 (right), the frequency of $\gamma\delta$ T cells was elevated. These $\gamma\delta$ T cells improved their ability to
860 secrete IL-5 and IL-13 following allergen challenge. The main effects resulting from the
861 activation of $\gamma\delta$ T are listed within the lung figure.

862 \uparrow = induction/augmentation; \downarrow = reduction/inhibition.

863

864 **Figure 2: iNKT cell immune response in asthma.** In experimental asthma models (left)
865 IL-4-and IL-13-producing iNKT2 cells will enhance IgE production, airway eosinophilia
866 and airway hyperreactivity (AHR), while IL-17-producing iNKT17 cells could promote
867 airway neutrophilia and AHR. The frequency of iNKT cells in BALF (bronchoalveolar
868 lavage fluid) from asthmatic patients (right) rang from 0.1% to 10%. The main effects
869 resulting from the activation of iNKT cells are listed within the lung figure.

870 \uparrow = induction/augmentation; ? = no functional effects described.

871

872 **Figure 3: MAIT cell immune response in asthma.** The studies concerning the implication
873 of MAIT cells in experimental asthma models (left) are currently in progress. IL-17-
874 producing MAIT cells are associated with severe exacerbations in asthmatic patients
875 (right) while their IFN γ -producing counterpart maybe involved in protection.

876 ? = no functional effects described.

877

878




Figure 1

879
880




Figure 2

881
882




Figure 3

883
884

Table 1. Major features of $\gamma\delta$ T, iNKT and MAIT cells.

	$\gamma\delta$ T		iNKT			MAIT	
TCR	<ul style="list-style-type: none"> • Mice: Vγ1, Vγ4, Vγ5, Vγ7 (TRGV1, 4, 5, 7). • Humans: Vδ1-3 (TRDV1-3), Vγ2-5, Vγ8, Vγ9, Vγ11 (TRDV2-5, 8, 9, 11). 		<ul style="list-style-type: none"> • Mice: Vα14-Jα18 (TRAV11-TRAJ18). • Humans: Vα24-Jα18 (TRAV10-TRAJ18). 			<ul style="list-style-type: none"> • Mice: Vα19-Jα33 (TRAV1-2-TRAJ33). • Humans: Vα7.2-Jα33 (TRAV1-2-TRAJ33). 	
Antigen-presenting molecules	BTN3A (as an intermediate)		CD1d			MR1	
Antigens	stress molecules (MICA and MICB), heat-shock proteins or synthetic phosphoantigens (pAg)		glycolipids (α -GalCer, iGb3, others)			Riboflavin derivatives (5-OP-RU)	
Major subsets	$\gamma\delta$ T IFN- γ ⁺	$\gamma\delta$ T17	iNKT1	iNKT2	iNKT17	MAIT1	MAIT17
Transcription factors	Tbet	RORC	Tbet, GATA-3	GATA-3	RORC	Tbet	RORC
Major cytokines produced	IFN- γ	IL-17	IFN- γ , IL-4	IL-4, IL-13	IL-17	IFN- γ	IL-17