



HAL
open science

Local null controllability of the penalized Boussinesq system with a reduced number of controls

Jon Asier Bárcena-Petisco, Kévin Le Balc'h

► **To cite this version:**

Jon Asier Bárcena-Petisco, Kévin Le Balc'h. Local null controllability of the penalized Boussinesq system with a reduced number of controls. 2020. hal-02913358v1

HAL Id: hal-02913358

<https://hal.science/hal-02913358v1>

Preprint submitted on 8 Aug 2020 (v1), last revised 7 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Local null controllability of the penalized Boussinesq system with a reduced number of controls

Jon Asier Bárcena-Petisco* Kévin Le Balch[†]

August 7, 2020

Abstract: In this paper we consider the Boussinesq system with homogeneous Dirichlet boundary conditions and defined in a regular domain $\Omega \subset \mathbb{R}^N$ for $N = 2$ and $N = 3$. The incompressibility condition of the fluid is replaced by its approximation by penalization with a small parameter $\varepsilon > 0$. We prove that our system is locally null controllable using a control with a restricted number of components, defined in an open set ω contained in Ω and whose cost is bounded uniformly when $\varepsilon \rightarrow 0$. The proof is based on a linearization argument and the null-controllability of the linearized system is obtained by proving a new Carleman estimate for the adjoint system. This observability inequality is obtained thanks to the coercivity of some second order differential operator involving crossed derivatives.

Key words: Carleman inequality, controllability, nonlinear system, penalized Stokes system

AMS subject classification: 35K40, 93B05, 93C20

Abbreviated title: Null controllability of the penalized Boussinesq system.

Acknowledgements: The first author was supported by grants from Région Ile-de-France. The second author was supported by the SysNum cluster of excellence University of Bordeaux.

*Sorbonne Université, Laboratoire Jacques-Louis Lions, F-75005, Paris, France, barcena@ljl11.math.upmc.fr

[†]Institut de Mathématiques de Bordeaux, 351 Cours de la Libération, 33400 Bordeaux, France, kevin.le-balch@math.u-bordeaux.fr

1 Introduction

1.1 Presentation of the system

For a given time $T > 0$, Ω a sufficiently smooth bounded open set of \mathbb{R}^N ($N = 2, 3$) and ω a nonempty open set contained in Ω , we consider the controlled penalized Boussinesq system, which is given by:

$$\begin{cases} y_t - \Delta y + (y \cdot \nabla)y + \frac{1}{2}(\nabla \cdot y)y + \nabla p = \theta e_N + \tilde{v}1_\omega & \text{in } Q, \\ \theta_t - \Delta \theta + y \cdot \nabla \theta + \frac{1}{2}(\nabla \cdot y)\theta = v_{N+1}1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, \theta = 0 & \text{on } \Sigma, \\ y(0) = y^0, \theta(0) = \theta^0 & \text{in } \Omega, \end{cases} \quad (1.1)$$

where e_N stands for the N -th vector of the canonical basis of \mathbb{R}^N , $Q := (0, T) \times \Omega$, $\Sigma := (0, T) \times \partial\Omega$. In the controlled system (1.1), $y = y(t, x)$ represents the velocity of the particles of the fluid, $\theta = \theta(t, x)$ their temperature and $v = (\tilde{v}, v_{N+1}) = (v_1, \dots, v_N, v_{N+1})(t, x)$ stands for the control which acts on ω .

System (1.1) approximates the classical Boussinesq system, which is the same as (1.1) with $\varepsilon = 0$ (which implies that $\frac{1}{2}(\nabla \cdot y)y = 0$ and $\frac{1}{2}(\nabla \cdot y)\theta = 0$). The classical Boussinesq system has been introduced by Joseph Boussinesq in 1877 and it models an incompressible fluid subjected to small variations of temperature. In addition, this way of approximating the incompressibility condition is called the penalty method and was introduced in [Tem68] in the Navier-Stokes case. As explained for instance in the survey [She97] and in the papers [Ber78] and [OJ84], this approximation procedure is widely used for numerical purpose.

We are going to focus on the small-time null-controllability of the system (1.1) with a reduced number of controls. In particular, we seek controls v that satisfy:

$$\tilde{v} = 0, \text{ if } N = 2, \quad v_1 = v_3 = 0, \text{ if } N = 3. \quad (1.2)$$

This choice matches with what is known about the controllability of the classical Boussinesq system when the control satisfies (1.2) (see [Car12]). From a modelling point of view in the two-dimensional case, we want to control both the velocity and the temperature of the fluid by acting only in the temperature equation. Similarly, in the three-dimensional case we want to control the velocity by acting on both the temperature and the velocity, but with a scalar force acting on the second component of the velocity. One of the main interest in studying the controllability of (1.1) is that the controls of (1.1) converge to the controls of the Boussinesq system as ε goes to 0. Thus, in order

to compute numerically the controls of the Boussinesq system, it suffices to compute the controls of (1.1) for $\varepsilon > 0$ small enough.

1.2 Main result

In this part, we precise mathematically the main result of the paper. For this purpose, we introduce three different geometrical assumptions on Ω and ω .

Hypothesis 1.1. For $\Omega \subset \mathbb{R}^2$, let $\sigma_1, \dots, \sigma_k$ be the arc-length parametrizations of the different connected components of $\partial\Omega$. We assume that for any $i \in \{1, \dots, k\}$ and for any $s \in [0, 1]$ such that $(\sigma_1^i)'(s) = 0$ or $(\sigma_2^i)'(s) = 0$, we have that $\kappa^i(s) \neq 0$ where $\kappa^i(s)$ is the curvature of $\partial\Omega$ at the point $\sigma^i(s)$.

Hypothesis 1.2. For $\Omega \subset \mathbb{R}^3$, we assume that there are $\Omega_0 \subset \mathbb{R}^2$ satisfying Hypothesis 1.1, $\delta > 0$ and $H_-, H_+ \in C(\overline{\Omega_0}; [\delta, +\infty))$ such that:

$$\Omega = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : (x_1, x_2) \in \Omega_0 \text{ and } -H_-(x_1, x_2) < x_3 < H_+(x_1, x_2)\}.$$

Hypothesis 1.3. For $\Omega \subset \mathbb{R}^3$, we assume that there are an interval $I \subset \mathbb{R}$ and a curve $\mathfrak{C} \subset \mathbb{R}^2$ such that $\Gamma := I \times \mathfrak{C}$ is a relative non-trivial open set of $\partial\Omega$. In addition, we consider a control domain $\omega \subset \Omega$ such that $\Gamma \cap \partial\omega$ contains a relatively non-trivial open set.

Example 1.1. We recall that, as proved in [BP20a], Hypothesis 1.1 is satisfied by Ω any strictly convex smooth domain and, if we allow rotations, by any regular domain. Moreover, we can easily construct regular domains Ω satisfying Hypothesis 1.2, by considering a cylinder and some cupolas. Finally, Hypothesis 1.3 is satisfied by any smooth domain Ω containing a cylindrical part on its boundary.

We now present the main result of the paper:

Theorem 1.2. *Let Ω, ω such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exists $\varepsilon_0 > 0$ such that for every time $T > 0$ there exist $\delta_T > 0$ and $C_T > 0$ such that for every $\varepsilon \in (0, \varepsilon_0)$ and every initial data $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$ satisfying $\|(y^0, \theta^0)\|_{L^2(\Omega)} \leq \delta_T$, there exist a control $v \in L^2(Q_\omega)^{N+1}$ satisfying (1.2),*

$$\|v\|_{L^2(Q_\omega)} \leq C_T, \tag{1.3}$$

and a weak solution (y, θ) of (1.1) verifying

$$(y, \theta)(T) = 0.$$

Here and in the whole paper we shorten the notation and denote $\|\cdot\|_V$ the norm $\|\cdot\|_{V^k}$ for V any Banach space and $k \in \mathbb{N}_*$. Indeed, the value of k is always easily deducible from the context.

In order to prove Theorem 1.2 we use the classical approach of proving first the controllability of the linearized system and then using a fixed-point theorem. In addition, we prove the controllability of the linearized system by proving a Carleman inequality for the homogeneous adjoint system and then using the approach of [LTT13]. The most difficult and original part is to prove the Carleman inequality and in particular to prove the coercivity of some 2nd order differential operator involving crossed derivatives. Before continuing, we make some comments on Theorem 1.2, its proof and related bibliography:

- For the definition of a weak solution of (1.1), one can adapt [Tem68, Section I.1.]. Remark that the existence of a weak solution is guaranteed by an adaptation of [Tem68, Théorème I.2.] but the uniqueness is only valid in the 2D case.
- Theorem 1.2 is a null-controllability result, uniform with respect to the parameter $\varepsilon > 0$ because of the estimate (1.3). So, by letting $\varepsilon \rightarrow 0$ in (1.1), we recover the results from [Car12] for less regular initial data but for more restrictive assumptions on Ω, ω .
- We use the geometric hypothesis for proving the coercivity of a 2nd order differential operator involving the crossed derivatives. For $\Omega \subset \mathbb{R}^2$, the assumption we make on the geometry of Ω , i.e. Hypothesis 1.1, has already been introduced in the recent paper [BP20a], dealing with the null-controllability of the linear penalized 2-D Stokes system. This is crucial for proving the null-controllability of such a system through one scalar control. Indeed, as shown in [Zua96, Theorem 1.2] and [BP20a, Section 2.1] some geometric hypothesis is needed because in the rhombus the linearised system around 0 cannot be controlled due to the fact that some eigenfunctions of the elliptic operator are not observable.
- At the heuristic level, it seems natural to obtain the controllability of the whole Boussinesq system (1.1) by acting only with $N - 1$ scalar controls. Indeed, v_{N+1} directly controls the component θ by the last equation of (1.1), then θ acts as an indirect control to control y_N in the N -th equation of (1.1)₁. In addition, if $N = 2$ the penalized divergence condition implies that for ε small enough y_2 acts as an indirect control to control y_1 . Similarly, if $N = 3$, y_1 is directly controlled by v_1 and y_2 is indirectly controlled by y_1 and y_3 by the penalized divergence condition for ε small enough.
- To prove the Carleman inequality, with the objective of highlighting the main ideas, we prioritize giving a clear proof, even at the expense of not getting the most optimal results. With that spirit, some recurrent operations are stated as lemmas (see Lemmas 2.5 and 2.7 below). These technical results can be useful for proving other Carleman inequalities in a different context.

- The results presented in Theorem 1.2 are original. Indeed, control problems with an approximation by penalization have first been studied in [IPY09, Section 4], where the penalized Stokes system was studied but without restriction on the control v . Next, the null controllability of the penalized Navier-Stokes system has been studied in [Bad11], but again without restriction on the control \tilde{v} . Finally, the null-controllability of the penalized Stokes system in the 2D-case with a scalar control has been established in [BP20a]. So, Theorem 1.2 is the first null-controllability result for a penalized system with a reduced number of controls in the nonlinear setting, see Section 4 for a similar result in the Navier-Stokes case. Moreover, both the linear and nonlinear results in the 3D-case in this paper are new.
- Considering other systems, the study of controllability problems in which the control has a reduced number of components has been an active topic of research recently. In particular, for the Stokes and Navier-Stokes systems we can consult for instance the following documents: [LZ96, FCGIP06, Gue07, CG09, CnG13, CL14, CnGG15, GM18, BP20b]. For more results on the controllability of linear parabolic systems with a reduced number of controls, see the survey [AKBGBdT11] and the references therein.

The rest of the paper is organized as follows: in Section 2 we prove the null controllability of the linearized system, Section 3 is devoted to the proof of the local null controllability of (1.1) i.e. we prove Theorem 1.2, in Section 4 we make some remarks and present some open problems.

2 Null-controllability of the linearized system

In this section we prove the null controllability of the linearized system of (1.1). We divide the proof as follows.

- In Section 2.1 we linearize the system (1.1) around 0, and we recall the equivalence between the null-controllability of the linearized system and the observability of the corresponding adjoint system.
- In Section 2.2 we recall some previous results about parabolic systems, elliptic systems and Carleman inequalities.
- In Sections 2.3 and 2.4 we obtain some Carleman inequalities for the adjoint system, for the 2D case and the 3D case respectively.
- In Section 2.5 we use the source term method to get the null-controllability of the linearized system and a source term, exponentially decreasing at $t = T$. This method has recently

been used for many other control systems (see, for instance, [DL19, MT18, FCLdM16, BM20, Tak17, LB18, HSLB19, GZ19]).

We also introduce the notations $Q_\omega := (0, T) \times \omega$, δ_{ij} is a constant that is 1 if $i = j$ and 0 otherwise, $f = (\tilde{f}, f_{N+1}) = (f_1, \dots, f_N, f_{N+1})$ denotes the source term, $H^{k_1, k_2}(Q) := H^{k_1}(0, T; L^2(\Omega)) \cap L^2(0, T; H^{k_2}(\Omega))$ and $H^{k_1, k_2}(\Sigma) := H^{k_1}(0, T; L^2(\partial\Omega)) \cap L^2(0, T; H^{k_2}(\partial\Omega))$ (for $k_1, k_2 \in \mathbb{R}^+$).

2.1 Linearization around 0, null-controllability results and observability estimates

When we linearize (1.1) around 0, we obtain:

$$\begin{cases} y_t - \Delta y + \nabla p = \tilde{f} + \theta e_N + \tilde{v}1_\omega & \text{in } Q, \\ \theta_t - \Delta \theta = f_{N+1} + v_{N+1}1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, \theta = 0 & \text{on } \Sigma, \\ y(0) = y^0, \theta(0) = \theta^0 & \text{in } \Omega, \end{cases} \quad (2.1)$$

for $f \in L^2(Q)^{N+1}$ decaying sufficiently fast (to be determined later on) at $t = T$. The goal is to obtain the following result:

Proposition 2.1. *Let Ω, ω such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist $\varepsilon_0 > 0$, $m \geq 1$ and $C > 0$ such that for every $T > 0$, $\varepsilon \in (0, \varepsilon_0)$ and $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$, there exists a control $v \in L^2(Q_\omega)^{N+1}$ satisfying (1.2), such that*

$$\|v\|_{L^2(Q_\omega)} \leq K(T) \|(y^0, \theta^0)\|_{L^2(\Omega)},$$

and

$$(y, \theta)(T) = 0,$$

for

$$K(T) := C \exp\left(\frac{C}{T^m}\right) \quad (2.2)$$

and for (y, θ) the solution of (2.1) with $f = 0$.

In order to prove Proposition 2.1, by the Hilbert Uniqueness Method (see, for instance, [Rus78],

[Lio88] and [Cor07]), it is equivalent to establish an observability estimate for the adjoint system:

$$\begin{cases} -\varphi_t - \Delta\varphi + \nabla\pi = 0 & \text{in } Q, \\ -\psi_t - \Delta\psi = \varphi_N & \text{in } Q, \\ \nabla \cdot \varphi = -\varepsilon\pi & \text{in } Q, \\ \varphi = 0, \psi = 0 & \text{on } \Sigma, \\ \varphi(T) = \varphi^T, \psi(T) = \psi^T & \text{in } \Omega, \end{cases} \quad (2.3)$$

for $(\varphi^T, \psi^T) \in L^2(\Omega)^{N+1}$.

Proposition 2.2. *Let Ω, ω such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist $\varepsilon_0 > 0$, $m \geq 1$ and $C > 0$ such that for every $T > 0$, $\varepsilon \in (0, \varepsilon_0)$ and $(\varphi^T, \psi^T) \in L^2(\Omega)^{N+1}$ the solution (φ, ψ) of (2.3) satisfies:*

$$\|\varphi(0, \cdot)\|_{L^2(\Omega)}^2 + \|\psi(0, \cdot)\|_{L^2(\Omega)}^2 \leq K(T) \iint_{Q_\omega} \delta_{3,N} |\varphi_1(t, x)|^2 + |\psi(t, x)|^2 dt dx, \quad (2.4)$$

with $K(T)$ as in (2.2).

To obtain the observability estimate (2.4), we will use Carleman estimates. More precisely, Proposition 2.2 will be a direct consequence of the Propositions 2.8, 2.10, 2.12, see below, and a classical dissipation argument.

2.2 Toolbox of elliptic, parabolic estimates and Carleman estimates

In this part, we recall elliptic and parabolic regularity estimates. We also present the Carleman estimates that will be useful in the sequel.

2.2.1 A parabolic and a elliptic result

Let us recall the regularity results of the Cauchy problem of the penalized Stokes system, which is given by the equations:

$$\begin{cases} u_t - \Delta u + \nabla q = \tilde{f} & \text{in } Q, \\ \varepsilon q + \nabla \cdot u = 0 & \text{in } Q, \\ u = 0 & \text{on } \Sigma, \\ u(0) = u^0 & \text{in } \Omega. \end{cases} \quad (2.5)$$

We have the following regularity estimate:

Lemma 2.3. *Let $k \in \mathbb{N}_*$ and $\Omega \subset \mathbb{R}^N$. Then, there are $\varepsilon_0 > 0$ and $C > 0$ such that if $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $u^0 \in H^{2k-1}(\Omega)^N$ and $\tilde{f} \in H^{k-1, 2k-2}(Q)^N$ such that (u^0, \tilde{f}) satisfy all the compatibility conditions that make sense, we have that the solution (u, q) of (2.5) belongs to $H^{k, 2k}(Q)^N \times H^{k-1, 2k-1}(Q)$ with the estimate:*

$$\|u\|_{H^{k, 2k}(Q)} + \|q\|_{H^{k-1, 2k-1}(Q)} \leq C \left(\|\tilde{f}\|_{H^{k-1, 2k-2}(Q)} + \|u^0\|_{H^{2k-1}(\Omega)} \right). \quad (2.6)$$

As explained in [BP20a, Lemma 2.5] (though it was done for the specific case $u^0 = 0$), the proof of Lemma 2.3 is mainly by induction. The initial case ($i = 1$) can be proved again by the Galerkin method. As for the inductive case, we get the regularity in the time variable by considering that u_t is a solution of (2.5) with (\tilde{f}, u^0) replaced by $(\tilde{f}_t, \Delta u^0 + \varepsilon^{-1} \nabla(\nabla \cdot u^0) + \tilde{f}(0, \cdot))$ and using again the Galerkin method. Moreover, we get the regularity in space by using the estimate for the steady Stokes problem given in [Tem77, Proposition I.2.2].

As for elliptic results, we recall the following result that is proved in [BP20a, Theorem 1.8] (considering the symmetry between the first and second variable):

Proposition 2.4. *Let Ω be a domain such that Hypothesis 1.1 holds. Then, for $a_0 > 0$ small enough, there is $C > 0$ such that for any function $w \in H^4(\Omega) \cap H_0^1(\Omega)$ and for any $a \in [0, a_0]$ we have that:*

$$\|\partial_{x_2} w\|_{C^0(\bar{\Omega})} \leq C (\|\partial_{x_1 x_2} w\|_{H^2(\Omega)} + \|L_a w\|_{H^1(\partial\Omega)}), \quad (2.7)$$

with:

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w. \quad (2.8)$$

Roughly, Proposition 2.4 states that the crossed derivative with some information on the second derivative on the boundary is coercive in $H^4(\Omega) \cap H_0^1(\Omega)$.

2.2.2 Classical Carleman estimates

We consider the following weights defined in Q :

$$\begin{aligned} \alpha(t, x) &:= \frac{e^{2\lambda\|\eta\|_\infty} - e^{\lambda\eta(x)}}{t^m(T-t)^m}, & \xi(t, x) &:= \frac{e^{\lambda\eta(x)}}{t^m(T-t)^m}, \\ \alpha^*(t) &:= \sup_{x \in (0, L)} \alpha(t, x), & \xi^*(t) &:= \inf_{x \in (0, L)} \xi(t, x), \end{aligned} \quad (2.9)$$

for $m \in \mathbb{R}^+$, $\lambda > 0$ to be fixed later and for η a C^2 function satisfying:

$$\eta = 0 \text{ on } \partial\Omega, \quad \eta > 0 \text{ in } \Omega, \quad \inf_{\Omega \setminus \omega_0} |\nabla \eta| > 0 \quad (2.10)$$

for some $\omega_0 \subset\subset \omega$. The existence of such a function η is proved in [FI96]. In fact, the weights (2.9) are also taken from [FI96]. We recall that from (2.9) and (2.10) we obtain on Σ that:

$$\alpha(t, x) = \alpha^*(t), \quad \xi(t, x) = \xi^*(t). \quad (2.11)$$

We also recall that for all $m > 0$ and $\delta > 0$ there is $C > 0$ such that for all $s \geq 0$:

$$s(\xi + \xi^* + \alpha + \alpha^*) \leq Ce^{s\delta\alpha}. \quad (2.12)$$

In addition, for all $\delta > 0$ there is $C > 0$ such that if $\lambda \geq C$ we have that:

$$\alpha^* \leq (1 + \delta)\alpha. \quad (2.13)$$

The weights (2.9) allow to prove very nice results when working with parabolic equations. For instance, they allow to estimate a function in a quantitative way with its derivatives and with a local term. From [CG09, Lemma 1], we can obtain the following result by an easy induction, recalling that we can deal with the local terms by integrating by parts and leaving just the lower and higher order terms:

Lemma 2.5. *Let $k \in \mathbb{N}$, $m \geq 1$ and $r \in \mathbb{R}$. Then, there are $C > 0$ and $\lambda_0 \geq 1$ such that if $T > 0$, $\lambda \geq \lambda_0$, $s \geq CT^{2m}$ and $u \in L^2(0, T; H^{k+1}(\Omega))$, we have:*

$$\begin{aligned} \sum_{i=0}^k s^{2+r-2i} \lambda^{3+r-2i} \iint_Q e^{-2s\alpha} \xi^{2+r-2i} |\nabla^i u|^2 &\leq C \left(s^{r-2k} \lambda^{1+r-2k} \iint_Q e^{-2s\alpha} \xi^{r-2k} |\nabla^{k+1} u|^2 \right. \\ &\left. + s^{2+r} \lambda^{3+r} \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^{2+r} |u|^2 + s^{2+r-2k} \lambda^{3+r-2k} \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^{2+r-2k} |\nabla^k u|^2 \right). \end{aligned} \quad (2.14)$$

To continue with, we recall the following Carleman estimate for the heat equation with nonhomogeneous boundary conditions.

Lemma 2.6. *There is $C > 0$ such that for all $u \in C^2(\overline{Q})$, $a \in (0, 1]$, $m \geq 1$, $r \in \mathbb{R}$, $\lambda \geq C$ and $s \geq Ce^{C\lambda}(T^m + T^{2m})$ we have the inequality:*

$$\begin{aligned} s^{3+r} \lambda^{4+r} \iint_Q e^{-2s\alpha} \xi^{3+r} |u|^2 + s^{1+r} \lambda^{2+r} \iint_Q e^{-2s\alpha} \xi^{1+r} |\nabla u|^2 \\ \leq Cs^r \lambda^r \left(\iint_Q e^{-2s\alpha} \xi^r |g|^2 + s^{3+r} \lambda^{4+r} \iint_{Q_{\omega_0}} \xi^{3+r} |u|^2 + s^{1+r} \lambda^{1+r} \iint_{\Sigma} e^{-2s\alpha} \xi^{1+r} |\partial_n u|^2 \right), \end{aligned} \quad (2.15)$$

for $g := au_t - \Delta u$. In addition, if $u \equiv 0$ on Σ , one can drop the trace term in the right-hand side of (2.15).

Lemma 2.6 is well-known. We get the case $i = 1$, $m \geq 1$ and $r \in \mathbb{R}$ by repeating all the steps in [FCGBGP06, Theorem 1], where the authors prove the case $(a, r, m) = (1, 0, 1)$. In addition, we can get the uniformity in a following the steps of, for instance, [FI96] or [Bad11, Lemma 4.1]. As for the case with Dirichlet boundary condition, this is the classical Carleman estimate for the heat equation, which is proved in [FI96].

To finish with, we consider the following technical result:

Lemma 2.7. *Let $k \in \mathbb{N}$, $m > 0$, $r \in \mathbb{R}$ and let us consider the weights given in (2.9). Then, there is $C > 0$ such that for all $\lambda \geq 1$ and $s \geq C(\lambda)(T^m + T^{2m})$ we have the estimate:*

$$\|s^r e^{-s\alpha^*} (\xi^*)^r \varphi\|_{H^{k,2k}(Q)} + \|s^r e^{-s\alpha^*} (\xi^*)^r \pi\|_{H^{k-1,2k-1}(Q)} \leq \|s^{r+k+k/m} e^{-s\alpha^*} (\xi^*)^{r+k+k/m} \varphi\|_{L^2(Q)}, \quad (2.16)$$

for φ the solution of:

$$\begin{cases} -\varphi_t - \Delta\varphi + \nabla\pi = 0 & \text{in } Q, \\ \varepsilon\pi + \nabla \cdot \varphi = 0 & \text{in } Q, \\ \varphi = 0 & \text{on } \Sigma, \\ \varphi(T) = \varphi^T & \text{in } \Omega. \end{cases} \quad (2.17)$$

As far as we know, very similar versions of Lemma 2.7 have been used to deal with the trace terms since [Ima01] and it is widely known by the control community of Navier-Stokes like system. However, this result has mainly been used for $k = 1$ or $k = 2$, and as far as we know, the proof of the general case is not written. Thus, for completeness, we give the proof in Appendix A.

2.3 Proof of the observability estimate in 2-D

In this section we prove the following Carleman inequality for the solution of the homogeneous 2-D Boussinesq system (2.3):

Proposition 2.8. *Let Ω be such that Hypothesis 1.1 holds, $\omega \subset \Omega$ be a nonempty open set and $m \geq 8$. Then, there are $\varepsilon_0 > 0$, $C > 0$ and $\lambda_0 \geq 1$ such that if $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $(\varphi^T, \psi^T) \in L^2(\Omega)^3$, $\lambda \geq \lambda_0$, and $s \geq e^{C\lambda}(T^m + T^{2m})$, we have:*

$$s^{12} \lambda^{13} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3 \lambda^4 \iint_Q e^{-6s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-4s\alpha} |\psi|^2. \quad (2.18)$$

for the weights defined in (2.9) and (φ, ψ) the solution of (2.3).

First, we recall from [BP20a, Theorem 1.7] (considering that there is a symmetry between the variable ϕ_1 and ϕ_2 and replacing s by $\frac{5}{2}s$) the following Carleman estimate for the penalized Stokes system with an observation only with the second component.

Proposition 2.9. *Let Ω, ω, m as in Proposition 2.8, $\omega_0 \subset\subset \omega$ be an open set such that $\inf_{\omega \setminus \omega_0} |\nabla \eta| > 0$ and let $m \geq 8$. Then, there are $\varepsilon_0 > 0$, $C > 0$ and $\lambda_0 \geq 1$ such that if $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $\varphi^T \in L^2(\Omega)^2$, $\lambda \geq \lambda_0$, and $s \geq e^{C\lambda}(T^m + T^{2m})$, we have:*

$$s^{15} \lambda^{16} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 \leq C s^{34} \lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2,$$

for the weights defined in (2.9) and φ the solution of (2.17)-

Proof of Proposition 2.8. Considering Proposition 2.9 and Lemma 2.7 for $k = 1$, $r = 6$ and replacing s by $5s/2$, we obtain the estimate:

$$\begin{aligned} s^{12} \lambda^{13} \int_0^T e^{-5s\alpha^*} (\xi^*)^{12} \|\varphi\|_{H^2(\Omega)}^2 + s^{12} \lambda^{13} \iint_Q e^{-5s\alpha^*} (\xi^*)^{12} |\varphi_t|^2 \\ \leq C s^{15} \lambda^{16} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 \leq C s^{34} \lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2. \end{aligned} \quad (2.19)$$

Using the classical Carleman inequality of the heat equation on the ψ variable with homogeneous Dirichlet boundary conditions, (by Lemma 2.6 with $3s$ instead of s) we find for $\lambda \geq \lambda_0$ and $s \geq e^{C\lambda}(T^m + T^{2m})$ that:

$$s^3 \lambda^4 \iint_Q e^{-6s\alpha} \xi^3 |\psi|^2 \leq C \left(\iint_Q e^{-6s\alpha} |\varphi_2|^2 + s^3 \lambda^4 \iint_{Q_\omega} e^{-6s\alpha} \xi^3 |\psi|^2 \right). \quad (2.20)$$

We deal with the local term in the right hand side of (2.19) as in [Car12]. We consider a positive function χ supported in ω such that $\chi = 1$ in ω_0 . We have for all $\delta > 0$ by integrating by parts, using Lemma A.2, (2.12), (2.13) and Young's inequality the following estimate:

$$\begin{aligned} s^{34} \lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2 &\leq s^{34} \lambda^{35} \iint_{Q_\omega} e^{-5s\alpha} \chi \xi^{34} |\varphi_2|^2 \\ &= -s^{34} \lambda^{35} \iint_{Q_\omega} e^{-5s\alpha} \chi \xi^{34} (\psi_t + \Delta \psi) \varphi_2 = s^{34} \lambda^{35} \iint_{Q_\omega} \psi (\partial_t - \Delta) (e^{-5s\alpha} \chi \xi^{34} \varphi_2) \\ &\leq C_\delta \iint_{Q_\omega} e^{-4s\alpha} |\psi|^2 + \delta s^{12} \lambda^{13} \left(\int_0^T e^{-5s\alpha^*} (\xi^*)^{12} \|\varphi\|_{H^2(\Omega)}^2 + \iint_Q e^{-5s\alpha^*} (\xi^*)^{12} |\varphi_t|^2 \right). \end{aligned} \quad (2.21)$$

Thus, if we combine (2.19)-(2.21), by choosing δ small enough we obtain (2.18). \square

2.4 Proof of the observability estimate in 3-D

In this section we prove the following Carleman inequality, for the solution of the homogeneous 3-D Boussinesq system assuming Hypothesis 1.2 or 1.3.

2.4.1 A Carleman inequality assuming Hypothesis 1.2

In this section we prove the following result:

Proposition 2.10. *Let Ω be such that Hypothesis 1.2 holds, $\omega \subset \Omega$ be a nonempty open set and let $m \geq 2$. Then, there are $\varepsilon_0 > 0$, $C > 0$ and $\lambda_0 \geq 1$ such that if $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $(\varphi^T, \psi^T) \in L^2(\Omega)^4$, $\lambda \geq \lambda_0$, and $s \geq e^{C\lambda}(T^m + T^{2m})$, we have:*

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-s\alpha} (|\psi|^2 + |\varphi_2|^2), \quad (2.22)$$

for the weights defined in (2.9) and (φ, ψ) the solution of (2.3).

A geometrical result. First of all we prove the following geometrical result:

Proposition 2.11. *Let Ω be such that Hypothesis 1.2 holds. Then, for $a_0 > 0$ small enough, there is $C > 0$ such that for any function $w \in H^4(\Omega) \cap H_0^1(\Omega)$ and for any $a \in [0, a_0]$ we have that:*

$$\|\partial_{x_1} w\|_{L^2(\Omega)} \leq C \left(\|\partial_{x_1 x_2} w\|_{H^2(\Omega)} + \|\partial_{x_1 x_3} w\|_{H^2(\Omega)} + \|L_a w\|_{H^1(\partial\Omega)} \right), \quad (2.23)$$

with

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w - a \partial_{x_3 x_3} w. \quad (2.24)$$

Proof. We recall that by Hypothesis 1.2 there are a domain $\Omega_0 \subset \mathbb{R}^2$ and $H_-, H_+ \in C^0(\overline{\Omega_0}, [\delta, +\infty))$ such that $\Omega = \{x : -H_-(x_1, x_2) < x_3 < H_+(x_1, x_2)\}$. Using Proposition 2.4 we obtain that there is $C > 0$ and $a_0 > 0$ such that for all $a \in [0, a_0]$ and $s \in (-\delta, \delta)$ we have that:

$$\|\partial_{x_1} w\|_{C^0(\overline{\Omega_0} \times \{s\})} \leq C \left(\|\partial_{x_1 x_2} w\|_{H^2(\Omega_0 \times \{s\})} + \|\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w\|_{H^1(\partial\Omega_0 \times \{s\})} \right). \quad (2.25)$$

Considering that $\partial_{x_3 x_3} w = 0$ on $\partial\Omega_0 \times (-\delta, \delta)$ (because of Dirichlet boundary conditions) we have that:

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w \quad \text{on } \Omega_0 \times (-\delta, \delta). \quad (2.26)$$

Thus, if we integrate (2.25) squared on $(-\delta, \delta)$ and consider (2.26), we obtain that:

$$\|\partial_{x_1} w\|_{L^2(\Omega_0 \times (-\delta, \delta))}^2 \leq C \left(\|\partial_{x_1 x_2} w\|_{H^2(\Omega_0 \times (-\delta, \delta))}^2 + \|L_a w\|_{H^1(\partial\Omega_0 \times (-\delta, \delta))}^2 \right). \quad (2.27)$$

Finally, considering that for all $x = (x_1, x_2, x_3) \in \Omega$ and $s \in (-\delta, \delta)$ we have that:

$$\partial_{x_1} w(x_1, x_2, x_3) = \partial_{x_1} w(x_1, x_2, s) + \int_s^{x_3} \partial_{x_1 x_3} w(x_1, x_2, s') ds'. \quad (2.28)$$

By integrating (2.28), using (2.27) and remarking that $\Omega_0 \times (-\delta, \delta) \subset \Omega$, we easily obtain (2.23). \square

Proof of Proposition 2.10. Because the proof is quite technical, we divide it into twelve steps.

- **Step 1:** We first use the coercivity estimate given in Proposition 2.11 for φ_1 .
- **Step 2:** We add terms involving φ_2 and φ_3 in both sides of the previous estimate.
- **Step 3:** We get rid of the terms involving φ_1 in the right-hand side of the previous estimate thanks to the equations satisfied by φ_2 and φ_3 .
- **Step 4:** We choose from this step forward to treat only one term in the right-hand side of the estimate to simplify. It will be $\partial_{x_2x_3}\varphi_2$, all the other terms can be treated in a similar way. In this step we use Lemma 2.5.
- **Step 5:** We begin with a Carleman estimate for the derivatives of $\partial_{x_2x_3}\varphi_2$, this Carleman estimate comes from Lemma 2.6.
- **Step 6:** The previous inequality leads to a global term involving the pressure. This is why we use also a Carleman estimate for the derivatives of $\partial_{x_2x_3x_2}\pi$.
- **Step 7:** We then get rid of the trace terms of $\partial_{x_2x_3}\varphi_2$ and $\partial_{x_2x_3x_2}\pi$ by using the regularity result stated in Lemma 2.7.
- **Step 8:** We eliminate the local term in pressure i.e. $\partial_{x_2x_3x_2}\pi$ by the second equation of the system.
- **Step 9:** We gather Steps 4, 5, 6, 7 and 8.
- **Step 10:** We first estimate the local term in the derivatives of $\partial_{x_2x_3}\varphi_2$ by a local term in $\partial_{x_2x_3}\varphi_2$ by using standard integration by parts. We then estimate the local term of $\partial_{x_2x_3}\varphi_2$ in function of a local term in φ_2 .
- **Step 11:** We apply the standard Carleman estimate in the equation satisfied by the temperature.
- **Step 12:** We estimate the local term in φ_3 by the equation satisfied by the temperature.

First, let us fix three nonempty open sets $\omega_1, \omega_2, \omega_3$ such that $\omega_0 \subset\subset \omega_1 \subset\subset \omega_2 \subset\subset \omega_3 \subset\subset \omega$.

Step 1: The equation on the boundary of φ_1 is given by

$$\partial_{x_1x_1}\varphi_1 - \frac{\varepsilon}{1+\varepsilon}\partial_{x_2x_2}\varphi_1 - \frac{\varepsilon}{1+\varepsilon}\partial_{x_3x_3}\varphi_1 = \frac{1}{1+\varepsilon}(\partial_{x_1x_2}\varphi_2 + \partial_{x_1x_3}\varphi_3). \quad (2.29)$$

Thus, using Proposition 2.11 and Poincaré inequality we have for every $t \in [0, T]$:

$$\begin{aligned} \|\varphi_1(t, \cdot)\|_{L^2(\Omega)} \leq C & \left(\|\partial_{x_1x_2}\varphi_1(t, \cdot)\|_{H^2(\Omega)} + \|\partial_{x_1x_3}\varphi_1(t, \cdot)\|_{H^2(\Omega)} \right. \\ & \left. + \|\partial_{x_1x_2}\varphi_2(t, \cdot)\|_{H^2(\Omega)} + \|\partial_{x_1x_3}\varphi_3(t, \cdot)\|_{H^2(\Omega)} \right). \quad (2.30) \end{aligned}$$

Thus, we have that:

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi_1|^2 \leq C \left(s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \right. \\ \left. + s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_3|^2) \right), \quad (2.31)$$

for α^* and ξ^* defined in (2.9).

Step 2: We deduce from (2.31) the following estimate

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C \left(s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \right. \\ \left. + s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_3|^2 + |\varphi_2|^2 + |\varphi_3|^2) \right). \quad (2.32)$$

Step 3: We consider the equations satisfied by φ_2 and φ_3 , which are

$$\begin{cases} \frac{\varepsilon}{1+\varepsilon} \partial_t \varphi_2 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_1 x_1} \varphi_2 - \partial_{x_2 x_2} \varphi_2 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_3 x_3} \varphi_2 - \frac{1}{1+\varepsilon} (\partial_{x_1 x_2} \varphi_1 + \partial_{x_2 x_3} \varphi_3) = 0, \\ \frac{\varepsilon}{1+\varepsilon} \partial_t \varphi_3 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_1 x_1} \varphi_3 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_2 x_2} \varphi_3 - \partial_{x_3 x_3} \varphi_3 - \frac{1}{1+\varepsilon} (\partial_{x_1 x_3} \varphi_1 + \partial_{x_2 x_3} \varphi_2) = 0. \end{cases} \quad (2.33)$$

So we easily deduce that

$$s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \\ \leq C s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left(|\nabla^i \partial_{x_2 x_3} \varphi_3|^2 + |\nabla^i \partial_t \varphi_2|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_2|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_2|^2 \right. \\ \left. + |\nabla^i \partial_{x_2 x_3} \varphi_2|^2 + |\nabla^i \partial_t \varphi_3|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_3|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_3|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_3|^2 \right). \quad (2.34)$$

Now we remember that φ_2, φ_3 satisfy two heat equations with a source term involving the pressure

$$\forall i \in \{2, 3\}, \partial_t \varphi_i - \Delta \varphi_i = -\partial_{x_i} \pi \text{ in } Q, \varphi_i = 0 \text{ on } \Sigma. \quad (2.35)$$

Therefore, we can actually replace the partial derivative in time of φ_2 and φ_3 by $\partial_{x_2} \pi$ and $\partial_{x_3} \pi$ in (2.34). That is to say, we obtain

$$s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \\ \leq C s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left(|\nabla^i \partial_{x_2 x_3} \varphi_3|^2 + |\nabla^i \partial_{x_2} \pi|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_2|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_2|^2 \right. \\ \left. + |\nabla^i \partial_{x_2 x_3} \varphi_2|^2 + |\nabla^i \partial_{x_3} \pi|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_3|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_3|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_3|^2 \right). \quad (2.36)$$

We now gather (2.36) with (2.31) to deduce that

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C \left(s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left(|\varphi_2|^2 + |\varphi_3|^2 + |\nabla^i \partial_{x_2} \pi|^2 + |\nabla^i \partial_{x_3} \pi|^2 \right. \right. \\ \left. \left. + \sum_{i=0}^2 \sum_{j=2,3} |\nabla^i \partial_{x_1 x_2} \varphi_j|^2 + |\nabla^i \partial_{x_2 x_3} \varphi_j|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_j|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_j|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_j|^2 \right) \right). \quad (2.37)$$

Step 4: As we have said before, we focus on the estimation of $\partial_{x_2 x_3} \varphi_2$. We apply Lemma 2.5 to $\partial_{x_2 x_3} \varphi_2$, this leads to

$$s^{55}\lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53}\lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51}\lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\ + \dots + s^7 \lambda^8 \iint_Q e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \\ \leq C \left(s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^{55} \lambda^{56} \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 \right. \\ \left. + s^7 \lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \right). \quad (2.38)$$

Step 5: We now apply the Carleman estimate given by Lemma 2.6 to the equation satisfied by $\nabla^{25} \partial_{x_2 x_3}$, we obtain the following inequality

$$s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \\ \leq C \left(s^2 \lambda^2 \iint_Q e^{-2s\alpha} |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right. \\ \left. + s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha} \xi^3 |\partial_n \nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right). \quad (2.39)$$

Step 6: In order to estimate the global term involving the pressure in the right-hand side of (2.39), we also employ a Carleman estimate for the pressure. Indeed, by taking the divergence in (2.3)₁ we can easily see that:

$$\varepsilon \partial_t \pi - (1 - \varepsilon) \Delta \pi = 0,$$

so in particular

$$\forall i \in \{2, 3\}, \quad \frac{\varepsilon}{1 - \varepsilon} \partial_t (\partial_{x_i} \pi) - \Delta (\partial_{x_i} \pi) = 0.$$

Consequently, from Lemma 2.6 we obtain that:

$$s^2 \lambda^3 \iint_Q e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + \lambda \iint_Q e^{-2s\alpha} |\nabla^{26} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\ \leq C \left(s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha} \xi^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \right). \quad (2.40)$$

Step 7: We now gather the estimates (2.39) and (2.40) and take λ, s sufficiently large to get

$$\begin{aligned}
& s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad s^2 \lambda^3 \iint_Q e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + \lambda \iint_Q e^{-2s\alpha} |\nabla^{26} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\
& \leq C \left(s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha} \xi^3 |\partial_n \nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad \left. + s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha} \xi^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \right). \quad (2.41)
\end{aligned}$$

From (2.11) and the regularity result stated in Lemma 2.7, we deduce that for $m \geq 2$,

$$\begin{aligned}
& \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \varphi\|_{H^{16,32}(Q)} + \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \pi\|_{H^{15,31}(Q)} \\
& \leq \|s^{3/2+16+16/m} e^{-s\alpha^*} (\xi^*)^{3/2+16+16/m} \varphi\|_{L^2(Q)} \leq \|s^{51/2} e^{-s\alpha^*} (\xi^*)^{51/2} \varphi\|_{L^2(Q)}. \quad (2.42)
\end{aligned}$$

Then, we obtain that

$$\begin{aligned}
& s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha} \xi^3 |\partial_n \nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha} \xi^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\
& \leq \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2. \quad (2.43)
\end{aligned}$$

So by taking λ sufficiently large, the trace terms of the right hand side of (2.41) can be absorbed by the left-hand side of (2.32).

Step 8: From (2.35) and (2.42), by integration by parts and cut-off arguments, we have that there exist $r, r' > 0$ such that

$$\begin{aligned}
& s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 = -s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} [\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi][(\partial_t - \Delta) \nabla^{25} \partial_{x_2 x_3} \varphi_2] \\
& \leq \|s^{1/2} e^{-s\alpha^*} (\xi^*)^{1/2} \pi\|_{H^{15,31}(Q)}^2 + s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha+2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha+2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2. \quad (2.44)
\end{aligned}$$

This proves that for λ sufficiently large, the local term of the pressure in the right hand side of (2.41) can be absorbed by the left-hand side of (2.32).

Step 9: By using (2.38), (2.41), (2.43) and (2.44), we have

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \left(\lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad + s^{55} \lambda^{56} \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^7 \lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad \left. + s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right). \quad (2.45)
\end{aligned}$$

Step 10: We now estimate the local term in φ_2 , we proceed by standard integrations by parts, as in [BP20a, Section 5, Step 3], to obtain

$$\begin{aligned}
& s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^7 \lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^3 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq \delta \left(s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad \left. + \cdots + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \right) \\
& \quad + C_\delta s^r \lambda^{r'} \iint_{Q_{\omega_2}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\partial_{x_2 x_3} \varphi_2|^2, \quad (2.46)
\end{aligned}$$

for some other parameters $r, r' > 0$. By taking δ sufficiently small in (2.46), we deduce from (2.45) the following estimate

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_2}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\partial_{x_2 x_3} \varphi_2|^2. \quad (2.47)
\end{aligned}$$

By doing integration by parts in the same spirit as before, we have

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\varphi|^2, \quad (2.48)
\end{aligned}$$

for other parameters $r, r' > 0$.

By using (2.37), (2.42), (2.48) and performing the same strategy from Step 5 to Step 10 for all the terms appearing in the right-hand side of (2.37), we obtain that there exist $r, r' > 0$ such that

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha+2s\alpha^*} \xi^r (|\varphi_2|^2 + |\varphi_3|^2). \quad (2.49)$$

Step 11: Now, we apply the classical Carleman estimate for the heat equation with homogenous Dirichlet boundary condition satisfied by ψ , i.e. we use Lemma 2.6,

$$s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 + s\lambda^2 \iint_Q e^{-3s\alpha} \xi |\nabla\psi|^2 \leq C \left(\iint_Q e^{-3s\alpha} |\varphi_3|^2 + s^3\lambda^4 \iint_{Q_{\omega_0}} e^{-3s\alpha} \xi^3 |\psi|^2 \right). \quad (2.50)$$

By using (2.13), we sum (2.49) and (2.50) to get

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 + s\lambda^2 \iint_Q e^{-3s\alpha} \xi |\nabla\psi|^2 \leq C \left(s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha+2s\alpha^*} \xi^r (|\varphi_2|^2 + |\varphi_3|^2) + s^3\lambda^4 \iint_{Q_{\omega_0}} e^{-3s\alpha} \xi^3 |\psi|^2 \right). \quad (2.51)$$

Step 12: To eliminate the local term in φ_3 in the right-hand side of (2.51), we proceed exactly as in the proof of Proposition 2.8 in Section 2.3. Therefore, we obtain the desired estimate (2.22). \square

2.4.2 A Carleman inequality assuming Hypothesis 1.3

The goal of this part is to prove the following result:

Proposition 2.12. *Let Ω, ω be such that Hypothesis 1.3 holds and let $m \geq 12$. Then, there are $\varepsilon_0 > 0$, $C > 0$ and $\lambda_0 \geq 1$ such that if $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $\varphi^T \in (L^2(\Omega))^3$, $\lambda \geq \lambda_0$, and $s \geq e^{C\lambda}(T^m + T^{2m})$, we have:*

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-s\alpha^*} (|\psi|^2 + |\varphi_2|^2), \quad (2.52)$$

for the weights defined in (2.9) and for (φ, ψ) the solution of (2.3).

To prove Proposition 2.12, we follow the strategy of [FCGIP06], which consists in getting a Carleman estimate with some local terms and then using that the observation domain touches the boundary. The fact that the coupling of the system is of order 2 is an additional difficulty with respect to the systems treated in [FCGIP06].

Proof of Proposition 2.12. First of all, we consider that, by taking a smaller control domain if necessary, we can suppose that there are $\gamma \in \langle e_2, e_3 \rangle$ and $\tilde{\Gamma} \subset \Gamma$ a relatively open set such that $\omega = \{x + \lambda\gamma : x \in \tilde{\Gamma}, \lambda \in (0, 1)\}$. In addition, we consider

$$\omega_1 := \{x + \lambda\gamma : x \in \hat{\Gamma}, \lambda \in (0, 1/2)\}, \quad (2.53)$$

for $\hat{\Gamma}$ a relatively open set contained in $\tilde{\Gamma}$. Finally, we consider ω_0 a non-empty open set compactly contained in ω_1 .

In order to prove the Carleman inequality (2.52), we first remark that $\partial_{x_1}\varphi_1$ satisfies the equation:

$$\left(\frac{\varepsilon}{1+\varepsilon}\partial_t - \Delta\right)\partial_{x_1}\varphi_1 = \frac{1}{1+\varepsilon}(\partial_{x_1x_1x_2}\varphi_2 + \partial_{x_1x_1x_3}\varphi_3 + \partial_{x_1x_2x_2}\varphi_1 + \partial_{x_1x_3x_3}\varphi_1). \quad (2.54)$$

To continue with, using Poincaré's inequality for φ_1 and Lemma 2.5 for $\partial_{x_1}\varphi_1$ with $k = 1$, we obtain that:

$$\begin{aligned} & s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*}(\xi^*)^{16}|\varphi_1|^2 + s^{16}\lambda^{17} \iint_Q e^{-2s\alpha}\xi^{16}|\partial_{x_1}\varphi_1|^2 + s^{14}\lambda^{15} \iint_Q e^{-2s\alpha}\xi^{14}|\nabla\partial_{x_1}\varphi_1|^2 \\ & \leq C\left(s^{12}\lambda^{13} \iint_Q e^{-2s\alpha}\xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 + s^{16}\lambda^{17} \iint_{Q_{\omega_0}} \xi^{16}|\partial_{x_1}\varphi_1|^2 \right. \\ & \quad \left. + s^{14}\lambda^{15} \iint_{Q_{\omega_0}} \xi^{14}|\nabla\partial_{x_1}\varphi_1|^2\right). \end{aligned} \quad (2.55)$$

In particular, by using (2.54) and the Carleman estimate coming from Lemma 2.6 for the terms of $\nabla^2\partial_{x_1}\varphi_1$ we obtain that:

$$\begin{aligned} s^{12}\lambda^{13} \iint_Q e^{-2s\alpha}\xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 & \leq C\left(s^9\lambda^9 \iint_Q e^{-2s\alpha}\xi^9|\nabla^2g|^2 + s^{12}\lambda^{13} \iint_{Q_{\omega_0}} \xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 \right. \\ & \quad \left. + s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha}\xi^{10}|\partial_n\nabla^2\partial_{x_1}\varphi_1|^2\right), \end{aligned} \quad (2.56)$$

for $g := \partial_{x_1x_1x_2}\varphi_2 + \partial_{x_1x_1x_3}\varphi_3 + \partial_{x_1x_2x_2}\varphi_1 + \partial_{x_1x_3x_3}\varphi_1$.

To continue with, we find from Poincaré's inequality and Lemma 2.5 with $k = 0$ that:

$$\begin{aligned} & s^{17}\lambda^{18} \iint_Q e^{-2s\alpha^*}(\xi^*)^{17}(|\varphi_2|^2 + |\varphi_3|^2) + s^{17}\lambda^{18} \iint_Q e^{-2s\alpha}\xi^{17}(|\partial_{x_1}\varphi_2|^2 + |\partial_{x_1}\varphi_3|^2) \\ & \leq C\left(s^{15}\lambda^{16} \iint_Q e^{-2s\alpha}\xi^{15}(|\nabla\partial_{x_1}\varphi_2|^2 + |\nabla\partial_{x_1}\varphi_3|^2) \right. \\ & \quad \left. + s^{17}\lambda^{18} \iint_{Q_{\omega_0}} e^{-2s\alpha}\xi^{17}(|\partial_{x_1}\varphi_2|^2 + |\partial_{x_1}\varphi_3|^2)\right). \end{aligned} \quad (2.57)$$

Next, we remark that we can follow the steps 3-11 of Section 2.4.1 and by using the estimates on the weights (2.12), (2.13), Lemmas 2.6 and 2.7 and (2.55), (2.56), (2.57), we deduce that:

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \left(\iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha} (|\psi|^2 + |\varphi_2|^2) \right. \\ \left. \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 + s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha} \xi^{10} |\partial_n \nabla^2 \partial_{x_1}\varphi_1|^2 \right). \quad (2.58)$$

Remark 2.13. It helps us to have only derivatives of $\partial_{x_1}\varphi_2$ and $\partial_{x_1}\varphi_2$ in the right-hand side of (2.57) because we eventually do a Carleman inequality for $\nabla^i \partial_{x_1}\varphi$ as in [BP20a, Section 5], so the local term that we get is of the form $\nabla^i \partial_{x_1}\varphi$. If instead of having $\partial_{x_1}\varphi_2$ and $\partial_{x_1}\varphi_2$ we had $\nabla\varphi_2$ and $\nabla\varphi_3$, after applying the Carleman estimate to $\nabla^i\varphi$ we would get a local term which have $\partial_{x_2}^i\varphi_1$ and $\partial_{x_3}^i\varphi_1$ terms from which we do not know how to get a Carleman inequality with a local term that only depends on φ_2 and ψ .

Let us now treat the boundary term of φ_1 as in Section 2.4.1. By interpolation and Lemma 2.7 we have:

$$s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha} \xi^{10} |\partial_n \nabla^2 \partial_{x_1}\varphi_1|^2 \\ \leq C \|s^{23/4}\lambda^{23/4} e^{-s\alpha^*} (\xi^*)^{23/4} \varphi\|_{L^2(0,T;H^4(\Omega))}^{1/2} \|s^{19/4}\lambda^{19/4} e^{-s\alpha^*} (\xi^*)^{19/4} \varphi\|_{L^2(0,T;H^6(\Omega))}^{3/2} \\ \leq C \|s^{31/4+2/m}\lambda^{23/4} e^{-s\alpha^*} (\xi^*)^{23/4} \varphi\|_{L^2(Q)}^{1/2} \|s^{31/4+3/m}\lambda^{19/4} e^{-s\alpha^*} (\xi^*)^{31/4+3/m} \varphi\|_{L^2(Q)}^{3/2}.$$

So, if $m \geq 12$, we have that the trace term in the right-hand side of (2.58) can be absorbed by the left-hand side of (2.58), then we obtain:

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \left(\iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha} (|\psi|^2 + |\varphi_2|^2) \right. \\ \left. + \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 \right). \quad (2.59)$$

Let us now estimate the third right-hand side term of (2.59). From (2.53), recalling that e_1 is tangent to every point of $\tilde{\Gamma}$ by Hypothesis 1.3, $\partial_{x_1}\varphi_1 \equiv 0$ in $\partial\omega_1 \cap \partial\Omega$, and from Poincaré's inequality we obtain that:

$$\iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 \leq C \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} (|\partial_{x_2x_1}\varphi_1|^2 + |\partial_{x_3x_1}\varphi_1|^2). \quad (2.60)$$

Consequently, by using the equations (2.33), we can estimate the local terms in the right-hand side of (2.60) in function of local terms of φ_2 and φ_3 . By putting this in (2.59), we bound the local term in $\partial_{x_1}\varphi_1$ by local terms in φ_2 and φ_3 . In addition, we can deal with the local term of φ_3 by using the equation satisfied by ψ as before. Hence, by straightforward computations, we obtain the expected estimate (2.52). \square

2.5 Source term method

In this section we adapt the source term method of [LTT13] to our case. More precisely, from Proposition 2.1 we have an estimate for the control cost in L^2 of system (2.1). We now fix $M > 0$ such that $K(T) \leq Me^{M/T^m}$, with $K(T)$ defined as in (2.2). In addition, we fix the values:

$$q \in (1, \sqrt[2m]{2}), \quad p > q^{2m}/(2 - q^{2m}), \quad (2.61)$$

and the weights:

$$\rho_0(t) := M^{-p} \exp\left(-\frac{Mp}{(q-1)^m(T-t)^m}\right), \quad (2.62)$$

$$\rho_{\mathcal{F}}(t) := M^{-1-p} \exp\left(-\frac{(1+p)q^{2m}M}{(q-1)^m(T-t)^m}\right). \quad (2.63)$$

In addition, we define associated spaces for the source term, the state and the control:

$$\mathcal{F} := \left\{ f \in L^2(Q)^{N+1} : \frac{f}{\rho_{\mathcal{F}}} \in L^2(Q)^{N+1} \right\}, \quad (2.64)$$

$$\mathcal{Y} := \left\{ (y, \theta) \in (L^2(Q))^{N+1} : \frac{(y, \theta)}{\rho_0} \in L^2(Q)^{N+1} \right\}, \quad (2.65)$$

$$\mathcal{V} := \left\{ v \in L^2(Q_\omega)^{N+1} : \frac{v}{\rho_0} \in L^2(Q_\omega)^{N+1} \text{ and } v \text{ satisfies (1.2)} \right\}. \quad (2.66)$$

From the proof of [LTT13, Proposition 2.3], we deduce the null controllability of (2.1):

Proposition 2.14. *Let Ω, ω be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist $\varepsilon_0 > 0$, $m \geq 1$ and $C > 0$ such that for every $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $f \in \mathcal{F}$, $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$, there exists a control $v \in \mathcal{V}$ such that the solution (y, θ) of (2.1) belongs to \mathcal{Y} and we have the following estimate:*

$$\|(y, \theta)/\rho_0\|_{C([0, T]; L^2(\Omega))} + \|v/\rho_0\|_{L^2(Q_\omega)} \leq C (\|(y^0, \theta^0)\|_{L^2(\Omega)} + \|f/\rho_{\mathcal{F}}\|_{L^2(Q)}). \quad (2.67)$$

In particular, $(y, \theta)(T) = 0$.

The next proposition gives more information on the regularity of the controlled trajectory obtained in Proposition 2.14. We consider the weight:

$$\rho(t) = \exp\left(-\frac{M\beta}{(q-1)^m(T-t)^m}\right), \text{ with } \frac{(1+p)q^{2m}}{2} < \beta < p.$$

We remark that $\rho(T) = 0$ and that ρ satisfies the inequalities:

$$\rho_0 \leq C\rho, \quad \rho_{\mathcal{F}} \leq C\rho, \quad |\rho'| \rho_0 \leq C\rho^2, \quad \rho^2 \leq C\rho_{\mathcal{F}}. \quad (2.68)$$

Proposition 2.15. *Let Ω, ω be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist $\varepsilon_0 > 0$, $m \geq 1$ and $C > 0$ such that for every $T > 0$, $\varepsilon \in (0, \varepsilon_0)$, $f \in \mathcal{F}$, $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$, there exists a control $v \in \mathcal{V}$ such that the solution (y, θ) of (2.1) satisfies:*

$$\|(y, \theta)/\rho\|_{L^2(0,T;H^2(\Omega)) \cap H^1(0,T;L^2(\Omega))} + \|v/\rho_0\|_{L^2(Q_\omega)} \leq C \left(\|(y^0, \theta^0)\|_{H_0^1(\Omega)} + \|f/\rho_{\mathcal{F}}\|_{L^2(Q)} \right). \quad (2.69)$$

In particular, $(y, \theta)(T) = 0$.

The proof of Proposition 2.15 is a straightforward adaptation of the proof of [LTT13, Proposition 2.8] by using a regularity estimate of the penalized Stokes system (see Lemma 2.3) and of the heat equation.

Remark 2.16. For each $f \in \mathcal{F}$ and $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$, by classical arguments (see [LTT13, Proposition 2.9]), we can fix a control $v \in \mathcal{V}$ such that (y, θ) and v satisfy (2.69) by choosing among those the unique minimizer of the functional $v \mapsto \|(y, \theta)/\rho\|_{L^2(0,T;H^2(\Omega)) \cap H^1(0,T;L^2(\Omega))}^2 + \|v/\rho_0\|_{L^2(Q_\omega)}^2$.

3 Local null-controllability of the nonlinear Boussinesq system

In this section we conclude the proof of Theorem 1.2. In particular, it is organised as follows:

- In Section 3.1 we use a fixed-point argument to get the local null-controllability of the nonlinear system (1.1) for initial data in $H_0^1(\Omega)^{N+1}$.
- In Section 3.2 we use a regularization argument to deduce the local null-controllability of (1.1) for initial data in $L^2(\Omega)^{N+1}$.

3.1 Fixed-point argument

In this section, we give the proof of Theorem 1.2 for initial data in $H_0^1(\Omega)^{N+1}$. In this part, C denotes a positive constant that depends on Ω, ω and T but independent of ε and varying from line to line. In addition, we denote:

$$\mathcal{F}_r := \{f \in \mathcal{F} : \|f/\rho_{\mathcal{F}}\|_{L^2(Q)} \leq r\}, \quad (3.1)$$

for $r > 0$ a small enough parameter independent of ε that will be determined later.

We now fix $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$ such that:

$$\|(y^0, \theta^0)\|_{H_0^1(\Omega)} \leq r. \quad (3.2)$$

It follows that we can define an operator \mathcal{N} acting on \mathcal{F}_r by:

$$\mathcal{N}(f) := \left(-(y \cdot \nabla)y - \frac{1}{2}(\nabla \cdot y)y, -y \cdot \nabla\theta - \frac{1}{2}(\nabla \cdot y)\theta \right), \quad (3.3)$$

where (y, θ) is the corresponding trajectory of (2.1) associated to the initial values (y^0, θ^0) , the force f and the control v given by Proposition 2.15 and Remark 2.16.

To conclude the proof of Theorem 1.2 for regular initial data it suffices to check that, for $r > 0$ small enough not depending on ε , \mathcal{N} is a contractive mapping from \mathcal{F}_r into itself and then apply the Banach fixed-point theorem.

Step 1: \mathcal{F}_r is invariant for \mathcal{N} provided that r is small enough. By using (2.68)₃ and the Sobolev embeddings $H^1(\Omega) \hookrightarrow L^4(\Omega)$ and $H^2(\Omega) \hookrightarrow W^{1,4}(\Omega)$ because $N \leq 3$, we have for almost every $t \in (0, T)$:

$$\begin{aligned} \left\| \frac{\mathcal{N}(f)}{\rho_{\mathcal{F}}}(t) \right\|_{L^2(\Omega)} &\leq C \left| \frac{\rho^2}{\rho_{\mathcal{F}}}(t) \right| \left(\left\| \frac{(y \cdot \nabla)y}{\rho^2}(t) \right\|_{L^2(\Omega)} + \left\| \frac{(\nabla \cdot y)y}{\rho^2}(t) \right\|_{L^2(\Omega)} \right. \\ &\quad \left. + \left\| \frac{y \cdot \nabla\theta}{\rho^2}(t) \right\|_{L^2(\Omega)} + \left\| \frac{(\nabla \cdot y)\theta}{\rho^2}(t) \right\|_{L^2(\Omega)} \right) \\ &\leq C \left(\left\| \frac{y}{\rho}(t) \right\|_{L^4(\Omega)} \left\| \frac{\nabla y}{\rho}(t) \right\|_{L^4(\Omega)} + \left\| \frac{y}{\rho}(t) \right\|_{L^4(\Omega)} \left\| \frac{\nabla\theta}{\rho}(t) \right\|_{L^4(\Omega)} \right. \\ &\quad \left. + \left\| \frac{\nabla y}{\rho}(t) \right\|_{L^4(\Omega)} \left\| \frac{\theta}{\rho}(t) \right\|_{L^4(\Omega)} \right) \\ &\leq C \left(\left\| \frac{y}{\rho}(t) \right\|_{H^1(\Omega)} \left\| \frac{y}{\rho}(t) \right\|_{H^2(\Omega)} + \left\| \frac{y}{\rho}(t) \right\|_{H^1(\Omega)} \left\| \frac{\theta}{\rho}(t) \right\|_{H^2(\Omega)} \right. \\ &\quad \left. + \left\| \frac{y}{\rho}(t) \right\|_{H^2(\Omega)} \left\| \frac{\theta}{\rho}(t) \right\|_{H^1(\Omega)} \right). \end{aligned}$$

Then, by integrating in the interval time $t \in (0, T)$ and by using (2.69), (3.2) and (3.1), we have:

$$\begin{aligned} \left\| \frac{\mathcal{N}(F)}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)} &\leq C \left(\left\| \frac{y}{\rho} \right\|_{C([0,T];H_0^1(\Omega))} \left(\left\| \frac{y}{\rho} \right\|_{L^2(0,T;H^2(\Omega))} + \left\| \frac{\theta}{\rho} \right\|_{L^2(0,T;H^2(\Omega))} \right) \right. \\ &\quad \left. + C \left\| \frac{\theta}{\rho} \right\|_{C([0,T];H_0^1(\Omega))} \left\| \frac{y}{\rho} \right\|_{L^2(0,T;H^2(\Omega))} \right) \\ &\leq C \left(\|(y^0, \theta^0)\|_{H_0^1(\Omega)}^2 + \|F/\rho_{\mathcal{F}}\|_{L^2(Q)}^2 \right) \\ &\leq Cr^2. \end{aligned}$$

So, for $r > 0$ small enough, \mathcal{N} stabilises \mathcal{F}_r .

Step 2: \mathcal{N} is contracting provided that r is small enough. Using the same kind of arguments, it is not difficult to obtain that:

$$\left\| \frac{\mathcal{N}(f^1) - \mathcal{N}(f^2)}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)} \leq Cr \left\| \frac{f^1 - f^2}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)}.$$

Consequently, by taking r sufficiently small, \mathcal{N} is a contracting mapping on the closed ball \mathcal{F}_r . Therefore, by the Banach fixed point theorem, \mathcal{N} has a unique fixed-point f . By denoting (y, θ, v) the associated trajectory to f , we find that (y, θ, v) satisfies the system (1.1) and $(y, \theta)(T) = 0$. Remark that r does not depend on ε , so the control is bounded uniformly when ε goes to 0. This concludes the proof.

3.2 Smoothing effect of the nonlinear Boussinesq system

In this section, we give the proof of Theorem 1.2 for data in $L^2(\Omega)^{N+1}$. This type of arguments have already been used in [CMS20] and [CSFCB⁺20]. The key remark is the following regularity lemma.

Lemma 3.1. *Let $T > 0$. There exists a positive constant $C_T > 0$ such that for every $\varepsilon > 0$, any $(y_0, \theta_0) \in L^2(\Omega)^{N+1}$ and any weak solution of (1.1) with control $v \equiv 0$,*

$$\exists t_0 \in [0, T], \|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq C_T \|(y_0, \theta_0)\|_{L^2(\Omega)}.$$

Proof. We multiply (1.1)₁ by y and (1.1)₂ by θ , then integrate in Ω , we find

$$\begin{aligned} \left(\frac{d}{dt} \int_{\Omega} y(t, \cdot)^2 \right) + \int_{\Omega} |\nabla y(t, \cdot)|^2 + \frac{1}{\varepsilon} \int_{\Omega} (\nabla \cdot y(t, \cdot))^2 &= \int_{\Omega} \theta(t, \cdot) y_N(t, \cdot), \\ \left(\frac{d}{dt} \int_{\Omega} \theta(t, \cdot)^2 \right) + \int_{\Omega} |\nabla \theta(t, \cdot)|^2 &= 0. \end{aligned}$$

We sum, use Young's inequality and Gronwall's lemma to obtain that there exists $C > 0$ such that for all $t \in [0, T]$:

$$\|(y, \theta)(t, \cdot)\|_{L^2(\Omega)}^2 + \int_0^t \|(y, \theta)(s, \cdot)\|_{H^1(\Omega)}^2 ds \leq C \|(y_0, \theta_0)\|_{L^2(\Omega)}^2.$$

In particular, this gives us the existence of a time $t_0 \in [0, T]$ such that

$$\|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq \sqrt{\frac{C}{T}} \|(y_0, \theta_0)\|_{L^2(\Omega)}.$$

This concludes the proof with $C_T = \sqrt{C/T}$. □

We now have all the tools to end the proof of Theorem 1.2.

Proof of Theorem 1.2. We can control the initial data with the following three steps:

Step 1: regularization of the data. By setting $v \equiv 0$ and by using Lemma 3.1, we deduce that there exists $t_0 \in (0, T/2)$ such that $\|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq C_{T/2} \|(y_0, \theta_0)\|_{L^2(\Omega)}$.

Step 2: local null-controllability result in time $T/2$. By taking $(y_0, \theta_0) \in L^2(\Omega)^{N+1}$ such that $C_{T/2} \|(y_0, \theta_0)\|_{L^2(\Omega)} \leq \delta_{T/2}$, where $\delta_{T/2}$ is the radius of local null-controllability of (1.1) for initial data in $H_0^1(\Omega)^{N+1}$, we obtain that there exists a control $v \in L^2((t_0, t_0 + T/2) \times \omega)^{N+1}$ satisfying (1.2) such that $(y, \theta)(t_0 + T/2, \cdot) = 0$.

Step 3: do nothing at the end of the interval time. We set $v \equiv 0$ in $(t_0 + T/2, T) \times \omega$, so $(y, \theta)(T, \cdot) = 0$. \square

4 Remarks and open problems

In this section, we make some remarks and formulate some open problems concerning the null-controllability of the penalized Boussinesq system (1.1).

- If $N = 3$ by symmetry and by adapting Hypotheses 1.2 or 1.3, we can construct controls which satisfy:

$$\lambda_1 v_1 + \lambda_2 v_2 = v_3 = 0$$

for any $\lambda_1, \lambda_2 \in \mathbb{R}$. Indeed, all the proofs in this paper can be adapted to these situations by a simple change of coordinates.

- We can prove analogue controllability results assuming Hypothesis 1.1, 1.2 or 1.3 for the penalized Navier-Stokes system, which we recall is given by:

$$\begin{cases} y_t - \Delta y + (y \cdot \nabla) y + \frac{1}{2} (\nabla \cdot y) y + \nabla p = v 1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, & \text{on } \Sigma, \\ y(0) = y^0, & \text{in } \Omega. \end{cases} \quad (4.1)$$

Indeed, by following the strategy of this paper and by omitting the steps related with the heat equation satisfied by θ , we can prove that (4.1) is locally null-controllable uniformly on ε with a control v with one null component. Obtaining the local null-controllability of (4.1) with a control with two null-components in 3-D is an interesting open problem. A good strategy seems to employ the return method in the spirit of [CL14].

- The exponents of s , λ and ξ and the constant m stated in Propositions 2.8, 2.10 and 2.12 are a bit arbitrary. In particular, by combining Lemmas 2.5 and 2.7 a sufficient number of times we can get an analogous result for any $m > 1$ by choosing the exponent of s , λ and ξ large enough. This implies that the cost of the null controllability of the linearised system without a source term is less than $Ce^{CT^{-m}}$ for all $m > 1$.

- Removing the geometrical hypothesis on Ω is an interesting open problem.
- There are still plenty of systems with penalizations which approximate the incompressibility condition (see [She97]) whose null controllability properties have not been studied yet.

A Proof of Lemma 2.7

Let us prove Lemma 2.7. The proof consists in two steps: first we use parabolic regularity estimates and then we estimate the weights.

As a first step, we prove the following result:

Lemma A.1. *Let $k \in \mathbb{N}$ and $h \in H_0^k(0, T)$. Then, there is $C > 0$ such that for all $\varphi^T \in L^2(\Omega)$:*

$$\|h\varphi\|_{H^{k,2k}(Q)} + \|h\pi\|_{H^{k-1,2k-1}(Q)} \mathbf{1}_{k \geq 1} \leq C \|h^{(k)}\varphi\|_{L^2(Q)}, \quad (\text{A.1})$$

for (φ, π) the solution of (2.17).

Proof of Lemma A.1. The proof of Lemma A.1 is done by induction. The base case, $k = 0$, is trivial. Let us now prove the inductive case. By hypothesis we have (A.1) for any function $h \in H_0^k(0, T)$ and we have to prove the estimate:

$$\|h\varphi\|_{H^{k+1,2(k+1)}(Q)} + \|h\pi\|_{H^{k,2k+1}(Q)} \leq C \|h^{(k+1)}\varphi\|_{L^2(Q)}, \quad (\text{A.2})$$

for any function $h \in H_0^{k+1}(0, T)$. We have that $(h\varphi, h\pi)$ satisfies:

$$\begin{cases} -(h\varphi)_t - \Delta(h\varphi) + \nabla(h\pi) = h'\varphi & \text{in } Q, \\ \varepsilon h\pi + \nabla \cdot (h\varphi) = 0 & \text{in } Q, \\ h\varphi = 0 & \text{on } \Sigma, \\ (h\varphi)(T, \cdot) = 0 & \text{in } \Omega. \end{cases}$$

Thus, since h and its derivatives get null on $t = T$, Lemma 2.3 implies that:

$$\|h\varphi\|_{H^{k+1,2(k+1)}(Q)} + \|h\pi\|_{H^{k,2k+1}(Q)} \leq C \|h'\varphi\|_{H^{k,2k}(Q)}. \quad (\text{A.3})$$

As $h' \in H_0^k(0, T)$ we can now use the inductive hypothesis (A.1) and obtain (A.2) from (A.3). \square

Using Lemma A.1 with $h = s^r e^{-s\alpha^*} (\xi^*)^r$ we find that:

$$\|s^r e^{-s\alpha^*} (\xi^*)^r \varphi\|_{H^{k,2k}(Q)} + \|s^r e^{-s\alpha^*} (\xi^*)^r \pi\|_{H^{k-1,2k-1}(Q)} \leq \|\partial_t^k (s^r e^{-s\alpha^*} (\xi^*)^r) \varphi\|_{L^2(Q)}. \quad (\text{A.4})$$

To conclude, we have to estimate the time derivative of the weights. We first recall the following result, which is classical, whose proof is sketched for completeness:

Lemma A.2. *Let $k \in \mathbb{N}_*$. Then, for all $s \geq C(\lambda)T^m$ we have the following estimate:*

$$|\partial_t^k \xi^*| + |\partial_t^k \alpha^*| \leq s^{k/m} (\xi^*)^{1+k/m}. \quad (\text{A.5})$$

Sketch of the proof of Lemma A.2. It is easy to prove by induction that:

$$\partial_t^k \xi^* = \frac{q(t, T)}{(t(T-t))^{m+k}}, \quad \partial_t^k \alpha^* = \frac{q(t, T)(e^{2\lambda\|\eta\|_\infty} - 1)}{(t(T-t))^{m+k}}. \quad (\text{A.6})$$

for q a homogeneous polynomial of degree k . Thus, considering that $t \leq T \leq (C(\lambda))^{-1}s^{1/m}$ as $s \geq C(\lambda)T^m$, we easily obtain (A.5) from (A.6). \square

Finally, (A.5) and $s\xi^* \geq 1$ imply the estimate:

$$\begin{aligned} \partial_t^k (s^r e^{-s\alpha^*} (\xi^*)^r) &\leq C s^r \sum_{i_1 + \dots + i_{k+1} = k} \left(\prod_{j=1, \dots, k: i_j \neq 0} s \partial_t^{i_j} \alpha^* \right) e^{-s\alpha^*} \partial_t^{i_{k+1}} [(\xi^*)^r] \\ &\leq s^{r+k+k/m} (\xi^*)^{r+k+k/m} e^{-s\alpha^*}. \end{aligned} \quad (\text{A.7})$$

Indeed, we get the maximum exponent for s and ξ^* in (A.7) by picking $j_1, \dots, j_k = 1$ and $j_{k+1} = 0$. Consequently, we obtain the desired estimate (2.16) from (A.4) and (A.7).

References

- [AKBGBdT11] F. Ammar-Khodja, A. Benabdallah, M. González Burgos, and L. de Teresa. Recent results on the controllability of linear coupled parabolic problems: a survey. *Math. Control Relat. F.*, 1(3):267–306, 2011.
- [Bad11] M. Badra. Global Carleman inequalities for Stokes and penalized Stokes equations. *Math. Control Relat. F.*, 1(2):149–175, 2011.
- [Ber78] M. Bercovier. Perturbation of mixed variational problems. application to mixed finite element methods. *ESAIM Math. Model. Num.*, 12(3):211–236, 1978.
- [BM20] K. Beauchard and F. Marbach. Unexpected quadratic behaviors for the small-time local null controllability of scalar-input parabolic equations. *J. Math. Pure. Appl.*, 2020.

- [BP20a] J. A. Bárcena-Petisco. Null controllability of a penalized Stokes problem in dimension two with one scalar control. *Asymptotic Anal.*, 117(3-4):161–198, 2020.
- [BP20b] J. A. Bárcena-Petisco. Uniform controllability of a stokes problem with a transport term in the zero-diffusion limit. *SIAM J. Control Optim.*, 58(3):1597–1625, 2020.
- [Car12] N. Carreño. Local controllability of the n-dimensional boussinesq system with n-1 scalar controls in an arbitrary control domain. *Math. Control Relat. F.*, 2(4):361–382, 2012.
- [CG09] J.-M. Coron and S. Guerrero. Null controllability of the N -dimensional Stokes system with $N - 1$ scalar controls. *J. Differ. Equations*, 246(7):2908–2921, 2009.
- [CL14] J.-M. Coron and P. Lissy. Local null controllability of the three-dimensional Navier-Stokes system with a distributed control having two vanishing components. *Invent. Math.*, 198(3):833–880, 2014.
- [CMS20] J.-M. Coron, F. Marbach, and F. Sueur. Small-time global exact controllability of the Navier-Stokes equation with Navier slip-with-friction boundary conditions. *J. Eur. Math. Soc. (JEMS)*, 22(5):1625–1673, 2020.
- [CnG13] N. Carreño and S. Guerrero. Local null controllability of the N -dimensional Navier-Stokes system with $N - 1$ scalar controls in an arbitrary control domain. *J. Math. Fluid Mech.*, pages 1–15, 2013.
- [CnGG15] N. Carreño, S. Guerrero, and M. Gueye. Insensitizing controls with two vanishing components for the three-dimensional Boussinesq system. *ESAIM: COCV*, 21(1):73–100, 2015.
- [Cor07] J.-M. Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2007.
- [CSFCB⁺20] F. W. Chaves-Silva, E. Fernández-Cara, K. Le Balc’h, J. L. F. Machado, and D. A. Souza. Small-time global exact controllability to the trajectories for the viscous boussinesq system. *arXiv eprint: 2006.01682*, 2020.
- [DL19] M. Duprez and P. Lissy. Bilinear local controllability to the trajectories of the Fokker-Planck equation with a localized control. *arXiv preprint arXiv:1909.02831*, 2019.
- [FCGBGP06] E. Fernández-Cara, M. González-Burgos, S. Guerrero, and J.-P. Puel. Null controllability of the heat equation with boundary Fourier conditions: the linear case. *ESAIM: COCV*, 12(3):442–465, 2006.

- [FCGIP06] E. Fernández-Cara, S. Guerrero, O. Yu. Imanuvilov, and J.-P. Puel. Some controllability results for the N -Dimensional Navier–Stokes and Boussinesq systems with $N - 1$ scalar controls. *SIAM J. Control Optim.*, 45(1):146–173, 2006.
- [FCLdM16] E. Fernández-Cara, J. Limaco, and S. B. de Menezes. Controlling linear and semilinear systems formed by one elliptic and two parabolic PDEs with one scalar control. *ESAIM:COCV*, 22(4):1017–1039, 2016.
- [FI96] A. V. Fursikov and O. Yu. Imanuvilov. *Controllability of evolution equations*. Number 34. Seoul National University, 1996.
- [GM18] S. Guerrero and C. Montoya. Local null controllability of the N -dimensional Navier–Stokes system with nonlinear Navier-slip boundary conditions and $N - 1$ scalar controls. *J. Math. Pure. Appl.*, 113:37–69, 2018.
- [Gue07] S. Guerrero. Controllability of systems of Stokes equations with one control force: existence of insensitizing controls. *Ann. I. H. Poincaré*, 24:1029–1054, 2007.
- [GZ19] B. Geshkovski and E. Zuazua. Controllability of one-dimensional viscous free boundary flows. *HAL preprint*, 2019.
- [HSLB19] V. Hernández-Santamaría and K. Le Balc’h. Local null-controllability of a nonlocal semilinear heat equation. *arXiv preprint arXiv:1912.08710*, 2019.
- [Ima01] O. Yu Imanuvilov. Remarks on exact controllability for the Navier-Stokes equations. *ESAIM:COCV*, 6:39–72, 2001.
- [IPY09] O. Yu. Imanuvilov, J.-P. Puel, and M. Yamamoto. Carleman estimates for parabolic equations with nonhomogeneous boundary conditions. *Chin. Ann. Math.*, 30(4):333–378, 2009.
- [LB18] K. Le Balc’h. Local controllability of reaction-diffusion systems around nonnegative stationary states. *arXiv e-prints:1809.05303*, to appear in *ESAIM Control Optim. Calc. Var.*, Sep 2018.
- [Lio88] J.-L. Lions. Contrôlabilité exacte, perturbations et stabilisation de systemes distribués, tome 1, RMA 8, 1988.
- [LTT13] Y. Liu, T. Takahashi, and M. Tucsnak. Single input controllability of a simplified fluid-structure interaction model. *ESAIM Control Optim. Calc. Var.*, 19(1):20–42, 2013.

- [LZ96] J.-L. Lions and E. Zuazua. A generique uniqueness result for the Stokes system and its control theoretical consequences. *Partial differential equations and applications: Collected Papers in Honor of Carlo Pucci*, 177:221–235, 1996.
- [MT18] S. Micu and T. Takahashi. Local controllability to stationary trajectories of a Burgers equation with nonlocal viscosity. *J. Differ. Equations*, 264(5):3664–3703, 2018.
- [OJ84] J. T. Oden and O.-P. Jacquotte. Stability of some mixed finite element methods for Stokesian flows. *Comput. Meth. Appl. Mat.*, 43(2):231–247, 1984.
- [Rus78] D. L. Russell. Controllability and stabilizability theory for linear partial differential equations: recent progress and open questions. *Siam Rev.*, 20(4):639–739, 1978.
- [She97] J. Shen. Pseudo-compressibility methods for the unsteady incompressible Navier-Stokes equations. *Proceedings of the 1994 Beijing symposium on nonlinear evolution equations and infinite dynamical systems*, pages 68–78, 1997.
- [Tak17] T. Takahashi. Boundary local null-controllability of the Kuramoto–Sivashinsky equation. *Math. Control Signal*, 29(1):2, 2017.
- [Tem68] R. Temam. Une méthode d’approximation de la solution des équations des Navier-Stokes. *Bull. Soc. Math. France*, 96:115–152, 1968.
- [Tem77] R. Temam. *Navier-Stokes equations. Theory and numerical analysis*. North Holland Publishing Company, first edition, 1977.
- [Zua96] E. Zuazua. A uniqueness result for the linear system of elasticity and its control theoretical consequences. *SIAM J. Control. and Optim.*, 34(5):1473–1495, 1996.