



**HAL**  
open science

# Local null controllability of the penalized Boussinesq system with a reduced number of controls

Jon Asier Bárcena-Petisco, Kévin Le Balc'H

► **To cite this version:**

Jon Asier Bárcena-Petisco, Kévin Le Balc'H. Local null controllability of the penalized Boussinesq system with a reduced number of controls. *Mathematical Control and Related Fields*, 2022, 12 (3), pp.641. 10.3934/mcrf.2021038 . hal-02913358v2

**HAL Id: hal-02913358**

**<https://hal.science/hal-02913358v2>**

Submitted on 7 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Local null controllability of the penalized Boussinesq system with a  
2 reduced number of controls

3 Jon Asier Bárcena-Petisco\*      Kévin Le Balc'h†

4 June 7, 2021

5 **Abstract:** In this paper we consider the Boussinesq system with homogeneous Dirichlet bound-  
6 ary conditions, defined in a regular domain  $\Omega \subset \mathbb{R}^N$  for  $N = 2$  and  $N = 3$ . The incompressibility  
7 condition of the fluid is replaced by its approximation by penalization with a small parameter  $\varepsilon > 0$ .  
8 We prove that our system is locally null controllable using a control with a restricted number of  
9 components, localized in an open set  $\omega$  contained in  $\Omega$ . We also show that the control cost is  
10 bounded uniformly with respect to  $\varepsilon \rightarrow 0$ . The proof is based on a linearization argument. The  
11 null controllability of the linearized system is obtained by proving a new Carleman estimate for  
12 the adjoint system. This inequality is derived by exploiting the coercivity of some second order  
13 differential operator involving crossed derivatives.

14  
15 **Key words:** Carleman inequality, controllability, nonlinear system, penalized Stokes system

16 **AMS subject classification:** 35K40, 93B05, 93C20

17 **Abbreviated title:** null controllability of the penalized Boussinesq system.

18 **Acknowledgements:** This project has received funding from the European Research Council  
19 (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agree-  
20 ment NO: 694126-DyCon) and from the French National Research Agency (ANR) in the frame of  
21 Programme IdEx Bordeaux - SysNum (ANR-10-IDEX-03-02). The two authors were also sup-  
22 ported by grants from Région Ile-de-France and “Numerical simulation and optimal control in view  
23 of temperature regulation in smart buildings” of the Nouvelle Aquitaine Region respectively.

---

\*[1] Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049 Madrid, Spain, [2] Chair of Computational Mathematics, Fundación Deusto, Avenida de las Universidades 24, 48007 Bilbao, Basque Country, Spain. [jon.barcena@uam.es](mailto:jon.barcena@uam.es)

†Institut de Mathématiques de Bordeaux, 351 Cours de la Libération, 33400 Bordeaux, France, [kevin.le-balch@math.u-bordeaux.fr](mailto:kevin.le-balch@math.u-bordeaux.fr)

# 1 Introduction

## 2 1.1 Presentation of the system

3 For a given time  $T > 0$ ,  $\Omega$  a sufficiently smooth bounded, connected, open set of  $\mathbb{R}^N$  ( $N = 2, 3$ )  
4 and  $\omega$  a nonempty open set contained in  $\Omega$ , we consider the controlled penalized Boussinesq system

$$\begin{cases} y_t - \Delta y + (y \cdot \nabla)y + \frac{1}{2}(\nabla \cdot y)y + \nabla p = \theta e_N + \tilde{v}1_\omega & \text{in } Q, \\ \theta_t - \Delta \theta + y \cdot \nabla \theta + \frac{1}{2}(\nabla \cdot y)\theta = v_{N+1}1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, \theta = 0 & \text{on } \Sigma, \\ y(0) = y^0, \theta(0) = \theta^0 & \text{in } \Omega, \end{cases} \quad (1.1)$$

5 where  $e_N$  stands for the  $N$ -th vector of the canonical basis of  $\mathbb{R}^N$ ,  $Q := (0, T) \times \Omega$ ,  $\Sigma := (0, T) \times \partial\Omega$ .  
6 In the controlled system (1.1),  $y = y(t, x)$  represents the velocity of the particles of the fluid,  
7  $\theta = \theta(t, x)$  their temperature and  $v = (\tilde{v}, v_{N+1}) = (v_1, \dots, v_N, v_{N+1})(t, x)$  stands for the control,  
8 acting on  $\omega$ .

9 System (1.1) approximates the classical incompressible Boussinesq system (replacing  $\nabla \cdot y = -\varepsilon p$   
10 by  $\nabla \cdot y = 0$  that implies that  $\frac{1}{2}(\nabla \cdot y)y = 0$  and  $\frac{1}{2}(\nabla \cdot y)\theta = 0$ ). This system has been introduced  
11 by Joseph Boussinesq in 1877 for modelling an incompressible fluid subjected to small variations  
12 of temperature. In addition, this way of approximating the incompressibility condition is called  
13 the penalty method and was introduced in [Tem68] in the Navier-Stokes case. As explained for  
14 instance in the survey [She97] and in the papers [Ber78] and [OJ84], this approximation procedure  
15 is widely used for numerical purpose.

16 We are going to focus on the small-time null controllability of the system (1.1) with a reduced  
17 number of controls. In particular, we seek controls  $v$  that satisfy:

$$\tilde{v} = 0, \text{ if } N = 2, \quad v_1 = v_3 = 0, \text{ if } N = 3. \quad (1.2)$$

18 This choice matches with what is known about the controllability of the classical Boussinesq system  
19 when the control satisfies (1.2) (see [Car12]). From a modelling point of view in the two-dimensional  
20 case, one wants to control both the velocity and the temperature of the fluid by acting only in the  
21 temperature equation. Similarly, in the three-dimensional case, one seeks to control the velocity  
22 by acting on both the temperature and the velocity, but with a scalar force acting on the second  
23 component of the velocity. One of the main interest in studying the controllability of (1.1) is that  
24 the controls of (1.1) converge to the controls of the Boussinesq system as  $\varepsilon$  goes to 0. Thus, in order

1 to compute numerically the controls of the Boussinesq system, it suffices to compute the controls  
 2 of (1.1) for  $\varepsilon > 0$  small enough.

### 3 1.2 Main result

4 In this part, we precise mathematically the main result of the paper. For this purpose, we introduce  
 5 three different geometrical assumptions on  $\Omega$  and  $\omega$ .

6 *Hypothesis 1.1.* For  $\Omega \subset \mathbb{R}^2$ , let  $\sigma^1, \dots, \sigma^k$  be the arc-length parametrizations of the different  
 7 connected components of  $\partial\Omega$ . We assume that for any  $i \in \{1, \dots, k\}$  and for any  $s \in [0, 1]$  such  
 8 that  $(\sigma_1^i)'(s) = 0$  or  $(\sigma_2^i)'(s) = 0$ , we have that  $\kappa^i(s) \neq 0$  where  $\kappa^i(s)$  is the curvature of  $\partial\Omega$  at the  
 9 point  $\sigma^i(s)$ .

*Hypothesis 1.2.* For  $\Omega \subset \mathbb{R}^3$ , we assume that there are  $\Omega_0 \subset \mathbb{R}^2$  satisfying Hypothesis 1.1,  $\delta > 0$   
 and  $H_-, H_+ \in C(\overline{\Omega_0}; [\delta, +\infty))$  such that:

$$\Omega = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 : (x_1, x_2) \in \Omega_0 \text{ and } -H_-(x_1, x_2) < x_3 < H_+(x_1, x_2)\}.$$

10 *Hypothesis 1.3.* For  $\Omega \subset \mathbb{R}^3$ , we assume that there are an interval  $I \subset \mathbb{R}$  and a curve  $\mathfrak{C} \subset \mathbb{R}^2$  such  
 11 that  $\Gamma := I \times \mathfrak{C}$  is a relative non trivial open set of  $\partial\Omega$ . In addition, we consider a control domain  
 12  $\omega \subset \Omega$  such that  $\Gamma \cap \partial\omega$  contains a relative non trivial open set.

13 *Example 1.1.* We recall that, as proved in [BP20a], Hypothesis 1.1 is satisfied by any strictly convex  
 14 smooth domain  $\Omega$ . Moreover, according to [BP20a, Lemma 1.3], for any smooth domain  $\Omega$ , one can  
 15 find a rotation (i.e. a linear application of the type  $U_\theta(x, y) = ((\cos \theta)x - (\sin \theta)y, (\sin \theta)x + (\cos \theta)y)$ )  
 16 that maps  $\Omega$  to a domain  $\Omega'$  satisfying Hypothesis 1.1. Furthermore, we can easily construct  
 17 regular domains  $\Omega$  satisfying Hypothesis 1.2, by considering a cylinder and some cupolas. Finally,  
 18 Hypothesis 1.3 is satisfied by any smooth domain  $\Omega$  containing a cylindrical part on its boundary.

19 We now present the main result of the paper:

20 **Theorem 1.2.** *Let  $\Omega, \omega$  be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exists  $\varepsilon_0 > 0$   
 21 such that for every time  $T > 0$  there exist  $\delta_T > 0$  and  $C_T > 0$  such that for every  $\varepsilon \in (0, \varepsilon_0)$   
 22 and every initial data  $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$  satisfying  $\|(y^0, \theta^0)\|_{L^2(\Omega)} \leq \delta_T$ , there exist a control  
 23  $v \in L^2(Q_\omega)^{N+1}$  satisfying (1.2),*

$$\|v\|_{L^2(Q_\omega)} \leq C_T, \tag{1.3}$$

and a weak solution  $(y, \theta)$  of (1.1) verifying

$$(y, \theta)(T) = 0.$$

24 Here and in the whole paper we shorten the notation and denote  $\|\cdot\|_V$  the norm  $\|\cdot\|_{V^k}$  for  $V$  any  
 25 Banach space and  $k \in \mathbb{N}_*$ . Indeed, the value of  $k$  is always easily deducible from the context.

1 In order to prove Theorem 1.2 we use the classical approach of proving first the controllability of  
 2 the linearized system and then using a fixed point theorem. In addition, we prove the controllability  
 3 of the linearized system by proving a Carleman inequality for the homogeneous adjoint system and  
 4 then using the approach of [LTT13]. The most difficult and original part is to prove the Carleman  
 5 inequality and in particular to prove the coercivity of some 2nd order differential operator involving  
 6 crossed derivatives. Before continuing, we make some comments on Theorem 1.2, its proof and  
 7 related bibliography:

- 8 • For the definition of a weak solution of (1.1), one can adapt [Tem68, Section I.1.]. Remark  
 9 that the existence of a weak solution is guaranteed by an adaptation of [Tem68, Théorème  
 10 I.2.] but the uniqueness is only valid in the 2D case.
- 11 • Theorem 1.2 is a null controllability result, uniform with respect to the parameter  $\varepsilon > 0$   
 12 because of the estimate (1.3). So, by letting  $\varepsilon \rightarrow 0$  in (1.1), we recover the results from  
 13 [Car12] for less regular initial data but for more restrictive assumptions on  $\Omega, \omega$ .
- 14 • We use the geometric hypothesis for proving the coercivity of a 2nd order differential operator  
 15 involving the crossed derivatives. For  $\Omega \subset \mathbb{R}^2$ , the assumption we make on the geometry of  
 16  $\Omega$ , i.e. Hypothesis 1.1, has already been introduced in the recent paper [BP20a], dealing  
 17 with the null controllability of the linear penalized 2-D Stokes system. This is crucial for  
 18 proving the null controllability of such a system with one scalar control. Indeed, as shown in  
 19 [Zua96, Theorem 1.2] and [BP20a, Section 2.1] some geometric hypothesis is needed because  
 20 the linearized system around 0 cannot be controlled due to the fact that some eigenfunctions  
 21 of the elliptic operator are not observable in a rhombus.
- 22 • At the heuristic level, it seems natural to obtain the controllability of the whole Boussinesq  
 23 system (1.1) by acting only with  $N - 1$  scalar controls. Indeed,  $v_{N+1}$  directly controls the  
 24 component  $\theta$  by the last equation of (1.1), then  $\theta$  acts as an indirect control to control  $y_N$  in  
 25 the  $N$ -th equation of (1.1)<sub>1</sub>. In addition, if  $N = 2$  the penalized divergence condition implies  
 26 that for  $\varepsilon$  small enough  $y_2$  acts as an indirect control to control  $y_1$ . Similarly, if  $N = 3$ ,  
 27  $y_1$  is directly controlled by  $v_1$  and  $y_2$  is indirectly controlled by  $y_1$  and  $y_3$  by the penalized  
 28 divergence condition for  $\varepsilon$  small enough.
- 29 • To prove the Carleman inequality, with the objective of highlighting the main ideas, we  
 30 prioritize giving a clear proof, even at the expense of not getting the most optimal results. In  
 31 that spirit, some recurrent operations are stated as lemmas (see Lemmas 2.5 and 2.7 below).  
 32 These technical results can be useful for proving other Carleman inequalities in a different  
 33 context.

- 1 • The results presented in Theorem 1.2 are original. Indeed, control problems with an approx-  
2 imation by penalization have first been studied in [IPY09, Section 4], where the penalized  
3 Stokes system was studied but without restriction on the control  $v$ . Next, the null controlla-  
4 bility of the penalized Navier-Stokes system has been studied in [Bad11], but again without  
5 restriction on the control  $\tilde{v}$ . Finally, the null controllability of the penalized Stokes system  
6 in the 2D-case with a scalar control has been established in [BP20a]. So, Theorem 1.2 is the  
7 first null controllability result for a penalized system with a reduced number of controls in  
8 the nonlinear setting, see Section 4 for a similar result in the Navier-Stokes case. Moreover,  
9 both the linear and nonlinear results in the 3D-case in this paper are new.
- 10 • Considering other systems, the study of controllability problems in which the control has a  
11 reduced number of components has been an active topic of research recently. In particular,  
12 for the Stokes and Navier-Stokes systems we can consult for instance the following papers:  
13 [LZ96, FCGIP06, Gue07, CG09, CnG13, CL14, CnGG15, GM18, BP20b]. For more results  
14 on the controllability of linear parabolic systems with a reduced number of controls, see the  
15 survey [AKBGBdT11] and the references therein.

16 The rest of the paper is organized as follows:

- 17 • In Section 2, we prove the null controllability of the linearized system.
- 18 • In Section 3, we prove the local null controllability of (1.1), i.e. we prove Theorem 1.2.
- 19 • In Section 4, we make some remarks and present some open problems.

## 20 **2 Null controllability of the linearized system**

21 In this section we prove the null controllability of the linearized system of (1.1). We divide the  
22 proof as follows:

- 23 • In Section 2.1 we linearize the system (1.1) around 0, and we recall the equivalence between  
24 the null controllability of the linearized system and the observability of the corresponding  
25 adjoint system.
- 26 • In Section 2.2 we recall some previous results about parabolic systems, elliptic systems and  
27 Carleman inequalities.
- 28 • In Sections 2.3 and 2.4 we obtain some Carleman inequalities for the adjoint system, for the  
29 2D case and the 3D case respectively.

- In Section 2.5 we use the source term method to get the null controllability of the linearized system and a source term, exponentially decreasing at  $t = T$ . This method has recently been used for many other control systems (see, for instance, [DL19, MT18, FCLdM16, BM20, Tak17, LB20, HSLB20, GZ21]).

We also introduce the notations  $Q_\omega := (0, T) \times \omega$ ,  $\delta_{ij}$  is a constant that is 1 if  $i = j$  and 0 otherwise,  $f = (\tilde{f}, f_{N+1}) = (f_1, \dots, f_N, f_{N+1})$  denotes the source term,  $H^{k_1, k_2}(Q) := H^{k_1}(0, T; L^2(\Omega)) \cap L^2(0, T; H^{k_2}(\Omega))$  and  $H^{k_1, k_2}(\Sigma) := H^{k_1}(0, T; L^2(\partial\Omega)) \cap L^2(0, T; H^{k_2}(\partial\Omega))$  (for  $k_1, k_2 \in \mathbb{R}^+$ ).

## 2.1 Linearization around 0, null controllability results and observability estimates

The linearization of (1.1) around 0 gives

$$\begin{cases} y_t - \Delta y + \nabla p = \theta e_N + \tilde{v}1_\omega & \text{in } Q, \\ \theta_t - \Delta \theta = v_{N+1}1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, \theta = 0 & \text{on } \Sigma, \\ y(0) = y^0, \theta(0) = \theta^0 & \text{in } \Omega. \end{cases} \quad (2.1)$$

The goal is to obtain the following result:

**Proposition 2.1.** *Let  $\Omega, \omega$  be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist  $\varepsilon_0 > 0$ ,  $m \geq 1$  and  $C > 0$  such that for every  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$  and  $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$ , there exists a control  $v \in L^2(Q_\omega)^{N+1}$  satisfying (1.2),*

$$\|v\|_{L^2(Q_\omega)} \leq K(T) \|(y^0, \theta^0)\|_{L^2(\Omega)}, \text{ where } K(T) := C \exp\left(\frac{C}{T^m}\right), \quad (2.2)$$

and such that the solution  $(y, \theta)$  of (2.1) satisfies  $(y, \theta)(T) = 0$ .

In order to prove Proposition 2.1, by the Hilbert Uniqueness Method (see, for instance, [Rus78], [Lio88] and [Cor07]), it is equivalent to establish an observability estimate for the adjoint system:

$$\begin{cases} -\varphi_t - \Delta \varphi + \nabla \pi = 0 & \text{in } Q, \\ -\psi_t - \Delta \psi = \varphi_N & \text{in } Q, \\ \nabla \cdot \varphi = -\varepsilon \pi & \text{in } Q, \\ \varphi = 0, \psi = 0 & \text{on } \Sigma, \\ \varphi(T) = \varphi^T, \psi(T) = \psi^T & \text{in } \Omega. \end{cases} \quad (2.3)$$

1 **Proposition 2.2.** *Let  $\Omega, \omega$  be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist  $\varepsilon_0 > 0$ ,*  
2  *$m \geq 1$  and  $C > 0$  such that for every  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$  and  $(\varphi^T, \psi^T) \in L^2(\Omega)^{N+1}$  the solution*  
3  *$(\varphi, \psi)$  of (2.3) satisfies:*

$$\|\varphi(0, \cdot)\|_{L^2(\Omega)}^2 + \|\psi(0, \cdot)\|_{L^2(\Omega)}^2 \leq K(T) \iint_{Q_\omega} \delta_{3,N} |\varphi_1(t, x)|^2 + |\psi(t, x)|^2 dt dx, \quad (2.4)$$

4 *with  $K(T)$  as in (2.2).*

5 To obtain the observability estimate (2.4), we will use Carleman estimates. More precisely,  
6 Proposition 2.2 will be a direct consequence of the Propositions 2.8, 2.10, 2.12, see below, and a  
7 classical dissipation argument.

## 8 2.2 Toolbox of elliptic, parabolic estimates and Carleman estimates

9 In this part, we recall elliptic and parabolic regularity estimates. We also present the Carleman  
10 estimates that will be useful in the sequel.

### 11 2.2.1 A parabolic and an elliptic result

12 Let us recall the regularity results of the Cauchy problem of the penalized Stokes system, which  
13 is given by the equations:

$$\begin{cases} u_t - \Delta u + \nabla q = \tilde{f} & \text{in } Q, \\ \varepsilon q + \nabla \cdot u = 0 & \text{in } Q, \\ u = 0 & \text{on } \Sigma, \\ u(0) = u^0 & \text{in } \Omega. \end{cases} \quad (2.5)$$

14 We have the following regularity estimate:

**Lemma 2.3.** *Let  $k \in \mathbb{N}_*$  and  $\Omega \subset \mathbb{R}^N$ . Then, there are  $\varepsilon_0 > 0$  and  $C > 0$  such that if  $T > 0$ ,  
 $\varepsilon \in (0, \varepsilon_0)$ ,  $u^0 \in H^{2k-1}(\Omega)^N$  and  $\tilde{f} \in H^{k-1, 2k-2}(Q)^N$  are such that  $(u^0, \tilde{f})$  satisfies the compatibility  
conditions*

$$g_0 := u^0 \in H_0^1(\Omega), \quad g_1 := \tilde{f}(0) - L_\varepsilon g_0 \in H_0^1(\Omega), \quad \dots, \quad g_{k-1} := \frac{d^{k-2} \tilde{f}}{dt^{k-2}}(0) - L_\varepsilon g_{k-2} \in H_0^1(\Omega),$$

15 *with  $L_\varepsilon g = -\Delta g - \varepsilon^{-1} \nabla(\nabla \cdot g)$ , we have that the solution  $(u, q)$  of (2.5) belongs to  $H^{k, 2k}(Q)^N \times$   
16  $H^{k-1, 2k-1}(Q)$  with the estimate:*

$$\|u\|_{H^{k, 2k}(Q)} + \|q\|_{H^{k-1, 2k-1}(Q)} \leq C \left( \|\tilde{f}\|_{H^{k-1, 2k-2}(Q)} + \|u^0\|_{H^{2k-1}(\Omega)} \right). \quad (2.6)$$

1 As explained in [BP20a, Lemma 2.5] (though it was done for the specific case  $u^0 = 0$ ), the proof of  
 2 Lemma 2.3 is mainly by induction. The initial case ( $k = 1$ ) can be proved again by the Galerkin  
 3 method. As for the inductive case, we get the regularity in the time variable by considering that  
 4  $u_t$  is a solution of (2.5) with  $(\tilde{f}, u^0)$  replaced by  $(\tilde{f}_t, \Delta u^0 + \varepsilon^{-1} \nabla(\nabla \cdot u^0) + \tilde{f}(0, \cdot))$  and using again  
 5 the Galerkin method (for instance, as in [Eva10, Theorem 7.1.6]). Moreover, we get the regularity  
 6 in space by using the estimate for the steady Stokes problem given in [Tem77, Proposition I.2.2].

7 As for elliptic results, we recall the following result that is proved in [BP20a, Theorem 1.8]  
 8 (considering the symmetry between the first and second variable):

9 **Proposition 2.4.** *Let  $\Omega$  be a domain such that Hypothesis 1.1 holds. Then, for  $a_0 > 0$  small  
 10 enough, there is  $C > 0$  such that for any function  $w \in H^4(\Omega) \cap H_0^1(\Omega)$  and for any  $a \in [0, a_0]$  we  
 11 have that:*

$$\|\partial_{x_2} w\|_{C^0(\bar{\Omega})} \leq C(\|\partial_{x_1 x_2} w\|_{H^2(\Omega)} + \|L_a w\|_{H^1(\partial\Omega)}), \quad (2.7)$$

12 *with:*

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w. \quad (2.8)$$

13 Roughly, Proposition 2.4 states that the crossed derivative with some information on the second  
 14 derivative on the boundary is coercive in  $H^4(\Omega) \cap H_0^1(\Omega)$ .

## 15 2.2.2 Classical Carleman estimates

16 We consider the following weights defined in  $Q$ :

$$\begin{aligned} \alpha(t, x) &:= \frac{e^{2\lambda\|\eta\|_\infty} - e^{\lambda\eta(x)}}{t^m(T-t)^m}, & \xi(t, x) &:= \frac{e^{\lambda\eta(x)}}{t^m(T-t)^m}, \\ \alpha^*(t) &:= \sup_{x \in (0, L)} \alpha(t, x), & \xi^*(t) &:= \inf_{x \in (0, L)} \xi(t, x), \end{aligned} \quad (2.9)$$

17 for  $m \in \mathbb{R}^+$ ,  $\lambda > 0$  that will be fixed later and for  $\eta \in C^2(\bar{\Omega}; \mathbb{R})$  satisfying:

$$\eta = 0 \text{ on } \partial\Omega, \quad \eta > 0 \text{ in } \Omega, \quad \inf_{\bar{\Omega} \setminus \omega_0} |\nabla \eta| > 0 \quad (2.10)$$

18 for some  $\omega_0 \subset\subset \omega$ . The existence of such a function  $\eta$  is proved in [FI96]. In fact, the weights (2.9)  
 19 are also taken from [FI96]. We recall that from (2.9) and (2.10) we obtain on  $\Sigma$  that:

$$\alpha(t, x) = \alpha^*(t), \quad \xi(t, x) = \xi^*(t). \quad (2.11)$$

20 We also recall that for all  $m > 0$  and  $\delta > 0$  there is  $C > 0$  such that for all  $s \geq 0$ :

$$s(\xi + \xi^* + \alpha + \alpha^*) \leq C e^{s\delta\alpha}. \quad (2.12)$$

21 In addition, for all  $\delta > 0$  there is  $C > 0$  such that if  $\lambda \geq C$  we have that:

$$\alpha^* \leq (1 + \delta)\alpha. \quad (2.13)$$

1 The weights (2.9) allow to prove very nice results when working with parabolic equations. For  
2 instance, they allow to estimate a function in a quantitative way with its derivatives and with  
3 a local term. From [CG09, Lemma 1], we can obtain the following result by an easy induction,  
4 recalling that we can deal with the local terms by integrating by parts and leaving just the lower  
5 and higher order terms:

**Lemma 2.5.** *Let  $k \in \mathbb{N}$ ,  $m \geq 1$  and  $r \in \mathbb{R}$ . Then, there are  $C > 0$  and  $\lambda_0 \geq 1$  such that if  $T > 0$ ,  
 $\lambda \geq \lambda_0$ ,  $s \geq CT^{2m}$  and  $u \in L^2(0, T; H^{k+1}(\Omega))$ , we have:*

$$\begin{aligned} \sum_{i=0}^k s^{2+r-2i} \lambda^{3+r-2i} \iint_Q e^{-2s\alpha \xi^{2+r-2i}} |\nabla^i u|^2 &\leq C \left( s^{r-2k} \lambda^{1+r-2k} \iint_Q e^{-2s\alpha \xi^{r-2k}} |\nabla^{k+1} u|^2 \right. \\ &\left. + s^{2+r} \lambda^{3+r} \iint_{Q_{\omega_0}} e^{-2s\alpha \xi^{2+r}} |u|^2 + s^{2+r-2k} \lambda^{3+r-2k} \iint_{Q_{\omega_0}} e^{-2s\alpha \xi^{2+r-2k}} |\nabla^k u|^2 \right). \end{aligned} \quad (2.14)$$

6 To continue with, we recall the following Carleman estimate for the heat equation with nonho-  
7 mogeneous boundary conditions.

**Lemma 2.6.** *There is  $C > 0$  such that for all  $u \in C^2(\overline{Q})$ ,  $a \in (0, 1]$ ,  $m \geq 1$ ,  $r \in \mathbb{R}$ ,  $\lambda \geq C$  and  
 $s \geq Ce^{C\lambda}(T^m + T^{2m})$  we have the inequality:*

$$\begin{aligned} s^{3+r} \lambda^{4+r} \iint_Q e^{-2s\alpha \xi^{3+r}} |u|^2 + s^{1+r} \lambda^{2+r} \iint_Q e^{-2s\alpha \xi^{1+r}} |\nabla u|^2 \\ \leq Cs^r \lambda^r \left( \iint_Q e^{-2s\alpha \xi^r} |g|^2 + s^{3+r} \lambda^{4+r} \iint_{Q_{\omega_0}} \xi^{3+r} |u|^2 + s^{1+r} \lambda^{1+r} \iint_{\Sigma} e^{-2s\alpha \xi^{1+r}} |\partial_n u|^2 \right), \end{aligned} \quad (2.15)$$

8 for  $g := -au_t - \Delta u$ . In addition, if  $u \equiv 0$  on  $\Sigma$ , one can drop the trace term in the right-hand side  
9 of (2.15).

10 Lemma 2.6 is well known. We get the case  $i = 1$ ,  $m \geq 1$  and  $r \in \mathbb{R}$  by repeating all the steps  
11 in [FCGBGP06, Theorem 1], where the authors prove the case  $(a, r, m) = (1, 0, 1)$ . In addition,  
12 we can get the uniformity in the parameter  $a \in (0, 1]$  following the steps of, for instance, [FI96]  
13 or [Bad11, Lemma 4.1]. As for the case with Dirichlet boundary condition, this is the classical  
14 Carleman estimate for the heat equation, which is proved in [FI96].

15 To finish with, we consider the following technical result:

**Lemma 2.7.** *Let  $k \in \mathbb{N}$ ,  $m > 0$ ,  $r \in \mathbb{R}$  and let us consider the weights given in (2.9). Then, there  
is  $C > 0$  such that for all  $\lambda \geq 1$  and  $s \geq e^{C\lambda}(T^m + T^{2m})$  the following estimate holds:*

$$\begin{aligned} \|s^r e^{-s\alpha^*} (\xi^*)^r \varphi\|_{H^{k, 2k}(Q)} + \|s^r e^{-s\alpha^*} (\xi^*)^r \pi\|_{H^{k-1, 2k-1}(Q)} \\ \leq \|s^{r+k+k/m} e^{-s\alpha^*} (\xi^*)^{r+k+k/m} \varphi\|_{L^2(Q)}, \end{aligned} \quad (2.16)$$

1 for  $\varphi$  the solution of:

$$\begin{cases} -\varphi_t - \Delta\varphi + \nabla\pi = 0 & \text{in } Q, \\ \varepsilon\pi + \nabla \cdot \varphi = 0 & \text{in } Q, \\ \varphi = 0 & \text{on } \Sigma, \\ \varphi(T) = \varphi^T & \text{in } \Omega. \end{cases} \quad (2.17)$$

2 As far as we know, very similar versions of Lemma 2.7 have been used to deal with the trace  
 3 terms since [Ima01] and it is widely known by the control community of Navier-Stokes like system.  
 4 However, this result has mainly been used for  $k = 1$  or  $k = 2$ , and as far as we know, the proof of  
 5 the general case is not written. Thus, for completeness, we give the proof in Appendix A.

### 6 2.3 Proof of the observability estimate in 2-D

7 In this section we prove the following Carleman inequality for the solution of the homogeneous 2-D  
 8 Boussinesq system (2.3):

9 **Proposition 2.8.** *Let  $\Omega$  be such that Hypothesis 1.1 holds,  $\omega \subset \Omega$  be a nonempty open set and  
 10  $m \geq 8$ . Then, there are  $\varepsilon_0 > 0$ ,  $C > 0$  and  $\lambda_0 \geq 1$  such that if  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  $(\varphi^T, \psi^T) \in L^2(\Omega)^3$ ,  
 11  $\lambda \geq \lambda_0$ , and  $s \geq e^{C\lambda}(T^m + T^{2m})$ , we have:*

$$s^{12}\lambda^{13} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-6s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-4s\alpha} |\psi|^2, \quad (2.18)$$

12 for the weights defined in (2.9) and  $(\varphi, \psi)$  the solution of (2.3).

13 First, we recall from [BP20a, Theorem 1.7] (considering that there is a symmetry between the  
 14 variable  $\phi_1$  and  $\phi_2$  and replacing  $s$  by  $\frac{5}{2}s$ ) the following Carleman estimate for the penalized Stokes  
 15 system with an observation only with the second component.

**Proposition 2.9.** *Let  $\Omega, \omega, m$  as in Proposition 2.8,  $\omega_0 \subset\subset \omega$  be an open set such that  $\inf_{\omega \setminus \omega_0} |\nabla\eta| >$   
 0 and let  $m \geq 8$ . Then, there are  $\varepsilon_0 > 0$ ,  $C > 0$  and  $\lambda_0 \geq 1$  such that if  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  
 $\varphi^T \in L^2(\Omega)^2$ ,  $\lambda \geq \lambda_0$ , and  $s \geq e^{C\lambda}(T^m + T^{2m})$ , we have:*

$$s^{15}\lambda^{16} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 \leq Cs^{34}\lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2,$$

16 for the weights defined in (2.9) and  $\varphi$  the solution of (2.17).

*Proof of Proposition 2.8.* Considering Proposition 2.9 and Lemma 2.7 for  $k = 1$ ,  $r = 6$  and replac-

ing  $s$  by  $5s/2$ , applied to the first equation of (2.3), we obtain the estimate:

$$\begin{aligned} s^{12}\lambda^{13} \int_0^T e^{-5s\alpha^*} (\xi^*)^{12} \|\varphi\|_{H^2(\Omega)}^2 + s^{12}\lambda^{13} \iint_Q e^{-5s\alpha^*} (\xi^*)^{12} |\varphi_t|^2 \\ \leq C s^{15}\lambda^{16} \iint_Q e^{-5s\alpha^*} (\xi^*)^{15} |\varphi|^2 \leq C s^{34}\lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2. \end{aligned} \quad (2.19)$$

- 1 Using the classical Carleman inequality of the heat equation on the  $\psi$  variable with homogeneous
- 2 Dirichlet boundary conditions for the second equation of (2.3), (by Lemma 2.6 with  $3s$  instead of
- 3  $s$ ) we find for  $\lambda \geq \lambda_0$  and  $s \geq e^{C\lambda(T^m + T^{2m})}$  that:

$$s^3\lambda^4 \iint_Q e^{-6s\alpha} \xi^3 |\psi|^2 \leq C \left( \iint_Q e^{-6s\alpha} |\varphi_2|^2 + s^3\lambda^4 \iint_{Q_\omega} e^{-6s\alpha} \xi^3 |\psi|^2 \right). \quad (2.20)$$

We deal with the local term in the right hand side of (2.19) as in [Car12]. We consider a positive function  $\chi$  supported in  $\omega$  such that  $\chi = 1$  in  $\omega_0$ . We have for all  $\delta > 0$  by integrating by parts, using Lemma A.2, (2.12), (2.13) and Young's inequality the following estimate:

$$\begin{aligned} s^{34}\lambda^{35} \iint_{Q_{\omega_0}} e^{-5s\alpha} \xi^{34} |\varphi_2|^2 &\leq s^{34}\lambda^{35} \iint_{Q_\omega} e^{-5s\alpha} \chi \xi^{34} |\varphi_2|^2 \\ &= -s^{34}\lambda^{35} \iint_{Q_\omega} e^{-5s\alpha} \chi \xi^{34} (\psi_t + \Delta\psi) \varphi_2 = s^{34}\lambda^{35} \iint_{Q_\omega} \psi (\partial_t - \Delta)(e^{-5s\alpha} \chi \xi^{34} \varphi_2) \\ &\leq C_\delta \iint_{Q_\omega} e^{-4s\alpha} |\psi|^2 + \delta s^{12}\lambda^{13} \left( \int_0^T e^{-5s\alpha^*} (\xi^*)^{12} \|\varphi\|_{H^2(\Omega)}^2 + \iint_Q e^{-5s\alpha^*} (\xi^*)^{12} |\varphi_t|^2 \right). \end{aligned} \quad (2.21)$$

- 4 Thus, if we combine (2.19)-(2.21), by choosing  $\delta$  small enough we obtain (2.18). □

## 5 2.4 Proof of the observability estimate in 3-D

- 6 In this section we prove the following Carleman inequality, for the solution of the homogeneous 3-D
- 7 Boussinesq system assuming Hypothesis 1.2 or 1.3.

### 8 2.4.1 A Carleman inequality assuming Hypothesis 1.2

- 9 In this section we prove the following result:

10 **Proposition 2.10.** *Let  $\Omega$  be such that Hypothesis 1.2 holds,  $\omega \subset \Omega$  be a nonempty open set and let*  
 11  *$m \geq 2$ . Then, there are  $\varepsilon_0 > 0$ ,  $C > 0$  and  $\lambda_0 \geq 1$  such that if  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  $(\varphi^T, \psi^T) \in L^2(\Omega)^4$ ,*  
 12  *$\lambda \geq \lambda_0$ , and  $s \geq e^{C\lambda(T^m + T^{2m})}$ , we have:*

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-s\alpha} (|\psi|^2 + |\varphi_2|^2), \quad (2.22)$$

- 13 for the weights defined in (2.9) and  $(\varphi, \psi)$  the solution of (2.3).

1 **A geometrical result.** First of all we prove the following geometrical result:

2 **Proposition 2.11.** *Let  $\Omega$  be such that Hypothesis 1.2 holds. Then, for  $a_0 > 0$  small enough, there*  
 3 *is  $C > 0$  such that for any function  $w \in H^4(\Omega) \cap H_0^1(\Omega)$  and for any  $a \in [0, a_0]$  we have that:*

$$\|\partial_{x_1} w\|_{L^2(\Omega)} \leq C \left( \|\partial_{x_1 x_2} w\|_{H^2(\Omega)} + \|\partial_{x_1 x_3} w\|_{H^2(\Omega)} + \|L_a w\|_{H^1(\partial\Omega)} \right), \quad (2.23)$$

4 *with*

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w - a \partial_{x_3 x_3} w. \quad (2.24)$$

5 *Proof.* We recall that by Hypothesis 1.2 there are a domain  $\Omega_0 \subset \mathbb{R}^2$  and  $H_-, H_+ \in C^0(\overline{\Omega_0}, [\delta, +\infty))$   
 6 such that  $\Omega = \{x : -H_-(x_1, x_2) < x_3 < H_+(x_1, x_2)\}$ . Using Proposition 2.4 we obtain that there  
 7 are  $C > 0$  and  $a_0 > 0$  such that for all  $a \in [0, a_0]$  and  $s \in (-\delta, \delta)$  we have that:

$$\|\partial_{x_1} w\|_{C^0(\overline{\Omega_0} \times \{s\})} \leq C \left( \|\partial_{x_1 x_2} w\|_{H^2(\Omega_0 \times \{s\})} + \|\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w\|_{H^1(\partial\Omega_0 \times \{s\})} \right). \quad (2.25)$$

8 Considering that  $\partial_{x_3 x_3} w = 0$  on  $\partial\Omega_0 \times (-\delta, \delta)$  (because of Dirichlet boundary conditions) we have  
 9 that:

$$L_a w = -\partial_{x_1 x_1} w - a \partial_{x_2 x_2} w \quad \text{on } \Omega_0 \times (-\delta, \delta). \quad (2.26)$$

10 Thus, if we integrate (2.25) squared on  $(-\delta, \delta)$  and consider (2.26), we obtain that:

$$\|\partial_{x_1} w\|_{L^2(\Omega_0 \times (-\delta, \delta))}^2 \leq C \left( \|\partial_{x_1 x_2} w\|_{H^2(\Omega_0 \times (-\delta, \delta))}^2 + \|L_a w\|_{H^1(\partial\Omega_0 \times (-\delta, \delta))}^2 \right). \quad (2.27)$$

11 Finally, considering that for all  $x = (x_1, x_2, x_3) \in \Omega$  and  $s \in (-\delta, \delta)$  we have that:

$$\partial_{x_1} w(x_1, x_2, x_3) = \partial_{x_1} w(x_1, x_2, s) + \int_s^{x_3} \partial_{x_1 x_3} w(x_1, x_2, s') ds'. \quad (2.28)$$

12 By integrating (2.28), using (2.27) and remarking that  $\Omega_0 \times (-\delta, \delta) \subset \Omega$ , we easily obtain (2.23).  $\square$

13 *Proof of Proposition 2.10.* Because the proof is quite technical, we divide it into twelve steps.

- 14 • **Step 1:** We first use the coercivity estimate given in Proposition 2.11 for  $\varphi_1$ . We transform  
 15 this coercivity estimate into a weighted estimate of  $\varphi_1$ .
- 16 • **Step 2:** We add global terms involving  $\varphi_2$  and  $\varphi_3$  in both sides of the previous estimate.
- 17 • **Step 3:** We get rid of the terms involving  $\varphi_1$  in the right-hand side of the previous estimate  
 18 thanks to the equations satisfied by  $\varphi_2$  and  $\varphi_3$ .
- 19 • **Step 4:** We choose from this step forward to treat only one term in the right-hand side of  
 20 the estimate to simplify, i.e.  $\partial_{x_2 x_3} \varphi_2$ , all the other terms can be treated in a similar way. In  
 21 this step we use Lemma 2.5.

- 1 • **Step 5:** We begin with a Carleman estimate for the derivatives of  $\partial_{x_2x_3}\varphi_2$ , this Carleman  
2 estimate comes from Lemma 2.6.
- 3 • **Step 6:** The previous inequality leads to a global term involving the pressure  $\pi$ . This is why  
4 we also use a Carleman estimate for the derivatives of  $\partial_{x_2x_3x_2}\pi$ .
- 5 • **Step 7:** We then get rid of the trace terms containing  $\partial_{x_2x_3}\varphi_2$  and  $\partial_{x_2x_3x_2}\pi$  by using the  
6 regularity result stated in Lemma 2.7.
- 7 • **Step 8:** We eliminate the local term involving the pressure i.e.  $\partial_{x_2x_3x_2}\pi$  by the second  
8 equation of the system.
- 9 • **Step 9:** We gather Steps 4, 5, 6, 7 and 8.
- 10 • **Step 10:** We first estimate the local term involving the derivatives of  $\partial_{x_2x_3}\varphi_2$  by a local term  
11 involving  $\partial_{x_2x_3}\varphi_2$  by using standard integration by parts. We then estimate the local term of  
12  $\partial_{x_2x_3}\varphi_2$  in function of a local term of  $\varphi_2$ .
- 13 • **Step 11:** We apply the standard Carleman estimate for the heat equation satisfied by  $\psi$ .
- 14 • **Step 12:** We estimate the local term of  $\varphi_3$  by the equation satisfied by  $\psi$ .

15 First, let us fix three nonempty open sets  $\omega_1, \omega_2, \omega_3$  such that  $\omega_0 \subset\subset \omega_1 \subset\subset \omega_2 \subset\subset \omega_3 \subset\subset \omega$ .

16 **Step 1:** From (2.3), we readily see that the equation of  $\varphi_1$  on the boundary is given by

$$-\partial_{x_1x_1}\varphi_1 - \frac{\varepsilon}{1+\varepsilon}\partial_{x_2x_2}\varphi_1 - \frac{\varepsilon}{1+\varepsilon}\partial_{x_3x_3}\varphi_1 = \frac{1}{1+\varepsilon}(\partial_{x_1x_2}\varphi_2 + \partial_{x_1x_3}\varphi_3) \text{ on } \Sigma. \quad (2.29)$$

Thus, using Proposition 2.11 and Poincaré inequality we have that for every  $t \in [0, T)$ :

$$\begin{aligned} \|\varphi_1(t, \cdot)\|_{L^2(\Omega)} \leq C \left( \|\partial_{x_1x_2}\varphi_1(t, \cdot)\|_{H^2(\Omega)} + \|\partial_{x_1x_3}\varphi_1(t, \cdot)\|_{H^2(\Omega)} \right. \\ \left. + \|\partial_{x_1x_2}\varphi_2(t, \cdot)\|_{H^2(\Omega)} + \|\partial_{x_1x_3}\varphi_3(t, \cdot)\|_{H^2(\Omega)} \right). \quad (2.30) \end{aligned}$$

Therefore, we deduce that:

$$\begin{aligned} s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi_1|^2 \leq C \left( s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1x_2}\varphi_1|^2 + |\nabla^i \partial_{x_1x_3}\varphi_1|^2) \right. \\ \left. + s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1x_2}\varphi_2|^2 + |\nabla^i \partial_{x_1x_3}\varphi_3|^2) \right), \quad (2.31) \end{aligned}$$

17 for  $\alpha^*$  and  $\xi^*$  defined in (2.9). Note that the previous exponents: 51, 52 are chosen sufficiently big  
18 to absorb the trace terms, see Step 7. They are not optimal.

**Step 2:** We deduce from (2.31) the following estimate

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C \left( s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \right. \\ \left. + s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_3|^2 + |\varphi_2|^2 + |\varphi_3|^2) \right). \quad (2.32)$$

1 **Step 3:** From (2.3), the equations satisfied by  $\varphi_2$  and  $\varphi_3$  are

$$\begin{cases} -\frac{\varepsilon}{1+\varepsilon} \partial_t \varphi_2 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_1 x_1} \varphi_2 - \partial_{x_2 x_2} \varphi_2 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_3 x_3} \varphi_2 - \frac{1}{1+\varepsilon} (\partial_{x_1 x_2} \varphi_1 + \partial_{x_2 x_3} \varphi_3) = 0 \text{ in } Q, \\ -\frac{\varepsilon}{1+\varepsilon} \partial_t \varphi_3 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_1 x_1} \varphi_3 - \frac{\varepsilon}{1+\varepsilon} \partial_{x_2 x_2} \varphi_3 - \partial_{x_3 x_3} \varphi_3 - \frac{1}{1+\varepsilon} (\partial_{x_1 x_3} \varphi_1 + \partial_{x_2 x_3} \varphi_2) = 0 \text{ in } Q. \end{cases} \quad (2.33)$$

So we easily deduce that

$$s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \\ \leq C s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left( |\nabla^i \partial_{x_2 x_3} \varphi_3|^2 + |\nabla^i \partial_t \varphi_2|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_2|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_2|^2 \right. \\ \left. + |\nabla^i \partial_{x_2 x_3} \varphi_2|^2 + |\nabla^i \partial_t \varphi_3|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_3|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_3|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_3|^2 \right). \quad (2.34)$$

2 From (2.3), we also have that  $\varphi_2, \varphi_3$  satisfy

$$\forall i \in \{2, 3\}, -\partial_t \varphi_i - \Delta \varphi_i = -\partial_{x_i} \pi \text{ in } Q, \quad \varphi_i = 0 \text{ on } \Sigma. \quad (2.35)$$

Therefore, for every  $i = 2, 3$ , we have that

$$|\partial_t \varphi_i|^2 \leq C (|\Delta \varphi_i|^2 + |\partial_{x_i} \pi|^2) \leq C (|\partial_{x_1 x_1} \varphi_i|^2 + |\partial_{x_2 x_2} \varphi_i|^2 + |\partial_{x_3 x_3} \varphi_i|^2 + |\partial_{x_i} \pi|^2),$$

so from (2.34), we obtain

$$s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} (|\nabla^i \partial_{x_1 x_2} \varphi_1|^2 + |\nabla^i \partial_{x_1 x_3} \varphi_1|^2) \\ \leq C s^{51}\lambda^{52} \sum_{i=0}^2 \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left( |\nabla^i \partial_{x_2 x_3} \varphi_3|^2 + |\nabla^i \partial_{x_2} \pi|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_2|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_2|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_2|^2 \right. \\ \left. + |\nabla^i \partial_{x_2 x_3} \varphi_2|^2 + |\nabla^i \partial_{x_3} \pi|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_3|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_3|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_3|^2 \right). \quad (2.36)$$

We now gather (2.36) with (2.31) to deduce that

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C \left( s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} \left( |\varphi_2|^2 + |\varphi_3|^2 + |\nabla^i \partial_{x_2} \pi|^2 + |\nabla^i \partial_{x_3} \pi|^2 \right. \right. \\ \left. \left. + \sum_{i=0}^2 \sum_{j=2,3} |\nabla^i \partial_{x_1 x_2} \varphi_j|^2 + |\nabla^i \partial_{x_2 x_3} \varphi_j|^2 + |\nabla^i \partial_{x_1 x_1} \varphi_j|^2 + |\nabla^i \partial_{x_2 x_2} \varphi_j|^2 + |\nabla^i \partial_{x_3 x_3} \varphi_j|^2 \right) \right). \quad (2.37)$$

**Step 4:** We focus on the estimation of  $\partial_{x_2x_3}\varphi_2$ . We apply Lemma 2.5 to  $\partial_{x_2x_3}\varphi_2$  with  $k = 24$ ,  $i = 53$ , this leads to

$$\begin{aligned} & s^{55}\lambda^{56} \iint_Q e^{-2s\alpha\xi^{55}} |\partial_{x_2x_3}\varphi_2|^2 + s^{53}\lambda^{54} \iint_Q e^{-2s\alpha\xi^{53}} |\nabla\partial_{x_2x_3}\varphi_2|^2 + s^{51}\lambda^{52} \iint_Q e^{-2s\alpha\xi^{51}} |\nabla^2\partial_{x_2x_3}\varphi_2|^2 \\ & \quad + \cdots + s^7\lambda^8 \iint_Q e^{-2s\alpha\xi^7} |\nabla^{24}\partial_{x_2x_3}\varphi_2|^2 \\ & \leq C \left( s^5\lambda^6 \iint_Q e^{-2s\alpha\xi^5} |\nabla^{25}\partial_{x_2x_3}\varphi_2|^2 + s^{55}\lambda^{56} \iint_{Q_{\omega_0}} e^{-2s\alpha\xi^{55}} |\partial_{x_2x_3}\varphi_2|^2 \right. \\ & \quad \left. + s^7\lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha\xi^7} |\nabla^{24}\partial_{x_2x_3}\varphi_2|^2 \right). \end{aligned} \quad (2.38)$$

**Step 5:** We now apply the Carleman estimate given by Lemma 2.6 to the equation satisfied by  $\nabla^{25}\partial_{x_2x_3}\varphi_2$ , i.e.

$$-\partial_t(\nabla^{25}\partial_{x_2x_3}\varphi_2) - \Delta(\nabla^{25}\partial_{x_2x_3}\varphi_2) = -\nabla^{25}\partial_{x_2x_3}\partial_{x_2}\pi \text{ in } Q, \quad (2.39)$$

we obtain the following inequality

$$\begin{aligned} & s^5\lambda^6 \iint_Q e^{-2s\alpha\xi^5} |\nabla^{25}\partial_{x_2x_3}\varphi_2|^2 + s^3\lambda^4 \iint_Q e^{-2s\alpha\xi^3} |\nabla^{25}\partial_{x_2x_3}\varphi_2|^2 \\ & \leq C \left( s^2\lambda^2 \iint_Q e^{-2s\alpha} |\nabla^{25}\partial_{x_2x_3}\partial_{x_2}\pi|^2 + s^5\lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha\xi^5} |\nabla^{25}\partial_{x_2x_3}\varphi_2|^2 \right. \\ & \quad \left. + s^3\lambda^3 \iint_{\Sigma} e^{-2s\alpha\xi^3} |\partial_n \nabla^{25}\partial_{x_2x_3}\varphi_2|^2 \right). \end{aligned} \quad (2.40)$$

**Step 6:** In order to estimate the global term involving the pressure in the right-hand side of (2.40), we also employ a Carleman estimate for the pressure. Indeed, by taking the divergence in (2.3)<sub>1</sub> we can easily see that:

$$-\varepsilon\partial_t\pi - (1-\varepsilon)\Delta\pi = 0 \text{ in } Q,$$

so in particular

$$\forall i \in \{2, 3\}, \quad -\frac{\varepsilon}{1-\varepsilon}\partial_t(\partial_{x_i}\pi) - \Delta(\partial_{x_i}\pi) = 0 \text{ in } Q.$$

Consequently, from Lemma 2.6 we obtain that:

$$\begin{aligned} & s^2\lambda^3 \iint_Q e^{-2s\alpha\xi^2} |\nabla^{25}\partial_{x_2x_3}\partial_{x_2}\pi|^2 + \lambda \iint_Q e^{-2s\alpha} |\nabla^{26}\partial_{x_2x_3}\partial_{x_2}\pi|^2 \\ & \leq C \left( s^2\lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha\xi^2} |\nabla^{25}\partial_{x_2x_3}\partial_{x_2}\pi|^2 + s^{-1}\lambda^{-1} \iint_{\Sigma} e^{-2s\alpha\xi^{-1}} |\partial_n \nabla^{25}\partial_{x_2x_3}\partial_{x_2}\pi|^2 \right). \end{aligned} \quad (2.41)$$

**Step 7:** We now gather the estimates (2.40) and (2.41) and take  $\lambda, s$  sufficiently large to get

$$\begin{aligned}
& s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad s^2 \lambda^3 \iint_Q e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + \lambda \iint_Q e^{-2s\alpha} |\nabla^{26} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\
& \leq C \left( s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha} \xi^3 |\partial_n \nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad \left. + s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha} \xi^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \right). \quad (2.42)
\end{aligned}$$

From (2.11) and the regularity result stated in Lemma 2.7 with  $k = 16$  and  $r = 3/2$ , we deduce that for  $m \geq 2$ ,

$$\begin{aligned}
& \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \varphi\|_{H^{16,32}(Q)} + \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \pi\|_{H^{15,31}(Q)} \\
& \leq \|s^{3/2+16+16/m} e^{-s\alpha^*} (\xi^*)^{3/2+16+16/m} \varphi\|_{L^2(Q)} \leq \|s^{51/2} e^{-s\alpha^*} (\xi^*)^{51/2} \varphi\|_{L^2(Q)}. \quad (2.43)
\end{aligned}$$

From (2.11) and (2.43), we then obtain that

$$\begin{aligned}
& s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha} \xi^3 |\partial_n \nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha} \xi^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\
& = s^3 \lambda^3 \iint_{\Sigma} e^{-2s\alpha^*} (\xi^*)^3 |\partial_n \nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 + s^{-1} \lambda^{-1} \iint_{\Sigma} e^{-2s\alpha^*} (\xi^*)^{-1} |\partial_n \nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 \\
& \leq \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \varphi\|_{H^{16,32}(Q)}^2 + \|s^{3/2} e^{-s\alpha^*} (\xi^*)^{3/2} \pi\|_{H^{15,31}(Q)}^2 \\
& \leq \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2. \quad (2.44)
\end{aligned}$$

- 1 So by taking  $\lambda$  sufficiently large, the trace terms of the right hand side of (2.42) can be absorbed
- 2 by the left-hand side of (2.32).

**Step 8:** From (2.35) and (2.43), by integration by parts and cut-off arguments, we have that there exist  $r, r' > 0$  such that

$$\begin{aligned}
& s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^2 |\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi|^2 = -s^2 \lambda^3 \iint_{Q_{\omega_0}} e^{-2s\alpha} [\nabla^{25} \partial_{x_2 x_3} \partial_{x_2} \pi][(\partial_t - \Delta) \nabla^{25} \partial_{x_2 x_3} \varphi_2] \\
& \leq \|s^{1/2} e^{-s\alpha^*} (\xi^*)^{1/2} \pi\|_{H^{15,31}(Q)}^2 + s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha+2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha+2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2. \quad (2.45)
\end{aligned}$$

- 3 So for  $\lambda$  sufficiently large, the local term of the pressure in the right hand side of (2.42) can be
- 4 absorbed by the left-hand side of (2.32).

**Step 9:** By using (2.38), (2.42), (2.44) and (2.45), we have

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \left( \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad + s^{55} \lambda^{56} \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^7 \lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad \left. + s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \right). \quad (2.46)
\end{aligned}$$

**Step 10:** We now estimate the local term in  $\varphi_2$ , we proceed by standard integrations by parts, as in [BP20a, Section 5, Step 3], to obtain

$$\begin{aligned}
& s^r \lambda^{r'} \iint_{Q_{\omega_1}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^7 \lambda^8 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^7 |\nabla^{24} \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + s^5 \lambda^6 \iint_{Q_{\omega_0}} e^{-2s\alpha} \xi^3 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq \delta \left( s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \right. \\
& \quad \left. + \cdots + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \right) \\
& \quad + C_\delta s^r \lambda^{r'} \iint_{Q_{\omega_2}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\partial_{x_2 x_3} \varphi_2|^2, \quad (2.47)
\end{aligned}$$

for some other parameters  $r, r' > 0$ . By taking  $\delta$  sufficiently small in (2.47), we deduce from (2.46) the following estimate

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_2}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\partial_{x_2 x_3} \varphi_2|^2. \quad (2.48)
\end{aligned}$$

By integration by parts in the same spirit as before, we have

$$\begin{aligned}
& s^{55} \lambda^{56} \iint_Q e^{-2s\alpha} \xi^{55} |\partial_{x_2 x_3} \varphi_2|^2 + s^{53} \lambda^{54} \iint_Q e^{-2s\alpha} \xi^{53} |\nabla \partial_{x_2 x_3} \varphi_2|^2 + s^{51} \lambda^{52} \iint_Q e^{-2s\alpha} \xi^{51} |\nabla^2 \partial_{x_2 x_3} \varphi_2|^2 \\
& \quad + \cdots + s^5 \lambda^6 \iint_Q e^{-2s\alpha} \xi^5 |\nabla^{25} \partial_{x_2 x_3} \varphi_2|^2 + s^3 \lambda^4 \iint_Q e^{-2s\alpha} \xi^3 |\nabla^{26} \partial_{x_2 x_3} \varphi_2|^2 \\
& \leq C \lambda^3 s^{51} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + C s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha + 2s\alpha^*} \xi^r |\varphi|^2, \quad (2.49)
\end{aligned}$$

1 for other parameters  $r, r' > 0$ .

2 By using (2.37), (2.43), (2.49) and performing the same strategy from Step 5 to Step 10 for  
 3 other terms appearing in the right-hand side of (2.37), we obtain that there exist  $r, r' > 0$  such  
 4 that

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 \leq C s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha+2s\alpha^*} \xi^r (|\varphi_2|^2 + |\varphi_3|^2). \quad (2.50)$$

**Step 11:** From (2.3), we apply the classical Carleman estimate for the heat equation with homogenous Dirichlet boundary condition satisfied by  $\psi$ , i.e. we use Lemma 2.6,

$$s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 + s\lambda^2 \iint_Q e^{-3s\alpha} \xi |\nabla \psi|^2 \leq C \left( \iint_Q e^{-3s\alpha} |\varphi_3|^2 + s^3\lambda^4 \iint_{Q_{\omega_0}} e^{-3s\alpha} \xi^3 |\psi|^2 \right). \quad (2.51)$$

By using (2.13), we sum (2.50) and (2.51) to get

$$s^{51}\lambda^{52} \iint_Q e^{-2s\alpha^*} (\xi^*)^{51} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-3s\alpha} \xi^3 |\psi|^2 + s\lambda^2 \iint_Q e^{-3s\alpha} \xi |\nabla \psi|^2 \leq C \left( s^r \lambda^{r'} \iint_{Q_{\omega_3}} e^{-4s\alpha+2s\alpha^*} \xi^r (|\varphi_2|^2 + |\varphi_3|^2) + s^3\lambda^4 \iint_{Q_{\omega_0}} e^{-3s\alpha} \xi^3 |\psi|^2 \right). \quad (2.52)$$

5 **Step 12:** To eliminate the local term in  $\varphi_3$  in the right-hand side of (2.52), we proceed  
 6 exactly as in the proof of Proposition 2.8 in Section 2.3. Therefore, we obtain the desired estimate  
 7 (2.22). □

### 8 2.4.2 A Carleman inequality assuming Hypothesis 1.3

9 The goal of this part is to prove the following result:

10 **Proposition 2.12.** *Let  $\Omega, \omega$  be such that Hypothesis 1.3 holds and let  $m \geq 12$ . Then, there are*  
 11  *$\varepsilon_0 > 0$ ,  $C > 0$  and  $\lambda_0 \geq 1$  such that if  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  $(\varphi^T, \psi^T) \in (L^2(\Omega))^4$ ,  $\lambda \geq \lambda_0$ , and*  
 12  *$s \geq e^{C\lambda}(T^m + T^{2m})$ , we have:*

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \iint_{Q_\omega} e^{-s\alpha^*} (|\psi|^2 + |\varphi_2|^2), \quad (2.53)$$

13 for the weights defined in (2.9) and for  $(\varphi, \psi)$  the solution of (2.3).

14 To prove Proposition 2.12, we follow the strategy of [FCGIP06], which consists in getting a  
 15 Carleman estimate with some local terms and then using that the observation domain touches the  
 16 boundary. The fact that the coupling of the system is of order 2 is an additional difficulty with  
 17 respect to the systems treated in [FCGIP06].

1 *Proof of Proposition 2.12.* First of all, we consider that, by taking a smaller control domain if  
2 necessary, we can suppose that there are  $\gamma \in \text{Span}(e_2, e_3)$  and  $\tilde{\Gamma} \subset \Gamma$  a relative open set such that  
3  $\omega = \{x + \lambda\gamma : x \in \tilde{\Gamma}, \lambda \in (0, 1)\}$ . In addition, we consider

$$\omega_1 := \{x + \lambda\gamma : x \in \hat{\Gamma}, \lambda \in (0, 1/2)\}, \quad (2.54)$$

4 for  $\hat{\Gamma}$  a relative open set contained in  $\tilde{\Gamma}$ . Finally, we consider  $\omega_0$  a non empty open set compactly  
5 contained in  $\omega_1$ .

6 In order to prove the Carleman inequality (2.53), using (2.3), we first remark that  $\partial_{x_1}\varphi_1$  satisfies  
7 the equation:

$$\left(-\frac{\varepsilon}{1+\varepsilon}\partial_t - \Delta\right)\partial_{x_1}\varphi_1 = \frac{1}{1+\varepsilon}(\partial_{x_1x_1x_2}\varphi_2 + \partial_{x_1x_1x_3}\varphi_3 + \partial_{x_1x_2x_2}\varphi_1 + \partial_{x_1x_3x_3}\varphi_1). \quad (2.55)$$

To continue with, using Poincaré's inequality for  $\varphi_1$  and Lemma 2.5 for  $\partial_{x_1}\varphi_1$  with  $k = 1$ , we obtain that:

$$\begin{aligned} & s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*}(\xi^*)^{16}|\varphi_1|^2 + s^{16}\lambda^{17} \iint_Q e^{-2s\alpha}\xi^{16}|\partial_{x_1}\varphi_1|^2 + s^{14}\lambda^{15} \iint_Q e^{-2s\alpha}\xi^{14}|\nabla\partial_{x_1}\varphi_1|^2 \\ & \leq C\left(s^{12}\lambda^{13} \iint_Q e^{-2s\alpha}\xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 + s^{16}\lambda^{17} \iint_{Q_{\omega_0}} \xi^{16}|\partial_{x_1}\varphi_1|^2 \right. \\ & \quad \left. + s^{14}\lambda^{15} \iint_{Q_{\omega_0}} \xi^{14}|\nabla\partial_{x_1}\varphi_1|^2\right). \end{aligned} \quad (2.56)$$

In particular, by using (2.55) and the Carleman estimate coming from Lemma 2.6 for the terms of  $\nabla^2\partial_{x_1}\varphi_1$  we obtain that:

$$\begin{aligned} s^{12}\lambda^{13} \iint_Q e^{-2s\alpha}\xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 & \leq C\left(s^9\lambda^9 \iint_Q e^{-2s\alpha}\xi^9|\nabla^2g|^2 + s^{12}\lambda^{13} \iint_{Q_{\omega_0}} \xi^{12}|\nabla^2\partial_{x_1}\varphi_1|^2 \right. \\ & \quad \left. + s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha}\xi^{10}|\partial_n\nabla^2\partial_{x_1}\varphi_1|^2\right), \end{aligned} \quad (2.57)$$

8 for  $g := \partial_{x_1x_1x_2}\varphi_2 + \partial_{x_1x_1x_3}\varphi_3 + \partial_{x_1x_2x_2}\varphi_1 + \partial_{x_1x_3x_3}\varphi_1$ .

To continue with, we find from Poincaré's inequality and Lemma 2.5 with  $k = 0$  that:

$$\begin{aligned} & s^{17}\lambda^{18} \iint_Q e^{-2s\alpha^*}(\xi^*)^{17}(|\varphi_2|^2 + |\varphi_3|^2) + s^{17}\lambda^{18} \iint_Q e^{-2s\alpha}\xi^{17}(|\partial_{x_1}\varphi_2|^2 + |\partial_{x_1}\varphi_3|^2) \\ & \leq C\left(s^{15}\lambda^{16} \iint_Q e^{-2s\alpha}\xi^{15}(|\nabla\partial_{x_1}\varphi_2|^2 + |\nabla\partial_{x_1}\varphi_3|^2) \right. \\ & \quad \left. + s^{17}\lambda^{18} \iint_{Q_{\omega_0}} e^{-2s\alpha}\xi^{17}(|\partial_{x_1}\varphi_2|^2 + |\partial_{x_1}\varphi_3|^2)\right). \end{aligned} \quad (2.58)$$

Next, we remark that we can follow the steps 3-11 of Section 2.4.1 and by using the estimates on the weights (2.12), (2.13), Lemmas 2.6 and 2.7 and (2.56), (2.57), (2.58), we deduce that:

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \left( \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha} (|\psi|^2 + |\varphi_2|^2) \right. \\ \left. \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 + s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha} \xi^{10} |\partial_n \nabla^2 \partial_{x_1}\varphi_1|^2 \right). \quad (2.59)$$

1 *Remark 2.13.* It helps us to have only derivatives of  $\partial_{x_1}\varphi_2$  and  $\partial_{x_1}\varphi_3$  in the right-hand side of  
2 (2.58) because we eventually do a Carleman inequality for  $\nabla^i \partial_{x_1}\varphi$  as in [BP20a, Section 5], so the  
3 local term that we get is of the form  $\nabla^i \partial_{x_1}\varphi$ . If instead of having  $\partial_{x_1}\varphi_2$  and  $\partial_{x_1}\varphi_3$  we had  $\nabla\varphi_2$   
4 and  $\nabla\varphi_3$ , after applying the Carleman estimate to  $\nabla^i\varphi$  we would get a local term which has  $\partial_{x_2}^i\varphi_1$   
5 and  $\partial_{x_3}^i\varphi_1$  terms from which we do not know how to get a Carleman inequality with a local term  
6 that only depends on  $\varphi_2$  and  $\psi$ .

Let us now treat the boundary term of  $\varphi_1$  as in Section 2.4.1. By interpolation and Lemma 2.7 we have:

$$s^{10}\lambda^{10} \iint_{\Sigma} e^{-2s\alpha} \xi^{10} |\partial_n \nabla^2 \partial_{x_1}\varphi_1|^2 \\ \leq C \|s^{23/4}\lambda^{23/4} e^{-s\alpha^*} (\xi^*)^{23/4} \varphi\|_{L^2(0,T;H^4(\Omega))}^{1/2} \|s^{19/4}\lambda^{19/4} e^{-s\alpha^*} (\xi^*)^{19/4} \varphi\|_{L^2(0,T;H^6(\Omega))}^{3/2} \\ \leq C \|s^{31/4+2/m}\lambda^{23/4} e^{-s\alpha^*} (\xi^*)^{23/4} \varphi\|_{L^2(Q)}^{1/2} \|s^{31/4+3/m}\lambda^{19/4} e^{-s\alpha^*} (\xi^*)^{31/4+3/m} \varphi\|_{L^2(Q)}^{3/2}.$$

So, if  $m \geq 12$ , we have that the trace term in the right-hand side of (2.59) can be absorbed by the left-hand side of (2.59), then we obtain:

$$s^{16}\lambda^{17} \iint_Q e^{-2s\alpha^*} (\xi^*)^{15} |\varphi|^2 + s^3\lambda^4 \iint_Q e^{-\frac{201}{100}s\alpha} \xi^3 |\psi|^2 \leq C \left( \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha} (|\psi|^2 + |\varphi_2|^2) \right. \\ \left. + \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 \right). \quad (2.60)$$

7 Let us now estimate the third right-hand side term of (2.60). From (2.54), recalling that  $e_1$   
8 is tangent to every point of  $\tilde{\Gamma}$  by Hypothesis 1.3,  $\partial_{x_1}\varphi_1 \equiv 0$  in  $\partial\omega_1 \cap \partial\Omega$ , and from Poincaré's  
9 inequality we obtain that:

$$\iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} |\partial_{x_1}\varphi_1|^2 \leq C \iint_{Q_{\omega_1}} e^{-\frac{31}{16}s\alpha^*} (|\partial_{x_2x_1}\varphi_1|^2 + |\partial_{x_3x_1}\varphi_1|^2). \quad (2.61)$$

10 Consequently, by using the equations (2.33), we can estimate the local terms in the right-hand side  
11 of (2.61) in function of local terms of  $\varphi_2$  and  $\varphi_3$ . By putting this in (2.60), we bound the local  
12 term in  $\partial_{x_1}\varphi_1$  by local terms in  $\varphi_2$  and  $\varphi_3$ . In addition, we can deal with the local term of  $\varphi_3$  by  
13 using the equation satisfied by  $\psi$  as before. Hence, by straightforward computations, we obtain the  
14 expected estimate (2.53).  $\square$

## 1 2.5 Source term method

2 In this section we adapt the source term method of [LTT13] to our case. It is worth mentioning  
3 that we use the uniform analyticity of the penalized semigroup, that come from uniform maximal  
4 regularity estimates of the linearized Boussinesq system (2.1).

5 From Proposition 2.1 we have an estimate for the control cost in  $L^2$  of system (2.1). We now fix  
6  $M > 0$  such that  $K(T) \leq Me^{M/T^m}$ , with  $K(T)$  defined as in (2.2). In addition, we fix the values:

$$q \in (1, \sqrt[2m]{2}), \quad p > q^{2m}/(2 - q^{2m}), \quad (2.62)$$

7 and the weights:

$$\rho_0(t) := M^{-p} \exp\left(-\frac{Mp}{(q-1)^m(T-t)^m}\right), \quad (2.63)$$

8

$$\rho_{\mathcal{F}}(t) := M^{-1-p} \exp\left(-\frac{(1+p)q^{2m}M}{(q-1)^m(T-t)^m}\right). \quad (2.64)$$

9 We introduce the linearized Boussinesq system, with source term  $f = (\tilde{f}, f_{N+1}) \in L^2(Q)^{N+1}$ ,

$$\begin{cases} y_t - \Delta y + \nabla p = \theta e_N + \tilde{f} + \tilde{v}1_\omega & \text{in } Q, \\ \theta_t - \Delta \theta = f_{N+1} + v_{N+1}1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, \theta = 0 & \text{on } \Sigma, \\ y(0) = y^0, \theta(0) = \theta^0 & \text{in } \Omega. \end{cases} \quad (2.65)$$

In addition, we define associated spaces for the source term, the state and the control, respectively:

$$\mathcal{F} := \left\{ f \in L^2(Q)^{N+1} : \frac{f}{\rho_{\mathcal{F}}} \in L^2(Q)^{N+1} \right\}, \quad (2.66)$$

$$\mathcal{Y} := \left\{ (y, \theta) \in L^2(Q)^{N+1} : \frac{(y, \theta)}{\rho_0} \in L^2(Q)^{N+1} \right\}, \quad (2.67)$$

$$\mathcal{V} := \left\{ v \in L^2(Q_\omega)^{N+1} : \frac{v}{\rho_0} \in L^2(Q_\omega)^{N+1} \text{ and } v \text{ satisfies (1.2)} \right\}. \quad (2.68)$$

10 From an adaptation of the proof of [LTT13, Proposition 2.3], since the linearized Boussinesq  
11 semigroup is analytic, we deduce the null controllability of the system (2.65) for source terms  $f$   
12 that exponentially decrease at  $t = T$ .

**Proposition 2.14.** *Let  $\Omega, \omega$  be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist  $\varepsilon_0 > 0$ ,  $m \geq 1$  and  $C > 0$  such that for every  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  $f \in \mathcal{F}$ ,  $(y^0, \theta^0) \in L^2(\Omega)^{N+1}$ , there exists a control  $v \in \mathcal{V}$  such that the solution  $(y, \theta)$  of (2.65) belongs to  $\mathcal{Y}$  and we have the following estimate:*

$$\|(y, \theta)/\rho_0\|_{C([0,T];L^2(\Omega))} + \|v/\rho_0\|_{L^2(Q_\omega)} \leq C (\|(y^0, \theta^0)\|_{L^2(\Omega)} + \|f/\rho_{\mathcal{F}}\|_{L^2(Q)}). \quad (2.69)$$

1 *In particular,  $(y, \theta)(T) = 0$ .*

The next proposition gives more information on the regularity of the controlled trajectory obtained in Proposition 2.14. We consider the weight:

$$\rho(t) = \exp\left(-\frac{M\beta}{(q-1)^m(T-t)^m}\right), \text{ with } \frac{(1+p)q^{2m}}{2} < \beta < p.$$

2 We remark that  $\rho(T) = 0$  and that  $\rho$  satisfies the inequalities:

$$\rho_0 \leq C\rho, \quad \rho_{\mathcal{F}} \leq C\rho, \quad |\rho'| \rho_0 \leq C\rho^2, \quad \rho^2 \leq C\rho_{\mathcal{F}}. \quad (2.70)$$

3 **Proposition 2.15.** *Let  $\Omega, \omega$  be such that Hypothesis 1.1, 1.2 or 1.3 holds. Then, there exist  $\varepsilon_0 > 0$ ,*  
 4  *$m \geq 1$  and  $C > 0$  such that for every  $T > 0$ ,  $\varepsilon \in (0, \varepsilon_0)$ ,  $f \in \mathcal{F}$ ,  $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$ , there exists*  
 5 *a control  $v \in \mathcal{V}$  such that the solution  $(y, \theta)$  of (2.65) satisfies:*

$$\|(y, \theta)/\rho\|_{L^2(0,T;H^2(\Omega)) \cap H^1(0,T;L^2(\Omega))} + \|v/\rho_0\|_{L^2(Q_\omega)} \leq C \left( \|(y^0, \theta^0)\|_{H_0^1(\Omega)} + \|f/\rho_{\mathcal{F}}\|_{L^2(Q)} \right). \quad (2.71)$$

6 *In particular,  $(y, \theta)(T) = 0$ .*

7 The proof of Proposition 2.15 is a straightforward adaptation of the proof of [LTT13, Proposition  
 8 2.8] by using uniform maximal regularity estimates of the penalized Stokes system (see Lemma 2.3)  
 9 and maximal regularity estimate of the heat equation.

10 *Remark 2.16.* For each  $f \in \mathcal{F}$  and  $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$ , by classical arguments (see [LTT13,  
 11 Proposition 2.9]), we can fix a control  $v \in \mathcal{V}$  such that  $(y, \theta)$  and  $v$  satisfy (2.71) by choosing  
 12 among those the unique minimizer of the functional  $v \mapsto \|(y, \theta)/\rho\|_{L^2(0,T;H^2(\Omega)) \cap H^1(0,T;L^2(\Omega))}^2 +$   
 13  $\|v/\rho_0\|_{L^2(Q_\omega)}^2$ .

### 14 **3 Local null controllability of the nonlinear Boussinesq system**

15 In this section we conclude the proof of Theorem 1.2. It is organized as follows:

- 16 • In Section 3.1 we use a fixed point argument to get the local null controllability of the nonlinear  
 17 system (1.1) for initial data in  $H_0^1(\Omega)^{N+1}$ .
- 18 • In Section 3.2 we use a regularization argument to deduce the local null controllability of  
 19 (1.1) for initial data in  $L^2(\Omega)^{N+1}$ .

### 1 3.1 Fixed point argument

2 In this section, we give the proof of Theorem 1.2 for initial data in  $H_0^1(\Omega)^{N+1}$ . In this part,  $C$   
3 denotes a positive constant that depends on  $\Omega$ ,  $\omega$  and  $T$  but independent of  $\varepsilon$  and varying from  
4 line to line. In addition, we denote:

$$\mathcal{F}_r := \{f \in \mathcal{F} : \|f/\rho_{\mathcal{F}}\|_{L^2(Q)} \leq r\}, \quad (3.1)$$

5 for  $r > 0$  a small enough parameter independent of  $\varepsilon$  that will be determined later.

6 We now fix  $(y^0, \theta^0) \in H_0^1(\Omega)^{N+1}$  such that:

$$\|(y^0, \theta^0)\|_{H_0^1(\Omega)} \leq r. \quad (3.2)$$

7 It follows that we can define an operator  $\mathcal{N}$  acting on  $\mathcal{F}_r$  by:

$$\mathcal{N}(f) := \left( -(y \cdot \nabla)y - \frac{1}{2}(\nabla \cdot y)y, -y \cdot \nabla\theta - \frac{1}{2}(\nabla \cdot y)\theta \right), \quad (3.3)$$

where  $(y, \theta)$  is the corresponding trajectory of (2.65) associated to the initial values  $(y^0, \theta^0)$ , the force  $f$  and the control  $v$  given by Proposition 2.15 and Remark 2.16.

To conclude the proof of Theorem 1.2 for regular initial data it suffices to check that, for  $r > 0$  small enough not depending on  $\varepsilon$ ,  $\mathcal{N}$  is a contractive mapping from  $\mathcal{F}_r$  into itself and then apply the Banach fixed point theorem.

**Step 1:  $\mathcal{F}_r$  is invariant for  $\mathcal{N}$  provided that  $r$  is small enough.** By using (2.70)<sub>4</sub> and the Sobolev embeddings  $H^1(\Omega) \hookrightarrow L^4(\Omega)$  and  $H^2(\Omega) \hookrightarrow W^{1,4}(\Omega)$  because  $N \leq 3$ , we have for almost every  $t \in (0, T)$ :

$$\begin{aligned} \left\| \frac{\mathcal{N}(f)}{\rho_{\mathcal{F}}} (t) \right\|_{L^2(\Omega)} &\leq C \left| \frac{\rho^2}{\rho_{\mathcal{F}}} (t) \right| \left( \left\| \frac{(y \cdot \nabla)y}{\rho^2} (t) \right\|_{L^2(\Omega)} + \left\| \frac{(\nabla \cdot y)y}{\rho^2} (t) \right\|_{L^2(\Omega)} \right. \\ &\quad \left. + \left\| \frac{y \cdot \nabla\theta}{\rho^2} (t) \right\|_{L^2(\Omega)} + \left\| \frac{(\nabla \cdot y)\theta}{\rho^2} (t) \right\|_{L^2(\Omega)} \right) \\ &\leq C \left( \left\| \frac{y}{\rho} (t) \right\|_{L^4(\Omega)} \left\| \frac{\nabla y}{\rho} (t) \right\|_{L^4(\Omega)} + \left\| \frac{y}{\rho} (t) \right\|_{L^4(\Omega)} \left\| \frac{\nabla\theta}{\rho} (t) \right\|_{L^4(\Omega)} \right. \\ &\quad \left. + \left\| \frac{\nabla y}{\rho} (t) \right\|_{L^4(\Omega)} \left\| \frac{\theta}{\rho} (t) \right\|_{L^4(\Omega)} \right) \\ &\leq C \left( \left\| \frac{y}{\rho} (t) \right\|_{H^1(\Omega)} \left\| \frac{y}{\rho} (t) \right\|_{H^2(\Omega)} + \left\| \frac{y}{\rho} (t) \right\|_{H^1(\Omega)} \left\| \frac{\theta}{\rho} (t) \right\|_{H^2(\Omega)} \right. \\ &\quad \left. + \left\| \frac{y}{\rho} (t) \right\|_{H^2(\Omega)} \left\| \frac{\theta}{\rho} (t) \right\|_{H^1(\Omega)} \right). \end{aligned}$$

Then, by integrating in the time interval  $t \in (0, T)$  and by using (2.71), (3.2) and (3.1), we have:

$$\begin{aligned} \left\| \frac{\mathcal{N}(f)}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)} &\leq C \left( \left\| \frac{y}{\rho} \right\|_{C([0, T]; H_0^1(\Omega))} \left( \left\| \frac{y}{\rho} \right\|_{L^2(0, T; H^2(\Omega))} + \left\| \frac{\theta}{\rho} \right\|_{L^2(0, T; H^2(\Omega))} \right) \right) \\ &\quad + C \left\| \frac{\theta}{\rho} \right\|_{C([0, T]; H_0^1(\Omega))} \left\| \frac{y}{\rho} \right\|_{L^2(0, T; H^2(\Omega))} \\ &\leq C \left( \|(y^0, \theta^0)\|_{H_0^1(\Omega)}^2 + \|f/\rho_{\mathcal{F}}\|_{L^2(Q)}^2 \right) \\ &\leq Cr^2. \end{aligned}$$

So, for  $r > 0$  small enough,  $\mathcal{N}$  stabilizes  $\mathcal{F}_r$ .

**Step 2:  $\mathcal{N}$  is contracting provided that  $r$  is small enough.** Using the same kind of arguments, it is not difficult to obtain that:

$$\left\| \frac{\mathcal{N}(f^1) - \mathcal{N}(f^2)}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)} \leq Cr \left\| \frac{f^1 - f^2}{\rho_{\mathcal{F}}} \right\|_{L^2(Q)}.$$

- 1 Consequently, by taking  $r$  sufficiently small,  $\mathcal{N}$  is a contracting mapping on the closed ball  $\mathcal{F}_r$ .
- 2 Therefore, by the Banach fixed point theorem,  $\mathcal{N}$  has a unique fixed point  $f$ . By denoting  $(y, \theta, v)$
- 3 the associated trajectory to  $f$ , we find that  $(y, \theta, v)$  satisfies the system (1.1) and  $(y, \theta)(T) = 0$ .
- 4 Remark that  $r$  does not depend on  $\varepsilon$ , so the control is bounded uniformly when  $\varepsilon$  goes to 0. This
- 5 concludes the proof.

### 6 3.2 Smoothing effect of the nonlinear Boussinesq system

- 7 In this section, we give the proof of Theorem 1.2 for initial data in  $L^2(\Omega)^{N+1}$ . This type of
- 8 arguments have already been used in [CMS20] and [CSFCB<sup>+</sup>20]. The key remark is the following
- 9 regularity lemma.

**Lemma 3.1.** *Let  $T > 0$ . There exists a positive constant  $C_T > 0$  such that for every  $\varepsilon > 0$ , any  $(y_0, \theta_0) \in L^2(\Omega)^{N+1}$  and any weak solution of (1.1) with control  $v \equiv 0$ ,*

$$\exists t_0 \in [0, T], \|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq C_T \|(y_0, \theta_0)\|_{L^2(\Omega)}.$$

*Proof.* We multiply (1.1)<sub>1</sub> by  $y$  and (1.1)<sub>2</sub> by  $\theta$ , then integrate in  $\Omega$ , we find

$$\begin{aligned} \frac{d}{dt} \left( \int_{\Omega} y(t, \cdot)^2 \right) + \int_{\Omega} |\nabla y(t, \cdot)|^2 + \frac{1}{\varepsilon} \int_{\Omega} (\nabla \cdot y(t, \cdot))^2 &= \int_{\Omega} \theta(t, \cdot) y_N(t, \cdot), \\ \frac{d}{dt} \left( \int_{\Omega} \theta(t, \cdot)^2 \right) + \int_{\Omega} |\nabla \theta(t, \cdot)|^2 &= 0. \end{aligned}$$

We sum, use Young's inequality and Gronwall's lemma to obtain that there exists  $C > 0$  such that for all  $t \in [0, T]$ :

$$\|(y, \theta)(t, \cdot)\|_{L^2(\Omega)}^2 + \int_0^t \|(y, \theta)(s, \cdot)\|_{H^1(\Omega)}^2 ds \leq C \|(y_0, \theta_0)\|_{L^2(\Omega)}^2.$$

In particular, this gives us the existence of a time  $t_0 \in [0, T]$  such that

$$\|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq \sqrt{\frac{C}{T}} \|(y_0, \theta_0)\|_{L^2(\Omega)}.$$

1 This concludes the proof with  $C_T = \sqrt{C/T}$ . □

2 We now have all the tools to end the proof of Theorem 1.2.

3 *Proof of Theorem 1.2.* We divide the control strategy into three steps.

4 **Step 1: regularization of the data.** By setting  $v \equiv 0$  and by using Lemma 3.1, we deduce  
5 that there exists  $t_0 \in (0, T/2)$  such that  $\|(y, \theta)(t_0, \cdot)\|_{H^1(\Omega)} \leq C_{T/2} \|(y_0, \theta_0)\|_{L^2(\Omega)}$ .

6 **Step 2: local null controllability result in time  $T/2$ .** By taking  $(y_0, \theta_0) \in L^2(\Omega)^{N+1}$   
7 such that  $C_{T/2} \|(y_0, \theta_0)\|_{L^2(\Omega)} \leq \delta_{T/2}$ , where  $\delta_{T/2}$  is the radius of local null controllability of (1.1)  
8 for initial data in  $H_0^1(\Omega)^{N+1}$ , we obtain that there exists a control  $v \in L^2((t_0, t_0 + T/2) \times \omega)^{N+1}$   
9 satisfying (1.2) such that  $(y, \theta)(t_0 + T/2, \cdot) = 0$ .

10 **Step 3: do nothing at the end of the time interval.** We set  $v \equiv 0$  in  $(t_0 + T/2, T) \times \omega$ ,  
11 so  $(y, \theta)(T, \cdot) = 0$ . □

## 12 4 Remarks and open problems

13 In this section, we make some remarks and formulate some open problems concerning the null  
14 controllability of the penalized Boussinesq system (1.1).

- 15 • If  $N = 3$  by symmetry and by adapting Hypotheses 1.2 or 1.3, we can construct controls  
16 which satisfy:

$$\lambda_1 v_1 + \lambda_2 v_2 = v_3 = 0$$

17 for any  $\lambda_1, \lambda_2 \in \mathbb{R}$ . Indeed, all the proofs in this paper can be adapted to these situations by  
18 a simple change of coordinates.

- We can prove analogue controllability results assuming Hypothesis 1.1, 1.2 or 1.3 for the penalized Navier-Stokes system, which we recall is given by:

$$\begin{cases} y_t - \Delta y + (y \cdot \nabla)y + \frac{1}{2}(\nabla \cdot y)y + \nabla p = v1_\omega & \text{in } Q, \\ \nabla \cdot y = -\varepsilon p & \text{in } Q, \\ y = 0, & \text{on } \Sigma, \\ y(0) = y^0, & \text{in } \Omega. \end{cases} \quad (4.1)$$

Indeed, by following the strategy of this paper and omitting the steps related to the heat equation satisfied by  $\theta$ , we can prove that (4.1) is locally null controllable uniformly on  $\varepsilon$  with a control  $v$  with one null component. Obtaining the local null controllability of (4.1) with a control with two null-components in 3-D is an interesting open problem. A good strategy seems to employ the return method in the spirit of [CL14].

- The exponents of  $s$ ,  $\lambda$  and  $\xi$  and the constant  $m$  stated in Propositions 2.8, 2.10 and 2.12 are a bit arbitrary. In particular, by combining Lemmas 2.5 and 2.7 a sufficient number of times we can get an analogous result for any  $m > 1$  by choosing the exponent of  $s$ ,  $\lambda$  and  $\xi$  large enough. This implies that the cost of the null controllability of the linearized system without a source term is less than  $Ce^{CT-m}$  for all  $m > 1$ .
- Removing the geometrical hypothesis on  $\Omega$  is an interesting open problem.
- There are still plenty of systems with penalizations which approximate the incompressibility condition (see [She97]) whose null controllability properties have not been studied yet.

## A Proof of Lemma 2.7

Let us prove Lemma 2.7. The proof consists in two steps: first we use parabolic regularity estimates and then we estimate the weights.

As a first step, we prove the following result:

**Lemma A.1.** *Let  $k \in \mathbb{N}$  and  $h \in H_0^k(0, T)$ . Then, there is  $C > 0$  such that for all  $\varphi^T \in L^2(\Omega)$ :*

$$\|h\varphi\|_{H^{k,2k}(Q)} + \|h\pi\|_{H^{k-1,2k-1}(Q)} \mathbf{1}_{k \geq 1} \leq C \|h^{(k)}\varphi\|_{L^2(Q)}, \quad (\text{A.1})$$

for  $(\varphi, \pi)$  the solution of (2.17).

- 1 *Proof of Lemma A.1.* The proof of Lemma A.1 is done by induction. The base case,  $k = 0$ , is trivial.  
2 Let us now prove the inductive case. By hypothesis we have (A.1) for any function  $h \in H_0^k(0, T)$   
3 and we have to prove the estimate:

$$\|h\varphi\|_{H^{k+1, 2(k+1)}(Q)} + \|h\pi\|_{H^{k, 2k+1}(Q)} \leq C \|h^{(k+1)}\varphi\|_{L^2(Q)}, \quad (\text{A.2})$$

for any function  $h \in H_0^{k+1}(0, T)$ . We have that  $(h\varphi, h\pi)$  satisfies:

$$\begin{cases} -(h\varphi)_t - \Delta(h\varphi) + \nabla(h\pi) = -h'\varphi & \text{in } Q, \\ \varepsilon h\pi + \nabla \cdot (h\varphi) = 0 & \text{in } Q, \\ h\varphi = 0 & \text{on } \Sigma, \\ (h\varphi)(T, \cdot) = 0 & \text{in } \Omega. \end{cases}$$

- 4 Thus, since  $h$  and its derivatives vanish at  $t = T$ , Lemma 2.3 implies that:

$$\|h\varphi\|_{H^{k+1, 2(k+1)}(Q)} + \|h\pi\|_{H^{k, 2k+1}(Q)} \leq C \|h'\varphi\|_{H^{k, 2k}(Q)}. \quad (\text{A.3})$$

- 5 As  $h' \in H_0^k(0, T)$  we can now use the inductive hypothesis (A.1) and obtain (A.2) from (A.3).  $\square$

- 6 Using Lemma A.1 with  $h = s^r e^{-s\alpha^*} (\xi^*)^r$  we find that:

$$\|s^r e^{-s\alpha^*} (\xi^*)^r \varphi\|_{H^{k, 2k}(Q)} + \|s^r e^{-s\alpha^*} (\xi^*)^r \pi\|_{H^{k-1, 2k-1}(Q)} \leq \|\partial_t^k (s^r e^{-s\alpha^*} (\xi^*)^r) \varphi\|_{L^2(Q)}. \quad (\text{A.4})$$

- 7 To conclude, we have to estimate the time derivative of the weights. We first recall the following  
8 result, which is classical, whose proof is sketched for completeness:

- 9 **Lemma A.2.** *Let  $k \in \mathbb{N}_*$ . Then, for all  $s \geq e^{C\lambda} T^m$  we have the following estimate:*

$$|\partial_t^k \xi^*| + |\partial_t^k \alpha^*| \leq s^{k/m} (\xi^*)^{1+k/m}. \quad (\text{A.5})$$

- 10 *Sketch of the proof of Lemma A.2.* It is easy to prove by induction that:

$$\partial_t^k \xi^* = \frac{q(t, T)}{(t(T-t))^{m+k}}, \quad \partial_t^k \alpha^* = \frac{q(t, T)(e^{2\lambda\|\eta\|_\infty} - 1)}{(t(T-t))^{m+k}}. \quad (\text{A.6})$$

- 11 for  $q$  a homogeneous polynomial of degree  $k$ . Thus, considering that  $t \leq T \leq e^{-C\lambda} s^{1/m}$  as  $s \geq$   
12  $e^{C\lambda} T^m$ , we easily obtain (A.5) from (A.6).  $\square$

Finally, (A.5) and  $s\xi^* \geq 1$  imply the estimate:

$$\begin{aligned} \partial_t^k (s^r e^{-s\alpha^*} (\xi^*)^r) &\leq C s^r \sum_{i_1+\dots+i_{k+1}=k} \left( \prod_{j=1,\dots,k:i_j \neq 0} s \partial_t^{i_j} \alpha^* \right) e^{-s\alpha^*} \partial_t^{i_{k+1}} [(\xi^*)^r] \\ &\leq s^{r+k+k/m} (\xi^*)^{r+k+k/m} e^{-s\alpha^*}. \end{aligned} \quad (\text{A.7})$$

- 1 Indeed, we get the maximum exponent for  $s$  and  $\xi^*$  in (A.7) by picking  $j_1, \dots, j_k = 1$  and  $j_{k+1} = 0$ .  
 2 Consequently, we obtain the desired estimate (2.16) from (A.4) and (A.7).

### 3 References

- 4 [AKBGBdT11] F. Ammar-Khodja, A. Benabdallah, M. González Burgos, and L. de Teresa. Recent  
 5 results on the controllability of linear coupled parabolic problems: a survey. *Math.*  
 6 *Control Relat. F.*, 1(3):267–306, 2011.
- 7 [Bad11] M. Badra. Global Carleman inequalities for Stokes and penalized Stokes equations.  
 8 *Math. Control Relat. F.*, 1(2):149–175, 2011.
- 9 [Ber78] M. Bercovier. Perturbation of mixed variational problems. Application to mixed  
 10 finite element methods. *ESAIM Math. Model. Num.*, 12(3):211–236, 1978.
- 11 [BM20] K. Beauchard and F. Marbach. Unexpected quadratic behaviors for the small-time  
 12 local null controllability of scalar-input parabolic equations. *J. Math. Pure. Appl.*,  
 13 136:22–91, 2020.
- 14 [BP20a] J. A. Bárcena-Petisco. Null controllability of a penalized Stokes problem in dimen-  
 15 sion two with one scalar control. *Asymptotic Anal.*, 117(3-4):161–198, 2020.
- 16 [BP20b] J. A. Bárcena-Petisco. Uniform controllability of a Stokes problem with a transport  
 17 term in the zero-diffusion limit. *SIAM J. Control Optim.*, 58(3):1597–1625, 2020.
- 18 [Car12] N. Carreño. Local controllability of the n-dimensional boussinesq system with n-1  
 19 scalar controls in an arbitrary control domain. *Math. Control Relat. F.*, 2(4):361–  
 20 382, 2012.
- 21 [CG09] J.-M. Coron and S. Guerrero. Null controllability of the  $N$ -dimensional Stokes  
 22 system with  $N - 1$  scalar controls. *J. Differ. Equations*, 246(7):2908–2921, 2009.
- 23 [CL14] J.-M. Coron and P. Lissy. Local null controllability of the three-dimensional Navier-  
 24 Stokes system with a distributed control having two vanishing components. *Invent.*  
 25 *Math.*, 198(3):833–880, 2014.

- 1 [CMS20] J.-M. Coron, F. Marbach, and F. Sueur. Small-time global exact controllability of  
2 the Navier-Stokes equation with Navier slip-with-friction boundary conditions. *J.*  
3 *Eur. Math. Soc.*, 22(5):1625–1673, 2020.
- 4 [CnG13] N. Carreño and S. Guerrero. Local null controllability of the  $N$ -dimensional Navier–  
5 Stokes system with  $N - 1$  scalar controls in an arbitrary control domain. *J. Math.*  
6 *Fluid Mech.*, 15:139–153, 2013.
- 7 [CnGG15] N. Carreño, S. Guerrero, and M. Gueye. Insensitizing controls with two van-  
8 ishing components for the three-dimensional Boussinesq system. *ESAIM:COCV*,  
9 21(1):73–100, 2015.
- 10 [Cor07] J.-M. Coron. *Control and nonlinearity*, volume 136 of *Mathematical Surveys and*  
11 *Monographs*. American Mathematical Society, Providence, RI, 2007.
- 12 [CSFCB<sup>+</sup>20] F. W. Chaves-Silva, E. Fernández-Cara, K. Le Balc’h, J. L. F. Machado, and D. A.  
13 Souza. Small-time global exact controllability to the trajectories for the viscous  
14 boussinesq system. *arXiv eprint: 2006.01682*, 2020.
- 15 [DL19] M. Duprez and P. Lissy. Bilinear local controllability to the trajectories of the  
16 Fokker-Planck equation with a localized control. *arXiv preprint arXiv:1909.02831*,  
17 2019.
- 18 [Eva10] L. C. Evans. *Partial Differential Equation*. American Mathematical Society, second  
19 edition, 2010.
- 20 [FCGBGP06] E. Fernández-Cara, M. González-Burgos, S. Guerrero, and J.-P. Puel. Null con-  
21 trollability of the heat equation with boundary Fourier conditions: the linear case.  
22 *ESAIM:COCV*, 12(3):442–465, 2006.
- 23 [FCGIP06] E. Fernández-Cara, S. Guerrero, O. Yu. Imanuvilov, and J.-P. Puel. Some control-  
24 lability results for the  $N$ -Dimensional Navier–Stokes and Boussinesq systems with  
25  $N - 1$  scalar controls. *SIAM J. Control Optim.*, 45(1):146–173, 2006.
- 26 [FCLdM16] E. Fernández-Cara, J. Limaco, and S. B. de Menezes. Controlling linear and semilin-  
27 ear systems formed by one elliptic and two parabolic PDEs with one scalar control.  
28 *ESAIM:COCV*, 22(4):1017–1039, 2016.
- 29 [FI96] A. V. Fursikov and O. Yu. Imanuvilov. *Controllability of evolution equations*.  
30 Number 34. Seoul National University, 1996.

- 1 [GM18] S. Guerrero and C. Montoya. Local null controllability of the  $N$ -dimensional  
2 Navier–Stokes system with nonlinear Navier-slip boundary conditions and  $N - 1$   
3 scalar controls. *J. Math. Pure. Appl.*, 113:37–69, 2018.
- 4 [Gue07] S. Guerrero. Controllability of systems of Stokes equations with one control force:  
5 existence of insensitizing controls. *Ann. I. H. Poincaré*, 24:1029–1054, 2007.
- 6 [GZ21] B. Geshkovski and E. Zuazua. Controllability of one-dimensional viscous free  
7 boundary flows. *SIAM J. Control Optim.*, 59(3):1830–1850, 2021.
- 8 [HSLB20] V. Hernández-Santamaría and K. Le Balc’h. Local null-controllability of a nonlocal  
9 semilinear heat equation. *Appl. Math. Opt.*, pages 1–49, 2020.
- 10 [Ima01] O. Yu. Imanuvilov. Remarks on exact controllability for the Navier-Stokes equa-  
11 tions. *ESAIM:COCV*, 6:39–72, 2001.
- 12 [IPY09] O. Yu. Imanuvilov, J.-P. Puel, and M. Yamamoto. Carleman estimates for  
13 parabolic equations with nonhomogeneous boundary conditions. *Chin. Ann. Math.*,  
14 30(4):333–378, 2009.
- 15 [LB20] K. Le Balc’h. Local controllability of reaction-diffusion systems around nonnegative  
16 stationary states. *ESAIM:COCV*, 26(55):1–32, 2020.
- 17 [Lio88] J.-L. Lions. Contrôlabilité exacte, perturbations et stabilisation de systemes dis-  
18 tribués, tome 1, RMA 8, 1988.
- 19 [LTT13] Y. Liu, T. Takahashi, and M. Tucsnak. Single input controllability of a simplified  
20 fluid-structure interaction model. *ESAIM:COCV*, 19(1):20–42, 2013.
- 21 [LZ96] J.-L. Lions and E. Zuazua. A generique uniqueness result for the Stokes system and  
22 its control theoretical consequences. *Partial differential equations and applications:  
23 Collected Papers in Honor of Carlo Pucci*, 177:221–235, 1996.
- 24 [MT18] S. Micu and T. Takahashi. Local controllability to stationary trajectories of a  
25 Burgers equation with nonlocal viscosity. *J. Differ. Equations*, 264(5):3664–3703,  
26 2018.
- 27 [OJ84] J. T. Oden and O.-P. Jacquotte. Stability of some mixed finite element methods  
28 for Stokesian flows. *Comput. Meth. Appl. Mat.*, 43(2):231–247, 1984.
- 29 [Rus78] D. L. Russell. Controllability and stabilizability theory for linear partial differential  
30 equations: recent progress and open questions. *Siam Rev.*, 20(4):639–739, 1978.

- 1 [She97] J. Shen. Pseudo-compressibility methods for the unsteady incompressible Navier-  
2 Stokes equations. *Proceedings of the 1994 Beijing symposium on nonlinear evolu-*  
3 *tion equations and infinite dynamical systems*, pages 68–78, 1997.
- 4 [Tak17] T. Takahashi. Boundary local null-controllability of the Kuramoto–Sivashinsky  
5 equation. *Math. Control Signal*, 29(1):1–21, 2017.
- 6 [Tem68] R. Temam. Une méthode d’approximation de la solution des équations des Navier-  
7 Stokes. *Bull. Soc. Math. France*, 96:115–152, 1968.
- 8 [Tem77] R. Temam. *Navier-Stokes equations. Theory and numerical analysis*. North Holland  
9 Publishing Company, first edition, 1977.
- 10 [Zua96] E. Zuazua. A uniqueness result for the linear system of elasticity and its control  
11 theoretical consequences. *SIAM J. Control. and Optim.*, 34(5):1473–1495, 1996.