



Note: The performance of CIPSI on the ground state electronic energy of benzene

Pierre-Francois Loos, Yann Damour, Anthony Scemama

► To cite this version:

Pierre-Francois Loos, Yann Damour, Anthony Scemama. Note: The performance of CIPSI on the ground state electronic energy of benzene. 2020. hal-02922278v1

HAL Id: hal-02922278

<https://hal.science/hal-02922278v1>

Preprint submitted on 3 Sep 2020 (v1), last revised 13 Oct 2020 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Note: The performance of CIPSI on the ground state electronic energy of benzene

Pierre-François Loos,^{1,*} Yann Damour,¹ and Anthony Scemama^{1,†}

¹Laboratoire de Chimie et Physique Quantiques (UMR 5626), Université de Toulouse, CNRS, UPS, France

Following the recent work of Eriksen *et al.* [arXiv:2008.02678 [physics.chem-ph]], we report the performance of the *Configuration Interaction using a Perturbative Selection made Iteratively* (CIPSI) method on the non-relativistic frozen-core correlation energy of the benzene molecule in the cc-pVDZ basis. Following our usual protocol, we obtain a correlation energy of $-863.4 \text{ m}E_h$ which agrees with the theoretical estimate of $-863 \text{ m}E_h$ proposed by Eriksen *et al.* using an extensive array of highly-accurate new electronic structure methods.

Although sometimes decried, one cannot deny the usefulness of benchmark sets and their corresponding reference data for the electronic structure community. These are indeed essential for the validation of existing theoretical models and to bring to light and subsequently understand their strengths and, more importantly, their weaknesses. In that regard, the previous benchmark datasets provided by the *Simons Collaboration on the Many-Electron Problem* have been extremely valuable.^{1–3} The same comment applies to the excited-state benchmark set of Thiel and coworkers.^{4–8} Following a similar goal, we have recently proposed a large set of highly-accurate vertical transition energies for various types of excited states thanks to the renaissance of selected configuration interaction (SCI) methods^{9–11} which can now routinely produce near full configuration interaction (FCI) quality excitation energies for small- and medium-sized organic molecules.^{12–16}

In a recent preprint,¹⁷ Eriksen *et al.* have proposed a blind test for a particular electronic structure problem inviting several groups around the world to contribute to this endeavour. In addition to coupled cluster theory with singles, doubles, triples, and quadruples (CCSDTQ),^{18,19} a large panel of highly-accurate, emerging electronic structure methods were considered: (i) the many-body expansion FCI (MBE-FCI),^{20–23} (ii) three SCI methods including a second-order perturbative correction (ASCI,^{24–26} iCI,²⁷ and SHCI^{28–30}), (iii) a selected coupled-cluster theory method which also includes a second-order perturbative correction (FCCR),³¹ (iv) the density-matrix renormalization group approach (DMRG),³² and (v) two flavors of FCI quantum Monte Carlo (FCIQMC),^{33,34} namely AS-FCIQMC³⁵ and CAD-FCIQMC.³⁶ We refer the interested reader to Ref. 17 and its supporting information for additional details on each method and the complete list of references. Soon after, Lee *et al.* reported phaseless auxiliary-field quantum Monte Carlo³⁷ (ph-AFQMC) correlation energies for the very same problem.³⁸

The target application is the non-relativistic frozen-core correlation energy of the ground state of the benzene molecule in the cc-pVDZ basis. The geometry of benzene has been computed at the MP2/6-31G* level and it can be found in the supporting information of Ref. 17 alongside its nuclear repulsion and Hartree-Fock energies. This corresponds to an active space of 30 electrons and 108 orbitals, *i.e.*, the Hilbert space of benzene is of the order of 10^{35} Slater determinants. Needless to say that this size of Hilbert space cannot be tackled by exact diagonalization with current architectures. The correlation energies reported in Ref. 17 are gathered in Table I alongside the best ph-AFQMC estimate from Ref. 38 based on a CAS(6,6)

TABLE I. The frozen-core correlation energy (in $\text{m}E_h$) of benzene in the cc-pVDZ basis set using various methods.

Method	E_c	Ref.
ASCI	-860.0	17
iCI	-861.1	17
CCSDTQ	-862.4	17
DMRG	-862.8	17
FCCR	-863.0	17
MBE-FCI	-863.0	17
CAD-FCIQMC	-863.4	17
AS-FCIQMC	-863.7	17
SHCI	-864.2	17
ph-AFQMC	-864.3(4)	38
CIPSI	-863.4	This work

trial wave function. The outcome of this work is nicely summarized in the abstract of Ref. 17: “*In our assessment, the evaluated high-level methods are all found to qualitatively agree on a final correlation energy, with most methods yielding an estimate of the FCI value around $-863 \text{ m}E_h$. However, we find the root-mean-square deviation of the energies from the studied methods to be considerable ($1.3 \text{ m}E_h$), which in light of the acclaimed performance of each of the methods for smaller molecular systems clearly displays the challenges faced in extending reliable, near-exact correlation methods to larger systems.*”

For the sake of completeness and our very own curiosity, we report in this Note the frozen-core correlation energy obtained with a fourth flavor of SCI known as *Configuration Interaction using a Perturbative Selection made Iteratively* (CIPSI),¹⁰ which also includes a second-order perturbative (PT2) correction. In short, the CIPSI algorithm belongs to the family of SCI+PT2 methods. The idea behind such methods is to avoid the exponential increase of the size of the CI expansion by retaining the most energetically relevant determinants only, thanks to the use of a second-order energetic criterion to select perturbatively determinants in the FCI space. However, performing SCI calculations rapidly becomes extremely tedious when one increases the system size as one hits the exponential wall inherently linked to these methods.

From a historical point of view, CIPSI is probably one of the oldest SCI algorithm. It was developed in 1973 by Huron, Rancurel, and Malrieu¹⁰ (see also Ref. 39). Recently, the determinant-driven CIPSI algorithm has been efficiently implemented^{40,41} in the open-source programming environment QUANTUM PACKAGE by our group enabling to perform mas-

sively parallel computations.^{42–44} In particular, we were able to compute highly-accurate calculations of ground- and excited-state energies for small- and medium-sized molecules (including benzene).^{12–16} CIPSI is also frequently used to provide accurate trial wave function for QMC calculations.^{40,41,45–54} The particularity of the current implementation is that the selection step and the PT2 correction are computed *simultaneously* via a hybrid semistochastic algorithm^{42,44} (which explains the statistical error associated with the PT2 correction in the following). Moreover, a renormalized version of the PT2 correction (dubbed rPT2 below) has been recently implemented for a more efficient extrapolation to the FCI limit.⁴⁴ We refer the interested reader to Ref. 44 where one can find all the details regarding the implementation of the CIPSI algorithm.

Being late to the party, we obviously cannot report blindly our CIPSI results. However, following the philosophy of Eriksen *et al.*¹⁷ and Lee *et al.*,³⁸ we will report our results with the most neutral tone, leaving the freedom to the reader to make up his/her mind. We then follow our usual “protocol”^{12–16,50–52} by performing a preliminary SCI calculation using Hartree-Fock orbitals in order to generate a SCI wave function with at least 10^7 determinants. Natural orbitals are then computed based on this wave function, and a new, larger SCI calculation is performed with this new set of orbitals. This has the advantage to produce a smoother and faster convergence of the SCI energy toward the FCI limit. The total SCI energy is defined as the sum of the variational energy $E_{\text{var.}}$ (computed via diagonalization of the CI matrix in the reference space) and a second-order perturbative correction $E_{(\text{r})\text{PT2}}$ which takes into account the external determinants, *i.e.*, the determinants which do not belong to the variational space but are linked to the reference space via a nonzero matrix element. The magnitude of $E_{(\text{r})\text{PT2}}$ provides a qualitative idea of the “distance” to the FCI limit. As mentioned above, SCI+PT2 methods rely heavily on extrapolation, especially when one deals with medium-sized systems. We then linearly extrapolate the total SCI energy to $E_{(\text{r})\text{PT2}} = 0$ (which effectively corresponds to the FCI limit). Note that, unlike excited-state calculations where it is important to enforce that the wave functions are eigenfunctions of the \hat{S}^2 spin operator,⁵⁵ the present wave functions do not fulfil this property as we aim for the lowest possible energy of a single state. We have found that $\langle \hat{S}^2 \rangle$ is, nonetheless, very close to zero ($\sim 5 \times 10^{-3}$ a.u.). The corresponding energies are reported in Table II as functions of the number of determinants in the variational space N_{det} .

A second run has been performed with localized orbitals. Starting from the same natural orbitals, a Boys-Foster localization procedure⁵⁶ was performed in several orbital windows: i) core, ii) valence σ , iii) valence π , iv) valence π^* , v) valence σ^* , vi) the higher-lying σ orbitals, and vii) the higher-lying π orbitals.⁵⁷ Like Pipek-Mezey,⁵⁸ this choice of orbital windows allows to preserve a strict σ - π separation in planar systems like benzene. As one can see from the energies of Table II, for a given value of N_{det} , the variational energy as well as the

PT2-corrected energies are much lower with localized orbitals than with natural orbitals. We, therefore, consider these energies more trustworthy, and we will base our best estimate of the correlation energy of benzene on these calculations. The convergence of the CIPSI correlation energy using localized orbitals is illustrated in Fig. 1, where one can see the behavior of the correlation energy, $\Delta E_{\text{var.}}$ and $\Delta E_{\text{var.}} + E_{(\text{r})\text{PT2}}$, as a function of N_{det} (left panel). The right panel of Fig. 1 is more instructive as it shows $\Delta E_{\text{var.}}$ as a function of $E_{(\text{r})\text{PT2}}$, and their corresponding four-point linear extrapolation curves that we have used to get our final estimate of the correlation energy. From this figure, one clearly sees that the rPT2-based correction behaves more linearly than its corresponding PT2 version, and is thus systematically employed in the following.

Our final numbers are gathered in Table III, where, following the notations of Ref. 17, we report, in addition to the final variational energies $\Delta E_{\text{var.}}$, the extrapolation distances, ΔE_{dist} , defined as the difference between the final computed energy, ΔE_{final} , and the extrapolated energy, $\Delta E_{\text{extrap.}}$, associated with ASCI, iCI, SHCI, DMRS, and CIPSI. The three flavours of SCI fall into an interval ranging from $-860.0 \text{ m}E_h$ (ASCI) to $-864.2 \text{ m}E_h$ (SHCI), while the other non-SCI methods yield correlation energies ranging from -863.7 to $-862.8 \text{ m}E_h$ (see Table I). Our final CIPSI number (obtained with localized orbitals and rPT2 correction via a four-point linear extrapolation) is $-863.4(5) \text{ m}E_h$, where the error reported in parenthesis represents the fitting error (not the extrapolation error for which it is much harder to provide a theoretically sound estimate). For comparison, the best post blind test SHCI estimate is $-863.3 \text{ m}E_h$, which agrees almost perfectly with our best CIPSI estimate, while the best post blind test ASCI and iCI correlation energies are -861.3 and $-864.15 \text{ m}E_h$, respectively (see Table III).

The present calculations have been performed on the AMD partition of GENCI’s Irene supercomputer. Each Irene’s AMD node is a dual-socket AMD Rome (Epyc) CPU@2.60 GHz with 256GiB of RAM, with a total of 64 physical CPU cores per socket. These nodes are connected via Infiniband HDR100. The first step of the calculation, *i.e.*, performing a CIPSI calculation up to $N_{\text{det}} \sim 10^7$ with Hartree-Fock orbitals in order to produce natural orbitals, takes roughly 24 hours on a single node, and reaching the same number of determinants with natural orbitals or localized orbitals takes roughly the same amount of time. A second 24-hour run on 10 distributed nodes was performed to push the selection to 8×10^7 determinants, and a third distributed run using 40 nodes was used to reach 16×10^7 determinants. In total, the present calculation has required 150k core hours, most of it being spent in the last stage of the computation.

We thank Janus Eriksen and Cyrus Umrigar for useful comments. This work was performed using HPC resources from GENCI-TGCC (2020-gen1738) and from CALMIP (Toulouse) under allocation 2020-18005.

* loos@irsamc.ups-tlse.fr

† scemama@irsamc.ups-tlse.fr




FIG. 1. Convergence of the CIPSI correlation energy of benzene using localized orbitals. Left: $\Delta E_{\text{var.}}$, $\Delta E_{\text{var.}} + E_{\text{PT2}}$, and $\Delta E_{\text{var.}} + E_{\text{rPT2}}$ (in mE_h) as functions of the number of determinants in the variational space N_{det} . Right: $\Delta E_{\text{var.}}$ (in mE_h) as a function of E_{PT2} or E_{rPT2} . The four-point linear extrapolation curves (dashed lines) are also reported. The theoretical estimate of $-863 mE_h$ from Ref. 17 is marked by a black line for comparison purposes.

TABLE II. Variational energy $E_{\text{var.}}$, second-order perturbative correction E_{PT2} and its renormalized version E_{rPT2} (in E_h) as a function of the number of determinants N_{det} for the ground-state of the benzene molecule computed in the cc-pVDZ basis set. The statistical error on $E_{(\text{r})\text{PT2}}$, corresponding to one standard deviation, are reported in parenthesis.

N_{det}	Natural orbitals			Localized orbitals		
	$E_{\text{var.}}$	$E_{\text{var.}} + E_{\text{PT2}}$	$E_{\text{var.}} + E_{\text{rPT2}}$	$E_{\text{var.}}$	$E_{\text{var.}} + E_{\text{PT2}}$	$E_{\text{var.}} + E_{\text{rPT2}}$
1 280	-230.978 056	-231.559 025(212)	-231.463 633(177)	-231.101 676	-231.519 522(149)	-231.472 224(132)
2 560	-231.043 712	-231.542 344(139)	-231.474 885(120)	-231.161 264	-231.515 577(155)	-231.482 477(140)
5 120	-231.115 142	-231.534 122(213)	-231.488 815(190)	-231.224 632	-231.516 375(191)	-231.495 022(177)
10 240	-231.188 813	-231.531 660(516)	-231.502 992(473)	-231.283 295	-231.520 907(271)	-231.507 708(255)
20 480	-231.260 065	-231.534 172(611)	-231.517 063(573)	-231.330 209	-231.526 433(586)	-231.518 045(561)
40 960	-231.321 906	-231.538 269(501)	-231.528 301(478)	-231.366 008	-231.532 288(303)	-231.526 639(293)
81 920	-231.366 895	-231.541 945(813)	-231.535 785(785)	-231.392 888	-231.536 578(614)	-231.532 575(597)
163 840	-231.392 866	-231.545 499(761)	-231.541 010(739)	-231.414 132	-231.541 400(624)	-231.538 378(609)
327 680	-231.407 802	-231.548 699(662)	-231.544 980(645)	-231.431 952	-231.545 873(557)	-231.543 532(545)
655 360	-231.418 752	-231.551 208(661)	-231.548 004(645)	-231.447 007	-231.548 856(498)	-231.547 043(489)
1 310 720	-231.428 852	-231.552 760(616)	-231.550 006(603)	-231.460 970	-231.552 137(453)	-231.550 723(446)
2 621 440	-231.439 324	-231.553 845(572)	-231.551 544(560)	-231.473 751	-231.555 261(403)	-231.554 159(397)
5 242 880	-231.450 156	-231.557 541(534)	-231.555 558(524)	-231.485 829	-231.558 303(362)	-231.557 451(358)
10 485 760	-231.461 927	-231.559 390(481)	-231.557 796(474)	-231.497 515	-231.562 568(322)	-231.561 901(319)
20 971 520	-231.474 019	-231.561 315(430)	-231.560 063(424)	-231.508 714	-231.564 707(275)	-231.564 223(273)
41 943 040	-231.487 978	-231.564 529(382)	-231.563 593(377)	-231.519 122	-231.567 419(240)	-231.567 069(238)
83 886 080	-231.501 334	-231.566 994(317)	-231.566 325(314)	-231.528 568	-231.570 084(199)	-231.569 832(198)
167 772 160	-231.514 009	-231.569 939(273)	-231.569 467(271)	-231.536 655	-231.571 981(175)	-231.571 804(174)

¹ J. P. F. LeBlanc, A. E. Antipov, F. Becca, I. W. Bulik, G. K.-L. Chan, C.-M. Chung, Y. Deng, M. Ferrero, T. M. Henderson, C. A. Jiménez-Hoyos, *et al.*, *Phys. Rev. X* **5**, 041041 (2015).

² M. Motta, D. M. Ceperley, G. K.-L. Chan, J. A. Gomez, E. Gull, S. Guo, C. A. Jiménez-Hoyos, T. N. Lan, J. Li, F. Ma, *et al.*, *Phys. Rev. X* **7**, 031059 (2017).

³ K. T. Williams, Y. Yao, J. Li, L. Chen, H. Shi, M. Motta, C. Niu, U. Ray, S. Guo, R. J. Anderson, *et al.*, *Phys. Rev. X* **10**, 011041 (2020).

⁴ S. P. A. Sauer, M. Schreiber, M. R. Silva-Junior, and W. Thiel, *J. Chem. Theory Comput.* **5**, 555 (2009).

⁵ M. Schreiber, M. R. Silva-Junior, S. P. A. Sauer, and W. Thiel, *J. Chem. Phys.* **128**, 134110 (2008).

⁶ M. R. Silva-Junior, M. Schreiber, S. P. A. Sauer, and W. Thiel, *J. Chem. Phys.* **129**, 104103 (2008).

⁷ M. R. Silva-Junior, S. P. A. Sauer, M. Schreiber, and W. Thiel, *Mol. Phys.* **108**, 453 (2010).

⁸ M. R. Silva-Junior, M. Schreiber, S. P. A. Sauer, and W. Thiel, *J. Chem. Phys.* **133**, 174318 (2010).

⁹ C. F. Bender and E. R. Davidson, *Phys. Rev.* **183**, 23 (1969).

¹⁰ B. Huron, J. P. Malrieu, and P. Rancurel, *J. Chem. Phys.* **58**, 5745 (1973).

TABLE III. Extrapolation distances, ΔE_{dist} , defined as the difference between the final computed energy, ΔE_{final} , and the extrapolated energy, ΔE_{extrap} , associated with ASCI, iCI, SHCI, DMRG, and CIPSI for the best blind-test and post-blind-test estimates of the correlation energy of benzene in the cc-pVDZ basis. The final variational energies ΔE_{var} , are also reported. See Ref. 17 for more details. All correlation energies are given in mE_h .

Method	ΔE_{var}	ΔE_{final}	$\Delta E_{\text{extrap.}}$	ΔE_{dist}
Best blind-test estimates				
ASCI	-737.1	-835.4	-860.0	-24.6
iCI	-730.0	-833.7	-861.1	-27.4
SHCI	-827.2	-852.8	-864.2	-11.4
DMRG	-859.2	-859.2	-862.8	-3.6
Best post-blind-test estimates				
ASCI	-772.4	-835.2	-861.3	-26.1
iCI	-770.7	-842.8	-864.2	-21.3
SHCI	-835.2	-854.9	-863.6	-8.7
CIPSI	-814.8	-850.2	-863.4	-13.2

- ¹¹ R. J. Buenker and S. D. Peyerimhoff, *Theor. Chim. Acta* **35**, 33 (1974).
- ¹² P. F. Loos, A. Scemama, A. Blondel, Y. Garniron, M. Caffarel, and D. Jacquemin, *J. Chem. Theory Comput.* **14**, 4360 (2018).
- ¹³ P.-F. Loos, M. Boggio-Pasqua, A. Scemama, M. Caffarel, and D. Jacquemin, *J. Chem. Theory Comput.* **15**, 1939 (2019).
- ¹⁴ P. F. Loos, A. Scemama, and D. Jacquemin, *J. Phys. Chem. Lett.* **11**, 2374 (2020).
- ¹⁵ P. F. Loos, F. Lippalini, M. Boggio-Pasqua, A. Scemama, and D. Jacquemin, *J. Chem. Theory Comput.* **16**, 1711 (2020).
- ¹⁶ P. F. Loos, A. Scemama, M. Boggio-Pasqua, and D. Jacquemin, *J. Chem. Theory Comput.* **16**, 3720 (2020).
- ¹⁷ J. J. Eriksen, T. A. Anderson, J. E. Deustua, K. Ghanem, D. Hait, M. R. Hoffmann, S. Lee, D. S. Levine, I. Magoulas, J. Shen, N. M. Tubman, K. B. Whaley, E. Xu, Y. Yao, N. Zhang, A. Alavi, G. K.-L. Chan, M. Head-Gordon, W. Liu, P. Piecuch, S. Sharma, S. L. Ten-no, C. J. Umrigar, and J. Gauss, “The ground state electronic energy of benzene,” (2020), arXiv:2008.02678 [physics.chem-ph].
- ¹⁸ N. Oliphant and L. Adamowicz, *J. Chem. Phys.* **95**, 6645 (1991).
- ¹⁹ S. A. Kucharski and R. J. Bartlett, *J. Chem. Phys.* **97**, 4282 (1992).
- ²⁰ J. J. Eriksen, F. Lippalini, and J. Gauss, *J. Phys. Chem. Lett.* **8**, 4633 (2017).
- ²¹ J. J. Eriksen and J. Gauss, *J. Chem. Theory Comput.* **14**, 5180 (2018).
- ²² J. J. Eriksen and J. Gauss, *J. Chem. Theory Comput.* **15**, 4873 (2019).
- ²³ J. J. Eriksen and J. Gauss, *J. Phys. Chem. Lett.* **27**, 7910 (2019).
- ²⁴ N. M. Tubman, J. Lee, T. Y. Takeshita, M. Head-Gordon, and K. B. Whaley, *J. Chem. Phys.* **145**, 044112 (2016).
- ²⁵ N. M. Tubman, D. S. Levine, D. Hait, M. Head-Gordon, and K. B. Whaley, “An efficient deterministic perturbation theory for selected configuration interaction methods,” (2018), arXiv:1808.02049 [cond-mat.str-el].
- ²⁶ N. M. Tubman, C. D. Freeman, D. S. Levine, D. Hait, M. Head-Gordon, and K. B. Whaley, *J. Chem. Theory Comput.* **16**, 2139 (2020).
- ²⁷ W. Liu and M. R. Hoffmann, *J. Chem. Theory Comput.* **12**, 1169 (2016).
- ²⁸ A. A. Holmes, H. J. Changlani, and C. J. Umrigar, *J. Chem. Theory Comput.* **12**, 1561 (2016).
- ²⁹ A. A. Holmes, C. J. Umrigar, and S. Sharma, *J. Chem. Phys.* **147**, 164111 (2017).
- ³⁰ S. Sharma, A. A. Holmes, G. Jeanmairet, A. Alavi, and C. J. Umrigar, *J. Chem. Theory Comput.* **13**, 1595 (2017).
- ³¹ E. Xu, M. Uejima, and S. L. Ten-no, *Phys. Rev. Lett.* **121**, 113001 (2018).
- ³² S. R. White, *Phys. Rev. Lett.* **69**, 2863 (1992).
- ³³ G. H. Booth, A. J. W. Thom, and A. Alavi, *J. Chem. Phys.* **131**, 054106 (2009).
- ³⁴ D. Cleland, G. H. Booth, and A. Alavi, *J. Chem. Phys.* **132**, 041103 (2010).
- ³⁵ K. Ghanem, A. Y. Lozovoi, and A. Alavi, *J. Chem. Phys.* **151**, 224108 (2019).
- ³⁶ J. E. Deustua, I. Magoulas, J. Shen, and P. Piecuch, *J. Chem. Phys.* **149**, 151101 (2018).
- ³⁷ M. Motta and S. Zhang, *WIREs Comput. Mol. Sci.* **8**, e1364 (2018).
- ³⁸ J. Lee, F. D. Malone, and D. R. Reichman, “Note: The performance of phaseless auxiliary-field quantum monte carlo on the ground state electronic energy of benzene,” (2020), arXiv:2008.04736 [physics.chem-ph].
- ³⁹ S. Evangelisti, J.-P. Daudey, and J.-P. Malrieu, *Chem. Phys.* **75**, 91 (1983).
- ⁴⁰ E. Giner, A. Scemama, and M. Caffarel, *Can. J. Chem.* **91**, 879 (2013).
- ⁴¹ E. Giner, A. Scemama, and M. Caffarel, *J. Chem. Phys.* **142**, 044115 (2015).
- ⁴² Y. Garniron, A. Scemama, P.-F. Loos, and M. Caffarel, *J. Chem. Phys.* **147**, 034101 (2017).
- ⁴³ Y. Garniron, A. Scemama, E. Giner, M. Caffarel, and P. F. Loos, *J. Chem. Phys.* **149**, 064103 (2018).
- ⁴⁴ Y. Garniron, T. Applencourt, K. Gasperich, A. Benali, A. Ferté, J. Paquier, B. Pradines, R. Assaraf, P. Reinhardt, J. Toulouse, P. Barbaresco, N. Renon, G. David, J.-P. Malrieu, M. Vérit, M. Caffarel, P.-F. Loos, E. Giner, and A. Scemama, *J. Chem. Theory Comput.* **15**, 3591 (2019).
- ⁴⁵ M. Caffarel, E. Giner, A. Scemama, and A. Ramírez-Solís, *J. Chem. Theory Comput.* **10**, 5286 (2014).
- ⁴⁶ M. Caffarel, T. Applencourt, E. Giner, and A. Scemama, *J. Chem. Phys.* **144**, 151103 (2016).
- ⁴⁷ “Recent Progress in Quantum Monte Carlo,” (2016), [Online; accessed 6. Jul. 2020].
- ⁴⁸ A. Scemama, E. Giner, T. Applencourt, and M. Caffarel, “QMC using very large configuration interaction-type expansions,” Pacifichem, Advances in Quantum Monte Carlo (2015).
- ⁴⁹ A. Scemama, T. Applencourt, E. Giner, and M. Caffarel, *J. Comput. Chem.* **37**, 1866 (2016).
- ⁵⁰ A. Scemama, Y. Garniron, M. Caffarel, and P.-F. Loos, *J. Chem. Theory Comput.* **14**, 1395 (2018).
- ⁵¹ A. Scemama, A. Benali, D. Jacquemin, M. Caffarel, and P.-F. Loos, *J. Chem. Phys.* **149**, 034108 (2018).
- ⁵² A. Scemama, M. Caffarel, A. Benali, D. Jacquemin, and P. F. Loos., *Res. Chem.* **1**, 100002 (2019).
- ⁵³ M. Dash, S. Moroni, A. Scemama, and C. Filippi, *J. Chem. Theory Comput.* **14**, 4176 (2018).
- ⁵⁴ M. Dash, J. Feldt, S. Moroni, A. Scemama, and C. Filippi, *J. Chem. Theory Comput.* **15**, 4896 (2019).
- ⁵⁵ T. Applencourt, K. Gasperich, and A. Scemama, “Spin adaptation with determinant-based selected configuration interaction,” (2018), arXiv:1812.06902 [physics.chem-ph].
- ⁵⁶ J. M. Foster and S. F. Boys, *Rev. Mod. Phys.* **32**, 300 (1960).
- ⁵⁷ MO indices for Boys-Foster localization procedure: core [1–6]; σ [7–18]; π [19–21]; π^* [22–24]; σ^* [25–36]; higher-lying π [39,41–43,46,49,50,53–57,71–74,82–85,87,92,93,98]; higher-lying σ [37,38,40,44,45,47,48,51,52,58–70,75–81,86,88–91,94–97,99–114].
- ⁵⁸ J. Pipek and P. G. Mezey, *J. Chem. Phys.* **90**, 4916 (1989).