



Biomechanical constraints remain major risk factors for low back pain : results from a prospective cohort study in French male employees

Aline Ramond-Roquin, Audrey Petit, Julie Bodin, C Sérazin, Isabelle Richard, Y. Roquelaure

► To cite this version:

Aline Ramond-Roquin, Audrey Petit, Julie Bodin, C Sérazin, Isabelle Richard, et al.. Biomechanical constraints remain major risk factors for low back pain : results from a prospective cohort study in French male employees. 22nd International Conference on Epidemiology in Occupational Health (EPICOH), Sep 2011, Oxford, United Kingdom. hal-02922833

HAL Id: hal-02922833

<https://hal.science/hal-02922833>

Submitted on 24 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biomechanical constraints remain major risk factors for low back pain

Prospective study among French workers

RAMOND-ROQUIN A^{1,2}, PETIT LE MANACH A¹, BODIN J¹, SERAZIN C¹, PAROT E^{1,3}, HA C⁴, ROQUELAURE Y^{1,5}

¹ Laboratory of Ergonomics and Epidemiology in Occupational Health, University of Angers ; ² Department of General Practice, University of Angers ;

³ Center of Clinical Research, University Hospital of Angers ; ⁴ Institute of Health Surveillance, Saint-Maurice ; ⁵ Department of Occupational Health,

University of Angers

aline.ramond@univ-angers.fr

INTRODUCTION

Low back pain (LBP) in France : A major public health problem



One in two adults suffer from LBP at least once a year¹



Direct costs for chronic LBP :
~ 715 € per patient every 6 months²




Impact on occupational settings :
LBP = the most frequent reason for sick leave³

OBJECTIVE

Prospective modelling of the risk of LBP in the French male working population

METHODS

Design : Cohort prospective study



Statistical analysis:

Multivariate and multi-step logistic modelling of the risk of LBP in the male workers.


Selection of statistically significant risk factors ($p < 0.05$) and confounding factors.

Ethical aspects :

Written consent from the workers - Authorization from the National Commission of Data Protection

RESULTS

Study population :



Final multivariate model (n=1255 due to missing data):

RISK FACTORS		OR (IC 95) [†]
Frequent bending (more than 2 hours daily)	No	1
	Forward	1.45 (1.07-1.97)
	Sideways	1.78 (0.74-4.27)
	Both	2.13 (1.52-3.00)
Driving industrial vehicles	Never	1
	Occasionally/Often	1.35 (1.00-1.81)
Working more hours than officially planned	No	1
	Yes	1.38 (1.05-1.81)
Low support from supervisors	No	1
	Yes	1.35 (1.02-1.79)
Low skill discretion	No	1
	Yes	1.28 (0.95-1.72)

[†] Odds-ratios for LBP and confidence intervals at 95%

DISCUSSION & CONCLUSION

Prospective study ; diversified cohort of male workers ; numerous risk factors

→ Useful and relevant to the development of **public health policy** for workers.

Current research on LBP focus on psychosocial factors.

However biomechanical constraints (frequent bending, driving industrial vehicles) remain major risk factors that deserve to be considered in primary prevention !

REFERENCES

- ¹Gourmelen J, Chastang JF, Ozguler A et al. Frequency of low back pain among men and women aged 30 to 64 years in France. Results of two national surveys. Ann Readapt Med Phys. 2007 Nov;50(8):640-4, 633-9
- ²Depont F, Hunsche E, Abouelfath A et al. Medical and non-medical direct costs of chronic low back pain in patients consulting primary care physicians in France. Fundam Clin Pharmacol. 2010 Feb;24(1):101-8.
- ³Caisse Nationale d'Assurance Maladie - Travailleurs Salariés.