



**HAL**  
open science

## Positive feedback between future climate change and the carbon cycle

Pierre Friedlingstein, Laurent Bopp, Philippe Ciais, Jean-Louis Dufresne, Laurent Fairhead, Hervé Letreut, Patrick Monfray, James Orr

### ► To cite this version:

Pierre Friedlingstein, Laurent Bopp, Philippe Ciais, Jean-Louis Dufresne, Laurent Fairhead, et al.. Positive feedback between future climate change and the carbon cycle. *Geophysical Research Letters*, 2001, 28 (8), pp.1543-1546. 10.1029/2000GL012015 . hal-02923899

**HAL Id: hal-02923899**

**<https://hal.science/hal-02923899>**

Submitted on 28 Oct 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Positive feedback between future climate change and the carbon cycle

Pierre Friedlingstein, Laurent Bopp, Philippe Ciais,  
IPSL/LSCE, CE-Saclay, 91191, Gif sur Yvette, France

Jean-Louis Dufresne, Laurent Fairhead, Hervé LeTreut,  
IPSL/LMD, Université Paris 6, 75252, Paris, France

Patrick Monfray, and James Orr  
IPSL/LSCE, CE-Saclay, 91191, Gif sur Yvette, France

**Abstract.** Future climate change due to increased atmospheric CO<sub>2</sub> may affect land and ocean efficiency to absorb atmospheric CO<sub>2</sub>. Here, using climate and carbon three-dimensional models forced by a 1% per year increase in atmospheric CO<sub>2</sub>, we show that there is a positive feedback between the climate system and the carbon cycle. Climate change reduces land and ocean uptake of CO<sub>2</sub>, respectively by 54% and 35% at 4 × CO<sub>2</sub>. This negative impact implies that for prescribed anthropogenic CO<sub>2</sub> emissions, the atmospheric CO<sub>2</sub> would be higher than the level reached if climate change does not affect the carbon cycle. We estimate the gain of this climate-carbon cycle feedback to be 10% at 2 × CO<sub>2</sub> and 20% at 4 × CO<sub>2</sub>. This translates into a 15% higher mean temperature increase.

### Introduction

Atmospheric CO<sub>2</sub> is expected to increase in the coming decades due to emissions of CO<sub>2</sub> by fossil fuel burning and land use changes. The rate of increase depends on anthropogenic emissions and on the capacity of the oceans and the land biosphere to take up CO<sub>2</sub> [Schimel *et al.*, 1995]. Current climate models predict a mean temperature increase of 1 to 4.5°C compared to the present for a doubling of atmospheric CO<sub>2</sub> [Kattenberg *et al.*, 1996]. Recent carbon cycle studies suggest that such climate change may reduce the uptake of CO<sub>2</sub> by the ocean [Maier-Reimer *et al.*, 1996; Sarmiento *et al.*, 1998; Matear and Hirst, 1999] or the land biosphere [Cao and Woodward, 1998; Meyer *et al.*, 1999; Cramer *et al.*, 2000]. It is thus necessary to account for the climate impact on the carbon cycle when translating anthropogenic emissions into CO<sub>2</sub> concentrations.

### Method

In this study, we used a model structure composed of a coupled ocean-atmosphere general circulation model (OAGCM), and models of land and ocean components of the carbon cycle, the carbon cycle models being forced by the climate fields of the OAGCM. Two climate simulations have been run with the OAGCM: the control run where the CO<sub>2</sub> is held constant at 350 ppmv and the transient climate run where the CO<sub>2</sub> increases at a rate of 1% per year from 350 ppmv up to 1400 ppmv (Figure 1a). We then performed

two carbon simulations. In the “constant climate” simulation the carbon models are forced by a CO<sub>2</sub> increase of 1% per year and the control climate from the OAGCM. In the “climate change” simulation, the carbon models are forced by the same 1%/yr CO<sub>2</sub> increase as well as the climate from the transient climate run.


In this experiment, atmospheric CO<sub>2</sub> and monthly averaged climate fields from the IPSL OAGCM [Braconnot *et al.*, 2000] are used to drive both terrestrial and oceanic carbon cycle models. These two models allow one to translate anthropogenic CO<sub>2</sub> emissions into atmospheric CO<sub>2</sub> concentration trajectories and vice-versa. The terrestrial carbon model (SLAVE) [Friedlingstein *et al.*, 1995; Ciais *et al.*, 1999] is driven by surface air temperature, precipitation, and solar radiation, and calculates net primary productivity (NPP) following a light use efficiency formulation [Field *et al.*, 1995] that is a function of temperature and water stress. NPP increases with CO<sub>2</sub> under a Michaelis-Menten beta factor formulation [Gifford, 1992], which has a global value of 0.5, in the upper range of experimental data [DeLucia *et al.*, 1999], although, nitrogen limitation and deposition as well as vegetation dynamics and land use changes are ignored in this study. The ocean carbon model (IPSL-OCCM1) [Aumont *et al.*, 1999; Le Quéré *et al.*, 1999], based on the HAMOCC3 biogeochemical scheme [Maier-Reimer, 1993] is driven by monthly mean global fields of oceanic circulation, temperature, salinity, and surface fields of winds, sea ice and water fluxes all issued from the OAGCM. Both land and ocean carbon models have been applied successfully to study seasonal, interannual and decadal characteristics of the carbon cycle over the historical period [Friedlingstein *et al.*, 1995; Ciais *et al.*, 1999; Aumont *et al.*, 1999; Le Quéré *et al.*, 1999].

**Table 1.** Changes in cumulated carbon budget at 2×CO<sub>2</sub> and 4×CO<sub>2</sub>

	2×CO <sub>2</sub>		4×CO <sub>2</sub>	
	Constant Clim.	Clim. Change	Constant Clim.	Clim. Change
Ocean uptake (GtC)	347	312 (−10%)	1002	800 (−20%)
Land uptake (GtC)	403	310 (−23%)	1195	808 (−32%)
Atmospheric Increase (GtC)	742	742	2226	2226
Anthropogenic emission (GtC)	1492	1364 (−8.5%)	4423	3834 (−13%)

Copyright 2001 by the American Geophysical Union.

Paper number 2000GL012015.  
0094-8276/01/2000GL012015\$05.00



**Figure 1.** Carbon budget. a) Atmospheric CO<sub>2</sub> scenario used as a forcing for the climate model (in ppmv) (*12*). b) Simulated annual biospheric CO<sub>2</sub> uptake (GtC/yr) for the constant climate simulation (red line), the climate change simulation (green line) and the difference between the two simulations, showing the climate change impact on reduction biospheric carbon uptake (blue line). c) same as b), but for the ocean. d) Annual rate of compatible anthropogenic CO<sub>2</sub> emissions calculated as the sum of atmospheric CO<sub>2</sub> growth rate and land plus ocean carbon uptakes (GtC/yr). Lines colors follows the same convention as in b).

### Climate Impact on Land Uptake


In the constant climate experiment, increasing CO<sub>2</sub> stimulates terrestrial NPP from 70 to 110 GtC/yr at 2 × CO<sub>2</sub>, and to 150 GtC/yr at 4 × CO<sub>2</sub>. These results fall within the range of previous model [Cao and Woodward, 1998; Meyer et al., 1999; Cramer et al., 2000]. The residence time of carbon in living and dead biomass induces a transient disequilibrium between NPP and the release due to oxidation of decaying material. A net biospheric uptake (NEP) grows as long as atmospheric CO<sub>2</sub> increases, reaching 9 GtC/yr at 2 × CO<sub>2</sub> and 12 GtC/yr at 4 × CO<sub>2</sub> (Figure 1b). When CO<sub>2</sub> stabilizes, so does the NPP, and the biosphere reaches a new equilibrium state. The climate change experiment, shows a much smaller NEP than the constant climate run (Figure 1b). Ten years before reaching 2 × CO<sub>2</sub>, NEP saturates at around 7 GtC/yr, and starts to decrease after 120 years despite increasing atmospheric CO<sub>2</sub>. When CO<sub>2</sub> reaches 4 × CO<sub>2</sub>, NEP only amounts to 5.5 GtC/yr, less than half of the value found at the same CO<sub>2</sub> level in the constant climate run. The cumulative land uptake in the climate change run is 310 GtC at 2 × CO<sub>2</sub> and 808 GtC at 4 × CO<sub>2</sub>, that is respectively 23% and 32% lower than in the constant climate simulation (Table 1).

The strong reduction of NEP induced by the climate change is mainly located in the subtropics (especially South


America) and caused by increase in soil aridity, due to a larger increase in evaporative demand than in precipitation (Figure 2). Qualitatively similar findings were found previously [Cao and Woodward, 1998; Cramer et al., 2000].

### Climate Impact on Ocean Uptake

For the constant climate run, rising atmospheric CO<sub>2</sub> also increases the oceanic uptake. At 2 × CO<sub>2</sub>, the ocean carbon sink reaches 7.5 GtC/yr and 10.5 GtC/yr at 4 × CO<sub>2</sub> (Figure 1c). After CO<sub>2</sub> stabilizes, the ocean uptake decreases as the ocean carbon tends toward a new equilibrium state. As for the land uptake, the oceanic uptake is always lower in the climate change simulation than under constant climate. After 80 years, oceanic uptake saturates around 7 GtC/yr, and shows a slight decrease during the last ten years of increasing atmospheric CO<sub>2</sub>. At 4 × CO<sub>2</sub>, the oceanic uptakes amounts to 5.7 GtC/yr, which is 35% lower than in the constant climate run. When cumulated, the climate induced decrease of oceanic uptake is 10% at 2 × CO<sub>2</sub>, and 20% at 4 × CO<sub>2</sub> (Table 1). The effect of the global warming scenario in reducing oceanic CO<sub>2</sub> uptake is, at 2 × CO<sub>2</sub> of the same order as that was previously found [Maier-Reimer et al., 1996; Sarmiento et al., 1998; Matear and Hirst, 1999]. As shown on Figure 3, the reduction in the oceanic uptake of carbon, as discussed in earlier studies, results from the combination of three effects: impact of increased sea-surface temperature on CO<sub>2</sub> solubility, impact of reduced vertical mixing on CO<sub>2</sub> transport from the surface to the deep ocean and impact of changes in the biogeochemical cycle of CO<sub>2</sub>. The combination of those three climatic feedbacks lead to a reduced oceanic uptake of CO<sub>2</sub>, principally located at high latitudes, and for its main part in the Southern Ocean. However, this effect might be over-evaluated in our experiments due to abnormally strong oceanic convection in the Southern Ocean.



**Figure 2.** Zonal mean difference between the climate change and the constant climate simulations at the time of 4 × CO<sub>2</sub> of a) annual surface land temperature (°C) (red line) and precipitation (mm/yr) (blue line), b) soil water content (mm), c) Net Primary Productivity (gC/m<sup>2</sup>/yr), and d) net carbon uptake (gC/m<sup>2</sup>/yr).



**Figure 3.** Zonal mean difference between the climate change and the constant climate simulations at the time of  $4\times\text{CO}_2$  of a) sea surface temperature ( $^{\circ}\text{C}$ ), b) depth of the mixed layer (m), c) export production ( $\text{gC}/\text{m}^2/\text{yr}$ ), and d) net carbon uptake ( $\text{gC}/\text{m}^2/\text{yr}$ ).


### Impact on Derived Emissions

Our two estimates of both terrestrial and oceanic carbon uptakes allow us to determine the compatible anthropogenic emissions with and without accounting for the climate change. In the constant climate simulation, in order to sustain a  $1\%/yr$  increase in atmospheric  $\text{CO}_2$ , the compatible emissions have to peak at  $50\text{ GtC}/\text{yr}$  at the time of  $4\times\text{CO}_2$ , whereas they would be lowered by  $10\text{ GtC}/\text{yr}$  in the climate change run (Figure 1d). When cumulated, the compatible emissions are respectively reduced by  $8\%$  and  $13\%$  at  $2\times\text{CO}_2$  and  $4\times\text{CO}_2$  when climate change is accounted for (Table 1). Thus, to achieve a given atmospheric  $\text{CO}_2$  trajectory any economic  $\text{CO}_2$  emission scenario needs to prescribe lower emissions if the climate impact on the carbon cycle is accounted for. Furthermore, the response of the carbon cycle to the warming being non-linear, reductions in emissions will then have to be increasingly stronger with time.

### Climate System – Carbon Cycle Feedback

In a more consistent study where the carbon cycle is forced by anthropogenic emissions, as in the real world, our results would translate into a faster atmospheric  $\text{CO}_2$  buildup as land and ocean efficiencies to sequester carbon decrease with time. That in turn would feed back in a more rapid climate change and may have further adverse impacts on terrestrial and oceanic processes and on the  $\text{CO}_2$  concentration.

In the following, we provide the first estimate of the magnitude of this positive feedback. Using a classical approach [Hansen *et al.*, 1984], we define the gain of the climate system carbon cycle feedback,  $g$ , as  $\partial^*T/\partial C \times \partial^*C/\partial T$  where the first term represents the overall physical sensitivity of temperature to atmospheric  $\text{CO}_2$ , and the second term represents the overall sensitivity of atmospheric  $\text{CO}_2$  to temperature. In our climate change simulation, the sensitivity of temperature to  $\text{CO}_2$  gradually decreases from  $0.007\text{ K}/\text{ppmv}$  at  $2\times\text{CO}_2$  to  $0.003\text{ K}/\text{ppmv}$  at  $4\times\text{CO}_2$  (Figure 4a). The  $\text{CO}_2$  sensitivity to temperature can be inferred from Figure 1, showing the impact of climate change on the carbon fluxes, and from the calculated airborne fraction. The  $\text{CO}_2$  sensitivity to temperature increases strongly from  $20\text{ ppmv}/\text{K}$  at  $2\times\text{CO}_2$  to  $60\text{ ppmv}/\text{K}$  at  $4\times\text{CO}_2$  (Figure 4b). The gain,  $g$ , defined above, amounts to  $0.11$  at  $2\times\text{CO}_2$  and reaches  $0.19$  at  $4\times\text{CO}_2$  (Figure 4c). The net feedback,  $f$ , which is the global warming amplification, defined as  $1/(1-g)$ , reaches  $1.12$  and  $1.23$  at  $2\times\text{CO}_2$  and  $4\times\text{CO}_2$  respectively (Figure 4d). Assuming that future emissions follow a trajectory compatible with today's climate (Figure 1d, constant climate simulation), one can approximate the  $\text{CO}_2$  levels and the climate change that would occur in a coupled climate-carbon cycle configuration. This analytical calculation gives a  $5.2^{\circ}\text{C}$  warming at a  $\text{CO}_2$  level of  $1560\text{ ppmv}$  after 140 years instead of a  $4.6^{\circ}\text{C}$  warming at  $1400\text{ ppmv}$ , as given by the uncoupled simulation.



**Figure 4.** Time evolution of a)  $\partial^*T/\partial C$ , the overall sensitivity of surface temperature to the atmospheric  $\text{CO}_2$  ( $10^3\text{ K}/\text{ppmv}$ ), b)  $\partial^*C/\partial T$ , the overall sensitivity of atmospheric  $\text{CO}_2$  to surface temperature ( $\text{ppmv}/\text{K}$ ), c)  $g$ , the gain of the climate system-carbon cycle feedback calculated as  $g = \partial^*T/\partial C \times \partial^*C/\partial T$ , and d)  $f$ , the global warming amplification calculated as  $f = 1/(1-g)$ .

## Conclusions

Our results suggest that the future climate change impact on the carbon cycle can be large, with a risk of seeing both ocean and biospheric capacity to absorb anthropogenic CO<sub>2</sub> significantly reduced as the Earth warms up, leaving larger CO<sub>2</sub> fraction in the atmosphere and therefore enhancing the climate change. In order to further explore these effects, it should be given high priority to develop comprehensive models where physical climate system and carbon cycle are explicitly coupled.

This study is a first attempt to quantify the climate-carbon feedback under elevated CO<sub>2</sub>. To help reduce uncertainties, and to identify the key processes controlling CO<sub>2</sub> and climate requires a better understanding of the observed historical trends. In future scenarios, one should also specifically account for changes in non-CO<sub>2</sub> greenhouse gases, in future land use and land cover, in vegetation-climate feedbacks controlled by stomatal conductance and canopy development, as well as for alterations in land and ocean ecosystem distribution, and in the cycling of nutrients. In addition, non-linear changes in the ocean-atmosphere dynamics [Manabe et al., 1992], could affect the magnitude of the feedback we have calculated here.

**Acknowledgments.** We thank O. Aumont, J. Jouzel and K. Rodgers for useful comments on the manuscript. J.-Y. Grandpeix developed the formalism used for our feedback analysis. The computer time was provided by IDRIS/CNRS. This work was part of the IPSL modelling pole, and is supported by PNEDC and PROOF.

## References

- Aumont, O., J. C. Orr, P. Monfray, G. Madec, and E. Maier-Reimer, Nutrient trapping in the equatorial pacific: The ocean circulation solution, *Global Biogeochemical Cycles*, **13**, 351–371, 1999.
- Braconnot, P., O. Marti, S. Joussaume, and Y. Leclainche, Ocean feedback in response to 6 kyr bp insolation., *Journal of Climate*, **13**(9), 1537 – 1553, 2000.
- Cao, M. and F. I. Woodward, Dynamic responses of terrestrial ecosystem carbon cycling to global climate change, *Nature*, **393**, 249–252, 1998.
- Ciais, P., P. Friedlingstein, D. S. Schimel, and P. P. Tans, A global calculation of the soil carbon  $\delta^{13}\text{C}$  value : Implications for the biospheric uptake of anthropogenic CO<sub>2</sub>, *Global Biogeochemical Cycles*, **13**, 519–530, 1999.
- Cramer, W., A. Bondeau, F. Woodward, I. Prentice, R. Betts, V. Brovkin, P. Cox, V. Fisher, J. Foley, A. Friend, C. Kucharik, M. Lomas, N. Ramankutty, S. Sitch, B. Smith, A. White, and C. Young-Molling, Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models, *Global Change Biology*, 2000.
- DeLucia, E., J. Hamilton, S. Naidu, R. Thomas, J. Andrews, A. Finzi, M. Lavine, R. Matamala, J. Mohan, G. Hendrey, and W. Schlesinger, Net primary production of a forest ecosystem with experimental CO<sub>2</sub> enrichment, *Science*, **284**, 1177–1179, 1999.
- Field, C. B., J. T. Randerson, and C. M. Malmström, Global net primary production: Combining ecology and remote sensing, *Remote Sensing of Environment*, **51**, 74–88, 1995.
- Friedlingstein, P., I. Y. Fung, E. A. Holland, J. G. John, G. P. Brasseur, D. J. Erickson, and D. S. Schimel, On the contribution of the biospheric CO<sub>2</sub> fertilization to the missing sink, *Global Biogeochemical Cycles*, **9**, 541–556, 1995.
- Gifford, R. M., Interaction of carbon dioxide with growth-limiting environmental factors in vegetation productivity: Implications for the global carbon cycle, *Advances in Bioclimatology*, **1**, 24–58, 1992.
- Hansen, J., A. Lacis, D. Rind, G. Russel, P. Stone, I. Fung, R. Ruedy, and J. Lerner, Climate sensitivity: Analysis of feedback mechanisms, in *Climate Processes and Climate Sensitivity*, *Geophysical Monograph* **29**, edited by J. Hansen and T. Takahashi, pp. pp. 130–163, American Geophysical Union, Washington, D. C., 1984.
- Kattenberg, A., F. Giorgi, H. Grassl, G. A. Meehl, J. F. B. Mitchell, R. J. Stouffer, T. Tokioka, A. J. Weaver, and T. M. L. Wigley, *Climate Change 95, The Science of Climate Change*, chapter 6. Climate Models - Projections of Future Climate, pp. 285–357, Intergovernmental Panel on Climate Change, Cambridge, England, 1996.
- Le Quéré, C., J. C. Orr, P. Monfray, O. Aumont, and G. Madec, Interannual variability of the oceanic sink of CO<sub>2</sub> from 1979 through 1997, *Global Biogeochemical Cycles*, in press, 1999.
- Maier-Reimer, E., Geochemical cycles in an ocean general circulation model. Preindustrial tracer distributions, *Global Biogeochemical Cycles*, **7**, 645–677, 1993.
- Maier-Reimer, E., U. Mikolajewicz, and A. Winguth, Future ocean uptake of CO<sub>2</sub>: interaction between ocean circulation and biology, *Climate Dynamics*, **12**, 711–721, 1996.
- Manabe, S., R. J. Stouffer, M. J. Spelman, and K. Bryan, Transient response of a coupled ocean-atmosphere model to gradual changes of atmospheric CO<sub>2</sub>. Part II: Annual mean response, *Journal of Climate*, **4**, 785–818, 1992.
- Matear, R. and A. Hirst, Climate change feedback on the future oceanic CO<sub>2</sub> uptake, *Tellus*, **51B**, 722–733, 1999.
- Meyer, R., F. Joos, G. Esser, M. Heimann, G. Hooss, G. Kohlmaier, W. Sauf, R. Voss, and U. Wittenberg, The substitution of high-resolution terrestrial biosphere models and carbon sequestration in response to changing CO<sub>2</sub> and climate, *Global Biogeochemical Cycles*, **13**, 785–802, 1999.
- Sarmiento, J. L., T. M. C. Hughes, R. J. Stouffer, and S. Manabe, Simulated response of the ocean carbon cycle to anthropogenic climate warming, *Nature*, **393**, 245–249, 1998.
- Schimel, D., I. Enting, M. Heimann, T. Wigley, D. Raynaud, D. Alves, and U. Siegenthaler, CO<sub>2</sub> and the carbon cycle, in *Climate Change 1994, Radiative Forcing of Climate Change and Evaluation of the IPCC IS92 Emission Scenarios*, edited by J. T. Houghton, L. G. Meira Filho, J. Bruce, Hoesung Lee, B. A. Callander, E. Haites, N. Harris, and K. Maskell, Cambridge University Press, New York, 1995.
- P. Friedlingstein, L. Bopp, P. Ciais, P. Monfray, and J. Orr, IPSL/LSCE, CE-Saclay, 91191, Gif sur Yvette, France. (email: pierre@lsce.saclay.cea.fr; bopp@lsce.saclay.cea.fr; ciais@cea.fr; monfray@cea.fr; orr@cea.fr)
- J.-L. Dufresne, L. Fairhead, and H. LeTreut, IPSL/LMD, Université Paris 6, 75252, Paris, France. (email: Jean-Louis.Dufresne@lmd.jussieu.fr; Laurent.Fairhead@lmd.jussieu.fr; Herve.Letreut@lmd.jussieu.fr)

(Received July 7, 2000; accepted November 27, 2000.)