



HAL
open science

GEOMETRY AND VOLUME PRODUCT OF FINITE DIMENSIONAL LIPSCHITZ-FREE SPACES

Matthew Alexander, Matthieu Fradelizi, Luis García-Lirola, Artem Zvavitch

► **To cite this version:**

Matthew Alexander, Matthieu Fradelizi, Luis García-Lirola, Artem Zvavitch. GEOMETRY AND VOLUME PRODUCT OF FINITE DIMENSIONAL LIPSCHITZ-FREE SPACES. *Journal of Functional Analysis*, 2021, 280 (4), pp.108849. 10.1016/j.jfa.2020.108849 . hal-02924433

HAL Id: hal-02924433

<https://hal.science/hal-02924433>

Submitted on 28 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GEOMETRY AND VOLUME PRODUCT OF FINITE DIMENSIONAL LIPSCHITZ-FREE SPACES

MATTHEW ALEXANDER, MATTHIEU FRADELIZI, LUIS C. GARCÍA-LIROLA, AND ARTEM ZVAVITCH

ABSTRACT. The goal of this paper is to study geometric and extremal properties of the convex body $B_{\mathcal{F}(M)}$, which is the unit ball of the Lipschitz-free Banach space associated with a finite metric space M . We investigate ℓ_1 and ℓ_∞ -sums, in particular we characterize the metric spaces such that $B_{\mathcal{F}(M)}$ is a Hanner polytope. We also characterize the finite metric spaces whose Lipschitz-free spaces are isometric. We discuss the extreme properties of the volume product $\mathcal{P}(M) = |B_{\mathcal{F}(M)}| \cdot |B_{\mathcal{F}(M)}^\circ|$, when the number of elements of M is fixed. We show that if $\mathcal{P}(M)$ is maximal among all the metric spaces with the same number of points, then all triangle inequalities in M are strict and $B_{\mathcal{F}(M)}$ is simplicial. We also focus on the metric spaces minimizing $\mathcal{P}(M)$, and on Mahler's conjecture for this class of convex bodies.

1. INTRODUCTION

Consider a metric space (M, d) containing a special designated point a_0 . Such a pair is usually called a pointed metric space. To this metric space we can associate the space $\text{Lip}_0(M)$ of Lipschitz functions $f : M \rightarrow \mathbb{R}$, with the special property $f(a_0) = 0$. We refer to [G, GK, Os, We] for many interesting facts about $\text{Lip}_0(M)$ and its geometry. It turns out that $\text{Lip}_0(M)$ is a Banach space with norm

$$\|f\|_{\text{Lip}_0(M)} = \sup \left\{ \frac{f(x) - f(y)}{d(x, y)}, \text{ where } x, y \in M, \text{ and } x \neq y \right\}.$$

This space is called the Lipschitz dual of M . The closed unit ball of the space $\text{Lip}_0(M)$ is compact for the topology of pointwise convergence on M , and therefore this space has a canonical predual which is called the Lipschitz-free space over M (also Arens–Eells space or transportation cost space) and is denoted by $\mathcal{F}(M)$. It is the closed subspace of $\text{Lip}_0(M)^*$ generated by the evaluation functionals $\{\delta(x) : x \in M\}$ defined by $\delta(x)(f) = f(x)$, for every $f \in \text{Lip}_0(M)$. The case in which M is a finite metric space was studied recently in [DKO, KMO, OO]. The goal of this paper is to study the geometry of the unit ball of $\mathcal{F}(M)$ when M is finite and, in particular, its volume product.

More precisely, assume our metric space $M = \{a_0, \dots, a_n\}$, is finite with metric d . Note that each function f in $\text{Lip}_0(M)$ is just a set of n values $f(a_1), \dots, f(a_n)$ and thus we can identify $\text{Lip}_0(M)$ with \mathbb{R}^n , assigning to a function $f \in \text{Lip}_0(M)$ a vector $f = (f(a_1), \dots, f(a_n)) \in \mathbb{R}^n$. Let us denote $d_{i,j} = d(a_i, a_j)$, $B_{\text{Lip}_0(M)}$ the unit ball of the $\text{Lip}_0(M)$ and $B_{\mathcal{F}(M)} = B_{\text{Lip}_0(M)}^\circ$, the unit ball of $\mathcal{F}(M)$. Let e_1, \dots, e_n be the standard basis of \mathbb{R}^n and let $e_0 = 0$. For $0 \leq i \neq j \leq n$ let

$$m_{i,j} = \frac{e_i - e_j}{d_{i,j}}.$$

We denote the set of elementary molecules $m_{i,j}$

$$\text{Mol}(M) = \{m_{i,j} : a_i, a_j \in M, i \neq j\}.$$

Then $\|f\|_{\text{Lip}_0(M)} = \max_{m \in \text{Mol}(M)} \langle f, m \rangle$. We note that $B_{\mathcal{F}(M)} = \text{conv}(\text{Mol}(M))$. We shall study the convex bodies in \mathbb{R}^n that can be obtained as unit ball of a Lipschitz-free space. For example, these bodies have at most $n(n+1)$ vertices, but this is not the only constraint as we shall see.

In section 2, we explain the correspondence between metric spaces and weighted graphs that is used throughout the article. To each finite metric space (M, d) one can associate a canonical weighted graph

Date: April, 2020.

2010 Mathematics Subject Classification. Primary: 52A38; Secondary: 52A21, 52A40, 54E45, 05C12.

Key words and phrases. Finite metric space, Volume product, Lipschitz-free space, Kantorovich-Rubinstein polytope, Lipschitz polytope.

The first author is supported in part by the Chateaubriand Fellowship of the Office for Science & Technology of the Embassy of France in the United States and The Centre National de la Recherche Scientifique funding visiting research at Université Paris-Est Marne-la-Vallée; the second author supported in part by the Agence Nationale de la Recherche, project GeMeCoD (ANR 2011 BS01 007 01); the third author is supported in part by the grants MTM2017-83262-C2-2-P and Fundación Séneca Región de Murcia 20906/PI/18 also supported by a postdoctoral grant from Fundación Séneca; the fourth author is supported in part by the U.S. National Science Foundation Grant DMS-1101636 and the Bzout Labex, funded by ANR, reference ANR-10-LABX-58.

$G(M, d)$, the weights on the edges being the distances between the vertices. Then the distance in M corresponds to the shortest path distance in the graph. After removing the unnecessary edges, the edges that appear in the graph model correspond exactly to the vertices of the unit ball of the Lipschitz-free space associated with it. This was recently proved by Aliaga and Guirao [AG] (see also [AP]) in a more general setting: a molecule $m_{i,j}$ is an extreme point of $B_{\mathcal{F}(M)}$ if and only if $d_{i,j} < d_{i,k} + d_{k,j}$ for every $k \notin \{i, j\}$. We give a simpler proof for finite sets in Section 2. We further extend this characterization by proving that the dimension of the face of $B_{\mathcal{F}(M)}$ which contains $m_{i,j}$ in its interior is exactly the number of points in the geodesics (or the metric segment $[i, j]$) joining i to j in the graph, this is our Theorem 2.4. We point out the connection with the so-called Kantorovich-Rubinstein polytopes (studied and popularized in [Ve, GP]).

In Section 3, we explore the conditions for a Lipschitz-free space to be decomposed into ℓ_1 -sum or ℓ_∞ sum of spaces. One way of producing an ℓ_1 -sum is to start from two metric spaces M, N and to consider $M \diamond N$ the metric space obtained by identifying the distinguished points of M and N . Then one has $\mathcal{F}(M \diamond N) = \mathcal{F}(M) \oplus_1 \mathcal{F}(N)$. In fact we prove in Theorem 3.3 that this is the only possible way to get an ℓ_1 -sum and that such a decomposition is unique. We also characterize ℓ_∞ -sums in Theorem 3.6 and use these theorems to describe exactly when a Lipschitz free space can be a Hansen-Lima space, that is a space obtained by taking ℓ_1 or ℓ_∞ sums of ℓ_1^n or ℓ_∞^n equivalently when their unit balls are *Hanner polytopes*.

In Section 4, we introduce the weighted incidence matrix associated with the canonical graph of a metric space. Its kernel is called the cycle space or first homology group of the graph. Using these notions, we prove that two finite metric spaces have isometric Lipschitz-free spaces if and only if their canonical graphs have the same number of vertices and edges and there is a bijection between their cycles which is constant on the 2-connected components.

In Section 5, we investigate the volume product of Lipschitz-free spaces. Recall that the polar body K° of a convex body K is defined by

$$K^\circ = \{y \in \mathbb{R}^n : \langle y, x \rangle \leq 1 \text{ for all } x \in K\}.$$

We note that in most cases in this paper we will assume K to be symmetric. We write $|A|$ to denote the k -dimensional Lebesgue measure (volume) of a measurable set $A \subset \mathbb{R}^n$, where $k = 1, \dots, n$ is the dimension of the minimal flat containing A . Then the volume product of a symmetric convex body K is defined by

$$\mathcal{P}(K) = |K||K^\circ|,$$

and the volume product is invariant under invertible linear transformations on \mathbb{R}^n . The maximum for the volume product in the class of symmetric convex bodies is provided by the Blaschke-Santaló inequality: $\mathcal{P}(K) \leq \mathcal{P}(B_2^n)$, for all symmetric convex bodies $K \subset \mathbb{R}^n$, where B_2^n is the Euclidean unit ball. It was conjectured by Mahler that the minimum of the volume product is attained at the cube in the class of symmetric convex bodies in \mathbb{R}^n , i.e. $\mathcal{P}(K) \geq \mathcal{P}(B_\infty^n) = \mathcal{P}(B_1^n) = \frac{4^n}{n!}$ for all symmetric convex bodies $K \subset \mathbb{R}^n$ where B_∞^n is the unit cube and $B_1^n = (B_\infty^n)^\circ$ is a cross-polytope. The case of $n = 2$ was confirmed by Mahler himself [Ma]. Very recently, Iriyeh and Shibata [IS] gave a positive solution for 3-dimensional symmetric case (see also [FHMRZ]). The conjecture was also proved to be true in several particular special cases, like, e.g., unconditional bodies [SR, Me, R2], convex bodies having hyperplane symmetries which fix only one common point [BF], zonoids [R1, GMR] and bodies of revolution [MR1]. Moreover it was also proved [St, RSW, GM] that bodies with some positive curvature assumption cannot be local minimizers of the volume product. An isomorphic version of the conjecture was proved by Bourgain and Milman [BM]: there is a universal constant $c > 0$ such that $\mathcal{P}(K) \geq c^n \mathcal{P}(B_2^n)$; see also different proofs in [Ku, Na, GPV]. For more information on Mahler's conjecture and the volume product in general, see expository articles [Sc, RZ]. In Section 5, we discuss the maximal and minimal properties of

$$\mathcal{P}(M) := \mathcal{P}(B_{\mathcal{F}(M)})$$

where M is a metric space of fixed number of elements. Our main conjecture here is the following.

Conjecture 1.1. Let M be a metric space with $n + 1$ points. Then $\mathcal{P}(M) \geq \mathcal{P}(B_1^n)$.

We will prove a partial result towards Conjecture 1.1: if $\mathcal{P}(M)$ is minimal and $B_{\mathcal{F}(M)}$ is a simplicial polytope, then M is a tree. It was proved by A. Godard in [Go] that M is a tree if and only if $B_{\mathcal{F}(M)}$ is an affine image of B_1^n , we give a simpler proof of this fact in Section 2. Thus, it would be tempting to say in the conjecture above that the equality is possible if and only if M is a tree. However, for every $n \geq 3$ we can find a metric space giving equality which is not a tree. The reason is the following. If $M = \{0, 1, 2, 3\}$ is the cycle graph, then $B_{\mathcal{F}(M)}$ is a linear image of B_∞^3 . Now, given a tree N with $n - 3$

points then $B_{\mathcal{F}(M \diamond N)} = \text{conv}(B_{\mathcal{F}(M)} \times \{0\}, \{0\} \times B_{\mathcal{F}(N)})$ and so we have that

$$\mathcal{P}(M \diamond N) = \frac{3!(n-4)!}{n!} \mathcal{P}(B_1^3) \mathcal{P}(B_1^{n-4}) = \frac{4^n}{n!}.$$

For the formula of the volume product of the direct sum of two convex bodies, see e.g. [RZ]. In the previous example, $\mathcal{F}(M \diamond N)$ is isometric to $\ell_\infty^3 \oplus_1 \ell_1^{n-4}$. Hanner polytopes are the conjectured minimizers for the volume product among symmetric convex bodies.

It is interesting to note that the maximal value for $\mathcal{P}(M)$ is also an extremely interesting and open problem. Indeed $B_{\mathcal{F}(M)} \neq B_2^n$ for finite M . Thus the maximal case will not follow from the Santaló inequality and we must look for other maximum(s) along with the possible conjectured minimizers. For metric spaces of three points, this maximum is attained in the case in which all the distances $d_{i,j}$ between different points coincide. However, this is no longer true for metric spaces with more than three points. This will follow from the result that the maximum of $\mathcal{P}(M)$ is attained at a metric space such that $B_{\mathcal{F}(M)}$ is a simplicial polytope. We will also prove that all triangle inequalities in M have to be strict for $\mathcal{P}(M)$ to be a maximum.

We will begin with a discussion on the connection between graphs and the Lipschitz-free space of the associated metric space, and we will discuss known results concerning these spaces and the relationship between special graphs and metric spaces. In Section 3 we will analyze when it is possible to decompose $\mathcal{F}(M)$ as an ℓ_1 or an ℓ_∞ -sum and we will apply that to characterize the metric spaces M such that $B_{\mathcal{F}(M)}$ is a Hanner polytope. In Section 4, we characterize the finite metric spaces such that the corresponding Lipschitz-free spaces are isometric. Section 5 is dedicated to the connections of the structure of metric spaces and the corresponding volume product. In particular, we use the *shadow movement* technique to show in Section 5.1 that the maximum of $\mathcal{P}(M)$ is attained at a metric space where all triangle inequalities are strict and $B_{\mathcal{F}(M)}$ is simplicial. In Section 5.2 we compute the volume product corresponding to the complete graph with equal weights. Finally, in Section 5.3 we focus on the metric spaces minimizing $\mathcal{P}(M)$.

2. RELATIONSHIP TO GRAPHS

There is a correspondence between metric spaces and weighted undirected connected graphs. Indeed, to any weighted undirected connected graph $G = (V, E, w)$, with vertices V , edges E and weight $w : E \rightarrow \mathbb{R}_+$ one can associate a finite metric space on its set of vertices V by using the shortest path distance. Reciprocally, to any finite metric space (M, d) , we can canonically associate a weighted undirected connected finite graph $G(M, d)$ as follows: we first consider the complete weighted graph on M with the weight on the edge between two points being their distance. Then one erases the edge between two points x, y if its weight is equal to the sum of the weights of the edges along a path joining the two points, i.e. if there is $z \neq x, y$ such that $d(x, y) = d(x, z) + d(z, y)$. We will say that $G(M, d)$ is the *canonical graph associated with the metric space* (M, d) . We sometimes abuse notation and say that M is a tree, a cycle, etc., meaning that $G(M, d)$ is a tree, a cycle, etc.

The following lemma describes which weighted graphs are the canonical graph of some metric space, it also yields a bijection between those graphs and finite metric spaces.

Lemma 2.1. *Let $G = (V, E, w)$ be a weighted connected graph. Then there is a metric space such that G is the canonical graph associated with the metric if and only if the following holds: for every $\{x, y\} \in E$ and $x_1, \dots, x_l \in V$ such that $x_1 = x, x_l = y$ and $\{x_k, x_{k+1}\} \in E$ for every k , we have $0 < w(x, y) < \sum_{k=1}^l w(x_k, x_{k+1})$.*

Proof. Assume first that there is a metric d on V such that $G = G(M, d)$. Then $0 < d(x, y) = w(x, y)$ for any $\{x, y\} \in E$ and by the triangle inequality $w(x, y) \leq \sum_{k=1}^l w(x_k, x_{k+1})$ for all x_1, \dots, x_l such that $x_1 = x, x_l = y$ and $\{x_k, x_{k+1}\} \in E$ for every k . Moreover, assume that $w(x, y) = \sum_{k=1}^l w(x_k, x_{k+1})$. Then we also have $w(x, y) = w(x, x_2) + w(x_2, y)$ and so $\{x, y\} \notin E$, a contradiction.

Conversely, assume that $G = (V, E)$ satisfies the condition in the statement. Define a metric on V as the shortest path distance:

$$d(x, y) = \min \left\{ \sum_{k=1}^l w(x_k, x_{k+1}) : x_1 = x, x_l = y, \{x_k, x_{k+1}\} \in E \forall k \in \{1, \dots, l\} \right\}$$

Clearly d is a metric on V . We should check that the graph $G' = (V, E')$ associated with (V, d) is G .

Fix $x, y \in V$, $x \neq y$. Then $d(x, y) = \sum_{k=1}^l w(x_k, x_{k+1})$ for some x_1, \dots, x_l such that $x_1 = x, x_l = y$, and $\{x_k, x_{k+1}\} \in E$ for all k . Assume that $\{x, y\} \notin E$. Then $l \geq 3$. Moreover, we should have $d(x_2, y) = \sum_{k=2}^l w(x_k, x_{k+1})$ as otherwise we would have a shorter path for the distance between x and y .

Therefore, $d(x, y) = d(x, x_2) + d(x_2, y)$. So $\{x, y\} \notin E'$. On the other hand, assume that $\{x, y\} \in E \setminus E'$. Then there is $z \in V \setminus \{x, y\}$ such that $d(x, y) = d(x, z) + d(z, y)$. Take $x_1, \dots, x_m, x_{m+1}, \dots, x_l$ such that $x_1 = x$, $x_m = z$, $x_l = y$, $d(x, z) = \sum_{k=1}^{m-1} w(x_k, x_{k+1})$, $d(z, y) = \sum_{k=m}^l w(x_k, x_{k+1})$ and $\{x_k, x_{k+1}\} \in E$ for all k . Then $d(x, y) = \sum_{k=1}^l w(x_k, x_{k+1})$. Moreover, $d(x, y) \leq w(x, y)$ by the definition of d . This is a contradiction with the hypothesis in the statement. Thus $E = E'$, and clearly the weight of the edges in E' is $d(x, y)$. Therefore $G = G(M, d)$. \square

This representation is very well adapted to our study because the edges that appear in the graph model are exactly corresponding to the vertices of the unit ball of the Lipschitz-free space. This was recently proved by Aliaga and Guirao [AG] for compact metric spaces, and even more recently by Aliaga and Pernecká [AP] for complete metric spaces. We give a simpler proof for finite sets below. We denote $\text{ext } K$ the set of extreme points of a convex body K , these are precisely the vertices of K when K is a polytope.

Proposition 2.2 ([AG]). *Let (M, d) be a pointed finite metric space, with $M = \{a_0, \dots, a_n\}$. Let $G = (M, E, w)$ be the canonical associated with (M, d) . The following are equivalent.*

- (i) $m_{i,j} \notin \text{ext } B_{\mathcal{F}(M)}$.
- (ii) There exists $k \notin \{i, j\}$ such that $d_{i,j} = d_{i,k} + d_{k,j}$.
- (iii) There exists $k \notin \{i, j\}$ such that $m_{i,j}$ belongs to the segment between $m_{i,k}$ and $m_{k,j}$.

Proof. (ii) \Rightarrow (iii). Let $k \notin \{i, j\}$ such that $d_{i,j} = d_{i,k} + d_{k,j}$. Then one has

$$m_{i,j} = \frac{e_i - e_j}{d_{i,j}} = \frac{d_{i,k}}{d_{i,k} + d_{k,j}} \frac{e_i - e_k}{d_{i,k}} + \frac{d_{k,j}}{d_{i,k} + d_{k,j}} \frac{e_k - e_j}{d_{k,j}} = \frac{d_{i,k}}{d_{i,k} + d_{k,j}} m_{i,k} + \frac{d_{k,j}}{d_{i,k} + d_{k,j}} m_{k,j}.$$

Hence $m_{i,j}$ belongs to the segment between $m_{i,k}$, and $m_{k,j}$.

(iii) \Rightarrow (i). Let $k \notin \{i, j\}$ such that $m_{i,j}$ belongs to the segment between $m_{i,k}$ and $m_{k,j}$. Since i, j, k are distinct, it follows that $m_{i,j} \neq m_{i,k}$ and $m_{i,j} \neq m_{k,j}$ thus $m_{i,j}$ is not an extreme point of $B_{\mathcal{F}(M)}$.

(i) \Rightarrow (ii). If $m_{i,j}$ is not an extreme point of $B_{\mathcal{F}(M)}$, then $m_{i,j}$ belongs to the relative interior of a face of $B_{\mathcal{F}(M)}$. Let $\mathcal{E} = \{(a_k, a_l) : 0 \leq k \neq l \leq n\}$. For $e = (a_k, a_l) \in \mathcal{E}$ we denote $d_e = d_{k,l}$ and $m_e = m_{k,l}$. Let $\gamma \subset \mathcal{E}$ be the subset of \mathcal{E} of smallest cardinality such that $m_{i,j} \in \text{conv}(m_e : e \in \gamma)$. By Carathéodory's theorem, $2 \leq r \leq n$, where $r = \text{card}(\gamma)$. Then $S_\gamma := \text{conv}(m_e : e \in \gamma)$ is an $(r-1)$ -dimensional simplex. There exists $(\lambda_e)_{e \in \gamma}$ such that $\lambda_e > 0$, for all $e \in \gamma$, $\sum_{e \in \gamma} \lambda_e = 1$ and

$$(1) \quad m_{i,j} = \frac{e_i - e_j}{d_{i,j}} = \sum_{e \in \gamma} \lambda_e m_e.$$

Note that if $(a_k, a_l) \in \gamma$ then $(a_l, a_k) \notin \gamma$. Indeed, otherwise by triangle inequality we will have

$$1 = \|m_{i,j}\| \leq |\lambda_{(a_k, a_l)} - \lambda_{(a_l, a_k)}| + 1 - \lambda_{(a_k, a_l)} - \lambda_{(a_l, a_k)}$$

and so $\lambda_{(a_k, a_l)} = 0$, contradicting the minimality of γ .

Let $M_\gamma = \{a_k \in M : \exists e \in \gamma; a_k \in e\}$ be the vertices of M which belongs to some of the edges in γ . Let $\gamma' = \gamma \cup \{(i, j)\}$. We want to prove that the graph $C_\gamma := (M_\gamma, \gamma')$ is a cycle. First, it is not difficult to see using the minimality of γ that the graph C_γ is connected. Since S_γ is an $(r-1)$ -dimensional simplex whose affine hull doesn't contain the origin, its r vertices are linearly independent. Since these vertices are differences of basis vectors, one has at least $r+1$ basis vectors involved in the vertices of S_γ . Thus, $\text{card}(M_\gamma) \geq r+1$. By linear independence, each $a_k \in M_\gamma$ belongs to at least two edges. By double counting one has

$$2(r+1) = 2 \text{card}(\gamma') = \sum_{a \in M_\gamma} \text{deg}(a) \geq 2 \text{card}(M_\gamma) \geq 2(r+1).$$

Hence $\text{card}(M_\gamma) = r+1$ and each vertex of C_γ has exactly degree two. Thus C_γ is a two-regular connected graph: it is the $(r+1)$ -cycle graph. Then it follows from (1) $d_{i,j} = \sum_{e \in \gamma} \lambda_e d_e$. Since $r \geq 2$ there exists $k \notin \{i, j\}$ such that $k \in e$ for some $e \in \gamma$. Then one has $d_{i,j} = d_{i,k} + d_{k,j}$. \square

One deduces easily the following corollary.

Corollary 2.3. *Let (M, d) be a pointed finite metric space, with $M = \{a_0, \dots, a_n\}$. Let $G = (M, E, w)$ be the canonical graph associated with (M, d) . Let $0 \leq i \neq j \leq n$. Then $m_{i,j}$ is an extreme point of $B_{\mathcal{F}(M)}$ if and only if $\{a_i, a_j\} \in E$. Moreover $B_{\mathcal{F}(M)} = \text{conv}(m_{i,j} : \{a_i, a_j\} \in E)$ and $B_{\mathcal{F}(M)}$ has exactly $2 \text{card}(E)$ vertices.*

Indeed, one can go a bit further and relate the number of points in the metric segment (or geodesic)

$$[i, j] = \{k \in \{0, \dots, n\} : d_{i,j} = d_{i,k} + d_{k,j}\}$$

with the dimension of the face F of $B_{\mathcal{F}(M)}$ such that $m_{i,j}$ is in the relative interior of F . The Proposition 2.2 says that the dimension of F is 0 precisely if $[i, j] = \{i, j\}$. In general, we have the following.

Theorem 2.4. *Let (M, d) be a pointed finite metric space, with $M = \{a_0, \dots, a_n\}$. Given $i \neq j \in \{0, \dots, n\}$, let F be the face of $B_{\mathcal{F}(M)}$ such that $m_{i,j}$ belongs to the relative interior of F . Then*

- a) $F = \text{conv}\{m_{u,v} : d_{i,j} = d_{i,u} + d_{u,v} + d_{v,j}\}$,
- b) the dimension of F coincides with the number of points in $[i, j] \setminus \{i, j\}$.

As a consequence, $m_{i,j}$ belongs to the interior of a facet of $B_{\mathcal{F}(M)}$ if and only if $k \in [i, j]$ for all $k \in \{0, \dots, n\}$. That is a particular case of the study of points of differentiability of the norm of $B_{\mathcal{F}(M)}$ done in [PR, Theorem 4.3] for uniformly discrete metric spaces.

To prove Theorem 2.4, we will need to check when a set of molecules is linearly independent. The following lemma does the work.

Lemma 2.5. *Let (M, d) be a metric space with $n + 1$ elements and $G(M, d) = (V, E, w)$ be its canonical graph. Let $E' \subset E$. Then $\{m_\gamma : \gamma \in E'\}$ is a basis of \mathbb{R}^n if and only if the subgraph with edges E' is a spanning tree of $G(M, d)$.*

Proof. Note that $\text{span}\{m_\gamma : \gamma \in E'\} = \text{span}\left\{\frac{e_i - e_j}{d_{i,j}} : \{a_i, a_j\} \in E'\right\} = \text{span}\{e_i - e_j : \{a_i, a_j\} \in E'\}$. Thus we may assume that $d_{i,j} = 1$ for all $\{a_i, a_j\} \in E'$. Let $G' = (V, E')$. Assume first that G' is a spanning tree of G . Then $\#E' = n$. Consider the (vertex-edge) incidence matrix of G , $Q(G')$, which is defined as follows. We consider that every edge is assigned an orientation, which is arbitrary but fixed. The rows and the columns of $Q(G')$ are indexed by V' and E' , respectively. The (i, j) -entry of $Q(G')$ is 0 if vertex i and edge γ_j are not incident, and otherwise it is 1 or -1 according as γ_j originates or terminates at i , respectively. The rank of $Q(G')$ is n since G' is connected (see e.g. Theorem 2.3 in [Ba]). Thus, the n molecules $\{m_e : e \in E'\}$ are linearly independent and so they are a basis of \mathbb{R}^n .

Conversely, assume that $\{m_e : e \in E'\}$ is a basis. Then $\#E' = n$. Moreover, any cycle on G' would provide a linear dependence relation among the molecules $\{m_e : e \in E'\}$. Thus, G' does not contain any cycle. Finally, G' is connected, since otherwise there would be a non-zero Lipschitz function vanishing on $\{m_e : e \in E'\}$. This shows that G' is a spanning tree of G . □

Proof of Theorem 2.4. Let $u \neq v$ be such that $d_{i,j} = d_{i,u} + d_{u,v} + d_{v,j}$. It is easy to check that every Lipschitz function attaining its norm on $m_{i,j}$ also attains its norm on $m_{u,v}$. Therefore, $m_{u,v}$ belongs to the intersection of all the exposed faces of $B_{\mathcal{F}(M)}$ containing $m_{i,j}$. That intersection is precisely F . Conversely, assume that $m_{u,v} \in F$. Consider the 1-Lipschitz function $f(t) := d(t, a_j)$. Then $\langle f, m_{i,j} \rangle = 1$ and so

$$m_{u,v} \in F \subset \{\mu \in B_{\mathcal{F}(M)} : \langle f, \mu \rangle = 1\}.$$

That is, $\langle f, m_{u,v} \rangle = 1$. So $d_{u,j} - d_{v,j} = d_{u,v}$. Now, consider the function $g(t) := \frac{d_{i,j}}{2} \frac{d(t, a_j) - d(t, a_i)}{d(t, a_j) + d(t, a_i)}$. This is a 1-Lipschitz function considered first in [IKW] that turns out to be very useful when studying the extremal structure of free spaces (see e.g. [GPR]), since it only peaks at points in the segment $[i, j]$. As before, we have $\langle g, m_{i,j} \rangle = 1$ and so

$$m_{u,v} \in F \subset \{\mu \in B_{\mathcal{F}(M)} : \langle g, \mu \rangle = 1\}.$$

That is, $\langle g, m_{u,v} \rangle = 1$. This implies that $d_{i,j} = d_{i,u} + d_{u,j}$. Then

$$d_{i,j} = d_{i,u} + d_{u,j} = d_{i,u} + d_{u,v} + d_{v,j}.$$

It follows that

$$F = \text{conv}\{m_{u,v} : m_{u,v} \in F\} = \text{conv}\{m_{u,v} : d_{i,j} = d_{i,u} + d_{u,v} + d_{v,j}\}.$$

In order to prove b), let $k = \dim F$ and note that $k = \dim \text{span}\{m_{u,v} - m_{i,j} : m_{u,v} \in F\}$. We claim that we only need to consider the vectors $m_{u,v} - m_{i,j}$ where $v \neq j$. Indeed, it also follows from a) that if $m_{u,v} \in F$ then $m_{i,u}, m_{v,j} \in F$. Note also that

$$d_{i,u}(m_{i,u} - m_{x,y}) + d_{u,v}(m_{u,v} - m_{x,y}) + d_{v,j}(m_{v,j} - m_{i,j}) = 0.$$

Thus, $\dim F = \dim \text{span}\{m_{u,v} - m_{i,j} : m_{u,v} \in F, v \neq j\}$. Since the vector $m_{i,j}$ does not belong to $\text{span}\{m_{u,v} : m_{u,v} \in F, v \neq j\}$, we have that $k = \dim \text{span} A$ where

$$A = \{m_{u,v} \in F : v \neq j\} = \{m_{u,v} : d_{i,j} = d_{i,u} + d_{u,v} + d_{v,j}, v \neq j\}.$$

Now, let G be the canonical graph associated with M . We claim that A contains $\#[i, j] \setminus \{i, j\}$ linearly independent vectors. Indeed, consider $V' = [i, j] \setminus \{j\}$. For any $v \in V'$, there is a path in G connecting x and v , note also that $m_{u,v} \in A$ for any u on that path. Let $G' = (V', E')$ be the subgraph of G obtaining by joining together those paths, and let $G'' = (V', E'')$ be a spanning tree of G' . Note that E'' contains $\#[i, j] \setminus \{i, j\}$ edges. Then $m_{u,v} \in A$ for any $(u, v) \in E''$ and Lemma 2.5 tells us that the set $\{m_{u,v}\}_{(u,v) \in E''}$ is linearly independent.

Finally, given a linearly independent set of molecules $\{m_{u,v} : \{u, v\} \in B\} \subset A$, we have that the graph with edges given by the set B does not contain any cycle. Since its nodes belong to the set $[i, j] \setminus \{j\}$, the cardinality of B is at most that of $[i, j] \setminus \{i, j\}$. This shows that there are at most $\#[i, j] \setminus \{i, j\}$ linearly independent vectors in A . □

The following nice relationship between trees and affine images of B_1^n was first proved by A. Godard in a more general setting in section 4 of [Go]. We give here a simpler proof in the case of finite metric spaces.

Proposition 2.6 ([Go]). *Let (M, d) be a pointed finite metric space, with $M = \{a_0, \dots, a_n\}$. Let $G = (M, E, w)$ be the canonical weighted undirected connected finite graph associated with (M, d) . Then G is a tree if and only if $B_{\mathcal{F}(M)}$ is the linear image of B_1^n .*

Proof. Recall that the graph G is a tree if and only if it is connected and acyclic, equivalently it is connected and $\text{card}(E) = \text{card}(M) - 1 = n$. Since our graphs are all connected and using Corollary 2.3, we get that G is a tree if and only if $B_{\mathcal{F}(M)}$ has $2n$ vertices. But we know that $B_{\mathcal{F}(M)}$ is full dimensional and is centrally symmetric so it has exactly $2n$ vertices if and only if it is an affine image of B_1^n . □

Given a metric space M and a subset N endowed with the inherited metric so that $a_0 \in N$, from $B_{\mathcal{F}(M)} = \text{conv}\{m_{i,j} : i, j \in M, i \neq j\}$ it is clear that $\mathcal{F}(N)$ is a subspace of $\mathcal{F}(M)$ and $B_{\mathcal{F}(N)}$ is a section of $B_{\mathcal{F}(M)}$, this will be useful later. The fact that $a_0 \in N$ is not important since the Lipschitz-free operation of a metric space (M, d) with two different chosen roots gives two Banach spaces which are isometric can be seen directly in the following way: Assume that $M = \{a_1, \dots, a_{n+1}\}$. Define

$$K = \text{conv} \left(\left\{ \frac{e_i - e_j}{d_{i,j}} : 1 \leq i \neq j \leq n+1 \right\} \right) \subset \left\{ x \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} x_i = 0 \right\}.$$

Then K is an n -dimensional convex body living in a hyperplane of \mathbb{R}^{n+1} . For $1 \leq i \leq n+1$, denote by $B_{\mathcal{F}_{a_i}(M)}$ the unit ball of the Lipschitz-free space $\mathcal{F}_{a_i}(M)$ pointed at a_i . Then $B_{\mathcal{F}_{a_i}(M)} = P_{e_i^\perp}(K)$ is the orthogonal projection of K on e_i^\perp and this projection is in fact bijective from $\{x \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} x_i = 0\}$ onto $\{x \in \mathbb{R}^{n+1} : x_i = 0\}$. Thus for different i and j , $B_{\mathcal{F}_{a_i}(M)}$ and $B_{\mathcal{F}_{a_j}(M)}$ are affine images of each other. It is also clear from this point that $B_{\mathcal{F}(N)}$ is (isometric to) a section of $B_{\mathcal{F}(M)}$ is $N \subset M$. It is interesting to note that there is a lot of literature dedicated to the above K , called the *fundamental polytope of the metric space* and also the *Kantorovich-Rubinstein polytope*. It was studied in combinatorics starting from the paper by Vershik [Ve], see e.g. [GP]. In the particular case of unweighted graphs, it is called the *symmetric edge polytope*, see [DDM] and references therein.

3. DECOMPOSITIONS OF LIPSCHITZ-FREE SPACES

3.1. ℓ_1 -decompositions. It is interesting to note that the Lipschitz-free Banach space associated with the series composition of two graphs is the ℓ_1 -sum of their Lipschitz-free Banach spaces, in particular, its unit ball is the convex hull of the two unit balls.

Definition 3.1. Let (M_i, d_i) , $i = 1, 2$, be pointed metric spaces. We denote $M_1 \diamond M_2$ the metric space obtained by connecting the graph representations of M_1 and M_2 by identification of their distinguished points.

Note that the definition of $M_1 \diamond M_2$ actually depends on the chosen roots. However, we prefer to avoid the cumbersome notation $(M_1, 0_1) \diamond (M_2, 0_2)$.

If M_i has $n_i + 1$ points, we get that $B_{\mathcal{F}(M_1 \diamond M_2)} = \text{conv}(B_{\mathcal{F}(M_1)} \times \{0\}, \{0\} \times B_{\mathcal{F}(M_2)}) \subset \mathbb{R}^{n_1+n_2}$ and so

$$(2) \quad \mathcal{P}(M_1 \diamond M_2) = \frac{n_1!n_2!}{n!} \mathcal{P}(M_1)\mathcal{P}(M_2),$$

this fact will be very useful when studying the extremal values of the volume product.

Given a finite metric space M , we will say that M is *decomposable* if we can find M_1 and M_2 such that $M = M_1 \diamond M_2$. Otherwise we say that M is *indecomposable*. The decomposability of the metric space is closely related to the biconnectedness of its canonical graph. A connected graph G is called *biconnected* if for any vertex v of G the subgraph obtained by removing v is still connected. A *biconnected component* of G is a maximal biconnected subgraph. Any connected graph decomposes into a tree of biconnected components.

Note that we can write $M = M_1 \diamond M_2$ if and only if the canonical graph associated with M is not biconnected. Moreover, in such a case M_1 and M_2 are the union of biconnected components of M . As a consequence, we have the following:

Proposition 3.2. *Let M be a finite pointed metric space. Then M can be decomposed as $M_1 \diamond \dots \diamond M_r$, where the metric spaces M_i are indecomposable. This decomposition is unique, up to the order of the metric spaces.*

It makes sense to wonder if that is the unique way of decomposing a Lipschitz-free space as an ℓ_1 -sum. It turns that this is the case.

Theorem 3.3. *Let $M = \{a_0, \dots, a_n\}$ be a finite pointed metric space. Assume that $\mathcal{F}(M)$ is isometric to $X_1 \oplus_1 X_2$. Then there exist metric spaces M_1, M_2 (which are obtained from the decomposition of M into its biconnected components) such that X_k is isometric to $\mathcal{F}(M_k)$.*

Proof. Let $G = (M, E)$ be the canonical graph associated with the metric space. Let $\phi: \mathcal{F}(M) \rightarrow X_1 \oplus_1 X_2$ be an isometric isomorphism. Note that

$$\phi(\text{ext } B_{\mathcal{F}(M)}) = \text{ext } B_{X_1} \cup \text{ext } B_{X_2}.$$

Thus, if we denote $E_k = \{\{a_i, a_j\} \in E : \phi(m_{i,j}) \in \text{ext } B_{X_k}\}$ then $\text{ext } B_{X_k} = \{\phi(m_{i,j}) : \{a_i, a_j\} \in E_k\}$, $E = E_1 \cup E_2$ and $E_1 \cap E_2 = \emptyset$.

Now, let $M = \tilde{M}_1 \diamond \dots \diamond \tilde{M}_r$ be the canonical decomposition of M into indecomposable metric spaces, and let \tilde{E}_i be the edges of the i -th biconnected component of M . We claim that each \tilde{E}_i is either contained in E_1 or contained in E_2 . Note that for every pair of distinct edges in \tilde{E}_i there is a cycle containing them. Thus, it suffices to check that every cycle in G is either contained in E_1 or contained in E_2 . Let C be a cycle in G and assume that $C \cap E_1 \neq \emptyset$. We have that

$$0 = \sum_{\{a_i, a_j\} \in C} d_{i,j} m_{i,j} = \sum_{\{a_i, a_j\} \in C \cap E_1} d_{i,j} m_{i,j} + \sum_{\{a_i, a_j\} \in C \cap E_2} d_{i,j} m_{i,j}.$$

Thus, $\sum_{\{a_i, a_j\} \in C \cap E_1} d_{i,j} m_{i,j} \in X_1 \cap X_2 = \{0\}$. This implies that the edges in $C \cap E_1$ form a cycle, and so $C \cap E_2 = \emptyset$. This proves the claim.

For $k = 1, 2$, let $I_k = \{i \in \{1, \dots, r\} : E_i \subset \tilde{E}_k\}$ and consider M_k the metric space obtained by joining the \tilde{M}_i where $i \in I_k$, for $k = 1, 2$ with the same identifications considered before. Clearly, $\mathcal{F}(M)$ is isometric to $\mathcal{F}(M_1) \oplus_1 \mathcal{F}(M_2)$. Moreover, the extreme points of $B_{\mathcal{F}(M_k)}$ correspond, via this isometry, with the molecules $m_{i,j}$ such that $\{i, j\} \in E_k$. Therefore, $\mathcal{F}(M_k)$ is isometric to X_k , and we are done. \square

The previous result can be written in an equivalent way that avoids the use of Lipschitz-free spaces:

Corollary 3.4. *Let $M = \{a_0, \dots, a_n\}$ be a finite pointed metric space. Assume that $\text{Lip}_0(M)$ is isometric to $X_1 \oplus_\infty X_2$. Then there exist metric spaces M_1, M_2 (which are obtained from the decomposition of M into its biconnected components) such that X_i is isometric to $\text{Lip}_0(M_i)$.*

Remark 3.5. Given $1 < p < \infty$, we have that $\lambda^{1/p} x_1 + (1 - \lambda)^{1/p} x_2$ is an extreme point of $B_{X_1 \oplus_p X_2}$ for every $\lambda \in [0, 1]$ and $x_k \in \text{ext } B_{X_k}$, $k = 1, 2$. Thus $B_{X_1 \oplus_p X_2}$ contains an infinite number of extreme points. Therefore, it is not possible to decompose $\mathcal{F}(M)$ as $X_1 \oplus_p X_2$ for finite M .

3.2. ℓ_∞ -decompositions. We say that a metric space $M = \{a_0, a_1, \dots, a_n\}$ is a *spiderweb* if $n = 1$ or if the canonical graph associated with M is the complete bipartite graph $K_{2, n-1}$ where all the edges have the same weight. Note that if M has only two points, then it is a (trivial) spiderweb. In addition, the cycle of length 4 with equal weights is also a spiderweb. The next result shows that for a fixed number of points the spiderweb is the only one metric space whose Lipschitz-free space can be decomposed as an ℓ_∞ -sum.

Theorem 3.6. *Let $M = \{a_0, \dots, a_n\}$ be a finite metric space with $n \geq 3$. Then:*

- a) *If M is a spiderweb, then $\mathcal{F}(M)$ is isometric to $\ell_1^{n-1} \oplus_\infty \mathbb{R}$.*

b) Let X_1, X_2 be Banach spaces with $\dim(X_1) \geq \dim(X_2) \geq 1$. Assume that $\mathcal{F}(M)$ is isometric to $X_1 \oplus_\infty X_2$. Then X_1 is isometric to ℓ_1^{n-1} , $X_2 = \mathbb{R}$ and M is a spiderweb.

The following lemma is the key to prove Theorem 3.6. It characterizes centrally symmetric faces of the ball of a free space. Given $x, y \in M$, let us denote

$$\text{Mid}(x, y) = \{z \in M : 2d(x, z) = 2d(z, y) = d(x, y)\}.$$

Note that M is a spiderweb precisely if there are $x, y \in M$ such that $\text{Mid}(x, y) = M \setminus \{x, y\}$ and for all $z, u \in M \setminus \{x, y\}$ we have $d(z, u) = d(x, y)$.

Lemma 3.7. *Let M be a finite pointed metric space. Assume that F is a centrally symmetric face of $B_{\mathcal{F}(M)}$ with $\dim F \geq 2$. Then one of the following hold:*

a) *There are distinct points $x, y, u, v \in M$ such that $d(x, y) = d(u, v) = d(x, v) = d(u, y)$ and*

$$F = \text{conv}\{m_{x,y}, m_{u,v}, m_{x,v}, m_{u,y}\}.$$

b) *There are $x, y \in M$, $x \neq y$, such that*

$$F = \text{conv}\{m_{x,z}, m_{z,y} : z \in \text{Mid}(x, y)\}.$$

Proof. Let $m_{x,y} \in \text{ext } F \subset \text{ext } B_{\mathcal{F}(M)}$ and denote $m_{u,v}$ the opposite vertex with respect to the center of F . Since $\dim F \geq 2$, there is another vertex $m_{x',y'}$ of F , and let $m_{u',v'}$ be its opposite with respect to the center of F . Note that the middle point of the segments $[m_{x,y}, m_{u,v}]$ and $[m_{x',y'}, m_{u',v'}]$ coincide. That is,

$$(3) \quad m_{x,y} + m_{u,v} = m_{x',y'} + m_{u',v'}$$

We distinguish two cases:

Case 1. $\{x, y\} \cap \{u, v\} = \emptyset$. Then, up to exchanging the role of $m_{x',y'}$ and $m_{u',v'}$, it follows by linear independence that $x' = x$ and $u' = u$. It follows then that $y' = v$ and $v' = y$, and $d(x, y) = d(u, v) = d(x', y') = d(u', v')$. Since $m_{x',y'}$ was an arbitrary vertex of F , we have $F = \text{conv}\{m_{x,y}, m_{u,v}, m_{x,v}, m_{u,y}\}$.

Case 2. $\{x, y\} \cap \{u, v\} \neq \emptyset$. First assume that $u = x$. It follows from (3) that $x' = u' = x$. That is,

$$m_{x,y} + m_{x,v} = m_{x,y'} + m_{x,v'}.$$

Then $\{y', v'\} = \{y, v\}$. Since $m_{x,y} \neq m_{x,v}$, we have $y' = v$ and $v' = y$. That is a contradiction. Therefore $u \neq x$. Analogously, $v \neq y$. So either $u = y$ or $v = x$. Let's assume that we are in the case $u = y$. Then $v \neq x$ since $m_{x,y}$ and $m_{y,x}$ cannot belong to the same face, and

$$m_{x,y} + m_{y,v} = m_{x',y'} + m_{u',v'}.$$

Assume first that $d(x, y) \neq d(y, v)$. Then $\text{supp}\{m_{x',y'} + m_{u',v'}\} = \text{supp}\{m_{x,y} + m_{y,v}\}$ has at most three elements and so $d(x', y') = d(u', v')$. But then we also have $d(x, y) = d(y, v)$, a contradiction.

Thus, $d(x, y) = d(y, v)$, and it follows that they are also equal to $d(x', y')$ and $d(u', v')$. Therefore

$$e_x - e_v = e_{x'} - e_{y'} + e_{u'} - e_{v'}$$

and so $\{m_{x',y'}, m_{u',v'}\} = \{m_{x,z}, m_{z,v}\}$ for some $z \in M$ satisfying that $d(x, z) = d(z, v) = d(x, y) = d(y, v)$. That happens for any vertex of F . Thus,

$$F = \text{conv}(\text{ext } F) \subset \text{conv}\{m_{x,z}, m_{z,v} : d(x, z) = d(z, v)\}$$

and $m_{x,z} \in F$ if and only if $m_{z,v} \in F$. Let $f \in S_{\text{Lip}_0(M)}$ be an exposing functional for F . Pick z such that $m_{x,z}, m_{z,v} \in F$. Then $f(x) - f(z) = d(x, z)$ and $f(z) - f(v) = d(z, v)$. Thus,

$$d(x, v) \geq f(x) - f(v) = (f(x) - f(z)) + (f(z) - f(v)) = d(x, z) + d(z, v)$$

and so $z \in \text{Mid}(x, y)$ and $f(x) - f(v) = d(x, v)$. This means that

$$F \subset \text{conv}\{m_{x,z}, m_{z,v} : z \in \text{Mid}(x, v)\}.$$

Moreover, for every $z \in \text{Mid}(x, y)$ we have

$$d(x, v) = f(x) - f(v) = (f(x) - f(z)) + (f(z) - f(v)) \leq d(x, z) + d(z, v) = d(x, v)$$

and so $m_{x,z}, m_{z,v} \in F$. This shows that

$$F = \text{conv}\{m_{x,z}, m_{z,v} : z \in \text{Mid}(x, v)\}$$

as desired. The case $v = x$ is analogous and we get that $F = \text{conv}\{m_{u,z}, m_{z,y} : z \in \text{Mid}(u, y)\}$. □

Note that the previous result shows that if $F_1 \subset F_2$ are centrally symmetric faces of $B_{\mathcal{F}(M)}$ and F_1 has at least 6 vertices, then $F_1 = F_2$.

Proof of Theorem 3.6. Assume first that M is a spiderweb, that is, there are $x, y \in M$ such that $\text{Mid}(x, y) = M \setminus \{x, y\}$ and that $m_{u,v} \notin \text{ext } B_{\mathcal{F}(M)}$ if $u, v \in M \setminus \{x, y\}$. Up to isometry, we may assume that x is the distinguished point of M and $d(x, y) = 2$. Consider the 1-Lipschitz function f given by $f(x) = 0$, $f(y) = 2$ and $f(z) = 1$ otherwise. Let $F = \{m \in B_{\mathcal{F}(M)} : \langle f, m \rangle = 1\}$. Note that

$$F = \text{conv}\{m_{z,x}, m_{y,z} : z \in M \setminus \{x, y\}\}.$$

Then $\text{ext } B_{\mathcal{F}(M)} \subset F \cup (-F)$, so F is a facet of $B_{\mathcal{F}(M)}$. Moreover,

$$\frac{m_{z,x} + m_{y,z}}{2} = m_{y,x}$$

for all $z \in M \setminus \{x, y\}$, so F is symmetric with respect to $m_{y,x}$.

Let $K = F - m_{y,x}$. Then $K = B_Y$ for some Banach space Y . Therefore,

$$B_{\mathcal{F}(M)} = \text{conv}(F \cup (-F)) = B_Y + [m_{x,y}, m_{y,x}]$$

and so $\mathcal{F}(M) = Y \oplus_{\infty} \mathbb{R}$. Finally, it is clear that $\#\text{ext}(B_Y) = \#\text{ext}(F) = 2(n-1)$ and so Y is isometric to ℓ_1^{n-1} . This proves a).

Now, assume that $\mathcal{F}(M)$ is isometric to $X_1 \oplus_{\infty} X_2$ and that $\dim(X_2) \geq 2$. Take $u, v \in \text{ext } B_{X_2}$ such that $[u, v]$ is an edge of B_{X_2} . Then both $B_{X_1} + u$ and $B_{X_1} + [u, v]$ are centrally symmetric facets of $B_{X_1 \oplus_{\infty} X_2}$.

Case 1. Assume $\dim(X_1) \geq 3$. Then $\#\text{ext}(B_{X_1} + u) = \#\text{ext}(B_{X_1}) \geq 6$. Therefore Lemma 3.7 yields $B_{X_1} + u = B_{X_1} + [u, v]$, a contradiction.

Case 2. Suppose now that $\dim(X_1) = \dim(X_2) = 2$ and so $n = 4$. Then $B_{X_1} + [u, v]$ is a facet of $B_{X_1 \oplus_{\infty} X_2}$ with at least 8 vertices. By Lemma 3.7, there are x, y in M such that the set

$$\text{conv}\{m_{x,z}, m_{z,y} : z \in \text{Mid}(x, y)\}$$

has 8 vertices. But this is impossible since $\#\text{Mid}(x, y) \leq 3$.

Therefore, $X_2 = \mathbb{R}$. Then we have $B_{\mathcal{F}(M)} = B_{X_1} + [-u_0, u_0]$, for a certain vector u_0 . Thus, $F = B_{X_1} + u_0$ is a centrally symmetric facet of $B_{\mathcal{F}(M)}$. Now, we distinguish again two cases:

Case 1. Assume $n = 3$, that is, M has four points. By Lemma 3.7, we have two possible cases:

Case 1.1. $F = \text{conv } A$, where $A = \{m_{x,y}, m_{u,v}, m_{x,v}, m_{u,y}\} \subset \text{ext } B_{\mathcal{F}(M)}$, and $d(x, y) = d(u, v) = d(x, v) = d(u, y)$. Then $M = \{x, y, u, v\}$ and $\#\text{ext } B_{X_1} = \#\text{ext } F = 4$. Therefore, $B_{\mathcal{F}(M)}$ has precisely 8 vertices, the ones in $A \cup (-A)$. That is, the edges of the canonical graph associated with M are $\{x, y\}$, $\{u, v\}$, $\{x, v\}$ and $\{u, y\}$, and all of them have the same weight. So M is a cycle of length 4 with equal weights, in particular, a spiderweb.

Case 1.2. There are $x, y \in M$ such that $F = \text{conv}\{m_{x,z}, m_{z,y} : z \in \text{Mid}(x, y)\}$. Since F has at least 4 vertices, we have that $\text{Mid}(x, y) = M \setminus \{x, y\}$ and $\#\text{ext } F = 4$. Then $\#\text{ext } B_{\mathcal{F}(M)} = 8$. It follows that M is again a cycle of length 4 with equal weights.

Case 2. Assume $n \geq 4$. We have $\#\text{ext } F = \#\text{ext } B_{X_1} \geq 2(n-1) \geq 6$. Then the first case in Lemma 3.7 does not hold. Therefore, there are $x, y \in M$ such that $F = \text{conv } A$, where $A = \{m_{x,z}, m_{z,y} : z \in \text{Mid}(x, y)\}$. Note that $\text{ext } F \subset A$ and $\text{ext } B_{\mathcal{F}(M)} \subset \text{ext } F \cup \text{ext } (-F)$. It follows that for every $z \in M \setminus \{x, y\}$ we have $\{x, z\}, \{z, y\} \in E$, and all the edges are of this form. Thus, M is a spiderweb. Moreover, it follows that $\#\text{ext } B_{X_1} = 2(n-1)$ and so X_1 is isometric to ℓ_1^{n-1} .

□

Combining Proposition 3.9 and Theorem 3.6, we get the following result.

Corollary 3.8. *Let $M = \{a_0, \dots, a_n\}$ be a finite pointed metric space. Then $\mathcal{F}(M)$ is isometric to ℓ_{∞}^n if and only if $n \leq 2$ and M is tree, or $n = 3$ and M is a cycle with equal weights.*

3.3. Zonotopes. Godard [Go] characterized the metric spaces M such that $\mathcal{F}(M)$ is isometric to a subspace of L_1 as those metric spaces which embed into an \mathbb{R} -tree. In the finite-dimensional setting, the embeddability into L_1 is equivalent to the fact that the dual ball $B_{\text{Lip}_0(M)}$ is a zonoid [Bo] (see also [Sc, Ko, RZ]). Let us recall that a convex body is said to be a *zonotope* if it is a finite Minkowski sum of segments, and it is said to be a *zonoid* if it is the limit, in the Hausdorff distance, of a sequence of zonotopes. Both notions coincide for polytopes. Zonoids enjoy very nice geometrical properties, for instance, all faces are also zonoids and thus centrally symmetric. We refer the reader to [Sc, GW] for more results about zonoids and zonotopes.

Zonoids satisfy Mahler's conjecture [R1, GMR]. Thus, it makes sense to wonder about when $B_{\mathcal{F}(M)}$ is a zonoid. Unfortunately, there are few cases when that happens, as the following result shows.

Proposition 3.9. *Let $M = \{a_0, \dots, a_n\}$ be a finite pointed metric space. The following are equivalent:*

- i) $B_{\mathcal{F}(M)}$ is a zonotope.
- ii) $n \leq 2$, or $n = 3$ and M is a cycle graph (of length 4) with equal weights.

Proof. ii) \Rightarrow i). Every 2-dimensional symmetric polytope is a zonotope. In addition, if $n = 3$ and the canonical graph associated with M is the cycle graph with equal weights then $B_{\mathcal{F}(M)}$ is a linear image of the cube B_∞^3 .

i) \Rightarrow ii). Let F be a (centrally symmetric) facet of $B_{\mathcal{F}(M)}$. Assume that $n \geq 4$. Since F is a zonoid of dimension $n - 1$, it has at least 2^{n-1} vertices. In addition, by Lemma 3.7, there are x, y such that $F = \text{conv}\{m_{x,z}, m_{z,y} : d(x, z) = d(z, y)\}$. Then

$$2^{n-1} \leq \#\text{ext } F \leq 2\#\{z \in M : d(x, z) = d(z, y)\} \leq 2(n-1),$$

a contradiction. Thus, $n \leq 3$. In the case $n = 2$ there is nothing to prove. Thus, we may assume that $n = 3$. Let $G = (V, E, w)$ be the canonical graph associated with M . We distinguish some cases:

Case 1. G has at least one leaf, say a_0 . Then $\mathcal{F}(M)$ is a ℓ_1 -sum of \mathbb{R} and another space. It is easy to check that then all the facets of $B_{\mathcal{F}(M)}$ are not centrally symmetric.

Case 2. There is a node in G with degree 3. We may assume that the node is a_0 . Consider the 1-Lipschitz function f given by $f(a_i) = d_{i,0}$. Note that if $i, j \neq 0$ are distinct then $|f(a_i) - f(a_j)| < d_{i,j}$. Thus, $\langle f, m_{i,j} \rangle = 1$ if, and only if, $m_{i,j} = m_{i,0}$ and $i \neq 0$. That is, the face of $B_{\mathcal{F}(M)}$ given by f has dimension 2 and an odd number of vertices, so it is not centrally symmetric, a contradiction.

Since the previous cases lead to a contradiction, we have that G is a cycle, so $B_{\mathcal{F}(M)}$ has 8 vertices. Thus, F has 4 vertices. It follows that, for any $\{x, y\} \in E$, either $m_{x,y} \in F$ or $m_{y,x} \in F$. Lemma 3.7 says that, in any case, the set $\{d(x, y) : \{x, y\} \in E\}$ is a singleton. That is, all the edges have the same weight, as desired. □

3.4. Hanner polytopes. A symmetric convex body K is called a *Hanner polytope* if K is one-dimensional, or it is the ℓ_1 or ℓ_∞ sum of two (lower dimensional) Hanner polytopes. They are the unit balls of the Hansen-Lima spaces [HL]. Hanner polytopes are the conjectured minimizers for the volume product, moreover they are known to be local minimizers [NPRZ, Ki]. We finish the section characterizing for which metric spaces the ball of the Lipschitz-free space is a Hanner polytope.

Theorem 3.10. *Let M be a finite pointed metric space. The following are equivalent:*

- i) $B_{\mathcal{F}(M)}$ is a Hanner polytope.
- ii) $M = M_1 \diamond \dots \diamond M_r$, where each of the M_i is a spiderweb.

Proof. ii) \Rightarrow i) follows from the fact that $B_{\mathcal{F}(M_i)}$ is a Hanner polytope provided M_i is a spiderweb and the connection between the volume product of $B_{\mathcal{F}(M)}$ and the one of the $B_{\mathcal{F}(M_i)}$ given in (2).

i) \Rightarrow ii). We prove the result by induction on $n = \dim \mathcal{F}(M)$. For $n \leq 2$, every Hanner polytope is a linear image of B_1^n . So M is a tree, which is a sum of spiderwebs consisting on two points. Now, assume $n \geq 3$ and that the result is true for metric spaces with at most n points. Write $\mathcal{F}(M) = X_1 \oplus_p X_2$, where $p \in \{1, \infty\}$ and B_{X_1}, B_{X_2} are Hanner polytopes. If $p = \infty$, then M is a spiderweb by Theorem 3.6, and we are done. Otherwise, $p = 1$. Let $M = M_1 \diamond \dots \diamond M_r$ be the decomposition of M into its biconnected components. Then by Theorem 3.3, there are metric spaces $N_1 = M_{i_1} \diamond \dots \diamond M_{i_s}$ and $N_2 = M_{j_1} \diamond \dots \diamond M_{j_{r-s}}$, with $\{1, \dots, r\} = \{i_1, \dots, i_s, j_1, \dots, j_{r-s}\}$, such that X_i is isometric to $\mathcal{F}(N_i)$ for $i = 1, 2$. The induction hypothesis says that both N_1 and N_2 are the sum of spiderwebs. Since the decomposition of N_1 and N_2 into their biconnected components is unique, it follows that each of the M_i is a spiderweb, as desired. □

4. ISOMETRIES OF LIPSCHITZ-FREE SPACES

It is known that if the canonical graph associated with a metric space M is a complete graph, and $\mathcal{F}(M)$ is isometric to $\mathcal{F}(M')$, then M is isometric to M' [We, Theorem 3.55], equivalently, $B_{\mathcal{F}(M)}$ is a linear image of $B_{\mathcal{F}(M')}$. This does not hold in general, for instance $\mathcal{F}(M) = \ell_1^n$ whenever M is a tree. Indeed, the Lipschitz-free spaces over two metric spaces M and M' are isometric whenever M and M' have the same decomposition into indecomposable metric spaces. Our next goal is to characterize when two finite metric spaces M, M' have isometric Lipschitz-free spaces.

First, given a metric space with canonical graph $G = (V, E, d)$, we identify $\mathcal{F}(M)$ with a quotient of $\ell_1(E)$. To this end, consider the operator $\partial : \mathbb{R}^E \rightarrow \mathbb{R}^V$ defined by $\partial \chi_e = m_e$ for any $e \in E$ and extended to \mathbb{R}^E by linearity, where χ denotes the indicator function, so for $u, v \in V$, $\chi_u : \mathbb{R}^V \rightarrow \mathbb{R}$ and $\chi_u(u) = 1$

and $\chi_u(v) = 0$ for $v \neq u$. The *cycle space* is the kernel of ∂ . This kernel is also called the first homology group $H_1(G) = H_1(G, \mathbb{R})$.

Theorem 4.1. *Let M be a finite metric space and $G = (V, E, d)$ be its canonical graph. Then $\mathcal{F}(M)$ is isometric to $\ell_1(E)/\text{Ker } \partial$.*

Proof. We start by fixing an orientation of the edges. This orientation means that each edge $e \in E$ has a vertex e_+ which is a target and a vertex e_- which is a source and one has $e = (e_-, e_+)$ and moreover the molecule associated with e is well defined, it is $m_e = \frac{\chi_{e_+} - \chi_{e_-}}{d(e)}$. Then

$$B_{\mathcal{F}(M)} = \text{conv}\{\pm m_e : e \in E\} \subset \mathbb{R}^V \cap \{\varphi \in \mathbb{R}^V : \sum_{v \in V} \varphi(v) = 0\} := H$$

where H is a hyperplane in \mathbb{R}^V . Moreover $B_{\mathcal{F}(M)}$ is a convex set of dimension $|V| - 1$, so of full dimension in this hyperplane and it has exactly $|E|$ vertices. Let us now see how one can compute the norm in $\mathcal{F}(M)$ in relation with the ℓ_1 norm in \mathbb{R}^E defined by $\|f\|_1 = \sum_{e \in E} |f(e)|$. For $\varphi \in \mathbb{R}^V$, since $B_{\mathcal{F}(M)} = \text{conv}\{\pm m_e; e \in E\}$, one has

$$\begin{aligned} \|\varphi\|_{\mathcal{F}(M)} &= \inf\{t > 0; \varphi \in tB_{\mathcal{F}(M)}\} = \inf\{t > 0; \varphi = t \sum_{e \in E} t_e m_e; \sum_{e \in E} |t_e| = 1\} \\ &= \inf\left\{\sum_{e \in E} |t_e|; \varphi = \sum_{e \in E} t_e m_e\right\} = \inf\left\{\sum_{e \in E} |t_e|; \varphi = \partial\left(\sum_{e \in E} t_e \chi_e\right)\right\} \\ &= \inf\{\|f\|_1; \varphi = \partial f\}. \end{aligned}$$

This implies that for every $f \in \mathbb{R}^E$ one has

$$\|\partial f\|_{\mathcal{F}(M)} = \inf\{\|g\|_1; \partial f = \partial g\} = \inf\{\|f + h\|_1; h \in \text{Ker } \partial\} = \|f\|_{\ell_1(E)/\text{Ker } \partial}.$$

Hence one has $\mathcal{F}(M) = \ell_1(E)/\text{Ker } \partial$. Another way of seeing this is to say that $B_{\mathcal{F}(M)}$ is the projection of B_1^E , the ℓ_1 ball in \mathbb{R}^E .

□

Note that the previous result also says that $\text{Lip}_0(M)$ is isometric to the subspace $(\text{Ker } \partial)^\perp$ of $\ell_\infty(E)$. Indeed, the isometry is given by the map

$$f \mapsto \left(\frac{f(e^+) - f(e^-)}{d(e)} \right)_{e \in E}$$

Now, we can characterize which finite metric spaces have the same Lipschitz-free space. Given two graphs G and G' , we say that a bijection between their sets of edges is *cyclic* if it induces a bijection between the cycles of G and the cycles of G' .

Theorem 4.2. *Let M, M' be finite metric spaces with canonical graphs $G = (V, E, d)$ and $G' = (V', E', d')$. The following are equivalent:*

- i) $\mathcal{F}(M)$ is isometric to $\mathcal{F}(M')$, that is, there is a linear isomorphism T such that $T B_{\mathcal{F}(M)} = B_{\mathcal{F}(M')}$.
- ii) $|V| = |V'|$, $|E| = |E'|$, and $H_1(G)$ and $H_1(G')$ are isometric to the same subspace of ℓ_1^m , where $m = |E| = |E'|$.
- iii) $|V| = |V'|$ and there is a cyclic bijection $\sigma: E \rightarrow E'$ such that the function $e \mapsto d(\sigma(e))/d(e)$ is constant on each 2-connected component of G .

Moreover, any isometry $T: \mathcal{F}(M) \rightarrow \mathcal{F}(M')$ is induced by a cyclic bijection $\sigma: E \rightarrow E'$, meaning that $T(m_e) = \pm m'_{\sigma(e)}$ for all $e \in E$.

Notice that, in particular, we get that two unweighted graphs have the same Lipschitz-free space if and only if there is a cyclic bijection between their edges. In such a case, the graphs are said to be *cyclically equivalent* or *2-isomorphic*. H. Whitney [Wh] showed that two graphs are cyclically equivalent if and only if one can pass from one to the other by the successive application of two operations which are:

- (a) Vertex gluing and vice versa. Vertex gluing of M and M' is the same as what we denoted by $M \diamond M'$. The opposite operation, the separation of a graph into two non-connected components, can be done only at a “separating vertex”.
- (b) Switching: starting from a pair (u, v) of separating vertices one exchanges u and v .

Proof of Theorem 4.2. ii) \Rightarrow i). By Theorem 4.1, $\mathcal{F}(M) = \ell_1(E)/\text{Ker } H_1(G)$ and $\mathcal{F}(M') = \ell_1(E')/\text{Ker } H_1(G')$, these spaces are isometric.

i) \Rightarrow ii) and iii). If $B_{\mathcal{F}(M')} = TB_{\mathcal{F}(M)}$ then $\text{conv}\{\pm m'_e; e' \in E'\} = \text{conv}\{\pm Tm_e; e \in E\}$, where $m_e = \frac{\chi_{e^+} - \chi_{e^-}}{d(e)}$, for $e \in E$ and $m_e = \frac{\chi_{e^+} - \chi_{e^-}}{d(e)}$, for $e \in E$. Hence these convex sets have the same dimension and the same number of vertices. This implies that $|V| = |V'|$, $|E| = |E'| = m$. Moreover since T establishes a bijection among the extreme points, there exists a unique bijection $\sigma : E \rightarrow E'$ and a map $\varepsilon : E \rightarrow \{-1, 1\}$ such that for any $e \in E$ one has $T(m_e) = \varepsilon(e)m'_{\sigma(e)}$. We claim that σ is a cyclic bijection. Indeed, let $\{e_1, \dots, e_k\}$ be a cycle in E . Then $\sum_{i=1}^k d(e_i)m_{e_i} = 0$ and so

$$(4) \quad 0 = \sum_{i=1}^k d(e_i)\varepsilon(e_i)m_{\sigma(e_i)} = \sum_{i=1}^k \frac{d(e_i)}{d(\sigma(e_i))}\varepsilon(e_i)(\chi_{e_i^+} - \chi_{e_i^-}).$$

Then the degree of any vertex in the subgraph of G' induced by $\{\sigma(e_1), \dots, \sigma(e_k)\}$ is odd and so $\{\sigma(e_1), \dots, \sigma(e_k)\}$ is union of cycles in G' . If $I \subset \{1, \dots, k\}$ is so that $\{\sigma(e_i) : i \in I\}$ is a cycle in G' , then applying the same argument with T^{-1} we get that $\{e_i : i \in I\}$ is union of cycles in G . Thus $I = \{1, \dots, k\}$ and the image of the cycle $\{e_1, \dots, e_k\}$ is a cycle in G' . From (4) we have that $d(e_i)/d(\sigma(e_i)) = d(e_j)/d(\sigma(e_j))$ for all i, j . Thus, the function $e \mapsto d(\sigma(e))/d(e)$ is constant on each cycle, and so on each 2-connected component of G .

Now we prove ii). We denote $\partial : \mathbb{R}^E \rightarrow \mathbb{R}^V$ defined by $\partial\chi_e = m_e$ and $\partial' : \mathbb{R}^{E'} \rightarrow \mathbb{R}^{V'}$ defined by $\partial'\chi_{e'} = m'_{e'}$. From Theorem 4.1 one has $\mathcal{F}(M) = \ell_1(E)/\text{Ker } \partial$ and $\mathcal{F}(M') = \ell_1(E')/\text{Ker } \partial'$. Let us define $\tilde{T} : \mathbb{R}^E \rightarrow \mathbb{R}^{E'}$ by $\tilde{T}(\chi_e) = \varepsilon(e)\chi_{\sigma(e)}$ and extended by linearity. Then for any $e \in E$ one has

$$T(\partial(\chi_e)) = T(m_e) = \varepsilon(e)m'_{\sigma(e)} = \varepsilon(e)\partial'(\chi_{\sigma(e)}) = \partial'(\varepsilon(e)\chi_{\sigma(e)}) = \partial'(\tilde{T}(\chi_e)).$$

Thus $T \circ \partial = \partial' \circ \tilde{T}$. And since T and \tilde{T} are bijective this implies that

$$H_1(G) = \text{Ker}(\partial) = \text{Ker}(T \circ \partial) = \text{Ker}(\partial' \circ \tilde{T}) = (\tilde{T})^{-1}(\text{Ker}(\partial')) = (\tilde{T})^{-1}(H_1(G')).$$

Thus $H_1(G') = \tilde{T}(H_1(G))$, where \tilde{T} is an isometry from $\ell_1(E)$ to $\ell_1(E')$.

iii) \Rightarrow i). Clearly σ preserves the 2-connected components of the graph. Since $\mathcal{F}(M)$ is isometric to $\mathcal{F}(M_1) \oplus_1 \dots \oplus_1 \mathcal{F}(M_r)$ provided that $M = M_1 \diamond \dots \diamond M_r$, we may assume that G is 2-connected and so there is $a > 0$ such that $d(\sigma(e)) = ad(e)$ for all $e \in E$. We claim that there is an orientation on the edges of G' such that “ σ preserves directed cycles”, i.e. for every $e_1, \dots, e_k \in E$ and $\varepsilon_1, \dots, \varepsilon_k \in \{-1, 1\}$ such that $\sum_{i=1}^k \varepsilon_i(\chi_{e_i^+} - \chi_{e_i^-}) = 0$, we also have $\sum_{i=1}^k \varepsilon_i(\chi_{\sigma(e_i)^+} - \chi_{\sigma(e_i)^-}) = 0$.

The 2-connectness of G and Whitney’s theorem imply that, up to a graph automorphism, every cyclic bijection is a composition of switchings. Thus, it suffices to prove the case in which σ is a switching on a pair (u, v) of separating vertices of G , that is, we can write $E = E_1 \cup E_2$ and $E' = E_1 \cup E'_2$, where $\{u, x\} \in E_2 \iff \{v, x\} \in E'_2$ and $\{v, x\} \in E_2 \iff \{u, x\} \in E'_2$. Define $\sigma(e)^+ := u$ if $e^+ = v$ and $e \in E_2$, $\sigma(e)^+ := v$ if $e^+ = u$ and $e \in E_2$, and $\sigma(e)^+ := e^+$ otherwise, this orientation does the work.

Let \tilde{E} be a subset of E giving a spanning tree. Since σ preserves cycles, $\sigma(\tilde{E})$ is a spanning tree for G' , and we can choose an orientation of G' in the same way. Now, we can define a linear map $T : \mathcal{F}(M) \rightarrow \mathcal{F}(M')$ given by $T(m_e) = m_{\sigma(e)}$ if $e \in \tilde{E}$ and extended by linearity. Since \tilde{E} and $\sigma(\tilde{E})$ are spanning trees of G and G' , $\{m_e : e \in \tilde{E}\}$ is a basis on $\mathcal{F}(M)$ and $\{m_{\sigma(e)} : e \in \tilde{E}\}$ is a basis of $\mathcal{F}(M')$. Thus, T is an isomorphism. It remains to check that every vertex of $B_{\mathcal{F}(M)}$ maps to a vertex of $B_{\mathcal{F}(M')}$. This is clearly true for the ones of the form $\pm m_e, e \in \tilde{E}$. Let $e \in E \setminus \tilde{E}$. Then there are $e_1, \dots, e_r \in \tilde{E}$ such that $\{e_1, \dots, e_r, e\}$ is a cycle in G and $\chi_{e^+} - \chi_{e^-} = \sum_{i=1}^r \varepsilon_i(\chi_{e_i^+} - \chi_{e_i^-})$. Then, thanks to the choice of the orientation of the edges in G' , we have $\chi_{\sigma(e)^+} - \chi_{\sigma(e)^-} = \sum_{i=1}^r \varepsilon_i(\chi_{\sigma(e_i)^+} - \chi_{\sigma(e_i)^-})$ and so

$$\begin{aligned} T(d(e)m_e) &= T(\chi_{e^+} - \chi_{e^-}) = \sum_{i=1}^r \varepsilon_i T(\chi_{e_i^+} - \chi_{e_i^-}) = \sum_{i=1}^r \varepsilon_i d(e_i) T(m_{e_i}) = \sum_{i=1}^r \varepsilon_i d(e_i) m_{\sigma(e_i)} \\ &= \sum_{i=1}^r \varepsilon_i \frac{d(e_i)}{d(\sigma(e_i))} (\chi_{\sigma(e_i)^+} - \chi_{\sigma(e_i)^-}) = \frac{1}{a} (\chi_{\sigma(e)^+} - \chi_{\sigma(e)^-}) = d(e)m_{\sigma(e)} \end{aligned}$$

and so $T(m_e) = m_{\sigma(e)}$, which is a vertex of $B_{\mathcal{F}(M')}$ as desired. \square

In the case of 3-connected graphs, the result of Whitney mentioned before yields every cycle bijection $\sigma: E \rightarrow E'$ is induced by a vertex isomorphism, that is, there is $f: V \rightarrow V'$ such that $\sigma(e) = (f(e^+), f(e^-))$ for all e . As a consequence, we obtain the following result.

Corollary 4.3. *Let (M, d) and (M', d) be metric spaces such that $TB_{\mathcal{F}(M)} = B_{\mathcal{F}(M')}$ for a linear isomorphism T . Assume that $G(M, d)$ is 3-connected. Then M and M' are isometric, moreover, there is an isometry $f: M \rightarrow M'$ inducing T , i.e. $T(m_e) = m'_{f(e^+), f(e^-)}$ for all any edge e in $G(M, d)$.*

Note that the previous corollary improves the result of Weaver [We, Theorem 3.55] in the case of finite metric spaces.

5. EXTREMAL PROPERTIES OF THE VOLUME PRODUCT

We will focus now on the maximal and the minimal value of $\mathcal{P}(M) = |B_{\mathcal{F}(M)}| \cdot |B_{\text{Lip}_0(M)}|$. It is a well-known fact that the volume product of convex bodies is invariant under linear isometries. We start with the easy observation that $\mathcal{P}(M)$ is invariant under dilations on M . Given two metric spaces (M, d) and (N, ρ) , we say that a map $f: M \rightarrow N$ is a *dilation* if it is a bijection and there is a constant $a > 0$ such that $\rho(f(x), f(y)) = ad(x, y)$ for every $x, y \in M$.

Proposition 5.1. *$\mathcal{P}(M)$ is invariant under dilations of M .*

Proof. Let (M, d) and (N, ρ) be finite metric spaces and denote $\delta_M: M \rightarrow \mathcal{F}(M)$ and $\delta_N: N \rightarrow \mathcal{F}(N)$ the canonical embeddings. Assume that $f: M \rightarrow N$ is a bijection such that $\rho(f(x), f(y)) = ad(x, y)$ for some $a > 0$ and every $x, y \in M$. By the fundamental property of Lipschitz-free spaces (see e.g. [We]), there is a bounded linear operator $\hat{f}: \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ such that $\hat{f} \circ \delta_M = \delta_N \circ f$. Therefore

$$\begin{aligned} \hat{f}(B_{\mathcal{F}(M)}) &= \hat{f} \left(\text{conv} \left\{ \frac{\delta_M(x) - \delta_M(y)}{d(x, y)} : x, y \in M, x \neq y \right\} \right) \\ &= \text{conv} \left\{ \frac{\hat{f}(\delta_M(x)) - \hat{f}(\delta_M(y))}{d(x, y)} : x, y \in M, x \neq y \right\} \\ &= a \text{conv} \left\{ \frac{\delta_N(f(x)) - \delta_N(f(y))}{\rho(f(x), f(y))} : x, y \in M, x \neq y \right\} = aB_{\mathcal{F}(N)} \end{aligned}$$

That is, $B_{\mathcal{F}(N)}$ is a linear image of $B_{\mathcal{F}(M)}$. Then $\mathcal{P}(N) = \mathcal{P}(M)$. □

The main idea behind a number of proofs of our results in this section is the shadow system technique: a *shadow system of convex sets* along a direction $\theta \in S^{n-1}$ is a family of convex sets $L_t \in \mathbb{R}^n$ which are defined by

$$L_t = \text{conv}\{x + \alpha(x)t\theta : x \in B\},$$

where $B \subset \mathbb{R}^n$ is a bounded set, called the basis of the shadow system, $\alpha: B \rightarrow \mathbb{R}$ is a bounded function, called the speed of the shadow system, and t belongs to an open interval in \mathbb{R} . We say that a shadow system is *non-degenerate*, if all the convex sets L_t have non-empty interior. Shadow systems were first introduced by Rogers and Shephard [RS]. Campi and Gronchi [CG] proved that if L_t is a symmetric shadow system then $t \mapsto |L_t^\circ|^{-1}$ is a convex function of t . In [MR2], Meyer and Reisner generalized this result to the non-symmetric case and studied the equality case. The following proposition summarize those results in symmetric case:

Proposition 5.2 ([CG, MR2]). *Let L_t , $t \in [-a, a]$, be a non-degenerate shadow system in \mathbb{R}^n , with direction $\theta \in S^{n-1}$, then $t \mapsto |L_t^\circ|^{-1}$ is a convex function on $[-a, a]$.*

If, moreover, $t \mapsto |L_t|$ is affine on $[-a, a]$ and $t \mapsto \mathcal{P}(L_t)$ is constant on $[-a, a]$, then there exists $w \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$, such that for every $t \in [-a, a]$, one has $L_t = A_t(I_0)$, where $A_t: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the affine map defined by

$$A_t(x) = x + t(w \cdot x + \alpha)\theta.$$

5.1. Maximal Case. We will show that the maximum of the volume product is attained at a metric space such that its canonical graph is a weighted complete graph. In dimension two we can say something more.

Theorem 5.3. *For all metric spaces of three elements, $\mathcal{P}(M) \leq \mathcal{P}(K_3)$, where K_3 is the metric space corresponding to a complete graph with equal weights.*

Proof. The result follows immediately from Theorem 4 in [AFZ], but here we present a self-contained proof. Let $M = \{0, 1, 2\}$ be a metric space of three elements. By Proposition 5.1 we may assume that $d_{1,2} = 1$, so $B_{\mathcal{F}(M)} = \text{conv}\{\pm \frac{e_1}{d_{1,0}}, \pm \frac{e_2}{d_{2,0}}, \pm(e_1 - e_2)\}$. Then after a linear transformation we get the body $L = \text{conv}\{\pm e_1, \pm e_2, \pm(d_{1,0}e_1 - d_{2,0}e_2)\}$ with the same volume product. Note that $|d_{1,0} - d_{2,0}| \leq 1$. Consider the shadow system

$$L_t = \text{conv}\left\{\pm e_1, \pm e_2, \pm\left(\frac{d_{1,0} + d_{2,0}}{2}(e_1 - e_2) + t(e_1 + e_2)\right)\right\}$$

Then $t \mapsto |L_t|$ is constant on $[-1/2, 1/2]$ and $|L_t^\circ| = |L_{-t}^\circ|$. Thus, $\mathcal{P}(L_t) \leq \mathcal{P}(L_0)$ for all $t \in [-1/2, 1/2]$. In particular, $\mathcal{P}(M) = \mathcal{P}(L) \leq \mathcal{P}(L_0)$. Now, direct computation shows that $|L_0| = 1 + 2r$ and $|L_0^\circ| = \frac{4r-1}{r^2-1}$, where $r = \frac{d_{1,0} + d_{2,0}}{2}$. By simple calculus, $\mathcal{P}(L_0) \leq \mathcal{P}(K_3) = 9$.

□

To deal with the maximum in general dimension we need two results about shadow systems. The following lemma is an extension of Lemma 3.2 of [AFZ], where a similar result is proved in the case in which F is a facet of K .

Lemma 5.4. *Let $K \subset \mathbb{R}^n$ be a convex body with $0 \in \text{int } K$ and F be a face of K of dimension $0 \leq k \leq n-1$. Let $\{F_i\}_{i=1}^m$ be the facets of K containing F , with unit normal vectors $\{u_i\}_{i=1}^m$. Let $x_F \in \text{relint } F$ and $v \in \mathbb{S}^{n-1}$ be such that $\langle v, u_i \rangle > 0$ for all $i = 1, \dots, m$. Take $x_t = x_F + tv$ and $K_t = \text{conv}(K, x_t)$. Then, if $t > 0$ is small enough,*

$$|K_t| = |K| + \frac{t}{n} \sum_{i=1}^m |F_i| \langle v, u_i \rangle$$

and $|K_t^\circ| = |K| + o(t^{k+1})$.

Proof. Let $\{F_i\}_{i=m+1}^l$ be the facets of K that do not contain F , with normal unit vectors $\{u_i\}_{i=m+1}^l$. Let us denote by $h_K(u) = \sup\{\langle x, u \rangle : x \in K\}$ the support function of K . Note that $\langle x_F, u_i \rangle < h_K(u_i)$ for every $i = m+1, \dots, l$ since $x_F \in \text{relint } F$. Thus, there is $\varepsilon > 0$ such that we have $\langle x_t, u_i \rangle < h_K(u_i)$ for every $i = m+1, \dots, l$ if $t \in [0, \varepsilon]$. In addition, note that

$$K_t = \text{conv}(\text{ext } K \cup \{x_t\})$$

and so $\text{ext } K_t \subset \text{ext } K \cup \{x_t\}$.

Claim 1. $K_t = K \cup \bigcup_{i=1}^m \text{conv}(F_i, x_t)$ if $t \in [0, \varepsilon]$.

It is clear that $K \cup \bigcup_{i=1}^m \text{conv}(F_i, x_t) \subset K_t$. Conversely, it suffices to check that every edge of K_t is contained in $K \cup \bigcup_{i=1}^m \text{conv}(F_i, x_t)$. Let p, q be such that the segment $[p, q]$ is an edge of K_t . Then $p, q \in \text{ext } K_t$. If both of them belong to K , then $[p, q] \subset K$. Otherwise, we may assume that $p = x_t$ and $q \in \text{ext } K$. Assume that $q \notin \bigcup_{i=1}^m F_i$. Then $\langle q, u_i \rangle < h_K(u_i)$ for all $i \in \{1, \dots, m\}$. Thus,

$$\langle \lambda x_t + (1 - \lambda)q, u_i \rangle < h_K(u_i) \quad \forall i \in \{1, \dots, m\}$$

when $\lambda > 0$ is small enough. Moreover,

$$\langle \lambda x_t + (1 - \lambda)q, u_i \rangle < h_K(u_i) \quad \forall i \in \{m+1, \dots, l\}$$

for all $0 < \lambda < 1$. This means that $[p, q] \cap \text{int } K \neq \emptyset$, so $[p, q]$ also intersects $\text{int } K_t$ and so it is not an edge. Thus, $q \in \bigcup_{i=1}^m F_i$. Therefore $[p, q] \subset \text{conv}(F_i, x_t)$ for some $i \in \{1, \dots, m\}$ as desired. This proves the claim.

Claim 2. $\text{conv}(F_i, x_t) \cap \text{conv}(F_j, x_t) = \text{conv}(F_i \cap F_j, x_t)$ if $i, j \in \{1, \dots, m\}$.

Let $y \in \text{conv}(F_i, x_t) \cap \text{conv}(F_j, x_t)$. We can write

$$y = \lambda x_t + (1 - \lambda)y_i = \mu x_t + (1 - \mu)y_j$$

where $y_i \in F_i$, $y_j \in F_j$, and $0 \leq \lambda, \mu \leq 1$. Clearly we may assume $\lambda, \mu < 1$. Assume first that $\mu > \lambda$. If $y_i \in F_j$, then $y \in \text{conv}(F_i \cap F_j, x_t)$ and we are done. So we may assume $\langle y_i, u_j \rangle < h_K(u_j)$. Then we have

$$\langle y, u_j \rangle = \lambda \langle x_t, u_j \rangle + (1 - \lambda) \langle y_i, u_j \rangle < \lambda \langle x_t, u_j \rangle + (1 - \lambda) h_K(u_j).$$

On the other hand,

$$\langle y, u_j \rangle = \mu \langle x_t, u_j \rangle + (1 - \mu) \langle y_j, u_j \rangle = \mu \langle x_t, u_j \rangle + (1 - \mu) h_K(u_j)$$

so $(\mu - \lambda) \langle x_t, u_j \rangle < (\mu - \lambda) h_K(u_j)$, which yields $\langle x_t, u_j \rangle < h_K(u_j)$, a contradiction. If $\mu < \lambda$, we reach to a similar contradiction multiplying by u_i . Finally, assume that $\mu = \lambda < 1$. Then $y_i = y_j \in F_i \cap F_j$ and we are done. This proves the claim.

It follows from Claim 2 that $\text{conv}(F_i, x_t) \cap \text{conv}(F_j, x_t)$ has empty interior if $i \neq j$. Thus,

$$|K_t| = |K| + \sum_{i=1}^m |\text{conv}(F_i, x_t)| = |K| + \frac{1}{n} \sum_{i=1}^m |F_i| \langle x_t, u_i \rangle - h_K(u_i) = |K| + \frac{t}{n} \sum_{i=1}^m |F_i| \langle v, u_i \rangle.$$

Now, we focus on the volume of the polar body. Note that the vertices of K° are the points $\{u_i/h_K(u_i)\}_{i=1}^l$. Let $F' = \{x \in K^\circ : \langle x, y \rangle = 1 \forall y \in F\}$ be the face of K° corresponding to F . Then F' is the convex hull of the vertices of K° which belong to F' and so $F' = \text{conv}(\{u_i/h_K(u_i)\}_{i=1}^m)$.

For each $i \in \{1, \dots, m\}$, let $\{v_s^i\}_{s=1}^{r_i}$ be the vertices of K° which are adjacent to $u_i/h_K(u_i)$ and which do not belong to $\{u_j/h_K(u_j)\}_{j=1}^l$. Let

$$\eta = \min\{1 - \langle v_s^i, x_F \rangle : i \in \{1, \dots, m\}, s \in \{1, \dots, r_i\}\} > 0.$$

Note that $u_i/h_K(u_i) \in K_t^\circ$ if and only if $\langle u_i/h_K(u_i), x_t \rangle \leq 1$ if and only if $i \in \{m+1, \dots, l\}$. This means that

$$K^\circ = K_t^\circ \cup \text{conv}(u_i/h_K(u_i), w_s^i : i \in \{1, \dots, m\}, s \in \{1, \dots, r_i\}),$$

where w_s^i is the unique point in the segment $[u_i/h_K(u_i), v_s^i]$ with $\langle w_s^i, x_t \rangle = 1$. Write

$$w_s^i = (1 - \lambda_s^i) \frac{u_i}{h_K(u_i)} + \lambda_s^i v_s^i$$

where $0 \leq \lambda_s^i \leq 1$. An easy computation shows that

$$\lambda_s^i = \frac{t \langle \frac{u_i}{h_K(u_i)}, v \rangle}{\langle \frac{u_i}{h_K(u_i)} - v_s^i, x_t \rangle} \leq \frac{t \langle \frac{u_i}{h_K(u_i)}, v \rangle}{1 - \langle v_s^i, x_F \rangle - \varepsilon \left| \frac{u_i}{h_K(u_i)} - v_s^i \right|} \leq \frac{t \langle u_i, v \rangle}{h_K(u_i)(\eta - \varepsilon \text{diam}(K^\circ))} \leq ct$$

for some constant $c > 0$, provided ε is small enough. Therefore,

$$|u_i/h_K(u_i) - w_s^i| = \lambda_s^i |u_i/h_K(u_i) - v_s^i| \leq c \text{diam}(K^\circ) t = c't$$

for each $i = 1, \dots, m$ and $s = 1, \dots, r_i$. This means that

$$K^\circ \setminus K_t^\circ \subset \text{conv} \left(\bigcup_{i=1}^m B(u_i/h_K(u_i), c't) \right) = \text{conv}(\{u_i/h_K(u_i)\}_{i=1}^m) + c'tB = F' + c'tB.$$

Now, by Steiner's formula (see e.g. [Sc]),

$$|F' + c'tB| = \sum_{j=0}^n (c't)^{n-j} \kappa_{n-j} V_j(F') = o(t^{k+1})$$

since the affine dimension of F' is $n - k + 1$ (see e.g. [Gr, pag. 50]). Thus, $|K^\circ \setminus K_t^\circ| = o(t^{k+1})$. \square

Lemma 5.5. *Let $K \subset \mathbb{R}^n$ be a symmetric convex body and F be a face of K of dimension $1 \leq k \leq n - 1$. Let $\{F_i\}_{i=1}^m$ be the facets of K containing F , with unit normal vectors $\{u_i\}_{i=1}^m$. Let $x_F \in \text{relint } F$ and $v \in \mathbb{S}^{n-1}$ be such that $\langle v, u_i \rangle > 0$ for all $i = 1, \dots, m$. Take $x_t = x_F + tv$ and $K_t = \text{conv}(K, x_t, -x_t)$. Then, $\mathcal{P}(K_t) > \mathcal{P}(K)$ if $t > 0$ is small enough.*

Proof. Let $L_t = \text{conv}(K, x_t)$. It is easy to check that for each $i, j \in \{1, \dots, m\}$, the set $\text{conv}(F_i, x_t) \cap \text{conv}(-F_j, -x_t)$ has empty interior. Now, the argument in the proof of the previous lemma gives that

$$(L_t \cap (-L_t)) \setminus K = \bigcup_{i,j=1}^m \text{conv}(F_i, x_t) \cap \text{conv}(-F_j, -x_t)$$

and so $|(L_t \cap (-L_t)) \setminus K| = 0$. By Lemma 5.4 we have

$$|K_t| = |K| + |L_t \setminus K| + |(-L_t) \setminus K| = |K| + 2 \frac{t}{n} \sum_{i=1}^m |F_i| \langle v, u_i \rangle.$$

On the other hand, $K_t^\circ = \{x \in K : |\langle x, x_t \rangle| \leq 1\} = L_t^\circ \cap (-L_t)^\circ$ and so

$$|K^\circ \setminus K_t^\circ| \leq |K^\circ \setminus L_t^\circ| + |K^\circ \setminus (-L_t)^\circ| = o(t^{k+1})$$

if $t > 0$ is small enough. Thus $|K_t^\circ| = |K^\circ| + o(t^{k+1})$. Therefore,

$$\mathcal{P}(K_t) = \mathcal{P}(K) + 2|K^\circ| \frac{t}{n} \sum_{i=1}^m |F_i| \langle v, u_i \rangle + |K| o(t^{k+1}) > \mathcal{P}(K)$$

when t approaches 0.

□

We remark that the previous lemma does not work when $k = 0$. Indeed, if K is a square in \mathbb{R}^2 and x_F is a vertex of K then we can have $\mathcal{P}(K_t) = \mathcal{P}(K)$ for all $t > 0$.

Now we can show that the maximum of $\mathcal{P}(M)$ is attained at a weighted complete graph.

Theorem 5.6. *Let $M = \{a_0, \dots, a_n\}$ be a metric space such that $\mathcal{P}(M)$ is maximal among the metric spaces with the same number of elements. Then $d_{i,j} < d_{i,k} + d_{k,j}$ for every distinct points a_i, a_j, a_k .*

Proof. Assume the contrary. Consider the set

$$A = \{(i, j) \in \{0, \dots, n\}^2 : i \neq j, d_{i,j} = d_{i,k} + d_{k,j} \text{ for some } k \in \{0, \dots, n\} \setminus \{i, j\}\}.$$

Take $(i, j) \in A$ such that $d_{i,j}$ is maximal. Consider d^t given by $d_{i,j}^t = \frac{d_{i,j}}{1+t}$ and $d_{u,v}^t = d_{u,v}$ otherwise. The maximality of (i, j) implies that $d_{i,j} > \max\{|d_{i,k} - d_{k,j}| : k \in \{0, \dots, n\} \setminus \{i, j\}\}$. Thus, d^t is a metric on M for t small enough. Note that $\frac{e_i - e_j}{d_{i,j}^t} = (1+t)m_{i,j}$. Thus,

$$B_{\mathcal{F}(M, d^t)} = \text{conv}(\{m_{u,v} : \{u, v\} \neq \{i, j\}\}, \pm(1+t)m_{i,j}) = \text{conv}(B_{\mathcal{F}(M)}, \pm(1+t)m_{i,j}).$$

Note that $m_{i,j}$ is not an extreme point of $B_{\mathcal{F}(M)}$ since $(i, j) \in A$. Thus, $m_{i,j} \in \text{relint } F$ for some face F of $B_{\mathcal{F}(M)}$ with dimension k , $1 \leq k \leq n-1$. Let F_1, \dots, F_m be the facets of $B_{\mathcal{F}(M)}$ containing F , with normal vectors $\{u_s\}_{s=1}^m$. Note that $\langle m_{i,j}, u_s \rangle = h_{B_{\mathcal{F}(M)}}(u_s) > 0$. Therefore, Lemma 5.5 with $x_F = v = m_{i,j}$ yields that $\mathcal{P}(M, d^t) > \mathcal{P}(M, d)$ if $t > 0$ is small enough, a contradiction.

□

It is natural to wonder if the maximum of the volume product is attained at the unweighted complete graph. We showed in Theorem 5.1 that this is the case for metric spaces with three points. However, that is no longer the case in higher dimensions. In order to prove that, we will show that the maximum of $\mathcal{P}(M)$ is attained at a metric space such that $B_{\mathcal{F}(M)}$ is simplicial.

The following can be proved by using the same arguments as in Theorem 3.4 in [AFZ].

Proposition 5.7. *Let \mathcal{K} be a family of centered convex polytopes and let $K \in \mathcal{K}$ be such that $\mathcal{P}(K)$ is maximal among the elements of \mathcal{K} . Assume that $\text{conv}(K, (1+t)x, -(1+t)x) \in \mathcal{K}$ for every vertex x of K and $t > 0$ small enough. Then K is simplicial.*

In the particular case of free spaces, we have the following.

Theorem 5.8. *Let $M = \{a_0, \dots, a_n\}$ be a finite pointed metric space such that $\mathcal{P}(M)$ is maximal among the metric spaces with the same number of elements. Then $B_{\mathcal{F}(M)}$ is simplicial.*

Proof. Note that every extreme point of $B_{\mathcal{F}(M)}$ is of the form $m_{i,j}$ for some $a_i, a_j \in M$, $i \neq j$. By Theorem 5.6, we have $d_{i,j} > \max\{|d_{i,k} - d_{k,j}| : k \in \{0, \dots, n\} \setminus \{i, j\}\}$. Thus, d^t given by $d_{u,v}^t = d_{u,v}$ if $\{u, v\} \neq \{i, j\}$ and $d_{i,j}^t = \frac{d_{i,j}}{1+t}$ defines a metric on M for t small enough. Moreover,

$$B_{\mathcal{F}(M, d^t)} = \text{conv}(\{m_{u,v} : \{u, v\} \neq \{i, j\}\}, \pm(1+t)m_{i,j}) = \text{conv}(B_{\mathcal{F}(M)}, \pm(1+t)m_{i,j}).$$

This shows that the hypotheses of Proposition 5.7 hold and so $B_{\mathcal{F}(M)}$ is simplicial.

□

5.2. The special case of complete graph with equal weights. Let K_{n+1} denote the metric space of $n+1$ elements such that $d_{i,j} = 1$ if $i \neq j$, i.e. the metric space associated with the complete graph where all the weights are equal to 1.

Note that $B_{\mathcal{F}(K_{n+1})}$ is not simplicial whenever $n \geq 3$. Indeed, consider the 1-Lipschitz function given by $f(a_i) = 1$ if $i \in \{2, \dots, n\}$ and $f(a_0) = f(a_1) = 0$. Note that the $2n-2$ molecules $m_{i,0}, m_{i,1}$, $i \in \{2, \dots, n\}$ are vertices of $B_{\mathcal{F}(K_{n+1})}$ that belong to the face of $B_{\mathcal{F}(K_{n+1})}$ exposed by f . Since $2n-2 > n$, this face is not a simplex. Therefore, it follows from Proposition 5.8 that K_{n+1} is not a metric space with maximum volume product if $n \geq 3$.

We also would like to provide a computation for the volume product of K_{n+1} . Note first that $B_{\mathcal{F}(K_{n+1})} = \text{conv}\{\pm e_i, \pm(e_i - e_j); 1 \leq i \neq j \leq n\}$ has exactly $n(n+1)$ vertices.

Let us describe more precisely the unit ball of $\text{Lip}_0(K_{n+1})$.

Claim 3.

$$B_{\text{Lip}_0(K_{n+1})} = \text{conv} \left\{ \pm \sum_{i \in I} e_i : I \subset \{1, \dots, n\} \right\}.$$

Proof. Denote by C the right hand side set. First let us prove that $C \subset B_{\text{Lip}_0(K_{n+1})}$. One has

$$B_{\text{Lip}_0(K_{n+1})} = B_{\mathcal{F}(K_{n+1})}^\circ = \{x \in \mathbb{R}^n : |x_i| \leq 1, |x_i - x_j| \leq 1 \text{ for all } 1 \leq i \neq j \leq n\}.$$

For any $I \subset \{1, \dots, n\}$ let us denote $x(I) = \sum_{i \in I} e_i$. For any $k, l \in \{1, \dots, n\}$ one has $x(I)_k \in \{0, 1\}$ and $x(I)_k - x(I)_l \in \{-1, 0, 1\}$ hence $x(I) \in B_{\mathcal{F}(K_{n+1})}^\circ$. Therefore $C \subset B_{\text{Lip}_0(K_{n+1})}$.

To show that $B_{\mathcal{F}(K_{n+1})}^\circ \subset C$ we consider $x \in B_{\text{Lip}_0(K_{n+1})}$. Then $|x_i| \leq 1$ and so we may assume, reordering our axes if necessary, that $-1 \leq x_1 \leq x_2 \leq \dots \leq x_{n-1} \leq x_n \leq 1$. Further, since $|x_i - x_j| \leq 1$ we get $x_n - x_1 \leq 1$. Now let us consider the indices with positive entries and negative entries separately. That is, we let k be the last negative index, choosing it to be 0 if no entries are negative, and to be n if all entries are negative. Then

$$\begin{aligned} x = (x_2 - x_1)(-e_1) + (x_3 - x_2)(-e_1 - e_2) + \dots + (x_k - x_{k-1}) \left(-\sum_{i \leq k-1} e_i \right) + (-x_k) \left(-\sum_{i \leq k} e_i \right) \\ + x_{k+1} \sum_{i \geq k+1} e_i + (x_{k+2} - x_{k+1}) \sum_{i \geq k+2} e_i + \dots + (x_n - x_{n-1}) e_n, \end{aligned}$$

thus $\frac{1}{x_n - x_1} x$ is a convex combination of points in C . Therefore $B_{\text{Lip}_0(K_{n+1})} \subset C$. □

Claim 4.

$$B_{\text{Lip}_0(K_{n+1})} = \frac{1}{2} B_\infty^n + \frac{1}{2} \left[-\sum_{i=1}^n e_i, \sum_{i=1}^n e_i \right].$$

Proof. Denote by D the zonotope on the right hand side. First let us take an extreme point $x \in D$ then

$$x = \frac{1}{2} \sum_{i=1}^n \varepsilon_i e_i + \frac{1}{2} \varepsilon_{n+1} \sum_{i=1}^n e_i = \sum_{i=1}^n \frac{\varepsilon_i + \varepsilon_{n+1}}{2} e_i.$$

where the $\varepsilon_i \in \{-1, 1\}$. Let $I = \{i \in \{1, \dots, n\} \mid \varepsilon_i = \varepsilon_{n+1}\}$. Then

$$x = \varepsilon_{n+1} \sum_{i \in I} e_i \in B_{\text{Lip}_0(K_{n+1})}.$$

To see that $B_{\text{Lip}_0(K_{n+1})} \subset D$ we simply reverse our previous observation. So if we take $x \in C$ so $x = \varepsilon_{n+1} \sum_{i \in I} e_i$ where $I \subset \{1, \dots, n\}$. Then we define $\varepsilon_i = \varepsilon_{n+1}$ if $i \in I$ and $\varepsilon_i = -\varepsilon_{n+1}$ if $i \notin I$. Then by our choices of ε_i we have $x = \frac{1}{2} \sum_{i=1}^n \varepsilon_i e_i + \frac{1}{2} \varepsilon_{n+1} \sum_{i=1}^n e_i \in D$ as desired. □

Remark 5.9. It follows from Claim 4 that $B_{\text{Lip}_0(K_{n+1})}$ is a zonotope. This can be seen in a different way (see [Go] and also [Os, Example 10.13]) by showing that K_{n+1} embeds into a tree with $n+2$ points or by showing that $B_{\mathcal{F}(K_{n+1})}$ is a section of ℓ_1^{n+1} . We also note that using the fact that the Mahler conjecture is true for zonotopes [GMR] we get $\mathcal{P}(K_{n+1}) \geq \mathcal{P}(B_1^n)$. We will show it as a direct computation below.

Let us compute the volume product of K_{n+1} .

Claim 5.

$$\mathcal{P}(K_{n+1}) = \frac{n+1}{n!} \binom{2n}{n}.$$

Proof. Let us first compute the volume of $B_{\text{Lip}_0(K_{n+1})}$. Let $e = \sum_{i=1}^n e_i$. Then $B_{\text{Lip}_0(K_{n+1})}$ is a zonotope which is the following sum of $n+1$ segments:

$$B_{\text{Lip}_0(K_{n+1})} = \frac{1}{2} \left(\sum_{i=1}^n [-e_i, e_i] + [-e, e] \right).$$

Thus one may use the following formula for the volume of zonotopes [Sh]: if $Z = \sum_{i=1}^m [0, u_i]$ is a zonotope which is the sum of m vectors u_1, \dots, u_m in \mathbb{R}^n , with $m \geq n$ then

$$|Z| = \sum_{\text{card}(I)=n} |\det(u_i)_{i \in I}|.$$

Since for all $k \in \{1, \dots, n\}$ one has $|\det(e, (e_i)_{i \neq k})| = 1$ we get $|B_{\text{Lip}_0(K_{n+1})}| = n+1$.

Now let us compute the volume of $B_{\mathcal{F}(K_{n+1})} = \text{conv}\{\pm e_i, \pm(e_i - e_j); 1 \leq i \neq j \leq n\}$. We decompose $B_{\mathcal{F}(K_{n+1})}$ using the partition of \mathbb{R}^n into 2^n parts defined according to the coordinate signs. For $I \subset \{1, \dots, n\}$ let

$$C_I = \{x \in \mathbb{R}^n : x_i > 0 \text{ for all } i \in I \text{ and } x_j < 0 \text{ for all } j \notin I\}.$$

Then

$$B_{\mathcal{F}(K_{n+1})} \cap C_I = \text{conv}(0; (e_i)_{i \in I}; (-e_j)_{j \notin I}; (e_i - e_j)_{i \in I, j \notin I}) = \text{conv}(0; (e_i)_{i \in I}) - \text{conv}(0; (e_j)_{j \notin I}).$$

Hence if we denote $k = \text{card}(I)$ then

$$|B_{\mathcal{F}(K_{n+1})} \cap C_I| = |\text{conv}(0; (e_i)_{i \in I})|_k |\text{conv}(0; (e_j)_{j \notin I})|_{n-k} = \frac{1}{\text{card}(I)! \text{card}(I^c)!} = \frac{1}{k!(n-k)!}.$$

Thus

$$|B_{\mathcal{F}(K_{n+1})}| = \sum_{I \subset \{1, \dots, n\}} \frac{1}{\text{card}(I)! \text{card}(I^c)!} = \sum_{k=0}^n \binom{n}{k} \frac{1}{k!(n-k)!} = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k}^2 = \frac{1}{n!} \binom{2n}{n}.$$

□

Remark 5.10. Note that a more general case is the metric space for which $d_{i,j} \in \mathbb{Z}$ for all i, j . In this case the polytope $B_{\text{Lip}_0(M)}$ has vertices in the lattice \mathbb{Z}^n . Indeed, a result of Farmer [Fa] ensures that a Lipschitz function f is an extreme point of $B_{\text{Lip}_0(M)}$ if and only if for all i, j there are k_0, k_1, \dots, k_l such that $k_0 = i$, $k_l = j$ and

$$|f(a_{k_r}) - f(a_{k_{r+1}})| = d(a_{k_r}, a_{k_{r+1}})$$

for all $r = 0, \dots, l-1$, and so $f(a_j) \in \mathbb{Z}$ for every $a_j \in M$.

If, moreover, all the edges of the canonical graph associated with M have weight 1, then the vertices of $B_{\mathcal{F}(M)}$ are of the form $e_i - e_j$ for some i, j and so $B_{\mathcal{F}(M)}$ is also a polytope with vertices in the lattice \mathbb{Z}^n . Thus it is interesting to ask for minimal and maximal volume product among lattice polytopes whose dual is also a lattice polytope.

5.3. The Minimal Case. In this section, we focus on Conjecture 1.1. Note that it is already known that this conjecture holds for certain metric spaces. Namely:

- If M embeds into a tree, then $B_{\text{Lip}_0(M)}$ is a zonoid [Go] and so the result follows from [R1, GMR].
- If M is a cycle, then $B_{\mathcal{F}(M)}$ has $2n + 2$ vertices and so the result follows from the recent paper [Ka].

We will show some more cases where Conjecture 2.3 holds.

Given a connected graph $G = (V, E)$, we say that an edge $e \in E$ is a *bridge* if the subgraph $(V, E \setminus \{e\})$ is disconnected.

Proposition 5.11. *Let (M, d) be a finite metric space such that $\mathcal{P}(M)$ is minimal. Let $m_{i,j}$ be a vertex of $B_{\mathcal{F}(M)}$. Assume that all the facets of $B_{\mathcal{F}(M)}$ containing $m_{i,j}$ are simplices. Then, the edge $\{a_i, a_j\}$ is a bridge in $G(M, d)$.*

Proof. Let $G = (V, E, w)$, be the graph of M . Let $G^t = (V, E, w^t)$ be the weighted graph obtained from G by replacing the weights by $w^t(\{a_i, a_j\}) = w(\{a_i, a_j\})/(1+t)$ and $w^t(\{u, v\}) = w(\{u, v\})$ for any $\{u, v\} \in E \setminus \{\{a_i, a_j\}\}$. By Lemma 2.1, if $|t|$ is small enough then there is a metric d^t on M such that $G^t = G(M, d^t)$. Moreover, $t \mapsto |B_{\mathcal{F}(M, d^t)}|$ is affine if $|t|$ is small enough since all the faces containing $m_{i,j}$ are simplices. By the minimality of $\mathcal{P}(M)$ and Lemma 2 from [FMZ], $B_{\mathcal{F}(G)}$ must be a double-cone with apex $m_{i,j}$. Thus, $\text{span}(\text{ext}(B_{\mathcal{F}(M)}) \setminus \{\pm m_{i,j}\})$ has dimension $n - 1$. We claim that this implies that $\{a_i, a_j\}$ is a bridge in G . Indeed, otherwise there would exist a spanning tree in G not containing the edge $\{a_i, a_j\}$, and the vector space generated by the corresponding molecules would be n -dimensional by Lemma 2.5.

□

Let us remark that it is also possible to give an alternative proof of Proposition 5.11 using the characterization of isometric Lipschitz-free spaces given in Theorem 4.2 and the properties of a shadow system. Indeed, by [MR2] we have that $B_{\mathcal{F}(M, d^t)}$ is a linear image of $B_{\mathcal{F}(M)}$. Then there is a cyclic bijection between the edges of $G(M, d)$ and the ones of $G(M, d^t)$ such that d_t/d is constant on each 2-connected component. Thus the edge $\{a_i, a_j\}$ is the only element of its 2-connected component, that is, it is a bridge.

Theorem 5.12. *Let (M, d) be a metric space with minimal volume product such that $B_{\mathcal{F}(M)}$ is simplicial. Then M is a tree.*

Proof. Apply Proposition 5.11 to get every edge in $G(M, d)$ is a bridge. □

We note that the minimal case for four points corresponds to the question on the minimality of volume product in \mathbb{R}^3 . That question was recently solved in [IS] and automatically gives $\mathcal{P}(M) \geq \mathcal{P}(B_1^3)$, for M being a metric space of four elements. Here we present a direct proof that uses the structure of polytope of $B_{\mathcal{F}(M)}$. To that end, we need the following simple observation.

Lemma 5.13. *Assume that there is a face of $B_{\mathcal{F}(M)}$ containing the molecules $m_{i,j}$ and $m_{j,k}$. Then $m_{i,k}$ is not a vertex of $B_{\mathcal{F}(M)}$.*

Proof. In that case there is a 1-Lipschitz function f such that $f(a_i) - f(a_j) = d_{i,j}$ and $f(a_j) - f(a_k) = d_{j,k}$. Thus

$$d_{i,j} + d_{j,k} = f(a_i) - f(a_k) \leq d_{i,k}$$

and so $m_{i,k}$ is not a vertex. □

Theorem 5.14. *Let (M, d) be a metric space with four points. Then $\mathcal{P}(M) \geq \mathcal{P}(B_1^3)$. Moreover, the equality holds if and only if M is a tree or a cycle with equal weights.*

Proof. Let $G = (V, E, w)$ be the canonical graph associated with M . Note first that if the graph G has a leaf, then $M = M_1 \diamond M_2$, where $\#M_1 = 1$ and $\#M_2 = 3$. Then $\mathcal{P}(M) \geq \mathcal{P}(B_1^3)$ follows from (2) and the two dimensional case of Mahler's conjecture. Moreover, if equality holds then M_2 has minimal volume product and so it is a tree, thus M is a tree too.

Thus, we may assume that G does not any leaf. Assume also that M has minimal volume product among all metric spaces with four elements. Then every vertex of $B_{\mathcal{F}(M)}$ belonging to a facet that contains at least four vertices. Indeed, otherwise Proposition 5.11 says that G has a bridge, and so it has a leaf. Therefore, every edge in G belongs to a cycle, which has length 4 by Lemma 5.13. So either G is a cycle or it is a complete graph. In the first case $B_{\mathcal{F}(M)}$, has 8 vertices that lie in two parallel facets. Then [FMZ] (see also [LR, Ka]) says that $\mathcal{P}(M) \geq \mathcal{P}(B_1^3)$, and if equality holds then either $B_{\mathcal{F}(M)}$ is a double cone or it is affinely isometric to B_∞^3 . In the first case, 6 of the vertices of $B_{\mathcal{F}(M)}$ would be coplanar and it is easy to check that is not possible. On the other hand, one can check that if $\mathcal{F}(M)$ is isometric to ℓ_∞^3 then M is a cycle with equal weights, this also follows from Corollary 3.8.

Thus, it remains to check the case in which G is a complete graph. By relabeling the points of M we may assume that

$$d_{1,0} + d_{2,3} \leq d_{2,0} + d_{1,3} \leq d_{3,0} + d_{1,2}.$$

Let F be a facet of $B_{\mathcal{F}(M)}$ containing $m_{3,0}$ and three other vertices. Thanks to Lemma 5.13, there are only two possible cases:

Case 1. F contains $m_{3,0}$, $m_{3,2}$, $m_{1,2}$ and $m_{1,0}$. Consider the following determinant.

$$\Delta = \begin{vmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & d_{1,2}^{-1} & d_{1,0}^{-1} \\ 0 & -d_{2,3}^{-1} & -d_{1,2}^{-1} & 0 \\ d_{3,0}^{-1} & d_{2,3}^{-1} & 0 & 0 \end{vmatrix} = \frac{d_{3,0} + d_{1,2} - d_{2,3} - d_{1,0}}{d_{1,0}d_{1,2}d_{3,0}d_{2,3}}$$

We have that $\Delta = 0$ since the four molecules $m_{3,0}$, $m_{3,2}$, $m_{1,2}$ and $m_{1,0}$ are coplanar, and so $d_{1,0} + d_{2,3} = d_{3,0} + d_{2,3}$.

Case 2. F contains $m_{3,0}$, $m_{3,1}$, $m_{2,1}$ and $m_{2,0}$. The same argument as before yields $d_{2,0} + d_{1,3} = d_{3,0} + d_{1,3}$.

We conclude that in any case $d_{2,0} + d_{1,3} = d_{3,0} + d_{1,3}$. Then M satisfies the four-point condition and so $B_{\text{Lip}_0(M)}$ is a zonoid [Go]. Therefore $\mathcal{P}(M) \geq \mathcal{P}(B_1^3)$. Moreover, equality does not hold in this case since that would imply that $B_{\mathcal{F}(M)}$ is affinely isomorphic to B_∞^3 [GMR], so it would have 8 vertices. However, $B_{\mathcal{F}(M)}$ has 12 vertices since G is complete. □

Acknowledgements: The authors thank Matthieu Meyer and Gideon Schechtman for their valuable suggestions.

REFERENCES

- [AFZ] M. ALEXANDER, M. FRADELIZI AND A. ZVAVITCH, *Polytopes of Maximal Volume Product*, Discrete Comput. Geom. 62 (2019), no. 3, 583-600.
- [AG] R. J. ALIAGA AND A. J. GUIRAO, *On the preserved extremal structure of Lipschitz-free Banach spaces*, Studia Math. 245(1) (2019) 1-14.
- [AP] R. J. ALIAGA AND E. PERNECKÁ, *Supports and extreme points in Lipschitz-free spaces*, to appear in Rev. Matem. Iberoam.
- [BF] F. BARTHE AND M. FRADELIZI, *The volume product of convex bodies with many hyperplane symmetries*, Amer. J. Math. 135 (2013), 1-37.
- [Ba] R. B. BAPAT, *Graphs and matrices*. Second edition. Universitext. Springer, London; Hindustan Book Agency, New Delhi, 2014.
- [BM] J. BOURGAIN AND V. MILMAN, *New volume ratio properties for convex symmetric bodies in \mathbb{R}^n* , Invent. Math. 88 (1987), 319-340.
- [Bo] E. D. BOLKER, *A class of convex bodies*, Trans. Amer. Math. Soc. 145 (1969), 323-345.
- [CG] S. CAMPI AND P. GRONCHI, *On volume product inequalities for convex sets*, Proc. Amer. Math. Soc. 134 (2006), no. 8, 2393-2402.
- [DDM] A. D'ALÌ, E. DELUCCHI, AND M. MICHAŁEK, *Many faces of symmetric edge polytopes*, arXiv:1910.05193
- [DKO] S. DILWORTH, D. KUTZAROVA, AND M. OSTROVSKII, *Lipschitz-free Spaces on Finite Metric Spaces*, to appear in Canad. J. Math.
- [Fa] J. FARMER, *Extreme points of the unit ball of the space of Lipschitz functions*. Proc. Amer. Math. Soc. 121 (1994), no. 3, 807-813.
- [FHMRZ] M. FRADELIZI, A. HUBARD, M. MEYER, E. ROLDÁN-PENSADO, A. ZVAVITCH, *Equipartitions and Mahler volumes of symmetric convex bodies*, arXiv:1904.10765.
- [FMZ] M. FRADELIZI, M. MEYER, AND A. ZVAVITCH, *An application of shadow systems to Mahler's conjecture*, Discrete Comput. Geom. 48 (2012), no. 3, 721-734.
- [GPR] L. GARCÍA-LIROLA, A. PROCHÁZKA, AND A. RUEDA ZOCA, *A characterisation of the Daugavet property in spaces of Lipschitz functions*, J. Math. Anal. Appl. 464 (2018), 473-492.
- [GPV] A. GIANNOPOULOS, G. PAOURIS, AND B. VRITSIOU, *The isotropic position and the reverse Santaló inequality*, Israel J. Math. 203 (2015), no. 1, 1-22.
- [Go] A. GODARD, *Tree metrics and their Lipschitz-free spaces*, Proc. Amer. Math. Soc. 138 (2010), no. 12, 4311-4320.
- [G] G. GODEFROY, *A survey on Lipschitz-free Banach spaces*, Comment. Math. 55 (2015), no. 2, 89-118.
- [GK] G. GODEFROY AND N. J. KALTON, *Lipschitz-free Banach spaces*. Dedicated to Professor Aleksander Pełczyński on the occasion of his 70th birthday. Studia Math. 159 (2003), no. 1, 121-141.
- [GW] P. R. GOODEY AND W. WEIL, *Zonoids and generalisations*. In Handbook of Convex Geometry (P. M. Gruber, J. M. Wills, eds), vol. B, pp. 1297-1326, North-Holland, Amsterdam, 1993.
- [GP] J. GORDON AND F. PETROV, *Combinatorics of the Lipschitz polytope*, Arnold Math. J. 3 (2017), no. 2, 205218.
- [GM] Y. GORDON AND M. MEYER, *On the Minima of the Functional Mahler Product*, Houston J. Math. 40 (2014), no. 2, 385-393.
- [GMR] Y. GORDON, M. MEYER, AND S. REISNER, *Zonoids with minimal volume product - a new proof*, Proc. Amer. Math. Soc. 104 (1988), 273-276.
- [Gr] B. GRÜNBAUM, *Convex Polytopes*, Graduate Texts in mathematics, 221, Springer, 2003.
- [HL] A. B. HANSEN AND A. LIMA, *The structure of finite-dimensional Banach spaces with the 3.2. intersection property*, Acta Math. 146 (1981), no. 1-2, 123.
- [IKW] Y. IVAKHNO, V. KADETS AND D. WERNER, *Corrigendum to: The Daugavet property for spaces of Lipschitz functions*, Math. Scand. 104 (2009), 319-319.
- [IS] H. IRIYEH AND M. SHIBATA, *Symmetric Mahler's conjecture for the volume product in the three dimensional case*, arXiv:1706.01749.
- [KMO] S. KHAN, M. MIM, AND M. OSTROVSKII, *Isometric copies of ℓ_∞^n and ℓ_1^n in transportation cost spaces on finite metric spaces*, arXiv:1907.01155
- [Ka] R. KARASEV, *Mahler's conjecture for some hyperplane sections*, arXiv:1902.08971.
- [Ki] J. KIM, *Minimal volume product near Hanner polytopes*, J. Funct. Anal. 266 (2014), no. 4, 23602402.
- [Ko] A. KOLDOBSKY, *Fourier analysis in convex geometry*. Mathematical Surveys and Monographs, 116. American Mathematical Society, Providence, RI, 2005.
- [Ku] G. KUPERBERG, *From the Mahler Conjecture to Gauss Linking Integrals*, Geometric And Functional Analysis, 18/ 3, (2008), 870-892.
- [LR] M. A. LOPEZ AND S. REISNER, *A special case of Mahler's conjecture*, Discrete Comput. Geom., 20, (1998), 2 163-177.
- [Ma] K. MAHLER, *Ein Minimalproblem für konvexe Polygone*, Mathematica (Zutphen) B7 (1939), 118-127.
- [Me] M. MEYER, *Une caractérisation volumique de certains espaces normés*, Israel J. Math. 55 (1986), 317-326.
- [MR1] M. MEYER AND S. REISNER, *Inequalities involving integrals of polar-conjugate concave functions*, Monatsh. Math. 125 (1998), 219-227.
- [MR2] M. MEYER AND S. REISNER, *Shadow systems and volumes of polar convex bodies*, Mathematika 53 (2006), no. 1, 129-148 (2007)
- [Na] F. NAZAROV, *The Hörmander proof of the Bourgain-Milman Theorem*, GAFA Seminar Notes, Lecture Notes in Mathematics, Vol. 2050, (2012), 335-343.
- [NPRZ] F. NAZAROV, F. PETROV, D. RYABOGIN, AND A. ZVAVITCH, *A remark on the Mahler conjecture: local minimality of the unit cube*, Duke Math. J. 154 (2010), 419-430.
- [OO] S. OSTROVSKA, M. OSTROVSKII, *On relations between transportation cost spaces and L_1* , arXiv:1910.03625.
- [Os] M. OSTROVSKII, *Metric embeddings. Bilipschitz and coarse embeddings into Banach spaces* De Gruyter Studies in Mathematics, 49. De Gruyter, Berlin, 2013. xii+372 pp.

- [PR] A. PROCHÁZKA AND A. RUEDA ZOCA, *A characterisation of octahedrality in Lipschitz-free spaces*, Ann. Inst. Fourier (Grenoble) 68 (2018), no. 2, 569588.
- [R1] S. REISNER, *Zonoids with minimal volume-product*, Math. Zeitschrift 192 (1986), 339-346.
- [R2] S. REISNER, *Minimal volume product in Banach spaces with a 1-unconditional basis*, J. London Math. Soc. 36 (1987), 126-136.
- [RSW] S. REISNER, C. SCHÜTT, AND E. M. WERNER, *Mahler's conjecture and curvature*, Int. Res. Not. 2012, no. 1, 1-16.
- [RS] C. A. ROGERS AND G. C. SHEPHARD, *Some external problems for convex bodies*, Mathematika 5 (1958), 93-102.
- [RZ] D. RYABOGIN AND A. ZVAVITCH, *Analytic methods in convex geometry*, IM PAN Lecture Notes, Vol. 2, Warsaw 2014.
- [SR] J. SAINT RAYMOND, *Sur le volume des corps convexes symétriques*. Séminaire d'Initiation à l'Analyse, 1980-81, Université Paris VI, 1981.
- [Sc] R. SCHNEIDER, *Convex bodies: the Brunn-Minkowski theory*. Second expanded edition. Encyclopedia of Mathematics and its Applications 151, Cambridge University Press, Cambridge, 2014.
- [St] A. STANCU, *Two volume product inequalities and their applications*, Canad. Math. Bull. 52 (2009), 464-472.
- [Sh] G. C. SHEPHARD, *Combinatorial properties of associated zonotopes*. Canad. J. Math. 26 (1974), 302-321.
- [Ve] A. Vershik, *Classification of finite metric spaces and combinatorics of convex polytopes*, Arnold Math. J. 1 (2015), no. 1, 7581.
- [We] N. WEAVER, *Lipschitz algebras*. World Scientific Publishing Co., Inc., River Edge, NJ, 1999.
- [Wh] H. WHITNEY, *2-Isomorphic Graphs*. Amer. J. Math. 55 (1933), no. 1-4, 245254.

DEPARTMENT OF MATHEMATICAL SCIENCES, KENT STATE UNIVERSITY, KENT, OH 44242, USA

E-mail address: `malexan5@kent.edu`

E-mail address: `lgarcial@kent.edu`

E-mail address: `zvavitch@math.kent.edu`

UNIVERSITÉ PARIS-EST MARNE-LA-VALLÉE, LABORATOIRE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES (UMR 8050)
CITÉ DESCARTES - 5, BD DESCARTES, CHAMPS-SUR-MARNE 77454 MARNE-LA-VALLÉE CEDEX 2, FRANCE

E-mail address: `matthieu.fradelizi@u-pem.fr`