



Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex

Christine H. Meyer-Friessem, Nadine Attal, Ralf Baron, Didier Bouhassira, Nanna Brix Finnerup, Rainer Freynhagen, Janne Gierthmühlen, Maija L. Haanpää, Per T. Hansson, Troels Jensen, et al.

► To cite this version:

Christine H. Meyer-Friessem, Nadine Attal, Ralf Baron, Didier Bouhassira, Nanna Brix Finnerup, et al.. Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex. *European Journal of Pain*, 2020, 24 (6), pp.1058-1071. <10.1002/ejp.1550>. <hal-02927422>

HAL Id: hal-02927422

<https://hal.science/hal-02927422v1>

Submitted on 7 Jun 2023


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex

Christine H. Meyer-Frießem^{1,2} | Nadine Attal^{3,4} | Ralf Baron⁵ | Didier Bouhassira^{3,4} |
Nanna B. Finnerup^{6,7} | Rainer Freynhagen^{8,9} | Janne Gierthmühlen⁵ | Maija Haanpää^{10,11} |
Per Hansson^{12,13} | Troels S. Jensen^{6,7} | Harriet Kemp¹⁴ | Donna Kennedy¹⁴ |
Anne-Sofie Leffler¹³ | Andrew S. C. Rice¹⁴ | Märta Segerdahl^{15,16} | Jordi Serra¹⁷ |
Soeren Sindrup¹⁸ | Roma Solà¹⁷ | Thomas Tölle¹⁹ | Sigrid Schuh-Hofer²⁰ |
Rolf-Detlef Treede²⁰ | Esther Pogatzki-Zahn²¹ | Christoph Maier² | Jan Vollert^{14,20} 

¹Department of Anesthesiology, Intensive Care, Palliative and Pain Medicine, University Hospital Bergmannsheil Bochum, Bochum, Germany

²Department of Pain Medicine, BG University Hospital Bergmannsheil GmbH, Ruhr-University Bochum, Bochum, Germany

³INSERM U-987, Centre d'Evaluation et de Traitement de la Douleur, CHU Ambroise Paré, Boulogne-Billancourt, France

⁴Université Versailles-Saint-Quentin, Versailles, France

⁵Division of Neurological Pain Research and Therapy, Department of Neurology, Universitätsklinikum Schleswig-Holstein, Campus Kiel, Kiel, Germany

⁶Department of Neurology, Aarhus University Hospital, Aarhus, Denmark

⁷Danish Pain Research Center, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

⁸Department of Anaesthesiology, Critical Care Medicine, Pain Therapy & Palliative Care, Pain Center Lake Starnberg, Benedictus Hospital Tutzing, Tutzing, Germany

⁹Anaesthesiological Clinic, Klinikum Rechts der Isar, Technische Universität München, Munich, Germany

¹⁰Department of Helsinki University Central Hospital, Helsinki, Finland

¹¹Etera Mutual Pension Insurance Company Helsinki, Helsinki, Finland

¹²Division of Emergencies and Critical Care, Department of Pain Management and Research, Oslo University Hospital, Oslo, Norway

¹³Department of Molecular Medicine and Surgery, Karolinska Institute, Stockholm, Sweden

¹⁴Pain Research, Department of Surgery and Cancer, Imperial College, London, UK

¹⁵H. Lundbeck A/S, Copenhagen, Denmark

¹⁶Department of Physiology and Pharmacology, Karolinska Institute, Stockholm, Sweden

¹⁷Neuroscience Technologies, Ltd., Barcelona, Spain

¹⁸Department of Neurology, Odense University Hospital, Odense, Denmark

¹⁹Department of Neurology, Klinikum Rechts der Isar, Technische Universität München, Munich, Germany

²⁰Center of Biomedicine and Medical Technology Mannheim CBTM, Medical Faculty Mannheim, Heidelberg University, Heidelberg, Germany

²¹Department of Anaesthesiology, Intensive Care and Pain Medicine, University Hospital Muenster, Muenster, Germany

Correspondence

Jan Vollert, Pain Research, Department of Surgery and Cancer, Imperial College London, Chelsea and Westminster Campus, 369 Fulham Rd, London SW10 9NH, UK.
Email: j.vollert@imperial.ac.uk

Abstract

Background and Aims: Healthy women have generally been found to have increased experimental pain perception and chronic pain has a higher prevalence in female as compared to male patients. However, no study has investigated whether pain intensity and pain perception thresholds are distinct or similar between sexes within various chronic pain entities. We investigated whether average pain intensities and

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *European Journal of Pain* published by John Wiley & Sons Ltd on behalf of European Pain Federation - EFIC ®.

Funding information

The IMI EUROPAIN project has received support from the Innovative Medicines Initiative Joint Undertaking under grant agreement No. 115007. The NEUROPAIN project was supported by an independent investigator-initiated research grant from Pfizer Ltd. German Research Network on Neuropathic pain (DFNS) is supported by the German Ministry of Education and Research (BMBF; Grant No. 01EM0902).

pain thresholds assessed using quantitative sensory testing (QST) differed between women and men suffering from three distinct chronic pain conditions: Complex Regional Pain Syndrome (CRPS type I), peripheral nerve injury (PNI) or polyneuropathy (PNP), as compared to paired healthy volunteers.

Methods: QST data of 1,252 patients (669 female, 583 male) with PNI ($n = 342$), PNP ($n = 571$) or CRPS ($n = 339$), and average pain intensity reports from previously published studies were included. Absolute and z -values (adjusted for age and body region) of cold, heat, pressure (PPT) and pinprick pain thresholds were compared in generalized linear models with aetiology, duration of underlying pain disease and average pain intensity as fixed effects.

Results: Average pain intensity during the past four weeks did not differ between women and men, in both mean and range. In women absolute pain thresholds for cold, heat and pinprick were lower than in males across all diagnoses ($p < .05$). However, after z -transformation these differences disappeared except for PPT in CRPS ($p = .001$).

Discussion: Pain thresholds in patients show only minor sex differences. However, these differences mimic those observed in healthy subjects and do not seem to be linked to specific pathophysiological processes.

Significance: Female healthy participants and female patients with neuropathic pain conditions or CRPS I report lower pain thresholds compared to males, but pain intensity is similar and there is no sex difference in the extent to which the thresholds are altered in neuropathic pain or CRPS. Thus, the sex differences observed in various chronic pain conditions mimic those obtained in healthy participants, indicating that these differences are not linked to specific pathophysiological processes and are of minor clinical relevance.

1 | INTRODUCTION

Since decades, gender differences have represented a major topic in pain research, although studies have found conflicting results (Bouhassira, Lantéri-Minet, Attal, Laurent, & Touboul, 2008; Breivik, Collett, Ventafridda, Cohen, & Gallacher, 2006; Fillingim, King, Ribeiro-Dasilva, Rahim-Williams, & Riley, 2009; Friessem, Willweber-Strumpf, & Zenz, 2009; Lamerato et al., 2016; Moulin, Clark, Speechley, & Morley-Forster, 2002; Reitsma, Tranmer, Buchanan, & VanDenKerkhof, 2012; Torrance, Smith, Bennett, & Lee, 2006). The overall consensus is that healthy women appear to be more pain sensitive and demonstrate lower pain thresholds than men (Bartley & Fillingim, 2013; Fillingim et al., 2009; Mogil, 2012). In a recent review of more than 120 studies in healthy volunteers subjected to various experimental pain models, differences in pain thresholds were only minor to moderate (Mogil, 2012), while the majority of studies found female participants to have significantly lower pain tolerance to supra-threshold stimuli (Mogil, 2012; Racine et al., 2012). Within the German Research Network on Neuropathic Pain (DFNS) reference database of healthy controls, women were found to be significantly more pain sensitive using Quantitative Sensory Testing (QST; Rolke et al., 2006), although the differences were small and could only be seen in group comparison. Consequently, it has been hypothesized that

healthy women are more pain sensitive than men. Furthermore, higher prevalence in some chronic pain conditions has been reported in women as compared to men for, e.g., migraine, fibromyalgia, temporomandibular disorders (Bartley & Fillingim, 2013; Breivik et al., 2006; Davis, Robinson, Le, & Xie, 2011; Fillingim et al., 2009; Friessem et al., 2009; Lamerato et al., 2016; Moulin et al., 2002; Reitsma et al., 2012), to a lesser extent complex regional pain syndrome (CRPS; Demir, Ozaras, Karamehmetoğlu, Karacan, & Aytekin, 2010; de Mos et al., 2007; Ott & Maihöfner, 2018; Pons, Shipton, Williman, & Mulder, 2015; Roh et al., 2014) and some neuropathic pain conditions (Attal, Lantéri-Minet, Laurent, Fermanian, & Bouhassira, 2011; Bouhassira et al., 2008; Torrance et al., 2006). It may therefore be hypothesized that women suffering from various chronic pain conditions have distinct QST-phenotypes in the form of allodynia and hyperalgesia as compared to men. However, while pain tolerance has been extensively studied in healthy volunteers and to some extent in chronic low back pain patients (Meints, Wang, & Edwards, 2018) and neuropathic pain patients (Arap, Siqueira, Silva, Teixeira, & Siqueira, 2010; Goswami, Anastakis, Katz, & Davis, 2016; Krämer, Rolke, Bickel, & Birklein, 2004; Schäfer et al., 2014; Selim et al., 2010), it has not been attempted to demonstrate if such gender or sex differences are based on variation in pathology or merely mirroring biological differences that can be found in healthy participants.

In this study, we investigated whether pain thresholds based on QST and self-reported average pain intensity differed between women and men suffering from three distinct chronic pain conditions: CRPS type I (CRPS I), peripheral nerve injury (PNI) and painful polyneuropathy (PNP) as compared to paired healthy controls. As secondary objectives, we investigated if these potential differences reproduced the results found in healthy participants and whether they were linked to the aetiology of the underlying pathology.

2 | MATERIALS AND METHODS

2.1 | Consortia

Three consortia, which prospectively collected patients using identical study protocols, were involved in this data analysis: The DFNS (<http://www.neuropathischer-schmerz.de>) founded in 2002 aimed at promoting research on mechanisms and treatments for neuropathic pain, as well as establish a large database for QST-data.

The IMI EUROPAIN project (<http://www.imieuropain.org>), founded in 2009, aims to improve the treatment of patients with long-term pain. It consists of academic study groups working on pain research from Germany, Denmark and the UK. A Spanish SME (small- and medium-sized enterprises) and researchers from EFPIA (European Federation of Pharmaceutical Industries and Associations) partners active in the pain research are additional contributors. The NEUROPAIN project is an investigator-initiated study consisting of several researchers in the field of Neuropathic Pain research within Europe aiming to characterize subgroups of patients with neuropathic pain. IMI EUROPAIN and NEUROPAIN have collected data at the same time and to the same study protocols, including data to the central database from day 1 and are thus homogeneous data. To additionally ensure comparability of data collection, an analysis of heterogeneity was conducted, showing that the data of all consortia can be analysed as a homogeneous database independently of consortia (Vollert, Attal, et al., 2016).

2.2 | Central database

Patients and healthy participants were assessed between 2002 and 2013. During this time, no changes to the standardized QST-protocol were implemented. Furthermore, all investigators had undergone certification training in the QST-protocol before examining healthy volunteers and patients, and there was a surveillance of maintained data quality during the data collection period (Vollert et al., 2015; Vollert, Attal, et al., 2016). The ethics committee of each participating centre approved the assessment protocol and data collection. Each study centre used the

computer-assisted program Neuroquast[®] (Statconsult) for local data entry. Study records were imputed monthly into the fully-integrated central database. All centres underwent strict quality control (Magerl et al., 2010; Vollert et al., 2015) and an analysis of heterogeneity showed that the database can be analysed as a homogeneous dataset (Vollert, Attal, et al., 2016). All patients from the database suffering from CRPS type I, polyneuropathy or peripheral nerve injury were included in this analysis. All data in this study have been published before, on healthy participants (Rolke et al., 2006) and CRPS (Maier et al., 2010) by the DFNS, polyneuropathy and peripheral nerve injury by all three consortia (Baron et al., 2017; Maier et al., 2010; Vollert, Kramer, et al., 2016; Vollert et al., 2017).

2.3 | Patients, healthy participants and sex

Sex of the participants was determined using self-report. As it was not part of the initial study planning, no information of cis or trans gender was recorded, to the best of our knowledge and certainly in the vast majority of the included participants, self-reported gender and biological sex will be identical, as also confirmed during the neurological examination.

Healthy participants were included on condition of an inconspicuous medical history (Rolke et al., 2006). Volunteers suffering from any pain condition were excluded, intake of any medication within the last 24 hr prior QST lead to exclusion as well (Rolke et al., 2006). A full list of in- and exclusion rules has been defined and agreed upon by the consortia (Gierthmühlen et al., 2015).

Adult patients with neuropathic pain due to PNI (i.e., history of traumatic nerve injury and presence of sensory abnormalities in the innervation and/or abnormal electro-neurography), CRPS I (i.e., according to clinical criteria [Harden et al., 2010a]) and PNP (i.e., abnormal electro-neurography or abnormally decreased vibration detection threshold at two of four sites less 5/8 at the lower limb, which could not be explained by another disease [England et al., 2005]) were included in the central database. Further details for inclusion of these patient groups can be found, e.g., in (Vollert, Attal, et al., 2016). All patients gave written informed consent for transfer of their data into the common central database. Exclusion criteria were missing informed consent, insufficient language skills, pain treatment by topical local anaesthetics in the last seven days or by topical capsaicin in the last three months, since this might affect pain thresholds (Baron et al., 2017). Current systemic treatment with pain medication did, in contrast to topical treatment at the site of examination, not lead to exclusion; however, patients with additional secondary painful conditions or neurological or psychiatric conditions treated with opioids, anticonvulsants or antidepressants were excluded. Details on the patients included are presented in Table 1.

TABLE 1 Clinical data of healthy subjects and patients

Characteristic	Male healthy participants ^a (<i>n</i> = 70, 38.9%)	Female healthy participants ^a (<i>n</i> = 110, 61.1%)	All (<i>n</i> = 180)	<i>p</i> -value
Consortia, <i>n</i> / % DFNS	70/100	110/100	180/100	
Age (years), mean ± <i>SD</i>	37.5 ± 13.0	38.9 ± 13.0	38.4 ± 13.0	.461
	Male patients (<i>n</i> = 583, 46.6%)	Female patients (<i>n</i> = 669, 53.4%)	All (<i>n</i> = 1,252)	<i>p</i> -value
Consortia, <i>n</i> / % DFNS	361/61.9	475/71.0	836/66.8	
Age (years), mean ± <i>SD</i>	54.8 ± 13.1	55.7 ± 14.7	55.2 ± 13.9	.256
CRPS, <i>n</i> / %	77/22.7	262/77.3	339	
Age (years), mean ± <i>SD</i>	51.8 ± 12.2	52.1 ± 14.0	52.1 ± 13.6	.864
Region of pain				
Hand, <i>n</i>	66	242	308	.075
Foot, <i>n</i>	11	20	31	
Duration of pain disease ^b				
≤ 1 year, <i>n</i>	41	138	179	.894
≥ 1 year, <i>n</i>	35	122	157	
Average pain intensity (NRS)				
Mean ± <i>SD</i> (<i>d</i>)	4.0 ± 3.3	4.6 ± 3.2	4.5 ± 3.2 (0.19)	.152
NRS ≤ 3	25	73	98	.433
NRS ≥ 7	12	49	61	.531
Peripheral nerve injury, <i>n</i> / %	183/53.5	159/46.5	342	
Age (years), mean ± <i>SD</i>	49.2 ± 12.2	50.8 ± 14.1	50.0 ± 13.1	.280
Region of pain				
Hand or arm, <i>n</i>	86	72	158	.428
Leg or foot, <i>n</i>	65	49	114	
Dorsal or ventral trunk, <i>n</i>	24	31	55	
Other (e.g. face), <i>n</i>	8	7	14	
Duration of pain disease				
≤ 1 year, <i>n</i>	39	37	76	.489
≥ 1 year, <i>n</i>	92	72	164	
unknown, <i>n</i>	52	50	102	
Average pain intensity (NRS)				
Mean ± <i>SD</i> (<i>d</i>)	5.7 ± 2.4	5.3 ± 2.9	5.5 ± 2.6 (0.15)	.164
NRS ≤ 3	10	23	33	.005
NRS ≥ 7	34	32	66	.718
Polyneuropathy, <i>n</i> / %	323/56.6	248/43.3	571	
Age (years), mean ± <i>SD</i>	58.7 ± 12.5	62.7 ± 13.1	60.5 ± 12.9	<.001
Region of pain ^c				
Hand, <i>n</i>	5	4	9	.954
Leg or foot, <i>n</i>	317	244	561	
Duration of pain disease				
≤ 1 year, <i>n</i>	36	30	66	.578
≥ 1 year, <i>n</i>	216	155	371	
unknown, <i>n</i>	71	63	134	

(Continues)

TABLE 1 (Continued)

	Male patients (<i>n</i> = 583, 46.6%)	Female patients (<i>n</i> = 669, 53.4%)	All (<i>n</i> = 1,252)	<i>p</i> -value
Aetiology				
Diabetic	43	44	87	.013
Fabry	17	8	25	
Toxic	10	4	14	
Chemo-induced	14	15	29	
HIV-related	12	0	12	
Idiopathic or unknown	227	177	404	
Average pain intensity (NRS)				
Mean \pm <i>SD</i> (<i>d</i>)	5.4 \pm 2.7	5.8 \pm 2.8	5.6 \pm 2.7 (0.15)	.085
NRS \leq 3	38	28	64	.861
NRS \geq 7	62	70	132	.011

Note: Definition of average pain intensity: persistent or paroxysmal spontaneous pain, pain attacks, and pain evoked by daily-life stimuli during the past four weeks on average.

Abbreviations: *d*, Cohen's *d* (effect size); NRS, numerical rating scale (0–10; 0: no pain; 10: worst pain imaginable); *SD*, standard deviation.

^aRolke et al. 2006.

^bunknown *n* = 3.

^cothers *n* = 2.

2.4 | Assessment of average pain intensity

Before QST, all patients filled out the painDETECT questionnaire (PD-Q; Freynhagen, Baron, Gockel, & Tölle, 2006), which collects information about average pain intensity. Based on this questionnaire, the question "How *severe* was your pain during the past four weeks on average?" with pain intensity rated on a numerical rating scale (NRS; 0: no pain, 10: worst imaginable pain) was taken as "average pain" for this analysis. Here, average pain referred to any type of pain including continuous or paroxysmal pain and pain evoked by daily-life stimuli during the past four weeks.

2.5 | Quantitative sensory testing

According to the DFNS protocol, QST consists of seven tests measuring 13 parameters which assess the function of small and large afferent nerve fibres or corresponding CNS pathways: cold detection threshold (CDT), warm detection threshold (WDT), thermal sensory limen (TSL), paradoxical heat sensation (PHS), cold pain threshold (CPT), heat pain threshold (HPT), mechanical (tactile) detection threshold (MDT), mechanical pain threshold (MPT), mechanical pain sensitivity (MPS), dynamic mechanical allodynia (DMA), wind-up ratio (WUR), vibration detection threshold (VDT) and pressure pain threshold (PPT). Mechanical pain sensitivity was rated on NRS (0–100). Rolke et al. provided a detailed description of parameters, protocol, evaluation and database of reference values (Rolke et al., 2006). Testing location was the most painful area in CRPS and PNI and the dorsolateral aspect of both feet in patients with

polyneuropathy. In the original cohort of the DFNS healthy participants all were assessed on the feet (Rolke et al., 2006).

2.6 | A note on semantics: sensitivity, thresholds and hyperalgesia

In testing perception thresholds to sensory stimuli, a gain of function, i.e., an increased sensitivity, usually corresponds to a decreased threshold: meaning a less intense stimulus than normally gives rise to the requested percept. A curious exception to this rule is the cold pain threshold, as indeed a perception threshold at a lower temperature than normal would indicate a person being less sensitive to painful cold stimuli. On the other hand, a painful percept below the normal threshold would be higher or further away from starting temperature. To increase readability and accessibility for anyone who is not an expert in the field, for the purpose of this paper, whenever the terms "decreased threshold" or "lower threshold" are used, including cold pain thresholds, they should be read as "lower stimulus intensity needed to give rise to the requested percept" and therefore depict an increased sensitivity. In this paper, the term "allodynia" is only used for dynamical mechanical allodynia (DMA), while decreased pain thresholds are generally described as hyperalgesia. While we acknowledge that a strongly decreased pain threshold would fall in the IASP definition under allodynia ("pain due to a stimulus that does not normally provoke pain"), a less pronounced change in threshold might also be labelled hyperalgesia, if it is still within the realm of stimuli

that are normally considered painful. Increased sensitivity through decreased thresholds (the *minimum* intensity of a stimulus that is perceived as painful [IASP, 2017]), is not

to be confused with increased sensitivity due to decreased pain tolerance (the *maximum* intensity of a pain-producing stimulus that a subject is willing to accept [IASP, 2017]).




FIGURE 1 Sex-specific average pain intensity in patients with CRPS I, peripheral nerve injury and polyneuropathy. (a) Pie charts of gender distribution of patients with CRPS I, PNI and PNP in percent. (b) Average pain intensity on numerical rating scale (NRS, 0–10) plotted with mean values and standard deviation in male and female patients with CRPS, PNI and PNP. (c) Percent of average pain intensity on numerical rating scale (NRS, 0–10) classified in mild (NRS ≤ 3), moderate (NRS 4–6) and severe (NRS ≥ 7) of male and female patients with CRPS I, PNI and PNP. CRPS, complex regional pain syndrome; PNI, peripheral nerve injury; PNP, polyneuropathy. Definition of average pain intensity: persistent or paroxysmal spontaneous pain, pain attacks, and pain evoked by daily-life stimuli during the past four weeks on average. None of the differences are significant ($p < .05$ corrected for multiple testing using the Benjamini-Hochberg procedure)




FIGURE 2 Sex-specific QST-evoked pain in healthy participants due to cold, heat, pinprick and pressure stimuli. (a) cold pain threshold (CPT), (b) heat pain threshold (HPT), (c) mechanical pain threshold (MPT) and (d) pressure pain threshold (PPT). Data from (Rolke et al., 2006). All differences are significant on a $p < .05$ level, corrected for multiple testing using the Benjamini-Hochberg procedure

2.7 | Statistical analysis

Sex differences in thermal and mechanical pain thresholds were tested twofold (a) as absolute pain thresholds and (b) as z -values. According to Rolke et al. (2006), absolute thermal values (e.g., CPT, HPT) are distributed normally, while assessments of mechanical thresholds, e.g., PPT and MPT are distributed log-normally and were analysed accordingly. To be able to compare QST-parameters independently of their physical dimension and to focus on disease-specific differences, a z -transformation was applied to the DFNS normative material (Magerl et al., 2010; Pfau et al., 2014; Rolke et al., 2006). In this normalization procedure, all values are transformed to a sex, age and body-region adjusted mean = 0 and standard deviation = 1 in healthy participants. To test for sex differences, modified z -scores only adjusted for age and body-region were calculated additionally. Thus, z -scores indicate if a patient has loss of function signs (i.e., hypaesthesia, hypoalgesia, z -values below zero) or gain of function (i.e., hyperaesthesia, hyperalgesia, z -values above zero) as described in (Rolke et al., 2006).

Absolute pain thresholds and z -values of pain and detection thresholds were analysed in general linear models, with

sex and aetiology as fixed effect. To control for confounding effects, duration of the underlying disease (under one year, one to five years, over five years), average pain intensity (NRS: 0–10: <3, 3 to <7, 7–10), and in addition for absolute pain thresholds only, body region (upper limb, lower limb, head, trunk) and age decade were included as fixed effects as well. All p -values presented in this manuscript, except for the demographic comparisons by chi-squared test in Table 1, result from these corrected models, and $p < .05$ were considered statistically significant after correction for multiple testing using the Benjamini-Hochberg procedure to adjust false discovery rate.

3 | RESULTS

3.1 | Healthy participants and study population

Demographic data of all patients and healthy participants are shown in Table 1. The dataset comprised QST-data of 1,252 patients (583 males, 669 [53.4%] females) with PNP ($n = 571$, 43.4% female), PNI ($n = 342$, 46.5% female) or CRPS I ($n = 339$, 77.3% female).




FIGURE 3 Sex-specific pain thresholds in patients with CRPS I, peripheral nerve injury and polyneuropathy (*z*-values adjusted for age and body region, not for sex differences). (a) cold pain threshold (CPT), (b) heat pain threshold (HPT), (c) mechanical pain threshold (MPT) and (d) pressure pain threshold (PPT). Presented are mean *z*-values (gender unspecific—corrected just for age decade and body region) and standard error of mean. CRPS, complex regional pain syndrome; PNI, peripheral nerve injury; PNP, polyneuropathy

3.2 | Average pain intensity

In general, duration of painful condition and intensity of average pain within the last four weeks were similar between female and male patients in all aetiologies (see Table 1 and Figure 1). Compared to females, less male patients with CRPS reported a low pain intensity ($p = .005$). Conversely women with PNP more often reported a severe pain intensity as compared to men ($p = .011$).

3.3 | Evoked pain assessed by QST

Healthy female participants showed lower pain thresholds for cold, heat, pinprick and pressure (all $p < .01$) in comparison to males, whereas MPS and DMA did not differ (Figure 2).

In patients, nearly all absolute pain thresholds were lower (i.e., closer to baseline) in female as compared to male patients for the included painful conditions: CPT ($p = .042$), HPT ($p = .000$), and MPT ($p = .018$) (Figure 3; Table 2). PPT was lower for female patients only in CRPS ($p = .001$). However, *z*-values of pain thresholds did not differ significantly between

male and female patients, except for aetiology-specific sex effects for PPT in CRPS patients ($p < .001$). Additional sex effects were found for WUR and MDT. QST-profiles, separately for each aetiology and sex, are displayed in Figure 4, *p*-values resulting from the general linear models can be found in Table 2.

4 | DISCUSSION AND CONCLUSIONS

We investigated whether pain intensity and thresholds are distinct or similar between women and men. We found subtly, yet significantly lower pain thresholds for cold, heat, pressure and pinprick for healthy females. Given that in the same cohort, detection thresholds did not differ (shown in [Rolke et al., 2006]), it is unlikely that this difference comes from variations in innervation density, but rather points towards differential central processing (Riley, Robinson, Wise, Myers, & Fillingim, 1998). For patients suffering from CRPS I or painful polyneuropathy or nerve injury, average pain intensity was similar in women and men and there were only minor sex differences in pain thresholds.

TABLE 2 Multivariate general linear models for absolute pain thresholds and z -transformed pain and detection thresholds

	Model	Sex	Aetiology	Sex*aetiology
Untransformed, corrected for age decade and test area				
CPT	<0.001	0.042	0.492	0.509
HPT	<0.001	<0.001	0.115	0.171
MPT	<0.001	0.018	0.412	0.094
PPT	<0.001	0.926	0.011	0.001
z -values				
CDT	<0.001	0.063	0.219	0.703
WDT	0.001	0.467	0.391	0.234
TSL	<0.001	0.614	0.013	0.090
CPT	<0.001	0.189	<0.001	0.820
HPT	<0.001	0.058	<0.001	0.204
PPT	<0.001	0.001	<0.001	<0.001
MPT	0.004	0.801	0.002	0.202
MPS	<0.001	0.493	<0.001	0.413
WUR	0.220	0.021	0.933	0.184
MDT	<0.001	0.001	0.461	0.189
VDT	0.130	0.131	0.447	0.484

Note: p -values generated in multivariate general linear models for absolute pain and z -transformed pain and detection thresholds in regard to sex and aetiology, corrected for duration of disease, pain intensity, and for absolute pain thresholds additionally body region and age decade. Values in bold red indicate significance at $p < .05$ after correction for multiple testing using the Benjamini-Hochberg procedure.

Abbreviations: CDT, cold detection threshold; CPT, cold pain threshold; HPT, heat pain threshold; MDT, mechanical detection threshold; MPS, mechanical pain sensitivity; MPT, mechanical pain thresholds; PPT, pressure pain threshold; TSL, thermal sensory limen; VDT, vibration detection threshold; WDT, warmth detection threshold; WUR, wind-up ratio.

Female patients reported lower pain thresholds compared to male patients. These sex differences observed in these chronic pain conditions mimic those obtained in healthy participants, indicating that these differences are not linked to specific pathophysiological processes. Interestingly, the variations observed in experimentally evoked pain thresholds within each sex group were much higher than those observed between sexes.

The only exception was CRPS I, where a more pronounced pressure evoked hyperalgesia could be demonstrated in women as compared to men. CRPS I was also significantly more frequent in women in our cohort, corroborated by existing literature (Ott & Maihöfner, 2018), but the reason for this is unknown. It is possible that women are more likely to require CRPS, or that they are more at risk to be suffer from fractures (e.g., for higher rates of osteoporosis) leading to higher incidence of CRPS. Deep pressure pain has been found to have a lower threshold in healthy women as compared to men in prior studies (Chesterton, Barlas, Foster, Baxter, & Wright, 2003; Pfau et al., 2014; Riley et al., 1998). In contrast

to pinprick hyperalgesia, blunt hyperalgesia is considered at least partly due to peripheral sensitization in different pain states (Enax-Krumova et al., 2017; Gierthmühlen et al., 2012; Kilo, Schmelz, Koltzenburg, & Handwerker, 1994; Maier et al., 2010; Mainka et al., 2014; Pfau, Rolke, Nickel, Treede, & Daublaender, 2009). It is a hallmark sign of CRPS (Gierthmühlen et al., 2012; Harden et al., 2010b; Mainka et al., 2014). Blunt pressure hyperalgesia is more pronounced in CRPS than in peripheral nerve injury, but sex differences have not been reported previously. In our multivariate analyses, the sex difference regarding PPT in CRPS was not related to the duration of pain or average pain intensity, so it cannot be explained by a more severe CRPS.

There is a long lasting debate on whether or not women are more sensitive to pain (Hashmi & Davis, 2014; Mogil, 2012; Racine et al., 2012), with inconclusive results throughout many qualitative and quantitative reviews (Bartley & Fillingim, 2013; Fillingim, 2000; Fillingim et al., 2009; Greenspan et al., 2007; Mogil, 2012; Racine et al., 2012; Riley et al., 1998). Biological factors (hormones, reproductive stage or genetics), psychosocial factors (education and gender roles or sociocultural) and ethnical drivers might influence pain (Bartley & Fillingim, 2013; Rahim-Williams, Riley, Williams, & Fillingim, 2012). Varying result interpretations, missing differentiation between pain thresholds, tolerance and pain intensity, lack standardization of outcome measures (Williamson et al., 2017), different terminology (Greenspan et al., 2007), and the usually ignored influence of the hormone status in females (Pogatzki-Zahn et al., 2019) as well as the investigator provider besides the patient itself (Kállai, Barke, & Voss, 2004; Meyer-Frießem, Szalaty, Zahn, & Pogatzki-Zahn, 2019) cause an additional variety of results (Bartley & Fillingim, 2013; Racine et al., 2012). Hence, subtle sex differences might be overestimated.

This study is the first major study focussing on sex differences for pain thresholds (the *minimum* intensity of a stimulus that is perceived as painful [IASP, 2017]), rather than pain tolerance (Mogil, 2012; the *maximum* intensity of a pain-producing stimulus that a subject is willing to accept [IASP, 2017]). Both relate to the subjective experience of the individual, as they rely on what the individual defines as “painful.” In a clinical setting, pain tolerance is more meaningful (as it relates more to the patient's problem); however, pain thresholds have been found to be more experimentally robust (Gelfand, 1964). It has also been reported that gender role expectations affect pain tolerance, but not pain thresholds (Defrin, Shramm, & Eli, 2009), and that pain tolerance could also be more influenced by cultural components than pain thresholds (Dawson & List, 2009). Subsequently there are more reports of sex effects on pain tolerance (Hashmi & Davis, 2014). Indeed in healthy participants, sex differences have mainly concerned supra-threshold pain responses (Mogil, 2012), although studies




a Complex regional pain syndrome**b Polyneuropathy****c Peripheral nerve injury**

FIGURE 4 Sex-specific somatosensory profiles of patients with (a) CRPS I, (b) peripheral nerve injury and (c) polyneuropathy (z-values adjusted for age and body region, and for sex differences found in healthy participants). Mean of z-values and standard error of mean (SE). If the resulting z-value exceeds 1.96/−1.96, it is outside the 95% confidence interval of the standard normal distribution. CDT, Cold detection threshold; CPT, cold pain threshold; CRPS, complex regional pain syndrome; DMA, dynamic mechanical allodynia; HPT, heat pain threshold; MDT, mechanical (tactile) detection threshold; MPS, mechanical pain sensitivity; MPT, mechanical pain threshold; PHS, paradoxical heat sensation; PNI, peripheral nerve injury; PNP, polyneuropathy; PPT, pressure pain threshold; TSL, thermal sensory limen; VDT, vibration detection threshold; WDT, warm detection threshold; WUR, wind-up ratio. * $p < .05$, ** $p < .01$, corrected for multiple testing using the Benjamini-Hochberg procedure

As this study is based on a database query, there are limitations to the level of data we could extract. We did not collect information about the menstrual cycle of the female patients (Craft, Mogil, & Aloisi, 2004; Iacovides, Avidon, & Baker, 2015; Pogatzki-Zahn et al., 2019). Neither was the sex of experimenters conducting the QST-protocol in our study systematically recorded and cannot be reconstructed at this point. It has been reported that experimental pain reports may depend on the sex of the experimenter (Aslaksen, Myrbakk, Høifødt, & Flaten, 2007; Gijsbers & Nicholson, 2005; Levine & De Simone, 1991; Meyer-Frießem et al., 2019; Vigil & Alcock, 2014). As the original focus of this database collection did not lie on sex differences, detailed evaluation of gender identity including gender roles were not performed in addition to collecting data on the sex of the subject.

In addition, we did not take into consideration other potential factors involved in pain, e.g., psychosocial issues, genetics, endogenous hormone levels and social factors. Since our hypothesis was about trait differences between males and females, we preferred to use 4-week average pain scores as retrospective pain ratings instead of current pain intensities for analysis because of day to day variability of pain scores.

It is largely recognized that patients suffering from neuropathic pain or CRPS I vary broadly in sensory phenotypes and underlying mechanisms (Baron et al., 2017; Maier et al., 2010; Üçeyler et al., 2018; Vollert et al., 2017, 2018). It cannot be excluded that distinct sensory phenotypes may be associated with sex differences. While non-painful detection thresholds were not part of the question of this analysis, more pronounced fibre loss could impede pain thresholds, thus influencing pain thresholds. Still, as the QST-profiles show (Figure 4) sensory thresholds do not differ largely between female and male patients and cannot explain the significant difference in pressure pain threshold for CRPS I patients.

Given that this analysis was relying on database data, we could not retrieve information about medication for the study

did not all clearly differentiate between threshold testing and response to suprathreshold pain stimuli (Racine et al., 2012). Based on a review including more than 120 studies in healthy individuals subjected to different human pain models, pain thresholds were generally similar or only moderately different among males and females (Racine et al., 2012). In contrast, 80% of the studies reported that healthy females have lower pain tolerance (in response to cold, heat, pressure, muscle or electrical stimulations; Bartley & Fillingim, 2013; Bartley et al., 2016; Fillingim et al., 2009; Mogil, 2012; Racine et al., 2012).

Several limitations of this study should be acknowledged.

population—neither to assess if and how treatment and gender interact with sensory profiles and pain ratings, nor, more importantly, to look for sex-treatment interactions that could hint towards not only variances in underlying pathology, but also hold the potential for a more nuanced, sex-specified treatment. In the small subset of data of the patient population (not covering CRPS patients) we have medication data on; however, we do not observe any clear pattern of differential prescription (data not shown). In addition, a recent review summarized: “There is a lack of robust evidence to support a gender-specific analgesic management” (Packiasabapathy & Sadhasivam, 2018).

Although our sample is fairly large, it is not epidemiologically representative since it depended on referral to the participating centres, who are highly specialized pain clinics, and will not see patients who have no complaints or are easily adjusted on first-line treatment. Furthermore, we collected information on two neuropathic pain conditions and CRPS I which may not reflect other potential pain conditions including nociceptive pain.

Commonly, female healthy participants and female patients with CRPS I, PNI and PNP demonstrate lower pain thresholds to cold, heat, pressure and pinprick than male patients and healthy participants. However, there is no sex difference in the extent to which these thresholds are altered in CRPS I or peripheral neuropathic pain states, with the only exception of CRPS I in female patients where PPT is significantly lowered. Therefore, from our data, we cannot suggest variance in mechanisms of pain pathophysiology, and gender differences in pain thresholds seem to be of minor clinical relevance and can be adjusted for by sex-specific reference data.

ACKNOWLEDGEMENTS

The IMI EUROPAIN project is a public-private partnership/EU-Project for understanding chronic pain and improving its treatment. The NEUROPAIN project for the characterization of subgroups of patients with neuropathic pain is an investigator-initiated European multicentre study with Prof. Dr. R. Baron as principle investigator and ten co-investigator sites. Data for these consortia was collected at the following sites: Johannes Gutenberg-University, Mainz, Germany, University of Schleswig-Holstein, Kiel, Germany, Ruhr University Bochum, Germany, Technical University Munich, Germany, Ludwig-Maximilians-University, Munich, Germany, University of Tübingen, Germany, University of Freiburg, Germany, University of Ulm, Germany, University of Erlangen, Germany, Benedictus Hospital Tutzing, Germany, Aarhus University, Denmark, University of Southern Denmark, Odense, Denmark, Karolinska Institute, Stockholm, Sweden, Imperial College London, UK, Neuroscience Technologies, Barcelona, Spain.

CONFLICT OF INTEREST

All authors declare that they have no conflicts of interest regarding the topic of this publication.

AUTHOR CONTRIBUTIONS

All authors discussed the results and commented on the manuscript (MS). The specific contribution of each author was: CMF evaluated and interpreted data and drafted the MS. NA contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. RB contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. DB contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. NBF contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. RF contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. JG contributed to patient collection, corrected parts of the MS, added critical literature review of sex and pain aspects, critically reviewed, read and agreed to the final version. MH contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. PH contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. TSJ contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. HK contributed to patient collection, corrected parts of the MS, added critical literature review of sex and pain aspects, critically reviewed, read and agreed to the final version. DK contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. ASL contributed to patient collection, corrected parts of the MS, added critical literature review of sex and pain aspects, critically reviewed, read and agreed to the final version. ASCR contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. MS contributed to study design, contributed to data interpretation, drafted parts of the manuscript, critically reviewed, read and agreed to the final version. JS contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. SS contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. RS contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. TT contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. SSH collected original data, contributed to data evaluation, critically read the manuscript and agreed to the final version. RDT contributed to study design and patient collection, corrected parts of the MS, critically reviewed,

read and agreed to the final version. EPZ proof-read the MS, added critical literature review of sex and pain aspects, critically reviewed, read and agreed to the final version. CM contributed to study design and patient collection, corrected parts of the MS, critically reviewed, read and agreed to the final version. JV designed the analyses, performed statistics and interpreted data, created the artwork, drafted and finalized the manuscript.

ORCID

Jan Vollert  <https://orcid.org/0000-0003-0733-5201>

REFERENCES

- Arap, A., Siqueira, S. R. D. T., Silva, C. B., Teixeira, M. J., & Siqueira, J. T. T. (2010). Trigeminal pain and quantitative sensory testing in painful peripheral diabetic neuropathy. *Archives of Oral Biology*, 55, 486–493. <https://doi.org/10.1016/j.archoralbio.2010.03.021>
- Aslaksen, P. M., Myrbakk, I. N., Høifødt, R. S., & Flaten, M. A. (2007). The effect of experimenter gender on autonomic and subjective responses to pain stimuli. *Pain*, 129, 260–268. <https://doi.org/10.1016/j.pain.2006.10.011>
- Attal, N., Lanteri-Minet, M., Laurent, B., Fermanian, J., & Bouhassira, D. (2011). The specific disease burden of neuropathic pain: Results of a French nationwide survey. *Pain*, 152, 2836–2843. <https://doi.org/10.1016/j.pain.2011.09.014>
- Baron, R., Maier, C., Attal, N., Binder, A., Bouhassira, D., Cruccu, G., ... Treede, R.-D. (2017). Peripheral neuropathic pain: A mechanism-related organizing principle based on sensory profiles. *Pain*, 158, 261–272. <https://doi.org/10.1097/j.pain.0000000000000753>
- Bartley, E. J., & Fillingim, R. B. (2013). Sex differences in pain: A brief review of clinical and experimental findings. *British Journal of Anaesthesia*, 111, 52–58. <https://doi.org/10.1093/bja/aet127>
- Bartley, E. J., King, C. D., Sibille, K. T., Cruz-Almeida, Y., Riley, J. L., Glover, T. L., ... Fillingim, R. B. (2016). Enhanced pain sensitivity among individuals with symptomatic knee osteoarthritis: Potential sex differences in central sensitization. *Arthritis Care & Research*, 68, 472–480. <https://doi.org/10.1002/acr.22712>
- Bouhassira, D., Lantéri-Minet, M., Attal, N., Laurent, B., & Touboul, C. (2008). Prevalence of chronic pain with neuropathic characteristics in the general population. *Pain*, 136, 380–387. <https://doi.org/10.1016/j.pain.2007.08.013>
- Breivik, H., Collett, B., Ventafridda, V., Cohen, R., & Gallacher, D. (2006). Survey of chronic pain in Europe: Prevalence, impact on daily life, and treatment. *European Journal of Pain*, 10, 287–333. <https://doi.org/10.1016/j.ejpain.2005.06.009>
- Chesterton, L. S., Barlas, P., Foster, N. E., Baxter, G. D., & Wright, C. C. (2003). Gender differences in pressure pain threshold in healthy humans. *Pain*, 101, 259–266. [https://doi.org/10.1016/S0304-3959\(02\)00330-5](https://doi.org/10.1016/S0304-3959(02)00330-5)
- Craft, R. M., Mogil, J. S., & Aloisi, A. M. (2004). Sex differences in pain and analgesia: The role of gonadal hormones. *European Journal of Pain*, 8, 397–411. <https://doi.org/10.1016/j.ejpain.2004.01.003>
- Davis, J. A., Robinson, R. L., Le, T. K., & Xie, J. (2011). Incidence and impact of pain conditions and comorbid illnesses. *Journal of Pain Research*, 4, 331–345. <https://doi.org/10.2147/JPR.S24170>
- Dawson, A., & List, T. (2009). Comparison of pain thresholds and pain tolerance levels between Middle Easterners and Swedes and between genders. *Journal of Oral Rehabilitation*, 36, 271–278. <https://doi.org/10.1111/j.1365-2842.2009.01943.x>
- de Mos, M., de Bruijn, A. G. J., Huygen, F. J. P. M., Dieleman, J. P., Stricker, B. H. C., & Sturkenboom, M. C. J. M. (2007). The incidence of complex regional pain syndrome: A population-based study. *Pain*, 129, 12–20. <https://doi.org/10.1016/j.pain.2006.09.008>
- Defrin, R., Shramm, L., & Eli, I. (2009). Gender role expectations of pain is associated with pain tolerance limit but not with pain threshold. *Pain*, 145, 230–236. <https://doi.org/10.1016/j.pain.2009.06.028>
- Demir, S. E., Ozaras, N., Karamehmetoğlu, S. S., Karacan, I., & Aytekin, E. (2010). Risk factors for complex regional pain syndrome in patients with traumatic extremity injury. *Ulusal Travma ve Acil Cerrahi Dergisi = Turkish Journal of Trauma and Emergency Surgery: TJTES*, 16, 144–148.
- Enax-Krumova, E. K., Lenz, M., Frettlöh, J., Höffken, O., Reinersmann, A., Schwarzer, A., ... Maier, C. (2017). Changes of the sensory abnormalities and cortical excitability in patients with complex regional pain syndrome of the upper extremity after 6 months of multimodal treatment. *Pain Medicine*, 18(1), 95–106.
- England, J. D., Gronseth, G. S., Franklin, G., Miller, R. G., Asbury, A. K., Carter, G. T., ... American Academy of Physical Medicine and Rehabilitation. (2005). Distal symmetric polyneuropathy: A definition for clinical research: Report of the American Academy of Neurology, the American Association of Electrodiagnostic Medicine, and the American Academy of Physical Medicine and Rehabilitation. *Neurology*, 64, 199–207. <https://doi.org/10.1212/01.WNL.0000149522.32823.EA>
- Fillingim, R. B. (2000). Sex, gender, and pain: Women and men really are different. *Current Review of Pain*, 4, 24–30. <https://doi.org/10.1007/s11916-000-0006-6>
- Fillingim, R. B., King, C. D., Ribeiro-Dasilva, M. C., Rahim-Williams, B., & Riley, J. L. (2009). Sex, gender, and pain: A review of recent clinical and experimental findings. *The Journal of Pain*, 10, 447–485. <https://doi.org/10.1016/j.jpain.2008.12.001>
- Freyenhagen, R., Baron, R., Gockel, U., & Tölle, T. R. (2006). pain-DETECT: A new screening questionnaire to identify neuropathic components in patients with back pain. *Current Medical Research and Opinion*, 22, 1911–1920. <https://doi.org/10.1185/030079906X132488>
- Friessem, C. H., Willweber-Strumpf, A., & Zenz, M. W. (2009). Chronic pain in primary care. German figures from 1991 and 2006. *BMC Public Health*, 9, 299. <https://doi.org/10.1186/1471-2458-9-299>
- Gelfand, S. (1964). The relationship of experimental pain tolerance to pain threshold. *Canadian Journal of Psychology/Revue Canadienne de Psychologie*, 18, 36–42. <https://doi.org/10.1037/h0083283>
- Gierthmühlen, J., Enax-Krumova, E. K., Attal, N., Bouhassira, D., Cruccu, G., Finnerup, N. B., ... Maier, C. (2015). Who is healthy? Aspects to consider when including healthy volunteers in QST-based studies—a consensus statement by the EUROPAIN and NEUROPAIN consortia. *Pain*, 156, 2203–2211. <https://doi.org/10.1097/j.pain.0000000000000227>
- Gierthmühlen, J., Maier, C., Baron, R., Tölle, T., Treede, R.-D., Birbaumer, N., ... German Research Network on Neuropathic Pain (DFNS) study group. (2012). Sensory signs in complex regional pain syndrome and peripheral nerve injury. *Pain*, 153, 765–774. <https://doi.org/10.1016/j.pain.2011.11.009>
- Gijssbers, K., & Nicholson, F. (2005). Experimental pain thresholds influenced by sex of experimenter. *Perceptual and Motor Skills*, 101, 803–807. <https://doi.org/10.2466/PMS.101.7.803-807>

- Goswami, R., Anastakis, D. J., Katz, J., & Davis, K. D. (2016). A longitudinal study of pain, personality, and brain plasticity following peripheral nerve injury. *Pain, 157*, 729–739. <https://doi.org/10.1097/j.pain.0000000000000430>
- Greenspan, J. D., Craft, R. M., LeResche, L., Arendt-Nielsen, L., Berkley, K. J., Fillingim, R. B., ... Traub, R. J. (2007). Studying sex and gender differences in pain and analgesia: A consensus report. *Pain, 132*, S26–S45. <https://doi.org/10.1016/j.pain.2007.10.014>
- Harden, R. N., Bruhl, S., Perez, R. S. G. M., Birklein, F., Marinus, J., Maihofner, C., ... Vatine, J.-J. (2010a). Validation of proposed diagnostic criteria (the “Budapest Criteria”) for complex regional pain syndrome. *Pain, 150*, 268–274. <https://doi.org/10.1016/j.pain.2010.04.030>
- Harden, R. N., Bruhl, S., Perez, R. S. G. M., Birklein, F., Marinus, J., Maihofner, C., ... Vatine, J.-J. (2010b). Development of a severity score for CRPS. *Pain, 151*, 870–876. <https://doi.org/10.1016/j.pain.2010.09.031>
- Hashmi, J. A., & Davis, K. D. (2014). Deconstructing sex differences in pain sensitivity. *Pain, 155*, 10–13. <https://doi.org/10.1016/j.pain.2013.07.039>
- Iacovides, S., Avidon, I., & Baker, F. C. (2015). Does pain vary across the menstrual cycle? A review. *European Journal of Pain, 19*, 1389–1405. <https://doi.org/10.1002/ejp.714>
- IASP. (2017). *IASP terminology*. Retrieved from <https://www.iasp-pain.org/Education/Content.aspx?ItemNumber=1698>
- Kállai, I., Barke, A., & Voss, U. (2004). The effects of experimenter characteristics on pain reports in women and men. *Pain, 112*, 142–147. <https://doi.org/10.1016/j.pain.2004.08.008>
- Kilo, S., Schmelz, M., Koltzenburg, M., & Handwerker, H. O. (1994). Different patterns of hyperalgesia induced by experimental inflammation in human skin. *Brain, 117*(Pt 2), 385–396. <https://doi.org/10.1093/brain/117.2.385>
- Krämer, H. H., Rolke, R., Bickel, A., & Birklein, F. (2004). Thermal thresholds predict painfulness of diabetic neuropathies. *Diabetes Care, 27*, 2386–2391. <https://doi.org/10.2337/diacare.27.10.2386>
- Lamerato, L. E., Dryer, R. D., Wolff, G. G., Hegeman-Dingle, R., Mardekian, J., Park, P. W., & Zlateva, G. (2016). Prevalence of chronic pain in a large integrated healthcare delivery system in the U.S.A. *Pain Practice, 16*, 890–898. <https://doi.org/10.1111/papr.12334>
- Levine, F. M., & De Simone, L. L. (1991). The effects of experimenter gender on pain report in male and female subjects. *Pain, 44*, 69–72. [https://doi.org/10.1016/0304-3959\(91\)90149-R](https://doi.org/10.1016/0304-3959(91)90149-R)
- Magerl, W., Krumova, E. K., Baron, R., Tölle, T., Treede, R.-D., & Maier, C. (2010). Reference data for quantitative sensory testing (QST): Refined stratification for age and a novel method for statistical comparison of group data. *Pain, 151*, 598–605. <https://doi.org/10.1016/j.pain.2010.07.026>
- Maier, C., Baron, R., Tölle, T. R., Binder, A., Birbaumer, N., Birklein, F., ... Treede, R.-D. (2010). Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): Somatosensory abnormalities in 1236 patients with different neuropathic pain syndromes. *Pain, 150*, 439–450. <https://doi.org/10.1016/j.pain.2010.05.002>
- Mainka, T., Bischoff, F. S., Baron, R., Krumova, E. K., Nicolas, V., Pennekamp, W., ... Maier, C. (2014). Comparison of muscle and joint pressure-pain thresholds in patients with complex regional pain syndrome and upper limb pain of other origin. *Pain, 155*, 591–597. <https://doi.org/10.1016/j.pain.2013.12.014>
- Meints, S. M., Wang, V., & Edwards, R. R. (2018). Sex and race differences in pain sensitization among patients with chronic low back pain. *The Journal of Pain, 19*, 1461–1470. <https://doi.org/10.1016/j.jpain.2018.07.001>
- Meyer-Frießem, C. H., Szalaty, P., Zahn, P. K., & Pogatzki-Zahn, E. M. (2019). A prospective study of patients' pain intensity after cardiac surgery and a qualitative review: Effects of examiners' gender on patient reporting. *Scandinavian Journal of Pain, 19*, 39–51. <https://doi.org/10.1515/sjpain-2018-0111>
- Mogil, J. S. (2012). Sex differences in pain and pain inhibition: Multiple explanations of a controversial phenomenon. *Nature Reviews Neuroscience, 13*, 859–866. <https://doi.org/10.1038/nrn3360>
- Moulin, D. E., Clark, A. J., Speechley, M., & Morley-Forster, P. K. (2002). Chronic pain in Canada—Prevalence, treatment, impact and the role of opioid analgesia. *Pain Research and Management, 7*, 179–184. <https://doi.org/10.1155/2002/323085>
- Ott, S., & Maihofner, C. (2018). Signs and symptoms in 1,043 patients with complex regional pain syndrome. *The Journal of Pain, 19*, 599–611. <https://doi.org/10.1016/j.jpain.2018.01.004>
- Packiasabapathy, S., & Sadhasivam, S. (2018). Gender, genetics, and analgesia: Understanding the differences in response to pain relief. *Journal of Pain Research, 11*, 2729–2739. <https://doi.org/10.2147/JPR.S94650>
- Pfau, D. B., Krumova, E. K., Treede, R.-D., Baron, R., Toelle, T., Birklein, F., ... Maier, C. (2014). Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): Reference data for the trunk and application in patients with chronic postherpetic neuralgia. *Pain, 155*, 1002–1015. <https://doi.org/10.1016/j.pain.2014.02.004>
- Pfau, D. B., Rolke, R., Nickel, R., Treede, R.-D., & Daublaender, M. (2009). Somatosensory profiles in subgroups of patients with myogenic temporomandibular disorders and fibromyalgia syndrome. *Pain, 147*, 72–83. <https://doi.org/10.1016/j.pain.2009.08.010>
- Pogatzki-Zahn, E. M., Mengersen, C., Englbrecht, J. S., Klein, T., Magerl, W., & Zahn, P. K. (2019). Progesterone relates to enhanced incisional acute pain and pinprick hyperalgesia in the luteal phase of female volunteers. *Pain, 160*(8), 1781–1793.
- Pons, T., Shipton, E. A., Williman, J., & Mulder, R. T. (2015). Potential risk factors for the onset of complex regional pain syndrome type 1: A systematic literature review. *Anesthesiology Research and Practice, 2015*, 956539. <https://doi.org/10.1155/2015/956539>
- Racine, M., Tousignant-Laflamme, Y., Kloda, L. A., Dion, D., Dupuis, G., & Choinière, M. (2012). A systematic literature review of 10 years of research on sex/gender and experimental pain perception—Part 1: Are there really differences between women and men? *Pain, 153*, 602–618. <https://doi.org/10.1016/j.pain.2011.11.025>
- Rahim-Williams, B., Riley, J. L., Williams, A. K. K., & Fillingim, R. B. (2012). A quantitative review of ethnic group differences in experimental pain response: Do biology, psychology, and culture matter? *Pain Medicine, 13*, 522–540. <https://doi.org/10.1111/j.1526-4637.2012.01336.x>
- Reitsma, M., Tranmer, J. E., Buchanan, D. M., & VanDenKerkhof, E. G. (2012). The epidemiology of chronic pain in Canadian men and women between 1994 and 2007: Longitudinal results of the National Population Health Survey. *Pain Research and Management, 17*, 166–172. <https://doi.org/10.1155/2012/875924>
- Riley, J. L., Robinson, M. E., Wise, E. A., Myers, C. D., & Fillingim, R. B. (1998). Sex differences in the perception of noxious experimental

- stimuli: A meta-analysis. *Pain*, 74, 181–187. [https://doi.org/10.1016/S0304-3959\(97\)00199-1](https://doi.org/10.1016/S0304-3959(97)00199-1)
- Roh, Y. H., Lee, B. K., Noh, J. H., Baek, J. R., Oh, J. H., Gong, H. S., & Baek, G. H. (2014). Factors associated with complex regional pain syndrome type I in patients with surgically treated distal radius fracture. *Archives of Orthopaedic and Trauma Surgery*, 134, 1775–1781. <https://doi.org/10.1007/s00402-014-2094-5>
- Rolke, R., Baron, R., Maier, C., Tölle, T. R., Treede, R.-D., Beyer, A., ... Wasserka, B. (2006). Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): Standardized protocol and reference values. *Pain*, 123, 231–243. <https://doi.org/10.1016/j.pain.2006.01.041>
- Schäfer, A. G. M., Hall, T. M., Rolke, R., Treede, R.-D., Lütke, K., Mallwitz, J., & Briffa, K. N. (2014). Low back related leg pain: An investigation of construct validity of a new classification system. *Journal of Back and Musculoskeletal Rehabilitation*, 27, 409–418. <https://doi.org/10.3233/BMR-140461>
- Selim, M. M., Wendelschafer-Crabb, G., Hodges, J. S., Simone, D. A., Foster, S.-X.-Y.-L., Vanhove, G. F., & Kennedy, W. R. (2010). Variation in quantitative sensory testing and epidermal nerve fiber density in repeated measurements. *Pain*, 151, 575–581. <https://doi.org/10.1016/j.pain.2010.06.034>
- Torrance, N., Smith, B. H., Bennett, M. I., & Lee, A. J. (2006). The epidemiology of chronic pain of predominantly neuropathic origin. Results from a general population survey. *The Journal of Pain*, 7, 281–289. <https://doi.org/10.1016/j.jpain.2005.11.008>
- Üçeyler, N., Vollert, J., Broll, B., Riediger, N., Langjahr, M., Saffer, N., ... Sommer, C. (2018). Sensory profiles and skin innervation of patients with painful and painless neuropathies. *Pain*, 159, 1867–1876. <https://doi.org/10.1097/j.pain.0000000000001287>
- Vigil, J. M., & Alcock, J. (2014). Tough guys or sensitive guys? Disentangling the role of examiner sex on patient pain reports. *Pain Research and Management*, 19, e9–e12. <https://doi.org/10.1155/2014/531396>
- Vollert, J., Attal, N., Baron, R., Freynhagen, R., Haanpää, M., Hansson, P., ... Maier, C. (2016). Quantitative sensory testing using DFNS protocol in Europe: An evaluation of heterogeneity across multiple centers in patients with peripheral neuropathic pain and healthy subjects. *Pain*, 157, 750–758. <https://doi.org/10.1097/j.pain.0000000000000433>
- Vollert, J., Kramer, M., Barroso, A., Freynhagen, R., Haanpää, M., Hansson, P., ... Baron, R. (2016). Symptom profiles in the painDETECT Questionnaire in patients with peripheral neuropathic pain stratified according to sensory loss in quantitative sensory testing. *Pain*, 157, 1810–1818. <https://doi.org/10.1097/j.pain.0000000000000588>
- Vollert, J., Magerl, W., Baron, R., Binder, A., Enax-Krumova, E. K., Geisslinger, G., ... Treede, R.-D. (2018). Pathophysiological mechanisms of neuropathic pain: Comparison of sensory phenotypes in patients and human surrogate pain models. *Pain*, 159, 1090–1102. <https://doi.org/10.1097/j.pain.0000000000001190>
- Vollert, J., Maier, C., Attal, N., Bennett, D. L. H., Bouhassira, D., Enax-Krumova, E. K., ... Baron, R. (2017). Stratifying patients with peripheral neuropathic pain based on sensory profiles: Algorithm and sample size recommendations. *Pain*, 158, 1446–1455. <https://doi.org/10.1097/j.pain.0000000000000935>
- Vollert, J., Mainka, T., Baron, R., Enax-Krumova, E. K., Hüllemann, P., Maier, C., ... Treede, R.-D. (2015). Quality assurance for Quantitative Sensory Testing laboratories: Development and validation of an automated evaluation tool for the analysis of declared healthy samples. *Pain*, 156, 2423–2430. <https://doi.org/10.1097/j.pain.0000000000000300>
- Williamson, P. R., Altman, D. G., Bagley, H., Barnes, K. L., Blazeby, J. M., Brookes, S. T., ... Young, B. (2017). The COMET handbook: Version 1.0. *Trials*, 18, 280. <https://doi.org/10.1186/s13063-017-1978-4>

How to cite this article: Meyer-Frießem CH, Attal N, Baron R, et al. Pain thresholds and intensities of CRPS type I and neuropathic pain in respect to sex. *Eur J Pain*. 2020;24:1058–1071. <https://doi.org/10.1002/ejp.1550>