



HAL
open science

Carbon Leakage through the Terrestrial-aquatic Interface: Implications for the Anthropogenic CO₂ Budget

Pierre Régnier, Ronny Lauerwald, Philippe Ciais

► **To cite this version:**

Pierre Régnier, Ronny Lauerwald, Philippe Ciais. Carbon Leakage through the Terrestrial-aquatic Interface: Implications for the Anthropogenic CO₂ Budget. *Procedia Earth and Planetary Science*, 2014, 10, pp.319-324. 10.1016/j.proeps.2014.08.025 . hal-02927894

HAL Id: hal-02927894

<https://hal.science/hal-02927894>

Submitted on 24 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geochemistry of the Earth's Surface meeting, GES-10

Carbon leakage through the terrestrial-aquatic interface: Implications for the anthropogenic CO₂ budget

Pierre Regnier^{a*}, Ronny Lauerwald^{a,b}, Philippe Ciais^b

^aUniversité Libre de Bruxelles, DSTE, 50, av. F.D. Roosevelt, 1050 Bruxelles, Belgium

^bIPSL-LSCE, Centre d'Etudes Orme des Merisiers, 91191 Gif sur Yvette, France

Abstract

This contribution reviews the role of the terrestrial-aquatic interface in the global carbon cycle. We highlight that carbon leakage through this interface has profound ramifications for the terrestrial carbon balance and for the anthropogenic CO₂ budget. Our budget analysis identifies the need for an integrated process-based quantitative understanding of the terrestrial-aquatic interface that includes its response to anthropogenic perturbations. Complementary observational and modeling efforts in this direction are presented.

© 2014 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of the Scientific Committee of GES-10

Keywords:

1. Introduction

Since the beginning of the Industrial Era, CO₂ emissions due to fossil fuel combustion and cement production (E_{FF}) and land-use change (E_{LUC}) have led to a rapid accumulation of the carbon mass in the atmosphere^[1-4]. For the decade 2003-2012, E_{FF} and E_{LUC} have been estimated at 8.6±0.4 PgC yr⁻¹ and 0.9±0.5 PgC yr⁻¹, respectively. Altogether, anthropogenic emissions thus amount to 9.5 PgC yr⁻¹. Over the same period, the atmospheric CO₂ concentrations have increased by about 20ppm, which translates in a mean annual growth rate G_{ATM} of 4.3±0.1 PgC yr⁻¹ [3]. This implies that about 45 % of the anthropogenic emissions accumulate in the atmosphere, the

* Pierre Regnier. Tel.: +32-26503671
E-mail address: pregnier@ulb.ac.be

remainder, about 55 %, being taken up by the land (S_{LAND}) and ocean (S_{OCEAN}) ‘CO₂ sinks’.

From mass balance, the following equation constrains the anthropogenic CO₂ budget:

$$E_{\text{FF}} + E_{\text{LUC}} = G_{\text{ATM}} + S_{\text{LAND}} + S_{\text{OCEAN}} \quad (1)$$

The land sink term S_{LAND} has traditionally been estimated from the closure of the CO₂ budget, a contribution referred to as the ‘residual land sink’. Based on the partitioning of anthropogenic carbon, the net storage on land is:

$$\Delta C_{\text{LAND}} = S_{\text{LAND}} - E_{\text{LUC}} \quad (2)$$

Equation 1 considers only anthropogenic CO₂ emissions and their distribution among the atmosphere, ocean and land. This analysis thus omits that carbon is continuously displaced along the land-ocean aquatic continuum (LOAC) comprising freshwaters, estuaries and coastal waters. A significant fraction of this lateral carbon flux is a steady state component of the pre-industrial carbon cycle that can be ignored in the above budget analysis, which focuses only on the anthropogenic perturbation. However, the remaining fraction, which corresponds to the change in LOAC carbon fluxes since pre-industrial conditions, is a perturbation that is relevant for the global carbon budget. The inclusion of these lateral fluxes of anthropogenic CO₂ will affect the estimates of S_{LAND} and S_{OCEAN} in Eq. (1).

2. The land carbon sink and terrestrial carbon storage

For the past five decades, terrestrial ecosystems have been absorbing 25–30% of anthropogenic CO₂ emissions, with a major contribution from the carbon accumulation in forest biomass and soils^[5,6]. In the classical view which ignores the LOAC carbon cycle, the land sink results from a disequilibrium between terrestrial NPP and soil respiration plus fire disturbance emissions of carbon, initiated by human-induced changes in environmental conditions (both direct effects such as regrowth after land use change, and indirect biogeochemical effects such as the effects of rising CO₂, climate change and nitrogen deposition), its persistence being related to the residence time of carbon in the terrestrial biosphere (from a few decades to several centuries). Several processes contribute to S_{LAND} , but their precise quantification, both at the regional and global scales, remain entailed with large uncertainties. In particular, the modeling of soil carbon dynamics is poorly constrained^[7], as witnessed by the large discrepancies in estimated present-day stocks (from about 500 and 3000 PgC) and predicted future changes (cumulative change of land carbon storage ignoring LOAC, ranging from -165 PgC to +500 PgC for the 21st century, according to the spread between different scenarios and different coupled carbon-climate models^[8]). Proposed mechanisms behind S_{LAND} include the fertilization effect of CO₂ and nitrogen on vegetation growth and increasing growing season length in high latitudes attributed to climate warming^[6,9]. Overall, S_{LAND} mitigates the anthropogenic increase of atmospheric CO₂ levels, and provides a negative feedback in the climate-carbon cycle system^[10].

Pan et al.^[5] have estimated forest carbon sources and sinks that allow to constrain ΔC_{LAND} (the contribution from croplands and grasslands to biomass accumulation is much smaller). Consecutive forest inventories indicate a substantial 4.0 PgC yr⁻¹ sink with 2.9 PgC yr⁻¹ in biomass, 0.9 PgC yr⁻¹ in litter and soil, and 0.2 PgC yr⁻¹ in harvest products. This sink is partially offset by emissions from gross deforestation of tropical forests (2.8 PgC yr⁻¹) and the net C increase in forest thus reduces to 1.2 PgC yr⁻¹. The partition of this net C increase (1.0 PgC yr⁻¹ without the accumulation into harvest products) between biomass, litter and soil can be calculated if one takes into account that soil carbon loss represents on average 25% of total gross deforestation emissions^[11]. This leads to about 0.8 PgC yr⁻¹ accumulation in biomass and 0.2 PgC yr⁻¹ accumulation in litter and soil^[11]. In addition, drainage of peatlands contributes to an estimated carbon loss of about 0.4 PgC yr⁻¹ ^[12-13] and the total present-day soil C reservoir (including peatlands) could thus lose about 0.2 PgC yr⁻¹. Altogether, this leads to a ‘bottom-up’ estimate of the net carbon sequestration in terrestrial ecosystems of about 0.8 PgC yr⁻¹.

3. A leakage in the land carbon sink

The contribution of the Land-Ocean Aquatic Continuum (LOAC) to the global carbon budget is synthesized in

Regnier et al.^[11]. In what follows, we summarize the main findings concerning the contribution of the LOAC to the CO₂ anthropogenic perturbation and briefly discuss the implications for the estimates of the net carbon storage on land. The anthropogenic C input from terrestrial ecosystems to freshwaters has been estimated at 1 PgC yr⁻¹. This input can be decomposed into four fluxes attributed to soil-derived C, chemical weathering of continental surfaces, sewage and net C fixation in inland waters. Altogether, these four perturbation fluxes imply that only a portion of the anthropogenic CO₂ taken up by land ecosystems (S_{LAND}) remain sequestered in soil and biomass pools, as 1 ± 0.5 PgC yr⁻¹ is exported to the LOAC. This land to hydrosphere carbon flux, F_{LH}, of magnitude comparable to E_{LUC}, is thus significant for the anthropogenic CO₂ budget.

The exported anthropogenic C accumulates to a large part in sediments of freshwaters, estuaries and coastal seas and this term can thus be viewed as the LOAC sink (S_{LOAC}, 0.55 ± 0.3 PgC yr⁻¹). Another fraction is released back to the atmosphere (E_{LOAC}, 0.35 ± 0.2 PgC yr⁻¹), the magnitude of this flux resulting from the combined effects of CO₂ outgassing in inland waters and CO₂ uptake in the coastal ocean^[11,14]. Finally, a small fraction of anthropogenic C displaced by the LOAC accumulates in the open ocean (0.1 ± > 0.05 PgC yr⁻¹).

The carbon budget following Eq.1 implicitly incorporates the fluxes from the LOAC with S_{LAND}. However, if Terrestrial Ecosystems and the LOAC are separated in the analysis, the residual sink attributed to terrestrial ecosystems alone is now larger (3.15 ± 0.9 PgC yr⁻¹) than the value of 2.8 PgC yr⁻¹ reported for S_{LAND} in the classical analysis, because this flux is partially offset by E_{LOAC}. Most importantly, because anthropogenic CO₂ taken up by land ecosystems is exported to the LOAC, the annual land carbon storage change (1.25 PgC yr⁻¹) is notably smaller than the net CO₂ uptake by land ecosystems as calculated by Equation 2 (1.9 PgC yr⁻¹), a significant fraction of the displaced carbon (0.65 PgC yr⁻¹) being stored in freshwater and coastal sediments (S_{LOAC}), and to a lesser extent, in the open ocean. With the LOAC included, we now have:

$$\Delta C_{\text{LAND}} = S_{\text{LAND}} - E_{\text{LUC}} - F_{\text{LH}} \quad (3)$$

The net biomass and soil sequestration estimate calculated according to Equation 3 is broadly consistent with the ‘bottom-up’ estimates derived from biomass and soil-carbon inventories of Pan et al.^[5], thus providing additional support for a significant soil to hydrosphere anthropogenic carbon leakage. This budget analysis clearly identifies the need for an integrated process-based quantitative understanding of the LOAC response to anthropogenic perturbations, which include factors as diverse as land-use and climate change, hydraulic management, and agricultural, domestic and industrial activities. The anthropogenic perturbations affect both, the soil to hydrosphere C leakage and the fate of terrestrial C within the LOAC. Hydraulic management, for instance, is capable of changing the amount of C stored in sediments and to change the net-exchange of CO₂ with the atmosphere. An integrated quantitative understanding of the LOAC response to the anthropogenic perturbations should rely on spatially-resolved contemporary budgets of lateral C transport, retention in sediments and loss through decomposition and degassing. Key environmental drivers of these fluxes need to be identified to assess their sensitivity with respect to global change and modeling efforts need to be pursued to support improved estimates of the anthropogenic perturbations to the LOAC carbon cycle. Ongoing research efforts in these directions are discussed in section 4 and 5 of this contribution.

4. High resolution maps of CO₂ evasion from rivers

Observations of fluvial fluxes of solutes are a useful integrative measure for Earth Surface processes within a river catchment. Measurements of fluvial carbon fluxes, in particular, allow reassessing the C balance of terrestrial ecosystems by including the lateral carbon fluxes. A monitoring at the outlet of a river basin is, however, not sufficient because organic C and CO₂ stemming from soil respiration are not conservative during lateral transport and net-fluxes of CO₂ from the rivers to the atmosphere are significant^[15]. At regional and global scales, a spatially explicit assessment of CO₂ evasion from the river is thus required to improve the contemporary terrestrial carbon budget. In a fully data driven approach, a dense network of water chemistry observations, from which CO₂ partial pressures (pCO₂) and vertical river-atmosphere CO₂ fluxes (FCO₂) could be calculated, would be necessary. Such monitoring network is currently not available as only industrialized countries have sufficient data coverage. Large parts of the globe remain only sparsely or not at all covered, including hotspot areas such as tropical Africa and Asia

and Siberia^[11].

In an ongoing study (Lauerwald et al. in prep), an empirical model for river pCO₂ was developed to produce a global river pCO₂ map at half a degree resolution. Compared to Raymond et al., 2013^[15], which identifies broad regional patterns in global CO₂ evasion, this research thus takes the next step forward. Combining the pCO₂ map with spatially explicit estimates on stream surface area (A_{river}) and gas exchange velocity (k) relying on empirical equations from the literature^[15,16], the first global FCO₂ map was created, at a resolution compatible with that of Earth System Models. The empirical pCO₂ model was trained on 1182 sampling locations with suitable time-series, covering all latitudes and biomes from the high north down to the central Tropics. The identified predictors of pCO₂ are average net primary production (NPP, g C m⁻²yr⁻¹), population density (PD, inhabitants per km²), slope gradient within the river catchment (S, in °), and average air temperature at the sampling location (T, in °C).

In contrast to the predictor S, which is negatively correlated to river pCO₂ and will not change significantly over the next centuries, the other three predictors have a positive effect on river pCO₂ and are predicted to change considerably during the 21st century. Projections of these drivers for the year 2100 can thus be used to investigate plausible alterations of the global river CO₂ evasion. Expected increases in PD and T, and partly NPP due to CO₂ fertilization, will likely promote an increase in river pCO₂. The projected increase in atmospheric pCO₂ FCO₂ could partly alleviate this effect by reducing the river-atmosphere pCO₂ gradient and FCO₂. Changes in river discharge resulting from modifications in precipitation patterns will entrain changes in A_{river}, which will also have a direct effect of FCO₂.

The predicted FCO₂ map produced for present-day conditions is used here to revisit the C budget of the Amazon River, as an example. The FCO₂ from streams and rivers is estimated at 95 ± 25 TgC yr⁻¹ (Lauerwald et al., in prep). In a recent study, Abril et al.^[17] estimated that floodplains of the central Amazon Basin contribute to a FCO₂ of 210 ± 60 TgC yr⁻¹. Together with the 29 TgC yr⁻¹ of DOC and 18 TgC yr⁻¹ POC exported to the Atlantic Ocean [Global NEWS2^[18]], the total export of C originally fixed by plants adds up to about 350 TgC yr⁻¹, which represents about 8.5 % of the terrestrial NPP within the Amazon basin. However, a significant proportion of the FCO₂ from the floodplains but also from adjacent rivers was reported to originate from the root respiration of semi-aquatic, emergent vegetation on the floodplains^[17], and the contribution of terrestrial NPP to FCO₂ is thus likely lower. Projections on NPP, T, PD, atmospheric pCO₂ and river discharge will be used to provide a first-order estimate of how FCO₂ in the Amazon Basin might change until 2100.




Fig 1: Modelled average river pCO₂, stream surface area (A_{river}), and CO₂ evasion from rivers (FCO₂) in the Amazon River Basin for the period 1990-2010. Units of FCO₂ refer to total continental areas, incl. inland waters.

5. Perspectives: Including fluvial C transport into the land surface scheme ORCHIDEE

Fluvial C transport is currently ignored in Earth System models (ESMs) and the terrestrial C balance thus only considers NPP, soil heterotrophic respiration and land carbon storage. The integration of C leakage in ESMs will improve the allocation of the land and ocean C sinks and will allow attributing changes in lateral C fluxes to environmental drivers. In addition, the modelled fluvial C exports could also be used as input for global ocean biogeochemistry models. This objective, complementary to the empirical approach described above, requires an update of the land surface scheme ORCHIDEE by integrating the fluvial transfer of dissolved organic C (DOC) and dissolved CO₂ from soils to the ocean, including DOC decomposition and CO₂ evasion to the atmosphere during lateral transport (Fig.2).




Fig. 2: Schematic representation of the modelled water fluxes in ORCHIDEE. Geometrically, the river routing scheme (a) is organized in grid boxes (ig), which are further subdivided into basins (ib). It attributes, for each basin, the influxes of water and dissolved C received from neighboring basins. For each basin, the different reservoirs store and exchange water are defined (b). Blue arrows represent water fluxes from or to the reservoirs. ET: evapotranspiration, E: evaporation, I: infiltration, P: precipitation.

ORCHIDEE was recently upgraded to better represent soil organic C (SOC) pools and SOC decomposition (Guenet and Camino-Serrano, pers.com.). In particular, the soil hydrology module was updated to include DOC and CO₂ export from soil drainage (fig. 2). The river routing scheme of ORCHIDEE has now been enabled for DOC and CO₂ transport with the water flow. For each basin, the routing scheme distinguishes three main dynamic reservoirs for water, DOC and dissolved CO₂: the 'stream reservoir' (the river itself), the 'fast reservoir' (surface runoff) and the 'slow reservoir' (soil drainage). The routing scheme also represents the water exchange between the stream and the floodplain reservoirs, which are important for river systems like the Amazon^[17]. At each time step, a fraction of the DOC stored in the reservoirs is converted to CO₂ based on a reservoir-specific decomposition rate. The CO₂ evasion from the rivers and floodplains is then calculated. The ORCHIDEE enabled for lateral C transport is currently calibrated for the Amazon, a basin for which the hydrology has already been calibrated and validated^[19]. Ultimately, ORCHIDEE enabled for lateral C transport could be used to estimate the cumulative global land carbon storage change since the beginning of the Industrial Era. In the 5th assessment report of the IPCC^[4], a value of 30 ± 45 PgC was reported for this accumulation. Although it is a robust measurement-based estimate calculated as the residual of the other terms (anthropogenic emissions, atmospheric CO₂ growth rate and measured ocean C content change), the cumulative land C storage could actually be significantly smaller because of the continuous displacement of anthropogenic carbon through the terrestrial-aquatic interface.

Acknowledgements

Ronny Lauerwald was funded by the French National Research Agency ("Investissement d'Avenir", n°ANR-10-LABX-0018").

References

1. Houghton, R.A. How well do we know the flux of CO₂ from land-use change ? *Tellus B* 62(5), 337-351 (2010).
2. Boden et al., Global, Regional, and National Fossil-Fuel CO₂ Emissions, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tenn., USA, (2013).
3. Le Quéré, C. et al. Global carbon budget 2013. *Earth Syst. Sci. Data* 6, 689–760 (2014).
4. Ciais, P. et al. Chapter 6: Carbon and Other Biogeochemical Cycles, in: *Climate Change 2013 The Physical Science Basis*, edited by: Stocker, T., Qin, D., and Plattner, G.-K., Cambridge University Press, Cambridge (2013).
5. Pan, Y. et al. A large and persistent carbon sink in the world's forests. *Science* 333, 988-993 (2011).
6. Reichstein, M. et al. Climate extremes and the carbon cycle. *Nature* 500, 287–295 (2013).
7. Todd-Brown, K.E.O. et al. Causes of variation in soil carbon simulations from CMIP5 Earth system models and comparison with observations, *Biogeosciences* 10, 1717–1736 (2013).
8. Nishina, K. et al. Quantifying uncertainties in soil carbon responses to changes in global mean temperature and precipitation. *Earth Syst. Dynam.* 5, 197–209 (2014).
9. Denman, K. et al., 2007 Couplings Between Changes in the Climate System and Biogeochemistry, Intergovernmental Panel on Climate Change, 499–587 (2007).
10. Friedlingstein, P. et al. Climate-carbon cycle feedback analysis: Results from the C4MIP model intercomparison, *J. Climate* 19, 3337–3353 (2006).
11. Regnier, P. et al. Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nature Geoscience* 6, 597–607 (2013).
12. Hooijer, A. et al. Current and future CO₂ emissions from drained peatlands in Southeast Asia. *Biogeosciences* 7, 1505-1514 (2010).
13. Yu, Z. Holocene carbon flux histories of the world's peatlands: Global carbon-cycle implications. *Holocene* 21, 761-774 (2011).
14. Bauer, J. E. et al. The changing carbon cycle of the coastal ocean. *Nature* 504, 61–70 (2013).
15. Raymond, P. A. et al. Global carbon dioxide emissions from inland waters. *Nature* 503, 355–359 (2013).
16. Raymond, P. A. et al. Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers. *Limnology & Oceanography: Fluids & Environments* 2, 41–53 (2012).
17. Abril, G. et al. Amazon River carbon dioxide outgassing fuelled by wetlands. *Nature* 505, 395–398, (2014).
18. Mayorga, E. et al. Global Nutrient Export from WaterSheds 2 (NEWS 2): Model development and implementation. *Environmental Modelling & Software* 25, 837–853 (2010).
19. Guimberteau, M. et al. Future changes in precipitation and impacts on extreme streamflow over Amazonian sub-basins. *Environmental Research Letters* 8, (2013).