



**HAL**  
open science

# Northern Hemisphere Storm Tracks in Present Day and Last Glacial Maximum Climate Simulations: A Comparison of the European PMIP Models\*

M. Kageyama, P. J. Valdes, G. Ramstein, C. Hewitt, U. Wyputta

► **To cite this version:**

M. Kageyama, P. J. Valdes, G. Ramstein, C. Hewitt, U. Wyputta. Northern Hemisphere Storm Tracks in Present Day and Last Glacial Maximum Climate Simulations: A Comparison of the European PMIP Models\*. *Journal of Climate*, 1999, 12 (3), pp.742-760. 10.1175/1520-0442(1999)0122.0.CO;2 . hal-02931782

**HAL Id: hal-02931782**

**<https://hal.science/hal-02931782>**

Submitted on 8 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.











8 hPa at around (45°N, 180°). The (shaded) region of variances above 5 hPa stretches from the eastern coast of Japan to the interior of the American continent. These features are well reproduced in the UGAMP T42, ECHAM3, LSCCELMD5H, and UKMO simulations, with maxima of 8 hPa for the first two and 9 hPa for the last two, but less so in the others: the variance maximum in the LSCCELMD5 run is above 8 hPa, but is situated too much to the west (45°N, 160°E) and the corresponding region of high variability does not extend enough to the east. Both the UGAMP T21 and LSCCELMD4 runs exhibit weaker maxima (slightly above 6 hPa, well located for the UGAMP T21 run but too much to the west for LSCCELMD4) and storm tracks that are too limited in the eastward direction.

The Atlantic storm track reaches its maximum of more than 9 hPa at around (50°N, 50°W) in the ERA, and the shaded region of variances above 5 hPa extends from the American to the European continents. Its structure is more complex than the Pacific one, since from the main core of the storm track situated over Newfoundland, there is an extension of the storm track northeastward, over Iceland, and a sharp decrease in the variability east of Iceland compared to the smooth decrease eastward of the maximum of the Pacific storm track. It therefore appears stronger but shorter than the Pacific storm track. In addition, there is a second maximum over eastern Greenland. The main core of the storm track is correctly located in all simulations. As for the Pacific storm track, its magnitude is quite well reproduced in the ECHAM3, UGAMP T42, UKMO, and LSCCELMD5H runs, although the value of the maximum in all these simulations, except in the ECHAM3 one, is a little weak (7 or 8 hPa). Even though the main core of the storm track is fairly well represented by these models, the finer structure is not well captured. The high variability over Greenland is absent or only suggested, whereas the northeastward extension of the storm track is generally present but it does not decrease as sharply east of Iceland. ECHAM3 is the only model simulating a strong enough Atlantic storm track compared to the Pacific one, but in this model, the mean sea level pressure variance does not abruptly decrease to the east of Iceland (see, e.g., the 7-hPa contour); its Atlantic storm track is a little too long. For the other runs the main core maximum is generally too weak (7 hPa for UGAMP T21, 6 hPa for LSCCELMD5, 5 hPa for LSCCELMD4) but correctly located, and as was observed for the Pacific storm track, the length of the region of maximum activity is also restricted in the eastward direction. These storm tracks are generally too weak for the finer structure of the observed storm track to be captured.

A comparison of the 500-hPa geopotential height variances (not shown) confirms that even though all models correctly simulate high variability off the east coast of the continents, the storm tracks are better simulated in their amplitude and longitudinal extension in the higher resolution simulations (ECHAM3, UGAMP T42,

LSCCELMD5H, UKMO). However, from this point of view too few models have the stronger Atlantic storm track shown by the ERA. In the end, only ECHAM3 has storm tracks reaching a correct amplitude with the right balance between them.

For the LGM case we first comment on the prescribed SST runs and then the computed SST ones, as the conditions for the development of the extratropical perturbations are different for these two types of simulations. In the first type of experiments the SSTs are set to the CLIMAP dataset in all models. In the second type of runs, they are allowed to vary, within the constraints of prescribed oceanic meridional heat fluxes. The prescribed ocean heat fluxes vary between the models. In all cases, they are chosen so that they well reproduce present day conditions. Over sea ice, the models are free to predict the surface temperature. In addition, the treatment of the ocean heat fluxes under the extensive sea ice varies between models. The resulting SSTs are shown and further discussed in section 4c but an important fact about them is that they are warmer in the northeastern Atlantic, hence being far less zonal than the CLIMAP distribution in this region.

In all the prescribed SST simulations, both storm tracks significantly shift and/or extend eastward from present day (Fig. 1) to LGM (Fig. 2). The Pacific storm track as described by the mean sea level pressure variance (Fig. 2) shows a larger shift than as described by the 500-hPa geopotential height variance (not shown). On the other hand, the Atlantic storm track undergoes a comparable shift to the east in both diagnostics from present day to LGM simulations, the maxima for the LGM being situated off the European coast, if not over Europe itself for the LSCCELMD4, LSCCELMD5H, and ECHAM3 runs. Thus, the response to the changes in boundary conditions is shallower over the Pacific than over the Atlantic: for the Pacific storm track, the main difference in the boundary conditions between the LGM and the control is in the SST distribution. Looking more closely to temperature and wind latitude–altitude sections for the western Pacific (not shown) it is evident that the lower half of the troposphere is more perturbed than the upper half, which is relatively unchanged. This will also be seen in the next section in Figs. 6–10. These show how the Atlantic jet stream is much more perturbed at LGM than the Pacific one. Indeed, in the Atlantic case, the presence of the Laurentide ice sheet modifies the flow already affected by changes over the Pacific. The resulting flow, which arrives on a colder Atlantic Ocean, differs from the present day one throughout the troposphere (the differences in SSTs are also larger). Hence, it is not surprising that the changes in transient activity over the Pacific occur more at the surface than at upper levels, while they occur at all levels over the Atlantic.

The computed SST simulations (Fig. 3) show less dramatic changes in the geographical position of the storm tracks from present day to LGM. The Pacific






FIG. 4. Length of the 65-m contour of the high-pass 500-hPa geopotential variance. The largest extension is considered to be of 150° long for both storm tracks. The errors bars given are for ± one standard deviation around the mean value. Circle: present day storm tracks, squares: LGM storm tracks; black symbol: Pacific storm track, white symbol: Atlantic storm track.

storm track does not move significantly but, as in the prescribed SST simulations, the Atlantic storm track shows a significant shift to the east at all levels (geopotential height variance at 500 hPa not shown), except in the LSCCLMD4 simulations where variances are weak and quite stable. Nevertheless this shift is not as large as in the prescribed SST case. Hence, the differences in the position of the storm tracks between present day and LGM simulations are more important in the prescribed SST simulations than in the computed SST ones. For the two models for which there is a computed and a prescribed SST run, the difference between those is more significant at LGM than at present day. These differences are detailed in Dong and Valdes (1998) for the UGAMP model, and will be discussed in section 4c.

Figure 4 summarizes the previous results by giving the average length of each storm track for each simulation. The longitudinal extension of the storm track has been retrieved for each DJF season by looking for the grid boxes where the high-frequency 500-hPa geopotential variance was higher than 65 m in the 30°–60°N, 105°E–105°W region for the Pacific storm track and in the 30°–70°N, 105°W–45°E region for the Atlantic storm track. Therefore, the mean but also the standard deviation of these results are presented, giving an indication of the variability of the storm tracks, which could not be shown on the previous figures. Figure 4 confirms that the higher resolution models' Pacific storm tracks lie within the range of the ERA, whereas for the Atlantic storm track, only the ECHAM model approaches the ERA results. For all the simulations, there is a clear separation between the lower (LSCCLMD4, LSCCLMD5, UGAMP T21) and higher (LSCCLMD5H, UKMO, UGAMP T42, ECHAM3) resolution models, with significantly shorter storm tracks in the former ones. All models except LSCCLMD4c show an increase in the Atlantic storm track from the present day to the

LGM, which appears quite robust, while the changes in the Pacific storm-track length seem less so, although the higher resolution models consistently give an increase of the storm track, except UGAMP T42p.

For both diagnostics discussed here, the differences in the storm tracks simulated in the present day runs are confirmed by the LGM ones. They seem to be mainly linked to the resolution of the models, as was suggested in previous studies (Boville 1991; Rind 1988; Senior 1995) and as is evident from the comparison of the UGAMP T42 and T21 runs and the LSCCLMD5H and LSCCLMD5 ones. However, resolution is not the only factor explaining the differences in the storm-track representation, which is proved by the differences between the UGAMP T42 and the ECHAM3 T42 simulations, and between the LSCCLMD5 and UKMO runs, which are nevertheless smaller than between models of different resolutions. What cannot be seen from Fig. 4 is the position of the storm track and its changes from present day to LGM simulations. Nearly all models give an eastward shift for both storm tracks, more marked for the Atlantic one. The two different diagnostics discussed here show consistent changes from present day to LGM, although these changes can be different at the surface and at 500 hPa, as exemplified by the Pacific storm-track case in the prescribed SST simulations. We can also note here that while the present day Atlantic storm-track maximum is (correctly) situated at the same location for all models, this position fluctuates for all other cases. The mechanism responsible for this feature must be different for the particular case of the present day Atlantic storm track. In the next section, the study of the three-dimensional **E** vectors allows us to diagnose the storm tracks at different stages of their life cycle and at their corresponding levels and to describe them in a more precise way. We also examine the relationship between the storm tracks, the mean flow, and the surface climatological characteristics.

#### 4. Differences between the storm tracks: Reasons and implications

In this section, we examine more closely the relationship between the storm tracks and the baroclinicity of the simulated mean general circulation, as described by the meridional gradients in surface temperatures and the vertical shear of the zonal mean flow. We then show how the characteristics of the precipitation patterns are linked to those of the storm tracks and associated stationary waves. After introducing observed results for the present climate, we concentrate on three points: the mechanisms responsible for the differences between models of different resolutions (UGAMP and LSCCLMD runs) in section 4a; the differences due to the representation of the physics in two models using the same representation of the dynamics, at the same resolution (UGAMP and ECHAM3) in section 4b; the differences due to the computation of the SSTs (com-



































