



HAL
open science

A note on the simultaneous edge coloring

Nicolas Bousquet, Bastien Durain

► **To cite this version:**

Nicolas Bousquet, Bastien Durain. A note on the simultaneous edge coloring. *Discrete Mathematics*, 2020, 343 (5), pp.111781. 10.1016/j.disc.2019.111781 . hal-02935482

HAL Id: hal-02935482

<https://hal.science/hal-02935482>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

A note on the simultaneous edge coloring ^{*}

Nicolas Bousquet¹ and Bastien Durain²

¹Univ. Grenoble Alpes, CNRS, Laboratoire G-SCOP, Grenoble-INP, Grenoble, France.†

²ENS Lyon, Département d'informatique, France. ‡

December 2, 2019

Abstract

Let $G = (V, E)$ be a graph. A (*proper*) k -edge-coloring is a coloring of the edges of G such that any pair of edges sharing an endpoint receive distinct colors. A classical result of Vizing [3] ensures that any simple graph G admits a $(\Delta(G) + 1)$ -edge coloring where $\Delta(G)$ denotes the maximum degree of G . Recently, Cabello raised the following question: given two graphs G_1, G_2 of maximum degree Δ on the same set of vertices V , is it possible to edge-color their (edge) union with $\Delta + 2$ colors in such a way the restriction of G to respectively the edges of G_1 and the edges of G_2 are edge-colorings? More generally, given ℓ graphs, how many colors do we need to color their union in such a way the restriction of the coloring to each graph is proper?

In this short note, we prove that we can always color the union of the graphs G_1, \dots, G_ℓ of maximum degree Δ with $\Omega(\sqrt{\ell} \cdot \Delta)$ colors and that there exist graphs for which this bound is tight up to a constant multiplicative factor. Moreover, for two graphs, we prove that at most $\frac{3}{2}\Delta + 4$ colors are enough which is, as far as we know, the best known upper bound.

1 Introduction

All along the paper, we only consider simple loopless graphs. In his seminal paper, Vizing proved in [3] that any simple graph G can be properly edge-colored using $\Delta(G) + 1$ colors (where $\Delta(G)$ denotes the maximum degree of G). The *union* of two graphs G_1 and G_2 on vertex set V is the (simple) graph G with vertex set V and where uv is an edge if and only if uv is an edge of G_1 or an edge of G_2 . An edge coloring of G is *simultaneous with respect to G_1 and G_2* if its restrictions to the edge set of G_1 and to the edge set of G_2 are proper edge-colorings. Recently, Cabello raised the following question¹: given two graphs G_1, G_2 of maximum degree Δ on the same set of vertices V , does it always exist a simultaneous $(\Delta + 2)$ -edge coloring with respect to G_1 and G_2 ? Cabello proved that this property is satisfied if the intersection of G_1 and G_2 is regular [1]. Using Vizing's theorem, one can easily notice that there exists a simultaneous $(2\Delta + 1)$ -edge coloring. From a lower bound perspective, no graph where $\Delta + 2$ colors are needed is known.

Cabello introduced a generalization of this notion. Let ℓ graphs G_1, G_2, \dots, G_ℓ and G be their (edge) union. In other words, uv is an edge of G if and only if uv is an edge of at least one graph G_i with $i \leq \ell$. An edge-coloring of G is *simultaneous with respect to G_1, \dots, G_ℓ* if its restriction to each graph G_i is a proper edge-coloring. Cabello asked how many colors are needed to ensure the existence of a simultaneous coloring of G with respect to each G_i . Let us denote by $\chi'(G_1, \dots, G_\ell)$ the minimum number of colors needed to obtain a simultaneous coloring. And let $\chi'(\ell, \Delta)$ be the largest integer k such that $k = \chi'(G_1, \dots, G_\ell)$ for some graphs G_1, \dots, G_ℓ of maximum degree (at most) Δ . Vizing's theorem ensures that $\chi'(\ell, \Delta) \leq \ell\Delta + 1$ and Cabello exhibit a graph for which $\chi'(3, \Delta) \geq \Delta + 5$ (with $\Delta = 10$) [1]. In this note, we prove that the order of magnitude of $\chi'(\ell, \Delta)$ is $\Theta(\sqrt{\ell}\Delta)$. More precisely, we prove that the following statement holds:

^{*}This work was supported by the ANR Project GrR (ANR-18-CE40-0032).

[†]nicolas.bousquet@grenoble-inp.fr

[‡]bastien.durain@ens-lyon.fr

¹BIRS Workshop: Geometric and Structural Graph Theory (2017). See e.g. <https://sites.google.com/site/sophiespirkl/birs-open-problems.pdf>

Theorem 1

$$\chi'(\ell, \Delta) \leq 2\sqrt{2\ell}\Delta - \sqrt{2\ell} + 2.$$

We claim that this bound is tight up to a constant multiplicative factor. Let $\ell \in \mathbb{N}$ and Δ be an even value. Let $G := S_{1, k\Delta}$ be the star with $k\Delta$ leaves, where $k = \lfloor \sqrt{\frac{\ell}{2}} \rfloor$. Partition the edges of G into $2k$ sets A_1, \dots, A_{2k} of size $\frac{\Delta}{2}$. For every pair i, j , create the graph $G_{i,j}$ with edge set $A_i \cup A_j$. Note that each graph $G_{i,j}$ has maximum degree Δ since by construction the set of edges A_i induces a graph of maximum degree $\Delta/2$. Moreover the total number of graphs $G_{i,j}$ is $2k(2k-1)/2 \leq \ell$. Finally, by construction, every pair of edges of G appears in at least one graph $G_{i,j}$. So in order to obtain a simultaneous coloring, we need to color all the edges of G with different colors (since all the edges are incident to the center of the star). So:

Proposition 2

$$\chi'(\ell, \Delta) \geq \left\lfloor \sqrt{\frac{\ell}{2}} \right\rfloor \Delta.$$

Note that, for $\ell = 3$ and the graph $S_{1, 3\lfloor \Delta/2 \rfloor}$, a similar construction ensures that $\chi'(3, \Delta) \geq 3\lfloor \frac{\Delta}{2} \rfloor$, improving the lower bound of $\Delta + 5$. Indeed, let us partition the edges of the star into three sets A_1, A_2, A_3 of size $\lfloor \Delta/2 \rfloor$. We similarly define for every $i \neq j$ the graph $G_{i,j}$ with edge set $A_i \cup A_j$. Each graph $G_{i,j}$ has maximum degree Δ and every pair of edges appears in at least one graph $G_{i,j}$. So an edge coloring of $S_{1, 3\lfloor \Delta/2 \rfloor}$ simultaneous with respect to $G_{1,2}, G_{1,3}$ and $G_{2,3}$ is a proper edge coloring of $S_{1, 3\lfloor \Delta/2 \rfloor}$.

When $\ell = 2$, a careful reading of the proof of Theorem 1 permits to remark that we can improve the trivial $(2\Delta + 1)$ upper bound into 2Δ . We prove the following much better upper bound with a different technique:

Theorem 3

$$\chi'(2, \Delta) \leq \left\lfloor \frac{3}{2}\Delta + 4 \right\rfloor.$$

As far as we know, it is the best known upper bound.

2 Proof of Theorem 1

Let G_1, \dots, G_ℓ be ℓ graphs of maximum degree Δ . Let us partition the set of edges of $G = \cup_{i=1}^{\ell} G_i$ into two sets (all along the paper, the notation \cup stands for edge union). The *multiplicity* of an edge e is the number of graphs G_i with $i \leq \ell$ on which e appears. For some fixed k , the set E_1 is the set of edges with multiplicity at least k and E_2 is the set of edges with multiplicity less than k . We will optimize the value of k later. (Note that we do not necessarily assume that k is an integer). For every $i \in \{1, 2\}$, let us denote by H_i the graph G restricted to the edges of E_i . Note that $G = H_1 \cup H_2$.

We claim that the graph H_1 has degree at most $\ell\Delta/k$. Indeed, let u be a vertex and $E_1(u)$ be the set of edges of H_1 incident to it. Since every edge of H_1 has multiplicity at least k and at most $\ell\Delta$ edges (with multiplicity) are incident to u in G , at most $\frac{\ell}{k}\Delta$ different edges are in $E_1(u)$. So H_1 has maximum degree $\frac{\ell}{k}\Delta$. By Vizing's theorem, H_1 can be properly edge-colored with $(\frac{\ell}{k}\Delta + 1)$ colors.

Let us now prove that H_2 can be simultaneously edge-colored with $2k(\Delta - 1) + 1$ colors. Let us prove it by induction on the number of edges of H_2 . The empty graph can indeed be edge-colored with $2k(\Delta - 1) + 1$ colors. Let $e = uv$ be an edge of H_2 . By induction, there exists a simultaneous coloring c' of $H_2 \setminus e$ with $2k(\Delta - 1) + 1$ colors. Let us prove that c' can be extended into a simultaneous coloring of H_2 . Without loss of generality, we can assume that e is an edge of the graphs G_1, \dots, G_r with $r < k$ and is not an edge of G_{r+1}, \dots, G_ℓ . Let F be the set of edges of G_1, \dots, G_r incident to u or to v distinct from e . By assumption, there are at most $2r(\Delta - 1)$ such edges ($2(\Delta - 1)$ in each graph). Since $r < k$, at most $2k(\Delta - 1)$ edges are in F . So there exists a color a that does not appear in F . The edge e can be colored with a without violating any constraints. It holds by choice of a for G_i with $i \leq r$ and it holds since $e \notin G_i$ for $i > r$.

So $\chi'(\ell, \Delta) \leq \frac{\ell}{k}\Delta + 2k(\Delta - 1) + 2$ colors. We finally optimize the integer k which minimize the number of colors. We want to minimize $\frac{\ell}{k} + 2k$ which is minimal when $k = \sqrt{\frac{\ell}{2}}$. It finally ensures that $\chi'(\ell, \Delta) \leq 2\sqrt{2\ell}\Delta - \sqrt{2\ell} + 2$, which completes the proof of Theorem 1.

3 Proof of Theorem 3

Let G_1, G_2 be two graphs of maximum degree Δ and let G be their union. Let E_2 be the edges that appear in both graphs and, for every $i \in \{1, 2\}$, let E_1^i be the set of edges that appears only in G_i . Let us denote by H_2 (resp. H_1^i) the graph restricted to the edges of E_2 (resp. E_1^i).

For every vertex v and every graph H , we denote by $\deg_H(v)$ the degree of v in H . Let H be a graph and f, g be two functions from $V(H)$ to \mathbb{R}^+ . A (g, f) -factor of H is an edge-subgraph H' of H such that every vertex v satisfies $g(v) \leq \deg_{H'}(v) \leq f(v)$. Kano and Saito proved in [2] that the graph H admits a (g, f) -factor if

- (i) f and g are two integer valued functions, and
- (ii) for every vertex v , $g(v) < f(v)$, and
- (iii) there exists a real number θ such that $0 \leq \theta \leq 1$ and for every vertex v , $g(v) \leq \theta \cdot \deg_H(v) \leq f(v)$.

Let $1 \leq i \leq 2$. We will extract from H_1^i a (g, f) -factor where $g(v) = \lceil \frac{\deg_{H_1^i}(v)}{2} - 1 \rceil$ and $f(v) = \lceil \frac{\deg_{H_1^i}(v)}{2} \rceil$. The points (i) and (ii) are satisfied. Moreover, by choosing $\theta = \frac{1}{2}$, (iii) is also satisfied. Thus by [2], the graphs H_1^1 and H_1^2 admit (g, f) -factors. For $i \leq 2$, let K_1^i be a (g, f) -factor of H_1^i . For every i , let $L_1^i = H_1^i \setminus K_1^i$. Let $L = L_1^1 \cup L_1^2$ and $R = H_2 \cup K_1^1 \cup K_1^2$. Note that $G = L \cup R$. Let us now color these two graphs.

Let us first prove that L can be colored with $\lfloor \frac{\Delta}{2} \rfloor + 2$ colors. For every i , the graph L_1^i has maximum degree at most $\lfloor (\Delta/2) + 1 \rfloor$ since every vertex v of K_1^i has degree at least $\lceil (\deg_{H_1^i}(v)/2) - 1 \rceil$. By Vizing's theorem, the graph L_1^i can be colored with at most $\lfloor \frac{\Delta}{2} + 1 \rfloor + 1 = \lfloor \frac{\Delta}{2} \rfloor + 2$ colors. Since the edges of L_1^1 and L_1^2 are disjoint, $L = L_1^1 \cup L_1^2$ can be simultaneously colored with $\lfloor \frac{\Delta}{2} \rfloor + 2$ colors (the same set of colors can be re-used for each graph).

Let us now color the graph R . Let v be a vertex of R . Let us denote by d the degree of v in H_2 . Since edges of H_2 are in both G_1 and G_2 , the vertex v has degree at most $\Delta - d$ in both graphs H_1^1 and H_1^2 . Since the graphs K_1^1 and K_1^2 are (g, f) -factors of respectively H_1^1 and H_1^2 , the degree of the vertex v is at most $\lceil \frac{\Delta-d}{2} \rceil$ in each graph. So the degree of v in the graph R is at most $d + 2\lceil \frac{\Delta-d}{2} \rceil \leq \Delta + 1$. By Vizing's theorem, the graph R can be colored using at most $\Delta + 2$ colors.

Since $G = L \cup R$, we can find a simultaneous edge-coloring with respect to G_1 and G_2 using at most $\lfloor \frac{\Delta}{2} \rfloor + 2 + \Delta + 2 = \lfloor \frac{3}{2}\Delta \rfloor + 4$ colors.

4 Conclusion

Theorem 1 and Proposition 2 ensures that the following holds:

$$\left\lfloor \sqrt{\frac{\ell}{2}} \right\rfloor \Delta \leq \chi'(\ell, \Delta) \leq 2\sqrt{2\ell}\Delta - \sqrt{2\ell} + 2.$$

Closing the multiplicative gap of 4 between lower and upper bound is an interesting open problem. For $\ell = 2$, we still do not know any graph for which $\chi'(2, \Delta) > \Delta + 1$. Cabello asked the following question that is still widely open despite the progress obtained in Theorem 3:

Question 4 (Cabello) *Is it true that*

$$\chi'(2, \Delta) \leq \Delta + 2 \quad ?$$

Acknowledgements. The authors want to thank Louis Esperet for fruitful discussions and suggestions.

References

- [1] S. Cabello. *private communication*, 2018.
- [2] M. Kano and A. Saito. $[a, b]$ -factors of graphs. *Discrete Mathematics*, 47:113–116, 1983.
- [3] V. Vizing. On an estimate of the chromatic class of a p -graph. *Diskret. Analiz*, 3(7):25–30, 1964.