



HAL
open science

Modeling chlordecone toxicokinetics data in growing pigs using a nonlinear mixed-effects approach

A. Fourcot, C. Feidt, Alain Bousquet-mélou, A.A. Ferran, Jean-Luc Gourdine, M. Bructer, C. Joaquim-Justo, G. Rychen, A. Fournier

► To cite this version:

A. Fourcot, C. Feidt, Alain Bousquet-mélou, A.A. Ferran, Jean-Luc Gourdine, et al.. Modeling chlordecone toxicokinetics data in growing pigs using a nonlinear mixed-effects approach. *Chemosphere*, 2020, 250, pp.126151. 10.1016/j.chemosphere.2020.126151 . hal-02935835

HAL Id: hal-02935835

<https://hal.science/hal-02935835>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Modeling chlordecone toxicokinetics data in growing pigs using a nonlinear mixed-
2 effects approach

3 A. Fourcot^a, C. Feidt^a, A. Bousquet-Mélou^b, A. Ferran^b, J.-L. Gourdine^c, M. Bructer^d,
4 C. Joaquim-Justo^e, G. Rychen^a, A. Fournier^{a,*}

5 ^a Université de Lorraine, INRAE, USC 340, UR AFPA, 2 Avenue de la Forêt de Haye, TSA 40602, 54518
6 Vandœuvre-lès-Nancy, France

7 ^b INTHERES, Université de Toulouse, INRAE, ENVT, 31000 Toulouse, France

8 ^c Unité de Recherches Zootechniques, INRAE, URZ UR143, Domaine Duclos, 97170 Petit-Bourg,
9 France

10 ^d Plateforme Tropicale d'Expérimentation sur l'Animal, INRAE, PTEA UE1294, 97170 Petit-Bourg,
11 France

12 ^e Université de Liège, LEAE-CART, Freshwater and Oceanic sciences Unit of reSearch (FOCUS), B6C,
13 Allée du 6 Août, 11, 4000 Liège, Belgium

14 * Corresponding author. E-mail address: agnes.fournier@univ-lorraine.fr

15

16 Abstract

17 The use of chlordecone (CLD), a chlorinated polycyclic pesticide used in the French Indies
18 banana fields between 1972 and 1993, resulted in a long-term pollution of agricultural areas.
19 It has been observed that this persistent organic pollutant (POP) can transfer from
20 contaminated soils to food chain. Indeed, CLD is considered almost fully absorbed after
21 involuntary ingestion of contaminated soil by outdoor reared animals. The aim of this study
22 was to model toxicokinetics (TKs) of CLD in growing pigs using both non-compartmental and
23 nonlinear mixed-effects approaches (NLME). In this study, CLD dissolved in cremophor was
24 intravenously administrated to 7 Creole growing pigs and 7 Large White growing pigs (1 mg
25 kg⁻¹ body weight). Blood samples were collected from time t=0 to time t=84 days. CLD
26 concentrations in serum were measured by GCMS/MS. Data obtained were modeled using
27 Monolix (2019R). Results demonstrated that a bicompartmental model best described CLD
28 kinetics in serum. The influence of covariates (breed, initial weight and average daily gain) was

29 simultaneously evaluated and showed that average daily gain is the main covariate explaining
30 inter-individual TKs parameters variability. Body clearance was of $76.7 \text{ mL kg}^{-1} \text{ d}^{-1}$ and
31 distribution volume at equilibrium was of 6 L kg^{-1} . This modeling approach constitutes the first
32 application of NLME to study CLD TKs in farm animals and will be further used for rearing
33 management practices in contaminated areas.

34 Keywords: Chlordecone, Toxicokinetics, Pigs, Modeling, Monolix

35 1. Introduction

36 Chlordecone (CLD) is a chlorinated polycyclic ketone pesticide which was extensively used in
37 the French West Indies (FWI) from 1972 until 1993 to control the banana black weevil
38 (*Cosmopolites sordidus*) population (Cabidoche et al., 2009). The use of this insecticide has
39 resulted in long-term soils pollution, this compound transfers from soil to food chain and local
40 population is thus exposed to CLD through food (Dubuisson et al., 2007). Human exposure to
41 CLD is therefore a public health concern since CLD is known for its neurotoxic and
42 carcinogenic properties (Multigner et al., 2010; Costet et al., 2015). To limit human exposure
43 to CLD related to self-produced or informal sector products (ANSES, 2017), risk of food
44 contamination has to be reduced. Meat from local outdoor reared pigs can contribute to CLD
45 human exposure. Thus, CLD distribution and elimination processes need to be characterized
46 in pigs to further propose rearing strategies aiming at removing CLD from contaminated
47 animals. Furthermore, a study was conducted in 2013-2014 on 742 FWI inhabitants to assess
48 CLD levels in serum (KANARRI study). Although CLD was detected in serum of 90% of the
49 participants, impregnation value could not be used to assess health risk due to lack of an
50 internal health-based guidance value (Dereumeaux et al., 2019). The currently existing chronic
51 toxicological reference value (TRV) is an external exposure value fixed at $0.5 \mu\text{g kg}^{-1} \text{ body}$
52 $\text{weight (BW) d}^{-1}$ (AFSSA, 2007). CLD toxicokinetics (TKs) in pigs could also be used to build a
53 PBPK model in human because pig is an animal model close to the human model. This model
54 would then allow to convert the external TRV in an internal reference dose and then to discuss
55 the impregnation level of population.

56 The main route of animal exposure to CLD is related to soil ingestion. Level of soil ingestion is
57 known in ruminants (Jurjanz et al., 2017; Collas et al., 2019). In pigs reared outdoor, level of
58 soil ingestion has been studied in temperate conditions (Jurjanz and Roinsard, 2014) but not
59 yet in tropical conditions. However, because of their exploratory behavior, risk of pig meat
60 contamination must be even higher than for ruminant meat. It has been demonstrated that soil
61 bound CLD is not retained in soil matrices during digestive processes (Bouveret et al., 2013;
62 Jurjanz et al., 2014) and CLD is considered almost fully absorbed after oral administration
63 (Fournier et al., 2017). Unlike other POPs, CLD has a peculiar distribution with a preferential
64 accumulation in liver and a lower one in fat tissues (Soine et al., 1983; Jondreville et al., 2014;
65 Lastel et al., 2016). Thus, because of different distribution profiles within the organism (Soine
66 et al., 1982; Soine et al., 1983; Lerch et al., 2016), TK extrapolations from other POPs to CLD
67 are not appropriate. Limited information is currently available concerning TKs of CLD in human
68 or in animals. Some data are available in rodents (Egle et al., 1978; Houston et al., 1981),
69 piglets (Soine et al., 1983), ruminants (Fournier et al., 2017; Saint-Hilaire et al., 2019) and
70 humans (Adir et al., 1978; Cohn et al., 1978; Fariss et al., 1980). Extrapolations from rodents
71 or piglets to growing pigs are not appropriate because of metabolic differences between
72 animals of different species or different physiological status (Houston et al., 1981). Therefore,
73 it appears as a major priority to characterize CLD behavior in growing pigs. In this study, the
74 two main breeds of pigs reared in the FWI will be compared.

75 Until now, TKs of CLD in animals have been studied by compartmental approaches that did
76 not consider parameters like breed or body weight (Fournier et al., 2017; Saint-Hilaire et al.,
77 2019). The population approach, initially developed for pharmacokinetics applications, was
78 used to evaluate pharmacokinetics variability between individuals. Nonlinear mixed-effects
79 (NLME) have not been used yet for the modeling of TKs CLD data. Advantages of such an
80 approach are that parameters are estimated in the population of individuals, variability within
81 the population is evaluated and factors that control that variability may be identified.

82 The aim of this study was to model TKs of CLD in growing pigs using both a non-compartmental
 83 approach and a NLME approach. Such data are essential to better characterize elimination
 84 mechanisms of CLD in pigs.

85 2. Material and methods

86 2.1. Animals and management

87 This study was performed at the experimental facilities of INRAE-PTEA in Guadeloupe (FWI;
 88 16°N, 61°W). The experimental protocol was approved by the Ethics Committee of Guyana
 89 and West Indies (CEMEA-AG) under the project number APAFIS#6070-2016070721289156.

90 A total of 14 growing pigs born in the herd of the INRAE experimental station of Duclos (Petit-
 91 Bourg, Guadeloupe) were placed in individual barns. To compare TK parameters between the
 92 two main pig breeds reared in the FWI, two different breeds were studied: Large White (n=7)
 93 and Creole (n=7). Large White breed is one of the mainstream pig breed used worldwide that
 94 has been selected for high growth performance in optimal conditions (ie specialist type of
 95 breed). On the other hand, Creole breed is a generalist type of breed that has not been
 96 genetically selected and this local tropical Caribbean breed is mainly characterized by early
 97 maturity, a higher fat deposition than Large White breed and a better ability to cope with harsh
 98 tropical conditions (heat, nutritional stress; Gourdine et al., 2018). Physiological and growth
 99 performance differences between the two groups may affect TKs of CLD.

100 Table 1: Average values of IBW, FBW and ADG for each breed

Parameters	Creole pigs (n=7) mean ± SE	Large White pigs (n=7) mean ± SE	All pigs (n=14) mean ± SE
IBW kg	68.8 ± 1.8 ^a	67.9 ± 3.1 ^a	68.3 ± 1.7
FBW kg	91.2 ± 2.8 ^a	95.6 ± 2.4 ^a	92.8 ± 1.9
ADG g d ⁻¹	248 ± 25 ^a	327 ± 13 ^a	288 ± 17

IBW: Initial Body Weight; FBW: Final Body Weight; ADG: Average Daily Gain; SE: Standard Error.
 Values not followed by the same superscript letter significantly differ (P < 0.05; Student test)

101

102 Table 1 gives average values of initial body weight (IBW), final body weight (FBW) and average
103 daily gain (ADG) for each breed. At the beginning of the experiment, pigs were 26 and 23
104 weeks old for Creole and Large White, respectively, to obtain comparable animals in term of
105 IBW. In order to limit ADG of growing pigs and to limit dilution effect, a daily feed ration
106 composed by a commercial feed was established to obtain an ADG of 250 g d⁻¹. Water was
107 delivered *ad libitum*. All pigs were individually weighted every 7 days during the 84-days
108 depuration period. ADG was calculated by the equation (1):

$$109 \text{ ADG} = (\text{FBW} - \text{IBW}) / \text{number of days} \quad (1)$$

110 To compare IBW, FBW and ADG between the two breeds, a Student *t* test was used. Before
111 apply this parametric test, normality of data and equality of variances has been checked with
112 a Shapiro-test and a F-test, respectively. Statistical analysis were performed using R version
113 3.5.0. software. The α level for statistical significance was set at 0.05.

114 2.2. Experimental design

115 One week before the experiment, a catheter was inserted in the jugular vein of each pig to
116 allow CLD administration and blood sampling. After this surgical intervention, pigs were
117 allowed a 7-days adaptation period in individual barns (metal-slatted pen of 0.85 x 1.50 m).
118 The experiment consisted in a single administration of CLD by intravenous (*i.v.*) route (1 mg
119 kg⁻¹ BW, Kepone 99.9%, Sigma Aldrich). This high dose was chosen to be able to follow the
120 decrease of CLD concentrations in serum and easily compare to other species. Indeed, this
121 dose has already be used in goats and ewes (Fournier et al., 2017; Saint-Hilaire et al., 2019).
122 For these studies, no side effects were observed.

123 CLD was dissolved in cremophor (polyethoxylated castor oil, Kolliphor[®], Sigma-Aldrich) at
124 room temperature by sonication (2 hours). Pigs received 0.04 g kg⁻¹ BW of contaminated oil
125 concentrated at 24.8 mg CLD g⁻¹ oil. The catheter was flushed several times after CLD
126 administration to avoid contamination of following blood samples.

127 At 0, 30, 60 min, 2, 3, 7, 12, 24 hours, 3, 8, 14, 21, 28, 49, 70 and 84 days after administration,
128 blood samples (4 x 10 mL) were taken using the *i.v.* catheter for the first weeks (in dry tubes)
129 and then, by a direct venipuncture on the jugular vein (BD vacutainer®, ref 368815), when the
130 sampling was less intensive. Collected blood samples were allowed clotting for 2 hours at room
131 temperature and at 4°C during 24 hours before centrifugation at 3000 rpm for 10 min to get
132 serum samples which were stored at -20°C until analysis. After 84-depuration days, pigs were
133 slaughtered (stunning followed by bleeding). Liver, peri-renal fat and shaft (diaphragm muscle)
134 were collected. Tissues were stored at -20°C until analysis.

135 2.3. Analytical methods

136 2.3.1. Analysis of CLD in serum

137 CLD analysis was performed by the Center for Analytical Research and Technology (LEAE-
138 CART, University of Liège, Belgium) according to Fournier et al. (2017) with slight
139 modifications. Briefly, 10-monohydrochlordecone kindly synthesized by Dr. P.L. Saaidi
140 (Génomique Métabolique, Evry, France) according to Chevallier et al. (2019) was used as
141 internal standard (recovery surrogate). This compound was never detected in previous studies
142 for CLD analysis in goat, lamb and ewe serum (Jurjanz et al., 2014; Lastel et al., 2016; Lerch
143 et al., 2016; Fournier et al., 2017). This internal standard was added to 2 mL of pig serum
144 samples. After proteins denaturation with trimethylamine and formic acid, CLD was extracted
145 with n-hexane/diethylether/acetonitrile/ethanol (80/15/4/1, v/v) using a solid-phase extraction
146 (SPE) on Supelclean Envi-C₁₈ microcartridges (Supelco, Bellafonte, PA, USA) and subjected
147 to acidic purification using concentrated sulfuric acid 98% w/v. The final extract (100 µL of
148 hexanic solution) was spiked with PCB-209 (100 pg µL⁻¹ in hexane) used as a volume internal
149 standard. As described in Fournier et al. (2017), all PCBs and organochlorine pesticides were
150 eluted in the first hexanic solution from C₁₈ microcartridge. CLD and related compounds (5b
151 monohydroCLD and 10-monohydroCLD) were recovered in the second elution
152 (hexane/diethylether; 85/15; v/v). Then, there was no PCBs in the CLD elution. Furthermore,
153 PCB-209 was not present in commercial PCB mixtures such as Aroclor used in the past.

154 However, this congener was reported in organic paint pigment (Hu and Hornbuckle, 2010;
155 Anezaki and Nakano, 2014). Moreover, Koh et al. (2015) reported traces of PCB-209 in human
156 serum in some areas in USA but at trace levels and concentrations measured (0.00 to 0.01 ng
157 mL⁻¹) were far below the LOQ of the present study (0.06 ng mL⁻¹). As a consequence, when
158 considering all these elements and as PCB-209 was added as internal volume standard just
159 prior the injection in HRGC MS/MS, this compound cannot come from environmental
160 contamination. For each series of 21 experimental samples there was a procedural blank, a
161 matrix blank (porcine serum from Sigma Aldrich chemie Gmbh, Steinheim, Germany) and a
162 quality control (QC) sample composed of porcine serum (Sigma Aldrich chemie Gmbh,
163 Steinheim, Germany) spiked with 0.1 or 5 or 10 ng CLD mL⁻¹. These different QC were
164 distributed among the series in a random manner. The CLD limit of detection was 0.02 ng mL⁻¹
165 in the serum with a limit of quantification (LOQ) at 0.06 ng mL⁻¹. The mean \pm standard
166 deviation CLD recoveries in the spiked porcine serum used as QCs were 87.0% \pm 2.3%, 99.0%
167 \pm 4.2% and 97.4% \pm 4.3% for 0.1 ng mL⁻¹, 5 ng mL⁻¹ and 10 ng mL⁻¹ QCs respectively.
168 Recovery rates were always between 60 and 140% according to the requirement of SANCO
169 (2014). The measurement uncertainty has been assessed from spiked CLD human serum
170 according to Eppe et al. (2017). The uncertainty was 4.66% for 0.05 ng mL⁻¹ CLD human
171 spiked serum, 5.82% for 2.5 ng mL⁻¹ CLD human spiked serum and 6.34% for 10 ng mL⁻¹ CLD
172 human spiked serum. The mean uncertainty was 5.61% \pm 0.7%.

173 The extracts were analyzed by High Resolution Gas Chromatophy coupled to an ion trap mass
174 spectrometer to confirm the identity and concentrations of CLD (Trace GC Ultra and ITQ 1100
175 from ThermoQuest, ThermoScientific, Waltham, MA, USA). GC-MS was equipped with a J&W
176 Scientific DB-XLB capillary column (30 m \times 0.25 mm i.d., 0.25 μ m film) purchased from Agilent
177 (Santa Clara, CA, USA), using helium as a carrier gas. Chromatographic conditions were
178 described elsewhere (Fournier et al., 2017). The extracts were injected into the column using
179 a split/splitless injector. The transfer line temperature was kept at 290°C and the ion trap

180 temperature was set at 250°C. The electron ionisation was performed at 70 eV and the ion
181 trap was operating in MS/MS mode. Compounds identification was performed with MS/MS
182 transitions of 272>237, 238>203 and 498>428 for CLD, 10-monohydrochlordecone and PCB-
183 209 respectively. The collision energy was 2.5 eV, for all the compounds. Concentrations were
184 measured by quantifying m/z 238 ion for 10_monohydrochlordecone, m/z 272 ion for
185 chlordecone and m/z 498 ion for PCB-209.

186 2.3.2. Analysis of CLD in tissues

187 CLD concentrations in liver, shaft and peri-renal fat were determined by Liquid
188 Chromatography coupled with a tandem Mass Spectrometry (LC-MS/MS) as described in
189 Dromard et al. (2017), using the official method of the French National Reference Laboratory
190 (ANSES), in LABOCEA laboratory (Quimper, France), labeled by the French Accreditation
191 Committee (COFRAC). Briefly, CLD C13 was used as internal standard and add to all samples.
192 CLD was extracted from tissues with a solution of solvents: acetonitrile/dichloromethane 75/25
193 (v/v) and hexane/acetone 75/25 (v/v) for adipose tissue and hexane/acetone 85/15 (v/v) for
194 liver and muscle. CLD was transformed into chlordecone hydrate with soda and the aqueous
195 phase was rinsed with hexane. CLD was then subjected to acidic purification using
196 concentrated sulfuric acid 60% and extracted with a solution of hexane/acetone 85/15 (v/v).
197 Concentrations of CLD in tissues were expressed in $\mu\text{g CLD kg}^{-1}$ fresh weight (FW). The LOQ
198 was $1.0 \mu\text{g CLD kg}^{-1}$ in muscle and liver tissues and $3.0 \mu\text{g CLD kg}^{-1}$ in adipose tissues.

199 2.4. Estimations of CLD amounts in tissues

200 CLD concentrations at slaughtering obtained in tissues were used to estimate CLD amounts
201 in liver, muscle and fat tissues. The liver was weighed after slaughtering. The weight of the
202 total muscle mass was estimated considering that muscle percentages of BW in Creole and
203 Large White pigs are 43.3% and 54.8%, respectively (Renaudeau et al., 2005). These
204 percentages were multiplied by the BW at slaughter to obtain weight of total muscle mass in

205 the organism. Weight of total fat mass was estimated in the same way considering that fat
206 percentages of BW in Creole and Large White pigs are 34.7% and 22.4%, respectively
207 (Renaudeau et al., 2005). Total amounts of CLD in matrices were obtained multiplying the total
208 weight of each matrix by corresponding concentrations at slaughtering.

209 2.5. Non-compartmental analysis

210 Non-compartmental analysis describes the evolution of CLD concentrations without any
211 assumptions. This analysis is mainly used to obtain basic TK parameters. The non-
212 compartmental analysis was performed using the PKanalix software (Lixoft, version 2019R1).

213 To compare TK parameters between different species, a non-compartmental analysis was
214 performed on CLD kinetic data in pigs and with data previously obtained on goats and ewes
215 (Fournier et al., 2017; Saint-Hilaire et al., 2019). Non-lactating and non-pregnant adult ewes
216 (n=5) and goats (n=4) of 65.8 ± 2.2 and 33.0 ± 3.5 kg (mean \pm standard error (SE)) BW
217 respectively were used in these studies.

218 For each species, four TK parameters were calculated with the non-compartmental analysis:
219 total body clearance (CL), steady-state distribution volume (V_{ss}), mean residence time (MRT)
220 and area under the serum curve (AUC). Area under the serum curve from zero to infinity was
221 calculated by the linear log up log down method.

222 To compare TK parameters obtained in the three species, an analysis of variance (ANOVA)
223 test was performed and followed by a Tukey test (R version 3.5.0. software). The α level for
224 statistical significance was set at 0.05.

225 2.6. Population toxicokinetic model development

226 Population TK analysis aims to provide TK parameters on a set of individuals and to assess
227 the variability between individuals.

228 2.6.1. Methods and software

229 The population TK analysis was performed using a NLME approach with the Monolix software
230 (Lixoft, version 2019R1). Monolix estimates TK parameters using the stochastic approximation
231 expectation maximization (SAEM) algorithm.

232 2.6.2. Model building

233 **Individual modelisation**

234 Experimental data were best described by a linear two-compartmental model with a
235 proportional error. The two-compartmental model is represented in Figure 1. In this model, the
236 body is divided into central and peripheral compartments. The central compartment consists
237 mainly of the plasma and well perfused organs where the distribution of the molecule is almost
238 instantaneous. The peripheral compartment consists of less perfused organs, where the
239 molecule is distributed slower.

240 TK parameters were calculated for each pig individually with the two-compartmental model.
241 Parameters chosen in the structural model were body clearance (CL), distribution volume of
242 the central compartment (V1), intercompartmental clearance (Q) and distribution volume of the
243 peripheral compartment (V2). Clearances and volumes of distribution were normalized to BW
244 with the IBW. TK parameters (CL, V1, Q and V2) were assumed to follow a lognormal
245 distribution in the population of pigs.

246 Mean residence time (MRT) and elimination half-time life ($t_{1/2elim}$) were calculated with individual
247 parameters estimates by Eq. (2) and (3):

$$248 \text{ MRT} = V_{ss} / \text{CL} \quad (2)$$

249 With $V_{ss} = V1 + V2$

$$250 t_{1/2elim} = \ln(2) / \beta \quad (3)$$

$$251 \text{ With } \beta = \frac{1}{2} ((k_{12} + k_{21} + k_{10}) - \sqrt{(k_{12} + k_{21} + k_{10})^2 - 4 \cdot k_{21} \cdot k_{10}})$$

252 **Population modelisation: full model and covariate analysis**

253 CLD concentration C_{ij} in an individual i at time t_{ij} can be expressed as a function of the dose,
 254 the time t_{ij} and the model individual parameters (CL_i , $V1_i$, Q_i and $V2_i$). The full model can be
 255 written in the general form:

256 $C_{ij} = F(\Phi_i, t_{ij}) \times (1 + \varepsilon_{ij}), j = 1, \dots, n_i$

257
$$\Phi_i = \begin{pmatrix} CL_i \\ V1_i \\ Q_i \\ V2_i \end{pmatrix} = \begin{pmatrix} \mu_{CL} \cdot e^{\eta_{CL,i}} \\ \mu_{V1} \cdot e^{\eta_{V1,i}} \\ \mu_Q \cdot e^{\eta_{Q,i}} \\ \mu_{V2} \cdot e^{\eta_{V2,i}} \end{pmatrix}$$

258 Where C_{ij} is the observed CLD serum concentration measured on the individual i at time t_{ij} , F
 259 is the function describing a two compartment model (bi-exponential), Φ_i is the vector of
 260 individual parameters, $F(\Phi_i, t_{ij})$ is the value of the predicted CLD serum concentration at time
 261 t_{ij} for an individual with parameters Φ_i , and ε_{ij} is independent random variable normally
 262 distributed around zero with a variance of one. $F(\Phi_i, t_{ij})$ is the structural model and ε_{ij} is
 263 describing the residual error model. ε_{ij} measures the difference between the predicted value
 264 and the observed value of C_{ij} and was assumed to describe a proportional residual error model.
 265 μ is the population median of a model parameter. The variability sources (e^η) between the
 266 individual parameters Φ_i can be explained by covariates. Each parameter of Φ_i is assumed to
 267 be log-normally distributed.

268 Once the structural model was built, covariates were added. One factor (genotype with two
 269 categories: Large White vs. Creole) and two continuous variables (IBW and ADG) were tested
 270 as potential covariates on the TK parameters. First, variables were added one by one on
 271 individual CL and V2 estimations and were selected if their addition was able to cause a drop
 272 of the log-likelihood estimate (LL). Bayesian Information Criteria (BIC) was calculated and
 273 models with lower values of BIC were considered as better fitted to the observed data. P-value
 274 < 0.05 after a Wald test implemented in Monolix was considered statistically significant to
 275 include the covariate in the final model. Two covariates were eliminated from the model in a
 276 step-by-step manner, only ADG was statistically significant and added as covariable to obtain
 277 the final full model.

278 2.6.3. Model evaluation

279 Evaluation of goodness-of-fit was based on the following plots: (i) individual (IPRED) and
280 population predicted (PRED) concentrations versus observed concentrations (DV), (ii)
281 individual plots (PRED and OBS vs. time), (iii) population weighted (PWRES) and individual
282 weighted (IWRES) residuals versus time and (iv) normalized prediction distribution errors
283 (NPDE) versus time, IPRED and PRED (NPDE has to be normally distributed with a mean of
284 0 and a standard deviation of 1). Normality of residuals were evaluated using density plots of
285 weighted residuals.

286 2.6.4. Comparison of toxicokinetic parameters between the two breeds

287 To compare TK parameters obtained by the NLME model, a Student *t* test was used (R version
288 3.5.0. software). The α level for statistical significance was set at 0.05.

289 3. Results and discussion

290 3.1. Global kinetic analysis

291 Figure 2 shows CLD TKs in pig's serum over the 84-depuration days following the single *i.v.*
292 administration of 1 mg CLD kg⁻¹ BW. The first part of the curve (until 24 hours) corresponds
293 mainly to a distribution phase. During the first day following *i.v.* administration of CLD, serum
294 concentrations declined rapidly compared to the remaining kinetics, because of the partitioning
295 of the molecule in the body (Toutain and Bousquet-Melou, 2004). The second part of the curve
296 corresponds mainly to an elimination phase.

297 3.2. Non-compartmental analysis

298 TK parameters obtained with the non-compartmental analysis for CLD in pigs, ewes and goats
299 are presented in Table 2. The dose of 1 mg CLD kg⁻¹ BW was already used in goats and ewes
300 (Fournier et al., 2017; Saint-Hilaire et al., 2019) and was chosen in this study to allow
301 comparisons of TK parameters between the three species. Body clearance is the total
302 clearance, it depends on the elimination of the parent compound and on CLD metabolism. In

303 pigs, the body clearance was $79.6 \pm 4.4 \text{ mL kg}^{-1} \text{ d}^{-1}$ (mean \pm SE). The body clearance in pigs
 304 is comparable with the clearance determined in goats but 1.6 higher than in ewes ($P < 0.05$).
 305 Higher CLD elimination capacities in pigs than ewes could be explained either by higher
 306 capabilities for CLD elimination in the parent form (via biliary excretion for example), or higher
 307 metabolic capabilities, or both. Indeed, the enzyme *chlordecone reductase* involved in the
 308 biotransformation of CLD is known to be present in pigs as in humans, gerbils, ewes, goats
 309 (Fariss et al., 1980; Houston et al., 1981; Soine et al., 1983; Molowa et al., 1986a; Molowa et
 310 al., 1986b; Saint-Hilaire, 2018). Further studies could be performed to compare enzymatic
 311 activity of *chlordecone reductase* between species.

312 Table 2: Comparison of CLD toxicokinetic parameters obtained with non-compartmental
 313 analysis between pigs, ewes and goats following a single intravenous administration of 1 mg
 314 CLD kg^{-1}

315

Parameters	Pigs (n=14)		Ewes (n=5)		Goats (n=4)	
	Mean \pm SE	CV (%)	Mean \pm SE	CV (%)	Mean \pm SE	CV (%)
CL $\text{mL kg}^{-1} \text{ d}^{-1}$	79.6 ± 4.4^a	5.5	51.0 ± 6.1^b	12.0	66.6 ± 16.2^{ab}	24.4
Vss mL kg^{-1}	$5,982 \pm 354^a$	5.9	$1,807 \pm 157^b$	8.7	$1,615 \pm 446^b$	27.6
MRT d	77 ± 5^a	6.6	36 ± 2^b	4.5	24 ± 1^b	3.9
AUC ng d mL^{-1}	$12,505 \pm 636^a$	5.1	$21,202 \pm 3,373^b$	15.9	$18,911 \pm 5,687^{ab}$	30.1

CL: Body clearance; Vss: Volume of distribution at the equilibrium; MRT: Mean Residence Time; AUC: Area Under the Curve;
 CV: coefficient of variation; SE: Standard Error.

Values not followed by the same superscript letter significantly differ ($P < 0.05$; Tukey test).

Toxicokinetic parameters calculated from data obtained in Saint-Hilaire et al. (2019) and Fournier et al. (2017) for ewes and goats, respectively.

316

317 Despite high body clearance values showing high elimination capacities, the mean residence
 318 time (MRT) in pigs is 2.1 and 3.2 times higher than those obtained in ewes and goats ($P <$
 319 0.05), respectively. The MRT represents the average time the molecule stays in the body. This
 320 parameter is a function of the Vss and CL. Because of the high distribution volume, the MRT
 321 is higher in pigs (MRT = 78 ± 3 days) compared to both other species (MRT = 36 and 24 days
 322 in ewes and goats, respectively; $P < 0.05$). CLD elimination in pigs is slower than in ewes and
 323 goats because the Vss is 3.3 and 3.7 times larger in pigs than in ewes and goats, respectively.
 324 This assumes that pigs may store more CLD into deeper peripheral compartments than ewes

325 and goats. Deeper peripheral compartments represent tissues from which the molecule
326 depletes slower than other tissues because of CLD bindings to lipid or protein components.

327 This non-compartmental analysis allowed comparing the main TK parameters between pigs,
328 ewes and goats. Coefficients of variation of estimated mean parameters are ranged from 5.1
329 to 6.6. TK parameters estimation would probably be improved by considering inter-individual
330 variability. Thus, a NLME approach in pigs was applied to estimate more precisely TK
331 parameters.

332 3.3. Population toxicokinetic analysis

333 **Model evaluation**

334 The two-compartmental model best described the observed data, as indicated by a relative
335 standard error < 20% for most of all the model parameters. Good correlations were found
336 between observed concentrations vs. population predicted (PRED) and individual predicted
337 (IPRED) concentrations. Plots of population weighted (PWRES) and individual weighted
338 residuals (IWRES) vs. time and IPRED show that for both types of residuals, points are
339 randomly distributed around the zero line and most points fall within ± 3 standard deviations.
340 Simulation plots of NPDE vs. time and IPRED do not reveal biases because points are
341 scattered randomly around the zero line. Thus, these data support the model stability.

342 TK parameters obtained with the NLME approach are comparable with results obtained with
343 the non-compartmental approach. However, TK parameters seem to be better estimated with
344 the NLME approach because coefficients of variation are lower (Table 2 and Table 3).

345 **CLD elimination**

346 Estimated parameters derived from the population TK model are summarized in Table 3. Body
347 clearance (CL) is the most important TK parameter; it expresses the overall ability of the blood
348 compartment to eliminate the molecule. For all pigs, the body clearance is 76.7 ± 3.2 mL kg⁻¹.
349 Average daily gain (ADG) has a very significant effect on CL ($P < 0.05$): the higher the ADG

350 is, the lower the CL is. Despite high CL, mean residence time and elimination half-life in pigs
 351 are quite long ($MRT = 78 \pm 3$ days and $t_{1/2elim} = 54.7 \pm 2.3$ days) when compared to other species
 352 due to the molecule distribution in organism.

353 Table 3: Population toxicokinetic model estimates for CLD following a single intravenous
 354 administration of 1 mg CLD kg^{-1} to 14 pigs
 355
 356

Parameters	Model estimate median \pm SE	P ¹	CV (%)
CL mL kg^{-1} d ⁻¹	76.7 \pm 3.2	0.0009	4.1
V1 mL kg^{-1}	604 \pm 195	-	32.4
Q mL kg^{-1} d ⁻¹	16,680 \pm 2,870	-	17.1
V2 mL kg^{-1}	4,840 \pm 262	0.0142	5.4
Vss mL kg^{-1}	6,002 \pm 280	-	4.6
MRT d	78 \pm 3	-	4.3
t_{1/2elim} d	54.7 \pm 2.3	-	4.2

CL: Body clearance; V1: Central compartment volume of distribution; Q: Intercompartmental clearance; V2: Peripheral compartment volume of distribution; Vss: Steady-state volume of distribution; MRT: Mean residence time; $t_{1/2elim}$: Elimination half-life; CV: Coefficient of variation; SE: Standard Error.

¹ P-value obtained after a Wald test, testing significant effect of Average Daily Gain (ADG) on CL and V2 parameters.

357

358 **CLD distribution in organism**

359 Intercompartmental clearance (Q) expresses ability of the peripheral compartment to mobilize
 360 the molecule. The high intercompartmental clearance (16.7 ± 2.9 L kg^{-1}) suggests that
 361 equilibrium between central compartment and peripheral compartment is quickly reached.
 362 These results are consistent with the distribution half-life obtained in ewes: in less than one
 363 hour and a half, the equilibrium between the two compartments is reached (Saint-Hilaire et al.,
 364 2019).

365 Peripheral volume of distribution (V2) is not a physiologic value but a proportionality factor
 366 between amount of CLD in the organism at a given time and blood concentration at that time
 367 (Toutain and Bousquet-Mélou, 2004). The peripheral distribution volume V2 largely exceeds

368 the central distribution volume V_1 . The steady-state volume of distribution (V_{ss}) is calculated
369 by adding up volumes of central compartment (V_1) and peripheral compartment (V_2). In this
370 study, the V_{ss} ($V_{ss} = 6,002 \pm 280 \text{ mL kg}^{-1}$) largely exceeds the total volume of body water
371 (around 0.6 L kg^{-1}). The ratio between V_{ss} and the total volume of body water expresses the
372 overall partition coefficient (K_p). K_p expresses the overall affinity of the body for the molecule
373 (Toutain and Bousquet-Mélou, 2004). K_p can also be defined as the ratio between the unbound
374 fraction in plasma and the unbound fraction outside plasma. In this study, the high K_p ($10.0 \pm$
375 1.7) supposes that most of CLD is stored in the peripheral compartment and extensively bound
376 to proteins in tissues or to adipocytes (Toutain and Bousquet-Mélou, 2004).

377 ADG has a very significant effect on peripheral distribution volume V_2 ($P < 0.05$): the higher
378 the ADG, the lower the V_{ss} . This may be explained by carcass composition, when the animal
379 gain weight, lean and fat composition of carcass may change over time.

380 **Breed, average daily gain, composition of carcass: which factor may really affect TKs** 381 **variability in growing pigs?**

382 Covariate analysis showed that ADG is an important cofactor and that inter-individual
383 differences in parameters are rather related to the ADG than to breed. However, these results
384 must be put into perspective because breed and growth performance factors are inextricably
385 interlinked. Moreover, comparing a covariate (ADG) and a factor (breed) induce statistical
386 biases. These results must be discussed to explore which factor may impact inter-individual
387 variability.

388 TK parameters obtained with the NLME approach were used to compare the two breeds.
389 Estimated parameters obtained for Creole and Large White pigs with the NLME approach are
390 summarized in Table 4. Significant differences between Creole and Large White pigs were
391 observed for CL, V_2 and V_{ss} . Differences in CLD distribution (V_2 and V_{ss}) and elimination
392 (CL) may be explained by percentage of lean meat in carcass or by carcass fatness. Indeed,
393 Large White pigs have a better growth potential, higher ADG, feed efficiency and lean carcass

394 content than Creole pigs and Creole pigs have higher carcass fatness and bigger adipocytes
395 than Large White pigs (Renaudeau et al., 2005). Recently, Pouillet et al. (2019) have confirmed
396 the difference in pig metabolism between Creole and Large white pigs, by observing a lower
397 N utilization efficiency in Creole breed with a higher level of blood urea nitrogen and a lower
398 level of globulin compared to Large White pigs. Moreover, age differential between the two
399 breeds may have amplified growth performance differences and fatness of carcass
400 (Renaudeau et al., 2006).

401 CLD has a different behavior from other POPs such as PCB because CLD is known to be more
402 concentrated in liver and muscle than in fat tissues (Lastel et al., 2016; Lastel et al., 2018).
403 High concentrations of CLD in liver and muscles may be explained by its affinity for plasmatic
404 proteins (Soine et al., 1982). CLD linked to plasma proteins would arrive faster and in greater
405 quantities in organs and tissues heavily irrigated (liver and muscles) than in those weakly
406 perfused (fat tissues). Slaughtering data were analyzed to characterize molecule distribution
407 in the three matrices analyzed (liver, muscle and fat) and to highlight possible differences in
408 CLD distribution between the two breeds. Estimated amounts of CLD into tissues after 84
409 depuration-days in Creole and Large White pigs are presented in Table 5. No significant
410 difference was observed for CLD amounts in liver and muscle between the two breeds.
411 However, repartition of CLD in fat is significantly different between the two breeds and Creole
412 pigs have higher carcass adiposity. Although CLD concentrations in fat tissue are lower than
413 in muscle or liver, fat tissues store an important portion of CLD body burden, as shown in Table
414 5. Thus, carcass fatness may impact volume of distribution. To explain inter-individual
415 variability between pigs, animal-specific features (age, breed, sex), growth performance and
416 carcass fatness seem to be interesting. Breed effects we found suggest that variability between
417 breeds (especially between specialist and generalist breeds) should be taken into account in
418 decision support for a healthy farming practice. Further studies are needed to better
419 understand CLD storage, depuration capacities between tissues and characterize
420 physiological factors that drive TK parameters.

421 Table 4: Comparison of toxicokinetic parameters obtained with the nonlinear mixed-effects
 422 approach between Creole and Large White pigs
 423

Parameters	Creole (n=7) median ± SE	Large White (n=7) median ± SE	P
CL mL kg ⁻¹ d ⁻¹	82.1 ± 4.6 ^a	68.8 ± 1.6 ^b	0.0081
V1 mL kg ⁻¹	686 ± 420 ^a	1,061 ± 329 ^a	0.5236
Q mL kg ⁻¹ d ⁻¹	15,896 ± 2,435 ^a	22,841 ± 2,940 ^a	0.9604
V2 mL kg ⁻¹	5,792 ± 405 ^a	4,433 ± 232 ^b	0.0128
Vss mL kg ⁻¹	6,714 ± 270 ^a	5,316 ± 392 ^b	0.0158
MRT d	79 ± 3 ^a	71 ± 6 ^a	0.4240
t_{1/2elim} d	55 ± 2 ^a	49 ± 4 ^a	0.4112

CL: Body clearance; V1: Central compartment volume of distribution; Q: Intercompartmental clearance; V2: Peripheral compartment volume of distribution; Vss: Steady-state volume of distribution; MRT: Mean residence time; t_{1/2elim}: Elimination half-life; SE: Standard Error. Values not followed by the same superscript letter significantly differ (Student test).

424

425 Table 5: Estimated amounts of CLD into tissues after 84 depuration-days

	Creole (n=7) mean ± SE	Large White (n=7) mean ± SE	P
Liver (mg)	2.2 ± 0.1 ^a	2.2 ± 0.2 ^a	0.5619
Muscle ¹ (mg)	8.9 ± 1.2 ^a	12.5 ± 1.8 ^a	0.9352
Fat ² (mg)	13.2 ± 0.5 ^a	10.0 ± 0.8 ^b	0.0039

¹ Considering CLD concentrations in shaft muscle and that muscle percentages of body weight in Creole and Large White pigs are 43.3% and 54.8%, respectively (Renaudeau et al., 2005)

² Considering CLD concentrations in peri-renal fat tissue and that fat percentages of body weight in Creole and Large White pigs are 34.7% and 22.4%, respectively (Renaudeau et al., 2005)

Values not followed by the same superscript letter significantly differ (Student test).

426 Conclusion

427 This study permitted to establish for the first time a NLME model of CLD in growing pigs. Major
 428 TK parameters like body clearance (76.7 mL kg⁻¹ d⁻¹) and distribution volume (6 L kg⁻¹) were
 429 determined. ADG is an important covariate explaining inter-individual TK parameters
 430 variability. This study provided relevant and original data to describe distribution and
 431 elimination of CLD in pigs. In the near future, these results may be used by the risk manager
 432 to assess adequate rearing management practices in CLD contaminated areas. Moreover,

433 since the pig is a valuable model for humans, these results may also be used to build a TK
434 model of CLD in humans.

435 Acknowledgements

436 The experiment was funded by ANR through research program INSSICCA (ANR-16-CE21-
437 0008-01). The authors thank Harry Archimède (URZ), David Beramice, Bruno Bocage, Katia
438 Benony, Laurent Dantec, Mario Giorgi, Felix Silou (PTEA), Valeriusse Calif, Yoann Félicité,
439 Dalila Feuillet, Lucien Philibert (URZ) and Pamela Hartmeyer (URAFPA) for their technical
440 support. They are grateful to Alain Chauvin and Francis Le Gouevéc (UEPR, Rennes) for
441 surgeries. They warmly thank Catherine Adam (LEAE-CART) for analysis and Pierre-Loïc
442 Saaidi (Génomique Métabolique, CEA, CNRS, Univ Evry, Université Paris-Saclay, Evry,
443 France) for the 10-monohydrochlordecone synthesis. They thank Jean-Pierre Thomé (LEAE-
444 CART) for his advices.

445 References

446 Adir, J., Caplan, Y.H., Thompson, B.C.,1978. Kepone® serum half-life in humans. *Life*
447 *Sciences*. 22, 699–702. [https://doi.org/10.1016/0024-3205\(78\)90494-0](https://doi.org/10.1016/0024-3205(78)90494-0).

448 AFSSA, 2007. Avis de l'agence française de sécurité sanitaire des aliments relatif à
449 l'actualisation des données scientifiques sur la toxicité du chlordécone en vue d'une éventuelle
450 révision des limites tolérables d'exposition proposées par l'AFSSA en 2003.
451 <https://www.anses.fr/fr/system/files/RCCP2007sa0305.pdf>.

452 Anezaki, K. and Nakano, T., 2014. Concentration levels and congener profiles of
453 polychlorinated biphenyls, pentachlorobenzene, and hexachlorobenzene in commercial
454 pigments. *Environmental Science and Pollution Research*. 21, 998-1009.
455 <https://doi.org/10.1007/s11356-013-1977-2>.

456 ANSES, 2017. Exposition des consommateurs des Antilles au chlordécone, résultats de
457 l'étude Kannari. <https://www.anses.fr/fr/system/files/ERCA2014SA0029Ra.pdf>.

458 Bouveret, C., Rychen, G., Lerch, S., Jondreville, C., Feidt, C., 2013. Relative bioavailability of
459 tropical volcanic soil-bound chlordecone in piglets. *Journal of Agricultural and Food Chemistry*.
460 61, 9269–9274. <https://doi.org/10.1021/jf400697r>.

461 Cabidoche, Y.-M., Achard, R., Cattan, P., Clermont-Dauphin, C., Massat, F., Sansoulet, J.,
462 2009. Long-term pollution by chlordecone of tropical volcanic soils in the French West Indies:
463 A simple leaching model accounts for current residue. *Environmental Pollution*. 157, 1697–
464 1705. <https://doi.org/10.1016/j.envpol.2008.12.015>.

465 Chevallier, M.L., Dela-Negra, O., Chaussonnerie, S., Barbance, A., Muselet, D., Lagarde, F.,
466 Darii, E., Ugarte, E., Lescaop, E., Fonknechten, N., Weissenbach, J., Woignier, T., Gallard, J.-
467 F., Vuilleumier, S., Imfeld, G., Le Paslier, D., Saaidi, P.-L., 2019. Natural chlordecone
468 degradation revealed by numerous transformation products characterized in key French West
469 Indies environmental compartments. *Environmental Science & Technology*. 53, 6133-6143.
470 <https://doi.org/10.1021/acs.est.8b06305>.

471 Cohn, W.J., Boylan, J.J., Blanke, R.V., Fariss, M.W., Howell, J.R., Guzelian, P.S., 1978.
472 Treatment of chlordecone (Kepone) toxicity with cholestyramine – Results of a controlled
473 clinical trial. *The New England Journal of Medicine*. 298, 243–48.
474 <https://doi.org/10.1056/NEJM197802022980504>.

475 Collas, C., Mahieu, M., Tricheur, A., Crini, N., Badot, P.-M., Archimède, H., Rychen, G., Feidt,
476 C., Jurjanz, S., 2019. Cattle exposure to chlordecone through soil intake. The case-study of
477 tropical grazing practices in the French West Indies. *Science of the Total Environment*. 668,
478 161-170. <https://doi.org/10.1016/j.scitotenv.2019.02.384>.

479 Costet, N., Pelé, F., Comets, E., Rouget, F., Monfort, C., Bodeau-Livinec, F., Liganiza E.M.,
480 Bataille, H., Kadhel, P., Multigner, L., Cordier, S., 2015. Perinatal exposure to chlordecone and
481 infant growth. *Environmental Research*. 142, 123–134.
482 <https://doi.org/10.1016/j.envres.2015.06.023>.

483 Dereumeaux, C., Saoudi, A., Guldner, L., Pecheux, M., Chesneau, J., Thomé, J.-P., Ledrans,
484 M., Le Tertre, A., Denys, S., Fillol, C., 2019. Chlordecone and organochlorine compound levels
485 in the French West Indies population in 2013-2014. *Environmental and Pollution Research*.
486 <https://doi.org/10.1007/s11356-019-07181-9>.

487 Dromard, C.R., Guéné, M., Bouchon-Navaro, Y., Lemoine, S., Cordonnier, S., Bouchon, C.,
488 2017. Contamination of marine fauna by chlordecone in Guadeloupe : evidence of a seaward
489 decreasing gradient. *Environmental Science and Pollution Research*. 25, 14294-14301.
490 <https://doi.org/10.1007/s11356-017-8924-6>.

491 Dubuisson, C., Héraud, F., Leblanc, J.-C., Gallotti, S., Flamand, C., Bateau, A., Quenel, P.,
492 Volatier, J.-L., 2007. Impact of subsistence production on the management options to reduce
493 the food exposure of the Martinican population to chlordecone. *Regulatory Toxicology and*
494 *Pharmacology*. 49, 5–16. <https://doi.org/10.1016/j.yrtph.2007.04.008>.

495 Eppe, G., Schaechtele, A., Haedrich, J., Fernandes, A., 2017. Guidance document on
496 measurement uncertainty for laboratories performing PCDD/F and PCB analysis using isotope
497 dilution mass spectrometry. Working Group for Measurement Uncertainty in PCDD/F and PCB
498 Analysis. [https://ec.europa.eu/food/sites/food/files/safety/docs/animal-feed-](https://ec.europa.eu/food/sites/food/files/safety/docs/animal-feed-guidance_document_pcdd-f_pcb_en.pdf)
499 [guidance_document_pcdd-f_pcb_en.pdf](https://ec.europa.eu/food/sites/food/files/safety/docs/animal-feed-guidance_document_pcdd-f_pcb_en.pdf). <https://doi.org/10.13140/RG.2.2.35446.09285>.

500 Egle, J.L., Fernandez, S.B, Guzelian, P.S., Borzelleca, J.F., 1978. Distribution and excretion
501 of chlordecone (Kepone) in the rat. *Drug Metabolism and Disposition*. 6, 91–95.

502 Fariss, M.W., Blanke, R.V., Saady, J.J., Guzelian, P.S., 1980. Demonstration of major
503 metabolic pathways for Chlordecone (Kepone) in Humans. *Drug Metabolism and Disposition*.
504 8, 434–438.

505 Fournier, A., Feidt, C., Lastel, M.-L., Archimede, H., Thomé, J.-P., Mahieu, M., Rychen, G.,
506 2017. Toxicokinetics of Chlordecone in goats: Implications for risk management in French
507 West Indies. *Chemosphere*. 171, 564–570.
508 <https://doi.org/10.1016/j.chemosphere.2016.12.054>.

509 Gourdine, J-L, Bambou, J-C., Giorgi, M., Loranger-Merciris, G., Archimède, H., 2018.
510 Performance of growing pigs reared indoors or outdoors in sweet-potato fields. *Revue*
511 *d'élevage et de médecine vétérinaire des pays tropicaux*. 71, 41-46.
512 <https://doi.org/10.19182/remvt.31347>.

513 Houston, T.E., Mutter, L.C., Blanke, R.V., Guzelian, P.S., 1981. Chlordecone alcohol formation
514 in the Mongolian gerbil (*Meriones Uniguiculatus*): a model for human metabolism of
515 Chlordecone (Kepone). *Fundamental and Applied Toxicology*. 1, 293–298.

516 Hu, D. and Hornbuckle, K.C., 2010. Inadvertent polychlorinated biphenyls in commercial paint
517 pigments. *Environmental Science and Technology*. 44 (8), 2822-2827.
518 <https://doi.org/10.1021/es902413k>.

519 Jondreville, C., Fournier, A., Mahieu, M., Feidt, C., Archimède, H., Rychen, G., 2014. Kinetic
520 study of chlordecone orally given to laying hens (*Gallus Domesticus*). *Chemosphere*. 114,
521 275–281. <https://doi.org/10.1016/j.chemosphere.2014.05.008>.

522 Jurjanz, S. and Roinsard, A., 2014. Valorisation de l'herbe par des truies plein-air.
523 ALTERAGRI. May 2014, 125, p. 25-28.

524 Jurjanz, S., Jondreville, C., Mahieu, M., Fournier, A., Archimède, H., Rychen, G., Feidt, C.,
525 2014. Relative bioavailability of soil-bound chlordecone in growing lambs. *Environmental*
526 *Geochemistry and Health*. 36, 911–917. <https://doi.org/10.1007/s10653-014-9608-5>.

527 Jurjanz, S., Collas, C., Lastel, M.-L., Godard, X., Archimède, H., Rychen, G., Mahieu, M., Feidt,
528 C., 2017. Evaluation of soil intake by growing Creole young bulls in common grazing systems
529 in humid tropical conditions. *Animal*. 11. <https://doi.org/10.1017/S1751731116002755>.

530 Lastel, M.-L., Lerch, S., Fournier, A., Jurjanz, S., Mahieu, M., Archimède, H., Feidt, C., Rychen,
531 G., 2016. Chlordecone disappearance in tissues of growing goats after a one month
532 decontamination period - Effect of body fatness on Chlordecone retention. *Environmental*
533 *Science and Pollution Research*. 23, 3176–3183. <https://doi.org/10.1007/s11356-015-5833-4>.

534 Lastel, M.-L., Fournier, A., Jurjanz, S., Thomé, J.-P., Joaquim-Justo, C., Archimède, H.,
535 Mahieu, M., Feidt, C., Rychen, G., 2018. Comparison of chlordecone and NDL-PCB
536 decontamination dynamics in growing male kids after cessation of oral exposure: is there a
537 potential to decrease the body levels of these pollutants by dietary supplementation of
538 activated carbon or paraffin oil ?. *Chemosphere*. 193, 100-107.
539 <https://doi.org/10.1016/j.chemosphere.2017.10.120>.

540 Lerch, S., Guidou, C., Thomé, J.-P., Jurjanz, S., 2016. Non-dioxin-like Polychlorinated
541 Biphenyls (PCBs) and chlordecone release from adipose tissue to blood in response to body
542 fat mobilization in Ewe (*Ovis aries*). *Journal of Agricultural and Food Chemistry*. 64, 1212–
543 1220. <https://doi.org/10.1021/acs.jafc.5b05817>.

544 Molowa, D.T., Shayne, A.G., Guzelian, P.S., 1986a. Purification and characterization of
545 chlordecone reductase from Human liver. *The Journal Of Biological Chemistry*. 261, 12624–
546 12627.

547 Molowa, D.T., Wrighton, S.A., Blanke, R.V., Guzelian, P.S., 1986b. Characterization of a
548 unique aldo-keto reductase responsible for the reduction of chlordecone in the liver of the gerbil
549 and man. *Journal of Veterinary Pharmacology and Therapeutics*. 17, 375–384.
550 <https://doi.org/10.1080/15287398609530832>.

551 Multigner, L., Ndong, J.-R., Giusti, A., Romana, M., Delacroix-Maillard, H., Cordier, S., Jégou,
552 B., Thomé, J.-P., Blanchet, P., 2010. Chlordecone exposure and risk of prostate cancer.
553 *Journal of Clinical Oncology*. 28, 3457–3462. <https://doi.org/10.1200/JCO.2009.27.2153>.

554 Pouillet, N., Bambou, J.-C., Loyau, T., Trefeu, C., Feuillet, D., Beramice, D., Bocage, B.,
555 Renaudeau, D., Gourdine, J.-L., 2019. Effect of feed restriction and refeeding on performance
556 and metabolism of European and Caribbean growing pigs in a tropical climate. *Scientific*
557 *Reports*. 9, 4878. <https://doi.org/10.1038/s41598-019-41145-w>.

558 Renaudeau, D., Hilaire, M., Mourot, J., 2005. A comparison of carcass and meat quality
559 characteristics of Creole and Large White pigs slaughtered at 150 days of age. *Animal*
560 *Research*. 54, 43-54. <https://doi.org/10.1051/animres:2004042>.

561 Renaudeau, D., Giorgi, M., Silou, F., Weisbecker, J.-L., 2006. Effect of breed (lean or fat pigs)
562 and sex on performance and feeding behavior of group housed growing pigs in a tropical
563 climate. *Asian-Australasian Journal of Animal Sciences*. 19, 593-600.
564 <https://doi.org/10.5713/ajas.2006.593>.

565 Saint-Hilaire, M., 2018. Toxicocinétique de la chlordécone chez la brebis. Université de
566 Lorraine. 250 p.

567 Saint-Hilaire, M., Rychen, G., Thomé, J.-P., Joaquim-Justo, C., Le Roux, Y., Feidt, C.,
568 Fournier, A., 2019. Linear toxicokinetic of chlordecone in ewe's serum. *Environmental Science*
569 *and Pollution Research*. <https://doi.org/10.1007/s11356-019-05800-z>.

570 SANCO, 2014. Guidance Document on Analytical Quality Control and Validation Procedures
571 for Pesticide Residues Analysis in Food and Feed. SANCO/12571/2013 Supersedes
572 SANCO/12495/2011, 48 p.. [http://www.eurl-](http://www.eurl-pesticides.eu/library/docs/allcrl/AqcGuidance_Sanco_2013_12571.pdf)
573 [pesticides.eu/library/docs/allcrl/AqcGuidance_Sanco_2013_12571.pdf](http://www.eurl-pesticides.eu/library/docs/allcrl/AqcGuidance_Sanco_2013_12571.pdf).

574 Soine, P.J., Blanke, R.V., Guzelian, P.S., Schwartz, C.C., 1982. Preferential binding of
575 chlordecone to the protein and high density lipoprotein fractions of plasma from humans and
576 other species. *Journal of Toxicology and Environmental Health*. 9, 107–118.
577 <https://doi.org/10.1080/15287398209530146>.

578 Soine, P.J., Blanke, R.V. and Schwartz, C.C, 1983. Chlordecone metabolism in the pig.
579 *Toxicology Letters*. 17, 35–41. [https://doi.org/10.1016/0378-4274\(83\)90032-2](https://doi.org/10.1016/0378-4274(83)90032-2).





580 Toutain, P. L. and Bousquet-Melou, A. 2004. Volumes of Distribution. *Journal of Veterinary*
581 *Pharmacology and Therapeutics*. 27, 441–53. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2885.2004.00602.x)
582 [2885.2004.00602.x](https://doi.org/10.1111/j.1365-2885.2004.00602.x).

583 Koh, W.X., Hornbuckle, K.C., Thorne, P.S., 2015. Human serum from urban and rural
584 adolescents and their mothers shows exposure to polychlorinated biphenyls not found in
585 commercial mixtures. *Environmental Science and Technology*. 49, 8105-8112.
586 <https://doi.org/10.1021/acs.est.5b01854>.

Figure 1: Two-compartmental modelisation of the equilibrium between the central compartment and the peripheral compartment after a single intravenous administration of chlordecone (adapted from Toutain and Bousquet-Melou, 2004)

Figure 2: Serum CLD concentration (ng.mL^{-1}) vs. time in growing pigs ($n=14$) after a single intravenous administration of CLD at $1 \text{ mg.kg}^{-1} \text{ BW}$: observed kinetics from $t=0$ to $t=84$ days
Top graph on the right corresponds to an enlargement of the main graph from $t=0$ to $t=1.4$ days.

Chlordecone
single intravenous administration



Clearance Q
 $Q = k_{12} \cdot V1 = k_{21} \cdot V2$




Table 1: Average values of IBW, FBW and ADG for each breed

Parameters	Creole pigs (n=7)	Large White pigs (n=7)	All pigs (n=14)
	mean \pm SE	mean \pm SE	mean \pm SE
IBW	68.8 \pm 1.8 ^a	67.9 \pm 3.1 ^a	68.3 \pm 1.7
kg			
FBW	91.2 \pm 2.8 ^a	95.6 \pm 2.4 ^a	92.8 \pm 1.9
kg			
ADG	248 \pm 25 ^a	327 \pm 13 ^a	288 \pm 17
g d ⁻¹			

IBW: Initial Body Weight; FBW: Final Body Weight; ADG: Average Daily Gain; SE: Standard Error. Values not followed by the same superscript letter significantly differ (P < 0.05; Student test)

Table 2: Comparison of CLD toxicokinetic parameters obtained with non-compartmental analysis between pigs, ewes and goats following a single intravenous administration of 1 mg CLD kg⁻¹

Parameters	Pigs (n=14)		Ewes (n=5)		Goats (n=4)	
	Mean ± SE	CV (%)	Mean ± SE	CV (%)	Mean ± SE	CV (%)
CL mL kg ⁻¹ d ⁻¹	79.6 ± 4.4 ^a	5.5	51.0 ± 6.1 ^b	12.0	66.6 ± 16.2 ^{ab}	24.4
Vss mL kg ⁻¹	5,982 ± 354 ^a	5.9	1,807 ± 157 ^b	8.7	1,615 ± 446 ^b	27.6
MRT d	77 ± 5 ^a	6.6	36 ± 2 ^b	4.5	24 ± 1 ^b	3.9
AUC ng d mL ⁻¹	12,505 ± 636 ^a	5.1	21,202 ± 3,373 ^b	15.9	18,911 ± 5,687 ^{ab}	30.1

CL: Body clearance; Vss: Volume of distribution at the equilibrium; MRT: Mean Residence Time; AUC: Area Under the Curve; CV: coefficient of variation; SE: Standard Error.

Values not followed by the same superscript letter significantly differ ($P < 0.05$; Tukey test).

Toxicokinetic parameters calculated from data obtained in Saint-Hilaire et al. (2019) and Fournier et al. (2017) for ewes and goats, respectively.

Table 3: Population toxicokinetic model estimates for CLD following a single intravenous administration of 1 mg CLD kg⁻¹ to 14 pigs

Parameters	Model estimate median ± SE	P ¹	CV (%)
CL mL kg ⁻¹ d ⁻¹	76.7 ± 3.2	0.0009	4.1
V1 mL kg ⁻¹	604 ± 195	-	32.4
Q mL kg ⁻¹ d ⁻¹	16,680 ± 2,870	-	17.1
V2 mL kg ⁻¹	4,840 ± 262	0.0142	5.4
Vss mL kg ⁻¹	6,002 ± 280	-	4.6
MRT d	78 ± 3	-	4.3
t_{1/2elim} d	54.7 ± 2.3	-	4.2

CL: Body clearance; V1: Central compartment volume of distribution; Q: Intercompartmental clearance; V2: Peripheral compartment volume of distribution; Vss: Steady-state volume of distribution; MRT: Mean residence time; t_{1/2elim}: Elimination half-life; CV: Coefficient of variation; SE: Standard Error.

¹P-value obtained after a Wald test, testing significant effect of Average Daily Gain (ADG) on CL and V2 parameters.

Table 4: Comparison of toxicokinetic parameters obtained with the nonlinear mixed-effects approach between Creole and Large White pigs

Parameters	Creole (n=7) median \pm SE	Large White (n=7) median \pm SE	P
CL mL kg ⁻¹ d ⁻¹	82.1 \pm 4.6 ^a	68.8 \pm 1.6 ^b	0.0081
V1 mL kg ⁻¹	686 \pm 420 ^a	1,061 \pm 329 ^a	0.5236
Q mL kg ⁻¹ d ⁻¹	15,896 \pm 2,435 ^a	22,841 \pm 2,940 ^a	0.9604
V2 mL kg ⁻¹	5,792 \pm 405 ^a	4,433 \pm 232 ^b	0.0128
Vss mL kg ⁻¹	6,714 \pm 270 ^a	5,316 \pm 392 ^b	0.0158
MRT d	79 \pm 3 ^a	71 \pm 6 ^a	0.4240
t_{1/2elim} d	55 \pm 2 ^a	49 \pm 4 ^a	0.4112

CL: Body clearance; V1: Central compartment volume of distribution; Q: Intercompartmental clearance; V2: Peripheral compartment volume of distribution; Vss: Steady-state volume of distribution; MRT: Mean residence time; t_{1/2elim}: Elimination half-life; SE: Standard Error. Values not followed by the same superscript letter significantly differ (Student test).

Table 5: Estimated amounts of CLD into tissues after 84 depuration-days

	Creole (n=7) mean ± SE	Large White (n=7) mean ± SE	P
Liver (mg)	2.2 ± 0.1 ^a	2.2 ± 0.2 ^a	0.5619
Muscle ¹ (mg)	8.9 ± 1.2 ^a	12.5 ± 1.8 ^a	0.9352
Fat ² (mg)	13.2 ± 0.5 ^a	10.0 ± 0.8 ^b	0.0039

¹ Considering CLD concentrations in shaft muscle and that muscle percentages of body weight in Creole and Large White pigs are 43.3% and 54.8%, respectively (Renaudeau et al., 2005)

² Considering CLD concentrations in peri-renal fat tissue and that fat percentages of body weight in Creole and Large White pigs are 34.7% and 22.4%, respectively (Renaudeau et al., 2005)

Values not followed by the same superscript letter significantly differ (Student test).