



HAL
open science

On the regularity of the Green current for semi-extremal endomorphisms of P^2

Christophe Dupont, Axel Rogue

► **To cite this version:**

Christophe Dupont, Axel Rogue. On the regularity of the Green current for semi-extremal endomorphisms of P^2 . *Discrete & Continuous Dynamical Systems - A*, 2020, Conference on Dynamics, Equations and Applications (DEA), 40 (12), pp.6767-6781. 10.3934/dcds.2020163 . hal-02936357

HAL Id: hal-02936357

<https://hal.science/hal-02936357>

Submitted on 17 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE REGULARITY OF THE GREEN CURRENT FOR SEMI-EXTREMAL ENDOMORPHISMS OF \mathbb{P}^2

CHRISTOPHE DUPONT AND AXEL ROGUE

Université de Rennes, CNRS, IRMAR - UMR 6625
F-35000 Rennes, France

ABSTRACT. We study the regularity of the Green current for semi-extremal endomorphisms of \mathbb{P}^2 . Under suitable assumptions, we show that the point-wise lower Radon-Nikodym derivative of stable slices with respect to the one dimensional Lebesgue measure is bounded at almost every point for the equilibrium measure. This provides a weak amount of metric regularity for the Green current along holomorphic discs.

1. Introduction. Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let T be the Green current and let $\mu := T \wedge T$ be the equilibrium measure of f , see [13, 26]. The measure μ is invariant and ergodic, we denote \mathcal{J} its support. The critical set of f satisfies $\mu(\mathcal{C}) = 0$. The Lyapunov exponents $\lambda_2 \leq \lambda_1$ of μ will play a central role in this article. They are bounded below by $\frac{1}{2} \log d$, see [8].

1.1. Extremal and semi-extremal endomorphisms. We say that f is *extremal* if $\lambda_1 = \lambda_2 = \frac{1}{2} \log d$. Those endomorphisms are characterized by the four equivalent properties: $\mu \ll \text{Leb}_{\mathbb{P}^2}$, $\dim_H \mu = 4$, T is a positive smooth $(1, 1)$ -form on some open subset of \mathbb{P}^2 , and f is a Lattès map, see [6, 4, 10]. We recall that a Lattès map on \mathbb{P}^k is the projection of an affine dilation on a complex k -torus by means of a finite galoisian covering σ , see [15] for examples. Similar characterizations were proved for rational maps on \mathbb{P}^1 , in which case $\lambda = \frac{1}{2} \log d$ and $T = \mu$, see [23, 24, 27].

We say that f is *semi-extremal* if $\lambda_1 > \lambda_2 = \frac{1}{2} \log d$. Examples are provided by endomorphisms f of \mathbb{P}^2 having an invariant pencil of lines on which f induces a one dimensional Lattès map, see [18]. A natural question is to find characterizations of semi-extremal endomorphisms, as we have for extremal ones. We already know that they satisfy $\dim_H \mu = 2 + \frac{\log d}{\lambda_1}$, see [7, 10, 17], and that f is semi-extremal once $\mu \ll T \wedge \omega$, see [14]. Here ω stands for the Fubini-Study $(1, 1)$ -form on \mathbb{P}^2 , so that $T \wedge \omega$ is the trace measure of the Green current. Dujardin asked in [14, Section 3] the following questions.

Question 1.1. *Does every semi-extremal endomorphism satisfy $\mu \ll T \wedge \omega$? If $\mu \ll T \wedge \omega$, does f contain (in some sense) a one-dimensional Lattès map?*

A strategy for the first question would to show that semi-extremal endomorphisms have an invariant pencil of lines. Indeed, in this case the radial Lyapunov exponent of f is $\geq \log d$ (by [18, Theorem 1.7]), hence the Lyapunov exponent of the action of f on the pencil of lines (which belongs to the spectrum of μ by [18,

2020 *Mathematics Subject Classification.* 37F10, 32U40, 28A15.

Key words and phrases. Holomorphic dynamics, Lyapunov exponents, Green current.

Theorem 1.5]) must be equal to $\frac{1}{2} \log d$ (semi-extremality is assumed). This implies that f induces a Lattès map on the pencil of lines, which ensures that $\mu \ll T \wedge \omega$ by [18, Corollary 1.3].

We can make a similar observation for the second question. If $\mu \ll T \wedge \omega$ then $\frac{1}{2} \log d$ is one of the Lyapunov exponents of μ by [14]. Using the same arguments as before (relying on [18]), the Lyapunov exponent of the action of f on the pencil of lines must be $\frac{1}{2} \log d$, ensuring that f induces a Lattès map on this pencil.

In the general case (without assuming the existence of an invariant pencil of lines), the condition $\mu \ll T \wedge \omega$, equivalent to $T \wedge T \ll T \wedge \omega$, formally says that $T \ll \omega$ with respect to T . A first step to exhibit such regular properties for T consists in establishing regular properties for slices. Theorems 1.3 and 1.4 below provide results in that direction: they bound the pointwise lower Radon-Nikodym derivative of stable slices of T at μ -almost every point. In Section 1.4 we shall examine the guideline given by polynomial lifts of Lattès mappings on \mathbb{P}^1 : such regular slices of course exist, moreover there exist regular slices which do not intersect the support of μ (see Remark 1.6).

1.2. Statements of the results.

Definition 1.2. Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that the Lyapunov exponents of the equilibrium measure of f are different, hence $\lambda_2 < \lambda_1$. Let $v_s(x) \in \mathbb{P}(T_x \mathbb{P}^2)$ denote the stable Oseledec direction, which is defined for every x in an invariant μ -generic Borel subset A . Let R be the set of points $x \in A$ such that there exists a holomorphic disc $\xi_x : \mathbb{D} \rightarrow \mathbb{P}^2$ satisfying $\xi_x(0) = x$, $[\xi'_x(0)] = v_s(x)$ and

$$\liminf_{r \rightarrow 0} \frac{\xi_x^* T(\mathbb{D}(r))}{\text{Leb}(\mathbb{D}(r))} < \infty. \quad (1.1)$$

In that definition $[v] \in \mathbb{P}(T_x \mathbb{P}^2)$ stands for the line of $T_x \mathbb{P}^2$ directed by v . Since $\mu(\mathcal{C}) = 0$, one can assume that A does not intersect $\cup_{n \in \mathbb{Z}} f^n(\mathcal{C})$. The relation $f^* T = dT$ then implies that R is totally f -invariant, in particular, $\mu(R) \in \{0, 1\}$ by ergodicity. We prove that $\mu(R) > 0$ under two different sets of assumptions related to semi-extremality.

Theorem 1.3. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that $\mu \ll T \wedge \omega$ and $\lambda_2 < \lambda_1 < 2\lambda_2$. Then $\mu(R) = 1$.*

Theorem 1.4. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that f is semi-extremal and that $\lambda_2 < \lambda_1 < 2\lambda_2$. Assume also that $\mathcal{J} \cap \mathcal{C} = \emptyset$ and that v_s is Hölder continuous on A . Then $\mu(R) = 1$.*

For the second result, we shall see in Section 2.2 that the Hölder continuity of v_s on A is satisfied when f is partially hyperbolic on \mathcal{J} .

1.3. Outline of the proofs. To prove Theorems 1.3 and 1.4, a first step will consist in showing

$$d^{n_j} e^{-2M_1} \leq \|D_x f^{n_j}(\vec{v}_s(x))\|^2 \leq d^{n_j} e^{2M_1} \quad (1.2)$$

for every x in a subset of positive μ -measure, where $(n_j)_j$ is an increasing sequence of integers depending on x and $\vec{v}_s(x) \in T_x \mathbb{P}^2$ is a unitary vector in $v_s(x) \in \mathbb{P}(T_x \mathbb{P}^2)$ defined in Definition 1.2. It is crucial that no exponential error term $e^{\pm n\epsilon}$ occur in (1.2). Such error terms are actually inherent to (our context of) non uniform hyperbolicity, a difficulty is thus to get rid of them. A second step will be to integrate

(1.2), namely to construct a holomorphic disc on which f^{n_j} is a multiplication by $d^{n_j/2}e^{\pm M_1}$. We will use for that purpose a normal form Theorem for the inverse branches of f . Theorem 3.1 (see Section 3) summarizes both steps, it provides workable sufficient conditions ensuring that a point belongs to R .

Let us specify how the estimates (1.2) will be obtained. The lower bound is actually true for every endomorphism of \mathbb{P}^2 satisfying $\lambda_2 \leq \lambda_1 < 2\lambda_2$, it does not need any semi-extremality assumption, see Theorem 2.7. The upper bound in (1.2) needs semi-extremality: for Theorem 1.3 we use arguments developed by Dujardin [14] to construct Fatou directions (see Section 4), and for Theorem 1.4 we use the Central Limit Theorem for the logarithm of the tangent map on the stable direction $v_s(x)$ (see Section 5).

Let us notice that the Central Limit Theorem was used in [27] to prove regularity properties for the equilibrium measure of rational maps on \mathbb{P}^1 and in [16] to provide a new proof of the fact that μ is absolutely continuous when f is extremal on \mathbb{P}^k .

1.4. Study of the lifts of one dimensional Lattès mappings. The polynomial lifts of Lattès mappings on \mathbb{P}^1 are the most simple examples of semi-extremal endomorphisms of \mathbb{P}^2 . In this section we verify for them $\mu(R) = 1$ and $\mu \ll T \wedge \omega$ by using normal forms for their Green function.

Proposition 1.5. *Let $f[z : w : t] := [P(z, w) : Q(z, w) : t^d]$, where $[P : Q]$ is a Lattès mapping of \mathbb{P}^1 of degree $d \geq 2$. Then $\mu(R) = 1$.*

Proof. The mapping f belongs to the class of regular polynomial endomorphisms of \mathbb{C}^2 studied by Bedford-Jonsson [2]. Let $L := (P, Q)$ and let $L^n = (P^n, Q^n)$ denote the n -th iterate of L . The Green function of the homogeneous polynomial mapping L is

$$G(z, w) := \lim_n \frac{1}{d^n} \log \|(P^n(z, w), Q^n(z, w))\|.$$

Let us denote by $\Omega := \{G < 0\}$ the basin of attraction of the origin in \mathbb{C}^2 , and by $\partial\Omega = \{G = 0\}$ the boundary of Ω . In the affine chart $\{t = 1\}$, the Green current T of f is equal to $dd^c G^+$, where $G^+ := \max\{G, 0\}$. Since $[P : Q]$ is a Lattès mapping, the support of the equilibrium measure $\mu = T \wedge T$ of f coincides with $\partial\Omega$.

The results [3, Theorem 1.2 and Lemma 4.2] due to Berteloot-Loeb provide normal forms for the function G when $[P : Q]$ is a Lattès map of \mathbb{P}^1 of degree d . Precisely, if x does not belong to the union S of a finite number of complex lines in \mathbb{C}^2 passing through the origin (which has zero μ -measure), then there exist a biholomorphism $p : (\mathbb{C}^2, (0, 0)) \rightarrow (\mathbb{C}^2, x)$ and $\delta \in \mathbb{R}$ such that

$$G \circ p(Z, W) = |Z|^2 + \Re(W) \quad \text{and} \quad p^{-1} \circ L \circ p(Z, W) = (e^{i\delta} \sqrt{d} Z, dW).$$

In particular, up to a holomorphic change of coordinates near x , $\partial\Omega$ is a piece of the 3-euclidian sphere. Now observe that the function $G \circ p$ is strictly subharmonic on the Z -axis (the unique complex line in the tangent space at $(0, 0)$ of the real hypersurface $\{G \circ p = 0\}$), which is included in $\{G \circ p \geq 0\}$. Hence G^+ is strictly subharmonic along the holomorphic disc

$$\xi_x(u) := p(u, 0).$$

The limit of $\frac{\xi_x^* T(\mathbb{D}(r))}{\text{Leb}(\mathbb{D}(r))}$ when r tends to 0 then exists and is equal to $dd^c(G^+ \circ \xi_x)(0) \in \mathbb{R}_+^*$. The fact that $[\xi'_x(0)] = v_s(x)$ comes from [3, Lemma 4.2], which lifts the Lattès commuting diagram to line bundles, the local coordinate in the fibers being W . We thus have verified that μ -almost every x belongs to R , as desired. \square

Remark 1.6. In the preceding proof, for every $W_0 \in (\mathbb{C}, 0)$ such that $\Re(W_0) > 0$, the function G^+ is strictly subharmonic along the holomorphic disc $\xi(u, W_0) := p(u, W_0)$. Such a disc is included in $\{G \circ p > 0\}$, which does not intersect the support of μ .

Proposition 1.7. *Let $f[z : w : t] := [P(z, w) : Q(z, w) : t^d]$, where $[P : Q]$ is a Lattès mapping of \mathbb{P}^1 of degree $d \geq 2$. Then $\mu \ll T \wedge \omega$.*

Proof. We take the notations of the proof of Proposition 1.5. It suffices to verify $\mu \ll T \wedge \omega$ at every point $x \in \partial\Omega \setminus S$, since $\mu(S) = 0$. We work with the local coordinates provided by p . Let $G_0(Z, W) := |Z|^2 + \Re(W)$, $T_0 := dd^c G_0^+$ and $\omega_0 := \frac{i}{2}dZ \wedge d\bar{Z} + \frac{i}{2}dW \wedge d\bar{W}$. For every test function φ on $(\mathbb{C}^2, (0, 0))$, we have by using and adapting [22, Proposition 6.5.5]:

$$\int_{(\mathbb{C}^2, (0,0))} \varphi T_0 \wedge \omega_0 = \int_{\{G_0=0\}} \varphi d^c G_0 \wedge \omega_0 + \int_{\{G_0 \geq 0\}} \varphi dd^c G_0 \wedge \omega_0, \tag{1.3}$$

$$\int_{(\mathbb{C}^2, (0,0))} \varphi T_0 \wedge T_0 = \int_{\{G_0=0\}} \varphi d^c G_0 \wedge dd^c G_0, \tag{1.4}$$

where the orientation on $\{G_0 = 0\}$ is induced by the one of $\{G_0 \leq 0\}$. A straightforward computation shows that

$$d^c G_0 \wedge \omega_0 = d^c G_0 \wedge dd^c G_0, \tag{1.5}$$

on the spherical 3-manifold $\{G_0 = 0\}$. Finally, one obtains $T_0 \wedge T_0 \ll T_0 \wedge \omega_0$ by using (1.5) and comparing (1.3) and (1.4). \square

Remark 1.8. The measure given by (1.5) is equal to $d\Im W \wedge \frac{i}{2}dZ \wedge d\bar{Z}$, which belongs to the Lebesgue measure class of the 3-manifold $\{G_0 = 0\}$.

2. Oseledec theorem, stable directions, normal forms.

2.1. Oseledec theorem. Let us state the Oseledec Theorem in our context.

Theorem 2.1. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Let $\lambda_2 \leq \lambda_1$ be the Lyapunov exponents of the equilibrium measure μ of f .*

1. *If $\lambda_1 = \lambda_2 = \lambda$, then there exists an invariant Borel subset A of full μ -measure and disjoint from $\cup_{n \in \mathbb{Z}} f^n(\mathcal{C})$ such that for every $x \in A$:*

$$\forall \vec{v} \in T_x \mathbb{P}^2 \setminus \{0\}, \quad \lim_{n \rightarrow +\infty} \frac{1}{n} \log \|D_x f^n(\vec{v})\| = \lambda.$$

2. *If $\lambda_1 > \lambda_2$, then there exists an invariant Borel subset A of full μ -measure and disjoint from $\cup_{n \in \mathbb{Z}} f^n(\mathcal{C})$ such that for every $x \in A$, there exists $v_s(x) \in \mathbb{P}(T_x \mathbb{P}^2)$ satisfying:*

$$\forall \vec{v} \in T_x \mathbb{P}^2 \setminus v_s(x), \quad \lim_{n \rightarrow +\infty} \frac{1}{n} \log \|D_x f^n(\vec{v})\| = \lambda_1.$$

$$\forall \vec{v} \in v_s(x) \setminus \{0\}, \quad \lim_{n \rightarrow +\infty} \frac{1}{n} \log \|D_x f^n(\vec{v})\| = \lambda_2.$$

Moreover v_s is measurable and satisfies $[D_x f](v_s(x)) = v_s(f(x))$ for every $x \in A$, where $[D_x f]$ is the projectivization of the tangent map $D_x f$.

Let us assume that $\lambda_2 < \lambda_1$. For every $x \in A$, we denote

$$\psi(x) := \log \|D_x f(\vec{v}_s(x))\|,$$

where $\vec{v}_s(x) \in \{\vec{v} \in v_s(x) \subset T_x \mathbb{P}^2, \|\vec{v}\| = 1\}$. We have $\psi \in L^1(\mu)$, see [26, Section 3.7].

Lemma 2.2. *For every $x \in A$ and every $n \geq 1$, we have*

$$\frac{1}{n} \sum_{i=0}^{n-1} \psi(f^i(x)) = \frac{1}{n} \log \|D_x f^n(\vec{v}_s(x))\|.$$

This implies $\int \psi d\mu = \lambda_2$.

Proof. The first formula follows from the definition of ψ and the relation $[D_x f](v_s(x)) = v_s(f(x))$. We obtain the second formula by taking limits μ -almost everywhere: the left hand side converges to $\int \psi d\mu$ by Birkhoff ergodic Theorem, and the right hand side converges to λ_2 by Theorem 2.1. \square

2.2. Hölder continuity of the stable direction. The following result, due to Brin, is [25, Proposition 3.9] adapted to our setting, see also [1, Section 5.3]. We use the notation dist for the distances on \mathbb{P}^2 and $\mathbb{P}(T\mathbb{P}^2)$.

Theorem 2.3. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that f is partially hyperbolic on \mathcal{J} : that means that there exist $0 < \lambda_- < \lambda_+$ and $c \geq 1$ such that for every $x \in \mathcal{J}$, there exist unitary vectors $\vec{w}_s(x), \vec{w}_u(x) \in T_x \mathbb{P}^2$ whose angle is uniformly bounded from below on \mathcal{J} and which satisfy*

$$\forall n \geq 1, \|D_x f^n(\vec{w}_s(x))\| \leq c\lambda_-^n, \|D_x f^n(\vec{w}_u(x))\| \geq c^{-1}\lambda_+^n. \tag{2.1}$$

Then the direction $w_s = [\vec{w}_s] : \mathcal{J} \rightarrow \mathbb{P}(T\mathbb{P}^2)$ is Hölder continuous. More precisely, if $b := \max\{1, \|f\|_{C^1}\}$ then for every $a > b^2$ there exists $D_a > 1$ such that:

$$\text{dist}(x, y) < D_a^{-1} \Rightarrow \text{dist}(w_s(x), w_s(y)) \leq 3c^2 \frac{\lambda_+}{\lambda_-} (D_a \text{dist}(x, y))^{\frac{\log(\lambda_+/\lambda_-)}{\log(a/\lambda_-)}}.$$

Remark 2.4. If f is partially hyperbolic on \mathcal{J} and if $\lambda_2 < \lambda_1$, then one verifies that the stable Oseledec direction $v_s(x)$ provided by Theorem 2.1 coincides with $w_s(x)$ for every $x \in A$, hence v_s is Hölder continuous on A by Theorem 2.3. In particular, every partially hyperbolic endomorphism satisfying $\lambda_2 = \frac{1}{2} \log d$, $\lambda_2 < \lambda_1 < 2\lambda_2$ and $\mathcal{J} \cap \mathcal{C} = \emptyset$ fulfills the assumptions of Theorem 1.4.

Actually it is not easy to find examples of semi-extremal endomorphisms partially hyperbolic on \mathcal{J} . For instance the lift $f[z : w : t] = [z^2 - 2w^2 : z^2 : t^2]$ of the Lattès mapping $L[z : w] = [z^2 - 2w^2 : z^2]$ is not partially hyperbolic on \mathcal{J} . To see this, observe that the Lyapunov exponents of (f, μ) are $\frac{1}{2} \log 2$ (coming from the Lattès mapping L) and $\log 2$ (coming from t^2). In particular, the modulus of the multiplier of every n -repulsive cycle of L not intersecting the critical values of σ is equal to $(\sqrt{2})^n$. On the other hand, that modulus is equal to 4 for the fixed point $[1 : 1]$ (this is a critical value of σ , see [3, Section 3]). Hence f has n -periodic points in \mathcal{J} with multipliers $\{(\sqrt{2})^n, 2^n\}$ and $\{4^n, 2^n\}$ respectively. It follows that no $\lambda_- < \lambda_+$ satisfy (2.1) over \mathcal{J} .

Examples of partially hyperbolic endomorphisms should be found among uniformly expanding endomorphisms. Jonsson [20] obtained examples in the family of polynomial skew products. For instance, $f(z, w) = (z^2, w^2 + zw + cz)$ is uniformly expanding on \mathcal{J} if c is large enough (see Theorems 4.8, 8.2 and Examples 9.2 of

this article). Corollary 8.3 of [20] moreover characterizes the Axiom A property in terms of the dynamics of the critical set: $f(z, w) = (z^2 - 6, w^2 + 3 - z)$ is uniformly expanding on \mathcal{J} . We refer to [19] for results on a family of Axiom A endomorphisms of \mathbb{P}^2 (called *s-hyperbolic*): basic sets, attractors and invariant currents are investigated there in Sections 3, 4 and 5.

To deduce partial hyperbolicity from uniform expansivity is also not straightforward. This should involve the Lyapunov exponents of μ : the stable Oseledec direction $v_s(x)$ is a natural candidate for $w_s(x)$. Nonetheless, that direction is only defined μ -a.e. on \mathcal{J} , and an other difficulty comes from the fact that the Lyapunov exponents only give an asymptotic growth rate. Hence, even if the Lyapunov exponents are distinct, the existence of $c \geq 1$ and $\lambda_- < \lambda_+$ satisfying (2.1) for every $n \geq 1$ is not obvious.

Before stating the next Proposition, let us recall Remark 2.4: if f is partially hyperbolic on \mathcal{J} and if $\lambda_2 < \lambda_1$, then the stable Oseledec direction v_s provided by Theorem 2.1 is Hölder continuous on A .

Proposition 2.5. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that $\lambda_2 < \lambda_1$ and that v_s is Hölder continuous on A .*

1. *Then v_s extends to a Hölder continuous map on \mathcal{J} , still denoted v_s .*
2. *If $\mathcal{J} \cap \mathcal{C} = \emptyset$, then $\psi(x) = \log \|D_x f(\vec{v}_s(x))\|$ is Hölder continuous on \mathcal{J} .*

Proof. The first item is classical by using Cauchy sequences and the fact that A is dense in \mathcal{J} . Let us write $\text{dist}(v_s(x), v_s(y)) \leq c'd(x, y)^\alpha$ on \mathcal{J} and prove the second item. By working in charts of $T\mathbb{P}^2$, let us fix a section of the unit tangent bundle $x \mapsto \vec{v}_s(x)$ such that $\vec{v}_s(x) \in v_s(x)$ and $\|\vec{v}_s(x) - \vec{v}_s(y)\| \leq c' \text{dist}(x, y)^\alpha$. We get

$$\begin{aligned} & \|D_x f(\vec{v}_s(x)) - D_y f(\vec{v}_s(y))\| \\ & \leq \|D_x f(\vec{v}_s(x)) - D_x f(\vec{v}_s(y))\| + \|D_x f(\vec{v}_s(y)) - D_y f(\vec{v}_s(y))\| \\ & \leq \|D_x f\| \|\vec{v}_s(x) - \vec{v}_s(y)\| + \|D_x f - D_y f\| \|\vec{v}_s(y)\| \\ & \leq \|f\|_{C^2} (1 + c')d(x, y)^\alpha. \end{aligned}$$

This implies that $x \mapsto \|D_x f(\vec{v}_s(x))\|$ is Hölder continuous. Since $\mathcal{J} \cap \mathcal{C} = \emptyset$, that function is bounded below by a constant $\rho > 0$ on \mathcal{J} . By using that \log is ρ^{-1} -Lipschitz on $[\rho, +\infty[$, we get that $\psi(x) = \log \|D_x f(\vec{v}_s(x))\|$ is Hölder continuous on \mathcal{J} . □

2.3. Normal forms. Recall that the subset A defined in Definition 1.2 is invariant, does not intersect $\cup_{n \in \mathbb{Z}} f^n(\mathcal{C})$ and satisfies $\mu(A) = 1$. Let

$$\hat{A} := \{ \hat{x} = (x_n)_{n \in \mathbb{Z}} \in A^{\mathbb{Z}}, x_{n+1} = f(x_n) \}$$

and let $\hat{f} : \hat{A} \rightarrow \hat{A}$ be the left shift. We denote $\hat{x}_n := \hat{f}^n(\hat{x})$ for every $n \in \mathbb{Z}$. A function $\varphi_\epsilon : \hat{A} \rightarrow]0, +\infty[$ is ϵ -tempered if $e^{-\epsilon} \varphi_\epsilon(\hat{x}) \leq \varphi_\epsilon(\hat{f}(\hat{x})) \leq e^\epsilon \varphi_\epsilon(\hat{x})$. Let $f_{\hat{x}_n}^{-n}$ be the inverse branch of f^n sending a neighbourhood of x_n to a neighbourhood of x_0 . The following result provides normal forms for those mappings, see [5, 21]. The measure $\hat{\mu}$ stands for the unique \hat{f} -invariant measure on \hat{A} such that $(\pi_0)_* \hat{\mu} = \mu$, where $\pi_0(\hat{x}) := x_0$. Let $B_x(r)$ denote the ball centered at x of radius r in \mathbb{P}^2 .

Theorem 2.6. [5, Proposition 4.3] *For every $\epsilon > 0$ there exist ϵ -tempered functions $\eta_\epsilon, \rho_\epsilon : \hat{A} \rightarrow]0, 1[$, $\beta_\epsilon, L_\epsilon, M_\epsilon : \hat{A} \rightarrow [1, +\infty[$ and a function $N : \hat{A} \rightarrow \mathbb{N}$ satisfying the following properties. There exist injective holomorphic mappings $\xi_{\hat{x}} : B_{x_0}(\eta_\epsilon(\hat{x})) \rightarrow \mathbb{D}^2(\rho_\epsilon(\hat{x}))$ satisfying for every $\hat{x} \in \hat{A}$:*

1. $\xi_{\hat{x}}(x_0) = 0$ and $D_{x_0}\xi_{\hat{x}}(v_s(x_0))$ is the vertical axis in \mathbb{C}^2 ,
 2. $\forall p, q \in B_{x_0}(\eta_\epsilon(\hat{x}))$, $\frac{1}{2} \text{dist}(p, q) \leq \|\xi_{\hat{x}}(p) - \xi_{\hat{x}}(q)\| \leq \beta_\epsilon(\hat{x}) \text{dist}(p, q)$,
- and such that the following diagram commutes for every $n \geq N(\hat{x})$:

$$\begin{array}{ccc}
 B_{x_0}(\eta_\epsilon(\hat{x})) & \xleftarrow{f_{\hat{x}_n}^{-n}} & B_{x_n}(\eta_\epsilon(\hat{x}_n)) \\
 \downarrow \xi_{\hat{x}} & & \downarrow \xi_{\hat{x}_n} \\
 \mathbb{D}^2(\rho_\epsilon(\hat{x})) & \xleftarrow{R_{n, \hat{x}_n}} & \mathbb{D}^2(\rho_\epsilon(\hat{x}_n))
 \end{array}$$

The mappings R_{n, \hat{x}_n} have the following form depending on (λ_1, λ_2) :

1. If $\lambda_1 = \lambda_2 = \lambda$, then R_{n, \hat{x}_n} is a linear mapping satisfying

$$e^{-n(\lambda+\epsilon)} \|(z, w)\| \leq \|R_{n, \hat{x}_n}(z, w)\| \leq e^{-n(\lambda-\epsilon)} \|(z, w)\|.$$

2. If $\lambda_1 = k\lambda_2$ for some $k \geq 2$ then $R_{n, \hat{x}_n}(z, w) = (\alpha_{n, \hat{x}_n} z, \beta_{n, \hat{x}_n} w) + (\gamma_{n, \hat{x}_n} w^k, 0)$.
3. If $\lambda_1 \notin \{k\lambda_2, k \geq 1\}$, then $R_{n, \hat{x}_n}(z, w) = (\alpha_{n, \hat{x}_n} z, \beta_{n, \hat{x}_n} w)$.

Moreover, in the cases 2 and 3, we have

$$e^{-n(\lambda_1+\epsilon)} \leq |\alpha_{n, \hat{x}_n}| \leq e^{-n(\lambda_1-\epsilon)}, \quad |\gamma_{n, \hat{x}_n}| \leq M_\epsilon(\hat{x})e^{-n(\lambda_1-2\epsilon)}$$

and

$$e^{-n(\lambda_2+\epsilon)} \leq |\beta_{n, \hat{x}_n}| \leq e^{-n(\lambda_2-\epsilon)}. \tag{2.2}$$

In particular, if $\lambda_2 < \lambda_1$, the second coordinate of R_{n, \hat{x}_n} has the form $w \mapsto \beta_{n, \hat{x}_n} w$.

2.4. Estimates from pluripotential theory. The following result holds for every endomorphism of \mathbb{P}^2 satisfying $\lambda_1 < 2\lambda_2$. The proof relies on Briend-Duval’s strategy [8] by taking into account $\lambda_2 \geq \frac{1}{2} \log d$: the arguments use the facts that T has continuous potentials, satisfies $f^*T = dT$ and defines μ by the Monge-Ampère equation $\mu = T \wedge T$. Let us set

$$B_n(\rho) := \{x \in \mathbb{P}^2, u \mapsto f^n \circ (x + (D_x f^n)^{-1}(u)) : \mathbb{D}^2(\rho) \rightarrow \mathbb{P}^2 \text{ is injective} \},$$

$$R_n(\tau) := \left\{ x \in \mathbb{P}^2, \|(D_x f^n)^{-1}\|^{-1} \geq \tau^{-1} d^{n/2} \right\}.$$

For the definition of $B_n(\rho)$, we used charts of $T\mathbb{P}^2$ so that the mapping $x + (D_x f^n)^{-1}$ is defined on $\mathbb{D}^2(\rho)$ and takes its values in a neighbourhood of x in \mathbb{P}^2 .

Theorem 2.7. [6, Propositions 1 and 2] *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. If $\lambda_1 < 2\lambda_2$, then for every $\beta > 0$,*

1. $\exists \rho > 0, \forall n \geq 1, \mu(B_n(\rho)) \geq 1 - \beta$,
2. $\exists \rho > 0, \forall n \geq 1, \forall \tau > 0, \mu(B_n(\rho) \cap R_n(\tau)) \geq 1 - \beta - (\rho\tau)^{-2}$.
3. In particular, $\forall a > 0, \exists \tau > 0, \forall n \geq 1, \mu(R_n(\tau)) \geq 1 - a$.

The following Corollary will be used to prove Proposition 5.2.

Corollary 2.8. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$ satisfying $\lambda_2 < \lambda_1 < 2\lambda_2$. For every $\beta > 0$ let $\rho > 0$ given by Theorem 2.7. There exists a Borel set H of full μ -measure satisfying for every $x \in H$, there exists $n(x) \geq 1$ such that*

$$\forall n \geq n(x), \quad x \notin B_n(\rho) \text{ or } \|D_x f^n(\vec{v}_s(x))\| \geq \frac{d^{n/2}}{n}.$$

Proof. According to Theorem 2.7, we have $\sum_{n \geq 1} \mu(B_n(\rho) \cap R_n(n)^c) \leq \rho^{-2} \sum_{n \geq 1} \frac{1}{n^2}$. By Borel-Cantelli's Lemma, there exists H such that $\mu(H) = 1$ and

$$\forall x \in H, \exists n(x) \geq 1, \forall n \geq n(x), \quad x \in B_n(\rho)^c \cup R_n(n).$$

Let $x \in H$. If $x \in B_n(\rho)^c$, the proof is complete. If not, $x \in R_n(n)$, and since $\vec{v}_s(x)$ is unitary we get $\|D_x f^n(\vec{v}_s(x))\| \geq \|(D_x f^n)^{-1}\|^{-1} \geq \frac{d^{n/2}}{n}$. \square

3. A criterion for bounded Radon-Nikodym derivatives. The following theorem gives sufficient conditions ensuring that a point x belongs to the set R defined in Definition 1.2. The integer $N(\hat{x})$ is defined in Theorem 2.6.

Theorem 3.1. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. Assume that $\lambda_1 > \lambda_2$ and let $\hat{x} \in \hat{A}$. If there exist $\eta_0, M_1, M_2 > 0$ and an increasing sequence $(n_j)_{j \in \mathbb{N}}$ of integers such that $n_0 = 0$ and $n_1 \geq N(\hat{x})$ satisfying:*

1. $\eta_\epsilon(\hat{x}_{n_j}) \geq 4\eta_0, \beta_\epsilon(\hat{x}_{n_j}) \leq e^{M_2}$,
2. $e^{-M_1} d^{n_j/2} \leq \|D_x f^{n_j}(\vec{v}_s(\pi_0(\hat{x})))\| \leq e^{M_1} d^{n_j/2}$,

then $x := \pi_0(\hat{x}) \in R$: there exists a holomorphic disc $\xi_x : \mathbb{D} \rightarrow \mathbb{P}^2$ satisfying $\xi_x(0) = x, [\xi'_x(0)] = v_s(x)$ and

$$\liminf_{r \rightarrow 0} \frac{\xi_x^* T(\mathbb{D}(r))}{\text{Leb}(\mathbb{D}(r))} < \infty. \tag{3.1}$$

Proof. Let us apply Theorem 2.6 which gives normal forms for the inverse branches of f . Since $\eta_\epsilon(\hat{x}_{n_j}) \geq 4\eta_0$, the image of $B_{x_{n_j}}(\eta_\epsilon(\hat{x}_{n_j}))$ by $\xi_{\hat{x}_{n_j}}$ contains $\mathbb{D}^2(2\eta_0)$. Similarly, the image of $B_{x_0}(\eta_\epsilon(\hat{x}))$ by $\xi_{\hat{x}}$ contains $\mathbb{D}^2(2\eta_0)$. Let V be the vertical disc

$$V : \begin{array}{ccc} \mathbb{D}(2\eta_0) & \rightarrow & \mathbb{D}(2\eta_0) \times \mathbb{D}(2\eta_0) \\ w & \mapsto & (0, w) \end{array},$$

and let us set

$$\tilde{\xi}_0 := (\xi_{\hat{x}})^{-1} \circ V \quad \text{and} \quad \tilde{\xi}_{n_j} := (\xi_{\hat{x}_{n_j}})^{-1} \circ V.$$

By pulling back $(f^{n_j})^* T = d^{n_j} T$ by $\tilde{\xi}_0$, we get

$$(f^{n_j} \circ \tilde{\xi}_0)^* T = d^{n_j} \tilde{\xi}_0^* T \text{ on } \mathbb{D}(\eta_0). \tag{3.2}$$

Let us observe that

$$f^{n_j} \circ \tilde{\xi}_0 = f^{n_j} \circ (\xi_{\hat{x}})^{-1} \circ V = (\xi_{\hat{x}_{n_j}})^{-1} \circ R_{n_j, \hat{x}_{n_j}}^{-1} \circ V,$$

where the first equality comes from the definition of $\tilde{\xi}_0$ and the second one from Theorem 2.6. If $\beta_{n_j} : \mathbb{C} \rightarrow \mathbb{C}$ denotes the multiplication by $\beta_{n_j, \hat{x}_{n_j}}$, we also have the relation

$$R_{n_j, \hat{x}_{n_j}}^{-1} \circ V = V \circ \beta_{n_j}^{-1},$$

since the second coordinate of $R_{n_j, \hat{x}_{n_j}}$ is linear. Hence we get:

$$f^{n_j} \circ \tilde{\xi}_0 = \tilde{\xi}_{n_j} \circ \beta_{n_j}^{-1} \text{ on } \beta_{n_j}(\mathbb{D}(\eta_0)).$$

Equation (3.2) restricted to $\beta_{n_j}(\mathbb{D}(\eta_0)) \subset \mathbb{D}(\eta_0)$ then implies:

$$\tilde{\xi}_{n_j}^* T = d^{n_j} (\beta_{n_j}^{-1})_* \tilde{\xi}_0^* T \text{ on } \mathbb{D}(\eta_0),$$

which gives

$$\tilde{\xi}_{n_j}^* T(\mathbb{D}(\eta_0)) = d^{n_j} \tilde{\xi}_0^* T(\mathbb{D}(\eta_0 \cdot |\beta_{n_j, \hat{x}_{n_j}}|)). \tag{3.3}$$

Now we bound from below the right hand side of Equation (3.3). This is where we use the second hypothesis of Theorem 3.1.

Lemma 3.2. *We have $|\beta_{n_j, \hat{x}_{n_j}}| \in [d^{-n_j/2}e^{-M}, d^{-n_j/2}e^M]$, where $M := M_1 + M_2 + \ln 2$.*

Proof. By Theorem 2.6, we can write $f^{n_j} = (\xi_{\hat{x}_{n_j}})^{-1} \circ R_{n_j, \hat{x}_{n_j}}^{-1} \circ \xi_{\hat{x}}$. We study the differential of that composition. By using

$$\forall \vec{v} \in T_{f^{n_j}(x)}\mathbb{P}^2, \quad \frac{1}{2} \|\vec{v}\| \leq \|D_{f^{n_j}(x)}\xi_{\hat{x}_{n_j}}(\vec{v})\| \leq e^{M_2} \|\vec{v}\|$$

for $i = 0$ and $i = j$ (provided by Theorem 2.6) and the fact that the differential of $\xi_{\hat{x}}$ (resp. $\xi_{\hat{x}_{n_j}}$) at x (resp. $f^{n_j}(x)$) sends the line $v_s(x) \subset T_x\mathbb{P}^2$ (resp. $v_s(f^{n_j}(x)) \subset T_{f^{n_j}(x)}\mathbb{P}^2$) on the vertical axis, we obtain

$$\|D_x f^{n_j}(\vec{v}_s(x))\| (2e^{M_2})^{-1} \leq \left| \beta_{n_j, \hat{x}_{n_j}} \right|^{-1} \leq \|D_x f^{n_j}(\vec{v}_s(x))\| (2e^{M_2}).$$

Now we use the assumption $e^{-M_1} d^{n_j/2} \leq \|D_x f^{n_j}(\vec{v}_s(x))\| \leq e^{M_1} d^{n_j/2}$ to deduce

$$d^{-n_j/2} e^{-M_1} (2e^{M_2})^{-1} \leq \left| \beta_{n_j, \hat{x}_{n_j}} \right| \leq d^{-n_j/2} e^{M_1} (2e^{M_2}),$$

which concludes the proof. □

The right hand side term of Equation (3.3) hence satisfies:

$$d^{n_j} \tilde{\xi}_0^* T(\mathbb{D}(\eta_0 \cdot |\beta_{n_j, \hat{x}_{n_j}}|)) \geq \frac{\tilde{\xi}_0^* T(\mathbb{D}(\eta_0 d^{-n_j/2} e^{-M}))}{d^{-n_j}} = c_0 \frac{\tilde{\xi}_0^* T(\mathbb{D}(\eta_0 d^{-n_j/2} e^{-M}))}{\pi(\eta_0 d^{-n_j/2} e^{-M})^2}, \tag{3.4}$$

where $c_0 = \pi(\eta_0 e^{-M})^2$. By setting $r_j = \eta_0 d^{-n_j/2} e^{-M}$, we obtain:

$$d^{n_j} \tilde{\xi}_0^* T(\mathbb{D}(\eta_0 \cdot |\beta_{n_j, \hat{x}_{n_j}}|)) \geq c_0 \frac{\tilde{\xi}_0^* T(\mathbb{D}(r_j))}{\text{Leb}(\mathbb{D}(r_j))}.$$

Now Lemma 3.3 below bounds the left hand side term of Equation (3.3), hence

$$\liminf_{r \rightarrow 0} \frac{\tilde{\xi}_0^* T(\mathbb{D}(r))}{\text{Leb}(\mathbb{D}(r))} < \infty.$$

To conclude the proof of Theorem 3.1, we set $\xi_x : \mathbb{D} \rightarrow \mathbb{P}^2$ by $\xi_x(z) := \tilde{\xi}_0(\eta_0 z)$. □

Lemma 3.3. *There exists a constant $M_3 > 0$ such that for every $j \geq 0$,*

$$\tilde{\xi}_{n_j}^* T(\mathbb{D}(\eta_0)) \leq M_3.$$

Proof. Let $(B_i)_{i \in I}$ be a finite open covering of \mathbb{P}^2 by small balls, such that T admits for every $i \in I$ a bounded continuous plurisubharmonic potential G_i on the $4\eta_0$ -tubular neighbourhood \tilde{B}_i of B_i . Let $\|G\|_\infty := \max_{i \in I} \|G_i\|_\infty$ and let $\psi : \mathbb{D}(2\eta_0) \rightarrow \mathbb{R}^+$ be a function with compact support which is equal to 1 on $\mathbb{D}(\eta_0)$. Let us fix $j \geq 0$ and let $i \in I$ be such that $x_{n_j} \in B_i$, hence $B_{x_{n_j}}(4\eta_0) \subset \tilde{B}_i$. Since $\tilde{\xi}_{n_j} = (\xi_{\hat{x}_{n_j}})^{-1} \circ V$, we get

$$\tilde{\xi}_{n_j}(\mathbb{D}(2\eta_0)) \subset (\xi_{\hat{x}_{n_j}})^{-1}(\mathbb{D}^2(2\eta_0)) \subset B_{x_{n_j}}(4\eta_0).$$

We deduce

$$\tilde{\xi}_{n_j}^* T(\mathbb{D}(\eta_0)) \leq \int_{\mathbb{D}(2\eta_0)} \Delta(G_\alpha \circ \tilde{\xi}_{n_j}) \psi = \int_{\mathbb{D}(2\eta_0)} G_\alpha \circ \tilde{\xi}_{n_j} \Delta \psi \leq \|G\|_\infty \max_{\mathbb{D}(2\eta_0)} \Delta \psi.$$

We complete the proof by setting $M_3 := \|G\|_\infty \max_{\mathbb{D}(2\eta_0)} \Delta \psi$. □

4. Proof of Theorem 1.3.

4.1. Decomposable currents. This subsection is borrowed from Dujardin's article [14]. Let ω be the Fubini-Study $(1, 1)$ -form on \mathbb{P}^2 and let S be a closed positive current of bidegree $(1, 1)$ on \mathbb{P}^2 . Let $\sigma_S = S \wedge \omega$ denote the trace measure of S , this is a positive measure on \mathbb{P}^2 . We shall denote $S(\varphi) = \int_{\mathbb{P}^2} S \wedge \varphi$ for every $(1, 1)$ -test form φ on \mathbb{P}^2 . Locally, one can write S as

$$S = \sum_{j,k \in \{1,2\}} S_{j,k} i dz_j \wedge d\bar{z}_k. \quad (4.1)$$

where $S_{1,1}$ and $S_{2,2}$ are positive measures and $S_{1,2} = \overline{S_{2,1}}$ are complex measures. Let (e_1, e_2) be a basis of \mathbb{C}^2 . Let us recall that a $(1, 1)$ -positive vector of \mathbb{C}^2 is an element of $\Lambda^{1,1}\mathbb{C}^2$ of the form $i \sum_{j,k \in \{1,2\}} t_{j,k} e_j \wedge \bar{e}_k$ where $(t_{j,k})_{(j,k)}$ is a positive hermitian matrix. A positive vector of \mathbb{C}^2 is decomposable if it is equal to $iu \wedge \bar{u}$ for some $u \in \mathbb{C}^2$. In Equation (4.1), since S is a positive current, the measures $|S_{1,2}|$ and $|S_{2,1}|$ are dominated by the trace measure $\sigma_S = S_{1,1} + S_{2,2}$. By the Radon-Nikodym Theorem, there exists a measurable field Λ_S of $(1, 1)$ -positive covectors of trace 1 such that

$$S = \Lambda_S \sigma_S.$$

By duality, there exists a measurable field t_S of $(1, 1)$ -positive vectors of trace 1 such that

$$\forall \eta \text{ (1, 1) - test form on } \mathbb{P}^2, \quad S(\eta) = \int_{\mathbb{P}^2} \langle t_S, \eta \rangle \sigma_S. \quad (4.2)$$

By the Lebesgue density Theorem, there exists $E \subset \text{Supp } S$ such that $S \wedge \omega(E) = 1$ and for every $x \in E$,

$$\lim_{r \rightarrow 0} \frac{1}{\sigma_S(B_x(r))} \int_{B_x(r)} t_S(y) \sigma_S(y) = t_S(x).$$

We say that S is decomposable at $x \in E$ if the $(1, 1)$ -positive vector $t_S(x)$ is decomposable. For instance, if $S = [L]$ where L is a smooth curve, then $t_S(x) = i\vec{u}_x \wedge \overline{\vec{u}_x}$ where \vec{u}_x is a unitary tangent vector of L at x . The laminar currents are also examples of decomposable currents.

In [14], Dujardin established the following nice property: if f is not a Lattès map, then the Green current T is decomposable σ_T -almost everywhere, in particular it carries a directional information. This is the content of the following result.

Theorem 4.1. [14, Theorem 3.3] *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$ and let T be its Green current. If f is not a Lattès map, then there exists $E' \subset E$ of full σ_T -measure satisfying the following property. For every $x \in E'$, there exists $\vec{v}_x \in T_x \mathbb{P}^2$ satisfying $\|\vec{v}_x\| = 1$ and*

$$t_T(x) = i\vec{v}_x \wedge \overline{\vec{v}_x}.$$

Moreover the vectors \vec{v}_x satisfy the following property.

Proposition 4.2. [14, Section 3.2] *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$. We assume that f is not a Lattès map (hence $t_T(x) = i\vec{v}_x \wedge \overline{\vec{v}_x}$ σ_T -almost everywhere, with $\|\vec{v}_x\| = 1$). Then,*

$$\forall n \geq 1, \quad \int_{\mathbb{P}^2} \|D_x f^n(\vec{v}_x)\|^2 d\sigma_T(x) \leq d^n.$$

Markov's inequality then implies:

$$\forall n \geq 1, \forall a > 0, \quad (\sigma_T) \left\{ x \in \mathbb{P}^2, \|D_x f^n(\vec{v}_x)\|^2 \geq ad^n \right\} \leq \frac{1}{a}.$$

Proof. For σ_T -almost every x , we have

$$\|D_x f^n(\vec{v}_x)\|^2 \leq \text{Trace} (D_x f^n)_*(i\vec{v}_x \wedge \overline{\vec{v}_x}).$$

We use $t_T(x) = i\vec{v}_x \wedge \overline{\vec{v}_x}$ and the definition of the trace to get:

$$\begin{aligned} \|D_x f^n(\vec{v}_x)\|^2 &\leq \langle (D_x f^n)_*(t_T(x)), \omega(f^n(x)) \rangle \\ &\leq \langle t_T(x), (D_x f^n)^* \omega(f^n(x)) \rangle \end{aligned}$$

since $D_x f^n$ is invertible σ_T -almost everywhere. By integrating this inequality on \mathbb{P}^2 :

$$\int_{\mathbb{P}^2} \|D_x f^n(\vec{v}_x)\|^2 \sigma_T(x) \leq \int_{\mathbb{P}^2} \langle t_T(x), (D_x f^n)^* \omega(f^n(x)) \rangle \sigma_T(x).$$

By using Equation (4.2), we obtain

$$\int_{\mathbb{P}^2} \|D_x f^n(\vec{v}_x)\|^2 \sigma_T(x) \leq \int_{\mathbb{P}^2} T \wedge (f^n)^* \omega = d^n,$$

where the last equality uses the fact that $(f^n)^* \omega$ is cohomologous to $d^n \omega$, see for instance Sections 1.2 and A.3 of [13]. \square

4.2. Application of Theorem 3.1 to prove Theorem 1.3.

Proposition 4.3. *Assume that f is not a Lattès map (hence $t_T(x) = i\vec{v}_x \wedge \overline{\vec{v}_x}$ σ_T -almost everywhere, with $\|\vec{v}_x\| = 1$). We also assume that $\mu \ll \sigma_T$. Then, there exist $q \geq 1$ and $\gamma > 0$ such that*

$$\forall n \geq 1, \quad \mu \left\{ x \in \mathbb{P}^2, \|D_x f^n(\vec{v}_x)\|^2 \leq \frac{4}{\gamma} d^n \right\} \geq \frac{\gamma}{4q}.$$

Proof. Let us write $\mu = \varphi \sigma_T$ where $\varphi \in L^1(\sigma_T)$ and $\varphi \geq 0$. Let us set $\gamma := \sigma_T \{ \varphi > 0 \}$. It satisfies $\gamma > 0$ since $1 = \mu(\mathbb{P}^2) = \mu \{ \varphi > 0 \} = \int_{\{ \varphi > 0 \}} \varphi \sigma_T$. Moreover $\sigma_T \left\{ \frac{1}{q} \leq \varphi \right\}$ tends to $\sigma_T \{ 0 < \varphi \} = \gamma$ by Lebesgue convergence Theorem. Let us set $q \geq 1$ such that

$$\sigma_T \left\{ \frac{1}{q} \leq \varphi \right\} \geq \gamma/2. \tag{4.3}$$

For every $n \geq 1$, we denote

$$A_n := \left\{ x \in \mathbb{P}^2, \|D_x f^n(\vec{v}_x)\|^2 \leq \frac{4}{\gamma} d^n \right\}.$$

We want to show $\mu(A_n) \geq \frac{\gamma}{4q}$. Proposition 4.2 gives with $a = 4/\gamma$

$$\forall n \geq 1, \quad \sigma_T(A_n) \geq 1 - \gamma/4. \tag{4.4}$$

Equations (4.3) and (4.4) imply $\sigma_T(A_n \cap \left\{ \frac{1}{q} \leq \varphi \right\}) \geq \frac{\gamma}{4}$. Coming back to μ ,

$$\mu(A_n) \geq \mu(A_n \cap \left\{ \frac{1}{q} \leq \varphi \right\}) = \int_{A_n \cap \{ \frac{1}{q} \leq \varphi \}} \varphi \sigma_T.$$

We deduce

$$\mu(A_n) \geq \frac{1}{q} (\sigma_T)(A_n \cap \left\{ \frac{1}{q} \leq \varphi \right\}) \geq \frac{1}{q} \frac{\gamma}{4},$$

which is the desired lower bound. \square

Now let us prove Theorem 1.3. We assume that $\mu \ll \sigma_T$ and $\lambda_2 < \lambda_1 < 2\lambda_2$. Let

$$B_n := \left\{ x \in \mathbb{P}^2, \frac{1}{\tau_0^2} d^n \leq \|D_x f^n(\vec{v}_x)\|^2 \leq \frac{4}{\gamma} d^n \right\}.$$

By combining Proposition 4.3 and the third item of Theorem 2.7 (with $a = \frac{\gamma}{8q}$ and using $\|D_x f^n(\vec{v}_x)\| \geq \|(D_x f^n)^{-1}\|^{-1}$), we get $\mu(B_n) \geq \frac{\gamma}{8q}$. The set $\hat{B}_n := \pi_0^{-1}(B_n)$ then satisfies $\hat{\mu}(\hat{B}_n) \geq \frac{\gamma}{8q}$. Let M_2 be large enough and η_0 be small enough such that

$$\hat{E} := \{ \beta_\epsilon(\hat{x}) \leq e^{M_2}, \eta_\epsilon(\hat{x}) \geq 4\eta_0 \}$$

satisfies $\hat{\mu}(\hat{E} \cap \hat{f}^{-n}(E)) \geq 1 - \frac{\gamma}{16q}$. Therefore the $\hat{\mu}$ -measure of $\hat{C}_n := \hat{B}_n \cap \hat{E} \cap \hat{f}^{-n}(\hat{E})$ is larger than $\frac{\gamma}{16q}$, and the same property holds for $\hat{C} := \limsup \hat{C}_n$.

Lemma 4.4. *Every $\hat{x} \in \hat{C}$ satisfies the hypothesis of Theorem 3.1.*

Proof. Let $\hat{x} \in \hat{C}$ and $x = \pi_0(\hat{x})$. By definition of \hat{C} , there exists $(n_j)_j$ depending on \hat{x} such that $n_0 = 0$, $n_1 \geq N(\hat{x})$ and $\hat{x} \in \hat{C}_{n_j}$ for every $j \geq 0$. In particular, $\eta_\epsilon(\hat{x}_{n_j}) \geq 4\eta_0$, $\beta_\epsilon(\hat{x}_{n_j}) \leq e^{M_2}$, and

$$\frac{1}{\tau_0^2} d^{n_j} \leq \|D_x f^{n_j}(\vec{v}_x)\|^2 \leq \frac{4}{\gamma} d^{n_j}.$$

Those estimates combined with the definition of the stable Oseledec direction given in Theorem 2.1 imply $v_s(x) = v_x$ (recall that $\lambda_2 < \lambda_1$). By taking M_1 large enough such that $e^{-2M_1} \leq \frac{1}{\tau_0^2} \leq \frac{4}{\gamma} \leq e^{2M_1}$, we get

$$e^{-M_1} d^{n_j/2} \leq \|D_x f^{n_j}(\vec{v}_s(x))\| \leq e^{M_1} d^{n_j/2},$$

which completes the proof of the Lemma. \square

Lemma 4.4 ensures that $C := \pi_0(\hat{C}) \subset R$. But C has positive μ -measure, since $\mu(C) \geq \hat{\mu}(\hat{C}) \geq \frac{\gamma}{16q}$. Finally $\mu(R) > 0$, completing the proof of Theorem 1.3.

5. Proof of Theorem 1.4.

5.1. Central Limit Theorem. The dynamical system (\mathbb{P}^2, f, μ) satisfies a Central Limit Theorem, see [9, 11, 12, 16].

Theorem 5.1. *Let f be a holomorphic endomorphism of \mathbb{P}^2 of degree $d \geq 2$ and let μ be its equilibrium measure. Let $\phi : \mathcal{J} \rightarrow \mathbb{R}$ be a Hölder continuous function such that $\int \phi d\mu = 0$ and let $S_n(\phi) := \sum_{i=0}^{n-1} \phi \circ f^i$. Then the asymptotic variance*

$$\sigma_\phi := \lim_n \frac{1}{\sqrt{n}} \|S_n(\phi)\|_2$$

exists and the following alternative holds:

1. if $\sigma_\phi = 0$ there exists $u \in L^2(\mu)$ such that $\phi = u \circ f - u$.
2. if $\sigma_\phi > 0$ then $\frac{1}{\sqrt{n}} S_n(\phi)$ converges in law towards the Gaussian $\mathcal{N}(0, \sigma_\phi)$.

5.2. **The asymptotic variance of ψ vanishes.** We assume that $\lambda_2 < \lambda_1$ and consider

$$\psi(x) = \log \|D_x f(\vec{v}_s(x))\|$$

where $v_s(x)$ is the stable direction at $x \in A$. By Proposition 2.5, if $\mathcal{J} \cap \mathcal{C} = \emptyset$ and if v_s is Hölder continuous on A , then ψ extends to a Hölder continuous function on \mathcal{J} . If moreover $\lambda_2 = \frac{1}{2} \log d$, then $\psi_0 := \psi - \frac{1}{2} \log d$ satisfies $\int \psi_0 d\mu = 0$ by Lemma 2.2.

Proposition 5.2. *Assume that $\lambda_2 = \frac{1}{2} \log d$ and that $\lambda_2 < \lambda_1 < 2\lambda_2$. Assume also that $\mathcal{J} \cap \mathcal{C} = \emptyset$ and that v_s is Hölder continuous on A . Then $\sigma_{\psi_0} = 0$. In particular, there exists $u \in L^2(\mu)$ such that for μ -almost every x :*

$$\log \|D_x f(\vec{v}_s(x))\| = \lambda_2 + u \circ f - u. \tag{5.1}$$

Proof. Assume to the contrary that $\sigma_{\psi_0} > 0$. By Lemma 2.2 one can write

$$\frac{1}{\sigma_{\psi_0} \sqrt{n}} S_n(\psi_0) = \frac{1}{\sigma_{\psi_0} \sqrt{n}} \left(\log \|D_x f^n(\vec{v}_s(x))\| - n \frac{\log d}{2} \right).$$

Theorem 5.1 ensures that $\frac{1}{\sigma_{\psi_0} \sqrt{n}} S_n(\psi_0)$ converges in law towards $\mathcal{N}(0, 1)$. By setting

$$\beta_0 := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{-1} e^{-u^2/2} du \quad \text{and} \quad G_n := \{x \in \mathcal{J} \mid S_n(\psi_0)(x) \leq -\sigma_{\psi_0} \sqrt{n}\},$$

we get

$$\lim_{n \rightarrow \infty} \mu(G_n) = \beta_0.$$

Let $N \geq 1$ such that $\mu(G_n) \geq \frac{\beta_0}{2}$ for every $n \geq N$. Let $\rho > 0$ provided by Theorem 2.7 with $\beta = \frac{\beta_0}{4}$. We have $\mu(B_n(\rho)) \geq 1 - \frac{\beta_0}{4}$ for every $n \geq 1$. Let us define

$$F_n := B_n(\rho) \cap G_n, \quad F := \limsup F_n.$$

For every $n \geq N$, we have $\mu(F_n) \geq \frac{\beta_0}{4}$. We also have $\mu(F) \geq \frac{\beta_0}{4}$. Let H be given by Corollary 2.8. Since $\mu(H) = 1$ and $\mu(F) > 0$, there exists $x \in F \cap H$. Let $(n_i)_i$ be an increasing sequence of integers such that $x \in F_{n_i}$ for every $i \in \mathbb{N}$. Since $F_{n_i} \subset G_{n_i}$, we have:

$$S_{n_i}(\psi_0)(x) = \log \|D_x f^{n_i}(\vec{v}_s(x))\| - n_i \frac{\log d}{2} \leq -\sigma_{\psi_0} \sqrt{n_i},$$

which implies

$$d^{n_i/2} e^{-\sigma_{\psi_0} \sqrt{n_i}} \geq \|D_x f^{n_i}(\vec{v}_s(x))\|. \tag{5.2}$$

But x belongs to H and to $F_{n_i} \subset B_{n_i}(\rho)$, hence Corollary 2.8 ensures for every $n_i > n(x)$:

$$\|D_x f^{n_i}(\vec{v}_s(x))\| \geq \frac{d^{n_i/2}}{n_i}. \tag{5.3}$$

By combining (5.2) and (5.3), we obtain $e^{-\sigma_{\psi_0} \sqrt{n_i}} \geq \frac{1}{n_i}$ for every i large enough. This contradiction implies $\sigma_{\psi} = 0$ and completes the proof of Proposition 5.2. \square

5.3. Application of Theorem 3.1 to prove Theorem 1.4. By summing Equation (5.1) over an orbit of length n and using Lemma 2.2, we obtain:

$$\forall x \mu - a.e., \forall n \geq 1, \quad \log \|D_x f^n(\vec{v}_s(x))\| = n\lambda_2 + u \circ f^n - u.$$

The μ -measure of $\{|u| \leq R\}$ tends to 1 when R tends to infinity. The same property holds for $\{|u| \leq R\} \cap \{|u \circ f^n| \leq R\}$ since μ is invariant. Therefore, there exists $R > 0$ such that

$$B_n := \left\{ x \in \mathbb{P}^2, d^{n/2} e^{-2R} \leq \|D_x f^n(\vec{v}_s(x))\| \leq d^{n/2} e^{2R} \right\}$$

satisfies $\mu(B_n) \geq 1/2$ for every $n \geq 1$. Hence $\hat{B}_n := \pi_0^{-1}(B_n)$ satisfies $\hat{\mu}(\hat{B}_n) \geq 1/2$. Now, to complete the proof of Theorem 1.4, it suffices to repeat the end of Section 4.2, namely to introduce the subsets

$$\hat{E} := \left\{ \beta_\epsilon(\hat{x}) \leq e^{M_2}, \eta_\epsilon(\hat{x}) \geq 4\eta_0 \right\},$$

$\hat{C}_n := \hat{B}_n \cap \hat{E} \cap \hat{f}^{-n}(\hat{E})$, $\hat{C} := \limsup \hat{C}_n$ and to apply similarly Theorem 3.1.

Acknowledgments. We thank the referees for their cautious readings. Their suggestions enable us to improve the exposition, with for instance the addition of Section 1.4 and the development of Section 2.2.

REFERENCES

- [1] L. Barreira and Y. Pesin, *Nonuniform Hyperbolicity*, Encyclopedia of Mathematics and its Applications, 115, Cambridge University Press, Cambridge, 2007.
- [2] E. Bedford and M. Jonsson, *Dynamics of regular polynomial endomorphisms of \mathbf{C}^k* , *Amer. J. Math.*, **122** (2000), 153–212.
- [3] F. Berteloot and J.-J. Loeb, *Spherical hypersurfaces and Lattès rational maps*, *J. Math. Pures Appl. (9)*, **77** (1998), 655–666.
- [4] F. Berteloot and J.-J. Loeb, *Une caractérisation géométrique des exemples de Lattès de \mathbb{P}^k* , *Bull. Soc. Math. France*, **129** (2001), 175–188.
- [5] F. Berteloot, C. Dupont and L. Molino, *Normalization of bundle holomorphic contractions and applications to dynamics*, *Ann. Inst. Fourier (Grenoble)*, **58** (2008), 2137–2168.
- [6] F. Berteloot and C. Dupont, *Une caractérisation des endomorphismes de Lattès par leur mesure de Green*, *Comment. Math. Helv.*, **80** (2005), 433–454.
- [7] I. Binder and L. DeMarco, *Dimension of pluriharmonic measure and polynomial endomorphisms of \mathbf{C}^n* , *Int. Math. Res. Not.*, **2003** (2003), 613–625.
- [8] J.-Y. Briend and J. Duval, *Exposants de Liapounoff et distribution des points périodiques d'un endomorphisme de \mathbf{CP}^k* , *Acta Math.*, **182** (1999), 143–157.
- [9] S. Cantat and S. Le Borgne, *Théorème limite central pour les endomorphismes holomorphes et les correspondances modulaires*, *Int. Math. Res. Not.*, **2005** (2005), 3479–3510.
- [10] T.-C. Dinh and C. Dupont, *Dimension de la mesure d'équilibre d'applications méromorphes*, *J. Geom. Anal.*, **14** (2004), 613–627.
- [11] T.-C. Dinh, V.-A. Nguyen and N. Sibony, *Exponential estimates for plurisubharmonic functions and stochastic dynamics*, *J. Differential Geom.*, **84** (2010), 465–488.
- [12] T.-C. Dinh and N. Sibony, *Decay of correlations and the central limit theorem for meromorphic maps*, *Comm. Pure Appl. Math.*, **59** (2006), 754–768.
- [13] T.-C. Dinh and N. Sibony, *Dynamics in several complex variables: Endomorphisms of projective spaces and polynomial-like mappings*, *Holomorphic Dynamical Systems*, Lecture Notes in Math., 1998, Springer, Berlin, 2010, 165–294.
- [14] R. Dujardin, *Fatou directions along the Julia set for endomorphisms of \mathbf{CP}^k* , *J. Math. Pures Appl. (9)*, **98** (2012), 591–615.
- [15] C. Dupont, *Exemples de Lattès et domaines faiblement sphériques de \mathbf{C}^n* , *Manuscripta Math.*, **111** (2003), 357–378.
- [16] C. Dupont, *Bernoulli coding map and almost sure invariance principle for endomorphisms of \mathbb{P}^k* , *Probab. Theory Related Fields*, **146** (2010), 337–359.

- [17] C. Dupont, [On the dimension of invariant measures of endomorphisms of \$\mathbb{C}\mathbb{P}^k\$](#) , *Math. Ann.*, **349** (2011), 509–528.
- [18] C. Dupont and J. Taffin, Dynamics of fibered endomorphisms of \mathbb{P}^k , to appear in *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, [arXiv:1811.06909](#).
- [19] J. E. Fornæss and N. Sibony, [Hyperbolic maps on \$\mathbb{P}^2\$](#) , *Math. Ann.*, **311** (1998), 305–333.
- [20] M. Jonsson, [Dynamics of polynomial skew products on \$\mathbb{C}^2\$](#) , *Math. Ann.*, **314** (1999), 403–447.
- [21] M. Jonsson and D. Varolin, [Stable manifolds of holomorphic diffeomorphisms](#), *Invent. Math.*, **149** (2002), 409–430.
- [22] M. Klimek, *Pluripotential Theory*, London Mathematical Society Monographs, New Series, 6, The Clarendon Press, Oxford University Press, New York, 1991.
- [23] F. Ledrappier, Quelques propriétés ergodiques des applications rationnelles, *C. R. Acad. Sci. Paris Sér. I Math.*, **299** (1984), 37–40.
- [24] V. Mayer, [Comparing measures and invariant line fields](#), *Ergodic Theory Dynam. Systems*, **22** (2002), 555–570.
- [25] Y. B. Pesin, *Lectures on Partial Hyperbolicity and Stable Ergodicity*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2004.
- [26] N. Sibony, Dynamique des applications rationnelles de \mathbb{P}^k , in *Dynamique et Géométrie Complexes (Lyon, 1997)*, Panor. Synthèses, 8, Soc. Math. France, Paris, 1999, 97–185.
- [27] A. Zdunik, [Parabolic orbifolds and the dimension of the maximal measure for rational maps](#), *Invent. Math.*, **99** (1990), 627–649.

Received June 2019; revised January 2020.

E-mail address: christophe.dupont@univ-rennes1.fr

E-mail address: axel.rogue@univ-rennes1.fr