

# Circulating IL-4, IFN $\gamma$ and IL-17 conventional and Innate-like T-cell producers in adult asthma

Maria Leite-De-Moraes, Renata Belo, Céline Dietrich, David Soussan, Michel

Aubier, Marina Pretolani

## ▶ To cite this version:

Maria Leite-De-Moraes, Renata Belo, Céline Dietrich, David Soussan, Michel Aubier, et al.. Circulating IL-4, IFN $\gamma$  and IL-17 conventional and Innate-like T-cell producers in adult asthma. Allergy, In press, 10.1111/all.14474 . hal-02937423

# HAL Id: hal-02937423 https://hal.science/hal-02937423

Submitted on 6 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Circulating IL-4, IFNy and IL-17 conventional and
2	Innate-like T cell producers in adult asthma
3	Maria Leite-de-Moraes, PhD <sup>a,b</sup> , Renata Belo, PharmD, PhD <sup>a,b</sup> , Céline Dietrich, MSc <sup>a,b</sup> ,
4	David Soussan, MSc <sup>c,d,e,f</sup> , Michel Aubier, MD <sup>c,d,e,f</sup> and Marina Pretolani, PharmD, PhD <sup>c,d,e,f</sup>
5	From:
6	<sup>a</sup> Laboratory of Immunoregulation and Immunopathology, Institut Necker-Enfants Malades
7	(INEM), CNRS UMR8253 and Inserm UMR1151, Paris, France;
8	<sup>b</sup> Université Paris Descartes Sorbonne Paris Cité, Université de Paris, Paris, France;
9	<sup>c</sup> Inserm UMR1152, Physiopathologie et Epidémiologie des Maladies Respiratoires, Paris,
10	France;
11	<sup>d</sup> Université Paris Diderot, Université de Paris, Faculté de Médecine, site Bichat, Paris,
12	France;
13	<sup>e</sup> Laboratory of Excellence INFLAMEX, Université Sorbonne Paris-Cité;
14	<sup>f</sup> Département Hospitalo-Universitaire FIRE.
15	Corresponding author:
16	Maria Leite-de-Moraes, Institut Necker-Enfants Malades, CNRS UMR 8253 and Inserm
17	UMR1151, Faculté de Médecine Necker, 156 rue de Vaugirard, 75015, Paris, France. e-mail:
18	maria.leite-de-moraes@parisdescartes.fr
19	
20	Short title: Conventional and ILT cells in adult asthma

# 21 Disclosure of potential conflict of interest:

- 22 M. Pretolani's institution has received funding from AstraZeneca and Roche. The other authors
- 23 declare that they have no relevant conflicts of interest.

24	Authorship Contributions:
25	M.L.M, M.A., M.P. designed the research.
26	M.L.M, M.P., R.B. performed the research.
27	M.L.M, M.P., R.B., D.S. analysed the data.
28	M.L.M, M.P. wrote the manuscript
29	
30	Text Count
31	Word count in the body of manuscript: 1075
32	Table number: 2
33	Supplementary number: 1 Method + References, 6 Tables, 4 Figures
34	
35	
36	KEY WORDS
37	Asthma, T cells, flow cytometry, clinical immunology, interleukins
38	

## 40 ABBREVIATIONS

41	ACT	Asthma control test
42	ALL	All asthmatics
43	$FEV_1$	Forced expiratory volume in one sec
44	FVE	Forced vital capacity
45	HD	Healthy donors
46	GINA	Global Initiative for Asthma
47	ICS	Inhaled corticosteroids
48	IFNγ	Interferon- $\gamma$
49	IL	Interleukin
50	IL-17	Interleukin-17A
51	ILT	innate-like T cells
52	iNKT	invariant Natural Killer T cells
53	MAIT	Mucosal-associated invariant T cells
54	MAIT17	IL-17-producing Mucosal-associated invariant T cells
55	MOD	mild-to-moderate asthmatics
56	OCS	Oral corticosteroids
57	SEV	Severe asthmatics
58	Tc1	Type 1 cytotoxic CD8+ T cells
59	Tc2	Type 2 cytotoxic CD8+ T cells
60	TCR	T cell receptor
61 62	Th	T helper
63		
64		

65 To the Editor,

Compelling evidence outlined the importance of conventional T-helper (Th)2 and Th17 CD4<sup>+</sup> 66 T cells and of innate-like T cells (ILT), namely invariant natural killer T (iNKT), γδ T and 67 mucosal-associated invariant T (MAIT) cells, as sources of inflammatory cytokines involved 68 in asthma pathology.<sup>1-5</sup> Ideally, the analysis of the major inflammatory cells implicated in 69 asthma severity should be performed in the airways. However, for several reasons, these studies 70 were realised in a limited number of patients.<sup>6,7</sup> Here, we systematically profiled in peripheral 71 blood Th1, Th2 and Th17, respectively identified as interferon-y (IFNy)-, interleukin (IL)-4-72 and IL-17A (hereafter referred as IL-17)-producing CD4<sup>+</sup> T cells. Type 1 and Type 2 cytotoxic 73 74 (Tc1 and Tc2) CD8<sup>+</sup> T cells (IFNy and IL-4 producers, respectively) and Foxp3<sup>+</sup>CD25<sup>+</sup>CD4<sup>+</sup> regulatory T cells (Treg) were also analysed. Among ILT cells, we focused on IFNy-, IL-4- and 75 IL-17-producing iNKT, MAIT and V $\delta 2^+ \gamma \delta$  T cells. The frequency of these T cells was 76 compared in peripheral blood of 50 healthy donors (HD), 28 mild-to-moderate (GINA steps 1-77 3) and 72 severe asthmatics (GINA step 4 = 33 and GINA step 5 = 39) who have been included 78 over 4 years, in the multicentre Cohort of Bronchial Obstruction and Asthma (COBRA) (Table 79 S1).<sup>8</sup> The group merging all asthmatics was referred as 'ALL'. 80

The proportion of CD4<sup>+</sup> (Figure S1A), but not CD8<sup>+</sup> (Figure S1B), T cells was lower in ALL 81 than in HD. However, this difference disappeared when asthmatic patients were separated into 82 83 GINA steps 1-3, 4 and 5 sub-groups (Table 1). Within the CD4<sup>+</sup> and CD8<sup>+</sup> T cell populations, Th1, Th2, Th17 and Tc1 were greater in ALL than in HD (Figure S1C-F), whereas Tc2 84 frequency was similar between these groups (Figure S1G). Th2 were significantly higher in 85 GINA steps 1-3 and 5, but lower in GINA step 4 than in step 3 (Table 1). A similar trend was 86 observed for Tc1 (Table 1), whereas Tc2 (P=0.08) and Th1 (P=0.06) tended to behave as Th2 87 and Tc1, respectively (Table 1). 88

89 Although there is a consensus showing an association between pro-Th2 immune responses and allergic asthma,<sup>3</sup> we found that Th2 or Tc2 were neither overrepresented in atopic asthmatics 90 (Table S2), nor they correlated with circulating IgE or blood eosinophils (Table 2). In contrast, 91 92 Th1 were negatively associated with blood eosinophils and with daily dose of ICS (Table 2). 93 ICS daily dose was also negatively correlated with Th2 (Table 2). To better understand the possible contribution of these cells in the context of ICS use and atopic markers, we analysed 94 the ratios Th1/Th2, Th2/Th17 and Th1/Th17. These ratios were similar between HD and ALL 95 (Figure S1H, I and J). However, Th2/Th17 and Th1/Th17 positively correlated with circulating 96 IgE, but not with ICS use (Table 2) indicating that low frequency of Th17 is a hallmark of 97

98 elevated IgE levels.

Mirroring the higher frequency of activated T cells, we found lower Treg in ALL than in HD (Figure S2A), whereas no significant differences where observed when the asthmatics were considered as separate groups, according to GINA steps (Table 1). Further, the ratios Th1/Treg, Th17/Treg and, to a lesser extent, Th2/Treg, showed that these activated T cells were overrepresented in ALL compared to HD (Figure S2B to D). This latter ratio also positively correlated with circulating IgE (Table 2). These findings indicate that elevated frequencies of activated T cells and a lower incidence of Treg discriminated asthmatic patients from HD.

106 Among  $\gamma\delta$  T cells, the frequency of V $\delta$ 2 was higher in ALL than in HD (Figure S3A). Further,

we observed an elevated proportion of IFN $\gamma$ - and IL-17-producing V $\delta$ 2 in ALL (Figure S3B and C). No significant differences were observed when the sub-groups of asthmatics were analysed according to GINA steps (Table 1), nor when IFN $\gamma$ - and IL-17-producing V $\delta$ 2 were

110 examined in the context of clinical parameters (Table 2). These findings indicate that the role

- 111 of V $\delta$ 2 on asthma pathology requires confirmation.
- 112 ALL and asthmatics patient sub-groups had similar proportions of iNKT than HD (Figure S3D

and Table 1) However, these cells highly and negatively correlated with circulating IgE (Table

114 2) and they tended to be lower in atopic than in non-atopic asthmatics (P=0.12) (Table S2).

115 These data suggest iNKT are mainly associated with non-atopic asthma.

Examination of activation status of iNKT showed no differences in IL-4 and IFN $\gamma$  production between HD and ALL (Figure S3E and F). However, IL-4-producing iNKT were higher in GINA steps 1-3 than HD and GINA step 5 (Table 1). Both IL-4 and IFN $\gamma$ -producing iNKT correlated negatively with the daily dose of ICS (Table 2), suggesting a possible influence of this treatment on their frequency. It is noteworthy that IL-4-producing iNKT were positively correlated with the ACT score (Table 2) indicating that this population may have an unexpected protective role in these patients.

- In agreement with previous published data,<sup>6</sup> MAIT were significantly lower in ALL than in HD (Figure S3G), but not between HD and asthmatic patient sub-groups (Table1). Whereas no differences were observed in the frequency of IFN $\gamma^+$ /MAIT (Figure S3H and Table 1), IL-17producing MAIT (MAIT17) were higher in ALL than in HD (Figure S3I). Given the absence of correlation between MAIT17 and ICS use (Table 2) and the lack of difference in their
- of correlation between MAIT17 and ICS use (Table 2) and the lack of difference in their
- 128 frequency in OCS-treated and -untreated asthmatics of the GINA 5 step (Table S3), it may be
- speculated that MAIT17 represent a steroid less sensitive cell component involved in asthma

pathology. Further, MAIT17 were negatively correlated with IgE (Table 2) and their frequency was higher in non-atopic than in atopic patients (Table S2). In addition, MAIT17 were positively correlated with ACT score (Table 2). Together, these data suggest that these cells may play a predominant role in non-atopic asthmatic patients. These results are in agreement with our recent observations suggesting a role of MAIT17 in paediatric asthma.<sup>7</sup> However, further studies are required to confirm, or infirm the precise implication of MAIT in asthma.

Overall, our findings indicate that distinct T cell populations and their ratio should be considered to better understand their role in distinct asthma endotypes. It is noteworthy that the positive correlation between MAIT17 and ACT score, associated with our previous studies in asthmatic children,<sup>7</sup> suggest that these cells may play distinct roles in both adult and paediatric asthma pathology.

Although descriptive, this study paves the way for future asthma analysis in large patient cohorts. Overall, we suggest that the Th1/Th2 ratio and the frequency of MAIT17 should be taken into consideration for better identifying distinct asthma endotypes and for testing the clinical benefit of specific immunotherapies.

#### 145 **References**

- Peters MC, Wenzel SE. Intersection of biology and therapeutics: type 2 targeted therapeutics for adult asthma. *Lancet* 2020;**395**(10221):371-383.
- Agache I, Rocha C, Beltran J, Song Y, Posso M, Sola I, et al. Efficacy and safety of treatment
   with biologicals (benralizumab, dupilumab and omalizumab) for severe allergic asthma.
   *Allergy* 2020, Feb 16. Online ahead of print.
- 151 3. Lloyd CM, Hessel EM. Functions of T cells in asthma: more than just T(H)2 cells. *Nat Rev* 152 *Immunol* 2010;10:838-848.
- 4. Boonpiyathad T, Sozener ZC, Satitsuksanoa P, Akdis CA. Immunologic mechanisms in
  asthma. *Semin Immunol* 2019;46:101333.
- 5. Victor JR, Lezmi G, Leite-de-Moraes M. New Insights into Asthma Inflammation: Focus on
   iNKT, MAIT, and gammadelta T Cells. *Clin Rev Allergy Immunol* 2020. Apr 3. Online ahead
   of print.
- 6. Hinks TS, Zhou X, Staples KJ, Dimitrov BD, Manta A, Petrossian T, et al. Innate and adaptive T cells in asthmatic patients: Relationship to severity and disease mechanisms. *J Allergy Clin Immunol* 2015;**136**(2):323-333.
- 161 7. Lezmi G, Abou-Taam R, Garcelon N, Dietrich C, Machavoine F, Delacourt C, et al.
  162 Evidence for a MAIT-17-high phenotype in children with severe asthma. *J Allergy Clin*163 *Immunol* 2019;144(6):1714-1716 e1716.
- 164 8. Pretolani M, Soussan D, Poirier I, Thabut G, Aubier M, Group CS, et al. Clinical and
  165 biological characteristics of the French COBRA cohort of adult subjects with asthma. *Eur*166 *Respir J* 2017;**50**(2).
- 167 9. Raundhal M, Morse C, Khare A, Oriss TB, Milosevic J, Trudeau J, et al. High IFN-gamma
  168 and low SLPI mark severe asthma in mice and humans. *J Clin Invest* 2015;**125**:3037-3050.
- 169
- 170
- 171
- 172
- 173 Acknowledgments
- 174
- 175 This work was supported by grants from Legs Poix, Chancellerie des Universités, Paris, France;
- 176 ANR-2010-MIDI-005; Inserm (Institut National de la Santé et de la Recherche Médicale) and
- 177 CNRS (Centre National de la Recherche Scientifique).
- 178
- 179 Dr. LEITE-DE-MORAES has nothing to disclose.
- 180 Dr. BELO has nothing to disclose.
- 181 Ms DIETRICH has nothing to disclose.
- 182 Dr. SOUSSAN has nothing to disclose.
- 183 Prof. AUBIER reports grants from AStraZeneca, GSK and Roche outside the submitted work.
- 184 Dr. PRETOLANI reports grants from AstraZeneca and from Roche outside the submitted work.

Call type (in %)	Controls	Asthmatics				
	Controls	GINA steps 1-3	GINA step 4	GINA step 5	F	
Number of subjects	50	28	33	39	-	
CD4+/CD3+	69.8 (65.2-76.5)	64.5 (61.1-71.4)	67.5 (61.8-73.9)	69.6 (61.2-75.1)	0.42	
CD8+/CD3+	24.3 (19.7-27.7)	24.6 (18.9-28.8)	23.5 (15.6-28.6)	23.7 (18.3-29.2)	0.96	
IL-4+/CD4+ (Th2)	2.1 (1.2-2.8)	3.6 (1.8-4.7) †	2.1 (1.0-2.9) ‡	3.2 (2.1-4.3) †§	0.04	
IL-4+/CD8+ (Tc2)	0.74 (0.28-1.49)	1.1 (0.8-1.8)	0.8 (0.5-1.2)	1.2 (0.6-2.1)	0.08	
IFNγ+/CD4+ (Th1)	11.7 (8.1-16.1)	21.6 (14.8-26.2)	16.8 (10.6-20.9)	14.9 (11.3-21.0)	0.06	
IFNγ+/CD8+ (Tc1)	22.8 (15.2-38.5)	54.3 (38.5-65.7) †	50.4 (33.9-60.9) †	40.1 (28.9-57.4) †	0.05	
IL-17+/CD4+ (Th17)	0.88 (0.49-1.23)	1.2 (0.7-1.9)	1.0 (0.6-1.3)	1.1 (0.7-1.7)	0.47	
Foxp3+/CD25+/CD4+ (Treg)	6.5 (5.2-7.8)	5.6 (4.7-6.3)	4.8 (4.4-6.1)	5.2 (3.9-6.5)	0.69	
Ratio Th1/Th2	6.3 (3.7-8.7)	8.2 (4.7-9.3)	6.9 (5.5-9.7)	5.6 (3.5-8.1)	0.06	
Ratio Th2/Th17	2.7 (1.3-4.2)	2.4 (1.7-3.3)	1.9 (1.3-2.9)	2.8 (1.6-3.8)	0.22	
Ratio Th1/Th17	12.9 (9.0-21.8)	17.0 (10.3-27.2)	16.0 (8.4-22.2)	13.3 (8.3-18.1)	0.44	
Ratio Th1/Treg	2.0 (1.5-3.1)	3.8 (2.6-5.1)	3.3 (2.3-3.9)	2.9 (1.8-4.3)	0.34	
Ratio Th2/ Treg	0.40 (0.25-0.62)	0.58 (0.41-0.75)	0.50 (0.17-0.78)	0.67 (0.39-0.92)	0.69	
Ratio Th17/Treg	0.12 (0.05-0.27)	0.24 (0.13-0.40)	0.21 (0.13-0.33)	0.23 (0.11-0.36)	0.36	
iNKT/CD3+	0.09 (0.06-0.16)	0.06 (0.03-0.12)	0.09 (0.03-0.33)	0.07 (0.03-0.19)	0.64	
IL-4+/iNKT	6.9 (2.2-16.5)	16.5 (7.7-28.4) †	5.2 (3.6-22.2) ‡	9.8 (4.4-17.4) ‡	0.05	
IFNγ+/iNKT	47.1 (36.7-61.1)	53.2 (44.5-69.1)	47.6 (32.3-61.5)	45.9 (23.4-64.2)	0.14	
MAIT/CD3+ <sup>b</sup>	3.2 (1.8-4.3)	0.7 (0.3-2.7)	1.6 (1.1-3.4)	1.1 (0.7-2.1)	0.18	
IFNγ+/MAIT <sup>c0.92)</sup>	45.9 (24.8-56.0)	56.9 (53.3-60.9)	38.5 (20.1-60.7)	48.0 (42.9-66.3)	0.38	
IL-17+/MAIT (MAIT17) <sup>c</sup>	0.93 (0.49-1.88)	3.4 (2.5-7.6)	1.0 (0.8-2.2)	1.6 (0.8-5.6)	0.33	
Vδ2+/CD3+ <sup>d</sup>	1.5 (0.7-2.4)	3.5 (2.4-4.0)	3.1 (1.7-5.3)	2.2 (0.9-3.3)	0.27	
IFNγ+/Vδ2+ <sup>e</sup>	49.6 (36.0-63.8)	65.4 (49.9-77.4)	69.1 (44.8-77.6)	82.2 (63.8-87.1)	0.29	
IL-17+/ Vδ2+ <sup>e</sup>	0.07 (0.04-0.14)	0.43 (0.31-0.55)	0.15 (0.10-0.41)	0.33 (0.07-1.04)	0.72	

Table 1: Frequency of immune cells in peripheral blood of controls and asthmatics, according to GINA stages

*bbreviations:* IL, interleukin; IFN<sub>γ</sub>, interferon-<sub>γ</sub>; iNKT, invariant natural killer T; MAIT, mucosal-associated invariant T cel lata are medians (25-75 interquartile range)

Kruskal-Wallis test

55 asthmatics were included in the analysis 27 asthmatics were included in the analysis 28 asthmatics were included in the analysis 24 asthmatics were included in the analysis

P<0.05, as compared to controls,  $\ddagger$  P<0.05, as compared to GINA 3 asthmatics; § P<0.05, as compared to  $\exists$ INA 4 asthmatics (Mann-Whitney U-test)

old denotes statistical significance

Cell type (in %)	Circulating	lgE (IU/mL)	Blood EOS	(per mm <sup>3</sup> )	Daily dose	of ICS (µg)	Number of e	exacerbations	ACT scor	e
	r	Р	r	Р	r	Р	r	Р	r	P
CD4+/CD3+	0.048	0.640	-0.034	0.736	0.119	0.239	-0.013	0.902	0.114	0.304
CD8+/CD3+	-0.048	0.834	0.050	0.621	-0.089	0.378	0.073	0.487	-0.192	0.082
IL-4+/CD4+ (Th2)	-0.030	0.775	-0.183	0.078	-0.209	0.043	-0.004	0.967	-0.121	0.289
IL-4+/CD8+ (Tc2)	-0.022	0.835	-0.165	0.112	-0.122	0.239	-0.012	0.912	-0.104	0.362
IFNγ+/CD4+ (Th1)	-0.121	0.250	-0.217	0.035	-0.285	0.005	-0.147	0.178	0.029	0.799
IFNγ+/CD8+ (Tc1)	-0.134	0.204	-0.158	0.131	-0.185	0.075	-0.115	0.293	-0.038	0.740
IL-17+/CD4+ (Th17)	-0.134	0.200	-0.145	0.163	-0.166	0.109	-0.061	0.578	-0.005	0.961
Foxp3+/CD25+/CD4+ (Treg)	-0.100	0.346	-0.001	0.926	-0.056	0.594	-0.140	0.202	-0.017	0.884
Ratio Th1/Th2	-0.030	0.747	-0.030	0.790	- 0.030	0.801	-0.060	0.587	0.140	0.219
Ratio Th2/Th17	0.280	0.007	-0.007	0.946	-0.079	0.447	-0.039	0.718	-0.045	0.690
Ratio Th1/Th17	0.218	0.036	-0.008	0.940	-0.106	0.307	-0.081	0.453	0.030	0.791
Ratio Th1/Treg	0.168	0.128	-0.188	0.084	-0.123	0.261	-0.116	0.307	0.062	0.604
Ratio Th2/ Treg	0.285	0.009	-0.190	0.080	-0.138	0.206	-0.0003	0.998	-0.069	0.563
Ratio Th17/Treg	-0.064	0.569	-0.143	0.192	-0.059	0.595	-0.072	0.526	0.084	0.480
iNKT/CD3+	-0.427	< 0.001	0.086	0.395	0.113	0.261	-0.049	0.644	0.106	0.342
IL-4+/iNKT	0.163	0.123	-0.185	0.078	-0.325	0.001	-0.054	0.625	0.255	0.025
IFNγ+/iNKT	-0.059	0.580	0.039	0.714	-0.234	0.026	-0.162	0.144	-0.026	0.821
MAIT/CD3+ b	-0.062	0.661	-0.082	0.551	0.218	0.110	0.093	0.515	-0.173	0.249
IFNγ+/MAIT <sup>c</sup>	0.279	0.167	-0.336	0.080	0.105	0.594	-0.276	0.202	0.295	0.194
IL-17+/MAIT (MAIT17) <sup>c</sup>	-0.439	0.025	0.120	0.543	-0.303	0.117	-0.113	0.607	0.451	0.040
Vδ2+/CD3+ c	-0.075	0.713	0.028	0.887	-0.010	0.956	-0.011	0.953	-0.037	0.858
IFNγ+/Vδ2+ <sup>c</sup>	-0.036	0.865	-0.198	0.331	0.310	0.135	0.216	0.310	-0.030	0.896
IL-17+/ Vδ2+ <sup>c</sup>	-0.239	0.260	-0.07	0.719	-0.133	0.515	-0.074	0.732	0.252	0.258

Abbreviations: IU, international units; EOS, eosinophils; ICS, inhaled corticosteroids; IL. interleukin; IFNy. interferon-y; iNKT. invariant natural killer T; MAIT. mucosal-associated invariant T cells

<sup>a</sup> Spearman's rank-order method (n = 100)
 <sup>b</sup> 56 asthmatics were included in the analysis
 <sup>c</sup> 28 asthmatics were included in the analysis
 <sup>d</sup> 24 asthmatics were included in the analysis
 Bold denotes statistical significance

Γable S1: Characteristics of mild-to-moderate and severe asthmati	s providing blood samples	s for determining the frequency	y of immune cells
---	---------------------------	---------------------------------	-------------------

		Asthmatics, accord			
Parameter	ALL	MOD SEV			Da
		GINA steps 1-3 <sup>b</sup>	GINA step 4	GINA step 5	— F -
No. of patients (%)	100	28	33	39	-
Women – no. (%)	58 (58.0)	16 (57.1)	18 (54.6)	24 (61.5)	0.84
Age – years	50.1 ± 15.2	50.5 ± 17.4	51.5 ± 16.5	48.5 ± 12.3	0.56
BMI – kg per m²	26.6 ± 4.5	27.5 ± 4.5	$26.4 \pm 4.0$	26.0 ± 5.0	0.22
Smoking history – former/never (%) <sup>c</sup>	30/66 (31/69)	9/18 (33/67)	7/23 (23/77)	14/25 (36/64)	0.52
Age of asthma onset – years	23.4 ± 18.9	$23.2 \pm 20.6$	25.7 ± 19.3	21.7 ± 17.4	0.69
Asthma duration – years	27.0 ± 15.8	27.6 ± 14.0	25.2 ± 16.0	28.2 ± 17.0	0.84
With history of atopy – no. (%)	67 (67.0)	23 (82.1)	18 (54.6)	26 (66.7)	0.08
Blood eosinophils counts – 10 <sup>9</sup> per Liter	290 (180-450)	285 (200-385)	290 (190-550)	308 (110-510)	0.63
With ≥ 300 blood eosinophils – no. (%)	46 (46.0)	11 (39.3)	15 (45.5)	20 (51.3)	0.63
Total serum IgE – International Units per milliliter	221 (88-600)	415 (76-812)	176 (82-309)	220 (92-679)	0.17
Respiratory function					
Pre-bronchodilator FEV1 – % of predicted	73.9 ± 20.9	88.8 ± 16.0	71.1 ± 19.5 †	65.5 ± 19.9 †	< 0.0001
Post-bronchodilator FEV1 - % of predicted	79.9 ± 21.2	96.9 ± 14.3	76.8 ± 19.1 †	70.3 ± 20.2 †	< 0.0001
Pre-bronchodilator FEV <sub>1</sub> /FVC – % of predicted	62.8 ± 12.3	66.8 ± 10.5	60.8 ± 12.2	61.6 ± 13.1	0.12
Post-bronchodilator FEV <sub>1</sub> /FVC – % of predicted	65.5 ± 12.4	70.5 ± 9.0	63.8 ± 12.9 †	63.5 ± 13.4 †	0.03
Treatments					
On SABA – number (%)	100 (100.0)	28 (100.0)	33 (100.0)	39 (100.0)	-
On LABA – number (%)	93 (93.0)	23 (82.1)	32 (97.0)	38 (97.4) †	0.03
Daily dose of ICS <sup>c</sup>	1809 ± 1006	696 ± 249	2139 ± 761 †	2328 ± 907 †	< 0.0001

On maintenance use of OCS – no. (%)	16 (16.0)	0	0	16 (41.0)	-
Daily dose of oral prednisone – mg	25.3 ± 12.3	0	0	25.3 ± 12.3	-
On anti-IgE	18 (18.0)	0	0	18 (46.2)	-

#### Table S1 (continued)

\_

Asthma control							
	Score on ACT	13.5 ± 6.0	18.4 ± 4.5	13.1 ± 6.1 †	10.3 ± 4.5 †‡	< 0.0001	
	Annual rate of severe exacerbations requiring OCS bursts	$3.0 \pm 0.4$	0.4 ± 0.2	2.5 ± 0.6 †	5.3 ± 0.8 †‡	< 0.0001	

Abbreviations: BMI, body mass index; FEV<sub>1</sub>, Forced Expiratory Volume in One second; FVC, Forced Vital Capacity; SABA, short-acting β2-agonists; LABA, long-acting β2-ago ICS, inhaled corticosteroids; OCS, oral corticosteroids; ACT, asthma control test Data are n (%), or means ± SD, or medians (interquartile range). Results on annual rates of severe exacerbations are expressed as means ± SE <sup>a</sup> ANOVA, or Kruskal-Wallis test or  $\chi^2$  test <sup>b</sup> Data are available for 2 GINA 1, 17 GINA 2 and 9 GINA 3 asthmatics

 clining of beclomethasone equivalents
 † P<0.05, as compared to GINA 1-3 asthmatics, ‡ P <0.05, as compared to GINA 4 asthmatics (Student's t test for unpaired value, Mann-Whitney U-test or Fisher test 2-tailed)</li> Bold denotes statistical significance

Cell type (in %)	Non atopic	Atopic	P <sup>a</sup>
Number of patients	32	67	-
CD4+/CD3+	64.6 (60.9-73.8)	67.4 (61.7-74.6)	0.55
CD8+/CD3+	23.5 (14.7-28.6)	24.6 (18.8-29.2)	0.39
IL-4+/CD4+ (Th2)	2.8 (1.0-3.9)	2.8 (1.8-4.0)	0.58
IL-4+/CD8+ (Tc2)	1.0 (0.6-2.0)	1.0 (0.6-1.7)	0.79
IFNγ+/CD4+ (Th1)	17.9 (11.3-24.0)	16.8 (12.1-22.4)	0.79
IFNγ+/CD8+ (Tc1)	53.5 (39.6-61.3)	40.9 (28.7-62.2)	0.21
IL-17+/CD4+ (Th17)	1.3 (0.7-1.9)	1.0 (0.7-1.7)	0.52
Foxp3+/CD25+/CD4+ (Treg)	5.6 (4.4-6.4)	5.2 (4.3-6.2)	0.38
Ratio Th1/Th2	7.6 (4.5-9.2)	6.4 (4.2-8.3)	0.38
Ratio Th2/Th17	1.9 (1.1-2.9)	2.7 (1.7-3.1)	0.09
Ratio Th1/Th17	12.6 (9.2-20.7)	15.9 (8.7-21.3)	0.58
Ratio Th1/Treg	3.3 (1.8-4.2)	3.6 (2.4-4.3)	0.73
Ratio Th2/ Treg	0.52 (0.23-0.76)	0.65 (0.34-0.84)	0.30
Ratio Th17/Treg	0.25 (0.12-0.35)	0.21 (0.12-0.38)	0.96
iNKT/CD3+	0.11 (0.06-0.20)	0.06 (0.03-0.18)	0.18
IL-4+/iNKT	6.1 (3.6-14.6)	12.5 (4.4-23.1)	0.12
IFNy+/iNKT	45.6 (32.7-64.5)	51.7 (32.4-65.5)	0.74
MAIT/CD3+ <sup>b</sup>	1.4 (0.7-4.0)	1.0 (0.5-2.0)	0.11
IFNγ+/MAIT <sup>c</sup>	43.4 (32.5-56.3)	53.2 (36.2-67.9)	0.31
IL-17+/MAIT (MAIT17) <sup>c</sup>	3.2 (1.6-6.5)	1.0 (0.6-2.7)	0.05
Vδ2+/CD3+ <sup>d</sup>	2.9 (1.1-3.9)	3.1 (1.2-3.6)	0.89
IFNγ+/Vδ2+ <sup>e</sup>	76.0 (59.1-87.1)	77.2 (51.7-79.7)	0.67
IL-17+/ Vδ2+ <sup>e</sup>	0.29 (0.15-0.75)	0.25 (0.10-0.58)	0.89

Table S2: Frequency of immune cells in peripheral blood of asthmatics according to atopy <sup>a</sup>

Abbreviations: IU, international units; IL, interleukin; IFNγ, interferon-γ; iNKT, invariant natural killer T; MAIT, mucosal-associated invariant T cells

Data are medians (25-75 interquartile range)

<sup>a</sup> Mann-Whitney U-test

- <sup>b</sup> 55 asthmatics were included in the analysis
- <sup>c</sup> 27 asthmatics were included in the analysis

<sup>d</sup> 28 asthmatics were included in the analysis

<sup>e</sup> 24 asthmatics were included in the analysis

Bold denotes statistical significance

Cell type (in %)	Without OCS	On regular OCS				
		All	P ª	≤ 20 mg/day	> 20 mg/day	P <sup>b</sup>
Number of patients	23	16	-	9	7	-
CD4+/CD3+	71.7 (64.7-75.1)	64.7 (59.7-74.9)	0.53	67.8 (63.7-84.7)	53.0 (44.0-56.0)	0.06
CD8+/CD3+	22.2 (18.3-26.7)	27.3 (17.9-29.7)	0.31	29.5 (13.8-29.7)	26.9 (24.6-28.3)	0.99
IL-4+/CD4+ (Th2)	3.2 (2.2-4.0)	3.4 (2.2-4.8)	0.73	2.9 (1.6-4.1)	4.3 (3.2-4.8)	0.61
IL-4+/CD8+ (Tc2)	1.2 (0.6-2.1)	1.3 (0.5-1.9)	0.84	1.0 (0.4-4.2)	1.7 (1.0-1.7)	0.99
IFNγ+/CD4+ (Th1)	14.4 (10.6-19.7)	16.9 (11.7-23.5)	0.51	14.5 (10.7-22.9)	19.3 (14.4-23.6)	0.69
IFNγ+/CD8+ (Tc1)	38.4 (26.5-53.5)	40.9 (33.9-65.8)	0.25	39.8 (34.5-57.6)	55.9 (33.3-72.2)	0.69
IL-17+/CD4+ (Th17)	1.2 (0.8-1.8)	1.0 (0.7-1.6)	0.51	1.0 (0.9-1.6)	0.9 (0.6-1.6)	0.61
Foxp3+/CD25+/CD4+ (Treg)	5.4 (4.0-6.1)	4.4 (4.0-6.7)	0.75	4.4 (3.5-5.2)	5.8 (4.3-4.3-8.5)	0.39
Ratio Th1/Th2	5.8 (3.4-8.1)	5.5 (3.6-7.4)	0.75	5.9 (4.7-8.1)	3.8 (3.6-5.3)	0.53
Ratio Th2/Th17	2.8 (1.6-3.1)	2.8 (1.8-3.4)	0.68	2.1 (1.9-2.9)	3.0 (2.1-5.8)	0.39
Ratio Th1/Th17	10.7 (7.4-16.6)	15;2 (11.8-18.2)	0.36	15.7 (10.8-17.1)	13.5 (11.9-24.3)	0.69
Ratio Th1/Treg	2.7 (1.6-4.2)	3.5 (1.6-4.5)	0.52	3.5 (2.0-4.5)	3.6 (1.3-5.2)	0.71
Ratio Th2/ Treg	0.67 (0.41-0.77)	0.63 (0.35-0.99)	0.73	0.43 (0.38-0.92)	0.92 (0.36-1.02)	0.90
Ratio Th17/Treg	0.24 (0.13-0.42)	0.17 (0.10-0.35)	0.44	0.26 (0.17-0.35)	0.12 (0.09-0.24)	0.31
iNKT/CD3+	0.08 (0.03-0.17)	0.07 (0.03-0.20)	0.99	0.05 (0.03-0.12)	0.07 (0.04-0.25)	0.60
IL-4+/iNKT	13.1 (4.6-17.8)	5.7 (2.9-11.5)	0.11	7.8 (4.3-12.2)	4.8 (2.9-9.7)	0.57
IFNγ+/iNKT	43.9 (21.4-63.6)	46.7 (30.1-63.9)	0.72	52.8 (37.3-72.3)	42.3 (30.1-57.2)	0.41
MAIT/CD3+	0.8 (0.4-1.8)	1.3 (0.7-2.5)	0.29	1.1 (0.8-2.4)	1.5 (0.7-2.4)	0.99
IFNγ+/MAIT (MAIT17) <sup>c</sup>	44.4 (42.5-73.6)	51.6 (44.4-63.6)	0.99	54.0 (49.2-58.8)	51.6 (43.2-59.4)	0.99
IL-17+/MAIT <sup>c</sup>	6.5 (1.2-6.7)	1.2 (0.7-2.0)	0.30	1.0 (0.8-1.1)	2.0 (1.2-2.4)	0.78
Vδ2+/CD3+ <sup>d</sup>	3.1 (1.7-3.3)	1.5 (1.0-1.0-3.2)	0.99	1.5 (1.2-2.9)	2.1 (0.9-4.2)	0.91
IFNγ+/Vδ2+ <sup>d</sup>	86.2 (81.9-88.0)	81.9 (58.2-83.2)	0.46	65.7 (31.0-82.5)	86.8 (76.0-01.7)	0.28
IL-17+/ Vδ2+d	0.05 (0.03-0.16)	0.48 (0.31-1.57)	0.08	0.48 (0.35-0.86)	0.94 (0.25-1.67)	0.91

Abbreviations: OCS, oral corticosteroids; IL, interleukin; IFNγ, interferon-γ; iNKT, invariant natural killer T; MAIT, mucosal-associated invariant T cells Data are medians (25-75 interquartile range) <sup>a</sup> Mann-Whitney U-test between severe OCS-untreated and –treated asthmatics

<sup>b</sup> Mann-Whitney U-test between severe asthmatics treated with  $\leq$  20 mg/day and  $\geq$  20 mg/day OCS

<sup>c</sup> 10 asthmatics were included in the analysis

<sup>d</sup> 12 asthmatics were included in the analysis

Cell types	HD		ALL	
	r	P	r	Р
Th2 versus Th1	0.476	< 0.001	0.534	< 0.001
Th2 versus Th17	0.213	0.146	0.474	< 0.001
Th2 versus Tc1	0.312	0.030	0.348	< 0.001
Th2 versus Tc2	0.498	< 0.001	0.460	< 0.001
Th1 versus Th17	0.393	0.005	0.419	< 0.001
Th1 <i>versus</i> Tc1	0.730	< 0.001	0.671	< 0.001
Th1 versus Tc2	0.208	0.156	0.320	< 0.001
Th17 versus Tc1	0.282	0.049	0.190	0.071
Th17 versus Tc2	0.099	0.503	0.238	0.020
Tc1 versus Tc2	0.303	0.036	0.239	0.022
Th1 versus Treg	-0.253	0.161	0.145	0.182
Th2 versus Treg	-0.263	0.146	0.125	0.251
Th17 versus Treg.	-0.067	0.716	0.022	0.836
Tc1 versus Treg.	0.079	0.667	0.095	0.385
Tc2 versus Treg.	0.058	0.750	0.119	0.275

Table S4. Correlation analyses between Th2, Th1, Th17, Tc1 and Tc2 in healthy donors (HD) and asthmatic patients (ALL)<sup>a</sup>

<sup>a</sup> Spearman's rank-order method (n = 100) Bold denotes statistical significance

Cell type (in %)	Without omalizumab	With omalizumab	P <sup>b</sup>
Number of patients	21	18	-
CD4+/CD3+	71.7 (60.6-79.7)	69.6 (63.0-72.7)	0.40
CD8+/CD3+	21.7 (14.7-29.7)	25.5 (20.9-28.4)	0.61
IL-4+/CD4+ (Th2)	3.7 (1.9-4.9)	3.2 (2.2-3.7)	0.53
IL-4+/CD8+ (Tc2)	1.3 (0.7-2.1)	1.1 (0.6-2.0)	0.66
IFNγ+/CD4+ (Th1)	14.6 (8.7-23.5)	15.2 (11.6-19.9)	0.84
IFNγ+/CD8+ (Tc1)	43.5 (28.1-58.9)	40.0 (30.7-53.5)	0.97
IL-17+/CD4+ (Th17)	1.2 (0.7-1.7)	1.0 (0.7-1.8)	0.98
Foxp3+/CD25+/CD4+ (Treg)	5.2 (4.0-6.3)	5.2 (4.0-6.5)	0.87
Ratio Th1/Th2	5.5 (3.4-8.1)	5.8 (3.9-7.8)	0.45
Ratio Th2/Th17	2.9 (1.8-3.0)	2.2 (1.6-3.2)	0.90
Ratio Th1/Th17	12.6 (7.4-16.7)	13.5 (8.9-18.5)	0.83
Ratio Th1/Treg	2.5 (1.5-4.4)	3.2 (2.3-4.2)	0.68
Ratio Th2/ Treg	0.69 (0.39-1.00)	0.51 (0.38-0.83)	0.82
Ratio Th17/Treg	0.22 (0.11-0.34)	0.27 (0.12-0.39)	0.61
NKT/CD3+	0.09 (0.03-0.20)	0.06 (0.03-0.17)	0.60
IL-4+/iNKT	5.7 (4.3-13.6)	13.1 (4.5-17.8)	0.40
IFN <sub>7</sub> +/iNKT	43.3 (26.6-63.7)	51.0 (21.4-63.6)	0.85
MAIT/CD3+ <sup>c</sup>	1.1 (0.7-2.5)	1.0 (0.7-1.7)	0.56
IFNγ+/MAIT <sup>d</sup>	43.4 (40.5-50.1)	57.6 (46.2-71.1)	0.46
IL-17+/MAIT (MAIT17) <sup>d</sup>	1.6 (1.0-3.2)	2.0 (0.8-5.7)	0.75
Vδ2+/CD3+ <sup>e</sup>	1.3 (0.7-3.3)	3.0 (2.4-3.2)	0.68
IFNγ+/Vδ2+ <sup>e</sup>	74.1 (51.6-87.6)	82.6 (80.8-84.9)	0.68
IL-17+/ Vδ2+ <sup>e</sup>	0.33 (0.06-0.80)	0.57 (0.22-1.04)	0.94

Table S5: Frequency of immune cells in peripheral blood of severe asthmatics, according to treatment with omalizumab <sup>a</sup>

Abbreviations: OCS, oral corticosteroids; IL, interleukin; IFN<sub>γ</sub>, interferon-γ; iNKT, invariant natural killer T; MAIT, mucosal-associated invariant T cells

Data are medians (25-75 interquartile range)

<sup>a</sup>Only severe asthmatics of GINA 5 step were included in the analysis (n=39)

<sup>b</sup> Mann-Whitney U-test

<sup>c</sup> 23 asthmatics were included in the analysis

<sup>d</sup> 10 asthmatics were included in the analysis

<sup>e</sup> 12 asthmatics were included in the analysis

Cell type (in %)	Never smokers	Former smokers	P <sup>b</sup>
Number of patients	67	31	-
CD4+/CD3+	67.4 (61.1-74.6)	67.6 (63.0-73.9)	0.69
CD8+/CD3+	22.8 (18.0-29.5)	25.6 (15.8-27.5)	0.76
IL-4+/CD4+ (Th2)	2.9 (1.7-4.5)	2.3 (1.4-3.6)	0.18
IL-4+/CD8+ (Tc2)	1.1 (0.6-1.8)	0.9 (0.5-1.7)	0.39
IFNγ+/CD4+ (Th1)	17.6 (11.6-22.6)	15.2 (11.6-25.4)	0.93
IFNγ+/CD8+ (Tc1)	47.7 (33.0-61.8)	39.5 (33.7-60.9)	0.62
IL-17+/CD4+ (Th17)	1.3 (0.7-2.0)	0.9 (0.7-1.2)	0.04
Foxp3+/CD25+/CD4+ (Treg)	5.3 (4.3-6.3)	4.8 (4.3-6.1)	0.67
Ratio Th1/Th2	6.1 (4.0-8.2)	7.6 (5.7-9.5)	0.13
Ratio Th2/Th17	2.1 (1.4-3.0)	2.9 (1.6-3.2)	0.41
Ratio Th1/Th17	12.7 (8,6-19.3)	16.9 (10.7-23.3)	0.12
Ratio Th1/Treg	3.5 (2.3-4.2)	2.9 (1.9-4.8)	0.83
Ratio Th2/ Treg	0.57 (0.37-0.83)	0.47 (0.25-0.77)	0.40
Ratio Th17/Treg	0.25 (0.12-0.40)	0.17 (0.12-0.31)	0.18
iNKT/CD3+	0.08 (0.03-0.20)	0.06 (0.02-0.16)	0.38
IL-4+/iNKT	11.9 (4.6-23.2)	6.9 (3.4-14.8)	0.10
IFNy+/INKT	51.0 (35.9-65.5)	45.9 (27.8-63.4)	0.66
MAIT/CD3+ <sup>c</sup>	1.1 (0.5-2.3)	1.6 (0.7-3.2)	0.57
IFNγ+/MAIT <sup>d</sup>	48.2 (36.2-63.6)	62.6 (46.2-81.6)	0.24
IL-17+/MAIT (MAIT17) <sup>d</sup>	2.3 (0.8-6.4)	1.1 (0.7-2.3)	0.34
Vδ2+/CD3+ <sup>d</sup>	3.2 (1.4-3.6)	3.0 (1.1-4.1)	0.92
IFNγ+/Vδ2+ <sup>e</sup>	69.6 (51.7-79.8)	81.9 (77.4-91.0)	0.14
IL-17+/ Vδ2+ <sup>e</sup>	0.29 (0.13-0.55)	0.27 (0.10-0.96)	0.98

Table S6: Frequency of immune cells in peripheral blood according to smoking history in all asthmatics <sup>a</sup>

Abbreviations: IL, interleukin; IFN $\gamma$ , interferon- $\gamma$ ; iNKT, invariant natural killer T; MAIT, mucosal-associated invariant T cells

Data are medians (25-75 interquartile range)

<sup>a</sup> Data are available for 98 out of the 100 asthmatics included in the study

<sup>b</sup> Mann-Whitney U-test

- <sup>c</sup> 55 asthmatics were included in the analysis
- <sup>d</sup> 27 asthmatics were included in the analysis
- <sup>e</sup> 23 asthmatics were included in the analysis

Bold denotes statistical significance