



**HAL**  
open science

# Virtual an arrow Temperley–Lieb algebras, Markov traces, and virtual link invariants

Luis Paris, Loïc Rabenda

► **To cite this version:**

Luis Paris, Loïc Rabenda. Virtual an arrow Temperley–Lieb algebras, Markov traces, and virtual link invariants. 2021. hal-02940188v2

**HAL Id: hal-02940188**

**<https://hal.science/hal-02940188v2>**

Preprint submitted on 10 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Virtual and arrow Temperley–Lieb algebras, Markov traces, and virtual link invariants

LUIS PARIS  
LOÏC RABENDA

**Abstract** Let  $R^f = \mathbb{Z}[A^{\pm 1}]$  be the algebra of Laurent polynomials in the variable  $A$  and let  $R^a = \mathbb{Z}[A^{\pm 1}, z_1, z_2, \dots]$  be the algebra of Laurent polynomials in the variable  $A$  and standard polynomials in the variables  $z_1, z_2, \dots$ . For  $n \geq 1$  we denote by  $\text{VB}_n$  the virtual braid group on  $n$  strands. We define two towers of algebras  $\{\text{VTL}_n(R^f)\}_{n=1}^{\infty}$  and  $\{\text{ATL}_n(R^a)\}_{n=1}^{\infty}$  in terms of diagrams. For each  $n \geq 1$  we determine presentations for both,  $\text{VTL}_n(R^f)$  and  $\text{ATL}_n(R^a)$ . We determine sequences of homomorphisms  $\{\rho_n^f : R^f[\text{VB}_n] \rightarrow \text{VTL}_n(R^f)\}_{n=1}^{\infty}$  and  $\{\rho_n^a : R^a[\text{VB}_n] \rightarrow \text{ATL}_n(R^a)\}_{n=1}^{\infty}$ , we determine Markov traces  $\{T_n^f : \text{VTL}_n(R^f) \rightarrow R^f\}_{n=1}^{\infty}$  and  $\{T_n^a : \text{ATL}_n(R^a) \rightarrow R^a\}_{n=1}^{\infty}$ , and we show that the invariants for virtual links obtained from these Markov traces are the  $f$ -polynomial for the first trace and the arrow polynomial for the second trace. We show that, for each  $n \geq 1$ , the standard Temperley–Lieb algebra  $\text{TL}_n$  embeds into both,  $\text{VTL}_n(R^f)$  and  $\text{ATL}_n(R^a)$ , and that the restrictions to  $\{\text{TL}_n\}_{n=1}^{\infty}$  of the two Markov traces coincide.

**AMS Subject Classification** 57K12, 57K14, 20F36

## 1 Introduction

Let  $S_1, \dots, S_\ell$  be a collection of  $\ell$  oriented circles smoothly immersed in the plane and having only double crossings. We assign to each crossing a value “positive”, “negative”, or “virtual”, that we indicate on the graphical representation of  $S_1 \cup \dots \cup S_\ell$  as in Figure 1.1. Such a figure is called a *virtual link diagram*. We consider the equivalence relation on the set of virtual link diagrams generated by isotopy and the so-called *Reidemeister virtual moves*, as described in Kauffman [6, 7]. An equivalence class of virtual link diagrams is called a *virtual link*.




Figure 1.1: Crossings in a virtual link diagram

Let  $b = (b_1, \dots, b_n)$  be a collection of  $n$  smooth paths in the plane  $\mathbb{R}^2$  satisfying the following properties.

- (a)  $b_i(0) = (0, i)$  for all  $i \in \{1, \dots, n\}$ , and there exists a permutation  $w \in \mathfrak{S}_n$  such that  $b_i(1) = (1, w(i))$  for all  $i \in \{1, \dots, n\}$ .
- (b) Let  $p_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$  be the projection on the first coordinate. Then  $p_1(b_i(t)) = t$  for all  $i \in \{1, \dots, n\}$  and all  $t \in [0, 1]$ .

(c) The union of the images of the  $b_i$ 's has only normal double crossings.

As with virtual link diagrams, we assign to each crossing a value “positive”, “negative”, or “virtual”, that we indicate on the graphical representation as in Figure 1.1. Such a figure is called a *virtual braid diagram* on  $n$  strands. We consider the equivalence relation on the set of virtual braid diagrams on  $n$  strands generated by isotopy and some *Reidemeister virtual moves*, as described in Kauffman [6]. An equivalence class of virtual braid diagrams on  $n$  strands is called a *virtual braid* on  $n$  strands. The virtual braids on  $n$  strands form a group, denoted  $\text{VB}_n$ , called *virtual braid group* on  $n$  strands. The group operation is induced by the concatenation.

We know from Kamada [4] and Vershinin [12] that  $\text{VB}_n$  admits a presentation with generators  $\sigma_1, \dots, \sigma_{n-1}, \tau_1, \dots, \tau_{n-1}$  and relations

$$\begin{aligned} \tau_i^2 &= 1 \quad \text{for } 1 \leq i \leq n-1, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i, \tau_i \tau_j = \tau_j \tau_i, \tau_i \sigma_j = \sigma_j \tau_i \quad \text{for } |i-j| \geq 2, \\ \sigma_i \sigma_j \sigma_i &= \sigma_j \sigma_i \sigma_j, \tau_i \tau_j \tau_i = \tau_j \tau_i \tau_j, \tau_i \tau_j \sigma_i = \sigma_j \tau_i \tau_j \quad \text{for } |i-j| = 1. \end{aligned}$$

The generators  $\sigma_i$  and  $\tau_i$  are illustrated in Figure 1.2.




Figure 1.2: Generators of  $\text{VB}_n$

Note that the subgroup of  $\text{VB}_n$  generated by  $\sigma_1, \dots, \sigma_{n-1}$  is the braid group  $B_n$  on  $n$  strands. On the other hand,  $\text{VB}_n$  may be viewed as a subgroup of  $\text{VB}_{n+1}$  via the monomorphism  $\text{VB}_n \hookrightarrow \text{VB}_{n+1}$  which sends  $\sigma_i$  to  $\sigma_i$  and  $\tau_i$  to  $\tau_i$  for all  $i \in \{1, \dots, n-1\}$ .

Using the same procedure as for classic braids, we can close a virtual braid  $\beta$  and obtain a virtual link,  $\hat{\beta}$ , called the *closure* of  $\beta$ . We know that each virtual link is the closure of a virtual braid, and we can say when two closed virtual braids are equivalent in terms of virtual Markov moves, as follows.

We denote by  $\text{VB} = \bigsqcup_{n=1}^{\infty} \text{VB}_n$  the disjoint union of all virtual braid groups. Let  $\beta_1, \beta_2 \in \text{VB}$ . We say that  $\beta_1$  and  $\beta_2$  are connected by a *virtual Markov move* if we are in one of the following four cases.

- There exist  $n \geq 1$  and  $\alpha \in \text{VB}_n$  such that  $\beta_1, \beta_2 \in \text{VB}_n$  and  $\beta_2 = \alpha \beta_1 \alpha^{-1}$ .
- There exist  $n \geq 1$  and  $u \in \{\sigma_n, \sigma_n^{-1}, \tau_n\}$  such that  $\beta_1 \in \text{VB}_n$ ,  $\beta_2 \in \text{VB}_{n+1}$  and  $\beta_2 = \beta_1 u$ , or vice versa.
- There exists  $n \geq 2$  such that  $\beta_1 \in \text{VB}_n$ ,  $\beta_2 \in \text{VB}_{n+1}$ , and  $\beta_2 = \beta_1 \sigma_n^{-1} \tau_{n-1} \sigma_n$ , or vice versa.
- There exists  $n \geq 2$  such that  $\beta_1 \in \text{VB}_n$ ,  $\beta_2 \in \text{VB}_{n+1}$ , and  $\beta_2 = \beta_1 \tau_n \tau_{n-1} \sigma_{n-1} \tau_n \sigma_{n-1}^{-1} \tau_{n-1} \tau_n$ , or vice versa.

**Theorem 1.1** (Kamada [4], Kauffman–Lambropoulou [9]) *Let  $\beta_1, \beta_2 \in \text{VB}$ . Then  $\hat{\beta}_1 = \hat{\beta}_2$  if and only if  $\beta_1$  and  $\beta_2$  are connected by a finite sequence of virtual Markov moves.*

Let  $R$  be a ring. For  $n \geq 1$  we denote by  $R[\text{VB}_n]$  the group  $R$ -algebra of  $\text{VB}_n$ . Notice that, since  $\text{VB}_n$  is a subgroup of  $\text{VB}_{n+1}$ ,  $R[\text{VB}_n]$  is a subalgebra of  $R[\text{VB}_{n+1}]$ . A sequence  $\{T_n : R[\text{VB}_n] \rightarrow R\}_{n=1}^{\infty}$  of  $R$ -linear forms is called a *Markov trace* if it satisfies the following properties.

- (a)  $T_n(xy) = T_n(yx)$  for all  $n \geq 1$  and all  $x, y \in R[\text{VB}_n]$ .
- (b)  $T_n(x) = T_{n+1}(x\sigma_n) = T_{n+1}(x\sigma_n^{-1}) = T_{n+1}(x\tau_n)$  for all  $n \geq 1$  and all  $x \in R[\text{VB}_n]$ .
- (c)  $T_n(x) = T_{n+1}(x\sigma_n^{-1}\tau_{n-1}\sigma_n)$  for all  $n \geq 2$  and all  $x \in R[\text{VB}_n]$ .
- (d)  $T_n(x) = T_{n+1}(x\tau_n\tau_{n-1}\sigma_{n-1}\tau_n\sigma_{n-1}^{-1}\tau_{n-1}\tau_n)$  for all  $n \geq 2$  and all  $x \in R[\text{VB}_n]$ .

Note that our definition of “Markov trace” is not the one that can be usually found in the literature (see Kauffman–Lambropoulou [9], for example), but all known definitions, including this one, are equivalent up to renormalization.

Let  $\mathcal{VL}$  be the set of virtual links. Thanks to Theorem 1.1, from a Markov trace  $\{T_n : R[\text{VB}_n] \rightarrow R\}_{n=1}^{\infty}$  we can define an invariant  $I : \mathcal{VL} \rightarrow R$  by setting  $I(\hat{\beta}) = T_n(\beta)$  for all  $n \geq 1$  and all  $\beta \in \text{VB}_n$ . Conversely, from any invariant  $I : \mathcal{VL} \rightarrow R$ , we can define a Markov trace  $\{T_n : R[\text{VB}_n] \rightarrow R\}_{n=1}^{\infty}$  by setting  $T_n(\beta) = I(\hat{\beta})$  for all  $n \geq 1$  and all  $\beta \in \text{VB}_n$ , and then extending  $T_n$  linearly to  $R[\text{VB}_n]$ .

A *tower of algebras* is a sequence  $\{A_n\}_{n=1}^{\infty}$  of algebras such that  $A_n$  is a subalgebra of  $A_{n+1}$  for all  $n \geq 1$ . A sequence  $\{\rho_n : R[\text{VB}_n] \rightarrow A_n\}_{n=1}^{\infty}$  of homomorphisms is said to be *compatible* if the restriction of  $\rho_{n+1}$  to  $R[\text{VB}_n]$  is equal to  $\rho_n$  for all  $n$ . Let  $\{A_n\}_{n=1}^{\infty}$  be a tower of algebras and let  $\{\rho_n : R[\text{VB}_n] \rightarrow A_n\}_{n=1}^{\infty}$  be a compatible sequence of homomorphisms. Set  $S_i = \rho_n(\sigma_i)$  and  $v_i = \rho_n(\tau_i)$  for all  $i \in \{1, \dots, n-1\}$ . A sequence  $\{T'_n : A_n \rightarrow R\}_{n=1}^{\infty}$  of linear forms is a *Markov trace* if it satisfies the following properties.

- (a)  $T'_n(xy) = T'_n(yx)$  for all  $n \geq 1$  and all  $x, y \in A_n$ .
- (b)  $T'_n(x) = T'_{n+1}(xS_n) = T'_{n+1}(xS_n^{-1}) = T'_{n+1}(xv_n)$  for all  $n \geq 1$  and all  $x \in A_n$ .
- (c)  $T'_n(x) = T'_{n+1}(xS_n^{-1}v_{n-1}S_n)$  for all  $n \geq 2$  and all  $x \in A_n$ .
- (d)  $T'_n(x) = T'_{n+1}(xv_nv_{n-1}S_{n-1}v_nS_{n-1}^{-1}v_{n-1}v_n)$  for all  $n \geq 2$  and all  $x \in A_n$ .

Clearly, in that case, the sequence  $\{T_n = T'_n \circ \rho_n : R[\text{VB}_n] \rightarrow R\}_{n=1}^{\infty}$  is a Markov trace, and therefore it determines an invariant for virtual links.

A “natural” strategy to build Markov traces on  $\{K[\text{VB}_n]\}_{n=1}^{\infty}$ , and therefore invariants for virtual links, would be to transit through Markov traces on compatible towers of algebras, as defined above. This strategy won its spurs in the classical theory of knots and links, in particular thanks to Jones’ definitions of the Jones polynomial [2] and of the HOMFLY-PT polynomial [3]. As far as we know, this strategy is poorly used in the theory of virtual knots and links. Actually, the only reference we found is Li–Lei–Li [10], where the authors define a tower of algebras in terms of diagrams, claim (with no proof) that their algebras are the same as the virtual Temperley–Lieb algebras of Zhang–Kauffman–Ge [13], and show that the  $f$ -polynomial can be obtained from a Markov trace on this tower of algebras. They also gave presentations for these algebras in terms of generators and relations, but we found that one of their

relations should be substituted by another one to get a correct presentation (see Proposition 2.2 and Proposition 2.7).

Our aim in the present paper is to describe two invariants for virtual links in terms of Markov traces: the  $f$ -polynomial, also known as the Jones–Kauffman polynomial, and the arrow polynomial. The  $f$ -polynomial is a version of the Jones polynomial for virtual links defined from the Kauffman bracket. This was introduced by Kauffman [6] in his seminal paper on virtual knots and links, and its construction closely follows Kauffman’s construction [5] of the Jones polynomial for classical links. The arrow polynomial is a refinement of the  $f$ -polynomial. It coincides with the Jones polynomial on classical links, but it is much more powerful for (non-classical) virtual links. In particular, it provides a lower bound for the number of virtual crossings. It was constructed by Miyazawa [11] and Dye–Kauffman [1] (see also Kauffman [8]).

Section 2 is dedicated to the construction of a Markov trace associated with the  $f$ -polynomial. Our approach is close to that of Li–Lei–Li [10], but, on the one hand, our study of the  $f$ -polynomial is needed in our study of the arrow polynomial, and, on the other hand, we complete the study of Li–Lei–Li [10] with correct presentations for virtual Temperley–Lieb algebras and other results. For each  $n \geq 1$  we define an algebra  $\text{VTL}_n(R^f)$  in terms of diagrams, so that the sequence  $\{\text{VTL}_n(R^f)\}_{n=1}^\infty$  is a tower of algebras. In Proposition 2.2 we determine a presentation for  $\text{VTL}_n(R^f)$  and in Proposition 2.7 we show that the presentation for  $\text{VTL}_n(R^f)$  given in Li–Lei–Li [10] cannot be correct. Actually, the relation  $E_i E_j E_i = E_i$  for  $|i - j| = 1$ , which is standard in Temperley–Lieb algebras, must be replaced by a “virtual relation” of the form  $E_i v_j E_i = E_i$ . Then we determine a compatible sequence of homomorphisms  $\{\rho_n^f : R^f[\text{VB}_n] \rightarrow \text{VTL}_n(R^f)\}_{n=1}^\infty$  (Theorem 2.9), we determine a Markov trace on the tower of algebras  $\{\text{VTL}_n(R^f)\}_{n=1}^\infty$  (Theorem 2.10), and we show that this construction leads to the  $f$ -polynomial (Theorem 2.12).

Section 3 is dedicated to the arrow polynomial. Our construction can be viewed as a labeled version of the construction of Section 2. For each  $n \geq 1$  we define an algebra  $\text{ATL}_n(R^a)$  in terms of labeled diagrams, so that  $\{\text{ATL}_n(R^a)\}_{n=1}^\infty$  is a tower of algebras. Intuitively speaking, a label represents the number of cusps in Kauffman sense that can be found on an arc. In Proposition 3.2 we determine a presentation for  $\text{ATL}_n(R^a)$ . This is a sort of labeled version of the presentation for  $\text{VTL}_n(R^f)$  given in Proposition 2.2. Then we proceed as in Section 2: we determine a compatible sequence of homomorphisms  $\{\rho_n^a : R^a[\text{VB}_n] \rightarrow \text{ATL}_n(R^a)\}_{n=1}^\infty$  (Theorem 3.10), we determine a Markov trace on the tower of algebras  $\{\text{ATL}_n(R^a)\}_{n=1}^\infty$  (Theorem 3.11), and we show that this construction leads to the arrow polynomial (Theorem 3.15).

It is known that the arrow polynomial coincides with the  $f$ -polynomial on classical links (see Miyazawa [11] and Dye–Kauffman [1]). We show that this fact has an interpretation in terms of Markov traces on Temperley–Lieb algebras. For  $n \geq 1$  we denote by  $\text{TL}_n$  the  $n$ -th standard Temperley–Lieb algebra. We show that  $\text{TL}_n$  embeds into both,  $\text{VTL}_n(R^f)$  (Proposition 2.8) and  $\text{ATL}_n(R^a)$  (Proposition 3.9), and that the restriction to  $\{\text{TL}_n\}_{n=1}^\infty$  of the Markov trace on  $\{\text{VTL}_n(R^f)\}_{n=1}^\infty$  coincides with the restriction to  $\{\text{TL}_n\}_{n=1}^\infty$  of the Markov trace on  $\{\text{ATL}_n(R^a)\}_{n=1}^\infty$  (Proposition 3.13).

**Acknowledgments** The first author is supported by the French project “AlMaRe” (ANR-19-CE40-0001-01) of the ANR.

## 2 Virtual Temperley–Lieb algebras and $f$ -polynomial

Two rings are involved in this section. The first is the ring  $R_0^f = \mathbb{Z}[z]$  of polynomials in the variable  $z$  with integer coefficients. The second is the ring  $R^f = \mathbb{Z}[A^{\pm 1}]$  of Laurent polynomials in the variable  $A$  with integer coefficients. We assume that  $R_0^f$  is embedded into  $R^f$  via the identification  $z = -A^2 - A^{-2}$ . Notice that the superscript  $f$  over  $R_0$  and  $R$  in this notation is to underline the fact that all the constructions in the present section concern the  $f$ -polynomial. In contrast, the rings in the next section, which concerns the arrow polynomial, will be denoted  $R_0^a$  and  $R^a$ .

We start recalling the definition of the  $f$ -polynomial, as it will help the reader to understand the constructions and definitions that follow after.

Define the *Kauffman bracket*  $\langle L \rangle \in R^f$  of a (non-oriented) virtual link diagram  $L$  as follows. If  $L$  has only virtual crossings, then  $\langle L \rangle = z^\ell = (-A^2 - A^{-2})^\ell$ , where  $\ell$  is the number of components of  $L$ . Suppose that  $L$  has at least one non-virtual crossing  $p$ . Then  $\langle L \rangle = A \langle L_1 \rangle + A^{-1} \langle L_2 \rangle$ , where  $L_1$  and  $L_2$  are identical to  $L$  except in a neighborhood of  $p$  where there are as shown in Figure 2.1. The *writhe* of an (oriented) virtual link diagram  $L$ , denoted  $w(L)$ , is the number of positive crossings minus the number of negative crossings. Then the  $f$ -polynomial of an (oriented) virtual link diagram  $L$  is  $f(L) = f(L)(A) = (-A^3)^{-w(L)} \langle L \rangle$ .




Figure 2.1: Relation in the Kauffman bracket

**Theorem 2.1** (Kauffman [6]) *If two virtual link diagrams  $L$  and  $L'$  are equivalent, then  $f(L) = f(L')$ .*

The  $f$ -polynomial of a virtual link  $L$ , denoted  $f(L)$ , is the  $f$ -polynomial of any of its diagrams. This is a well-defined invariant thanks to Theorem 2.1.

Our goal now is to define a Markov trace whose associated invariant is the  $f$ -polynomial. We proceed as indicated in the introduction: we pass through a tower of algebras, the tower of virtual Temperley–Lieb algebras.

Let  $n \geq 1$  be an integer. A *flat virtual  $n$ -tangle* is a collection of  $n$  disjoint pairs in  $\{0, 1\} \times \{1, \dots, n\}$ , that is, a partition of  $\{0, 1\} \times \{1, \dots, n\}$  into pairs. Let  $E = \{\alpha_1, \dots, \alpha_n\}$  be a flat virtual  $n$ -tangle. Then we graphically represent  $E$  on the plane by connecting the two ends of each  $\alpha_i$  with an arc. For example, Figure 2.2 represents the flat virtual 3-tangle  $\{\alpha_1, \alpha_2, \alpha_3\}$ , where  $\alpha_1 = \{(0, 1), (0, 2)\}$ ,  $\alpha_2 = \{(0, 3), (1, 2)\}$  and  $\alpha_3 = \{(1, 3), (1, 1)\}$ .

We denote by  $\mathcal{E}_n$  the set of flat virtual  $n$ -tangles, and by  $\text{VTL}_n$  the free  $R_0^f$ -module freely generated by  $\mathcal{E}_n$ . We define a multiplication in  $\text{VTL}_n$  as follows. Let  $E$  and  $E'$  be two flat virtual  $n$ -tangles. By concatenating the diagrams of  $E$  and  $E'$  we get a family of closed curves and  $n$  arcs. These  $n$  arcs determine a partition of  $\{0, 1\} \times \{1, \dots, n\}$  into pairs, that is, a flat virtual  $n$ -tangle that we denote by  $E * E'$ . Let  $m$  be the number of obtained closed curves. Then we set  $EE' = z^m (E * E')$ . It is easily





Figure 2.2: Flat virtual tangle

checked that  $\text{VTL}_n$  endowed with this multiplication is an (unitary and associative) algebra that we call the  $n$ -th *virtual Temperley–Lieb algebra*.

**Example** On the left hand side of Figure 2.3 are illustrated the diagrams of two flat virtual 4-tangles,  $E$  and  $E'$ , and on the right hand side a diagram of  $E * E'$ . In this example, by concatenating the diagrams of  $E$  and  $E'$  we get only one closed curve, hence  $m = 1$  and  $EE' = z(E * E')$ .

Figure 2.3: Multiplication in  $\text{VTL}_n$ 

**Remark** It is easily seen that the embedding  $\mathcal{E}_n \rightarrow \mathcal{E}_{n+1}$  which sends each  $E \in \mathcal{E}_n$  to  $E \cup \{(0, n + 1), (1, n + 1)\}$  induces an injective homomorphism  $\text{VTL}_n \hookrightarrow \text{VTL}_{n+1}$ , for all  $n \geq 1$ . So, we have a tower of algebras  $\{\text{VTL}_n\}_{n=1}^\infty$ .

**Proposition 2.2** *Let  $n \geq 2$ . Then  $\text{VTL}_n$  has a presentation with generators  $E_1, \dots, E_{n-1}, v_1, \dots, v_{n-1}$  and relations*


$$\begin{aligned} E_i^2 &= zE_i, \quad v_i^2 = 1, \quad E_i v_i = v_i E_i = E_i, \quad \text{for } 1 \leq i \leq n-1, \\ E_i E_j &= E_j E_i, \quad v_i v_j = v_j v_i, \quad v_i E_j = E_j v_i, \quad \text{for } |i-j| \geq 2, \\ E_i v_j E_i &= E_i, \quad v_i v_j v_i = v_j v_i v_j, \quad v_i v_j E_i = E_j v_i v_j, \quad \text{for } |i-j| = 1. \end{aligned}$$

The generators  $E_i$  and  $v_i$  are illustrated in Figure 2.4.

The next four lemmas are preliminaries to the proof of Proposition 2.2. Let  $A_n$  be the algebra over  $R_0^f$  defined by the presentation with generators  $X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}$  and relations

$$\begin{aligned} X_i^2 &= zX_i, \quad y_i^2 = 1, \quad X_i y_i = y_i X_i = X_i, \quad \text{for } 1 \leq i \leq n-1, \\ X_i X_j &= X_j X_i, \quad y_i y_j = y_j y_i, \quad y_i X_j = X_j y_i, \quad \text{for } |i-j| \geq 2, \\ X_i y_j X_i &= X_i, \quad y_i y_j y_i = y_j y_i y_j, \quad y_i y_j X_i = X_j y_i y_j, \quad \text{for } |i-j| = 1. \end{aligned}$$

It is easily checked using diagrammatic calculation that there is a homomorphism  $\varphi : A_n \rightarrow \text{VTL}_n$  which sends  $X_i$  to  $E_i$  and  $y_i$  to  $v_i$  for all  $i \in \{1, \dots, n-1\}$ .

Figure 2.4: Generators of  $VTL_n$ 

**Lemma 2.3** *The following relations hold in  $A_n$ .*

$$\begin{aligned} X_i X_j y_i y_j &= X_i \quad \text{for } |i - j| = 1, \\ X_i X_j X_i &= X_i \quad \text{for } |i - j| = 1, \\ X_i X_j &= y_j y_i X_j \quad \text{for } |i - j| = 1, \\ y_{i+1} y_{i+2} y_i y_{i+1} X_i X_{i+2} &= X_i X_{i+2} \quad \text{for } 1 \leq i \leq n - 3. \end{aligned}$$

**Proof** Let  $i, j \in \{1, \dots, n - 1\}$  such that  $|i - j| = 1$ . Then

$$\begin{aligned} X_i X_j y_i y_j &= X_i y_i y_j X_i = X_i y_j X_i = X_i, \\ X_i X_j X_i &= X_i X_j y_i y_j y_i X_i = X_i y_j y_i X_i = X_i y_j X_i = X_i, \\ X_i X_j &= y_j y_i y_i y_j X_i X_j = y_j y_i X_j y_i y_j X_j = y_j y_i X_j y_i X_j = y_j y_i X_j. \end{aligned}$$

Let  $i \in \{1, \dots, n - 3\}$ . Then

$$\begin{aligned} y_{i+1} y_{i+2} y_i y_{i+1} X_i X_{i+2} &= y_{i+1} y_{i+2} X_{i+1} X_i X_{i+2} = X_{i+2} X_{i+1} X_i X_{i+2} = \\ &X_{i+2} X_{i+1} X_{i+2} X_i = X_{i+2} X_i = X_i X_{i+2}. \end{aligned}$$

□

We denote by  $\mathcal{U}(A_n)$  the group of units of  $A_n$  and by  $\mathfrak{S}_n$  the  $n$ -th symmetric group. We have a homomorphism  $\iota : \mathfrak{S}_n \rightarrow \mathcal{U}(A_n)$  which sends  $s_i$  to  $y_i$  for all  $i \in \{1, \dots, n - 1\}$ , where  $s_i = (i, i + 1)$ . Let  $n_0$  be the integer part of  $\frac{n}{2}$ . Let  $p \in \{0, 1, \dots, n_0\}$ . We set  $B_p = X_1 X_3 \cdots X_{2p-1}$  if  $p \neq 0$ , and  $B_p = B_0 = 1$  if  $p = 0$ . We denote by  $U_{1,p}$  the set of  $w \in \mathfrak{S}_n$  satisfying

$$\begin{aligned} w(1) &< w(3) < \cdots < w(2p - 1), \\ w(2i - 1) &< w(2i) \quad \text{for } 1 \leq i \leq p, \\ w(2p + 1) &< w(2p + 2) < \cdots < w(n), \end{aligned}$$

and we denote by  $U_{2,p}$  the set of  $w \in \mathfrak{S}_n$  satisfying

$$\begin{aligned} w(1) &< w(3) < \cdots < w(2p - 1), \\ w(2i - 1) &< w(2i) \quad \text{for } 1 \leq i \leq p. \end{aligned}$$

Then we set

$$\mathcal{B}_p = \{\iota(w_1) B_p \iota(w_2^{-1}) \mid w_1 \in U_{1,p}, w_2 \in U_{2,p}\},$$



for  $0 \leq p \leq n_0$ , and

$$\mathcal{B} = \bigcup_{p=0}^{n_0} \mathcal{B}_p.$$

Let  $E \in \mathcal{E}_n$ . We can write  $E$  in the form  $E = \{\alpha_1, \dots, \alpha_p, \alpha'_1, \dots, \alpha'_p, \beta_1, \dots, \beta_q\}$ , where

- each  $\alpha_i$  is of the form  $\alpha_i = \{(0, a_i), (0, b_i)\}$ , with  $1 \leq a_1 < a_2 < \dots < a_p \leq n$ , and  $a_i < b_i$  for all  $i \in \{1, \dots, p\}$ ;
- each  $\alpha'_i$  is of the form  $\alpha'_i = \{(1, a'_i), (1, b'_i)\}$ , with  $1 \leq a'_1 < a'_2 < \dots < a'_p \leq n$ , and  $a'_i < b'_i$  for all  $i \in \{1, \dots, p\}$ ;
- each  $\beta_j$  is of the form  $\beta_j = \{(0, c_j), (1, d_j)\}$ , with  $1 \leq c_1 < c_2 < \dots < c_q \leq n$ , and  $2p + q = n$ .

We define  $w_1 \in \mathfrak{S}_n$  by  $w_1(2i - 1) = a_i$  and  $w_1(2i) = b_i$  for all  $i \in \{1, \dots, p\}$ , and  $w_1(2p + j) = c_j$  for all  $j \in \{1, \dots, q\}$ . Similarly, we define  $w_2 \in \mathfrak{S}_n$  by  $w_2(2i - 1) = a'_i$  and  $w_2(2i) = b'_i$  for all  $i \in \{1, \dots, p\}$ , and  $w_2(2p + j) = d_j$  for all  $j \in \{1, \dots, q\}$ . We see that  $w_1 \in U_{1,p}$ ,  $w_2 \in U_{2,p}$ , and  $E = \varphi(\iota(w_1)B_p \iota(w_2^{-1}))$ . Moreover, such a form is unique for each  $E \in \mathcal{E}_n$ , and we have  $\varphi(Y) \in \mathcal{E}_n$  for all  $Y \in \mathcal{B}$ , hence  $\varphi$  restricts to a bijection from  $\mathcal{B}$  to  $\mathcal{E}_n$ .

So, in order to prove Proposition 2.2, it suffices to show that  $\mathcal{B}$  spans  $A_n$  as a  $R_0^f$ -module. We denote by  $\mathcal{M}$  the submonoid of  $A_n$  generated by  $X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}$ , that is, the set of finite products of elements in  $\{X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}\}$ . By definition  $\mathcal{M}$  spans  $A_n$  as a  $R_0^f$ -module, hence we only need to show that  $\mathcal{M}$  is contained in the  $R_0^f$ -submodule  $\text{Span}_{R_0^f}(\mathcal{B})$  of  $A_n$  spanned by  $\mathcal{B}$ .

**Lemma 2.4** *Let  $w_1, w_2 \in \mathfrak{S}_n$  and  $p \in \{0, 1, \dots, n_0\}$ . Then  $\iota(w_1)B_p \iota(w_2^{-1}) \in \mathcal{B}$ .*

**Proof** Let  $i \in \{1, \dots, p\}$  such that  $w_1(2i - 1) = b_i > w_1(2i) = a_i$ . By applying the relation  $y_{2i-1}X_{2i-1} = X_{2i-1}$  we can replace  $w_1$  with  $w_1s_{2i-1}$ , and then  $w_1(2i - 1) = a_i < w_1(2i) = b_i$ . So, we can assume that  $w_1(2i - 1) < w_1(2i)$  for all  $i \in \{1, \dots, p\}$ . Let  $i \in \{1, \dots, p - 1\}$  such that  $w_1(2i - 1) = a_{i+1} > w_1(2i + 1) = a_i$ . By applying the relation  $y_{2i}y_{2i-1}y_{2i+1}y_{2i}X_{2i-1}X_{2i+1} = X_{2i-1}X_{2i+1}$  we can replace  $w_1$  with  $w_1s_{2i}s_{2i-1}s_{2i+1}s_{2i}$ , and then we have  $w_1(2i - 1) = a_i < w_1(2i + 1) = a_{i+1}$  while keeping the inequalities  $w_1(2i - 1) < w_1(2i)$  and  $w_1(2i + 1) < w_1(2i + 2)$ . So, we can also assume that  $w_1(1) < w_1(3) < \dots < w_1(2p - 1)$ . Set  $q = n - 2p$ . Let  $j \in \{1, \dots, q - 1\}$  such that  $w_1(2p + j) = c_{j+1} > w_1(2p + j + 1) = c_j$ . By applying the relations  $y_{2p+j}X_{2i-1} = X_{2i-1}y_{2p+j}$  for  $i \in \{1, \dots, p\}$  we can replace  $w_1$  with  $w_1s_{2p+j}$  and  $w_2$  with  $w_2s_{2p+j}$ , and then  $w_1(2p + j) = c_j < w_1(2p + j + 1) = c_{j+1}$ . So, we can also assume that  $w_1(2p + 1) < w_1(2p + 2) < \dots < w_1(n)$ , that is,  $w_1 \in U_{1,p}$ . We use the same argument to show that  $w_2$  can be replaced with some  $w'_2 \in U_{2,p}$ . So,  $\iota(w_1)B_p \iota(w_2^{-1}) \in \mathcal{B}$ .  $\square$

**Lemma 2.5** *Let  $a, b \in \{1, \dots, n - 1\}$ ,  $a \leq b$ , and  $p \in \{0, 1, \dots, n_0\}$ . Then  $X_a y_{a+1} \dots y_b B_p \in \text{Span}_{R_0^f}(\mathcal{B})$ .*

**Proof** Suppose  $a \geq 2p + 1$  (which is always true if  $p = 0$ ). Then

$$\begin{aligned} X_a y_{a+1} \cdots y_b B_p &= X_a B_p y_{a+1} \cdots y_b = \\ &X_a (y_{a-1} y_a) (y_{a-2} y_{a-1}) \cdots (y_{2p+1} y_{2p+2}) (y_{2p+2} y_{2p+1}) \cdots (y_{a-1} y_{a-2}) (y_a y_{a-1}) B_p y_{a+1} \cdots y_b = \\ &(y_{a-1} y_a) (y_{a-2} y_{a-1}) \cdots (y_{2p+1} y_{2p+2}) X_{2p+1} B_p (y_{2p+2} y_{2p+1}) \cdots (y_{a-1} y_{a-2}) (y_a y_{a-1}) y_{a+1} \cdots y_b = \\ &(y_{a-1} y_a) (y_{a-2} y_{a-1}) \cdots (y_{2p+1} y_{2p+2}) B_{p+1} (y_{2p+2} y_{2p+1}) \cdots (y_{a-1} y_{a-2}) (y_a y_{a-1}) y_{a+1} \cdots y_b = \\ &\iota(w_1) B_{p+1} \iota(w_2^{-1}) \in \mathcal{B}, \end{aligned}$$

where  $w_1 = (s_{a-1} s_a) \cdots (s_{2p+1} s_{2p+2})$  and  $w_2 = s_b \cdots s_{a+1} (s_{a-1} s_a) \cdots (s_{2p+1} s_{2p+2})$ . Suppose  $a \leq 2p$ ,  $a$  is odd, and  $a = b$ . Let  $c \in \{1, \dots, p\}$  such that  $a = 2c - 1$ . Then

$$\begin{aligned} X_a y_{a+1} \cdots y_b B_p &= X_{2c-1}^2 X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &z X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = z B_p \in \text{Span}_{R'_0}(\mathcal{B}). \end{aligned}$$

Suppose  $a \leq 2p$ ,  $a$  is odd, and  $a < b$ . Let  $c \in \{1, \dots, p\}$  such that  $a = 2c - 1$ . Then

$$\begin{aligned} X_a y_{a+1} \cdots y_b B_p &= X_{2c-1} y_{2c} \cdots y_b X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &X_{2c-1} y_{2c} X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c-1} X_{2c+1} \cdots X_{2p-1} = \iota(s_{2c+1} \cdots s_b) B_p \in \mathcal{B}. \end{aligned}$$

Suppose  $a \leq 2p$  and  $a$  is even. Let  $c \in \{1, \dots, p\}$  such that  $a = 2c$ . Then

$$\begin{aligned} X_a y_{a+1} \cdots y_b B_p &= X_{2c} y_{2c+1} \cdots y_b X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &X_{2c} X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &y_{2c-1} y_{2c} X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &y_{2c-1} y_{2c} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c-1} X_{2c+1} \cdots X_{2p-1} = \iota(s_{2c-1} s_{2c} s_{2c+1} \cdots s_b) B_p \in \mathcal{B}. \end{aligned}$$

□

**Lemma 2.6** Let  $p \in \{0, 1, \dots, n_0\}$  and  $w \in \mathfrak{S}_n$ . Then  $X_1 \iota(w) B_p \in \text{Span}_{R'_0}(\mathcal{B})$ .

**Proof** There exist  $a \in \{1, \dots, n-1\}$ ,  $b \in \{0, 1, \dots, n-1\}$  and  $w_1 \in \langle s_3, \dots, s_{n-1} \rangle$  such that  $w = w_1 s_2 s_3 \cdots s_a s_1 s_2 \cdots s_b$ . Suppose  $a \leq b$ . Then

$$\begin{aligned} X_1 \iota(w) B_p &= \iota(w_1) X_1 y_2 \cdots y_a y_1 \cdots y_{a-1} y_a y_{a+1} \cdots y_b B_p = \\ &\iota(w_1) X_1 (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) y_a y_{a+1} \cdots y_b B_p = \\ &\iota(w_1) (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) X_a y_a y_{a+1} \cdots y_b B_p = \\ &\iota(w_1 (s_2 s_1) (s_3 s_2) \cdots (s_a s_{a-1})) X_a y_{a+1} \cdots y_b B_p. \end{aligned}$$

We know by Lemma 2.5 that  $X_a y_{a+1} \cdots y_b B_p \in \text{Span}_{R'_0}(\mathcal{B})$ , hence, by Lemma 2.4,  $X_1 \iota(w) B_p \in \text{Span}_{R'_0}(\mathcal{B})$ . Suppose  $a > b$ . Then

$$\begin{aligned} X_1 \iota(w) B_p &= \iota(w_1) X_1 y_2 \cdots y_b y_{b+1} \cdots y_a y_1 \cdots y_b B_p = \\ &\iota(w_1) X_1 (y_2 y_1) (y_3 y_2) \cdots (y_{b+1} y_b) y_{b+2} \cdots y_a B_p = \\ &\iota(w_1) (y_2 y_1) (y_3 y_2) \cdots (y_{b+1} y_b) X_{b+1} y_{b+2} \cdots y_a B_p = \\ &\iota(w_1 (s_2 s_1) (s_3 s_2) \cdots (s_{b+1} s_b)) X_{b+1} y_{b+2} \cdots y_a B_p. \end{aligned}$$

We know by Lemma 2.5 that  $X_{b+1}y_{b+2}\cdots y_a B_p \in \text{Span}_{R_0^f}(\mathcal{B})$ , hence, by Lemma 2.4,  $X_1 \iota(w) B_p \in \text{Span}_{R_0^f}(\mathcal{B})$ .  $\square$

**Proof of Proposition 2.2** As pointed out above, it suffices to show that the monoid  $\mathcal{M}$  is contained in  $\text{Span}_{R_0^f}(\mathcal{B})$ . Let  $Y \in \mathcal{M}$ . By using the relations  $y_i y_j X_i = X_j y_i y_j$  for  $|i - j| = 1$ , we see that  $Y$  can be written in the form  $Y = \iota(w_0) X_1 \iota(w_1) X_1 \cdots X_1 \iota(w_k)$  where  $k \geq 0$  and  $w_0, w_1, \dots, w_k \in \mathfrak{S}_n$ . We prove that  $Y \in \text{Span}_{R_0^f}(\mathcal{B})$  by induction on  $k$ . The case  $k = 0$  is trivial and the case  $k = 1$  follows from Lemma 2.4. So, we can assume that  $k \geq 2$  and that the inductive hypothesis holds. By the inductive hypothesis  $\iota(w_1) X_1 \cdots X_1 \iota(w_k) \in \text{Span}_{R_0^f}(\mathcal{B})$ , thus we just need to prove that  $\iota(w_0) X_1 \iota(w'_1) B_p \iota(w'_2)^{-1} \in \text{Span}_{R_0^f}(\mathcal{B})$  for all  $p \in \{0, 1, \dots, n_0\}$  and all  $w'_1, w'_2 \in \mathfrak{S}_n$ . We know by Lemma 2.6 that  $X_1 \iota(w'_1) B_p \in \text{Span}_{R_0^f}(\mathcal{B})$ , hence, by Lemma 2.4,  $\iota(w_0) X_1 \iota(w'_1) B_p \iota(w'_2)^{-1} \in \text{Span}_{R_0^f}(\mathcal{B})$ . This concludes the proof of Proposition 2.2.  $\square$

We have seen that the relation  $E_i E_j E_i = E_i$  holds for  $|i - j| = 1$  in  $\text{VTL}_n$  (see Lemma 2.3). However, we cannot replace the relation  $E_i v_j E_i = E_i$  with the relation  $E_i E_j E_i = E_i$  in the presentation of  $\text{VTL}_n$ . Indeed:

**Proposition 2.7** *Let  $n \geq 3$  and let  $\text{VTL}'_n$  be the algebra over  $R_0^f$  defined by the presentation with generators  $E'_1, \dots, E'_{n-1}, v'_1, \dots, v'_{n-1}$  and relations*

$$\begin{aligned} E_i'^2 &= z E_i', \quad v_i'^2 = 1, \quad E_i' v_i' = v_i' E_i' = E_i', \quad \text{for } 1 \leq i \leq n-1, \\ E_i' E_j' &= E_j' E_i', \quad v_i' v_j' = v_j' v_i', \quad v_i' E_j' = E_j' v_i', \quad \text{for } |i - j| \geq 2, \\ E_i' E_j' E_i' &= E_i', \quad v_i' v_j' v_i' = v_j' v_i' v_j', \quad v_i' v_j' E_i' = E_j' v_i' v_j', \quad \text{for } |i - j| = 1. \end{aligned}$$

Let  $\varphi : \text{VTL}'_n \rightarrow \text{VTL}_n$  be the homomorphism that sends  $E_i'$  to  $E_i$  and  $v_i'$  to  $v_i$  for all  $i \in \{1, \dots, n-1\}$ . Then  $\varphi$  is surjective but not injective.

**Proof** By definition  $E_1, \dots, E_{n-1}, v_1, \dots, v_{n-1}$  belong to the image of  $\varphi$ . Since these elements generate  $\text{VTL}_n$ , the homomorphism  $\varphi$  is surjective. Let  $C_2 = \{\pm 1\}$  be the cyclic group of order 2 and let  $\mathbb{Z}[C_2]$  be the group algebra of  $C_2$ . It is easily checked with the presentation of  $\text{VTL}'_n$  that there is a ring homomorphism  $\theta : \text{VTL}'_n \rightarrow \mathbb{Z}[C_2]$  satisfying  $\theta(E_i') = -1$  for all  $i \in \{1, \dots, n-1\}$ ,  $\theta(v_i') = 1$  for all  $i \in \{1, \dots, n-1\}$ , and  $\theta(z) = -1$ . Let  $i, j \in \{1, \dots, n-1\}$  such that  $|i - j| = 1$ . Then  $\theta(E_i' v_j' E_i') = 1$  and  $\theta(E_i') = -1$ , hence the relation  $E_i' v_j' E_i' = E_i'$  does not hold in  $\text{VTL}'_n$ . So,  $\varphi$  is not injective.  $\square$

Let  $n \geq 1$ . Recall that  $V_n = \{0, 1\} \times \{1, \dots, n\}$  is ordered by  $(0, 1) < (0, 2) < \cdots < (0, n) < (1, n) < \cdots < (1, 2) < (1, 1)$ . Let  $E$  be a flat virtual  $n$ -tangle. Let  $\gamma_1 = \{x_1, y_1\}, \gamma_2 = \{x_2, y_2\} \in E$  such that  $x_1 < y_1, x_2 < y_2$ , and  $x_1 < x_2$ . We say that  $\gamma_1$  *crosses*  $\gamma_2$  if  $x_1 < x_2 < y_1 < y_2$ . We say that  $E$  is *non-crossing* if there are no two elements in  $E$  that cross. Equivalently, a flat  $n$ -tangle is non-crossing if and only if it has a graphical representation with  $n$  disjoint arcs. We denote by  $\mathcal{E}_n^0$  the set of non-crossing flat virtual  $n$ -tangles.

Let  $n \geq 2$ . Recall that the *Temperley–Lieb algebra*  $\mathrm{TL}_n$  is the algebra over  $R_0^f$  defined by the presentation with generators  $E_1, \dots, E_{n-1}$  and relations

$$\begin{aligned} E_i^2 &= zE_i \text{ for } 1 \leq i \leq n-1, \quad E_iE_j = E_jE_i \text{ for } |i-j| \geq 2, \\ E_iE_jE_i &= E_i \text{ for } |i-j| = 1. \end{aligned}$$

The following is proved in Kauffman [5].

**Proposition 2.8** (Kauffman [5]) *Let  $n \geq 2$ . The homomorphism  $\mathrm{TL}_n \rightarrow \mathrm{VTL}_n$  which sends  $E_i$  to  $E_i$  for all  $i \in \{1, \dots, n-1\}$  is injective and its image is the  $R_0^f$ -submodule of  $\mathrm{VTL}_n$  freely generated by  $\mathcal{E}_n^0$ .*

Recall that  $R^f = \mathbb{Z}[A^{\pm 1}]$  denotes the algebra of Laurent polynomials in the variable  $A$ , and that  $R_0^f = \mathbb{Z}[z]$  is a subalgebra of  $R^f$  via the identification  $z = -A^2 - A^{-2}$ . For each  $n \geq 1$  we set  $\mathrm{VTL}_n(R^f) = R^f \otimes \mathrm{VTL}_n$ . This is a  $R^f$ -algebra and it is a free  $R^f$ -module freely generated by  $\mathcal{E}_n$ .

**Theorem 2.9** *Let  $n \geq 1$ . There exists a homomorphism  $\rho_n^f : R^f[\mathrm{VB}_n] \rightarrow \mathrm{VTL}_n(R^f)$  which sends  $\sigma_i$  to  $-A^{-2}1 - A^{-4}E_i$  and  $\tau_i$  to  $v_i$  for all  $i \in \{1, \dots, n-1\}$ .*

**Proof** We set  $S_i = -A^{-2}1 - A^{-4}E_i$ . We have

$$\begin{aligned} (-A^{-2}1 - A^{-4}E_i)(-A^{-2}1 - A^{-4}E_i) &= 1 + (A^2 + A^{-2})E_i + E_i^2 = \\ &= 1 + (A^2 + A^{-2})E_i + (-A^2 - A^{-2})E_i = 1. \end{aligned}$$

So,  $S_i$  is invertible and its inverse is  $-A^21 - A^4E_i$ . The element  $v_i$  is also invertible since  $v_i^2 = 1$ . It remains to verify that the following relations hold.

$$\begin{aligned} v_i^2 &= 1, \quad \text{for } 1 \leq i \leq n-1, \\ S_iS_j &= S_jS_i, \quad v_iv_j = v_jv_i, \quad v_iS_j = S_jv_i, \quad \text{for } |i-j| \geq 2, \\ S_iS_jS_i &= S_jS_iS_j, \quad v_iv_jv_i = v_jv_iv_j, \quad v_iv_jS_i = S_jv_iv_j, \quad \text{for } |i-j| = 1. \end{aligned}$$

The only of these relations which is not trivial is  $S_iS_jS_i = S_jS_iS_j$  for  $|i-j| = 1$ . Suppose  $|i-j| = 1$ . Then

$$\begin{aligned} S_iS_jS_i &= (-A^{-2}1 - A^{-4}E_i)(-A^{-2}1 - A^{-4}E_j)(-A^{-2}1 - A^{-4}E_i) = \\ &= -A^{-6}1 - A^{-8}E_i - A^{-8}E_j - A^{-10}E_iE_j - A^{-8}E_i - A^{-10}E_i^2 - A^{-10}E_jE_i - A^{-12}E_iE_jE_i = \\ &= -A^{-6}1 - 2A^{-8}E_i - A^{-8}E_j - A^{-10}E_iE_j - A^{-10}(-A^2 - A^{-2})E_i - A^{-10}E_jE_i - A^{-12}E_i = \\ &= -A^{-6}1 - A^{-8}E_i - A^{-8}E_j - A^{-10}E_iE_j - A^{-10}E_jE_i. \end{aligned}$$

By symmetry we also have  $S_jS_iS_j = -A^{-6}1 - A^{-8}E_i - A^{-8}E_j - A^{-10}E_iE_j - A^{-10}E_jE_i$ , hence  $S_iS_jS_i = S_jS_iS_j$ .  $\square$

**Remark** (1) Let  $B_n$  be the braid group on  $n$  strands and let  $\mathrm{TL}_n$  be the  $n$ -th Temperley–Lieb algebra. Then  $\rho_n^f(\beta) \in \mathrm{TL}_n(R^f)$  for all  $\beta \in B_n$ , where  $\mathrm{TL}_n(R^f) = R^f \otimes \mathrm{TL}_n \subset \mathrm{VTL}_n(R^f)$ .

(2) The sequence of homomorphisms  $\{\rho_n^f : R^f[\mathrm{VB}_n] \rightarrow \mathrm{VTL}_n(R^f)\}_{n=1}^\infty$  is compatible with the tower of algebras  $\{\mathrm{VTL}_n(R^f)\}_{n=1}^\infty$ .

- (3) Setting  $\rho_n^f(\sigma_i) = -A^{-2}1 - A^{-4}E_i$  instead of  $\rho_n^f(\sigma_i) = A1 + A^{-1}E_i$ , as an informed reader may expect, allows to include in  $\rho_n^f$  the corrective with the writhe and to define directly the  $f$ -polynomial without passing through the Kauffman bracket.

Let  $E$  be a flat virtual  $n$ -tangle. By connecting with an arc the point  $(0, i)$  with the point  $(1, i)$  for all  $i \in \{1, \dots, n\}$  in a diagram of  $E$  we obtain a family of closed curves that we call the *closure of the diagram* of  $E$ . We denote by  $t_n(E)$  the number of closed curves in this family, and we set  $T_n^{ff}(E) = z^{t_n(E)} = (-A^2 - A^{-2})^{t_n(E)}$ . Then we define  $T_n^{ff} : \text{VTL}_n(\mathcal{R}^f) \rightarrow \mathcal{R}^f$  by extending linearly the map  $T_n^{ff} : \mathcal{E}_n \rightarrow \mathcal{R}^f$ .

**Example** In Figure 2.5 is illustrated the closure of the flat virtual tangle  $E$  of Figure 2.2. In this case we have  $t_n(E) = 1$ , and therefore  $T_n^{ff}(E) = z = -A^2 - A^{-2}$ .




Figure 2.5: Closure of a flat virtual tangle

**Theorem 2.10** The sequence  $\{T_n^{ff} : \text{VTL}_n(\mathcal{R}^f) \rightarrow \mathcal{R}^f\}_{n=1}^{\infty}$  is a Markov trace.

**Proof** For each  $n \geq 2$  and each  $i \in \{1, \dots, n-1\}$  we set  $S_i = -A^{-2}1 - A^{-4}E_i$ . We have to show that the following equalities hold.

- (1)  $T_n^{ff}(xy) = T_n^{ff}(yx)$  for all  $n \geq 1$  and all  $x, y \in \text{VTL}_n(\mathcal{R}^f)$ .
- (2)  $T_n^{ff}(x) = T_{n+1}^{ff}(xS_n) = T_{n+1}^{ff}(xS_n^{-1}) = T_{n+1}^{ff}(xv_n)$  for all  $n \geq 1$  and all  $x \in \text{VTL}_n(\mathcal{R}^f)$ .
- (3)  $T_n^{ff}(x) = T_{n+1}^{ff}(xS_n^{-1}v_{n-1}S_n)$  for all  $n \geq 2$  and all  $x \in \text{VTL}_n(\mathcal{R}^f)$ .
- (4)  $T_n^{ff}(x) = T_{n+1}^{ff}(xv_nv_{n-1}S_{n-1}v_nS_{n-1}^{-1}v_{n-1}v_n)$  for all  $n \geq 2$  and all  $x \in \text{VTL}_n(\mathcal{R}^f)$ .

*Proof of (1).* We can assume that  $x = E$  and  $y = E'$  are flat virtual  $n$ -tangles. We fix graphical representations of  $E$  and  $E'$ . By concatenating the graphical representation of  $E$  on the left with that of  $E'$  on the right and then connecting with an arc the point  $(0, i)$  of  $E$  to the point  $(1, i)$  of  $E'$  for all  $i \in \{1, \dots, n\}$  we get a family of closed curves immersed in the plane, denoted  $\widehat{E \sqcup E'}$ . If  $m$  is the number of closed curves in this family, then  $T_n^{ff}(EE') = z^m$ . We can choose the  $n$  arcs connecting the points  $(0, i)$  of  $E$  to the point  $(1, i)$  of  $E'$  pairwise disjoint and disjoint from the concatenation of  $E$  and  $E'$ . In that case,  $\widehat{E \sqcup E'}$  is isotopic to  $\widehat{E' \sqcup E}$ , hence  $\widehat{E' \sqcup E}$  has the same number of closed curves as  $\widehat{E \sqcup E'}$ , and therefore  $T_n^{ff}(E'E) = z^m = T_n^{ff}(EE')$ .

*Proof of (2).* We can assume that  $x = E$  is a flat virtual  $n$ -tangle. We see in Figure 2.6 that the following equalities hold

$$T_{n+1}^{ff}(E) = zT_n^{ff}(E), \quad T_{n+1}^{ff}(EE_n) = T_n^{ff}(E), \quad T_{n+1}^{ff}(Ev_n) = T_n^{ff}(E).$$

Recall that  $S_n^{-1} = -A^2 1 - A^4 E_n$  (see the proof of Theorem 2.9). We saw in Figure 2.6 that  $T_{n+1}^{ff}(Ev_n) = T_n^{ff}(E)$ . On the other hand,

$$\begin{aligned} T_{n+1}^{ff}(ES_n) &= -A^{-2}T_{n+1}^{ff}(E) - A^{-4}T_{n+1}^{ff}(EE_n) = (-A^{-2}z - A^{-4})T_n^{ff}(E) = T_n^{ff}(E), \\ T_{n+1}^{ff}(ES_n^{-1}) &= -A^2T_{n+1}^{ff}(E) - A^4T_{n+1}^{ff}(EE_n) = (-A^2z - A^4)T_n^{ff}(E) = T_n^{ff}(E). \end{aligned}$$




Figure 2.6:  $n + 1$ -closures of  $E$ ,  $EE_n$  and  $Ev_n$

*Proof of (3).* We can again assume that  $x = E$  is a flat virtual  $n$ -tangle. We have

$$\begin{aligned} ES_n^{-1}v_{n-1}S_n &= E(-A^2 1 - A^4 E_n)v_{n-1}(-A^{-2} 1 - A^{-4} E_n) = \\ &= Ev_{n-1} + A^2 EE_n v_{n-1} + A^{-2} Ev_{n-1} E_n + EE_n v_{n-1} E_n = \\ &= Ev_{n-1} + A^2 EE_n v_{n-1} + A^{-2} Ev_{n-1} E_n + EE_n. \end{aligned}$$

Hence, by the above

$$\begin{aligned} T_{n+1}^{ff}(ES_n^{-1}v_{n-1}S_n) &= \\ T_{n+1}^{ff}(Ev_{n-1}) + A^{-2}T_{n+1}^{ff}(Ev_{n-1}E_n) + A^2T_{n+1}^{ff}(EE_n v_{n-1}) + T_{n+1}^{ff}(EE_n) &= \\ zT_n^{ff}(Ev_{n-1}) + A^{-2}T_n^{ff}(Ev_{n-1}) + A^2T_{n+1}^{ff}(v_{n-1}EE_n) + T_n^{ff}(E) &= \\ (-A^2 - A^{-2})T_n^{ff}(Ev_{n-1}) + A^{-2}T_n^{ff}(Ev_{n-1}) + A^2T_n^{ff}(v_{n-1}E) + T_n^{ff}(E) &= \\ (-A^2 - A^{-2})T_n^{ff}(Ev_{n-1}) + A^{-2}T_n^{ff}(Ev_{n-1}) + A^2T_n^{ff}(Ev_{n-1}) + T_n^{ff}(E) &= T_n^{ff}(E). \end{aligned}$$

*Proof of (4).* Again, we can assume that  $x = E$  is a flat virtual  $n$ -tangle. We have

$$\begin{aligned} Ev_n v_{n-1} S_{n-1} v_n S_{n-1}^{-1} v_{n-1} v_n &= Ev_n v_{n-1} (-A^{-2} 1 - A^{-4} E_{n-1}) v_n (-A^2 1 - A^4 E_{n-1}) v_{n-1} v_n = \\ &= (Ev_n v_{n-1} v_n v_{n-1} v_n) + A^2 (Ev_n v_{n-1} v_n E_{n-1} v_{n-1} v_n) + \\ &= A^{-2} (Ev_n v_{n-1} E_{n-1} v_n v_{n-1} v_n) + (Ev_n v_{n-1} E_{n-1} v_n E_{n-1} v_{n-1} v_n) = \\ &= (Ev_{n-1} v_n v_{n-1} v_{n-1} v_n) + A^2 (Ev_n v_{n-1} v_n E_{n-1} v_n) + \\ &= A^{-2} (Ev_n E_{n-1} v_n v_{n-1} v_n) + (Ev_n E_{n-1} v_n E_{n-1} v_n) = \\ &= (Ev_{n-1}) + A^2 (Ev_n v_{n-1} v_{n-1} E_n v_{n-1}) + A^{-2} (Ev_{n-1} E_n v_{n-1} v_{n-1} v_n) + (Ev_n E_{n-1} v_n) = \\ &= (Ev_{n-1}) + A^2 (EE_n v_{n-1}) + A^{-2} (Ev_{n-1} E_n) + (Ev_{n-1} E_n v_{n-1}). \end{aligned}$$

Hence, by the above

$$\begin{aligned}
& T_{n+1}^{ff}(Ev_n v_{n-1} S_{n-1} v_n S_{n-1}^{-1} v_{n-1} v_n) = \\
& T_{n+1}^{ff}(Ev_{n-1}) + A^2 T_{n+1}^{ff}(EE_n v_{n-1}) + A^{-2} T_{n+1}^{ff}(Ev_{n-1} E_n) + T_{n+1}^{ff}(Ev_{n-1} E_n v_{n-1}) = \\
& z T_n^{ff}(Ev_{n-1}) + A^2 T_{n+1}^{ff}(v_{n-1} EE_n) + A^{-2} T_n^{ff}(Ev_{n-1}) + T_{n+1}^{ff}(v_{n-1} Ev_{n-1} E_n) = \\
& (-A^2 - A^{-2}) T_n^{ff}(Ev_{n-1}) + A^2 T_n^{ff}(v_{n-1} E) + A^{-2} T_n^{ff}(Ev_{n-1}) + T_n^{ff}(v_{n-1} Ev_{n-1}) = \\
& (-A^2 - A^{-2}) T_n^{ff}(Ev_{n-1}) + A^2 T_n^{ff}(Ev_{n-1}) + A^{-2} T_n^{ff}(Ev_{n-1}) + T_n^{ff}(Ev_{n-1} v_{n-1}) = T_n^{ff}(E). \quad \square
\end{aligned}$$

**Corollary 2.11** For each  $n \geq 1$  we set  $T_n^f = T_n^{ff} \circ \rho_n^f : R^f[\text{VB}_n] \rightarrow R^f$ . Then  $\{T_n^f : R^f[\text{VB}_n] \rightarrow R^f\}_{n=1}^\infty$  is a Markov trace.

Recall that  $\mathcal{VL}$  denotes the set of virtual links. To complete the study of this section it remains to prove the following.

**Theorem 2.12** Let  $f^f : \mathcal{VL} \rightarrow R^f$  be the invariant defined from the Markov trace of Corollary 2.11. Then  $f^f$  coincides with the  $f$ -polynomial.

**Proof** Let  $\beta$  be a virtual braid on  $n$  strands and let  $\hat{\beta}$  be its closure. Observe that the relation  $\langle L \rangle = A \langle L_1 \rangle + A^{-1} \langle L_2 \rangle$  in the definition of the Kauffman bracket corresponds in terms of closed virtual braids to replacing each  $\sigma_i$  with  $A 1 + A^{-1} E_i$  and each  $\sigma_i^{-1}$  with  $A^{-1} 1 + A E_i$ . Once we have replaced each  $\sigma_i$  with  $A 1 + A^{-1} E_i$  and each  $\sigma_i^{-1}$  with  $A^{-1} 1 + A E_i$ , we get a linear combination  $\sum_{i=1}^\ell a_i E^{(i)}$ , where  $E^{(i)} \in \mathcal{E}_n$  and  $a_i \in R^f$ . For each  $i \in \{1, \dots, \ell\}$  we denote by  $m_i = t_n(E^{(i)})$  the number of closed curves in the closure of  $E^{(i)}$ . We see that

$$\langle \hat{\beta} \rangle = \sum_{i=1}^\ell a_i z^{m_i}.$$

Recall that  $w : \mathcal{VL} \rightarrow \mathbb{Z}$  denotes the writhe. Let  $\omega : \text{VB}_n \rightarrow \mathbb{Z}$  be the homomorphism which sends  $\sigma_i$  to 1 and  $\tau_i$  to 0 for all  $i \in \{1, \dots, n-1\}$ . Then  $w(\hat{\beta}) = \omega(\beta)$  and therefore  $f(\hat{\beta}) = (-A^3)^{-\omega(\beta)} \langle \hat{\beta} \rangle$ . So, in the above procedure, if we replace each  $\sigma_i$  with  $(-A^3)^{-1}(A 1 + A^{-1} E_i) = -A^{-2} 1 - A^{-4} E_i$  and each  $\sigma_i^{-1}$  with  $(-A^3)(A^{-1} 1 + A E_i) = -A^2 1 - A^4 E_i$ , then we get directly  $f(\hat{\beta})$ . It is clear that this procedure also leads to  $T_n^f(\beta)$ .  $\square$

### 3 Arrow Temperley–Lieb algebras and arrow polynomial

Throughout the section we consider the infinite families of variables  $\mathcal{Z} = \{z_k\}_{k=0}^\infty$  and  $\mathcal{Z}^* = \{z_k\}_{k=1}^\infty = \mathcal{Z} \setminus \{z_0\}$ , and we consider the algebra  $R_0^a = \mathbb{Z}[\mathcal{Z}]$  of polynomials in the variables in  $\mathcal{Z}$ , and the algebra  $R^a = \mathbb{Z}[A^{\pm 1}, \mathcal{Z}^*]$  of Laurent polynomials in the variable  $A$  and standard polynomials in the variables in  $\mathcal{Z}^*$ . We also assume that the algebra  $R_0^a$  is embedded into  $R^a$  via the identification  $z_0 = -A^2 - A^{-2}$ . Following the same strategy as in Section 2, we start by recalling the definition of the arrow polynomial, so that the reader will understand easier the constructions that will follow after.

Let  $S_1, \dots, S_\ell$  be a collection of  $\ell$  circles smoothly immersed in the plane and having only a finite number of double crossings. We assume that each circle  $S_i$  has an even number  $m_i$  of marked points

outside the crossings that we call *cusps*. We assume also that each segment between two successive cusps is oriented so that the orientations of the two segments adjacent to a given cusp are opposite. So, each cusp is either a *sink* or a *source*, according to the orientations of the segments adjacent to it (see Figure 3.1). If  $m_i = 0$ , then  $S_i$  is assumed to have a (unique) orientation. In addition, each cusp has a privileged side that we indicate with a small segment like in Figure 3.1. Finally, as for the virtual link diagrams, we assign a value “positive”, “negative”, or “virtual” to each crossing, that we indicate in its graphical representation as in Figure 3.2. Such a figure is called an *arrow virtual link diagram* with  $\ell$  components. Note that the virtual link diagrams are the arrow virtual link diagrams with no cusps.



Figure 3.1: Sink and source in an arrow virtual link diagram




Figure 3.2: Crossings in a virtual link diagram

**Example** Figure 3.3 shows an arrow virtual link diagram with two components. One component has two cusps and the other has no cusp.




Figure 3.3: Arrow virtual link diagram

Let  $L$  be an arrow virtual link diagram with only virtual crossings. Let  $S$  be a component of  $L$ . If  $S$  has two consecutive cusps  $p$  and  $q$  having the same privileged side, then we remove the two cusps and orient the new arc with the same orientation as that of the arc adjacent to  $p$  different from  $[p, q]$ . In the particular case where  $p$  and  $q$  are the only cusps of  $S$ , then we can choose any of the orientations of  $S$ . This operation is called a *reduction* of  $S$  and is illustrated in Figure 3.4. We apply such a reduction as many times as needed to get an *irreducible component*,  $S'$ . If  $2c$  is the number of cusp of  $S'$ , then  $c$  is called the *number of zigzags* of  $S$  and is denoted by  $\zeta(S) = c$ . If  $S_1, \dots, S_\ell$  are the components of  $L$ , then we set

$$\langle\langle L \rangle\rangle = \prod_{i=1}^{\ell} z_{\zeta(S_i)}.$$

This is a monomial of  $R_0^a$ .






Figure 3.4: Reduction

We define the *arrow Kauffman bracket*  $\langle\langle L \rangle\rangle \in R^a$  of any arrow virtual link diagram  $L$  as follows. If  $L$  has only virtual crossings, then  $\langle\langle L \rangle\rangle$  is the monomial  $\prod_{i=1}^{\ell} z_{\zeta(S_i)}$  defined above. Suppose that  $L$  has at least one non-virtual crossing at a point  $p$ . If the crossing is positive, then we set  $\langle\langle L \rangle\rangle = A\langle\langle L_1 \rangle\rangle + A^{-1}\langle\langle L_2 \rangle\rangle$ , and, if the crossing is negative, then we set  $\langle\langle L \rangle\rangle = A^{-1}\langle\langle L_1 \rangle\rangle + A\langle\langle L_2 \rangle\rangle$ , where  $L_1$  and  $L_2$  are identical to  $L$  except in a small neighborhood of  $p$  where there are as shown in Figure 3.5. As for the virtual link diagrams, the *writhe* of an arrow virtual link diagram  $L$ , denoted  $w(L)$ , is the number of positive crossings minus the number of negative crossings. Then the *arrow polynomial* of an arrow virtual link diagram  $L$  is defined by  $\vec{f}(L) = (-A^3)^{-w(L)}\langle\langle L \rangle\rangle$ .




Figure 3.5: Relation in the arrow Kauffman bracket

**Theorem 3.1** (Miyazawa [11], Dye–Kauffman [1]) (1) If two virtual link diagrams  $L$  and  $L'$  are equivalent, then  $\vec{f}(L) = \vec{f}(L')$ .

(2) If  $L$  is a diagram of a classical link, then  $\vec{f}(L) = f(L) \in R^f = \mathbb{Z}[A^{\pm 1}]$ .

**Remark** There is a notion of “equivalence” between arrow virtual link diagrams and Theorem 3.1 holds in this framework (see Miyazawa [11]), but the topic of the present paper are the virtual links, hence we state the theorem only for virtual link diagrams.

The *arrow polynomial* of a virtual link  $L$ , denoted  $\vec{f}(L)$ , is defined to be the arrow polynomial of any of its diagrams. This is a well-defined invariant thanks to Theorem 3.1.

Our aim now is to construct a Markov trace whose associated invariant is the arrow polynomial. We proceed with the same strategy as in Section 2 for the  $f$ -polynomial: we pass through a tower of algebras,  $\{\text{ATL}_n\}_{n=1}^{\infty}$ , that we will call arrow Temperley–Lieb algebras. We will also give a new proof/interpretation of Theorem 3.1 (2) in terms of Markov traces.

For the remainder of the section we need a more combinatorial definition of the multiplication in  $\text{VTL}_n$ . Recall that  $V_n = \{0, 1\} \times \{1, \dots, n\}$  is ordered by  $(0, 1) < (0, 2) < \dots < (0, n) < (1, n) < \dots < (1, 2) < (1, 1)$ . Let  $E, E' \in \mathcal{E}_n$ . An *arc* of length  $\ell$  in  $E \sqcup E'$  is a  $\ell$ -tuple  $\hat{\alpha} = (\alpha_1, \dots, \alpha_{\ell})$  in  $E \sqcup E'$ , where  $\alpha_i = \{(a_i, b_{i-1}), (c_i, b_i)\}$  with  $a_i, c_i \in \{0, 1\}$  and  $b_i \in \{1, \dots, n\}$ , satisfying the following properties.

- (a) If  $\alpha_i \in E$  and  $i < \ell$ , then  $c_i = 1$ ,  $\alpha_{i+1} \in E'$  and  $a_{i+1} = 0$ .
- (b) If  $\alpha_i \in E'$  and  $i < \ell$ , then  $c_i = 0$ ,  $\alpha_{i+1} \in E$  and  $a_{i+1} = 1$ .

- (c)  $a_0 = 0$  if  $\alpha_1 \in E$ ,  $a_0 = 1$  if  $\alpha_1 \in E'$ ,  $c_\ell = 0$  if  $\alpha_\ell \in E$ ,  $c_\ell = 1$  if  $\alpha_\ell \in E'$ , and  $(a_1, b_0) < (c_\ell, b_\ell)$ .

The *boundary* of  $\hat{\alpha}$  is  $\partial\hat{\alpha} = \{(a_1, b_0), (c_\ell, b_\ell)\}$ . There are  $n$  arcs in  $E \sqcup E'$  and their boundaries form a flat virtual  $n$ -tangle, denoted  $E * E'$ .

Let  $E, E' \in \mathcal{E}_n$ . A *cycle* of length  $2p \geq 2$  in  $E \sqcup E'$  is a  $2p$ -tuple  $\hat{\gamma} = (\gamma_1, \dots, \gamma_{2p})$  in  $E \sqcup E'$ , where  $\gamma_i = \{(a_i, b_{i-1}), (c_i, b_i)\}$  with  $a_i, c_i \in \{0, 1\}$  and  $b_i \in \{1, \dots, n\}$ , satisfying the following properties.

- (a)  $\gamma_i \in E$  and  $a_i = c_i = 1$ , if  $i$  is odd.  
(b)  $\gamma_i \in E'$  and  $a_i = c_i = 0$ , if  $i$  is even.  
(c)  $b_0 = b_{2p} < b_i$  for all  $i \in \{1, \dots, 2p - 1\}$ .

Let  $m$  be the number of cycles in  $E \sqcup E'$ . Then  $EE' = z^m(E * E')$ .

We can now define our algebra  $\text{ATL}_n$ . Let  $n \geq 1$ . An *arrow flat  $n$ -tangle* is a flat virtual  $n$ -tangle  $E$  endowed with a labeling  $f : E \rightarrow \mathbb{Z}$  such that, for  $\alpha = \{(a, b), (c, d)\} \in E$ ,  $f(\alpha)$  is odd if  $a = c$ , and  $f(\alpha)$  is even if  $a \neq c$ . Recall that  $\mathcal{Z} = \{z_k\}_{k=0}^\infty$  and  $R_0^a = \mathbb{Z}[\mathcal{Z}]$ . We denote by  $\mathcal{F}_n$  the set of arrow flat  $n$ -tangles and by  $\text{ATL}_n$  the free  $R_0^a$ -module freely generated by  $\mathcal{F}_n$ .

**Interpretation** Instead of labeling the arcs we could endow each arc with marked points (cusps) and each cusp with a privileged side that we indicate with a small segment, like for arrow virtual link diagrams, so that two consecutive cusps have different privileged sides. The number of cusps on an arc  $\alpha$  would be equal to  $|f(\alpha)|$ . Consider the order of  $V_n$  defined above. If we travel on the arc from its smallest extremity to its largest one, we set  $f(\alpha) > 0$  if the privileged side of the first encountered cusp is on the left hand side, and we set  $f(\alpha) < 0$  otherwise. An arrow flat tangle and its version with cusps are illustrated in Figure 3.6.




Figure 3.6: Arrow flat tangle

We now define the multiplication in  $\text{ATL}_n$ . Let  $F = (E, f)$  and  $F' = (E', f')$  be two arrow flat  $n$ -tangles. To simplify our notation we set  $f^*(\alpha) = f(\alpha)$  if  $\alpha \in E$  and  $f^*(\alpha) = f'(\alpha)$  if  $\alpha \in E'$ . The *parity* of an element  $\alpha = \{(a, b), (c, d)\} \in E \sqcup E'$  is  $\varpi(\alpha) = -1$  if  $a = c$ , and  $\varpi(\alpha) = 1$  if  $a \neq c$ . Let  $\hat{\alpha} = (\alpha_1, \dots, \alpha_\ell)$  be an arc of  $E \sqcup E'$ . Let  $i \in \{1, \dots, \ell\}$ . As in the above definition of arc, we set  $\alpha_i = \{(a_i, b_{i-1}), (c_i, b_i)\}$  for all  $i$ , where  $a_i, c_i \in \{0, 1\}$  and  $b_i \in \{1, \dots, n\}$ . We define the *cumulated parity* of  $\alpha_i$  relative to  $\hat{\alpha}$  by  $\varpi^c(\alpha_i) = \prod_{j=1}^{i-1} \varpi(\alpha_j)$  if  $(a_i, b_{i-1}) < (c_i, b_i)$ , and  $\varpi^c(\alpha_i) = \prod_{j=1}^i \varpi(\alpha_j)$  if  $(c_i, b_i) < (a_i, b_{i-1})$ . Then we set

$$g(\partial\hat{\alpha}) = g(\hat{\alpha}) = \sum_{i=1}^{\ell} \varpi^c(\alpha_i) f^*(\alpha_i).$$

At this stage we have an arrow flat  $n$ -tangle  $F * F' = (E * E', g)$ . Let  $\hat{\gamma} = (\gamma_1, \dots, \gamma_\ell)$  be a cycle of  $E \sqcup E'$ . Again, we write  $\gamma_i = \{(a_i, b_{i-1}), (c_i, b_i)\}$  for all  $i$ , where  $a_i, c_i \in \{0, 1\}$  and  $b_i \in \{1, \dots, n\}$ . As for an arc, we define the *cumulated parity* of  $\gamma_i$  relative to  $\hat{\gamma}$  by  $\varpi^c(\gamma_i) = \prod_{j=1}^{i-1} \varpi(\gamma_j)$  if  $(a_i, b_{i-1}) < (c_i, b_i)$ , and by  $\varpi^c(\gamma_i) = \prod_{j=1}^i \varpi(\gamma_j)$  if  $(c_i, b_i) < (a_i, b_{i-1})$ . We set

$$h(\hat{\gamma}) = \sum_{i=1}^{\ell} \varpi^c(\gamma_i) f^*(\gamma_i).$$

Observe that  $h(\hat{\gamma})$  is an even number. The *number of zigzags* of  $\hat{\gamma}$  is defined by  $\zeta(\hat{\gamma}) = \frac{|h(\hat{\gamma})|}{2}$ . Let  $\hat{\gamma}_1, \dots, \hat{\gamma}_m$  be the cycles of  $E \sqcup E'$ . Then the product of  $F$  and  $F'$  is

$$F F' = z_{\zeta(\hat{\gamma}_1)} \cdots z_{\zeta(\hat{\gamma}_m)}(F * F').$$

It is easily checked that  $\text{ATL}_n$  endowed with this multiplication is an (associative and unitary) algebra. We call it the  $n$ -th *arrow Temperley–Lieb algebra*.

**Example** On the left hand side of Figure 3.7 are illustrated two arrow flat tangles  $F$  and  $F'$ , and  $F * F'$  is illustrated on the right hand side. Here we have a unique cycle in  $E \sqcup E'$ ,  $\hat{\gamma}_1$ , and  $h(\hat{\gamma}_1) = 4$ , hence  $\zeta(\hat{\gamma}_1) = 2$  and  $F F' = z_2(F * F')$ .




Figure 3.7: Multiplication in  $\text{ATL}_n$

**Remark** Let  $n \geq 1$ . For  $F = (E, f) \in \mathcal{F}_n$  we define  $F^\sharp = (E^\sharp, f^\sharp) \in \mathcal{F}_{n+1}$  by setting  $E^\sharp = E \cup \{(0, n+1), (1, n+1)\}$ ,  $f^\sharp(\alpha) = f(\alpha)$  for all  $\alpha \in E$ , and  $f^\sharp(\{(0, n+1), (1, n+1)\}) = 0$ . Then the map  $\mathcal{F}_n \rightarrow \mathcal{F}_{n+1}$ ,  $F \mapsto F^\sharp$ , is an embedding which induces an injective homomorphism  $\text{ATL}_n \hookrightarrow \text{ATL}_{n+1}$ . So, we have a tower of algebras  $\{\text{ATL}_n\}_{n=1}^\infty$ .

**Proposition 3.2** *Let  $n \geq 2$ . Then  $\text{ATL}_n$  has a presentation with generators*

$$F_1, \dots, F_{n-1}, w_1, \dots, w_{n-1}, t_1, \dots, t_n, t_1^{-1}, \dots, t_n^{-1},$$

and relations

$$\begin{aligned}
t_i t_i^{-1} &= t_i^{-1} t_i = 1 \text{ for } 1 \leq i \leq n, & t_i t_j &= t_j t_i \text{ for } 1 \leq i < j \leq n, \\
w_i^2 &= 1 \text{ for } 1 \leq i \leq n-1, & w_i w_j &= w_j w_i \text{ for } |i-j| \geq 2, \\
w_i w_j w_i &= w_j w_i w_j \text{ for } |i-j| = 1, & w_i t_i &= t_{i+1} w_i \text{ for } 1 \leq i \leq n-1, \\
w_i t_{i+1} &= t_i w_i \text{ for } 1 \leq i \leq n-1, & w_i t_j &= t_j w_i \text{ for } j \neq i, i+1, \\
F_i t_i^m F_i &= z_{|m|} F_i \text{ for } 1 \leq i \leq n-1 \text{ and } m \in \mathbb{Z}, & F_i w_i &= F_i t_i = F_i t_{i+1}^{-1} \text{ for } 1 \leq i \leq n-1, \\
w_i F_i &= t_i^{-1} F_i = t_{i+1} F_i \text{ for } 1 \leq i \leq n-1, & F_i F_j &= F_j F_i \text{ for } |i-j| \geq 2, \\
F_i w_j &= w_j F_i \text{ for } |i-j| \geq 2, & F_i t_j &= t_j F_i \text{ for } j \neq i, i+1, \\
F_i w_j F_i &= F_i \text{ for } |i-j| = 1, & w_i w_j F_i &= F_j w_i w_j \text{ for } |i-j| = 1.
\end{aligned}$$

The generators  $F_i$ ,  $w_i$  and  $t_j$  are illustrated in Figure 3.8.




Figure 3.8: Generators of  $\text{ATL}_n$

The next six lemmas are preliminaries to the proof of Proposition 3.2. Let  $A_n$  be the  $R_0^a$ -algebra defined by a presentation with generators

$$X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}, u_1, \dots, u_n, u_1^{-1}, \dots, u_n^{-1},$$

and relations

$$\begin{aligned}
u_i u_i^{-1} &= u_i^{-1} u_i = 1 \text{ for } 1 \leq i \leq n, & u_i u_j &= u_j u_i \text{ for } 1 \leq i < j \leq n, \\
y_i^2 &= 1 \text{ for } 1 \leq i \leq n-1, & y_i y_j &= y_j y_i \text{ for } |i-j| \geq 2, \\
y_i y_j y_i &= y_j y_i y_j \text{ for } |i-j| = 1, & y_i u_i &= u_{i+1} y_i \text{ for } 1 \leq i \leq n-1, \\
y_i u_{i+1} &= u_i y_i \text{ for } 1 \leq i \leq n-1, & y_i u_j &= u_j y_i \text{ for } j \neq i, i+1, \\
X_i u_i^m X_i &= z_{|m|} X_i \text{ for } 1 \leq i \leq n-1 \text{ and } m \in \mathbb{Z}, & X_i y_i &= X_i u_i = X_i u_{i+1}^{-1} \text{ for } 1 \leq i \leq n-1, \\
y_i X_i &= u_i^{-1} X_i = u_{i+1} X_i \text{ for } 1 \leq i \leq n-1, & X_i X_j &= X_j X_i \text{ for } |i-j| \geq 2, \\
X_i y_j &= y_j X_i \text{ for } |i-j| \geq 2, & X_i u_j &= u_j X_i \text{ for } j \neq i, i+1, \\
X_i y_j X_i &= X_i \text{ for } |i-j| = 1, & y_i y_j X_i &= X_j y_i y_j \text{ for } |i-j| = 1.
\end{aligned}$$

It is easily checked using diagrammatic calculation that there is a homomorphism  $\varphi : A_n \rightarrow \text{ATL}_n$  which sends  $X_i$  to  $F_i$  for  $i \in \{1, \dots, n-1\}$ ,  $y_i$  to  $w_i$  for  $i \in \{1, \dots, n-1\}$ , and  $u_i^{\pm 1}$  to  $t_i^{\pm 1}$  for  $i \in \{1, \dots, n\}$ .

**Lemma 3.3** *The following equalities hold in  $A_n$ .*

$$\begin{aligned} X_i X_{i+1} y_i y_{i+1} &= X_i u_{i+2}^{-1} = u_{i+2}^{-1} X_i \text{ for } 1 \leq i \leq n-2, \\ y_{i+1} y_i X_{i+1} X_i &= u_{i+2} X_i = X_i u_{i+2} \text{ for } 1 \leq i \leq n-2, \\ X_i X_{i-1} y_i y_{i-1} &= X_i u_{i-1} = u_{i-1} X_i \text{ for } 2 \leq i \leq n-1, \\ y_{i-1} y_i X_{i-1} X_i &= u_{i-1}^{-1} X_i = X_i u_{i-1}^{-1} \text{ for } 2 \leq i \leq n-1. \end{aligned}$$

**Proof** We prove the first equality. The other three can be proved in the same way. Let  $i \in \{1, \dots, n-2\}$ . Then

$$\begin{aligned} X_i X_{i+1} y_i y_{i+1} &= X_i y_i y_{i+1} X_i = X_i u_{i+1}^{-1} y_{i+1} X_i = X_i y_{i+1} u_{i+2}^{-1} X_i = X_i y_{i+1} X_i u_{i+2}^{-1} = \\ &X_i u_{i+2}^{-1} = u_{i+2}^{-1} X_i. \end{aligned}$$

□

**Lemma 3.4** *Let  $i, j \in \{1, \dots, n-1\}$  such that  $|i-j| = 1$ . Then  $X_i X_j X_i = X_i$ .*

**Proof** We suppose that  $j = i+1$ . The case  $j = i-1$  can be proved in the same way.

$$\begin{aligned} X_i X_{i+1} X_i &= X_i X_{i+1} y_i y_{i+1} y_{i+1} y_i X_i = X_i u_{i+2}^{-1} y_{i+1} u_{i+1} X_i = X_i u_{i+2}^{-1} u_{i+2} y_{i+1} X_i = \\ &X_i y_{i+1} X_i = X_i. \end{aligned}$$

□

**Lemma 3.5** *Let  $i \in \{1, \dots, n-3\}$ . Then  $y_{i+1} y_{i+2} y_i y_{i+1} X_i X_{i+2} = X_i X_{i+2}$ .*

**Proof**

$$\begin{aligned} y_{i+1} y_{i+2} y_i y_{i+1} X_i X_{i+2} &= y_{i+1} y_{i+2} y_i y_{i+1} X_i u_{i+2} u_{i+2}^{-1} X_{i+2} = \\ y_{i+1} y_{i+2} y_i y_{i+1} y_i X_{i+1} X_i u_{i+2}^{-1} X_{i+2} &= y_{i+1} y_{i+2} X_{i+1} X_i u_{i+2}^{-1} X_{i+2} = \\ y_{i+1} y_{i+2} X_{i+1} u_{i+3} u_{i+3}^{-1} X_i u_{i+3} X_{i+2} &= y_{i+1} y_{i+2} y_{i+2} y_{i+1} X_{i+2} X_{i+1} X_i X_{i+2} = \\ X_{i+2} X_{i+1} X_{i+2} X_i &= X_{i+2} X_i = X_i X_{i+2}. \end{aligned}$$

□

Consider the action of the symmetric group  $\mathfrak{S}_n$  on  $\mathbb{Z}^n$  by permutations of the coordinates, and set  $G = \mathfrak{S}_n \ltimes \mathbb{Z}^n$ . Let  $\{e_1, \dots, e_n\}$  be the standard basis of  $\mathbb{Z}^n$  and let  $\{s_1, \dots, s_{n-1}\}$  be the standard set of generators of  $\mathfrak{S}_n$ . Recall that  $s_i$  is the transposition  $(i, i+1)$  for  $i \in \{1, \dots, n-1\}$ . We use multiplicative notation for the operation in  $\mathbb{Z}^n$  and we denote by  $1_{\mathbb{Z}^n}$  its neutral element. Let  $\mathcal{U}(A_n)$  be the group of units of  $A_n$ . We have a homomorphism  $\iota : G \rightarrow \mathcal{U}(A_n)$  which sends  $s_i$  to  $y_i$  for all  $i \in \{1, \dots, n-1\}$  and  $e_j$  to  $u_j$  for all  $j \in \{1, \dots, n\}$ .

Let  $n_0$  be the integer part of  $\frac{n}{2}$ . For  $p \in \{1, \dots, n_0\}$  we set  $B_p = X_1 X_3 \cdots X_{2p-1}$ , and for  $p = 0$  we set  $B_p = B_0 = 1$ . We denote by  $U_{1,p}$  the subset of  $G$  formed by the elements of the form  $g = wh$  where  $w \in \mathfrak{S}_n$  satisfies

$$\begin{aligned} w(1) < w(3) < \cdots < w(2p-1), \quad w(2i-1) < w(2i) \text{ for } 1 \leq i \leq p, \\ w(2p+1) < w(2p+2) < \cdots < w(n), \end{aligned}$$

and  $h = e_1^{\nu_1} e_2^{\nu_2} \cdots e_n^{\nu_n} \in \mathbb{Z}^n$  satisfies

$$\nu_i = 0 \text{ for } i \in \{2, 4, \dots, 2p, 2p+1, 2p+2, \dots, n\}.$$

On the other hand, we denote by  $U_{2,p}$  the subset of  $G$  formed by the elements of the form  $g = wh$  where  $w \in \mathfrak{S}_n$  satisfies

$$w(1) < w(3) < \cdots < w(2p-1), \quad w(2i-1) < w(2i) \text{ for } 1 \leq i \leq p,$$

and  $h = e_1^{\nu_1} e_2^{\nu_2} \cdots e_n^{\nu_n} \in \mathbb{Z}^n$  satisfies

$$\nu_i = 0 \text{ for } i \in \{2, 4, \dots, 2p\}.$$

Then we set

$$\mathcal{B}_p = \{\iota(g_1) B_p \iota(g_2^{-1}) \mid g_1 \in U_{1,p}, g_2 \in U_{2,p}\},$$

for  $0 \leq p \leq n_0$ , and

$$\mathcal{B} = \bigcup_{p=0}^{n_0} \mathcal{B}_p.$$

Let  $F = (E, f) \in \mathcal{F}_n$ . We can write  $E$  in the form  $E = \{\alpha_1, \dots, \alpha_p, \alpha'_1, \dots, \alpha'_p, \beta_1, \dots, \beta_q\}$ , where

- each  $\alpha_i$  is of the form  $\alpha_i = \{(0, a_i), (0, b_i)\}$ , with  $a_1 < a_2 < \cdots < a_p$ , and  $a_i < b_i$  for all  $i \in \{1, \dots, p\}$ ;
- each  $\alpha'_i$  is of the form  $\alpha'_i = \{(1, a'_i), (1, b'_i)\}$ , with  $a'_1 < a'_2 < \cdots < a'_p$ , and  $a'_i < b'_i$  for all  $i \in \{1, \dots, p\}$ ;
- each  $\beta_j$  is of the form  $\beta_j = \{(0, c_j), (1, d_j)\}$ , with  $c_1 < c_2 < \cdots < c_q$ , and  $2p + q = n$ .

We define

- $w_1 \in \mathfrak{S}_n$  by  $w_1(2i-1) = a_i$  and  $w_1(2i) = b_i$  for all  $i \in \{1, \dots, p\}$ , and  $w_1(2p+j) = c_j$  for all  $j \in \{1, \dots, q\}$ ;
- $w_2 \in \mathfrak{S}_n$  by  $w_2(2i-1) = a'_i$  and  $w_2(2i) = b'_i$  for all  $i \in \{1, \dots, p\}$ , and  $w_2(2p+j) = d_j$  for all  $j \in \{1, \dots, q\}$ ;
- $h_1 = e_1^{\nu_1} e_2^{\nu_2} \cdots e_n^{\nu_n} \in \mathbb{Z}^n$  by  $\nu_{1,2i-1} = \frac{f(\alpha_i)-1}{2}$  and  $\nu_{1,2i} = 0$  for  $i \in \{1, \dots, p\}$ , and  $\nu_{1,2p+j} = 0$  for  $j \in \{1, \dots, q\}$ ;
- $h_2 = e_1^{\nu_2} e_2^{\nu_2} \cdots e_n^{\nu_n} \in \mathbb{Z}^n$  by  $\nu_{2,2i-1} = \frac{f(\alpha'_i)-1}{2}$  and  $\nu_{2,2i} = 0$  for  $i \in \{1, \dots, p\}$ , and  $\nu_{2,2p+j} = -\frac{f(\beta_j)}{2}$  for  $j \in \{1, \dots, q\}$ ;
- $g_1 = w_1 h_1$  and  $g_2 = w_2 h_2$ .

Then  $g_1 \in U_{1,p}$ ,  $g_2 \in U_{2,p}$ , and  $F = \varphi(\iota(g_1) B_p \iota(g_2^{-1}))$ . Moreover, such an expression is unique, and  $\varphi(Y) \in \mathcal{F}_n$  for all  $Y \in \mathcal{B}$ . So,  $\varphi$  restricts to a bijection from  $\mathcal{B}$  to  $\mathcal{F}_n$ .

So, in order to prove Proposition 3.2, it suffices to show that  $\mathcal{B}$  spans  $A_n$  as a  $R_0^a$ -module. Let  $\mathcal{M}$  be the submonoid of  $A_n$  generated by  $X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}, u_1^{\pm 1}, \dots, u_n^{\pm 1}$ , that is, the set of finite products of elements in  $\{X_1, \dots, X_{n-1}, y_1, \dots, y_{n-1}, u_1^{\pm 1}, \dots, u_n^{\pm 1}\}$ . By definition  $\mathcal{M}$  spans  $A_n$  as a  $R_0^a$ -module, hence we only need to show that  $\mathcal{M}$  is contained in the  $R_0^a$ -submodule  $\text{Span}_{R_0^a}(\mathcal{B})$  of  $A_n$  spanned by  $\mathcal{B}$ .

**Lemma 3.6** Let  $g_1, g_2 \in G$  and  $p \in \{0, 1, \dots, n_0\}$ . Then  $\iota(g_1)B_p \iota(g_2^{-1}) \in \mathcal{B}$ .

**Proof** We write  $g_1 = w_1 h_1$  and  $g_2 = w_2 h_2$  with  $w_1, w_2 \in \mathfrak{S}_n$  and  $h_1, h_2 \in \mathbb{Z}^n$ . Let  $i \in \{1, \dots, p\}$  such that  $w_1(2i-1) = b_i > w_1(2i) = a_i$ . By using the relation  $y_i X_i = u_i^{-1} X_i$ , we can replace  $w_1$  with  $w_1 s_i$  and  $h_1$  with  $s_i(h_1 e_i) s_i \in \mathbb{Z}^n$ , and then  $w_1(2i-1) = a_i < w_1(2i) = b_i$ . So, we can assume that  $w_1(2i-1) < w_1(2i)$  for all  $i \in \{1, \dots, p\}$ . Let  $i \in \{1, \dots, p-1\}$  such that  $w_1(2i-1) = a_{i+1} > w_1(2i+1) = a_i$ . By Lemma 3.5 we can replace  $w_1$  with  $w_1 s_{2i} s_{2i-1} s_{2i+1} s_{2i}$  and  $h_1$  with  $(s_{2i} s_{2i+1} s_{2i-1} s_{2i}) h_1 (s_{2i} s_{2i-1} s_{2i+1} s_{2i}) \in \mathbb{Z}^n$ . Then we have  $w_1(2i-1) = a_i < w_1(2i+1) = a_{i+1}$  while keeping the inequalities  $w_1(2i-1) < w_1(2i)$  and  $w_1(2i+1) < w_1(2i+2)$ . Thus, we can also assume that  $w_1(1) < w_1(3) < \dots < w_1(2p-1)$ . Let  $j \in \{1, \dots, q-1\}$  such that  $w_1(2p+j) = c_{j+1} > w_1(2p+j+1) = c_j$ . By applying the relations  $y_{2p+j} X_{2i-1} = X_{2i-1} y_{2p+j}$  for  $i \in \{1, \dots, p\}$ , we can replace  $w_1$  with  $w_1 s_{2p+j}$ ,  $h_1$  with  $s_{2p+j} h_1 s_{2p+j} \in \mathbb{Z}^n$ , and  $g_2$  with  $g_2 s_{2p+j}$ , and then  $w_1(2p+j) = c_j < w_1(2p+j+1) = c_{j+1}$ . So, we can also assume that  $w_1(2p+1) < w_1(2p+2) < \dots < w_1(n)$ . We set  $h_1 = e_1^{\nu_{1,1}} \dots e_n^{\nu_{1,n}}$  and  $h_2 = e_1^{\nu_{2,1}} \dots e_n^{\nu_{2,n}}$ . Let  $i \in \{1, \dots, p\}$ . By applying the relation  $u_{2i-1}^{-1} X_{2i-1} = u_{2i} X_{2i-1}$ , we can replace  $\nu_{1,2i}$  with 0 and  $\nu_{1,2i-1}$  with  $\nu_{1,2i-1} - \nu_{1,2i}$ . So, we can also assume that  $\nu_{1,2i} = 0$  for all  $i \in \{1, \dots, p\}$ . Let  $j \in \{1, \dots, q\}$ . By applying the relations  $u_{2p+j} X_{2i-1} = X_{2i-1} u_{2p+j}$  for  $i \in \{1, \dots, p\}$ , we can replace  $\nu_{1,2p+j}$  with 0 and  $\nu_{2,2p+j}$  with  $\nu_{2,2p+j} - \nu_{1,2p+j}$ . Thus, we can also assume that  $\nu_{1,2p+j} = 0$  for all  $j \in \{1, \dots, q\}$ . In conclusion, we can assume that  $g_1 \in U_{1,p}$ .

We can use the same argument to show that  $g_2$  can be replaced with some  $g_2' \in U_{2,p}$ . So,  $\iota(g_1)B_p \iota(g_2^{-1}) \in \mathcal{B}$ .  $\square$

**Lemma 3.7** Let  $p \in \{0, 1, \dots, n_0\}$ ,  $a, b \in \{1, \dots, n-1\}$  and  $m \in \mathbb{Z}$ , such that  $a \leq b$ . Then  $X_a u_a^m y_{a+1} \dots y_b B_p \in \text{Span}_{\mathbb{R}_0}(\mathcal{B})$ .

**Proof** Suppose  $a \geq 2p+1$ . Then

$$\begin{aligned} X_a u_a^m y_{a+1} \dots y_b B_p &= X_a B_p u_a^m y_{a+1} \dots y_b = \\ &X_a (y_{a-1} y_a) (y_{a-2} y_{a-1}) \dots (y_{2p+1} y_{2p+2}) (y_{2p+2} y_{2p+1}) \dots (y_{a-1} y_{a-2}) (y_a y_{a-1}) B_p u_a^m y_{a+1} \dots y_b = \\ &(y_{a-1} y_a) (y_{a-2} y_{a-1}) \dots (y_{2p+1} y_{2p+2}) X_{2p+1} B_p (y_{2p+2} y_{2p+1}) \dots (y_{a-1} y_{a-2}) (y_a y_{a-1}) u_a^m y_{a+1} \dots y_b = \\ &(y_{a-1} y_a) (y_{a-2} y_{a-1}) \dots (y_{2p+1} y_{2p+2}) B_{p+1} (y_{2p+2} y_{2p+1}) \dots (y_{a-1} y_{a-2}) (y_a y_{a-1}) u_a^m y_{a+1} \dots y_b = \\ &\iota(g_1) B_{p+1} \iota(g_2^{-1}) \in \mathcal{B}, \end{aligned}$$

where

$$\begin{aligned} g_1 &= (s_{a-1} s_a) (s_{a-2} s_{a-1}) \dots (s_{2p+1} s_{2p+2}), \\ g_2 &= s_b \dots s_{a+1} e_a^{-m} (s_{a-1} s_a) (s_{a-2} s_{a-1}) \dots (s_{2p+1} s_{2p+2}). \end{aligned}$$

Suppose  $a \leq 2p$  and  $a$  is even. Let  $c$  such that  $a = 2c$ . Then

$$\begin{aligned} X_a u_a^m y_{a+1} \dots y_b B_p &= X_{2c} u_{2c+1}^{-m} y_{2c+1} \dots y_b X_{2c-1} X_1 \dots X_{2c-3} X_{2c+1} \dots X_{2p-1} = \\ &X_{2c} X_{2c-1} u_{2c+1}^{-m} y_{2c+1} \dots y_b X_1 \dots X_{2c-3} X_{2c+1} \dots X_{2p-1} = \\ &y_{2c-1} y_{2c} y_{2c} y_{2c-1} X_{2c} X_{2c-1} u_{2c+1}^{-m} y_{2c+1} \dots y_b X_1 \dots X_{2c-3} X_{2c+1} \dots X_{2p-1} = \\ &y_{2c-1} y_{2c} u_{2c+1} X_{2c-1} u_{2c+1}^{-m} y_{2c+1} \dots y_b X_1 \dots X_{2c-3} X_{2c+1} \dots X_{2p-1} = \\ &\iota(s_{2c-1} s_{2c} e_{2c+1}^{1-m} s_{2c+1} \dots s_b) B_p \in \mathcal{B}. \end{aligned}$$

Suppose  $a \leq 2p$ ,  $a$  is odd, and  $a = b$ . Let  $c$  such that  $a = 2c - 1$ . Then

$$\begin{aligned} X_a u_a^m y_{a+1} \cdots y_b B_p &= X_{2c-1} u_{2c-1}^m X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &= z_{|m|} X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = z_{|m|} B_p \in \text{Span}_{R_0^a}(\mathcal{B}). \end{aligned}$$

Suppose  $a \leq 2p$ ,  $a$  is odd, and  $a < b$ . Let  $c$  such that  $a = 2c - 1$ . Then

$$\begin{aligned} X_a u_a^m y_{a+1} \cdots y_b B_p &= X_{2c-1} u_{2c-1}^m y_{2c} y_{2c+1} \cdots y_b X_{2c-1} X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &= X_{2c-1} u_{2c}^{-m} y_{2c} X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &= X_{2c-1} y_{2c} u_{2c+1}^{-m} X_{2c-1} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &= X_{2c-1} y_{2c} X_{2c-1} u_{2c+1}^{-m} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \\ &= X_{2c-1} u_{2c+1}^{-m} y_{2c+1} \cdots y_b X_1 \cdots X_{2c-3} X_{2c+1} \cdots X_{2p-1} = \iota(e_{2c+1}^{-m} s_{2c+1} \cdots s_b) B_p \in \mathcal{B}. \end{aligned}$$

□

**Lemma 3.8** *Let  $g \in G$  and  $p \in \{0, 1, \dots, n_0\}$ . Then  $X_1 \iota(g) B_p \in \text{Span}_{R_0^a}(\mathcal{B})$ .*

**Proof** We write  $g$  in the form  $g = hw$  with  $h = e_1^{\nu_1} \cdots e_n^{\nu_n} \in \mathbb{Z}^n$  and  $w \in \mathfrak{S}_n$ . We have

$$X_1 \iota(g) B_p = X_1 u_1^{\nu_1} u_2^{\nu_2} \cdots u_n^{\nu_n} \iota(w) B_p = \iota(e_3^{\nu_3} \cdots e_n^{\nu_n}) X_1 \iota(e_1^{\nu_1 - \nu_2} w) B_p.$$

Thus, by Lemma 3.6, we can assume that  $h = e_1^m$  with  $m \in \mathbb{Z}$ . There exist  $w_1 \in \langle s_3, \dots, s_{n-1} \rangle$ ,  $a \in \{1, \dots, n-1\}$ , and  $b \in \{0, 1, \dots, n-1\}$  such that  $w = w_1 s_2 s_3 \cdots s_a s_1 s_2 \cdots s_b$ . We have

$$X_1 \iota(g) B_p = X_1 u_1^m \iota(w_1) y_2 \cdots y_a y_1 \cdots y_b B_p = \iota(w_1) X_1 u_1^m y_2 \cdots y_a y_1 \cdots y_b B_p.$$

Thus, by Lemma 3.6, we can assume that  $w = s_2 \cdots s_a s_1 \cdots s_b$ . Suppose  $a \leq b$ . Then

$$\begin{aligned} X_1 \iota(g) B_p &= X_1 u_1^m y_2 \cdots y_a y_1 \cdots y_b B_p = \\ &= X_1 u_1^m (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) y_a y_{a+1} \cdots y_b B_p = \\ &= (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) X_a u_a^m y_a y_{a+1} \cdots y_b B_p = \\ &= (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) X_a y_a u_{a+1}^m y_{a+1} \cdots y_b B_p = \\ &= (y_2 y_1) (y_3 y_2) \cdots (y_a y_{a-1}) X_a u_{a+1}^{m-1} y_{a+1} \cdots y_b B_p = \\ &= \iota((s_2 s_1) (s_3 s_2) \cdots (s_a s_{a-1})) X_a u_a^{1-m} y_{a+1} \cdots y_b B_p \in \text{Span}_{R_0^a}(\mathcal{B}). \end{aligned}$$

The last inclusion follows from Lemma 3.6 and Lemma 3.7. Suppose  $a > b$ . Then

$$\begin{aligned} X_1 \iota(g) B_p &= X_1 u_1^m y_2 \cdots y_a y_1 \cdots y_b B_p = \\ &= X_1 u_1^m (y_2 y_1) (y_3 y_2) \cdots (y_{b+1} y_b) y_{b+2} \cdots y_a B_p = \\ &= \iota((s_2 s_1) (s_3 s_2) \cdots (s_{b+1} s_b)) X_{b+1} u_{b+1}^m y_{b+2} \cdots y_a B_p \in \text{Span}_{R_0^a}(\mathcal{B}). \end{aligned}$$

Again, the last inclusion follows from Lemma 3.6 and Lemma 3.7. □

**Proof of Proposition 3.2** As pointed out before, it suffices to prove that  $\mathcal{M}$  is contained in  $\text{Span}_{R_0^a}(\mathcal{B})$ . Let  $Y \in \mathcal{M}$ . By using the relations  $y_i y_j X_i = X_j y_i y_j$  for  $|i - j| = 1$ , we see that  $Y$  can be written in the form  $Y = \iota(g_0) X_1 \iota(g_1) X_1 \cdots X_1 \iota(g_k)$ , where  $k \geq 0$  and  $g_0, g_1, \dots, g_k \in G$ . We argue by induction on  $k$ . The cases  $k = 0$  and  $k = 1$  follow directly from Lemma 3.6. So, we can assume that  $k \geq 2$  and that



the inductive hypothesis holds. By the inductive hypothesis,  $\iota(g_1)X_1 \cdots X_1 \iota(g_k) \in \text{Span}_{R_0^a}(\mathcal{B})$ . Thus, we just have to show that  $\iota(g_0)X_1 \iota(g'_1)B_p \iota(g'_2)^{-1} \in \text{Span}_{R_0^a}(\mathcal{B})$  for all  $p \in \{0, 1, \dots, n_0\}$  and all  $g'_1, g'_2 \in G$ . By Lemma 3.8 we have  $X_1 \iota(g'_1)B_p \in \text{Span}_{R_0^a}(\mathcal{B})$ , hence, by Lemma 3.6,  $\iota(g_0)X_1 \iota(g'_1)B_p \iota(g'_2)^{-1} \in \text{Span}_{R_0^a}(\mathcal{B})$ .  $\square$

Recall that the Temperley–Lieb algebra  $\text{TL}_n$  is the algebra over  $R_0^f = \mathbb{Z}[z]$  defined by the presentation with generators  $E_1, \dots, E_{n-1}$  and relations

$$E_i^2 = zE_i \text{ for } 1 \leq i \leq n-1, \quad E_i E_j = E_j E_i \text{ for } |i-j| \geq 2, \quad E_i E_j E_i = E_i \text{ for } |i-j| = 1.$$

We see in the presentation given in Proposition 3.2 that the relations  $F_i^2 = z_0 F_i$ , for  $1 \leq i \leq n-1$ , and  $F_i F_j = F_j F_i$ , for  $|i-j| \geq 2$ , hold in  $\text{ATL}_n$ . We also know that the relations  $F_i F_j F_i = F_i$ , for  $|i-j| = 1$ , hold (see Lemma 3.4). So, we have a ring homomorphism  $\iota : \text{TL}_n \rightarrow \text{ATL}_n$  which sends  $z$  to  $z_0$ , and  $E_i$  to  $F_i$  for all  $i \in \{1, \dots, n-1\}$ .

**Proposition 3.9** *Let  $n \geq 2$ . Then the above defined homomorphism  $\iota : \text{TL}_n \rightarrow \text{ATL}_n$  is injective.*

**Proof** We see from the presentations of  $\text{VTL}_n$  and  $\text{ATL}_n$  that there is a ring homomorphism  $\varphi : \text{ATL}_n \rightarrow \text{VTL}_n$  which sends  $z_m$  to  $z$  for all  $m \in \mathbb{N}$ ,  $F_i$  to  $E_i$  for all  $i \in \{1, \dots, n-1\}$ ,  $w_i$  to  $v_i$  for all  $i \in \{1, \dots, n-1\}$ , and  $t_j^{\pm 1}$  to 1 for all  $j \in \{1, \dots, n\}$ . By Proposition 2.8 the composition  $\varphi \circ \iota : \text{TL}_n \rightarrow \text{VTL}_n$  is injective, hence  $\iota : \text{TL}_n \rightarrow \text{ATL}_n$  is also injective.  $\square$

Recall that  $\mathcal{Z}^* = \{z_k\}_{k=1}^\infty$ ,  $R^a = \mathbb{Z}[A^{\pm 1}, \mathcal{Z}^*]$ , and that  $R_0^a$  is embedded into  $R^a$  via the identification  $z_0 = -A^2 - A^{-2}$ . For each  $n \geq 1$  we set  $\text{ATL}_n(R^a) = R^a \otimes \text{ATL}_n$ . This is a  $R^a$ -algebra and a free  $R^a$ -module freely generated by  $\mathcal{F}_n$ .

**Theorem 3.10** *Let  $n \geq 1$ . There exists a homomorphism  $\rho_n^a : R^a[\text{VB}_n] \rightarrow \text{ATL}_n(R^a)$  which sends  $\sigma_i$  to  $-A^{-2}1 - A^{-4}F_i$  and  $\tau_i$  to  $w_i$  for all  $i \in \{1, \dots, n-1\}$ .*

**Proof** The proof is almost identical to that of Theorem 2.9. We set  $S_i = -A^{-2}1 - A^{-4}F_i$ . It is easily checked as in the proof of Theorem 2.9 that  $(-A^{-2}1 - A^{-4}F_i)(-A^21 - A^4F_i) = 1$ , hence  $S_i$  is invertible and  $S_i^{-1} = -A^21 - A^4F_i$ . For  $i \in \{1, \dots, n-1\}$  we have  $w_i^2 = 1$ , hence  $w_i$  is also invertible. It remains to see that the following relations hold.

$$\begin{aligned} w_i^2 &= 1, \quad \text{for } 1 \leq i \leq n-1, \\ S_i S_j &= S_j S_i, \quad w_i w_j = w_j w_i, \quad w_i S_j = S_j w_i, \quad \text{for } |i-j| \geq 2, \\ S_i S_j S_i &= S_j S_i S_j, \quad w_i w_j w_i = w_j w_i w_j, \quad w_i w_j S_i = S_j w_i w_j, \quad \text{for } |i-j| = 1. \end{aligned}$$

The only relation which does not follow directly from the presentation of  $\text{ATL}_n(R^a)$  is  $S_i S_j S_i = S_j S_i S_j$ , for  $|i-j| = 1$ . But the latter can be proved in the same way as in the proof of Theorem 2.9.  $\square$

**Remark** (1) The sequence  $\{\rho_n^a : R^a[\text{VB}_n] \rightarrow \text{ATL}_n(R^a)\}_{n=1}^\infty$  is compatible with the tower of algebras  $\{\text{ATL}_n(R^a)\}_{n=1}^\infty$ .



- (1)  $T_n^{la}(xy) = T_n^{la}(yx)$  for all  $n \geq 1$  and all  $x, y \in \text{ATL}_n(\mathbb{R}^a)$ ;
- (2)  $T_n^{la}(x) = T_{n+1}^{la}(xS_n) = T_{n+1}^{la}(xS_n^{-1}) = T_{n+1}^{la}(xw_n)$  for all  $n \geq 1$  and all  $x \in \text{ATL}_n(\mathbb{R}^a)$ ;
- (3)  $T_n^{la}(x) = T_{n+1}^{la}(xS_n^{-1}w_{n-1}S_n)$  for all  $n \geq 2$  and all  $x \in \text{ATL}_n(\mathbb{R}^a)$ ;
- (4)  $T_n^{la}(x) = T_{n+1}^{la}(xw_nw_{n-1}S_{n-1}w_nS_{n-1}^{-1}w_{n-1}w_n)$  for all  $n \geq 2$  and all  $x \in \text{ATL}_n(\mathbb{R}^a)$ .

*Proof of (1).* We can assume that  $x = F = (E, f)$  and  $y = F' = (E', f')$  are arrow flat  $n$ -tangles. As in the definition of the multiplication in  $\text{ATL}_n$ , for  $\alpha \in E \sqcup E'$ , we set  $f^*(\alpha) = f(\alpha)$  if  $\alpha \in E$  and  $f^*(\alpha) = f'(\alpha)$  if  $\alpha \in E'$ . A long cycle of length  $2p$  in  $E \sqcup E'$  is a  $2p$ -tuple  $\hat{\gamma} = (\gamma_1, \dots, \gamma_{2p})$  in  $E \sqcup E'$ , where  $\gamma_i = \{(a_i, b_{i-1}), (c_i, b_i)\}$  with  $a_i, c_i \in \{0, 1\}$  and  $b_i \in \{1, \dots, n\}$ , satisfying the following properties.

- (a)  $\gamma_i \in E$  if  $i$  is odd,  $\gamma_i \in E'$  if  $i$  is even,  $a_{i+1} = 1$  if  $c_i = 0$ ,  $a_{i+1} = 0$  if  $c_i = 1$  (The indices are taken in  $\{1, \dots, 2p\}$  modulo  $2p$ . In particular,  $b_{2p} = b_0$ ).
- (b)  $(a_1, b_0) < (a_i, b_{i-1})$  for all  $i \in \{3, 5, \dots, 2p-1\}$  and  $(a_1, b_0) < (c_i, b_i)$  for all  $i \in \{1, 3, \dots, 2p-1\}$ .

Now, we define the number of zigzags for a long cycle exactly in the same way as we did for cycles. The *cumulated parity* of  $\gamma_i$  relative to  $\hat{\gamma}$  is  $\varpi^c(\gamma_i) = \prod_{j=1}^{i-1} \varpi(\gamma_j)$  if  $(a_i, b_{i-1}) < (c_i, b_i)$  and  $\varpi^c(\gamma_i) = \prod_{j=1}^i \varpi(\gamma_j)$  if  $(a_i, b_{i-1}) > (c_i, b_i)$ . We set

$$h(\hat{\gamma}) = \sum_{i=1}^{2p} \varpi^c(\gamma_i) f^*(\gamma_i).$$

We see that  $h(\hat{\gamma})$  is an even number. Then we define the *number of zigzags* of  $\hat{\gamma}$  by  $\zeta(\hat{\gamma}) = \frac{|h(\hat{\gamma})|}{2}$ . Let  $\hat{\gamma}_1, \dots, \hat{\gamma}_m$  be the long cycles of  $E \sqcup E'$ . Then

$$T_n^{la}(FF') = z_{\zeta(\hat{\gamma}_1)} z_{\zeta(\hat{\gamma}_2)} \cdots z_{\zeta(\hat{\gamma}_m)}.$$

Let  $\hat{\gamma} = (\gamma_1, \dots, \gamma_{2p})$  be a long cycle of  $E \sqcup E'$ . There exists a unique long cycle  $\hat{\gamma}'$  of  $E' \sqcup E$  of one the following forms

$$(\gamma_i, \gamma_{i+1}, \dots, \gamma_{2p}, \gamma_1, \gamma_2, \dots, \gamma_{i-1}) \text{ or } (\gamma_i, \gamma_{i-1}, \dots, \gamma_1, \gamma_{2p}, \gamma_{2p-1}, \dots, \gamma_{i+1}),$$


with  $i \in \{2, 4, \dots, 2p\}$ . In addition, each long cycle of  $E' \sqcup E$  is of this form, and  $h(\hat{\gamma}') = \pm h(\hat{\gamma})$ , hence  $\zeta(\hat{\gamma}') = \zeta(\hat{\gamma})$ . We conclude that  $T_n^{la}(FF') = T_n^{la}(F'F)$ .

*Proof of (2).* From now on the proof of Theorem 3.11 is almost identical to that of Theorem 2.10. We can assume that  $x = F = (E, f)$  is an arrow flat  $n$ -tangle. We see in Figure 3.10 that the following equalities hold.

$$T_{n+1}^{la}(F) = z_0 T_n^{la}(F), \quad T_{n+1}^{la}(FF_n) = T_n^{la}(F), \quad T_{n+1}^{la}(Fw_n) = T_n^{la}(F).$$

Recall that  $S_n^{-1} = -A^2 1 - A^4 F_n$  (see the proof of Theorem 3.10). It follows that

$$\begin{aligned} T_{n+1}^{la}(FS_n) &= -A^{-2} T_{n+1}^{la}(F) - A^{-4} T_{n+1}^{la}(FF_n) = (-A^{-2} z_0 - A^{-4}) T_n^{la}(F) = T_n^{la}(F), \\ T_{n+1}^{la}(FS_n^{-1}) &= -A^2 T_{n+1}^{la}(F) - A^4 T_{n+1}^{la}(FF_n) = (-A^2 z_0 - A^4) T_n^{la}(F) = T_n^{la}(F). \end{aligned}$$

Figure 3.10:  $n + 1$ -closure of  $F$ ,  $FF_n$ , and  $Fw_n$ 

*Proof of (3).* We can again assume that  $x = F = (E, f)$  is a flat  $n$ -tangle. We have

$$\begin{aligned} FS_n^{-1}w_{n-1}S_n &= F(-A^2 1 - A^4 F_n)w_{n-1}(-A^{-2} 1 - A^{-4} F_n) = \\ &Fw_{n-1} + A^2 FF_n w_{n-1} + A^{-2} Fw_{n-1} F_n + FF_n w_{n-1} F_n = \\ &Fw_{n-1} + A^2 FF_n w_{n-1} + A^{-2} Fw_{n-1} F_n + FF_n. \end{aligned}$$

By the above, it follows that

$$\begin{aligned} T_{n+1}^{ja}(FS_n^{-1}w_{n-1}S_n) &= \\ T_{n+1}^{ja}(Fw_{n-1}) + A^{-2}T_{n+1}^{ja}(Fw_{n-1}F_n) + A^2T_{n+1}^{ja}(FF_n w_{n-1}) + T_{n+1}^{ja}(FF_n) &= \\ z_0 T_n^{ja}(Fw_{n-1}) + A^{-2}T_n^{ja}(Fw_{n-1}) + A^2T_{n+1}^{ja}(w_{n-1}FF_n) + T_n^{ja}(F) &= \\ (-A^2 - A^{-2})T_n^{ja}(Fw_{n-1}) + A^{-2}T_n^{ja}(Fw_{n-1}) + A^2T_n^{ja}(w_{n-1}F) + T_n^{ja}(F) &= \\ (-A^2 - A^{-2})T_n^{ja}(Fw_{n-1}) + A^{-2}T_n^{ja}(Fw_{n-1}) + A^2T_n^{ja}(Fw_{n-1}) + T_n^{ja}(F) &= T_n^{ja}(F). \end{aligned}$$

*Proof of (4).* We can again assume that  $x = F = (E, f)$  is an arrow flat  $n$ -tangle. Then

$$\begin{aligned} Fw_n w_{n-1} S_{n-1} w_n S_{n-1}^{-1} w_{n-1} w_n &= \\ Fw_n w_{n-1} (-A^{-2} 1 - A^{-4} F_{n-1}) w_n (-A^2 1 - A^4 F_{n-1}) w_{n-1} w_n &= \\ (Fw_n w_{n-1} w_n w_{n-1} w_n) + A^2 (Fw_n w_{n-1} w_n F_{n-1} w_{n-1} w_n) + \\ A^{-2} (Fw_n w_{n-1} F_{n-1} w_n w_{n-1} w_n) + (Fw_n w_{n-1} F_{n-1} w_n F_{n-1} w_{n-1} w_n) &= \\ (Fw_{n-1} w_n w_{n-1} w_{n-1} w_n) + A^2 (Fw_n F_n w_{n-1} w_n w_{n-1} w_n) + \\ A^{-2} (Fw_n w_{n-1} w_n w_{n-1} F_n w_n) + (Fw_n w_{n-1} F_{n-1} w_{n-1} w_n) &= \\ (Fw_{n-1}) + A^2 (Fw_n F_n w_{n-1} w_{n-1} w_n w_{n-1}) + \\ A^{-2} (Fw_{n-1} w_n w_{n-1} w_{n-1} F_n w_n) + (Fw_n t_{n-1}^{-1} F_{n-1} t_{n-1} w_n) &= \\ (Fw_{n-1}) + A^2 (Fw_n F_n w_n w_{n-1}) + A^{-2} (Fw_{n-1} w_n F_n w_n) + (Ft_{n-1}^{-1} w_n F_{n-1} w_n t_{n-1}) &= \\ (Fw_{n-1}) + A^2 (Ft_n^{-1} F_n t_n w_{n-1}) + A^{-2} (Fw_{n-1} t_n^{-1} F_n t_n) + (Ft_{n-1}^{-1} w_{n-1} F_n w_{n-1} t_{n-1}). \end{aligned}$$

By the above, it follows that

$$\begin{aligned}
& T_{n+1}'^a(Fw_n w_{n-1} S_{n-1} w_n S_{n-1}^{-1} w_{n-1} w_n) = \\
& T_{n+1}'^a(Fw_{n-1}) + A^2 T_{n+1}'^a(Ft_n^{-1} F_n t_n w_{n-1}) + \\
& A^{-2} T_{n+1}'^a(Fw_{n-1} t_n^{-1} F_n t_n) + T_{n+1}'^a(Ft_{n-1}^{-1} w_{n-1} F_n w_{n-1} t_{n-1}) = \\
& z_0 T_n'^a(Fw_{n-1}) + A^2 T_{n+1}'^a(t_n w_{n-1} F t_n^{-1} F_n) + \\
& A^{-2} T_{n+1}'^a(t_n F w_{n-1} t_n^{-1} F_n) + T_{n+1}'^a(w_{n-1} t_{n-1} F t_{n-1}^{-1} w_{n-1} F_n) = \\
& (-A^2 - A^{-2}) T_n'^a(Fw_{n-1}) + A^2 T_n'^a(t_n w_{n-1} F t_n^{-1}) + \\
& A^{-2} T_n'^a(t_n F w_{n-1} t_n^{-1}) + T_n'^a(w_{n-1} t_{n-1} F t_{n-1}^{-1} w_{n-1}) = \\
& (-A^2 - A^{-2}) T_n'^a(Fw_{n-1}) + A^2 T_n'^a(F t_n^{-1} t_n w_{n-1}) + \\
& A^{-2} T_n'^a(F w_{n-1} t_n^{-1} t_n) + T_n'^a(F t_{n-1}^{-1} w_{n-1} w_{n-1} t_{n-1}) = \\
& (-A^2 - A^{-2}) T_n'^a(Fw_{n-1}) + A^2 T_n'^a(Fw_{n-1}) + A^{-2} T_n'^a(Fw_{n-1}) + T_n'^a(F) = T_n'^a(F). \quad \square
\end{aligned}$$

**Corollary 3.12** For each  $n \geq 1$  we set  $T_n^a = T_n'^a \circ \rho_n^a : R^a[\text{VB}_n] \rightarrow R^a$ . Then the sequence  $\{T_n^a : R^a[\text{VB}_n] \rightarrow R^a\}_{n=1}^\infty$  is a Markov trace.

Recall that  $R^f = \mathbb{Z}[A^{\pm 1}]$  is a subring of  $R^a = \mathbb{Z}[A^{\pm 1}, \mathcal{Z}^*]$  and that, for  $n \geq 1$ , we have an embedding  $\iota_n : \text{TL}_n(R^f) \rightarrow \text{ATL}_n(R^a)$  which sends  $E_i$  to  $F_i$  for all  $i \in \{1, \dots, n-1\}$  (see Proposition 3.9). The next proposition is a version of Theorem 3.1 (2) in terms of Temperley–Lieb algebras.

**Proposition 3.13** Let  $n \geq 1$ . Then  $T_n'^a(\iota_n(x)) = T_n'^f(x) \in R^f$  for all  $x \in \text{TL}_n(R^f)$ .

We denote by  $\mathcal{E}_n^\nu$  the set of flat virtual  $n$ -tangles  $E \in \mathcal{E}_n$  such that, for each  $\alpha = \{(a, b), (c, d)\} \in E$ ,  $d-b$  is odd if  $a = c$ , and  $d-b$  is even if  $a \neq c$ . We denote by  $\text{VTL}_n^\nu$  the  $R_0^f$ -submodule of  $\text{VTL}_n$  spanned by  $\mathcal{E}_n^\nu$ . For each  $E \in \mathcal{E}_n^\nu$  we define  $\iota_n^\nu(E) = (E, f) \in \mathcal{F}_n$  as follows. Let  $\alpha = \{(a, b), (c, d)\} \in E$ . If  $a = c$ , we set  $f(\alpha) = |d-b|$ , and if  $a = 0$  and  $c = 1$ , we set  $f(\alpha) = d-b$ . We define  $\iota_n^\nu : \text{VTL}_n^\nu \rightarrow \text{ATL}_n$  by identifying  $z$  with  $z_0$  and extending linearly the map  $\iota_n^\nu : \mathcal{E}_n^\nu \rightarrow \mathcal{F}_n$ . The key point in the proof of Proposition 3.13 is the following.

**Lemma 3.14** Let  $n \geq 1$ .

- (1)  $\text{VTL}_n^\nu$  is a subalgebra of  $\text{VTL}_n$  and  $\iota_n^\nu : \text{VTL}_n^\nu \rightarrow \text{ATL}_n$  is a ring homomorphism.
- (2) Let  $\text{VTL}_n^\nu(R^f) = R^f \otimes \text{VTL}_n^\nu$ , and let  $\iota_n^\nu : \text{VTL}_n^\nu(R^f) \rightarrow \text{ATL}_n(R^a)$  be the homomorphism induced by  $\iota_n^\nu : \text{VTL}_n^\nu \rightarrow \text{ATL}_n$ . Then  $T_n'^a(\iota_n^\nu(x)) = T_n'^f(x) \in R^f$  for all  $x \in \text{VTL}_n^\nu(R^f)$ .

**Proof** Let  $E, E' \in \mathcal{E}_n^\nu$ . Set  $\iota_n^\nu(E) = (E, f)$ ,  $\iota_n^\nu(E') = (E', f')$ , and  $\iota_n^\nu(E) * \iota_n^\nu(E') = (E * E', g)$ . We first show that  $E * E' \in \mathcal{E}_n^\nu$  and that  $(E * E', g) = \iota_n^\nu(E * E')$ . Let  $\hat{\alpha}$  be an arc in  $E \sqcup E'$  of length  $\ell$ , and let  $\partial \hat{\alpha} = \{x', y'\}$ , with  $x' = (a, b) < y' = (c, d)$ .

Suppose  $a = c = 0$  and  $\ell = 1$ . Then  $\partial \hat{\alpha} = \alpha_1 = \{(0, b), (0, d)\}$ ,  $d-b$  is odd, and  $g(\partial \hat{\alpha}) = d-b = |d-b|$ .

Suppose  $a = c = 0$  and  $\ell > 1$ . Then  $\ell$  is odd, say  $\ell = 2p+1$ . There exists a sequence  $a_0, a_1, \dots, a_{2p+1}$  in  $\{1, \dots, n\}$  such that  $\hat{\alpha}$  is of the form  $\hat{\alpha} = (\alpha_0, \beta_1, \gamma_2, \dots, \gamma_{2p-2}, \beta_{2p-1}, \alpha_{2p})$ ,

where  $\alpha_0 = \{(0, a_0), (1, a_1)\} \in E$ ,  $\beta_i = \{(0, a_i), (0, a_{i+1})\} \in E'$  for all  $i \in \{1, 3, \dots, 2p-1\}$ ,  $\gamma_i = \{(1, a_i), (1, a_{i+1})\} \in E$  for all  $i \in \{2, 4, \dots, 2p-2\}$ , and  $\alpha_{2p} = \{(1, a_{2p}), (0, a_{2p+1})\} \in E$ . We also have  $x' = (0, a_0)$ ,  $y' = (0, a_{2p+1})$ , and  $a_0 = b < a_{2p+1} = d$ . The numbers  $a_1 - a_0$  and  $a_{2p+1} - a_{2p}$  are even, and the number  $a_{i+1} - a_i$  is odd for every  $i \in \{1, \dots, 2p-1\}$ , hence  $a_{2p+1} - a_0 = d - b$  is odd. We have  $\varpi^c(\alpha_0) = 1$ , hence  $\varpi^c(\alpha_0)f^*(\alpha_0) = a_1 - a_0$ . Let  $i \in \{1, 3, \dots, 2p-1\}$ . We have  $\varpi^c(\beta_i) = 1$  if  $a_i < a_{i+1}$  and  $\varpi^c(\beta_i) = -1$  if  $a_i > a_{i+1}$ . In both cases we have  $\varpi^c(\beta_i)f^*(\beta_i) = a_{i+1} - a_i$ . Let  $i \in \{2, 4, \dots, 2p-2\}$ . We have  $\varpi^c(\gamma_i) = 1$  if  $a_i < a_{i+1}$  and  $\varpi^c(\gamma_i) = -1$  if  $a_i > a_{i+1}$ . In both cases we have  $\varpi^c(\gamma_i)f^*(\gamma_i) = a_{i+1} - a_i$ . We have  $\varpi^c(\alpha_{2p}) = -1$ , hence  $\varpi^c(\alpha_{2p})f^*(\alpha_{2p}) = a_{2p+1} - a_{2p}$ . So,  $g(\hat{\alpha}) = g(\partial\hat{\alpha}) = a_{2p+1} - a_0 = d - b = |d - b|$ .

Suppose  $a = c = 1$  and  $\ell = 1$ . Then  $\partial\hat{\alpha} = \alpha_1 = \{(1, b), (1, d)\}$ ,  $b - d$  is odd, and  $g(\partial\hat{\alpha}) = b - d = |b - d|$ . Suppose  $a = c = 1$  and  $\ell > 1$ . Then we show in the same way as for the case  $a = c = 0$  and  $\ell > 1$  that  $b - d$  is odd and  $g(\hat{\alpha}) = g(\partial\hat{\alpha}) = b - d = |b - d|$ .

Suppose  $a = 0$  and  $c = 1$ . Then  $\ell \geq 2$  and  $\ell$  is even, say  $\ell = 2p + 2$ . There exists a sequence  $a_0, a_1, \dots, a_{2p+2}$  in  $\{1, \dots, n\}$  such that  $\hat{\alpha}$  is of the form  $\hat{\alpha} = (\alpha_0, \beta_1, \gamma_2, \dots, \beta_{2p-1}, \gamma_{2p}, \delta_{2p+1})$ , where  $\alpha_0 = \{(0, a_0), (1, a_1)\} \in E$ ,  $\beta_i = \{(0, a_i), (0, a_{i+1})\} \in E'$  for all  $i \in \{1, 3, \dots, 2p-1\}$ ,  $\gamma_i = \{(1, a_i), (1, a_{i+1})\} \in E$  for all  $i \in \{2, 4, \dots, 2p\}$ , and  $\delta_{2p+1} = \{(0, a_{2p+1}), (1, a_{2p+2})\} \in E'$ . We also have  $x' = (0, b) = (0, a_0)$  and  $y' = (1, d) = (1, a_{2p+2})$ . The numbers  $a_1 - a_0$  and  $a_{2p+2} - a_{2p+1}$  are even and  $a_{i+1} - a_i$  is odd for each  $i \in \{1, \dots, 2p\}$ , hence  $d - b = a_{2p+2} - a_0$  is even. We have  $\varpi^c(\alpha_0) = 1$ , hence  $\varpi^c(\alpha_0)f^*(\alpha_0) = a_1 - a_0$ . Let  $i \in \{1, 3, \dots, 2p-1\}$ . We have  $\varpi^c(\beta_i) = 1$  if  $a_i < a_{i+1}$  and  $\varpi^c(\beta_i) = -1$  if  $a_i > a_{i+1}$ . In both cases we have  $\varpi^c(\beta_i)f^*(\beta_i) = a_{i+1} - a_i$ . Let  $i \in \{2, 4, \dots, 2p\}$ . We have  $\varpi^c(\gamma_i) = 1$  if  $a_i < a_{i+1}$  and  $\varpi^c(\gamma_i) = -1$  if  $a_i > a_{i+1}$ . In both cases we have  $\varpi^c(\gamma_i)f^*(\gamma_i) = a_{i+1} - a_i$ . We have  $\varpi^c(\delta_{2p+1}) = 1$ , hence  $\varpi^c(\delta_{2p+1})f^*(\delta_{2p+1}) = a_{2p+2} - a_{2p+1}$ . So,  $g(\hat{\alpha}) = g(\partial\hat{\alpha}) = a_{2p+2} - a_0 = d - b$ .

The above shows that  $E * E' \in \mathcal{E}_n^\nu$  and  $\iota_n^\nu(E * E') = \iota_n^\nu(E) * \iota_n^\nu(E')$ .

Let  $\hat{\gamma}$  be a cycle in  $E \sqcup E'$  of length  $\ell$ . Then  $\ell \geq 2$  and  $\ell$  is even, say  $\ell = 2p$ . There exists a sequence  $a_0, a_1, \dots, a_{2p}$  in  $\{1, \dots, n\}$  such that  $\hat{\gamma}$  is of the form  $\hat{\gamma} = (\alpha_1, \beta_2, \dots, \alpha_{2p-1}, \beta_{2p})$ , where  $\alpha_i = \{(1, a_{i-1}), (1, a_i)\} \in E$  for all  $i \in \{1, 3, \dots, 2p-1\}$  and  $\beta_i = \{(0, a_{i-1}), (0, a_i)\} \in E'$  for all  $i \in \{2, 4, \dots, 2p\}$ . We also have  $a_{2p} = a_0$ . Let  $i \in \{1, 3, \dots, 2p-1\}$ . We have  $\varpi^c(\alpha_i) = 1$  if  $a_{i-1} > a_i$  and  $\varpi^c(\alpha_i) = -1$  if  $a_{i-1} < a_i$ . In both cases we have  $\varpi^c(\alpha_i)f^*(\alpha_i) = a_{i-1} - a_i$ . Let  $i \in \{2, 4, \dots, 2p\}$ . We have  $\varpi^c(\beta_i) = 1$  if  $a_{i-1} > a_i$  and  $\varpi^c(\beta_i) = -1$  if  $a_{i-1} < a_i$ . In both cases we have  $\varpi^c(\beta_i)f^*(\beta_i) = a_{i-1} - a_i$ . Thus,  $h(\hat{\gamma}) = a_0 - a_{2p} = 0$ , hence  $\zeta(\hat{\gamma}) = 0$ . So, if  $m$  is the number of cycles in  $E \sqcup E'$ , then  $\iota_n^\nu(E) \iota_n^\nu(E') = z_0^m (\iota_n^\nu(E) * \iota_n^\nu(E'))$ , hence  $\iota_n^\nu(E) \iota_n^\nu(E') = \iota_n^\nu(EE')$ . This concludes the proof of the first part of the lemma.

Let  $E \in \mathcal{E}_n^\nu$  and let  $(E, f) = \iota_n^\nu(E)$ . In order to prove the second part of the lemma, it suffices to show that, if  $\hat{\gamma}$  is a cycle of  $\hat{E}$ , then  $\zeta(\hat{\gamma}) = 0$ , that is,  $h(\hat{\gamma}) = 0$ . Let  $\hat{\gamma}$  be a cycle of  $\hat{E}$ . Then  $\hat{\gamma}$  is of the form

$$\hat{\gamma} = (\alpha_{1,1}, \dots, \alpha_{1,p_1}, \beta_1, \gamma_{1,1}, \dots, \gamma_{1,q_1}, \delta_1, \dots, \alpha_{r,1}, \dots, \alpha_{r,p_r}, \beta_r, \gamma_{r,1}, \dots, \gamma_{r,q_r}, \\ \delta_r, \alpha_{r+1,1}, \dots, \alpha_{r+1,p_{r+1}}),$$

where

$$\alpha_{i,j} = \{(0, a_{i,j-1}), (1, a_{i,j})\}, \beta_i = \{(0, b_{i,0}), (0, b_{i,1})\}, \gamma_{i,j} = \{(1, c_{i,j-1}), (0, c_{i,j})\}, \\ \delta_i = \{(1, d_{i,0}), (1, d_{i,1})\}, a_{i,p_i} = b_{i,0}, b_{i,1} = c_{i,0}, c_{i,q_i} = d_{i,0}, d_{i,1} = a_{i+1,0}, a_{r+1,p_{r+1}} = a_{1,0}.$$

We have  $\varpi^c(\alpha_{i,j}) = 1$ , hence  $\varpi^c(\alpha_{i,j})f(\alpha_{i,j}) = a_{i,j} - a_{i,j-1}$ . We have  $\varpi^c(\beta_i) = -1$  if  $b_{i,0} > b_{i,1}$  and  $\varpi^c(\beta_i) = 1$  if  $b_{i,0} < b_{i,1}$ . In both cases we have  $\varpi^c(\beta_i)f(\beta_i) = b_{i,1} - b_{i,0}$ . We have  $\varpi^c(\gamma_{i,j}) = -1$ , hence  $\varpi^c(\gamma_{i,j})f(\gamma_{i,j}) = c_{i,j} - c_{i,j-1}$ . We have  $\varpi^c(\delta_i) = -1$  if  $d_{i,0} > d_{i,1}$  and  $\varpi^c(\delta_i) = 1$  if  $d_{i,0} < d_{i,1}$ . In both cases we have  $\varpi^c(\delta_i)f(\delta_i) = d_{i,1} - d_{i,0}$ . Combining these equalities we get  $h(\hat{\gamma}) = a_{r+1,p_{r+1}} - a_{1,0} = 0$ , hence  $\zeta(\hat{\gamma}) = 0$ .  $\square$

**Proof of Proposition 3.13** We have  $E_i \in \text{VTL}_n^\nu(R^f)$  for all  $i \in \{1, \dots, n-1\}$ ,  $\text{TL}_n(R^f)$  is generated by  $E_1, \dots, E_{n-1}$ , and  $\text{VTL}_n^\nu(R^f)$  is a subalgebra of  $\text{VTL}_n(R^f)$ , hence  $\text{TL}_n(R^f) \subset \text{VTL}_n^\nu(R^f)$ . Moreover, since  $\iota_n(E_i) = \iota_n^\nu(E_i)$  for all  $i \in \{1, \dots, n-1\}$ , we have  $\iota_n(x) = \iota_n^\nu(x)$  for all  $x \in \text{TL}_n(R^f)$ . We conclude from Lemma 3.14 that  $T_n^a(\iota_n(x)) = T_n^f(x) \in R^f$  for all  $x \in \text{TL}_n(R^f)$ .  $\square$

Now, it remains to prove the following.

**Theorem 3.15** *Let  $I^a : \mathcal{VL} \rightarrow R^a$  be the invariant defined from the Markov trace of Corollary 3.12. Then  $I^a$  coincides with the arrow polynomial.*

**Proof** The proof is similar to that of Theorem 2.12. Let  $\beta$  be a virtual braid on  $n$  strands and let  $\hat{\beta}$  be its closure. It is easily seen that the relations  $\langle\langle L \rangle\rangle = A\langle\langle L_1 \rangle\rangle + A^{-1}\langle\langle L_2 \rangle\rangle$  and  $\langle\langle L \rangle\rangle = A^{-1}\langle\langle L_1 \rangle\rangle + A\langle\langle L_2 \rangle\rangle$  in the definition of the arrow Kauffman bracket corresponds in terms of closed virtual braids to replacing each  $\sigma_i$  with  $A1 + A^{-1}F_i$  and each  $\sigma_i^{-1}$  with  $A^{-1}1 + AF_i$ . Once we have replaced each  $\sigma_i$  with  $A1 + A^{-1}F_i$  and each  $\sigma_i^{-1}$  with  $A^{-1}1 + AF_i$ , we get a linear combination  $\sum_{i=1}^{\ell} a_i F^{(i)}$ , where  $F^{(i)} = (E^{(i)}, f^{(i)}) \in \mathcal{F}_n$  and  $a_i \in R^a$  for all  $i$ . For  $i \in \{1, \dots, \ell\}$  let  $\hat{\gamma}_{i,1}, \dots, \hat{\gamma}_{i,m_i}$  be the cycles of  $\hat{E}^{(i)}$ . Then

$$\langle\langle \hat{\beta} \rangle\rangle = \sum_{i=1}^{\ell} a_i z_{\zeta(\hat{\gamma}_{i,1})} z_{\zeta(\hat{\gamma}_{i,2})} \cdots z_{\zeta(\hat{\gamma}_{i,m_i})}.$$

Recall that  $w : \mathcal{VL} \rightarrow \mathbb{Z}$  denotes the writhe. Let  $\omega : \text{VB}_n \rightarrow \mathbb{Z}$  be the homomorphism which sends  $\sigma_i$  to 1 and  $\tau_i$  to 0 for all  $i \in \{1, \dots, n-1\}$ . Then  $w(\hat{\beta}) = \omega(\beta)$ , and therefore  $\vec{f}(\hat{\beta}) = (-A^3)^{-\omega(\beta)} \langle\langle \hat{\beta} \rangle\rangle$ . So, in the above procedure, if we replace each  $\sigma_i$  with  $(-A^3)^{-1}(A1 + A^{-1}F_i) = -A^{-2}1 - A^{-4}F_i$  and each  $\sigma_i^{-1}$  with  $(-A^3)(A^{-1}1 + AF_i) = -A^21 - A^4F_i$ , then we get directly  $\vec{f}(\hat{\beta})$ . It is clear that this procedure also leads to  $T_n^a(\beta)$ .  $\square$

## References

- [1] **H A Dye, L H Kauffman**, *Virtual crossing number and the arrow polynomial*, J. Knot Theory Ramifications 18 (2009), no. 10, 1335–1357.
- [2] **V F R Jones**, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. (N.S.) 12 (1985), no. 1, 103–111.
- [3] **V F R Jones**, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) 126 (1987), no. 2, 335–388.
- [4] **S Kamada**, *Braid presentation of virtual knots and welded knots*, Osaka J. Math. 44 (2007), no. 2, 441–458.

- [5] **L H Kauffman**, *State models and the Jones polynomial*, *Topology* 26 (1987), no. 3, 395–407.
- [6] **L H Kauffman**, *Virtual knot theory*, *European J. Combin.* 20 (1999), no. 7, 663–690.
- [7] **L H Kauffman**, *A survey of virtual knot theory*, *Knots in Hellas '98 (Delphi)*, 143–202, Ser. Knots Everything, 24, World Sci. Publ., River Edge, NJ, 2000.
- [8] **L H Kauffman**, *An extended bracket polynomial for virtual knots and links*, *J. Knot Theory Ramifications* 18 (2009), no. 10, 1369–1422.
- [9] **L H Kauffman, S Lambropoulou**, *Virtual braids and the L-move*, *J. Knot Theory Ramifications* 15 (2006), no. 6, 773–811.
- [10] **Z Li, F Lei, J Li**, *Virtual braids, virtual Temperley-Lieb algebra and f-polynomial*, *Chin. Ann. Math. Ser. B* 38 (2017), no. 6, 1275–1286.
- [11] **Y Miyazawa**, *A multi-variable polynomial invariant for virtual knots and links*, *J. Knot Theory Ramifications* 17 (2008), no. 11, 1311–1326.
- [12] **V V Vershinin**, *On homology of virtual braids and Burau representation*, *Knots in Hellas '98, Vol. 3 (Delphi)*, *J. Knot Theory Ramifications* 10 (2001), no. 5, 795–812.
- [13] **Y Zhang, L H Kauffman, M-L Ge**, *Virtual Extension of Temperley–Lieb Algebra*, preprint, 2006, arXiv:math-ph/0610052.

Luis Paris, *IMB, UMR 5584, CNRS, Univ. Bourgogne Franche-Comté, 21000 Dijon, France*

Loïc Rabenda, *IMB, UMR 5584, CNRS, Univ. Bourgogne Franche-Comté, 21000 Dijon, France*

[lparis@u-bourgogne.fr](mailto:lparis@u-bourgogne.fr), [loic.rabenda@gmail.com](mailto:loic.rabenda@gmail.com)