



**HAL**  
open science

## **Economic impact of human-induced shrinkage of Posidonia oceanica meadows on coastal fisheries in the Gabes Gulf (Tunisia, Southern Mediterranean Sea)**

Radhouan El Zrelli, Lotfi Rabaoui, Rubén Roa-Ureta, Nicola Gallai, Sylvie Castet, Michel Grégoire, Nejla Bejaoui, Pierre Courjault-Radé

► **To cite this version:**

Radhouan El Zrelli, Lotfi Rabaoui, Rubén Roa-Ureta, Nicola Gallai, Sylvie Castet, et al.. Economic impact of human-induced shrinkage of Posidonia oceanica meadows on coastal fisheries in the Gabes Gulf (Tunisia, Southern Mediterranean Sea). *Marine Pollution Bulletin*, 2020, 155, pp.111124. 10.1016/j.marpolbul.2020.111124 . hal-02949095

**HAL Id: hal-02949095**

**<https://hal.science/hal-02949095>**

Submitted on 20 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**Economic impact of human-induced shrinkage of *Posidonia oceanica* meadows on coastal fisheries in the Gabes Gulf (Tunisia, Southern Mediterranean Sea)**

Radhouan El Zrelli<sup>a,b\*</sup>, Lotfi Rabaoui<sup>c</sup>, Rubén H. Roa-Ureta<sup>c</sup>, Nicola Gallai<sup>d</sup>, Sylvie Castet<sup>a</sup>, Michel Grégoire<sup>a</sup>, Nejla Bejaoui<sup>b</sup> & Pierre Courjault-Radé<sup>a</sup>

<sup>a</sup> *Géosciences Environnement Toulouse (GET), Université de Toulouse, UMR 5563 CNRS/UPS/IRD/CNES, 14 Avenue Edouard Belin, 31400 Toulouse, France*

<sup>b</sup> *Marine Studies Section, Center for Environment and Water, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia*

<sup>c</sup> *Institut National Agronomique de Tunis (INAT), Université de Carthage, 43 Avenue Charles Nicolle, 1082 Tunis Maharajène, Tunisia*

<sup>d</sup> *LEREPS, ENFA, Université Fédérale Toulouse Midi-Pyrénées, Toulouse Cedex, F 31042, France*

\* Corresponding author: Radhouan El Zrelli (Email: [radhouan.elzrelli@gmail.com](mailto:radhouan.elzrelli@gmail.com))

## **Abstract**

In early XX<sup>th</sup> century, the Gulf of Gabes (SE Tunisia) used to host the most extended *Posidonia oceanica* seagrass beds in the Mediterranean Sea, and a highly productive hotspot of benthic/demersal biodiversity. Sponge harvesting and seabed trawling provoked a first step of seagrass degradation. Subsequently, phosphogypsum releases from Gabes Industrial Complex, since mid-1970s, accelerated the decline of the remaining patches. A sharp reduction of coastal fisheries landings took place with the establishment of the last industrial plant units in 1985. The decrease in coastal commercial species landings was found to be directly correlated with *P. oceanica* decline. The trophic web system switched from a ‘benthic-dominated’ to a ‘pelagic-dominated’ system. The economic loss related to coastal fisheries was estimated at ~60 million € in 2014 and the 1990-2014 cumulated loss exceeded €750 million. This first economic valuation of the only direct-use consumptive value of the coastal fishing service provided by *P. oceanica* in Gabes Gulf is a first step towards the assessment of the environmental cost of the negative externalities caused by the local phosphate industry. It may be used as a preliminary decision-making aid to consider alternative industrial solutions.

**Keywords:** Gabes Gulf; *Posidonia oceanica*; Ecosystem services; Coastal fisheries; Economic valuation; Phosphogypsum.

## 1. Introduction

Over 40% of the oceans are impacted by pollution (Halpern et al., 2015). In the last fifty years, the number of marine dead zones has doubled with each passing decade (Diaz and Rosenberg, 2008; Limburg et al., 2020). Marine ecosystems contaminated by pollutants such as toxic runoff, dredged materials, oil spills, trash and debris, industrial wastes and sewage sludge, are persistently disrupted (Rabaoui et al., 2020). The Mediterranean basin is among the most impacted regions in the world together with the North Sea, South and East China Seas, east coast of North America, Red Sea and Arabian Gulf (Halpern et al., 2008, 2015, 2019). Consequently, the socio-economic status of some coastal fisheries has become threatened by habitat degradation (Pedro et al., 2020). Such an environmental and socio-economic status regarding coastal fisheries is the rule for all countries bordering the Mediterranean basin, where more than 250 million people live on the coasts (UNEP, 2012) along with more than 230 million tourists per year (Pons et al., 2009). The United Nations Environment Program (UNEP) has estimated that 650 million tons of sewages, 129 thousand tons of mineral oil, 60 thousand tons of mercury, 36 thousand tons of phosphates and 380 tons of lead are dumped into the Mediterranean Sea each year (UNEP, 2012).

The good health status of essential fish habitats like seagrass ecosystems is an essential factor for the maintenance of Mediterranean coastal fishing economic activity because of its ecosystemic functions for benthic and demersal commercial species (Unsworth et al., 2018). Seagrass meadows are among the most valuable coastal ecosystems worldwide in terms of goods and services they provide (Costanza et al., 1997; Hemminga and Duarte, 2000). According to Waycott et al. (2009), almost one third of the areal extent of seagrass has disappeared globally since the end of the XIX<sup>th</sup> century. In the Mediterranean basin, *Posidonia oceanica* is the most important endemic seagrass species and it forms meadows or

beds covering an area between 21 and 55 thousand km<sup>2</sup> (Pasqualini et al., 1998). Among the ecosystemic services provided by *P. oceanica* is the fishing income (Costanza et al., 1997; Campagne et al., 2015; Vasallo et al., 2015; Boudouresque et al., 2016). Direct monetary assessment of coastal fisheries through transformation of benthic and demersal species volumes into financial value testify of the concrete human benefits derived from ecological functions supplied by the seagrass meadows. Consequently, they can be considered as an ecosystem service *sensu* Liqueste et al. (2016) and as "supporting services" *sensu* MEA (2005). A first global valuation of ecosystemic services of all types of seagrass at the worldwide scale has been estimated at about 19,000 US\$ ha<sup>-1</sup> y<sup>-1</sup> (Costanza et al., 1997). After this founding work, economic valuations of these systems have been sparse (Dewsbury et al., 2016). This is all the truer in the case of the Mediterranean Sea, where few papers have addressed the value of ecosystem services provided by *P. oceanica* seagrass (Vasallo et al. 2013; Campagne et al., 2015).

The seagrass meadows of the Gulf of Gabes (SE Tunisia; Fig. 1) used to be considered as the most extensive beds in the Mediterranean Sea (De Gaillande, 1970a). But since the beginning of XX<sup>th</sup> century, they declined drastically because of multiple anthropogenic pressures (e.g. Ben Brahim et al., 2010; El Zrelli et al., 2017). According to Zaouali et al. (1993), the decline of *P. oceanica* meadows from the Gulf of Gabes has occurred in three main phases: the first phase started with the development of the destructive dredge (or *gangava*), used to collect sponges, which had induced a first local degradation of the meadows leading to the initiation of a slow silting process in the central area of the Gulf (Poizat, 1970). The second phase happened between the mid and late sixties when a major shrinkage of *Posidonia* beds took place following the development of bottom trawling techniques for shrimping. This led to limit the *Posidonia*-covered areas to coastal strips only and to a depth rarely exceeding 20-25 m, and consequently has triggered the silting process

from the centre of the Gulf to the coast (Poizat, 1970; Ktari-Chakroun and Azzouz, 1971). As for the third phase, it began in the mid-seventies when the remaining coastal strip of *P. oceanica* beds gradually disappeared around the first unit of Gabes phosphate fertilizer plant, put into operation in 1972 (Darmoul et al., 1980; Darmoul, 1988). By the second half of the seventies, the large quantities of dumped phosphogypsum wastes ( $\approx 14\,400\text{ T. d}^{-1}$  of dry PG; El Zrelli et al., 2018a) in the surrounding marine environment were responsible for degradation of *Posidonia* beds in front of the industrial complex due to the increase of turbidity coupled with the radioactive and chemical contamination of seawater (Darmoul et al., 1980; Darmoul, 1988; El Zrelli et al., 2018b; El Zrelli et al., 2019a). Several studies carried out across the central area of Gabes have confirmed a continuous degradation of the coastal belt of *P. oceanica* triggered by phosphogypsum releases (Pergent and Kempf, 1993; Zaouali, 1993; El Zrelli et al., 2017).

This continuous anthropogenic-induced regression of *P. oceanica* meadows has led to significant changes in the biodiversity of the impacted coastal area over the last few decades, causing thus significant changes in the catch composition and incomes of the coastal fisheries sector. In this context, several studies reported that the fisheries resources of the Gulf of Gabes are highly exploited (Hattab et al., 2013; Halouani et al., 2016, 2019), thus threatening the socioeconomic situation of Gabes coastal fisheries (FAO ArtFiMed, 2011a). In 1993, ~39 thousand people were directly employed in the coastal fisheries segment and only ~20 thousand in 2009 (FAO ArtFiMed, 2011a). Production varied from ~36 thousand tons in 1986-1990 (Pergent and Kempf, 1993) to less than 14 thousand tons in 2009 (FAO ArtFiMed, 2011a). No studies have been conducted so far on the changes on coastal fisheries neither on the valuation of economic changes related to the losses in *P. oceanica* seagrass habitats. This work was conducted with the aims *i*) to describe the decadal changes in the catch and composition of coastal fisheries in relation to the decline of *P. oceanica* beds and increase of

phosphogypsum wastes discharges, and *ii*) to assess the losses of economic value proceedings from goods and benefits provided by coastal fisheries in connection with the decline of *P. oceanica* meadows in the Gulf of Gabes.

## 2. Materials and Methods

### 2.1. Study area

The Gulf of Gabes extends between Ras Kaboudia in the north and the Libyan/Tunisian boundary in the south (Fig. 1a). It is the widest area of continental shelf in the Mediterranean Sea together with the North-Central Adriatic shelf. It covers about 36 thousand km<sup>2</sup> (Hattab et al., 2013). The study area concerns the central part of Gulf of Gabes which extends between the north-western coast of Djerba Island and El Akarit covering more than 100 km of coastline (Fig. 1). This area hosts three main landing sites: Ghannouch, Gabes and Zaratt (Fig. 1a). Among these, Gabes landing site is the most important since its commercial landings represent more than 80% of the total commercial landings from the 3 sites. They are subdivided into “*small pelagics*” and “*coastal fisheries*”. Since 1984, the small pelagics group represents an average of  $\approx 80\%$  of the commercial landings of Gabes governorate.

### 2.2. Fisheries Data collection

Annual volumes of landed catches of coastal and small pelagic fisheries were collected, over 36 years extending from 1980 to 2016. The 1980-1993 data were collected from the General Fisheries Commission in Tunisia (*Commission Générale des Pêches de Tunisie*; CGP, 1996); while the 1994-2015 data were provided by the Tunisian Directorate-General for Fisheries and Aquaculture (*Direction Générale de la Pêche et de l’Aquaculture*: DGPA; <http://www.onagri.nat.tn>). Data collected were checked from various reports and since the

catch volumes of coastal fisheries were not solid enough for the two years 1984 and 1986, these two records were excluded from the economic analyses ([Appendix I](#)).

Unfortunately, no official records of coastal and small pelagic fisheries catches were available before the eighties. According to [FAO ArtFiMed \(2011b\)](#), the coastal benthic/demersal landed volumes, before the eighties, were low and mostly composed of cephalopods (e.g. the common octopus, *Octopus vulgaris*) sold mainly in local markets ([M'Rabet et al., 2011](#)). At that time, there were no estimations for volumes sold in local fish markets or indirectly by fishermen.

### 2.3. Economic data collection

Official economic data were obtained from the CGP, between 1990 and 2014. The starting year of 1990 was determined by (i) uncertain reliability of data before this date, (ii) its coincidence with the beginning of the second phase of landed volumes evolution ([Appendix I](#)), and (iii) with the end of the quick degradation phase of *Posidonia oceanica* meadows. The market prices of fish sold in Gabes market were in Tunisian Dinars (TND). These prices were updated (<http://mon-convertisseur.fr>) and converted to Euros ([www.xe.com/fr/currencyconverter/](http://www.xe.com/fr/currencyconverter/)).

### 2.4. Prediction of the Area of *Posidonia* meadows over the period 1970 to 2014

The quantification of marine area covered by *Posidonia* meadows since the 60s has been established for four years using cartographic data of previous works ([Appendix II](#)). *Posidonia* maps have been digitized and their areas has been estimated using ArcMap software (version 10.2.2, [Fig. 1b-e](#)). Although this is a small sample, it is informative because it covers a long period of time and it has a clear trend of linear shrinking ([Fig. 2a](#)). Taking advantage of this informative sample, we fitted a linear regression model of *Posidonia* meadows area versus the year to predict the extension of these meadows in the years without

measurements. These predictions and their prediction standard errors were then used for determining the effect of the *Posidonia* meadows shrinkage on fisheries output.

### *2.5. Relationship between *P. oceanica* meadow extension and discharges of phosphogypsum*

We built a database with the annual meadow extension prediction and discharge of phosphogypsum in megatons into the Gulf of Gabes spanning the period of 1972 to 2016. Assuming that discharges of phosphogypsum have an additive effect, we examined the relation between cumulative annual discharges and the extension of *Posidonia* meadows with generalized linear models (*glm*). Taking into account that *Posidonia* meadow extension is a right-skewed positive random variable, we fit the *glm* assuming a gamma distribution with the log link function. The Gamma distribution model is appropriate for positive, continuous and right-skewed data with a variance that is nearly constant on the log scale.

Since meadow extension was not directly measured data but rather a prediction from the linear model described in §2.4, it was necessary to account for the prediction error when fitting the generalized linear model. To that effect we fitted the *glm* to each one of 1000 realizations of the predicted meadow extension, with each realization coming from a truncated normal distribution defined by a mean ( $\mu$ ) equal to the predicted extension, a standard deviation ( $\sigma$ ) equal to the predicted standard error of prediction, and bounds of  $[\mu-0.25\sigma, \mu+0.25\sigma]$ . This marginalization of the distribution of the meadow extension data accounted for some, though not all, the uncertainty from replacing actual meadow extension data with the predictions from the linear regression between year and the four available direct measurements of meadow extension.

### *2.6. Economic analysis*

Our goal is to assess the economic impact on commercial coastal fisheries in the central part of Gulf of Gabes over the last twenty-five years in relation to the gradual

shrinkage of *P. oceanica* seagrass meadows. We value the provisioning services by *P. oceanica* meadows sold on official Tunisian market using the market price-based approach (Silvis and Van der Heid, 2013), where the main output of the service offered by *P. oceanica* is the production of benthic and demersal commercial fishery stocks. This method consisted in valuation of the part of benthic/demersal species value dependent on *Posidonia*. We assume that the production in tons of benthic/demersal species at a time  $t$  ( $Q_t$ ) is as follows:

$$Q_t = a_t p_t = b_t z_t \quad (1)$$

where  $p$  is the area of *P. oceanica* per hectare (ha) at time  $t$ ,  $z$  the set of ecological factors that contribute to the production of benthic/demersal species per ha,  $a$  and  $b$  are the corresponding yield coefficients. The total value of *Posidonia* corresponds to  $Q_t$  multiplied by the market price of benthic/demersal at time  $t$  ( $m_t$ ),

$$V_t = m_t Q_t \quad (2)$$

From the equation 2, we can determine the hectare value of *Posidonia* at time  $t$ :

$$v_t = m_t a_t \quad (3)$$

We also measured the economic losses due to the degradation of *Posidonia* (ELDP) meadows in the Gulf of Gabes. For that, we compared the maximum potential quantity of *Posidonia*,  $P$ , with the quantity  $p_t$ , which is the quantity of *Posidonia* at time  $t$ . Then the loss value ( $l_t$ ) can be measured by the following formula:

$$l_t = v_t (P - p_t) \quad (4)$$

## 2.7. Fishing effort modelling (2000-2016)

Fishing effort data, including the number of boats and number of fishermen, were collected on yearly basis from the annual reports of fisheries statistics published by DGPA. To relate fisheries output to *P. oceanica* meadows extension, we built a database consisting of annual fisheries output in tons of landings, fishing effort in number of boats operating per

year, the prediction of meadows extension and the prediction standard error. The fishery effort and output were split between two fleets operating in the Gulf of Gabes, the coastal fleet and the small pelagic fish fleet. This database covered the years between 2000 and 2016. The limiting factor was the unavailability of fishing effort of both fleet components before 2000. Generally, the raw fishing yield data show differences between the coastal and the small pelagic fish fleets (Fig. 2b). The former shows a fairly constant annual yield around 2-3 thousand tons and a rise in effort over the last few years, while the latter shows a decreasing yield and constant effort (Fig. 2c).

To test the effect of meadows extension, we fitted a *glm* with the fisheries output as response variable and the fishing effort and the meadows extension as potential predictor variables, each fleet component separately. The fishing output data was assumed to have a gamma probability distribution with a log link function. As in the previous analyses, we accounted for the uncertainty in the prediction of meadows cover by marginalizing over 1000 realizations of the annual series of meadow shrinkage with the same parameters used to generate the data for the modelling of meadows extension versus cumulative phosphogypsum discharge.

To test the effect of the meadow extension on the fishing output, we fitted a *glm* where both the fishing effort and the meadow extension were included as predictors versus a model where the fishing effort was the single predictor. We then calculated the Akaike Information Criterion (AIC) difference between the model with both predictors versus the model with fishing effort as single predictor. Under the hypothesis that the extension of meadows did not contribute to the fishing output in addition to the effect of fishing effort, the distribution of the AIC difference across the 1000 realizations would be symmetrical and centered on 2 (Sakamoto et al., 1986). Conversely, under the hypothesis of the effect of the meadow

extension on fishing output the distribution of the AIC difference would be skewed to the left, and the stronger the effect the stronger the skewness.

### 3. Results

#### 3.1. Historical decline of *P. oceanica*

*P. oceanica* meadows have shrunk drastically along the last 5 decades. In spite of having only four data points the decline was well parameterized by the linear regression prediction, with rather narrow confidence bands (Fig. 2a). The reasons for this success with just four data points were that these points are well spaced in time and they are falling neatly on the linear trend (Fig. 2a). At the estimated rate of shrinkage, the meadows will disappear any year now, although the precision of this prediction allows for residual patches well into the future.

#### 3.2. Economic valuation

Table 1 summarizes the main results of the economic valuation of *Posidonia* in the Gulf of Gabes in 1990 and 2014. First, we observed that the value has increased by 263% in the period, from €2 million to more than €7.5 million. This increase is due to the high increase in market price of benthic species combined with the rise of benthic species production. In the same periods, the yield of fish has varied from 0.024 to 0.17 per hectare of *Posidonia*. Thus, the value of *Posidonia* per hectare was €48 in 1990 and more than €500 in 2014. Between 1990 and 2014, the loss of cover of *Posidonia* meadows in the Gulf of Gabes has resulted in a loss of value from €4 million initially (1990) to almost €60 million currently (2014).

#### 3.3. The phosphogypsum-*Posidonia*-fisheries connection

The hypothesis of the effect of cumulative phosphogypsum discharge on predicted *Posidonia* cover is overwhelmingly more supported than the alternative of no effect (Fig. 3a).

This is noted in much lower AIC values for the hypothesis that included the cumulative phosphogypsum discharge effect versus the null model that did not include that effect (Fig. 3a). In addition, the largest  $p$ -value of the null hypothesis of no cumulative phosphogypsum discharge effect across 1000 realization of the predicted *Posidonia* cover was very close to zero. Examination of the coefficients showed that every new megaton of phosphogypsum discharged into the Gulf of Gabes reduces *Posidonia* meadow cover by 1 hectare on average (Fig. 3b). Further, in the period from 2000 to 2016, the hypothesis of the effect of *Posidonia* meadows shrinkage is not supported for the coastal fishery but it is strongly supported for the small pelagics fishery (Fig. 3c & d). In the latter, the loss of 1 hectare of *Posidonia* meadows is related on average the loss of 1 tonne of fisheries yield.

#### 4. Discussion

Since the early 70s, the central part of the Gulf of Gabes has been severely impacted by high anthropogenic pressures of industrial origin (Mosbahi et al., 2019; El Kateb et al., 2020; Fourati et al., 2018), represented mainly by the huge untreated phosphogypsum quantities released directly in the coastal environment since 1972 (El Zrelli et al., 2015, 2018a). We have shown in this work that over the same period seagrass cover loss occurred linearly and connected to the cumulative effect of phosphogypsum discharges. The most likely mechanism of decline was a widespread process of siltation of the sea bottom. Further, Cyanophyceae proliferation increased markedly because of nutrients enrichment discharged by the phosphate processing industries directly into sea water (Zaouali, 1993). The benthic ecosystem became dominated by invertebrates of muddy substrata enriched with organic matter (Zaouali, 1993). This transformation of the benthic/demersal environment can be considered as a shift in the biocenosis of the central Gulf of Gabes which occurred between 1984/1985 and 1990 (Zaouali, 1993; Guillaumont et al., 1995).

The results showed that *Posidonia* meadows have an important contribution to the local economy with a per hectare value of €511 and a total loss of value of ~ €60 million per year (Table 1). As previously mentioned, very few papers addressed this value in the scientific literature especially for the endemic Mediterranean *Posidonia*. For example, Campagne et al. (2015) valued the *Posidonia* seagrass at €513 per hectare and a total of €55 million. But these values have been evaluated for several services only one of which was the contribution to the coastal fisheries. In addition, it is valid only in the socioeconomic French context (Campagne et al., 2015). Nevertheless, an attempt to determine the economic value of *Cymodocea nodosa* seagrass meadows for local fisheries at the Island of Gran Canaria using standard market prices has been carried out giving a monetary value at €866 per ha (Tuya et al., 2014) which is comparable to our valuation.

The loss estimate is an indicator that demonstrates the potential of economic gains in developing the *Posidonia* meadows. Indeed, the degradation of *Posidonia* meadows seems constant at least since 1990. If we assume that this degradation is linear and so is the increase in loss, a cumulated loss from 1990 to 2014 would give a total loss of more than €750 million. Such a monetary loss should be considered by stakeholders that depend essentially on the fishing activity.

Comparing the economic losses due to the degradation of *P. oceanica* from the Gulf of Gabes to the added values of Gabes Industrial Complex (represented mainly by the GCT (*Groupe Chimique Tunisien*) factories), during the 1990-2014 period, we found that *Posidonia* meadows of Gabes Gulf are economically more important (Table 2). In fact, the economic losses related to the shrinkage of *P. oceanica* represented, in 2014, 39% and 64% of the added value of all Tunisian GCT factories and those of Gabes only, respectively. The cumulative economic losses, over the 1990-2014 period (15 years), reached ~ €780 million, so that the quarter of added values of Tunisian GCT factories and ~ 43% of the added values

of Gabes GCT factories (Table 2). According to the study conducted by the European Community in 2017 on the economic losses related to the industrial pollution in Gabes (EC, 2017), the estimated annual cost of environmental degradation in Gabes Gulf was found to reach a maximum of €34 million ( $\approx 76$  million Tunisian Dinar), which represents nearly 27% of the added values of Tunisian GCT factories. However, it is worth noting that this latter economic valuation did not take into consideration the impact of phosphogypsum pollution on tourism and fisheries sector in the Gulf of Gabes in isolation, but also the impact of air pollution on other agricultural sectors, tourism, and human health. The fisheries sector *per se* represents  $\sim 47\%$  ( $\sim \text{€}16$  million) of this latter estimated amount (EC, 2017). Since the economic loss estimated during this study was found to be close to €59 million in 2014 (Table 2), it appeared that the EC (2017) study largely underestimated the economic losses in Gabes area ( $\text{€}16$  million *versus*  $\text{€}59$  million), although it took into account a wider range of economic sectors.

Ecological aspects of the *Posidonia* meadows shrinkage concern the life cycle of benthic/demersal coastal species, the role of *Posidonia* as primary producer and the morphological particularities of the central Gulf of Gabes. The strictly linear relationship applied between landing volumes of coastal fisheries and *Posidonia* meadows area is the most critical approximation used to carry out the evaluation. An estimation of the degree of residence of benthic commercial species has been proposed to improve the economic evaluation accuracy of *Posidonia* ecosystem services (Jackson et al., 2015). According to the available literature, the residence degree can be evaluated for the principal commercial landed species in the Gabes governorate (FAO, 2016). More than  $\sim 90\%$  of the benthic and demersal commercial species landed in the Gabes governorate basically depend on the presence of *Posidonia* meadows (FAO ArtFiMed, 2011b; FAO, 2016). Consequently, the linear

approximation retained to qualify the link between landed volumes and the seagrass beds is not far off with regards of the economic evaluation.

Phytoplankton constitutes the first step of the marine food chain sustaining all the aquatic life (e.g. [Field et al., 1998](#)) and especially oceanic life ([Duarte and Cebrian, 1996](#)). It feeds on mineral elements, mostly nitrogen and phosphorus. They are used to produce organic material via the photosynthesis process. The organic matter is reusable by zooplankton and then consumed by a wide variety of filter-feeding organisms. Nevertheless, Mediterranean waters are meso-oligotrophic to oligotrophic i.e. more or less depleted of mineral nutrients ([Estrada et al., 1985](#); [Herut et al., 2005](#)). In this situation of overall nutrient deficiency, the eastern Mediterranean basin is the most oligotrophic region ([Berman et al., 1984](#)) and the Gulf of Gabes is known as the most oligotrophic region of the eastern basin ([Drira et al., 2008](#)) because of weakness of mineral stock inherited from arid hinterlands and limited intrusions of Atlantic waters into the Gulf ([Bel Hassen et al., 2008](#)). Therefore, coastal waters of the Gulf offer very low levels of nutrients and little to sustain organisms which should result in low levels of biological productivity. Nevertheless, the Gulf of Gabes has the peculiarity of being highly productive. This contrast is a matter of debate. Some authors emphasise the role played by certain types of phytoplankton and zooplankton to explain the productivity of the Gabes Gulf ([Hamdi et al., 2015](#); [Drira et al., 2009](#); [Béjaoui et al., 2020](#)). An alternative view could be formulated in terms of the food web system. The combination of the well-established role of primary producer of *Posidonia* coupled with the great extent of the continental shelf covered by these seagrass meadows may be a pertinent approach to the paradoxical situation – ultra-oligotrophy vs. high productivity - of the Gulf of Gabes. The hypothesis herein proposed is to view the shallow-water (<50m) of the central part of Gabes Gulf food web as benthic-dominated rather than plankton-dominated.

Firstly, *Posidonia* seagrass are nursery habitats because the leafy underwater canopy they create provides shelter for small invertebrates, small fish and juveniles of larger fish species (Jehly et al., 2010). Other invertebrates grow nestled between the blades or in the sediments. The accumulation of smaller organisms amongst and on the *Posidonia* blades, as well as the *Posidonia* itself, attracts bigger animals. Thus, *Posidonia* can be home to many types of fish and invertebrates.

Secondly, *Posidonia* meadows contribute to global nitrogen cycling (Costanza et al., 1997; Duarte et al., 2010; Unsworth and Cullen, 2010). The role of *Posidonia* is essential because it retains mineral nutrients by reducing the water flow under the canopy of leaves, thus enabling nutrients to be trapped (Jehly et al., 2010). Consequently, the *Posidonia* biocenosis favoured expansion of phytoplankton populations as a permanent source of nutrients like nitrogen, phosphorous, and organic carbon used by them (Jehly et al., 2010). Nevertheless, the primary productivity of *Posidonia* is not the major direct source of carbon and nitrogen for higher trophic levels of the food web and most *Posidonia* meadow invertebrate herbivores utilize epiphytic species (Unsworth and Cullen, 2010). *Posidonia* is an adequate support for epiphytic species (Boudouresque et al., 2006; Duarte et al., 2004) and many species of algae and microalgae, bacteria and invertebrates that grow as “epiphytes” directly on living *Posidonia* leaves. Small invertebrates, such as crustaceans and snails, feed on epiphytes. They are in turn consumed by larger crustaceans and fish and are important links in the coastal food web (Duarte et al., 2004; Vizzini, 2009; Vizzini et al., 2002). Further, some epiphytic bacteria can extract nitrogen from the environment and make it available to larger animals (Jehly et al., 2010). Epiphytes are commonly associated to *Posidonia* leaves in the Gulf of Gabes (Ben Brahim et al., 2010).

Lastly, dead *Posidonia* leaves also play an important role in coastal ecosystems. When the leaves die, they decay on the sediment or are washed onto the beach, supporting a diverse community of decomposers that prosper on rotting material.

Taking into account the situation of small pelagic fisheries, it is worth noting that the efficiency of the purse seine technique coupled with the benthic to pelagic biocenosis shift hypothesis has strongly favoured the development of small pelagic fishing activity. Nevertheless, its decline at the end of the eighties (Fig. 2b & c) most probably corresponded to the effects of pollution load of the marine environments which reached a larger area in the central part of Gabes Gulf (Rabaoui et al., 2015; El Zrelli et al., 2017, 2019b & c). The high proportion of toxic phytoplankton indicates that quality of marine waters is still poor (Feki et al., 2013; Abdennadher, 2014). Nowadays, part of small pelagic catch by coastal fisheries techniques (as beach seine for example) has increased, especially the sardinella volume (Fig. 3a & b). This could be the consequence of the high nutrients amounts in coastal sea water. It is probably the reason why the coastal fisheries volumes are irregular from one year to the next and can be, from times to times, relatively high (Fig. 2b & c).

A first step towards a better understanding of the vulnerability of the Gabes Gulf could consist in introducing an economic measure of this vulnerability in terms of producer surplus. The producer surplus loss measures the difference between the amounts a producer of a good receives, the market price, and the minimum amount the producer is willing to accept for the good. This minimum amount is called the marginal cost. The producer surplus is represented in Fig. 4, where we assume that the demand for fish of the Gulf of Gabes is perfectly elastic. The degradation of the *Posidonia* seagrass increases the marginal cost for the fishermen. This is translated into a move of the supply curve of fish from  $S_0$  to  $S_1$  or  $S_2$  (Fig. 4). The producer surplus loss is estimated through the shaded area in Fig. 4: the more the marginal costs increase, the more the producer surplus losses increase.

Our indicator value and loss can be integrated in this figure. The value is represented by the rectangle  $pOQ_1S_1$ , when considering a loss of *Posidonia* from  $Q_0$  to  $Q_1$ , and the rectangle  $pOQ_2S_2$ , when considering a loss of *Posidonia* from  $Q_0$  to  $Q_2$ . The loss is represented by the rectangle  $S_1Q_1Q_0S_0$ , when considering a loss of *Posidonia* from  $Q_0$  to  $Q_1$ , and the rectangle  $S_2Q_2Q_0S_0$ , when considering a loss of *Posidonia* from  $Q_0$  to  $Q_2$ . The producer surplus can be considered as much accurate to value the *Posidonia* seagrass when the loss is important. Indeed, if we consider to change from  $S_0$  to  $S_2$  and use our value indicator, we can observe graphically in [Fig. 4](#) that the value will be very low compared to the real producer surplus loss. However, the producer surplus method is not easy to apply in a large territory because of the need of data on marginal cost or production from fishermen which is difficult to obtain.

Furthermore, we focus our analysis on the marketed direct use value of the seagrass. Thus, we studied the supply side and the impact of *Posidonia* seagrass on the fishermen benefits. However, the economic value of *Posidonia* seagrass is not limited to this as seen in [Campagne et al. \(2015\)](#). Indeed, it provides benefits through the regulation and maintenance services, water purification value, water oxygenation and cultural services. In order to complete our economic valuation of *Posidonia* seagrass in the Gulf of Gabes and give more accurate information to decision makers, a future line of research would be to realize these valuations, including interviewing various stakeholders, other than fishermen, living around the Gulf of Gabes and value the benefits of the cultural service.

## 5. Conclusion

A gradual intensification of sponge, then shrimp fishing activities during the first half of the 20<sup>th</sup> century has triggered the process of *Posidonia* meadows area reduction in the central Gulf of Gabes. Subsequently, between 1972 and 1985, successive commissioning of phosphate fertilizer plant units in Gabes has induced a high environmental stress via

phosphogypsum releases into coastal marine environments. Furthermore, the gradual disappearance of the seagrass beds caused the gradual silting up of the central part of the Gulf which progressed from centre to coast. This silting process contributed to the waters transparency decrease and thus to gradual disappearance of the seagrass beds. From an economic point of view, the loss of economic value provided by coastal fisheries income correlated to the decline of the *Posidonia* meadows has been valued at about €60 million in 2014 and a cumulated loss of more than €750 million from 1990 to 2014.

### **Acknowledgments**

The authors would like to thank the editor and the reviewer who contributed to improve the quality of the manuscript through their constructive comments and suggestions. We are also grateful to all those who helped in data collection and analysis. This study is dedicated to the memory of Habib Chairat (09.11.1960/29.10.2018) for his environmental protection efforts in the coastal oasis of Chatt Sidi Abd Essalam.

### **References**

- Abdennadher, M., Hamza, A., Fekih, W., Hannachi, I., Zouari Bellaaj, A., Bradai, M.N., Aleya, L. 2014. Factors determining the dynamics of toxic blooms of *Alexandrium minutum* during a 10-year study along the shallow southwestern Mediterranean coasts. Estuar. Coast. Shelf Sci. 106, 102-111.
- Béjaoui, B., Ben Ismail, S., Othmani, A., Ben Abdallah-Ben Hadj Hamid, O., Chevalier, C., Feki-Sahnoun, W., Harzallah, A., Ben Hadj Hamida, N., Bouaziz, R., Dahech, S., Diaz, F., Tounsi, K., Sammari, C., Pagano, M., Bel Hassen, M. 2019. Synthesis review of the Gulf of Gabes (eastern Mediterranean Sea, Tunisia): Morphological, climatic, physical

- oceanographic, biogeochemical and fisheries features. *Estuar. Coast. Shelf Sci.* 219, 395-408.
- Bel Hassen, M., Drira, Z., Hamza, A., Ayadi, H., Akrouf, F., Issaoui, H., 2008. Summer phytoplankton pigments and community composition related to water mass properties in the Gulf of Gabes. *Estuar. Coast. Shelf Sci.* 77 (4), 645–656.
- Ben Brahim M., Hamza, A., Hannachi, I., Rebai, A., Jarboui, O., Bouain, A., Aleya, L. 2010. Variability in the structure of epiphytic assemblages of *Posidonia oceanica* in relation to human interferences in the Gulf of Gabes, Tunisia. *Mar. Environ. Res.* 70, 411-421.
- Berman T., Townsend D.W., Elsayed S.Z., Trees C.C., Azov, Y. 1984. Optical transparency, chlorophyll and primary productivity in the Eastern Mediterranean near the Israeli coast. *Oceanol. Acta* 7 (3), 367-372.
- Boudouresque, C.F., Mayot, N., Pergent, G. 2006. The outstanding traits of the functioning of the *Posidonia oceanica* seagrass ecosystem. *Biol. Mar. Medit.* 13 (4), 109-113.
- Boudouresque, C.F., Pergent, G., Pergent-Martini, C., Ruitton, S., Thibaut, T., Verlaque, M. 2016. The necromass of the *Posidonia oceanica* seagrass meadow: fate, role, ecosystem services and vulnerability. *Hydrobiologia* 781, 25–42.
- Campagne, C.S., Salles, J.M., Boissery, P., Deter, J., 2015. The seagrass *Posidonia oceanica*: Ecosystem services identification and economic evaluation of goods and benefits. *Mar. Pollut. Bull.* 97, 391–400.
- CGP (Commission Générale des Pêches), 1996. *Annuaire des statistiques des pêches en Tunisie*. Ministère de l'Agriculture, Tunisie.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M., 1997. The value of the world's ecosystem services and natural capital. *Nature* 387, 253–260.

- Darmoul, B., 1988. Pollution dans le Golfe de Gabès (Tunisie) : bilan de six années de surveillance. Bull. Inst. Nat. Sci. Tech. Océanogr. Pêche Salammbô 1 (15), 61-83.
- Darmoul, B., Hadj Ali Salem, M., Vitiello, P., 1980. Effets des rejets industriels de la région de Gabès sur le milieu récepteur. Bull. Inst. Nat. Sci. Tech. Océanogr. Pêche Salammbô 7, 5-61.
- De Gaillande, D., 1970a. Note sur les peuplements de la zone centrale du Golfe de Gabès (Campagne Calypso 1965). Téthys 2(1), 131-138.
- De Gaillande, D., 1970b. Peuplements benthiques de l'herbier de *Posidonia oceanica* (Delile), de la pelouse à *Caulerpa prolifera* Lamouroux du large du Golfe de Gabès. Téthys,2(2), 373-383.
- Deudero, S., Alomar, C., 2015. Mediterranean marine biodiversity under threat: Reviewing influence of marine litter on species. Mar. Pollut. Bull. 98 (1-2), 58–68.
- Dewsbury, B.M., Bhat, M., Fourqurean, J.W., 2016. A review of seagrass economic valuations: Gaps and progress in valuation approaches. Ecosyst. Serv. 18, 68-77.
- Diaz, R.J., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems. Science 321 (5891), 926-929.
- Drira, Z., Bel Hassen, M., Hamza, A., Rebai, A., Bouain, A., Ayadi, H., Aleya, L., 2009 Spatial and temporal variations of microphytoplankton composition related to hydrographic conditions in the Gulf of Gabès. J. Mar. Biol. Assoc. U.K. 89 (8), 1559–1569.
- Drira, Z., Hamza, A., Bel Hassen, A., Ayadi, M., Bouaïn, H., Aleya, L., 2008. Dynamics of dinoflagellates and environmental factors during the summer in the Gulf of Gabès (Tunisia, Eastern Mediterranean Sea). Sci. Mar. 72 (1), 59–71.

- Duarte, C.M., Borum, J., Short, F.T., Walker, D.I., 2004. Seagrass ecosystems: their global status and prospects. In: Polunin, N.V.C. (Ed.), *Aquatic Ecosystems: Trends and Global Prospects*. Cambridge University Press, Cambridge, pp. 281-294.
- Duarte, C.M., Cebrián, J., 1996. The fate of marine autotrophic production. *Limnol. Oceanogr.* 41(8), 1758-1766.
- Duarte, C.M., Marbà, N., Gacia E., Fourquaran, J.W., Beggins, J., Barrón, C., Apostolaki, E.T., 2010. Seagrass community metabolism: Assessing the carbon sink capacity of seagrass meadows. *Global Biogeochem. Cycles* 24, GB4032. doi:10.1029/2010GB003793.
- EC (The European Commission), 2017. Etude d'impact de la pollution industrielle sur l'économie de la région de Gabès, rapport final.
- El Kateb, A., Staldera, C., Martínez-Colón, M., Mateu-Vicens, G., Francescangeli, F., Coletti, G., Stainbank, S., Spezzaferri, S. 2020. Foraminiferal-based biotic indices to assess the ecological quality status of the Gulf of Gabes (Tunisia): Present limitations and future perspectives. *Ecol. Ind.* 111, 105962.
- El Zrelli, R., Courjault-Radé, P., Rabaoui, L., Castet, S., Michel, S., Bejaoui, N., 2015. Heavy metal contamination and ecological risk assessment in the surface sediments of the coastal area surrounding the industrial complex of Gabes city Gulf of Gabes, SE Tunisia. *Mar Poll Bull* 101: 922-929
- El Zrelli, R., Courjault-Radé, P., Rabaoui, L., Daghbouj, N., Mansour, L., Balti, R., Castet, S., Attia, F., Michel, S., Bejaoui, N., 2017. Biomonitoring of coastal pollution in the Gulf of Gabes (SE, Tunisia): use of *Posidonia oceanica* seagrass as a bioindicator and its mat as an archive of coastal metallic contamination. *Environ Sci Pollut Res* 24: 22214–22225

- El Zrelli, R., Rabaoui, L., Daghbouj, N., Abda, H., Castet, S., Josse, C., van Beek, P., Souhaut, M., Michel, S., Bejaoui, N., Courjault-Radé, P., 2018a. Characterization of phosphate rock and phosphogypsum from Gabes phosphate fertilizer factories (SE Tunisia): high mining potential and implications for environmental protection. *Environ Sci Poll Res* DOI: 10.1007/s11356-018-1648-4.
- El Zrelli, R., Rabaoui, L., Ben Alaya, M., Daghbouj, N., Castet, S., Besson, P., Michel, S., Bejaoui, N., Courjault-Radé, P., 2018b. Seawater quality assessment and identification of pollution sources along the central coastal area of Gabes Gulf (SE Tunisia): Evidence of industrial impact and implications for marine environment protection. *Mar Poll Bull* 127: 445-452
- El Zrelli, R., Rabaoui, L., Van Beek, P., Castet, S., Souhaut, M., Grégoire, M., Courjault-Radé, P., 2019a. Natural radioactivity and radiation hazard assessment of industrial wastes from the coastal phosphate treatment plants of Gabes (Tunisia, Southern Mediterranean Sea), *Marine Pollution Bulletin*, 146 : 454-461.
- El Zrelli, R., Rabaoui, L., Abda, H., Daghbouj, N., Pérez-López, R., Castet, S., Aigouy, T., Bejaoui, N., Courjault-Radé, P., 2019b. Characterization of the role of phosphogypsum foam in the transport of metals and radionuclides in the Southern Mediterranean Sea, *J Hazard Mater*, 363: 258-267.
- El Zrelli, R., Rabaoui, L., Ben Alaya, M., Castet, S., Zouiten, C., Bejaoui, N., Courjault-Radé, P., 2019c. Decadal effects of solid industrial wastes on the coast : Gulf of Gabes (Tunisia, Southern Mediterranean Sea) as an example, *Estuarine, Coastal and Shelf Science*, 224 : 281-288.
- Estrada, M., Vives, F., Alcaraz, M., 1985. Life and productivity of the open sea. In: Margalef, R. (Ed.): *Western Mediterranean*. Pergamon Press, Oxford, pp. 148-197.

- FAO 2016. The state of Mediterranean and Black Sea fisheries. General Fisheries Commission for the Mediterranean, Rome, Italy, 42 p.
- FAO ArtFiMed. 2011a. La pêche artisanale à Ghannouch (Tunisie). Passé, présent, avenir. FAO-ArtFiMed Développement durable de la pêche artisanale méditerranéenne au Maroc et en Tunisie. Malaga. 42 p.
- FAO ArtFiMed. 2011b. Guide des principales espèces débarquées par la pêche artisanale à Ghannouch et Akarit (golfe de Gabès, Tunisie). FAO-ArtFiMed Développement durable de la pêche artisanale méditerranéenne au Maroc et en Tunisie. Malaga, Espagne, 52 p.
- Feki, W., Hamza, A., Frossard, V., Abdennadher, M., Hannachi, I., Jacquot, M., Belhassen, M., Aleya, L. 2013. What are the potential drivers of blooms of the toxic dinoflagellate *Karenia selliformis*? A 10-year study in the Gulf of Gabes, Tunisia, southwestern Mediterranean Sea. Harmful Algae 23: 8-18.
- Field, C.B., Behrenfeld, M.J., Randerson, J.T., Falkowski, P., 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. Science 281 (5374), 237–240.
- Fourati, R., Tedetti, M., Guigue, C., Goutx, M., Garcia, N., Zaghden, H., Sayadi, S., Elleuch, B. 2018. Sources and spatial distribution of dissolved aliphatic and polycyclic aromatic hydrocarbons in surface coastal waters of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea). Progress in Oceanography 163, 232-247.
- Guillaumont, B., Ben Mustapha, S., Ben Moussa, H., Zaouali, J., Soussi, N., Ben Mammou, A., Cariou, C. 1995. Pollution impact study in Gabes Gulf (Tunisia) using remote sensing data. Mar. Technol. Soc. J. 29 (2) 46-58.
- Halpern, B.S., Frazier, M., Afflerbach, J., Lowndes, J.S., Micheli, F., O’Hara, C., Scarborough, C., Selkoe, K.A. 2019. Recent pace of change in human impact on the world’s ocean. *Sci Rep* **9**, 11609.

- Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Cotton Rockwood, R., Selig, E.R., Solkoe, K.A., Walbridge, S., 2015. Spatial and temporal changes in cumulative human impacts on the world's ocean. *Nat. Comm.* 6, 7615.
- Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.V., Micheli, F., D'Agrosa, C., Bruno, J.F., Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P., Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global map of human impact on marine ecosystems. *Science* 319 (5865), 948-952.
- Halouani, G., Le Loc'h, F., Shin, Y.J., Velez, L., Hattab, T., Romdhane, M.S., Ben Rais Lasram, F. 2019. An end-to-end model to evaluate the sensitivity of ecosystem indicators to track fishing impacts. *Ecol. Indic.* 98, 121-130.
- Halouani, G., Abdou, K., Hattab, T., Romdhane, M.S., Ben Rais Lasram, F., Le loc'h, F. 2016. A spatio-temporal ecosystem model to simulate fishing management plans: A case of study in the Gulf of Gabes (Tunisia). *Mar. Pol.* 69, 62-72.
- Hamdi, I., M. Denis, M., Bellaaj-Zouari, A., Khemakhem, H., Bel Hassen, M., Hamza, A., A. Barani, A., Bezac, C., Maalej, S., 2015. The characterisation and summer distribution of ultraphytoplankton in the Gulf of Gabès (Eastern Mediterranean Sea, Tunisia) by using flow cytometry. *Cont. Shelf Res.* 93, 27–38.
- Hattab, T., Ben Rais Lasram, F., Albouy, C., Romdhane, M.S., Jarboui, O., Halouani, G., Cury, P., Le Loc'h, F. 2013. An ecosystem model of an exploited southern Mediterranean shelf region (Gulf of Gabes, Tunisia) and a comparison with other Mediterranean ecosystem model properties. *J. Marine Syst.* 128, 159–174.
- Hemminga, M.A., Duarte, C.M., 2000. *Seagrass ecology*. Cambridge University Press, Cambridge.

- Herut, B., Zohary, T., Krom, M.D., Mantoura, R.F.C., Pitta, P., Psarra, S., Rassoulzadegan, F., Tanaka, T., Thingstad, T.F., 2005. Response of East Mediterranean surface water to Saharan dust: On-board microcosm experiment and field observations. *Deep Sea Res Part 2 Top Stud Oceanogr.* 52 (22-23), 3024-3040.
- Jabeur, C., Gobert, B., Missaoui, H., 2001. Typologie de la flottille de pêche côtière dans le golfe de Gabès (Tunisie). *Aquat. Living Resour.* 13 (6), 421-428.
- Jackson, E.L., Rees, S.E., Wilding, C., Attrill, M.J., 2015. Use of a seagrass residency index to apportion commercial fishery landing values and recreation fisheries expenditure to seagrass habitat service. *Conserv. Biol.* 29 (3), 899–909.
- Jehly, A., Moutafian, A., Sagot, T., Shaughnessy, S., 2010. Phytoplankton composition off the coast of Point Revellata, Corsica, France. UCSC Biology Field Quarter, Fall 2010, BIOE159.
- Ktari-Chakroun, F., Azzouz, A., 1971. Les fonds chalutables de la région sud-est de la Tunisie (golfe de Gabès). *Bull. Inst. Nat. Sci. Tech. Pêche Salammbô* 2(1), 5-47.
- Laurans, Y., Rankovic, A., Billé, R., Pirard, R., Mermet, L., 2013. Use of ecosystem services economic valuation for decision making: Questioning a literature blindspot. *J. Environ. Manage.* 119, 208–219.
- Limburg, K.E., Breitburg, D., Swaney, D.P., Jacinto, J. 2020. Ocean Deoxygenation: A Primer. *One Earth* 2 (1), 24-29.
- Liquete, C., Udias, A., Conte, G., Grizzetti, B., Masi, F., 2016. Integrated valuation of a nature-based solution for water pollution control: Highlighting hidden benefits. *Ecosyst. Serv.* 22, 392–401.
- MEA (Millennium Ecosystem Assessment), 2005. *Ecosystems and human well-being: Synthesis.* Island Press, Washington, DC.

- Mosbahi, Serbaji, M.M., Pezy, J.P., Neifar, L., Dauvin, J.C. 2019. Response of benthic macrofauna to multiple anthropogenic pressures in the shallow coastal zone south of Sfax (Tunisia, central Mediterranean Sea). *Environ. Pollut.* 253, 474-487.
- Pasqualini, V., Pergent-Martini, C., Clabaut, P., Pergent, G., 1998. Mapping of *Posidonia oceanica* using aerial photograph and side-scan sonar: Application off the Island of Corsica (France). *Est. Coast. Shelf Sci.* 47 (3), 359-367.
- Pedro, L., Henriques, S., Pérez-Ibarra, I., Trujillo, M., García-Charton, J.A., Vasconcelos, R.P. 2020. Shifting baselines in a Mediterranean small-scale fishery. *Ocean Coast. Manage.* 183, 104985.
- Pergent, G., Kempf, M., 1993. L'environnement marin côtier en Tunisie. IFREMER, Brest, France. <https://archimer.ifremer.fr/doc/00105/21592/>
- Poizat, C., 1970. Hydrodynamisme et sédimentation dans le golfe de Gabès (Tunisie). *Téthys* 2(1), 267-296.
- Pons, P.O., Crang, M., Travlou, P., 2009. Culture of mass tourism - Doing the Mediterranean in the age of banal mobilities. Routledge, Taylor and Francis Group, London and New York.
- Rabaoui, L., El Zrelli, R., Ben Mansour, M., Balti, R., Mansour, L., Tlig-Zouari, S., Guerfel, M., 2015. On the relationship between the diversity and structure of benthic macroinvertebrate communities and sediment enrichment with heavy metals in Gabes gulf Tunisia. *J. Mar. Biol. Assoc. U.K.* 95, 233–245.
- Rabaoui, L., Cusack, M., Saderne, V., Krishnakumar, P.K., Lin, Y.J., Shemsi, A.M., El Zrelli, R., Arias-Ortiz, A., Masqué, P., Duarte, C.M, Qurban, M.A., 2020. Anthropogenic-induced acceleration of elemental burial rates in blue carbon repositories of the Arabian Gulf. *Sci. Total Environ.* 719, 135177.

- Sakamoto, Y., Ishiguro, M., Kitagawa, G., 1986. Akaike information criterion statistics. KTK Scientific Publishers, Tokyo.
- Silvis, H.J., van der Heide, C.M., 2013. Economic viewpoints on ecosystem services. Wageningenur, Wettelijke Onderzoekstaken Natuur & Milieu. WOt-rapport 123. 68 p.
- Tuya, F., Haroun, R., Espino, F., 2014. Economic assessment of ecosystem services: Monetary value of seagrass meadows for coastal fisheries. *Ocean Coast. Manage.* 96, 181-187.
- UNEP (United Nations Environment Program), 2012. État de l'environnement marin et côtier de la Méditerranée. Mediterranean Action Plan (Barcelona Convention), UNEP, Athens.
- Unsworth, R.K.F., Cullen, L.C., 2010. Recognising the necessity for Indo-Pacific seagrass conservation. *Conserv. Lett.* 3 (2), 63–73.
- Unsworth, R.K.F., Nordlund, L.M., Cullen-Unsworth, L.C., 2018. Seagrass meadows support global fisheries production. *Conserv. Lett.* e12566. <https://doi.org/10.1111/conl.12566>
- Vassallo, P., Paoli, C., Rovere, A., Montefalcone, M., Morri, C., Bianchi, C.N., 2013. The value of the seagrass *Posidonia oceanica*: A natural capital assessment. *Mar. Pollut. Bull.* 75 (1–2), 157–167.
- Vizzini, S., 2009. Analysis of the trophic role of Mediterranean seagrasses in marine coastal ecosystems: A review. *Bot. Mar.* 52 (5), 383-393.
- Vizzini, S., Sarà, G., Michener, R.H., Mazzola, A., 2002. The role and contribution of the seagrass *Posidonia oceanica* (L.) Delile organic matter for secondary consumers as revealed by carbon and nitrogen stable isotope analysis. *Acta Oecol.* 23 (4), 277–285.
- Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck Jr., K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Short, F.T., Williams, S.L., 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *PNAS* 106 (30), 12377–12381.

Zaouali, J., 1993. Les peuplements benthiques de la petite Syrte, golfe de Gabès-Tunisie.

Résultats de la campagne de prospection du mois de juillet 1996. Étude préliminaire :  
biocénoses et thanatocénoses récentes. Mar. Life 3 (1-2), 47-60.

**Table 1.** Economic value of *Posidonia oceanica* meadows.

<b>Year</b>	<b>1990</b>	<b>2014</b>
Market price (€)	1 965	2 960
Quantity (tons)	1 062	2 560
<i>Posidonia oceanica</i> meadows (ha) and {in percent of total land area}	43 563 {33.51}	14 826 {11.41}
Total value of <i>Posidonia oceanica</i> (€)	2 086 640	7 576 473
Yield of ton of benthic species per hectare of <i>Posidonia oceanica</i>	0.024	0.173
Per hectare value of <i>Posidonia</i> (€)	48	511
Economic losses due to the degradation of <i>Posidonia oceanica</i> (€)	4 140 278	58 854 676

**Table 2.** Economic losses linked to the *Posidonia oceanica* degradation versus the added values of the GCT (*Groupe Chimique Tunisien*) factories at the national (Tunisia) and regional (Gabes) scales, during the 1990-2014 period. <sup>(1)</sup> estimates made based on the economic valuation carried out during the present study; <sup>(2)</sup> data collected from the European Community study conducted in 2017 (EC, 2017).

<b>Years / Period</b>		<b>1990</b>	<b>2014</b>	<b>1990-2014</b>
Economic losses due to the <i>Posidonia oceanica</i> degradation (ELDPO, M€) <sup>(1)</sup>		4	59	780
Added values of all GCT factories in Tunisia (AVTF) <sup>(2)</sup>	Added Values (M€)	-24	152	3316
	Ratio ELDPO/AVTF (%)	117	39	24
Added values of Gabes GCT factories (AVGF) <sup>(2)</sup>	Added Values (M€)	-12	92	1824
	Ratio ELDPO/AVGF (%)	132	64	43




## Figures Captions

**Fig. 1.** Location of the Gabes Gulf (a) and the historical decline of *P. oceanica* (right panel: b. 1965 (De Gaillande, 1970b), c. 1981 (Darmoul, 1988), d. 1990 (Zaouali, 1993), e. 2014 (El Zrelli et al., 2017)).

**Fig. 2.** Historical trends of observed (asterisks) and predicted *Posidonia* cover in the Gulf of Gabes, from a linear model of annual decay plus two standard errors (a), and historical variations of “coastal” (b) and “small pelagic” (c) landing volumes and fishing efforts as well as the *Posidonia* area, during 1970-2015.

**Fig. 3.** Hypotheses testing using the Akaike Information Criterion (AIC) for the effect of fishing effort and predicted *Posidonia* cover on the (a & b) fishing yield by two fleets in the Gulf of Gabes and (c & d) coastal and pelagic fishing fleets.

**Fig. 4.** Estimate of the producer surplus loss after the *Posidonia oceanica* seagrass service degradation.



*Posidonia oceanica*

