

Non-contact rheology of finite-size air-water interfaces – Supplementary Material –

Vincent Bertin,^{1,2,*} Zaicheng Zhang,^{1,*} Rodolphe Boisgard,¹ Christine
Grauby-Heywang,¹ Elie Raphaël,² Thomas Salez,^{1,3,†} and Abdelhamid Maali^{1,‡}

¹*Univ. Bordeaux, CNRS, LOMA, UMR 5798, 33405 Talence, France.*

²*UMR CNRS Gulliver 7083, ESPCI Paris, PSL Research University, 75005 Paris, France.*

³*Global Station for Soft Matter, Global Institution for Collaborative Research and Education,
Hokkaido University, Sapporo, Hokkaido 060-0808, Japan.*

(Dated: November 6, 2020)

GENERAL MODEL

The modulus $|Z^*|$ of the amplitude Z^* of the vertical displacement Z of the sphere is assumed to be much smaller than the average sphere-bubble distance D , so that we can neglect the contribution of Z in the non-linear ($\propto h^3$) factor within the Reynolds equation (see Eq. (4) in the main text). The complex version of the latter becomes:

$$12\eta r i \omega \left[Z^* + u^*(r) \right] = \frac{d}{dr} \left[r \left(D + \frac{r^2}{2R_{\text{eff}}} \right)^3 \frac{dp^*(r)}{dr} \right], \quad (\text{S1})$$

with identical notations as in the main text. The excess pressure and deformation fields are set to zero above a given cut-off radius b , *i.e.*:

$$p^*(r) = 0, \quad u^*(r) = 0, \quad r > b. \quad (\text{S2})$$

We then introduce the discrete Hankel transform. The amplitudes of the excess pressure and deformation fields are expanded in terms of Fourier-Bessel series [1]. For $r \in [0, b]$, one thus has:

$$p^*(r) = \sum_{k=1}^{\infty} p_k J_0(\phi_k r), \quad u^*(r) = \sum_{k=1}^{\infty} u_k J_0(\phi_k r), \quad (\text{S3})$$

where J_0 is the Bessel function of the first kind with index 0, and $b\phi_k$ the k th root of J_0 . The series coefficients are related to the amplitude of the deformation field via the inversion relation:

$$u_k = \frac{2}{b^2 J_1^2(\phi_k b)} \int_0^b dr r u^*(r) J_0(\phi_k r), \quad (\text{S4})$$

where J_1 is the Bessel function of the first kind with index 1. We further introduce the dimensionless variables:

$$x = \frac{r}{\sqrt{2R_{\text{eff}}D}}, \quad U^* = u^*/Z^*, \quad P^* = \frac{p^*}{\frac{\eta R_{\text{eff}} Z^* \omega}{D^2}}, \quad B = \frac{b}{\sqrt{2R_{\text{eff}}D}}, \quad \psi_k = \phi_k \sqrt{2R_{\text{eff}}D}, \quad (\text{S5})$$

so that Eq. (S1) and the linearized Young-Laplace equation (Eq. (5) in the main text) respectively become:

$$24ix \left(1 + U^* \right) = \frac{d}{dx} \left[x (1 + x^2)^3 \frac{dP^*}{dx} \right], \quad (\text{S6})$$

and:

$$\frac{1}{8} \left(\frac{D_c}{D} \right) P^* = \frac{1}{x} \frac{d}{dx} \left(x \frac{dU^*}{dx} \right), \quad (\text{S7})$$

where $D_c = 16R_{\text{eff}}^2 \eta \omega / \gamma$ is a typical viscocapillary distance that emerges from the rescaling. By integrating Eq. (S6) with respect to x , and invoking the Fourier-Bessel decomposition, we find:

$$-x(1+x^2)^3 \sum_{k=1}^{\infty} \psi_k P_k J_1(\psi_k x) = 12ix^2 + 24i \sum_{k=1}^{\infty} U_k \frac{J_1(\psi_k x)}{\psi_k} x, \quad (\text{S8})$$

where $U_k = u_k/Z^*$ and $P_k = p_k D^2 / (\eta R_{\text{eff}} Z^* \omega)$. Using Eq. (S7), one can show that:

$$\psi_k^2 U_k = -\frac{1}{8} \left(\frac{D_c}{D} \right) P_k . \quad (\text{S9})$$

Injecting the last relation in Eq. (S8), one gets:

$$-\sum_{l=1}^{\infty} \psi_l P_l J_1(\psi_l x) = \frac{12ix}{(1+x^2)^3} + \frac{3i}{(1+x^2)^3} \left(\frac{D_c}{D} \right) \sum_{l=1}^{\infty} P_l \frac{J_1(\psi_l x)}{\psi_l^3} . \quad (\text{S10})$$

Then, we perform the Fourier-Bessel inversion by applying to Eq. (S10) the operator $\frac{2}{B^2 J_1^2(\psi_k B)} \int_0^B dy y J_1(\psi_k y) (*)$, where $(*)$ stands for a function of y , and we find:

$$P_k = A_k + \sum_{l=1}^{\infty} M_{k,l} P_l , \quad \forall k \in \mathbb{N}^* , \quad (\text{S11})$$

where A_k and $M_{k,l}$ are defined as:

$$A_k = -12i \frac{2}{B^2 J_1^2(\psi_k B)} \int_0^B dy \frac{y^2}{(1+y^2)^3} \frac{J_1(\psi_k y)}{\psi_k} , \quad (\text{S12})$$

and:

$$M_{k,l} = -3i \left(\frac{D_c}{D} \right) \frac{2}{B^2 J_1^2(\psi_k B)} \int_0^B dy \frac{y}{(1+y^2)^3} \frac{J_1(\psi_k y)}{\psi_k} \frac{J_1(\psi_l y)}{\psi_l^3} . \quad (\text{S13})$$

Finally, the amplitude of the hydrodynamic force is $F_h^* = 2\pi \int_0^b dr r p^*(r)$. The mechanical impedance $G^* = -F^*/Z^*$ can thus be expressed as [2]:

$$\begin{aligned} G^* &= -\frac{4\pi\eta\omega R_{\text{eff}}^2}{D} \int_0^B x P^*(x) dx \\ &= -\frac{4\pi\eta\omega R_{\text{eff}}^2}{D} \sum_{k=1}^{\infty} P_k \frac{B J_1(\psi_k B)}{\psi_k} . \end{aligned} \quad (\text{S14})$$

The mechanical impedance can be estimated numerically from this last equation, restricting the summation to *e.g.* $k \leq 1000$. We checked the robustness of the results against a change of that arbitrary upper bound.

LARGE-DISTANCE ASYMPTOTIC MODEL

We assume that the ratio D_c/D is a small parameter, and we expand the amplitude of the dimensionless excess pressure field as:

$$P^*(x) \simeq P_0^*(x) + \frac{D_c}{D} P_1^*(x) . \quad (\text{S15})$$

The leading-order term P_0^* is given by Eq. (S6) in the undeformed-bubble limit, and reads:

$$P_0^*(x) = -\frac{3i}{(1+x^2)^2} . \quad (\text{S16})$$

Similarly, the leading-order amplitude U_1^* of the dimensionless deformation field can be found by inserting P_0^* in Eq. (S7), and by invoking the boundary condition $U_1^*(x=B) = 0$. We find:

$$U_1^*(x) = -\frac{3i}{32} \frac{D_c}{D} \left[\log(1+B^2) - \log(1+x^2) \right] . \quad (\text{S17})$$

Then, introducing U_1^* in Eq. (S6) allows us to express the next-order correction in pressure:

$$P_1^*(x) = \frac{3}{32} \left[-3 \frac{\log(1+B^2)}{(1+x^2)^2} - \pi^2 + \frac{3}{(1+x^2)^2} + \frac{3}{(1+x^2)} + 2 \frac{\log(1+x^2)}{1+x^2} - \log^2(1+x^2) - 2\text{Li}_2(-x^2) \right], \quad (\text{S18})$$

where Li_2 denotes the dilogarithm function [3]. After some algebra, we find that:

$$\mathcal{G}' = \mathcal{R}[\mathcal{G}^*] \simeq \frac{3}{32} \left(\frac{D_c}{D} \right)^2 \left[-1 + \log(1+B^2) \right], \quad (\text{S19})$$

which is the dimensionless equivalent of Eq. (6) in the main text.

* These authors contributed equally.

† Electronic address: thomas.salez@u-bordeaux.fr

‡ Electronic address: abdelhamid.maali@u-bordeaux.fr

[1] G. N. Watson, *A treatise on the theory of Bessel functions* (Cambridge university press, 1995).

[2] S. Leroy and E. Charlaix, *Journal of Fluid Mechanics* **674**, 389 (2011).

[3] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, vol. 55 (US Government printing office, 1948).