



HAL
open science

Observation of Multiplicity Dependent Prompt $\chi_{c1}(3872)$ and $\psi(2S)$ Production in pp Collisions

Roel Aaij, Carlos Abellán Beteta, Thomas Ackernley, Bernardo Adeva, Marco Adinolfi, Hossein Afsharnia, Christine Angela Aidala, Salvatore Aiola, Ziad Ajaltouni, Simon Akar, et al.

► To cite this version:

Roel Aaij, Carlos Abellán Beteta, Thomas Ackernley, Bernardo Adeva, Marco Adinolfi, et al.. Observation of Multiplicity Dependent Prompt $\chi_{c1}(3872)$ and $\psi(2S)$ Production in pp Collisions. Phys.Rev.Lett., 2021, 126 (9), pp.092001. <10.1103/PhysRevLett.126.092001>. <hal-02953164>

HAL Id: hal-02953164

<https://hal.science/hal-02953164v1>

Submitted on 21 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Observation of multiplicity-dependent prompt $\chi_{c1}(3872)$ and $\psi(2S)$ production in pp collisions

LHCb collaboration[†]

Abstract

The production of $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons is studied as a function of charged particle multiplicity in pp collisions at a center-of-mass energy of 8 TeV, corresponding to an integrated luminosity of 2 fb^{-1} . For both states, the fraction that is produced promptly at the collision vertex is found to decrease as charged particle multiplicity increases. The ratio of $\chi_{c1}(3872)$ to $\psi(2S)$ cross-sections for promptly produced particles is also found to decrease with multiplicity, while no significant dependence on multiplicity is observed for the equivalent ratio of particles produced away from the collision vertex in b -hadron decays. This behavior is consistent with a calculation that models the $\chi_{c1}(3872)$ structure as a compact tetraquark. Implications for the binding energy of the $\chi_{c1}(3872)$ state are discussed.

Published in Phys. Rev. Lett. 126 (2021) 092001

© 2021 CERN for the benefit of the LHCb collaboration. CC BY 4.0 licence.

[†]Authors are listed at the end of this Letter.

In recent years, multiple new resonances containing heavy quarks have been observed that do not fit into the framework of conventional hadrons, see Ref. [1] for a recent review. The most studied of these exotic hadrons is the $\chi_{c1}(3872)$ state, also known as $X(3872)$. It was first discovered in the mass spectrum of $J/\psi\pi^+\pi^-$ in B -meson decays by the Belle collaboration [2], and has since been confirmed by multiple other experiments [3–6]. Despite intense scrutiny, the exact nature of the $\chi_{c1}(3872)$ state is still unclear.

Multiple explanations of the $\chi_{c1}(3872)$ structure have been proposed. Shortly after its discovery, it was considered as one of several possible charmonium states [7]. However, LHCb has since measured the quantum numbers to be $J^{PC} = 1^{++}$ [8], which disfavors its assignment as conventional charmonium because no compatible charmonium states with these quantum numbers are expected near the measured mass [9]. Other models consider the $\chi_{c1}(3872)$ state to be a tetraquark, which may have further substructure, composed of a diquark-antidiquark bound state [10–12] or a hadrocharmonium state where two light quarks orbit a charmonium core [13]. Mixtures of various exotic and conventional states have also been studied [14–17]. The remarkable proximity of the $\chi_{c1}(3872)$ mass to the sum of the D^0 and \bar{D}^{*0} meson masses have led to the consideration of its structure as a hadronic molecule, a state comprising these two mesons bound via pion exchange [18, 19]. In this case the binding energy of the $\chi_{c1}(3872)$ hadron would be small, as the mass difference $(M_{D^0} + M_{\bar{D}^{*0}}) - M_{\chi_{c1}(3872)} = 0.07 \pm 0.12 \text{ MeV}/c^2$ is consistent with zero [20]. Consequently, these models assign the $\chi_{c1}(3872)$ state a large radius of $O(10 \text{ fm})$ [17, 21]. Results from recent LHCb studies of the $\chi_{c1}(3872)$ line shape are compatible with the molecular interpretation but do not exclude other possibilities [20, 22].

Techniques developed to study quarkonium production in proton-nucleus (pA) collisions can be used to probe the binding energy of hadrons. Measurements of charmonium production in pA collisions at fixed target experiments [23, 24] and colliders [25–30] showed that $\psi(2S)$ production is suppressed more than J/ψ production in rapidity regions where a relatively large number of charged particles are produced. Similarly, measurements of Υ production at the Large Hadron Collider (LHC) revealed that the $\Upsilon(2S)$ and $\Upsilon(3S)$ states are suppressed more than the $\Upsilon(1S)$ [31, 32]. As the effects governing heavy quark production and transport through the nucleus are assumed to be similar for states with the same quark content, the mechanism for the suppression of excited states is expected to occur in the late stages of the collision, after the heavy quark pair has hadronized into a final state. Models incorporating final-state effects, such as heavy quark pair breakup via interactions with comoving hadrons, describe the relative suppression of excited quarkonium states in pA collisions [33–37]. Similar final-state effects can also disrupt formation of the $\chi_{c1}(3872)$ state via interactions with pions produced in the underlying event [38], and would be especially significant if the $\chi_{c1}(3872)$ structure is a large, weakly bound hadronic molecule.

High-multiplicity pp collisions provide a hadronic environment that approaches heavy ion collisions in many respects. Recently, phenomena typically thought only to occur in collisions of large nuclei have been observed in high-multiplicity pp collisions, including a near-side ridge in two-particle angular correlations [39], strangeness enhancement [40], and collective flow [41]. Multiplicity-dependent modification of Υ production has also been observed [42]. Therefore, high-multiplicity pp collisions provide a testing ground for examining final-state effects observed on quarkonium in pA and AA collisions. Measurements of such effects can provide new constraints on the structure of the $\chi_{c1}(3872)$ [43].

In this Letter, measurements of the fractions of $\chi_{c1}(3872)$ and $\psi(2S)$ states, f_{prompt} ,

that are produced promptly at the pp collision vertex as a function of charged particle multiplicity are presented. The $\chi_{c1}(3872)$ and $\psi(2S)$ states are compared by measuring the ratio of the $\chi_{c1}(3872)$ to $\psi(2S)$ cross-sections, as a function of multiplicity. The $\chi_{c1}(3872)$ and $\psi(2S)$ candidates are reconstructed through their decays to the $J/\psi\pi^+\pi^-$ final state, where the J/ψ meson subsequently decays to $\mu^+\mu^-$ pairs. This study uses data collected with the LHCb detector at a center-of-mass energy $\sqrt{s} = 8$ TeV, corresponding to an integrated luminosity of 2 fb^{-1} .

The LHCb detector [44, 45] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector elements comprise a silicon-strip vertex detector (VELO) surrounding the pp interaction region that allows b hadrons to be identified from their characteristically long flight distance; a tracking system that provides a measurement of the momentum, p , of charged particles; two ring-imaging Cherenkov (RICH) detectors that discriminate between different species of charged hadrons, and a series of tracking detectors interleaved with hadron absorbers for identifying muons. In this analysis, multiplicity is represented by the number of charged particle tracks reconstructed in the VELO, $N_{\text{tracks}}^{\text{VELO}}$. The VELO track-reconstruction efficiency has been measured to be about 99% [46].

Simulation is required to model the effects of the detector acceptance and the imposed selection requirements. In the simulation, pp collisions are generated using PYTHIA [47, 48] with a specific LHCb configuration [49]. Decays of unstable particles are described by EVTGEN [50]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [51] as described in Ref. [52].

Events considered in this analysis are selected by a set of triggers designed to record events containing the decay $J/\psi \rightarrow \mu^+\mu^-$. Tracks from triggered events that are identified as good muon candidates are retained. The muons are required to have momentum $p > 10\text{ GeV}/c$ and a momentum component transverse to the beam direction $p_{\text{T}} > 650\text{ MeV}/c$. Candidate J/ψ mesons are formed from a pair of oppositely charged muons with an invariant mass within $\pm 39\text{ MeV}/c^2$ (corresponding to 3 times the resolution on the mass) of the known J/ψ mass and combined $p_{\text{T}} > 3\text{ GeV}/c$. Charged pion candidates are selected using particle identification information from the RICH detectors. They are required to have $p > 3\text{ GeV}/c$ to ensure that the pions are above threshold in one of the RICH detectors, and have $p_{\text{T}} > 500\text{ MeV}/c$ to reduce combinatorial background.

Selected $\mu^+\mu^-\pi^+\pi^-$ combinations that form a good-quality common vertex are fitted with kinematic constraints that require all tracks to originate from a common vertex and constrain the dimuon mass to the known J/ψ mass [53]. The decay kinematics are required to satisfy $M_{J/\psi\pi^+\pi^-} - M_{J/\psi} - M_{\pi^+\pi^-} < 300\text{ MeV}/c^2$ and the candidates must have $p_{\text{T}} > 5\text{ GeV}/c$ and be within the pseudorapidity range $2 < \eta < 4.5$. The resulting $J/\psi\pi^+\pi^-$ invariant-mass spectrum is shown in Fig. 1.

To avoid multiplicity biases arising from tracks produced in multiple collisions that occur in the same beam crossing, only events with a single reconstructed collision vertex are considered. The position of collision vertices is restricted to a range along the beam direction $-60 < z < 120\text{ mm}$, to avoid biases from missing tracks that fall outside the VELO acceptance.

Both $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons can be produced promptly at the pp collision vertex, either directly or in strong decays of higher charmonia states, or in the decays of b hadrons, which travel several millimeters before decaying. The prompt component of the signal is separated from the component originating from b decays by performing

a simultaneous fit to the $J/\psi\pi^+\pi^-$ invariant-mass spectrum and the pseudo decay-time spectrum. The pseudo decay-time t_z is defined as

$$t_z \equiv \frac{(z_{\text{decay}} - z_{\text{PV}}) \times M}{p_z}, \quad (1)$$

where $z_{\text{decay}} - z_{\text{PV}}$ is the difference between the positions of the reconstructed vertex of the $J/\psi\pi^+\pi^-$ and the collision vertex along the beam axis, M is the known mass [53] of the reconstructed $\psi(2S)$ or $\chi_{c1}(3872)$ candidate, and p_z is the candidate's momentum along the beam axis. The signal in the t_z spectrum is fit with a delta function representing the prompt component and an exponential decay function representing the component from b decays, which are convolved with a double Gaussian resolution function. Two different parameterizations of the t_z background components using mass sidebands above and below the mass peak of interest are employed. The first is an empirically determined analytical function as was done in Ref. [54] and the second directly uses the t_z shape templates taken from the mass sidebands in the data.

In the fit to the invariant-mass spectrum, the $\psi(2S)$ peak is represented by a sum of two Crystal Ball functions, as in a previous LHCb analysis at 7 TeV [55]. The measured $\chi_{c1}(3872)$ peak is well described by a Gaussian function. The background contribution is studied by examining the invariant-mass spectrum constructed by like-sign pion pairs,




Figure 1: The $J/\psi\pi^+\pi^-$ invariant-mass spectrum. The inset shows the region of the $\chi_{c1}(3872)$ resonance.




Figure 2: The $\psi(2S)$ (left) invariant-mass and (right) t_z spectrum in the p_T and multiplicity ranges $p_T > 5 \text{ GeV}/c$ and $60 < N_{\text{tracks}}^{\text{VELO}} < 80$, with the simultaneous fit superimposed.

and is found to be well described by third-order Chebyshev polynomials; this shape is used to represent the background when fitting the $J/\psi\pi^+\pi^-$ mass spectra. The invariant mass and t_z spectra are divided into bins of $N_{\text{tracks}}^{\text{VELO}}$, and the fit is performed in each bin, separately for the $\psi(2S)$ meson and $\chi_{c1}(3872)$ state. An example is shown in Fig. 2.

The total yield and the measured fraction of the inclusive signal that is produced at the collision vertex for each state is determined by a fit. Due to the different production mechanisms, the observed $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons from these sources have different momentum distributions, which may lead to differences in acceptance and reconstruction efficiencies. To account for this effect, the p_T distributions of prompt and displaced signal candidates are extracted from the data using the *sPlot* technique [56], and the p_T distributions of the simulated particles are weighted to match those of the data. Since these measurements are binned in multiplicity and effects of multiplicity-dependent breakup may depend on p_T , the simulation is reweighted to match p_T distributions extracted from low- and high-multiplicity samples. The difference in the acceptance and reconstruction efficiencies found using these different parameterizations of the p_T distributions is taken as a systematic uncertainty. Corrections are applied to account for the relative acceptance of the LHCb spectrometer between particles produced at the primary vertex and in b decays $\varepsilon_{\text{prompt}}^{\text{acc}}/\varepsilon_{\text{b}}^{\text{acc}}$, which is determined via simulation to be 1.00 ± 0.01 for the $\psi(2S)$ state and 1.02 ± 0.01 for the $\chi_{c1}(3872)$ state, and for the relative reconstruction and selection efficiencies $\varepsilon_{\text{prompt}}^{\text{reco}}/\varepsilon_{\text{b}}^{\text{reco}}$, which are 0.99 ± 0.03 for the $\psi(2S)$ state and 1.11 ± 0.04 for the $\chi_{c1}(3872)$ state. The central value of f_{prompt} is taken as the average of the values obtained from the two fitting methods, while the difference is taken as a systematic uncertainty, which ranges from 1% to 2% for the $\psi(2S)$ fits and from 2% to 6% for the $\chi_{c1}(3872)$ fits. This uncertainty is uncorrelated between bins and is comparable to the statistical uncertainty on the $\psi(2S)$ data, while the statistical uncertainty dominates on the $\chi_{c1}(3872)$ data. The uncertainty on the relative efficiency is taken as a systematic uncertainty on f_{prompt} , which is correlated between the data points for each species. The resulting values




Figure 3: The fraction, f_{prompt} , of promptly produced $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons, as a function of the number of tracks reconstructed in the VELO. The vertical error bars (boxes) represent the uncorrelated (correlated) uncertainties, while the horizontal error bars indicate bin widths.

of f_{prompt} as a function of multiplicity are shown in Fig. 3, up to $N_{\text{tracks}}^{\text{VELO}} = 200$. The fraction of events with $N_{\text{tracks}}^{\text{VELO}} > 200$ is negligible and is not included in the analysis. The horizontal position of each point is the average value of $N_{\text{tracks}}^{\text{VELO}}$ for signal events within that bin.

A clear decrease of f_{prompt} is seen as the multiplicity increases, for both the $\psi(2S)$ and $\chi_{c1}(3872)$ hadrons. This could be due to a combination of several effects: the average multiplicity is higher in events containing a $b\bar{b}$ pair due to their fragmentation into hadrons and subsequent decays [57, 58]; or the suppression of prompt $\psi(2S)$ and $\chi_{c1}(3872)$ production via interactions with other particles produced at the vertex, which decreases the prompt production in high-multiplicity events, but does not affect production in b decays.

The prompt and b -decay components are examined directly by calculating the ratio of the $\chi_{c1}(3872)$ and $\psi(2S)$ cross-sections, $\sigma_{\chi}/\sigma_{\psi}$, times their respective branching fractions to the $J/\psi\pi^+\pi^-$ final state, \mathcal{B}_{χ} and \mathcal{B}_{ψ} . This ratio is given by

$$\frac{\sigma_\chi \mathcal{B}_\chi}{\sigma_\psi \mathcal{B}_\psi} = \frac{N_\chi f_{\text{prompt}}^\chi}{N_\psi f_{\text{prompt}}^\psi} \frac{\varepsilon_\psi^{\text{acc}}}{\varepsilon_\chi^{\text{acc}}} \frac{\varepsilon_\psi^{\text{reco}}}{\varepsilon_\chi^{\text{reco}}} \frac{\varepsilon_\psi^{\text{PID}}}{\varepsilon_\chi^{\text{PID}}}. \quad (2)$$

Here, N is the signal yield, f_{prompt} is the prompt fraction and the ε terms represent various efficiency corrections of the corresponding state. The ratio of cross-sections from b decays is found by replacing f_{prompt} with $(1-f_{\text{prompt}})$ in Eq. 2. Correlated systematic uncertainties largely cancel in the ratio, and the result is dominated by uncorrelated uncertainties. The ratio of efficiencies for four charged decay products to fall within the LHCb acceptance, $\varepsilon_\psi^{\text{acc}}/\varepsilon_\chi^{\text{acc}}$, is found via simulation to be consistent with one with an uncertainty of approximately 1% that is determined by varying the p_T distributions of the simulated $\psi(2S)$ and $\chi_{c1}(3872)$ hadrons. Control samples of identified muons and pions obtained from data are used to measure the ratio of muon and pion identification efficiencies, $\varepsilon_\psi^{\text{PID}}/\varepsilon_\chi^{\text{PID}}$, which is near one with an uncertainty of about 1% due to the finite size of the control sample. The only relative efficiency that has a significant deviation from unity is the ratio of reconstruction efficiencies, $\varepsilon_\psi^{\text{reco}}/\varepsilon_\chi^{\text{reco}}$, which is found via simulation to be 0.58 ± 0.02 (0.65 ± 0.04) for particles that are produced promptly (in b -decays). This is due to the different kinematic properties of the pion pair produced in the decays: pions from $\chi_{c1}(3872)$ hadron decays proceed through an intermediate $\rho^0(770)$ resonance [59] and have a higher reconstruction efficiency than pions from the $\psi(2S)$ decay due to their higher p_T . The uncertainty on the ratio of reconstruction efficiencies is taken from the variations observed when weighting the p_T distributions of the simulated $\psi(2S)$ and $\chi_{c1}(3872)$ hadrons to match those in the data in different multiplicity bins, as previously discussed.

The ratio of cross-sections is shown in Fig. 4. A decrease in the prompt production of $\chi_{c1}(3872)$ hadrons relative to prompt $\psi(2S)$ mesons is observed as the charged particle multiplicity increases. To illustrate this effect, a linear fit to this data, which considers only the uncorrelated uncertainties, is performed and returns a negative slope that differs from zero by 5 standard deviations.

After preliminary LHCb results on multiplicity-dependent $\chi_{c1}(3872)$ production were presented [60], calculations of these observables based on the Comover Interaction Model [34, 35] were performed [43]. In this model, promptly produced $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons interact with other produced particles, with a breakup cross-section σ_{br} that is determined by their radius and binding energy. The model assumes no interactions at low multiplicity, and the calculations are normalized to the data in the lowest multiplicity bin. A purely molecular $\chi_{c1}(3872)$ has a large radius and correspondingly high σ_{br} , and is quickly dissociated as multiplicity increases. If coalescence provides an additional formation mechanism for molecular $\chi_{c1}(3872)$, the ratio $\sigma_{\chi_{c1}(3872)}/\sigma_{\psi(2S)}$ rises with multiplicity. Neither of these calculations are consistent with the data. A compact tetraquark $\chi_{c1}(3872)$ has a slightly larger radius and σ_{br} than the $\psi(2S)$, and in this scenario $\sigma_{\chi_{c1}(3872)}/\sigma_{\psi(2S)}$ gradually decreases with multiplicity, matching the measured trend.

In contrast to the prompt data, the ratio of cross-sections for production in b decays shows a slight increase, which is not statistically significant. A linear fit to these data points, again without considering the correlated systematic uncertainty, gives a positive slope that is consistent with zero within 1.6 standard deviations. Since these hadrons originate from displaced decay vertices of b hadrons, they are not subject to suppression via interactions with other particles produced at the primary vertex. Consequently,




Figure 4: The ratio of the $\chi_{c1}(3872)$ and $\psi(2S)$ cross-sections measured in the $J/\psi\pi^+\pi^-$ channel as a function of the number of tracks reconstructed in the VELO. The point-to-point uncorrelated (correlated) uncertainties are shown as vertical error bars (boxes) and the bin widths are shown as horizontal error bars. See text for details on calculations from Ref. [43]

this ratio is set only by the branching fractions of b decays to $\chi_{c1}(3872)$ and $\psi(2S)$ hadrons. The multiplicity dependence of b hadron production has not been studied in detail and modification of the b hadron admixture could affect $\chi_{c1}(3872)$ production, as different b hadron species may have different decay probabilities to $\chi_{c1}(3872)$ states [61,62]. However, the uncertainties preclude drawing any firm conclusions on multiplicity-dependent modifications of b hadronization from this data.

In conclusion, the prompt $\chi_{c1}(3872)$ and prompt $\psi(2S)$ production cross-sections decrease relative to their production via b decays as the charged particle multiplicity increases in pp collisions at 8 TeV. A comparison between the $\chi_{c1}(3872)$ and $\psi(2S)$ states shows that, in contrast to production from b -decays, which display no significant dependence on multiplicity, prompt production of $\chi_{c1}(3872)$ is suppressed relative to prompt $\psi(2S)$ production as multiplicity increases. This observation is an important ingredient to obtain a full understanding of the nature of the $\chi_{c1}(3872)$ state.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW

and NCN (Poland); MEN/IFA (Romania); MSHE (Russia); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); A*MIDEX, ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, Thousand Talents Program, and Sci. & Tech. Program of Guangzhou (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom).

References

- [1] S. L. Olsen, T. Skwarnicki, and D. Zieminska, *Nonstandard heavy mesons and baryons: Experimental evidence*, Rev. Mod. Phys. **90** (2018) 015003, [arXiv:1708.04012](#).
- [2] Belle collaboration, S. K. Choi *et al.*, *Observation of a narrow charmonium - like state in exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ decays*, Phys. Rev. Lett. **91** (2003) 262001, [arXiv:hep-ex/0309032](#).
- [3] CDF collaboration, D. Acosta *et al.*, *Observation of the narrow state $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV*, Phys. Rev. Lett. **93** (2004) 072001, [arXiv:hep-ex/0312021](#).
- [4] D0 collaboration, V. M. Abazov *et al.*, *Observation and properties of the $X(3872)$ decaying to $J/\psi \pi^+ \pi^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, Phys. Rev. Lett. **93** (2004) 162002, [arXiv:hep-ex/0405004](#).
- [5] BaBar collaboration, B. Aubert *et al.*, *Study of the $B \rightarrow J/\psi K^- \pi^+ \pi^-$ decay and measurement of the $B \rightarrow X(3872) K^-$ branching fraction*, Phys. Rev. **D71** (2005) 071103, [arXiv:hep-ex/0406022](#).
- [6] LHCb collaboration, R. Aaij *et al.*, *Observation of $X(3872)$ production in pp collisions at $\sqrt{s} = 7$ TeV*, Eur. Phys. J. **C72** (2012) 1972, [arXiv:1112.5310](#).
- [7] E. J. Eichten, K. Lane, and C. Quigg, *Charmonium levels near threshold and the narrow state $X(3872) \rightarrow \pi^+ \pi^- J/\psi$* , Phys. Rev. **D69** (2004) 094019, [arXiv:hep-ph/0401210](#).
- [8] LHCb collaboration, R. Aaij *et al.*, *Determination of the $X(3872)$ meson quantum numbers*, Phys. Rev. Lett. **110** (2013) 222001, [arXiv:1302.6269](#).
- [9] T. Barnes, S. Godfrey, and E. S. Swanson, *Higher charmonia*, Phys. Rev. **D72** (2005) 054026, [arXiv:hep-ph/0505002](#).

- [10] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, *Diquark-antidiquarks with hidden or open charm and the nature of $X(3872)$* , Phys. Rev. **D71** (2005) 014028, [arXiv:hep-ph/0412098](#).
- [11] R. D. Matheus, S. Narison, M. Nielsen, and J. M. Richard, *Can the $X(3872)$ be a 1^{++} four-quark state?*, Phys. Rev. **D75** (2007) 014005, [arXiv:hep-ph/0608297](#).
- [12] S. Dubnicka, A. Z. Dubnickova, M. A. Ivanov, and J. G. Korner, *Quark model description of the tetraquark state $X(3872)$ in a relativistic constituent quark model with infrared confinement*, Phys. Rev. **D81** (2010) 114007, [arXiv:1004.1291](#).
- [13] P. C. Wallbott, G. Eichmann, and C. S. Fischer, *$X(3872)$ as a four-quark state in a Dyson-Schwinger/Bethe-Salpeter approach*, Phys. Rev. **D100** (2019) 014033, [arXiv:1905.02615](#).
- [14] P. Colangelo, F. De Fazio, and S. Nicotri, *$X(3872) \rightarrow D\bar{D}\gamma$ decays and the structure of $X(3872)$* , Phys. Lett. B **650** (2007) 166, [arXiv:hep-ph/0701052](#).
- [15] R. D. Matheus, F. S. Navarra, M. Nielsen, and C. M. Zanetti, *QCD Sum Rules for the $X(3872)$ as a mixed molecule-charmonium state*, Phys. Rev. **D80** (2009) 056002, [arXiv:0907.2683](#).
- [16] M. Butenschoen, Z.-G. He, and B. A. Kniehl, *Deciphering the $X(3872)$ via its polarization in prompt production at the CERN LHC*, Phys. Rev. Lett. **123** (2019) 032001, [arXiv:1906.08553](#).
- [17] S. Coito, G. Rupp, and E. van Beveren, *$X(3872)$ is not a true molecule*, Eur. Phys. J. **C73** (2013) 2351, [arXiv:1212.0648](#).
- [18] N. A. Tornqvist, *Isospin breaking of the narrow charmonium state of Belle at 3872-MeV as a deuson*, Phys. Lett. **B590** (2004) 209, [arXiv:hep-ph/0402237](#).
- [19] E. Braaten and M. Kusunoki, *Exclusive production of the $X(3872)$ in B meson decay*, Phys. Rev. **D71** (2005) 074005, [arXiv:hep-ph/0412268](#).
- [20] LHCb collaboration, R. Aaij *et al.*, *Study of the $\psi_2(3823)$ and $\chi_c(3872)$ states in $B^+ \rightarrow (J/\psi\pi^+\pi^-)K^+$ decays*, JHEP **08** (2020) 123, [arXiv:2005.13422](#).
- [21] M. Cardoso, G. Rupp, and E. van Beveren, *Unquenched quark-model calculation of $X(3872)$ electromagnetic decays*, Eur. Phys. J. **C75** (2015) 26, [arXiv:1411.1654](#).
- [22] LHCb collaboration, R. Aaij *et al.*, *Study of the line shape of the $\chi_c(3872)$ meson*, [arXiv:2005.13419](#), submitted to Phys. Rev. D.
- [23] NuSea collaboration, M. J. Leitch *et al.*, *Measurement of differences between J/ψ and ψ' suppression in p-A collisions at 800-GeV/c*, Phys. Rev. Lett. **84** (2000) 3256, [arXiv:nucl-ex/9909007](#).
- [24] NA50 collaboration, B. Alessandro *et al.*, *J/psi and psi-prime production and their normal nuclear absorption in proton-nucleus collisions at 400-GeV*, Eur. Phys. J. **C48** (2006) 329, [arXiv:nucl-ex/0612012](#).

- [25] PHENIX collaboration, A. Adare *et al.*, *Nuclear Modification of ψ' , χ_c , and J/ψ Production in $d+Au$ Collisions at $\sqrt{s_{NN}} = 200$ GeV*, Phys. Rev. Lett. **111** (2013) 202301, arXiv:1305.5516.
- [26] ALICE collaboration, B. B. Abelev *et al.*, *Suppression of $\psi(2S)$ production in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV*, JHEP **12** (2014) 073, arXiv:1405.3796.
- [27] LHCb collaboration, R. Aaij *et al.*, *Study of $\psi(2S)$ production cross-sections and cold nuclear matter effects in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV*, JHEP **03** (2016) 133, arXiv:1601.07878.
- [28] PHENIX collaboration, A. Adare *et al.*, *Measurement of the relative yields of $\psi(2S)$ to $\psi(1S)$ mesons produced at forward and backward rapidity in $p+p$, $p+Al$, $p+Au$, and ^3He+Au collisions at $\sqrt{s_{NN}} = 200$ GeV*, Phys. Rev. **C95** (2017) 034904, arXiv:1609.06550.
- [29] CMS collaboration, A. M. Sirunyan *et al.*, *Measurement of prompt $\psi(2S)$ production cross sections in proton-lead and proton-proton collisions at $\sqrt{s_{NN}} = 5.02$ TeV*, Phys. Lett. **B790** (2019) 509, arXiv:1805.02248.
- [30] ALICE collaboration, S. Acharya *et al.*, *Measurement of nuclear effects on $\psi(2S)$ production in p -Pb collisions at $\sqrt{s_{NN}} = 8.16$ TeV*, JHEP **07** (2020) 237, arXiv:2003.06053.
- [31] CMS collaboration, S. Chatrchyan *et al.*, *Event Activity Dependence of $Y(nS)$ Production in $\sqrt{s_{NN}}=5.02$ TeV pPb and $\sqrt{s}=2.76$ TeV pp Collisions*, JHEP **04** (2014) 103, arXiv:1312.6300.
- [32] LHCb collaboration, R. Aaij *et al.*, *Study of Υ production in pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV*, JHEP **11** (2018) 194, arXiv:1810.07655.
- [33] A. Capella, A. Kaidalov, A. Kouider Akil, and C. Gerschel, *J/ψ and ψ' suppression in heavy ion collisions*, Phys. Lett. **B393** (1997) 431, arXiv:hep-ph/9607265.
- [34] A. Capella, E. G. Ferreira, and A. B. Kaidalov, *Nonsaturation of the J/ψ suppression at large transverse energy in the comovers approach*, Phys. Rev. Lett. **85** (2000) 2080, arXiv:hep-ph/0002300.
- [35] E. G. Ferreira, *Excited charmonium suppression in proton-nucleus collisions as a consequence of comovers*, Phys. Lett. **B749** (2015) 98, arXiv:1411.0549.
- [36] Y.-Q. Ma, R. Venugopalan, K. Watanabe, and H.-F. Zhang, *$\psi(2S)$ versus J/ψ suppression in proton-nucleus collisions from factorization violating soft color exchanges*, Phys. Rev. **C97** (2018) 014909, arXiv:1707.07266.
- [37] E. G. Ferreira and J.-P. Lansberg, *Is bottomonium suppression in proton-nucleus and nucleus-nucleus collisions at LHC energies due to the same effects?*, JHEP **10** (2018) 094, Erratum ibid. **03** (2019) 063, arXiv:1804.04474.
- [38] E. Braaten, H.-W. Hammer, and T. Mehen, *Scattering of an Ultrasoft Pion and the $X(3872)$* , Phys. Rev. **D82** (2010) 034018, arXiv:1005.1688.

- [39] CMS collaboration, V. Khachatryan *et al.*, *Observation of Long-Range Near-Side Angular Correlations in Proton-Proton Collisions at the LHC*, JHEP **09** (2010) 091, arXiv:1009.4122.
- [40] ALICE collaboration, J. Adam *et al.*, *Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions*, Nature Phys. **13** (2017) 535, arXiv:1606.07424.
- [41] CMS collaboration, V. Khachatryan *et al.*, *Evidence for collectivity in pp collisions at the LHC*, Phys. Lett. **B765** (2017) 193, arXiv:1606.06198.
- [42] CMS collaboration, A. M. Sirunyan *et al.*, *Investigation into the event-activity dependence of $\Upsilon(nS)$ relative production in proton-proton collisions at $\sqrt{s} = 7$ TeV*, arXiv:2007.04277.
- [43] A. Esposito *et al.*, *The nature of $X(3872)$ from high-multiplicity pp collisions*, arXiv:2006.15044.
- [44] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [45] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, arXiv:1412.6352.
- [46] R. Aaij *et al.*, *Performance of the LHCb Vertex Locator*, JINST **9** (2014) P09007, arXiv:1405.7808.
- [47] T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, arXiv:0710.3820.
- [48] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, arXiv:hep-ph/0603175.
- [49] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [50] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [51] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [52] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [53] Particle Data Group, M. Tanabashi *et al.*, *Review of particle physics*, Phys. Rev. **D98** (2018) 030001.
- [54] LHCb collaboration, R. Aaij *et al.*, *Measurement of $\psi(2S)$ production cross-sections in proton-proton collisions at $\sqrt{s} = 7$ and 13 TeV*, Eur. Phys. J. **C80** (2020) 185, arXiv:1908.03099.

- [55] LHCb collaboration, R. Aaij *et al.*, *Measurement of $\psi(2S)$ meson production in pp collisions at $\sqrt{s} = 7$ TeV*, Eur. Phys. J. **C72** (2012) 2100, Erratum *ibid.* **C80** (2020) 49, [arXiv:1204.1258](#).
- [56] M. Pivk and F. R. Le Diberder, *sPlot: A statistical tool to unfold data distributions*, Nucl. Instrum. Meth. **A555** (2005) 356, [arXiv:physics/0402083](#).
- [57] P. C. Rowson *et al.*, *Charged Multiplicity of Hadronic Events Containing Heavy Quark Jets*, Phys. Rev. Lett. **54** (1985) 2580.
- [58] OPAL collaboration, G. Abbiendi *et al.*, *Charged particle multiplicities in heavy and light quark initiated events above the Z0 peak*, Phys. Lett. **B550** (2002) 33, [arXiv:hep-ex/0211007](#).
- [59] CDF collaboration, A. Abulencia *et al.*, *Measurement of the dipion mass spectrum in $X(3872) \rightarrow J/\psi\pi^+\pi^-$ decays.*, Phys. Rev. Lett. **96** (2006) 102002, [arXiv:hep-ex/0512074](#).
- [60] J. Matthew Durham for the LHCb collaboration, *LHCb measurements of the exotic tetraquark candidate $\chi_{c1}(3872)$ in high-multiplicity pp and pPb collisions*, in *28th International Conference on Ultrarelativistic Nucleus-Nucleus Collisions*, 2020, [arXiv:2002.01551](#).
- [61] ATLAS collaboration, M. Aaboud *et al.*, *Measurements of $\psi(2S)$ and $X(3872) \rightarrow J/\psi\pi^+\pi^-$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, JHEP **01** (2017) 117, [arXiv:1610.09303](#).
- [62] CMS collaboration, A. M. Sirunyan *et al.*, *Observation of the $B_s^0 \rightarrow X(3872)\phi$ decay*, [arXiv:2005.04764](#).

LHCb collaboration

R. Aaij³¹, C. Abellán Beteta⁴⁹, T. Ackernley⁵⁹, B. Adeva⁴⁵, M. Adinolfi⁵³, H. Afsharnia⁹, C.A. Aidala⁸⁴, S. Aiola²⁵, Z. Ajaltouni⁹, S. Akar⁶⁴, J. Albrecht¹⁴, F. Alessio⁴⁷, M. Alexander⁵⁸, A. Alfonso Alberio⁴⁴, Z. Aliouche⁶¹, G. Alkhazov³⁷, P. Alvarez Cartelle⁴⁷, S. Amato², Y. Amhis¹¹, L. An²¹, L. Anderlini²¹, A. Andreianov³⁷, M. Andreotti²⁰, F. Archilli¹⁶, A. Artamonov⁴³, M. Artuso⁶⁷, K. Arzymatov⁴¹, E. Aslanides¹⁰, M. Atzeni⁴⁹, B. Audurier¹¹, S. Bachmann¹⁶, M. Bachmayer⁴⁸, J.J. Back⁵⁵, S. Baker⁶⁰, P. Baladron Rodriguez⁴⁵, V. Balagura¹¹, W. Baldini²⁰, J. Baptista Leite¹, R.J. Barlow⁶¹, S. Barsuk¹¹, W. Barter⁶⁰, M. Bartolini^{23,h}, F. Baryshnikov⁸⁰, J.M. Basels¹³, G. Bassi²⁸, B. Batsukh⁶⁷, A. Battig¹⁴, A. Bay⁴⁸, M. Becker¹⁴, F. Bedeschi²⁸, I. Bediaga¹, A. Beiter⁶⁷, V. Belavin⁴¹, S. Belin²⁶, V. Bellee⁴⁸, K. Belous⁴³, I. Belov³⁹, I. Belyaev³⁸, G. Bencivenni²², E. Ben-Haim¹², A. Berezhnoy³⁹, R. Bernet⁴⁹, D. Berninghoff¹⁶, H.C. Bernstein⁶⁷, C. Bertella⁴⁷, E. Bertholet¹², A. Bertolin²⁷, C. Betancourt⁴⁹, F. Betti^{19,e}, M.O. Bettler⁵⁴, Ia. Bezshyiko⁴⁹, S. Bhasin⁵³, J. Bhom³³, L. Bian⁷², M.S. Bieker¹⁴, S. Bifani⁵², P. Billoir¹², M. Birch⁶⁰, F.C.R. Bishop⁵⁴, A. Bizzeti^{21,r}, M. Bjørn⁶², M.P. Blago⁴⁷, T. Blake⁵⁵, F. Blanc⁴⁸, S. Blusk⁶⁷, D. Bobulska⁵⁸, J.A. Boelhauve¹⁴, O. Boente Garcia⁴⁵, T. Boettcher⁶³, A. Boldyrev⁸¹, A. Bondar^{42,u}, N. Bondar³⁷, S. Borghi⁶¹, M. Borisyak⁴¹, M. Borsato¹⁶, J.T. Borsuk³³, S.A. Bouchiba⁴⁸, T.J.V. Bowcock⁵⁹, A. Boyer⁴⁷, C. Bozzi²⁰, M.J. Bradley⁶⁰, S. Braun⁶⁵, A. Brea Rodriguez⁴⁵, M. Brodski⁴⁷, J. Brodzicka³³, A. Brossa Gonzalo⁵⁵, D. Brundu²⁶, A. Buonaura⁴⁹, C. Burr⁴⁷, A. Bursche²⁶, A. Butkevich⁴⁰, J.S. Butter³¹, J. Buytaert⁴⁷, W. Byczynski⁴⁷, S. Cadeddu²⁶, H. Cai⁷², R. Calabrese^{20,g}, L. Calefice¹⁴, L. Calero Diaz²², S. Cali²², R. Calladine⁵², M. Calvi^{24,i}, M. Calvo Gomez⁸³, P. Camargo Magalhaes⁵³, A. Camboni⁴⁴, P. Campana²², D.H. Campora Perez⁴⁷, A.F. Campoverde Quezada⁵, S. Capelli^{24,i}, L. Capriotti^{19,e}, A. Carbone^{19,e}, G. Carboni²⁹, R. Cardinale^{23,h}, A. Cardini²⁶, I. Carli⁶, P. Carniti^{24,i}, K. Carvalho Akiba³¹, A. Casais Vidal⁴⁵, G. Casse⁵⁹, M. Cattaneo⁴⁷, G. Cavallero⁴⁷, S. Celani⁴⁸, J. Cerasoli¹⁰, A.J. Chadwick⁵⁹, M.G. Chapman⁵³, M. Charles¹², Ph. Charpentier⁴⁷, G. Chatzikonstantinidis⁵², C.A. Chavez Barajas⁵⁹, M. Chefdeville⁸, C. Chen³, S. Chen²⁶, A. Chernov³³, S.-G. Chitic⁴⁷, V. Chobanova⁴⁵, S. Cholak⁴⁸, M. Chrzaszcz³³, A. Chubykin³⁷, V. Chulikov³⁷, P. Ciambrone²², M.F. Cicala⁵⁵, X. Cid Vidal⁴⁵, G. Ciezarek⁴⁷, P.E.L. Clarke⁵⁷, M. Clemencic⁴⁷, H.V. Cliff⁵⁴, J. Closier⁴⁷, J.L. Cobbledick⁶¹, V. Coco⁴⁷, J.A.B. Coelho¹¹, J. Cogan¹⁰, E. Cogneras⁹, L. Cojocariu³⁶, P. Collins⁴⁷, T. Colombo⁴⁷, L. Congedo¹⁸, A. Contu²⁶, N. Cooke⁵², G. Coombs⁵⁸, G. Corti⁴⁷, C.M. Costa Sobral⁵⁵, B. Couturier⁴⁷, D.C. Craik⁶³, J. Crkovská⁶⁶, M. Cruz Torres¹, R. Currie⁵⁷, C.L. Da Silva⁶⁶, E. Dall'Occo¹⁴, J. Dalseno⁴⁵, C. D'Ambrosio⁴⁷, A. Danilina³⁸, P. d'Argent⁴⁷, A. Davis⁶¹, O. De Aguiar Francisco⁶¹, K. De Bruyn⁷⁷, S. De Capua⁶¹, M. De Cian⁴⁸, J.M. De Miranda¹, L. De Paula², M. De Serio^{18,d}, D. De Simone⁴⁹, P. De Simone²², J.A. de Vries⁷⁸, C.T. Dean⁶⁶, W. Dean⁸⁴, D. Decamp⁸, L. Del Buono¹², B. Delaney⁵⁴, H.-P. Dembinski¹⁴, A. Dendek³⁴, V. Denysenko⁴⁹, D. Derkach⁸¹, O. Deschamps⁹, F. Desse¹¹, F. Dettori^{26,f}, B. Dey⁷², P. Di Nezza²², S. Didenko⁸⁰, L. Dieste Maronas⁴⁵, H. Dijkstra⁴⁷, V. Dobishuk⁵¹, A.M. Donohoe¹⁷, F. Dordei²⁶, A.C. dos Reis¹, L. Douglas⁵⁸, A. Dovbnya⁵⁰, A.G. Downes⁸, K. Dreimanis⁵⁹, M.W. Dudek³³, L. Dufour⁴⁷, V. Duk⁷⁶, P. Durante⁴⁷, J.M. Durham⁶⁶, D. Dutta⁶¹, M. Dziewiecki¹⁶, A. Dziurda³³, A. Dzyuba³⁷, S. Easo⁵⁶, U. Egede⁶⁸, V. Egorychev³⁸, S. Eidelman^{42,u}, S. Eisenhardt⁵⁷, S. Ek-In⁴⁸, L. Eklund⁵⁸, S. Ely⁶⁷, A. Ene³⁶, E. Eppe⁶⁶, S. Escher¹³, J. Eschle⁴⁹, S. Esen³¹, T. Evans⁴⁷, A. Falabella¹⁹, J. Fan³, Y. Fan⁵, B. Fang⁷², N. Farley⁵², S. Farry⁵⁹, D. Fazzini^{24,i}, P. Fedin³⁸, M. Féo⁴⁷, P. Fernandez Declara⁴⁷, A. Fernandez Prieto⁴⁵, F. Ferrari^{19,e}, L. Ferreira Lopes⁴⁸, F. Ferreira Rodrigues², S. Ferreres Sole³¹, M. Ferrillo⁴⁹, M. Ferro-Luzzi⁴⁷, S. Filippov⁴⁰, R.A. Fini¹⁸, M. Fiorini^{20,g}, M. Firlej³⁴, K.M. Fischer⁶², C. Fitzpatrick⁶¹, T. Fiutowski³⁴, F. Fleuret^{11,b}, M. Fontana⁴⁷, F. Fontanelli^{23,h}, R. Forty⁴⁷, V. Franco Lima⁵⁹, M. Franco Sevilla⁶⁵, M. Frank⁴⁷, E. Franzoso²⁰,

G. Frau¹⁶, C. Frei⁴⁷, D.A. Friday⁵⁸, J. Fu²⁵, Q. Fuehring¹⁴, W. Funk⁴⁷, E. Gabriel³¹,
 T. Gaintseva⁴¹, A. Gallas Torreira⁴⁵, D. Galli^{19,e}, S. Gambetta⁵⁷, Y. Gan³, M. Gandelman²,
 P. Gandini²⁵, Y. Gao⁴, M. Garau²⁶, L.M. Garcia Martin⁵⁵, P. Garcia Moreno⁴⁴,
 J. García Pardiñas⁴⁹, B. Garcia Plana⁴⁵, F.A. Garcia Rosales¹¹, L. Garrido⁴⁴, D. Gascon⁴⁴,
 C. Gaspar⁴⁷, R.E. Geertsema³¹, D. Gerick¹⁶, L.L. Gerken¹⁴, E. Gersabeck⁶¹, M. Gersabeck⁶¹,
 T. Gershon⁵⁵, D. Gerstel¹⁰, Ph. Ghez⁸, V. Gibson⁵⁴, M. Giovannetti^{22,j}, A. Gioventù⁴⁵,
 P. Gironella Gironell⁴⁴, L. Giubega³⁶, C. Giugliano^{20,g}, K. Gizdov⁵⁷, E.L. Gkougkousis⁴⁷,
 V.V. Gligorov¹², C. Göbel⁶⁹, E. Golobardes⁸³, D. Golubkov³⁸, A. Golutvin^{60,80}, A. Gomes^{1,a},
 S. Gomez Fernandez⁴⁴, M. Goncerz³³, G. Gong³, P. Gorbounov³⁸, I.V. Gorelov³⁹, C. Gotti²⁴,
 E. Govorkova³¹, J.P. Grabowski¹⁶, R. Graciani Diaz⁴⁴, T. Grammatico¹²,
 L.A. Granado Cardoso⁴⁷, E. Graugés⁴⁴, E. Graverini⁴⁸, G. Graziani²¹, A. Grecu³⁶,
 L.M. Greeven³¹, P. Griffith²⁰, L. Grillo⁶¹, S. Gromov⁸⁰, L. Gruber⁴⁷, B.R. Gruber Cazon⁶²,
 C. Gu³, M. Guarise²⁰, P. A. Günther¹⁶, E. Gushchin⁴⁰, A. Guth¹³, Y. Guz^{43,47}, T. Gys⁴⁷,
 T. Hadavizadeh⁶⁸, G. Haefeli⁴⁸, C. Haen⁴⁷, J. Haimberger⁴⁷, S.C. Haines⁵⁴,
 T. Halewood-leagas⁵⁹, P.M. Hamilton⁶⁵, Q. Han⁷, X. Han¹⁶, T.H. Hancock⁶²,
 S. Hansmann-Menzemer¹⁶, N. Harnew⁶², T. Harrison⁵⁹, C. Hasse⁴⁷, M. Hatch⁴⁷, J. He⁵,
 M. Hecker⁶⁰, K. Heijhoff³¹, K. Heinicke¹⁴, A.M. Hennequin⁴⁷, K. Hennessy⁵⁹, L. Henry^{25,46},
 J. Heuel¹³, A. Hicheur², D. Hill⁶², M. Hilton⁶¹, S.E. Hollitt¹⁴, P.H. Hopchev⁴⁸, J. Hu¹⁶, J. Hu⁷¹,
 W. Hu⁷, W. Huang⁵, X. Huang⁷², W. Hulsbergen³¹, R.J. Hunter⁵⁵, M. Hushchyn⁸¹,
 D. Hutchcroft⁵⁹, D. Hynds³¹, P. Ibis¹⁴, M. Idzik³⁴, D. Ilin³⁷, P. Ilten⁵², A. Inglessi³⁷,
 A. Ishteev⁸⁰, K. Ivshin³⁷, R. Jacobsson⁴⁷, S. Jakobsen⁴⁷, E. Jans³¹, B.K. Jashal⁴⁶,
 A. Jawahery⁶⁵, V. Jevtic¹⁴, M. Jezabek³³, F. Jiang³, M. John⁶², D. Johnson⁴⁷, C.R. Jones⁵⁴,
 T.P. Jones⁵⁵, B. Jost⁴⁷, N. Jurik⁴⁷, S. Kandybei⁵⁰, Y. Kang³, M. Karacson⁴⁷, J.M. Kariuki⁵³,
 N. Kazeev⁸¹, M. Kecke¹⁶, F. Keizer^{54,47}, M. Kenzie⁵⁵, T. Ketel³², B. Khanji⁴⁷, A. Kharisova⁸²,
 S. Kholodenko⁴³, K.E. Kim⁶⁷, T. Kirn¹³, V.S. Kirsebom⁴⁸, O. Kitouni⁶³, S. Klaver³¹,
 K. Klimaszewski³⁵, S. Koliiev⁵¹, A. Kondybayeva⁸⁰, A. Konoplyannikov³⁸, P. Kopciwicz³⁴,
 R. Kopecna¹⁶, P. Koppenburg³¹, M. Korolev³⁹, I. Kostiuk^{31,51}, O. Kot⁵¹, S. Kotriakhova^{37,30},
 P. Kravchenko³⁷, L. Kravchuk⁴⁰, R.D. Krawczyk⁴⁷, M. Kreps⁵⁵, F. Kress⁶⁰, S. Kretzschmar¹³,
 P. Krokovny^{42,u}, W. Krupa³⁴, W. Krzemien³⁵, W. Kucewicz^{85,33,k}, M. Kucharczyk³³,
 V. Kudryavtsev^{42,u}, H.S. Kuindersma³¹, G.J. Kunde⁶⁶, T. Kvaratskheliya³⁸, D. Lacarrere⁴⁷,
 G. Lafferty⁶¹, A. Lai²⁶, A. Lampis²⁶, D. Lancierini⁴⁹, J.J. Lane⁶¹, R. Lane⁵³, G. Lanfranchi²²,
 C. Langenbruch¹³, J. Langer¹⁴, O. Lantwin^{49,80}, T. Latham⁵⁵, F. Lazzari^{28,s}, R. Le Gac¹⁰,
 S.H. Lee⁸⁴, R. Lefèvre⁹, A. Leflat³⁹, S. Legotin⁸⁰, O. Leroy¹⁰, T. Lesiak³³, B. Leverington¹⁶,
 H. Li⁷¹, L. Li⁶², P. Li¹⁶, X. Li⁶⁶, Y. Li⁶, Y. Li⁶, Z. Li⁶⁷, X. Liang⁶⁷, T. Lin⁶⁰, R. Lindner⁴⁷,
 V. Lisovskyi¹⁴, R. Litvinov²⁶, G. Liu⁷¹, H. Liu⁵, S. Liu⁶, X. Liu³, A. Loi²⁶, J. Lomba Castro⁴⁵,
 I. Longstaff⁵⁸, J.H. Lopes², G. Loustau⁴⁹, G.H. Lovell⁵⁴, Y. Lu⁶, D. Lucchesi^{27,l}, S. Luchuk⁴⁰,
 M. Lucio Martinez³¹, V. Lukashenko³¹, Y. Luo³, A. Lupato⁶¹, E. Luppi^{20,g}, O. Lupton⁵⁵,
 A. Lusiani^{28,q}, X. Lyu⁵, L. Ma⁶, S. Maccolini^{19,e}, F. Machefert¹¹, F. Maciuc³⁶, V. Macko⁴⁸,
 P. Mackowiak¹⁴, S. Maddrell-Mander⁵³, O. Madejczyk³⁴, L.R. Madhan Mohan⁵³, O. Maev³⁷,
 A. Maevskiy⁸¹, D. Maisuzenko³⁷, M.W. Majewski³⁴, S. Malde⁶², B. Malecki⁴⁷, A. Malinin⁷⁹,
 T. Maltsev^{42,u}, H. Malygina¹⁶, G. Manca^{26,f}, G. Mancinelli¹⁰, R. Manera Escalero⁴⁴,
 D. Manuzzi^{19,e}, D. Marangotto^{25,n}, J. Maratas^{9,t}, J.F. Marchand⁸, U. Marconi¹⁹,
 S. Mariani^{21,47,21}, C. Marin Benito¹¹, M. Marinangeli⁴⁸, P. Marino⁴⁸, J. Marks¹⁶,
 P.J. Marshall⁵⁹, G. Martellotti³⁰, L. Martinazzoli⁴⁷, M. Martinelli^{24,i}, D. Martinez Santos⁴⁵,
 F. Martinez Vidal⁴⁶, A. Massafferri¹, M. Materok¹³, R. Matev⁴⁷, A. Mathad⁴⁹, Z. Mathe⁴⁷,
 V. Matiunin³⁸, C. Matteuzzi²⁴, K.R. Mattioli⁸⁴, A. Mauri³¹, E. Maurice^{11,b}, J. Mauricio⁴⁴,
 M. Mazurek³⁵, M. McCann⁶⁰, L. McConnell¹⁷, T.H. Mcgrath⁶¹, A. McNab⁶¹, R. McNulty¹⁷,
 J.V. Mead⁵⁹, B. Meadows⁶⁴, C. Meaux¹⁰, G. Meier¹⁴, N. Meinert⁷⁵, D. Melnychuk³⁵,
 S. Meloni^{24,i}, M. Merk^{31,78}, A. Merli²⁵, L. Meyer Garcia², M. Mikhasenko⁴⁷, D.A. Milanese⁷³,
 E. Millard⁵⁵, M.-N. Minard⁸, L. Minzoni^{20,g}, S.E. Mitchell⁵⁷, B. Mitreska⁶¹, D.S. Mitzel⁴⁷,

A. Mödden¹⁴, R.A. Mohammed⁶², R.D. Moise⁶⁰, T. Mombächer¹⁴, I.A. Monroy⁷³, S. Monteil⁹,
 M. Morandin²⁷, G. Morello²², M.J. Morello^{28,q}, J. Moron³⁴, A.B. Morris⁷⁴, A.G. Morris⁵⁵,
 R. Mountain⁶⁷, H. Mu³, F. Muheim⁵⁷, M. Mukherjee⁷, M. Mulder⁴⁷, D. Müller⁴⁷, K. Müller⁴⁹,
 C.H. Murphy⁶², D. Murray⁶¹, P. Muzzetto²⁶, P. Naik⁵³, T. Nakada⁴⁸, R. Nandakumar⁵⁶,
 T. Nanut⁴⁸, I. Nasteva², M. Needham⁵⁷, I. Neri^{20,g}, N. Neri^{25,n}, S. Neubert⁷⁴, N. Neufeld⁴⁷,
 R. Newcombe⁶⁰, T.D. Nguyen⁴⁸, C. Nguyen-Mau⁴⁸, E.M. Niel¹¹, S. Nieswand¹³, N. Nikitin³⁹,
 N.S. Nolte⁴⁷, C. Nunez⁸⁴, A. Oblakowska-Mucha³⁴, V. Obraztsov⁴³, D.P. O'Hanlon⁵³,
 R. Oldeman^{26,f}, C.J.G. Onderwater⁷⁷, A. Ossowska³³, J.M. Otalora Goicochea²,
 T. Ovsianikova³⁸, P. Owen⁴⁹, A. Oyanguren⁴⁶, B. Pagare⁵⁵, P.R. Pais⁴⁷, T. Pajero^{28,47,q},
 A. Palano¹⁸, M. Palutan²², Y. Pan⁶¹, G. Panshin⁸², A. Papanestis⁵⁶, M. Pappagallo^{18,d},
 L.L. Pappalardo^{20,g}, C. Pappenheimer⁶⁴, W. Parker⁶⁵, C. Parkes⁶¹, C.J. Parkinson⁴⁵,
 B. Passalacqua²⁰, G. Passaleva²¹, A. Pastore¹⁸, M. Patel⁶⁰, C. Patrignani^{19,e}, C.J. Pawley⁷⁸,
 A. Pearce⁴⁷, A. Pellegrino³¹, M. Pepe Altarelli⁴⁷, S. Perazzini¹⁹, D. Pereima³⁸, P. Perret⁹,
 K. Petridis⁵³, A. Petrolini^{23,h}, A. Petrov⁷⁹, S. Petrucci⁵⁷, M. Petruzzo²⁵, A. Philippov⁴¹,
 L. Pica²⁸, M. Piccini⁷⁶, B. Pietrzyk⁸, G. Pietrzyk⁴⁸, M. Pili⁶², D. Pinci³⁰, J. Pinzino⁴⁷,
 F. Pisani⁴⁷, A. Piucci¹⁶, Resmi P.K¹⁰, V. Placinta³⁶, S. Playfer⁵⁷, J. Plews⁵², M. Plo Casasus⁴⁵,
 F. Polci¹², M. Poli Lener²², M. Poliakova⁶⁷, A. Poluektov¹⁰, N. Polukhina^{80,c}, I. Polyakov⁶⁷,
 E. Polycarpo², G.J. Pomery⁵³, S. Ponce⁴⁷, A. Popov⁴³, D. Popov^{5,47}, S. Popov⁴¹,
 S. Poslavskii⁴³, K. Prasanth³³, L. Promberger⁴⁷, C. Prouve⁴⁵, V. Pugatch⁵¹, A. Puig Navarro⁴⁹,
 H. Pullen⁶², G. Punzi^{28,m}, W. Qian⁵, J. Qin⁵, R. Quagliani¹², B. Quintana⁸, N.V. Raab¹⁷,
 R.I. Rabadan Trejo¹⁰, B. Rachwal³⁴, J.H. Rademacker⁵³, M. Rama²⁸, M. Ramos Pernas⁵⁵,
 M.S. Rangel², F. Ratnikov^{41,81}, G. Raven³², M. Reboud⁸, F. Redi⁴⁸, F. Reiss¹²,
 C. Remon Alepuz⁴⁶, Z. Ren³, V. Renaudin⁶², R. Ribatti²⁸, S. Ricciardi⁵⁶, D.S. Richards⁵⁶,
 K. Rinnert⁵⁹, P. Robbe¹¹, A. Robert¹², G. Robertson⁵⁷, A.B. Rodrigues⁴⁸, E. Rodrigues⁵⁹,
 J.A. Rodriguez Lopez⁷³, A. Rollings⁶², P. Roloff⁴⁷, V. Romanovskiy⁴³, M. Romero Lamas⁴⁵,
 A. Romero Vidal⁴⁵, J.D. Roth⁸⁴, M. Rotondo²², M.S. Rudolph⁶⁷, T. Ruf⁴⁷, J. Ruiz Vidal⁴⁶,
 A. Ryzhikov⁸¹, J. Ryzka³⁴, J.J. Saborido Silva⁴⁵, N. Sagidova³⁷, N. Sahoo⁵⁵, B. Saitta^{26,f},
 D. Sanchez Gonzalo⁴⁴, C. Sanchez Gras³¹, C. Sanchez Mayordomo⁴⁶, R. Santacesaria³⁰,
 C. Santamarina Rios⁴⁵, M. Santimaria²², E. Santovetti^{29,j}, D. Saranin⁸⁰, G. Sarpis⁶¹,
 M. Sarpis⁷⁴, A. Sarti³⁰, C. Satriano^{30,p}, A. Satta²⁹, M. Saur⁵, D. Savrina^{38,39}, H. Sazak⁹,
 L.G. Scantlebury Smead⁶², S. Schael¹³, M. Schellenberg¹⁴, M. Schiller⁵⁸, H. Schindler⁴⁷,
 M. Schmelling¹⁵, T. Schmelzer¹⁴, B. Schmidt⁴⁷, O. Schneider⁴⁸, A. Schopper⁴⁷, M. Schubiger³¹,
 S. Schulte⁴⁸, M.H. Schune¹¹, R. Schwemmer⁴⁷, B. Sciascia²², A. Sciubba³⁰, S. Sellam⁴⁵,
 A. Semennikov³⁸, M. Senghi Soares³², A. Sergi^{52,47}, N. Serra⁴⁹, J. Serrano¹⁰, L. Sestini²⁷,
 A. Seuthe¹⁴, P. Seyfert⁴⁷, D.M. Shangase⁸⁴, M. Shapkin⁴³, I. Shchemerov⁸⁰, L. Shchutska⁴⁸,
 T. Shears⁵⁹, L. Shekhtman^{42,u}, V. Shevchenko⁷⁹, E.B. Shields^{24,i}, E. Shmanin⁸⁰,
 J.D. Shupperd⁶⁷, B.G. Siddi²⁰, R. Silva Coutinho⁴⁹, G. Simi²⁷, S. Simone^{18,d}, I. Skiba^{20,g},
 N. Skidmore⁷⁴, T. Skwarnicki⁶⁷, M.W. Slater⁵², J.C. Smallwood⁶², J.G. Smeaton⁵⁴,
 A. Smetkina³⁸, E. Smith¹³, M. Smith⁶⁰, A. Snoch³¹, M. Soares¹⁹, L. Soares Lavra⁹,
 M.D. Sokoloff⁶⁴, F.J.P. Soler⁵⁸, A. Solovev³⁷, I. Solovyev³⁷, F.L. Souza De Almeida²,
 B. Souza De Paula², B. Spaan¹⁴, E. Spadaro Norella^{25,n}, P. Spradlin⁵⁸, F. Stagni⁴⁷, M. Stahl⁶⁴,
 S. Stahl⁴⁷, P. Stefko⁴⁸, O. Steinkamp^{49,80}, S. Stemmler¹⁶, O. Stenyakin⁴³, H. Stevens¹⁴,
 S. Stone⁶⁷, M.E. Stramaglia⁴⁸, M. Straticiu³⁶, D. Strelakina⁸⁰, S. Strokov⁸², F. Suljik⁶²,
 J. Sun²⁶, L. Sun⁷², Y. Sun⁶⁵, P. Svihra⁶¹, P.N. Swallow⁵², K. Swientek³⁴, A. Szabelski³⁵,
 T. Szumlak³⁴, M. Szymanski⁴⁷, S. Taneja⁶¹, Z. Tang³, T. Tekampe¹⁴, F. Teubert⁴⁷,
 E. Thomas⁴⁷, K.A. Thomson⁵⁹, M.J. Tilley⁶⁰, V. Tisserand⁹, S. T'Jampens⁸, M. Tobin⁶,
 S. Tolk⁴⁷, L. Tomassetti^{20,g}, D. Torres Machado¹, D.Y. Tou¹², M. Traill⁵⁸, M.T. Tran⁴⁸,
 E. Trifonova⁸⁰, C. Trippel⁴⁸, A. Tsaregorodtsev¹⁰, G. Tuci^{28,m}, A. Tully⁴⁸, N. Tuning³¹,
 A. Ukleja³⁵, D.J. Unverzagt¹⁶, A. Usachov³¹, A. Ustyuzhanin^{41,81}, U. Uwer¹⁶, A. Vagner⁸²,
 V. Vagnoni¹⁹, A. Valassi⁴⁷, G. Valenti¹⁹, N. Valls Canudas⁴⁴, M. van Beuzekom³¹,

H. Van Hecke⁶⁶, E. van Herwijnen⁸⁰, C.B. Van Hulse¹⁷, M. van Veghel⁷⁷, R. Vazquez Gomez⁴⁵, P. Vazquez Regueiro⁴⁵, C. Vázquez Sierra³¹, S. Vecchi²⁰, J.J. Velthuis⁵³, M. Veltri^{21,o}, A. Venkateswaran⁶⁷, M. Veronesi³¹, M. Vesterinen⁵⁵, D. Vieira⁶⁴, M. Vieites Diaz⁴⁸, H. Viemann⁷⁵, X. Vilasis-Cardona⁸³, E. Vilella Figueras⁵⁹, P. Vincent¹², G. Vitali²⁸, A. Vollhardt⁴⁹, D. Vom Bruch¹², A. Vorobyev³⁷, V. Vorobyev^{42,u}, N. Voropaev³⁷, R. Waldi⁷⁵, J. Walsh²⁸, C. Wang¹⁶, J. Wang³, J. Wang⁷², J. Wang⁴, J. Wang⁶, M. Wang³, R. Wang⁵³, Y. Wang⁷, Z. Wang⁴⁹, D.R. Ward⁵⁴, H.M. Wark⁵⁹, N.K. Watson⁵², S.G. Weber¹², D. Websdale⁶⁰, C. Weisser⁶³, B.D.C. Westhenry⁵³, D.J. White⁶¹, M. Whitehead⁵³, D. Wiedner¹⁴, G. Wilkinson⁶², M. Wilkinson⁶⁷, I. Williams⁵⁴, M. Williams^{63,68}, M.R.J. Williams⁵⁷, F.F. Wilson⁵⁶, W. Wislicki³⁵, M. Witek³³, L. Witola¹⁶, G. Wormser¹¹, S.A. Wotton⁵⁴, H. Wu⁶⁷, K. Wyllie⁴⁷, Z. Xiang⁵, D. Xiao⁷, Y. Xie⁷, H. Xing⁷¹, A. Xu⁴, J. Xu⁵, L. Xu³, M. Xu⁷, Q. Xu⁵, Z. Xu⁵, Z. Xu⁴, D. Yang³, Y. Yang⁵, Z. Yang³, Z. Yang⁶⁵, Y. Yao⁶⁷, L.E. Yeomans⁵⁹, H. Yin⁷, J. Yu⁷⁰, X. Yuan⁶⁷, O. Yushchenko⁴³, K.A. Zarebski⁵², M. Zavertyaev^{15,c}, M. Zdybal³³, O. Zenaiev⁴⁷, M. Zeng³, D. Zhang⁷, L. Zhang³, S. Zhang⁴, Y. Zhang⁴⁷, A. Zhelezov¹⁶, Y. Zheng⁵, X. Zhou⁵, Y. Zhou⁵, X. Zhu³, V. Zhukov^{13,39}, J.B. Zonneveld⁵⁷, S. Zucchelli^{19,e}, D. Zuliani²⁷, G. Zunica⁶¹.

¹ *Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*

² *Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*

³ *Center for High Energy Physics, Tsinghua University, Beijing, China*

⁴ *School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

⁵ *University of Chinese Academy of Sciences, Beijing, China*

⁶ *Institute Of High Energy Physics (IHEP), Beijing, China*

⁷ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China*

⁸ *Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France*

⁹ *Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*

¹⁰ *Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France*

¹¹ *Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*

¹² *LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*

¹³ *I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*

¹⁴ *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*

¹⁵ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*

¹⁶ *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*

¹⁷ *School of Physics, University College Dublin, Dublin, Ireland*

¹⁸ *INFN Sezione di Bari, Bari, Italy*

¹⁹ *INFN Sezione di Bologna, Bologna, Italy*

²⁰ *INFN Sezione di Ferrara, Ferrara, Italy*

²¹ *INFN Sezione di Firenze, Firenze, Italy*

²² *INFN Laboratori Nazionali di Frascati, Frascati, Italy*

²³ *INFN Sezione di Genova, Genova, Italy*

²⁴ *INFN Sezione di Milano-Bicocca, Milano, Italy*

²⁵ *INFN Sezione di Milano, Milano, Italy*

²⁶ *INFN Sezione di Cagliari, Monserrato, Italy*

²⁷ *Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*

²⁸ *INFN Sezione di Pisa, Pisa, Italy*

²⁹ *INFN Sezione di Roma Tor Vergata, Roma, Italy*

³⁰ *INFN Sezione di Roma La Sapienza, Roma, Italy*

³¹ *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*

³² *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*

³³ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*

³⁴ *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*

³⁵ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*

- ³⁶ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³⁷ *Petersburg Nuclear Physics Institute NRC Kurchatov Institute (PNPI NRC KI), Gatchina, Russia*
- ³⁸ *Institute of Theoretical and Experimental Physics NRC Kurchatov Institute (ITEP NRC KI), Moscow, Russia, Moscow, Russia*
- ³⁹ *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ⁴⁰ *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia*
- ⁴¹ *Yandex School of Data Analysis, Moscow, Russia*
- ⁴² *Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia*
- ⁴³ *Institute for High Energy Physics NRC Kurchatov Institute (IHEP NRC KI), Protvino, Russia, Protvino, Russia*
- ⁴⁴ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁵ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁶ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴⁷ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁸ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁹ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁵⁰ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵¹ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵² *University of Birmingham, Birmingham, United Kingdom*
- ⁵³ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁵⁴ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁵ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵⁶ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵⁷ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵⁸ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁹ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁶⁰ *Imperial College London, London, United Kingdom*
- ⁶¹ *Department of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁶² *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁶³ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁶⁴ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁵ *University of Maryland, College Park, MD, United States*
- ⁶⁶ *Los Alamos National Laboratory (LANL), Los Alamos, United States*
- ⁶⁷ *Syracuse University, Syracuse, NY, United States*
- ⁶⁸ *School of Physics and Astronomy, Monash University, Melbourne, Australia, associated to ⁵⁵*
- ⁶⁹ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁷⁰ *Physics and Micro Electronic College, Hunan University, Changsha City, China, associated to ⁷*
- ⁷¹ *Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ³*
- ⁷² *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³*
- ⁷³ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹²*
- ⁷⁴ *Universität Bonn - Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to ¹⁶*
- ⁷⁵ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹⁶*
- ⁷⁶ *INFN Sezione di Perugia, Perugia, Italy, associated to ²⁰*
- ⁷⁷ *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³¹*
- ⁷⁸ *Universiteit Maastricht, Maastricht, Netherlands, associated to ³¹*
- ⁷⁹ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³⁸*
- ⁸⁰ *National University of Science and Technology "MISIS", Moscow, Russia, associated to ³⁸*
- ⁸¹ *National Research University Higher School of Economics, Moscow, Russia, associated to ⁴¹*
- ⁸² *National Research Tomsk Polytechnic University, Tomsk, Russia, associated to ³⁸*
- ⁸³ *DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain, associated to ⁴⁴*
- ⁸⁴ *University of Michigan, Ann Arbor, United States, associated to ⁶⁷*
- ⁸⁵ *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*

^a *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

- ^b*Laboratoire Leprince-Ringuet, Palaiseau, France*
- ^c*P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*
- ^d*Università di Bari, Bari, Italy*
- ^e*Università di Bologna, Bologna, Italy*
- ^f*Università di Cagliari, Cagliari, Italy*
- ^g*Università di Ferrara, Ferrara, Italy*
- ^h*Università di Genova, Genova, Italy*
- ⁱ*Università di Milano Bicocca, Milano, Italy*
- ^j*Università di Roma Tor Vergata, Roma, Italy*
- ^k*AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*
- ^l*Università di Padova, Padova, Italy*
- ^m*Università di Pisa, Pisa, Italy*
- ⁿ*Università degli Studi di Milano, Milano, Italy*
- ^o*Università di Urbino, Urbino, Italy*
- ^p*Università della Basilicata, Potenza, Italy*
- ^q*Scuola Normale Superiore, Pisa, Italy*
- ^r*Università di Modena e Reggio Emilia, Modena, Italy*
- ^s*Università di Siena, Siena, Italy*
- ^t*MSU - Iligan Institute of Technology (MSU-IIT), Iligan, Philippines*
- ^u*Novosibirsk State University, Novosibirsk, Russia*