

Weighted energy estimates for the incompressible Navier-Stokes equations and applications to axisymmetric solutions without swirl

Pedro Gabriel Fernández-Dalgo, Pierre Gilles Lemarié-Rieusset

► To cite this version:

Pedro Gabriel Fernández-Dalgo, Pierre Gilles Lemarié-Rieusset. Weighted energy estimates for the incompressible Navier-Stokes equations and applications to axisymmetric solutions without swirl. 2020. hal-02955438v1

HAL Id: hal-02955438 https://hal.science/hal-02955438v1

Preprint submitted on 1 Oct 2020 (v1), last revised 15 Jun 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Weighted energy estimates for the incompressible Navier-Stokes equations and applications to axisymmetric solutions without swirl

Pedro Gabriel Fernández-Dalgo^{*†}, Pierre Gilles Lemarié-Rieusset^{‡§}

Abstract

We consider a family of weights which permit to generalize the Leray procedure to obtain weak suitable solutions of the 3D incompressible Navier–Stokes equations with initial data in weighted L^2 spaces. Our principal result concerns the existence of regular global solutions when the initial velocity is an axisymmetric vector field without swirl such that both the initial velocity and its vorticity belong to $L^2((1+r^2)^{-\frac{\gamma}{2}}dx)$, with $r = \sqrt{x_1^2 + x_2^2}$ and $\gamma \in (0, 2)$.

Keywords : Navier–Stokes equations, axisymmetric vector fields, swirl, Muckenhoupt weights, energy balance

AMS classification : 35Q30, 76D05.

1 Introduction

In 1934, Leray [7] proved global existence of weak solutions for the 3D incompressible Navier–Stokes equations

$$(NS) \begin{cases} \partial_t \mathbf{u} = \Delta \mathbf{u} - (\mathbf{u} \cdot \nabla) \mathbf{u} - \nabla p \\ \nabla \cdot \mathbf{u} = 0, \qquad \mathbf{u}(0, .) = \mathbf{u}_0 \end{cases}$$

[§]e-mail : pierregilles.lemarierieusset@univ-evry.fr

^{*}Université Paris-Saclay, CNRS, Univ Evry, Laboratoire de Mathématiques et Modélisation d'Evry, 91037, Evry-Courcouronnes, France

[†]e-mail : pedro.fernandez@univ-evry.fr

[‡]Université Paris-Saclay, CNRS, Univ Evry, Laboratoire de Mathématiques et Modélisation d'Evry, 91037, Evry-Courcouronnes, France

in the case of a fluid filling the whole space whose initial velocity \mathbf{u}_0 is in L^2 . Leray's strategy is to regularize the initial value and to mollify the nonlinearity through convolution with a bump function : let $\theta_{\epsilon}(x) = \frac{1}{\epsilon^3}\theta(\frac{x}{\epsilon})$, where $\theta \in \mathcal{D}(\mathbb{R}^3)$, θ is non-negative and radially decreasing and $\int \theta \, dx = 1$; the mollified equations are then

$$(NS_{\epsilon}) \begin{cases} \partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} - ((\theta_{\epsilon} * \mathbf{u}_{\epsilon}) \cdot \nabla) \mathbf{u}_{\epsilon} - \nabla p_{\epsilon} \\ \nabla \cdot \mathbf{u}_{\epsilon} = 0, \qquad \mathbf{u}_{\epsilon}, (0, .) = \theta_{\epsilon} * \mathbf{u}_{0}. \end{cases}$$

Standard methods give existence of a smooth solution on an interval $[0, T_{\epsilon}]$ where $T_{\epsilon} \approx \epsilon^3 \|\theta_{\epsilon} * \mathbf{u}_0\|_2^{-2}$. Then, the energy equality

$$\|\mathbf{u}_{\epsilon}(t,.)\|_{2}^{2}+2\int_{0}^{t}\|\nabla\otimes\mathbf{u}_{\epsilon}\|_{2}^{2}ds=\|\theta_{\epsilon}\ast\mathbf{u}_{0}\|_{2}^{2}$$

allows one to extend the existence time and to get a global solution \mathbf{u}_{ϵ} ; moreover, the same energy equality allows one to use a compactness argument and to get a subsequence \mathbf{u}_{ϵ_k} that converges to a solution \mathbf{u} of the Navier– Stokes equations (NS) which satisfies the energy *inequality*

$$\|\mathbf{u}(t,.)\|_{2}^{2} + 2 \int_{0}^{t} \|\nabla \otimes \mathbf{u}\|_{2}^{2} ds \le \|\mathbf{u}_{0}\|_{2}^{2}.$$

Weak solutions of equations (NS) that satisfy this energy inequality are called Leray solutions.

Recently, Bradshaw, Kukavica and Tsai [2] and Fernández-Dalgo and Lemarié-Rieusset [3] used Leray's procedure to find a global weak solution to the equations (NS) when \mathbf{u}_0 is no longer assumed to have finite energy but only to satisfy the weaker assumption

$$\int |\mathbf{u}_0(x)|^2 \, \frac{dx}{1+|x|^2} < +\infty.$$

The solutions then satisfy, for every finite positive T,

$$\sup_{0 \le t \le T} \int |\mathbf{u}(t,x)|^2 \, \frac{dx}{1+|x|^2} + \int_0^T \int |\nabla \otimes \mathbf{u}(t,x)|^2 \, \frac{dx}{1+|x|^2} < +\infty.$$

Whereas the cases of finite energy and of infinite energy sound very similar, this similarity breaks down when we consider higher regularity. Indeed, if we assume that both the initial velocity \mathbf{u}_0 and the initial vorticity $\omega_0 = \nabla \wedge \mathbf{u}_0$ are in L^2 (so that the divergence-free \mathbf{u}_0 belongs to H^1), we find that the Leray solution is unique on some interval [0, T] and remains in H^1 . This is based on the energy equality for the vorticity ω :

$$\|\omega(t,.)\|_{2}^{2} + 2\int_{0}^{t} \|\nabla \otimes \omega\|_{2}^{2} ds = \|\omega_{0}\|_{2}^{2} + 2\int_{0}^{t} \int \omega \cdot (\omega \cdot \nabla) \mathbf{u} \, dx \, ds.$$

The key point is the interpolation inequality

$$\left|\int \omega \cdot (\omega \cdot \nabla) \mathbf{u} \, dx\right| \le C \|\omega\|_3^3 \le C' \|\omega\|_2^{3/2} \|\nabla \otimes \omega\|_2^{3/2}$$

Then the Young inequality gives

$$\|\omega(t,.)\|_{2}^{2} + \int_{0}^{t} \|\nabla \otimes \omega\|_{2}^{2} ds \leq \|\omega_{0}\|_{2}^{2} + C'' \int_{0}^{t} \|\omega\|_{2}^{6} ds.$$

We find that for some positive $T \approx \|\omega_0\|_2^{-4}$ we have

$$\sup_{0 \le t'T} \|\omega\|_2^2 + \int_0^T \|\nabla \otimes \omega\|_2^2 \, ds < +\infty.$$

This strategy fails if we only assume that

$$\int |\mathbf{u}_0(x)|^2 \, \frac{dx}{1+|x|^2} + \int |\omega_0(x)|^2 \, \frac{dx}{1+|x|^2} < +\infty.$$

Indeed, the energy estimate one might hope would be

$$\begin{split} \|\frac{1}{\sqrt{1+|x|^2}}\omega(t,.)\|_2^2 &+ 2\int_0^t \|\frac{1}{\sqrt{1+|x|^2}}\nabla\otimes\omega\|_2^2 \,ds \\ &\leq \|\frac{1}{\sqrt{1+|x|^2}}\omega_0\|_2^2 + 4(\int_0^t \|\frac{1}{\sqrt{1+|x|^2}}\omega\|_2^2 \,ds)^{1/2}(\int_0^t \|\frac{1}{\sqrt{1+|x|^2}}\nabla\otimes\omega\|_2^2 \,ds)^{1/2} \\ &+ C\int_0^t \|\frac{1}{\sqrt{1+|x|^2}}\omega\|_3^2 \|\frac{1}{\sqrt{1+|x|^2}}\mathbf{u}\|_3 \,ds \\ &+ C\int_0^t \int \frac{1}{1+|x|^2}|\omega|^2|\nabla\otimes\mathbf{u}| \,dx \,ds. \end{split}$$

We cannot control the last term due to the lack of integrability : if we want to use interpolation inequalities, we should deal with $\int \frac{1}{(1+|x|^2)^{3/2}} |\omega|^2 |\nabla \otimes \mathbf{u}| dx$ instead of $\int \frac{1}{1+|x|^2} |\omega|^2 |\nabla \otimes \mathbf{u}| dx$.

In this paper, we show that this strategy may work in the case of an axisymmetric flow with no swirl when we consider an axisymmetric weight $\Phi(x) = \frac{1}{(1+r)^{\gamma}}$ (with $0 \leq \gamma < 2$) where $x = (x_1, x_2, x_3)$ and $r = \sqrt{x_1^2 + x_2^2}$. If,

in cylindrical coordinates, $\mathbf{u}_0 = u_{0,r}(r, z) \mathbf{e}_r + u_{0,z}(r, z) \mathbf{e}_z$ and if we assume that \mathbf{u}_0 is in $H^1(\Phi dx)$, i.e.

$$\int |\mathbf{u}_0(x)|^2 \, \Phi(x) \, dx + \int |\omega_0(x)|^2 \, \Phi(x) \, dx < +\infty,$$

we shall obtain the energy estimate

$$\begin{split} \|\sqrt{\Phi}\,\omega(t,.)\|_{2}^{2} &+ 2\int_{0}^{t} \|\sqrt{\Phi}\,\nabla\otimes\omega\|_{2}^{2}\,ds\\ &\leq \|\sqrt{\Phi}\,\omega_{0}\|_{2}^{2} + 2(\int_{0}^{t}\sqrt{\Phi}\,\omega\|_{2}^{2}\,ds)^{1/2}(\int_{0}^{t}\|\sqrt{\Phi}\,\nabla\otimes\omega\|_{2}^{2}\,ds)^{1/2}\\ &+ C\int_{0}^{t}\|\sqrt{\Phi}\,\omega\|_{3}^{2}\|\sqrt{\Phi}\,\mathbf{u}\|_{3}\,ds\\ &+ C\int_{0}^{t}\|\sqrt{\Phi}\,\omega\|_{3}^{3}\,ds. \end{split}$$

This will allow us to find a local-in-time solution in $H^1(\Phi dx)$. Moreover, we shall easily adapt Ladyzhenskaya's result [5] on global existence of axisymmetric solutions in H^1 and find a global solution in $H^1(\Phi dx)$. Remark that, in contrast with the case of H^1 , we cannot prove uniqueness of these regular solutions.

2 Main results.

We shall first prove global existence in the weighted L^2 setting, in dimension d with $2 \leq d \leq 4$ when the weight Φ satisfies some basic assumptions that allow the use of Leray's projection operator and of energy estimates :

Definition 2.1 An adapted weight function Φ on \mathbb{R}^d ($2 \le d \le 4$) is a continuous Lipschitz function Φ such that :

- $(H1) \ 0 < \Phi \le 1.$
- (H2) There exists $C_1 > 0$ such that $|\nabla \Phi| \le C_1 \Phi^{\frac{3}{2}}$
- (H3) There exists $r \in (1, 2]$ such that $\Phi^r \in \mathcal{A}_r$ (where \mathcal{A}_r is the Muckenhoupt class of weights). In the case d = 4, we require r < 2 as well.
- (H4) There exists $C_2 > 0$ such that $\Phi(x) \le \Phi(\frac{x}{\lambda}) \le C_2 \lambda^2 \Phi(x)$, for all $\lambda \ge 1$.

Examples of adapted weights can easily be given by radial slowly decaying functions :

- d = 2, $\Phi(x) = \frac{1}{(1+|x|)^{\gamma}}$ where $0 \le \gamma < 2$
- d = 3 or d = 4, $\Phi(x) = \frac{1}{(1+|x|)^{\gamma}}$ where $0 \le \gamma \le 2$
- d = 3, $\Phi(x) = \frac{1}{(1+r)^{\gamma}}$ where $r = \sqrt{x_1^2 + x_2^2}$ and $0 \le \gamma < 2$.

The following result concerns the existence of weak suitable solutions belonging to a weighted L^2 space, where the weight permits to consider initial data with a weak decay at infinity.

Theorem 1 Let $d \in \{2, 3, 4\}$. Consider a weight Φ satisfying $(\mathbf{H}1) - (\mathbf{H}4)$. Let \mathbf{u}_0 be a divergence free vector field, such that \mathbf{u}_0 belongs to $L^2(\Phi dx, \mathbb{R}^d)$. Then, there exists a global solution \mathbf{u} of the problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

such that

- *u* belongs to L[∞]((0, T), L²(Φdx)) and ∇⊗*u* belongs to L²((0, T), L²(Φdx)), for all T > 0,
- $p = \sum_{1 \le i,j \le d} \mathcal{R}_i \mathcal{R}_j(u_i u_j),$
- the map $t \in [0, +\infty) \mapsto \boldsymbol{u}(t, .)$ is weakly continuous from $[0, +\infty)$ to $L^2(\Phi dx)$, and is strongly continuous at t = 0,
- u satisfies the local energy inequality : there exists a locally finite nonnegative measure μ such that

$$\partial_t(\frac{|\boldsymbol{u}|^2}{2}) = \Delta(\frac{|\boldsymbol{u}|^2}{2}) - |\nabla \otimes \boldsymbol{u}|^2 - \nabla \cdot \left(\frac{|\boldsymbol{u}|^2}{2}\boldsymbol{u}\right) - \nabla \cdot (p\boldsymbol{u}) - \mu,$$

(remark : $\mu = 0$ when d = 2).

If we consider the problem of higher regularity, the case of dimension d = 2 is easy, while, in the case d = 3, one must restrict the study to the case of axisymmetric flows with no swirl (to circumvent the stretching effect in the evolution of the vorticity).

Theorem 2 (Case d = 2.) Let Φ be a weight satisfying $(\mathbf{H}1) - (\mathbf{H}4)$. Let \mathbf{u}_0 be a divergence free vector field, such that $\mathbf{u}_0, \nabla \otimes \mathbf{u}_0$ belong to $L^2(\Phi dx)$. Then there exists a global solution \mathbf{u} of the problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

such that

- \boldsymbol{u} and $\nabla \otimes \boldsymbol{u}$ belong to $L^{\infty}((0,T), L^2(\Phi \, dx))$ and $\Delta \boldsymbol{u}$ belongs to $L^2((0,T), L^2(\Phi \, dx))$, for all T > 0,
- the maps $t \in [0, +\infty) \mapsto \boldsymbol{u}(t, .)$ and $t \in [0, +\infty) \mapsto \nabla \otimes \boldsymbol{u}(t, .)$ are weakly continuous from $[0, +\infty)$ to $L^2(\Phi dx)$, and are strongly continuous at t = 0,

Theorem 3 (Case d = 3.) Let Φ be a weight satisfying $(\mathbf{H}1) - (\mathbf{H}4)$. Let \mathbf{u}_0 be a divergence free axisymmetric vector field without swirl, such that $\mathbf{u}_0, \nabla \otimes \mathbf{u}_0$ belong to $L^2(\Phi dx)$. Assume moreover that Φ depends only on $r = \sqrt{x_1^2 + x_2^2}$. Then there exists a time T > 0, and a local solution \mathbf{u} on (0,T) of the problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

such that

- u is axisymmetric without swirl, u and ∇⊗u belong to L[∞]((0, T), L²(Φ dx)) and Δu belongs to L²((0, T), L²(Φ dx)),
- the maps t → u(t,.) and t → ∇u(t,.) are weakly continuous from [0,T) to L²(Φ dx), and are strongly continuous at t = 0,

An extra condition on the weight permits to obtain a global existence result. Moreover, if the vorticity is more integrable at time t = 0, it will remain so in positive times. The next theorem precise these conditions on the weight.

Theorem 4 (Case d = 3.) Let Φ be a weight satisfying $(\mathbf{H}_1) - (\mathbf{H}_4)$. Assume moreover that Φ depends only on $r = \sqrt{x_1^2 + x_2^2}$. Let Ψ be another continuous weight (that depends only on r) such that $\Phi \leq \Psi \leq 1$, $\Psi \in \mathcal{A}_2$ and there exists $C_1 > 0$ such that

$$|\nabla \Psi| \leq C_1 \sqrt{\Phi} \Psi \text{ and } |\Delta \Psi| \leq C_1 \Phi \Psi.$$

Let u_0 be a divergence free axisymmetric vector field without swirl, such that u_0 , belongs to $L^2(\Phi dx)$ and $\nabla \otimes u_0$ belongs to $L^2(\Psi dx)$. Then there exists a global solution \boldsymbol{u} of the problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

such that

- **u** is axisymmetric without swirl, **u** belongs to $L^{\infty}((0,T), L^2(\Phi dx)), \nabla \otimes$ \boldsymbol{u} belong to $L^{\infty}((0,T), L^2(\Psi dx))$ and $\Delta \boldsymbol{u}$ belongs to $L^2((0,T), L^2(\Psi dx))$, for all T > 0,
- the maps $t \in [0, +\infty) \mapsto \boldsymbol{u}(t, .)$ and $t \in [0, +\infty) \mapsto \nabla \otimes \boldsymbol{u}(t, .)$ are weakly continuous from $[0, +\infty)$ to $L^2(\Phi dx)$ and to $L^2(\Psi dx)$ respectively, and are strongly continuous at t = 0,

Example : we can take $\Phi(x) = \frac{1}{(1+r)^{\gamma}}$ and $\Psi(x) = \frac{1}{(1+r^2)^{\delta/2}}$ with $0 \le \delta \le 1$ $\gamma < 2.$

Some lemmas on weights. 3

Let us first recall the definition of Muckenhoupt weights : for $1 < q < +\infty$, a positive weight w belongs to $\mathcal{A}_q(\mathbb{R}^d)$ if and only if

$$\sup_{x \in \mathbb{R}^{d}, \rho > 0} \left(\frac{1}{|B(x,\rho)|} \int_{B(x,\rho)} \Phi \, dx \right)^{\frac{1}{q}} \left(\frac{1}{|B(x,\rho)|} \int_{B(x,\rho)} \Phi^{-\frac{1}{q-1}} \, dx \right)^{1-\frac{1}{q}} < +\infty.$$
(1)

Due to the Hölder inequality, we have $\mathcal{A}_q(\mathbb{R}^d) \subset \mathcal{A}_r(\mathbb{R}^d)$ if $q \leq r$. One easily checks that $w_{\gamma} = \frac{1}{(1+|x|)^{\gamma}}$ belongs to $\mathcal{A}_q(\mathbb{R}^d)$ if and only if

$$-d(q-1) < \gamma < d.$$

Thus, $\Phi = w_{\gamma}$ is an adapted weight if and only if $0 \leq \gamma \leq 2$ and $\gamma < d$.

One may of course replace in inequality (1) the balls $B(x, \rho)$ by the cubes $Q(x,\rho) =]x_1 - \rho, x_1 + \rho[\times \cdots \times]x_d - \rho, x_d + \rho[$. Thus, we can see that, if $\Phi(x) = \Psi(x_1, x_2)$ and $1 < q < +\infty$, then $\Phi \in \mathcal{A}_q(\mathbb{R}^3)$ if and only if $\Psi \in \mathcal{A}_q(\mathbb{R}^2)$. In particular, $\Phi(x) = \frac{1}{(1+r)^{\gamma}}$ is an adapted weight on \mathbb{R}^3 if and only if $0 \leq \gamma < 2$.

Lemma 3.1 Let Φ satisfy (H1) and (H2) and let $1 \leq r < +\infty$. Then : a) $\sqrt{\Phi}f \in H^1$ if and only if $f \in L^2(\Phi dx)$ and $\nabla f \in L^2(\Phi dx)$; moreover we have

$$\|\sqrt{\Phi}f\|_{H^1} \approx \left(\int \Phi(|f|^2 + |\nabla f|^2) \, dx\right)^{1/2}$$

b) $\Phi f \in W^{1,r}$ if and only if $f \in L^r(\Phi^r dx)$ and $\nabla f \in L^r(\Phi^r dx)$; moreover we have

$$\|\Phi f\|_{W^{1,r}} \approx \left(\int \Phi^r (|f|^r + |\nabla f|^r) \, dx\right)^1$$

Proof. This is obvious since $|\nabla \Phi| \leq C_1 \Phi^{3/2} \leq C_1 \Phi$ and $|\nabla(\sqrt{\Phi})| = \frac{1}{2} \frac{|\nabla \Phi|}{\Phi} \sqrt{\Phi} \leq \frac{1}{2} C_1 \sqrt{\Phi}.$

Lemma 3.2 If $\Phi \in \mathcal{A}_s$ then we have for all $\theta \in (0,1)$, $\Phi^{\theta} \in \mathcal{A}_p$ with $\theta = \frac{p-1}{s-1}$. In particular, if a weight Φ satisfies (**H**3), we obtain $\Phi \in \mathcal{A}_p$ with $p = 1 + \frac{r-1}{r} = 2 - \frac{1}{r} < 2$, and so $\Phi \in \mathcal{A}_2$.

Proof. As $\frac{1}{p} = \frac{1}{s} + \frac{s-p}{ps}$, we find by the Hölder inequality

$$\begin{split} (\int_{Q} \Phi^{\frac{p-1}{s-1}} dx)^{\frac{1}{p}} (\int_{Q} \Phi^{-(\frac{p-1}{s-1})(\frac{1}{p-1})} dx)^{1-\frac{1}{p}} \\ &= (\int_{Q} (\Phi^{\frac{1}{s}} (\Phi^{-\frac{1}{s-1}})^{\frac{s-p}{ps}})^{p} dx)^{\frac{1}{p}} (\int_{Q} \Phi^{-(\frac{p-1}{s-1})(\frac{1}{p-1})} dx)^{1-\frac{1}{p}} \\ &\leq (\int_{Q} \Phi dx)^{\frac{1}{s}} (\int_{Q} \Phi^{-\frac{1}{s-1}} dx)^{\frac{1}{p}-\frac{1}{s}+1-\frac{1}{p}} \end{split}$$

Let us recall that for a weight $w \in \mathcal{A}_q$ $(1 < q < +\infty)$, the Riesz transforms and the Hardy–Littlewood maximal function are bounded on $L^q(w \, dx)$. We thus have the following inequalities :

 \diamond

Lemma 3.3 Let Φ satisfy (H1), (H2) and (H3). Then :

a) for j = 1, ..., d, the Riesz transforms R_j satisfy that $\|\sqrt{\Phi}R_j f\|_2 \leq C \|\sqrt{\Phi}f\|_2$ and $\|\sqrt{\Phi}R_j f\|_{H^1} \leq C \|\sqrt{\Phi}f\|_{H^1}$;

b) for j = 1, ..., d, the Riesz transforms R_j satisfy that $\|\Phi R_j f\|_r \leq C \|\Phi f\|_r$ and $\|\Phi R_j f\|_{W^{1,r}} \leq C \|\Phi f\|_{W^{1,r}}$;

c) if \mathbb{P} is the Leray projection operator on divergence-free vector fields, then for a vector field \mathbf{u} we have $\|\sqrt{\Phi}\mathbb{P}\mathbf{u}\|_2 \leq C\|\sqrt{\Phi}\mathbf{u}\|_2$ and $\|\sqrt{\Phi}\mathbb{P}\mathbf{u}\|_{H^1} \leq C\|\sqrt{\Phi}\mathbf{u}\|_{H^1}$;

d) if $d \in \{2, 3, 4\}$, then for a vector field \boldsymbol{u} we have

$$\|\sqrt{\Phi} \, \boldsymbol{u}\|_{H^1} pprox \|\sqrt{\Phi} \, \boldsymbol{u}\|_2 + \|\sqrt{\Phi}
abla \cdot \boldsymbol{u}\|_2 + \|\sqrt{\Phi}
abla \wedge \boldsymbol{u}\|_2.$$

e) Let $\theta_{\epsilon}(x) = \frac{1}{\epsilon^d} \theta(\frac{x}{\epsilon})$, where $\theta \in \mathcal{D}(\mathbb{R}^d)$, θ is non-negative and radially decreasing and $\int \theta \, dx = 1$. Then we have $\|\sqrt{\Phi} (\theta_{\epsilon} * f)\|_2 \leq C \|\sqrt{\Phi} f\|_2$ and $\|\sqrt{\Phi} (\theta_{\epsilon} * f)\|_{H^1} \leq C(\|\sqrt{\Phi} f\|_{L^2} + \|\sqrt{\Phi} \nabla f\|_{L^2})$ (where the constant C does not depend on ϵ nor f).

Proof. a) is a consequence of $\Phi \in \mathcal{A}_2$ and of Lemma 3.1 (since $\partial_k(R_j f) = R_j(\partial_k f)$). Similarly, b) is a consequence of $\Phi^r \in \mathcal{A}_r$ and of Lemma 3.1.

c) is a consequence of a) : if $\mathbf{v} = \mathbb{P}\mathbf{u}$, then $v_j = \sum_{k=1}^d R_j R_k(u_k)$.

d) is a consequence of a) : if $\mathcal{R} = (R_1, \ldots, R_d)$, we have the identity

$$-\Delta \mathbf{u} =
abla \wedge (
abla \wedge \mathbf{u}) -
abla (
abla \cdot \mathbf{u})$$

so that

$$\partial_k \mathbf{u} = R_k \mathcal{R} \wedge (\nabla \wedge \mathbf{u}) - R_k \mathcal{R} (\nabla \cdot \mathbf{u}).$$

e) is a consequence of $\Phi \in \mathcal{A}_2$ and of Lemma 3.1 : a classical inequality [4] states that we have $|\theta_{\epsilon} * f| \leq \mathcal{M}_f$ (where \mathcal{M}_f is the Hardy–Littlewood maximal function of f) and, similarly, $|\partial_k(\theta_{\epsilon} * f)| \leq \mathcal{M}_{\partial_k f}$.

A final lemma states that Φ is slowly decaying at infinity :

Lemma 3.4 Let Φ satisfy (H1) and (H2). Then there exists a constant C_3 such that

$$\frac{1}{(1+|x|)^2} \le C_3 \Phi.$$

If d = 3 and Φ depends only on $r = \sqrt{x_1^2 + x_2^2}$, then

$$\frac{1}{(1+|r|)^2} \le C_3 \Phi$$

Proof. We define $x_0 = \frac{1}{|x|}x$ and $g(\lambda) = \Phi(\lambda x_0)$. We have

$$g'(\lambda) = x_0 \cdot \nabla \Phi(\lambda x_0) \ge -C_1(\Phi(\lambda x_0))^{3/2} = -C_1 g(\lambda)^{3/2}.$$

Thus

$$C_1 \lambda \ge -\int_0^\lambda g'(\mu)g(\mu)^{-3/2} d\mu = 2(g(\lambda)^{-1/2} - g(0)^{-1/2})$$

and we get

$$\Phi(x)^{-1/2} \le \Phi(0) + \frac{C_1}{2}|x| \le \sqrt{C_3}(1+|x|).$$

If Φ depends only on r, we find that

$$\frac{1}{(1+|r|)^2} \le C_3 \Phi(x_1, x_2, 0) = C_3 \Phi(x).$$

	۰.	
٤.		1
		۲

4 Proof of Theorem 1 (the case of $L^2(\Phi dx)$)

4.1 A priori controls

Let $\phi \in \mathcal{D}(\mathbb{R}^d)$ be a real-valued test function which is equal to 1 in a neighborhood of 0 and let $\phi_{\epsilon}(x) = \phi(\epsilon x)$. Let

$$\mathbf{u}_{0,\epsilon} = \mathbb{P}(\phi_{\epsilon}\mathbf{u}_0)$$

Thus, $\mathbf{u}_{0,\epsilon}$ is divergence free and converges to \mathbf{u}_0 in $L^2(\Phi \, dx)$ since $\Phi \in \mathcal{A}^2$.

Let $\theta_{\epsilon}(x) = \frac{1}{\epsilon^d} \theta(\frac{x}{\epsilon})$, where $\theta \in \mathcal{D}(\mathbb{R}^d)$, θ is non-negative and radially decreasing and $\int \theta \, dx = 1$. We denote $\mathbf{b}_{\epsilon} = \mathbf{u}_{\epsilon} * \theta_{\epsilon}$. Let \mathbf{u}_{ϵ} be the unique global solution of the problem

$$(NS_{\epsilon}) \begin{cases} \partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} - (\mathbf{b}_{\epsilon} \cdot \nabla) \mathbf{u}_{\epsilon} - \nabla p_{\epsilon} \\ \nabla \cdot \mathbf{u}_{\epsilon} = 0, \qquad \mathbf{u}_{\epsilon}, (0, .) = \mathbf{u}_{0, .} \end{cases}$$

which belongs to $\mathcal{C}([0, +\infty), L^2(\mathbb{R}^d)) \cap L^2((0, +\infty), \dot{H}^1(\mathbb{R}^d)).$

We want to demonstrate that

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}(t)\|_{L^{2}}^{2} &+ \int_{0}^{t} \|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}}^{2} ds \\ &\leq \|\sqrt{\Phi}\mathbf{u}_{0,\epsilon}\|_{L^{2}}^{2} + C_{\Phi} \int_{0}^{t} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}^{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}^{2d} ds, \end{aligned}$$

$$(2)$$

where C_{Φ} does not depend on ϵ nor on \mathbf{u}_0 . (When d = 4, the inequality will hold only if $\|\sqrt{\Phi}\mathbf{u}_{\epsilon}(t)\|_{L^2}$ remains small enough).

Since $\sqrt{\Phi}$, $\nabla\sqrt{\Phi} \in L^{\infty}$, pointwise multiplication by $\sqrt{\Phi}$ maps boundedly H^1 to H^1 and H^{-1} to H^{-1} . Thus, $\sqrt{\Phi}\mathbf{u}_{\epsilon} \in L^2H^1$ and $\sqrt{\Phi}\partial_t\mathbf{u}_{\epsilon} \in L^2H^{-1}$, we can calculate $\int \partial_t \mathbf{u}_{\epsilon} \cdot \mathbf{u}_{\epsilon} \Phi dx$ and obtain :

$$\int \frac{|\mathbf{u}_{\epsilon}(t,x)|^{2}}{2} \Phi \, dx + \int_{0}^{t} \int |\nabla \otimes \mathbf{u}_{\epsilon}|^{2} \, \Phi dx \, ds$$
$$= \int \frac{|\mathbf{u}_{0,\epsilon}(x)|^{2}}{2} \Phi \, dx - \int_{0}^{t} \int (\nabla \otimes \mathbf{u}_{\epsilon}) \cdot (\nabla \Phi \otimes \mathbf{u}_{\epsilon}) \, dx \, ds \quad (3)$$
$$+ \int_{0}^{t} \int (\frac{|\mathbf{u}_{\epsilon}|^{2}}{2} \mathbf{b}_{\epsilon} + p \mathbf{u}_{\epsilon}) \cdot \nabla \Phi \, dx \, ds.$$

We use the fact that $|\nabla \Phi| \le C_0 \Phi^{\frac{3}{2}} \le C_0 \Phi$, in order to control the following term

$$\left| -\int_0^t \int (\nabla \otimes \mathbf{u}_{\epsilon}) \cdot (\nabla \Phi \otimes \mathbf{u}_{\epsilon}) dx \, ds \right| \leq \frac{1}{8} \int_0^t \|\sqrt{\Phi} \nabla \otimes \mathbf{u}\|_{L^2(\Phi dx)}^2 + C \int_0^t \|\sqrt{\Phi} \, \mathbf{u}\|_{L^2(\Phi dx)}^2$$

Now, we analyze the integrals containing the pressure term. We distinguish two cases :

• Case 1: d = 2 and $r \in (1, 2]$, or d = 3 and $r \in [\frac{6}{5}, 2]$, or d = 4 and $r \in [\frac{4}{3}, 2)$. For those values of d and r we have

$$0 \leq \frac{d}{2} - \frac{d}{2r} \leq 1$$
 and $\dot{H}^{\frac{d}{2} - \frac{d}{2r}} \subset L^{2r}$

and

$$0 \le \frac{d}{r} - \frac{d}{2} \le 1$$
 and $\dot{H}^{\frac{d}{r} - \frac{d}{2}} \subset L^{\frac{r}{r-1}}$.

Using the continuity of the Riesz transforms on $L^r(\Phi^r dx)$,

$$\int_0^t \int \left(\frac{|\mathbf{u}_{\epsilon}|^2 |\mathbf{b}_{\epsilon}|}{2} + |p| |\mathbf{u}_{\epsilon}|\right) |\nabla \Phi| \, dx \, ds \leq \int_0^t \|\Phi(|\mathbf{u}_{\epsilon}| |\mathbf{b}_{\epsilon}| + |p|)\|_r \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{\frac{r}{r-1}} \\ \leq C \int_0^t \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{2r} \|\sqrt{\Phi} \mathbf{b}_{\epsilon}\|_{2r} \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{\frac{r}{r-1}} \, ds$$

Using the Sobolev embedding $\dot{H}^{\frac{d}{2}-\frac{d}{2r}} \subset L^{2r}$, the fact that $|\nabla\sqrt{\Phi}| \leq C\sqrt{\Phi}$, and the continuity of the maximal function operator on $L^2(\Phi dx)$, we have

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2r} \\ &\leq C\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}^{1-(\frac{d}{2}-\frac{d}{2r})}\|\nabla\otimes(\sqrt{\Phi}\mathbf{b}_{\epsilon})\|_{2}^{\frac{d}{2}-\frac{d}{2r}} \\ &\leq C'\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}^{1-(\frac{d}{2}-\frac{d}{2r})}(\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}+\|\sqrt{\Phi}\nabla\otimes\mathbf{b}_{\epsilon}\|_{2})^{\frac{d}{2}-\frac{d}{2r}} \\ &\leq C''\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{1-(\frac{d}{2}-\frac{d}{2r})}(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}+\|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{2})^{\frac{d}{2}-\frac{d}{2r}}, \end{aligned}$$

and

$$\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2r} \leq C \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{1-(\frac{d}{2}-\frac{d}{2r})} (\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2} + \|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{2})^{\frac{d}{2}-\frac{d}{2r}}.$$

Using the embedding $\dot{H}^{\frac{d}{r}-\frac{d}{2}} \subset L^{\frac{r}{r-1}}$, we also have

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\frac{r}{r-1}} \\ &\leq C\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{1-(\frac{d}{r}-\frac{d}{2})}\|\nabla\otimes(\sqrt{\Phi}\mathbf{u}_{\epsilon})\|_{L^{2}}^{\frac{d}{r}-\frac{d}{2}} \\ &\leq C\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{1-(\frac{d}{r}-\frac{d}{2})}(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}+\|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}})^{\frac{d}{r}-\frac{d}{2}}.\end{aligned}$$

Hence, we find

$$\int_0^t \int \left(\frac{|\mathbf{u}_{\epsilon}|^2 |\mathbf{b}_{\epsilon}|}{2} + |p| |\mathbf{u}_{\epsilon}|\right) |\nabla \Phi| \, dx \, ds$$

$$\leq C \int_0^t \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2^{3-\frac{d}{2}} (\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2 + \|\sqrt{\Phi} \nabla \otimes \mathbf{u}_{\epsilon}\|_{L^2})^{\frac{d}{2}} \, ds.$$

Using the Young inequality, we then find for d = 2 or d = 3

$$\int_{0}^{t} \int \left(\frac{|\mathbf{u}_{\epsilon}|^{2}|\mathbf{b}_{\epsilon}|}{2} + |p||\mathbf{u}_{\epsilon}|\right) |\nabla\Phi| \, dx \, ds$$

$$\leq \frac{1}{8} \int_{0}^{t} \|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}}^{2} \, ds + C_{\Phi} \int_{0}^{t} \|\sqrt{\Phi}\mathbf{u}_{0}\|_{L^{2}}^{2} + \|\sqrt{\Phi}\mathbf{u}_{0}\|_{L^{2}}^{\frac{12-2d}{4-d}} \, ds,$$

where, as $d \in \{2, 3\}$, we have $\frac{12-2d}{4-d} = 2d$.

When d = 4, provided that $\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2 < \epsilon_0$ with $C\epsilon_0 < \frac{1}{8}$ we find

$$\int_0^t \int \left(\frac{|\mathbf{u}_{\epsilon}|^2 |\mathbf{b}_{\epsilon}|}{2} + |p| |\mathbf{u}_{\epsilon}|\right) |\nabla \Phi| \, dx \, ds$$
$$\leq \frac{1}{8} \int_0^t \|\sqrt{\Phi} \nabla \otimes \mathbf{u}_{\epsilon}\|_{L^2}^2 \, ds + \frac{1}{8} \int_0^t \|\sqrt{\Phi} \mathbf{u}_0\|_{L^2}^2 \, ds$$

• Case 2: d = 3 and $r \in (1, \frac{6}{5})$, or d = 4 and $r \in (1, \frac{4}{3})$. Let $q = \frac{dr}{d-r}$; for those values of d, r and q, we have

$$W^{1,r} \subset L^q$$

$$0 \leq \frac{d}{2} - \frac{d}{2r} \leq 1 \text{ and } \dot{H}^{d(1-\frac{1}{r})} \subset L^{\frac{2r}{2-r}}$$

and

$$0 \le \frac{d}{r} - \frac{d}{2} - 1 \le 1$$
 and $\dot{H}^{\frac{d}{r} - \frac{d}{2} - 1} \subset L^{\frac{q}{q-1}}$.

Using the continuity of the Riesz transforms on $L^r(\Phi^r dx)$, we have

$$\int_{0}^{t} \int \left(\frac{|\mathbf{u}_{\epsilon}|^{2}|\mathbf{b}_{\epsilon}|}{2} + |p||\mathbf{u}_{\epsilon}|\right) |\nabla\Phi| \, dx \, ds$$

$$\leq \int_{0}^{t} \|\Phi|\mathbf{u}_{\epsilon}|^{2}\|_{q} \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\frac{q}{q-1}} \, ds + \int_{0}^{t} \|\Phi p\|_{q} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\frac{q}{q-1}} \, ds$$

$$\leq C \int_{0}^{t} \|\Phi|\mathbf{u}_{\epsilon}|^{2}\|_{W^{1,r}} \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\frac{q}{q-1}} \, ds + \sum_{ij} \int_{0}^{t} \|\Phi b_{\epsilon,i}u_{\epsilon,j}\|_{W^{1,r}} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\frac{q}{q-1}} \, ds.$$

We have

$$\begin{split} \|\Phi b_{\epsilon,i}u_{\epsilon,j}\|_{W^{1,r}} \\ &\leq \|\Phi b_{\epsilon,i}u_{\epsilon,j}\|_{r} + \sum_{k} (\|b_{\epsilon,i}u_{\epsilon,j}\partial_{k}\Phi\|_{L^{r}} + \|\Phi b_{\epsilon,i}\partial_{k}u_{\epsilon,j}\|_{L^{r}} + \|\Phi u_{\epsilon,i}\partial_{k}b_{\epsilon,j}\|_{L^{r}}) \\ &\leq C(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\frac{2r}{2-r}}\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2} + \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\frac{2r}{2-r}}\|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\frac{2r}{2-r}}\|\sqrt{\Phi}\nabla\otimes\mathbf{b}_{\epsilon}\|_{2}), \\ &\leq C'(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}} + \|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}})(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{\dot{H}^{d(1-\frac{1}{r})}} + \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\dot{H}^{d(1-\frac{1}{r})}}). \end{split}$$

We have

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\dot{H}^{d(1-\frac{1}{r})}} \\ &\leq C\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}^{1-(d-\frac{d}{r})}\|\nabla\otimes(\sqrt{\Phi}\mathbf{b}_{\epsilon})\|_{2}^{d-\frac{d}{r}} \\ &\leq C'\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}^{1-(d-\frac{d}{r})}(\|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}+\|\sqrt{\Phi}\nabla\otimes\mathbf{b}_{\epsilon}\|_{2})^{d-\frac{d}{r}} \\ &\leq C''\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{1-(d-\frac{d}{r})}(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}+\|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}})^{d-\frac{d}{r}}, \end{aligned}$$

and finally we get

$$\sum_{i,j} \|\Phi b_{\epsilon,i} u_{\epsilon,j}\|_{W^{1,r}} + \|\Phi |u_{\epsilon}|^{2}\|_{W^{1,r}}$$

$$\leq C \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{2}^{1-(d-\frac{d}{r})} (\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{L^{2}} + \|\sqrt{\Phi} \nabla \otimes \mathbf{u}_{\epsilon}\|_{L^{2}})^{1+d-\frac{d}{r}}.$$

On the other hand, we have

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{\frac{q}{q-1}} \\ &\leq C \|\sqrt{\Phi}\mathbf{b}_{\epsilon}\|_{2}^{2-(\frac{d}{r}-\frac{d}{2})} \|\nabla\otimes(\sqrt{\Phi}\mathbf{b}_{\epsilon})\|_{2}^{\frac{d}{r}-\frac{d}{2}-1} \\ &\leq C' \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{2-(\frac{d}{r}-\frac{d}{2})} (\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}} + \|\sqrt{\Phi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}})^{\frac{d}{r}-\frac{d}{2}-1}. \end{aligned}$$

Hence, we find again

$$\int_0^t \int \left(\frac{|\mathbf{u}_{\epsilon}|^2 |\mathbf{b}_{\epsilon}|}{2} + |p| |\mathbf{u}_{\epsilon}|\right) |\nabla \Phi| \, dx \, ds$$

$$\leq C \int_0^t \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2^{3-\frac{d}{2}} (\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2 + \|\sqrt{\Phi} \nabla \otimes \mathbf{u}_{\epsilon}\|_{L^2})^{\frac{d}{2}} \, ds.$$

and we conclude in the same way as for the first case.

In the Case 1 and Case 2, we have found

$$\int_{0}^{t} \int \left(\frac{|\mathbf{u}_{\epsilon}|^{2}|\mathbf{b}_{\epsilon}|}{2} + |p||\mathbf{u}_{\epsilon}|\right) |\nabla\Phi| \, dx \, ds$$

$$\leq \frac{1}{8} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}^{2} + C_{\Phi} \int_{0}^{t} \|\sqrt{\Phi}\mathbf{u}_{0}\|_{L^{2}}^{2} + \|\sqrt{\Phi}\mathbf{u}_{0}\|_{L^{2}}^{2d} \, ds.$$

From these controls, we get inequality (3), and thus inequality (2). Inequality (2) gives us a control on the size of $\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_2$ on an interval of time that does not depend on ϵ :

Lemma 4.1 If α is a continuous non-negative function on [0,T) which satisfies, for three constants $A, B \in (0, +\infty)$ and $b \in [1, \infty)$,

$$\alpha(t) \le A + B \int_0^t \alpha(s) + \alpha(s)^b \, ds.$$

Let $0 < T_1 < T$ and $T_0 = \min(T_1, \frac{1}{3^b(A^{b-1} + (BT_1)^{b-1})})$. We have, for every $t \in [0, T_0], \alpha(t) \leq 3A$.

Proof. We try to estimate the first time $T^* < T_1$ (if it exists) for which we have

$$\alpha(T^*) = 3A$$

We have

$$\alpha \le \frac{A}{BT_1} + (\frac{BT_1}{A})^{b-1} \alpha^b.$$

We thus find

$$\alpha(T^*) \le 2A + T^*(3A)^b (1 + (\frac{BT_1}{A})^{b-1})$$

and thus

$$T^*3^b(A^{b-1} + (BT_1)^{b-1}) \ge 1.$$

 \diamond

By Lemma 4.1 and (2), we thus find that there exists a constant $C_{\Phi} \ge 1$ such that if T_0 satisfies

• if
$$d = 2$$
, $C_{\Phi} \left(1 + \|\mathbf{u}_0\|_{L^2(\Phi dx)}^2 \right) T_0 \le 1$
• if $d = 3$, $C_{\Phi} \left(1 + \|\mathbf{u}_0\|_{L^2(\Phi dx)}^2 \right)^2 T_0 \le 1$

• if
$$d = 4$$
 and $\|\mathbf{u}_0\|_{L^2(\Phi \, dx)} \le \frac{1}{C_{\Phi}}, C_{\Phi} T_0 \le 1$

then

$$\sup_{0 \le t \le T_0} \| \mathbf{u}_{\epsilon}(t,.) \|_{L^2(\Phi dx)}^2 + \int_0^{T_0} \| \nabla \otimes \mathbf{u}_{\epsilon} \|_{L^2(\Phi dx)}^2 \, ds \le C_{\Phi} (1 + \| \mathbf{u}_0 \|_{L^2(\Phi dx)}^2).$$
(4)

4.2 Passage to the limit and local existence

We know that \mathbf{u}_{ϵ} is bounded in $L^{\infty}((0, T_0), L^2(\Phi \, dx))$ and $\nabla \otimes \mathbf{u}_{\epsilon}$ is bounded in $L^2((0, T_0), L^2(\Phi \, dx))$. This will alow us to use a simple variant of the Aubin-Lions theorem :

Lemma 4.2 (Aubin–Lions theorem) Let s > 0, 1 < q and $\sigma < 0$. Let (f_n) be a sequence of functions on $(0,T) \times \mathbb{R}^d$ such that, for all $T_0 \in (0,T)$ and all $\varphi \in \mathcal{D}(\mathbb{R}^d)$,

- φf_n is bounded in $L^2((0,T_0),H^s)$
- $\varphi \partial_t f_n$ is bounded in $L^q((0,T_0), H^{\sigma})$.

Then, there exists a subsequence (f_{n_k}) such that f_{n_k} is strongly convergent in $L^2_{\text{loc}}([0,T] \times \mathbb{R}^d)$. More precisely : if we denote f_{∞} the limit, then for all $T_0 \in (0,T)$ and all $R_0 > 0$,

$$\lim_{n_k \to +\infty} \int_0^{T_0} \int_{|x| \le R_0} |f_{n_k} - f_\infty|^2 \, dx \, dt = 0.$$

For a proof of the Lemma, see [1, 6].

We want to verify that $\varphi \partial_t \mathbf{u}_{\epsilon}$ is bounded in $L^{\alpha}((0, T_0), H^{-s})$ for some $s \in (-\infty, 0)$ and some $\alpha > 1$.

In Case 1, we have that $\Phi \mathbf{b}_{\epsilon} \otimes \mathbf{u}_{\epsilon}$ and $\Phi p_{\epsilon} = \sum_{i=1}^{3} \sum_{j=1}^{3} R_i R_j(b_{\epsilon,i} u_{\epsilon,j})$ are bounded in $L^{\alpha_1}((0,T_0), L^r)$, where $\alpha_1 = \frac{2r}{dr-d}$, so that $\alpha_1 \in [2,\infty)$ if d=2, $\alpha_1 \in [\frac{4}{3}, 4]$ if d=3 and $\alpha_1 \in (1,2]$ if d=4.

In Case 2, we have that $\Phi \mathbf{b}_{\epsilon} \otimes \mathbf{u}_{\epsilon}$ and Φp_{ϵ} are bounded in $L^{\alpha_2}((0, T_0), W^{1,r})$, where $\alpha_2 = \frac{2r}{r+dr-d}$ and thus it is bounded in $L^{\alpha_2}L^q$, with $q = \frac{dr}{d-r}$. We have $\alpha_2 \in (\frac{4}{3}, 2)$ if d = 3 and $\alpha_2 \in (1, 2)$ if d = 4.

Let $\varphi \in \mathcal{D}(\mathbb{R}^d)$. We have that $\varphi \mathbf{u}_{\epsilon}$ is bounded in $L^2((0, T_0), H^1)$; moreover, writing

$$\partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} - \left(\sum_{j=1}^3 \partial_j (b_{\epsilon,j} \mathbf{u}_{\epsilon}) + \nabla p_{\epsilon}\right)$$

and using the embeddings $L^r \subset \dot{H}^{\frac{d}{2}-\frac{d}{r}} \subset H^{-1}$ (in Case 1) or $L^{\frac{dr}{d-r}} \subset H^{-(\frac{d}{r}-\frac{d}{2}-1)} \subset H^{-1}$ (in Case 2) we see that $\varphi \partial_t \mathbf{u}_{\epsilon}$ is bounded in $L^{\alpha_i}((0,T_0), H^{-2})$.

Thus, by the Aubin–Lions theorem, there exist **u** and a sequence $(\epsilon_k)_{k\in\mathbb{N}}$ converging to 0 such that \mathbf{u}_{ϵ_k} converges strongly to **u** in $L^2_{\text{loc}}([0, T_0) \times \mathbb{R}^3)$:

for every $\tilde{T} \in (0, T_0)$ and every R > 0, we have

$$\lim_{k \to +\infty} \int_0^{\tilde{T}} \int_{|y| < R} |\mathbf{u}_{\epsilon_k} - \mathbf{u}|^2 \, dx \, ds = 0.$$

Then, we have that \mathbf{u}_{ϵ_k} converge *-weakly to \mathbf{u} in $L^{\infty}((0, T_0), L^2(\Phi dx))$, $\nabla \otimes \mathbf{u}_{\epsilon_k}$ converges weakly to $\nabla \otimes \mathbf{u}$ in $L^2((0, T_0), L^2(\Phi dx))$, and \mathbf{u}_{ϵ_k} converges weakly to \mathbf{u} in $L^3((0, T_0), L^3(\Phi^{\frac{3}{2}}dx))$. We deduce that $\mathbf{b}_{\epsilon_k} \otimes \mathbf{u}_{\epsilon_k}$ is weakly convergent in $(L^{6/5}L^{6/5})_{\text{loc}}$ to $\mathbf{b} \otimes \mathbf{u}$ and thus in $\mathcal{D}'((0, T_0) \times \mathbb{R}^d)$; as in Case 1, it is bounded in $L^{\alpha_1}((0, T_0), L^r)$, and in Case 2 it is bounded in $L^{\alpha_2}((0, T_0), W^{1,r})$, it is weakly convergent in these spaces respectively (as \mathcal{D} is dense in their dual spaces).

By the continuity of the Riesz transforms on $L^r(\Phi^r dx)$ and on $W^{1,r}(\Phi^r dx)$, we find that in the Case 1 and Case 2, p_{ϵ_k} is convergent to the distribution $p = \sum_{i=1}^3 \sum_{j=1}^3 R_i R_j(u_i u_j)$. We have obtained

$$\partial_t \mathbf{u} = \Delta \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} - \nabla p.$$

Moreover, we have seen that $\partial_t \mathbf{u}$ is locally in $L^1 H^{-2}$, and thus \mathbf{u} has representative such that $t \mapsto \mathbf{u}(t, .)$ is continuous from $[0, T_0)$ to $\mathcal{D}'(\mathbb{R}^d)$ and coincides with $\mathbf{u}(0, .) + \int_0^t \partial_t \mathbf{u} \, ds$.

In the sense of distributions, we have

$$\mathbf{u}(0,.) + \int_0^t \partial_t \mathbf{u} \, ds = \mathbf{u} = \lim_{k \to +\infty} \mathbf{u}_{\epsilon_k} = \lim_{k \to +\infty} \mathbf{u}_{0,\epsilon_k} + \int_0^t \partial_t \mathbf{u}_{n_k} \, ds = \mathbf{u}_0 + \int_0^t \partial_t \mathbf{u} \, ds,$$

hence, $\mathbf{u}(0, .) = \mathbf{u}_0$, and \mathbf{u} is a solution of (NS).

Now, we want to prove the energy balance. In the case of dimension 2, we remark that, since $\sqrt{\Phi}\mathbf{u} \in L^{\infty}L^2 \cap L^2H^1$, we have by interpolation that $\sqrt{\Phi}\mathbf{u} \in L^4L^4$, and then we can define $((\mathbf{u} \cdot \nabla)\mathbf{u}) \cdot \mathbf{u}$. The equality

$$\partial_t(\frac{|\mathbf{u}|^2}{2}) = \Delta(\frac{|\mathbf{u}|^2}{2}) - |\nabla \mathbf{u}|^2 - \nabla \cdot \left(\frac{|\mathbf{u}|^2}{2}\mathbf{u}\right) - \nabla \cdot (p\mathbf{u})$$

is then easy to prove.

Let us consider the case $d \ge 3$. We define

$$A_{\epsilon} = -\partial_t (\frac{|\mathbf{u}_{\epsilon}|^2}{2}) + \Delta(\frac{|\mathbf{u}_{\epsilon}|^2}{2}) - \nabla \cdot \left(\frac{|\mathbf{u}_{\epsilon}|^2}{2}\mathbf{u}_{\epsilon}\right) - \nabla \cdot (p_{\epsilon}\mathbf{u}_{\epsilon}) = |\nabla \otimes \mathbf{u}_{\epsilon}|^2.$$

As u_{ϵ_k} is locally strongly convergent in $L^2 L^2$; and locally bounded in $L^{\infty} L^2$, it is then locally strongly convergent in $L^{p'}L^2$, with $p' < \infty$. Then, as $\sqrt{\Phi} \nabla \otimes$ \mathbf{u}_{ϵ} is bounded in $L^2((0,T), L^2)$, by the Gagliardo-Nirenberg interpolation inequalities we obtain \mathbf{u}_{ϵ_k} is locally strongly convergent in $L^{p'}L^{q'}$ with $\frac{2}{p'} + \frac{3}{q'} > \frac{d}{2}$.

In Case 1, we know that p_{ϵ_k} is locally weakly convergent in $L^{\alpha}L^r$ and we know that \mathbf{u}_{ϵ_k} is locally strongly convergent in $L^{\frac{\alpha}{\alpha-1}}L^{\frac{r}{r-1}}$, and hence $p_{\epsilon_k}\mathbf{u}_{\epsilon_k}$ converges in the sense of distributions.

In Case 2, we know that p_{ϵ_k} is locally weakly convergent in $L^{\beta}L^q$ and we know that \mathbf{u}_{ϵ_k} is locally strongly convergent in $L^{\frac{\beta}{\beta-1}}L^{\frac{q}{q-1}}$, and hence $p_{\epsilon_k}\mathbf{u}_{\epsilon_k}$ converges in the sense of distributions.

Thus, A_{ϵ_k} is convergent in $\mathcal{D}'((0,T) \times \mathbb{R}^d)$ to

$$A = -\partial_t \left(\frac{|\mathbf{u}|^2}{2}\right) + \Delta\left(\frac{|\mathbf{u}|^2}{2}\right) - \nabla \cdot \left(\frac{|\mathbf{u}|^2}{2}\mathbf{u}\right) - \nabla \cdot (p\mathbf{u}),$$

and $A = \lim_{k \to +\infty} |\nabla \otimes \mathbf{u}_{\epsilon_k}|^2$. If $\theta \in \mathcal{D}((0,T) \times \mathbb{R}^d)$ is non-negative, we have that $\sqrt{\theta} \nabla \otimes \mathbf{u}_{\epsilon_k}$ is weakly convergent in $L^2 L^2$ to $\sqrt{\theta} \nabla \otimes \mathbf{u}$, so that

$$\iint A\theta \, dx \, ds = \lim_{\epsilon_k \to +\infty} \iint A_{\epsilon_k} \theta \, dx \, ds = \lim_{k \to +\infty} \iint |\nabla \otimes \mathbf{u}_{\epsilon_k}|^2 \theta \, dx \, ds \ge \iint |\nabla \otimes \mathbf{u}|^2 \theta \, dx \, ds.$$

Hence, there exists a non-negative locally finite measure μ on $(0,T) \times \mathbb{R}^3$ such that $A = |\nabla \mathbf{u}|^2 + \mu$, i.e. such that

$$\partial_t \left(\frac{|\mathbf{u}|^2}{2}\right) = \Delta\left(\frac{|\mathbf{u}|^2}{2}\right) - |\nabla \mathbf{u}|^2 - \nabla \cdot \left(\frac{|\mathbf{u}|^2}{2}\mathbf{u}\right) - \nabla \cdot (p\mathbf{u}) - \mu.$$

4.3 Convergence to the initial data

In order to take the limit to $k \to \infty$, first we introduce a test function $0 \le \varphi \le 1$ which is equal to 1 in a neighborhood of 0 and we let $\varphi_R(x) = \varphi(\frac{x}{R})$. We have,

$$\int \frac{|\mathbf{u}_{\epsilon_k}(t,x)|^2}{2} \varphi_R \Phi \, dx + \int_0^t \int |\nabla \otimes \mathbf{u}_{\epsilon_k}|^2 \, \varphi_R \Phi \, dx \, ds$$

= $\int \frac{|\mathbf{u}_{0,\epsilon}(x)|^2}{2} \varphi_R \Phi \, dx - \sum_{1 \le i \le d} \int_0^t \int \partial_i \mathbf{u}_{\epsilon_k} \cdot \mathbf{u}_{\epsilon_k} \, (\partial_i \varphi_R \, \Phi + \partial_i \Phi \, \varphi_R) \, dx \, ds$
+ $\sum_i \int_0^t \int (\frac{|\mathbf{u}_{\epsilon_k}|^2}{2} \mathbf{b}_{\epsilon_k,i} + p_\epsilon \mathbf{u}_{\epsilon_k,i}) \, (\partial_i \varphi_R \, \Phi + \partial_i \Phi \, \varphi_R) \, dx \, ds$

and then we find

$$\begin{split} &\limsup_{k \to +\infty} \int \frac{|\mathbf{u}_{\epsilon_k}(t,x)|^2}{2} \varphi_R \Phi \, dx + \int_0^t \int |\nabla \otimes \mathbf{u}_{\epsilon_k}|^2 \, \varphi_R \Phi \, dx \, ds \\ &= \int \frac{|\mathbf{u}_0(x)|^2}{2} \varphi_R \Phi \, dx - \sum_{1 \le i \le d} \int_0^t \int \partial_i \mathbf{u} \cdot \mathbf{u} \, (\partial_i \varphi_R \, \Phi + \partial_i \Phi \, \varphi_R) \, dx \, ds \\ &+ \sum_i \int_0^t \int (\frac{|\mathbf{u}|^2}{2} + p) \mathbf{u}_i \, (\partial_i \varphi_R \, \Phi + \partial_i \Phi \, \varphi_R) \, dx \, ds. \end{split}$$

Since $\mathbf{u}_{\epsilon_k} = \mathbf{u}_{0,\epsilon_k} + \int_0^t \partial_t \mathbf{u}_{\epsilon_k} \, ds$, we see that $\mathbf{u}_{\epsilon_k}(t, .)$ is convergent to $\mathbf{u}(t, .)$ in $\mathcal{D}'(\mathbb{R}^d)$, hence is weakly convergent in L^2_{loc} (as it is bounded in $L^2(\Phi dx)$), so that :

$$\int \frac{|\mathbf{u}(t,x)|^2}{2} \varphi_R \Phi \, dx \leq \limsup_{k \to +\infty} \int \frac{|\mathbf{u}_{\epsilon_k}(t,x)|^2}{2} \varphi_R \Phi \, dx.$$

On the other hand, as $\nabla \otimes \mathbf{u}_{\epsilon_k}$ is weakly convergent in $L^2 L^2(\Phi dx)$, we have

$$\int_0^t \int \frac{|\nabla \otimes \mathbf{u}|^2}{2} \varphi_R \Phi \, dx \, ds \leq \limsup_{k \to +\infty} \int_0^t \int \frac{|\nabla \otimes \mathbf{u}_{\epsilon_k}|^2}{2} \varphi_R \Phi \, dx \, ds.$$

Thus, letting R go to $+\infty$, we find by dominated convergence for every $t \in (0, T_0)$,

$$\|\mathbf{u}(t,.)\|_{L^{2}(\Phi dx)}^{2} + 2\int_{0}^{t} \|\nabla \otimes \mathbf{u}(s,.)\|_{L^{2}(\Phi dx)}^{2} ds$$

$$\leq \|\mathbf{u}_{0}\|_{L^{2}(\Phi dx)}^{2} - \int_{0}^{t} \int \nabla (|\mathbf{u}|^{2}) \cdot \nabla \Phi \, dx \, ds + \int_{0}^{t} \int (|\mathbf{u}|^{2}\mathbf{u} + 2p\mathbf{u}) \cdot \nabla \Phi \, dx \, ds$$

Letting t go to 0, we find

$$\limsup_{t \to 0} \|\mathbf{u}(t,.)\|_{L^2(\Phi dx)}^2 \le \|\mathbf{u}_0\|_{L^2(\Phi dx)}^2.$$

As **u** is weakly continuous in $L^2(\Phi dx)$, we also have

$$\|\mathbf{u}_0\|_{L^2(\Phi dx)}^2 \le \liminf_{t \to 0} \|\mathbf{u}(t,.)\|_{L^2(\Phi dx)}^2$$

This gives $\|\mathbf{u}_0\|_{L^2(\Phi dx)}^2 = \lim_{t\to 0} \|\mathbf{u}(t,.)\|_{L^2(\Phi dx)}^2$, which allows to turn the weak convergence into a strong convergence.

4.4 Global existence using a scaling argument

Let $\lambda > 0$, then \mathbf{u}_{ϵ} is a solution of the Cauchy initial value problem for the approximated Navier–Stokes equations (NS_{ϵ}) on (0, T) with initial value $\mathbf{u}_{0,\epsilon}$ if and only if $\mathbf{u}_{\epsilon,\lambda}(t,x) = \lambda \mathbf{u}_{\epsilon}(\lambda^2 t, \lambda x)$ is a solution for the approximated Navier–Stokes equations $(NS_{\lambda\epsilon})$ on $(0, T/\lambda^2)$ with initial value $\mathbf{u}_{0,\epsilon,\lambda}(x) = \lambda \mathbf{u}_{0,\epsilon}(\lambda x)$. We shall write $\mathbf{u}_{0,\lambda} = \lambda \mathbf{u}_0(\lambda x)$.

We have seen that

$$\begin{aligned} \|\sqrt{\Phi}\mathbf{u}_{\epsilon,\lambda}(t)\|_{L^2}^2 &+ \int_0^t \|\sqrt{\Phi}\nabla \otimes \mathbf{u}_{\epsilon,\lambda}\|_{L^2}^2 \\ &\leq \|\sqrt{\Phi}\mathbf{u}_{0,\epsilon,\lambda}\|_{L^2}^2 + C_{\Phi}\int_0^t \|\sqrt{\Phi}\mathbf{u}_{\epsilon,\lambda}\|_{L^2}^2 + \|\sqrt{\Phi}\mathbf{u}_{\epsilon,\lambda}\|_{L^2}^{2d} ds \end{aligned}$$

(under the extra condition, when d = 4, that $\|\sqrt{\Phi}\mathbf{u}_{\epsilon,\lambda}(t)\|_{L^2}$ remains smaller than ϵ_0).

By Lemma 4.1, we thus found that there exists a constant $C_{\Phi} \geq 1$ such that if T_{λ} satisfies

• if
$$d = 2$$
, $C_{\Phi} \left(1 + \| \mathbf{u}_{0,\lambda} \|_{L^{2}(\Phi dx)}^{2} \right) T_{\lambda} = 1$

• if
$$d = 3$$
, $C_{\Phi} \left(1 + \| \mathbf{u}_{0,\lambda} \|_{L^{2}(\Phi dx)}^{2} \right)^{2} T_{\lambda} = 1$

• if
$$d = 4$$
 and $||u_{0,\lambda}||_{L^2(\Phi \, dx)} \le \frac{1}{C_{\Phi}}, C_{\Phi} T_{\lambda} = 1$

then

$$\sup_{0 \le t \le T_{\lambda}} \| \mathbf{u}_{\epsilon,\lambda}(t,.) \|_{L^{2}(\Phi dx)}^{2} + \int_{0}^{T_{\lambda}} \| \nabla \otimes \mathbf{u}_{\epsilon,\lambda} \|_{L^{2}(\Phi dx)}^{2} ds \le C_{\Phi}(1 + \| \mathbf{u}_{0,\lambda} \|_{L^{2}(\Phi dx)}^{2}).$$

$$\tag{5}$$

It gives that the solutions \mathbf{u}_{ϵ} are controlled, uniformly in ϵ , on $(0, \lambda^2 T_{\lambda})$ since for $t \in (0, T_{\lambda})$,

$$\int |\mathbf{u}_{\epsilon,\lambda}(t,x)|^2 \Phi(x) \, dx = \int |\mathbf{u}_{\epsilon}(\lambda^2 t,y)|^2 \Phi(\frac{y}{\lambda}) \lambda^{2-d} \, dy \ge \lambda^{2-d} \int |\mathbf{u}_{\epsilon}(\lambda^2 t,x)|^2 \Phi(x) \, dx$$

and

$$\int_{0}^{T_{\lambda}} \int |\nabla \otimes \mathbf{u}_{\epsilon,\lambda}(t,x)|^{2} \Phi(x) \, dx \, dt = \int_{0}^{\lambda^{2} T_{\lambda}} \int |\nabla \otimes \mathbf{u}_{\epsilon,\lambda}(s,y)|^{2} \Phi(\frac{y}{\lambda}) \lambda^{2-d} \, dy \, ds$$
$$\geq \lambda^{2-d} \int_{0}^{\lambda^{2} T_{\lambda}} \int |\nabla \otimes \mathbf{u}_{\epsilon}(t,x)|^{2} \Phi(x) \, dx \, dt$$

$$\int_0^{T_\lambda} \int |\nabla \otimes \mathbf{u}_{\epsilon,\lambda}(t,x)|^2 \Phi(x) \, dx \, dt \ge C_\lambda \int_0^{\lambda^2 T_\lambda} \|\nabla \otimes \mathbf{u}_\epsilon\|_{L^2(\Phi dx)}^2 \, ds.$$

Moreover, we have $\lim_{\lambda \to +\infty} \|\mathbf{u}_{0,\lambda}\|_{L^2(\Phi \, dx)} = 0$ when d = 4 and $\lim_{\lambda \to +\infty} \lambda^2 T_{\lambda} = +\infty$ when $2 \leq d \leq 4$. Indeed, we have

$$\int \lambda^2 |\mathbf{u}_0(\lambda x)|^2 \Phi(x) \, dx = \lambda^{2-d} \int |\mathbf{u}_0(x)|^2 \Phi(\frac{x}{\lambda}) \, dx = \lambda^{4-d} \int |\mathbf{u}_0(x)|^2 \frac{\Phi(\frac{x}{\lambda})}{\lambda^2 \Phi(x)} \Phi(x) \, dx$$

Since $\frac{\Phi(\frac{x}{\lambda})}{\lambda^2 \Phi(x)} \leq \min\{C_2, \frac{1}{\lambda^2 \Phi(x)}\}$ by hypothesis (**H**4), we find by dominated convergence that $\|\mathbf{u}_{0,\lambda}\|_{L^2(\Phi dx)} = o(\lambda^{\frac{4-d}{2}})$ and thus $\lim_{\lambda \to +\infty} \lambda^2 T_{\lambda} = +\infty$.

Thus, if we consider a finite time T and a sequence ϵ_k , we may choose λ such that $\lambda^2 T_{\lambda} > T$ (and such that $\|\mathbf{u}_{0,\lambda}\|_{L^2(\Phi dx)} < \epsilon_0$ if d = 4); we have a uniform control of $\mathbf{u}_{\epsilon,\lambda}$ and of $\nabla \otimes \mathbf{u}_{\epsilon,\lambda}$ on $(0, T_{\lambda})$, hence a uniform control of \mathbf{u}_{ϵ} and of $\nabla \otimes \mathbf{u}_{\epsilon,\lambda}$ on $(0, T_{\lambda})$, hence a uniform control of \mathbf{u}_{ϵ} and of $\nabla \otimes \mathbf{u}_{\epsilon}$ on (0, T). We may exhibit a solution on (0, T) using the Rellich-Lions theorem by extracting a subsequence ϵ_{k_n} . A diagonal argument permits then to obtain a global solution.

 \diamond

Theorem 1 is proved.

5 Proof of Theorem 2 (the case d = 2).

In the case of dimension d = 2, the Navier–Stokes equations are well-posed in H^1 and we don't need to mollify the equations. Thus, we may approximate the Navier–Stokes equations with

$$(NS_{\epsilon}) \begin{cases} \partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} - (\mathbf{u}_{\epsilon} \cdot \nabla) \mathbf{u}_{\epsilon} - \nabla p_{\epsilon} \\ \nabla \cdot \mathbf{u}_{\epsilon} = 0, \qquad \mathbf{u}_{\epsilon}, (0, .) = \mathbf{u}_{0, \epsilon} \end{cases}$$

with

$$\mathbf{u}_{0,\epsilon} = \mathbb{P}(\phi_{\epsilon}\mathbf{u}_0).$$

Then the vorticity ω_{ϵ} is solution of

$$\begin{cases} \partial_t \omega_{\epsilon} = \Delta \omega_{\epsilon} - (\mathbf{u}_{\epsilon} \cdot \nabla) \omega_{\epsilon} \\ \nabla \cdot \omega_{\epsilon} = 0, \qquad \omega_{\epsilon}, (0, .) = \omega_{0, \epsilon} \end{cases}$$

with

$$\omega_{0,\epsilon} = \nabla \wedge (\phi_{\epsilon} \mathbf{u}_0).$$

 $\mathbf{u}_{0,\epsilon}$ belongs to H^1 , so we know that we have a global solution \mathbf{u}_{ϵ} . We then just have to prove that, for every finite time T_0 , we have a uniform control of the norms $\|\omega_{\epsilon}\|_{L^{\infty}((0,T_0),L^2(\Phi dx))}$ and $\|\nabla \omega_{\epsilon}\|_{L^2((0,T_0),L^2(\Phi dx))}$. We can calculate $\int \partial_t \omega_\epsilon \cdot \omega_\epsilon \Phi \, dx$ so that

$$\int \frac{|\omega_{\epsilon}(t,x)|^2}{2} \Phi \, dx + \int_0^t \int |\nabla \omega_{\epsilon}|^2 \, \Phi \, dx \, ds$$

=
$$\int \frac{|\omega_{0,\epsilon}(x)|^2}{2} \Phi \, dx - \int_0^t \int \nabla (\frac{|\omega_{\epsilon}|^2}{2}) \, \cdot \nabla \Phi \, dx \, ds$$

+
$$\int_0^t \int \frac{|\omega_{\epsilon}|^2}{2} \mathbf{u}_{\epsilon} \cdot \nabla \Phi \, dx \, ds.$$

As

$$\begin{aligned} \int_0^t \int \frac{|\omega_{\epsilon}|^2}{2} \mathbf{u}_{\epsilon} \cdot \nabla \Phi \, dx \, ds &\leq \int_0^t \|\sqrt{\Phi}\omega_{\epsilon}\|_{L^{\frac{8}{3}}}^2 \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^4} \\ &\leq \int_0^t (\|\sqrt{\Phi}\omega_{\epsilon}\|_{L^2}^{3/4} \|\nabla(\sqrt{\Phi}\omega_{\epsilon})\|_{L^2}^{1/4})^2 \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^4} \end{aligned}$$

we obtain

$$\|\sqrt{\Phi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} + \int_{0}^{t} \|\sqrt{\Phi}\nabla\omega_{\epsilon}\|_{L^{2}}^{2} \leq \|\sqrt{\Phi}\omega_{0,\epsilon}\|_{L^{2}}^{2} + C_{\Phi}\int_{0}^{t} \|\sqrt{\Phi}\omega_{\epsilon}\|_{L^{2}}^{2} (1 + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{4}}^{4}) \, ds$$

 \diamond

We can conclude that, for all T > 0 and for all $t \in (0, T)$,

$$\|\sqrt{\Phi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} + \int_{0}^{t} \|\sqrt{\Phi}\nabla\omega_{\epsilon}\|_{L^{2}}^{2} \leq \|\sqrt{\Phi}\omega_{0,\epsilon}\|_{L^{2}}^{2} e^{C_{\Phi}\sup_{\epsilon>0}\int_{0}^{t}(1+\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{4}})^{\frac{4}{3}} ds$$

Thus, we have uniform controls on (0, T).

Proof of Theorems 3 and 4 (the axisym-6 metric case)

Axisymmetry. 6.1

In \mathbb{R}^3 , we consider the usual coordinates (x_1, x_2, x_3) and the cylindrical coordinates (r, θ, z) given by the formulas $x_1 = r \cos \theta$, $x_2 = r \sin \theta$ and $x_3 = z$.

We denote $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ the usual canonical basis

$$\mathbf{e}_1 = (1, 0, 0), \ \mathbf{e}_2 = (0, 1, 0), \ \mathbf{e}_3 = (0, 0, 1).$$

We attach to the point x (with $r \neq 0$) another orthonormal basis

$$\mathbf{e}_r = \frac{\partial x}{\partial r} = \cos\theta \,\mathbf{e}_1 + \sin\theta \,\mathbf{e}_2, \ \mathbf{e}_\theta = \frac{1}{r} \frac{\partial x}{\partial \theta} = -\sin\theta \,\mathbf{e}_1 + \cos\theta \,\mathbf{e}_2, \ \mathbf{e}_z = \frac{\partial x}{\partial z} = \mathbf{e}_3.$$

For a vector field $\mathbf{u} = (u_1, u_2, u_3) = u_1 \mathbf{e}_1 + u_2 \mathbf{e}_2 + u_3 \mathbf{e}_3$, we can see that

$$\mathbf{u} = (u_1 \cos \theta + u_2 \sin \theta) \mathbf{e}_r + (-u_1 \sin \theta + u_2 \cos \theta) \mathbf{e}_\theta + u_3 \mathbf{e}_z.$$

We will denote $(u_r, u_\theta, u_z)_p$ the coordinates of **u** in the basis $(\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{e}_z)$. We will consider axially symmetric (axisymmetric) vector fields **u** without swirl and axisymmetric scalar functions a, which means that

$$\mathbf{u} = u_r(r, z) \mathbf{e}_r + u_z(r, z) \mathbf{e}_z$$
 and $a = a(r, z)$.

6.2 The H^1 case.

We will use the following well known results of Ladyzhensaya [5, 6].

Proposition 6.1 Let u_0 be a divergence free axisymmetric vector field without swirl, such that u_0 belongs to H^1 . Then, the following problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

has a unique solution $\mathbf{u} \in \mathcal{C}([0, +\infty), H^1)$. This solution is axisymmetric without swirl. Moreover, $\mathbf{u}, \nabla \otimes \mathbf{u}$ belong to $L^{\infty}((0, +\infty), L^2)$, and $\nabla \otimes \mathbf{u}, \Delta \mathbf{u}$ belong to $L^2((0, +\infty), L^2)$.

If $u_0 \in H^2$, we have the inequality

$$\int \frac{|\omega(t)|^2}{r^2} dx \le \int \frac{|\omega_0|^2}{r^2} \le \|\nabla \otimes \omega_0\|_2^2.$$

6.3 A priori controls

Let $\phi \in \mathcal{D}(\mathbb{R}^2)$ be a real-valued radial function which is equal to 1 in a neighborhood of 0 and let $\phi_{\epsilon}(x) = \phi(\epsilon(x_1, x_2))$. For $\epsilon \in (0, 1]$, let

$$\mathbf{u}_{0,\epsilon} = \mathbb{P}(\phi_{\epsilon}\mathbf{u}_0).$$

Thus, $\mathbf{u}_{0,\epsilon}$ is a divergence free axisymmetric without swirl vector field which belongs to H^1 . As we have

$$\omega_{0,\epsilon} = \nabla \wedge \mathbf{u}_{0,\epsilon} = \nabla \wedge (\phi_{\epsilon} \mathbf{u}_{0}) = \phi_{\epsilon} \omega_{0} + \epsilon (\nabla \phi)(\epsilon x) \wedge \mathbf{u}_{0},$$

using $\Phi \in \mathcal{A}_2$ and $|\epsilon \nabla \phi(\epsilon x)| \leq C \frac{1}{r} \mathbb{1}_{r \geq \frac{1}{C\epsilon}} \leq C' \mathbb{1}_{r \geq \frac{1}{C\epsilon}} \sqrt{\Phi}$, we can see that

$$\lim_{\varepsilon \to 0} \|\mathbf{u}_0 - \mathbf{u}_{0,\epsilon}\|_{L^2(\Phi \, dx)} + \|\omega_0 - \omega_{0,\epsilon}\|_{L^2(\Psi \, dx)} = 0$$

Let \mathbf{u}_ϵ be the global solution of the problem

$$(NS_{\epsilon}) \begin{cases} \partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} - (\mathbf{u}_{\epsilon} \cdot \nabla) \mathbf{u}_{\epsilon} - \nabla p_{\epsilon} \\ \nabla \cdot \mathbf{u}_{\epsilon} = 0, \qquad \mathbf{u}_{\epsilon}(0, .) = \mathbf{u}_{0, \epsilon} \end{cases}$$

given by the Proposition 6.1. We denote $\omega_{\epsilon} = \nabla \wedge \mathbf{u}_{\epsilon}$, then

$$\partial_t \mathbf{u}_{\epsilon} = \Delta \mathbf{u}_{\epsilon} + (\mathbf{u}_{\epsilon} \cdot \nabla) \mathbf{u}_{\epsilon} - \nabla p_{\epsilon}$$
(6)

and

$$\partial_t \omega_\epsilon = \Delta \omega_\epsilon + (\omega_\epsilon \cdot \nabla) \mathbf{u}_\epsilon - (\mathbf{u}_\epsilon \cdot \nabla) \omega_\epsilon \tag{7}$$

As $\sqrt{\Psi}\omega_{\epsilon} \in L^{2}H^{1}$ (because $\sqrt{\Psi}, \nabla\sqrt{\Psi} \in L^{\infty}$) and $\sqrt{\Psi}\partial_{t}\omega_{\epsilon} \in L^{2}H^{-1}$, we can calculate $\int \partial_{t}\omega_{\epsilon} \cdot \omega_{\epsilon}\Psi dx$ using (7) so that

$$\begin{split} \int \frac{|\omega_{\epsilon}(t,x)|^2}{2} \Psi \, dx + \int_0^t \int |\nabla \otimes \omega_{\epsilon}|^2 \Psi dx \, ds \\ &= \int \frac{|\omega_{0,\epsilon}(x)|^2}{2} \Psi \, dx - \int_0^t \int \nabla (\frac{|\omega_{\epsilon}|^2}{2}) \cdot \nabla \Psi dx \, ds \\ &+ \int_0^t \int \frac{|\omega_{\epsilon}|^2}{2} \mathbf{u}_{\epsilon} \cdot \nabla \Psi - (\omega_{\epsilon} \cdot \mathbf{u}_{\epsilon}) \omega_{\epsilon} \cdot \nabla \Psi \, dx \\ &- \int_0^t \int ((\omega_{\epsilon} \cdot \nabla) \omega_{\epsilon}) \cdot \mathbf{u}_{\epsilon} \Psi \, dx \, ds \\ &\leq \int \frac{|\omega_{0,\epsilon}(x)|^2}{2} \Psi \, dx + \frac{1}{8} \int_0^t \int |\nabla \otimes \omega_{\epsilon}|^2 \Psi \, dx \, ds + C \int_0^t \|\sqrt{\Psi} \, \omega_{\epsilon}\|_2^2 \, ds \\ &+ C \int_0^t \|\sqrt{\Psi} \, \omega_{\epsilon}\|_2 \|\sqrt{\Psi} \, \omega_{\epsilon}\|_6 \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_3 \, ds \\ &- \int_0^t \int ((\omega_{\epsilon} \cdot \nabla) \omega_{\epsilon}) \cdot \mathbf{u}_{\epsilon} \Psi \, dx \, ds \\ &\leq \int \frac{|\omega_{0,\epsilon}(x)|^2}{2} \Psi \, dx + \frac{1}{4} \int_0^t \int |\nabla \otimes \omega_{\epsilon}|^2 \Psi \, dx \, ds + C \int_0^t \|\sqrt{\Psi} \, \omega_{\epsilon}\|_2^2 \, ds \\ &+ C' \int_0^t \|\sqrt{\Psi} \, \omega_{\epsilon}\|_2^2 (\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_3 + (\|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_3^4) \, ds \\ &- \int_0^t \int ((\omega_{\epsilon} \cdot \nabla) \omega_{\epsilon}) \cdot \mathbf{u}_{\epsilon} \Psi \, dx \, ds \end{split}$$

As $\omega_{\epsilon} = \omega_{\epsilon,\theta} \mathbf{e}_{\theta}$, we have

$$\omega_{\epsilon} \cdot \nabla \omega_{\epsilon} = -\frac{\omega_{\epsilon,\theta}^2}{r} \,\mathbf{e}_r.$$

In order to control $\mathbf{u}_{\epsilon} \cdot (\omega_{\epsilon} \cdot \nabla \omega_{\epsilon})$, we split the domain of integration in a domain where r is small and a domain where r is large. The support of ϕ_1 is contained in $\{x \mid r < R\}$ for some $R > 0\}$, and the support of $1 - \phi_1$ is contained in $\{x \mid r > R_0\}$ for some $R_0 > 0\}$. We have

$$\inf_{r < R} \Phi(x) = \inf_{\sqrt{x_1^2 + x_2^2} < R} \Phi(x_1, x_2, 0) > 0$$

and similarly

$$\inf_{r < R} \Psi(x) = \inf_{\sqrt{x_1^2 + x_2^2} < R} \Psi(x_1, x_2, 0) > 0.$$

On the other hand, we have

$$\inf_{r>R_0} r^2 \Phi(x) = \inf_{\sqrt{x_1^2 + x_2^2} > R_0} (x_1^2 + x_2)^2 \Phi(x_1, x_2, 0) \ge \inf_{|x| > R_0} |x|^2 \Phi(x) > 0.$$

We then write :

$$\begin{split} &-\int_0^t \int ((\omega_{\epsilon} \cdot \nabla)\omega_{\epsilon}) \cdot \mathbf{u}_{\epsilon} \ \Psi \, dx \, ds \\ &= \int_0^t \int \phi_1((\omega_{\epsilon} \cdot \nabla)\mathbf{u}_{\epsilon}) \cdot \omega_{\epsilon}) \Psi \, dx \, ds + \int_0^t \int (\omega_{\epsilon} \cdot \mathbf{u}_{\epsilon})(\omega_{\epsilon} \cdot \nabla\phi_1) \Psi \, dx \, ds \\ &+ \int_0^t \int \phi_1(\omega_{\epsilon} \cdot \mathbf{u}_{\epsilon})\omega_{\epsilon} \cdot \nabla\Psi \, dx \, ds \\ &- \int_0^t \int (1 - \phi_1)(\mathbf{u}_{\epsilon} \cdot (\omega_{\epsilon} \cdot \nabla\omega_{\epsilon})) \Psi \, dx \, ds \\ &\leq C \int_0^t \int |\omega_{\epsilon}|^2 |\nabla \otimes \mathbf{u}_{\epsilon}| \Psi^{3/2} \, dx \, ds + C \int_0^t \int |\omega_{\epsilon}|^2 |\mathbf{u}_{\epsilon}| \sqrt{\Phi} \, \Psi \, dx \, ds. \end{split}$$

As $\Psi \in \mathcal{A}_2$, we have $\|\sqrt{\Psi}\nabla \otimes \mathbf{u}_{\epsilon}\|_2 \approx \|\sqrt{\Psi}\omega_{\epsilon}\|_2$; moreover,

$$\|\nabla \otimes (\sqrt{\Phi}\mathbf{u}_{\epsilon})\|_{2} \leq C(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2} + \|\sqrt{\Psi}\omega_{\epsilon}\|_{2})$$

and

$$\|\nabla \otimes (\sqrt{\Psi}\omega_{\epsilon})\|_{2} \leq C(\|\sqrt{\Psi}\omega_{\epsilon}\|_{2} + \|\sqrt{\Psi}\nabla \otimes \omega_{\epsilon}\|_{2}),$$

and thus we get

$$\begin{split} &-\int_{0}^{t}\int ((\omega_{\epsilon}\cdot\nabla)\omega_{\epsilon})\cdot\mathbf{u}_{\epsilon} \Psi\,dx\,ds\\ &\leq C\int_{0}^{t}\|\sqrt{\Psi}\nabla\otimes\mathbf{u}_{\epsilon}\|_{L^{2}}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{3}}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{6}}\,ds+C\int_{0}^{t}\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{6}}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{3}}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}\,ds\\ &\leq C'\int_{0}^{t}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}^{\frac{3}{2}}(\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}+\|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{L^{2}})^{\frac{3}{2}}\,ds\\ &+C'\int_{0}^{t}\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}^{\frac{3}{2}}(\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}+\|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{L^{2}})^{\frac{1}{2}}\,ds\\ &\leq C''\int_{0}^{t}(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}+\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{\frac{3}{2}})\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{2}+\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{3}+\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{6}\,ds\\ &+\frac{1}{8}\int_{0}^{t}\|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{2}^{2}\,ds\end{split}$$

We finally find that

$$\begin{aligned} \|\sqrt{\Psi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} + \int_{0}^{t} \|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{L^{2}}^{2} ds \\ &\leq \|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}}^{2} + C \int (1 + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{3} + (\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{3}^{4/3})\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{2} ds \\ &+ C \int_{0}^{t} (\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{4/3})\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{2} + \|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{3} + \|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{6} ds \end{aligned}$$
(8)
$$&\leq \|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}}^{2} \\ &+ C' \int_{0}^{t} (1 + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{2}^{4/3})\|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{2} + \|\sqrt{\Psi}\omega_{\epsilon}\|_{2}^{6} ds \end{aligned}$$

We already know that $\|\sqrt{\Phi}\mathbf{u}_{\epsilon}(t)\|_{L^2}$ remains bounded (independently of ϵ) on every bounded interval, so that we may again use Lemma 4.1 and control $\sup_{0 \le t \le T_0} \|\omega_{\epsilon}(t,.)\|_{L^2(\Psi dx)}^2 + \int_0^{T_0} \|\nabla \omega\|_{L^2(\Psi dx)}^2 ds$ for some T_0 , where both T_0 and the control don't depend on ϵ . The control is then transferred to the limit ω since $\omega = \lim \omega_{\epsilon_k} = \lim \nabla \wedge \mathbf{u}_{\epsilon_k}$. This proves local existence of a regular solution and Theorem 3 is proved.

6.4 The case of a very regular initial value.

We present a result apparently more restrictive that our main Theorem (Theorem 4), but we will see that it implies almost directly our main Theorem. **Proposition 6.2** Let Φ be a weight satisfying $(\mathbf{H}_1) - (\mathbf{H}_4)$. Assume moreover that Φ depends only on $r = \sqrt{x_1^2 + x_2^2}$. Let Ψ be another continuous weight (that depends only on r) such that $\Phi \leq \Psi \leq 1$, $\Psi \in \mathcal{A}_2$ and there exists $C_1 > 0$ such that

$$|\nabla \Psi| \leq C_1 \sqrt{\Phi \Psi} \text{ and } |\Delta \Psi| \leq C_1 \Phi \Psi.$$

Let \mathbf{u}_0 be a divergence free axisymmetric vector field without swirl, such that \mathbf{u}_0 , belongs to $L^2(\Phi dx)$, $\nabla \otimes \mathbf{u}_0$ and $\Delta \mathbf{u}_0$ belong to $L^2(\Psi dx)$. Then there exists a global solution \mathbf{u} of the problem

$$(NS) \begin{cases} \partial_t \boldsymbol{u} = \Delta \boldsymbol{u} - (\boldsymbol{u} \cdot \nabla) \boldsymbol{u} - \nabla p \\ \nabla \cdot \boldsymbol{u} = 0, \qquad \boldsymbol{u}(0, .) = \boldsymbol{u}_0 \end{cases}$$

such that

- u is axisymmetric without swirl, u belongs to L[∞]((0, T), L²(Φ dx)), ∇⊗ u belong to L[∞]((0, T), L²(Ψ dx)) and Δu belongs to L²((0, T), L²(Ψ dx)), for all T > 0,
- the maps $t \in [0, +\infty) \mapsto \boldsymbol{u}(t, .)$ and $t \in [0, +\infty) \mapsto \nabla \otimes \boldsymbol{u}(t, .)$ are weakly continuous from $[0, +\infty)$ to $L^2(\Phi dx)$ and to $L^2(\Psi dx)$ respectively, and are strongly continuous at t = 0,

Proof.

Ladyzenskaya's inequality for axisymmetric fields with no swirl (Proposition 6.1) gives

$$\int \frac{|\omega_{\epsilon}(t)|^2}{r^2} dx \le \int \frac{|\omega_{0,\epsilon}|^2}{r^2} dx.$$
(9)

As we have

$$\partial_i \omega_{0,\epsilon} = \phi_{\epsilon} \partial_i \omega_0 + \epsilon \partial_i \phi(\epsilon x) \omega_0 + \epsilon (\nabla \phi)(\epsilon x) \wedge \partial_i \mathbf{u}_0 + \epsilon^2 (\nabla \partial_i \phi)(\epsilon x) \wedge \mathbf{u}_0,$$

we can see that

$$\lim_{\epsilon \to 0} \|\nabla \otimes \omega_{0,\epsilon} - \nabla \otimes \omega_0\|_{L^2(\Psi \, dx)} = 0.$$

As

$$\int \frac{|\omega_{0,\epsilon} - \omega_0|^2}{r^2} dx \le C(\int_{0 < r < 1} |\nabla \otimes \omega_{0,\epsilon} - \nabla \otimes \omega_0|^2 \Psi \, dx + \int_{1 < r < +\infty} |\omega_{0,\epsilon} - \omega_0|^2 \Psi \, dx),$$

we also have

$$\lim_{\epsilon \to 0} \int \frac{|\omega_{0,\epsilon} - \omega_0|^2}{r^2} dx = 0.$$

We know that

$$\int \frac{|\omega_{\epsilon}(t,x)|^{2}}{2} \Psi \, dx + \int_{0}^{t} \int |\nabla \otimes \omega_{\epsilon}|^{2} \Psi \, dx \, ds$$

=
$$\int \frac{|\omega_{0,\epsilon}(x)|^{2}}{2} \Psi \, dx - \int_{0}^{t} \int \nabla (\frac{|\omega_{\epsilon}|^{2}}{2}) \cdot \nabla \Psi \, dx \, ds$$

+
$$\int_{0}^{t} \int \frac{|\omega_{\epsilon}|^{2}}{2} \mathbf{u}_{\epsilon} \cdot \nabla \Psi \, dx \, ds$$

-
$$\int_{0}^{t} \int (\omega_{\epsilon} \cdot \mathbf{u}_{\epsilon}) \omega_{\epsilon} \cdot \nabla \Psi \, dx \, ds - \int_{0}^{t} \int \mathbf{u}_{\epsilon} (\omega_{\epsilon} \cdot \nabla \omega_{\epsilon}) \Psi \, dx \, ds$$

which implies

$$\begin{aligned} \|\sqrt{\Psi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} + 2\int_{0}^{t} \|\sqrt{\Psi}\nabla\omega_{\epsilon}\|_{L^{2}}^{2} \\ &\leq \|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}}^{2} + 2\int_{0}^{t} \|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}^{2} \|\sqrt{\Psi}\nabla\omega_{\epsilon}\|_{L^{2}}^{2} \\ &+ \int_{0}^{t} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{3}}^{2} \|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{3}}^{2} \\ &+ \int_{0}^{t} \frac{1}{r} |\mathbf{u}_{r,\epsilon}| |\omega_{\epsilon}|^{2} \Psi \, dx \, ds \end{aligned}$$

Furthermore, we have

$$\int_0^t \int \frac{1 - \phi_1(x)}{r} |\mathbf{u}_{r,\epsilon}| |\omega_{\epsilon}|^2 \Psi \, dx \, ds \le \int_0^t \|\sqrt{\Phi} \mathbf{u}_{\epsilon}\|_{L^3} \|\sqrt{\Psi} \omega_{\epsilon}\|_{L^3}^2$$

and

$$\int_0^t \int \frac{\phi_1(x)}{r} |\mathbf{u}_{\epsilon,r}| |\omega_{\epsilon}|^2 dx \, ds \le C \int_0^t \left\| \frac{\omega_{\epsilon}}{r} \right\|_{L^2} \|\sqrt{\Psi} \mathbf{u}_{\epsilon}\|_{L^{\infty}} \|\sqrt{\Psi} \omega_{\epsilon}\|_{L^2},$$

where

$$\begin{aligned} \|\frac{\omega_{\epsilon}}{r}\|_{L^{2}} &\leq C \|\frac{\omega_{0,\epsilon}}{r}\|_{L^{2}} \leq C(\|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}} + \|\sqrt{\Psi}\nabla\otimes\omega_{0,\epsilon}\|_{L^{2}}) \\ &\leq C'(\|\sqrt{\Phi}\mathbf{u}_{0}\|_{L^{2}} + \|\sqrt{\Psi}\omega_{0}\|_{L^{2}} + \|\sqrt{\Psi}\nabla\otimes\omega_{0}\|_{L^{2}}) \end{aligned}$$

and

$$\|\sqrt{\Psi}\mathbf{u}_{\epsilon}\|_{L^{\infty}}^{2} \leq C\|\nabla\otimes(\sqrt{\Psi}\mathbf{u}_{\epsilon})\|_{2}\|\Delta(\sqrt{\Psi}\mathbf{u}_{\epsilon})\|_{2} \leq C'(\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}+\|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}+\|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{L^{2}})^{2}.$$

Then, if we denote $A_0 = \|\sqrt{\Phi}\mathbf{u}_0\|_{L^2} + \|\sqrt{\Psi}\omega_0\|_{L^2} + \|\sqrt{\Psi}\nabla \otimes \omega_0\|_{L^2}$, we have

$$\begin{aligned} \|\sqrt{\Psi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} &+ \int_{0}^{t} \|\sqrt{\Psi}\nabla\otimes\omega_{\epsilon}\|_{L^{2}}^{2} \\ &\leq \|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}}^{2} + C\int_{0}^{t} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}^{2} \\ &+ C_{\Phi}\int_{0}^{t} \|\sqrt{\Psi}\omega_{\epsilon}\|_{L^{2}}^{2} (1 + A_{0} + A_{0}^{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{3}}^{2} + \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{3}}^{2}) \, ds \end{aligned}$$

We can then conclude that, for all T > 0 and for all $t \in (0, T)$,

$$\begin{aligned} \|\sqrt{\Psi}\omega_{\epsilon}(t)\|_{L^{2}}^{2} &+ \int_{0}^{t} \|\sqrt{\Psi}\nabla \otimes \omega_{\epsilon}\|_{L^{2}}^{2} \\ \leq &(\|\sqrt{\Psi}\omega_{0,\epsilon}\|_{L^{2}}^{2} + C_{\Phi}\sup_{\epsilon>0}\int_{0}^{T} \|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{2}}^{2})e^{C_{\Phi}\sup_{\epsilon>0}\int_{0}^{t}(1+A_{0}^{2}+\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{3}}^{2}+\|\sqrt{\Phi}\mathbf{u}_{\epsilon}\|_{L^{3}}^{2})ds \end{aligned}$$

Then, we can obtain a solution on (0, T) using the Aubin–Lions Theorem and finish with a diagonal argument to get a global solution. \diamond

6.5 End of the proof.

We begin by consider a local solution \mathbf{v} on $(0, T_0)$ with initial value \mathbf{u}_0 given by Theorem 3, which is continuous from $(0, T_0)$ to \mathcal{D}' . We take $T_1 \in (0, T_0)$ such that $\nabla \otimes (\nabla \wedge \mathbf{v})(T_1, .) \in L^2(\Phi dx)$. We consider a solution \mathbf{w} on $(T_1, +\infty)$ and initial value $\mathbf{v}(T_1)$ given by Proposition 6.2. Our global solution is defined as $\mathbf{u} = \mathbf{v}$ on $(0, T_1)$ and $\mathbf{u} = \mathbf{w}$ on $(T_1, +\infty)$.

References

- F. Boyer and P. Fabrie, Mathematical Tools for the Study of the Incompressible Navier-Stokes Equations and Related Models, Applied Mathematical Sciences book series 183, Springer, USA, 2012.
- [2] Z. Bradshaw, I. Kukavica and T.P. Tsai, Existence of global weak solutions to the Navier-Stokes equations in weighted spaces, arXiv:1910.06929v1
- [3] P.G Fernández-Dalgo and P.G. Lemarié-Rieusset, Weak solutions for Navier-Stokes equations with initial data in weighted L² spaces. Arch Rational Mech Anal 237, 347–382 (2020).

- [4] L. Grafakos, Modern Fourier Analysis (2nd ed.), Graduate Texts in Mathematics 250, Springer, USA, 2009.
- [5] O. Ladyzhenskaya, Unique solvability in the large of a threedimensional Cauchy problem for the Navier-Stokes equations in the presence of axial symmetry, Zap. Nauchn. Sem. LOMI 7 (1968), 155– 177.
- [6] P.G. Lemarié-Rieusset, *The Navier-Stokes problem in the 21st century*, CRC Press, Boca Raton, FL, 2016.
- [7] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, Acta Math. 63 (1934), 193-248.