



**HAL**  
open science

# Decoding Dishevelled-Mediated Wnt Signaling in Vertebrate Early Development

De-Li Shi

► **To cite this version:**

De-Li Shi. Decoding Dishevelled-Mediated Wnt Signaling in Vertebrate Early Development. *Frontiers in Cell and Developmental Biology*, 2020, 8, 10.3389/fcell.2020.588370 . hal-02955701

**HAL Id: hal-02955701**

**<https://hal.science/hal-02955701>**

Submitted on 2 Oct 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

**Decoding Dishevelled-mediated Wnt signaling in vertebrate early development**

De-Li Shi

Developmental Biology Laboratory, CNRS-UMR7622, IBPS, Sorbonne University, 75005 Paris,  
France

Correspondence

Email: [de-li.shi@upmc.fr](mailto:de-li.shi@upmc.fr)

Tel: (33) 1 44272772; Fax: (33) 1 44273445

Short title: Dishevelled specificity in development

## **Abstract**

Dishevelled proteins are key players of Wnt signaling pathways. They transduce Wnt signals and perform cellular functions through distinct conserved domains. Due to the presence of multiple paralogs, the abundant accumulation of maternal transcripts, and the activation of distinct Wnt pathways, their regulatory roles during vertebrate early development and the mechanism by which they dictate the pathway specificity have been enigmatic and attracted much attention in the past decades. Extensive studies in different animal models have provided significant insights into the structure-function relationship of conserved Dishevelled domains in Wnt signaling and the implications of Dishevelled isoforms in early developmental processes. Notably, intra- and inter-molecular interactions and Dishevelled dosage may be important in modulating the specificity of Wnt signaling. There are also distinct and redundant functions among Dishevelled isoforms in development and disease, which may result from differential spatiotemporal expression patterns and biochemical properties and post-translational modifications. This review presents the advances and perspectives in understanding Dishevelled-mediated Wnt signaling during gastrulation and neurulation in vertebrate early embryos.

Key words: Wnt signaling; Dishevelled; Wnt/ $\beta$ -catenin; Wnt/PCP; embryonic axis; gastrulation; neurulation; convergence and extension; morphogenetic movements; mouse, *Xenopus*; zebrafish

## Introduction

Wnt signaling plays critical roles in a wide variety of biological process, including embryonic axis formation, cell proliferation, differentiation and migration, polarity establishment, and stem cell self-renewal (Steinhart and Angers, 2018; Wiese et al., 2018). Upon stimulation by Wnt ligands, membrane receptors (Frizzled) and co-receptors (LRP5/6, glypican-3/4, ROR, and RYK) assemble into complexes to activate divergent pathways (Niehrs, 2012; Green et al., 2014; Stricker et al., 2017). The activity of Frizzled receptors is further controlled by an auxiliary regulatory system involving RSPO1-4, LGR4/5/6 and ZNRF3/RNF43 (Jiang and Cong, 2016; Lehoczsky and Tabin, 2018). The canonical Wnt pathway (Wnt/ $\beta$ -catenin) regulates target gene transcription through stabilization and nuclear accumulation of  $\beta$ -catenin by inhibition of its destruction complex, consisting of Axin-GSK3 $\beta$ -APC, whereas the non-canonical Wnt pathway (Wnt/planar cell polarity or Wnt/PCP) is implicated in polarized cellular orientation and asymmetric cell movements through activation of major regulators of the cytoskeleton. Aberrant signaling of both pathways leads to tumorigenesis and metastasis of multiple cancer types, as well as human birth defects (Clevers and Nusse 2012; Butler and Wallingford, 2017; Humphries and Mlodzik 2018), but how they are regulated in development and disease remains elusive.

Dishevelled (Dvl or Dsh in *Drosophila*) is a family of proteins that function as common intracellular conductors of both Wnt/ $\beta$ -catenin and Wnt/PCP pathways (Boutros and Mlodzik 1999; Wallingford and Habas 2005; Gao and Chen 2010). *Drosophila dsh* alleles were first identified in genetic mutants with disruptions of hair and bristle polarity (Wallingford and Habas 2005). Vertebrates possess three highly conserved *Dvl* genes. Extensive studies in mouse, *Xenopus* and zebrafish have revealed their critical roles in germ layer specification and morphogenetic movements, which require Wnt/ $\beta$ -catenin and Wnt/PCP signaling, respectively. There is accumulating evidence that Dvl isoforms display both distinct and redundant functions (Gentzel and Schambony, 2017). However, a number of important questions regarding Dvl-mediated Wnt signaling during development remain enigmatic (Mlodzik 2016), such as Dvl-regulated switch of distinct Wnt pathways, the specific functions of Dvl isoforms in Wnt signaling and development, the post-translational modifications of Dvl functions, and the maternal

contributions of Dvl to early developmental events. Fortunately, structure-function and mutational analyses have significantly advanced our understanding of Dvl-regulated Wnt signaling in development and disease. This review focuses on progresses made in this fascinating research field by using complementary vertebrate animal models.

### **Dvl functional domains in Wnt signaling**

Dvl proteins contain several highly conserved domains required for activating different Wnt pathways, including in particular the N-terminal DIX (Dishevelled and  $\alpha$ Xin) domain, the central PDZ (Post-synaptic density protein-95, Disc large tumor suppressor, Zonula occludens-1) domain, and the C-terminal DEP (Dishevelled, Egl-10 and Pleckstrin) domain (Fig. 1). The DIX domain is involved in Wnt-induced dynamic Dvl homo- and hetero-oligomerization that is important for Wnt/ $\beta$ -catenin signaling (Kishida et al., 1999; Kan et al., 2020; Ma et al., 2020). The PDZ domain interacts with a conserved KTxxxW motif located immediately after the seventh transmembrane domain of Frizzled receptors (Umbhauer et al., 2000; Wong et al., 2003), and with a wide variety of binding partners that function either as agonists or antagonists of Wnt signaling (Wallingford and Habas 2005; Sharma et al., 2018). Biochemical and functional analyses suggest that it participates in both Wnt/ $\beta$ -catenin and Wnt/PCP signaling (Habas et al., 2001; Lee et al., 2015). The DEP domain plays a major role in Dvl membrane recruitment by Frizzled receptors (Axelrod et al., 1998; Rothbacher et al., 2000; Wong et al., 2000; Pan et al., 2004; Park et al., 2005). It functions in the Wnt/PCP pathway either with the PDZ domain to activate Rho/ROCK or by direct interaction with Rac to trigger JNK activation. However, more recent evidence suggests that Dvl dimerization triggered by the N-terminal region of DEP domain is required for Wnt/ $\beta$ -catenin signaling (Gammons et al., 2016a, b; Paclíková et al., 2017).

The C-terminal region beyond the DEP domain also shows evolutionary conservation but displays unique features among Dvl isoforms. Specifically, Dvl3 contains histidine-single amino acid repeats required for Wnt5a-stimulated activation of nuclear factor of activated T cells (NF-AT) and possesses proline-rich domains likely involved in the interaction with other Dvl isoforms (Ma et al., 2010; Wang and Malbon 2012). The extreme 13 amino acids that are conserved in all

Dvl isoforms binds to the third intracellular loop of Frizzled receptors and stabilizes Frizzled-Dvl interaction in Wnt/ $\beta$ -catenin signaling (Tauriello et al., 2012). Moreover, the last 3 residues represent a type II PDZ-binding (PDZ-B) motif that can occupy the peptide-binding pocket of the PDZ domain, inducing Dvl to adopt a closed conformation and an auto-inhibited state (Lee et al., 2015; Qi et al., 2017). Dvl variants with an opened conformation show efficient membrane recruitment and reduced activity in Wnt/ $\beta$ -catenin signaling but display increased activity in Wnt/PCP signaling (Qi et al., 2017; Harnoř et al., 2019). The function of Dvl C-terminal region in Wnt signaling is further demonstrated in autosomal-dominant Robinow syndrome caused by de novo frameshift mutations in human *DVL1* and *DVL3* genes, which delete and replace the C-terminal region after the DEP domain (Bunn et al., 2015; White et al., 2015, 2016; Danyel et al., 2018). In vitro analysis suggests that *DVL1* lacking the C-terminal region displays reduced activity in Wnt/ $\beta$ -catenin signaling (Bunn et al., 2015). These findings suggest an importance of the C-terminus in intra- or inter-molecular interaction, which may be subjected to regulation by other partners to switch pathway specificity. Indeed, recent studies show that casein kinase 1 $\epsilon$  (CK1 $\epsilon$ ) and NIMA-related kinase 2 (NEK2) function as scaffold proteins and regulate the dynamics of Dvl conformational changes by phosphorylation of the PDZ domain and modulation of its interaction with the extreme C-terminal tail (Harnoř et al., 2019; Hanáková et al., 2019).

### **Dvl dosage effect on morphogenetic movements and cell fate specification during development in mice**

The three *Dvl* genes (*Dvl1*, *Dvl2* and *Dvl3*) in mice are broadly expressed throughout early development. Extensive analyses of mutant phenotypes have uncovered both unique and redundant functions for these genes. Mice deficient in *Dvl1* show reduced social interaction and abnormal sensorimotor gating (Lijam et al., 1997). This abnormal behavior is caused by defective Wnt/ $\beta$ -catenin signaling that may impair central nervous system functions (Belinson et al., 2016). Mice deficient in *Dvl2* exhibit more severe phenotypes, with defective cardiac morphogenesis, somite segmentation, and neural tube closure (Hamblet et al., 2002). *Dvl3* functions redundantly with *Dvl1* and *Dvl2* in several processes, including cardiac outflow tract, cochlea and neural tube

development (Etheridge et al., 2008). These works reveal a sensitivity of Wnt/PCP signaling to *Dvl* dosage because most defective phenotypes in *Dvl* mutants are related to impaired Wnt/PCP signaling, in particular the defective outflow tract morphogenesis (Sinha et al., 2012). Thus, Dvl isoforms are critically required for morphogenetic movements. Particularly, they mediate Wnt/PCP in CE movements during neurulation. Dvl2 plays a predominant role in neural tube closure, but Dvl1 and Dvl3 are also involved in this process (Wang et al., 2006). The functional importance of Dvl isoforms in neural tube formation has been confirmed by the identification of rare mutations in all three human *DVL* genes, which disturb normal functions of DVL isoforms in non-canonical Wnt signaling and cause neural tube defects (De Marco et al., 2013; Liu et al., 2020). By comparison, low levels of Dvl expression from a single allele may be sufficient to normally support those developmental processes triggered by the Wnt/ $\beta$ -catenin pathway (Soares et al., 2005; Wynshaw-Boris, 2012). However, deletion of all six *Dvl* alleles causes absence of mesoderm gene expression and mesoderm formation that are dependent on Wnt/ $\beta$ -catenin signaling (Ngo et al., 2020).

### **Distinct and redundant Dvl functions during *Xenopus* development**

Dvl function in vertebrates was first studied during *Xenopus* development. Overexpression of Dvl2 (Xdsh) in the ventral region of early embryos induced the formation of a complete secondary axis reminiscent of activation of maternal Wnt/ $\beta$ -catenin signaling (Sokol et al., 1995). However, dorsal overexpression of Xdd1, a truncated form of Dvl2 that lacks the PDZ domain and interferes with Wnt/ $\beta$ -catenin signaling triggered by Wnt ligands, did not affect dorsoventral axis formation (Sokol, 1996). Because maternal Wnt/ $\beta$ -catenin signaling is required for dorsal fate specification by activating the transcription of target genes in the Spemann organizer (Carron and Shi, 2016), the absence of an inhibitory effect by Xdd1 implies that Dvl function may be dispensable for the activation of maternal Wnt/ $\beta$ -catenin signaling. Consistently, simultaneous depletion of maternally expressed *Dvl2* and *Dvl3* from oocytes did not affect the expression of maternal Wnt/ $\beta$ -catenin target genes and the formation of dorsal axis (Tadjuidje et al., 2011). However, it is possible that low levels of Dvl proteins are still present in the oocytes due to incomplete depletion of maternal

*Dvl* mRNA. Thus, the requirement of maternal *Dvl* for dorsal axis formation in *Xenopus* requires a complete loss-of-function study. Nevertheless, a recent study suggests that activation of maternal Wnt/ $\beta$ -catenin pathway and formation of dorsal axis may be achieved through a Dvl-independent mechanism (Yan et al., 2018). During organogenesis, it seems that Dvl isoforms display less functional redundancy in developmental processes that involve Wnt/ $\beta$ -catenin signaling, which may be due to their differential expression patterns. For example, Dvl1 and Dvl2 are required for neural crest cell specification and somite segmentation, while Dvl3 maintains gene expression in the myotome (Gray et al., 2009).

Maternal Dvl2 and Dvl3 likely display distinct and redundant functions during CE movements. In these coordinated processes, lateral cells converge toward the dorsal region to narrow the germ layers, while dorsal midline cells undergo mediolateral intercalation by polarized protrusive behaviors to lengthen the embryo along the anteroposterior axis (Keller and Sutherland, 2020). Previous studies show that Dvl regulates CE movements through Wnt/PCP signaling (Tada and Smith 2000; Djiane et al., 2000; Wallingford and Harland 2001; Habas et al., 2001, 2003). Moreover, inhibition of maternal Dvl2 or Dvl3 function suggests that they exhibit a non-redundant but an additive effect on CE movements (Tadjuidje et al., 2011). More recent works reveal that Dvl1, but not Dvl2 or Dvl3, activates the Wnt/ $\text{Ca}^{2+}$  pathway (another branch of the non-canonical pathway) during CE movements (Gentzel et al., 2015), further supporting the distinct functions of Dvl isoforms in morphogenetic movements. As in mice, Dvl function is also required for neural tube closure by regulating CE movements of the midline and coordinating polarity among epithelial cells (Wallingford and Harland 2002; Seo et al., 2017). In addition, Dvl2-mediated activation of Rac1 through the DEP domain also controls the protrusive activity of neural crest cells during migration (Kratzer et al., 2020). Because cellular polarization is tightly dependent on the asymmetric activation of Wnt/PCP signaling, both reduced and increased levels of Dvl disrupts asymmetric movements. However, the cellular behaviors are completely different. Increasing the activity of Dvl perturbs cell polarity by randomizing the formation of cellular protrusions, whereas reducing the activity of Dvl prevents cellular protrusions (Wallingford et al., 2000; Cheng et al., 2017).



## **Maternal contributions of Dvl proteins to axis patterning and morphogenetic processes during zebrafish development**

The zebrafish genome contains five *Dvl* paralogs: *dvl1a*, *dvl1b*, *dvl2*, *dvl3a*, and *dvl3b*. In the early embryos, *dvl2* and *dvl3a* are maternally expressed and represent about 98% of the total pool, whereas the transcript levels of the other *Dvl* genes are negligible (Harvey et al., 2013). Knockout of *Dvl* genes reveal both distinct and redundant functions in embryonic axis specification and morphogenetic movements (Xing et al., 2018). Maternal-zygotic mutants for *dvl1a*, *dvl1b*, *dvl3a*, and *dvl3b* are phenotypically normal and fertile. In sharp contrast, maternal-zygotic *dvl2* mutants display strongly impaired CE movements during gastrulation and develop severe craniofacial defects with a “bulldog” facial phenotype, reminiscent of impaired Wnt/PCP signaling in midline structures (Kimmel et al., 2001). Zygotic *dvl2* mutant embryos are essentially normal, but only about half of them can survive to adulthood, and all male individuals show absence of courtship behavior. This suggests that there may be defects in central nervous system functions as mice *Dvl1* mutants (Lijam et al., 1997). The highest level of *dvl2* expression during early development may explain at least partly the most severe phenotypes of *dvl2* mutants.

The specification of dorsal axis in zebrafish also requires maternal  $\beta$ -catenin signaling to trigger the expression of organizer genes (Kelly et al., 2000; Bellipanni et al., 2006; Fuentes et al., 2020). However, the involvement of upstream regulators has not been conclusively established. Importantly, maternal-zygotic *dvl2* and *dvl3a* double mutants, which were unresponsive to stimulation by Wnt ligands, show normal specification of dorsal cell fate, suggesting that components of Wnt signaling upstream of  $\beta$ -catenin may be dispensable for its stabilization. The activation of maternal Wnt/ $\beta$ -catenin signaling independent of Dvl activity is further confirmed in zebrafish *huluwa* mutants. Maternal depletion of *huluwa* impairs Wnt/ $\beta$ -catenin signaling and causes loss of dorso-anterior structures. Mechanistically, Huluwa protein accumulates in the cell membrane at the dorsal region and functions independently of Wnt ligands and Frizzled receptors to promote tankyrase-mediated degradation of Axin, thereby stabilizing  $\beta$ -catenin (Yan et al., 2018).

Mutational analyses of Dvl functions also reveal a major contribution of maternal Dvl to zygotic events and confirm the importance of Dvl dosage in Wnt/PCP signaling (Fig. 2). This dosage effect is particularly reflected by the requirement of Dvl2 and Dvl3a for CE movements. Although Dvl2 plays a predominant role, Dvl3a exerts a strong synergistic effect on the loss of Dvl2 function, and progressive removal of Dvl2 and Dvl3a maternal or zygotic products increasingly aggravates CE defects and reduces the elongation of anteroposterior axis. Furthermore, zygotic *dv/2* and *dv/3a* double mutants only display a shortened body length. However, maternal-zygotic *dv/2* and *dv/3a* double mutants show most strongly impaired CE movements and completely lack axis extension. They also develop severe trunk and posterior deficiencies associated with down-regulation of zygotic Wnt/ $\beta$ -catenin target genes (Xing et al., 2018). Because zygotic Wnt/ $\beta$ -catenin signaling has an opposite effect with respect to maternal Wnt/ $\beta$ -catenin signaling and functions to specify the posterior region (Carron and Shi, 2016), these findings highlight the importance of maternal Dvl in setting up zygotic morphogenetic and patterning processes. They support the view that maternally expressed gene products perform essential functions after zygotic genome activation (Marlow, 2020; Solnica-Krezel 2020). Thus, both maternal and zygotic Dvl dosages are important for proper cell movements and embryonic axis patterning that occur during gastrulation.

### **Perspectives**

Dvl conserved domains in Wnt pathways have attracted much attention. Although the function of DIX, PDZ and DEP domains is relatively understood, how they cooperate to switch pathway specificity remains elusive. Detailed analysis of other conserved domains or isoform-specific regions, such as the basic region preceding the PDZ domain, the proline-rich region and histidine-single amino acid repeats in the C-terminal region beyond the DEP domain, and the extreme C-terminus, may provide insights into Dvl-mediated signal transduction. Because Dvl post-translational modifications, in particular phosphorylation and ubiquitination, and Dvl interaction partners are important for subcellular localizations and specific functions of Dvl proteins (Sharma et al., 2018; Harrison et al., 2020), it is of interest to understand how these

modulate Dvl activity and dictate signaling outcomes in key developmental processes. Indeed, dysregulation of Dvl phosphorylation impairs both Wnt/ $\beta$ -catenin and Wnt/PCP signaling during zebrafish and *Xenopus* embryogenesis (Shimizu et al., 2014; Rauschenberger et al., 2017). The specificity of Dvl isoforms also merits investigations because tissue-specific expression patterns and differential biochemical properties may contribute to their particular functions. Another intriguing question is the sensibility of Wnt/PCP signaling, but not Wnt/ $\beta$ -catenin signaling, to Dvl dosage, which is observed in all vertebrates. It suggests that Wnt/PCP-dependent developmental processes critically require Dvl function to activate downstream effectors. Consequently, moderate diminution of Dvl dosage could significantly affect polarized cellular behaviors and cell polarity. By comparison,  $\beta$ -catenin may be stabilized independently of upstream Wnt signaling components, and target genes of Wnt/ $\beta$ -catenin signaling may be regulated by other factors, at least during dorsal fate specification in zebrafish and *Xenopus* (Li et al., 2015; Yan et al., 2018). Thus, tissue morphogenesis regulated by Wnt/PCP signaling is more sensitive to Dvl dysfunction, as a result, many human congenital disorders, such as neural tube defects and Robinow syndrome, are associated with mutations in *DVL* genes.

### **Author contributions**

DLS performed the literature analysis, prepared the figures, and wrote the manuscript.

### **Conflicts of interest**

The author has no conflict of interest to declare.

### **Acknowledgments**

This work was supported by the Centre National de la Recherche Scientifique (CNRS) and the Sorbonne University.

### **References**

- Axelrod, J. D., Miller, J. R., Shulman, J. M., Moon, R. T., and Perrimon, N. (1998). Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes Dev.* 12, 2610-2622. doi: 10.1101/gad.12.16.2610
- Belinson, H., Nakatani, J., Babineau, B. A., Birnbaum, R. Y., Ellegood, J., Bershteyn, M., et al. (2016). Prenatal  $\beta$ -catenin/Brn2/Tbr2 transcriptional cascade regulates adult social and stereotypic behaviors. *Mol. Psychiatry* 21, 1417-1433. doi: 10.1038/mp.2015.207

- Bellipanni, G., Varga, M., Maegawa, S., Imai, Y., Kelly, C., Myers, A. P., et al. (2006). Essential and opposing roles of zebrafish beta-catenins in the formation of dorsal axial structures and neurectoderm. *Development* 133, 1299-1309. doi: 10.1242/dev.02295.
- Boutros, M., and Mlodzik, M. (1999). Dishevelled: at the crossroads of divergent intracellular signaling pathways. *Mech. Dev.* 83, 27-37. doi: 10.1016/s0925-4773(99)00046-5
- Bunn, K. J., Daniel, P., Rösken, H. S., O'Neill, A. C., Cameron-Christie, S. R., Morgan, T., et al. (2015). Mutations in DVL1 cause an osteosclerotic form of Robinow syndrome. *Am. J. Hum. Genet.* 96, 623-630. doi: 10.1016/j.ajhg.2015.02.010
- Butler, M. T., and Wallingford, J. B. (2017). Planar cell polarity in development and disease. *Nat. Rev. Mol. Cell Biol.* 18, 375-388. doi: 10.1038/nrm.2017.11
- Carron, C., and Shi, D. L. (2016). Specification of anteroposterior axis by combinatorial signaling during *Xenopus* development. *Wiley Interdiscip. Rev. Dev. Biol.* 5, 150-168. doi: 10.1002/wdev.217
- Cheng, X. N., Shao, M., Li, J. T., Wang, Y. F., Qi, J., Xu, Z. G., et al. (2017). Leucine repeat adaptor protein 1 interacts with Dishevelled to regulate gastrulation cell movements in zebrafish. *Nat. Commun.* 8:1353. doi: 10.1038/s41467-017-01552-x
- Clevers, H., and Nusse, R. (2012). Wnt/ $\beta$ -catenin signaling and disease. *Cell* 149, 1192-1205. doi: 10.1016/j.cell.2012.05.012
- Danyel, M., Kortüm, F., Dathe, K., Kutsche, K., and Horn, D. (2018). Autosomal dominant Robinow syndrome associated with a novel DVL3 splice mutation. *Am. J. Med. Genet. A* 176, 992-996. doi: 10.1002/ajmg.a.38635
- De Marco, P., Merello, E., Consales, A., Piatelli, G., Cama, A., Kibar, Z., et al. (2013). Genetic analysis of disheveled 2 and disheveled 3 in human neural tube defects. *J. Mol. Neurosci.* 49, 582-588. doi: 10.1007/s12031-012-9871-9
- Djiane, A., Riou, J., Umbhauer, M., Boucaut, J., and Shi, D. L. (2000). Role of frizzled 7 in the regulation of convergent extension movements during gastrulation in *Xenopus laevis*. *Development* 127, 3091-3100.
- Etheridge, S. L., Ray, S., Li, S., Hamblet, N. S., Lijam, N., Tsang, M., et al. (2008). Murine dishevelled 3 functions in redundant pathways with dishevelled 1 and 2 in normal cardiac outflow tract, cochlea, and neural tube development. *PLoS Genet.* 4:e1000259. doi: 10.1371/journal.pgen.1000259
- Fuentes, R., Tajer, B., Kobayashi, M., Pelliccia, J. L., Langdon, Y., Abrams, E. W., et al. (2020). The maternal coordinate system: Molecular-genetics of embryonic axis formation and patterning in the zebrafish. *Curr. Top. Dev. Biol.* 140, 341-389. doi: 10.1016/bs.ctdb.2020.05.002
- Gammons, M. V., Renko, M., Johnson, C. M., Rutherford, T. J., and Bienz, M. (2016a). Wnt signalosome assembly by DEP domain swapping of Dishevelled. *Mol. Cell* 64, 92-104. doi: 10.1016/j.molcel.2016.08.026
- Gammons, M. V., Rutherford, T. J., Steinhart, Z., Angers, S., and Bienz, M. (2016b). Essential role of the Dishevelled DEP domain in a Wnt-dependent human-cell-based complementation assay. *J. Cell Sci.* 129, 3892-3902. doi: 10.1242/jcs.195685
- Gao, C., and Chen, Y. G. (2010). Dishevelled: the hub of Wnt signaling. *Cell. Signal.* 22, 717-727. doi: 10.1016/j.cellsig.2009.11.021

- Gentzel, M., and Schambony, A. (2017). Dishevelled paralogs in vertebrate development: redundant or distinct? *Front. Cell Dev. Biol.* 5:59. doi: 10.3389/fcell.2017.00059
- Gentzel, M., Schille, C., Rauschenberger, V., and Schambony, A. (2015). Distinct functionality of dishevelled isoforms on Ca<sup>2+</sup>/calmodulin-dependent protein kinase 2 (CamKII) in *Xenopus* gastrulation. *Mol. Biol. Cell* 26, 966-977. doi: 10.1091/mbc.E14-06-1089
- Gray, R. S., Bayly, R. D., Green, S. A., Agarwala, S., Lowe, C. J., and Wallingford, J. B. (2009). Diversification of the expression patterns and developmental functions of the dishevelled gene family during chordate evolution. *Dev. Dyn.* 238, 2044-2057. doi: 10.1002/dvdy.22028
- Green, J., Nusse, R., and van Amerongen, R. (2014). The role of Ryk and Ror receptor tyrosine kinases in Wnt signal transduction. *Cold Spring Harb Perspect. Biol.* 6:a009175. doi: 10.1101/cshperspect.a009175
- Habas, R., Dawid, I. B., and He, X. (2003). Coactivation of Rac and Rho by Wnt/Frizzled signaling is required for vertebrate gastrulation. *Genes Dev.* 17, 295-309. doi: 10.1101/gad.1022203
- Habas, R., Kato, Y., and He, X. (2001). Wnt/Frizzled activation of Rho regulates vertebrate gastrulation and requires a novel Formin homology protein Daam1. *Cell* 107, 843-854. doi: 10.1016/s0092-8674(01)00614-6
- Hamblet, N. S., Lijam, N., Ruiz-Lozano, P., Wang, J., Yang, Y., Luo, Z., et al. (2002). Dishevelled 2 is essential for cardiac outflow tract development, somite segmentation and neural tube closure. *Development* 129, 5827-5838. doi: 10.1242/dev.00164
- Hanáková, K., Bernatík, O., Kravec, M., Micka, M., Kumar, J., Harnoš, J., et al. (2019). Comparative phosphorylation map of Dishevelled 3 links phospho-signatures to biological outputs. *Cell Commun. Signal.* 17:170. doi: 10.1186/s12964-019-0470-z
- Harnoš, J., Cañizal, M. C. A., Jurásek, M., Kumar, J., Holler, C., Schambony, A., et al. (2019). Dishevelled-3 conformation dynamics analyzed by FRET-based biosensors reveals a key role of casein kinase 1. *Nat. Commun.* 10:1804. doi: 10.1038/s41467-019-09651-7
- Harrison, C., Shao, H., Strutt, H., and Strutt, D. (2020). Molecular mechanisms mediating asymmetric subcellular localisation of the core planar polarity pathway proteins. *Biochem. Soc. Trans.* 48, 1297-1308. doi: 10.1042/BST20190404
- Harvey, S. A., Sealy, I., Kettleborough, R., Fenyes, F., White, R., Stemple, D., et al. (2013). Identification of the zebrafish maternal and paternal transcriptomes. *Development* 140, 2703-2710. doi: 10.1242/dev.095091
- Humphries, A. C., and Mlodzik, M. (2018). From instruction to output: Wnt/PCP signaling in development and cancer. *Curr. Opin. Cell Biol.* 51, 110-116. doi: 10.1016/j.ceb.2017.12.005
- Jiang, X., and Cong, F. (2016). Novel regulation of Wnt signaling at the proximal membrane level. *Trends Biochem. Sci.* 41, 773-783. doi: 10.1016/j.tibs.2016.06.003
- Kan, W., Enos, M. D., Korkmazhan, E., Muennich, S., Chen, D. H., Gammons, M. V., et al. (2020). Limited dishevelled/Axin oligomerization determines efficiency of Wnt/ $\beta$ -catenin signal transduction. *elife* 9:e55015. doi: 10.7554/eLife.55015
- Keller, R., and Sutherland, A. (2020). Convergent extension in the amphibian, *Xenopus laevis*. *Curr. Top. Dev. Biol.* 136, 271-317. doi: 10.1016/bs.ctdb.2019.11.013


- Kelly, C., Chin, A. J., Leatherman, J. L., Kozlowski, D. J., and Weinberg, E. S. (2000). Maternally controlled (beta)-catenin-mediated signaling is required for organizer formation in the zebrafish. *Development* 127, 3899-3911
- Kimmel, C. B., Miller, C. T., and Moens, C. B. (2001). Specification and morphogenesis of the zebrafish larval head skeleton. *Dev. Biol.* 233, 239-257. doi: 10.1006/dbio.2001.0201
- Kishida, S., Yamamoto, H., Hino, S., Ikeda, S., Kishida, M., and Kikuchi, A. (1999). DIX domains of Dvl and axin are necessary for protein interactions and their ability to regulate beta-catenin stability. *Mol. Cell. Biol.* 19, 4414-4422. doi: 10.1128/mcb.19.6.4414
- Kratzer, M. C., Becker, S. F. S., Grund, A., Merks, A., Harnoš, J., Bryja, V., et al. (2020). The Rho guanine nucleotide exchange factor Trio is required for neural crest cell migration and interacts with Dishevelled. *Development* 147:dev186338. doi: 10.1242/dev.186338
- Lee, H. J., Shi, D. L., and Zheng, J. J. (2015). Conformational change of Dishevelled plays a key regulatory role in the Wnt signaling pathways. *elife* 4:e08142. doi: 10.7554/eLife.08142
- Lehoczky, J. A., and Tabin, C. J. (2018). Rethinking WNT signalling. *Nature* 557, 495-496. doi: 10.1038/d41586-018-04820-y
- Li, H. Y., El Yakoubi, W., and Shi, D. L. (2015). Direct regulation of siamois by VegT is required for axis formation in *Xenopus* embryo. *Int. J. Dev. Biol.* 59, 443-451. doi: 10.1387/ijdb.150040ds
- Lijam, N., Paylor, R., McDonald, M. P., Crawley, J. N., Deng, C. X., Herrup, K., et al. (1997). Social interaction and sensorimotor gating abnormalities in mice lacking Dvl1. *Cell* 90, 895-905. doi: 10.1016/s0092-8674(00)80354-2
- Liu, L., Liu, W., Shi, Y., Gao Y., Lei Y., Finnel, R., et al. (2020). *DVL* mutations identified from human neural tube defects and Dandy-Walker malformation obstruct the Wnt signaling pathway. *J. Genet. Genomics.* doi: 10.1016/j.jgg.2020.06.003
- Ma, L., Wang, Y., Malbon, C. C., and Wang, H. Y. (2010). Dishevelled-3 C-terminal His single amino acid repeats are obligate for Wnt5a activation of non-canonical signaling. *J. Mol. Signal.* 5:19. doi: 10.1186/1750-2187-5-19
- Ma, W., Chen, M., Kang, H., Steinhart, Z., Angers, S., He, X., et al. (2020). Single-molecule dynamics of Dishevelled at the plasma membrane and Wnt pathway activation. *Proc. Natl. Acad. Sci. U.S.A.* 117, 16690-16701. doi: 10.1073/pnas.1910547117
- Marlow, F. L. (2020). Setting up for gastrulation in zebrafish. *Curr. Top. Dev. Biol.* 136, 33-83. doi: 10.1016/bs.ctdb.2019.08.002
- Mlodzik, M. (2016). The Dishevelled protein family: Still rather a mystery after over 20 years of molecular studies. *Curr. Top. Dev. Biol.* 117, 75-91. doi: 10.1016/bs.ctdb.2015.11.027
- Ngo, J., Hashimoto, M., Hamada, H., and Wynshaw-Boris, A. (2020). Deletion of the Dishevelled family of genes disrupts anterior-posterior axis specification and selectively prevents mesoderm differentiation. *Dev. Biol.* 464, 161-175. doi: 10.1016/j.ydbio.2020.05.010
- Niehrs, C. (2012). The complex world of WNT receptor signalling. *Nat. Rev. Mol. Cell Biol.* 13, 767-779. doi: 10.1038/nrm3470

- Pačlíková, P., Bernatík, O., Radaszkiewicz, T. W., and Bryja, V. (2018). The N-terminal part of the Dishevelled DEP domain is required for Wnt/ $\beta$ -Catenin signaling in mammalian cells. *Mol. Cell. Biol.* 37:e00145-17. doi: 10.1128/MCB.00145-17
- Pan, W. J., Pang, S. Z., Huang, T., Guo, H. Y., Wu, D., and Li, L. (2004). Characterization of function of three domains in dishevelled-1: DEP domain is responsible for membrane translocation of dishevelled-1. *Cell Res.* 14, 324-330. doi: 10.1038/sj.cr.7290232
- Park, T. J., Gray, R. S., Sato, A., Habas, R., and Wallingford, J. B. (2005). Subcellular localization and signaling properties of dishevelled in developing vertebrate embryos. *Curr. Biol.* 15, 1039-1044. doi: 10.1016/j.cub.2005.04.062
- Qi, J., Lee, H. J., Saquet, A., Cheng, X. N., Shao, M., Zheng, J. J., et al. (2017). Autoinhibition of Dishevelled protein regulated by its extreme C terminus plays a distinct role in Wnt/ $\beta$ -catenin and Wnt/planar cell polarity (PCP) signaling pathways. *J. Biol. Chem.* 292, 5898-5908. doi: 10.1074/jbc.M116.772509
- Rothbächer, U., Laurent, M. N., Dearnorff, M. A., Klein, P. S., Cho, K. W., and Fraser, S. E. (2000). Dishevelled phosphorylation, subcellular localization and multimerization regulate its role in early embryogenesis. *EMBO J.* 19, 1010-1022. doi: 10.1093/emboj/19.5.1010
- Seo, H. S., Habas, R., Chang, C., and Wang, J. (2017). Bimodal regulation of Dishevelled function by Vangl2 during morphogenesis. *Hum. Mol. Genet.* 26, 2053-2061. doi: 10.1093/hmg/ddx095
- Sharma, M., Castro-Piedras, I., Simmons, G. E. Jr., and Pruitt, K. (2018). Dishevelled: A masterful conductor of complex Wnt signals. *Cell. Signal.* 47, 52-64. doi: 10.1016/j.cellsig.2018.03.004
- Shimizu, N., Ishitani, S., Sato, A., Shibuya, H., and Ishitani, T. (2014). Hipk2 and PP1c cooperate to maintain Dvl protein levels required for Wnt signal transduction. *Cell Rep.* 8, 1391-1404. doi: 10.1016/j.celrep.2014.07.040
- Sinha, T., Wang, B., Evans, S., Wynshaw-Boris, A., and Wang, J. (2012). Dishevelled mediated planar cell polarity signaling is required in the second heart field lineage for outflow tract morphogenesis. *Dev. Biol.* 370, 135-144. doi: 10.1016/j.ydbio.2012.07.023
- Soares, M. L., Haraguchi, S., Torres-Padilla, M. E., Kalmar, T., Carpenter, L., Bell, G., et al. (2005). Functional studies of signaling pathways in peri-implantation development of the mouse embryo by RNAi. *BMC Dev. Biol.* 5:28. doi: 10.1186/1471-213X-5-28
- Sokol, S. Y. (1996). Analysis of Dishevelled signalling pathways during *Xenopus* development. *Curr. Biol.* 6, 1456-1467. doi: 10.1016/s0960-9822(96)00750-6
- Sokol, S. Y., Klingensmith, J., Perrimon, N., and Itoh, K. (1995). Dorsalizing and neuralizing properties of Xdsh, a maternally expressed *Xenopus* homolog of dishevelled. *Development* 121, 1637-1647
- Solnica-Krezel, L. (2020). Maternal contributions to gastrulation in zebrafish. *Curr. Top. Dev. Biol.* 140, 391-427. doi: 10.1016/bs.ctdb.2020.05.001
- Steinhart, Z., and Angers, S. (2018). Wnt signaling in development and tissue homeostasis. *Development* 145:dev146589. doi: 10.1242/dev.146589
- Stricker, S., Rauschenberger, V., and Schambony, A. (2017). ROR-family receptor tyrosine kinases. *Curr. Top. Dev. Biol.* 123, 105-142. doi: 10.1016/bs.ctdb.2016.09.003


- Tada, M., and Smith, J. C. (2000). Xwnt11 is a target of Xenopus Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. *Development* 127, 2227-2238.
- Tadjuidje, E., Cha, S. W., Louza, M., Wylie, C., and Heasman, J. (2011). The functions of maternal Dishevelled 2 and 3 in the early Xenopus embryo. *Dev. Dyn.* 240, 1727-1736. doi: 10.1002/dvdy.22671
- Tauriello, D. V., Jordens, I., Kirchner, K., Sloatstra, J. W., Kruitwagen, T., Bouwman, B. A., et al. (2012). Wnt/ $\beta$ -catenin signaling requires interaction of the Dishevelled DEP domain and C terminus with a discontinuous motif in Frizzled. *Proc. Natl. Acad. Sci. U.S.A.* 109, E812-E820. doi: 10.1073/pnas.1114802109
- Umbhauer, M., Djiane, A., Goisset, C., Penzo-Méndez, A., Riou, J. F., Boucaut, J. C., et al. (2000). The C-terminal cytoplasmic Lys-Thr-X-X-X-Trp motif in frizzled receptors mediates Wnt/ $\beta$ -catenin signalling. *EMBO J.* 19, 4944-4954. doi: 10.1093/emboj/19.18.4944
- Rauschenberger, V., Bernkopf, D. B., Krenn, S., Jalal, K., Heller, J., Behrens, J., et al. (2017). The phosphatase Pgam5 antagonizes Wnt/ $\beta$ -Catenin signaling in embryonic anterior-posterior axis patterning. *Development* 144, 2234-2247. doi: 10.1242/dev.144477
- Wallingford, J. B., and Habas, R. (2005). The developmental biology of Dishevelled: an enigmatic protein governing cell fate and cell polarity. *Development* 132, 4421-4436. doi: 10.1242/dev.02068
- Wallingford, J. B., and Harland, R. M. (2001). Xenopus Dishevelled signaling regulates both neural and mesodermal convergent extension: parallel forces elongating the body axis. *Development* 128, 2581-2592
- Wallingford, J. B., and Harland, R. M. (2002). Neural tube closure requires Dishevelled-dependent convergent extension of the midline. *Development* 129, 5815-5825. doi: 10.1242/dev.00123
- Wallingford, J. B., Rowning, B. A., Vogeli, K. M., Rothbächer, U., Fraser, S. E., and Harland, R. M. (2000). Dishevelled controls cell polarity during Xenopus gastrulation. *Nature* 405, 81-85. doi: 10.1038/35011077
- Wang, H. Y., and Malbon, C. C. (2012). Dishevelled C-terminus: prolyl and histidinyll motifs. *Acta Physiol. (Oxf)* 204, 65-73. doi: 10.1111/j.1748-1716.2011.02291.x
- Wang, J., Hamblet, N. S., Mark, S., Dickinson, M. E., Brinkman, B. C., Segil, N., et al. (2006). Dishevelled genes mediate a conserved mammalian PCP pathway to regulate convergent extension during neurulation. *Development* 133, 1767-1778. doi: 10.1242/dev.02347
- White, J. J., Mazzeu, J. F., Coban-Akdemir, Z., Bayram, Y., Bahrambeigi, V., Hoischen, A., et al. (2015). DVL1 frameshift mutations clustering in the penultimate exon cause autosomal-dominant Robinow syndrome. *Am. J. Hum. Genet.* 96, 612-622. doi: 10.1016/j.ajhg.2015.02.015
- White, J. J., Mazzeu, J. F., Hoischen, A., Bayram, Y., Withers, M., Gezdirici, A., et al. (2016). DVL3 alleles resulting in a -1 frameshift of the last exon mediate autosomal-dominant Robinow syndrome. *Am. J. Hum. Genet.* 98, 553-561. doi: 10.1016/j.ajhg.2016.01.005
- Wiese, K. E., Nusse, R., and van Amerongen, R. (2018). Wnt signalling: conquering complexity. *Development* 145:dev165902. doi: 10.1242/dev.165902
- Wong, H. C., Bourdelas, A., Krauss, A., Lee, H. J., Shao, Y., Wu, D., et al. (2003). Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in the C-terminal region of Frizzled. *Mol. Cell* 12, 1251-1260. doi: 10.1016/s1097-2765(03)00427-1



- Wong, H. C., Mao, J., Nguyen, J. T., Srinivas, S., Zhang, W., Liu, B., et al. (2000). Structural basis of the recognition of the dishevelled DEP domain in the Wnt signaling pathway. *Nat. Struct. Biol.* 7, 1178-1184. doi: 10.1038/82047
- Wynshaw-Boris, A. (2012). Dishevelled: in vivo roles of a multifunctional gene family during development. *Curr. Top. Dev. Biol.* 101, 213-235. doi: 10.1016/B978-0-12-394592-1.00007-7
- Xing, Y. Y., Cheng, X. N., Li, Y. L., Zhang, C., Saquet, A., Liu, Y. Y., et al. (2018). Mutational analysis of dishevelled genes in zebrafish reveals distinct functions in embryonic patterning and gastrulation cell movements. *PLoS Genet.* 14:e1007551. doi: 10.1371/journal.pgen.1007551
- Yan, L., Chen, J., Zhu, X., Sun, J., Wu, X., Shen, W., et al. (2018). Maternal Huluwa dictates the embryonic body axis through  $\beta$ -catenin in vertebrates. *Science* 362:eaat1045. doi: 10.1126/science.aat1045



**Fig. 1.** Structure-function relationship of conserved Dvl domains in the activation of Wnt/ $\beta$ -catenin and Wnt/PCP signaling (see text for details).



**Fig. 2.** Maternal and zygotic Dvl dosage effect on embryonic axis elongation in zebrafish embryos at 5 days post-fertilization (Xing et al., 2018). (A) Wild-type sibling embryo. (B) Zygotic *dv12* mutant. (C) The “bulldog” facial phenotype of maternal-zygotic *dv12* mutant with cyclopia. (D) Maternal-zygotic *dv13a* mutant. (E) Triallelic *dv12* and *dv13a* mutant. (F) Zygotic *dv12* and *dv13a* double mutant. Scale bar (A-F): 400  $\mu$ m.