



**HAL**  
open science

## On the existence of optimal shapes in architecture

Michael Hinz, Frédéric Magoulès, Anna Rozanova-Pierrat, Marina Rynkovskaya, Alexander Teplyaev

► **To cite this version:**

Michael Hinz, Frédéric Magoulès, Anna Rozanova-Pierrat, Marina Rynkovskaya, Alexander Teplyaev. On the existence of optimal shapes in architecture. *Applied Mathematical Modelling*, 2021, 94, pp.676-687. 10.1016/j.apm.2021.01.041 . hal-02956458v2

**HAL Id: hal-02956458**

**<https://hal.science/hal-02956458v2>**

Submitted on 15 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the existence of optimal shapes in architecture

Michael Hinz<sup>\*</sup>      Frédéric Magoulès<sup>†</sup>      Anna Rozanova-Pierrat<sup>‡</sup>  
Marina Rynkovskaya<sup>§</sup>      Alexander Teplyaev<sup>¶</sup>

## Abstract

We consider shape optimization problems for elasticity systems in architecture. A typical objective in this context is to identify a structure of maximal stability that is close to an initially proposed one. For structures without external forces on varying parts, classical methods allow proving the existence of optimal shapes within well-known classes of bounded uniformly Lipschitz domains. We discuss this for maximally stable roof structures. We then introduce a more general framework that includes external forces on varying parts (for instance, caused by loads of snow on roofs) and prove the existence of optimal shapes, now in a subclass of bounded uniformly Lipschitz domains, endowed with generalized surface measures on their boundaries. These optimal shapes realize the infimum of the corresponding energy of the system. Generalizing further to yet another, very new framework, now involving classes of bounded uniform domains with fractal measures on their boundaries, we finally prove the existence of optimal architectural shapes that actually realize the minimum of the energy. As a by-product we establish the well-posedness of the elasticity system on such domains. In an auxiliary result we show the convergence of energy functionals along a sequence of suitably converging domains. This result is helpful for an efficient approximation of an optimal shape by shapes that can be constructed in practice.

**Keywords:** Elasticity system; shape optimization; trace and extension; fractals; Mosco convergence; stable roof structures under snow loads

## 1 Introduction

The identification of a maximally stable (or most lightweight etc.) structure is a natural task in shape optimization. The main idea is to vary a shape within a chosen class of domains, all satisfying certain industrial constraints (such as the quantity of material, spatial and dimensional restrictions, or some similarity to a fixed form), and to search for an optimal shape that minimizes a given target (energy) functional. The found optimal shape must also belong to the chosen class of domains. There is a huge body of literature on different numerical methods [4, 14, 20, 34, 45, 46] and on implemented packages [44] suitable for simulations.

From the numerical point of view, finite-dimensional shape optimization problems are always solvable: In a finite-dimensional model, a discretized domain is determined by a finite mesh, so there are only finitely many possible changes of its boundary which respect all given constraints, and one can find at least one optimal shape. Non-existence results and other counterexamples usually come from the passage to the limit as the mesh becomes finer and finer. If one abandons the discretized perspective and adopts a more theoretical point of view, where the domains are subsets of  $\mathbb{R}^n$  and the target functional is defined on

---

<sup>\*</sup>Fakultät für Mathematik, Universität Bielefeld, Postfach 100131, 33501 Bielefeld, Germany.

<sup>†</sup>CentraleSupélec, Université Paris Saclay, France and University of Pécs, Pécs, Hungary

<sup>‡</sup>CentraleSupélec, Université Paris-Saclay, France.

<sup>§</sup>Peoples' Friendship University of Russia (RUDN University), Russian Federation

<sup>¶</sup>Department of Mathematics, University of Connecticut, Storrs, CT 06269-3009 USA.

infinite-dimensional space, the existence of an optimal shape is no longer trivial at all. There are different classical examples of shape optimization problems for which no optimal shape exists [22, Section 4.2].

We study a linear elasticity system (system (2) below) whose solution models the displacement of an architectural structure such as a roof or a bridge. The domain is subject to two types of boundary conditions: a Neumann (or 'stress') boundary condition that models external forces (tractions, effects of loads, etc.) affecting a part of the boundary of the structure, and a homogeneous Dirichlet boundary condition on another part of the boundary that is clamped and therefore cannot be displaced. The problem to find an optimal shape that minimizes the elastic energy of the system (cf. Remark 4) had been introduced in [14] under the additional assumption that the Neumann boundary condition is homogeneous on those parts of the boundary that can vary in the shape optimization. From a modeling perspective, homogeneous Neumann conditions on the varying parts are adequate if there is no significant external force acting on the structure's surface's corresponding parts. The existence of an optimal shape for this problem had so far remained open. Here our first contribution is proof of the existence of optimal shapes for this problem; we formulate this for the concrete example of maximally stable roofs, Theorem 3. (We emphasize that roofs are considered only as an illustrative example. The same method could be applied to other structures.) To provide the existence proof, we invoke classical results on shape optimization for bounded Lipschitz domains, mainly due to Chenaï [11]. We refer to [22] for more details and further references and to [12] for a survey on related well-posedness results for elasticity systems on bounded Lipschitz domains. The assumption of homogeneous Neumann conditions on the varying part of the boundary is essential for the classical methods [11] to work.

In practice, the 'varying' parts of the surface may be subject to significant forces that should be taken into account when searching for an optimal shape. For instance, in the case of a roof, the upper surface could be subject to the weight of a heavy load of snow. Such external forces correspond to inhomogeneous Neumann conditions on the varying parts of the boundary. If such conditions are imposed, care is needed because boundary integrals with respect to surface measures appear in the variational formulation (5) of the problem. Varying the shape, the natural notion of convergence for these measures is weak convergence. Mathematically this is delicate in the sense that the weak limit of a sequence of codimension one Hausdorff measures (which are the 'natural' surface measures on the boundary of a Lipschitz domain) is not necessarily a Hausdorff measure itself [28]. This fact motivates us to impose an additional condition on the Hausdorff measures on the boundaries (see (20)). This condition ensures that the limit measure, even if it may not be Hausdorff, is still diffuse enough to allow a variational analysis. Since we cannot guarantee that it is Hausdorff, an optimal shape may not be an element of the domains' initial class. Therefore it can realize only the infimum of energies over this class, but not their minimum. This approach is based on the methods recently introduced in the article [28], where the reader can find a detailed discussion in the framework of linear acoustics. Here we apply it to the linear elasticity system (2) and prove the existence of an optimal shape that realizes the infimum of the target (energy) functional, Theorem 4.

It is desirable to know that there are optimal shapes realizing the minimum of the energy, not just the infimum. As just discussed, this cannot be achieved in the context of bounded (uniformly) Lipschitz domains with natural Hausdorff measures on their boundaries. We, therefore, describe yet another framework. It is based on results in [23] and allows classes of shapes that are (generally non-Lipschitz) bounded uniform domains with measures on their boundaries that may have fractal features. The boundaries may have different parts of different Hausdorff dimensions. In general, the measures we consider on these boundaries are very different from Hausdorff measures, see [23] for a detailed discussion. For classes of such general domains, we can prove the existence of optimal shapes that realize the minimum of the energy, Theorem 5. In the particular case of roof shape optimization, this means that if such general designs are permitted, maximally stable roof shapes can always be seen to exist, Corollary 1. It may hardly be possible to construct such geometrically complicated

structures. However, numerical experiments conducted for other problems suggest that optimal shapes can have a multiscale nature, see for instance [29, Section 5] (and also [39]) for shape optimization in the absorption of the acoustical energy for the Helmholtz equation [23, 28]. (It was even observed that, in order to absorb an infinite range of frequencies 'almost optimally' [39, Section 3.4.1], the shape has to have obstacles at infinitely many different scales.) These experiments indicate that non-classical shapes capture the physical reality better than classical (smooth or Lipschitz) shapes in certain situations. Therefore, it seems reasonable to allow such shapes, even if the reasons for their appearance may not be so obvious. An irregular non-classical optimal shape can then be approximated by simpler shapes that can actually be constructed but still retain unusual scaling properties. (This is similar to the approximation of self-similar fractal boundaries by Lipschitz prefractals [13, 26].) Theorem 6 ensures that if this approximation is performed in suitable topologies for domains, the energy, as a function of a domain, behaves continuously and follows this approximation. Shapes that are approximately fractal have practical applications in civil engineering [1]. A discussion of multiscale features in architecture can be found in [37]. Our research is connected to several types of theoretical analysis in fractal domains, such as [7, 8, 10, 27, 35, 41, and references therein].

In Theorem 2 we provide the well-posedness result for the elasticity system on uniform bounded domains with fractal boundary measures. It is needed for the shape optimization results mentioned above and relies on fairly general boundary trace and extension results from [2, 5, 9, 25, 43] and on Korn's inequality for uniform domains [16].

The complexity of the considered shape optimization problem for the elasticity system is caused by exotic measures on the boundaries. This may be viewed as a geometric complication. In other specific results on elasticity difficulties may arise for different reasons. Often the system itself is of high complexity, see for instance [30–32] for studies of di- and micropolar thermoelastic porous materials and [42] for elasticity results for multibody systems at high velocities and under high loads.

In Section 2 we collect some notation. In Section 3 we introduce the elasticity system, discuss trace and extension methods, Theorem 1, and verify the validity of a norm equivalence with uniform constants, Lemma 1. We then define weak solutions (5) and obtain the well-posedness result, Theorem 2. Shape optimization problems for the simple case of homogeneous Neumann boundary conditions on varying parts are discussed in Section 4 using the well-known results from [11] and [22]. As a practical example, we address the existence question for optimal shapes of a roof, Theorem 3, and we comment on a related problem formulated in [14]. In Section 5, we follow the method of [28] to study a less simple generalization with possibly inhomogeneous Neumann boundary conditions on varying parts. Section 6 addresses the uniform domain setup and states the existence of energy, minimizing optimal shapes. In A we provide generalized Green's formulas that justify our definition of weak solution. In B we briefly comment on a technical detail in the proof of Lemma 1. C contains an auxiliary result, namely the Mosco convergence of energy functionals for Robin type problems along a sequence of suitably converging uniform domains.

## Acknowledgements

M.H. gratefully acknowledges financial support by the DFG IRTG 2235 and the DFG CRC 1283. A.T. was supported in part by NSF grant DMS-1613025. A. R.-P. thanks her student Antoine Verdon for his preliminary work in the subject. M.R. thanks for the financial support CentraleSupélec.

## 2 Some notation

For the Euclidean scalar product of two vectors  $\xi, \eta \in \mathbb{R}^N$  we write  $\xi \cdot \eta$ . By  $\mathcal{M}_N$  we denote the vector space of  $N \times N$ -matrices with real entries, and by  $A : B$  we denote the full

contraction

$$A : B = \text{tr } A^t B = \sum_{i,j=1}^N a_{ij} b_{ij} \quad (1)$$

of two matrices  $A = (a_{ij})_{1 \leq i,j \leq N}$  and  $B = (b_{ij})_{1 \leq i,j \leq N}$  from  $\mathcal{M}_N$ . Note that the full contraction (1) provides a natural inner product on  $\mathcal{M}_N$ . Let  $\mathcal{M}_N^s$  be the subspace of  $\mathcal{M}_N$  consisting of symmetric matrices and given  $\alpha > 0$  and  $\beta > 0$  let  $\mathcal{M}_N^s(\alpha, \beta)$  denote the subspace of all invertible  $M \in \mathcal{M}_N^s$  such that  $M\xi \cdot \xi \geq \alpha|\xi|^2$  and  $M^{-1}\xi \cdot \xi \geq \beta|\xi|^2$  for all  $\xi \in \mathbb{R}^N$ .

Given a domain  $\Omega \subset \mathbb{R}^N$  and a vector field  $\mathbf{v} \in W^{1,2}(\Omega)^N$  we denote the symmetric part of its gradient by

$$e(\mathbf{v}) = \frac{1}{2} (\nabla \mathbf{v} + (\nabla \mathbf{v})^t).$$

The assumption on  $\mathbf{v}$  implies that  $e(\mathbf{v}) \in L^2(\Omega, \mathcal{M}_N^s)$ . Here  $W^{1,2}(\Omega)$  is the classical Sobolev space, and product norms on  $W^{1,2}(\Omega)^N$  are defined in the natural way. For an element  $T = (T_{ij})_{1 \leq i,j \leq N}$  of  $W^{1,2}(\Omega)^{N \times N}$  we define  $\text{div } T \in L^2(\Omega)^N$  as the vector field  $(\sum_{j=1}^N \partial_{x_j} T_{1j}, \dots, \sum_{j=1}^N \partial_{x_j} T_{Nj})$ .

By  $B(x, r)$  we denote the Euclidean open ball centered in  $x$  of the radius  $r > 0$ . We use the symbol  $\lambda^N$  for the  $N$ -dimensional Lebesgue measure and the symbol  $\mathcal{H}^N$  for the  $N$ -dimensional Hausdorff measure.

### 3 Variational formulation and well-posedness

We assume  $N \geq 2$ . Let  $\Omega \subset \mathbb{R}^N$  be a bounded domain and let  $\Gamma_{\text{Dir}}$  and  $\Gamma_{\text{Neu}}$  be subsets of its boundary  $\partial\Omega$ . We assume that they are 'almost disjoint' (in the sense that their overlap has zero measure), this will be made precise below. Let  $A \in L^\infty(\Omega, \mathcal{M}_N^s(\alpha, \beta))$  and write  $\sigma(\mathbf{v}) = Ae(\mathbf{v})$ ,  $\mathbf{v} \in W^{1,2}(\Omega)^N$ . We are interested in solutions  $\mathbf{u} \in W^{1,2}(\Omega)^N$  to problems of type

$$\begin{cases} -\text{div } \sigma(\mathbf{u}) = \mathbf{f} & \text{in } \Omega, \\ \mathbf{u} = 0 & \text{on } \Gamma_{\text{Dir}}, \\ \sigma(\mathbf{u}) \cdot \mathbf{n} = \mathbf{g} & \text{on } \Gamma_{\text{Neu}}. \end{cases} \quad (2)$$

Here  $\mathbf{n}$  denotes the outward unit normal. The domain  $\Omega$  is 'clamped' at  $\Gamma_{\text{Dir}}$ . The part of  $\Gamma_{\text{Neu}}$  on which  $\mathbf{g}$  is not zero is subjected to traction represented by the force field  $\mathbf{g}$ . The vector field  $\mathbf{f}$  represents a force field experienced inside  $\Omega$ . The prospective solution  $\mathbf{u}$  is the (unknown) displacement vector field,  $e(\mathbf{u})$  is the strain tensor, and  $\sigma(\mathbf{u})$  is the stress tensor, determined by the given coefficient  $A$ , which is minus the elasticity tensor in Hooke's law.

#### Remark 1

- (i) A more common choice for  $\sigma$  within the linear elasticity theory for isotropic materials is  $\sigma(\mathbf{v}) = \lambda(\text{tr } e(\mathbf{v}))I + 2\mu e(\mathbf{v})$ , where  $\mu > 0$  and  $\lambda > -2\mu/N$  are the Lamé coefficients of the material. See for instance [12, Section 6.2]. With insubstantial modifications our results also apply to this case.
- (ii) Note that from (2) one can recover the system considered in [14, Subsection 3.1, formula (1)] if  $\Gamma_{\text{Neu}}$  is split into a part where  $\mathbf{g}$  is nonzero and one where it is zero.

We will give a rigorous meaning to (2) in terms of a variational formulation. This works well if the boundary  $\partial\Omega$  is the support of a suitable measure so that Sobolev functions, fields, or tensors on  $\Omega$  have well-defined traces on  $\partial\Omega$ .

Recall that given  $d > 0$ , a Borel measure  $\mu_F$  on  $\mathbb{R}^N$  with  $F = \text{supp } \mu_F$  is said to be *upper  $d$ -regular* if there is a constant  $c_d > 0$  such that

$$\mu_F(B(x, r)) \leq c_d r^d, \quad x \in F, \quad 0 < r \leq 1. \quad (3)$$

This property is well-known and widely used, we refer to [2] and [19]. Recall also that a domain  $\Omega \subset \mathbb{R}^N$  is said to be an  $W^{1,2}$ -extension domain if there is a bounded linear extension operator  $E : W^{1,2}(\Omega) \rightarrow W^{1,2}(\mathbb{R}^N)$ , [24, 38]. We state a trace result that follows from a special case of [23, Theorem 5.1] and the finiteness of the measure on  $\partial\Omega$ . The result is based on [9, Corollaries 7.3 and 7.4], which in turn use [2, Theorems 7.2.2 and 7.3.2].

**Theorem 1** *Let  $\Omega \subset \mathbb{R}^N$  be a bounded  $W^{1,2}$ -extension domain. Suppose that  $\mu_{\partial\Omega}$  is a Borel measure with  $\text{supp } \mu_{\partial\Omega} = \partial\Omega$  and such that (3) holds with some  $N - 2 < d \leq N$ .*

- (i) *There are a compact linear operator  $\text{Tr} : W^{1,2}(\Omega) \rightarrow L^2(\partial\Omega, \mu_{\partial\Omega})$  and a constant  $c_{\text{Tr}} > 0$ , depending only on  $N, \varepsilon, d$  and  $c_d$ , such that*

$$\|\text{Tr } f\|_{L^2(\partial\Omega, \mu_{\partial\Omega})} \leq c_{\text{Tr}} \|f\|_{W^{1,2}(\Omega)}, \quad f \in W^{1,2}(\Omega).$$

*Endowed with the norm*

$$\|\varphi\|_{\text{Tr}(W^{1,2}(\Omega))} := \inf\{\|g\|_{W^{1,2}(\Omega)} \mid \varphi = \text{Tr } g\},$$

*the image  $\text{Tr}(W^{1,2}(\Omega))$  becomes a Hilbert space. The embedding*

$$\text{Tr}(W^{1,2}(\Omega)) \subset L^2(\partial\Omega, \mu_{\partial\Omega})$$

*is compact.*

- (ii) *There is a linear operator  $H_{\partial\Omega} : \text{Tr}(W^{1,2}(\Omega)) \rightarrow W^{1,2}(\Omega)$  of norm one such that  $\text{Tr}(H_{\partial\Omega}\varphi) = \varphi$  for all  $\varphi \in \text{Tr}(W^{1,2}(\Omega))$ .*

The trace of vector fields or tensors is understood in the component-wise sense.

**Remark 2**

- (i) *The trace operator is defined by  $\text{Tr } f := \tilde{g}$ , where*

$$\tilde{g}(x) = \lim_{r \rightarrow 0} \frac{1}{\lambda^N(B(x, r))} \int_{B(x, r)} g(y) dy \quad (4)$$

*is the pointwise redefinition of an extension  $g \in W^{1,2}(\mathbb{R}^N)$  of  $f$ . By the Lebesgue differentiation theorem this limit exists outside a  $\lambda^N$ -null set, but due to the Sobolev regularity of  $g$  it exists outside a much smaller set, as can be made precise using capacities, [2, 33]. Since  $\mu_{\partial\Omega}$  satisfies (3) with  $d$  as stated, the set of points of  $\partial\Omega$  where this limit exists is of full  $\mu_{\partial\Omega}$ -measure, [2, Section 7]. The independence of the chosen extension is proved in [9, Theorem 6.1], another proof is given in [43, Theorem 1].*

- (ii) *The extension operator  $H_{\partial\Omega}$  is defined as the 1-harmonic extension: Given  $\varphi \in \text{Tr}(W^{1,2}(\Omega))$ ,  $H_{\partial\Omega}\varphi$  is the (unique) element  $w$  of  $W^{1,2}(\Omega)$  such that  $\text{Tr } w = \varphi$  and  $\Delta w = w$  in  $\Omega$  in the weak sense. Details can be found in [23, Section 5].*

- (iii) *Clearly the Hausdorff dimension of  $\partial\Omega$  is at least  $N - 1$ . If the maximal possible exponent  $d$  in (3) is less than  $N - 1$  then  $\mu_{\partial\Omega}$  is singular with respect to the  $(N - 1)$ -dimensional Hausdorff measure  $\mathcal{H}^{N-1}$  and assigns positive mass also to some parts of  $\partial\Omega$  that have Hausdorff dimension less than  $N - 1$ .*

Let  $\Omega$  and  $\mu_{\partial\Omega}$  be as in Theorem 1. For a fixed Borel subset  $\Gamma_{\text{Dir}}$  of  $\partial\Omega$  we also consider the closed subspace

$$V(\Omega, \Gamma_{\text{Dir}}) = \{v \in W^{1,2}(\Omega) \mid \text{Tr } v = 0 \text{ } \mu_{\partial\Omega}\text{-a.e. on } \Gamma_{\text{Dir}}\}$$

of  $W^{1,2}(\Omega)$ .

Now suppose that  $\Gamma_{\text{Dir}}$  and  $\Gamma_{\text{Neu}}$  are two Borel subsets of  $\partial\Omega$  such that  $\partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}$  and  $\mu_{\partial\Omega}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0$ . Then a rigorous meaning can be given to the system (2) in terms of the following variational formulation. We say that a vector field  $\mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N$  is a *weak solution* to (2) with data  $\mathbf{f} \in L^2(\Omega)^N$  and  $\mathbf{g} \in L^2(\Gamma_{\text{Neu}})^N$  if

$$\int_{\Omega} \sigma(\mathbf{u}) : e(\theta) dx = \int_{\Omega} \mathbf{f} \cdot \theta dx + \int_{\Gamma_{\text{Neu}}} \mathbf{g} \cdot \text{Tr } \theta \, d\mu_{\partial\Omega}, \quad \theta \in V(\Omega, \Gamma_{\text{Dir}})^N. \quad (5)$$

A generalized Green formula guarantees that in cases when  $\Omega$ ,  $A$ ,  $\mathbf{f}$  and  $\mathbf{g}$  are smooth and classical solutions exist, these are also weak solutions, see formula (26) of Proposition 2 in A.

We establish the existence of weak solutions for a specific class of  $W^{1,2}$ -extension domains. Recall that a domain  $\Omega$  of  $\mathbb{R}^N$  is an  $(\varepsilon, \infty)$ -domain,  $\varepsilon > 0$ , if for any  $x, y \in \Omega$  there is a rectifiable arc  $\gamma \subset \Omega$  with length  $\ell(\gamma)$  joining  $x$  to  $y$  and satisfying

1.  $\ell(\gamma) \leq \frac{|x-y|}{\varepsilon}$  and
2.  $d(z, \partial\Omega) \geq \varepsilon|x-z| \frac{|y-z|}{|x-y|}$  for  $z \in \gamma$ .

By [24, Theorem 1] any  $(\varepsilon, \infty)$ -domain is a  $W^{1,2}$ -extension domain.

We quote the following special case of the Korn inequality proved in [16, Theorem 2.1]. A more standard version for bounded Lipschitz domains can be found in [12, Theorem 6.3-3].

**Proposition 1** *Let  $\Omega \subset \mathbb{R}^n$  be a bounded  $(\varepsilon, \infty)$ -domain. There is a constant  $C_K(\varepsilon, N, \text{diam}(\Omega)) > 0$  depending only on  $\varepsilon$ ,  $N$  and  $\text{diam}(\Omega)$  such that*

$$\|\mathbf{u}\|_{W^{1,2}(\Omega)^N} \leq C_K(\varepsilon, N, \text{diam}(\Omega)) (\|\mathbf{u}\|_{L^2(\Omega)^N} + \|e(\mathbf{u})\|_{L^2(\Omega)^{N \times N}}) \quad (6)$$

for all  $\mathbf{u} \in W^{1,2}(\Omega)^N$ . The norm  $\mathbf{u} \rightarrow \left( \|\mathbf{u}\|_{L^2(\Omega)^N}^2 + \|e(\mathbf{u})\|_{L^2(\Omega)^{N \times N}}^2 \right)^{\frac{1}{2}}$  is equivalent to  $\|\cdot\|_{W^{1,2}(\Omega)^N}$  on  $W^{1,2}(\Omega)^N$ .

If  $\Omega$  is a bounded  $(\varepsilon, \infty)$ -domain then the embedding  $W^{1,2}(\Omega) \rightarrow L^2(\Omega)$  is compact. (This is actually true for all bounded  $W^{1,2}$ -extension domains [6, 40].) Combining the compactness of this embedding and (6) one obtains the following equivalence of the norms on  $V(\Omega, \Gamma_{\text{Dir}})^N$ .

**Lemma 1**

- (i) *Let  $\Omega \subset \mathbb{R}^N$  be a bounded  $(\varepsilon, \infty)$ -domain and  $\mu_{\partial\Omega}$  a Borel measure with  $\text{supp } \mu_{\partial\Omega} = \partial\Omega$  and such that (3) holds with some  $N-2 < d \leq N$ . Suppose that  $\Gamma_{\text{Dir}}$  is a Borel subset of  $\partial\Omega$  and  $\mu_{\partial\Omega}(\Gamma_{\text{Dir}}) > 0$ . Then there is a constant  $c_K(\Omega, \mu_{\partial\Omega}, \Gamma_{\text{Dir}}) > 0$  such that*

$$c_K(\Omega, \mu_{\partial\Omega}, \Gamma_{\text{Dir}}) \|\mathbf{u}\|_{W^{1,2}(\Omega)^N} \leq \|e(\mathbf{u})\|_{L^2(\Omega)^{N \times N}} \leq \|\mathbf{u}\|_{W^{1,2}(\Omega)^N} \quad (7)$$

for all  $\mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N$ .

- (ii) *Let  $\Omega \subset D \subset \mathbb{R}^N$  be bounded  $(\varepsilon, \infty)$ -domains and  $\mu_{\partial D}$ ,  $\mu_{\partial\Omega}$  Borel measures with  $\text{supp } \mu_{\partial D} = \partial D$  and  $\text{supp } \mu_{\partial\Omega} = \partial\Omega$  and such that for both (3) holds with some  $N-2 < d \leq N$ . Suppose that  $\Gamma_{\text{Dir}}$  is a Borel subset of  $\partial D \cap \partial\Omega$  with  $\mu_{\partial D}(\Gamma_{\text{Dir}}) > 0$  and  $\mu_{\partial\Omega}|_{\Gamma_{\text{Dir}}} = \mu_{\partial D}|_{\Gamma_{\text{Dir}}}$ . Then the constant  $c_K(\Omega, \mu_{\partial\Omega}, \Gamma_{\text{Dir}})$  in (7) can be replaced by another constant  $c_K(\varepsilon, N, D, \mu_{\partial D}, \Gamma_{\text{Dir}}) > 0$  depending only on  $\varepsilon$ ,  $N$ ,  $D$ ,  $\mu_{\partial D}$  and  $\Gamma_{\text{Dir}}$ . Moreover, the Poincaré inequality*

$$\int_{\Omega} \mathbf{u} \cdot \mathbf{u} \, dx \leq C_P(\varepsilon, N, D, \mu_{\partial D}, \Gamma_{\text{Dir}}) \int_{\Omega} \nabla \mathbf{u} : \nabla \mathbf{u} \, dx \quad (8)$$

holds for all  $\mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N$ , where  $C_P(\varepsilon, N, D, \mu_{\partial D}, \Gamma_{\text{Dir}}) > 0$  is a constant depending only on  $\varepsilon$ ,  $N$ ,  $D$ ,  $\mu_{\partial D}$  and  $\Gamma_{\text{Dir}}$ .

**Proof :** Statement (i) can be proved by the same arguments as used in [12, Theorem 6.3-4, step (iii) in the proof]. To see (ii) recall that according to [24, Theorem 2] there exists an extension operator  $E_\Omega$  taking  $\mathbf{u}$  into a locally integrable function  $E_\Omega \mathbf{u}$  on  $\mathbb{R}^N$  with gradient in  $L^2(\mathbb{R}^N)^{N \times N}$  such that

$$\|\nabla E_\Omega \mathbf{u}\|_{L^2(D)^{N \times N}} \leq \|\nabla E_\Omega \mathbf{u}\|_{L^2(\mathbb{R}^N)^{N \times N}} \leq C_{\text{Ext}}(\varepsilon, N) \|\nabla \mathbf{u}\|_{L^2(\Omega)^{N \times N}}, \quad (9)$$

where  $C_{\text{Ext}}(\varepsilon, N)$  is a constant depending only on  $\varepsilon$  and  $N$ . By adding two inequalities we can see the same holds with  $e(\mathbf{u})$  in place of  $\nabla \mathbf{u}$ . The pointwise redefinition  $\widetilde{E}_\Omega \mathbf{u}$  of  $E_\Omega \mathbf{u}$  in the sense of (4) satisfies

$$\widetilde{E}_\Omega \mathbf{u} = 0 \quad \mu_{\partial D}\text{-a.e. on } \Gamma_{\text{Dir}}, \quad (10)$$

see B for details of the brief argument. Using (10) and the boundedness of  $D$  one can then follow a standard pattern, see [17, Proposition 7.1] or [18], to obtain the Poincaré inequality

$$\int_D \text{Ext}_\Omega \mathbf{u} \cdot \text{Ext}_\Omega \mathbf{u} \, dx \leq C_P(\varepsilon, N, D, \mu_{\partial D}, \Gamma_{\text{Dir}}) \int_D \nabla \text{Ext}_\Omega \mathbf{u} : \nabla \text{Ext}_\Omega \mathbf{u} \, dx \quad (11)$$

for all  $\mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N$ . An application of (i) to  $D$  yields

$$\begin{aligned} c_K(D, \mu_{\partial D}, \Gamma_{\text{Dir}}) \|\nabla \text{Ext}_\Omega \mathbf{u}\|_{L^2(D)^{N \times N}} &\leq \|e(\text{Ext}_\Omega \mathbf{u})\|_{L^2(D)^{N \times N}} \\ &\leq C_{\text{Ext}}(\varepsilon, N) \|e(\mathbf{u})\|_{L^2(\Omega)^{N \times N}}, \end{aligned}$$

and combining with (11), we arrive at

$$\|\mathbf{u}\|_{L^2(\Omega)^N} \leq \frac{C_P(D, \mu_{\partial D}, \Gamma_{\text{Dir}})^{1/2} C_{\text{Ext}}(\varepsilon, N)}{c_K(D, \mu_{\partial D}, \Gamma_{\text{Dir}})} \|e(\mathbf{u})\|_{L^2(\Omega)^{N \times N}}.$$

Plugging this into (6), we obtain the first claim in (ii). Slight modifications of the preceding arguments also give (8). See also ([15, Theorem 10]) for a similar scalar result.  $\square$

We obtain the following well-posedness result:

**Theorem 2** *Let  $\Omega \subset \mathbb{R}^N$  be a bounded  $(\varepsilon, \infty)$ -extension domain. Suppose that  $\mu_{\partial\Omega}$  is a Borel measure with  $\text{supp } \mu_{\partial\Omega} = \partial\Omega$  and such that (3) holds with some  $N - 2 < d \leq N$ . Suppose that  $\Gamma_{\text{Dir}}$  and  $\Gamma_{\text{Neu}}$  are Borel subsets of  $\partial\Omega$  such that  $\partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}$ ,  $\mu_{\partial\Omega}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0$  and  $\mu_{\partial\Omega}(\Gamma_{\text{Dir}}) > 0$ .*

*Then for all  $\mathbf{g} \in [\text{Tr}(W^{1,2}(\Omega))]^N$  and  $\mathbf{f} \in L^2(\Omega)^N$  there is a unique weak solution  $\mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N$  of (5). Moreover, there is a constant  $C > 0$  such that*

$$\|\mathbf{u}\|_{W^{1,2}(\Omega)^N} \leq C (\|\mathbf{f}\|_{L^2(\Omega)^N} + \|\mathbf{g}\|_{[\text{Tr}(W^{1,2}(\Omega))]^N}). \quad (12)$$

**Proof :** Theorem 2 is a direct consequence of the Lax-Milgram Lemma. Indeed, by the definition of  $\sigma(\mathbf{u})$  and (7),

$$\int_\Omega \sigma(\mathbf{v}) : e(\mathbf{v}) \, dx \geq \alpha \int_\Omega e(\mathbf{v}) : e(\mathbf{v}) \, dx \geq \alpha c^{-1} \|\mathbf{v}\|_{W^{1,2}(\Omega)^N}^2, \quad \mathbf{v} \in V(\Omega, \Gamma_{\text{Dir}})^N.$$

Also the continuity of the bilinear form is seen easily. The continuity of the linear functional  $\theta \mapsto \int_\Omega \mathbf{f} \cdot \theta \, dx + \int_{\Gamma_{\text{Neu}}} \mathbf{g} \cdot \text{Tr } \theta \, d\mu_{\partial\Omega}$  on  $V(\Omega, \Gamma_{\text{Dir}})^N$  follows using Theorem 1, and by the same theorem we obtain estimate (12).  $\square$

**Remark 3** *If  $\Omega$  in Theorem 2 satisfies the hypotheses of Lemma 1 (ii) then (12) holds with a constant  $C > 0$  depending only on  $\varepsilon, N, D, \mu_{\partial D}$  and  $\Gamma_{\text{Dir}}$ .*



## 4 Lipschitz optimal shapes for homogeneous Neumann conditions

We proceed to the existence of optimal shapes for the elasticity system (2). In this section, we prove it for a practical example within a well-known setup involving Lipschitz domains and explain possible generalizations and the difficulties involved. This may be seen as motivation for our new existence results in Sections 5 and 6.

For  $A$  as in (2), and  $\Omega$ ,  $\mu_{\partial\Omega}$ ,  $\Gamma_{\text{Dir}}$ ,  $\Gamma_{\text{Neu}}$  as in Theorem 2 we can define the functional

$$J(\Omega, \mu_{\partial\Omega}, \mathbf{v}) := c_1 \int_{\Omega} |\mathbf{v}|^2 dx + c_2 \int_{\Omega} Ae(\mathbf{v}) : e(\mathbf{v}) dx, \quad \mathbf{v} \in V(\Omega, \Gamma_{\text{Dir}})^N, \quad (13)$$

where  $c_1$  and  $c_2$  are fixed nonnegative constants. Suppose that  $\mathbf{f}$  and  $\mathbf{g}$  are fixed data, that we can find a class of domains within which  $\Omega$  can vary but  $\Gamma_{\text{Dir}}$  is kept fixed and that for each  $\Omega$  from that class Theorem 2 yields a unique weak solution  $\mathbf{u}(\Omega, \mu_{\partial\Omega})$  to (2) with the same data  $\mathbf{f}$  and  $\mathbf{g}$ . Then we may interpret

$$\Omega \mapsto J(\Omega, \mu_{\partial\Omega}, \mathbf{u}(\Omega, \mu_{\partial\Omega})) \quad (14)$$

as a functional on this class of domains, and we may attempt to minimize it.

### Remark 4

- (i) For  $c_1 = 0$  and  $c_2 = 1$  the value  $J(\Omega, \mu_{\partial\Omega}, \mathbf{u}(\Omega, \mu_{\partial\Omega}))$  is the elastic energy stored in  $\Omega$  (the compliance of  $\Omega$ ), [14, Subsection 3.1]. For  $c_1 = 1$  and  $c_2 = 0$  the value  $J(\Omega, \mu_{\partial\Omega}, \mathbf{u}(\Omega, \mu_{\partial\Omega}))$  is the square of the  $L^2$ -norm of the displacement field, and the minimization of this norm may be viewed as a mathematically tractable substitute for the (intractable) minimization of the  $L^\infty$ -norm.
- (ii) For a mixed boundary value problem for the Helmholtz equation functionals similar to (14), but with an additional boundary term, are studied in [23, Section 7].

We discuss a practical application. Suppose that we are searching for the most stable (strongest) roof for a building and model it by a domain  $\Omega \subset \mathbb{R}^3$ . We assume that the moving part  $\Gamma_{\text{Neu}}$  of the boundary  $\partial\Omega$  is the union of two disjoint parts  $\Gamma_{\text{upper}}$  and  $\Gamma_{\text{lower}}$  which model the upper and the lower side of the roof. The set  $\Gamma_{\text{Dir}}$  models vertical boundary parts of the roof at which it is fixed by a structural connection to other parts of the building. See Fig. 1. We impose the condition that the volume  $c_v > 0$  of the roof itself remains fixed and that  $\Gamma_{\text{upper}}$  and  $\Gamma_{\text{lower}}$  are always parallel with a vertical distance

$$h_z(\Omega) = \frac{c_v}{\mu(\Gamma_{\text{lower}})}. \quad (15)$$

Since  $c_v$  is kept constant,  $h_z(\Omega)$  will typically change when  $\Gamma_{\text{upper}}$  and  $\Gamma_{\text{lower}}$  move. We assume that  $\mathbf{g} = 0$  and set

$$\mathbf{f} = \rho_0 h_z(\Omega) \mathbf{e}_z, \quad (16)$$

where  $\rho_0 > 0$  is a constant (mass) density and  $\mathbf{e}_z = (0, 0, 1)$ . The field  $\mathbf{f}(\Omega)$  represents the weight of the roof  $\Omega$ .

To introduce suitable classes of domains let us recall the following from [3, 11]. Given  $\varepsilon > 0$ , a domain  $\Omega \subset \mathbb{R}^N$  is said to have the  $\varepsilon$ -cone property if for all  $x \in \partial\Omega$  there exists  $\xi_x \in \mathbb{R}^N$  with  $\|\xi_x\| = 1$  such that for all  $y \in \overline{\Omega} \cap B(x, \varepsilon)$

$$C(y, \xi_x, \varepsilon) = \{z \in \mathbb{R}^N \mid (z - y, \xi_x) \geq \cos(\varepsilon) \|z - y\| \text{ and } 0 < \|z - y\| < \varepsilon\} \subset \Omega.$$

It is well-known that a domain  $\Omega$  with bounded boundary  $\partial\Omega$  has the  $\varepsilon$ -cone property for some  $\varepsilon > 0$  if and only if it is a Lipschitz domain, [22, Theorem 2.4.7]. Now let  $D \subset \mathbb{R}^3$  be a bounded Lipschitz domain and  $\varepsilon > 0$ . Somewhat similarly to [22, Section 2.4] we write

$$\mathcal{O}(D, \varepsilon) := \{\Omega \subset D \mid \Omega \text{ is a domain satisfying the } \varepsilon\text{-cone property}\}. \quad (17)$$




Figure 1: The boundary parts  $\Gamma_{\text{upper}}$  and  $\Gamma_{\text{lower}}$  remain parallel to each other with distance  $h_z$  as defined in (16). The lower part  $\Gamma_{\text{lower}}$  is supposed to stay within the closure of a fixed open set  $G$ . We assume that all possible shapes  $\Omega$  (and therefore also the set  $G$ ) are subsets of a fixed open set  $D$ .

Let  $G$  be a nonempty open proper subset of  $D$ ,  $\Gamma_{\text{Dir}}$  a compact subset of  $\partial D$  with  $\mathcal{H}^2(\Gamma_{\text{Dir}}) > 0$ , let  $\varepsilon > 0$ ,  $c_v > 0$  and  $0 < \ell_0 < \ell_1$ . We define a class of admissible shapes by

$$\begin{aligned} U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}}) &= \{\Omega \in \mathcal{O}(D, \varepsilon) \mid \lambda^3(\Omega) = c_v, \ell_0 \leq \mathcal{H}^2(\partial\Omega) \leq \ell_1, \\ &\partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}, \mathcal{H}^2(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0, \Gamma_{\text{Dir}} = \partial\Omega \cap \partial D, \\ &\Gamma_{\text{Neu}} = \Gamma_{\text{lower}} \cup \Gamma_{\text{upper}}, \Gamma_{\text{lower}} \subset \overline{G}, \Gamma_{\text{upper}} = \Gamma_{\text{lower}} + h_z \mathbf{e}_z\}. \end{aligned} \quad (18)$$

In this case the existence of an optimal shape realizing the minimum of the functional on  $U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$  follows from the results of Chenais [11]. By [22, Theorem 2.4.10] the collection  $\mathcal{O}(D, \varepsilon)$  of domains having the  $\varepsilon$ -cone property and contained in  $D$  is compact with respect to the convergence in the sense of characteristic functions, the convergence in the Hausdorff sense and the convergence in the sense of compacts. For any sequence  $\Omega_n \rightarrow \Omega$  of sets in this class covering in all three senses, their boundaries  $\partial\Omega_n \rightarrow \partial\Omega$  and also their closures  $\overline{\Omega}_n \rightarrow \overline{\Omega}$  converge in the Hausdorff sense. The set of domains  $U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$  defined in (18) is a closed subset of the class  $\mathcal{O}(D, \varepsilon)$  and therefore compact. Note also that each  $\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$ , together with  $\mu_{\partial\Omega} = \mathcal{H}^2|_{\partial\Omega}$ , satisfies the hypotheses of Theorem 2. Proceeding as in [22, Theorem 4.3.1] we obtain the following result for the case of zero Neumann data.

**Theorem 3** *Let  $U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$  be as in (18),  $\rho_0 > 0$ ,  $c_1 \geq 0$  and  $c_2 \geq 0$ . For each  $\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$  let  $\mathbf{u}(\Omega, \mathcal{H}^2)$  denote the unique weak solution to (2) on  $\Omega$  with  $\mathbf{g} \equiv 0$  and  $\mathbf{f} \equiv \mathbf{f}(\Omega)$  defined as in (16). Let  $J$  be the energy functional defined in (13).*

*Then there is an optimal shape  $\Omega_{\text{opt}} \in U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$  realizing the minimum of (14) over  $U_{\text{ad}}(D, G, \varepsilon, c_v, \ell_0, \ell_1, \Gamma_{\text{Dir}})$ .*

**Remark 5**

- (i) *A similar example of a shape optimization problem is studied in [14]. There the authors propose to minimize (14) with  $c_1 = 0$  and  $c_2 = 1$  in (13) in a setup where  $\Gamma_{\text{Dir}}$  and a*

part  $\Gamma_{\text{Neu,fix}}$  of  $\Gamma_{\text{Neu}}$  are fixed and only  $\Gamma := \Gamma_{\text{Neu}} \setminus \Gamma_{\text{Neu,fix}}$ , on which  $\mathbf{g}$  is assumed to be zero, can move. On the fixed part  $\Gamma_{\text{Neu,fix}}$  the function  $\mathbf{g}$  can be nonzero. They keep the volume of  $\Omega$  constant and introduce a penalization term in the functional which restricts the area of the possible changes of  $\Gamma$ . This term is based on the (signed) distance to an initially given shape  $\Gamma_0$ .

- (ii) The idea to restrict the possible shapes to a neighborhood of a given shape is equivalent to fixing a compact set  $\overline{G}$  within which the moving boundary part is required to stay,  $\Gamma \subset \overline{G}$ . In [28], the latter condition is shown to be necessary to ensure the existence of an optimal shape. A comparable condition is also used in [23], where the domains  $\Omega$  are required to contain another fixed open set  $D_0$ .

The sketched roof shape optimization problem can easily be generalized in the sense that (14) could be minimized over

$$U_{\text{ad}}(D, G, \varepsilon, c_v, \Gamma_{\text{Dir}}, \Gamma_{\text{Neu,fix}}) = \{\Omega \in \mathcal{O}(D, \varepsilon) \mid \lambda^N(\Omega) = c_v, \\ \partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}, \mathcal{H}^{N-1}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0, \Gamma_{\text{Dir}} = \partial\Omega \cap \partial D, \\ \Gamma_{\text{Neu,fix}} \subset \Gamma_{\text{Neu}}, \Gamma_{\text{Neu}} \setminus \Gamma_{\text{Neu,fix}} \subset \overline{G}\}, \quad (19)$$

where  $D, G, \varepsilon, c_v, \Gamma_{\text{Dir}}$  are similarly as before. In this setup  $\Gamma_{\text{Neu,fix}} \subset \Gamma_{\text{Neu}}$  is kept fixed and only  $\Gamma = \Gamma_{\text{Neu}} \setminus \Gamma_{\text{Neu,fix}}$  can move, as in [14]. If  $\mathbf{g}$  can be nonzero only on the fixed part  $\Gamma_{\text{Neu,fix}}$  (but not on the moving part  $\Gamma$ ) then one can still prove the existence of an optimal shape by the method of [22, Theorem 4.3.1].

As mentioned in the introduction, it may be desirable to allow nonzero Neumann data  $\mathbf{g}$  also on the moving part  $\Gamma$  of the boundary (for instance, if we wish to include an external force on the roof surface caused by the weight of snow). In the case of nonzero Neumann data we need to follow our recent studies [23, 28] to obtain results on compactness and the existence of optimal shapes. The main additional difficulty consists in the fact that nonzero Neumann data on the moving part  $\Gamma$  requires correct handling of the 'area measure'  $\mu_{\partial\Omega}|_{\Gamma}$  on this part, because the variational formulation (5) involves integrals with respect to it. A sequence of domains from (19) converges to a domain in that class in the three senses discussed above. However, the natural type of convergence of measures on the boundaries is their weak convergence, and the weak limit of Hausdorff measures is not necessarily a Hausdorff measure, [28, Section 3].

In the next two sections, we present two results regarding this issue. The first, Theorem 4, states that for classes of admissible shapes built on  $\mathcal{O}(D, \varepsilon)$  we can find optimal shapes that realize the infimum of the functional (14). The second, Theorem 5, considers wider parametrized classes of admissible shapes, namely certain uniform domains whose boundaries carry measures satisfying (3). It guarantees the existence of optimal shapes which then indeed realize the minima of (14) over such classes. This may be viewed as a relaxation of the Lipschitz boundary case.

## 5 Lipschitz optimal shapes realizing the infimum of the energy

We follow the method in [28]. Given a bounded Lipschitz domain  $D \subset \mathbb{R}^N$  and  $\varepsilon > 0$  we write, similarly as before,  $\mathcal{O}(D, \varepsilon)$  for the collection of all domains  $\Omega \subset D$  that satisfy the  $\varepsilon$ -cone property. Now suppose that  $G$  is a nonempty open proper subset of  $D$ ,  $\Gamma_{\text{Dir}}$  a compact subset of  $\partial D$  with  $\mathcal{H}^{N-1}(\Gamma_{\text{Dir}}) > 0$  and  $\varepsilon > 0, c_v > 0, \hat{c} > 0$  are given constants.

We define the class

$$\begin{aligned} U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}}) &= \{\Omega \in \mathcal{O}(D, \varepsilon) \mid \lambda^N(\Omega) = c_v, \\ \partial\Omega &= \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}, \mathcal{H}^{N-1}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0, \Gamma_{\text{Dir}} = \partial\Omega \cap \partial D, \Gamma_{\text{Neu}} \subset \bar{G} \\ &\text{and } \mathcal{H}^{N-1}(\Gamma_{\text{Neu}} \cap B(x, r)) \leq \hat{c}r^{N-1} \text{ for any } x \in \Gamma_{\text{Neu}} \text{ and } r > 0\}. \end{aligned} \quad (20)$$

**Remark 6** Since  $D$  is bounded there is some  $\ell_1 > 0$ , depending on  $\hat{c} > 0$ , such that for all  $\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  we have  $\mathcal{H}^{N-1}(\Gamma_{\text{Neu}}) \leq \ell_1$ , and a smaller  $\hat{c}$  leads to a smaller bound  $\ell_1$ . Clearly there is also some uniform lower bound  $\ell_0$  for  $\mathcal{H}^{N-1}(\Gamma_{\text{Neu}})$ , and consequently  $\hat{c}$  must be chosen large enough to ensure the class  $\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  contains more than one domain.

Now suppose that  $\mathbf{f} \in L_2(D)^N$  and  $\mathbf{g} \in [\text{Tr}(W^{1,2}(D))]^N$  are given. For any domain  $\Omega$  and Borel measure  $\mu_{\partial\Omega}$  on  $\partial\Omega$  satisfying the hypotheses of Theorem 2, let  $\mathbf{u}(\Omega, \mu_{\partial\Omega})$  denote the unique weak solution for (2) on  $\Omega$  with  $\mu_{\partial\Omega}$  on  $\partial\Omega$ . Note that in particular each  $\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  satisfies these hypotheses with  $\mu_{\partial\Omega} = \mathcal{H}^{N-1}|_{\partial\Omega}$ , and similarly as before we write  $\mathbf{u}(\Omega, \mathcal{H}^{N-1})$  for the corresponding weak solution.

We consider the functional (14) with  $c_1 \geq 0$  and  $c_2 \geq 0$  in (13). Ideally we would like to minimize  $J(\Omega, \mathcal{H}^{N-1}, \mathbf{u}(\Omega, \mathcal{H}^{N-1}))$  on  $U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$ . However, under the present hypotheses one can only prove the following.

**Theorem 4** Let  $U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  be as in (20),  $c_1 \geq 0$ ,  $c_2 \geq 0$ ,  $\mathbf{f} \in L_2(D)^N$  and  $\mathbf{g} \in [\text{Tr}(W^{1,2}(D))]^N$ .

Then there are a domain  $\Omega_{\text{opt}} \in U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  and a finite Borel measure  $\mu_{\partial\Omega_{\text{opt}}}$  on  $\partial\Omega_{\text{opt}}$ , equivalent to  $\mathcal{H}^{N-1}|_{\partial\Omega_{\text{opt}}}$ , such that  $\mathcal{H}^{N-1}(\Gamma_{\text{Neu,opt}}) \leq \mu_{\partial\Omega_{\text{opt}}}(\Gamma_{\text{Neu,opt}})$  and

$$\begin{aligned} J(\Omega_{\text{opt}}, \mu_{\partial\Omega_{\text{opt}}}, \mathbf{u}(\Omega_{\text{opt}}, \mu_{\partial\Omega_{\text{opt}}})) \\ = \inf_{\Omega \in U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})} J(\Omega, \mathcal{H}^{N-1}, \mathbf{u}(\Omega, \mathcal{H}^{N-1})). \end{aligned}$$

If  $\mu_{\partial\Omega_{\text{opt}}} = \mathcal{H}^{N-1}|_{\partial\Omega_{\text{opt}}}$ , then the infimum is actually a minimum.

**Proof:** Insubstantial modifications of [28, Lemma 3.1] guarantee that for fixed parameters the class  $U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  is compact with respect to the convergence in the sense of characteristic functions, the Hausdorff sense and the sense of compacts. Moreover, each sequence  $(\Omega_n)_n$  in  $U_{\text{ad}}(D, G, \varepsilon, c_v, \hat{c}, \Gamma_{\text{Dir}})$  has a subsequence  $(\Omega_{n_k})_k$  converging to a limit domain  $\Omega_*$  in the Hausdorff sense, the sense of characteristic functions and the sense of compacts and for which the Hausdorff measures  $\mu_{\partial\Omega_{n_k}} = \mathcal{H}^{N-1}|_{\partial\Omega_{n_k}}$  converge weakly to a measure  $\mu_{\partial\Omega_*}$  which is equivalent to  $\mathcal{H}^{N-1}|_{\partial\Omega_*}$  and satisfies

$$\mathcal{H}^{N-1}(\Gamma_{\text{Neu},*}) \leq \mu_{\partial\Omega_*}(\Gamma_{\text{Neu},*}). \quad (21)$$

Here we write  $\Gamma_{\text{Neu},*}$  for the part of  $\partial\Omega_*$  that corresponds to  $\Gamma_{\text{Neu}}$  in (20), below we will use a similar notation with analogous meaning.

Now suppose that this sequence  $(\Omega_n)_n \subset \hat{U}_{\text{ad}}(D, G, \varepsilon, \hat{c}, c_v, \ell_1)$  is a minimizing sequence of the energy functional (14), this minimizing sequence exists since the functional is nonnegative. We relabel and denote the subsequence  $(\Omega_{n_k})_k$  above by  $(\Omega_k)_k$ . Let  $\mathbf{u}_k := \mathbf{u}(\Omega_k, \mathcal{H}^{N-1})$  denote the unique weak solution to (2) on  $\Omega_k$  and let  $\mathbf{u}_* := \mathbf{u}(\Omega_*, \mu_{\partial\Omega_*})$  denote that on  $\Omega_*$  with measure  $\mu_{\partial\Omega_*}$  on the boundary.

Since all domains  $\Omega_k$  have Lipschitz boundaries of finite Hausdorff measures, there are extension operators  $E : W^{1,2}(\Omega_k)^N \rightarrow W^{1,2}(\mathbb{R}^N)^N$  and a constant  $C_E > 0$  independent of  $k$  such that  $\|E\mathbf{u}_k\|_{W^{1,2}(D)^N} \leq C_E \|\mathbf{u}_k\|_{W^{1,2}(\Omega_k)^N}$  for all  $k$ , see [11]. By Lemma 1 (ii) and Theorem 2 (together with Theorem 1) we can find a constant  $C > 0$ , independent of  $k$ , such that

$$\sup_k \|\mathbf{u}_k\|_{W^{1,2}(\Omega_k)^N} \leq C (\|\mathbf{f}\|_{L_2(D)^N} + \|\mathbf{g}\|_{\text{Tr}(W^{1,2}(D))^N}).$$

This means that  $(E\mathbf{u}_k|_D)_k$  is bounded in  $W^{1,2}(D)^N$ , and consequently there exists  $\mathbf{u}^* \in W^{1,2}(D)^N$  such that  $E\mathbf{u}_k|_D \rightharpoonup \mathbf{u}^*$  in  $W^{1,2}(D)^N$ .

We define linear functionals  $F_k$  and  $F$  on  $V(\Omega, \Gamma_{\text{Dir}})^N$  by

$$F_k[\phi] = \int_{\Omega_k} \sigma(\mathbf{u}_k) : e(\phi) dx - \int_{\Omega_k} \mathbf{f} \cdot \phi - \int_{\Gamma_{\text{Neu},k}} \mathbf{g} \cdot \phi d\mathcal{H}^{N-1},$$

$\phi \in V(\Omega, \Gamma_{\text{Dir}})^N$ , and

$$F[\phi] = \int_{\Omega} \sigma(\mathbf{u}^*) : e(\phi) dx - \int_{\Omega_*} \mathbf{f} \cdot \phi - \int_{\Gamma_{\text{Neu},*}} \mathbf{g} \cdot \phi d\mu^*.$$

Following the proof of [28, Theorem 3.2] one can see that  $F[\phi] = \lim_k F_k[\phi] = 0$  for all  $\phi \in V(D, \Gamma_{\text{Dir}})^N$ , which implies that  $F[\phi] = 0$ ,  $\phi \in V(D, \Gamma_{\text{Dir}})^N$ . By the uniqueness established in Theorem 2 we must have  $\mathbf{u}^*|_{\Omega_*} = \mathbf{u}_*$ .

Using (5) it is not difficult to show that

$$\lim_k \|\mathbf{1}_{\Omega_k} \nabla E\mathbf{u}_k\|_{L^2(D)^{N \times N}} = \|\mathbf{1}_{\Omega} \nabla E\mathbf{u}^*\|_{L^2(D)^{N \times N}}.$$

Since by the Rellich-Kondrachov theorem also  $\lim_k E\mathbf{u}_k = \mathbf{u}^*$  strongly in  $L^2(D)^N$ , it follows that  $\lim_k J(\Omega_k, \mathcal{H}^{N-1}, \mathbf{u}_k) = J(\Omega_*, \mu_{\partial\Omega_*}, \mathbf{u}_*)$ , so that  $\Omega_{\text{opt}} := \Omega_*$  and  $\mu_{\Omega_{\text{opt}}} = \mu_{\partial\Omega_*}$  are as desired. See [28] for more details.  $\square$

## 6 Relaxation and optimal shapes realizing the energy minimum

Choosing a larger class of admissible shapes, we can ensure the existence of an optimal shape which actually realizes the minimum of (14). Following [23] we relax the restriction on the domains  $\Omega$  to be Lipschitz domains and the restriction on the measures on the boundaries to be Hausdorff measures.

Let  $D \subset \mathbb{R}^N$  be a bounded Lipschitz domain and  $\varepsilon > 0$ . By  $\hat{\mathcal{O}}(D, \varepsilon)$  we denote the collection of all  $(\varepsilon, \infty)$ -domains  $\Omega \subset D$ . We require the measures on the boundary to satisfy the upper regularity condition (3) and a second, somewhat complementary scaling condition. Given  $s > 0$  and a Borel measure  $\mu_F$  on  $\mathbb{R}^N$  with  $F = \text{supp } \mu_F$  we say that  $\mu_F$  is *lower s-regular in the closed ball sense* if there is a constant  $\bar{c}_s > 0$  such that

$$\mu_F(\overline{B(x, r)}) \geq \bar{c}_s r^s, \quad x \in F, \quad 0 < r \leq 1. \quad (22)$$

Now let  $D_0$  be a non-empty Lipschitz domain and a proper subset of  $D$  and suppose that  $\Gamma_{\text{Dir}} \subset \partial D \cap \partial D_0$  has positive measure  $\mathcal{H}^{N-1}(\Gamma_{\text{Dir}}) > 0$ . Given  $\varepsilon > 0$ ,  $N-1 \leq s < N$ ,  $0 \leq d \leq s$ ,  $\bar{c}_s > 0$  and  $c_d > 0$  let

$$\begin{aligned} U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}}) &:= \{(\Omega, \mu_{\partial\Omega}) \mid \Omega \in \hat{\mathcal{O}}(D, \varepsilon), D_0 \subset \Omega, \\ \lambda^N(\Omega) &= c_v, \partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}, \Gamma_{\text{Dir}} = \partial\Omega \cap \partial D, \mu_{\partial\Omega} = \mathcal{H}^{N-1}|_{\Gamma_{\text{Dir}}} + \mu_{\Gamma_{\text{Neu}}}, \\ \mu_{\Gamma_{\text{Neu}}} &\text{ with } \text{supp } \mu_{\Gamma_{\text{Neu}}} = \Gamma_{\text{Neu}} \text{ satisfies (3) and (22), } \mu_{\partial\Omega}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0\} \end{aligned} \quad (23)$$

Using [23, Theorem 3] it can be seen that  $U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  is compact with respect to the convergence in the Hausdorff sense, the sense of compacts, the sense of characteristic functions and the sense of weak convergence of the measures  $\mu_{\partial\Omega}$  on the boundaries. By Lemma 1 (ii) and Theorem 2 we have (12) with a uniform constant for all  $(\Omega, \mu_{\partial\Omega}) \in U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$ . This, together with uniform estimates for the norms of extension operators, allows to obtain the following result by similar arguments as in the Lipschitz case.

**Theorem 5** Let  $U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  be as in (23),  $c_1 \geq 0$  and  $c_2 \geq 0$  in (13). Let  $\mathbf{f} \in L_2(D)^N$  and  $\mathbf{g} \in [\text{Tr}(W^{1,2}(D))]^N$ .

Then there is  $(\Omega_{\text{opt}}, \mu_{\partial\Omega_{\text{opt}}}) \in U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  such that

$$\begin{aligned} J(\Omega_{\text{opt}}, \mu_{\partial\Omega_{\text{opt}}}, \mathbf{u}(\Omega_{\partial\Omega_{\text{opt}}}, \mu_{\partial\Omega_{\text{opt}}})) \\ = \min_{(\Omega, \mu_{\partial\Omega}) \in U_{\text{ad}}(D, D_0, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})} J(\Omega, \mu_{\partial\Omega}, \mathbf{u}(\Omega, \mu_{\partial\Omega})). \end{aligned}$$

Slight modifications of Theorems 4 and 5 yield generalizations of Theorem 3 on optimal roof shapes, such as the following. Let  $N = 3$  and let  $D \subset \mathbb{R}^3$  be a bounded Lipschitz domain,  $G$  a nonempty open proper subset of  $D$  and  $\Gamma_{\text{Dir}}$  a compact subset of  $\partial D$  with  $\mathcal{H}^2(\Gamma_{\text{Dir}}) > 0$  as in Section 4. Consider the class

$$\begin{aligned} U_{\text{ad}}(D, G, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}}) &= \{(\Omega, \mu_{\partial\Omega}) \mid \Omega \in \hat{\mathcal{O}}(D, \varepsilon), \lambda^3(\Omega) = c_\nu, \\ &\partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma_{\text{Neu}}, \Gamma_{\text{Dir}} = \partial\Omega \cap \partial D, \mu_{\partial\Omega} = \mathcal{H}^{N-1}|_{\Gamma_{\text{Dir}}} + \mu_{\Gamma_{\text{Neu}}}, \\ &\mu_{\Gamma_{\text{Neu}}} \text{ with supp } \mu_{\Gamma_{\text{Neu}}} = \Gamma_{\text{Neu}} \text{ satisfies (3) and (22), } \mu_{\partial\Omega}(\Gamma_{\text{Dir}} \cap \Gamma_{\text{Neu}}) = 0, \\ &\Gamma_{\text{Neu}} = \Gamma_{\text{lower}} \cup \Gamma_{\text{upper}}, \Gamma_{\text{lower}} \subset \bar{G}, \Gamma_{\text{upper}} = \Gamma_{\text{lower}} + h_z \mathbf{e}_z\}. \end{aligned} \quad (24)$$

This class is compact with respect to the same four types of convergences as (23). (We point out that [23, Theorem 3] can be used without problems because the domains  $\Omega$  are nonempty by construction.) The following result, which roughly speaking is a corollary of Theorem 5, generalizes Theorem 3 to the case of possibly nonzero Neumann data  $\mathbf{g}$  on the moving parts.

**Corollary 1** Let  $U_{\text{ad}}(D, G, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  be as in (24),  $\rho_0 > 0$ ,  $c_1 \geq 0$  and  $c_2 \geq 0$ . For each  $(\Omega, \mu_{\partial\Omega}) \in U_{\text{ad}}(D, G, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  let  $\mathbf{u}(\Omega, \mu_{\partial\Omega})$  denote the unique weak solution to (2) with  $\mathbf{g} \in [\text{Tr}(W^{1,2}(D))]^3$  and  $\mathbf{f} \equiv \mathbf{f}(\Omega)$  defined as in (16). Let  $J$  be the energy functional defined in (13).

Then there is an optimal shape  $(\Omega_{\text{opt}}, \mu_{\partial\Omega_{\text{opt}}}) \in U_{\text{ad}}(D, G, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$  realizing the minimum of (14) over  $U_{\text{ad}}(D, G, \varepsilon, c_v, s, d, \bar{c}_s, c_d, \Gamma_{\text{Dir}})$ .

## 7 Conclusions

For the linear elasticity system (2) with homogeneous Neumann conditions on the varying parts, optimal (energy minimizing) roof shapes always exist in classes (18) based on the well-known classes (17) of bounded Lipschitz domains, Theorem 3. This result had not been known before, but it relies on classical methods. Suppose inhomogeneous Neumann conditions on the varying parts are imposed. In that case, optimal shapes realizing the infimum of the energy always exist in classes (20) of bounded Lipschitz domains with generalized boundary measures. This result is new, and it was obtained using the novel methods in [28]. To ensure the existence of optimal shapes that realize the minimum of the energy under inhomogeneous Neumann conditions, we need to allow fractal boundaries and discuss measures (not only shapes). If we do so, energy minimizing optimal shapes in classes (23) of bounded uniform domains always exist, Theorem 5, and in particular, optimal roof shapes exist, Corollary 1. This result is new, and it is based on the completely novel approach introduced in [23]. Future work will include more specific numerical experiments and also theoretical generalizations, for instance, existence results for optimal shapes for Robin boundary value problems in elasticity.

## A Green's formulas

We provide versions of Green's formulas in the context of Theorem 1. They are not used explicitly in our results, but they provide the correct justification of the variational formu-

lation (2). Consider the space

$$H(\operatorname{div}, \Omega) = \{\mathbf{u} = (u_1, \dots, u_N)^t : \Omega \rightarrow \mathbb{R}^N \mid u_1, \dots, u_N \in L^2(\Omega), \operatorname{div} \mathbf{u} \in L^2(\Omega)\},$$

it is a Hilbert space corresponding to the norm

$$\|\mathbf{u}\|_{\operatorname{div}, \Omega} = \left( \|\mathbf{u}\|_{L^2(\Omega)^N}^2 + \|\operatorname{div} \mathbf{u}\|_{L^2(\Omega)}^2 \right)^{\frac{1}{2}}.$$

**Proposition 2** *Let  $\Omega \subset \mathbb{R}^N$  be a bounded  $W^{1,2}$ -extension domain. Suppose that  $\mu_{\partial\Omega}$  is a Borel measure with  $\operatorname{supp} \mu_{\partial\Omega} = \partial\Omega$  and such that (3) holds with some  $N - 2 < d \leq N$ . For all  $\mathbf{u} \in H(\operatorname{div}, \Omega)$  we can define a bounded linear functional  $\mathbf{u} \cdot \mathbf{n} \in (\operatorname{Tr}(W^{1,2}(\Omega)))'$  by the identity*

$$\langle \mathbf{u} \cdot \mathbf{n}, \operatorname{Tr} w \rangle_{(\operatorname{Tr}(W^{1,2}(\Omega)))', \operatorname{Tr}(W^{1,2}(\Omega))} = \int_{\Omega} \mathbf{u} \cdot \nabla w \, dx - \int_{\Omega} (\operatorname{div} \mathbf{u}) w \, dx, \quad (25)$$

$w \in W^{1,2}(\Omega)$ .

For all  $T \in H(\operatorname{div}, \Omega)^N$  we can define a bounded linear functional  $T \cdot \mathbf{n} \in [(\operatorname{Tr}(W^{1,2}(\Omega)))']^N$  by

$$\langle T \cdot \mathbf{n}, \operatorname{Tr} \theta \rangle_{[(\operatorname{Tr}(W^{1,2}(\Omega)))']^N, [\operatorname{Tr}(W^{1,2}(\Omega))]^N} = \int_{\Omega} T : \nabla \theta \, dx - \int_{\Omega} (\operatorname{div} T) \cdot \theta \, dx, \quad (26)$$

$\theta \in W^{1,2}(\Omega)^N$ .

For the special case that  $\Omega$  is a Lipschitz domain and  $\mu_{\partial\Omega}$  equals  $\mathcal{H}^{N-1}$  a proof of formula (25) can be found in [21, Theorem 2.5 and formula (2.17)]. The same proof, combined with Theorem 1, yields (25). (Note also that a particular version of (25) had been proved in [13] for a specific domain  $\Omega$ .) Formula (26) is just its matrix form and an immediate consequence of (25). See [12, p. 69] for a version of this formula for smooth domains and tensors.

## B A technical remark

We comment briefly on how to see the claimed identity (10) in the proof of Lemma 1, i.e., that  $\widetilde{E_{\Omega} \mathbf{u}} = 0$   $\mu_{\partial D}$ -a.e. on  $\Gamma_{\operatorname{Dir}}$ . One can argue as follows: If  $u_i$  are the components of  $\mathbf{u}$  then the vector fields  $\mathbf{u}^{(M)}$ ,  $M \geq 1$ , with components  $u_i^{(M)} := (-M) \vee (u_i \wedge M)$  satisfy  $\lim_{M \rightarrow \infty} \|\mathbf{u}^{(M)} - \mathbf{u}\|_{W^{1,2}(\Omega)^N} = 0$ , what by (9) is inherited to their extensions. Therefore (and by Theorem 1) it suffices to show (10) for  $\mathbf{u}$  with bounded components. If  $\chi$  is a smooth cut-off function with compact support inside an open ball  $B \supset \overline{D}$  then we have  $(\chi E_{\Omega} \mathbf{u})^{\sim} = \widetilde{E_{\Omega} \mathbf{u}}$  on  $\overline{D}$ . Since  $L^{\infty}$ -functions with finite Dirichlet integral form an algebra and Poincaré's inequality holds for  $B$ , the product  $\chi E_{\Omega} \mathbf{u}$  is an element of  $\dot{W}^{1,2}(B)$ , and we can use standard capacity arguments based on this space instead of  $W^{1,2}(\mathbb{R}^N)$ . In particular, adequate modifications of [9, Theorems 5.2 and 5.4] now allow to conclude (10) similarly as in [9, Theorem 6.1].

## C On the Mosco convergence of energy functionals

In this auxiliary section we consider energy functionals for Robin problems of type

$$\begin{cases} -\operatorname{div} e(\mathbf{u}) &= \mathbf{f} & \text{in } \Omega, \\ e(\mathbf{u}) \cdot \mathbf{n} + \alpha \mathbf{1}_{\Gamma} \mathbf{u} &= 0 & \text{on } \partial\Omega, \end{cases}$$

where  $\alpha \geq 0$  and  $\Gamma$  is a subset of  $\partial\Omega$ . We show that if a sequence of domains converges in a suitable sense then these energy functionals on the domains converge in the sense of Mosco [36].

Let  $H$  be a Hilbert space. Recall from [36, Definition 2.1.1] that a sequence of quadratic forms  $a_n(\cdot, \cdot) : H \times H \rightarrow (-\infty, +\infty]$  is said to  $M$ -converge on  $H$  to a quadratic form  $a(\cdot, \cdot) : H \times H \rightarrow (-\infty, +\infty]$  if

1. For every  $u \in H$  there is a sequence  $(u_n)_n \subset H$ , convergent to  $u$  and such that  $\underline{\lim}_{n \rightarrow \infty} a_n(u_n, u_n) \leq a(u, u)$ .
2. For every sequence  $(v_n)_n \subset H$  converging weakly to  $u \in H$  we have  $\underline{\lim}_{n \rightarrow \infty} a_n(v_n, v_n) \geq a(u, u)$ .

Let  $N \geq 2$  and let  $\Omega \subset \mathbb{R}^N$  be a bounded domain. Suppose that  $\mu_{\Gamma_{\text{Dir}}}$  and  $\mu_{\Gamma}$  are Borel measures with  $\text{supp } \mu_{\Gamma_{\text{Dir}}} = \Gamma_{\text{Dir}}$  and  $\text{supp } \mu_{\Gamma} = \Gamma$  such that  $\partial\Omega = \Gamma_{\text{Dir}} \cup \Gamma$  and  $\Gamma_{\text{Dir}} \cap \Gamma$  is a zero set for both  $\mu_{\Gamma_{\text{Dir}}}$  and  $\mu_{\Gamma}$ . For each  $n \geq 1$  let  $\Omega_n \subset \mathbb{R}^N$  a domain and let  $\mu_{\Gamma_n}$  be a Borel measure with  $\text{supp } \mu_{\Gamma_n} = \Gamma_n$  such that  $\partial\Omega_n = \Gamma_{\text{Dir}} \cup \Gamma_n$  and  $\Gamma_{\text{Dir}} \cap \Gamma_n$  is a zero set for both  $\mu_{\Gamma_{\text{Dir}}}$  and  $\mu_{\Gamma_n}$ . If all domains are  $W^{1,2}$ -extension domains contained in a bounded Lipschitz domain  $D$  and all  $\mu_{\Gamma_{\text{Dir}}}$ ,  $\mu_{\Gamma}$  and  $\mu_{\Gamma_n}$  satisfy (3) with  $N - 2 < d \leq N$  then for any non-negative real numbers  $\alpha$  and  $\alpha_n$  we can define quadratic forms on  $L^2(D)^N$  by

$$a_n(\mathbf{u}, \mathbf{u}) = \begin{cases} \int_{\Omega_n} e(\mathbf{u}) : e(\mathbf{u}) dx + \int_{\Omega_n} |\mathbf{u}|^2 dx + \alpha_n \int_{\Gamma_n} |\text{Tr} \mathbf{u}|^2 d\mu_{\Gamma_n}, & \mathbf{u} \in V(\Omega_n, \Gamma_{\text{Dir}})^N, \\ +\infty, & \text{otherwise} \end{cases}$$

and

$$a(\mathbf{u}, \mathbf{u}) = \begin{cases} \int_{\Omega} e(\mathbf{u}) : e(\mathbf{u}) dx + \int_{\Omega} |\mathbf{u}|^2 dx + \alpha \int_{\Gamma} |\text{Tr} \mathbf{u}|^2 d\mu_{\Gamma}, & \mathbf{u} \in V(\Omega, \Gamma_{\text{Dir}})^N, \\ +\infty, & \text{otherwise.} \end{cases}$$

The following convergence result can be obtained by the same arguments as [23, Theorem 8].

**Theorem 6** *Let  $\varepsilon > 0$ , let  $(\Omega_n)_n$  be a sequence of  $(\varepsilon, \infty)$ -domains  $\Omega_n \subset D$ . Let  $\Gamma_{\text{Dir}}$ ,  $\Gamma_n$  and  $\mu_{\Gamma_{\text{Dir}}}$ ,  $\mu_{\Gamma_n}$  be as above and assume that all  $\mu_{\Gamma_n}$  satisfy (3) with the same exponent  $N - 1 < d \leq N$  and the same constant  $c_d$ .*

*If  $\lim_n \Omega_n = \Omega$  in the Hausdorff sense and in the sense of characteristic functions and  $\lim_n \alpha_n \mu_n = \alpha \mu$  in the sense of weak convergence, then we have*

$$\lim_n a_n = a$$

*in the sense of  $M$ -convergence on  $L^2(D)$ .*

**Remark 7** *For simplicity, and because it is sufficient for many practical purposes,  $\Gamma_{\text{Dir}}$  is assumed to be fixed in the above setup. It would be sufficient to keep the set  $\Gamma_{\text{Dir}} \cap \Gamma_n$  fixed, see [15, Theorem 10].*

## References

- [1] *Fractal wall (TM)*. Brevet Colas - École Polytechnique 03/00881, Noise abatement wall, US patent 7308965B2.
- [2] D. R. ADAMS AND L. I. HEDBERG, *Function Spaces and Potential Theory*, vol. 314 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 1996.
- [3] S. AGMON, *Lectures on Elliptic Boundary Value Problems*, Van Nostrand Math. Studies, Van Nostrand, Princeton, 1965.
- [4] G. ALLAIRE, *Conception Optimale de Structures*, vol. 58 of Mathématiques et Applications, Springer, Berlin, 2007.



- [5] W. ARENDT AND M. WARMA, *The Laplacian with Robin boundary conditions on arbitrary domains*, Pot. Anal., 19 (2003), pp. 341–363.
- [6] K. ARFI AND A. ROZANOVA PIERRAT, *Dirichlet-to-Neumann or Poincaré-Steklov operator on fractals described by  $d$ -sets*, Discrete and Continuous Dynamical Systems - S, 12 (2019), pp. 1–26.
- [7] J. AZZAM, S. HOFMANN, J. M. MARTELL, K. NYSTRÖM, AND T. TORO, *A new characterization of chord-arc domains*, J. Eur. Math. Soc. (JEMS), 19 (2017), pp. 967–981.
- [8] C. BARDOS, D. GREBENKOV, AND A. ROZANOVA-PIERRAT, *Short-time heat diffusion in compact domains with discontinuous transmission boundary conditions*, Math. Models Methods Appl. Sci., 26 (2016), pp. 59–110.
- [9] M. BIEGERT, *On traces of Sobolev functions on the boundary of extension domains*, Proc. of the Amer. Math. Soc., 137 (2009), pp. 4169–4176.
- [10] K. BURDZY, Z.-Q. CHEN, D. MARSHALL, AND K. RAMANAN, *Obliquely reflected Brownian motion in nonsmooth planar domains*, Ann. Probab., 45 (2017), pp. 2971–3037.
- [11] D. CHENAIS, *On the existence of a solution in a domain identification problem*, Journal of Mathematical Analysis and Applications, 52 (1975), pp. 189–219.
- [12] P. G. CIARLET, *Mathematical Elasticity, vol.1. Three-dimensional Elasticity*, Studies in Mathematics and its Applications, North-Holland, Amsterdam, 1988.
- [13] S. CREO, M. R. LANCIA, P. VERNOLE, M. HINZ, AND A. TEPLYAEV, *Magnetostatic problems in fractal domains*, in Analysis, Probability and Mathematical Physics on Fractals, P. Alonso-Ruiz, J. P. Chen, L. G. Rogers, R. S. Strichartz, and A. Teplyaev, eds., vol. 5 of Fractals and Dynamics in Mathematics, Science, and the Arts: Theory and Applications, World Scientific, 2020, pp. 477–502.
- [14] C. DAPOGNY, A. FAURE, G. MICHAILIDIS, G. ALLAIRE, A. COUVELAS, AND R. ESTEVEZ, *Geometric constraints for shape and topology optimization in architectural design*, Computational Mechanics, 59 (2017), pp. 933–965.
- [15] A. DEKKERS, A. ROZANOVA PIERRAT, AND A. TEPLYAEV, *Mixed boundary valued problem for linear and nonlinear wave equations in domains with fractal boundaries*, Submitted. Preprint hal-02514311, (2020).
- [16] R. DURAN AND M. MUSCHIETTI, *The Korn inequality for Jones domains*, Electron. J. Differential Equations, [electronic only] (2004). Paper No. 127, 10 p.
- [17] M. EGERT, R. HALLER-DINTELMANN, AND J. REHBERG, *Hardy’s inequality for functions vanishing on a part of the boundary*, Pot. Anal., 43 (2015), pp. 49–78.
- [18] L. C. EVANS, *Partial Differential Equations*, vol. 19 of Graduate Studies in Mathematics, Amer. Math. Soc., Providence, second ed., 2010.
- [19] K. J. FALCONER, *The Geometry of Fractal Sets*, vol. 85 of Cambridge Tracts Math., Cambridge Univ. Press, Cambridge, 1985.
- [20] J. GERO, *Architectural optimization - a review*, Engineering Optimization, 1 (1975), pp. 189–199.
- [21] V. GIRAULT AND P.-A. RAVIART, *Finite Element Methods for the Navier-Stokes Equations, Theory and Algorithms*, Springer, New York, 1986.

- [22] A. HENROT AND M. PIERRE, *Shape Variation and Optimization*, vol. 28 of EMS Tracts in Mathematics, European Mathematical Society (EMS), Zürich, 2018.
- [23] M. HINZ, A. ROZANOVA PIERRAT, AND A. TEPLYAEV, *Non-Lipschitz uniform domain shape optimization in linear acoustics*, SIAM J. Control Optim., in press, (2020).
- [24] P. W. JONES, *Quasi conformal mappings and extendability of functions in Sobolev spaces*, Acta Mathematica, 147 (1981), pp. 71–88.
- [25] A. JONSSON, *Besov spaces on closed sets by means of atomic decomposition*, Complex Variables and Elliptic Equations, 54 (2009), pp. 585–611.
- [26] M. R. LANCIA, *A Transmission Problem with a Fractal Interface*, Zeitschrift für Analysis und ihre Anwendungen, 21 (2002), pp. 113–133.
- [27] M. L. LAPIDUS AND C. POMERANCE, *Counterexamples to the modified Weyl-Berry conjecture on fractal drums*, Math. Proc. Cambridge Philos. Soc., 119 (1996), pp. 167–178.
- [28] F. MAGOULÈS, T. P. K. NGUYEN, P. OMNÈS, AND A. ROZANOVA PIERRAT, *Optimal absorption of acoustical waves by a boundary*, SIAM J. Control Optim., 59 (2021), pp. 561–583.
- [29] F. MAGOULÈS, P. NGYUEN, P. OMNES, AND A. ROZANOVA PIERRAT, *Optimal and efficient shapes in acoustic boundary absorption*, Preprint hal-02543993, (2020).
- [30] M. MARIN, *Some basic theorems in elastostatics of micropolar materials with voids*, J. Comp. Appl. Math., 70 (1996), pp. 115–126.
- [31] M. MARIN, I. ABBAS, S. VLASE, AND E. CRACIUN, *A study of deformations in a thermoelastic dipolar body with voids*, Symmetry, 12 (2020), p. 267.
- [32] M. MARIN, E. CRACIUN, AND N. POP, *Considerations on mixed initial-boundary value problems for micropolar porous bodies*, Dyn. Syst. Appl., 25 (2016), pp. 175–196.
- [33] V. MAZ’JA, *Sobolev Spaces*, Springer Ser. Sov. Math., Springer-Verlag, Berlin, 1985.
- [34] J. MICHALEK, R. CHOUDHARY, AND P. PAPALAMBROS, *Architectural layout design optimization*, Engineering Optimization, 34 (2002), pp. 461–484.
- [35] S. MOLCHANOV AND B. VAINBERG, *On spectral asymptotics for domains with fractal boundaries*, Comm. Math. Phys., 183 (1997), pp. 85–117.
- [36] U. MOSCO, *Convergence of convex sets and of solutions of variational inequalities*, Advances in Mathematics, 3 (1969), pp. 510–585.
- [37] M. OSTWALD, J. VAUGHAN, AND C. TUCKER, *Characteristic Visual Complexity: Fractal dimensions in the architecture of Frank Lloyd Wright and Le Corbusier*, in Nexus VII: Architecture and Mathematics, K. Williams, ed., Kim Williams Books, Turin, 2008, pp. 217–231.
- [38] L. G. ROGERS, *Degree-independent Sobolev extension on locally uniform domains*, Journal of Functional Analysis, 235 (2006), pp. 619–665.
- [39] A. ROZANOVA PIERRAT, *Wave propagation and fractal boundary problems: mathematical analysis and applications*, HDR, Université Paris-Saclay, France, 2020.
- [40] ———, *Generalization of Rellich-Kondrachov theorem and trace compactness in the framework of irregular and fractal boundaries*, in Fractals in engineering: Theoretical aspects and Numerical approximations, M. R. Lancia and A. Rozanova Pierrat, eds., vol. 8 of ICIAM 2019 SEMA SIMAI Springer Series, Springer Intl. Publ., 2021.

- [41] A. ROZANOVA-PIERRAT, D. S. GREBENKOV, AND B. SAPOVAL, *Faster diffusion across an irregular boundary*, Phys. Rev. Lett., 108 (2012), p. 240602.
- [42] S. VLASE, M. MARIN, A. ÖCHSNER, AND M. L. SCUTARO, *Motion equation for a flexible one-dimensional element used in the dynamical analysis of a multibody system*, Continuum. Mech. Thermodyn., 31 (2019), pp. 715–724.
- [43] H. WALLIN, *The trace to the boundary of Sobolev spaces on a snowflake*, Manuscripta Math., 73 (1991), pp. 117–125.
- [44] T. WORTMANN, *Model-based Optimization for Architectural Design: Optimizing Daylight and Glare in Grasshopper*, Technology | Architecture + Design, 1 (2017), pp. 176–185.
- [45] C. ZANCHETTIN, T. B. LUDERMIR, AND L. M. ALMEIDA, *Hybrid Training Method for MLP: Optimization of Architecture and Training*, IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics), 41 (2011), pp. 1097–1109.
- [46] M. ZAWIDZKI, *Discrete Optimization in Architecture*, Springer, Singapore, 2017.