



HAL
open science

OSL chronology of socio-ecological systems during the mid-Holocene in the eastern coast of the Sultanate of Oman (Arabian Peninsula)

Mailys Richard, Norbert Mercier, Vincent Charpentier, Jean-François Berger

► To cite this version:

Mailys Richard, Norbert Mercier, Vincent Charpentier, Jean-François Berger. OSL chronology of socio-ecological systems during the mid-Holocene in the eastern coast of the Sultanate of Oman (Arabian Peninsula). *Journal of Archaeological Science: Reports*, 2020, 33, pp.102465. 10.1016/j.jasrep.2020.102465 . hal-02966603

HAL Id: hal-02966603

<https://hal.science/hal-02966603>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **OSL chronology of socio-ecological systems during the mid-Holocene in the eastern**
2 **coast of the Sultanate of Oman (Arabian Peninsula)**

3

4 Richard M.¹, Mercier N.¹, V. Charpentier², Berger J.-F.³

5

6 ¹ Institut de Recherche sur les Archéomatériaux - Centre de Recherche en Physique Appliquée
7 à l'Archéologie, UMR 5060 CNRS, Université Bordeaux-Montaigne, 33607 Pessac, France

8 ² Institut national de recherches archéologiques préventives, ARSCAN, UMR 7041 CNRS,
9 Université Paris X, 92023 Nanterre, France

10 ³ CNRS, Environnement Ville Société - Institut de Recherches Géographiques, UMR 5600
11 CNRS, Université de Lyon, 69676 BRON, France

12

13

14 **Abstract**

15

16 The eastern coast of the Arabian Peninsula is a key region to explore past climatic and
17 environmental changes and their impact on the coastal ecosystems and Neolithic communities
18 during the mid-Holocene (~8000-5000 years ago). Indeed, it is located at the frontier zone
19 between dry and wet tropical systems, where the monsoon has reduced its activity and
20 gradually shifted southward from the 6th millennium BC. Past human adaptation to new
21 ecological conditions (more arid) and resources is reflected in the archaeological record, as a
22 response to important climatic and eustatic fluctuations.

23 In order to correlate human occupation during the Neolithic with the local climatic
24 and palaeoenvironmental records, sediments were sampled at two archaeological sites and
25 surrounding sabkhas, located in the Ja'alan region, on the eastern coast of the Sultanate of
26 Oman. Optically and infrared stimulated luminescence dating were applied to quartz and
27 feldspar grains extracted from sediments coming from both archaeological (Ruways 1 and
28 Suwayh 1 shell middens) and palaeoenvironmental (current sabkhas) contexts in order to
29 provide, together with radiocarbon dating, a chronology for this area. The results obtained
30 allowed constraining human occupation during the Neolithic between 8340 ± 480 and $5390 \pm$
31 290 years (~6800-3100 years BC) at Ruways 1, and between 6180 ± 310 and 5480 ± 270
32 years BP (~4500-3200 years BC) at Suwayh 1. These occupations coincide with the
33 development of a lagoon/mangrove, recorded in sediments from cores drilled under the

34 current sabkha, whose last stage of development is dated to 5320 ± 590 (C2) and 5220 ± 570
35 (C1) years (3900-2600 years BC) at Khor al Jaramah.

36

37 **1. Introduction**

38

39 The study of past groups of coastal fisher-gatherer-herders in their palaeoenvironmental
40 context is of great interest to understand the resilience of these populations facing important
41 climatic changes during the mid-Holocene in the Arabian Peninsula. The eastern coast of the
42 Arabian Peninsula, and more specifically the Ja'alan region in the Sultanate of Oman (Fig. 1),
43 presents several important archaeological sites documenting human occupation starting from
44 the middle of the 6th millennium BC (~8000-7000 years; Charpentier 2008; Berger et al.,
45 2005). During the mid-Holocene, this territory suffered climatic (Indian-Arabian monsoon),
46 eustatic and tectonic fluctuations, which might have directly affected Neolithic populations
47 that exploited marine, lagoon and deltaic resources (Berger et al., 2005; Berger et al., 2013).

48 More specifically, along the Oman coast, a particular Neolithic economy flourished
49 during the first half of the mid-Holocene, without pottery and agriculture but with a
50 significant component of marine resources associated with the presence of a few domesticated
51 animals (Crassard and Dreschler, 2013). Coastal environments and marine resources have
52 played a major role in the Neolithic development of eastern Arabia, as an apparent resilience
53 to climatic variations during the mid-Holocene. After the onset of the humid period around
54 9000 years (end of the early Holocene), this sub-arid to arid zone suffered rapidly changing
55 environmental conditions, with past latitudinal monsoon circulation and upwelling activity
56 changes through time (Jung et al., 2004, Fleitmann et al., 2007, Parker et al., 2008) that
57 impacted coastal paleoenvironments and biomass productivity (Lézine et al. 2010; Berger et
58 al. 2013; Bassinot et al. 2011).

59 The effect of climate change on Neolithic populations over a long time, inducing
60 environmental and economic stress, can be investigated by cross-checking data from human
61 occupation in shell middens with palaeoenvironmental records. The evolution of coastal
62 environments can be reconstructed through the study of transects in sections and cores in
63 archaeological sites and surrounding sabkhas. This study focuses on establishing the
64 chronology of the occupation phases of stratified shell middens that were produced by the
65 interaction of aeolian and anthropogenic processes (Ruways 1 and Suwayh 1 archaeological
66 sites), and the chronology of coastal environmental archives that are preserved under the

67 current surrounding sabkhas (core drilling), with the goal of studying past human behaviour in
68 its palaeoenvironmental context.

69

70 **2. Archaeological and palaeoenvironmental contexts**

71 Ruways 1 (RWY-1) and Suwayh 1 (SWY-1) (governorate of al-Sharqqiyah, Oman)
72 are located along the coast of the Arabian sea, in a vast plain comprised of sabkhas, which are
73 seasonally fed by wadis coming from large quaternary alluvial fans at the bottom of the Jebel
74 Khamis (Fig. 1). RWY-1 is a site featuring a long sequence of occupations (dated by
75 radiocarbon between 6380 and 3400 years BC) with one potentially dating to the beginning of
76 the middle Neolithic, which is remarkable in a stratified context in Ja'alan (Berger et al.,
77 2020). The RWY-1 and SWY-1 sites present great potential for reconstructing the economic
78 strategies of Neolithic coastal populations, strongly oriented towards the exploitation of
79 lagoons and mangrove ecosystems (Martin, 2005; Berger et al., 2020).

80 Under the current sabkhas, preserved palaeo-mangroves and lagoons seem to be
81 correlated with mid-Holocene fossil deltas and estuaries. Episodes of marine transgression
82 alternated with mangrove and/or lagoon phases are recorded in stratigraphic profiles at the
83 Suwayh sabkha (Berger et al., 2005). In particular, during the 5th millennium BC (around
84 7000-6000 years), Neolithic sites are associated to an optimum period for mangrove
85 development, due to sea-level stability (Berger et al., 2013). Around 5000-4500 years, an arid
86 period led to the expansion of sabkhas and the regression of mangroves (Berger et al., 2013).
87 This climatic setting may have affected both settlement networks and economic strategies of
88 the Neolithic populations that occupied the area. To investigate the impact of such
89 environmental and climatic changes on past human populations, cores were drilled at the
90 nearby modern sabkhas of Ruways, Daffah and Khor al Jaramah (Fig. 1).
91 Palaeoenvironmental and climatic data extracted from these cores were used to reconstruct the
92 landscape at the time of occupation of the Ruways 1 and Suwayh 1 shell middens.

93

94 **2.1. Ruways 1 and Suways 1 shell middens**

95 Ruways 1 (RWY-1) is located at the center of a wide, open and flat coastline of the
96 Arabian Sea. Oriented NE–SW, this coastline associates wadi estuaries with vast sabkhas,
97 where barchanoid dunes were deposited after the disappearance of the mid-Holocene coastal
98 lagoon and mangrove systems. The current system shows peri-desert arid conditions since the
99 increasing aridification of the Late Holocene. The archaeological site extends over 8 hectares
100 and was discovered in 1996 by Mauro Cremaschi (University of Milan) and later prospected

101 by a French expedition. Ruways 1 was used as a ballast quarry at the beginning of 2010,
102 leading to the destruction of half of the site but revealing continuous, semi-circular
103 stratigraphic sections over nearly 80 m in length and 3 m in height in the eastern part of the
104 shell midden.

105 Ten sedimentary units were identified in the sequence. At the base, cemented gravels
106 from fluvial-deltaic origin provide the substrate upon which lies the shell midden. Sediment
107 deposition follows vertical and lateral aggradation phases. Erosion caused by aeolian
108 processes was identified throughout the section in the form of truncations of some of the older
109 anthropogenic deposits (Fig. 2). According to two radiocarbon dates obtained on charcoal and
110 marine shells, the settlement started during the late centuries of the 7th millennium BC
111 (around 8300 years BP; Berger et al., 2020). At the surface, archaeological remains indicate
112 an occupation until the 4th millennium BC (around 6000-5000 years, Berger et al., 2020).
113 Besides providing information on the subsistence strategy and technological behaviour of the
114 population, this site contains burials from the Middle Neolithic period that document funeral
115 rituals and periods of stress recorded in human skeletons. Both lithic and shell technology
116 have been used to manufacture stone tools, beads, net sinkers and hooks (Berger et al., 2020).
117 The recovered faunal remains belonged to vertebrates and molluscs, indicating the use of
118 terrestrial, marine and lagoon/mangrove food resources. Among the species identified,
119 mammalian remains can be attributed to sheep and goat, while reptile and turtle remains have
120 also been recovered (Berger et al., 2020). Among the marine shells, the clam *Marcia recens* is
121 the most abundant and indicates that Neolithic populations exploited species from the silty-
122 sand of the lagoon. Most of the fish remains were attributed to species living in brackish
123 waters, such as mullet (*Mugilidae*), sardine (*Clupeiformes*) and sea bream (*Sparidae*). These
124 species, along with catfish (*Ariidae*), indicate the occurrence of a paleo-lagoon and a
125 mangrove in the vicinity of Ruways 1 which were preferentially exploited by the Neolithic
126 populations (Berger et al., 2020). Botanical macro-remains also allow reconstructing the local
127 environment. *Avicennia marina* is the dominant species identified among the charred wood in
128 the upper part of the sequence with, to a lesser extent, *Rhizophora cf mucronata* attesting the
129 existence of a mangrove, today absent. Semi-desert taxa such as *Chenopodiaceae* were also
130 documented (Berger et al., 2020).

131 Suwayh 1 (SWY-1) is a Neolithic shell midden covering ca. 25,000 m² of surface area.
132 It is located along an old wadi estuary, between modern sabkhas and the Arabian Sea (Fig. 1).
133 Similar to Ruways 1, the site was partially destroyed when it was used as a ballast quarry 20
134 years ago. The excavations conducted during the 2000's allowed documenting a stratigraphic

135 sequence of 2 m thick (Fig. 3), with human occupation levels from the 6th and 5th millennia
136 (~8000-6000 years BP) and a necropolis in the recent portion of the site (Charpentier, et al.
137 2003). Seven test pits were opened for documenting the phases of the occupation. Unlike the
138 tombs, these levels are rich in archaeological material, mainly comprised of fishing
139 instruments such as stone net sinkers, shellfish hooks and jewellery. Dwellings were
140 uncovered in different phases of the site (Charpentier et al., forthcoming). SWY-1 is dated to
141 the Middle Neolithic period (Charpentier, 2008) and is a settlement specialised in shark
142 fishing (Marrast et al., 2020). The palaeoecological data available for this site indicate that,
143 starting from ~6500 years ago (~4500 years BC), species from mangroves (e.g., *Terebralia*
144 *palustris*) are represented and preceded species collected from the lagoon environment
145 (*Amiantis umbonella*) (Martin 2005, Berger et al., 2005).

146 The sedimentation rate of these two shell middens is highly variable and depends on
147 the rate of aeolian sedimentation (and truncation events) and anthropogenic deposition. It is
148 primarily the human accumulation of shells and rocks/pebbles that allowed the construction of
149 these coastal sedimentary anomalies. They favour the trapping of sand particles during human
150 occupation and protect it afterwards. The lithostratigraphic and sedimentological study has
151 revealed sets of sedimentary units that are clearly aeolian (facies with more or less pure
152 allogenic sand) or clearly anthropogenic (facies rich in ash, charcoal, accumulations of shells
153 or fish bones linked to past human occupations) and intermediate facies, revealing the more or
154 less strong reworking of anthropogenic deposits by aeolian processes (Berger et al. 2020).
155 Sedimentological, magnetic and geochemical analyses are ongoing.

156

157 **2.3. Cores**

158

159 In 2017 and 2018, a systematic off-site geoarchaeological and geophysical survey was
160 performed across the modern sabkhas, associated with electrical resistivity tomography (ERT)
161 measurements and thermic mechanical coring to depths up to 3-5 m (with a Cobra corer).
162 Three cores were drilled in the following localities: Ruways, Daffah and Khor al Jaramah
163 (Fig. 1). Sedimentological data extracted from the cores were used to reconstruct the middle
164 Holocene landscape: they indicate that the lagoons and mangroves were located under the
165 modern sabkhas.

166 Four main sedimentary units were distinguished in the upper 10 m: 1) at the base, a
167 deep and probably late Quaternary alluvial formation (not reached by cores); 2) coarse sandy
168 to gravelly alluvial pre-mid-Holocene deposits organised in a succession of paleochannels; 3)

169 fine fluvial-lagoon silty-sand formations associated with the maximum marine transgression,
170 4) red silts and gypsum interbedded with aeolian sands corresponding to the current sabkha
171 (Berger et al. 2020). OSL dating was carried out on this latter sedimentary unit (upper part of
172 the core), allowing to date the latest episode of lagoon/mangrove development.

173

174 **3. Materials and Methods**

175 **3.1. Sampling and preparation**

176 Sediment samples ($n = 21$) were collected at different localities (Ruways, RWY,
177 Suwayh, SWR, Daffah, DAF and Khor al Jaramah, KAJ; Tab. 1) during the 2017 expedition
178 (SOM Figs. S1 to S13). For the archaeological sites, sediments were mostly sampled using
179 lightproof bags during the night after cleaning the section to avoid any contamination by
180 newly bleached grains. Regarding the cores, sediments were sampled using a lightproof tube
181 that was inserted either in the section of the test pit dug near the core (Ruways and Daffah,
182 SOM Figs. S10 and S11) or directly in the core (Khor al Jaramah; SOM Figs. S12 and S13).
183 For Ruways 1, sediments were sampled from the base to the top of the sequence (phases I, II,
184 VII, VIII and IX, SOM Figs. S1 to S8); for Suwayh 1, only the upper levels were sampled
185 from a test pit (SOM Figs. S9); for the cores, sampling was focused on the lagoon/mangrove
186 levels.

187 Sample preparation and OSL measurements were conducted at the IRAMAT-CRP2A
188 (Institut de Recherche sur les Archéomatériaux-Centre de Recherche en Physique Appliquée à
189 l'Archéologie) at Bordeaux Montaigne University. Wet sieving was performed to extract the
190 most abundant granulometric fraction, 100-120, 120-160 or 160-200 μm depending on the
191 sample (Tab. 1). Chemical treatment included HCl (10%) to dissolve carbonates and H_2O_2
192 (30%) to break down organic matter. Feldspars were dissolved using H_2SiF_6 (31%) over one
193 week. Finally, quartz grains were rinsed with HCl (10%) and sieved again (using 100, 120 or
194 160 μm sieves for the 100-120, 120-160 or 160-200 μm granulometric fractions, respectively)
195 to eliminate partially dissolved feldspar grains.

196

197 **3.2. Equivalent dose determination**

198 Equivalent doses (D_e) were determined by applying the single aliquot regeneration
199 (SAR) protocol (Murray and Wintle, 2000) to multi-grain aliquots (1 mm \varnothing) analysed with a
200 lexsys SMART TL/OSL reader (Richter et al., 2015) for the quartz samples, and with a
201 lexsys research TL/OSL reader (Richter et al., 2013) for the feldspar samples. D_e values were

202 obtained on quartz using the protocol described in Tab. 2. For BDX 19229, whose signal
203 reached saturation, and for BDX 19235, pIRIR₂₂₅ measurements were performed (Tab. 3).

204 Regarding the measurement procedure, green LEDs were used for stimulation
205 (stimulation power density of 50 mW·cm⁻²) and the OSL signal was detected with a
206 combination of UV filters (Schott BG-3, 3 mm in conjunction with a Delta BP365/50 EX).
207 Aliquots were irradiated using the inbuilt ⁹⁰Sr/⁹⁰Y beta-sources fitted to the readers. An
208 infrared stimulation at 50°C for 60 s was performed before each OSL measurement to avoid
209 potential contamination from remaining feldspars. We observed a dominant fast decaying
210 OSL signal component for our quartz samples (Fig. 4).

211 For BDX19229 and BDX19235, the pIRIR₂₂₅ protocol (Thomsen et al., 2008; Buylaert
212 et al., 2011) was applied (Tab. 3). Infrared laser diodes were used for stimulation (power
213 density: 120 mW·cm⁻²) and the signal was detected with a combination of optical filters
214 (Schott KG3, 3 mm + Chroma D410/30 x for the research reader).

215 Preheat plateau tests were conducted on natural aliquots from samples BDX 19231
216 and 19232 (Ruways 1) and BDX 19238, 19239 and 19240 (Suwayh 1). For Ruways 1
217 samples, measurements were done between 180°C and 300°C (cutheat = 200°C) with 20°C
218 steps; for Suwayh 1 samples, from 170°C to 250°C (cutheat = 160°C) with 20°C steps. Since
219 low D_e values were observed after preliminary tests, the preheat temperature tests were
220 conducted on lower temperatures for Suwayh 1 samples.

221 Dose recovery tests (DRT) were performed on each sample, on three to five bleached
222 aliquots (*n* = 74; Tab. 4), using the preheat and cutheat temperatures mentioned above. The
223 aliquots were bleached in a solar simulator (Hölne SOL500) for 1 min. The irradiation dose
224 applied was similar to the one expected on natural aliquot.

225 All data were processed using the software Analyst v.4.52 (Duller, 2007). The signal
226 was integrated using the first 1.5 s for quartz and the first 5 s for feldspars, while the
227 background was subtracted from the last 15 s for quartz and the last 30 s for feldspars. The
228 following criteria were applied for D_e values selection: a recycling ratio limit of 10%; a
229 maximum recuperation of 5% of the natural signal; a test dose signal more than 3 sigma
230 above background. Equivalent doses were obtained by fitting an exponential function through
231 the normalised luminescence signals. The statistic treatment of the D_e values was done using
232 either the Central Age Model (CAM, Galbraith et al., 1999) or the Finite Mixture Model
233 (FMM, Galbraith and Green, 1990) for samples exhibiting a large scatter of the D_e values.

234

235

236 **3.3. Dose rate determination**

237 **3.3.1. Alpha and beta dose rates**

238 A portion of the sediment sampled from the section or the core (see Tab. 1) was dried in an
239 oven for one week at 40 °C and the water content (wet weight %) measured. ²³⁸U, ²³²Th and K
240 contents were determined using a laboratory gamma-ray spectrometer equipped with a high-
241 resolution, broad energy Ge (BEGe) detector. Since no HF treatment was applied, both alpha
242 and beta dose rates were derived from these values taking into account the conversion factors
243 of Guérin et al. (2011).

244 **3.3.2. Gamma dose rate**

245 For all samples except those collected directly from the KAJ core (BDX 19245 and 19246;
246 SOM Figs. S12 and S13) and for samples BDX 19232 and 19237 from Ruways 1, whose
247 location did not allow to safely dig a 30 cm deep hole in the section (SOM Figs. S4 and S8),
248 the gamma dose rate was measured *in situ* using a portable gamma-ray multichannel analyser
249 connected to a NaI(Tl) detector. The data were processed following the “threshold” technique
250 (Mercier and Falguères, 2007). For the other samples, the gamma dose rate was derived from
251 their ²³⁸U, ²³²Th and K contents deduced from BEGe gamma spectrometric analyses.

252 **3.3.3. Cosmic dose rate**

253 The depth of the sampling position was measured from the surface level (Tab. 1). This value
254 was used to calculate the cosmic dose rate according to Prescott and Hutton (1988).

255

256 **4. Results**

257 OSL ages on quartz and pIRIR₂₂₅ ages on feldspar aliquots were calculated with the 1
258 σ error range, taking into account the alpha attenuation and beta absorption factors of Brennan
259 et al. (1991) and Guérin et al. (2012) respectively. Water content values used for age
260 calculation range from 3% to 27% depending on the context (measured values, Tab. 5).

261

262 **4.1. Equivalent doses**

263 **4.1.1. Preheat plateau test**

264 Results obtained do not show temperature dependence between 180°C and 280°C for
265 Ruways 1 samples (SOM Fig. S14), and between 170°C and 250°C for Suwayh 1 samples
266 (SOM Fig. S15). Measurements were thus conducted using a 260°C preheat temperature and
267 200°C cutheat temperatures for Ruways 1 samples, and using lower temperatures for Suwayh
268 1 (210°C preheat; 160°C cutheat) and cores samples (190°C preheat; 160°C cutheat).

269

270 **4.1.2. Dose recovery tests**

271 For Ruways 1 and Suwayh 1 samples, dose recovery ratios range from 0.96 ± 0.03 to
272 1.04 ± 0.05 (Tab. 4). For the core samples, dose recovery ratios are close to 1 except for BDX
273 19243 (Daffah C2) that yields a higher recovery ratio of 1.44 ± 0.16 . This high value can be
274 related to the low irradiation dose applied (0.5 Gy), which is similar to the natural dose, and to
275 a relatively high residual signal after bleaching.

276

277 **4.2. Interpreting D_e values distribution**

278 The distribution of D_e values is heterogeneous for most of the samples (Tab. 6), as
279 shown in Fig. 5 (Ruways 1 phase II). Indeed, when applying the Central Age Model (CAM,
280 Galbraith et al., 1999), overdispersion (OD) values are higher than 20% for half of the
281 samples, and 5 out of 21 D_e values have OD >50% (Tab. S1). This indicates that several grain
282 populations are present, which may result from differential bleaching and/or microdosimetry
283 effects. The superposition of these populations can be described using the finite mixture
284 model (FMM, Galbraith and Green, 1990). However, this model must be applied with caution
285 on data obtained from the analysis of multi-grain aliquots because averaging effects may
286 occur, even for small aliquots (Arnold and Roberts, 2009; Galbraith and Roberts, 2012).
287 However, considering that only a small percentage of grains give rise to the OSL signal
288 (typically ~5% produce 95% of the total light sum, Duller et al. (2000)), the FMM model is
289 applicable when only a few grains have been measured, which is the case here with our 1 mm
290 \varnothing aliquots (e.g., Preusser et al., 2011). FMM modelling was thus conducted for samples with a
291 number of measured aliquots ($n \geq 22$ and OD >15% (a sigma-b value of 0.10 was assumed) ;
292 the components (central value and proportion) are presented in Tab. S2. For samples with n
293 aliquots < 22, only the CAM was applied, including the feldspar samples from Ruways 1
294 (BDX 19229 and 19235) and some of the quartz samples from the cores (BDX 19242, 19244,
295 19245 and 19246) (Tab. 6). For pIRIR₂₂₅ ages, no residual dose nor fading corrections were
296 applied. However, considering that the IRSL age for BDX 19235 is in agreement with OSL
297 ages obtained for Phase II (BDX 19226, 19227, 19228 and 19236, Tab. 6), the effect of these
298 two variables might have been compensated for this sample.

299 D_e values obtained either using the FMM or CAM model are presented in Tab. 6 and
300 Tab. S1. FMM D_e values are systematically lower than CAM D_e values, but considering the
301 associated uncertainties, most of the values are indistinguishable (Fig. 6). Nevertheless, the
302 use of the main FMM component (Tab. S2) help to discard incomplete bleached grains which,

303 when taken into account (as it is the case with the CAM model), induce an overestimation of
304 the mean D_e values.

305 For Ruways 1, CAM D_e values range from 0.29 ± 0.05 to 8.55 ± 0.53 Gy for the
306 quartz samples and from 14.78 ± 1.55 to 520.66 ± 79.78 Gy for the feldspar samples, with OD
307 values between 9 ± 2 and 98 ± 13 %. FMM D_e values range from 0.13 ± 0.01 to 8.12 ± 0.17
308 Gy for the quartz samples.

309 For Suwayh 1, D_e values range from 5.91 ± 0.13 to 7.43 ± 0.25 using the CAM (OD
310 values between 13 ± 2 and 20 ± 2 %). The D_e distribution is thus more homogeneous for these
311 samples in comparison with most of the samples collected in a similar context at Ruways 1.
312 Indeed, some aliquots likely include partially bleached grains, especially BDX 19238 and
313 19240 (Fig. 7) and the use of the FMM allowed to extract the main D_e component whose
314 proportion represents $\geq 89\%$ (Tab. S2). The resulting D_e values range from 5.76 ± 0.15 to
315 7.08 ± 0.15 Gy.

316 For the cores, CAM D_e values vary from 1.68 ± 0.51 to 5.39 ± 0.44 Gy (5.01 ± 0.26
317 Gy for the FMM) for Ruways-C1, from 0.57 ± 0.06 to 0.70 ± 0.08 Gy (0.66 ± 0.06 Gy for the
318 FMM) for Daffah-C2 and equal 7.92 ± 0.43 and 8.46 ± 0.38 Gy for the Khor al Jaramah-C1
319 and C2 samples. Consequently, these values indicate either a large range of dose rates or ages
320 significantly different for similar contexts.

321

322 4.3. Dose rates

323 The main contribution to the dose rate comes from the beta dose rate, except for the
324 feldspar samples (Fig. 8). Alpha dose rates were taken into account for all samples since the
325 grains have not been etched. They are low and represent maximum 3% of the total dose rate.
326 Both were derived from ^{238}U , ^{232}Th and K contents of the sediment samples (Tab. 5). For
327 Ruways 1, the ^{238}U content ranges from 0.53 ± 0.01 to 3.20 ± 0.04 ppm and for 7 out of 12
328 samples, the ^{232}Th content is lower, ranging from 0.57 ± 0.02 to 4.62 ± 0.04 ppm. These low
329 ^{232}Th values for the Ruways 1 sediments are not well understood and further investigations
330 are required to explain the mechanisms. For the Suwayh 1, Ruways-C1, Daffah-C2 and Khor
331 al Jaramah-C1/C2 samples, the ^{232}Th content is systematically higher than ^{238}U content. K
332 contents vary also considerably, ranging from 0.16 ± 0.01 (RWY, BDX 19226) to 1.26 ± 0.02
333 (KAJ-C2, BDX 19246). The highest K contents were measured in the core samples where the
334 sediments were essentially composed of clayey silt.

335 The gamma dose rate was measured *in situ* for most of the samples (Tab. 6). In-situ
336 measurements values range from 210 ± 10 to 405 ± 20 $\mu\text{Gy}\cdot\text{a}^{-1}$ for Ruways 1, from 207 ± 10

337 to $345 \pm 17 \mu\text{Gy}\cdot\text{a}^{-1}$ for Suwayh 1, from 331 ± 17 to $372 \pm 19 \mu\text{Gy}\cdot\text{a}^{-1}$ for Daffah-C1/C2 and
338 overlap for Ruways-C1 (475 ± 24 to $478 \pm 24 \mu\text{Gy}\cdot\text{a}^{-1}$). The gamma dose rates calculated for
339 BDX 19232 and 19237 using the U, Th and K contents (444 ± 3 and $276 \pm 2 \mu\text{Gy}\cdot\text{a}^{-1}$), fall
340 within the same range than those measured *in situ*. For Khor al Jaramah samples, the gamma
341 doses derived from radioelements contents (481 ± 6 and $494 \pm 4 \mu\text{Gy}\cdot\text{a}^{-1}$) are similar to those
342 measured using the portable gamma-probe at Ruways-C1 (around $475 \mu\text{Gy}\cdot\text{a}^{-1}$); this is in
343 agreement with the environmental dose rates that were not measured *in situ* for these samples
344 (SOM Figs. S11 and S12).

345 For the feldspar samples, a value of $10 \pm 1\%$ was taken into account for the internal K
346 content of the feldspar grains. The associated internal dose rate, of $500 \pm 50 \mu\text{Gy}\cdot\text{a}^{-1}$,
347 represents up to 42% of the dose rate (Fig. 8), thus being an important parameter for the age
348 calculation.

349 The cosmic dose was calculated considering the present thickness of sediments, then
350 leading to minimum values. They represent between 14% and 27% of the total dose rate (Fig.
351 8). However, for the top of the sequence at Ruways 1 (< 60 cm from the top of the section),
352 human occupation is less frequent and erosion/reworking processes by deflation (sandstorm)
353 may have occurred, impacting the cosmic dose rate that can therefore not be estimated with
354 precision.

355

356 **5. Age interpretation and discussion**

357

358 FMM ages were used for age interpretation for samples with OD values $\geq 15\%$ and n
359 aliquots measured ≥ 22 aliquots. Indeed, for these samples, CAM D_e values appear
360 overestimated likely due to the presence of partially bleached grains (Tab. S3). CAM ages
361 were used for interpretation for samples with OD values $< 15\%$ and/or n aliquots measured $<$
362 22 aliquots.

363 Ages obtained for the archaeological sequence of the Ruways 1 shell midden range
364 from 440 ± 76 ka to 100 ± 10 years, indicating a deposition time from the Middle Pleistocene
365 to the late Holocene (Tab. 6 and Fig. 9). Phase I, corresponding to a gravelly substrate,
366 deposited by a river, is dated to 440 ± 76 ka using pIRIR₂₂₅. Phase II, corresponding to the
367 first Neolithic occupation, is dated between 8340 ± 480 and 7740 ± 580 years (6800-5200
368 years BC). A weighted mean age of 7930 ± 240 years is obtained for this unit. Phases VII and
369 VIII are characterised by aeolian sand sedimentary structures where deflation forms
370 predominate. Phase VII is dated to 5930 ± 370 years (4280-3540 years BC) and phase VIII,

371 from 5830 ± 320 to 5390 ± 290 years (3100-4100 years BC). For this phase, OD values are
372 low (9-13%) indicating that quartz grains were well bleached before deposition and leading to
373 indistinguishable CAM and FMM ages. Finally, ages obtained for phase IX, composed of
374 sediment that fills a deflation corridor interlocking phase VI, range from 3410 ± 190 to $100 \pm$
375 10 years. They indicate that the sediment at the top of the sequence was disturbed, which led
376 to a recent resetting of the OSL signal. The age of 3410 ± 190 years should thus be considered
377 as a minimum age, the two other samples from this unit being dated to around 250 and 100
378 years. Consequently, human occupation at Ruways 1 spans the mid-Holocene: OSL ages
379 indicate a deposition time from the 6th millennium BC (phase II) to the 4th millennium BC
380 (phase VIII), with the main occupation occurring during the 4th millennium BC (phases VII
381 and VIII).

382 At Suwayh 1, age results range from 6180 ± 310 to 5480 ± 270 years (4500-3200
383 years BC) and are contemporaneous with phases VII and VIII at Ruways 1. However, OSL
384 ages tend to be slightly younger than radiocarbon ages obtained on shells, by 500-700 years
385 (calibrated with a $\Delta=0$, Marrast et al., 2020; Charpentier et al., forthcoming). Two hypotheses
386 could explain this phenomenon: ^{14}C ages might be overestimated due to the reservoir effect
387 that is not well constrained in the region, and/or OSL ages are underestimated because of the
388 post-burial increase in U in the sediments caused by the proximity to the mouth of the wadi.

389 With regard to the cores, ages range from 3110 ± 370 to 970 ± 290 years for Ruways-
390 C1, from 540 ± 70 to 420 ± 50 years for Daffah-C2 and are 5220 ± 570 and 5320 ± 590 years
391 for Khor al Jaramah-C1 and C2, respectively. These age results indicate that the top of the
392 Ruways-C1 and Daffah-C2 cores are recent deposits that cannot be correlated with the
393 occupations at the shell middens. Indeed, these deposits belong to the post-Neolithic hyper-
394 arid phase. However, the ages for BDX 19245 and 19246 (Khor al Jaramah-C1 and C2)
395 indicate that a phase of lagoon/mangrove development occurred ~5900-4600 years ago (3900-
396 2600 years BC) and are contemporaneous with human occupation at Ruways 1 (phases
397 VII/VIII) and Suwayh 1 (Fig. 9). Considering the depth of the sampling (90-100 cm) and that
398 earlier paleolagoon episodes were identified in a 480 cm-deep core drilled in Ruways (core 4)
399 and dated using ^{14}C on shells from ~6700 years BP (5000-5400 years BC) (Berger et al.,
400 2020), the dates obtained at Khor al Jaramah could be related to the end of the
401 lagoon/mangrove development.

402 Previous palynological and micropalaeontological analyses and ^{14}C ages on shells
403 obtained from cores drilled under the modern sabkhas at Suwayh indicate that an optimum
404 period for mangrove development occurred ~6000 years ago (Lézine et al., 2002). This

405 episode was followed by two phases of seawater incursion ca. 5100 and 4500 years ago that
406 led to the formation of a lagoon (Lézine et al., 2002). Based on geological and environmental
407 data from the Suwayh and Alashkara sabkhas, Berger et al. (2005, 2013) highlighted at least
408 five sea-level transgressions that occurred between ~6800-6400 and ~4400-4200 years ago.
409 The resolution of the OSL ages obtained in this study does not allow to correlate the lagoon
410 episodes recorded in the studied cores with a specific phase. However, both
411 palaeoenvironmental and palaeoclimatic data extracted from the Khor al Jaramah cores can be
412 correlated with human occupation phases at Ruways 1 and Suwayh 1, based on our
413 chronological data.

414

415 **6. Conclusions**

416 Luminescence dating (OSL and pIRIR₂₂₅) was performed on sediment sampled from
417 both archaeological and geological contexts, in order to correlate human occupation at
418 Ruways 1 and Suwayh 1 shell middens with palaeoclimatic and palaeoenvironmental data
419 preserved in sediments under the modern sabkhas at Ruways, Daffah and Khor al Jaramah.
420 This work aimed at establishing the timeline of Neolithic occupation phases at the two shell
421 middens, as well as providing a chronology for lagoon/mangrove phases that are linked to sea
422 level variations and important climate change during the mid-Holocene.

423 For most of the samples, the large scatter of the D_e values suggested the presence of
424 several grain populations characterised by different levels of bleaching at the time of
425 deposition, and/or submitted to different dose rates (microdosimetry heterogeneities). For
426 such samples, the FMM model was applied and the D_e values were compared with those
427 calculated using the CAM model. The ages are in agreement with the general stratigraphic
428 order, allowing the establishment of an OSL chronology for the Holocene levels, especially
429 for levels with few or no organic remains that can be dated using ^{14}C .

430 Age results obtained in this study allowed dating human occupation at Ruways 1 from
431 8340 ± 480 (6320 ± 480 years BC) for the phase II to 5390 ± 290 years (3370 ± 290 years
432 BC) for the phase VIII, and confirm the antiquity of the settlement, as suggested by ^{14}C ages
433 (6th-7th millennium BC, Berger et al., 2020). However, OSL ages obtained for the top of the
434 sequence are likely underestimated, since two out of three ages obtained are younger than 300
435 years, indicating a recent resetting of the OSL signal corresponding to reworked sediments.
436 Ages obtained for the intermediate phases VII and VIII suggest an occupation time spanning
437 from ca. 6000 to 5000 years BP (~4000-3000 years BC). Chronological data obtained on the
438 sediment from the Khor al Jaramah cores indicate that the formation of the lagoon/mangrove

439 occurred around ~5900-4700 years ago (3900-2600 years BC), coinciding with the Neolithic
440 occupation at the shell midden.

441

442

443 **Acknowledgments**

444 This study is part of the NeoArabia project (Dir. J.F. Berger), funded by the French
445 National Research Agency (ANR-16-CE03-0007), dealing with the very long span of the
446 Arabian Neolithic (6200-2800 BC). We thank all the team members of NeoArabia for their
447 help in the field and for discussing the data.

448 We thank the Ministry of Heritage and Culture of the Sultanate of Oman for granting
449 permission for fieldwork along the Omani Arabian sea shores, with specific reference to H.E.
450 Salim Mohammed Almahruqi, Undersecretary for Heritage; Mr Sultan Said Al-Bakri,
451 Director General for Archaeology; Mr Khamis Al-Asmi, Director of the Department of
452 Excavations and Archaeological Studies.

453 V. Charpentier would like to thank the Consultative Commission for Excavations
454 Abroad of the Ministry of European and Foreign Affairs, for granting the French
455 Archaeological Mission.

456 M. Richard and N. Mercier are grateful to LaScArBx (Bordeaux Archaeological
457 Sciences Labex) administrated by ANR (Agence Nationale de la Recherche) with the
458 reference ANR-10-LABX-52.

459 We thank the two anonymous reviewers for their constructive remarks on this
460 manuscript.

461

462

463 **References**

464

465 Arnold, L.J., Demuro, M., Ruiz, M.N., 2012. Empirical insights into multi-grain averaging
466 effects from 'pseudo' single-grain OSL measurements, *Radiation Measurements* 47,
467 652-658.

468 Bassinot, F.C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard, F., Bopp,
469 L., 2011. Holocene evolution of summer winds and marine productivity in the tropical
470 Indian Ocean in response to insolation forcing: data-model comparison, *Climate of the*
471 *Past* 7, 815-829.

- 472 Berger, J.-F., Davtian, G., Cleuziou, S., 2005. Évolution paléogéographique du Ja'alan
473 (Oman) à l'Holocène moyen : impact sur l'évolution des paléomilieus littoraux et les
474 stratégies d'adaptation des communautés humaines. *Paléorient* 31, 46-63.
- 475 Berger, J.F., Charpentier, V., Crassard, R., Martin, C., Davtian, G., López-Sáez, J.A., 2013.
476 The dynamics of mangrove ecosystems, changes in sea level and the strategies of
477 Neolithic settlements along the coast of Oman (6000–3000 cal. BC). *Journal of*
478 *Archaeological Science* 40, 3087-3104.
- 479 Berger, J.F., Guilbert-Berger, R., Marast, A., Munoz, O., Guy, H., Barra, A., López-Sáez,
480 J.A., Pérez-Díaz, S., Mashkour, M., Debue, K., Lefèvre, C., Gosselin, M., Brugnaux,
481 G., Mougne, C., Thorin, S., Nisbet, R., Oberlin, C., Mercier, N., Richard, M., Depreux,
482 B., Bearez, P., 2020. First contribution of the excavation and the chronostratigraphic
483 study of Ruways 1 Neolithic shell midden (Oman), in terms of Neolithization,
484 palaeoeconomy, social-environmental interactions and site formation processes,
485 *Arabian Archaeology and Epigraphy*, 32-49.
- 486 Brennan, B.J., Lyons, R.G., Phillips, S.W., 1991. Attenuation of alpha particle track dose for
487 spherical grains. *International Journal of Radiation Applications and Instrumentation.*
488 *Part D. Nuclear Tracks and Radiation Measurements* 18, 249-253.
- 489 Buylaert, J.-P., Thiel, C., Murray, A., Vandenberghe, D., Yi, S., Lu, H., 2011. IRSL and post-
490 IR IRSL residual doses recorded in modern dust samples from the Chinese Loess
491 Plateau. 38, 432.
- 492 Charpentier, V., Marquis, P., Pellé, É., 2003. La nécropole et les derniers horizons Ve
493 millénaire du site de Gorbat al-Mahar (Suwayh, SWY-1, Sultanat d'Oman) : premiers
494 résultats. *Proceedings of the Seminar for Arabian Studies* 33, 11-19.
- 495 Charpentier, V., 2008. Hunter-gathers of the “empty quarter of the early Holocene” to the last
496 Neolithic societies: chronology of the late prehistory off South-Eastern Arabia (8000-
497 3100 BC). *Proceedings of the Seminar for Arabian Studies* 38, 59-82.
- 498 Charpentier, V. Lacaze, M. Marquis, P. Marrast, A. Mashkour, M. Munoz, O. , E. Pellé
499 (Forthcoming). First Neolithic communities in the Arabian Sea. The excavation of the
500 6th-5th millennium BCE Suwayh 1 settlement, (Sultanate of Oman). *Arabian*
501 *Archaeology and Epigraphy*.
- 502 Crassard, R., Drechsler, P., 2013. Towards new paradigms: multiple pathways for the Arabian
503 Neolithic, *Arabian Archaeology and Epigraphy* 24, 3-8.
- 504 Duller, G.A., 2007. Assessing the error on equivalent dose estimates derived from single
505 aliquot regenerative dose measurements. *Ancient TL* 25, 15-24.

506 Duller, G.A., Bøtter-Jensen, L., Murray, A.S., 2000. Optical dating of single sand-sized grains
507 of quartz: sources of variability, *Radiation Measurements* 32, 453-457.

508 Fleitmann, D., Burns, S.J., Mangini, A., Mudelsee, M., Kramers, J., Villa, I., Neff, U., Al-
509 Subbary, A.A., Buettner, A., Hippler, D., Matter, A., 2007. Holocene ITCZ and Indian
510 monsoon dynamics recorded in stalagmites from Oman and Yemen (Socotra),
511 *Quaternary Science Reviews* 26, 170-188.

512 Galbraith, R.F., Green, P.F., 1990. Estimating the component ages in a finite mixture.
513 *International Journal of Radiation Applications and Instrumentation. Part D. Nuclear*
514 *Tracks and Radiation Measurements* 17, 197-206.

515 Galbraith, R.F., Roberts, R.G., Laslett, G.M., Yoshida, H., Olley, J.M., 1999. Optical dating
516 of single and multiple grains of quartz from Jinnium rock shelter, northern Australia:
517 Part I, experimental design and statistical models. *Archaeometry* 41, 339-364.

518 Galbraith, R.F., Roberts, R.G., 2012. Statistical aspects of equivalent dose and error
519 calculation and display in OSL dating: An overview and some recommendations,
520 *Quaternary Geochronology* 11, 1-27.

521 Guérin, G., Mercier, N., Adamiec, G., 2011. Dose-rate conversion factors: update. *Ancient TL*
522 29, 5-8.

523 Guérin, G., Mercier, N., Nathan, R., Adamiec, G., Lefrais, Y., 2012. On the use of the infinite
524 matrix assumption and associated concepts: A critical review. *Radiation Measurements*
525 47, 778-785.

526 Jung, S.J.A., Davies, G.R., Ganssen, G.M., Kroon, D., 2004. Synchronous Holocene sea
527 surface temperature and rainfall variations in the Asian monsoon system, *Quaternary*
528 *Science Reviews* 23, 2207-2218.

529 Lézine, A.-M., Saliège, J.-F., Mathieu, R., Tagliatela, T.-L., Mery, S., Charpentier, V.,
530 Cleuziou, S., 2002. Mangroves of Oman during the late Holocene; climatic implications
531 and impact on human settlements. *Vegetation History and Archaeobotany* 11, 221-232.

532 Lézine, A.-M., Robert, C., Cleuziou, S., Inizan, M.-L., Braemer, F., Saliège, J.-F., Sylvestre,
533 F., Tiercelin, J.-J., Crassard, R., Méry, S., Charpentier, V., Steimer-Herbet, T., 2010.
534 Climate change and human occupation in the Southern Arabian lowlands during the last
535 deglaciation and the Holocene, *Global and Planetary Change* 72, 412-428.

536 Marrast, A., Béarez, P., Charpentier, V., 2020. Sharks in the lagoon? Fishing exploitation at
537 the Neolithic site of Suwayh 1 (Ash Sharqiyah region, Arabian Sea, Sultanate of Oman),
538 *Arabian Archaeology and Epigraphy* 31, 178-193.

539 Martin, C., 2005. Stratégies et statut de la collecte des mollusques marins sur les sites côtiers
540 d'Oman du Néolithique à l'âge du Bronze: Apport des sites de suwayh 1, ra's al-
541 Khabbah 1 et Ra's al-Jinz 2, Paléorient, 169-175.

542 Mercier, N., Falguères, C., 2007. Field gamma dose-rate measurement with a NaI(Tl)
543 detector: re-evaluation of the "threshold" technique. *Ancient TL* 25, 1-4.

544 Murray, A.S., Wintle, A.G., 2000. Luminescence dating of quartz using an improved single-
545 aliquot regenerative-dose protocol. *Radiation Measurements* 32, 57-73.

546 Parker, A.G., Rose, J.I., 2008. Climate change and human origins in southern Arabia,
547 *Proceedings of the Seminar for Arabian Studies* 38, 25-42.

548 Prescott, J.R., Hutton, J.T., 1988. Cosmic ray and gamma ray dosimetry for TL and ESR.
549 *International Journal of Radiation Applications and Instrumentation. Part D. Nuclear*
550 *Tracks and Radiation Measurements* 14, 223-227.

551 Preusser, F., Schmitt, L., Delile, H., Grosprêtre, L., 2011. Optically Stimulated Luminescence
552 (OSL) dating of the sedimentation history of the Yzeron Basin (Chaudanne sub-
553 catchment), Rhône Valley, France, *Quaternaire* 22, 73-83.

554 Richter, D., Richter, A., Dornich, K., 2013. Lexsyg — A new system for luminescence
555 research. *Geochronometria* 40, 220-228.

556 Richter, D., Richter, A., Dornich, K., 2015. Lexsyg smart - a luminescence detection system
557 for dosimetry, material research and dating application. *Geochronometria* 42, 202-209.

558 Thomsen, K.J., Murray, A.S., Jain, M., Bøtter-Jensen, L., 2008. Laboratory fading rates of
559 various luminescence signals from feldspar-rich sediment extracts. *Radiation*
560 *Measurements* 43, 1474-1486.

561

562

563

564 **Figures**

565

566 Fig. 1: Location of shell middens (Ruways 1 and Suwayh 1, red squares) and cores (Ruways,
567 Daffah and Khor al Jaramah, red circles) in the eastern coast of the Sultanate of Oman.
568 Modified after Berger et al. (2005: Figure 2).

569

570 Fig. 2: Ruways 1. Stratigraphic profile (east-west) and depositional phases.

571

572 Fig. 3: Suwayh 1. Stratigraphic profile. Drawing by J-F. Berger and G. Davtian. After
573 Charpentier (2008: Figure 3).

574

575 Fig. 4: Shine-down OSL curve (signal versus stimulation time) of BDX 19226.

576

577 Fig. 5: Ruways 1. a. Abanico plots of equivalent doses obtained for BDX 19226 (a), 19227
578 (b) and 19228 (c) (Phase II). Overdispersion values are $23 \pm 3\%$ (a), $35 \pm 4\%$ (b) and $32 \pm 4\%$
579 (c) respectively.

580

581 Fig. 6: Graphic comparison of D_e values obtained using the FMM and CAM. Samples number
582 follows the stratigraphic order (depth and layer attribution from Tabs. 1 and 6). For D_e values
583 < 1 Gy, error bars are not visible due to the scale. For BDX 19234, the FMM and CAM D_e
584 values are indistinguishable.

585


586 Fig. 7: Suwayh 1: Histogram of equivalent doses obtained for BDX 19238 (a), 19239 (b) and
587 19240 (c). Overdispersion values are $20 \pm 2\%$ (a), $13 \pm 2\%$ (b) and $20 \pm 2\%$ (c).


588

589 Fig. 8: Dose rate distribution (%).


590




591 Fig. 9: Graphic representation of luminescence ages obtained as a function of depth. For
592 young ages (< 1 ka), error bars are not visible due to the scale. The samples that fall within
593 the grey shade can be correlated: the formation of the lagoon in Khor al Jaramah is
594 contemporaneous with human occupation at Ruways 1 (phases VII/VIII) and Suwayh 1.






BDX 19226









- Ruways 1 (FMM) ◆ Suwayh 1 (FMM) ■ Ruways-core 1 (FMM) ▲ Daffah-Core 2 (FMM)
- Ruways-1 (CAM) ◇ Suwayh 1 (CAM) □ Ruways-core 1 (CAM) △ Daffah-Core 2 (CAM)


a.







b.



c.







Lab #	Site	Sample #	Depth (cm)	Context	Sample type	Sedimentology	Material ^a	Granulometric fraction
19226	Ruways 1	RWY-1-OSL-1	240	Section	Tube	Fine sand	Q	160-200
19227	Ruways 1	RWY-1-OSL-2	195	Section	Tube	Fine sand	Q	160-200
19228	Ruways 1	RWY-1-OSL-3	150	Section	Tube	Fine sand	Q	160-200
19229	Ruways 1	RWY-1-OSL-4	195	Section	Bag	Sand	F	160-200
19230	Ruways 1	RWY-1-OSL-5	60	Section	Bag	Sandy lens	Q	160-200
19231	Ruways 1	RWY-1-OSL-6	40	Section	Bag	Sandy lens	Q	160-200
19232	Ruways 1	RWY-1-OSL-7	20	Section	Bag	Sandy lens	Q	160-200
19233	Ruways 1	RWY-1-OSL-8	50	Section	Bag	Sand	Q	160-200
19234	Ruways 1	RWY-1-OSL-9	25	Section	Bag	Sand	Q	160-200
19235	Ruways 1	RWY-1-OSL-10	250	Section	Bag	Sand	F	160-200
19236	Ruways 1	RWY-1-OSL-11	210	Section	Bag	Sand	Q	160-200
19237 ^b	Ruways 1	RWY-1-OSL-12	20	Section	Bag	Sand	Q	120-160
19238	Suwayh 1	SWY-OSL-1	120	Section	Bag	Silt	Q	120-160
19239	Suwayh 1	SWY-OSL-2	45	Section	Bag	Silt	Q	120-160
19240	Suwayh 1	SWY-OSL-3	15	Section	Bag	Silt	Q	120-160
19241	Ruways-C1	RWY-C1-OSL-1	50	Core	Tube	Silt	Q	100-120
19242	Ruways-C1	RWY-C1-OSL-2	20	Core	Tube	Coarse sand	Q	100-120
19243	Daffah-C2	DAF-C2-OSL-1	85	Core	Tube	Silt	Q	120-160
19244	Daffah-C2	DAF-C2-OSL-2	20	Core	Tube	Silt	Q	100-120
19245 ^b	Khor al Jaramah-C1	KAJ-C1-OSL-1	90-100	Core	Tube	Silt	Q	100-120
19246 ^b	Khor al Jaramah-C2	KAJ-C2-OSL-1	90-100	Core	Tube	Clayey-silt	Q	100-120

Tab. 1. List of samples analysed in this study. Sample numbers match those provided in SOM.

^a Material: quartz (Q) / feldspars (F)

^b no in-situ gamma dosimetry measurements were performed for these samples (see text).

Step	Treatment
1	Given dose
2	Preheat (190-260°C for 10 s)
3	IR stimulation for 60 s at 50°C
4 (L _x)	Green stimulation for 100 s at 125°C
5	Given test dose
6	Cutheat (160-200°C for 0 s)
7	IR stimulation for 60 s at 50°C
8 (T _x)	Green stimulation for 100 s at 125°C
9	Return to 1

Tab. 2. OSL SAR measurement protocol applied to quartz.

Step	Treatment
1	Given dose
2	Preheat (250°C for 60 s)
3	IR stimulation for 200 s at 50°C
4 (L _x)	IRSL stimulation for 200 s at 225°C
5	Given test dose
6	Preheat (250°C for 60 s)
7	IR stimulation for 200 s at 50°C
8 (T _x)	IRSL stimulation for 200 s at 225°C
9	IRSL stimulation for 60 s at 325°C
10	Return to 1

Tab. 3. pIRIR₂₂₅ SAR measurement protocol applied to feldspars.

Lab #	Site	<i>n</i> aliquots	DRT	Dose (Gy)
19226	Ruways 1	3	1.00 ± 0.03	5.4
19227	Ruways 1	3	1.00 ± 0.06	5.4
19228	Ruways 1	3	0.99 ± 0.08	5.4
19230	Ruways 1	3	1.04 ± 0.05	0.9
19231	Ruways 1	3	1.01 ± 0.01	0.9
19232	Ruways 1	3	0.96 ± 0.02	5.4
19233	Ruways 1	3	0.98 ± 0.06	5.4
19234	Ruways 1	3	1.00 ± 0.03	5.4
19236	Ruways 1	3	0.99 ± 0.02	5.4
19237	Ruways 1	3	0.96 ± 0.03	5.4
19238	Suwayh 1	5	0.99 ± 0.02	5.3
19239	Suwayh 1	5	0.99 ± 0.03	5.3
19240	Suwayh 1	4	1.02 ± 0.02	5.3
19241	Ruways-C1	5	0.97 ± 0.07	3.5
19242	Ruways-C1	5	1.08 ± 0.23	1.8
19243	Daffah-C2	5	1.44 ± 0.16	0.5
19244	Daffah-C2	5	1.25 ± 0.19	0.5
19245	Khor al Jaramah-C1	5	0.89 ± 0.06	7.1
19246	Khor al Jaramah-C2	5	0.92 ± 0.03	7.1

Tab. 4. Dose recovery test results obtained on quartz. The dose (Gy) is the artificial dose applied after bleaching the samples in a solar simulator (see text).

Lab #	Site	^{238}U (ppm)	^{232}Th (ppm)	K (%)	Water content (weight%)
19226	Ruways 1	2.36 ± 0.02	0.57 ± 0.02	0.16 ± 0.01	3
19227	Ruways 1	2.25 ± 0.02	0.83 ± 0.02	0.25 ± 0.01	3
19228	Ruways 1	2.25 ± 0.02	0.60 ± 0.02	0.19 ± 0.01	3
19229	Ruways 1	0.53 ± 0.01	1.12 ± 0.01	0.30 ± 0.01	3
19230	Ruways 1	3.16 ± 0.04	4.18 ± 0.04	0.36 ± 0.01	3
19231	Ruways 1	3.20 ± 0.04	4.62 ± 0.04	0.28 ± 0.01	3
19232 ^a	Ruways 1	2.42 ± 0.02	2.65 ± 0.02	0.27 ± 0.01	3
19233	Ruways 1	2.36 ± 0.03	0.88 ± 0.03	0.20 ± 0.01	3
19234	Ruways 1	2.31 ± 0.03	0.92 ± 0.03	0.24 ± 0.01	3
19235	Ruways 1	1.57 ± 0.02	3.91 ± 0.02	0.85 ± 0.01	3
19236	Ruways 1	2.19 ± 0.02	0.86 ± 0.02	0.25 ± 0.01	3
19237 ^a	Ruways 1	1.55 ± 0.02	1.11 ± 0.02	0.25 ± 0.01	3
19238	Suwayh 1	1.28 ± 0.02	1.91 ± 0.02	0.62 ± 0.01	9
19239	Suwayh 1	1.18 ± 0.02	1.79 ± 0.02	0.51 ± 0.01	9
19240	Suwayh 1	2.67 ± 0.02	3.28 ± 0.02	0.44 ± 0.01	9
19241	Ruways-C1	1.46 ± 0.02	3.14 ± 0.02	1.15 ± 0.01	19
19242	Ruways-C1	4.07 ± 0.04	6.46 ± 0.04	0.56 ± 0.01	12
19243	Daffah-C2	1.20 ± 0.03	1.50 ± 0.03	0.80 ± 0.01	17
19244	Daffah-C2	1.18 ± 0.01	2.06 ± 0.01	0.92 ± 0.01	13
19245 ^a	Khor al Jaramah-C1	1.07 ± 0.03	4.34 ± 0.03	1.02 ± 0.01	17
19246 ^a	Khor al Jaramah-C2	1.13 ± 0.02	4.48 ± 0.05	1.26 ± 0.02	27

Tab. 5. Radioisotopes content measured in the sediment samples using BEGe gamma-ray spectrometry and water content used for age calculation.

^a no in-situ gamma dosimetry measurements were performed for these samples. The gamma dose rates were derived using U, Th and K content.

Lab #	Site	n	Dose rate ($\mu\text{Gy}\cdot\text{a}^{-1}$)					Total	D_e FMM (Gy)	Age FMM (years)	D_e CAM (Gy)	Age CAM (years)	Age (years BC)
			Alpha	Beta	Gamma	Cosmic	Internal						
19226 (Q)	Ruways 1 (Phase II)	32	16 ± 3	404 ± 4	329 ± 16	152 ± 10		901 ± 17	6.98 ± 0.37	7 740 ± 580	7.91 ± 0.32	8 780 ± 590	5 730 ± 580
19227 (Q)	Ruways 1 (Phase II)	33	16 ± 3	463 ± 5	333 ± 17	161 ± 10		973 ± 18	8.12 ± 0.16	8 340 ± 480	8.55 ± 0.53	8 780 ± 720	6 320 ± 480
19228 (Q)	Ruways 1 (Phase II)	33	16 ± 3	418 ± 5	291 ± 15	171 ± 10		896 ± 16	6.80 ± 0.21	7 590 ± 470	7.56 ± 0.43	8 440 ± 650	5 570 ± 470
19229 (F)	Ruways 1 (Phase I)	8	14 ± 2	298 ± 5	210 ± 10	161 ± 20	500 ± 50	1183 ± 51	-	-	520.66 ± 79.78	440 000 ± 76 000	438 000 ± 76 000
19230 (Q)	Ruways 1 (Phase IX)	33	30 ± 5	721 ± 10	325 ± 16	194 ± 10		1271 ± 19	0.13 ± 0.01	100 ± 10	0.29 ± 0.05	230 ± 40	-
19231 (Q)	Ruways 1 (Phase IX)	38	32 ± 5	677 ± 9	405 ± 20	207 ± 10		1321 ± 23	0.33 ± 0.01	250 ± 20	0.53 ± 0.08	400 ± 60	-
19232 (Q)*	Ruways 1 (Phase IX)	40	22 ± 3	533 ± 5	444 ± 3	236 ± 10		1235 ± 7	4.21 ± 0.09	3 410 ± 190	3.63 ± 0.32	2 940 ± 290	1 390 ± 190
19233 (Q)	Ruways 1 (Phase VII)	24	17 ± 3	440 ± 8	280 ± 14	200 ± 10		937 ± 16	5.56 ± 0.19	5 930 ± 370	6.46 ± 0.45	6 890 ± 590	3 910 ± 370
19234 (Q)	Ruways 1 (Phase VIII)	22	17 ± 3	465 ± 10	289 ± 14	227 ± 10		998 ± 17	5.82 ± 0.13	5 830 ± 330	5.82 ± 0.12	5 830 ± 320	3 810 ± 320
19235 (F)	Ruways 1 (Phase II)	5	43 ± 8	866 ± 9	323 ± 16	150 ± 20	500 ± 50	1882 ± 54	-	-	14.78 ± 1.55	7 850 ± 1 040	5 830 ± 1 040
19236 (Q)	Ruways 1 (Phase II)	37	16 ± 3	454 ± 6	291 ± 15	158 ± 10		919 ± 16	7.37 ± 0.15	8 020 ± 460	8.37 ± 0.59	9 110 ± 800	6 000 ± 460
19237 (Q)*	Ruways 1 (Phase VIII)	27	17 ± 3	392 ± 4	276 ± 2	236 ± 10		921 ± 5	4.82 ± 0.1	5 230 ± 270	4.96 ± 0.13	5 390 ± 290	3 370 ± 290
19238 (Q)	Suwayh 1	37	16 ± 2	600 ± 7	241 ± 12	178 ± 10		1035 ± 14	6.40 ± 0.13	6 180 ± 310	6.74 ± 0.23	6 510 ± 370	4 160 ± 310
19239 (Q)	Suwayh 1	38	14 ± 2	513 ± 8	207 ± 10	203 ± 10		937 ± 13	5.76 ± 0.15	6 140 ± 320	5.91 ± 0.13	6 300 ± 320	4 290 ± 320
19240 (Q)	Suwayh 1	38	31 ± 5	672 ± 6	345 ± 17	246 ± 10		1293 ± 19	7.08 ± 0.15	5 480 ± 270	7.43 ± 0.25	5 750 ± 320	3 460 ± 270
19241 (Q)	Ruways-C1	24	21 ± 3	913 ± 9	475 ± 24	200 ± 10		1609 ± 26	5.01 ± 0.26	3 110 ± 370	5.39 ± 0.44	3 350 ± 450	1 100 ± 370
19242 (Q)	Ruways-C1	7	57 ± 8	959 ± 8	478 ± 24	236 ± 10		1730 ± 27	-	-	1.68 ± 0.51	970 ± 290	-
19243 (Q)	Daffah-C2	22	13 ± 2	643 ± 13	372 ± 19	187 ± 10		1214 ± 23	0.66 ± 0.05	540 ± 70	0.70 ± 0.08	580 ± 90	-
19244 (Q)	Daffah-C2	14	17 ± 2	769 ± 8	331 ± 17	236 ± 10		1352 ± 19	-	-	0.57 ± 0.06	420 ± 50	-
19245 (Q)*	Khor al Jaramah-C1	9	22 ± 3	829 ± 15	481 ± 6	184 ± 10		1516 ± 16	-	-	7.92 ± 0.42	5 220 ± 570	3 210 ± 570
19246 (Q)*	Khor al Jaramah-C2	7	20 ± 3	893 ± 9	494 ± 4	184 ± 10		1591 ± 10	-	-	8.46 ± 0.38	5 320 ± 590	3 300 ± 590

Tab. 6. OSL ages on quartz (Q) and pIRIR₂₂₅ ages on feldspar (F). Ages were rounded to ten years, except for BDX 19229, rounded to thousand years. The error on the age takes into account the systematic error. For samples with less than 22 aliquots measured, no FMM ages were calculated. CAM overdispersion values are presented in SOM Tab. S1 FMM components are provided in SOM Tab. S2. Age values used for interpretation are shown in bold text and are converted in years BC in the last column.

*no in-situ gamma dosimetry measurements were performed for these samples (see text).