



HAL
open science

Sign choices in the AGM for genus two theta constants

Jean Kieffer

► **To cite this version:**

Jean Kieffer. Sign choices in the AGM for genus two theta constants. Publications Mathématiques de Besançon : Algèbre et Théorie des Nombres, 2022, pp.37-58. 10.5802/pmb.45 . hal-02967220v2

HAL Id: hal-02967220

<https://hal.science/hal-02967220v2>

Submitted on 9 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sign choices in the AGM for genus two theta constants

Jean Kieffer

October 8, 2022

Abstract

Existing algorithms to compute genus 2 theta constants in quasi-linear time use Borchartd sequences, an analogue of the arithmetic-geometric mean for four complex numbers. In this paper, we show that these Borchartd sequences are only given by good choices of square roots, as in the genus 1 case. This removes the sign indeterminacies when computing genus 2 theta constants without relying on numerical integration.

Résumé

Les algorithmes existants pour le calcul de thêta-constantes en genre 2 en temps quasilinéaire utilisent des suites de Borchartd, un analogue de la moyenne arithmético-géométrique pour quatre nombres complexes. Dans cet article, nous montrons que ces suites de Borchartd sont constituées uniquement de bons choix de signes, comme c'est le cas en genre 1. Ce résultat permet de lever les indéterminations de signes lors du calcul de thêta-constantes en genre 2 sans recours à l'intégration numérique.

Keywords: Theta functions, Genus 2, Algorithms, Borchartd mean
Subject classification (2010): 11Y35; 11Y16; 11F41; 11F27

1 Introduction

Denote by \mathcal{H}_g the Siegel half space of principally polarized abelian varieties of dimension g , consisting of all matrices $\tau \in M_g(\mathbb{C})$ such that τ is symmetric and $\text{Im}(\tau)$ is positive definite; for instance, \mathcal{H}_1 is the usual upper half plane. The *theta constants* are the holomorphic functions on \mathcal{H}_g defined by

$$\theta_{a,b}(\tau) = \sum_{m \in \mathbb{Z}^g} \exp \left(i\pi \left(\left(m + \frac{a}{2} \right)^t \tau \left(m + \frac{a}{2} \right) + \left(m + \frac{a}{2} \right)^t b \right) \right), \quad (1)$$

where a and b run through $\{0, 1\}^g$ (by convention, vectors in formula (1) are written vertically). Theta constants have a fundamental importance in the

theory of Siegel modular forms, as every scalar-valued Siegel modular function of any weight on \mathcal{H}_g has an expression in terms of quotients of theta constants [15, Thm. 9 p. 222]. Moreover, for $1 \leq g \leq 3$, then the stronger result that every Siegel modular form is a polynomial in the theta constants holds [13, 14, 9].

In numerical algorithms manipulating modular forms, the following operations are therefore very common: first, given (quotients of) theta constants at a given $\tau \in \mathcal{H}_g$, compute τ ; second, given $\tau \in \mathcal{H}_g$, compute the theta constants $\theta_{a,b}(\tau)$. For instance, these operations are important building blocks in algorithms computing modular polynomials [7, 20, 21] or Hilbert class polynomials [6, 8, 25] via complex approximations.

The arithmetic-geometric mean (AGM) [1, 3, 2, 16] gives an algorithm to find τ given its theta constants. This algorithm is quasi-linear in terms of the required precision. In order to compute theta constants in quasi-linear time as well, a well-studied strategy is to combine the AGM with Newton iterations. This strategy was first described in [5] in the genus 1 case, in [4] in the genus 2 case, and later extended to theta *functions*, in opposition to theta *constants*, in [18, 19]. These references also outline extensions to higher genus.

The genus 1 case. Let us detail the genus 1 case to convey the general idea. After reducing the argument $\tau \in \mathcal{H}_1$ using Gauss's algorithm [25, §6.1], we can assume that τ belongs to the classical fundamental domain under the action of $\mathrm{SL}_2(\mathbb{Z})$, denoted by \mathcal{F}_1 .

First assume that theta quotients at $\tau \in \mathcal{F}_1$ are given. Then the sequence

$$B(\tau) = \left(\frac{\theta_{0,0}^2(2^n \tau)}{\theta_{0,0}^2(\tau)}, \frac{\theta_{0,1}^2(2^n \tau)}{\theta_{0,0}^2(\tau)} \right)_{n \geq 0}$$

is an *AGM sequence*, meaning that each term is obtained from the previous one by means of the transformation

$$(x, y) \mapsto \left(\frac{x+y}{2}, \sqrt{x}\sqrt{y} \right)$$

for some choice of the square roots. This is a consequence of the duplication formula [23, p. 221], the correct square roots being the theta quotients themselves. In the algorithm, the sign ambiguity is easily removed using the fact that \sqrt{x} and \sqrt{y} should lie in a common open quarter plane [5, Thm. 2]: we say that the sequence $B(\tau)$ is given by *good sign choices*. It converges quadratically to $1/\theta_{0,0}^2(\tau)$, as the series expansion (1) shows.

It turns out that the sequence $B(-1/\tau)$ is also an AGM sequence with good sign choices [5, Prop. 7]. Its first term can be computed from theta quotients at τ using the transformation formulas for theta constants under $\mathrm{SL}_2(\mathbb{Z})$. The limit of $B(-1/\tau)$ is $1/\theta_{0,0}^2(-1/\tau)$. Finally, we can recover τ using the formula

$$\theta_{0,0}^2 \left(\frac{-1}{\tau} \right) = -i\tau\theta_{0,0}^2(\tau). \quad (2)$$

Since AGM sequences with good sign choices converge quadratically, this gives an algorithm to *invert* theta functions on \mathcal{F}_1 with quasi-linear complexity in the output precision, at least for fixed τ . This method was already known to Gauss [10, X.1, pp. 184–206], and we recommend [3, §3C] for a historical exposition of Gauss’s works on the AGM and elliptic functions.

In order to compute theta functions at a given $\tau \in \mathcal{F}_1$, the most efficient known method is to build a Newton scheme [5], using the AGM method to invert theta constants. This yields a quasi-linear algorithm to compute genus 1 theta constants, whose complexity can be made uniform in $\tau \in \mathcal{F}_1$ [5, Thm. 5].

The genus 2 case. A similar strategy can be applied to theta functions in genus 2, using *Borchardt sequences*, a generalization of AGM sequences for four complex numbers [1, 2, 16]. Let us refer to §2 for the definition of Borchardt sequences, the numbering of genus 2 theta constants, and the definition of the matrices $\gamma_k \in \mathrm{Sp}_4(\mathbb{Z})$ for $0 \leq k \leq 3$. The Borchardt sequences we consider are the sequences $B(\gamma_k \tau)$ for $0 \leq k \leq 3$, where

$$B(\tau) = \left(\frac{\theta_0^2(2^n \tau)}{\theta_0^2(\tau)}, \frac{\theta_1^2(2^n \tau)}{\theta_0^2(\tau)}, \frac{\theta_2^2(2^n \tau)}{\theta_0^2(\tau)}, \frac{\theta_3^2(2^n \tau)}{\theta_0^2(\tau)} \right)_{n \geq 0}$$

for every $\tau \in \mathcal{H}_2$. Their first terms are given by different combinations of theta quotients at τ (see Corollary 3.3). It is known that for a given τ , all but a finite number of sign choices in these Borchardt sequences are good, and the other sign choices can be determined using certified computations of hyperelliptic integrals at relatively low precision: see the discussion before Prop. 3.3 in [19], and [22] for an algorithm that provides this input. However, the required precision and the cost of the numerical integration algorithms depend heavily on τ .

Actually, when τ belongs to the usual fundamental domain \mathcal{F}_2 under the action of $\mathrm{Sp}_4(\mathbb{Z})$, practical experiments suggest that *all* sign choices are good in the genus 2 algorithm as well [4, Conj. 9.1], [8, Conj. 9]. The goal of this paper is to prove this fact. More precisely, we define in §2 a subset $\mathcal{F}' \subset \mathcal{H}_2$ containing \mathcal{F}_2 , and prove the following result.

Theorem 1.1. *For every $\tau \in \mathcal{F}'$, every $0 \leq k \leq 3$ and every $n \geq 0$, the theta constants*

$$\theta_j(2^n \gamma_k \tau) \quad \text{for } 0 \leq j \leq 3$$

are contained in a common open quarter plane.

Dupont [4, Prop. 9.1] proved this result in the particular case of $\gamma_0 = I_4$.

As a consequence, we can invert genus 2 theta constants in quasi-linear time by using only Borchardt sequences with good sign choices. On the practical side, this result reduces the effort needed to invert genus 2 theta constants with controlled precision losses; see for instance [4, §7.4.2] for an analysis of precision losses when computing limits of Borchardt sequences. On the theoretical side, we hope that our result can be a first step towards removing other heuristic assumptions when computing genus 2 theta constants (in particular, the assumption [4, §10.2] that the function used in the Newton scheme is analytic

with invertible Jacobian), and obtaining algorithms with uniform complexity in $\tau \in \mathcal{F}_2$.

This document is organized as follows. In Section 2, we introduce our notational conventions. In Section 3, we use the action of the symplectic group to bring the matrices $2^n \gamma_k \tau \in \mathcal{H}_2$ closer to the cusp at infinity: this is critical to obtain accurate information from the series expansion (1). We give estimates on genus 2 theta constants in Section 4, and we finish the proof of the main theorem in Section 5.

Acknowledgement. The author would like to thank Aurel Page and the anonymous referees for their careful reading and helpful suggestions to improve the exposition.

2 Theta constants and Borchartd sequences

We define a *Borchartd sequence* to be a sequence of complex numbers

$$(s_b^{(n)})_{b \in (\mathbb{Z}/2\mathbb{Z})^2, n \geq 0}$$

with the following property: for every $n \geq 0$, there exist $t_b^{(n)}$ for $b \in (\mathbb{Z}/2\mathbb{Z})^2$ such that $t_b^{(n)}$ is a square root of $s_b^{(n)}$, and

$$s_b^{(n+1)} = \frac{1}{4} \sum_{b_1+b_2=b} t_{b_1}^{(n)} t_{b_2}^{(n)} \quad \text{for each } b \in (\mathbb{Z}/2\mathbb{Z})^2.$$

The duplication formula [23, p. 221] states that for every $\tau \in \mathcal{H}_2$, the sequence

$$B(\tau) = (\theta_{0,b}^2(2^n \tau))_{b \in \{0,1\}^2, n \geq 0}$$

is a Borchartd sequence; the choice of square roots at each step is given by the theta constants $\theta_{0,b}(2^n \tau)$ themselves. By the series expansion (1), we have

$$\theta_{0,b}(2^n \tau) = \sum_{m \in \mathbb{Z}^2} \exp(-2^n \pi m^t \operatorname{Im}(\tau) m) \exp(i\pi (2^n m^t \operatorname{Re}(\tau) m + m^t b)).$$

When n tends to infinity, all the terms except $m = 0$ converge rapidly to zero, because $\operatorname{Im}(\tau)$ is positive definite. Therefore the Borchartd sequence $B(\tau)$ converges to $(1, 1, 1, 1)$.

We say that a set of complex numbers is *in good position* when it is included in an open quarter plane seen from the origin, i.e. a set of the form

$$\{r \exp(i(\alpha_0 + \alpha)) \mid r > 0 \text{ and } 0 < \alpha < \pi/2\}$$

for some $\alpha_0 \in \mathbb{R}$. The property of being in good position is invariant by nonzero complex scaling. A Borchartd sequence is given by *good sign choices* if for every $n \geq 0$, the complex numbers $t_b^{(n)}$ for $b \in (\mathbb{Z}/2\mathbb{Z})^2$ are in good position.

Let us now detail the algorithm to recover $\tau \in \mathcal{H}_2$ from its theta quotients. We first introduce the matrices $\gamma_k \in \mathrm{Sp}_4(\mathbb{Z})$ alluded to in the introduction. Let

$$S_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, S_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, S_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

and define the matrix $\gamma_k \in \mathrm{Sp}_4(\mathbb{Z})$ for $0 \leq k \leq 3$ by

$$\gamma_0 = I_4, \text{ and } \gamma_k = \begin{pmatrix} -I_2 & -S_k \\ S_k & -I + S_k^2 \end{pmatrix} \text{ for } 1 \leq k \leq 3.$$

For convenience, we also introduce a numbering of theta constants [4, §6.2]:

$$\theta_{(a_0, a_1), (b_0, b_1)} =: \theta_j \quad \text{where } j = b_0 + 2b_1 + 4a_0 + 8a_1 \in \llbracket 0, 15 \rrbracket.$$

Assuming that the choices of square roots in the sequences $B(\gamma_k \tau)$ can be determined, we can compute $\tau \in \mathcal{F}_2$ from its theta quotients as follows.

Algorithm 2.1 ([4, §9.2.3]).

Input: The projective vector of squares of theta constants $\theta_j^2(\tau)$ for $j \in \llbracket 0, 15 \rrbracket$, for some $\tau \in \mathcal{H}_2$.

Output: The matrix τ .

1. For each $0 \leq k \leq 3$, compute the first term of the sequence $B(\gamma_k \tau)/\theta_0^2(\gamma_k \tau)$ using the transformation formulas for theta constants under $\mathrm{Sp}_4(\mathbb{Z})$ (see Igusa [15, Thm. 2 p. 175 and Cor. p. 176], or Corollary 3.3);
2. For each $0 \leq k \leq 3$, compute $1/\theta_0^2(\gamma_k \tau)$ as the limit of the Borchartd sequence $B(\gamma_k \tau)/\theta_0^2(\gamma_k \tau)$;
3. Use the input and the newly computed $\theta_0^2(\gamma_0 \tau) = \theta_0^2(\tau)$ to compute all squares of theta constants at τ ;
4. Recover $\tau = \begin{pmatrix} z_1 & z_3 \\ z_3 & z_2 \end{pmatrix}$ using the relations given in [4, §6.3.1]:

$$\theta_0^2(\gamma_1 \tau) = -iz_1 \theta_4^2(\tau), \quad \theta_0^2(\gamma_2 \tau) = -iz_2 \theta_8^2(\tau), \quad \theta_0^2(\gamma_3 \tau) = -\det(\tau) \theta_0^2(\tau).$$

In the sequel, we use the following notational conventions. For $\tau \in \mathcal{H}_2$, we write

$$\tau = \begin{pmatrix} z_1(\tau) & z_3(\tau) \\ z_3(\tau) & z_2(\tau) \end{pmatrix} \quad \text{and} \quad \begin{cases} x_j(\tau) = \mathrm{Re} z_j(\tau) \\ y_j(\tau) = \mathrm{Im} z_j(\tau) \end{cases} \quad \text{for } 1 \leq j \leq 3.$$

For $1 \leq j \leq 3$, we also write

$$q_j(\tau) = \exp(-\pi y_j(\tau)).$$

We denote by $\lambda_1(\tau)$ the smallest eigenvalue of $\mathrm{Im}(\tau)$, and define

$$r(\tau) = \min \left\{ \lambda_1(\tau), \frac{y_1(\tau)}{2}, \frac{y_2(\tau)}{2} \right\}.$$

We often omit the argument τ to ease notation. We define \mathcal{F}' to be the set of all $\tau \in \mathcal{H}_2$ such that the following conditions are satisfied:

$$\begin{aligned} |x_j(\tau)| &\leq \frac{1}{2} \quad \text{for each } 1 \leq j \leq 3, \\ 2|y_3(\tau)| &\leq y_1(\tau) \leq y_2(\tau), \\ y_1(\tau) &\geq \frac{\sqrt{3}}{2}, \\ |z_j(\tau)| &\geq 1 \quad \text{for } j \in \{1, 2\}. \end{aligned} \tag{3}$$

The domain \mathcal{F}' contains the classical fundamental domain \mathcal{F}_2 for the action of $\mathrm{Sp}_4(\mathbb{Z})$ on \mathcal{H}_2 [17, Prop. 3 p. 33]. Assumptions similar to (3) are usual when giving analytic estimates on theta constants: for instance, the domain \mathcal{B} in [25] is defined by the first three inequalities of (3).

Finally, for each $\tau \in \mathcal{H}_2$, we write

$$\begin{aligned} \xi_{4,6}(\tau) &= 2 \exp\left(i\pi \frac{z_1(\tau)}{4}\right), \\ \xi_{8,9}(\tau) &= 2 \exp\left(i\pi \frac{z_2(\tau)}{4}\right), \\ \xi_0(\tau) &= 1 + 2 \exp(i\pi z_1(\tau)) + 2 \exp(i\pi z_2(\tau)), \\ \xi_{0,2}(\tau) &= 1 + 2 \exp(i\pi z_1(\tau)), \\ \xi_{0,1}(\tau) &= 1 + 2 \exp(i\pi z_2(\tau)), \quad \text{and} \\ \xi_{12}(\tau) &= \exp\left(i\pi \frac{z_1(\tau) + z_2(\tau)}{4}\right) \left(\exp\left(i\pi \frac{z_3(\tau)}{2}\right) + \exp\left(-i\pi \frac{z_3(\tau)}{2}\right) \right). \end{aligned} \tag{4}$$

These complex numbers correspond to the first term(s) of the series defining theta constants at τ . For instance, $\xi_{4,6}(\tau)$ approximates both $\theta_4(\tau)$ and $\theta_6(\tau)$. We will recall the definitions (4) before using them in the computations of §4.

3 Other expressions for theta constants at $2^n \gamma_k \tau$

For every $n \geq 0$, we define

$$\begin{aligned} \eta_1^{(n)} &= \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 2^n & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & \eta_2^{(n)} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 2^n \end{pmatrix}, \\ \eta_3^{(n)} &= \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 2^n & 0 \\ 1 & 0 & 0 & 2^n \end{pmatrix}, & \text{and } \eta_4^{(n)} &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \eta_3^{(n)}. \end{aligned}$$

Lemma 3.1. *Let $n \geq 0$.*

1. *For every $1 \leq k \leq 4$, the matrix $\eta_k^{(n)}$ belongs to $\mathrm{Sp}_4(\mathbb{Z})$.*

2. For every $\tau = \begin{pmatrix} z_1 & z_3 \\ z_3 & z_2 \end{pmatrix} \in \mathcal{H}_2$, we have

$$\begin{aligned}\tau_1^{(n)} &:= \eta_1^{(n)}(2^n \gamma_1 \tau) = \begin{pmatrix} 2^{-n} z_1 & z_3 \\ z_3 & 2^n z_2 \end{pmatrix}, \\ \tau_2^{(n)} &:= \eta_2^{(n)}(2^n \gamma_2 \tau) = \begin{pmatrix} 2^n z_1 & z_3 \\ z_3 & 2^{-n} z_2 \end{pmatrix}, \\ \tau_3^{(n)} &:= \eta_3^{(n)}(2^n \gamma_3 \tau) = 2^{-n} \tau, \quad \text{and} \\ \tau_4^{(n)} &:= \eta_4^{(n)}(2^n \gamma_3 \tau) = \begin{pmatrix} -2^n/z_1 & -z_3/z_1 \\ -z_3/z_1 & 2^{-n}(z_2 - z_3^2/z_1) \end{pmatrix}.\end{aligned}\tag{5}$$

Proof. 1. The lines of each $\eta_k^{(n)}$ define a symplectic basis of \mathbb{Z}^4 .

2. The action of $\mathrm{Sp}_4(\mathbb{Z})$ on \mathcal{H}_2 extends to an action of the larger group

$$\mathrm{GSp}_4(\mathbb{Q}) = \left\{ \gamma \in \mathrm{GL}_4(\mathbb{Q}) \mid \exists \mu \in \mathbb{Q}^\times, \gamma^t \begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix} \gamma = \mu \begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix} \right\}.$$

The matrix $2^n \gamma_k \tau$ is the image of τ under

$$\begin{pmatrix} -2^n I_2 & -2^n S_k \\ S_k & -I + S_k^2 \end{pmatrix} \in \mathrm{GSp}_4(\mathbb{Q}).$$

When we multiply this matrix by $\eta_k^{(n)}$ on the left, we obtain

$$\begin{aligned}\mathrm{Diag}(-1, -2^n, -2^n, -1) & \quad \text{for } k = 1, \\ \mathrm{Diag}(-2^n, -1, -1, -2^n) & \quad \text{for } k = 2, \text{ and} \\ \mathrm{Diag}(-1, -1, -2^n, -2^n) & \quad \text{for } k = 3. \quad \square\end{aligned}$$

We recall the transformation formulas for theta constants in genus 2. For a square matrix m , we denote by m_0 the column vector containing the diagonal of m .

Proposition 3.2 ([15, Thm. 2 p.175 and Cor. p.176]). *Let $a, b \in \{0, 1\}^2$, and let*

$$\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}_4(\mathbb{Z}).$$

Define

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} = \gamma^t \begin{pmatrix} a - (CD^t)_0 \\ b - (AB^t)_0 \end{pmatrix}.$$

Then, for every $\tau \in \mathcal{H}_2$, we have

$$\theta_{a,b}(\gamma\tau) = \kappa(\gamma) \zeta_8^{\varepsilon(\gamma,a,b)} \det(C\tau + D)^{1/2} \theta_{\alpha',\beta'}(\tau)$$

where

$$\zeta_8 = e^{i\pi/4}, \quad \begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} \alpha \\ \beta \end{pmatrix} \pmod{2},$$

$$\varepsilon(\gamma, a, b) = 2(B\alpha)^t(C\beta) - (D\alpha)^t(B\alpha) - (C\beta)^t(A\beta) + 2((AB^t)_0)^t(D\alpha - C\beta),$$

and $\kappa(\gamma)$ is an eighth root of unity depending only on γ , with a sign ambiguity coming from the choice of a holomorphic square root of $\det(C\tau + D)$.

Corollary 3.3. *For every $\tau \in \mathcal{H}_2$, we have the following equalities of projective tuples:*

$$\begin{aligned} (\theta_j(2^n \gamma_1 \tau))_{0 \leq j \leq 3} &= \begin{cases} (\theta_4(\tau) : \theta_0(\tau) : \theta_6(\tau) : \theta_2(\tau)) & \text{if } n = 0, \\ (\theta_0(\tau_1^{(n)}) : \theta_4(\tau_1^{(n)}) : \theta_2(\tau_1^{(n)}) : \theta_6(\tau_1^{(n)})) & \text{if } n \geq 1, \end{cases} \\ (\theta_j(2^n \gamma_2 \tau))_{0 \leq j \leq 3} &= \begin{cases} (\theta_8(\tau) : \theta_9(\tau) : \theta_0(\tau) : \theta_1(\tau)) & \text{if } n = 0, \\ (\theta_0(\tau_2^{(n)}) : \theta_1(\tau_2^{(n)}) : \theta_8(\tau_2^{(n)}) : \theta_9(\tau_2^{(n)})) & \text{if } n \geq 1, \end{cases} \\ (\theta_j(2^n \gamma_3 \tau))_{0 \leq j \leq 3} &= (\theta_0(\tau_3^{(n)}) : \theta_8(\tau_3^{(n)}) : \theta_4(\tau_3^{(n)}) : \theta_{12}(\tau_3^{(n)})) \quad \text{for every } n \geq 0, \\ (\theta_j(2^n \gamma_3 \tau))_{0 \leq j \leq 3} &= (\theta_0(\tau_4^{(n)}) : \theta_8(\tau_4^{(n)}) : \theta_1(\tau_4^{(n)}) : \theta_9(\tau_4^{(n)})) \quad \text{for every } n \geq 0, \end{aligned}$$

where the $\tau_j^{(n)}$ are defined as in (5).

Proof. Apply Proposition 3.2 to the matrices $\eta_i^{(n)}$. □

When $\tau \in \mathcal{F}'$, the real and imaginary parts of $\tau_k^{(n)}$ for $1 \leq k \leq 3$ are easy to study: for instance, from the second inequality in (3) we always have

$$y_3(\tau_k^{(n)})^2 \leq \frac{1}{4} y_1(\tau_k^{(n)}) y_2(\tau_k^{(n)}).$$

Such estimates are less obvious for the matrices $\tau_4^{(n)}$.

Lemma 3.4. *Let $\tau \in \mathcal{F}'$. Then, for every $n \geq 0$, we have*

$$\begin{aligned} |y_3(\tau_4^{(n)})| &\leq \frac{3}{2^{n+2}} y_1(\tau_4^{(n)}), \\ y_3(\tau_4^{(n)})^2 &\leq \frac{3}{7} y_1(\tau_4^{(n)}) y_2(\tau_4^{(n)}), \quad \text{and} \\ |x_2(\tau_4^{(n)})| &\leq \frac{9}{2^{n+3}}. \end{aligned}$$

Proof. Write z_1 for $z_1(\tau)$, etc. We have

$$y_3(\tau_4^{(n)}) = \text{Im}(-z_3/z_1) = \frac{1}{|z_1|^2} (x_3 y_1 - y_3 x_1),$$

so

$$|y_3(\tau_4^{(n)})| \leq \frac{3y_1}{4|z_1|^2} = \frac{3}{2^{n+2}} y_1(\tau_4^{(n)}),$$

since $y_1(\tau_4^{(n)}) = 2^n y_1 / |z_1|^2$ by (5). For the second inequality, we have

$$\operatorname{Im}(\tau_4^{(n)}) = \begin{pmatrix} 2^{-n} z_1 & -2^{-n} z_3 \\ 0 & 1 \end{pmatrix}^{-t} (2^{-n} \operatorname{Im} \tau) \begin{pmatrix} 2^{-n} \bar{z}_1 & -2^{-n} \bar{z}_3 \\ 0 & 1 \end{pmatrix}^{-1}$$

so

$$\det \operatorname{Im}(\tau_4^{(n)}) = \frac{1}{|z_1|^2} \det \operatorname{Im} \tau.$$

Moreover $\det \operatorname{Im} \tau \geq \frac{3}{4} y_1^2$, so

$$\frac{y_3(\tau_4^{(n)})^2}{y_1(\tau_4^{(n)}) y_2(\tau_4^{(n)})} \leq \frac{y_3(\tau_4^{(n)})^2}{y_3(\tau_4^{(n)})^2 + \frac{3y_1^2}{4|z_1|^2}} \leq \frac{1}{1 + \frac{4}{3}|z_1|^2} \leq \frac{3}{7}.$$

For the last inequality, we compute

$$2^n x_2(\tau_4^{(n)}) = x_2 - \frac{1}{|z_1|^2} ((x_3^2 - y_3^2)x_1 + 2x_3 y_3 y_1)$$

and

$$\left| \frac{1}{|z_1|^2} (x_3^2 - y_3^2)x_1 \right| \leq \frac{1}{2} \max\{x_3^2, \frac{y_3^2}{|z_1|^2}\} \leq \frac{1}{8},$$

so

$$|2^n x_2(\tau_4^{(n)})| \leq \frac{1}{2} + \frac{1}{8} + \frac{1}{2} = \frac{9}{8}. \quad \square$$

4 Bounds on theta constants

Typically, when $\tau \in \mathcal{H}_2$ is close enough to the cusp at infinity (more precisely when $\operatorname{Im} z_1(\tau)$, $\operatorname{Im} z_2(\tau)$, and $\det \operatorname{Im}(\tau)$ are large), useful information on theta constants at τ can be obtained from the series expansion (1). Our computations are similar in spirit to those found in [17, pp. 116–117], [4, §6.2], [11, §5.1]. All our estimates are based on the following key lemma.

Lemma 4.1. *Let $f: \mathbb{N} \rightarrow \mathbb{R}$ be a strictly increasing function, and assume that $f(k+2) - f(k+1) \geq f(k+1) - f(k)$ for every $k \geq 0$. Let $0 < q < 1$. Then*

$$\sum_{k=0}^{\infty} q^{f(k)} \leq \frac{q^{f(0)}}{1 - q^{f(1)-f(0)}}.$$

Proof. Use that $f(k) \geq f(0) + k(f(1) - f(0))$ for all k . □

Lemma 4.2. *Let $k \geq 1$, and let $\tau \in \mathcal{H}_2$ such that*

$$y_3(\tau)^2 \leq \frac{1}{4} y_1(\tau) y_2(\tau) \quad \text{and} \quad k |y_3(\tau)| \leq y_2(\tau).$$

Define

$$\xi_{4,6}(\tau) = 2 \exp\left(i\pi \frac{z_1(\tau)}{4}\right)$$

and

$$\begin{aligned} \rho_{4,6}^{(k)}(q_1, q_2) &= \frac{q_1^2}{1 - q_1^4} + \frac{q_2^{1-\frac{1}{k}}}{1 - q_2^{3-\frac{1}{k}}} + \frac{q_2^{1+\frac{1}{k}}}{1 - q_2^{3+\frac{1}{k}}} \\ &\quad + \frac{q_1^{7/8} q_2^{1/2}}{(1 - q_2^{3/2})(1 - q_1^2)} + \frac{q_1^{25/8} q_2^{3/2}}{(1 - q_2^{9/2})(1 - q_1^6)}. \end{aligned}$$

Then for $j \in \{4, 6\}$, we have

$$\left| \frac{\theta_j(\tau)}{\xi_{4,6}(\tau)} - 1 \right| \leq \rho_{4,6}^{(k)}(q_1(\tau), q_2(\tau)).$$

Proof. Write $u = \begin{pmatrix} 1/2 \\ 0 \end{pmatrix}$. Using the definition, we obtain

$$\left| \frac{\theta_j(\tau)}{\xi_{4,6}(\tau)} - 1 \right| \leq \frac{1}{2} q_1^{-1/4} \sum_{\substack{m \in \mathbb{Z}^2 \\ m \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix}}} \exp(-\pi(m+u)^t \operatorname{Im}(\tau)(m+u)).$$

We split this sum in two parts, according to whether the second coordinate of m is zero or not. The first part gives

$$q_1^{-1/4} \sum_{m \in \mathbb{N} + \frac{3}{2}} q_1^{m^2} \leq q_1^{-1/4} \frac{q_1^{9/4}}{1 - q_1^4} = \frac{q_1^2}{1 - q_1^4}$$

by Lemma 4.1. The second part is

$$q_1^{-1/4} \sum_{m_1 \in \mathbb{N} + \frac{1}{2}} \sum_{m_2 \geq 1} q_1^{m_1^2} q_2^{m_2^2} \cdot 2 \cosh(2\pi y_3 m_1 m_2).$$

We use the fact that for every $(m_1, m_2) \in \mathbb{R}_+^2$,

$$|2y_3 m_1 m_2| \leq \frac{y_1}{2} m_1^2 + \frac{y_2}{2} m_2^2.$$

When $m_1 = 1/2$, we use the following bound instead:

$$|2y_3 m_1 m_2| = |y_3 m_2| \leq \frac{y_2 m_2}{k}.$$

Therefore the total contribution of the second part is bounded by

$$\begin{aligned} & q_1^{-1/4} \sum_{m_2 \geq 1} q_1^{1/4} q_2^{m_2^2} \cdot 2 \cosh\left(\pi \frac{y_2}{k} m_2\right) \\ & + q_1^{-1/4} \sum_{m_1 \in \mathbb{N} + \frac{3}{2}} \sum_{m_2 \geq 1} q_1^{m_1^2} q_2^{m_2^2} \cdot 2 \cosh\left(\pi \left(\frac{y_1}{2} m_1^2 + \frac{y_2}{2} m_2^2\right)\right) \\ & \leq \frac{q_2^{1-\frac{1}{k}}}{1 - q_2^{3-\frac{1}{k}}} + \frac{q_2^{1+\frac{1}{k}}}{1 - q_2^{3+\frac{1}{k}}} + \frac{q_1^{7/8} q_2^{1/2}}{(1 - q_2^{3/2})(1 - q_1^2)} + \frac{q_1^{25/8} q_2^{3/2}}{(1 - q_2^{9/2})(1 - q_1^6)} \end{aligned}$$

by other applications of Lemma 4.1. \square

Lemma 4.3. Let $k \geq 1$, and let $\tau \in \mathcal{H}_2$ such that

$$y_3(\tau)^2 \leq \frac{1}{4}y_1(\tau)y_2(\tau) \quad \text{and} \quad k|y_3(\tau)| \leq y_1(\tau).$$

Define

$$\xi_{8,9}(\tau) = 2 \exp\left(i\pi \frac{z_2(\tau)}{4}\right),$$

and

$$\begin{aligned} \rho_{8,9}^{(k)}(q_1, q_2) &= \frac{q_2^2}{1 - q_2^4} + \frac{q_1^{1 - \frac{1}{k}}}{1 - q_1^{3 - \frac{1}{k}}} + \frac{q_1^{1 + \frac{1}{k}}}{1 - q_1^{3 + \frac{1}{k}}} \\ &\quad + \frac{q_2^{7/8} q_1^{1/2}}{(1 - q_1^{3/2})(1 - q_2^2)} + \frac{q_2^{25/8} q_1^{3/2}}{(1 - q_1^{9/2})(1 - q_2^6)}. \end{aligned}$$

Then for $j \in \{8, 9\}$, we have

$$\left| \frac{\theta_j(\tau)}{\xi_{8,9}(\tau)} - 1 \right| \leq \rho_{8,9}^{(k)}(q_1(\tau), q_2(\tau)).$$

Proof. We proceed in a similar fashion as in the proof of Lemma 4.2 by switching the roles of q_1 and q_2 . \square

Lemma 4.4. Let $\tau \in \mathcal{H}_2$ such that

$$y_3(\tau)^2 \leq \frac{1}{4}y_1(\tau)y_2(\tau).$$

Define

$$\begin{aligned} \xi_0(\tau) &= 1 + 2 \exp(i\pi z_1(\tau)) + 2 \exp(i\pi z_2(\tau)), \\ \xi_{0,2}(\tau) &= 1 + 2 \exp(i\pi z_1(\tau)), \\ \xi_{0,1}(\tau) &= 1 + 2 \exp(i\pi z_2(\tau)), \end{aligned}$$

and

$$\rho_0(q_1, q_2) = \frac{2q_1^4}{1 - q_1^5} + \frac{2q_2^4}{1 - q_2^5} + \frac{2q_1^{1/2} q_2^{1/2}}{(1 - q_1^{3/2})(1 - q_2^{3/2})} + \frac{2q_1^{3/2} q_2^{3/2}}{(1 - q_1^{9/2})(1 - q_2^{9/2})}.$$

Then we have

$$\begin{aligned} |\theta_0(\tau) - \xi_0(\tau)| &\leq \rho_0(q_1(\tau), q_2(\tau)), \\ |\theta_j(\tau) - \xi_{0,2}(\tau)| &\leq \rho_0(q_1(\tau), q_2(\tau)) + 2q_2(\tau) && \text{for } j \in \{0, 2\}, \\ |\theta_j(\tau) - \xi_{0,1}(\tau)| &\leq \rho_0(q_1(\tau), q_2(\tau)) + 2q_1(\tau) && \text{for } j \in \{0, 1\}, \text{ and} \\ |\theta_j(\tau) - 1| &\leq \rho_0(q_1(\tau), q_2(\tau)) + 2q_1(\tau) + 2q_2(\tau) && \text{for } 0 \leq j \leq 3. \end{aligned}$$

Proof. We proceed again in a similar fashion as in the proof of Lemma 4.2. The terms of $\rho_0(q_1, q_2)$ are obtained by considering the following subsets of indices $m \in \mathbb{Z}^2$:

$$\left\{ \binom{m_1}{0} \mid |m_1| \geq 2 \right\}, \quad \left\{ \binom{0}{m_2} \mid |m_2| \geq 2 \right\},$$

and

$$\left\{ \binom{m_1}{m_2} \mid |m_1| \geq 1, |m_2| \geq 1 \right\}. \quad \square$$

Lemma 4.5. *Let $\tau \in \mathcal{H}_2$ such that*

$$|x_3(\tau)| \leq \frac{1}{2} \quad \text{and} \quad 2|y_3(\tau)| \leq \min\{y_1(\tau), y_2(\tau)\}.$$

Write

$$\xi_{12}(\tau) = \exp\left(i\pi \frac{z_1(\tau) + z_2(\tau)}{4}\right) \left(\exp\left(i\pi \frac{z_3(\tau)}{2}\right) + \exp\left(-i\pi \frac{z_3(\tau)}{2}\right) \right),$$

and

$$\begin{aligned} \rho_{12}(q_1, q_2) &= \frac{q_1^{3/2}}{1 - q_1^{7/2}} + \frac{q_1^{5/2}}{1 - q_1^{9/2}} + \frac{q_2^{3/2}}{1 - q_2^{7/2}} + \frac{q_2^{5/2}}{1 - q_2^{9/2}} \\ &\quad + \frac{q_1^{7/8} q_2^{7/8}}{(1 - q_1^2)(1 - q_2^2)} + \frac{q_1^{25/8} q_2^{25/8}}{(1 - q_1^6)(1 - q_2^6)}. \end{aligned}$$

Then we have

$$\left| \frac{\theta_{12}(\tau)}{\xi_{12}(\tau)} - 1 \right| \leq \rho_{12}(q_1(\tau), q_2(\tau)).$$

Proof. By (1), we have

$$\begin{aligned} \theta_{12}(\tau) &= 2 \sum_{m_1 \in \mathbb{N} + \frac{1}{2}} \sum_{m_2 \in \mathbb{N} + \frac{1}{2}} \exp(i\pi(m_1^2 z_1 + m_2^2 z_2)) \\ &\quad \cdot (\exp(2\pi i m_1 m_2 z_3) + \exp(-2\pi i m_1 m_2 z_3)). \end{aligned}$$

We leave the term corresponding to $(m_1, m_2) = (\frac{1}{2}, \frac{1}{2})$ aside, and write

$$\begin{aligned} &\left| \frac{\theta_{12}(\tau)}{2 \exp(i\pi(z_1 + z_2)/4)} - (\exp(i\pi z_3/2) + \exp(-i\pi z_3/2)) \right| \\ &\leq \sum_{\substack{(m_1, m_2) \in (\mathbb{N} + \frac{1}{2})^2 \\ (m_1, m_2) \neq (\frac{1}{2}, \frac{1}{2})}} q_1^{m_1^2 - \frac{1}{4}} q_2^{m_2^2 - \frac{1}{4}} \cdot 2 \cosh(2\pi m_1 m_2 y_3). \end{aligned}$$

Since $|x_3| \leq \frac{1}{2}$, the absolute value of the argument of $\exp(i\pi z_3/2)$ is at most $\pi/4$. Therefore,

$$|\exp(i\pi z_3/2) + \exp(-i\pi z_3/2)| \geq \exp(\pi |y_3|/2).$$

We obtain

$$\left| \frac{\theta_{12}(\tau)}{\xi_{12}(\tau)} - 1 \right| \leq \sum_{\substack{(m_1, m_2) \in (\mathbb{N} + \frac{1}{2})^2 \\ (m_1, m_2) \neq (\frac{1}{2}, \frac{1}{2})}} q_1^{m_1^2 - \frac{1}{4}} q_2^{m_2^2 - \frac{1}{4}} \cdot 2 \cosh(2\pi(m_1 m_2 - \frac{1}{4})y_3).$$

We separate the terms corresponding to $m_2 = \frac{1}{2}$. Since $2|y_3| \leq y_1$, their contribution is bounded by

$$\sum_{m_1 \in \mathbb{N} + \frac{3}{2}} \left(q_1^{m_1^2 - \frac{1}{2}m_1} + q_1^{m_1^2 + \frac{1}{2}m_1 - \frac{1}{2}} \right) \leq \frac{q_1^{3/2}}{1 - q_1^{7/2}} + \frac{q_1^{5/2}}{1 - q_1^{9/2}}.$$

Similarly, the contribution from the terms with $m_1 = 1/2$ is bounded by

$$\frac{q_2^{3/2}}{1 - q_2^{7/2}} + \frac{q_2^{5/2}}{1 - q_2^{9/2}}.$$

For the remaining terms, we use the majoration

$$|2\pi(m_1 m_2 - \frac{1}{4})y_3| \leq |2\pi m_1 m_2 y_3| \leq \pi |y_3| (m_1^2 + m_2^2) \leq \frac{\pi}{2} (m_1^2 y_1 + m_2^2 y_2).$$

Thus, the rest of the sum is bounded by

$$\begin{aligned} & \sum_{m_1, m_2 \in \mathbb{N} + \frac{3}{2}} q_1^{m_1^2 - \frac{1}{4}} q_2^{m_2^2 - \frac{1}{4}} \cdot 2 \cosh\left(\frac{\pi}{2}(m_1^2 y_1 + m_2^2 y_2)\right) \\ & \leq \sum_{m_1, m_2 \in \mathbb{N} + \frac{3}{2}} \left(q_1^{\frac{1}{2}m_1^2 - \frac{1}{4}} q_2^{\frac{1}{2}m_2^2 - \frac{1}{4}} + q_1^{\frac{3}{2}m_1^2 - \frac{1}{4}} q_2^{\frac{3}{2}m_2^2 - \frac{1}{4}} \right) \\ & \leq \frac{q_1^{7/8} q_2^{7/8}}{(1 - q_1^2)(1 - q_2^2)} + \frac{q_1^{25/8} q_2^{25/8}}{(1 - q_1^6)(1 - q_2^6)}. \quad \square \end{aligned}$$

We give another version of these estimates that we will use for $\tau_4^{(n)}$.

Lemma 4.6. *Let $k \geq 2$, and let $\tau \in \mathcal{H}_2$ such that*

$$y_3(\tau)^2 \leq \frac{3}{7} y_1(\tau) y_2(\tau) \quad \text{and} \quad k |y_3(\tau)| \leq y_1(\tau).$$

Let $\alpha = \sqrt{3/7}$. Define

$$\begin{aligned} \rho_{0,1}^{(k)}(q_1, q_2) &= \frac{2q_2^4}{1 - q_2^5} + \frac{2q_1}{1 - q_1^3} + \frac{2q_1^{1 - \frac{2}{k}} q_2}{1 - q_1^{3 - \frac{2}{k}}} + \frac{2q_1^{1 + \frac{2}{k}} q_2}{1 - q_1^{3 + \frac{2}{k}}} \\ &+ \frac{2q_1^{1 - \alpha} q_2^{4(1 - \alpha)}}{(1 - q_1^{3(1 - \alpha)})(1 - q_2^{5(1 - \alpha)})} + \frac{2q_1^{1 + \alpha} q_2^{4(1 + \alpha)}}{(1 - q_1^{3(1 + \alpha)})(1 - q_2^{5(1 + \alpha)})} \end{aligned}$$

and

$$\begin{aligned} \rho'_{8,9}(q_1, q_2) &= \frac{q_2^2}{1 - q_2^4} + \frac{q_1^{1-\frac{1}{k}}}{1 - q_1^{3-\frac{1}{k}}} + \frac{q_1^{1+\frac{1}{k}}}{1 - q_1^{3+\frac{1}{k}}} \\ &+ \frac{q_2^{2-\frac{3}{4}\alpha} q_1^{1-\alpha}}{(1 - q_2^{4(1-\alpha)})(1 - q_1^{3(1-\alpha)})} + \frac{q_2^{2+\frac{3}{4}\alpha} q_1^{1+\alpha}}{(1 - q_2^{4(1+\alpha)})(1 - q_1^{3(1+\alpha)})}. \end{aligned}$$

Then we have

$$|\theta_j(\tau) - \xi_{0,1}(\tau)| \leq \rho'_{0,1}(\tau) \quad \text{for } j \in \{0, 1\}$$

and

$$\left| \frac{\theta_j(\tau)}{\xi_{8,9}(\tau)} - 1 \right| \leq \rho'_{8,9}(\tau) \quad \text{for } j \in \{8, 9\}.$$

Proof. We bound the cross-product terms by

$$\begin{aligned} |2y_3 m_1 m_2| &\leq \alpha y_1 m_1^2 + \alpha y_2 m_2^2, \\ |2y_3 m_1 m_2| &\leq \frac{1}{k} y_1 m_1 && \text{if } m_2 = \frac{1}{2}, \quad \text{and} \\ |2y_3 m_1 m_2| &\leq \frac{2}{k} y_1 m_1 && \text{if } m_2 = 1. \end{aligned}$$

For $j \in \{0, 1\}$, we separate the terms with $|m_2| \leq 1$ or $m_1 = 0$, and obtain

$$\begin{aligned} |\theta_j(\tau) - \xi_{0,1}(\tau)| &\leq 2 \sum_{m_2 \geq 2} q_2^{m_2^2} + 2 \sum_{m_1 \geq 1} q_1^{m_1^2} + 2 \sum_{m_2 \geq 1} q_2 (q_1^{m_1^2 - \frac{2}{k} m_1} + q_1^{m_1^2 + \frac{2}{k} m_1}) \\ &+ 2 \sum_{m_1 \geq 1} \sum_{m_2 \geq 2} q_1^{m_1^2} q_2^{m_2^2} \cdot 2 \cosh(\alpha(y_1 m_1^2 + y_2 m_2^2)) \\ &\leq \frac{2q_2^4}{1 - q_2^5} + \frac{2q_1}{1 - q_1^3} + \frac{2q_1^{1-\frac{2}{k}} q_2}{1 - q_1^{3-\frac{2}{k}}} + \frac{2q_1^{1+\frac{2}{k}} q_2}{1 - q_1^{3+\frac{2}{k}}} \\ &+ \frac{2q_1^{1-\alpha} q_2^{4(1-\alpha)}}{(1 - q_1^{3(1-\alpha)})(1 - q_2^{5(1-\alpha)})} + \frac{2q_1^{1+\alpha} q_2^{4(1+\alpha)}}{(1 - q_1^{3(1+\alpha)})(1 - q_2^{5(1+\alpha)})}. \end{aligned}$$

For $j \in \{8, 9\}$, we separate the terms with $|m_2| = \frac{1}{2}$ or $m_1 = 0$. We obtain

$$\begin{aligned} \left| \frac{\theta_j(\tau)}{\xi_{8,9}(\tau)} - 1 \right| &\leq q_2^{-1/4} \sum_{m_2 \in \mathbb{N} + \frac{3}{2}} q_2^{m_2^2} + \sum_{m_1 \geq 1} (q_1^{m_1^2 - \frac{1}{k} m_1} + q_1^{m_1^2 + \frac{1}{k} m_1}) \\ &+ q_2^{-1/4} \sum_{m_2 \in \mathbb{N} + \frac{3}{2}} \sum_{m_1 \geq 1} q_2^{m_2^2} q_1^{m_1^2} \cdot 2 \cosh(\alpha(y_1 m_1^2 + y_2 m_2^2)) \\ &\leq \frac{q_2^2}{1 - q_2^4} + \frac{q_1^{1-\frac{1}{k}}}{1 - q_1^{3-\frac{1}{k}}} + \frac{q_1^{1+\frac{1}{k}}}{1 - q_1^{3+\frac{1}{k}}} \\ &+ \frac{q_2^{2-\frac{3}{4}\alpha} q_1^{1-\alpha}}{(1 - q_2^{4(1-\alpha)})(1 - q_1^{3(1-\alpha)})} + \frac{q_2^{2+\frac{3}{4}\alpha} q_1^{1+\alpha}}{(1 - q_2^{4(1+\alpha)})(1 - q_1^{3(1+\alpha)})}. \quad \square \end{aligned}$$

Finally, when n is large, we will show that the theta constants $\theta_j(2^n \gamma_k \tau)$ for $0 \leq j \leq 3$ are in good position using the following lemma. Recall the definition of $r(\tau)$ and $\lambda_1(\tau)$ from §2.

Lemma 4.7. *Let $\tau \in \mathcal{H}_2$.*

1. *If $r(\tau) \geq 0.4$, then the $\theta_j(\tau)$ for $0 \leq j \leq 3$ are in good position.*
2. *If $\lambda_1(\tau) \geq 0.6$, then the $\theta_j(\tau)$ for $0 \leq j \leq 3$ are in good position.*

Proof. 1. Write

$$q = \exp(-\pi r(\tau)).$$

For $0 \leq j \leq 3$, we have

$$\begin{aligned} |\theta_j(\tau) - 1| &\leq 4q^2 + \sum_{n \in \mathbb{Z}^2, \|n\|^2 \geq 2} \exp(-\pi \lambda_1(\tau) \|n\|^2) \\ &\leq 8q^2 + 4q^4 + 8q^5 + 4q^8 + 4 \frac{1+q}{(1-q)^2} q^9. \end{aligned} \quad (6)$$

In this inequality, the first term $4q^2$ comes from the four vectors $n \in \mathbb{Z}^2$ with $\|n\| = 1$. Then we separate the terms $n = \begin{pmatrix} n_1 \\ n_2 \end{pmatrix}$ such that $|n_1| \geq 3$ and $|n_2| \geq 3$; this accounts for the term $4q^9(1+q)/(1-q)^2$, as in the proof of [4, Prop. 6.1]. We leave the remaining terms as they are.

If $q \leq 0.287$, then the quantity on the right hand side of (6) is less than $\sqrt{2}/2$, and the $\theta_j(\tau)$ are contained in a disk which is itself contained in a quarter plane. We have $q \leq 0.287$ when $r(\tau) \geq 0.4$.

2. Write

$$q = \exp(-\pi \lambda_1(\tau)).$$

Then for $0 \leq j \leq 3$, we have

$$|\theta_j(\tau) - 1| \leq 4q + 4q^2 + 4q^4 + 8q^5 + 4q^8 + 4 \frac{1+q}{(1-q)^2} q^9.$$

This quantity is less than $\sqrt{2}/2$ when $\lambda_1(\tau) \geq 0.6$. □

We conclude this section with lower bounds on r or λ_1 at $\gamma_k \tau$ for $\tau \in \mathcal{F}'$ and $1 \leq k \leq 3$.

Lemma 4.8. *For every $\tau \in \mathcal{F}'$, we have*

$$r(\gamma_1 \tau) \geq \frac{9y_1(\tau)}{34|z_1(\tau)|^2}, \quad r(\gamma_2 \tau) \geq \frac{9y_2(\tau)}{34|z_2(\tau)|^2}, \quad \text{and} \quad \lambda_1(\gamma_3 \tau) \geq \frac{9}{44y_2(\tau)}.$$

Proof. We have

$$\text{Im}(\gamma_1 \tau) = \begin{pmatrix} z_1 & z_3 \\ 0 & -1 \end{pmatrix}^{-t} \text{Im}(\tau) \begin{pmatrix} \bar{z}_1 & \bar{z}_3 \\ 0 & -1 \end{pmatrix}^{-1} = \frac{1}{|z_1|^2} \begin{pmatrix} y_1 & \alpha \\ \alpha & \beta \end{pmatrix}$$

with $\alpha = y_1 x_3 - y_3 x_1$, so $|\alpha| \leq \frac{3}{4} y_1$. Moreover,

$$\det \operatorname{Im}(\gamma_1 \tau) = \frac{1}{|z_1|^2} \det \operatorname{Im}(\tau)$$

and $\det \operatorname{Im}(\tau) \geq 9/16$, so

$$\beta \leq \frac{|z_1|^2}{y_1} \det \operatorname{Im}(\tau) + \frac{9}{16} y_1 \quad \text{and} \quad \beta \geq \frac{|z_1|^2}{y_1} \det \operatorname{Im}(\tau) \geq \frac{9}{16} y_1.$$

Therefore,

$$\lambda_1(\gamma_1 \tau) \geq \frac{\det \operatorname{Im}(\gamma_1 \tau)}{\operatorname{Tr} \operatorname{Im}(\gamma_1 \tau)} \geq \frac{y_1}{|z_1|^2} \frac{1}{1 + \frac{25}{16} \frac{y_1^2}{|z_1|^2 \det \operatorname{Im}(\tau)}} \geq \frac{9 y_1}{34 |z_1|^2}.$$

We did not use the property that $y_1 \leq y_2$, so the same proof works for $\gamma_2 \tau$. Finally, we consider $\gamma_3 \tau$. We have

$$\operatorname{Im}(\gamma_3 \tau) = \frac{1}{|\det \tau|^2} \begin{pmatrix} \beta_1 & \alpha \\ \alpha & \beta_2 \end{pmatrix}$$

with

$$\begin{aligned} \beta_1 &= y_1 |z_3|^2 + y_2 |z_1|^2 - y_3 (z_1 \bar{z}_3 + z_3 \bar{z}_1), \\ \beta_2 &= y_1 |z_2|^2 + y_2 |z_3|^2 - y_3 (z_2 \bar{z}_3 + z_3 \bar{z}_2). \end{aligned}$$

We compute

$$|\det \tau|^2 \operatorname{Tr} \operatorname{Im}(\gamma_3 \tau) = \beta_1 + \beta_2 \leq y_1 y_2^2 + y_1^2 y_2 + \frac{1}{2} (y_1 + y_2 + |y_3|) \leq \frac{11}{3} y_1 y_2^2$$

because $y_1 y_2 \geq 3/4$. Therefore,

$$\lambda_1(\gamma_3 \tau) \geq \frac{3 \det \operatorname{Im}(\tau)}{11 y_1 y_2^2} \geq \frac{9}{44 y_2}. \quad \square$$

5 Proof of the main theorem

In this final section, we prove Theorem 1.1 by separating different cases according to the value of n . If n is large enough, then Lemmas 4.7 and 4.8 are enough to conclude; if n is smaller, then we apply the theta transformation formula (Proposition 3.2) and the bounds on other theta constants given in §4.

In the proofs, we use numerical calculations, typically in order to show that a given angle $\alpha(q)$ is smaller than $\pi/2$ for certain values of q . Such calculations are easily certified using interval arithmetic, since the functions $\alpha(q)$ we consider are simple: they are either increasing or convex functions of q .

In order to help the reader visualize the estimates, we created pictures using GeoGebra [12].

Proposition 5.1. *Let $\tau \in \mathcal{F}'$. Then for every $n \geq 0$, the theta constants $\theta_j(2^n\tau)$ for $0 \leq j \leq 3$ are in good position.*

Proof. For every $n \geq 0$, we have

$$r(2^n\tau) = 2^n r(\tau) \geq \sqrt{3}/4 \geq 0.4,$$

so the result follows from Lemma 4.7. \square

Lemma 5.2. *Let $\tau \in \mathcal{F}'$.*

1. *For every $n \geq 0$ such that $2^n \leq 8.77y_1(\tau)$, the theta constants $\theta_j(\tau_1^{(n)})$ for $j \in \{0, 2, 4, 6\}$ are in good position.*
2. *For every $n \geq 0$ such that $2^n \leq 8.77y_2(\tau)$, the theta constants $\theta_j(\tau_2^{(n)})$ for $j \in \{0, 1, 8, 9\}$ are in good position.*

Proof. We only prove the first statement, the second one being symmetric. We separate three cases: $n = 0$, $n = 1$, and $n \geq 2$.

Case 1: $n = 0$. Then $\tau_1^{(n)} = \tau$. By [24, Prop. 7.7], we have

$$\begin{aligned} |\theta_j(\tau) - 1| &\leq 0.405 && \text{for } j \in \{0, 1, 2, 3\}, \text{ and} \\ \left| \frac{\theta_j(\tau)}{\xi_{4,6}(\tau)} - 1 \right| &\leq 0.348 && \text{for } j \in \{4, 6\}. \end{aligned}$$

The absolute value of the argument of $\xi_{4,6}(\tau)$ is at most $\pi/8$. Therefore the angle between any two $\theta_j(\tau)$ for $j \in \{0, 1, 2, 3, 4, 6\}$ is at most

$$\frac{\pi}{8} + \arcsin(0.348) + \arcsin(0.405) < \frac{\pi}{2}.$$

Case 2: $n = 1$. We study the relative positions of $\xi_{0,2}$ and $\xi_{4,6}$ at $\tau_1^{(1)}$. Since $|2^{-n}x_1(\tau)| \leq 1/4$, the absolute value of the argument of $\xi_{4,6}(\tau_1^{(1)})$ is bounded above by $\pi/16$. Moreover,

$$|\xi_{0,2}(\tau_1^{(1)})| \geq 1, \quad |\arg(\xi_{0,2}(\tau_1^{(1)}))| \leq \arctan\left(\frac{2q_1 \sin(\pi/4)}{1 + 2q_1 \cos(\pi/4)}\right),$$

and the arguments of $\xi_{0,2}$ and $\xi_{4,6}$ have the same sign. Therefore the angle between any two $\theta_j(\tau_1^{(1)})$ for $j \in \{0, 2, 4, 6\}$ is at most

$$\max\left\{\frac{\pi}{16}, \arctan\left(\frac{2q_1 \sin(\pi/4)}{1 + 2q_1 \cos(\pi/4)}\right)\right\} + \arcsin \rho_{4,6}^{(4)}(q_1, q_2) + \arcsin(\rho_0(q_1, q_2) + 2q_2)$$

by Lemmas 4.2 and 4.4. This quantity is less than $\pi/2$ because

$$q_2(\tau_1^{(1)}) \leq \exp(-\pi\sqrt{3}) \quad \text{and} \quad q_1(\tau_1^{(1)}) \leq \exp(-\pi\sqrt{3}/8).$$

Case 3: $n \geq 2$. We proceed as in Case 2, but we now have


$$q_2(\tau_1^{(n)}) \leq \exp(-2\pi\sqrt{3}), \quad 8|y_3(\tau_1^{(n)})| \leq y_2(\tau_1^{(n)}), \quad \text{and} \quad |x_1(\tau_1^{(n)})| \leq \frac{1}{8}.$$

Therefore the angle between the $\theta_j(\tau_1^{(n)})$ for $j \in \{0, 2, 4, 6\}$ is bounded by

$$\begin{aligned} & \max\left\{\frac{\pi}{32}, \arctan\left(\frac{2q_1 \sin(\pi/8)}{1 + 2q_1 \cos(\pi/8)}\right)\right\} \\ & + \arcsin(\rho_0(q_1, \exp(-2\pi\sqrt{3})) + 2 \exp(-2\pi\sqrt{3})) \\ & + \arcsin \rho_{4,6}^{(8)}(q_1, \exp(-2\pi\sqrt{3})). \end{aligned} \quad (7)$$

This angle remains less than $\pi/2$ when $q_1(\tau_1^{(n)}) \leq 0.699$. This is the case when $2^n \geq 8.77y_1(\tau)$. \square

The geometric situation in Case 3 of Lemma 5.2 can be represented as follows.



In this picture, we take $q_1 = 0.699$, and represent two complex numbers $\xi_{0,2}$ and $\xi_{4,6}$ with modulus one, separated by an angle of

$$\max\left\{\frac{\pi}{32}, \arctan\left(\frac{2q_1 \sin(\pi/8)}{1 + 2q_1 \cos(\pi/8)}\right)\right\} \simeq 0.22.$$

Then we draw disks centered in $\xi_{0,2}$ and $\xi_{4,6}$ with radii $\rho_0(q_1, \exp(-2\pi\sqrt{3}))$ and $\rho_{4,6}^{(8)}(q_1, \exp(-2\pi\sqrt{3}))$ respectively. Finally we represent the smallest angular sector seen from the origin containing these two disks. The green angle is equal to the quantity (7), and is indeed smaller than $\pi/2$.

Proposition 5.3. *Let $\tau \in \mathcal{F}'$.*

1. For every $n \geq 0$, the theta constants $(\theta_j(2^n \gamma_1 \tau))_{0 \leq j \leq 3}$ are in good position.
2. For every $n \geq 0$, the theta constants $(\theta_j(2^n \gamma_2 \tau))_{0 \leq j \leq 3}$ are in good position.

Proof. By Lemma 4.8, we have

$$r(\gamma_1 \tau) \geq \frac{9 y_1}{34 |z_1|^2} \geq \frac{9 y_1}{34(1/4 + y_1^2)} \geq \frac{0.205}{y_1(\tau)}$$

because $y_1(\tau) \geq \sqrt{3}/2$. By Lemma 4.7, the $\theta_j(2^n \gamma_1 \tau)$ for $0 \leq j \leq 3$ are in good position when $2^n r(\gamma_1 \tau) \geq 0.4$. This is the case when $2^n \geq 1.96 y_1$. On the other hand, Lemma 5.2 applies when $2^n \leq 8.77 y_1$. The second statement is proved in the same way. \square

Lemma 5.4. *Let $\tau \in \mathcal{F}'$. Then, for every $n \geq 0$ such that $2^n \leq 1.66 y_1$, the theta constants $\theta_j(\tau_3^{(n)})$ for $j \in \{0, 4, 8, 12\}$ are in good position.*

Proof. Write $q = q_1(\tau_3^{(n)})$ for short. We separate two cases: $n \geq 1$, and $n = 0$.

Case 1: $n \geq 1$. In this case, we have

$$|x_j(\tau_3^{(n)})| \leq 1/4 \quad \text{for each } 1 \leq j \leq 3.$$

Therefore, given the expressions of ξ_0 , $\xi_{4,6}$, $\xi_{8,9}$ and ξ_{12} (see (4)), and by Lemmas 4.2 to 4.5,

- The angle between $\theta_4(\tau_3^{(n)})$ and $\theta_8(\tau_3^{(n)})$ is bounded by

$$\frac{\pi}{8} + 2 \arcsin \rho_{4,6}^{(2)}(q, q).$$

- The angle between $\theta_4(\tau_3^{(n)})$ (or θ_8) and $\theta_0(\tau_3^{(n)})$ is bounded by

$$\frac{\pi}{16} + \arcsin \rho_{4,6}^{(2)}(q, q) + 2q \sin(\pi/4) + \arcsin \rho_0(q, q).$$

- The angle between $\theta_{12}(\tau_3^{(n)})$ and $\theta_4(\tau_3^{(n)})$ (or θ_8) is bounded by

$$\frac{3\pi}{16} + \arcsin \rho_{12}(q, q) + \arcsin \rho_{4,6}^{(2)}(q, q).$$

- The angle between $\theta_{12}(\tau_3^{(n)})$ and $\theta_0(\tau_3^{(n)})$ is bounded by

$$\frac{\pi}{4} + \arcsin \rho_{12}(q, q) + \arcsin \rho_0(q, q).$$

All these quantities remain less than $\pi/2$ when $q \leq 0.151$. This is the case when $2^n \leq 1.66 y_1$.

Case 2: $n = 0$. In this case, we have $q \leq \exp(-\pi\sqrt{3}/2)$. Therefore,

- The angle between θ_4 and θ_8 is bounded by

$$\frac{\pi}{4} + 2 \arcsin \rho_{4,6}^{(2)}(q, q) < \frac{\pi}{2}.$$


- The angle between θ_4 (or θ_8) and θ_0 is bounded by

$$\frac{\pi}{8} + \arcsin \rho_{4,6}^{(2)}(q, q) + \arcsin(\rho_0(q, q) + 4q) < \frac{\pi}{2}.$$

- The angle between θ_{12} and θ_4 (or θ_8) is bounded by

$$\frac{3\pi}{8} + \arcsin \rho_{12}(q, q) + \arcsin \rho_{4,6}^{(2)}(q, q) < \frac{\pi}{2}.$$

These estimations can be represented as follows, with similar conventions as in the picture after Lemma 5.2:



We finally study the angle between θ_{12} and θ_0 . The argument of $\xi_{12}(\tau)$ is $x_1/4 + x_2/4 + \beta$ with $\beta = \arg(\exp(i\pi z_3/2) + \exp(-i\pi z_3/2))$. Up to conjugating, we may assume that $y_3 \geq 0$ and $x_3 \geq 0$. Then

$$\exp(i\pi z_3/2) + \exp(-i\pi z_3/2) = \exp(-i\pi z_3/2)(1 + \exp(i\pi z_3))$$

so

$$\beta + \frac{\pi x_3}{2} \geq \arctan\left(\frac{q_3 \sin(\pi x_3)}{1 + q_3}\right) \geq \arctan\left(\frac{2x_3 q_3}{1 + q_3}\right).$$

In general, we have

$$|\beta| \leq \frac{\pi}{4} - \arctan\left(\frac{q^{1/2}}{1 + q^{1/2}}\right).$$

On the other hand,

$$1 \leq \operatorname{Re}(\xi_0(\tau)) \leq 1 + 4q, \quad \operatorname{Im}(\xi_0(\tau)) = 2q_1 \sin(\pi x_1) + 2q_2 \sin(\pi x_2).$$

We discuss two cases according to the signs of x_1 and x_2 :

- If x_1 and x_2 have opposite signs, then the angle between θ_{12} and θ_0 is at most

$$\frac{3\pi}{8} + \arctan(2q) + \arcsin \rho_{12}(q, q) + \arcsin \rho_0(q, q).$$

- If x_1 and x_2 have the same sign, say positive, then

$$\frac{x_1 + x_2}{4} - \arg \xi_0(\tau) \leq \frac{x_1 + x_2}{4}.$$

Therefore the angle between θ_{12} and θ_0 is at most

$$\frac{\pi}{2} - \arctan\left(\frac{q^{1/2}}{1 + q^{1/2}}\right) + \arcsin \rho_{12}(q, q) + \arcsin \rho_0(q, q).$$

This function of q is not increasing, but it is convex.

A numerical investigation shows that both quantities remain less than $\pi/2$ when $q \leq \exp(\sqrt{3}/2)$. \square

Lemma 5.5. *Let $\tau \in \mathcal{F}'$, and let $n_0 \in \mathbb{N}$ such that $2^{n_0} > 1.66y_1$. Then, for every $n \geq n_0$ such that $2^n \leq 4.2y_2(\tau)$, the theta constants $\theta_j(\tau_4^{(n)})$ for $j \in \{0, 1, 8, 9\}$ are in good position.*

Proof. By assumption, we have $y_1(\tau_4^{(n)}) \geq \frac{3}{4} \cdot 1.66 \geq 1.24$, so $q_1(\tau_4^{(n)}) \leq 0.021$. Moreover we must have $n \geq 1$, so by Lemma 3.4, $|x_2(\tau_4^{(n)})| \leq 9/16$, and

$$|y_3(\tau_4^{(n)})| \leq \frac{3}{8}y_1(\tau_4^{(n)}).$$

Therefore, we can apply Lemma 4.6 with $k = 8/3$: we have

$$\begin{aligned} |\theta_j(\tau_4^{(n)}) - \xi_{0,1}(\tau_4^{(n)})| &\leq \rho_{0,1}^{(8/3)}(0.021, q_2(\tau_4^{(n)})) \quad \text{for } j \in \{0, 1\}, \\ \left| \frac{\theta_j(\tau_4^{(n)})}{\xi_{8,9}(\tau_4^{(n)})} - 1 \right| &\leq \rho_{8,9}^{(8/3)}(0.021, q_2(\tau_4^{(n)})) \quad \text{for } j \in \{8, 9\}. \end{aligned}$$

Let us investigate the difference between the arguments of $\xi_{8,9}(\tau_4^{(n)})$ and $\xi_{0,1}(\tau_4^{(n)})$. Both have the sign of $x_2(\tau_4^{(n)})$, which we may assume to be positive. If the argument of $\xi_{8,9}$ is the largest, then the difference is bounded by

$$\arg \xi_{8,9}(\tau_4^{(n)}) \leq \frac{9\pi}{64}.$$

If the argument of $\xi_{0,1}$ is the largest, we distinguish two cases. If $x_2(\tau_4^{(n)}) \geq \frac{3\pi}{8}$, then

$$\arg \xi_{0,1}(\tau_4^{(n)}) - \arg \xi_{8,9}(\tau_4^{(n)}) \leq \arctan\left(\frac{2q_2}{1 + 2q_2 \cos(9\pi/16)}\right) - \frac{3\pi}{32}.$$

On the other hand, if $x_2(\tau) \leq 3\pi/8$, then

$$\arg \xi_{0,1}(\tau_4^{(n)}) - \arg \xi_{8,9}(\tau_4^{(n)}) \leq \arg \xi_{0,1}(\tau_4^{(n)}) \leq \arctan\left(\frac{2q_2 \sin(3\pi/8)}{1 + 2q_2 \cos(3\pi/8)}\right)$$

Note that $|\xi_{0,1}(\tau_4^{(n)})|$ is always greater than $\cos(\pi/16)$. Therefore the angle between the $\theta_j(\tau_4^{(n)})$ for $j \in \{0, 1, 8, 9\}$ is at most

$$\begin{aligned} & \max \left\{ \frac{9\pi}{64}, \arctan\left(\frac{2q_2}{1 + 2q_2 \cos(9\pi/16)}\right) - \frac{3\pi}{32}, \arctan\left(\frac{2q_2 \sin(3\pi/8)}{1 + 2q_2 \cos(3\pi/8)}\right) \right\} \\ & + \arcsin \rho_{8,9}'^{(8/3)}(0.021, q_2) + \arcsin \frac{\rho_{0,1}'^{(8/3)}(0.021, q_2)}{\cos(\pi/16)}. \end{aligned}$$

This quantity is less than $\pi/2$ when $q_2(\tau_4^{(n)}) \leq 0.38$. Since $y_2(\tau_4^{(n)}) \geq \frac{3}{2^{n+2}}y_2(\tau)$ by Lemma 3.4, this is the case when $2^n \leq 2.43y_2(\tau)$.

On the other hand, if $2^n > 2.43y_2(\tau)$, then we must have $n \geq 2$. Moreover,

$$y_1(\tau_4^{(n)}) > 2.43 \frac{y_1(\tau)y_2(\tau)}{|z_1(\tau)^2|} > 1.82,$$

so $q_1(\tau_4^{(n)}) < 0.0033$. Then, the angle bound improves to

$$\begin{aligned} & \max \left\{ \frac{9\pi}{128}, \arctan\left(\frac{2q_2 \sin(9\pi/32)}{1 + 2q_2 \cos(9\pi/32)}\right) \right\} \\ & + \arcsin \rho_{8,9}'^{(16/3)}(0.0033, q_2) + \arcsin \rho_{0,1}'^{(16/3)}(0.0033, q_2). \end{aligned}$$

This quantity is less than $\pi/2$ when $q_2(\tau_4^{(n)}) \leq 0.571$, and the latter inequality holds when $2^n \leq 4.2y_2(\tau)$. \square

Proposition 5.6. *Let $\tau \in \mathcal{F}'$. Then, for every $n \geq 0$, the theta constants $\theta_j(2^n \gamma_3 \tau)$ for $0 \leq j \leq 3$ are in good position.*

Proof. By Lemma 4.8, we have

$$\lambda_1(\gamma_3 \tau) \geq \frac{9}{44y_2(\tau)}.$$

Therefore, by Lemma 4.7, the theta constants are in good position as soon as

$$2^n \frac{9}{44y_2(\tau)} \geq 0.6, \quad \text{or} \quad 2^n \geq 2.94y_2(\tau).$$

When n is smaller, we use the transformation formulas. Lemma 5.4 applies when $2^n \leq 1.66y_1(\tau)$, and Lemma 5.5 applies when $1.66y_1(\tau) < 2^n \leq 4.2y_2(\tau)$. \square

Propositions 5.1, 5.3 and 5.6 together imply Theorem 1.1.

References

- [1] C. W. Borchardt. Theorie des arithmetisch-geometrisches Mittels aus vier Elementen. In *Gesammelte Werke*, pages 373–431. Reimer, Berlin, 1888.
- [2] J.-B. Bost and J.-F. Mestre. Moyenne arithmético-géométrique et périodes de courbes de genre 1 et 2. *Gaz. Math.*, 38:36–64, 1988.
- [3] D. A. Cox. The arithmetic-geometric mean of Gauss. *Enseign. Math.*, 30:275–330, 1984.
- [4] R. Dupont. *Moyenne arithmético-géométrique, suites de Borchardt et applications*. PhD thesis, École polytechnique, 2006.
- [5] R. Dupont. Fast evaluation of modular functions using Newton iterations and the AGM. *Math. Comp.*, 80(275):1823–1847, 2011.
- [6] A. Enge. The complexity of class polynomial computation via floating point approximations. *Math. Comp.*, 78(266):1089–1107, 2009.
- [7] A. Enge. Computing modular polynomials in quasi-linear time. *Math. Comp.*, 78(267):1809–1824, 2009.
- [8] A. Enge and E. Thomé. Computing class polynomials for abelian surfaces. *Exp. Math.*, 23:129–145, 2014.
- [9] E. Freitag and R. Salvati Manni. On the variety associated to the ring of theta constants in genus 3. *Amer. J. Math.*, 141(3):705–732, 2019.
- [10] C. F. Gauss. *Werke*. Dietrich, Göttingen, 1868.
- [11] P. Habegger and F. Pazuki. Bad reduction of genus 2 curves with CM Jacobian varieties. *Compos. Math.*, 153(12):2534–2576, 2017.
- [12] M. Hohenwarter, M. Borchers, G. Ancsin, B. Bencze, M. Blossier, J. Éliás, K. Frank, L. Gál, A. Hofstätter, F. Jordan, B. Karacsony, Z. Konečný, Z. Kovács, W. Küllinger, E. Lettner, S. Lizfelner, B. Parisse, C. Solyom-Gecse, and M. Tomaschko. GeoGebra 6.0.588.0, 2020.
- [13] J.-I. Igusa. On the graded ring of theta-constants. *Amer. J. Math.*, 86(1):219–246, 1964.
- [14] J.-I. Igusa. On the graded ring of theta-constants (II). *Amer. J. Math.*, 88(1):221–236, 1966.
- [15] J.-I. Igusa. *Theta functions*. Springer-Verlag, 1972.
- [16] F. Jarvis. Higher genus arithmetic-geometric means. *Ramanujan J.*, 17(1):1–17, 2008.

- [17] H. Klingen. *Introductory lectures on Siegel modular forms*, volume 20 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 1990.
- [18] H. Labrande. Computing Jacobi's θ in quasi-linear time. *Math. Comp.*, 87:1479–1508, 2018.
- [19] H. Labrande and E. Thomé. Computing theta functions in quasi-linear time in genus 2 and above. *LMS J. Comp. Math.*, special issue: Algorithmic Number Theory Symposium (ANTS XII), 19:163–177, 2016.
- [20] E. Milio. A quasi-linear time algorithm for computing modular polynomials in dimension 2. *LMS J. Comput. Math.*, 18:603–632, 2015.
- [21] E. Milio and D. Robert. Modular polynomials on Hilbert surfaces. *J. Number Theory*, 216:403–459, 2020.
- [22] P. Molin and C. Neurohr. Computing period matrices and the Abel-Jacobi map of superelliptic curves. *Math. Comp.*, 88(316):847–888, 2019.
- [23] D. Mumford. *Tata lectures on theta. I*, volume 28 of *Progr. Math.* Birkhäuser, Boston, 1983.
- [24] M. Streng. *Complex multiplication of abelian surfaces*. PhD thesis, Universiteit Leiden, 2010.
- [25] M. Streng. Computing Igusa class polynomials. *Math. Comp.*, 83:275–309, 2014.

Jean Kieffer
Institut de Mathématiques de Bordeaux
351 cours de la Libération, 33400 Talence, France