



HAL
open science

From Steklov to Neumann via homogenisation

Alexandre Girouard, Antoine Henrot, Jean Lagacé

► **To cite this version:**

Alexandre Girouard, Antoine Henrot, Jean Lagacé. From Steklov to Neumann via homogenisation. Archive for Rational Mechanics and Analysis, 2020, 239, pp.981-1023. 10.1007/s00205-020-01588-2 . hal-02969275

HAL Id: hal-02969275

<https://hal.science/hal-02969275>

Submitted on 16 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FROM STEKLOV TO NEUMANN VIA HOMOGENISATION

ALEXANDRE GIROUARD, ANTOINE HENROT, AND JEAN LAGACÉ

ABSTRACT. We study a new link between the Steklov and Neumann eigenvalues of Euclidean domains. This is obtained through an homogenisation limit of the Steklov problem on a periodically perforated domain, converging to a family of eigenvalue problems with dynamical boundary conditions. For this problem, the spectral parameter appears both in the interior of the domain and on its boundary. This intermediary problem interpolates between Steklov and Neumann eigenvalues of the domain. As a corollary, we recover some isoperimetric type bounds for Neumann eigenvalues from known isoperimetric bounds for Steklov eigenvalues. The interpolation also leads to the construction of planar domains with first perimeter-normalized Stekov eigenvalue that is larger than any previously known example. The proofs are based on a modification of the energy method. It requires quantitative estimates for norms of harmonic functions. An intermediate step in the proof provides a homogenisation result for a transmission problem.

1. Introduction

Let $\Omega \subset \mathbb{R}^d$ be a bounded and connected domain with smooth boundary $\partial\Omega$. Consider on Ω the Neumann eigenvalue problem

$$\begin{cases} -\Delta f = \mu f & \text{in } \Omega, \\ \partial_\nu f = 0 & \text{on } \partial\Omega, \end{cases} \quad (1)$$

as well as the Steklov eigenvalue problem

$$\begin{cases} \Delta u = 0 & \text{in } \Omega, \\ \partial_\nu u = \sigma u & \text{on } \partial\Omega. \end{cases} \quad (2)$$

Here Δ is the Laplacian, and ∂_ν is the outward pointing normal derivative. Both problems consist in finding the eigenvalues μ and σ such that there exist non-trivial smooth solutions to the boundary value problems (1) and (2). For both problems, the spectra form discrete unbounded sequences

$$0 = \mu_0 < \mu_1 \leq \mu_2 \leq \dots \nearrow \infty$$

and

$$0 = \sigma_0 < \sigma_1 \leq \sigma_2 \leq \dots \nearrow \infty,$$

where each eigenvalue is repeated according to multiplicity. The corresponding eigenfunctions $\{f_k\}$ and $\{u_k\}$ have natural normalisations as orthonormal bases of $L^2(\Omega)$ and $L^2(\partial\Omega)$, respectively.

1.1. From Steklov to Neumann : heuristics. Let us start by painting with broad brushes the relationships between the Neumann and Steklov eigenvalue problems. They exhibit many similar features, and it is not a surprise that they do so. Indeed, in both cases the eigenvalues are those of a differential or pseudo-differential operator, namely the

Laplacian and the Dirichlet-to-Neumann map, whose kernels consist of constant functions. Moreover, in both cases, the natural isoperimetric type problem consists in maximizing μ_k and σ_k (instead of minimizing it as is usual for the Dirichlet problem). The relation between the two boundary value problem is not solely heuristic and incidental. Indeed, it is known from the works of Arrieta–Jiménez-Casas–Rodríguez-Bernal [2] and Lamberti–Provenzano [25, 26] that one can recover the Steklov problem as a limit of weighted Neumann problems

$$\begin{cases} -\Delta f = \mu \rho_\varepsilon f & \text{in } \Omega, \\ \partial_\nu f = 0 & \text{on } \partial\Omega, \end{cases}$$

where ρ_ε is a density function whose support converges to the boundary as $\varepsilon \rightarrow 0$. If we are to interpret the Neumann problem as finding the frequencies and modes of vibrations of a free boundary membrane, this means that the Steklov problem represents the frequencies and modes of a membrane whose mass is concentrated at the boundary.

Our primary goal in this paper is to establish a link in the reverse direction, by realizing the Neumann problem as a limit of appropriate Steklov problems. This is achieved in two steps. The first one is to accumulate uniformly distributed boundary elements inside the domain Ω . This is done by perforating the interior of the domain with small holes that are uniformly distributed. On these new boundary components, we consider the Steklov boundary conditions; this step is known as *the homogenisation process*. We assume that the ratio of the radii of the holes to the distance between them is at a Cioranescu–Murat type critical regime. Then, the eigenvalues and eigenfunctions of the Steklov problem converge to those of a *dynamical eigenvalue problem*

$$\begin{cases} -\Delta U = A_d \beta \Sigma U & \text{in } \Omega, \\ \partial_\nu U = \Sigma U & \text{on } \partial\Omega, \end{cases} \quad (3)$$

where $\beta \geq 0$ is the critical regime parameter and A_d is the area of the unit sphere in \mathbb{R}^d . Its eigenvalues form a discrete unbounded sequence:

$$\Sigma_{0,\beta} < \Sigma_{1,\beta} \leq \Sigma_{2,\beta} \leq \dots \nearrow \infty,$$


and once again the functions associated to the eigenvalue $\Sigma_{0,\beta} = 0$ are constant.

Remark 1. In [35], Joachim von Below and Gilles François studied an eigenvalue problem that is equivalent to Problem (3), which stems from a parabolic equation with dynamical boundary conditions. Indeed, they study the eigenvalue problem

$$\begin{cases} -\Delta u = \lambda u & \text{in } \Omega, \\ \partial_\nu u = \lambda \alpha u & \text{on } \partial\Omega. \end{cases} \quad (4)$$

If λ is an eigenvalue of Problem (4), then $\Sigma = \alpha^{-1} \lambda$ is an eigenvalue of Problem (3) with parameter $\beta = \frac{1}{\alpha A_d}$.

The parameter β in (3) can be interpreted as a weight on the interior of the domain, with the boundary $\partial\Omega$ having constant weight 1. In order to recover the Neumann problem, the second step will therefore be to send the parameter β to ∞ , putting all the weight inside the domain. Under an appropriate normalisation, eigenvalues and eigenfunctions of Problem (3) converge to those of Problem (1), completing the circle for the relation between the Steklov and the Neumann problems.

FIGURE 1. The domain Ω_ε

1.2. The homogenisation process. Consider a family of problems obtained by removing periodically placed balls from the domain Ω . More precisely, given $0 < \varepsilon < 1$, and $\mathbf{k} \in \mathbb{Z}^d$, define the cube

$$Q_{\mathbf{k}}^\varepsilon := \varepsilon \mathbf{k} + \left[-\frac{\varepsilon}{2}, \frac{\varepsilon}{2}\right]^d \subset \mathbb{R}^d,$$

and define the set of indices

$$I^\varepsilon := \left\{ \mathbf{k} \in \mathbb{Z}^d : Q_{\mathbf{k}}^\varepsilon \subset \Omega \right\}.$$

Let r_ε be an increasing positive function of ε with $r_\varepsilon < \varepsilon/2$. For $\mathbf{k} \in \mathbb{Z}^d$, define

$$T_{\mathbf{k}}^\varepsilon := B(\varepsilon \mathbf{k}, r_\varepsilon) \subset Q_{\mathbf{k}}^\varepsilon$$

and set

$$T^\varepsilon := \bigcup_{\mathbf{k} \in I^\varepsilon} T_{\mathbf{k}}^\varepsilon \subset \Omega.$$

Consider the family of perforated domains

$$\Omega^\varepsilon = \Omega \setminus \overline{T^\varepsilon}.$$

The Steklov eigenvalues of Ω^ε are written as $\sigma_k^\varepsilon := \sigma_k(\Omega^\varepsilon)$, and we write $\{u_k^\varepsilon\}$ for a corresponding complete sequence of eigenfunctions, normalized by

$$\int_{\partial\Omega^\varepsilon} (u_k^\varepsilon)^2 \, dA = 1.$$

Our first main result is the following critical regime homogenisation theorem for the Steklov problem.

Theorem 2. *Suppose that $r_\varepsilon^{d-1} \varepsilon^{-d} \rightarrow \beta$ for some $\beta \in [0, \infty)$, as $\varepsilon \searrow 0$. Then σ_k^ε converges to the eigenvalue $\Sigma_{k,\beta}$ of (3). The functions $U_k^\varepsilon \in H^1(\Omega)$ obtained by harmonic extension of a normalized Steklov eigenfunction u_k^ε over the holes $T^\varepsilon \subset \Omega$ form a sequence which weakly converges in $H^1(\Omega)$ to a solution U_k associated with $\Sigma_{k,\beta}$ of (3).*

Remark 3. If the eigenvalue $\Sigma_k = \Sigma_{k,\beta}$ is multiple of multiplicity m , i.e.

$$\Sigma_{k-1} < \Sigma_k = \dots = \Sigma_{k+m-1} < \Sigma_{k+m},$$

the convergence statement in the previous theorem is understood in the following sense. Given a basis U_k, \dots, U_{k+m-1} for the eigenspace associated with Σ_k , there is a family of $m \times m$ orthogonal matrices $M(\varepsilon)$ such that

$$M(\varepsilon) \begin{pmatrix} U_k^\varepsilon \\ \vdots \\ U_{k+m-1}^\varepsilon \end{pmatrix} \rightarrow \begin{pmatrix} U_k \\ \vdots \\ U_{k+m-1} \end{pmatrix}$$

as $\varepsilon \rightarrow 0$. One could also be content with the weaker statement that if the eigenvalues are multiple, the convergence statement of theorem 2 is only true up to taking a subsequence.

Remark 4. Literature on homogenisation theory is often concerned with the situation where holes are proportional to their reference cell. That is, $r_\varepsilon = c\varepsilon$ for some constant $c \in (0, 1/2)$. In this case one has $r_\varepsilon^{d-1} \varepsilon^{-d} \rightarrow \infty$. It follows from [10] that $\sigma_k^\varepsilon \rightarrow 0$. Indeed it is proved there that any bounded domain $\Omega \subset \mathbb{R}^d$ satisfies

$$\sigma_k(\Omega) |\partial\Omega|^{\frac{1}{d-1}} \leq C_{d,k},$$

where the number $C_{d,k} > 0$ depend only on the dimension d and index k . The hypothesis that $r_\varepsilon^{d-1} \varepsilon^{-d} \rightarrow \infty$ implies that $|\partial\Omega^\varepsilon| \rightarrow \infty$, which forces $\sigma_k^\varepsilon \rightarrow 0$, as claimed. Note that this also corresponds to the homogenisation regime which was studied by Vanninathan in [34] for a slightly different problem, for which the Dirichlet boundary condition was imposed on $\partial\Omega$ and the Steklov condition on ∂T^ε .

The regime that we consider in Theorem 2 is the critical regime for the Steklov problem, where we observe a change of behaviour in the limiting problem. This is akin to the situation studied by Rauch–Taylor [31] and Cioranescu–Murat [9].

1.3. Convergence to the Neumann problem and spectral comparison theorems. The β parameter in Problem (3) can be interpreted as a relative weight between the interior of Ω and its boundary $\partial\Omega$ for the behaviour of that problem, see Section 2.2 for details on this interpretation. Our second main result is the following theorem, describing the specific dependence on β in (3).

Theorem 5. For each $k \in \mathbb{N}$, the eigenvalue $\Sigma_{k,\beta}$ depends continuously on $\beta \in [0, \infty)$ and satisfies

$$\lim_{\beta \rightarrow \infty} A_d \beta \Sigma_{k,\beta} = \mu_k.$$

The eigenfunctions $\{U_{k,\beta}\}$ satisfy

$$\beta^{1/2} U_{k,\beta} \rightarrow f_k$$

weakly in $H^1(\Omega)$ as $\beta \rightarrow \infty$, where f_k is the k th non-trivial Neumann eigenfunction.

Remark 6. We make the second observation that this convergence cannot be uniform in k , as that would contradict [35, Theorem 4.4].

The relationships between isoperimetric type problems for the Neumann and Steklov eigenvalue problems have been investigated for the first few eigenvalues in [17, 18] from the point of view of the Robin problem. Our methods also allow us to investigate the relationship between these isoperimetric problems for every eigenvalue rank k .

The combination of Theorem 5 and Theorem 2 allows the transfer of known bounds for Steklov eigenvalues to bounds for Neumann eigenvalues. For instance, we can combine these two theorems with [10, Theorem 1.3], asserting that for bounded Euclidean domains with smooth boundary

$$\sigma_k(\Omega)|\partial\Omega|^{\frac{2}{d-1}} \leq C(d)k^{2/d}. \quad (5)$$

This leads to the following.

Corollary 7. *The Neumann eigenvalues of a bounded domain $\Omega \subset \mathbb{R}^d$ satisfy*

$$\mu_k(\Omega)|\Omega|^{2/d} \leq C(d)k^{2/d}, \quad (6)$$

where the constant $C(d)$ is exactly that of [10, Theorem 1.3].

Remark 8. *The existence of a constant depending only on the dimension in inequality (6) is already known. In fact, Kröger obtained a better constant in [24]. However, it follows from Corollary 7 that any improvement to the bound (5) will transfer to bounds on Neumann eigenvalues.*

One of the original motivation for this project was the study of the following quantity:

$$\hat{\sigma}_k^* := \sup \{ \sigma_k(\Omega)|\partial\Omega| : \Omega \subset \mathbb{R}^2 \text{ bounded with smooth boundary} \}.$$

In dimension $d = 2$, we are able to get a stronger version of Corollary 7 in the sense that we obtain a direct link between $\hat{\sigma}_k^*$ and

$$\hat{\mu}_k^* := \sup \{ \mu_k(\Omega)|\Omega| : \Omega \subset \mathbb{R}^2 \text{ bounded with smooth boundary} \}.$$

In that case, we obtain.

Theorem 9. *For $d = 2$ and every $k \in \mathbb{N}$,*

$$\hat{\mu}_k^* \leq \hat{\sigma}_k^*.$$

Remark 10. *From [15] we have that*

$$\sigma_1(\Omega)|\partial\Omega| < 4\pi.$$

It follows from Theorem 2 and Theorem 5 that

$$\mu_1(\Omega)|\Omega| \leq 4\pi.$$

Of course this bound is already known. Indeed the optimal upper bound is given by the famous Szegő–Weinberger theorem:

$$\mu_1(\mathbb{D})\pi \approx 3.39\pi.$$

One can remark, however, that 4π corresponds to the Pólya bound for the first Neumann eigenvalue. We can now also see that obtaining bounds of a similar type for σ_k would also transfer to μ_k .

The previous discussion also yields the following corollary.

Corollary 11.

$$\hat{\sigma}_1^* \geq \mu_1(\mathbb{D}) \times \pi \approx 3.39\pi.$$

Indeed, by the Szego-Weinberger inequality, we have that

$$\hat{\sigma}_1^* \geq \pi \mu_1(\mathbb{D}) \approx 3.39\pi.$$

Furthermore, this number will be approached as close as desired in homogenisation sequence of pierced unit disks with large enough parameter β . Of course, it is known

from [16] that if one is allowed to optimise amongst surfaces rather than Euclidean domains, then there is a sequence of surfaces such that $\sigma_1(\Omega_n) |\partial\Omega_n| \rightarrow 4\pi$. This leads to the natural conjecture

Conjecture.

$$\hat{\sigma}_1^* = \mu_1(\mathbb{D}) \times \pi \approx 3.39\pi.$$

Note that the previous best known lower bound for $\hat{\sigma}_1^*$ was attained on some concentric annulus, whose first normalised Steklov eigenvalue is approximately 2.17π , see [20]. We also observe that a similar analysis yields that for any $\Omega \subset \mathbb{R}^2$ bounded with smooth boundary,

$$\hat{\sigma}_k^* \geq \mu_k(\Omega) |\Omega|.$$

In particular, it follows from Weyl's law for Neumann eigenvalues that there exists a sequence $a_k \sim 4\pi k$ such that

$$\hat{\sigma}_k^* \geq a_k.$$

1.4. Discussion. Homogenisation theory is a young branch of mathematics which started around the 1960's. Its general goal is to describe macroscopic properties of materials through their microscopic structure. To the best of our knowledge, the first papers to study periodically perforated domains from a rigorous mathematical point of view are those of Marchenko and Khruslov from the early 1960's (e.g. [27]) leading to their influential book [28] in 1974. The topic became widely known in the west with the work of Rauch and Taylor on the *crushed ice problem* [31] in 1975 and then with the publication in 1982 of [9] by Cioranescu and Murat. Many of these early results were concerned with the Poisson problem $\Delta u_\varepsilon = f$ under Dirichlet boundary conditions $u_\varepsilon = 0$ on $\partial\Omega_\varepsilon$. The limiting behaviour of the solution u_ε depends on the rate at which $r_\varepsilon \searrow 0$. Three regimes are considered. If the size of the holes r_ε tends to zero very fast, then in the limit the solutions tend to those of the Poisson problem on the original domain Ω , while if the size of the holes are big enough the solutions tend to zero. The main interest comes from the critical regime, in which case the solutions tend to solutions of a new elliptic problem.

In this paper we are concerned with the much less studied Steklov spectral problem (2). From the point of view of homogenisation theory, this problem is atypical in the sense that the function spaces which occur for different values of the parameter $\varepsilon > 0$ are not naturally related. This means in particular that this problem does not yield to any of the usual general frameworks used in homogenisation theory (such as [22, Chapter 11]). Nevertheless, several authors have considered homogenisation for this problem, using ad hoc methods depending on the specific situation considered. The behaviour of Steklov eigenvalues under singular perturbations such as the perforation of a single hole has also been studied in [21, 30].

Our main inspiration for this work is the paper [34]. Several papers have also considered homogenisation in the situation where the Dirichlet condition is imposed on the outer boundary while the Steklov condition is considered only on the boundary of the holes [7, 13]. In these papers the holes are proportional to the size of the reference cell. The novelty of our homogenisation result in the case of the Steklov problem is that consider holes that are shrinking much faster than that, in a critical regime where the limiting problem is fundamentally different. They are in fact shrinking at the precise rate which makes their total surface area (or perimeter in dimension 2) is asymptotically comparable with the volume of the domain. This is similar to the work of Rauch–Taylor [31] for the Neumann problem, Cioranescu–Murat [9] for the Dirichlet problem and Kaizu for the Robin problem [23].

The *energy method* of Tartar (see [1, Section 1.3] for an exposition) has been used extensively in the study of homogenisation problems at critical regimes, see [9, 23] and more recently [6], where they obtain norm-resolvent convergence for the Dirichlet, Neumann and Robin problem. That method uses an auxiliary function, satisfying some energy-minimising PDE in the fundamental cells, in order to derive convergence of the problem in the weak formulation. The method, in its Robin or Neumann form, is boundary-condition agnostic and as such is ill-suited for the Steklov problem, where the normalisation is with respect to $L^2(\partial\Omega^\varepsilon)$. Indeed, while the technique could be used to obtain some form of convergence, it will not be able to transfer boundary estimates to $L^2(\Omega)$, and ensure that the limit solution doesn't degenerate to the trivial one.

Nevertheless, one can interpret our technique as a variation on the energy method, adapted for problems defined on the boundary. We are also using an auxiliary PDE in order to derive convergence, but it does not stem from compensated compactness. The main difference, however, is that we can deduce interior estimates from those on the boundary of the periodic holes from our auxiliary problem, see Lemma 26.

1.5. Structure of the proof and plan of the paper. In Section 2 we formally describe properties of the various eigenvalue problems that we study as well as the functions spaces over which they are defined. While they are well-known, the notation used for all of them often collides. In that section, we fix notation once and for all for the remainder of the paper for definedness and ease of references.

In Section 3 we study one of the main technical tools in this paper, properties of harmonic extensions of functions on annuli to the interior disk. Results are separated in two categories: those that rely on the fact that the functions satisfy a Robin-type boundary condition, and more general results that do not rely on such a thing. Most of our results will be obtained by considering the Fourier expansion of a function

$$u(\rho, \boldsymbol{\theta}) = \sum_{\ell, m} a_\ell^m(\rho) Y_\ell^m(\boldsymbol{\theta})$$

in spherical harmonics, and obtaining our inequalities term by term for every a_ℓ^m .

Section 4 is the *pièce de résistance* of this paper. It is where we show Theorem 2 and the proof proceeds in many steps. We first prove that the family of harmonic extensions $U^\varepsilon := U_k^\varepsilon$ is bounded in the Sobolev space $H^1(\Omega)$ hence there exists a subsequence $\varepsilon_n \searrow 0$ such that U^{ε_n} weakly converges to a function $U \in H^1(\Omega)$. This allows us to consider properties of the weak limit U , and of an associated limit Σ to the eigenvalue sequence $\sigma_k^{\varepsilon_n}$.

It is then not so hard to show that, using the weak formulations of Problems (2) and (3) that the limit of the homogenised Steklov problem contains terms corresponding to the limit dynamical eigenvalue problem, plus some spurious terms that must be shown to converge to zero. This is done by studying two representations of the weak formulation using Green's identity either towards the inside of the holes or the inside of the domains Ω^ε . The first one is used to show that the functionals that arise in the study of the homogenisation problem are uniformly bounded, hence we can use smooth test functions. In the second representation, we are therefore allowed to use smooth test functions, which allows us to recover convergence to zero of the spurious terms. A key step in this argument is to understand the limit behaviour of an auxiliary homogenisation problem for the transmission eigenvalue problem (see Proposition 20).

Once we have established convergence to a solution of Problem 3, we end up showing that the limit eigenpair (Σ, U) does not degenerate to the trivial function. Using

variational characterisation of eigenvalues and eigenfunctions, we can also show that we get complete spectral convergence, and that subsequences are not needed.

Finally, in Section 5 we show convergence to the Neumann problem as $\beta \rightarrow \infty$. The method is similar to the one used at the end of the previous section, but many of the inequalities are more subtle. We also show the comparison theorems between Steklov and Neumann eigenvalues in this section.

In this paper, we use c and C to mean constants whose precise value is not important to our argument, and whose exact value may change from line to line. We use the notations $f = O(g)$ and $f \ll g$ interchangeably to mean that there exists a constant C such that $|f(x)| \leq Cg(x)$.

2. Notation and function spaces

Four different eigenvalue problems will be used. The goal of this section is to introduce them and fix the relevant notation. Throughout the paper, we use real valued functions.

2.1. The Steklov problem on Ω and Ω^ε . Given a bounded domain Ω whose boundary $\partial\Omega$ is smooth, the Dirichlet-to-Neumann operator (DtN map) Λ acts on $C^\infty(\partial\Omega)$ as

$$\Lambda f = \partial_\nu \widehat{f},$$

where \widehat{f} is the harmonic extension of f to the interior of Ω . The DtN map is an elliptic, positive, self-adjoint pseudodifferential operator of order 1. Because $\partial\Omega$ is compact, it follows from standard theory of such operators, see e.g [32], that the eigenvalues form a non-negative unbounded sequence $\{\sigma_k : k \in \mathbb{N}_0\}$ and that there exists an orthonormal basis of (f_k) of $L^2(\partial\Omega)$ such that $\Lambda f_k = \sigma_k f_k$. The harmonic extensions $u_k = \widehat{f}_k$ satisfy the Steklov problem

$$\begin{cases} -\Delta u_k = 0 & \text{in } \Omega, \\ \partial_\nu u_k = \sigma_k u_k & \text{on } \partial\Omega. \end{cases}$$

In general, we use the same symbol u_k for the function on Ω and for its trace on the boundary $\partial\Omega$. The eigenvalue sequence for Ω^ε is denoted $\{\sigma_k^\varepsilon : k \in \mathbb{N}_0\}$, with corresponding eigenfunctions u_k^ε . The eigenfunctions u_k , respectively u_k^ε , form an orthonormal basis with respect to the inner products

$$(f, g)_\partial := \int_{\partial\Omega} f g \, dA,$$

respectively

$$(f, g)_{\partial^\varepsilon} := \int_{\partial\Omega^\varepsilon} f g \, dA.$$

The k -th nonzero eigenvalue σ_k is characterised by

$$\sigma_k = \inf \left\{ \frac{\int_\Omega |\nabla u|^2 \, d\mathbf{x}}{\int_{\partial\Omega} u^2 \, dA} : u \in H^1(\Omega) \text{ and } (u, u_j)_\partial = 0 \text{ for } 0 \leq j < k \right\}. \quad (7)$$

The eigenvalues σ_k^ε have the same characterisation, integrating over Ω^ε and $\partial\Omega^\varepsilon$ respectively instead, and with the orthogonality being with respect to $(\cdot, \cdot)_{\partial^\varepsilon}$.

2.2. Dynamical eigenvalue problem. For $\beta \in (0, \infty)$, consider the eigenvalue problem:

$$\begin{cases} -\Delta U = A_d \beta \Sigma U & \text{in } \Omega, \\ \partial_\nu U = \Sigma U & \text{on } \partial\Omega. \end{cases} \quad (8)$$

where A_d is the area of the unit sphere in \mathbb{R}^d . Problem (8) was introduced with a slightly different normalization in [35], where it is called a *dynamical eigenvalue problem*. The eigenvalues and eigenfunctions are those of the operator

$$P := \begin{pmatrix} -(A_d \beta)^{-1} \Delta & 0 \\ 0 & \partial_\nu \end{pmatrix}.$$

This unbounded operator is defined on an appropriate domain in the space $L^2_{A_d \beta}(\Omega) \times L^2(\partial\Omega)$ which consists simply of $L^2(\Omega) \times L^2(\partial\Omega)$ equipped with the inner product defined by

$$(f, g)_\beta := A_d \beta \int_\Omega f g \, dx + \int_{\partial\Omega} f g \, dA,$$

The dynamical eigenvalue problem (8) has a discrete sequence of eigenvalues

$$0 = \Sigma_{0,\beta} < \Sigma_{1,\beta} \leq \Sigma_{2,\beta} \leq \dots \nearrow \infty.$$

Let $X \subset L^2_{A_d \beta}(\Omega) \times L^2(\partial\Omega)$ be the subspace defined by

$$X := \{U = (u, \tau u) : u \in H^1(\Omega)\},$$

where $\tau : H^1(\Omega) \rightarrow L^2(\partial\Omega)$ is the trace operator. The eigenfunctions $U_{k,\beta}$ associated to $\Sigma_{k,\beta}$ form a basis of X , but not of $L^2_{A_d \beta}(\Omega) \times L^2(\partial\Omega)$. The eigenvalues $\Sigma_{k,\beta}$ are characterised by

$$\Sigma_{k,\beta} = \inf \left\{ \frac{\int_\Omega |\nabla U|^2 \, dx}{A_d \beta \int_\Omega U^2 \, dx + \int_{\partial\Omega} U^2 \, dA} : U \in H^1(\Omega) \text{ and } (U, U_{j,\beta})_\partial = 0 \text{ for } 0 \leq j < k \right\}.$$

2.3. The Neumann eigenvalue problem. We will also make use of the classical Neumann eigenvalue problem

$$\begin{cases} -\Delta f = \mu f & \text{in } \Omega, \\ \partial_\nu f = 0 & \text{on } \partial\Omega. \end{cases}$$

The Neumann eigenvalues form an increasing sequence

$$0 = \mu_0 < \mu_1 \leq \mu_2 \leq \dots \infty,$$

with associated eigenfunctions f_k , orthonormal with respect to the $L^2(\Omega)$ inner product

$$(f, g) := \int_\Omega f g \, dx.$$

The eigenvalues are characterised by

$$\mu_k = \inf \left\{ \frac{\int_\Omega |\nabla f|^2 \, dx}{\int_\Omega f^2 \, dx} : f \in H^1(\Omega) \text{ and } (f, f_j) = 0 \text{ for } j = 0, 1, \dots, k-1 \right\}.$$

3. Comparison theorems

In this section, we derive comparison inequalities that will be used repeatedly. For $0 < r < R$, set

$$A_{r,R} = B(0, R) \setminus \overline{B(0, r)}.$$

For any set X where it is defined, the Dirichlet energy of a function $f : X \rightarrow \mathbb{R}$ is

$$\mathcal{D}(f) := \int_X |\nabla f|^2 \, dx.$$

The Dirichlet energy of f on a subset $Y \subset X$ is written $\mathcal{D}(f; Y)$. Note that since every problem under consideration is self-adjoint no generality is lost by studying real-valued functions.

3.1. Comparison theorems for functions satisfying a Steklov boundary condition.

Lemma 12. *Fix a positive real number $\sigma > 0$. For any $0 < r < R \leq 1$, let $u \in C^\infty(\overline{A_{r,R}})$ be such that*

$$\begin{cases} \Delta u = 0 & \text{in } A_{r,R}, \\ \partial_\nu u = \sigma u & \text{on } \partial B(0, r). \end{cases}$$

Consider the function $h : B(0, r) \rightarrow \mathbb{R}$ defined by

$$\begin{cases} h = u & \text{on } \partial B(0, r), \\ \Delta h = 0 & \text{in } B(0, r). \end{cases}$$

The following inequalities holds as the ratio r/R goes to 0.

I. *Dirichlet energy comparison :*

$$\mathcal{D}(h) \leq 5\mathcal{D}(u) \left(\frac{r}{R} \right)^d \left(1 + O\left(\left(\frac{r}{R} \right)^d \right) \right). \quad (9)$$

II. *L^2 comparisons at the boundary : if $R \geq cr^{\frac{d-1}{d}}$, then there is a constant C , depending only on c and on the dimension such that*

$$\|u\|_{L^2(\partial B(0,r))} \leq C \|u\|_{L^2(A_{r,R})}.$$

III. *Comparison between $L^2(B(0, r))$ and $L^2(A_{r,R})$: there is a constant C depending only on the dimension such that*

$$\|h\|_{L^2(B(0,r))}^2 \leq C \left(\frac{r}{R} \right)^{d+1} \mathcal{D}(u).$$

IV. *L^∞ bounds : Suppose that $R < c < 1$. Then, there is a constant C depending only on the dimension and on c such that for r small enough,*

$$\|u\|_{L^\infty(A_{r,R})} \leq C \|u\|_{L^2(\partial B(0,r))}.$$

Proof. For every $\ell \geq 0$, we denote by N_ℓ the dimension of the space H_ℓ of spherical harmonics of order ℓ and denote by $Y_\ell^m(\boldsymbol{\theta})$, $1 \leq m \leq N_\ell$ the standard orthonormal basis of spherical harmonics on the unit sphere. On $A_{r,R}$, the function u admits a Fourier decomposition in spherical harmonics

$$u(\rho, \boldsymbol{\theta}) = \sum_{\substack{\ell \geq 0 \\ 1 \leq m \leq N_\ell}} a_\ell^m(\rho) Y_\ell^m(\boldsymbol{\theta}).$$

We start by studying the form of the coefficients a_ℓ^m .

Case $d > 2$. The harmonicity condition on u implies that the radial parts $a_\ell^m(\rho)$ are given by

$$a_\ell^m(\rho) = c_\ell^m \rho^\ell + c_{-\ell}^m \rho^{-\ell+2-d}. \quad (10)$$

By convention the coefficients c_0^1 and c_{-0}^1 are assumed to be different, the minus sign referring as for the other coefficients to the solution blowing up at the origin. The Steklov condition for u on $\partial B(0, r)$ along with the orthogonality of the spherical harmonics Y_ℓ^m imply

$$-\partial_\rho a_\ell^m(r) = \sigma a_\ell^m(r),$$

which yields the relations

$$c_{-\ell}^m = \overbrace{\left(\frac{\ell + r\sigma}{\ell - 2 + d - r\sigma} \right)}^{M:=} r^{2\ell+d-2} c_\ell^m. \quad (11)$$

In turns this yields the following explicit expression for the radial functions

$$a_\ell^m(\rho) = c_\ell^m \rho^\ell \left(1 + M \left(\frac{r}{\rho} \right)^{2\ell+d-2} \right). \quad (12)$$

For $r \leq \frac{d-2}{2\sigma}$, it follows that

$$\frac{\ell}{\ell - 2 + d} \leq M \leq 1. \quad (13)$$

Case $d = 2$. Equations (10) holds except at $\ell = 0$, in which case,

$$a_0^1(\rho) = c_0^1 + c_{-0}^1 \log \rho.$$

In that case, (12) becomes

$$\begin{aligned} a_0^1 &= c_0^1 \left(1 + \frac{r\sigma}{1 + r \log r} \log(1/\rho) \right) \\ &= c_0^1 (1 + M \log(1/\rho)). \end{aligned}$$

Observe for the sequel that when $d = 2$ and $\ell = 0$, then $M = O(r\sigma)$, and if $\ell > 0$, then

$$M = 1 + O(r\sigma). \quad (14)$$

We first prove inequality I. Because inequality (9) is invariant under scaling, it is sufficient to prove the case $R = 1$ and to let $r \rightarrow 0$. The harmonic extension of u to $B(0, r)$ is given by

$$h(\rho, \boldsymbol{\theta}) = \sum_{\substack{\ell \geq 0 \\ 1 \leq m \leq N_\ell}} a_\ell^m(r) \frac{\rho^\ell}{r^\ell} Y_\ell^m(\boldsymbol{\theta}).$$

The Dirichlet energy of h is

$$\mathcal{D}(h) = \sum_{\substack{\ell \geq 1 \\ 1 \leq m \leq N_\ell}} \ell a_\ell^m(r)^2 r^{d-2}. \quad (15)$$

On the other hand, the Dirichlet energy of u is given, from Green's identity, and the Steklov condition on $\partial B(0, r)$

$$\mathcal{D}(u) = \sum_{\ell, m} \sigma a_\ell^m(r)^2 r^{d-1} + \sum_{\ell, m} a_\ell^m(1) \partial_\rho a_\ell^m(1). \quad (16)$$

Our goal is now to find a bound on (15) in terms of (16). It is sufficient to show that for each $\ell \geq 1$,

$$\ell a_\ell^m(r)^2 \leq 4r^2 a_\ell^m(1) \partial_\rho a_\ell^m(1) \left(1 + O\left(r^d\right)\right).$$

Suppose without loss of generality that $c_\ell^m = 1$. The substitution of (11) into (10) imply that

$$a_\ell^m(r)^2 = r^{2\ell} (1 + M)^2 \leq 5r^2, \quad (17)$$

and that

$$a_\ell^m(1) \partial_\rho a_\ell^m(1) = \ell \left(1 + \frac{(2-d)M}{\ell} r^{2\ell+d-2} + \frac{(2-d-\ell)M^2}{\ell} r^{2(2\ell+d-2)}\right) = \ell + O\left(r^d\right). \quad (18)$$

Hence, dividing (17) by (18), and using the bounds on M from (13) and (14) we have that for $\ell \geq 1$,

$$\ell a_\ell^m(r)^2 \leq 5a_\ell^m(1) \partial_\rho a_\ell^m(1) r^2 \left(1 + O\left(r^d\right)\right).$$

This proves inequality I. To prove inequality II, note that

$$\|u\|_{L^2(\partial B(0,r))} = \sum_{\ell,m} a_\ell^m(r)^2 r^{d-1}$$

while

$$\|u\|_{L^2(A_{r,R})} = \sum_{\ell,m} \int_r^R a_\ell^m(\rho)^2 \rho^{d-1} d\rho.$$

Once again, we will proceed term-wise. Supposing once again that $c_\ell^m = 1$, (17) still holds and

$$\begin{aligned} \int_r^R a_\ell^m(\rho)^2 d\rho &= \int_r^R \left(\rho^\ell + Mr^{2\ell+d-2} \rho^{2-\ell-d}\right)^2 \rho^{d-1} d\rho \\ &= \frac{\rho^{2\ell+d}}{2\ell+d} + Mr^{2\ell+d-2} \rho^2 + \frac{M^2 r^{2(2\ell+d-2)} \rho^{-2\ell-d+4}}{4-2\ell-d} \Big|_{\rho=r}^R, \\ &\geq \frac{R^{2\ell+d}}{2\ell+d} + O\left(r^{2\ell+d}\right), \end{aligned}$$

with the last term in the second line replaced by $M^2 r^{2(2\ell+d-2)} \log \rho$ if $2\ell+d = 4$. It follows from $R \geq cr^{\frac{d-1}{d}}$ that

$$\frac{\|a_\ell^m Y_\ell^m\|_{L^2(A_{r,R})}}{\|a_\ell^m Y_\ell^m\|_{L^2(\partial B(0,r))}} \geq \frac{c}{2\ell+d} r^{-2\ell} + O(r).$$

Since the right hand side goes to ∞ as $\ell \rightarrow \infty$ when $r < 1$, we obtain inequality II. **For inequality III, we have**

$$\|h\|_{L^2(B(0,r))}^2 = \sum_{\ell,m} \frac{a_\ell^m(r)^2}{2\ell+d-1} r^{d-1}.$$

Term by term and from equation (15), observe that

$$\|h\|_{L^2(B(0,r))}^2 \leq r \mathcal{D}(h).$$

Inequality III then follows from inequality I.

Finally, for inequality IV, we have from equation (12) that there is a constant C such that $a_\ell^m(\rho) \leq Cc_\ell^m \rho^\ell$. On the other hand, it is known that L^∞ bounds for spherical harmonics are saturated by the zonal harmonics [36, Section 10.2.1], and that

$$\|Y_\ell^m\|_{L^\infty(\mathbb{S}^{d-1})} = O\left(\ell^{\frac{d-1}{2}}\right).$$

Combining the bounds on a_ℓ^m and Y_ℓ^m , and the fact that $R < 1$ we obtain

$$\begin{aligned} \|a_\ell^m Y_\ell^m\|_{L^\infty(A_{r,R})} &\leq C c_\ell^m R^\ell \ell^{\frac{d-1}{2}} \\ &\leq C' c_\ell^m. \end{aligned}$$

Summability of the coefficients c_ℓ^m finishes the proof of inequality IV. \square

3.2. General H^1 comparison theorems on balls. The next two lemmas do not depend on any specific boundary condition. The first one gives bounds for Sobolev constants of annuli.

Lemma 13. *For $0 < r < R < 1$, define*

$$\gamma(r, R) := \inf \left\{ \frac{\int_{A_{r,R}} |\nabla u|^2 + u^2 \, dx}{\int_{\partial B(0,r)} u^2 \, dA} : u \in H^1(A_{r,R}), u|_{\partial B(0,r)} \neq 0 \right\}. \quad (19)$$

Suppose that $R \geq cr^{\frac{d-1}{d}} \geq 2r$ for some $c > 0$. Then, there is a constant C depending only on the dimension and on c such that

$$\gamma(r, R) \geq C \min \left\{ R^d r^{1-d}, r^{\frac{1}{d}-1} \right\}.$$

Proof. As earlier, write a function $u \in H^1(A_{r,R})$ as

$$u(\rho, \boldsymbol{\theta}) = \sum_{\ell, m} a_\ell^m(\rho) Y_\ell^m(\boldsymbol{\theta}).$$

Using the notation $u_\boldsymbol{\theta}$ for the tangential gradient, the Dirichlet energy of u is expressed as

$$\begin{aligned} \mathcal{D}(u) &= \int_r^R \int_{\mathbb{S}^{d-1}} (u_\rho^2 + \rho^{-2} u_\boldsymbol{\theta}^2) \rho^{d-1} \, d\boldsymbol{\theta} \, d\rho \\ &\geq \int_r^R \int_{\mathbb{S}^{d-1}} (u_\rho^2) \rho^{d-1} \, d\boldsymbol{\theta} \, d\rho \\ &= \sum_{\ell, m} \int_r^R (a_\ell^m(\rho)')^2 \rho^{d-1} \, d\rho. \end{aligned}$$

On the other hand, the denominator in (19) is given by

$$\int_{\partial B(0,r)} u^2 \, dA = r^{d-1} \sum_{\ell, m} a_\ell^m(r)^2.$$

Combining these last two expressions in (19) and defining the density $w(\rho) = \left(\frac{\rho}{r}\right)^{d-1}$, we see it is enough to prove that

$$\sum_{\ell, m} \int_r^R \left((\partial_\rho a_\ell^m(\rho))^2 + a_\ell^m(\rho)^2 \right) w(\rho) \, d\rho \geq C \min \left\{ R^d r^{1-d}, r^{\frac{1}{d}-1} \right\} \sum_{\ell, m} a_\ell^m(r)^2. \quad (20)$$

Indeed, working term by term, we prove that any smooth function $f : [r, R] \rightarrow \mathbb{R}$ satisfies

$$\int_r^R (f'(\rho)^2 + f(\rho)^2) w(\rho) \, d\rho \geq C \min \left\{ R^d r^{1-d}, r^{\frac{1}{d}-1} \right\} f(r)^2.$$

To this end, assume without loss of generality that $f(r) = 1$. Following a strategy that was used in [12] and in [8], consider the two following situations.

Let $t \in (r, R)$, to be fixed later.

Case a. Suppose first that for all $\rho \in (t, R)$, $|f(\rho)| \geq 1/2$. It follows from monotonicity and explicit integration that

$$\int_r^R |f(\rho)|^2 w(\rho) d\rho \geq \frac{r^{1-d}}{4d} (R^d - t^d).$$

Case b. Suppose there exists $\rho_0 \in (t, R)$ such that $f(\rho_0) < 1/2$. Splitting the integral, using the fact that $w(\rho) \geq 1$ and is increasing for all ρ together with the Cauchy–Schwarz inequality leads to

$$\begin{aligned} \int_r^R f'(\rho)^2 w(\rho) d\rho &\geq \int_r^t f'(\rho)^2 + \int_t^{\rho_0} f'(\rho)^2 w(\rho) d\rho \\ &\geq \frac{1}{t-r} \left(\int_r^t f'(\rho) d\rho \right)^2 + \left(\frac{t}{r} \right)^{d-1} \frac{1}{\rho_0 - t} \left(\int_t^{\rho_0} f'(\rho) d\rho \right)^2. \end{aligned} \quad (21)$$

By hypothesis $R < 1$ so that $\frac{1}{\rho_0 - t} > 1$. This leads to

$$\int_r^R f'(\rho)^2 w(\rho) d\rho \geq \frac{1}{2} \min \left\{ \frac{1}{t-r}, \left(\frac{t}{r} \right)^{d-1} \right\} \left(\int_r^{\rho_0} f'(\rho) d\rho \right)^2,$$

Choosing $t = \min \left\{ r^{\frac{d-1}{d}}, R/2 \right\}$ guarantees that $\min \left\{ \frac{1}{t-r}, \left(\frac{t}{r} \right)^{d-1} \right\} = \left(\frac{t}{r} \right)^{d-1}$ so that

$$\int_r^R |f'(\rho)|^2 w(\rho) d\rho \geq \frac{1}{2} \left(\frac{t}{r} \right)^{d-1} \left(\int_r^{\rho_0} f'(\rho) d\rho \right)^2.$$

It follows from the definition of t that

$$\left(\frac{t}{r} \right)^{d-1} \geq \min \left\{ 1, \left(\frac{C}{2} \right)^{d-1} \right\} r^{\frac{1-d}{d}}.$$

We can bound asymptotically the last integral in (21) as

$$\begin{aligned} \int_r^R f'(\rho)^2 w(\rho) d\rho &\geq C r^{\frac{1-d}{d}} \left(\int_r^{\rho_0} f'(\rho) d\rho \right)^2 \\ &\geq \frac{A}{4} r^{\frac{1-d}{d}}. \end{aligned}$$

This also ensures that $R^d - t^d \geq (1 - 2^{-d})R^d$. Since both situations are exclusive, inequality (20) holds, finishing the proof. \square

The next Lemma compares L^2 norms on $B(0, r)$ with H^1 norms on $B(0, R)$.

Lemma 14. *For $0 < r < R \leq 1$, if $R \geq cr^{\frac{d-1}{d}}$ for some $c > 0$, there is a constant C depending only on c and on the dimension such that for all $u \in H^1(B(0, R))$,*

$$\|u\|_{L^2(B(0,r))} \leq Cr^{1/2} \|u\|_{H^1(B(0,R))}.$$

Proof. Let $u \in H^1(B(0, R))$. Given $r \in (0, R)$,

$$\int_{B(0,r)} u^2 d\mathbf{x} = \int_0^r \rho^{d-1} \int_{S^{d-1}} u^2 d\theta d\rho = \int_0^r \|u\|_{L^2(\partial B(0,\rho))}^2 d\rho$$

It follows from the definition of γ in Lemma 13 that

$$\|u\|_{\partial B(0,\rho)}^2 \leq \frac{1}{\gamma(\rho, R)} \|u\|_{H^1(A_{\rho,R})}^2 \leq \frac{1}{\gamma(\rho, R)} \|u\|_{H^1(B(0,R))}^2.$$

Substitution in the above leads to

$$\int_{B(0,r)} u^2 \, d\mathbf{x} \leq \|u\|_{H^1(B(0,R))}^2 \int_0^r \frac{1}{\gamma(\rho, R)} \, d\rho.$$

It follows from Lemma 13 that

$$\begin{aligned} \int_0^r \frac{1}{\gamma(\rho, R)} \, d\rho &\leq \frac{1}{C} \int_0^r \left(\frac{\rho^{d-1}}{R^d} + \rho^{1-\frac{1}{d}} \right) \, d\rho \\ &= \frac{1}{C} \left(\frac{r^d}{dR^d} + \frac{r^{2-\frac{1}{d}}}{2-\frac{1}{d}} \right) \\ &\leq \frac{1}{C} \left(\frac{r}{cd} + \frac{r^{2-\frac{1}{d}}}{2-\frac{1}{d}} \right) \quad \leftarrow \text{since } R^d \geq cr^{d-1} \\ &\leq \tilde{C}r. \end{aligned}$$

□

Finally, we require the following lemma about the behaviour of the boundary trace operator as a domain gets shrunk.

Lemma 15. *Let $\Omega \subset \mathbb{R}^d$ be a bounded open set and denote*

$$\gamma(\Omega) := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^2 + u^2 \, d\mathbf{x}}{\int_{\partial\Omega} u^2 \, dA} : u \in H^1(\Omega), u|_{\partial\Omega} \neq 0 \right\}.$$

Then, the following inequality holds as $\varepsilon \rightarrow 0$:

$$\gamma(\varepsilon\Omega) \geq \varepsilon\gamma(\Omega).$$

Proof. Consider $u \in H^1(\varepsilon\Omega)$, $u|_{\partial\varepsilon\Omega} \neq 0$. Then, with the change of variable $\mathbf{y} = \varepsilon^{-1}\mathbf{x}$,

$$\begin{aligned} \int_{\varepsilon\Omega} |\nabla u(\mathbf{x})|^2 + u(\mathbf{x})^2 \, d\mathbf{x} &= \varepsilon^d \int_{\Omega} \varepsilon^{-2} |\nabla u(\mathbf{y})|^2 + u(\mathbf{y})^2 \, d\mathbf{y} \\ &\geq \varepsilon^d \int_{\Omega} \gamma(\Omega) \int_{\partial\Omega} u(\mathbf{y})^2 \, dA \\ &= \varepsilon\gamma(\Omega) \int_{\partial\varepsilon\Omega} u(\mathbf{x})^2 \, dA. \end{aligned}$$

□

3.3. Uniform bounds on Steklov eigenfunctions. In order to obtain convergence of the eigenfunctions we will need that their L^∞ norm stays bounded. In this subsection, since we will need to understand specific interplay between boundary surface area and volume, we will diverge from our convention and denote the area of a boundary $\partial\Omega$ by $\mathcal{H}^{d-1}(\partial\Omega)$. In [4][Theorem 3.1], the authors prove that for any Steklov eigenfunction u of a domain Ω with eigenvalue σ ,

$$\|u\|_{L^\infty(\Omega)} \leq C \|u\|_{L^2(\partial\Omega)}$$

where C depends continuously on the dimension d , σ , $|\Omega|$ and the norm of the trace application

$$T : \text{BV}(\Omega) \rightarrow L^1(\partial\Omega).$$

It is clear that in our situation, the dimension and $|\Omega|$ stay bounded. The eigenvalues will be shown later to also stay bounded, but we need to control the norm of T . In [4][Proposition 5.1], they give the following condition under which said norm stays

bounded for a family of domains. If $\{\Omega^\varepsilon\}$ is a family of open bounded domains with $\mathcal{H}^{d-1}(\partial\Omega^\varepsilon) < \infty$ and such that $\Omega^\varepsilon \subset K$ for some bounded set K , and if there exists constants Q and δ such that for every $x \in \partial\Omega^\varepsilon$

$$\sup \left\{ \frac{\mathcal{H}^{d-1}(\partial^* E \cap \partial^* \Omega^\varepsilon)}{\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)} : E \subset \Omega^\varepsilon \cap B_\delta(x), \text{Per}(E, \Omega_n) < \infty \right\} \leq Q, \quad (22)$$

then the norm of $T : \text{BV}(\Omega^\varepsilon) \rightarrow L^1(\partial\Omega^\varepsilon)$ is uniformly bounded in ε . Here, $\partial^* E$ denotes the reduced boundary of E , which in general may be much smaller than the topological boundary.

The next lemma is inspired by [4][Example 2]. Note that in their example, $r_\varepsilon = o\left(\varepsilon^{\frac{2d-1}{d-1}}\right)$, which means that the radius of the holes is one order of magnitude in ε smaller than the critical level. Parts of the proof are in a similar spirit but we need more precise estimates separately around every boundary component.

Lemma 16. *For all ε , the norm of $T : \text{BV}(\Omega^\varepsilon) \rightarrow L^1(\partial\Omega^\varepsilon)$ is uniformly bounded in ε .*

Proof. Following (22), for any $x_\varepsilon \in \partial\Omega^\varepsilon$, we want to bound the ratio

$$\frac{\mathcal{H}^{d-1}(\partial^* E \cap \partial\Omega^\varepsilon)}{\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)} \quad (23)$$

for sets $E \subset B_\delta(x_\varepsilon) \cap \Omega^\varepsilon$. Let us make the observation that

$$\#\{\mathbf{n} \in I^\varepsilon : Q_\mathbf{n}^\varepsilon \cap B_\delta(x_\varepsilon) \neq \emptyset\} \leq 2\omega_d \left(\frac{\delta}{\varepsilon}\right)^d$$

and for some M , $\mathcal{H}^{d-1}(\partial\Omega \cap B_\delta(x_\varepsilon)) \leq M\delta^{d-1}$ so that for any $E \subset \Omega^\varepsilon \cap B_\delta(x_\varepsilon)$ of finite perimeter,

$$\mathcal{H}^{d-1}(\partial^* E \cap \partial\Omega^\varepsilon) \leq \mathcal{H}^{d-1}(\partial\Omega^\varepsilon \cap B_\delta(x_\varepsilon)) \leq \left(M + \frac{2d}{\beta}\delta\right)\delta^{d-1}. \quad (24)$$

In particular, we can assume without loss of generality that this previous bound is also one for $\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)$, otherwise the ratio (23) is smaller than 1.

By definition of I^ε , for all $\mathbf{n} \in I^\varepsilon$ we have that $\partial\Omega^\varepsilon \cap Q_\mathbf{n}^\varepsilon = \partial T_\mathbf{n}^\varepsilon$. From this, we can decompose

$$\mathcal{H}^{d-1}(\partial^* E \cap \partial\Omega^\varepsilon) = \mathcal{H}^{d-1}(\partial^* E \cap \partial\Omega) + \sum_{\mathbf{n} \in I^\varepsilon} \mathcal{H}^{d-1}(\partial^* E \cap \partial T_\mathbf{n}^\varepsilon). \quad (25)$$

We first observe that if δ is chosen such that $2\omega_d\delta^d \leq |\Omega|$, the relative isoperimetric inequality [14][Theorem 5.6.2 (ii)] implies that there is a constant C such that

$$\begin{aligned} \mathcal{H}^{d-1}(\partial^* E \cap \partial\Omega) &\leq C\mathcal{H}^{d-1}(\partial^* E \cap \Omega) \\ &= C(\mathcal{H}^{d-1}(\partial^* E \cap (\partial\Omega^\varepsilon \setminus \partial\Omega)) + \mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)). \end{aligned} \quad (26)$$

This means that we only need to find bounds for the sum over I^ε in (25).

The following construction is a localised version of the sets E_t and F_t defined in [4, Page 18]. For $\mathbf{n} \in I^\varepsilon$, define

$$E_{\mathbf{n},t} := E \cap \{x \in Q_\mathbf{n}^\varepsilon : \text{dist}_{\partial T_\mathbf{n}^\varepsilon}(x) > t\}$$

and

$$F_{\mathbf{n},t} := E \setminus \{x : \text{dist}_{\partial T_\mathbf{n}^\varepsilon}(x) > t\}.$$

Assume that there is $t \in (0, \varepsilon/4)$ such that

$$\mathcal{H}^{d-1}(\partial^* F_{\mathbf{n},t} \cap Q_\mathbf{n}^\varepsilon) \leq \mathcal{H}^{d-1}(\partial^* E \cap Q_\mathbf{n}^\varepsilon). \quad (27)$$

Since projections on convex sets are nonexpansive and $F_{\mathbf{n},t} \subset Q_{\mathbf{n}}^\varepsilon$, we have that

$$\begin{aligned} \mathcal{H}^{d-1}(\partial^* E \cap \partial T_{\mathbf{n}}^\varepsilon) &= \mathcal{H}^{d-1}(\partial^* F_{\mathbf{n},t} \cap \partial T_{\mathbf{n}}^\varepsilon) \\ &\leq \mathcal{H}^{d-1}(\partial^* F_{\mathbf{n},t} \cap Q_{\mathbf{n}}^\varepsilon) \\ &\leq \mathcal{H}^{d-1}(\partial^* E \cap Q_{\mathbf{n}}^\varepsilon). \end{aligned} \quad (28)$$

Let $J^\varepsilon := \{\mathbf{n} \in I^\varepsilon : \text{equation (27) holds}\}$. We assume that $J^\varepsilon \neq I^\varepsilon$, since if it is the case our previous arguments is sufficient to finish the proof. Let $\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon$.

From the coarea formula and the isoperimetric inequality (Add a bit more details, see [4][page 18]) we have that there is a dimensional constant C_d such that

$$\begin{aligned} |E \cap Q_{\mathbf{n}}^\varepsilon| &\geq C_d \varepsilon^d = C_d |Q_{\mathbf{n}}^\varepsilon| \\ &= \frac{C_d}{A_d \beta} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon). \end{aligned} \quad (29)$$

Using once again the isoperimetric inequality as well as equations (26) and (28) to find C (whose exact value may change line to line), depending only on the dimension and on Ω

$$\begin{aligned} \sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} |E \cap Q_{\mathbf{n}}^\varepsilon| &\leq C \mathcal{H}^{d-1}(\partial^* E)^{\frac{d}{d-1}} \\ &= C \left(\mathcal{H}^{d-1}(\partial^* E \cap \partial \Omega) + \mathcal{H}^{d-1}(\partial^* E \cap \partial T^\varepsilon) + \mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon) \right)^{\frac{d}{d-1}} \\ &\leq C \left(\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon) + \sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon) \right)^{\frac{d}{d-1}}. \end{aligned} \quad (30)$$

Combining equations (29) and (30), we obtain that there is a constant C depending only on the dimension, Ω and β such that

$$1 \leq C \left(\sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon) \right)^{\frac{1}{d-1}} \left(1 + \frac{\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)}{\sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon)} \right)^{\frac{d}{d-1}}.$$

It follows from (24) that we can choose δ small enough depending on Ω , the dimension, β but not on ε such that

$$C \left(\sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon) \right)^{\frac{1}{d-1}} \leq \frac{1}{2}.$$

This implies that there is a constant c such that for all $\varepsilon > 0$,

$$c \leq \frac{\mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon)}{\sum_{\mathbf{n} \in I^\varepsilon \setminus J^\varepsilon} \mathcal{H}^{d-1}(\partial T_{\mathbf{n}}^\varepsilon)}. \quad (31)$$

Combining (26), (28) and (31), we have that there is a constant C such that

$$\mathcal{H}^{d-1}(\partial^* E \cap \partial \Omega^\varepsilon) \leq C \mathcal{H}^{d-1}(\partial^* E \cap \Omega^\varepsilon),$$

finishing the proof. \square

Remark 17. When $r_\varepsilon = o\left(\varepsilon^{\frac{d}{d-1}}\right)$, (i.e. in the subcritical regime), the previous result along with [4][Theorem 4.1] implies convergence of the eigenvalues of the Steklov problem on Ω^ε to the eigenvalues of the Steklov problem on Ω .

4. Homogenisation of the Steklov problem

Let us first establish some basic facts related to the geometry of the homogenisation problem, under the assumption that $r_\varepsilon^{d-1}\varepsilon^{-d} \rightarrow \beta \in [0, \infty)$ as $\varepsilon \rightarrow 0$. The number of holes $N(\varepsilon) = \#I^\varepsilon$ satisfies

$$N(\varepsilon) \sim |\Omega|\varepsilon^{-d} \quad \text{as } \varepsilon \rightarrow 0.$$

This implies,

$$|T^\varepsilon| = \sum_{\mathbf{k} \in I^\varepsilon} |T_{\mathbf{k}}^\varepsilon| = O(r_\varepsilon) \quad \text{and} \quad |\partial T^\varepsilon| = \sum_{\mathbf{k} \in I^\varepsilon} |\partial T_{\mathbf{k}}^\varepsilon| \sim A_d \beta |\Omega|.$$

The remainder of the section is split into three parts. In the first one we extend the functions u_k^ε to the whole of Ω , in order to obtain weak H^1 convergence, up to taking subsequences. In the second part, we prove that those converging subsequences converge to solutions of Problem (3). Finally, we prove in the third part that the only functions they can converge to are the corresponding eigenfunction in (3), implying convergence as $\varepsilon \rightarrow 0$, with this understood in the sense of Remark 3 if the limit problem has eigenvalues that are not simple.

4.1. Extension of eigenfunctions. For $k \geq 1$, recall that $u_k^\varepsilon : \Omega^\varepsilon \rightarrow \mathbb{R}$ is the k 'th Steklov eigenfunction on Ω^ε :

$$\begin{cases} \Delta u_k^\varepsilon = 0 & \text{in } \Omega^\varepsilon, \\ \partial_\nu u_k^\varepsilon = \sigma_k^\varepsilon u_k^\varepsilon & \text{on } \partial\Omega^\varepsilon. \end{cases}$$

Recall also that the eigenfunctions u_k^ε are normalized by requiring that

$$\int_{\partial\Omega^\varepsilon} (u_k^\varepsilon)^2 \, dA = 1.$$

Define the function $U_k^\varepsilon \in H^1(\Omega)$ to be the harmonic extension of u_k^ε to the interior of the holes:

$$\begin{cases} U_k^\varepsilon = u_k^\varepsilon & \text{in } \overline{\Omega^\varepsilon}, \\ \Delta U_k^\varepsilon = 0 & \text{in } T^\varepsilon. \end{cases}$$

Lemma 18. *There is a sequence $\varepsilon_n \rightarrow 0$ such that $U_k^{\varepsilon_n}$ has a weak limit in $H^1(\Omega)$.*

Proof. It suffices to show that $\{U_k^\varepsilon : 0 < \varepsilon \leq 1\}$ is bounded in $H^1(\Omega)$. Recall that

$$\|U_k^\varepsilon\|_{H^1}^2 = \|U_k^\varepsilon\|_{L^2}^2 + \mathcal{D}(U_k^\varepsilon).$$

Let us start by an estimation of the Dirichlet energy:

$$\mathcal{D}(U_k^\varepsilon; \Omega) = \mathcal{D}(u_k^\varepsilon; \Omega^\varepsilon) + \mathcal{D}(U_k^\varepsilon; T^\varepsilon) = \sigma_k^\varepsilon + \mathcal{D}(U_k^\varepsilon; T^\varepsilon). \quad (32)$$

It follows from inequality I in Lemma 12 and monotonicity of the Dirichlet energy, that the contribution from the holes is

$$\begin{aligned} \mathcal{D}(U_k^\varepsilon; T^\varepsilon) &= \sum_{\mathbf{k} \in I^\varepsilon} \mathcal{D}(U_k^\varepsilon; T_{\mathbf{k}}^\varepsilon) \\ &\leq \sum_{\mathbf{k} \in I^\varepsilon} 5 \left(\frac{r_\varepsilon}{\varepsilon}\right)^d \mathcal{D}(u_k^\varepsilon; Q_{\mathbf{k}}^\varepsilon) \left(1 + O\left(\left(\frac{r_\varepsilon}{\varepsilon}\right)^d\right)\right) \\ &\leq 5 \left(\frac{r_\varepsilon}{\varepsilon}\right)^d \mathcal{D}(u_k^\varepsilon; \Omega^\varepsilon) \left(1 + O\left(\left(\frac{r_\varepsilon}{\varepsilon}\right)^d\right)\right) \\ &\leq C \sigma_k(\Omega^\varepsilon), \end{aligned} \quad (33)$$

for some constant C . It is known from [11] that there is a constant $C_{d,k}$ such that

$$\sigma_k^\varepsilon \leq C_{d,k} |\partial\Omega^\varepsilon|^{\frac{1}{1-d}},$$

which implies that σ_k^ε is bounded above independently of ε . Inserting that bound in equations (32) and (33) gives

$$\mathcal{D}(U_k^\varepsilon) \ll |\partial\Omega^\varepsilon|^{\frac{1}{1-d}}.$$

Let us now bound the L^2 norm of U_k^ε . Let λ be the first eigenvalue of the following Robin problem on Ω :

$$\begin{cases} -\Delta u = \lambda u & \text{in } \Omega, \\ \partial_\nu u = -u & \text{on } \partial\Omega. \end{cases}$$

It is well known (see *e.g.* [3]) that $\lambda > 0$ and that it admits the following characterization:

$$\lambda = \inf_{v \in H^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 \, d\mathbf{x} + \int_{\partial\Omega} v^2 \, dS}{\int_\Omega v^2 \, d\mathbf{x}}.$$

Applying this to $v = U_k^\varepsilon$ leads to

$$\begin{aligned} \int_\Omega (U_k^\varepsilon)^2 \, d\mathbf{x} &\leq \frac{1}{\lambda} \left(\int_\Omega |\nabla U_k^\varepsilon|^2 \, d\mathbf{x} + \int_{\partial\Omega} (U_k^\varepsilon)^2 \, dA \right) \\ &\leq \frac{1}{\lambda} (\mathcal{D}(U_k^\varepsilon) + 1). \end{aligned}$$

Together this leads to

$$\|U_k^\varepsilon\|_{H^1(\Omega)}^2 \leq \frac{1}{\lambda} \left(\frac{C}{|\partial\Omega|^{1/(d-1)}} + 1 \right)$$

for some constant C , finishing the proof. \square

From now on we will abuse notation and relabel that sequence $\varepsilon_n \rightarrow 0$ along which U_k^ε has a weak limit as $\varepsilon \rightarrow 0$ again.

4.2. Establishing the limit problem. Our aim by the end of this subsection is to prove the following weaker version of Theorem 2.

Proposition 19. *Let $k \in \mathbb{N}$. As $\varepsilon \rightarrow 0$, the pairs $(\sigma_k^\varepsilon, U_k^\varepsilon)$ converge to a solution (Σ, U) of (3), the convergence of the functions U_k^ε being weak in $H^1(\Omega)$.*

Up to choosing a subsequence, we assume that σ_k^ε converges to some number Σ and also that $\{U_k^\varepsilon\} \subset H^1(\Omega)$ is weakly converging in $H^1(\Omega)$ to some $U \in H^1(\Omega)$, from which we also get strong convergence to U in $L^2(\Omega)$. Considering the real-valued test function $V \in H^1(\Omega)$, we see that

$$\begin{aligned} \int_\Omega \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x} &= \int_{\Omega^\varepsilon} \nabla u_k^\varepsilon \cdot \nabla V \, d\mathbf{x} + \int_{T^\varepsilon} \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x} \\ &= \sigma_k^\varepsilon \int_{\partial\Omega^\varepsilon} u_k^\varepsilon V \, d\mathbf{x} + \int_{T^\varepsilon} \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x} \\ &= \sigma_k^\varepsilon \int_{\partial\Omega} u_k^\varepsilon V \, dA + \sigma_k^\varepsilon \int_{\partial T^\varepsilon} u_k^\varepsilon V \, dA + \int_{T^\varepsilon} \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x}. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$ leads, if the limits exist, to

$$\int_\Omega \nabla U \cdot \nabla V \, d\mathbf{x} - \Sigma \int_{\partial\Omega} UV \, dA = \Sigma \lim_{\varepsilon \rightarrow 0} \int_{\partial T^\varepsilon} u_k^\varepsilon V \, dA + \lim_{\varepsilon \rightarrow 0} \int_{T^\varepsilon} \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x}.$$

It follows from the Cauchy–Schwarz inequality that

$$\int_{T^\varepsilon} \nabla U_k^\varepsilon \cdot \nabla V \, d\mathbf{x} \leq \left(\int_{T^\varepsilon} |\nabla U_k^\varepsilon|^2 \, d\mathbf{x} \int_{T^\varepsilon} |\nabla V|^2 \, d\mathbf{x} \right)^{1/2},$$

which tends to 0 according to inequality I in Lemma 12. It follows that

$$\int_{\Omega} \nabla U \cdot \nabla V \, d\mathbf{x} - \Sigma \int_{\partial\Omega} UV \, dA = \Sigma \lim_{\varepsilon \rightarrow 0} \int_{\partial T^\varepsilon} u_k^\varepsilon V \, dA,$$

and all that is left to do is to analyse the last term.

Proposition 20. *Suppose that $\varepsilon^{-d} r_\varepsilon^{d-1} \rightarrow \beta \geq 0$. Then, for each $V \in H^1(\Omega)$ the following holds,*

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial T^\varepsilon} u_k^\varepsilon V \, dA = A_d \beta \int_{\Omega} UV \, d\mathbf{x}.$$

Remark 21. *The functional $V \mapsto \int_{\partial T^\varepsilon} u_k^\varepsilon V$ is bounded on $H^1(\Omega)$. By the Riesz–Fréchet representation theorem, there exists a function $\xi^\varepsilon \in H^1(\Omega)$ such that*

$$\int_{\partial T^\varepsilon} u_k^\varepsilon V = \int_{\Omega} \nabla \xi^\varepsilon \cdot \nabla V + \xi^\varepsilon V \, dx \quad \forall V \in H^1(\Omega).$$

Using appropriate test functions shows that ξ^ε is the weak solution of the following transmission problem:

$$\begin{cases} \Delta \xi^\varepsilon = 0 & \text{in } \Omega^\varepsilon \cup T^\varepsilon, \\ \partial_\nu \xi^\varepsilon_+ + \partial_\nu \xi^\varepsilon_- = u_k^\varepsilon & \text{on } \partial T^\varepsilon, \\ \partial_\nu \xi^\varepsilon = 0 & \text{on } \partial\Omega. \end{cases}$$

Proposition 20 is an homogenisation result for this problem. It means that in the limit as $\varepsilon \rightarrow 0$, the solution converges to that of the following problem:

$$\begin{cases} -\Delta \Xi + (1 - A_d \beta) \Xi = 0 & \text{in } \Omega, \\ \partial_\nu \Xi = 0 & \text{on } \partial\Omega. \end{cases}$$

Transmission problems have recently been the subject of investigation through means of homogenisation, see for example [29].

Proof of Proposition 20. Define the family of bounded functionals $L_\varepsilon : H^1(\Omega) \rightarrow \mathbb{R}$ by

$$L_\varepsilon(V) := \int_{\partial T^\varepsilon} V \, dA.$$

We will give two representations of those functionals.

Step 1: Inner representation of L_ε .

Define $\varphi_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$\varphi_\varepsilon(\mathbf{x}) = \begin{cases} \frac{|\mathbf{x}|^2}{2r_\varepsilon} & \text{for } \mathbf{x} \in B(0, r_\varepsilon), \\ 0 & \text{elsewhere.} \end{cases} \quad (34)$$

By periodizing along $\varepsilon \mathbb{Z}^d$ we obtain the function $\Phi_\varepsilon : \mathbb{R}^d \rightarrow \mathbb{R}$ given by

$$\Phi_\varepsilon(\mathbf{x}) := \sum_{\mathbf{k} \in I^\varepsilon} \varphi_\varepsilon(\mathbf{x} - \varepsilon \mathbf{k}).$$

Lemma 22. *The functional $L_\varepsilon : H^1(\Omega) \rightarrow \mathbb{R}$ admits the following representation:*

$$L_\varepsilon(V) = \frac{d}{r_\varepsilon} \int_{T^\varepsilon} V \, d\mathbf{x} + \int_{T^\varepsilon} \nabla \Phi_\varepsilon \cdot \nabla V \, d\mathbf{x}.$$

Proof. It is straightforward to check that

$$\begin{cases} \Delta \Phi_\varepsilon = \frac{d}{r_\varepsilon} & \text{in } T^\varepsilon, \\ \partial_\nu \Phi_\varepsilon = 1 & \text{on } \partial T^\varepsilon, \\ \Phi = 0 & \text{in } \Omega^\varepsilon. \end{cases}$$

The function Φ_ε therefore satisfies the weak identity

$$\int_{T^\varepsilon} \nabla \Phi_\varepsilon \cdot V = -\frac{d}{r_\varepsilon} \int_{T^\varepsilon} V + \int_{\partial T^\varepsilon} V, \quad \forall V \in H^1(\Omega).$$

□

For each $k \in \mathbb{N}$ and $\varepsilon > 0$, the functional $\widetilde{L}_\varepsilon : H^1(\Omega) \rightarrow \mathbb{R}$ is defined by

$$\widetilde{L}_\varepsilon(V) := L(u_k^\varepsilon V).$$

Lemma 23. *There is an $\varepsilon_0 > 0$ such that the family $\{\widetilde{L}_\varepsilon\}_{\varepsilon > 0} \subset (H^1(\Omega))^*$ is uniformly bounded for $0 < \varepsilon \leq \varepsilon_0$.*

Proof. Given $V \in H^1(\Omega)$, it follows from the Lemma 22 that

$$L_\varepsilon(U_k^\varepsilon V) = \frac{d}{r_\varepsilon} \int_{T^\varepsilon} U_k^\varepsilon V \, dx + \int_{T^\varepsilon} \nabla \Phi_\varepsilon \cdot \nabla(U_k^\varepsilon V) \, dx. \quad (35)$$

To bound the first term, start by using the Cauchy–Schwarz inequality:

$$\left| \frac{d}{r_\varepsilon} \int_{T^\varepsilon} U_k^\varepsilon V \, dx \right| \leq \frac{d}{r_\varepsilon} \|U_k^\varepsilon\|_{L^2(T^\varepsilon)} \|V\|_{L^2(T^\varepsilon)}. \quad (36)$$

It follows from Lemma 14 that

$$\begin{aligned} \|V\|_{L^2(T^\varepsilon)} &\leq r_\varepsilon^{1/2} \|V\|_{H^1(\Omega)}, \\ \|U_k^\varepsilon\|_{L^2(T^\varepsilon)} &\leq r_\varepsilon^{1/2} \|U_k^\varepsilon\|_{H^1(\Omega)}, \end{aligned} \quad (37)$$

so that

$$\sup_{\varepsilon \in (0,1)} \left| \frac{d}{r_\varepsilon} \int_{T^\varepsilon} U_k^\varepsilon V \, dx \right| < \infty.$$

Inequality III of Lemma 12 and monotonicity of the Dirichlet energy lead to

$$\begin{aligned} \|U^\varepsilon\|_{L^2(T^\varepsilon)}^2 &= \sum_{k \in I^\varepsilon} \|U^\varepsilon\|_{L^2(T_k^\varepsilon)}^2 \\ &\leq C \left(\frac{r_\varepsilon}{\varepsilon}\right)^{d+1} \sum_{k \in I^\varepsilon} \mathcal{D}(U^\varepsilon; B(\varepsilon \mathbf{k}, \varepsilon) \setminus B(\varepsilon \mathbf{k}, r_\varepsilon)) \\ &\leq C \left(\frac{r_\varepsilon}{\varepsilon}\right)^{d+1} \sigma_k(\Omega_\varepsilon). \end{aligned}$$

We have already seen that $\sigma_k(\Omega_\varepsilon)$ is bounded above independently of ε , according to [10]. Moreover,

$$\frac{1}{r_\varepsilon} \left(\frac{r_\varepsilon}{\varepsilon}\right)^{d+1} = \frac{r_\varepsilon^{d-1}}{\varepsilon^d} \frac{r_\varepsilon}{\varepsilon} \rightarrow 0 \quad \text{as } \varepsilon \searrow 0.$$

It follows that

$$r_\varepsilon^{-1/2} \|U_k^\varepsilon\|_{L^2(T^\varepsilon)} < \infty.$$

Substitution in (36) together with (37) lead to

$$\sup_{\varepsilon \in (0,1)} \left| \frac{d}{r_\varepsilon} \int_{T^\varepsilon} U_k^\varepsilon V \, dx \right| < \infty. \quad (38)$$

To bound the second term in (35), the generalised Hölder inequality leads to

$$\begin{aligned}
\left| \int_{T^\varepsilon} \nabla \Phi_\varepsilon \cdot \nabla (U_k^\varepsilon V) \, d\mathbf{x} \right| &\leq \left| \int_{T^\varepsilon} U_k^\varepsilon \nabla \Phi_\varepsilon \cdot \nabla V + V \nabla \Phi_\varepsilon \cdot \nabla U_k^\varepsilon \, d\mathbf{x} \right| \\
&\leq \|U_k^\varepsilon\|_{L^2(T^\varepsilon)} \|\nabla V\|_{L^2(T^\varepsilon)} \|\nabla \Phi_\varepsilon\|_{L^\infty(T^\varepsilon)} \\
&\quad + \|\nabla U_k^\varepsilon\|_{L^2(T^\varepsilon)} \|V\|_{L^2(T^\varepsilon)} \|\nabla \Phi_\varepsilon\|_{L^\infty(T^\varepsilon)} \\
&\leq \left(\|U_k^\varepsilon\|_{L^2(T^\varepsilon)} + \|\nabla U_k^\varepsilon\|_{L^2(T^\varepsilon)} \right) \|\nabla V\|_{H^1(\Omega)} \|\nabla \Phi_\varepsilon\|_{L^\infty(T^\varepsilon)} \\
&\leq \|U_k^\varepsilon\|_{H^1(T^\varepsilon)} \|\nabla V\|_{H^1(\Omega)}.
\end{aligned}$$

In the last inequality we have used $\|\nabla \Phi_\varepsilon\|_{L^\infty(T^\varepsilon)} = 1$, which follows from (34). This quantity is uniformly bounded as $\varepsilon \searrow 0$ since we have shown in the proof of Lemma 18 that U_k^ε is bounded in $H^1(\Omega)$. Together with (38) this proves for each $V \in H^1(\Omega)$ the existence of a constant C such that $|L_\varepsilon(U_k^\varepsilon V)| \leq C\|V\|$ for each ε , and the conclusion follows from the Banach–Steinhaus theorem. \square

Step 2: Outer representation of L_ε . Consider the torus $\mathcal{C} = \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ and introduce the fundamental cell \mathcal{C}^ε as the perforated torus

$$\mathcal{C}^\varepsilon := \mathcal{C} \setminus \overline{B(0, \rho_\varepsilon)},$$

where $\rho_\varepsilon := \varepsilon^{-1} r_\varepsilon$ is the renormalised radius. Following [34], we define the function $\psi_\varepsilon \in H^1(\mathcal{C}^\varepsilon)$ through the weak variational problem:

$$\int_{\mathcal{C}^\varepsilon} \nabla \psi_\varepsilon \cdot \nabla V = -c_\varepsilon \int_{\mathcal{C}^\varepsilon} V + \int_{\partial B(0, \rho_\varepsilon)} V. \quad (39)$$

By taking $V \equiv 1$, one sees that the necessary and sufficient condition for existence of a solution (see e.g. [33, Theorem 5.7.7]) is

$$c_\varepsilon = \frac{A_d \rho_\varepsilon^{d-1}}{|\mathcal{C}^\varepsilon|} \sim A_d (\rho_\varepsilon)^{d-1}.$$

Uniqueness of the solution is guaranteed by requiring that ψ_ε be orthogonal to constants on \mathcal{C}^ε . Therefore, ψ_ε is the unique function such that

$$\begin{cases} \Delta \psi_\varepsilon = c_\varepsilon & \text{in } \mathcal{C}^\varepsilon \\ \partial_\nu \psi_\varepsilon = 1 & \text{on } \partial B(0, \rho_\varepsilon) \end{cases} \quad \text{and} \quad \int_{\mathcal{C}^\varepsilon} \psi_\varepsilon = 0.$$

Consider the union of all cells strictly contained in Ω ,

$$\tilde{\Omega}^\varepsilon := \bigcup_{\mathbf{k} \in I^\varepsilon} Q_{\mathbf{k}}^\varepsilon \subset \Omega^\varepsilon.$$

Define the function $\Psi_\varepsilon : \mathbb{R}^d \setminus \bigcup_{\mathbf{k} \in \mathbb{Z}^d} T_{\mathbf{k}}^\varepsilon \rightarrow \mathbb{R}$ as the scaled lift of ψ_ε . That is, if $q : \mathbb{R}^d \rightarrow \mathbb{R}^d / \mathbb{Z}^d$ is the covering map, then

$$\Psi_\varepsilon(\varepsilon \mathbf{x}) := \psi_\varepsilon(q(\mathbf{x})).$$

This function satisfies

$$\begin{cases} \Delta \Psi_\varepsilon = \varepsilon^{-2} c_\varepsilon & \text{in } \tilde{\Omega}^\varepsilon, \\ \partial_\nu \Psi_\varepsilon = \varepsilon^{-1} & \text{on } \partial T^\varepsilon. \end{cases}$$

Lemma 24. *The functional $L_\varepsilon : H^1(\Omega) \rightarrow \mathbb{R}$ admits the following representation:*

$$L_\varepsilon(V) = \varepsilon \int_{\tilde{\Omega}^\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla V \, d\mathbf{x} + \varepsilon \int_{\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} V \partial_\nu \Psi_\varepsilon \, dA + \varepsilon^{-1} c_\varepsilon \int_{\tilde{\Omega}^\varepsilon} V \, dA.$$

The proof is immediate since for $V \in H^1(\Omega)$ the following holds:

$$\int_{\tilde{\Omega}^\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla V \, d\mathbf{x} = -\varepsilon^{-2} c_\varepsilon \int_{\tilde{\Omega}^\varepsilon} V \, d\mathbf{x} + \varepsilon^{-1} \int_{\partial T^\varepsilon} V \, dA + \int_{\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} V \partial_\nu \Psi_\varepsilon \, dA.$$

We establish the following claim concerning ψ_ε .

Lemma 25. *There is a constant C , depending only on the dimension and on β , such that*

$$\|\psi_\varepsilon\|_{H^1(\mathcal{C}^\varepsilon)} \leq C \varepsilon^{\frac{1}{2} + \frac{1}{d}}.$$

Furthermore, for any $s > 1$, any compact set $K \subset \mathcal{C}$, containing the origin in its interior, there is a constant C' depending only on the dimension, β , s , and on K such that

$$\|\psi_\varepsilon\|_{H^s(\mathcal{C}^\varepsilon \setminus K)} \leq C' \varepsilon^{\frac{1}{2} + \frac{1}{d}}. \quad (40)$$

In particular, this implies $\|D^\alpha \psi_\varepsilon\|_{L^\infty(\mathcal{C}^\varepsilon \setminus K)}$ decays as $\varepsilon^{\frac{1}{2} + \frac{1}{d}}$ for any multi-index α .

Proof. Observe that since ψ_ε has mean 0 on \mathcal{C}^ε , the Poincaré–Wirtinger inequality implies

$$\mu_1(\mathcal{C}^\varepsilon) \|\psi_\varepsilon\|_{L^2(\mathcal{C}^\varepsilon)}^2 \leq \|\nabla \psi_\varepsilon\|_{L^2(\mathcal{C}^\varepsilon)}^2,$$

where $\mu_1(\mathcal{C}^\varepsilon)$ is the first non-zero Neumann eigenvalue of \mathcal{C}^ε . Observe that

$$\mu_1(\mathcal{C}^\varepsilon) \rightarrow 4\pi^2 \quad \text{as } \varepsilon \rightarrow 0.$$

Indeed, $\mu_1(\mathcal{C}^\varepsilon)$ is the first non-zero Neumann eigenvalue of a punctured d -dimensional torus, which is known to converge to the first nonzero eigenvalue of the torus itself as $\varepsilon \searrow 0$. See [5, Chapter 9] for instance.

Take $V = \psi_\varepsilon$ in the variational characterisation (39) of ψ_ε , and consider ε to be small enough that $\mu_1(\mathcal{C}^\varepsilon) \geq 1$. Using the Cauchy–Schwarz inequality yields

$$\begin{aligned} \|\psi_\varepsilon\|_{H^1(\mathcal{C}^\varepsilon)}^2 &\leq 2 \int_{\mathcal{C}^\varepsilon} |\nabla \psi_\varepsilon|^2 \, d\mathbf{x} \\ &\leq 2 \int_{\partial B(0, \rho_\varepsilon)} \psi_\varepsilon \, dA \\ &\leq 2 \sqrt{A_d \rho_\varepsilon^{d-1}} \|\psi_\varepsilon\|_{L^2(\partial B(0, \rho_\varepsilon))} \\ &\leq 2 \sqrt{A_d \rho_\varepsilon^{d-1}} \|\tau_\varepsilon\| \|\psi_\varepsilon\|_{H^1(\mathcal{C}^\varepsilon)}, \end{aligned} \quad (41)$$

where τ_ε is the trace operator $H^1(\mathcal{C}^\varepsilon) \rightarrow L^2(\partial B(0, \rho_\varepsilon))$. From the definition of τ_ε and monotonicity of the involved integrals, we have that

$$\|\tau_\varepsilon\| \leq \gamma \left(\varepsilon^{\frac{1}{d-1}}, 1 \right)^{-1},$$

where γ is defined in Lemma 13. We therefore deduce that $\|\tau_\varepsilon\| \ll \varepsilon^{1/d}$, the implicit constant depending only on the dimension and on β . Finally, dividing both sides in (41) by $\|\psi_\varepsilon\|_{H^1}$, observing that $\rho_\varepsilon^{d-1} \sim \beta \varepsilon$ and inserting the bound for $\|\tau_\varepsilon\|$ in (41) finishes the proof of the first inequality.

For the second one, we have from [19, Theorem 8.10] that there is a constant \tilde{C} depending only on the dimension, on K , and on s such that

$$\|\psi_\varepsilon\|_{H^s(\mathcal{C}^\varepsilon \setminus K)} \leq \tilde{C} \left(\|\psi_\varepsilon\|_{H^1(\mathcal{C}^\varepsilon)} + \|c_\varepsilon\|_{H^s(\mathcal{C}^\varepsilon)} \right).$$

The second term is of order $\rho_\varepsilon^{d-1} \sim \varepsilon$, and bounds for the H^1 norm of ψ_ε that were obtained in (41) conclude the proof of the second inequality.

As for the remark considering the L^∞ bounds on derivatives of ψ_ε , we have from inequality (40) that

$$\|D^\alpha \psi_\varepsilon\|_{H^{s-|\alpha|}(\mathcal{C}^\varepsilon \setminus K)} \leq \|\psi_\varepsilon\|_{H^s(\mathcal{C}^\varepsilon \setminus K)} \leq C' \varepsilon^{\frac{1}{2} + \frac{1}{d}}.$$

Choosing $s > |\alpha| + \frac{d+1}{2}$, and ε small enough that $\mathcal{C}^\varepsilon \setminus K = \mathcal{C} \setminus K$, we have that

$$H^{s-|\alpha|}(\mathcal{C}^\varepsilon \setminus K) \hookrightarrow L^\infty(\mathcal{C}^\varepsilon \setminus K)$$

with norm independent of ε , concluding the proof. \square

Step 3: Computing the limit problem. It follows from Lemma 23 that it is sufficient to verify convergence for a test function $V \in C^\infty(\overline{\Omega})$. From the outer representation for L_ε we have that

$$\begin{aligned} \tilde{L}_\varepsilon(V) &= \int_{\partial T^\varepsilon} u_k^\varepsilon V \\ &= \varepsilon \int_{\tilde{\Omega}^\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla(u_k^\varepsilon V) + \int_{\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} u_k^\varepsilon V \partial_\nu \Psi_\varepsilon + \underbrace{\varepsilon^{-1} c_\varepsilon \int_{\tilde{\Omega}^\varepsilon} u_k^\varepsilon V}_{-Ad\beta \int_\Omega UV}, \end{aligned} \quad (42)$$

where convergence of the last term stems from strong L^2 convergence. We now show that the two other terms converge to 0. For the first one, the Cauchy–Schwarz inequality gives

$$\int_{\tilde{\Omega}^\varepsilon} \varepsilon \nabla \Psi_\varepsilon \cdot \nabla(u_k^\varepsilon V) \leq \left(\int_{\tilde{\Omega}^\varepsilon} \varepsilon^2 |\nabla \Psi_\varepsilon|^2 \int_{\tilde{\Omega}^\varepsilon} |\nabla(u_k^\varepsilon V)|^2 \right)^{1/2}.$$

Let us first observe that since V is in $C^\infty(\overline{\Omega})$, we have that

$$\begin{aligned} \int_{\tilde{\Omega}^\varepsilon} |\nabla(u_k^\varepsilon V)|^2 &\leq 2 \sup_{\mathbf{x} \in \tilde{\Omega}^\varepsilon} (|V(\mathbf{x})|^2 + |\nabla V(\mathbf{x})|^2) \|U_k^\varepsilon\|_{H_1(\Omega)} \\ &= O(1). \end{aligned}$$

On to the other term, note that

$$\begin{aligned} \int_{\tilde{\Omega}^\varepsilon} \varepsilon^2 |\nabla \Psi_\varepsilon|^2 &= \varepsilon^2 \sum_{\mathbf{k} \in I^\varepsilon} \int_{Q_{\mathbf{k}}^\varepsilon} |\nabla \Psi_\varepsilon|^2 \\ &= \sum_{\mathbf{k} \in I^\varepsilon} \varepsilon^d \int_{\mathcal{C}^\varepsilon} |\nabla \Psi_\varepsilon|^2 \\ &\sim |\Omega| \int_{\mathcal{C}^\varepsilon} |\nabla \Psi_\varepsilon|^2. \end{aligned}$$

It follows from Lemma 25 that

$$\int_{\mathcal{C}^\varepsilon} |\nabla \Psi_\varepsilon|^2 \, d\mathbf{x} = O\left(\varepsilon^{1 + \frac{2}{d}}\right),$$

hence that term indeed goes to 0. For the second term in (42), we have from the generalised Hölder inequality that

$$\varepsilon \int_{\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} u_k^\varepsilon V \partial_\nu \Psi_\varepsilon \, dA \leq \varepsilon \|V\|_{L^2(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} \|u_k^\varepsilon\|_{L^2(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} \|\partial_\nu \Psi_\varepsilon\|_{L^\infty(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)},$$

we now analyse each of those norms.

First, by scaling of derivatives we have

$$\begin{aligned} \|\varepsilon \partial_\nu \Psi_\varepsilon\|_{L^\infty(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} &\leq \sup_{|\alpha|=1} \|D^\alpha \psi_\varepsilon\|_{L^\infty(\mathcal{C}^\varepsilon \setminus B(0,1/4))} \\ &\ll \varepsilon^{\frac{1}{2} + \frac{1}{d}}, \end{aligned}$$

where the last bound holds by Lemma 25. We move on to $\|u_k^\varepsilon\|_{L^2(\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)}$. Denote by \tilde{I}^ε the set of indices $\mathbf{n} \in \mathbb{Z}^d$ such that $\partial Q_{\mathbf{n}}^\varepsilon \cap \partial\tilde{\Omega}^\varepsilon \neq \emptyset$. One can see that

$$\|u_k^\varepsilon\|_{L^2(\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} \leq \sum_{\mathbf{n} \in \tilde{I}^\varepsilon} \|U_k^\varepsilon\|_{L^2(\partial Q_{\mathbf{n}}^\varepsilon)}.$$

On the other hand, it follows from Lemma 15 that there is a constant C (in fact, the trace constant of the unit cube) such that

$$\begin{aligned} \sum_{\mathbf{n} \in \tilde{I}^\varepsilon} \|U_k^\varepsilon\|_{L^2(\partial Q_{\mathbf{n}}^\varepsilon)} &\leq C\varepsilon^{-1/2} \sum_{\mathbf{n} \in \tilde{I}^\varepsilon} \|U_k^\varepsilon\|_{H^1(Q_{\mathbf{n}}^\varepsilon)} \\ &\leq C\varepsilon^{-1/2} \|U_k^\varepsilon\|_{H^1(\Omega)}. \end{aligned}$$

Finally, since V is fixed and smooth we have that

$$\begin{aligned} \|V\|_{L^2(\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} &\leq |\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon|^{1/2} \sup_{\mathbf{x}} V(\mathbf{x}) \\ &\leq C|\partial\Omega|^{1/2} \sup_{\mathbf{x}} V(\mathbf{x}). \end{aligned}$$

All in all, this implies that the second term in (42) is bounded by a constant times $\varepsilon^{1/d}$, hence goes to 0 as $\varepsilon \rightarrow 0$, finishing the proof of Lemma 20. \square

We are now ready to complete the proof of the main result of this subsection.

Proof of Proposition 19. We know from [35] that we only need to show that the solutions converge to a solution (Σ, U) of the weak formulation of Problem 3,1 which is that for any test function V ,

$$\int_{\Omega} \nabla U \cdot \nabla V \, d\mathbf{x} = \Sigma \left(A_d \beta \int_{\Omega} UV \, d\mathbf{x} + \int_{\partial\Omega} UV \, dA \right).$$

The weak formulation of the Steklov problem on Ω^ε is that for all test functions V ,

$$\int_{\Omega^\varepsilon} \nabla u_k^\varepsilon \cdot \nabla V \, d\mathbf{x} = \sigma_k^\varepsilon \left(\int_{\partial\Omega} u_k^\varepsilon V \, dA + \int_{\partial T^\varepsilon} u_k^\varepsilon V \, dA \right).$$

The convergence of the gradient terms follows from weak convergence in H^1 of U_k^ε to U and Lemma 12. We already have that $\sigma_k^\varepsilon \rightarrow \Sigma$. The integrals on $\partial\Omega$ converge by weak convergence in H^1 and compactness of the trace operator on $\partial\Omega$. Finally, convergence of the interior term comes from Lemma 20. \square

4.3. Spectral convergence of the problem. We need the following technical lemma.

Lemma 26. *As $\varepsilon \rightarrow 0$, we have that*

$$\tilde{L}_\varepsilon(u_k^\varepsilon) \rightarrow A_d \beta \int_{\Omega} U^2 \, d\mathbf{x}.$$

Remark 27. *Observe that this situation is specific to the sequence u_k^ε . Indeed, there are sequences $\{v_\varepsilon\}$ converging weakly to some $v \in H^1(\Omega)$ such that*

$$\lim_{\varepsilon \rightarrow 0} \tilde{L}_\varepsilon(v_\varepsilon) \neq A_d \beta \int_{\Omega} Uv \, d\mathbf{x}.$$

Proof. From the outer representation (42), we have that

$$\begin{aligned} \tilde{L}_\varepsilon(u_k^\varepsilon) &= \varepsilon \int_{\tilde{\Omega}^\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla (u_k^\varepsilon)^2 \, d\mathbf{x} + \varepsilon \int_{\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} (u_k^\varepsilon)^2 \partial_\nu \Psi_\varepsilon \, d\mathbf{x} + \varepsilon^{-1} c_\varepsilon \int_{\tilde{\Omega}^\varepsilon} (u_k^\varepsilon)^2 \, d\mathbf{x}, \\ &= \varepsilon \int_{\tilde{\Omega}^\varepsilon} 2\nabla \Psi_\varepsilon \cdot u_k^\varepsilon \nabla u_k^\varepsilon \, d\mathbf{x} + \varepsilon \int_{\partial\tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} (u_k^\varepsilon)^2 \partial_\nu \Psi_\varepsilon \, d\mathbf{x} + \varepsilon^{-1} c_\varepsilon \int_{\tilde{\Omega}^\varepsilon} (u_k^\varepsilon)^2 \, d\mathbf{x}. \end{aligned}$$

The last term converges towards the desired $A_d \beta \int_{\Omega} U^2 \, d\mathbf{x}$ once again by strong L^2 convergence of the sequence U_k^ε . To study the first term, let us now introduce the sets

$$\omega^\varepsilon := \bigcup_{\mathbf{k} \in I^\varepsilon} B\left(\varepsilon \mathbf{k}, \frac{\varepsilon}{4}\right) \setminus B(\varepsilon \mathbf{k}, r_\varepsilon) \subset \tilde{\Omega}^\varepsilon,$$

and decompose

$$\varepsilon \int_{\tilde{\Omega}^\varepsilon} \nabla \Psi_\varepsilon \cdot \nabla (u_k^\varepsilon)^2 \, d\mathbf{x} = 2 \left(\int_{\omega^\varepsilon} + \int_{\tilde{\Omega}^\varepsilon \setminus \omega^\varepsilon} \right) \varepsilon \nabla \Psi_\varepsilon \cdot u_k^\varepsilon \nabla u_k^\varepsilon \, d\mathbf{x}.$$

For the integral over ω^ε , we use the L^∞ norm estimate of u_k^ε from Lemma 12 and the scaled H^1 norm estimate for $\varepsilon \nabla \Psi_\varepsilon$ from Lemma 25 and the L^2 boundedness of ∇u_k^ε from Lemma 18 in the generalised Hölder inequality to obtain

$$\begin{aligned} \int_{\omega^\varepsilon} 2\varepsilon \nabla \Psi_\varepsilon \cdot u_k^\varepsilon \nabla u_k^\varepsilon \, d\mathbf{x} &\leq 2 \|\varepsilon \nabla \Psi_\varepsilon\|_{L^2(\Omega^\varepsilon)^d} \|\nabla u_k^\varepsilon\|_{L^2(\Omega^\varepsilon)^d} \|u_k^\varepsilon\|_{L^\infty(\omega^\varepsilon)} \\ &\ll \varepsilon^{\frac{1}{2} + \frac{1}{d}}. \end{aligned}$$

For the integral over $\tilde{\Omega}^\varepsilon \setminus \omega^\varepsilon$, observe that scaling Lemma 25 yields

$$\begin{aligned} \|\varepsilon \nabla \Psi_\varepsilon\|_{L^\infty(\tilde{\Omega}^\varepsilon \setminus \omega^\varepsilon)^d} &= \sup_{|\alpha|=1} \|D^\alpha \psi_\varepsilon\|_{L^\infty(\varepsilon \mathcal{C}^\varepsilon \setminus B(0, 1/4))} \\ &\ll \varepsilon^{\frac{1}{2} + \frac{1}{d}}. \end{aligned}$$

Inserting that bound into the generalised Hölder inequality, along with H^1 boundedness of u_k^ε yields

$$\begin{aligned} \int_{\omega^\varepsilon} 2\varepsilon \nabla \Psi_\varepsilon \cdot u_k^\varepsilon \nabla u_k^\varepsilon \, d\mathbf{x} &\leq 2 \|\varepsilon \nabla \Psi_\varepsilon\|_{L^\infty(\tilde{\Omega}^\varepsilon \setminus \omega^\varepsilon)^d} \|\nabla u_k^\varepsilon\|_{L^2(\Omega^\varepsilon)^d} \|u_k^\varepsilon\|_{L^2(\Omega^\varepsilon)} \\ &\ll \varepsilon^{\frac{1}{2} + \frac{1}{d}}. \end{aligned}$$

Finally, for the integral on the boundary $\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon$, we have from Hölder that

$$\varepsilon \int_{\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon} (u_k^\varepsilon)^2 \partial_\nu \Psi_\varepsilon \, dA \leq \varepsilon \|\partial_\nu \Psi_\varepsilon\|_{L^\infty(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} \|u_k^\varepsilon\|_{L^2(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)}^2 \quad (43)$$

We have as in the proof of Lemma 20 that the L^2 norm of u_k^ε is uniformly bounded, from uniform boundedness of the trace operator. From Lemma 25 we have that

$$\varepsilon \|\partial_\nu \Psi_\varepsilon\|_{L^\infty(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)} = O\left(\varepsilon^{\frac{1}{2} + \frac{1}{d}}\right),$$

and we have that

$$\|u_k^\varepsilon\|_{L^2(\partial \tilde{\Omega}^\varepsilon \setminus \partial T^\varepsilon)}^2 = O(\varepsilon^{-1/2}),$$

implying again that the product in (43) is going to 0 as $\varepsilon \rightarrow 0$, concluding the proof. \square

Until now, we have shown that the harmonic extensions to the holes in Ω^ε of Steklov eigenpairs $(\sigma_k^\varepsilon, u_k^\varepsilon)$ converge weakly in H^1 and strongly in L^2 to a solution (Σ, U) to the problem

$$\begin{cases} -\Delta U = A_d \beta \Sigma U & \text{in } \Omega, \\ \partial_\nu U = \Sigma U & \text{on } \partial \Omega. \end{cases}$$

It remains to be shown that the convergence is to the ‘‘right’’ eigenpair $(\Sigma_{k,\beta}, U_k)$. For the remainder of this section, as β is fixed, we will write simply Σ_k . We first start by the following lemma, showing that the limit function U does not degenerate to the 0 function.

Lemma 28. *Let U be such that $U_k^\varepsilon \rightarrow U$ weakly in $H^1(\Omega)$. Then,*

$$(U, U)_\beta = 1.$$

Proof. By compactness of the trace operator on $\partial\Omega$, we have that

$$\int_{\partial\Omega} (U_k^\varepsilon)^2 \, d\mathbf{x} \rightarrow \int_{\partial\Omega} U^2 \, d\mathbf{x}.$$

as $\varepsilon \rightarrow 0$. Moreover, by Lemma 26, we have that

$$\int_{\partial T^\varepsilon} (U_k^\varepsilon)^2 \, d\mathbf{x} \rightarrow A_d \beta \int_{\Omega} U^2 \, d\mathbf{x}$$

as $\varepsilon \rightarrow 0$. Hence $(U_k^\varepsilon, U_k^\varepsilon)_{\partial^\varepsilon} \rightarrow (U, U)_\beta$. Since U_k^ε has been normalised to $L^2(\partial\Omega^\varepsilon)$ norm 1, this concludes the proof. \square

We are now ready to complete the proof of our first main result.

Proof of Theorem 2. We first show that all the eigenvalues converge. We proceed by induction on the eigenvalue rank k . The case $k = 0$ is trivial. Indeed, we then have that the eigenvalues $\sigma_0^\varepsilon \equiv 0$ obviously converge to $\Sigma_0 = 0$ and the normalised eigenfunctions $U_0^\varepsilon(\mathbf{x}) = |\partial\Omega^\varepsilon|^{-1/2}$, which converges to the constant function

$$U_0(\mathbf{x}) = (|\partial\Omega| + A_d \beta |\Omega|)^{-1/2}.$$

Suppose now that for all $0 \leq j \leq k-1$, we have that U_j^ε converges to U_j weakly in $H^1(\Omega)$. We first show that

$$\Sigma_k \geq \sigma_k^\varepsilon + o(1). \quad (44)$$

In order to do so, we will show that the eigenfunctions U_k are good approximations to appropriate test functions for the variational characterisation (7) of σ_k^ε .

Observe that by compactness of the trace operator on $\partial\Omega$ and by Lemma 20

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial\Omega^\varepsilon} u_j^\varepsilon U_k \, dA = \int_{\partial\Omega} U_k U_j \, dA + A_d \beta \int_{\Omega} U_k U_j \, d\mathbf{x} = 0$$

for all $0 \leq j \leq k-1$. Hence, we can write

$$U_k = V^\varepsilon + \sum_{j=0}^{k-1} \eta_j^\varepsilon u_j^\varepsilon \, dA \quad (45)$$

where for all $0 \leq j \leq k-1$ and all $\varepsilon > 0$,

$$\int_{\partial\Omega^\varepsilon} V^\varepsilon u_j^\varepsilon = 0$$

and $\eta_j^\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$. Now, we have that

$$\begin{aligned} \Sigma_k &= \int_{\Omega} |\nabla U_k|^2 \, d\mathbf{x} \\ &\geq \int_{\Omega^\varepsilon} |\nabla V^\varepsilon|^2 + \sum_{j=0}^{k-1} (\eta_j^\varepsilon)^2 \nabla u_j^\varepsilon \cdot \nabla V^\varepsilon + \sum_{j,l=0}^{k-1} \eta_j^\varepsilon \eta_l^\varepsilon \nabla u_j^\varepsilon \cdot \nabla u_l^\varepsilon \, d\mathbf{x}. \end{aligned}$$

Since the u_j^ε and V^ε are bounded in $H^1(\Omega)$, the integral of the two sums in the previous equation go to 0 as $\varepsilon \rightarrow 0$. On the other hand, we have that for all $\varepsilon > 0$, V^ε is an appropriate test function for σ_k^ε . Hence,

$$\int_{\Omega^\varepsilon} |\nabla V^\varepsilon|^2 \, d\mathbf{x} \geq \sigma_k^\varepsilon \int_{\partial\Omega^\varepsilon} (V^\varepsilon)^2 \, dA.$$

It follows from the decomposition (45) and the fact that by Lemma 20,

$$\int_{\partial\Omega^\varepsilon} U_k^2 \, dA \rightarrow \int_{\partial\Omega} U_k^2 \, dA + A_d \beta \int_{\Omega} U_k^2 \, d\mathbf{x} = 1$$

that

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial\Omega^\varepsilon} (V^\varepsilon)^2 \, d\mathbf{x} = 1,$$

implying that indeed $\Sigma_k \geq \sigma_k^\varepsilon + o(1)$. We now show that

$$\Sigma_k \leq \sigma_k^\varepsilon.$$

Let (Σ, U) be the limit eigenpair for $(\sigma_k^\varepsilon, u_k^\varepsilon)$ and suppose that

$$(\Sigma, U) = (\Sigma_j, U_j)$$

for some $0 \leq j \leq k-1$. We have that

$$\begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} \int_{\partial\Omega^\varepsilon} u_k^\varepsilon \, dA \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\partial\Omega^\varepsilon} u_k^\varepsilon U_j \, dA + \int_{\partial\Omega^\varepsilon} u_k^\varepsilon (u_j^\varepsilon - U_j) \, dA \end{aligned}$$

The first term converges to 1 by the assumption that $(U, U_j)_\beta = 1$ and Lemma 20. As for the second term, we have by the Cauchy–Schwarz inequality and the normalisation $\|u_k^\varepsilon\|_{L^2(\partial\Omega^\varepsilon)} = 1$ that

$$\begin{aligned} \int_{\partial\Omega^\varepsilon} u_k^\varepsilon (u_j^\varepsilon - U_j) \, dA &\leq \|u_j^\varepsilon - U_j\|_{L^2(\partial\Omega^\varepsilon)} \\ &\rightarrow 0 \end{aligned}$$

as $\varepsilon \rightarrow 0$ by Lemma 26. This results in a contradiction. We therefore deduce that σ_k^ε converges to some eigenvalue Σ of problem 3 that is larger than Σ_j for all $j \leq k-1$. Combining this with the bound (44) implies that σ_k^ε converges to Σ_k , and convergence of the eigenfunction therefore follows, when the eigenvalues are simple.

Otherwise, if Σ_k has multiplicity m , *i.e.*

$$\Sigma_{k-1} < \Sigma_k = \dots = \Sigma_{k+m-1} < \Sigma_{k+m},$$

observe that the above argument still yields convergence of σ_j^ε to Σ_j for all $k \leq j < k+m$. For the eigenfunctions, start by fixing a basis U_k, \dots, U_{k+m-1} of the eigenspace associated with Σ_k . Observe that along any subsequence there is a further subsequence such that all the eigenfunctions U_j^ε , converge simultaneously to solutions of Problem 3. Since for all ε , the functions U_j^ε were $L^2(\partial\Omega^\varepsilon)$ -orthogonal, in the limit they are still orthogonal, this time with respect to $(\cdot, \cdot)_\beta$. This implies that in the limit they span the eigenspace associated with Σ_k . As such, the projection on the span of $\{U_j^\varepsilon : k \leq j < k+m\}$ converges to the projections on the span of $\{U_j : k \leq j < k+m\}$. Since this was true along any subsequence, it is also true for the whole sequence, proving convergence of the projections in the sense alluded to in Remark 3. \square

5. Dynamical boundary conditions with large parameter

The goal of this final section is to understand the limit as β becomes large of the eigenvalues $\Sigma_{k,\beta}$ and of the corresponding eigenfunctions $U_{k,\beta}$, normalized by

$$1 = (U_{k,\beta}, U_{k,\beta})_\beta = \int_{\partial\Omega} U_{k,\beta}^2 \, dA + A_d \beta \int_{\Omega} U_{k,\beta}^2 \, d\mathbf{x}.$$

Recall that the Neumann eigenvalues of Ω are

$$0 = \mu_0 \leq \mu_1 \leq \mu_2 \leq \dots \nearrow \infty.$$

We are now ready to prove our second main result.

Proof of Theorem 5. For k fixed, we start by showing that $\beta\Sigma_{k,\beta}$ is bounded. Consider the min–max characterisation of $\Sigma_{k,\beta}$ to obtain

$$\tilde{\Sigma}_{k,\beta} := \beta\Sigma_{k,\beta} = \min_{\substack{E \subset H^1(\Omega) \\ \dim(E)=k+1}} \max_{f \in E \setminus \{0\}} \frac{\int_{\Omega} |\nabla f|^2 \, d\mathbf{x}}{\frac{1}{\beta} \int_{\partial\Omega} f^2 \, dA + A_d \int_{\Omega} f^2 \, d\mathbf{x}}. \quad (46)$$

The quotient on the righthand side of (46) is clearly bounded uniformly in β for any $k+1$ dimensional subspace of smooth functions on Ω . We can therefore suppose that a subsequence in β of $\tilde{\Sigma}_{k,\beta}$ converges, say to $\tilde{\Sigma}_{k,\infty}$. Let us now prove that $\tilde{U}_{k,\beta} := \beta^{1/2}U_{k,\beta}$ is a bounded family in $H^1(\Omega)$. The normalisation on $U_{k,\beta}$ implies that

$$1 \geq A_d \beta \int_{\Omega} U_{k,\beta}^2 \, d\mathbf{x} = A_d \int_{\Omega} \tilde{U}_{k,\beta}^2 \, d\mathbf{x}.$$

For the Dirichlet energy, we have that

$$\int_{\Omega} |\nabla \tilde{U}_{k,\beta}|^2 \, d\mathbf{x} = \beta\Sigma_{k,\beta},$$

which was already shown to be bounded. Therefore there is also a weakly convergent subsequence in $H^1(\Omega)$ as $\beta \rightarrow \infty$, converging to say $\tilde{U}_{k,\infty}$. We take the subsequence to coincide with the one for $\tilde{\Sigma}_{k,\beta}$. Observe as well that the normalisation condition on $U_{k,\beta}$ prevents the limit $\tilde{U}_{k,\infty}$ from vanishing identically.

The functions $\tilde{U}_{k,\beta}$ satisfy the following weak variational characterisation for any element $V \in H^1(\Omega)$

$$\int_{\Omega} \nabla \tilde{U}_{k,\beta} \cdot \nabla V \, d\mathbf{x} = \tilde{\Sigma}_{k,\beta} \left(\beta^{-1} \int_{\partial\Omega} \tilde{U}_{k,\beta} V \, dA + A_d \int_{\Omega} \tilde{U}_{k,\beta} V \, d\mathbf{x} \right).$$

Letting $\beta \rightarrow \infty$, weak convergence of $\tilde{U}_{k,\beta}$ in $H^1(\Omega)$ implies that the limit satisfies the weak identity

$$\int_{\Omega} \nabla \tilde{U}_{k,\infty} \cdot \nabla V \, d\mathbf{x} = \tilde{\Sigma}_{k,\infty} A_d \int_{\Omega} \tilde{U}_{k,\infty} V \, d\mathbf{x}.$$

In other words, $\tilde{U}_{k,\infty}$ is a solution to the Neumann eigenvalue problem with eigenvalue $\mu = \tilde{\Sigma}_{k,\infty} A_d$.

We now proceed by recursion on k to show convergence to the right eigenpair. Once again, the statement is trivial for $k=0$ and the constant eigenfunction. Assume that we have convergence for the first $k-1$ eigenpairs. We now proceed in a similar fashion as in the proof of the spectral convergence to Problem (3). We repeat the argument because the inequalities are more subtle. We first show that

$$\tilde{\mu}_k(\Omega) := \frac{\mu_k(\Omega)}{A_d} \geq \tilde{\Sigma}_{k,\beta}(1 + o(1)). \quad (47)$$

Write

$$f_k = F_{\beta} + \sum_{j=0}^{k-1} (f_k, U_{j,\beta})_{\beta} U_{j,\beta},$$

with $F_{\beta} \perp_{\beta} U_{j,\beta}$ for $0 \leq j < k$. We have that

$$(f_k, U_{j,\beta})_{\beta} = (f_k, \beta^{-1/2} f_j)_{\beta} + (f_k, U_{j,\beta} - \beta^{-1/2} f_j)_{\beta}. \quad (48)$$

The first inner product develops as

$$(f_k, \beta^{-1/2} f_j)_\beta = \beta^{-1/2} \int_{\partial\Omega} f_k f_j \, dA + A_d \beta^{1/2} \int_{\Omega} f_k f_j \, d\mathbf{x}.$$

The first term clearly goes to 0 as $\beta \rightarrow \infty$, and the second one vanishes by orthogonality of the Neumann eigenfunctions in $L^2(\Omega)$. We now turn our attention to the second inner product in (48). We have that

$$\lim_{\beta \rightarrow \infty} \beta^{-1/2} \int_{\partial\Omega} f_k (\tilde{U}_{j,\beta} - f_j) \, dA = 0$$

by weak H^1 convergence of $\tilde{U}_{j,\beta}$ and compactness of the trace operator. On the other hand, strong L^2 convergence implies that

$$A_d \beta^{1/2} \int_{\Omega} f_k (\tilde{U}_{j,\beta} - f_j) \, d\mathbf{x} = o(\beta^{1/2}). \quad (49)$$

All in all, this implies that

$$(f_k, U_{j,\beta})_\beta = o(\beta^{1/2})$$

for all $0 \leq j < k$. We now write

$$\begin{aligned} \tilde{\mu}_k &= \frac{1}{A_d} \int_{\Omega} |\nabla f_k|^2 \, d\mathbf{x} \\ &\geq \frac{1}{A_d} \int_{\Omega} |\nabla F_\beta|^2 \, d\mathbf{x} - \frac{1}{A_d} \sum_{j=0}^{k-1} (f_k, U_{j,\beta})_\beta^2 \Sigma_{j,\beta}. \end{aligned} \quad (50)$$

Since $\beta \Sigma_{j,\beta}$ is bounded and equation (49) implies that $(f_k, U_{j,\beta})_\beta^2 = o(\beta)$, we deduce that the last term in (50) goes to 0. We now observe that by the variational characterisation of $\Sigma_{k,\beta}$,

$$\begin{aligned} \frac{1}{A_d} \int_{\Omega} |\nabla F_\beta|^2 \, d\mathbf{x} &\geq \frac{\Sigma_{k,\beta}}{A_d} \left(\int_{\partial\Omega} F_\beta^2 + A_d \beta \int_{\Omega} F_\beta^2 \right), \\ &= \tilde{\Sigma}_{k,\beta} \left(\frac{1}{A_d \beta} \int_{\partial\Omega} F_\beta^2 \, dA + \int_{\Omega} F_\beta^2 \, d\mathbf{x} \right). \end{aligned}$$

In the same way as we obtained the bound on the last term in (50), the first integral is $o(\beta)$. As for the second one, we write

$$\begin{aligned} \int_{\Omega} F_\beta^2 \, d\mathbf{x} &= \int_{\Omega} \left(f_k - \sum_{j=0}^{k-1} (f_k, U_{j,\beta})_\beta U_{j,\beta}^2 \right)^2 \, d\mathbf{x} \\ &= 1 - \sum_{j=0}^{k-1} \int_{\Omega} (f_k, U_{j,\beta})_\beta f_j U_{j,\beta} \, d\mathbf{x} + \sum_{j,\ell=0}^{k-1} \int_{\Omega} (f_k, U_{j,\beta})_\beta (f_k, U_{\ell,\beta})_\beta U_{j,\beta} U_{\ell,\beta} \, d\mathbf{x}. \end{aligned}$$

Strong $L^2(\Omega)$ convergence of $\beta^{1/2} U_{j,\beta}$ and the fact that $(f_k, U_{j,\beta}) = o(\beta^{1/2})$ imply that the last two integrals converge to 0 as $\beta \rightarrow \infty$.

We have therefore obtained that

$$\tilde{\mu}_k(\Omega) \geq \tilde{\Sigma}_{k,\beta} (1 + o(1)),$$

i.e. we have indeed proven assertion (47).

Suppose now that $(\tilde{\Sigma}_{k,\beta}, \tilde{U}_{k,\beta})$ converge to a Neumann eigenpair $(\tilde{\mu}_j, f_j)$ for some $j < k$. Then, we have that

$$\begin{aligned} 1 &= \lim_{\beta \rightarrow \infty} \int_{\Omega} f_j \tilde{U}_{k,\beta} \, d\mathbf{x} \\ &= \lim_{\beta \rightarrow \infty} \int_{\Omega} (f_j - \tilde{U}_{j,\beta}) \tilde{U}_{k,\beta} \, d\mathbf{x} + \int_{\Omega} \tilde{U}_{k,\beta} U_{j,\beta} \, d\mathbf{x} \\ &\leq \lim_{\beta \rightarrow \infty} \|f_j - \tilde{U}_{j,\beta}\|_{L^2(\Omega)} \|\tilde{U}_{k,\beta}\|_{L^2(\Omega)} + \left| \int_{\Omega} U_{k,\beta} U_{j,\beta} \, d\mathbf{x} \right| \\ &= 0, \end{aligned}$$

where the limit comes from strong convergence in $L^2(\partial\Omega)$ to 0 of $U_{j,\beta}$ and our recursion hypothesis. This is a contradiction, hence the convergence is to the correct eigenpair. This implies convergence of the whole sequence if the Neumann spectrum of Ω is simple. If there are eigenvalues with multiplicity, the same procedure as for the homogenisation problem yields once again convergence.

As for continuity in β , the same proof goes through in exactly the same way, except for the fact that we do not need to show the boundedness results in β . \square

We can now prove the comparison results between Steklov and Neumann eigenvalues.

Proof of Corollary 7. It is proved in [10, Theorem 1.4] that any bounded domain $\Omega \subset \mathbb{R}^d$ satisfies

$$\sigma_k(\Omega) |\partial\Omega| \leq C(d) |\Omega|^{\frac{d-2}{d}} k^{2/d},$$

where $C(d)$ is a constant which depends only on the dimension. When applied to Ω^ε this leads, after taking $\varepsilon \rightarrow 0$, to

$$\Sigma_{k,\beta} (|\partial\Omega| + A_d \beta |\Omega|) \leq C(d) |\Omega|^{\frac{d-2}{d}} k^{2/d}.$$

Taking the limit $\beta \rightarrow \infty$ leads to

$$\mu_k |\Omega| \leq C(d) |\Omega|^{\frac{d-2}{d}} k^{2/d}.$$

This is equivalent to

$$\mu_k |\Omega|^{2/d} \leq C(d) k^{2/d}.$$

\square

Finally, all is left to do is to prove the previous theorem has the following corollary in dimension $d = 2$. It allows one to transform universal bounds for Steklov eigenvalues into universal bounds for Neumann eigenvalues.

We write Ω_β^ε for a domain Ω^ε as constructed earlier whose holes are exactly of radius $r_\varepsilon^{d-1} = \beta \varepsilon^d$.

Proof of Theorem 9. We have from Theorem 2 that

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \sigma_k \left(\Omega_\beta^\varepsilon \right) \left| \partial\Omega_\beta^\varepsilon \right| &= \Sigma_{k,\beta}(\Omega) (|\partial\Omega| + A_d \beta |\Omega|) \\ &= \frac{|\partial\Omega|}{\beta} \tilde{\Sigma}_{k,\beta} + A_d |\Omega| \tilde{\Sigma}_{k,\beta}. \end{aligned}$$

Now, the first term clearly goes to 0 as $\beta \rightarrow \infty$ while, by Theorem 5, we have that

$$\lim_{\beta \rightarrow \infty} A_d |\Omega| \tilde{\Sigma}_{k,\beta} = \mu_k(\Omega) |\Omega|.$$

□

Acknowledgements. The authors would like to thank Iosif Polterovich for several very useful conversations. We would also like to thank Valeri Smyshlyaev and Ilya Kamot-ski for pointing us towards many classical results in homogenisation theory. We would also like to thank Almut Burchard for possible further related questions. The authors also would like to thank Gérard Philippin, who was helpful in the early stage of this project. A.G. acknowledges support from NSERC. A.H. wants also to thank the Centre de Recherche Mathématiques de Montréal and the Département de Mathématiques of Université Laval for grants and fruitful atmosphere during his stay there in 2018. J.L. was supported by the NSERC postdoctoral fellowship, and EPSRC grant number EP/P024793/1.

REFERENCES

- [1] G. Allaire. *Shape optimization by the homogenization method*, volume 146 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2002.
- [2] J. Arrieta, Á. Jiménez-Casas, and A. Rodríguez-Bernal. Flux terms and Robin boundary conditions as limit of reactions and potentials concentrating at the boundary. *Rev. Mat. Iberoam.*, 24(1):183–211, 2008.
- [3] D. Bucur, P. Freitas, and J. Kennedy. The Robin problem. In *Shape optimization and spectral theory*, pages 78–119. De Gruyter Open, Warsaw, 2017.
- [4] D. Bucur, A. Giacomini, and P. Trebeschi. L^∞ bounds of Steklov eigenfunctions and spectrum stability under domain variations. cvgmt preprint.
- [5] I. Chavel. *Isoperimetric inequalities*, volume 145 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2001. Differential geometric and analytic perspectives.
- [6] K. Cherednichenko, P. Dondl, and F. Rösler. Norm-resolvent convergence in perforated domains. *Asymptot. Anal.*, 110(3-4):163–184, 2018.
- [7] V. Chiado Piat, S. S. Nazarov, and A. L. Piatnitski. Steklov problems in perforated domains with a coefficient of indefinite sign. *Netw. Heterog. Media*, 7(1):151–178, 2012.
- [8] D. Cianci and A. Girouard. Large spectral gaps for Steklov eigenvalues under volume constraints and under localized conformal deformations. *Ann. Global Anal. Geom.*, 54(4):529–539, 2018.
- [9] D. Cioranescu and F. Murat. Un terme étrange venu d’ailleurs. In *Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. II (Paris, 1979/1980)*, volume 60 of *Res. Notes in Math.*, pages 98–138, 389–390. Pitman, Boston, Mass.-London, 1982.
- [10] B. Colbois, A. El Soufi, and A. Girouard. Isoperimetric control of the Steklov spectrum. *J. Funct. Anal.*, 261(5):1384–1399, 2011.
- [11] B. Colbois, A. El Soufi, and A. Girouard. Isoperimetric control of the spectrum of a compact hypersurface. *J. Reine Angew. Math.*, 683:49–65, 2013.
- [12] B. Colbois, A. El Soufi, and A. Girouard. Compact manifolds with fixed boundary and large Steklov eigenvalues. *Proc. Amer. Math. Soc.*, 147(9):3813–3827, 2019.
- [13] H. Douanla. Homogenization of Steklov spectral problems with indefinite density function in perforated domains. *Acta Appl. Math.*, 123:261–284, 2013.
- [14] Lawrence C. Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. Textbooks in Mathematics. CRC Press, Boca Raton, FL, revised edition, 2015.
- [15] A. Fraser and R. Schoen. Minimal surfaces and eigenvalue problems. In *Geometric analysis, mathematical relativity, and nonlinear partial differential equations*, volume 599 of *Contemp. Math.*, pages 105–121. Amer. Math. Soc., Providence, RI, 2013.
- [16] A. Fraser and R. Schoen. Sharp eigenvalue bounds and minimal surfaces in the ball. *Invent. Math.*, 203(3):823–890, 2016.
- [17] P. Freitas and R. S. Laugesen. From Neumann to Steklov and beyond, via Robin: the Weinberger way. *Amer. J. Math.*, to appear.
- [18] P. Freitas and R. S. Laugesen. From Steklov to Neumann and beyond, via Robin: the Szegő way. *Canad. J. Math.*, to appear.

- [19] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [20] A. Girouard and I. Polterovich. Spectral geometry of the Steklov problem (survey article). *J. Spectr. Theory*, 7(2):321–359, 2017.
- [21] S. Gryshchuk and M. Lanza de Cristoforis. Simple eigenvalues for the Steklov problem in a domain with a small hole. A functional analytic approach. *Math. Methods Appl. Sci.*, 37(12):1755–1771, 2014.
- [22] V. V. Jikov, S. M. Kozlov, and O. A. Oleinik. *Homogenization of differential operators and integral functionals*. Springer-Verlag, Berlin, 1994. Translated from the Russian by G. A. Yosifian [G. A. Iosif'yan].
- [23] S. Kaizu. The Robin problems on domains with many tiny holes. *Proc. Japan Acad. Ser. A Math. Sci.*, 61(2):39–42, 1985.
- [24] P. Kröger. Upper bounds for the Neumann eigenvalues on a bounded domain in Euclidean space. *J. Funct. Anal.*, 106(2):353–357, 1992.
- [25] P.-D. Lamberti and L. Provenzano. Viewing the Steklov eigenvalues of the Laplace operator as critical Neumann eigenvalues. In *Current trends in analysis and its applications*, Trends Math., pages 171–178. Birkhäuser/Springer, Cham, 2015.
- [26] P.-D. Lamberti and L. Provenzano. Neumann to Steklov eigenvalues: asymptotic and monotonicity results. *Proc. Roy. Soc. Edinburgh Sect. A*, 147(2):429–447, 2017.
- [27] V. A. Marčenko and E. Ya. Khruslov. Boundary-value problems with fine-grained boundary. *Mat. Sb. (N.S.)*, 65 (107):458–472, 1964.
- [28] V. A. Marčenko and E. Ya. Khruslov. *Boundary value problems in domains with a fine-grained boundary*. Izdat. Naukova Dumka, Kiev, 1974.
- [29] G. Dal Maso, G. Franzina, and D. Zucco. Transmission conditions obtained by homogenisation. *Nonlinear Analysis*, 177:361 – 386, 2018. Nonlinear PDEs and Geometric Function Theory, in honor of Carlo Sbordone on his 70th birthday.
- [30] S. A. Nazarov. Asymptotic expansions of eigenvalues of the Steklov problem in singularly perturbed domains. *Algebra i Analiz*, 26(2):119–184, 2014.
- [31] J. Rauch and M. E. Taylor. Potential and scattering theory on wildly perturbed domains. *J. Funct. Anal.*, 18:27–59, 1975.
- [32] M. A. Shubin. *Pseudodifferential operators and spectral theory*. Springer-Verlag, Berlin, second edition, 2001. Translated from the 1978 Russian original by Stig I. Andersson.
- [33] M. E. Taylor. *Partial differential equations I. Basic theory*, volume 115 of *Applied Mathematical Sciences*. Springer, New York, second edition, 2011.
- [34] M. Vanninathan. Homogenization of eigenvalue problems in perforated domains. *Proc. Indian Acad. Sci. Math. Sci.*, 90(3):239–271, 1981.
- [35] J. von Below and G. François. Spectral asymptotics for the Laplacian under an eigenvalue dependent boundary condition. *Bull. Belg. Math. Soc. Simon Stevin*, 12(4):505–519, 2005.
- [36] S. Zelditch. *Eigenfunctions of the Laplacian on a Riemannian Manifold*. CBMS Regional Conference Series in Mathematics. Conference Board of the Mathematical Sciences, 2017.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, PAVILLON ALEXEANDRE-VACHON, UNIVERSITÉ LAVAL,
QUÉBEC, QC, G1V 0A6, CANADA

Email address: alexandre.girouard@mat.ulaval.ca

UNIVERSITÉ DE LORRAINE, CNRS, INSTITUT ELIE CARTAN DE LORRAINE, 54506 VANDŒUVRE-LÈS-NANCY
CEDEX, FRANCE

Email address: antoine.henrot@univ-lorraine.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, WC1E 6BT,
UNITED KINGDOM

Email address: j.lagace@ucl.ac.uk