



**HAL**  
open science

## Overweight in the pluri-ethnic adolescent population of New Caledonia: Dietary patterns, sleep duration and screen time

Stephane Frayon, Guillaume Wattelez, Emilie Paufigue, Akila Nedjar-Guerre, Christophe Serra-Mallol, Olivier Galy

### ► To cite this version:

Stephane Frayon, Guillaume Wattelez, Emilie Paufigue, Akila Nedjar-Guerre, Christophe Serra-Mallol, et al.. Overweight in the pluri-ethnic adolescent population of New Caledonia: Dietary patterns, sleep duration and screen time. *The Lancet Regional Health - Western Pacific*, 2020, 2, 10.1016/j.lanwpc.2020.100025 . hal-02970510

**HAL Id: hal-02970510**

**<https://hal.science/hal-02970510>**

Submitted on 14 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

**Overweight in the pluri-ethnic adolescent population of New Caledonia:  
dietary patterns, sleep duration and screen time**

Stéphane Frayon<sup>1</sup>, Guillaume Wattelez<sup>1</sup>, Emilie Paufique<sup>1</sup>, Akila Nedjar-Guerre<sup>1</sup>, Christophe Serra-Mallol<sup>2</sup>,  
Olivier Galy<sup>1</sup>

<sup>1</sup>Interdisciplinary Laboratory for Research in Education, EA 7483, School of Education, University of New Caledonia, New Caledonia

<sup>2</sup>Centre on Work Organizations and Policies (CERTOP), UMR 5044 CNRS - University of Toulouse Jean Jaurès, 5 allées Antonio Machado, 31058 Toulouse cedex 9, France

Address correspondence to: Dr Stéphane Frayon, BP R4, Avenue James Cook, 98851 Noumea Cedex, New Caledonia. Email: [stephanefrayon@hotmail.com](mailto:stephanefrayon@hotmail.com), ORCID ID: 0000-0001-9767-4148

## **Abstract**

### **Background**

A high prevalence of overweight and obesity has been found in adolescents of New Caledonia and other Pacific Island Countries and Territories. Although Westernization may contribute to the weight gain in populations of Oceanian, Non-European, Non-Asian ancestry (ONENA), little is known about the sociodemographic and lifestyle factors associated with overweight in the Melanesian and Polynesian adolescents of New Caledonia.

### **Methods**

In this cross-sectional study, a pluri-ethnic sample of New Caledonian adolescents (N = 954; age M = 13.2 years) completed a survey to estimate sleep duration, screen time, and dietary pattern using a food frequency questionnaire. Demographic data (gender, ethnicity, socioeconomic status: SES, area of residence) were collected, and anthropometric measures were used to compute weight status.

### **Findings**

We found a higher risk for being overweight in Melanesian (OR = 1.67) and Polynesian (OR = 5.40) adolescents compared with European adolescents, even after controlling for age, SES, area of residence, dietary pattern, sleep duration and screen time. We also found that low SES (OR = 3.43) and sleep duration (OR = .65 per hour) were independently associated with overweight status in the European but not in ONENA adolescents.

### **Interpretation**

In this study, the main contribution to being overweight was ethnic background, i.e. being Melanesian or Polynesian. The hypothesis of a genetic influence thus seems attractive and merits further analyses.

### **Funding**

This project was funded by the University of New Caledonia and the Fondation Nestlé France.

**Keywords:** Obesity; Melanesian; Polynesian; Lifestyle; Pacific; nutrition; food frequency; ethnicity; Westernization; rural; urban.

## Introduction

The prevalence of obesity is high in both adults and adolescents of the Pacific Island Countries and Territories (PICTs).<sup>1-3</sup> As in other countries, overweight in Pacific adolescents has been associated with gender (being female), sugar sweetened beverage consumption, sedentary behaviours<sup>4</sup> and sociocultural influences.<sup>5,6</sup> Overweight and obesity have been correlated with several noncommunicable diseases, such as type 2 diabetes, cardiovascular events, cancer and hypertension.<sup>7-9</sup> Moreover, overweight also seems to be associated with psychological issues like depression,<sup>10-13</sup> body dissatisfaction,<sup>14,15</sup> disordered eating behaviours,<sup>16</sup> low self-esteem,<sup>14,17-21</sup> and insufficient sleep duration.<sup>22,23</sup>

New Caledonia is a French archipelago in the South Pacific whose pluri-ethnic population has been affected by obesity and the issues surrounding it. The lifestyles of the New Caledonian populations differ greatly among the main ethnic groups, which are distinguished both culturally and socioeconomically.<sup>24</sup> The Melanesians (45% of the population) were the first inhabitants before the colonial era. Today, some of them still follow the traditional Pacific lifestyle, whereas others have also taken on a more Western lifestyle while still staying close to traditional customs. Other communities of European and Polynesian origins (31% and 12%, respectively) have adopted a frankly Western way of life, even though all Polynesian communities still have different cultural practices than Europeans. A high prevalence of overweight has been noted in New Caledonian adolescents,<sup>25,26</sup> with a higher prevalence in Melanesian and Polynesian adolescents.<sup>24</sup> However, to date, little is known about the factors associated with overweight in these adolescents.

The major cause of overweight/obesity is often a long-term mismatch between energy intake and expenditure.<sup>27,28</sup> This mismatch has become more prominent today in relation with lifestyle changes, which have been striking in populations of Oceanian, Non-European, Non-Asian (ONENA) ancestry.<sup>29</sup> Indeed, unhealthy eating habits, physical inactivity, screen watching and sleeping patterns may contribute to the current obesity epidemic in PICTs,<sup>30</sup> but these trends have never been studied in New Caledonian adolescents.

It has been well documented that dietary energy intake is one of the major contributors to positive energy balance in children and adolescents.<sup>31</sup> In PICTs, traditional foods like yams, taro, fish, and indigenous fruits and vegetables have been replaced by imported sugar, soft drinks, rice, canned foods and calorie-dense snack foods.<sup>6,32–35</sup> Adding Western foods to the traditional diet can contribute to an increased risk of obesity even in rural areas where lifestyles and diets have remained largely traditional.<sup>36</sup> More generally, dietary change was observed in five Pacific Island nations between 1961 and 2000 and was found to consist of greater food and energy intake with a notable increase in the consumption of fatty foods and meat,<sup>34</sup> which may be correlated with the BMI increase observed in the same period. Peer influence and greater autonomy about food choices in adolescence may accentuate the unhealthy diets and weight gain in this group.<sup>37,38</sup>

Modernization may also contribute to weight gain, as it has meant easier access to technologies like televisions (TV), personal computers (PC), smartphones and game consoles. Higher screen time has been linked to higher weight status, probably due to decreased energy expenditure, more physical inactivity, increased caloric intake while using screen devices, and increased food-advertising exposure.<sup>39–41</sup> In Vanuatu, the increased measures of adiposity and risk for obesity were associated with ownership of electronic devices.<sup>42</sup> A shift in adolescents' preferences for devices seems to have occurred recently, and the use of smartphones and small connected devices is rising.<sup>43,44</sup> As a consequence, adolescents owning a smartphone are more likely to have high screen time compared with non-owner adolescents.<sup>45</sup> In the Pacific, Maori children were 1.3 times and Pacific children were 1.1 times more likely to watch over 2 hours of television per day compared with non-Maori and non-Pacific children.<sup>46</sup> Moreover, the association between high fatness levels and high TV watching was found in Maori and Pacific Island youths in New Zealand.<sup>47</sup>

In addition to the putative effect of screen time on weight status, screen use was also associated with shorter sleep duration.<sup>48</sup> Short sleep duration is also a risk factor for obesity, and the relationship between high screen time exposure and the more frequent consumption of foods high in fat, free sugars or salt has been demonstrated.<sup>49</sup> To date, the exact mechanism through which sleep duration affects overweight/obesity is not completely understood. Little is known about adolescent sleep duration in the South Pacific

population, with most of the studies conducted in New Zealand. Among New Zealander adults,<sup>50,51</sup> children,<sup>52</sup> and adolescents,<sup>53</sup> the Maori and Pacific Island populations showed shorter sleep durations than the Europeans.

Given that the prevalence of overweight is much higher in the Polynesian and Melanesian adolescents of New Caledonia, we hypothesized that differences would be found between these groups and the adolescents with an European background, who showed a lower prevalence of overweight. The aim of this study was to look for putative links between overweight/obesity and sleep duration, dietary patterns and screen use in the pluri-ethnic adolescent population of New Caledonia.

## **Materials and methods**

The sample for this study consisted in 954 adolescents (488 boys, 466 girls) recruited in New Caledonia. Participant ages ranged from 10.5 to 16.1 years ( $M = 13.22$ ,  $SD = 1.23$ ). Full descriptive information about the sample is presented in Table S1 and Table S2.

Ethics approval for this project was obtained from the Consultative Ethics Committee of New Caledonia (CCE 2018-06 001) and the project was conducted in accordance with the requirements of the Declaration of Helsinki. Eight secondary public schools were randomly selected to obtain a representative repartition between rural and urban areas, which is 63% and 37%, respectively, in New Caledonia. The selection criterion was school size ( $N > 150$ ) to ensure sufficient data in a single field trip. One or two classes were then randomly selected in each of four grades (levels) by a staff member, for a total of approximately 150 adolescents (6 groups with a mean of 25 students per division). We obtained ~~only~~ 89% of the expected data due to parental refusal, no return of the parental agreement or absence of the participants on the days of intervention. Participants completed two anonymous online questionnaires consisting of the measures described above and additional measures on body image or self-esteem, which are not reported here. Data were collected over two consecutive days in order not to disrupt the students' schedules too much. There were missing data because some participants were absent on at least one of the two consecutive days or because they used a wrong anonymous number. Adolescents with missing data ( $N = 108$ ; 10.2%) were then

excluded from analyses (Figure S1). Parents gave informed written consent prior to the adolescents' participation in the study, and all participants received written debriefing information at the end of the study.

In the absence of a validated food frequency questionnaire (FFQ) for New Caledonian adolescents, we used a short FFQ previously used for Aboriginal and Torres Strait Islanders.<sup>54</sup> This FFQ was adopted from another tool developed in Australia to assess dietary intake in children and adolescents up to 16 years old.<sup>55</sup> The short questions on this FFQ are able to discriminate between different categories of food intake and provide information on relative intakes.<sup>54</sup> Moreover, this FFQ has been validated in a pluri-ethnic population composed with non-indigenous Australians (with European background) and Torres Strait Islanders (with a Melanesian background), and it was previously used in New Caledonia.<sup>56</sup> Minor modifications were made by the research team to include foods identified as important in the Melanesian diet. For example, tubers like cassava, yams and taro are consumed instead of white potatoes. The FFQ contains 28 questions on food and beverage intake. The intake frequencies for the individual food items of the FFQ were first converted to daily frequency equivalents (DFEs), calculated by allocating proportional values to the original frequency categories with reference to a base value of 1.0, equivalent to once a day<sup>57</sup>. The scores were calculated as follows: DFE score of 0: never ; .07: less than once per week; .14: once per week; 0.21: 1-2 times per week; 0.29: 1-3 times per week; .36: 2–3 times per week; .50: 3-4 times per week; .71: 4-6 times per week; .79: 5-6 times per week; 1: once a day; 2: 2 times per day; 2.5: 2-3 times per day; 4.5: 4-5 times per day; and 6: more than 6 times per day. The questions were asked on tablets distributed by the team and sometimes computerized in class so that the answers were automatically recorded in the database.

The sleep duration was determined with the following four questions: “What time do you fall asleep on school days?”, “What time do you fall asleep on the weekend?”, “What time do you wake up in the school week?” and “What time do you wake up on the weekend?” There were 13 available categories for the time an adolescent might fall asleep, from “Around 9 pm or before” to “Around 3 am or later” with a 30-minute interval between each category. There were 15 available categories for the wake-up time from “Around 5 am or before” to “Around midday or later” with a 30-minute interval between each category. Answers were converted to numerical values by using the median value of the time interval in the



categorized answer or by using 30 minutes before (respectively, after) for the first (respectively, the last) category. The final sleep duration was the difference between the wake-up time and the falling-asleep time. Answers about sleep duration during the school week and the weekend were first separately processed, and then both factors were combined to get a total sleep duration for the full week as follows:  $\text{Mean Sleep} = 5 \times \text{Sleep (Weekdays)} + 2 \times \text{Sleep (Weekend)}$  divided by 7.

The screen time was determined with the following two questions: “How much time do you spend on a computer or tablet on the weekend?” and “How much time do you spend on computer or tablet during the school day?” There were four available categories for the amount of time an adolescent might use a computer, laptop or tablet screen on school days from “Less than one hour” to “More than 3 hours”. There were six available categories for the time an adolescent might use a computer, laptop or tablet screen on weekends from “Less than one hour” to “More than 5 hours”. Answers were converted to numerical values by using the median value of the time interval in the categorized answer or by using 30 minutes before for the first category or 30 minutes after for the last category. Answers about screen time during the school week and the weekend were first separately processed, and then both factors were combined to get a total screen time during the full week as follows:  $\text{Mean screen time use} = 5 \times \text{screen use (Weekdays)} + 2 \times \text{screen use (Weekend)}$  divided by 7. The same procedure was used for smartphone use with two questions: “If you have a mobile phone, how long are you connected to your phone during a day of the week?” and “If you have a mobile phone, how long are you connected to your phone during a day of the weekend?”

Ethnicity was self-reported by the adolescents and categorized as recommended by the INSERM report on New Caledonia,<sup>58</sup> with the exception that we did not allow participants to choose more than one ethnic group. Participants from minor ethnic groups (approximately 5% of the whole sample) were then clustered in the category “Other”.

Socioeconomic status (SES) was indexed on the basis of the occupation of the household reference person (defined as the householder with the highest income) using the National Statistics Socio-Economic Classification.<sup>59</sup> In this study, three categories were retained: managerial and professional occupations (high SES), intermediate occupations (intermediate SES), and routine and manual occupations (low SES).

Based on data from the last census in New Caledonia,<sup>60</sup> area of residence was determined using a European standard.<sup>61</sup> Densely populated areas comprising at least 50,000 inhabitants in a continuous zone with more than 500 inhabitants per km<sup>2</sup> were classified as urban. Areas with more than 50,000 inhabitants in a continuous area of over 100 inhabitants per km<sup>2</sup> or areas adjacent to an urban area were classified as semi-urban. Rural areas were defined as those areas that did not fulfil the conditions required to qualify as urban or semi-urban.

All anthropometric measurements were recorded by a member of the research team (trained staff). Height was measured to the nearest 0.1 cm using a portable stadiometer (Leicester Tanita HR 001, Tanita Corporation, Tokyo, Japan). Weight was determined using a scale (Tanita HA 503, Tanita Corporation) to the nearest 0.1 kg, with adolescents weighed in light clothing. Body mass index (BMI) was then calculated by dividing weight (in kilograms) by height (in meters) squared.

For weight status classifications, three BMI-based definitions were used: from the International Obesity Task Force: IOTF BMI-based references;<sup>62</sup> the World Health Organization: WHO BMI-based references;<sup>63</sup> and the French BMI-based references.<sup>64</sup> The BMI z-scores (BMIz) were calculated using the LMS reference values for the IOTF BMI-based classification,<sup>65</sup> because this reference showed the best relationship with body fat in New Caledonian adolescents.<sup>25</sup>

All analyses were conducted using IBM SPSS v.26, with *p*-values < .05 indicating statistical significance. For preliminary demographic analysis, gender differences were assessed with independent-samples t-tests (continuous variables) and chi-squared tests (categorical variables).

The principal component analysis (PCA) was run with all 24 food groups entered as raw frequencies using varimax orthogonal rotation. Factors were retained on the basis of eigenvalues > 1.0, a scree plot of the eigenvalues, and the interpretability of components. Variables with factor loadings  $\geq .4$  and  $\leq -.4$  were considered significant when naming patterns. The patterns produced were transformed to remove skew and converted to z-scores, so factor scores had a standardised mean of zero. Data factorability was assessed using the Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy and Bartlett's test of sphericity. Cronbach's alpha coefficient was used to determine the internal consistency. All participants received a

score for all five patterns, with positive scores indicating high consumption of foods associated with that pattern and negative scores indicating lower consumption. For example, the higher the Fruits and vegetables score, the higher an individual's frequency of consuming fruits, vegetables, tubers, soup-and fish.

To determine which factors were significantly associated with each dietary pattern, we computed linear mixed models with each dietary pattern score as the criterion variable. To account for intra-class correlation, school were included and treated as random effects in the linear mixed models with no predictors (null model). Results from this model were used to calculate the intra-class correlation coefficient (ICC), which is the proportion of between-school variance to total variance. Second, independent standardized variables (gender, ethnicity, SES, area of residence, BMIz and age) were added to the models as covariables and treated as fixed effects. Ethnicity, gender, area of residence, and SES were previously categorized into groups by creating dummy variables.

Differences in BMIz, sleep duration and screen time were assessed using a 4 x 2 analysis of variance (ANOVA), with ethnicity and gender treated as independent variables, and post-hoc Tukey tests used to follow-up significant differences. Linear mixed models were the used to determine the factors significantly associated with sleep duration or screen time using the method described above.

Last, we used multivariable logistic regression analyses to identify the factors associated with overweight. The variables in the models were age, dietary pattern scores, screen use and sleep duration for continuous variables and SES (low, intermediate and high), ethnicity (European, Melanesian, Polynesian and Other), gender (boy and girl) and area of residence (rural and urban) for categorical variables.

To take into account a possible selection bias because of the missing data for the participants, we also ran sensitivity analysis using varying sample sizes to ensure the robustness of the analyses. Because no measure had no more than 5% of the data missing (Figure S2), we did not perform any imputation methods.

### **Role of funding Source**

The funding sources (University of New Caledonia and the Fondation Nestlé France) had no role in study design, data collection, data analysis, interpretation and writing of the report.

## Results

### Sociodemographic and anthropometric characteristics

The study population comprised 954 adolescents between 10.5 and 16.1 years old, with 51.2% boys ( $13.2 \pm 1.2$  years) and 48.8% girls ( $13.3 \pm 1.2$  years). The boys and girls did not significantly differ regarding age, but the boys were significantly taller and heavier than the girls (Table S1). No significant difference was found concerning ethnicity, SES or area of residence repartition according to gender.

When the data were stratified by ethnicity, no difference regarding age or gender repartition was found (Table S2). However, ethnic subgroups differed significantly concerning height, weight, SES and area of residence. Notably, Polynesian and Melanesian adolescents were more likely to live in a rural area and have lower SES than their European counterparts.

### Overweight prevalence

In our sample, overweight/obesity concerned between 25.4% (French reference) and 35.5% (WHO reference) of the adolescents, with a 32.3% prevalence for the IOTF reference (Table 1). Sensitivity analysis for all the participants with measured height and weight ( $N = 1011$ ) showed no significant variation in weight status repartition ( $p > .005$  using  $\chi^2$ ). Whatever the reference, Polynesian and Melanesian adolescents showed higher overweight/obesity prevalence than adolescents with a European background. Table S3 shows the BMIz average according to ethnicity and gender. The results of a 4 x 2 ANOVA showed no significant interaction between ethnicity and gender,  $F(3, 954) = 2.16, p = .091, \eta_p^2 < .01$ . However, there was a significant main effect of gender [ $F(1, 954) = 4.86, p = .028, \eta_p^2 < .01$ ] and ethnicity [ $F(3, 954) = 53.01, p < .001, \eta_p^2 = .14$ ]. Tukey testing indicated that Melanesian and Polynesian adolescents had significantly higher BMIz averages than adolescents from European and other ethnic backgrounds ( $p < .001$  in all cases), while adolescents from European and other origins presented no difference in BMIz average.

Table 1: Weight status of New Caledonian adolescents stratified by ethnic background and gender

Weight status according to reference used*		Whole sample N=954		Melanesian N=429				European N=368				Polynesian N=97				Other N=60			
				Boy N=214		Girl N=215		Boy N=196		Girl N=172		Boy N=49		Girl N=48		Boy N=29		Girl N=31	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
IOTF ref.	U	74	7.8	10	4.7	7	3.3	20	10.2	31	18.0	1	2.0	0	0.0	2	6.9	3	9.7
	N	594	62.3	135	63.1	122	56.7	146	74.5	113	65.7	18	36.7	21	43.8	17	58.6	22	71.0
	OV	187	19.6	40	18.7	63	29.3	22	11.2	23	13.4	10	20.4	17	35.4	6	20.7	6	19.4
	OB	99	10.4	29	13.6	23	10.7	8	4.1	5	2.9	20	40.8	10	20.8	4	13.8	0	0.0
French ref.	U	17	1.8	2	0.9	2	0.9	5	2.6	7	4.1	1	2.0	0	0.0	0	0.0	0	0.0
	N	695	72.9	149	69.6	146	67.9	163	83.2	143	83.1	18	36.7	28	58.3	20	69.0	28	90.3
	OV	242	25.4	63	29.4	67	31.2	28	14.3	22	12.8	30	61.2	20	41.7	9	31.0	3	9.7
WHO ref.	U	18	1.9	2	0.9	2	0.9	5	2.6	8	4.7	1	2.0	0	0.0	0	0.0	0	0.0
	N	598	62.7	123	57.5	117	54.4	152	77.6	131	76.2	16	32.7	16	33.3	18	62.1	25	80.6
	OV	202	21.2	47	22.0	64	29.8	27	13.8	24	14.0	7	14.3	21	43.8	6	20.7	6	19.4
	OB	136	14.3	42	19.6	32	14.9	12	6.1	9	5.2	25	51.0	11	22.9	5	17.2	0	0.0

\*ref: reference, U: underweight, N: normal weight, OV: overweight, OB: obese

## Dietary pattern

Results of PCA yielded KMO = 0.877 and the items had sufficiently high correlations for PCA (Bartlett's test of sphericity =  $\chi^2$  (276) = 4860,  $p < 0.001$ ). Furthermore, the 24 items used in the PCA showed good consistency (Cronbach's alpha = .73). The number of components decided upon was confirmed by an examination of the scree plot, which showed an inflection at the sixth component. PCA identified five dietary patterns that explained 47.3% of the variance in the respondents' diets (see Table 2).

**Table 2: Factor loadings of the food groups in the dietary patterns (principal components) identified in New Caledonian adolescents**

	Principal Component				
	Meats	Fast Food	Fruits and vegetables ONENA traditional	Sweets	Dairies and breakfast
<b>% variance explained</b>	<b>11.0</b>	<b>10.6</b>	<b>9.3</b>	<b>8.8</b>	<b>7.6</b>
Pork	0.642				
Red meat	0.627				
Chicken meat	0.588				
Canned meat	0.573				
Delicatessen	0.458				
Pasta & rice	0.456			0.433	
Fish	0.452		0.442		
Beans	0.404				
Fast food (hamburger, pizza, take-away food)		0.716			
French fries		0.707			
Salty snack/Potato chips		0.618			
Egg		0.537			
Fruits			0.729		
Vegetables			0.712		
Tubers			0.600		
Soup			0.433		
Candy		0.417		0.621	
Cake & pastries				0.606	
Sugar sweet beverage				0.577	
Bread				0.504	
Cheese					0.597
Milk					0.577

Breakfast cereal					0.566
Yoghurt					0.558

The label assigned to the components was based on items with high loadings within each component and the interpretability of factors. These included: (1) Meats, (2) Fast food, (3) Fruits and vegetables, (4) Sweets, and (5) Dairies. It should be noted that the third component, Fruits and vegetables, included fruits and vegetables as well as tubers, soups-and fish, and it seems to represent the traditional ONENA diet. Also, the fifth pattern Dairies included cheese and yoghurt, as well as breakfast cereal with milk, and it could thus be interpreted as Dairies and breakfast pattern. Sensitivity analysis was conducted for all participants who responded to the FFQ (N = 1031) and similar dietary patterns were found (Table S4).

**Table 3: Mixed linear models of dietary patterns and background factors**

	Meats		Fast Food		Fruits and vegetables traditional ONENA diet		Sweets		Dairies and breakfast	
	$\beta$	p	$\beta$	p	$\beta$	p	$\beta$	p	$\beta$	p
Age (years)	.040	.216	-.084	.008	-.036	.230	.121	.001	-.115	<.001
Rural area\$	-.028	.501	.079	.089	.104	.004	.035	.624	-.080	.031
Girl†	-.172	<.001	-.040	.202	-.007	.828	.033	.299	-.112	<.001
Melanesian‡	.100	.021	.161	<.001	.261	<.001	.205	<.001	-.214	<.001
Polynesian‡	.061	.103	.028	.442	.003	.933	.098	.009	-.088	.012
Other‡	.034	.313	.047	.152	.003	.918	.012	.723	-.018	.579
Low SES§	.040	.334	.148	<.001	.039	.316	.018	.657	-.005	.904
Mid SES§	-.001	.981	-.015	.676	-.004	.904	.010	.769	.003	.935
BMIz	.047	.178	-.050	.139	.116	<.001	-.116	.001	-.072	.031
ICC		.001		.049		.092		.039		.047

ICC: Intra-class correlation coefficient. Reference is: † boy, ‡ European, \$ urban area, § high SES.

Mixed linear regressions of dietary patterns and background factors are shown in Table 3. In all mixed models, the intra-class correlation between schools was poor (ICC < .01). The first pattern, Meats, explained 11.0% of the variation in the adolescent diets, and high scores were associated with ethnic background (being Melanesian) and gender (being a boy), but not with SES, age, area of residence or BMIz. The second pattern, Fast food, explained 10.6% of the variation in the adolescent diets, and high scores were associated with younger age, being Melanesian and having low SES. In the third dietary pattern, Fruit and vegetables, the traditional Pacific diet, high scores were associated with being Melanesian, living in a rural

area, and high BMIz. The score for the fourth pattern, Sweets, was positively associated with ethnic background (being Melanesian or Polynesian) and older age and negatively associated with BMIz. The score for the last pattern, Dairies and breakfast, was negatively associated with older age, rural area, being a girl, being Melanesian or Polynesian, and BMIz, but not with SES.

### Sleep duration

Table 4 shows the sleep duration average on weekdays and weekends according to ethnicity and gender. The results of a 4 x 2 ANOVA for average sleep duration showed no significant interaction between ethnicity and gender,  $F(3, 954) = 0.52, p = .783, \eta_p^2 < .01$ . However, there were significant main effects of ethnicity [ $F(3, 954) = 13.33, p = <.001, \eta_p^2 < .01$ ] and gender [ $F(1, 954) = 8.09, p = .005, \eta_p^2 < .01$ ]. Tukey testing indicated that Polynesian adolescents had significantly lower sleep duration than Europeans ( $p < .001$ ), Melanesians ( $p = .007$ ) and adolescents from other ethnic backgrounds ( $p = .047$ ). Melanesian adolescents showed lower sleep duration than Europeans ( $p < .001$ ) but higher than Polynesian adolescents ( $p = .007$ ). No difference was found between adolescents from other origins and Melanesian or European counterparts. Sensitivity analysis was conducted for all participants who responded to this part of the survey ( $N = 1031$ ) and similar results were found (Table S5), except that no significant effects of gender were found ( $p = .253$ ).

**Table 4: Sleep durations for New Caledonian adolescents stratified by ethnic background and gender**

Ethnicity	Gender	Sleep duration on weekdays (hours/day)		Sleep duration on weekends (hours/day)		Average sleep duration (hours/day)		N
		Mean	SD	Mean	SD	Mean	SD	
Melanesian	Boy	8.05	1.48	8.48	2.07	8.17	1.35	214
	Girl	8.14	1.22	9.01	1.75	8.39	1.08	215
	Total	8.10	1.36	8.74	1.94	8.28	1.23	429
European	Boy	8.44	1.16	8.84	2.00	8.55	1.12	196
	Girl	8.51	1.23	9.33	1.50	8.74	1.07	172
	Total	8.47	1.19	9.07	1.79	8.64	1.10	368
Polynesian	Boy	7.69	2.01	7.46	2.20	7.63	1.73	49
	Girl	7.94	1.08	8.39	1.93	8.07	1.05	48
	Total	7.81	1.62	7.92	2.11	7.84	1.45	97
Other	Boy	8.26	1.27	7.91	2.48	8.16	1.40	29
	Girl	8.32	1.28	9.10	1.80	8.54	1.15	31
	Total	8.29	1.27	8.53	2.22	8.36	1.28	60
Total	Boy	8.18	1.43	8.49	2.12	8.27	1.34	488



	Girl	8.27	1.23	9.07	1.70	8.50	1.10	466
	Total	8.22	1.34	8.77	1.95	8.38	1.23	954

SD: Standard deviation

After controlling for age, SES, area of residence and BMIz in a mixed linear regression (schools ICC = .012), the association between sleep time average and gender remained significant ( $\beta = .097, p = .002$ ). Similarly, the ethnicity effect on sleep duration was confirmed for Polynesians ( $\beta = -.149, p < .001$ ) and Melanesians ( $\beta = -.090, p = .038$ ). Moreover, a significant negative relationship was found between age and sleep duration ( $\beta = -.126, p < .001$ ) and between BMIz and sleep duration ( $\beta = -.098, p = .004$ ).

### Screen use

Figure 1 and Table S6 show the screen time (PC/tablet and smartphone) average on weekdays and weekends according to ethnicity and gender. The results of a 4 x 2 ANOVA for PC/tablet screen time showed no significant interaction between ethnicity and gender,  $F(3, 954) = 0.94, p = .418, \eta_p^2 < .01$ . Moreover, the main effect of gender was not significant,  $F(1, 954) = 0.07, p = .798, \eta_p^2 < .01$ , but the main effect of ethnicity was [ $F(3, 954) = 24.72, p < .001, \eta_p^2 < .01$ ]. Tukey testing indicated that the adolescents of other origins had significantly higher screen time than European ( $p = .016$ ), Melanesian ( $p < .001$ ) and Polynesian ( $p = .011$ ) adolescents. Melanesian adolescents showed lower screen time than all the other groups ( $p < .001$ ). No difference was found between European and Polynesian adolescents. Sensitivity analysis using a higher number of adolescents ( $N = 1010$ ) showed the same results (Table S7).

Concerning smartphone use, the results showed no significant interaction between ethnicity and gender,  $F(3, 954) = 0.23, p = .878, \eta_p^2 < .01$ . Moreover, the main effect of ethnicity was not significant [ $F(1, 954) = 2.46, p = .061, \eta_p^2 < .01$ ], but the main effect of gender was [ $F(3, 954) = 7.19, p = .023, \eta_p^2 < .01$ ]. Student's t-test indicated that the gender difference was only significant between Melanesian boys and girls. Sensitivity analysis (Table S7) showed a significant effect of gender ( $p = .021$ ) and ethnicity ( $p = .008$ ) with a higher smartphone use for adolescents from other origins compared with Melanesian adolescents ( $p = .036$ ).

For total screen time (smartphone + other screen use), the results showed a significant main effect of ethnicity,  $F(3, 954) = 12.50, p < .001, \eta_p^2 = .04$ . However, there was no significant main effect of gender [ $F(1, 954) = 1.40, p = .236, \eta_p^2 < .01$ ] and no significant interaction between ethnicity and gender [ $F(3, 954) = 0.65, p = .582, \eta_p^2 < .01$ ]. Tukey testing indicated that Melanesian adolescents had significantly lower total screen time on average than European ( $p < .001$ ) and Polynesian ( $p = .005$ ) adolescents and those from other origins ( $p < .001$ ). No differences were found between the other ethnic groups of adolescents. Sensitivity analysis (Table S7) showed the same results except that Tukey post-hoc analysis indicated that European had lower total screen time than adolescents from other origins ( $p = .004$ ).

When controlling for age, SES, area of residence and BMIz in a mixed linear regression (schools ICC = .047), the association between total screen time and gender remained non-significant and a positive association was found for adolescents from other ethnicities ( $\beta = .067, p = .038$ ) compared to Europeans. Moreover, a positive correlation was found with age ( $\beta = .278, p < .001$ ) but no correlation was found with SES, living area or BMIz.

### **Overweight predictors**

Results of the multiple logistic regression examining the relationship between the adolescents' weight status (being overweight) and the five identified dietary patterns, screen time and sleep duration, controlled for ethnic background, age, gender, SES and area of residence, are presented in Table 5. For the whole sample, Polynesian and Melanesian adolescents were more likely to be overweight than their European counterparts (OR = 5.40 and OR = 1.67, respectively), and less likely to be underweight (Table S8, OR = .07 and OR = .27 respectively). Moreover, adolescents with lower SES were more likely to be overweight (OR = 2.16), but this trend was found in the whole sample and the European adolescent subgroup only. Melanesian adolescents with higher screen time were less likely to be overweight (OR = .74), and this trend was not found in the other ethnic subgroups or the whole sample. In the European adolescent subgroup, higher sleep duration was correlated with a lower risk of being overweight (OR = .65), and higher risk of being underweight (Table S8, OR = 1.55).

Dietary pattern analysis showed that the Meats and Fast food patterns were not independently associated with overweight in the whole sample or any of the ethnic subgroups. However, adolescents with high scores in the Fruits and vegetables pattern were more likely to be overweight in the European subgroup and the whole sample (OR = 2.06 and OR = 1.19, respectively). Correlations between the Sweets and the Dairies and breakfast patterns and being overweight were found: participants with higher scores for the Sweets pattern were less likely to be overweight in the whole sample (OR = .78) and the European (OR = .60) and Melanesian (OR = .81) adolescents. Similarly, participants scoring high in the Dairies and breakfast pattern were less likely to be overweight in the whole sample (OR = .83) and the European (OR = .52) and Polynesian (OR = .49) adolescents. No correlation was found between dietary patterns and being underweight in the whole sample or the European and Melanesian subgroup (Table S8).

**Table 5: Multiple logistic regression for predictors of being overweight**

		Whole sample N=954 R <sup>2</sup> =.134		Melanesian N=429 R <sup>2</sup> =.069		European N=368 R <sup>2</sup> =.165		Polynesian N=97 R <sup>2</sup> =.131	
		OR (95%CI)	p	OR (95%CI)	p	OR (95%CI)	p	OR (95%CI)	p
Ethnicity	Melanesian	1.67 (1.07-2.61)	.023						
	Polynesian	5.40 (3.13-9.30)	<.001						
	Other	1.75 (.90-3.40)	.101						
	European	1.00							
SES	Low	2.16 (1.42-3.30)	<.001	1.38 (.70-2.72)	.351	3.43 (1.67-7.07)	<.001	1.08 (.29-4.01)	.913
	Mid	1.39 (.80-2.41)	.262	1.02 (.42-2.48)	.972	.77 (.26-2.29)	.635	2.49 (.39-15.91)	.334
	High	1.00		1.00		1.00		1.00	
Area of residence	Urban	.86 (.59-1.26)	.445	.70 (.37-1.33)	.279	1.63 (.75-3.51)	.215	1.11 (.37-3.35)	.851
	Rural	1.00		1.00		1.00		1.00	
Gender	Boy	.87 (.64-1.19)	.382	.66 (.43-1.02)	.060	1.05 (.54-2.07)	.880	1.40 (.53-3.69)	.499
	Girl	1.00		1.00		1.00		1.00	
Age (years)		1.14 (1.00-1.29)	.048	1.14 (.96-1.36)	.139	1.16 (.88-1.53)	.280	.88 (.60-1.29)	.523
Sleeping duration (hours/day)		.93 (.83-1.05)	.267	1.07 (.90-1.27)	.464	.65 (.48-.87)	.004	.97 (.70-1.36)	.866
Screen use (hours/day)		.88 (.76-1.01)	.069	.74 (.59-0.92)	.008	0.96 (.73-1.26)	.742	0.86 (.58-1.28)	.462
Nutrition	Pattern 1 factor score: Meats	1.06 (.91-1.23)	.434	1.14 (.94-1.38)	.178	1.47 (.97-2.21)	.067	.83 (.52-1.32)	.427

Pattern 2 factor score: Fast food	.93 (.80-1.109)	.388	0.84 (.69-1.02)	.072	1.33 (.77-2.30)	.305	.90 (.55-1.48)	.678
Pattern 3 factor score: Fruits and vegetables	1.19 (1.02-1.40)	.032	1.05 (.85-1.30)	.657	2.06 (1.36-3.10)	.001	1.32 (.81-2.15)	.270
Pattern 4 factor score: Sweets	.78 (.66-.92)	.004	.81 (.66-.99)	.040	.60 (.38-.97)	.036	1.25 (.75-2.08)	.396
Pattern 5 factor score: Dairies and breakfast	.83 (.71-.98)	.029	.97 (.77-1.21)	.766	.52 (.35-.78)	.002	.49 (.30-.82)	.006

*OR: adjusted odds ratio; CI: confidence interval*

## Discussion

This study in a pluri-ethnic population of adolescents living in New Caledonia revealed five dominant dietary patterns, with some of them found to be associated with overweight. Sleep duration and screen time were evaluated and correlations with weight status were explored. Results showed that ethnicity and low SES remained the main factors for being overweight in New Caledonia.

To our knowledge, this is the first study examining the dietary patterns of New Caledonian adolescents from 10.5 to 16 years old. Five main food consumption patterns were identified: Meats, Fast food, Fruits and vegetables, Sweets, and Dairies and breakfast. In most studies, an unhealthy Western dietary pattern has been associated with higher intakes of red and processed meats, refined grains, sweets and desserts, high-fat dairy products and fast food, while a healthy pattern has been found to consist of higher intakes of fruits, vegetables, whole grains, fish, poultry, and low-fat dairy products.<sup>66-69</sup> In our sample, the Meats (including red meat and canned meat), Fast food and Sweets patterns seemed to represent unhealthy or Western-like patterns, whereas the Fruits and vegetables and Dairies and breakfast patterns may have represented healthy patterns.

The Fruits and vegetables pattern was dense in fruits, vegetables, tubers, soup and fish, and it therefore could be considered as the traditional ONENA diet.<sup>70</sup> In rural areas, high scores for traditional ONENA Fruits and vegetables pattern and low scores for the Dairies and breakfast pattern were observed. These trends can be partly explained by the availability and affordability of these products.<sup>56,71</sup> Fruits and vegetables are commonly cultivated in rural areas by traditional Melanesian family farms or bigger

European farms and thus are widely available and affordable. Food can be purchased but is also available for subsistence consumption, comprising subsistence production plus gifts and exchanges.<sup>72</sup> In contrast, cheese and yoghurt are not produced in rural areas of New Caledonia and may be expensive and not always available. Melanesian adolescents showed a major difference in dietary patterns compared with their European counterparts. They consumed more meat, fast food, fruits and vegetables and sweets and consumed less dairy, even after controlling SES, age and place of residence. The high consumption of fruits and vegetables may reflect the traditional Pacific diet of the Melanesian adolescents, but the high consumption of meats, fast food and sweets compared with their European counterparts was surprising and needs further investigation. Polynesian adolescents showed some differences with their European counterparts, with a higher score in the Sweets pattern and a lower score for the Dairies and breakfast pattern. Last, girls showed lower scores for the Meats and Dairies patterns than boys. It should be noted that the study design and the FFQ we used did not allow us to estimate the total energy intake of the participants. It was difficult to determine whether a given ethnic group's higher consumption in a specific pattern was associated with a higher energy intake or whether the higher and lower scores in some patterns balanced each other. The Melanesian adolescents' scores were especially higher on average in four patterns (Meats, Fast food, Fruits and vegetables and sweets), whereas they had a lower score on average in the last pattern (Dairies).

Our results showed that the European adolescents scoring high in the Fruits and vegetables pattern were more likely to be overweight. No causal relationship could be determined because of the cross-sectional study design; however, an explanation may be that the European adolescents who were already overweight may have increased fruits and vegetables uptake or declarations of uptake. These behaviours may be a strategy to avoid weight gain or to lose weight. In this case, this strategy was not used by Polynesian or Melanesian adolescents, because the fruits and vegetables consumption of these subgroups of adolescents was not associated with the overweight odds ratio, possibly because of less weight consciousness, less parental intervention or less motivation to avoid weight gain. Similarly, adolescents scoring high in the Sweets pattern were less likely to be overweight in the whole sample and in the

Melanesian and European subgroups, but not in the Polynesian subgroup. Once again, this might be a strategy to avoid weight gain for overweight adolescents. Another trend is that the European and Polynesian adolescents with lower scores in the Dairies pattern were more likely to be overweight. This pattern consists of cheese and yoghurt but also milk and breakfast cereals and, as previously explained, overweight adolescents may avoid eating high-fat products like cheese. Moreover, dairy products are not typical of the ONENA traditional diet.<sup>70</sup> Another explanation is that this pattern may reflect the relationship between breakfast skipping and overweight, previously described in New Caledonian adolescents.<sup>24</sup> No specific dietary patterns were found to be associated with underweight condition in our samples.

We showed that Melanesian and Polynesian adolescents had lower sleep durations (20 min and 40 min less, respectively) than their European counterparts, who slept about 8 hours and half per night. These results were very similar to those obtained by Galland et al.,<sup>53</sup> who showed that New Zealand Europeans slept about 8 hours and 20 min and that bedtimes were later for Maori and Pacific adolescents (15 and 41 min, respectively). We found a negative correlation between age and sleep duration, and this trend was observed by other authors in the Pacific area.<sup>53</sup> Short sleep durations have been associated with a greater risk of developing overweight/obesity and a significant change in BMIz at all ages.<sup>23,73,74</sup> However, a recent report showed that the relationship between sleep time and weight status should be interpreted with caution, as higher BMI may precede less sleep.<sup>75</sup> and thus we do not know if one is the cause of the other or the other way around. In our sample, the association between sleep duration and overweight was only found in the European subgroup, with the odds of being overweight increased by 50% per hour of less sleep. Interestingly, no association was found between sleep duration and weight status in the Melanesian or Polynesian adolescents. These results are in line with observations by Reither et al.,<sup>76</sup> in a large survey (n = 30,133) that showed that the relationship between sleep duration and BMI in the U.S. adolescents may differ across ethnicities and genders. They found a negative association between BMI and sleep duration in white, Hispanics and Asian boys; no such association in black boys and white, Hispanic and Asian girls; and a positive association in black girls.<sup>76</sup>

Although screen time showed ethnic differences in our study, no correlation between screen time and being overweight was found in the whole sample. Melanesian adolescents seem to have lower screen time than their European counterparts. Moreover, Melanesians with higher screen time were less likely to be overweight than the Europeans, but the magnitude of this association was weak (OR=0.74). It should be noted that most studies report a positive relationship between TV watching and overweight,<sup>77</sup> but we only examined PC/tablet and smartphone screen time in our survey. Concerning computer use/game time, van Erkis found no evidence for a relationship with BMI/BMIz.<sup>77</sup> Moreover, a recent review revealed insufficient evidence of an association between overall screen time or non-television screen time and greater adiposity.<sup>78</sup>

In this study, the main factor for overweight was ethnic background, i.e. being Melanesian (OR=1.67) or Polynesian (OR=5.40), which makes attractive the hypothesis of a genetic influence. Recent studies about the relationship between the CREBRF gene variant and BMI gain,<sup>79–81</sup> indicated that the hypothesis of a thrifty gene should be explored for people of Polynesian background in New Caledonia. The absence of this variant in the Melanesian population of Solomon Islands,<sup>81</sup> and the particularity of the gut microbiome in the rural Melanesian population of Papua New Guinea,<sup>82</sup> suggest another possible mechanism to explain weight gain. The high consumption of the traditional ONENA diet (fruits, vegetables, tubers, fish) of the Melanesian adolescents in this study and a genetic influence,<sup>82</sup> might have led to a specific microbiome with higher energy extraction from carbohydrate-rich diets. The modernization of New Caledonia has led to the higher consumption of processed food (fast food and sweets) by Melanesian adolescents, which might then have led to overweight because of this specific microbiome.

Another possible way to explain our results was to consider the physical activity (PA) of the adolescents. It has been shown that low PA may be correlated with overweight in adolescent populations.<sup>1,38,40,56</sup> PA may differ across ethnic subgroups and thus explain the disparities in overweight prevalence. Indeed, relatively low PA was found in New Caledonian Melanesian adolescents,<sup>83</sup> with a rural/urban difference.<sup>84</sup> Future investigations are needed in New Caledonia to determine whether PA is lower in Melanesian and Polynesian adolescents compared with their European counterparts.

While our study has strengths, including the anthropometric measurement of all participants, there are limitations to consider when interpreting our findings. First, the data are cross-sectional, and thus no causal relationships could be determined. Due to the size of the sample, the sociodemographic characteristics of the participants might be slightly different from the total school-going adolescent population. Concerning the short FFQ used, social desirability bias may have led overweight adolescents to overreport or underreport food consumption.<sup>85</sup> Moreover, the design of the FFQ did not allow us to determine the total energy intake. This FFQ was also validated in a specific population in Australia, so it may not reflect some of the specificities of the New Caledonian adolescent diet. Also, we did not assess PA, which can have an impact on weight status. Concerning sleep time duration, FFQ and screen use, we obtained self-declarative data and the results should thus be considered with caution. Despite these limitations, this is the first study to examine concomitantly the impacts of sleep duration, screen time and diet on overweight in a pluri-ethnic Pacific population of adolescents.

## **Conclusion**

The prevalence of overweight/obesity is high in New Caledonian adolescents, particularly Polynesian and Melanesian adolescents. Regardless of the potential modifiable behaviours, specific diet interventions targeting adolescents may be useful to prevent obesity in New Caledonia. Interventions to diminish the consumption of fast food and sweets in Melanesian and Polynesian adolescents may also be useful, as would interventions targeting breakfast intake in Polynesian and European adolescents. Although sleep duration was correlated with overweight only in European adolescents, low sleep duration may have other consequences for well-being. Thus, interventions targeting sleep duration may be useful for all adolescents. However, our results are compatible with a genetic influence on weight gain, although research concerning genetic variant and/or microbiome influence remains challenging.

## **Contributors**

SF and OG conceived and designed the experiment. EP, ANG, CSM, SF, GW and OG collected the data. SF and GW analysed the data. SF wrote the original draft. All the authors collected the data and reviewed, edited and approved the final manuscript.



**Declaration of interests**

The authors declare that they have no competing interests.

**Acknowledgments**

We would like to thank the school teaching teams and administrative staffs for their help and support in our investigations, especially the Vice-Rectorat of New Caledonia. We would like to thank Paul Zongo, Pierre-Yves Le Roux and Fabrice Wacalie for their help for collecting the data.

**Data sharing statement**

The datasets collected during the current study are available from the corresponding author on reasonable request.

## Reference

- 1 Kessaram T, McKenzie J, Girin N, *et al.* Overweight, obesity, physical activity and sugar-sweetened beverage consumption in adolescents of Pacific islands: results from the Global School-Based Student Health Survey and the Youth Risk Behavior Surveillance System. *BMC Obes* 2015; **2**: 34.
- 2 Pacific islanders pay heavy price for abandoning traditional diet. *Bull World Health Organ* 2010; **88**: 484–5.
- 3 Hughes RG, Lawrence MA. Globalization, food and health in Pacific Island countries. *Asia Pac J Clin Nutr* 2005; **14**: 298–306.
- 4 Pengpid S, Peltzer K. Dietary health behaviour and beliefs among university students from 26 low, middle and high income countries. *Asia Pac J Clin Nutr* 2015; **24**: 744–52.
- 5 Craig PL, Swinburn BA, Matenga-Smith T, Matangi H, Vaughn G. Do Polynesians still believe that big is beautiful? Comparison of body size perceptions and preferences of Cook Islands, Maori and Australians. *N Z Med J* 1996; **109**: 200–3.
- 6 Pollock NJ, Institute for Polynesian Studies L. These roots remain. Food habits in islands of the central and eastern Pacific since western contact. 1992. <https://agris.fao.org/agris-search/search.do?recordID=XF2016055909> (accessed June 19, 2020).
- 7 Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass index and risk of 22 specific cancers: a population-based cohort study of 5·24 million UK adults. *Lancet Lond Engl* 2014; **384**: 755–65.
- 8 Dwivedi AK, Dubey P, Cistola DP, Reddy SY. Association Between Obesity and Cardiovascular Outcomes: Updated Evidence from Meta-analysis Studies. *Curr Cardiol Rep* 2020; **22**: 25.
- 9 Seravalle G, Grassi G. Obesity and hypertension. *Pharmacol Res* 2017; **122**: 1–7.
- 10 Luppino FS, Wit LM de, Bouvy PF, *et al.* Overweight, Obesity, and Depression: A Systematic Review and Meta-analysis of Longitudinal Studies. *Arch Gen Psychiatry* 2010; **67**: 220–9.
- 11 Mannan M, Mamun A, Doi S, Clavarino A. Prospective Associations between Depression and Obesity for Adolescent Males and Females- A Systematic Review and Meta-Analysis of Longitudinal Studies. *PLoS ONE* 2016; **11**. DOI:10.1371/journal.pone.0157240.
- 12 Mühlig Y, Antel J, Föcker M, Hebebrand J. Are bidirectional associations of obesity and depression already apparent in childhood and adolescence as based on high-quality studies? A systematic review. *Obes Rev* 2016; **17**: 235–49.
- 13 Quek Y-H, Tam WWS, Zhang MWB, Ho RCM. Exploring the association between childhood and adolescent obesity and depression: a meta-analysis. *Obes Rev* 2017; **18**: 742–54.
- 14 Wardle J, Cooke L. The impact of obesity on psychological well-being. *Best Pract Res Clin Endocrinol Metab* 2005; **19**: 421–40.
- 15 Weinberger N-A, Kersting A, Riedel-Heller SG, Luck-Sikorski C. Body Dissatisfaction in Individuals with Obesity Compared to Normal-Weight Individuals: A Systematic Review and Meta-Analysis. *Obes Facts* 2017; **9**: 424–41.

- 16Micali N, Stavola BD, Ploubidis G, Simonoff E, Treasure J, Field AE. Adolescent eating disorder behaviours and cognitions: Gender-specific effects of child, maternal and family risk factors. *Br J Psychiatry* 2015; **207**: 320–7.
- 17Danielsen YS, Stormark KM, Nordhus IH, *et al.* Factors Associated with Low Self-Esteem in Children with Overweight\*. *Obes Facts* 2012; **5**: 722–33.
- 18French SA, Story M, Perry CL. Self-Esteem and Obesity in Children and Adolescents: A Literature Review. *Obes Res* 1995; **3**: 479–90.
- 19Griffiths LJ, Parsons TJ, Hill AJ. Self-esteem and quality of life in obese children and adolescents: a systematic review. *Int J Pediatr Obes IJPO Off J Int Assoc Study Obes* 2010; **5**: 282–304.
- 20Miller CT, Downey KT. A Meta-Analysis of Heavyweight and Self-Esteem: *Personal Soc Psychol Rev* 2016; published online Dec 21. DOI:10.1207/s15327957pspr0301\_4.
- 21Sikorski C, Luppia M, Luck T, Riedel-Heller SG. Weight stigma ‘gets under the skin’-evidence for an adapted psychological mediation framework: a systematic review. *Obes Silver Spring Md* 2015; **23**: 266–76.
- 22Arora T, Broglia E, Pushpakumar D, Lodhi T, Taheri S. An Investigation into the Strength of the Association and Agreement Levels between Subjective and Objective Sleep Duration in Adolescents. *PLoS ONE* 2013; **8**. DOI:10.1371/journal.pone.0072406.
- 23Miller MA, Kruisbrink M, Wallace J, Ji C, Cappuccio FP. Sleep duration and incidence of obesity in infants, children, and adolescents: a systematic review and meta-analysis of prospective studies. *Sleep* 2018; **41**. DOI:10.1093/sleep/zsy018.
- 24Frayon S, Cherrier S, Cavaloc Y, *et al.* Nutrition behaviors and sociodemographic factors associated with overweight in the multi-ethnic adolescents of New Caledonia. *Ethn Health* 2017; **0**: 1–17.
- 25Frayon S, Cherrier S, Cavaloc Y, Wattelez G, Lerrant Y, Galy O. Relationship of body fat and body mass index in young Pacific Islanders: a cross-sectional study in European, Melanesian and Polynesian groups. *Pediatr Obes* 2018; **13**: 357–64.
- 26Tubert-Jeannin S, Pichot H, Rouchon B, Pereira B, Hennequin M. Common risk indicators for oral diseases and obesity in 12-year-olds: a South Pacific cross sectional study. *BMC Public Health* 2018; **18**: 112.
- 27Chaput J-P, Doucet E, Tremblay A. Obesity: a disease or a biological adaptation? An update. *Obes Rev Off J Int Assoc Study Obes* 2012; **13**: 681–91.
- 28Mozaffarian D. Dietary and Policy Priorities for Cardiovascular Disease, Diabetes, and Obesity: A Comprehensive Review. *Circulation* 2016; **133**: 187–225.
- 29Tarantola A, Horwood PF, Goarant C, *et al.* Counting Oceanians of Non-European, Non-Asian Descent (ONENA) in the South Pacific to Make Them Count in Global Health. *Trop Med Infect Dis* 2019; **4**: 114.
- 30Krueger PM, Reither EN. Mind the gap: race/ethnic and socioeconomic disparities in obesity. *Curr Diab Rep* 2015; **15**: 95.
- 31Walker JL, Ardouin S, Burrows T. The validity of dietary assessment methods to accurately measure energy intake in children and adolescents who are overweight or obese: a systematic review. *Eur J Clin Nutr* 2018; **72**: 185–97.

- 32 Dancause KN, Dehuff C, Soloway LE, *et al.* Behavioral changes associated with economic development in the South Pacific: Health transition in Vanuatu. *Am J Hum Biol* 2011; **23**: 366–76.
- 33 Hawley NL, McGarvey ST. Obesity and diabetes in Pacific Islanders: the current burden and the need for urgent action. *Curr Diab Rep* 2015; **15**: 29.
- 34 Ulijaszek S. Modernisation, migration and nutritional health of Pacific Island populations. *Environ Sci Int J Environ Physiol Toxicol* 2005; **12**: 167–76.
- 35 Wattlez G, Frayon S, Cavaloc Y, Cherrier S, Lerrant Y, Galy O. Sugar-Sweetened Beverage Consumption and Associated Factors in School-Going Adolescents of New Caledonia. *Nutrients* 2019; **11**. DOI:10.3390/nu11020452.
- 36 Dancause KN, Vilar M, Wilson M, *et al.* Behavioral risk factors for obesity during health transition in Vanuatu, South Pacific. *Obes Silver Spring Md* 2013; **21**: E98–104.
- 37 Neumark-Sztainer D, Story M, Perry C, Casey MA. Factors influencing food choices of adolescents: findings from focus-group discussions with adolescents. *J Am Diet Assoc* 1999; **99**: 929–37.
- 38 Salvy S-J, Bowker JC. Peers and Obesity during Childhood and Adolescence: A Review of the Empirical Research on Peers, Eating, and Physical Activity. *J Obes Weight Loss Ther* 2014; **4**. DOI:10.4172/2165-7904.1000207.
- 39 Lissak G. Adverse physiological and psychological effects of screen time on children and adolescents: Literature review and case study. *Environ Res* 2018; **164**: 149–57.
- 40 Maher C, Olds TS, Eisenmann JC, Dollman J. Screen time is more strongly associated than physical activity with overweight and obesity in 9- to 16-year-old Australians. *Acta Paediatr Oslo Nor 1992* 2012; **101**: 1170–4.
- 41 Mhrshahi S, Drayton BA, Bauman AE, Hardy LL. Associations between childhood overweight, obesity, abdominal obesity and obesogenic behaviors and practices in Australian homes. *BMC Public Health* 2017; **18**. DOI:10.1186/s12889-017-4595-y.
- 42 Sun C, Pomer A, Dancause KN, *et al.* Ownership of consumer electronics is associated with measures of adiposity during health transition in Vanuatu. *Am J Hum Biol Off J Hum Biol Counc* 2017; **29**. DOI:10.1002/ajhb.22928.
- 43 Chassiakos Y (Linda) R, Radesky J, Christakis D, Moreno MA, Cross C, Media C on CA. Children and Adolescents and Digital Media. *Pediatrics* 2016; **138**. DOI:10.1542/peds.2016-2593.
- 44 Lauricella AR, Wartella E, Rideout VJ. Young children’s screen time: The complex role of parent and child factors. *J Appl Dev Psychol* 2015; **36**: 11–7.
- 45 Lemola S, Perkinson-Gloor N, Brand S, Dewald-Kaufmann JF, Grob A. Adolescents’ Electronic Media Use at Night, Sleep Disturbance, and Depressive Symptoms in the Smartphone Age. *J Youth Adolesc* 2015; **44**: 405–18.
- 46 Anderson YC, Wynter LE, Grant CC, *et al.* Physical activity is low in obese New Zealand children and adolescents. *Sci Rep* 2017; **7**: 1–7.
- 47 Sluyter JD, Scragg RKR, Plank LD, Waqa GD, Fotu KF, Swinburn BA. Sizing the association between lifestyle behaviours and fatness in a large, heterogeneous sample of youth of multiple ethnicities from 4 countries. *Int J Behav Nutr Phys Act* 2013; **10**: 115.

- 48 Hale L, Guan S. Screen Time and Sleep among School-Aged Children and Adolescents: A Systematic Literature Review. *Sleep Med Rev* 2015; **21**: 50–8.
- 49 Börnhorst C, Wijnhoven TM, Kunešová M, *et al.* WHO European Childhood Obesity Surveillance Initiative: associations between sleep duration, screen time and food consumption frequencies. *BMC Public Health* 2015; **15**: 442.
- 50 Lee CH, Sibley CG. Sleep duration and psychological well-being among New Zealanders. *Sleep Health* 2019; **5**: 606–14.
- 51 Paine S-J, Gander PH. Explaining ethnic inequities in sleep duration: a cross-sectional survey of Māori and non-Māori adults in New Zealand. *Sleep Health* 2016; **2**: 109–15.
- 52 Vaipuna TFW, Williams SM, Farmer VL, *et al.* Sleep patterns in children differ by ethnicity: cross-sectional and longitudinal analyses using actigraphy. *Sleep Health* 2018; **4**: 81–6.
- 53 Galland BC, de Wilde T, Taylor RW, Smith C. Sleep and pre-bedtime activities in New Zealand adolescents: differences by ethnicity. *Sleep Health* 2020; **6**: 23–31.
- 54 Gwynn JD, Flood VM, D’Este CA, *et al.* The reliability and validity of a short FFQ among Australian Aboriginal and Torres Strait Islander and non-Indigenous rural children. *Public Health Nutr* 2011; **14**: 388–401.
- 55 Flood VM, Webb K, Rangan AM. Recommendations for short questions to assess food consumption in children for the NSW Health Surveys. 2005.
- 56 Galy O, Paufique E, Nedjar-Guerre A, *et al.* Living in Rural and Urban Areas of New Caledonia: Impact on Food Consumption, Sleep Duration and Anthropometric Parameters Among Melanesian Adolescents. *Nutrients* 2020; **12**. DOI:10.3390/nu12072047.
- 57 Jayasinghe SN, Breier BH, McNaughton SA, *et al.* Dietary Patterns in New Zealand Women: Evaluating Differences in Body Composition and Metabolic Biomarkers. *Nutrients* 2019; **11**: 1643.
- 58 INSERM. Situation sociale et comportements de santé des jeunes en Nouvelle-Calédonie. 2008; published online March.
- 59 Krieger N. Review of A Researcher’s Guide to the National Statistics Socio-Economic Classification. *J Public Health Policy* 2003; **24**: 467–73.
- 60 ISEE. Structure de la population et évolutions. 2014.  
<http://www.isee.nc/population/recensement/structure-de-la-population-et-evolutions> (accessed Dec 14, 2015).
- 61 European Communities. The European Union labour force survey - Methods and definitions - 2001 (Eurostat, Guidelines, Labour statistics, Survey). 2003.  
<http://unstats.un.org/unsd/EconStatKB/KnowledgebaseArticle10230.aspx> (accessed Dec 14, 2015).
- 62 Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity: Extended international BMI cut-offs. *Pediatr Obes* 2012; **7**: 284–94.
- 63 de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2007; **85**: 660–7.
- 64 Rolland-Cachera MF, Cole TJ, Sempé M, Tichet J, Rossignol C, Charraud A. Body Mass Index variations: centiles from birth to 87 years. *Eur J Clin Nutr* 1991; **45**: 13–21.

- 65 Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity: Extended international BMI cut-offs. *Pediatr Obes* 2012; **7**: 284–94.
- 66 Gregório MJ, Rodrigues AM, Eusébio M, *et al.* Dietary Patterns Characterized by High Meat Consumption Are Associated with Other Unhealthy Life Styles and Depression Symptoms. *Front Nutr* 2017; **4**. DOI:10.3389/fnut.2017.00025.
- 67 Newby PK, Tucker KL. Empirically Derived Eating Patterns Using Factor or Cluster Analysis: A Review. *Nutr Rev* 2004; **62**: 177–203.
- 68 Roman G, Rusu A, Graur M, *et al.* DIETARY PATTERNS AND THEIR ASSOCIATION WITH OBESITY: A CROSS-SECTIONAL STUDY. *Acta Endocrinol Buchar* 2019; **15**: 86–95.
- 69 Wang X, Ding N, Tucker KL, *et al.* A Western Diet Pattern Is Associated with Higher Concentrations of Blood and Bone Lead among Middle-Aged and Elderly Men. *J Nutr* 2017; **147**: 1374–83.
- 70 Shintani TT, Hughes CK. Traditional diets of the Pacific and coronary heart disease. *J Cardiovasc Risk* 1994; **1**: 16–20.
- 71 Miller V, Yusuf S, Chow CK, *et al.* Availability, affordability, and consumption of fruits and vegetables in 18 countries across income levels: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet Glob Health* 2016; **4**: e695–703.
- 72 Serra Mallol C. Monetary income, public funds, and subsistence consumption: the three components of the food supply in French Polynesia — a comparative study of Tahiti and Rapa Iti islands. *Rev Agric Food Environ Stud* 2018; **99**: 37–55.
- 73 Fatima Y, Doi S a. R, Mamun AA. Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias-adjusted meta-analysis. *Obes Rev* 2015; **16**: 137–49.
- 74 Seo SH, Shim YS. Association of Sleep Duration with Obesity and Cardiometabolic Risk Factors in Children and Adolescents: A Population-Based Study. *Sci Rep* 2019; **9**: 1–10.
- 75 Sokol RL, Grummon AH, Lytle LA. Sleep duration and body mass: direction of the associations from adolescence to young adulthood. *Int J Obes* 2020; **44**: 852–6.
- 76 Reither EN, Krueger PM, Hale L, Reiter EM, Peppard PE. Ethnic Variation in the Association between Sleep and Body Mass among U.S. Adolescents. *Int J Obes* 2005 2014; **38**: 944–9.
- 77 van Ekris E, Altenburg TM, Singh AS, Proper KI, Heymans MW, Chinapaw MJM. An evidence-update on the prospective relationship between childhood sedentary behaviour and biomedical health indicators: a systematic review and meta-analysis. *Obes Rev Off J Int Assoc Study Obes* 2016; **17**: 833–49.
- 78 Stiglic N, Viner RM. Effects of screentime on the health and well-being of children and adolescents: a systematic review of reviews. *BMJ Open* 2019; **9**: e023191.
- 79 Krishnan M, Major TJ, Topless RK, *et al.* Discordant association of the CREBRF rs373863828 A allele with increased BMI and protection from type 2 diabetes in Māori and Pacific (Polynesian) people living in Aotearoa/New Zealand. *Diabetologia* 2018; **61**: 1603–13.
- 80 Minster RL, Hawley NL, Su C-T, *et al.* A thrifty variant in CREBRF strongly influences body mass index in Samoans. *Nat Genet* 2016; **48**: 1049–54.
- 81 Naka I, Furusawa T, Kimura R, *et al.* A missense variant, rs373863828-A (p.Arg457Gln), of CREBRF and body mass index in Oceanic populations. *J Hum Genet* 2017; **62**: 847–9.

- 82 Greenhill AR, Tsuji H, Ogata K, *et al.* Characterization of the Gut Microbiota of Papua New Guineans Using Reverse Transcription Quantitative PCR. *PLoS ONE* 2015; **10**. DOI:10.1371/journal.pone.0117427.
- 83 Galy O, Yacef K, Caillaud C. Improving Pacific Adolescents' Physical Activity Toward International Recommendations: Exploratory Study of a Digital Education App Coupled With Activity Trackers. *JMIR MHealth UHealth* 2019; **7**: e14854.
- 84 Zongo P, Frayon S, Antoine-Jonville S, *et al.* Anthropometric Characteristics and Physical Fitness in Rural and Urban 11- to 16-Year-Old Melanesian Adolescents: A Cross-sectional Study in New Caledonian Schools. *Asia Pac J Public Health* 2017; **29**: 589–98.
- 85 Miller TM, Abdel-Maksoud MF, Crane LA, Marcus AC, Byers TE. Effects of social approval bias on self-reported fruit and vegetable consumption: a randomized controlled trial. *Nutr J* 2008; **7**: 18.

**Figure 1: Total screen time (in hours/day) for New Caledonian adolescents stratified by ethnic background and gender**

