



Analysis of a cross-diffusion model for rival gangs interaction in a city

Alethea B. T. Barbaro, Nancy Rodriguez, Havva Yoldaş, Nicola Zamponi

► To cite this version:

Alethea B. T. Barbaro, Nancy Rodriguez, Havva Yoldaş, Nicola Zamponi. Analysis of a cross-diffusion model for rival gangs interaction in a city. *Communications in Mathematical Sciences*, 2021, 19 (8), pp.2139 - 2175. 10.4310/CMS.2021.v19.n8.a4 . hal-02973956

HAL Id: hal-02973956

<https://hal.science/hal-02973956>

Submitted on 21 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Analysis of a cross-diffusion model for rival gangs interaction in a city

Alethea B. T. Barbaro* Nancy Rodriguez[†] Havva Yoldaş[‡] Nicola Zamponi[§]

October 5, 2020

Abstract

We study a two-species cross-diffusion model that is inspired by a system of convection-diffusion equations derived from an agent-based model on a two-dimensional discrete lattice. The latter model has been proposed to simulate gang territorial development through the use of graffiti markings. We find two energy functionals for the system that allow us to prove a weak-stability result and identify equilibrium solutions. We show that under the natural definition of weak solutions, obtained from the weak-stability result, the system does not allow segregated solutions. Moreover, we present a result on the long-term behavior of solutions in the case when the product of the mass of the densities are less than one. This result is complemented with numerical experiments.

Contents

1	Introduction	2
2	Energy functionals and a-priori estimates	5
2.1	Two energy functionals	6
2.2	A-priori estimates	7
2.2.1	Estimates from “natural” energy balance equation (10).	7
2.2.2	Additional estimates from Maxwell-Boltzmann energy balance equation (12). . .	10
3	Existence analysis	11
3.1	Local-in-time existence of strong solutions.	11
3.2	Weak solutions	12
3.3	Weak stability analysis	13
4	Stationary states	25
4.1	Linear stability analysis	26
5	Long-time behavior	27
6	Numerical results	31

*Department of Mathematics, Applied Mathematics & Statistics, Case Western Reserve University, 10900 Euclid Avenue, Yost Hall, Cleveland, Ohio 44106-7058, USA. abb71@case.edu

[†]Engineering Center, ECOT 225, 526 UCB, Boulder, CO 80309-0526 nrod@unc.edu

[‡]Camille Jordan Institute, Claude Bernard University of Lyon 1, Batiment Jean Braconnier, 21 Avenue Claude Bernard, 69622 Villeurbanne Cedex France. yoldas@math.univ-lyon1.fr

[§]University of Mannheim, School of Business Informatics and Mathematics, B6, 28, 68159 Mannheim, Germany. nzamponi@mail.uni-mannheim.de

1 Introduction

This article is devoted to the study of a two-population model with cross-diffusion:

$$\begin{cases} \partial_t \rho_A(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_A(t, x, y) + 2\beta c \rho_A(t, x, y) \nabla \rho_B(t, x, y)), & x, y \in \Omega, t > 0, \\ \partial_t \rho_B(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_B(t, x, y) + 2\beta c \rho_B(t, x, y) \nabla \rho_A(t, x, y)), & x, y \in \Omega, t > 0, \end{cases} \quad (1)$$

complemented with the initial data

$$\rho_A(0, x, y) = \rho_A^{in}(x, y) \text{ and } \rho_B(0, x, y) = \rho_B^{in}(x, y), \quad x, y \in \Omega, \quad (2)$$

and the homogeneous Neumann boundary conditions

$$\partial_\nu \rho_A(t, x, y) = \partial_\nu \rho_B(t, x, y) = 0 \quad x, y \in \partial\Omega, t > 0. \quad (3)$$

In system (1)-(3), β and c are positive parameters and $\Omega \subset \mathbb{R}^2$ a bounded domain. Such a system can arise, for example, by considering the following two-species segregation model involving two densities of agents, ρ_A and ρ_B , along with respective marking densities g_A and g_B , introduced in [1]:

$$\begin{cases} \partial_t g_A(t, x, y) = c \rho_A(t, x, y) - g_A(t, x, y), & x, y \in \Omega, t > 0, \\ \partial_t g_B(t, x, y) = c \rho_B(t, x, y) - g_B(t, x, y), & x, y \in \Omega, t > 0, \\ \partial_t \rho_A(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_A(t, x, y) + 2\beta \rho_A(t, x, y) \nabla g_B(t, x, y)), & x, y \in \Omega, t > 0, \\ \partial_t \rho_B(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_B(t, x, y) + 2\beta \rho_B(t, x, y) \nabla g_A(t, x, y)), & x, y \in \Omega, t > 0, \end{cases} \quad (4)$$

with homogeneous Neumann boundary conditions. System (4) models the dynamics of two competing groups that mark their territory, *e.g.* with graffiti, and whose movement strategies is a combination of passive diffusion and directed movement towards the gradients of the marking densities of the competing groups. To arrive at the reduced system (1)-(3) from system (4), we assume that the marking densities equilibrate much more rapidly than the population densities. Hence we assume

$$\partial_t g_A(t, x, y) = \partial_t g_B(t, x, y) = 0.$$

However, we remark that system (1)-(3) can also be seen as a more general model of cross diffusion system where the inter-specific interactions can lead to segregation.

The notion of cross-diffusion was initially motivated by Morisita's theory of environmental density [19, 20], which brings to the forefront the influence that a population pressure has on the dispersal of a population due to the interface between individuals. In [21], Shigesada, Kawasaki, and Teramoto introduced a behavioral model for the movement of individuals based on Morisita's observations. According to Morisita's theory, the movement of individuals is influenced by the following three factors: (i) random movement; (ii) population pressure due to mutual interference between individuals; and (iii) movement toward favorable places. The population pressure due to the competing population leads to the cross-diffusion. To see this from a mathematical point of view, we consider our two populations, ρ_A and ρ_B . Under the assumption of Fickian diffusion, we obtain a system of two equations:

$$\begin{cases} \partial_t \rho_A = \operatorname{div} J_A(\rho_A, \rho_B), \\ \partial_t \rho_B = \operatorname{div} J_B(\rho_A, \rho_B), \end{cases}$$

where J_A, J_B are the flows of the populations A and B , respectively. The flow proposed by Shigesada, Kawasaki, and Teramoto has the form:

$$J_A = \nabla \cdot ((c_A + a_{AA}\rho_A + a_{AB}\rho_B)\rho_A) \quad \text{and} \quad J_B = \nabla \cdot ((c_B + a_{BA}\rho_A + a_{BB}\rho_B)\rho_B)$$

with $a_{AA}, a_{AB}, a_{BA}, a_{BB}, c_A, c_B \geq 0$.

Soon after, Busenberg and Travis introduced some epidemic models with migration that also include cross-diffusion in [5]. In their model, the authors assume that the population flow J_i , $i \in \{A, B\}$, are proportional to the gradient of a potential function, Ψ , that only depends on the total population $P = \rho_A + \rho_B$. The proportion is assumed to be the portion of the subpopulation i , which leads to the following form of the flow:

$$J_i = a \frac{\rho_i}{P} \nabla \Psi(P), \text{ for } i \in \{A, B\}.$$

In [16], Gurtin and Pipkin introduced the potential function $\Psi(s) = s^2/2$, which yields that:

$$J_i = a \rho_i \nabla (\rho_A + \rho_B), \text{ for } i \in \{A, B\}. \quad (5)$$

In [14], Galiano and Selgas considered a more general version of the Gurtin and Pipkin model, where potential function depends on a general linear combination of the population densities, with the addition of random movement and environmental effects. The most general version of the system they consider is:

$$J_i(\rho_A, \rho_B) = \rho_i \nabla (a_{iA} \rho_A + a_{iB} \rho_B + b_i V) + c_i \nabla \rho_i,$$

where V is the environmental potential. In [14], the existence of weak solutions for non-negative initial data in L^∞ was proved in two parameter cases. The first was under the condition that

$$4a_{AA}a_{BB} - (a_{AB} + a_{BA})^2 > a_0,$$

for some $a_0 > 0$. This condition implies an ellipticity condition on the matrix $(a_{ij})_{i,j \in \{A,B\}}$ and can be relaxed. The authors were also able to prove the existence of solutions in the case when $a = a_{ij}$ for all $i, j \in \{A, B\}$, with $a > 0$.

System (1)-(3), which we consider here, is a special case of this general model, where only passive diffusion and cross-diffusion are considered. Thus, the populations does not take into account the population pressure due to their own group. In particular, we assume that a_{AA}, a_{BB}, b_A, b_B are all equal to zero. Thus, this case falls outside of the two cases considered in [14]. It is worth noting that systems with local self- and cross-diffusion have found many applications, for example in, illicit trade of drugs [12]; epidemic models with diffusion of polymorphic populations [5]; models for overcrowding effect with nonuniform ease of dispersal for different individuals [16]; opinion dynamics [22]; and biochemical reactions [23].

Many analytical results for cross-diffusion systems are available in the literature. For example, in [6] Chen et al. consider a reaction-cross-diffusion model for an arbitrary number of competing populations which, in the case of linear transition rates, extends the two-species SKT model presented in [21]. Existence of global-in-time weak solutions to the model in a bounded domain with homogeneous Neumann boundary conditions is shown via an entropy method and an approximation scheme. Crucial conditions on the diffusion matrix are either weak cross-diffusion or detailed balance. Another cross-diffusion system where the diffusion matrix depends linearly on the densities is the two-species ion transport model through narrow membrane channels studied by Burger et al. in [4]. There the authors prove global existence of weak solutions to the equations in a bounded domain with no-flux boundary conditions via an entropy method, as well as global existence of strong solutions near the equilibrium. The result is generalized in [15] to the case of an arbitrary number of species with different specific electrical charges and mixed Dirichlet-Neumann boundary condition. The systems considered in [4] and [15] also present a degeneracy in the entropy structure, that is, some gradient estimates are lost when the densities approach some critical region. Other degenerate cross-diffusion models have been recently studied in literature; for example, a large class of population models with degenerate cross-diffusion was analyzed in [24], while the combination of degenerate cross-diffusion and nonlocal interaction is a feature of both models considered in [3] and [11]. The problem of degenerate cross-diffusion in a moving domain was

considered in [2], while the interplay between singularity and degeneracy was a major feature of the model studied in [7]. Reaction-cross-diffusion systems with Laplacian structure have been considered by Desvillettes et al. in [8, 9, 10].

Cross-diffusion equations can be seen as a large class of nonlinear, strongly coupled evolution PDEs with the structure:

$$\partial_t \rho = \operatorname{div} (A(\rho) \nabla \rho) \equiv \sum_{i=1}^n \partial_{x_i} (A(\rho) \partial_{x_i} \rho), \quad x \in \mathbb{R}^n, \quad t > 0. \quad (6)$$

The unknown of the system, $\rho = \rho(t, x) \in \mathbb{R}^n$, usually represents a vector of densities or concentrations. Therefore, it should be nonnegative to be consistent with the physics; sometimes it is also required to be (uniformly) bounded for the same reason. Quite often, in cross-diffusion systems coming from the applied sciences the matrix $A(\rho) \in \mathbb{R}^{n \times n}$, the so-called *diffusion matrix*, is neither symmetric nor positive semidefinite, which means that the standard coercivity-based approaches to the analysis of (6) are ineffective. Moreover maximum/minimum principles are also usually unavailable due to the fact that A is full and lacks a suitable structure. For these reasons, the analytical study of cross-diffusion equations is in general quite challenging.

A useful method, the so-called *boundedness-by-entropy method*, in the analysis of (reaction-)cross-diffusion systems has been developed by Jüngel and collaborators (see e.g. [17] for a comprehensive review) after an idea found in Burger et al. [4]. This method is suitable for systems of evolution PDEs presenting a formal gradient flow structure, or an *entropy structure*. Specifically, it works for systems which can be written in the following form:

$$\partial_t \rho = \operatorname{div} \left(\mathbb{M} \nabla \frac{\delta \mathcal{H}[\rho]}{\delta \rho} \right), \quad t > 0, \quad (7)$$

where $\mathbb{M}$ is a positive semidefinite (often also symmetric) matrix and $\frac{\delta \mathcal{H}[\rho]}{\delta \rho}$ is the Frechét derivative of the convex functional $\mathcal{H}$, which is called the *mathematical entropy* of the system. In many cases $\mathcal{H}[\rho]$ has the form

$$\mathcal{H}[\rho] = \int_{\Omega} h(\rho) dx,$$

where h a scalar convex function called *entropy density*, then the object $\frac{\delta \mathcal{H}[\rho]}{\delta \rho}$ can be identified, via Riesz representation theorem, with the gradient of h :

$$\frac{\delta \mathcal{H}[\rho]}{\delta \rho} \simeq Dh(\rho),$$

which is referred to as *entropy variable*. A first consequence of this formulation is that the functional $\mathcal{H}$ is a Lyapounov functional for (7), that is; it is nonincreasing in time along the solutions of (7):

$$\frac{d}{dt} \mathcal{H}[\rho(t)] = - \int_{\Omega} \nabla Dh(\rho) \cdot \mathbb{M} \nabla Dh(\rho) dx \leq 0, \quad t > 0,$$

since $\mathbb{M}$ is positive semidefinite by assumption¹. Furthermore, if $Dh : \mathcal{D} \rightarrow \mathbb{R}^n$ is a globally invertible mapping, then the physical variable ρ can be written in terms of the entropy variable $w = Dh(\rho)$ via $\rho = (Dh)^{-1}(w)$. As a consequence $\rho \in \mathcal{D}$ whenever $w \in \mathbb{R}^n$. So, if (7) can be written and solved in terms of w , then the constraint $\rho(x, t) \in \mathcal{D}$ will hold whenever $w(x, t)$ is finite (that is, for

¹Here we assumed homogeneous Neumann boundary conditions and no reaction terms, that is the right-hand side of (6) is zero. If any of these conditions are not verified, the quantity $\frac{d}{dt} \mathcal{H}[\rho(t)]$ might be positive. However, suitable compatibility conditions usually ensure that the entropy $\mathcal{H}[\rho(t)]$ remains at every time upper bounded via a Gronwall argument.

a.e. x, t , provided that w is integrable). In particular, if $\mathcal{D} \subset \mathbb{R}^n$ is bounded, then $\rho \in L^\infty$ with bounds that only depend on $\mathcal{D}$; similarly, if $\mathcal{D} \subset \mathbb{R}_+^n$, then ρ has nonnegative components. These ideas can be exploited to formulate an existence argument which proceeds roughly in three steps: (i) writing an approximate scheme which yields a sequence of approximate solutions to (7); (ii) deriving an entropy balance inequality which yields gradient estimates for the approximate solution; (iii) showing via suitable compactness result that the approximate sequence has a converging subsequence and taking the limit in the approximate system to recover a weak solution to (7).

Unfortunately, this program cannot be straightforwardly carried out when studying (1)-(3) because of its extremely degenerate structure. Indeed, a standard entropy (formal gradient flow) structure requires the existence of a *convex* entropy functional, which cannot be the case for (1)-(3). Precisely, a necessary condition for a cross-diffusion system (6) to admit a convex entropy is the normal ellipticity of the differential operator $\rho \mapsto \operatorname{div}(A(\rho)\nabla\rho)$, that is, the property that the real part of every eigenvalue of $A(\rho)$ is nonnegative [18, Lemma 3.2]. This property is not verified by (1)-(3); as a matter of fact, the diffusion matrix $A(\rho)$ in (1)-(3) has one positive and one negative eigenvalue in the region $\{(\rho_A, \rho_B) \in \mathbb{R}_+^2 \mid \rho_A \rho_B > 1\}$. Consistently with this fact, the only Lyapounov functional that is known for (1)-(3) is nonconvex. Furthermore, the property that the mapping $Dh : \mathbb{R}_+^2 \rightarrow \mathbb{R}^2$ being invertible also fails for (1)-(3), since such a mapping is not even one-to-one.

A search for a workaround to counter these difficulties and obtain nonetheless some global-in-time existence result for (1)-(3) has been unsuccessful, and only a local-in-time existence result is available for (1)-(3), which comes from Amann's theory [18, Thr. 3.1] and holds under the assumption that the initial datum is $W^{1,p}$ with $p > d = 2$ and takes values in the region $\{(\rho_A, \rho_B) \in \mathbb{R}_+^2 \mid \rho_A \rho_B < 1\}$. However, we are also able to provide a weak-stability result, which is a key step in the proof of the global well-posedness and provides evidence that the system is not likely to be ill-posed. Unfortunately, finding an approximation to (1)-(3) for which we can prove existence and then apply the weak stability result has been a challenge and remains an open problem.

The paper is organized as follows: In Section 2, we give two energy functionals that the system (1)-(3) attains. These energy functionals are the key tools that help us obtain complementary estimates on the solutions. We also present these complementary a-priori estimates in Section 2. A Maxwell-Boltzmann entropy functional holds under some constraints on the solutions, mainly that $\rho_A \rho_B < 1$. Moreover, in Section 4, we are interested in understanding the stationary states of (1)-(3), as they outline the possible long-term behavior of the evolution problem. Section 3 is dedicated to the existence analysis. An important question to consider is whether segregated steady states, which are physical in many situations, arise. Based on the natural definition of weak solutions, obtained from the weak stability result, we show that all steady states must be constant. This implies that perhaps the use of the entropy structure is not suitable to study segregated solutions. An alternative is that the model actually does not capture the physical property of segregation. Our final result is on the long-term behavior of solutions to (1)-(3), in the case when the product of the population densities ρ_A, ρ_B are small; see Section 5. We complement these results with some numerical simulations in Section 6.

2 Energy functionals and a-priori estimates

For the sake of simplicity we assume in this section that $2\beta c = 1$. Note that this does not influence the existence analysis as it follows from a simple rescaling of the system. In the following we denote $\rho = (\rho_A, \rho_B)$, and $\Omega \subset \mathbb{R}^2$ is an open, bounded set with Lipschitz boundary.

2.1 Two energy functionals

In this section we present two energy functionals that will be useful in obtaining bounds for ρ_A and ρ_B . Let us define the first energy functional $\mathcal{H}[\rho]$:

$$\mathcal{H}[\rho] = \int_{\Omega} h(\rho_A(x), \rho_B(x)) dx, \quad (8)$$

where

$$h(\rho_A, \rho_B) = \rho_A \log \rho_A - \rho_A + \rho_B \log \rho_B - \rho_B + \rho_A \rho_B.$$

System (1)-(3) is a formal gradient flow with respect to $\mathcal{H}$:

$$\partial_t \rho = \operatorname{div} \left(\mathbb{M} \nabla \frac{\delta \mathcal{H}[\rho]}{\delta \rho} \right), \quad \text{with } \mathbb{M} = \begin{pmatrix} \rho_A & 0 \\ 0 & \rho_B \end{pmatrix}, \quad (9)$$

where $\frac{\delta \mathcal{H}[\rho]}{\delta \rho}$ is the Frechét derivative of $\mathcal{H}$, which can be identified, via Riesz representation theorem, with the gradient of h :

$$\frac{\delta \mathcal{H}[\rho]}{\delta \rho} \simeq Dh(\rho_A, \rho_B) = (\log \rho_A + \rho_B, \log \rho_B + \rho_A)^\top.$$

The matrix $\mathbb{M}$ is positive semidefinite in $\mathbb{R}_+^2 \equiv [0, \infty)^2$. Testing (9) against $Dh(\rho_A, \rho_B)$ yields the energy balance equation:

$$\frac{d\mathcal{H}[\rho]}{dt} + \int_{\Omega} \{ \rho_A |\nabla (\log \rho_A + \rho_B)|^2 + \rho_B |\nabla (\log \rho_B + \rho_A)|^2 \} dx = 0. \quad (10)$$

On the other hand, system (1)-(3) also admits another gradient-flow structure under some restrictions on its solution. Let us define the open set:

$$\mathcal{D} = \{ (\rho_A, \rho_B) \in \mathbb{R}_+^2 \mid \rho_A \rho_B < 1 \},$$

and the Maxwell-Boltzmann entropy functional:

$$\mathcal{H}_{MB}[\rho] = \int_{\Omega} h_{MB}(\rho_A(x), \rho_B(x)) dx,$$

where

$$h_{MB}(\rho_A, \rho_B) = \rho_A \log \rho_A - \rho_A + \rho_B \log \rho_B - \rho_B.$$

Then (1)-(3) can be rewritten as

$$\partial_t \rho = \operatorname{div} \left(\mathbb{M}' \frac{\mathcal{H}_{MB}[\rho]}{\delta \rho} \right), \quad \text{with } \mathbb{M}' = \begin{pmatrix} \rho_A & \rho_A \rho_B \\ \rho_A \rho_B & \rho_B \end{pmatrix}, \quad (11)$$

where

$$\frac{\delta \mathcal{H}_{MB}[\rho]}{\delta \rho} \simeq Dh_{MB}(\rho_A, \rho_B) = (\log \rho_A, \log \rho_B)^\top.$$

We remark that $\mathbb{M}'$ is positive semi-definite on $\overline{\mathcal{D}}$. Testing (11) against $Dh_{MB}(\rho_A, \rho_B)$ yields the balance equation for $\mathcal{H}_{MB}[\rho]$:

$$\begin{aligned}
\frac{d}{dt} \mathcal{H}_{MB}[\rho] &= \int_{\Omega} (\log \rho_A \partial_t \rho_A + \log \rho_B \partial_t \rho_B) dx \\
&= - \int_{\Omega} (\rho_A^{-1} \nabla \rho_A \cdot (\nabla \rho_A + \rho_A \nabla \rho_B) + \rho_B^{-1} \nabla \rho_B \cdot (\nabla \rho_B + \rho_B \nabla \rho_A)) dx \\
&= - \int_{\Omega} (\rho_A^{-1} |\nabla \rho_A|^2 + \rho_B^{-1} |\nabla \rho_B|^2 + 2 \nabla \rho_A \cdot \nabla \rho_B) dx \\
&= -4 \int_{\Omega} (|\nabla \sqrt{\rho_A}|^2 + |\nabla \sqrt{\rho_B}|^2 + 2 \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_A} \cdot \nabla \sqrt{\rho_B}) dx \\
&= -4 \int_{\Omega} (\sqrt{\rho_A \rho_B} |\nabla (\sqrt{\rho_A} + \sqrt{\rho_B})|^2 + (1 - \sqrt{\rho_A \rho_B}) (|\nabla \sqrt{\rho_A}|^2 + |\nabla \sqrt{\rho_B}|^2)) dx.
\end{aligned}$$

Summarizing up gives the following:

$$\frac{d}{dt} \mathcal{H}_{MB}(\rho) + 4 \int_{\Omega} (\sqrt{\rho_A \rho_B} |\nabla (\sqrt{\rho_A} + \sqrt{\rho_B})|^2 + (1 - \sqrt{\rho_A \rho_B}) (|\nabla \sqrt{\rho_A}|^2 + |\nabla \sqrt{\rho_B}|^2)) dx = 0. \quad (12)$$

We point out that (12) is only useful if $\rho_A \rho_B \leq 1$, otherwise we obtain terms we cannot control. Moreover, (12) represents an improvement of the previous entropy inequality, as the factor $1 - \sqrt{\rho_A \rho_B}$ is larger than $(1 - \sqrt{\rho_A \rho_B})^2$ when $\sqrt{\rho_A \rho_B} < 1$.

2.2 A-priori estimates

In this section, we give a-priori estimates on the agent densities ρ_A and ρ_B . The estimates are obtained from energy balance equations (10) and (12).

Throughout the section we assume that the initial datum $\rho^{in} \in L^2(\Omega)$, where $\Omega \in \mathbb{R}^2$ is an open, bounded domain with Lipschitz boundary. As a consequence $\mathcal{H}_{MB}[\rho^{in}] \leq \mathcal{H}[\rho^{in}] < \infty$. Also, we denote $\Omega_T \equiv \Omega \times (0, T)$ for every $T > 0$.

Lemma 2.1 (mass conservation). *System (1)-(3) conserves mass. In particular we have the following estimate:*

$$\|\rho_i\|_{L^\infty(0,T;L^1(\Omega))} = \|\rho_i^{in}\|_{L^1(\Omega)}, \quad i \in \{A, B\}. \quad (13)$$

Proof. Integrating (1)-(3) in Ω yields

$$\int_{\Omega} \rho_i(t) dx = \int_{\Omega} \rho_i^{in} dx, \quad i \in \{A, B\}, \quad t > 0. \quad (14)$$

Thus (13) holds. $\square$

2.2.1 Estimates from “natural” energy balance equation (10).

Lemma 2.2. *We obtain the following estimates for ρ :*

$$\|(1 - \sqrt{\rho_A \rho_B}) \nabla \sqrt{\rho_i}\|_{L^2(\Omega_T)} \leq C, \quad i \in \{A, B\}, \quad (15)$$

$$\|(1 + \sqrt{\rho_A \rho_B}) \nabla (\sqrt{\rho_A} + \sqrt{\rho_B})\|_{L^2(\Omega_T)} \leq \tilde{C}, \quad (16)$$

$$\|(\sqrt{\rho_A \rho_B} - 1)^2\|_{L^{4/3}(0,T;W^{1,4/3}(\Omega))} \leq C_T, \quad (17)$$

where $C, \tilde{C}, C_T > 0$ are some constants, C_T depending on $T > 0$. Moreover, the following estimates hold true for $\sqrt{\rho_A} + \sqrt{\rho_B}$:

$$\|\sqrt{\rho_A} + \sqrt{\rho_B}\|_{L^2(0,T;H^1(\Omega))} \leq \tilde{C}_T, \quad (18)$$

$$\|\sqrt{\rho_A} + \sqrt{\rho_B}\|_{L^4(\Omega_T)} \leq \hat{C}_T, \quad (19)$$

where $\tilde{C}, \hat{C}_T > 0$ some constant depending on $T > 0$.

Proof. Integrating (10) in the time interval $[0, T]$ with $T > 0$ arbitrary leads to

$$\mathcal{H}[\rho(T)] + 4 \int_{\Omega_T} (|\nabla \sqrt{\rho_A} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_B}|^2 + |\nabla \sqrt{\rho_B} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_A}|^2) dx dt \leq \mathcal{H}[\rho^{in}]. \quad (20)$$

However, since $2(x^2 + y^2) \geq (x \pm y)^2$ for every $x, y \in \mathbb{R}$, we deduce

$$\begin{aligned} 2|\nabla \sqrt{\rho_A} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_B}|^2 + 2|\nabla \sqrt{\rho_B} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_A}|^2 &\geq |(1 + \sqrt{\rho_A \rho_B}) \nabla (\sqrt{\rho_A} + \sqrt{\rho_B})|^2 \\ &\geq |\nabla (\sqrt{\rho_A} + \sqrt{\rho_B})|^2, \end{aligned}$$

and

$$2|\nabla \sqrt{\rho_A} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_B}|^2 + 2|\nabla \sqrt{\rho_B} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_A}|^2 \geq |(1 - \sqrt{\rho_A \rho_B}) \nabla (\sqrt{\rho_A} - \sqrt{\rho_B})|^2,$$

and so (15)-(16) hold.

The definition of $\mathcal{H}$ and (20) lead to

$$\|\rho_A \log \rho_A\|_{L^\infty(0,T;L^1(\Omega))} + \|\rho_B \log \rho_B\|_{L^\infty(0,T;L^1(\Omega))} \leq C. \quad (21)$$

Then (18) follows. The following Gagliardo-Nirenberg inequality holds since $\Omega \subset \mathbb{R}^2$:

$$\|u\|_{L^4(\Omega)} \leq C_{GN} \|u\|_{L^2(\Omega)}^{1/2} \|u\|_{H^1(\Omega)}^{1/2}, \text{ for all } u \in H^1(\Omega). \quad (22)$$

Choosing $u = \sqrt{\rho_A} + \sqrt{\rho_B}$ in the above inequality and integrating it in time leads to

$$\int_0^T \|\sqrt{\rho_A} + \sqrt{\rho_B}\|_{L^4(\Omega)}^4 dt \leq C_{GN}^4 \left(\sup_{t \in [0,T]} \|\sqrt{\rho_A}(t) + \sqrt{\rho_B}(t)\|_{L^2(\Omega)}^2 \right) \int_0^T \|\sqrt{\rho_A}(t) + \sqrt{\rho_B}(t)\|_{H^1(\Omega)}^2 dt,$$

which, thanks to (13), (18), leads to (19).

From (15), (19) and the identity

$$\begin{aligned} \frac{1}{2} \nabla [(\sqrt{\rho_A \rho_B} - 1)^2] &= (\sqrt{\rho_A \rho_B} - 1) \nabla \sqrt{\rho_A \rho_B} \\ &= \sqrt{\rho_B} (\sqrt{\rho_A \rho_B} - 1) \nabla \sqrt{\rho_A} + \sqrt{\rho_A} (\sqrt{\rho_A \rho_B} - 1) \nabla \sqrt{\rho_B} \end{aligned}$$

we deduce that

$$\|\nabla [(\sqrt{\rho_A \rho_B} - 1)^2]\|_{L^{4/3}(\Omega_T)} \leq C_T.$$

Since $(\sqrt{\rho_A \rho_B} - 1)^2 \leq C(1 + \rho_A^2 + \rho_B^2)$, from the above estimate and (19), as well as Poincaré's Lemma, we obtain (17). □

Lemma 2.3 (estimate on $\rho_A \rho_B$). *We have the following estimate for the product of the agent densities ρ_A and ρ_B*

$$\|\rho_A \rho_B\|_{L^{3/2}(\Omega_T)} \leq C_T, \quad (23)$$

where $C_T > 0$ is a constant depending on $T > 0$.

Proof. We give the proof by using the so-called H^{-1} method, i.e. by testing (1)-(3) against $\psi \approx (-\Delta)^{-1}(\rho_A + \rho_B)$. For $t \in (0, T)$ define the function $\psi(t)$ as the only solution to

$$\begin{cases} -\Delta\psi(t) = \rho_A(t) + \rho_B(t) - \langle(\rho_A(t) + \rho_B(t))\rangle, & \text{in } \Omega, \\ \partial_\nu\psi(t) = 0 & \text{on } \partial\Omega, \end{cases} \quad (24)$$

and

$$\int_{\Omega} \psi(t) dx = 0,$$

where

$$\langle\rho\rangle \equiv \int_{\Omega} \frac{\rho}{|\Omega|} dx.$$

We remark that $\langle\rho(t)\rangle = \langle\rho^{in}\rangle$ is constant in time thanks to (14). Let us first compute the following:

$$\frac{d}{dt} \int_{\Omega} \frac{1}{2} |\nabla\psi|^2 dx = \int_{\Omega} \nabla\psi \cdot \nabla\partial_t\psi dx = \int_{\Omega} \psi\partial_t(-\Delta\psi) dx = \int_{\Omega} \psi\partial_t(\rho_A + \rho_B) dx.$$

Therefore, testing each equation in (1)-(3) against ψ and summing the equations leads to

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} \frac{1}{2} |\nabla\psi|^2 dx &= - \int_{\Omega} (\nabla\rho_A + \rho_A\nabla\rho_B + \nabla\rho_B + \rho_B\nabla\rho_A) \cdot \nabla\psi dx \\ &= - \int_{\Omega} \nabla(\rho_A + \rho_B + \rho_A\rho_B) \cdot \nabla\psi dx \\ &= - \int_{\Omega} (\rho_A + \rho_B + \rho_A\rho_B) (\rho_A + \rho_B - \langle\rho_A + \rho_B\rangle) dx. \end{aligned}$$

Thanks to the mass conservation (14), the energy balance (10) and the fact that $\rho_A\rho_B \leq C + h(\rho_A, \rho_B)$, we deduce

$$\frac{d}{dt} \int_{\Omega} \frac{1}{2} |\nabla\psi|^2 dx + \int_{\Omega} (\rho_A + \rho_B + \rho_A\rho_B) (\rho_A + \rho_B) dx \leq C,$$

and integrating the above inequality in $[0, T]$ yields

$$\int_{\Omega} |\nabla\psi(T)|^2 dx + \int_0^T \int_{\Omega} (\rho_A + \rho_B + \rho_A\rho_B) (\rho_A + \rho_B) dx dt \leq C_T + \int_{\Omega} |\nabla\psi(0)|^2 dx.$$

Testing (24) against $\psi(t)$ and exploiting Poincaré's Lemma (remember that $\int_{\Omega} \psi(t) dx = 0$) leads to

$$\begin{aligned} \|\nabla\psi(t)\|_{L^2(\Omega)}^2 &\leq \int_{\Omega} (\rho_A(t) + \rho_B(t))\psi(t) dx \leq \|\rho_A(t) + \rho_B(t)\|_{L^2(\Omega)} \|\psi(t)\|_{L^2(\Omega)} \\ &\leq C_P \|\rho_A(t) + \rho_B(t)\|_{L^2(\Omega)} \|\nabla\psi(t)\|_{L^2(\Omega)} \end{aligned}$$

which means

$$\|\nabla\psi(t)\|_{L^2(\Omega)} \leq C_P \|\rho_A(t) + \rho_B(t)\|_{L^2(\Omega)}, \quad t \in [0, T].$$

In particular, since $\rho^{in} \in L^2(\Omega)$ by assumption, it follows that $\|\nabla\psi(0)\|_{L^2(\Omega)} \leq C$, so we conclude that

$$\int_{\Omega} |\nabla\psi(T)|^2 dx + \int_0^T \int_{\Omega} (\rho_A + \rho_B + \rho_A\rho_B) (\rho_A + \rho_B) dx dt \leq C_T.$$

It follows

$$\int_{\Omega_T} (\rho_A + \rho_B) \rho_A \rho_B dx dt \leq C_T,$$

which, by Young's inequality, $\rho_A + \rho_B \geq 2\sqrt{\rho_A\rho_B}$, leads to (23). $\square$

Lemma 2.4 (estimate on the fluxes). *We have the following estimate for the fluxes*

$$\|\nabla \rho_A + \rho_A \nabla \rho_B\|_{L^{4/3}(\Omega_T)} + \|\nabla \rho_B + \rho_B \nabla \rho_A\|_{L^{4/3}(\Omega_T)} \leq C_T. \quad (25)$$

where C_T is a constant depending on $T > 0$.

Proof. Since

$$\begin{aligned} \nabla \rho_A + \rho_A \nabla \rho_B &= 2\sqrt{\rho_A} (\nabla \sqrt{\rho_A} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_B}) \\ &= 2\sqrt{\rho_A} (\nabla(\sqrt{\rho_A} + \sqrt{\rho_B}) + (\sqrt{\rho_A \rho_B} - 1) \nabla \sqrt{\rho_B}), \end{aligned}$$

from (16), (15), (19) it follows

$$\begin{aligned} \|\nabla \rho_A + \rho_A \nabla \rho_B\|_{L^{4/3}(\Omega_T)} &\leq 2\|\sqrt{\rho_A}\|_{L^4(\Omega_T)} (\|\nabla(\sqrt{\rho_A} + \sqrt{\rho_B})\|_{L^2(\Omega_T)} + \|(\sqrt{\rho_A \rho_B} - 1) \nabla \sqrt{\rho_B}\|_{L^2(\Omega_T)}) \\ &\leq C_T. \end{aligned}$$

Since a similar argument can be done for $\nabla \rho_B + \rho_B \nabla \rho_A$, we obtain (25). $\square$

Lemma 2.5 (estimate on $\partial_t \rho_A$ and $\partial_t \rho_B$). *We have the following estimate on the time derivative of ρ_A and ρ_B :*

$$\|\partial_t \rho_A\|_{L^{4/3}(0,T;W^{1,4}(\Omega)')} + \|\partial_t \rho_B\|_{L^{4/3}(0,T;W^{1,4}(\Omega)')} \leq C_T. \quad (26)$$

where C_T is a constant depending on $T > 0$.

Proof. Given any test function ψ , bound (25) yields

$$\langle \partial_t \rho_A, \psi \rangle = - \int_0^T \int_{\Omega} \nabla \psi \cdot (\nabla \rho_A + \rho_A \nabla \rho_B) dx dt \leq C \|\nabla \psi\|_{L^4(\Omega_T)},$$

which means that $\partial_t \rho_A$ is bounded in $L^{4/3}(0, T; W^{1,4}(\Omega)')$. In the same way one proves the same bound for $\partial_t \rho_B$ and obtain (26). $\square$

2.2.2 Additional estimates from Maxwell-Boltzmann energy balance equation (12).

In this subsection we assume that the solution ρ to (1)-(3) satisfies $\rho \in \mathcal{D}$ a.e. in Ω_T . This means that (12) holds. On the other hand, we wish to point out that ρ fulfills also (10), which implies that the estimates derived in the previous subsection are additionally satisfied.

Lemma 2.6. *We have the following estimates on ρ :*

$$\|(1 - \sqrt{\rho_A \rho_B})^{1/2} \nabla \sqrt{\rho_i}\|_{L^2(0,T;L^2(\Omega))} \leq C_T, \quad i \in \{A, B\}, \quad (27)$$

$$\|(1 - \sqrt{\rho_A \rho_B})^{3/2}\|_{L^{4/3}(0,T;W^{1,4/3}(\Omega))} \leq \tilde{C}_T, \quad (28)$$

where $C_T, \tilde{C}_T > 0$ are some constants depending on $T > 0$.

Proof. Integrating (12) in the time interval $[0, T]$ leads to the following

$$\mathcal{H}_{MB}[\rho(T)] + 4 \int_{\Omega_T} (\sqrt{\rho_A \rho_B} |\nabla(\sqrt{\rho_A} + \sqrt{\rho_B})|^2 + (1 - \sqrt{\rho_A \rho_B}) (|\nabla \sqrt{\rho_A}|^2 + |\nabla \sqrt{\rho_B}|^2)) dx dt = \mathcal{H}_{MB}[\rho^{in}], \quad (29)$$

and thus (27). From (19), (27) and the identity

$$\begin{aligned} -\frac{2}{3}\nabla[(1 - \sqrt{\rho_A\rho_B})^{3/2}] &= (1 - \sqrt{\rho_A\rho_B})^{1/2}\nabla\sqrt{\rho_A\rho_B} \\ &= \sqrt{\rho_B}(1 - \sqrt{\rho_A\rho_B})^{1/2}\nabla\sqrt{\rho_A} + \sqrt{\rho_A}(1 - \sqrt{\rho_A\rho_B})^{1/2}\nabla\sqrt{\rho_B} \end{aligned}$$

we deduce that

$$\|\nabla[(1 - \sqrt{\rho_A\rho_B})^{3/2}]\|_{L^{4/3}(\Omega_T)} \leq C_T.$$

From the above estimate and the $L^\infty(\Omega_T)$ bound coming from the assumption $\rho_A\rho_B \leq 1$ a.e. in Ω we obtain (28). $\square$

Remark 2.7. Estimate (27) is an improvement on (15) (since the gradient of $\sqrt{\rho_i}$ is less degenerate in the region $\{\rho_A\rho_B = 1\}$). Similarly, estimate (28) is better than (17) as the bounds for $\nabla\sqrt{\rho_A}$, $\nabla\sqrt{\rho_B}$ are less degenerate.

Remark 2.8. Note that (4) satisfies the following:

$$\begin{aligned} \frac{d\mathcal{H}}{dt} + \frac{1}{4} \int_{\Omega} [\rho_A |\nabla(\log \rho_A + 2\beta g_B)|^2 + \rho_B |\nabla(\log \rho_B + 2\beta g_A)|^2] dx + 2\beta \int_{\Omega} (\rho_A g_B + \rho_B g_A) dx \\ = 4\beta c \int_{\Omega} \rho_A \rho_B dx \\ \leq 2\beta c \int_{\Omega} (\rho_A^2 + \rho_B^2) dx \\ = 2\beta c \left(\|\sqrt{\rho_A}\|_{L^4(\Omega)}^4 + \|\sqrt{\rho_B}\|_{L^4(\Omega)}^4 \right). \end{aligned}$$

3 Existence analysis

In this section we provide results on local-in time existence of strong solutions, define the notion of weak solutions and perform a weak stability analysis.

We consider the scaled equations with homogeneous Neumann boundary conditions in a bounded, open $\Omega \subset \mathbb{R}^2$ with Lipschitz boundary:

$$\begin{cases} \partial_t \rho_A = \operatorname{div}(\nabla \rho_A + \rho_A \nabla \rho_B), & \text{in } \Omega \times (0, \infty), \\ \rho_A(0) = \rho_A^{in}, & \text{in } \Omega. \end{cases} \quad (30)$$

$$\begin{cases} \partial_t \rho_B = \operatorname{div}(\nabla \rho_B + \rho_B \nabla \rho_A), & \text{in } \Omega \times (0, \infty), \\ \rho_B(0) = \rho_B^{in}, & \text{in } \Omega, \end{cases} \quad (31)$$

with

$$\partial_\nu \rho_A = \partial_\nu \rho_B = 0 \quad \text{on } \partial\Omega \times (0, \infty). \quad (32)$$

An analytical study of (30)–(32) is the content of the next part.

3.1 Local-in-time existence of strong solutions.

The diffusion matrix of (30) and (31) is given by

$$A(\rho) = \begin{pmatrix} 1 & \rho_A \\ \rho_B & 1 \end{pmatrix}, \quad \rho_A, \rho_B \geq 0,$$

has eigenvalues $\lambda_{\pm}(\rho) = 1 \pm \sqrt{\rho_A \rho_B}$. Therefore, $\lambda_+(\rho) \geq \lambda_-(\rho) > 0$ for $\rho = (\rho_A, \rho_B) \in \mathcal{D}$, where

$$\mathcal{D} = \{\rho \in \mathbb{R}_+^2 \mid \rho_A \rho_B < 1\}.$$

Applying [18, Thr. 3.1] to (30), (31) yields the following

Lemma 3.1 (Local-in-time existence). *Let $\rho_A^{in}, \rho_B^{in} \in W^{1,p}(\Omega; \mathbb{R}^2)$ for some $p > 2$. Assume that there exists $\epsilon_0 > 0$ such that*

$$\min \{\rho_A^{in}(x), \rho_B^{in}(x), 1 - \rho_A^{in}(x)\rho_B^{in}(x)\} \geq \epsilon_0 \quad a.e. \ x \in \Omega.$$

Then there exists a unique maximal solution ρ to (30)–(32) satisfying $\rho \in C^0([0, T^), W^{1,p}(\Omega; \mathbb{R}^2)) \cap C^\infty((0, T^*); \mathbb{R}^2)$, with $0 < T^* \leq \infty$, and there exists $\epsilon_1 > 0$ such that*

$$\min \{\rho_A(t, x), \rho_B(t, x), 1 - \rho_A(t, x)\rho_B(t, x)\} \geq \epsilon_1 \quad x \in \Omega, \quad t \in (0, T^*).$$

This means that the solution exists as long as its value remains far away from the border of the region $\mathcal{D}$. Unfortunately, it is not clear how to guarantee such property for arbitrary large times.

3.2 Weak solutions

We first give a definition of a weak solution to (30)–(32).

Definition 3.2 (Weak solution). A Lebesgue-measurable function $\rho : \Omega \times (0, T) \rightarrow \mathbb{R}_+^2$ is called a weak solution to (30)–(32) if (and only if) the following properties are satisfied.

(i) It has the regularity:

$$\rho_A, \rho_B \text{ and } \rho_A \rho_B \in L^\infty(0, T; L^1(\Omega)), \quad \partial_t \rho_A, \partial_t \rho_B \in L^{4/3}(0, T; W^{1,4}(\Omega)'), \quad (33)$$

$$(1 + \sqrt{\rho_A \rho_B}) \nabla [\sqrt{\rho_A} + \sqrt{\rho_B}] \in L^2(\Omega \times (0, T)), \quad (34)$$

$$\frac{\sqrt{\rho_A \rho_B} - 1}{1 + \sqrt{\rho_A} + \sqrt{\rho_B}} \in L^2(0, T; H^1(\Omega)), \quad (35)$$

$$\begin{aligned} & f(\sqrt{\rho_A}, \sqrt{\rho_B}) \in L^2(0, T; H^1(\Omega)), \text{ for all } f \in C^1(\mathbb{R}_+^2) \text{ such that} \\ & |f(u_A, u_B)| \leq C(1 + u_A^2 + u_B^2), \text{ and } |Df(u_A, u_B)| \leq C|u_A u_B - 1|, \text{ for all } u_A, u_B \geq 0. \end{aligned} \quad (36)$$

(ii) The following weak formulation of (30)–(32) holds for all $T > 0$:

$$\int_0^T \langle \partial_t \rho_A, \phi \rangle dt + 2 \int_0^T \int_\Omega \nabla \phi \cdot \sqrt{\rho_A} \cdot \zeta_A dx dt = 0 \quad \text{for all } \phi \in L^4(0, T; W^{1,4}(\Omega)), \quad (37)$$

$$\int_0^T \langle \partial_t \rho_B, \phi \rangle dt + 2 \int_0^T \int_\Omega \nabla \phi \cdot \sqrt{\rho_B} \cdot \zeta_B dx dt = 0 \quad \text{for all } \phi \in L^4(0, T; W^{1,4}(\Omega)), \quad (38)$$

$$\rho_A(t) \rightarrow \rho_A^{in}, \quad \rho_B(t) \rightarrow \rho_B^{in} \text{ strongly in } W^{1,4}(\Omega)' \text{ as } t \rightarrow 0, \quad (39)$$

where the quantities $\zeta_A, \zeta_B \in L^2(\Omega \times (0, T))$ are identified by the relations, holding in the sense of distributions,

$$\zeta_A + \zeta_B = (1 + \sqrt{\rho_A \rho_B}) \nabla (\sqrt{\rho_A} + \sqrt{\rho_B}), \quad (40)$$

$$\begin{aligned}\Psi(\sqrt{\rho_A \rho_B})(\zeta_A - \zeta_B) &= \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})(\sqrt{\rho_A} - \sqrt{\rho_B})] \\ &\quad - (\sqrt{\rho_A} - \sqrt{\rho_B}) \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})],\end{aligned}\tag{41}$$

$$\begin{aligned}2 \frac{\Phi(\sqrt{\rho_A \rho_B})}{\sqrt{\rho_A}} g(\sqrt{\rho_A} + \sqrt{\rho_B}) \zeta_A &= \nabla [\Phi(\sqrt{\rho_A \rho_B}) g(\sqrt{\rho_A} + \sqrt{\rho_B}) (\log \rho_A + \rho_B)] \\ &\quad - \frac{1}{2} (\log \rho_A + \rho_B) g(\sqrt{\rho_A} + \sqrt{\rho_B}) \frac{\Phi'(\sqrt{\rho_A \rho_B})}{\sqrt{\rho_A \rho_B} - 1} \nabla [(\sqrt{\rho_A \rho_B} - 1)^2] \\ &\quad - (\log \rho_A + \rho_B) \Phi(\sqrt{\rho_A \rho_B}) g'(\sqrt{\rho_A} + \sqrt{\rho_B}) \nabla [\sqrt{\rho_A} + \sqrt{\rho_B}],\end{aligned}\tag{42}$$

$$\begin{aligned}2 \frac{\Phi(\sqrt{\rho_A \rho_B})}{\sqrt{\rho_B}} g(\sqrt{\rho_A} + \sqrt{\rho_B}) \zeta_B &= \nabla [\Phi(\sqrt{\rho_A \rho_B}) g(\sqrt{\rho_A} + \sqrt{\rho_B}) (\log \rho_B + \rho_A)] \\ &\quad - \frac{1}{2} (\log \rho_B + \rho_A) g(\sqrt{\rho_A} + \sqrt{\rho_B}) \frac{\Phi'(\sqrt{\rho_A \rho_B})}{\sqrt{\rho_A \rho_B} - 1} \nabla [(\sqrt{\rho_A \rho_B} - 1)^2] \\ &\quad - (\log \rho_B + \rho_A) \Phi(\sqrt{\rho_A \rho_B}) g'(\sqrt{\rho_A} + \sqrt{\rho_B}) \nabla [\sqrt{\rho_A} + \sqrt{\rho_B}],\end{aligned}\tag{43}$$

with $g(s) = (1 + s^4)^{-1}$ for $s \geq 0$, for every $\Psi, \Phi \in C^1 \cap W^{1,\infty}(\mathbb{R}_+)$ such that there exists $C > 0$:

$$|\Psi(s)| \leq C|s - 1|, \quad |\Phi(s)| \leq Cs, \quad |\Phi'(s)| \leq Cs|s - 1|, \quad \text{for all } s > 0.$$

(iii) The mass of each species is conserved:

$$\int_{\Omega} \rho_A(t) dx = \int_{\Omega} \rho_A^{in} dx, \quad \int_{\Omega} \rho_B(t) dx = \int_{\Omega} \rho_B^{in} dx, \quad t > 0,\tag{44}$$

and the *integrated energy balance* is satisfied:

$$\mathcal{H}[\rho(T)] + 4 \int_{\Omega_T} (|\zeta_A| + |\zeta_B|^2) dx dt \leq \mathcal{H}[\rho^{in}], \quad \text{for all } T > 0.\tag{45}$$

Remark 3.3. We point out that, if $\sqrt{\rho_A}, \sqrt{\rho_B} \in L^2(0, T; H^1(\Omega))$, which is the nondegenerate case, then $\zeta_A = \nabla \sqrt{\rho_A} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_B}$, $\zeta_B = \nabla \sqrt{\rho_B} + \sqrt{\rho_A \rho_B} \nabla \sqrt{\rho_A}$, as one would expect.

3.3 Weak stability analysis

In this section, we prove the following result.

Lemma 3.4 (Weak stability). *Let $\rho^{in} = (\rho_A^{in}, \rho_B^{in}) : \Omega \rightarrow \mathbb{R}_+^2$ such that $\rho_A^{in}, \rho_B^{in} \in L^2(\Omega)$. Moreover, let $\rho^n = (\rho_A^n, \rho_B^n)$ be a sequence of weak solutions to (30)–(32) having ρ^{in} as initial datum according to Definition 3.2. Assume furthermore that $\sqrt{\rho_A^n}, \sqrt{\rho_B^n} \in L^2(0, T; H^1(\Omega))$ for every $n \in \mathbb{N}$. Then ρ^n converges (up to subsequences) strongly in $L^1(\Omega \times (0, T))$ for every $T > 0$ to a weak solution $\rho = (\rho_A, \rho_B) : \Omega \times (0, \infty) \rightarrow \mathbb{R}_+^2$ to (30)–(32) in the sense of Definition 3.2.*

Remark 3.5. The Lemma implies that the weak solutions described in Definition 3.2 are limit points of standard, *nondegenerate* weak solutions to the system: notice that the assumption $\sqrt{\rho_A^n}, \sqrt{\rho_B^n} \in L^2(0, T; H^1(\Omega))$ for every $n \in \mathbb{N}$, which is not true in general for weak solutions as for Def. 3.2.

Remark 3.6. Another perspective into Lemma 3.4 is the notion that the entropy structure of the system is robust, that is, the estimates provided by the entropy balance inequality are sufficient to show compactness of a suitable sequence of approximated solutions and prove that a limit point of such approximating sequence is a weak solution to the system in the sense of Def. 3.2. Unfortunately, no approximating sequence with the required regularity is known, this is why the global-in-time existence of weak solutions to the system is an open problem.

Proof. By assumption, for every $T > 0$, the approximate solution ρ^n satisfies

$$\int_0^T \langle \partial_t \rho_A^n, \phi \rangle dt + 2 \int_0^T \int_{\Omega} \nabla \phi \cdot \sqrt{\rho_A^n} (\nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}) dx dt = 0, \text{ for all } \phi \in C_c^1(\Omega_T), \quad (46)$$

$$\int_0^T \langle \partial_t \rho_B^n, \phi \rangle dt + 2 \int_0^T \int_{\Omega} \nabla \phi \cdot \sqrt{\rho_B^n} (\nabla \sqrt{\rho_B^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_A^n}) dx dt = 0, \text{ for all } \phi \in C_c^1(\Omega_T), \quad (47)$$

$$\rho_A^n(t) \rightarrow \rho_A^{in}, \quad \rho_B^n(t) \rightarrow \rho_B^{in} \text{ strongly in } W^{-1,4/3}(\Omega) \text{ as } t \rightarrow 0, \quad (48)$$

as well as (20), and therefore also (16)–(26).

Notation. Given a sequence f_n in some Banach space X , that is weakly (or weak-*) convergent in X , we denote with $\overline{f_n}$ the weak (or weak-*) limit of f . Moreover, we define $\mathbb{R}_+ = [0, \infty)$.

The proof is divided into four steps.

Step 1: strong convergence of $\rho_A^n + \rho_B^n$.

Let $f \in W^{1,\infty}(\mathbb{R}_+, \mathbb{R}_+)$ function such that $|f'(x)| \leq C|1-x|$ for $x \geq 0$. Let us define the vector fields

$$\begin{aligned} U_A^n &= (\rho_A^n, -\nabla \rho_A^n - \rho_A^n \nabla \rho_B^n), \\ U_B^n &= (\rho_B^n, -\nabla \rho_B^n - \rho_B^n \nabla \rho_A^n), \\ V^n &= (f(\sqrt{\rho_A^n \rho_B^n}), 0, 0, 0). \end{aligned} \quad (49)$$

From (19), (25) we deduce that for $i \in \{A, B\}$, U_i^n is bounded in $L^{4/3}(\Omega_T)$, while (1)–(3) means that $\operatorname{div}_{(t,x)} U_i^n = 0$ (a fortiori $\operatorname{div}_{(t,x)} U_i^n$ is relatively compact in $W^{-1,r}(\Omega_T)$ for every $r > 1$). On the other hand V^n is bounded in $L^\infty(\Omega_T)$ and the antisymmetric part $\operatorname{curl}_{(t,x)} V^n$ of its Jacobian can be estimated as

$$\begin{aligned} |\operatorname{curl}_{(t,x)} V^n| &\leq C |\nabla f(\sqrt{\rho_A^n \rho_B^n})| \leq C |f'(\sqrt{\rho_A^n \rho_B^n})| (\sqrt{\rho_A^n} |\nabla \sqrt{\rho_B^n}| + \sqrt{\rho_B^n} |\nabla \sqrt{\rho_A^n}|) \\ &\leq C (\sqrt{\rho_A^n} |\sqrt{\rho_A^n \rho_B^n} - 1| |\nabla \sqrt{\rho_B^n}| + \sqrt{\rho_B^n} |\sqrt{\rho_A^n \rho_B^n} - 1| |\nabla \sqrt{\rho_A^n}|), \end{aligned}$$

which means, thanks to (15), (19), that $\operatorname{curl}_{(t,x)} V^n$ is bounded in $L^{4/3}(\Omega_T)$, and a fortiori relatively compact in $W^{-1,r}(\Omega_T)$ for some $r > 1$. Therefore, the Div-Curl Lemma [Theorem 10.21 in [13]] implies that

$$\overline{U^n \cdot V^n} = \overline{U^n} \cdot \overline{V^n} \quad \text{a.e. in } \Omega_T.$$

Therefore, for $i \in \{A, B\}$ and for every $f \in W^{1,\infty}(\mathbb{R}_+, \mathbb{R}_+)$,

$$\overline{\rho_i^n f(\sqrt{\rho_A^n \rho_B^n})} = \overline{\rho_i^n} \overline{f(\sqrt{\rho_A^n \rho_B^n})}, \text{ a.e. in } \Omega_T, \text{ such that } |f'(x)| \leq C|1-x| \text{ for } x \geq 0. \quad (50)$$

Let us consider (50) with $f(x) = \min\{k, F(x)\}$ where $k \geq 1$ is arbitrary and $F \in L_{loc}^1(\mathbb{R}_+, \mathbb{R}_+)$ such that $F' \in L^\infty(\mathbb{R}_+)$, $F(x) \leq C(1+x)$, $|F'(x)| \leq C|1-x|$ for $x \geq 0$. Let us now estimate the quantity

$$\begin{aligned} &\|\overline{\rho_i^n (\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\})} - \overline{F(\sqrt{\rho_A^n \rho_B^n})}\|_{L^1(\Omega_T)} \\ &\leq \liminf_{n \rightarrow \infty} \|\rho_i^n (\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\}) - F(\sqrt{\rho_A^n \rho_B^n})\|_{L^1(\Omega_T)} \\ &\leq \sup_{n \in \mathbb{N}} \|\rho_i^n\|_{L^2(\Omega_T)} \|\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n})\|_{L^2(\Omega_T)}, \end{aligned}$$

where we used Fatou's Lemma in the first inequality. From (19) it follows

$$\begin{aligned} \|\overline{\rho_i^n(\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n}))}\|_{L^1(\Omega_T)}^2 &\leq C \sup_{n \in \mathbb{N}} \int_{\Omega_T \cap \{F(\sqrt{\rho_A^n \rho_B^n}) > k\}} |F(\sqrt{\rho_A^n \rho_B^n})|^2 dx dt \\ &\leq \frac{C}{k} \sup_{n \in \mathbb{N}} \int_{\Omega_T \cap \{F(\sqrt{\rho_A^n \rho_B^n}) > k\}} |F(\sqrt{\rho_A^n \rho_B^n})|^3 dx dt \\ &\leq \frac{C}{k} \sup_{n \in \mathbb{N}} \int_{\Omega_T} (1 + \sqrt{\rho_A^n \rho_B^n})^3 dx dt, \end{aligned}$$

which, thanks to (23), implies

$$\|\overline{\rho_i^n(\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n}))}\|_{L^1(\Omega_T)} \leq \frac{C}{\sqrt{k}}, \quad i \in \{A, B\}, \quad k \geq 1.$$

In a similar way one shows that

$$\|\overline{\rho_i^n(\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n}))}\|_{L^1(\Omega_T)} \leq \frac{C}{\sqrt{k}}, \quad i \in \{A, B\}, \quad k \geq 1.$$

From the above inequalities and (50) we deduce

$$\begin{aligned} &\|\overline{\rho_i^n F(\sqrt{\rho_A^n \rho_B^n})} - \overline{\rho_i^n} \overline{F(\sqrt{\rho_A^n \rho_B^n})}\|_{L^1(\Omega_T)} \\ &\leq \|\overline{\rho_i^n(\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n}))}\|_{L^1(\Omega_T)} + \|\overline{\rho_i^n}(\min\{k, F(\sqrt{\rho_A^n \rho_B^n})\} - F(\sqrt{\rho_A^n \rho_B^n}))\|_{L^1(\Omega_T)} \\ &\leq \frac{C}{\sqrt{k}}, \quad i \in \{A, B\}, \quad k \geq 1, \end{aligned}$$

implying that for $i \in \{A, B\}$,

$$\overline{\rho_i^n F(\sqrt{\rho_A^n \rho_B^n})} = \overline{\rho_i^n} \overline{F(\sqrt{\rho_A^n \rho_B^n})}, \quad \text{a.e. in } \Omega_T, \quad (51)$$

and for every such $F \in L_{loc}^1(\mathbb{R}_+, \mathbb{R}_+)$,

$$F' \in L^\infty(\mathbb{R}_+), \quad F(x) \leq C(1+x), \quad |F'(x)| \leq C|1-x| \text{ for } x \geq 0. \quad (52)$$

Let us now choose $F = F_\delta$ in (51)-(52), where $0 < \delta < 1$ and

$$F_\delta(s) = \begin{cases} 0, & s \leq 1, \\ (s-1)^2, & 1 < s \leq 1+\delta, \\ s + \delta^2 - 1 - \delta, & s > 1+\delta. \end{cases}$$

Let us estimate (similar idea as before)

$$\begin{aligned} \|\overline{\rho_i^n(F_\delta(\sqrt{\rho_A^n \rho_B^n}) - (\sqrt{\rho_A^n \rho_B^n} - 1)_+)}\|_{L^1(\Omega_T)}^2 &\leq \sup_{n \in \mathbb{N}} \|\rho_i^n\|_{L^2(\Omega_T)}^2 \|F_\delta(\sqrt{\rho_A^n \rho_B^n}) - (\sqrt{\rho_A^n \rho_B^n} - 1)_+\|_{L^2(\Omega_T)}^2 \\ &\leq C\delta, \end{aligned}$$

where the last step comes from the fact that $|F_\delta(s) - (s-1)_+| \leq C\delta$ for every $s \geq 0$. We can deduce that (51)-(52) holds with $F(s) = (s-1)_+$, that is

$$\overline{\rho_i^n(\sqrt{\rho_A^n \rho_B^n} - 1)_+} = \overline{\rho_i^n} \overline{(\sqrt{\rho_A^n \rho_B^n} - 1)_+}, \quad i \in \{A, B\} \quad \text{a.e. in } \Omega_T. \quad (53)$$

In a similar way, by writing (51)-(52) with $F = G_\delta$, $0 < \delta < 1$,

$$G_\delta = \begin{cases} s - \delta^2 - 1 + \delta, & 0 \leq s \leq 1 - \delta, \\ -(s - 1)^2, & 1 - \delta < s \leq 1, \\ 0, & s > 1. \end{cases}$$

One deduces that

$$\overline{\rho_i^n (\sqrt{\rho_A^n \rho_B^n} - 1)_-} = \overline{\rho_i^n (\sqrt{\rho_A^n \rho_B^n} - 1)_-}, \quad i \in \{A, B\}, \quad \text{a.e. in } \Omega_T. \quad (54)$$

Summing (53) and (54) allows us to conclude

$$\overline{\rho_i^n \sqrt{\rho_A^n \rho_B^n}} = \overline{\rho_i^n} \overline{\sqrt{\rho_A^n \rho_B^n}}, \quad i \in \{A, B\}, \quad \text{a.e. in } \Omega_T. \quad (55)$$

Let $k \in \mathbb{N}$ arbitrary. Define the vector field

$$Z^n = (\min\{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2, k^2\}, 0, 0, 0), \quad n \in \mathbb{N}.$$

Clearly Z^n is bounded in $L^\infty(\Omega_T)$, while

$$|\operatorname{curl}_{(t,x)} Z^n| \leq C |\nabla [\min\{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2, k^2\}]| \leq Ck |\nabla (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})|,$$

so thanks to (16) $\operatorname{curl}_{(t,x)} Z^n$ is bounded in $L^2(\Omega_T)$ and therefore relatively compact in $W^{-1,r}(\Omega_T)$ for some $r > 1$. The Div-Curl Lemma allows us once again to deduce

$$\overline{U_i^n \cdot Z^n} = \overline{U_i^n} \cdot \overline{Z^n} \quad i \in \{A, B\}, \quad \text{a.e. in } \Omega_T,$$

which is equivalent to

$$\overline{\rho_i^n \min\{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2, k^2\}} = \overline{\rho_i^n} \overline{\min\{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2, k^2\}} \quad i \in \{A, B\}, \quad \text{a.e. in } \Omega_T. \quad (56)$$

Let us define $v^{n,k} = \sqrt{\rho_A^n} + \sqrt{\rho_B^n} - \min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\} = (\sqrt{\rho_A^n} + \sqrt{\rho_B^n} - k)_+$. For $t \in [0, T]$ and $k \geq 2$ let us estimate

$$\begin{aligned} \|v^{n,k}(t)\|_{L^2(\Omega)}^2 &= \int_{\Omega} \left(\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} - k \right)_+^2 dx \\ &\leq \int_{\Omega \cap \{\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} > k\}} \left(\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} \right)^2 dx \\ &\leq \frac{1}{\log k} \int_{\Omega \cap \{\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} > k\}} \left(\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} \right)^2 \log \left(\sqrt{\rho_A^n(t)} + \sqrt{\rho_B^n(t)} \right) dx \\ &\leq \frac{C}{\log k} \int_{\Omega} (\rho_A^n(t) + \rho_B^n(t)) (1 + \log(\rho_A^n(t) + \rho_B^n(t))) dx. \end{aligned}$$

From (13), (21) it follows

$$\|v^{n,k}\|_{L^\infty(0,T;L^2(\Omega))} \leq \frac{C}{\sqrt{\log k}}, \quad n, k \in \mathbb{N}, \quad k \geq 2. \quad (57)$$

From Gagliardo-Nirenberg inequality (22) applied with $u = v^{n,k}$ it follows

$$\int_0^T \|v^{n,k}\|_{L^4(\Omega)}^4 dx dt \leq C \left(\sup_{t \in [0,T]} \|v^{n,k}\|_{L^2(\Omega)}^2 \right) \int_0^T \|v^{n,k}\|_{H^1(\Omega)}^2 dx dt,$$

which, thanks to (57), leads to

$$\|v^{n,k}\|_{L^4(\Omega_T)}^4 \leq \frac{C}{\log k} \|\sqrt{\rho_A^n} + \sqrt{\rho_B^n}\|_{L^2(0,T;H^1(\Omega))}^2.$$

Bound (18) and the definition of $v^{n,k}$ allow us to deduce

$$\|\sqrt{\rho_A^n} + \sqrt{\rho_B^n} - \min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\}\|_{L^4(\Omega_T)} \leq \frac{C}{(\log k)^{1/4}}, \quad n, k \in \mathbb{N}, \quad k \geq 2,$$

which, together with Cauchy-Schwartz inequality

$$\begin{aligned} & \|(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2\|_{L^2(\Omega_T)} \\ & \leq \|\sqrt{\rho_A^n} + \sqrt{\rho_B^n} - \min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\}\|_{L^4(\Omega_T)} \|\sqrt{\rho_A^n} + \sqrt{\rho_B^n} + \min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\}\|_{L^4(\Omega_T)} \end{aligned}$$

and (19), allows us to conclude

$$\|(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2\|_{L^2(\Omega_T)} \leq \frac{C}{(\log k)^{1/4}}, \quad n, k \in \mathbb{N}, \quad k \geq 2. \quad (58)$$

Let $\mathcal{M}(\overline{\Omega_T}) = C(\overline{\Omega_T})'$ the space of Radon measures on $\overline{\Omega_T}$. Since $(\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2$ is bounded in $L^1(\Omega_T)$ (thanks to (19)), then (up to subsequences) it is weak-* convergent in $\mathcal{M}(\overline{\Omega_T})$. Since (56) holds, we can write

$$\begin{aligned} & \|(\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2\|_{\mathcal{M}(\overline{\Omega_T})} \\ & \leq \|(\rho_A^n + \rho_B^n)[(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2]\|_{\mathcal{M}(\overline{\Omega_T})} \\ & \quad + \|(\rho_A^n + \rho_B^n)[(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2]\|_{\mathcal{M}(\overline{\Omega_T})} =: J_1 + J_2. \end{aligned} \quad (59)$$

Let us bound the terms J_1, J_2 . Since the norm in $\mathcal{M}(\overline{\Omega_T})$ is weak-* lower semicontinuous, it follows

$$\begin{aligned} J_1 & \leq \liminf_{n \rightarrow \infty} \|(\rho_A^n + \rho_B^n)[(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2]\|_{\mathcal{M}(\overline{\Omega_T})} \\ & \leq \sup_{n \in \mathbb{N}} \|(\rho_A^n + \rho_B^n)[(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2]\|_{\mathcal{M}(\overline{\Omega_T})}. \end{aligned}$$

Since $L^1(\Omega_T) \hookrightarrow \mathcal{M}(\overline{\Omega_T})$, it follows

$$\begin{aligned} J_1 & \leq C \sup_{n \in \mathbb{N}} \|(\rho_A^n + \rho_B^n)[(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2]\|_{L^1(\Omega_T)} \\ & \leq C \sup_{n \in \mathbb{N}} \|\rho_A^n + \rho_B^n\|_{L^2(\Omega_T)} \|(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\min\{\sqrt{\rho_A^n} + \sqrt{\rho_B^n}, k\})^2\|_{L^2(\Omega_T)}. \end{aligned}$$

Since (19), (58) hold, we obtain

$$J_1 \leq \frac{C}{(\log k)^{1/4}}, \quad n, k \in \mathbb{N}, \quad k \geq 2.$$

In a similar way one can show

$$J_2 \leq \frac{C}{(\log k)^{1/4}}, \quad n, k \in \mathbb{N}, \quad k \geq 2.$$

From the previous two bounds and (59) one concludes

$$\|(\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - (\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2\|_{\mathcal{M}(\overline{\Omega_T})} \leq \frac{C}{(\log k)^{1/4}}, \quad k \geq 2,$$

which means

$$(\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 = \overline{(\rho_A^n + \rho_B^n)} \overline{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2} \quad \text{in } \mathcal{M}(\overline{\Omega_T}).$$

For every $f \in \mathcal{M}(\overline{\Omega_T})$, $\phi \in C(\overline{\Omega_T})$, let $\langle f, \phi \rangle$ be the dual product between f , ϕ (i.e. $\langle f, \phi \rangle$ is the result of the application of the linear, bounded functional f to ϕ). It follows

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\Omega_T} (\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 dx dt &= \langle (\rho_A^n + \rho_B^n)(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2, 1 \rangle \\ &= \int_{\Omega_T} \overline{(\rho_A^n + \rho_B^n)} \overline{(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2} dx dt. \end{aligned} \quad (60)$$

On the other hand, summing (55) in $i \in \{A, B\}$, multiplying it times 2 and integrating it in Ω_T leads to

$$\lim_{n \rightarrow \infty} \int_{\Omega_T} (\rho_A^n + \rho_B^n) 2\sqrt{\rho_A^n \rho_B^n} dx dt = \int_{\Omega_T} \overline{(\rho_A^n + \rho_B^n)} 2\sqrt{\rho_A^n \rho_B^n} dx dt. \quad (61)$$

Taking the difference between (60) and (61) yields

$$\lim_{n \rightarrow \infty} \int_{\Omega_T} (\rho_A^n + \rho_B^n)^2 dx dt = \int_{\Omega_T} (\overline{\rho_A^n + \rho_B^n})^2 dx dt,$$

which means (thanks to [13, Thr. 10.20]) that $\rho_A^n + \rho_B^n$ is strongly convergent in $L^2(\Omega_T)$.

Step 2: strong convergence of $\rho_A^n \rho_B^n$. Let

$$\phi(s) = \begin{cases} e^{1+1/(s^2-1)} & 0 \leq s < 1 \\ 0 & s \geq 1 \end{cases}.$$

For every $r > 0$, $u \in \mathbb{R}_+^2$, let us define the function

$$f_{(r,u)}(\rho) = \phi\left(\frac{|\rho - u|}{r}\right), \quad \rho \in \mathbb{R}^2.$$

We define also

$$\begin{aligned} \Gamma_{cr} &= \{(\rho_1, \rho_2) \in \mathbb{R}_+^2 \mid \rho_1 \rho_2 = 1\}, \\ \mathcal{F} &= \left\{ f_{(r,u)} \mid r \in \mathbb{Q} \cap (0, \infty), u \in (\mathbb{Q} \cap [0, \infty))^2, \overline{B_r(u)} \cap \Gamma_{cr} = \emptyset \right\}. \end{aligned}$$

We point out that $\mathcal{F}$ is a countable family of $C_c^\infty(\mathbb{R}^2)$ functions whose gradient vanishes in a neighbourhood of Γ_{cr} . This fact, together with (15), easily implies that

$$\|f_{(r,u)}(\rho^n)\|_{L^2(0,T;H^1(\Omega))} \leq C(r, u, T). \quad (62)$$

Moreover,

$$\begin{aligned} \frac{1}{2} \partial_t f_{(r,u)}(\rho^n) &= \frac{1}{2} \frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_A} \partial_t \rho_A^n + \frac{1}{2} \frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_B} \partial_t \rho_B^n \\ &= \operatorname{div} \left(\frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_A} \sqrt{\rho_A^n} (\nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}) + \frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_B} \sqrt{\rho_B^n} (\nabla \sqrt{\rho_B^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_A^n}) \right) \\ &\quad - \nabla \frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_A} \cdot \sqrt{\rho_A^n} (\nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}) - \nabla \frac{\partial f_{(r,u)}(\rho^n)}{\partial \rho_B} \cdot \sqrt{\rho_B^n} (\nabla \sqrt{\rho_B^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_A^n}) \\ &=: \operatorname{div} \left(\mathcal{J}_{(r,u)}^n \right) + \Xi_{(r,u)}^n. \end{aligned}$$

Once again, the assumptions on $f_{(r,u)}$ as well as (15) imply that

$$\|\mathcal{J}_{(r,u)}^n\|_{L^2(\Omega_T)} + \|\Xi_{(r,u)}^n\|_{L^1(\Omega_T)} \leq C(r, u, T),$$

which leads to

$$\|\partial_t f_{(r,u)}(\rho^n)\|_{L^1(0,T;W^{-1,1}(\Omega))} \leq C(r, u, T). \quad (63)$$

We are therefore allowed to apply Aubin-Lions Lemma (and the uniform $L^\infty(\Omega_T)$ bound for $f_{(r,u)}(\rho^n)$) to deduce the strong convergence of $f_{(r,u)}(\rho^n)$ in $L^q(\Omega_T)$ for every $q < \infty$. In particular,

For all $f_{(r,u)} \in \mathcal{F}$, there exists $(\rho^{n_k(r,u)})_{k \in \mathbb{N}} \subset (\rho^n)_{n \in \mathbb{N}}$ such that $f_{(r,u)}(\rho^{n_k(r,u)})$ is a.e. convergent in Ω_T .

However, since $\mathcal{F}$ is countable, a Cantor diagonal argument allows us to find a subsequence (not relabeled) of ρ^n such that

$$\text{For all } f_{(r,u)} \in \mathcal{F}, \quad f_{(r,u)}(\rho^n) \rightarrow \xi_{(r,u)} \quad \text{a.e. in } \Omega_T.$$

Let $(x, t) \in \Omega_T$ be a point where such convergence holds true. Since (from Step 1) $\rho_A^n + \rho_B^n$ is (up to subsequences) strongly convergent in $L^2(\Omega_T)$, we can assume w.l.o.g. that $\rho^n(x, t)$ is bounded in $\mathbb{R}^2$. There are two cases.

Case 1: There exists $f_{(r_1, u_1)} \in \mathcal{F}$ such that $\xi_{(r_1, u_1)}(x, t) > 0$.

Since $f_{(r,u)}(\rho) = \phi(|u - \rho|/r)$, and $\phi|_{[0,1]}$ is one-to-one and continuous, then $|\rho^n(x, t) - u_1| \rightarrow d_1 \in [0, r)$. By continuity there exist $f_{(r_2, u_2)}, f_{(r_3, u_3)} \in \mathcal{F}$ such that u_1, u_2, u_3 are not aligned and $\xi_{(r_2, u_2)}(x, t), \xi_{(r_3, u_3)}(x, t) > 0$. Since $f_{(r,u)}(\rho)$ is a one-to-one function of $|\rho - u|$ for $|\rho - u| \leq 1$, the limits $d_i = \lim_{n \rightarrow \infty} |\rho^n(x, t) - u_i|$, $i = 1, 2, 3$, exist finite. As a consequence, each accumulation point of $\rho^n(x, t)$ will fall at the intersection of three circles $(\partial B_{d_i}(u_i))$, $i = 1, 2, 3$ with mutually not aligned centers, which can only consist of at most one point. This means that all accumulation points coincide with this point, i.e. the sequence $\rho^n(x, t)$ is convergent to that point. A fortiori $\rho_A^n(x, t)\rho_B^n(x, t)$ is also convergent.

Case 2: For all $f_{(r,u)} \in \mathcal{F}$ it holds $\xi_{(r,u)}(x, t) = 0$.

Consider a generic subsequence $\rho^{n_m}(x, t)$ of $\rho^n(x, t)$. Since it is bounded, it has a sub-subsequence $\rho^{n_{m_k}}(x, t)$ that is convergent to some limit $\ell \in \mathbb{R}_+^2$. However, since $f_{(r,u)}(\ell) = 0$ for every $f_{(r,u)} \in \mathcal{F}$, the only possibility is that $\ell \in \Gamma_{cr}$. In particular $\lim_{k \rightarrow \infty} \rho_A^{n_{m_k}} \rho_B^{n_{m_k}}(x, t) = 1$. Being the subsequence $\rho^{n_m}(x, t)$ arbitrary, this means that $\rho_A^n \rho_B^n(x, t) \rightarrow 1$ as $n \rightarrow \infty$.

Summarizing up, we have proved that $\rho_A^n \rho_B^n$ is a.e. convergent in Ω_T . Bound (23) implies that $\rho_A^n \rho_B^n$ is strongly convergent in $L^{3/2-\eta}(\Omega_T)$ for every $\eta \in (0, 1/2]$. Furthermore, we have also showed that $\rho_A^n \rightarrow \rho_A, \rho_B^n \rightarrow \rho_B$ a.e. in E , where

$$E = \left\{ (x, t) \in \Omega_T \mid \lim_{n \rightarrow \infty} \rho_A^n(x, t) \rho_B^n(x, t) \neq 1 \right\}.$$

This also implies (together with (19)) that $\rho_A^n \rightarrow \rho_A, \rho_B^n \rightarrow \rho_B$ strongly in $L^{2-\eta}(E)$, for every $\eta \in (0, 1]$.

We also point out that $|\sqrt{\rho_A^n} \pm \sqrt{\rho_B^n}| = \sqrt{\rho_A^n + \rho_B^n \pm 2\sqrt{\rho_A^n \rho_B^n}}$ is a.e. convergent in Ω_T , and so is $|\rho_A^n - \rho_B^n| = |\sqrt{\rho_A^n} + \sqrt{\rho_B^n}| |\sqrt{\rho_A^n} - \sqrt{\rho_B^n}|$.

Step 3: strong convergence of ρ_A^n, ρ_B^n . Now we must prove that ρ_A^n, ρ_B^n are a.e. convergent also in $E^c = \{\lim_{n \rightarrow \infty} \rho_A^n \rho_B^n = 1\}$. To this aim, let $\psi \in C^2(\mathbb{R})$ be a cutoff such that

$$\psi(s) = \begin{cases} 0 & s < \frac{1}{3}, \\ 1 & s > \frac{2}{3}, \\ \text{nondecreasing} & \frac{1}{3} \leq s \leq \frac{2}{3}. \end{cases} \quad (64)$$

Moreover define

$$g(s) = \frac{1}{1+s^4}, \quad s \geq 1, \quad (65)$$

$$f_A(\rho^n) = \psi(\sqrt{\rho_A^n \rho_B^n})g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_A^n + \rho_B^n), \quad (66)$$

$$f_B(\rho^n) = \psi(\sqrt{\rho_A^n \rho_B^n})g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_B^n + \rho_A^n). \quad (67)$$

Since $\psi(s)s^{-\alpha}$ is bounded for every $\alpha \geq 0$ and $|\log s| \leq C(s^{-1/8} + s)$, it holds

$$\begin{aligned} |f_A(\rho^n)| &\leq \frac{\psi(\sqrt{\rho_A^n \rho_B^n})g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})}{(\rho_A^n \rho_B^n)^{1/4}} ((\rho_B^n)^{1/4}(\rho_A^n)^{1/4}|\log \rho_A^n| + (\rho_A^n)^{1/4}(\rho_B^n)^{5/4}) \\ &\leq C \frac{(\rho_B^n)^{1/4}[(\rho_A^n)^{1/8} + (\rho_A^n)^{5/4}] + (\rho_A^n)^{1/4}(\rho_B^n)^{5/4}}{1 + (\rho_A^n)^2 + (\rho_B^n)^2} \\ &\leq C, \end{aligned} \quad (68)$$

which means that $f_A(\rho^n)$ is bounded in $L^\infty(\Omega \times (0, T))$. Similarly one shows the same bound for $f_B(\rho^n)$. Let us now consider

$$\nabla f_A(\rho^n) = I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &:= \psi(\sqrt{\rho_A^n \rho_B^n})g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})\nabla(\log \rho_A^n + \rho_B^n) \\ I_2 &:= \psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n)\nabla g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}), \\ I_3 &:= g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_A^n + \rho_B^n)\nabla \psi(\sqrt{\rho_A^n \rho_B^n}). \end{aligned}$$

First we give an estimate to I_1 . From (20) it follows

$$\begin{aligned} \|I_1\|_{L^2(\Omega_T)} &= 2 \left\| \frac{\psi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n \rho_B^n}} g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \sqrt{\rho_B^n} (\nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}) \right\|_{L^2(\Omega_T)} \\ &\leq 2 \left\| \frac{\psi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n \rho_B^n}} \right\|_{L^\infty(\Omega_T)} \|g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \sqrt{\rho_B^n}\|_{L^\infty(\Omega_T)} \|\nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}\|_{L^2(\Omega_T)} \\ &\leq C_T. \end{aligned}$$

Now, we consider I_2 . Since g'/g is bounded in $\mathbb{R}_+$ while $f_A(\rho^n)$ is bounded in $L^\infty(\Omega_T)$,

$$\|I_2\|_{L^2(\Omega_T)} = \left\| f_A(\rho^n) \frac{g'(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})}{g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})} \nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \right\|_{L^2(\Omega)} \leq C \|\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})\|_{L^2(\Omega_T)} \leq C_T$$

where the last inequality comes from (16).

Finally, let us consider I_3 . Since $|\psi'(s)| \leq Cs^{1/4}|1-s|$ for $s \geq 0$ one obtains

$$\begin{aligned} \|I_3\|_{L^2(\Omega_T)} &= \|g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_A^n + \rho_B^n)\psi'(\sqrt{\rho_A^n \rho_B^n})\nabla(\sqrt{\rho_A^n \rho_B^n})\|_{L^2(\Omega_T)} \\ &\leq \|g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_A^n + \rho_B^n)(\sqrt{\rho_A^n \rho_B^n})^{1/4}|1 - \sqrt{\rho_A^n \rho_B^n}|(\sqrt{\rho_A^n}|\nabla \sqrt{\rho_B^n}| + \sqrt{\rho_B^n}|\nabla \sqrt{\rho_A^n}|)\|_{L^2(\Omega_T)} \\ &\leq \|g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})(\log \rho_A^n + \rho_B^n)(\rho_A^n \rho_B^n)^{1/8}\|_{L^\infty(\Omega_T)} \\ &\quad \times \|(1 - \sqrt{\rho_A^n \rho_B^n})(|\nabla \sqrt{\rho_A^n}| + |\nabla \sqrt{\rho_B^n}|)\|_{L^2(\Omega_T)} \leq C_T, \end{aligned}$$

where we used (15), (19). We conclude that $\nabla f_A(\rho^n)$ is bounded in $L^2(\Omega_T)$. Similarly, one can show that $\nabla f_B(\rho^n)$ is bounded in $L^2(\Omega \times (0, T))$, too. This means that the vector fields

$$Y_i^n = (f_i(\rho^n), 0, 0, 0), \quad i \in \{A, B\},$$

are bounded in $L^\infty(\Omega)$ and the antisymmetric part of their Jacobian is bounded in $L^2(\Omega_T)$, thus relatively compact in $W^{-1,r}(\Omega_T)$ for some $r > 1$. Once again, the Div-Curl Lemma (applied to $U_A^n - U_B^n$ defined in (49) and $Y_A^n - Y_B^n$) leads to

$$\overline{(\rho_A^n - \rho_B^n)(f_A(\rho^n) - f_B(\rho^n))} = \overline{(\rho_A^n - \rho_B^n)} \overline{(f_A(\rho^n) - f_B(\rho^n))},$$

which means

$$\frac{(\rho_A^n - \rho_B^n)\psi(\rho_A^n \rho_B^n)}{1 + (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^4} \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right) = \overline{(\rho_A^n - \rho_B^n)} \frac{\psi(\rho_A^n \rho_B^n)}{1 + (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^4} \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right). \quad (69)$$

Let

$$\begin{aligned} \mu^n &= \min\{\rho_A^n, \rho_B^n\} = \frac{1}{2}(\rho_A^n + \rho_B^n) - \frac{1}{2}|\rho_A^n - \rho_B^n|, \\ M^n &= \max\{\rho_A^n, \rho_B^n\} = \frac{1}{2}(\rho_A^n + \rho_B^n) + \frac{1}{2}|\rho_A^n - \rho_B^n|, \end{aligned}$$

and

$$\sigma^n = \begin{cases} 1, & \rho_A^n > \rho_B^n, \\ 0, & \rho_A^n = \rho_B^n, \\ -1, & \rho_A^n < \rho_B^n. \end{cases}$$

Since both $\rho_A^n + \rho_B^n$ and $|\rho_A^n - \rho_B^n|$ are a.e. convergent in $\Omega \times (0, T)$, then also μ^n, M^n are a.e. convergent in $\Omega \times (0, T)$ (and strongly convergent in $L^{4-\eta}(\Omega \times (0, T))$ for every $\eta > 0$) towards some nonnegative functions μ, M (respectively). On the other hand $|\sigma^n| \leq 1$ a.e. in $\Omega \times (0, T)$ and so $\sigma^n \rightharpoonup^* \sigma$ weakly* in $L^\infty(\Omega \times (0, T))$.

Let us point out that

$$(\rho_A^n - \rho_B^n) \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right) = (M^n - \mu^n) \left(\log \frac{M^n}{\mu^n} + \mu^n - M^n \right)$$

which implies that $(\rho_A^n - \rho_B^n) \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right)$ is a.e. convergent on $\{\mu > 0\}$. On the other hand, since $\psi(\rho_A^n \rho_B^n) > 0$ only on $\{\rho_A^n \rho_B^n \geq 1/3\} = \{1/\mu^n \leq 3M^n\}$, it follows (via Lagrange's theorem) that for a suitable point $\lambda^n \in [\mu^n, M^n]$

$$\psi(\sqrt{M^n \mu^n})(M^n - \mu^n) \log \frac{M^n}{\mu^n} = \psi(\sqrt{M^n \mu^n}) \frac{(M^n - \mu^n)^2}{\lambda^n} \leq \psi(\sqrt{M^n \mu^n}) \frac{(M^n - \mu^n)^2}{\mu^n} \leq C|M^n|^3.$$

Since M^n, μ^n are bounded in $L^2(\Omega \times (0, T))$, it follows (by dominated convergence) that

$$\frac{(\rho_A^n - \rho_B^n)\psi(\rho_A^n \rho_B^n)}{1 + (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^4} \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right) \rightarrow \frac{(M - \mu)\psi(M\mu)}{1 + (\sqrt{M} + \sqrt{\mu})^4} \left(\log \frac{M}{\mu} + \mu - M \right) \text{ strongly in } L^1(\Omega_T).$$

A similar argument allows us to obtain

$$\frac{\psi(M^n \mu^n)}{1 + (\sqrt{M^n} + \sqrt{\mu^n})^4} \left(\log \frac{M^n}{\mu^n} + \mu^n - M^n \right) \rightarrow \frac{\psi(M\mu)}{1 + (\sqrt{M} + \sqrt{\mu})^4} \left(\log \frac{M}{\mu} + \mu - M \right) \text{ strongly in } L^1(\Omega_T).$$

As a consequence,

$$\rho_A^n - \rho_B^n = \sigma^n(M^n - \mu^n) \rightharpoonup \sigma(M - \mu) \text{ weakly in } L^1(\Omega \times (0, T)),$$

$$\begin{aligned} \frac{\psi(\rho_A^n \rho_B^n)}{1 + (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^4} \left(\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \right) &= \sigma^n \frac{\psi(M^n \mu^n)}{1 + (\sqrt{M^n} + \sqrt{\mu^n})^4} \left(\log \frac{M^n}{\mu^n} + \mu^n - M^n \right) \\ &\rightharpoonup \sigma \frac{\psi(M\mu)}{1 + (\sqrt{M} + \sqrt{\mu})^4} \left(\log \frac{M}{\mu} + \mu - M \right) \text{ weakly in } L^1(\Omega \times (0, T)). \end{aligned}$$

From the above relations and (69) as well as the fact that

$$M\mu = 1, \quad \log \frac{M}{\mu} + \mu - M = 0 \quad \Leftrightarrow \quad M = \mu = 1,$$

we deduce that $|\sigma|^2 = 1$ in $\{M > \mu, M\mu = 1\}$. However, since $|\sigma^n| \leq 1$ a.e. in Ω_T , $n \in \mathbb{N}$, this means that

$$\overline{(\sigma^n)^2} \leq 1 = |\sigma|^2 = \overline{(\sigma^n)^2} \quad \text{a.e. in } \{M > \mu, M\mu = 1\}.$$

However, being $x \mapsto x^2$ strictly convex, it follows from [13, Thr. 10.20] that $\sigma^n \rightarrow \sigma$ a.e. in $\{M > \mu, M\mu = 1\}$. This allows us to deduce that $\rho_A^n - \rho_B^n = \sigma^n(M^n - \mu^n)$ is a.e. convergent in $\{M\mu = 1\} = E^c$, and so are $\rho_A^n = \frac{1}{2}(\rho_A^n + \rho_B^n) + \frac{1}{2}(\rho_A^n - \rho_B^n)$ and $\rho_B^n = \frac{1}{2}(\rho_A^n + \rho_B^n) - \frac{1}{2}(\rho_A^n - \rho_B^n)$ (because we already know that $\rho_A^n + \rho_B^n$ is a.e. convergent in Ω_T). Since we already knew that ρ_A^n, ρ_B^n are a.e. convergent in E , we conclude that ρ_A^n, ρ_B^n are a.e. convergent in Ω_T and therefore by dominated convergence (and (19)) ρ_A^n, ρ_B^n are also strongly convergent in $L^{2-\delta}(\Omega_T)$ for every $\delta \in (0, 1]$.

Step 4: limit in the equations. Now we show that (37), (38) hold for the limit functions ρ_A, ρ_B . We first study the convergence of the expressions

$$\zeta_A^n = \nabla \sqrt{\rho_A^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_B^n}, \text{ and } \zeta_B^n = \nabla \sqrt{\rho_B^n} + \sqrt{\rho_A^n \rho_B^n} \nabla \sqrt{\rho_A^n}.$$

Since ζ_A^n, ζ_B^n are bounded in $L^2(\Omega_T)$, it holds (up to subsequences)

$$\zeta_A^n \rightharpoonup \zeta_A, \quad \zeta_B^n \rightharpoonup \zeta_B \quad \text{weakly in } L^2(\Omega_T).$$

We want to show that (40)–(43) hold. Let us consider

$$\zeta_A^n + \zeta_B^n = (\sqrt{\rho_A^n \rho_B^n} + 1) \nabla (\sqrt{\rho_A^n} + \sqrt{\rho_B^n}).$$

We know that $\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})$ is bounded in $L^2(\Omega \times (0, T))$ and that $\sqrt{\rho_A^n} + \sqrt{\rho_B^n} \rightarrow \sqrt{\rho_A} + \sqrt{\rho_B}$ strongly in $L^2(\Omega \times (0, T))$, therefore

$$\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \rightharpoonup \nabla(\sqrt{\rho_A} + \sqrt{\rho_B}) \text{ weakly in } L^2(\Omega_T).$$

Moreover, thanks to (16), $J_A^n + J_B^n$ is bounded in $L^2(\Omega_T)$, while $\sqrt{\rho_A^n \rho_B^n} \rightarrow \sqrt{\rho_A \rho_B}$ strongly in $L^{3-\delta}(\Omega_T)$ for every $\delta \in (0, 2]$ (thanks to (23)). Therefore

$$\zeta_A^n + \zeta_B^n \rightharpoonup (\sqrt{\rho_A \rho_B} + 1) \nabla(\sqrt{\rho_A} + \sqrt{\rho_B}) \text{ weakly in } L^2(\Omega_T).$$

Therefore, (40) holds. Let us now turn our attention to

$$\zeta_A^n - \zeta_B^n = (1 - \sqrt{\rho_A^n \rho_B^n}) \nabla(\sqrt{\rho_A^n} - \sqrt{\rho_B^n}).$$

Let us consider a generic function $\Psi \in C^1 \cap W^{1,\infty}(\mathbb{R}_+)$ such that $|\Psi(s)| \leq C|s-1|$ for $s \geq 0$. It holds

$$\begin{aligned} \Psi(\sqrt{\rho_A^n \rho_B^n})(\zeta_A^n - \zeta_B^n) &= \nabla[\Psi(\sqrt{\rho_A^n \rho_B^n})(1 - \sqrt{\rho_A^n \rho_B^n})(\sqrt{\rho_A^n} - \sqrt{\rho_B^n})] \\ &\quad - (\sqrt{\rho_A^n} - \sqrt{\rho_B^n}) \nabla[(1 - \sqrt{\rho_A^n \rho_B^n})\Psi(\sqrt{\rho_A^n \rho_B^n})]. \end{aligned}$$

However, due to (19), (23) we have

$$\Psi(\sqrt{\rho_A^n \rho_B^n})(1 - \sqrt{\rho_A^n \rho_B^n})(\sqrt{\rho_A^n} - \sqrt{\rho_B^n}) \rightharpoonup \Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})(\sqrt{\rho_A} - \sqrt{\rho_B}) \text{ weakly in } L^{12/7}(\Omega_T),$$

while, thanks to (15), (19) we have

$$\nabla[(1 - \sqrt{\rho_A^n \rho_B^n})\Psi(\sqrt{\rho_A^n \rho_B^n})] \rightharpoonup \nabla[(1 - \sqrt{\rho_A \rho_B})\Psi(\sqrt{\rho_A \rho_B})] \text{ weakly in } L^{4/3}(\Omega_T),$$

so it follows that

$$\begin{aligned} \Psi(\sqrt{\rho_A \rho_B})(\zeta_A - \zeta_B) &= \text{weak} \lim_{n \rightarrow \infty} \Psi_\varepsilon(\sqrt{\rho_A^n \rho_B^n})(\zeta_A^n - \zeta_B^n) \\ &= \nabla[\Psi_\varepsilon(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})(\sqrt{\rho_A} - \sqrt{\rho_B})] - (\sqrt{\rho_A} - \sqrt{\rho_B}) \nabla[(1 - \sqrt{\rho_A \rho_B})\Psi_\varepsilon(\sqrt{\rho_A \rho_B})], \end{aligned}$$

implying that (41) holds.

Let now consider $\Phi \in C^1 \cap W^{1,\infty}(\mathbb{R}_+)$ such that $|\Phi(s)| \leq Cs$ for $s \geq 0$ and $s \in [0, \infty) \mapsto \frac{\Phi'(s)}{s(s-1)} \in \mathbb{R}$ is continuous and bounded. The function $\Phi(\sqrt{\rho_A^n \rho_B^n})g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})/\sqrt{\rho_A^n}$ (with g defined in (65)) is bounded in $L^\infty(\Omega_T)$ since

$$0 \leq \frac{\Phi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n}} g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) = \frac{\Phi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n \rho_B^n}} \frac{\sqrt{\rho_B^n}}{1 + (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^4} \leq C.$$

Therefore, it is strongly convergent (in $L^q(\Omega)$ for every $q < \infty$) to $\psi(\sqrt{\rho_A \rho_B})g(\sqrt{\rho_A} + \sqrt{\rho_B})/\sqrt{\rho_A}$, so that

$$\frac{\Phi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n}} g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \zeta_A^n \rightharpoonup \frac{\Phi(\sqrt{\rho_A \rho_B})}{\sqrt{\rho_A}} g(\sqrt{\rho_A} + \sqrt{\rho_B}) \zeta_A \text{ weakly in } L^2(\Omega_T). \quad (70)$$

On the other hand, (66) implies

$$\begin{aligned} \frac{\Phi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n}} g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \zeta_A^n &= \Phi(\sqrt{\rho_A^n \rho_B^n}) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \nabla(\log \rho_A^n + \rho_B^n) \\ &= \nabla[\Phi(\sqrt{\rho_A^n \rho_B^n}) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) (\log \rho_A^n + \rho_B^n)] \\ &\quad - (\log \rho_A^n + \rho_B^n) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \nabla \Phi(\sqrt{\rho_A^n \rho_B^n}) \\ &\quad - (\log \rho_A^n + \rho_B^n) \Phi(\sqrt{\rho_A^n \rho_B^n}) \nabla g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \\ &= \nabla[\Phi(\sqrt{\rho_A^n \rho_B^n}) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) (\log \rho_A^n + \rho_B^n)] \\ &\quad - (\log \rho_A^n + \rho_B^n) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \frac{\Phi'(\sqrt{\rho_A^n \rho_B^n})}{2(\sqrt{\rho_A^n \rho_B^n} - 1)} \nabla(\sqrt{\rho_A^n \rho_B^n} - 1)^2 \\ &\quad - (\log \rho_A^n + \rho_B^n) \Phi(\sqrt{\rho_A^n \rho_B^n}) g'(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}). \end{aligned} \quad (71)$$

The term $\Phi(\sqrt{\rho_A^n \rho_B^n}) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) (\log \rho_A^n + \rho_B^n)$ can be estimated as in (68) to show that it is bounded in $L^\infty(\Omega)$, so thanks to the strong convergence of ρ^n and the continuity of the involved functions we deduce

$$\begin{aligned} \nabla[\Phi(\sqrt{\rho_A^n \rho_B^n}) g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) (\log \rho_A^n + \rho_B^n)] &\rightharpoonup \nabla[\Phi(\sqrt{\rho_A \rho_B}) g(\sqrt{\rho_A} + \sqrt{\rho_B}) (\log \rho_A + \rho_B)] \\ &\text{weakly in } L^q(0, T; W^{-1,q}(\Omega)), \text{ for all } q \in (1, \infty). \end{aligned}$$

Moreover

$$\nabla(\sqrt{\rho_A^n \rho_B^n} - 1)^2 \rightharpoonup \nabla(\sqrt{\rho_A \rho_B} - 1)^2 \text{ weakly in } L^{4/3}(\Omega_T).$$

Furthermore, repeating the estimates in (68) with $\Phi'(\sqrt{\rho_A^n \rho_B^n})/2(\sqrt{\rho_A^n \rho_B^n} - 1)$ in place of $\psi(\sqrt{\rho_A^n \rho_B^n})$ allows us to deduce that the factor $(\log \rho_A^n + \rho_B^n)g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})\Phi'(\sqrt{\rho_A^n \rho_B^n})/2(\sqrt{\rho_A^n \rho_B^n} - 1)$ is bounded in $L^\infty(\Omega_T)$, and therefore

$$\begin{aligned} (\log \rho_A^n + \rho_B^n)g(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})\frac{\Phi'(\sqrt{\rho_A^n \rho_B^n})}{2(\sqrt{\rho_A^n \rho_B^n} - 1)} &\rightarrow (\log \rho_A + \rho_B)g(\sqrt{\rho_A} + \sqrt{\rho_B})\frac{\Phi'(\sqrt{\rho_A \rho_B})}{2(\sqrt{\rho_A \rho_B} - 1)} \\ &\text{strongly in } L^q(\Omega_T), \text{ for all } q < \infty. \end{aligned}$$

Finally, repeating the estimates in (68) with g replaced by g' allows us to deduce that $(\log \rho_A^n + \rho_B^n)\Phi(\sqrt{\rho_A^n \rho_B^n})g'(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})$ is bounded in $L^\infty(\Omega_T)$, so

$$\begin{aligned} (\log \rho_A^n + \rho_B^n)\Phi(\sqrt{\rho_A^n \rho_B^n})g'(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) &\rightarrow (\log \rho_A + \rho_B)\Phi(\sqrt{\rho_A \rho_B})g'(\sqrt{\rho_A} + \sqrt{\rho_B}) \\ &\text{strongly in } L^q(\Omega_T), \text{ for all } q < \infty, \end{aligned}$$

while we already know that

$$\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \rightharpoonup \nabla(\sqrt{\rho_A} + \sqrt{\rho_B}) \text{ weakly in } L^2(\Omega_T).$$

Taking the limit $n \rightarrow \infty$ in (71) yields (42). Relation (43) is obtained in a similar way.

From (26) it follows

$$\partial_t \rho_i^n \rightharpoonup \partial_t \rho_i \text{ weakly in } L^{4/3}(0, T; W^{-1,4/3}(\Omega)), \quad i \in \{A, B\},$$

and via the compact Sobolev embedding $W^{1,4/3}(0, T; W^{1,4}(\Omega)') \hookrightarrow C_{weak}([0, T], W^{1,4}(\Omega)')$

$$\rho_i^n \rightarrow \rho_i \text{ in } C_{weak}([0, T], W^{1,4}(\Omega)'), \quad i \in \{A, B\}. \quad (72)$$

Since $\zeta_i^n \rightharpoonup \zeta_i$ weakly in $L^2(\Omega_T)$, $i \in \{A, B\}$, while $\sqrt{\rho_A^n} \rightarrow \sqrt{\rho_A}$, $\sqrt{\rho_B^n} \rightarrow \sqrt{\rho_B}$ strongly in $L^{4-\delta}(\Omega_T)$ for every $\delta \in (0, 3]$, it follows that one can take the limit in (46), (47) and conclude that $\rho \equiv (\rho_A, \rho_B)$ satisfies (37), (38).

Let us now make sure that (39) is satisfied by ρ . Given any constant in time $\phi \in W^{1,4}(\Omega)$ and any $t \in (0, T)$, from (26) (and the fundamental theorem of calculus, which holds since $\rho^n \in W^{1,4/3}(0, T; W^{1,4}(\Omega)')$ and therefore $t \mapsto \langle \rho^n(t), \phi \rangle$ is in $W^{1,4/3}(0, T)$) it follows

$$\begin{aligned} \left| \int_{\Omega} \rho_i^n(t) \phi dx - \int_{\Omega} \rho_i^{in} \phi dx \right| &\leq \int_0^t |\langle \partial_t \rho_i^n(t'), \phi \rangle| dt' \leq \|\partial_t \rho_i^n\|_{L^{4/3}(0,t; W^{1,4}(\Omega)')} \|\phi\|_{L^4(0,t; W^{1,4}(\Omega))} \\ &\leq C t^{1/4} \|\phi\|_{W^{1,4}(\Omega)}. \end{aligned}$$

Since (72) holds, it follows that $\int_{\Omega} \rho_i^n(t) \phi dx \rightarrow \int_{\Omega} \rho_i(t) \phi dx$ as $n \rightarrow \infty$, so

$$\left| \int_{\Omega} \rho_i(t) \phi dx - \int_{\Omega} \rho_i^{in} \phi dx \right| \leq C t^{1/4} \|\phi\|_{W^{1,4}(\Omega)}, \quad \text{for all } \phi \in W^{1,4}(\Omega),$$

which means

$$\|\rho_i(t) - \rho_i^{in}\|_{W^{1,4}(\Omega)'} \leq C t^{1/4}.$$

Being $t \in (0, T)$ arbitrary and $C > 0$ independent of t , we deduce that (39) holds. Therefore, ρ is a weak solution to (30)–(32) according to Definition 3.2. This finishes the proof. $\square$

4 Stationary states

In this section, we study the steady states of (1)-(3), i.e. the constant-in-time solutions to (1)-(3). First, we provide a definition.

Definition 4.1 (Steady state). A *steady state* of (1)-(3) is a weak solution to (1)-(3) in the sense of Def. 3.2 that is constant in time.

In the following we prove that the only allowed steady states for the system are constant.

Proposition 4.2. *Let $\Omega \in \mathbb{R}^2$ open, bounded and connected with Lipschitz boundary. Every steady state of (1)-(3) is constant in Ω .*

Proof. According to the definition of a weak solution, the integrated entropy balance (45) must hold. Being a steady state constant in time, this implies that $\zeta_A = \zeta_B = 0$ a.e. in Ω . From (40) it follows immediately that $\sqrt{\rho_A} + \sqrt{\rho_B} = k_1$ is constant in Ω . In particular, $\rho_A, \rho_B \in L^\infty(\Omega)$. From (41) we deduce

$$\nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})(\sqrt{\rho_A} - \sqrt{\rho_B})] - (\sqrt{\rho_A} - \sqrt{\rho_B}) \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})] = 0$$

for every $\Psi \in C^1 \cap W^{1,\infty}(\mathbb{R}_+)$ such that $|\Psi(s)| \leq C|s - 1|$, for $s > 0$.

Since $\sqrt{\rho_A} + \sqrt{\rho_B}$ is constant it also holds trivially that

$$\nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})(\sqrt{\rho_A} + \sqrt{\rho_B})] - (\sqrt{\rho_A} + \sqrt{\rho_B}) \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})] = 0.$$

Putting the two previous equations together yields

$$\nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})\sqrt{\rho_i}] - \sqrt{\rho_i} \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})] = 0, \quad i \in \{A, B\}. \quad (73)$$

Multiplying the equation above times $\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})\sqrt{\rho_j} \in H^1(\Omega)$ (thanks to (36) and the boundedness of ρ_A, ρ_B) and summing for $i, j = A, B, i \neq j$, lead to

$$\nabla [\Psi(\sqrt{\rho_A \rho_B})^2(1 - \sqrt{\rho_A \rho_B})^2\sqrt{\rho_A \rho_B}] = 2\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})\sqrt{\rho_A \rho_B} \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})],$$

which is equivalent to

$$\begin{aligned} & \Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B}) \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})\sqrt{\rho_A \rho_B}] \\ &= \Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})\sqrt{\rho_A \rho_B} \nabla [\Psi(\sqrt{\rho_A \rho_B})(1 - \sqrt{\rho_A \rho_B})] \quad \text{a.e. in } \Omega, \end{aligned}$$

for every $\Psi \in C^1 \cap W^{1,\infty}(\mathbb{R}_+)$ such that $|\Psi(s)| \leq C|s - 1|$, for $s > 0$. Let us now choose

$$\Psi(s) = \begin{cases} (s - 1)^2, & s \leq 2, \\ 2s - 3, & s > 2. \end{cases}$$

The function $\eta(s) = \Psi(s)(1 - s)$ is strictly decreasing in $\mathbb{R}_+$, so it is invertible. Defining $v = \eta(\sqrt{\rho_A \rho_B})$, $\xi(u) = u\eta^{-1}(u)$ for $u \in (-\infty, 1]$, we deduce

$$v \nabla \xi(v) - \xi(v) \nabla v = \nabla \tilde{\xi}(v) = 0 \quad \text{a.e. in } \Omega,$$

where $\tilde{\xi}(u) \equiv \int_c^u (w \xi'(w) - \xi(w)) dw$, $u \leq 1$. It follows that $\tilde{\xi}(v)$ is constant in Ω . However, $\tilde{\xi}'(u) = u \xi'(u) - \xi(u) = u^2 \frac{d}{du} \left(\frac{\xi(u)}{u} \right) = u^2 \frac{d}{du} \eta^{-1}(u) < 0$ for $u \leq 1$. Therefore, $\tilde{\xi}$ is one-to-one, implying that $v = \eta(\sqrt{\rho_A \rho_B})$ is constant. Being η also a one-to-one function, we deduce that $\sqrt{\rho_A \rho_B} = k_2$ is constant in Ω . We distinguish two cases, according to the value of k_2 .

Case 1: $k_2 \neq 1$. In this situation (73) immediately yields that ρ_A, ρ_B are constant, provided that one chooses Ψ such that $\Psi(k_2) \neq 0$.

Case 2: $k_2 = 1$. In this case we must consider (42), (43), which immediately imply that $\log \rho_A + \rho_B$, $\log \rho_B + \rho_A$ are constant in Ω . Therefore, also their difference is constant, i.e.

$$\rho_A - \rho_B + \log \frac{\rho_B}{\rho_A} = k_3 \quad \text{a.e. in } \Omega.$$

Since $\rho_A \rho_B = 1$ by assumption, it follows

$$F(\rho_A) \equiv \rho_A - \frac{1}{\rho_A} - 2 \log \rho_A = k_3 \quad \text{a.e. in } \Omega.$$

However, $F'(s) = 1 + \frac{1}{s^2} - \frac{2}{s} = \frac{(s-1)^2}{s^2} > 0$ for $s \neq 1$, which means that F is strictly monotone. We conclude that ρ_A is constant, implying that also ρ_B is constant. This finishes the proof of the Proposition. $\square$

The result might mean that the class of solutions we considered is perhaps too small, as segregated states are ruled out. On the other hand, the definition arises naturally from the weak stability argument and only employs the entropy structure of the equations. It is entirely possible that different analytical tools might yield segregated steady states.

4.1 Linear stability analysis

We now consider the stability of the steady states of the system (1)-(3), copied here for reference:

$$\begin{cases} \partial_t \rho_A(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_A(t, x, y) + 2\beta c \rho_A(t, x, y) \nabla \rho_B(t, x, y)), \\ \partial_t \rho_B(t, x, y) = \frac{1}{4} \nabla \cdot (\nabla \rho_B(t, x, y) + 2\beta c \rho_B(t, x, y) \nabla \rho_A(t, x, y)). \end{cases}$$

In order to better understand the system, we perform a linear stability analysis around the uniformly distributed steady state,

$$\begin{cases} \bar{\rho}_A(x) = N_1, & 0 \leq x \leq L, \\ \bar{\rho}_B(x) = N_2, & 0 \leq x < L, \end{cases} \quad (74)$$

where $N_1, N_2 > 0$ are the equilibrium densities and $L \in \mathbb{R}_+$.

To this end, we consider perturbations of the form $\epsilon = \delta_i e^{\alpha t} e^{ikx}$ with $\delta_i \ll 1$ where $i \in \{A, B\}$.

$$\rho_A(x) = \bar{\rho}_A + \delta_A e^{\alpha t + ikx}, \quad 0 \leq x \leq L, \quad (75)$$

$$\rho_B(x) = \bar{\rho}_B + \delta_B e^{\alpha t + ikx}, \quad 0 \leq x < L. \quad (76)$$

Lemma 4.3. *The uniform steady state solution (74) of system (1)-(3) is linearly stable if the following condition holds true:*

$$\beta c \leq \frac{1}{2\sqrt{\bar{\rho}_A \bar{\rho}_B}}. \quad (77)$$

Proof. We plug solutions (75) into system (1)-(3):

$$\begin{cases} \frac{\partial}{\partial t}(\bar{\rho}_A + \delta_A e^{\alpha t + ikx}) = \frac{1}{4} \nabla \cdot (\nabla(\bar{\rho}_A + \delta_A e^{\alpha t + ikx}) + 2\beta c(\bar{\rho}_A + \delta_A e^{\alpha t + ikx}) \nabla(\bar{\rho}_B + \delta_B e^{\alpha t + ikx})), \\ \frac{\partial}{\partial t}(\bar{\rho}_B + \delta_B e^{\alpha t + ikx}) = \frac{1}{4} \nabla \cdot (\nabla(\bar{\rho}_B + \delta_B e^{\alpha t + ikx}) + 2\beta c(\bar{\rho}_B + \delta_B e^{\alpha t + ikx}) \nabla(\bar{\rho}_A + \delta_A e^{\alpha t + ikx})), \end{cases}$$

and obtain

$$\begin{cases} \alpha \delta_A e^{\alpha t + i k x} = \frac{1}{4} \nabla \cdot ((i k \delta_A e^{\alpha t + i k x}) + 2 \beta c (\bar{\rho}_A + \delta_A e^{\alpha t + i k x})(i k \delta_B e^{\alpha t + i k x})), \\ \alpha \delta_B e^{\alpha t + i k x} = \frac{1}{4} \nabla \cdot ((i k \delta_B e^{\alpha t + i k x}) + 2 \beta c (\bar{\rho}_B + \delta_B e^{\alpha t + i k x})(i k \delta_A e^{\alpha t + i k x})). \end{cases}$$

This implies

$$\begin{cases} \alpha \delta_A = \frac{-k^2}{4} \delta_A - \frac{k^2}{2} \delta_B \beta c \bar{\rho}_A + \mathcal{O}(\delta_A \delta_B), \\ \alpha \delta_B = \frac{-k^2}{4} \delta_B - \frac{k^2}{2} \delta_A \beta c \bar{\rho}_B + \mathcal{O}(\delta_A \delta_B). \end{cases}$$

Writing the linear part of this in matrix-vector form $(M - \alpha I) \vec{\delta} = 0$, we have

$$\begin{bmatrix} -(\frac{k^2}{4} + \alpha) & -\frac{k^2}{2} \beta c \bar{\rho}_A \\ -\frac{k^2}{2} \beta c \bar{\rho}_B & -(\frac{k^2}{4} + \alpha) \end{bmatrix} \begin{bmatrix} \delta_A \\ \delta_B \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Hence, it follows from the characteristic polynomial

$$\left(\frac{k^2}{4} + \alpha \right)^2 - \frac{k^4}{4} (\beta c)^2 \bar{\rho}_A \bar{\rho}_B = 0,$$

that

Therefore, the uniform steady state solution will be linearly stable when $\beta c \leq \frac{1}{2\sqrt{\bar{\rho}_A \bar{\rho}_B}}$. $\square$

5 Long-time behavior

In this section we give our result on the long-time behaviour of solutions. We denote

$$\oint_{\Omega} \equiv |\Omega|^{-1} \int_{\Omega}.$$

Theorem 5.1 (Convergence to steady state). *Let $\rho : \Omega \times (0, T) \rightarrow \mathbb{R}_+^2$ be a weak solution to (30)–(32) according to Definition 3.2. Define the constant steady state associated to ρ as*

$$\rho^\infty \equiv (\rho_A^\infty, \rho_B^\infty), \quad \rho_i^\infty = \oint_{\Omega} \rho_i(t) dx = \oint_{\Omega} \rho_i^{in} dx \quad i \in \{A, B\}, \quad t > 0,$$

and assume that $\rho_i^\infty > 0$ for $i \in \{A, B\}$. Define the relative entropy functional as

$$\mathcal{H}(\rho \mid \rho^\infty) = \int_{\Omega} h^*(\rho \mid \rho^\infty) dx,$$

where

$$\begin{aligned} h^*(\rho \mid \rho^\infty) &= h(\rho) - h(\rho^\infty) - h'(\rho^\infty) \cdot (\rho - \rho^\infty), \\ h(\rho) &= \rho_A \log \rho_A + \rho_B \log \rho_B + \rho_A \rho_B. \end{aligned}$$

Then $\mathcal{H}(\rho(t) \mid \rho^\infty) \rightarrow 0$ as $t \rightarrow \infty$.

Furthermore, if $\rho_A^\infty \leq 1$ and $\rho_B^\infty \leq 1$, then $\rho(t) \rightarrow \rho^\infty$ strongly in $L^1(\Omega)$ as $t \rightarrow \infty$.

Remark 5.2. In the physical variables, the constraint on the steady state is $\rho_i^\infty \leq (2\beta c)^{-1}$, $i \in \{A, B\}$.

Proof. The proof is divided into two parts. First we prove that $\lim_{t \rightarrow \infty} \mathcal{H}(\rho(t) \mid \rho^\infty) = 0$, then we show that, if both masses are not larger than 1, then $\rho(t) \rightarrow \rho^\infty$ as $t \rightarrow \infty$ strongly in $L^1(\Omega)$.

Step 1: Show that $\lim_{t \rightarrow \infty} \mathcal{H}(\rho(t) \mid \rho^\infty) = 0$.

Remark 5.3. In the following we will identify the quantities ζ_A, ζ_B with $\nabla\sqrt{\rho_A} + \sqrt{\rho_A\rho_B}\nabla\sqrt{\rho_B}$, $\nabla\sqrt{\rho_B} + \sqrt{\rho_A\rho_B}\nabla\sqrt{\rho_A}$, respectively. Albeit this identification is not known to hold exactly for nondegenerate weak solutions (as the latter expressions are not clearly defined), the present theorem could be proved also by using the properties (40)–(43) and proceeding in a similar way as in the proof of Prop. 4.2. We chose to omit technical details for the sake of a simple exposition.

From (20) it follows that

$$\int_0^\infty \int_\Omega ((1 + \sqrt{\rho_A\rho_B})^2 |\nabla(\sqrt{\rho_A} + \sqrt{\rho_B})|^2 + (1 - \sqrt{\rho_A\rho_B})^2 |\nabla(\sqrt{\rho_A} - \sqrt{\rho_B})|^2) dx dt \leq C.$$

As a consequence there exists an increasing sequence of time instants $t_n \rightarrow \infty$ such that

$$(1 + \sqrt{\rho_A\rho_B})\nabla(\sqrt{\rho_A} + \sqrt{\rho_B})|_{t=t_n} \rightarrow 0, \quad (1 - \sqrt{\rho_A\rho_B})\nabla(\sqrt{\rho_A} - \sqrt{\rho_B})|_{t=t_n} \rightarrow 0 \text{ strongly in } L^2(\Omega), \quad (78)$$

as $n \rightarrow \infty$.

Define $\rho_i^n \equiv \rho_i(t_n)$ for $i \in \{A, B\}$, $n \in \mathbb{N}$. In particular, $\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n})$ is bounded in $L^2(\Omega)$. However by mass conservation $\sqrt{\rho_A^n} + \sqrt{\rho_B^n}$ is bounded in $L^2(\Omega)$, and so $\sqrt{\rho_A^n} + \sqrt{\rho_B^n}$ is bounded in $H^1(\Omega)$. By Sobolev embedding (in 2 space dimensions) $\sqrt{\rho_A^n} + \sqrt{\rho_B^n}$ is bounded in $L^p(\Omega)$ for every $p < \infty$.

From (78) we deduce that $\nabla(\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) \rightarrow 0$ strongly in $L^2(\Omega)$. Poincaré-Wirtinger Lemma yields

$$\sqrt{\rho_A^n} + \sqrt{\rho_B^n} - \int_\Omega (\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) dx \rightarrow 0 \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty. \quad (79)$$

From (78) we also deduce

$$(1 - \sqrt{\rho_A^n\rho_B^n})\nabla\rho_i^n \rightarrow 0 \text{ strongly in } L^2(\Omega), \quad i \in \{A, B\}. \quad (80)$$

The above relation and the uniform L^p bound for ρ_i^n lead to (also, thanks to the definition of weak solution (3.2), $(\sqrt{\rho_A^n\rho_B^n} - 1)^3\sqrt{\rho_i^n} \in H^1(\Omega)$ is an admissible test function, $i \in \{A, B\}$)

$$\frac{1}{5}\nabla((\sqrt{\rho_A^n\rho_B^n} - 1)^5) = (\sqrt{\rho_A^n\rho_B^n} - 1)^4(\sqrt{\rho_A^n}\nabla\sqrt{\rho_B^n} - \sqrt{\rho_B^n}\nabla\sqrt{\rho_A^n}) \rightarrow 0 \text{ strongly in } L^{2-\epsilon}(\Omega), \quad (81)$$

for every $\epsilon > 0$. Again, by Poincaré Lemma one deduce that

$$(\sqrt{\rho_A^n\rho_B^n} - 1)^5 - \int_\Omega (\sqrt{\rho_A^n\rho_B^n} - 1)^5 dx \rightarrow 0 \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty. \quad (82)$$

Since ρ_A^n, ρ_B^n are bounded in $L^p(\Omega)$ for every $p < \infty$, then up to subsequences

$$\int_\Omega (\sqrt{\rho_A^n} + \sqrt{\rho_B^n}) dx \rightarrow c_1, \quad \int_\Omega (\sqrt{\rho_A^n\rho_B^n} - 1)^5 dx \rightarrow \tilde{c}_2.$$

From the above relation and (79), (82) we get

$$\sqrt{\rho_A^n} + \sqrt{\rho_B^n} \rightarrow c_1, \quad (\sqrt{\rho_A^n\rho_B^n} - 1)^5 \rightarrow \tilde{c}_2 \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty.$$

However, being $x \in \mathbb{R} \mapsto (x-1)^5 \in \mathbb{R}$ globally invertible, it follows

$$\sqrt{\rho_A^n} + \sqrt{\rho_B^n} \rightarrow c_1, \quad \sqrt{\rho_A^n \rho_B^n} \rightarrow c_2 := 1 + c_2^{1/5} \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty. \quad (83)$$

As a consequence

$$\rho_A^n + \rho_B^n = (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - 2\sqrt{\rho_A^n \rho_B^n} \rightarrow c_1^2 - 2c_2 \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty.$$

The above relation and mass conservation implies

$$\rho_A^n + \rho_B^n \rightarrow \rho_A^\infty + \rho_B^\infty \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty. \quad (84)$$

Moreover,

$$|\sqrt{\rho_A^n} - \sqrt{\rho_B^n}|^2 = (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})^2 - 4\sqrt{\rho_A^n \rho_B^n} \rightarrow c_1^2 - 4c_2 \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty,$$

and so

$$|\rho_A^n - \rho_B^n| = (\sqrt{\rho_A^n} + \sqrt{\rho_B^n})|\sqrt{\rho_A^n} - \sqrt{\rho_B^n}| \rightarrow c_3 := c_1\sqrt{c_1^2 - 4c_2} \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty. \quad (85)$$

In particular, since $2 \max\{x, y\} = x + y + |x - y|$, $2 \min\{x, y\} = x + y - |x - y|$ for every $x, y \geq 0$, it follows

$$M^n := \max\{\rho_A^n, \rho_B^n\} \rightarrow M, \quad \mu^n := \min\{\rho_A^n, \rho_B^n\} \rightarrow \mu, \text{ strongly in } L^p(\Omega), \text{ for all } p < \infty, \quad (86)$$

and M, μ are constants. Notice that $\sqrt{M\mu} = \lim_{n \rightarrow \infty} \sqrt{\rho_A^n \rho_B^n}$ a.e. in Ω .

From (80), (81) it follows

$$\nabla [(\sqrt{\rho_A^n \rho_B^n} - 1)^5 \rho_i^n] \rightarrow 0 \text{ strongly in } L^{2-\epsilon}(\Omega), \quad i \in \{A, B\},$$

that is

$$(\sqrt{\rho_A^n \rho_B^n} - 1)^5 \rho_i^n \rightarrow \theta_i \text{ strongly in } L^{2-\epsilon}(\Omega), \quad i \in \{A, B\},$$

for some constants θ_A, θ_B . We distinguish two cases:

Case 1: $M\mu \neq 1$.

In this case ρ_i^n is a.e. convergent in Ω to a constant which, due to mass conservation and uniform $L^p(\Omega)$ bounds, must be equal to ρ_i^∞ . It follows

$$\rho_i^n \rightarrow \rho_i^\infty \text{ strongly in } L^q(\Omega), \text{ for all } q < \infty, \quad i \in \{A, B\}. \quad (87)$$

Case 2: $M\mu = 1$.

In this case let us observe that relation (78) can be rewritten as

$$\sqrt{\rho_A^n} \nabla (\log \rho_A^n + \rho_B^n) \rightarrow 0, \quad \sqrt{\rho_B^n} \nabla (\log \rho_B^n + \rho_A^n) \rightarrow 0, \text{ strongly in } L^2(\Omega). \quad (88)$$

Let ψ like in (64). Since

$$\begin{aligned} \nabla(\psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n)) &= \psi'(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n) \nabla \sqrt{\rho_A^n \rho_B^n} + \psi(\sqrt{\rho_A^n \rho_B^n}) \nabla (\log \rho_A^n + \rho_B^n) \\ &= \sqrt{\rho_B^n} \frac{\psi'(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n \rho_B^n}(1 - \sqrt{\rho_A^n \rho_B^n})} \sqrt{\rho_A^n} (\log \rho_A^n + \rho_B^n) (1 - \sqrt{\rho_A^n \rho_B^n}) \nabla \sqrt{\rho_A^n \rho_B^n} \\ &\quad + \sqrt{\rho_B^n} \frac{\psi(\sqrt{\rho_A^n \rho_B^n})}{\sqrt{\rho_A^n \rho_B^n}} \sqrt{\rho_A^n} \nabla (\log \rho_A^n + \rho_B^n) \end{aligned}$$

from (78), (88) it follows that

$$\nabla(\psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n)) \rightarrow 0 \text{ strongly in } L^{2-\epsilon}(\Omega), \text{ for all } \epsilon > 0.$$

Being $\int_{\Omega} \psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n) dx$ bounded, we deduce

$$\psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_A^n + \rho_B^n) \rightarrow c_4 \text{ strongly in } L^{2-\epsilon}(\Omega), \text{ for all } \epsilon > 0.$$

In a similar way,

$$\psi(\sqrt{\rho_A^n \rho_B^n})(\log \rho_B^n + \rho_A^n) \rightarrow c_5 \text{ strongly in } L^{2-\epsilon}(\Omega), \text{ for all } \epsilon > 0.$$

In particular, since $\sqrt{\rho_A^n \rho_B^n} \rightarrow 1$ a.e. in Ω , then

$$\log \frac{\rho_A^n}{\rho_B^n} + \rho_B^n - \rho_A^n \rightarrow c_6 := c_4 - c_5 \quad \text{a.e. in } \Omega.$$

Let $\sigma^n = \frac{\rho_A^n - \rho_B^n}{|\rho_A^n - \rho_B^n|}$ on $\{M^n > \mu^n\}$, $\sigma^n = 0$ on $\{\rho_A^n = \rho_B^n\}$. We just proved

$$\sigma^n \left(\log \frac{M^n}{\mu^n} + \mu^n - M^n \right) \rightarrow c_6 \quad \text{a.e. in } \Omega.$$

However, we know from (86) that $\log \frac{M^n}{\mu^n} + \mu^n - M^n \rightarrow \log \frac{M}{\mu} + \mu - M$ a.e. in Ω , with M, μ constants such that $\sqrt{M\mu} = 1$, so

$$\log \frac{M^n}{\mu^n} + \mu^n - M^n \rightarrow 2 \log M + \frac{1}{M} - M \quad \text{a.e. in } \Omega.$$

Since the function $x \in (0, \infty) \mapsto 2 \log x + \frac{1}{x} - x \in \mathbb{R}$ is one-to-one (strictly decreasing), it vanishes only at $x = 1$. If $M = \mu = 1$ then $\rho_A^n - \rho_B^n \rightarrow 0$ a.e. in Ω and so (87) holds. Let us therefore assume $M > \mu$. In this case

$$\sigma^n \rightarrow c_7 := c_6 \left(2 \log M + \frac{1}{M} - M \right)^{-1} \quad \text{a.e. in } \Omega.$$

It follows that $\rho_A^n - \rho_B^n = \sigma^n (M^n - \mu^n)$ is a.e. convergent in Ω towards a constant, i.e. (87) holds. The uniform $L^p(\Omega)$ bound, valid for every $p < \infty$, implies $\lim_{n \rightarrow \infty} \mathcal{H}(\rho(t_n) \mid \rho^\infty) = 0$. However, since $t \mapsto \mathcal{H}(\rho(t) \mid \rho^\infty)$ is nonincreasing in time, we conclude that $\lim_{t \rightarrow \infty} \mathcal{H}(\rho(t) \mid \rho^\infty) = 0$.

Step 2: Show that $\lim_{t \rightarrow \infty} \rho(t) = \rho^\infty$ strongly in $L^1(\Omega)$.

Assume $\rho_A^\infty, \rho_B^\infty \leq 1$.

We aim to prove that there exists $R > 0, \gamma > 0$ such that if $\rho_A + \rho_B \geq R$ then

$$h^*(\rho \mid \rho^\infty) \geq \gamma(\rho_A + \rho_B), \quad (89)$$

$$h^*(\rho \mid \rho^\infty) > 0 \quad \text{for all } \rho \in \mathbb{R}_+^2 \setminus \{\rho^\infty\}. \quad (90)$$

If (89), (90) hold, then from Step 1 one easily concludes that $\lim_{t \rightarrow \infty} \rho(t) = \rho^\infty$ strongly in $L^1(\Omega)$.

Let us start with (89). We have

$$\begin{aligned} h^*(\rho \mid \rho^\infty) &= \rho_A \log \left(\frac{\rho_A}{\rho_A^\infty} \right) + \rho_A^\infty - \rho_A + \rho_B \log \left(\frac{\rho_B}{\rho_B^\infty} \right) + \rho_B^\infty - \rho_B + (\rho_A - \rho_A^\infty)(\rho_B - \rho_B^\infty) \\ &= (\rho_A + \rho_B) \log(\rho_A + \rho_B) + \rho_A \rho_B + \left[\frac{\rho_A}{\rho_A + \rho_B} \log \left(\frac{\rho_A}{\rho_A + \rho_B} \right) + \frac{\rho_B}{\rho_A + \rho_B} \log \left(\frac{\rho_B}{\rho_A + \rho_B} \right) \right. \\ &\quad \left. - (1 + \rho_B^\infty) \frac{\rho_A}{\rho_A + \rho_B} - (1 + \rho_A^\infty) \frac{\rho_B}{\rho_A + \rho_B} \right] (\rho_A + \rho_B) + \rho_A^\infty + \rho_B^\infty + \rho_A^\infty \rho_B^\infty, \end{aligned}$$

from which (89) follows straightforwardly (as all the terms in the square brackets are bounded).

Let us now show (90). We begin by proving that $h^*(\rho \mid \rho^\infty) > 0$ for $\min\{\rho_A, \rho_B\} = 0$. Since $h^*(0 \mid \rho^\infty) > 0$ trivially, let us consider the case $\rho_A = 0, \rho_B > 0$ (the complementary case $\rho_B = 0, \rho_A > 0$ is treated in an analogous way). We need to study the function

$$f(\rho_B) \equiv h^*(\rho \mid \rho^\infty)|_{\rho_A=0} = \rho_A^\infty + \rho_B \log \left(\frac{\rho_B}{\rho_B^\infty} \right) + \rho_B^\infty - \rho_B - \rho_A^\infty (\rho_B - \rho_B^\infty), \quad \rho_B > 0.$$

Clearly f is a convex function that is positive for $\rho_B = 0$ and tends to infinity when $\rho_B \rightarrow \infty$. Its point of absolute minimum is $\rho_B = \rho_B^\infty \exp(\rho_A^\infty)$, which means

$$f(\rho_B) \geq f(\rho_B^\infty \exp(\rho_A^\infty)) = \rho_A^\infty + \rho_B^\infty (1 + \rho_A^\infty - \exp(\rho_A^\infty)), \quad \rho_B \geq 0.$$

The function $s \in [0, 1] \mapsto \exp(s) - 1 - 2s \in \mathbb{R}$ is strictly convex, vanishes at zero and equals $e - 3 < 0$ at 1. It follows that it is negative in $(0, 1]$, that is, $\exp(s) < 1 + 2s$ for $0 < s \leq 1$. We deduce

$$f(\rho_B) \geq f(\rho_B^\infty \exp(\rho_A^\infty)) > \rho_A^\infty (1 - \rho_B^\infty) \geq 0, \quad \rho_B \geq 0.$$

Therefore, $h^*(\rho \mid \rho^\infty) > 0$ for $\min\{\rho_A, \rho_B\} = 0$.

Let us assume by contradiction that a point $\rho' \in \mathbb{R}_+^2 \setminus \{\rho^\infty\}$ exists such that $h^*(\rho' \mid \rho^\infty) \leq 0$. From (89) we deduce that $h^*(\rho \mid \rho^\infty) > 0$ for $\rho_A + \rho_B \geq R$, so $\rho'_A + \rho'_B < R$. Furthermore, since $h^*(\rho \mid \rho^\infty) > 0$ for $\min\{\rho_A, \rho_B\} = 0$, it follows that $\rho'_A > 0$ as $\rho'_B > 0$. We deduce that the function $\rho \mapsto h^*(\rho \mid \rho^\infty)$ achieves local minimum inside the open region $\{\rho_A > 0, \rho_B > 0, \rho_A + \rho_B < R\}$ in a point $\tilde{\rho} \neq \rho^\infty$; in particular $Dh^*(\tilde{\rho} \mid \rho^\infty) = 0$, i.e. $Dh(\tilde{\rho}) = Dh(\rho^\infty)$. Let us now show that the only solution $\rho \in (0, \infty)^2$ to $Dh(\rho) = Dh(\rho^\infty)$ is $\rho = \rho^\infty$. The equation rewrites as

$$\log \rho_A + \rho_B = \log \rho_A^\infty + \rho_B^\infty, \quad \log \rho_B + \rho_A = \log \rho_B^\infty + \rho_A^\infty,$$

which leads to

$$\rho_A = \rho_A^\infty \exp(\rho_B^\infty - \rho_B), \quad g(\rho_B) = \log \rho_B^\infty + \rho_A^\infty, \quad g(s) \equiv \log(s) + \rho_A^\infty \exp(\rho_B^\infty - s).$$

Since $\rho_A^\infty \leq 1$ and $\rho_B^\infty \leq 1$ it holds

$$g'(s) = \frac{1}{s} - \rho_A^\infty \exp(\rho_B^\infty - s) \geq \frac{1 - s \exp(1 - s)}{s} > 0 \quad \text{for } s > 0, \quad s \neq 1,$$

since $s \mapsto s \exp(1 - s)$ achieves its strict maximum as $s = 1$. This means that g is strictly increasing and therefore the equation $g(\rho_B) = \log \rho_B^\infty + \rho_A^\infty$ has exactly one solution (i.e. $\rho_B = \rho_B^\infty$).

We conclude that (90) holds. This finishes the proof. $\square$

6 Numerical results

In this section we present numerical simulations illustrating Theorem 5.1. In Figure 1 panel (c) we observe the long-term solutions to system (1)-(3) with initial data $\rho_A(0, x) = .5 + e^{-(x-1)^2}$ and $\rho_B(0, x) = .1 + e^{-(x+1)^2}$ which is seen in panel (a). As expected from Theorem 5.1, the solutions converge to the constant equilibrium solutions. Note that ρ_A has an initial mass that is larger than the mass of ρ_B and thus the constant equilibrium solution observed at time $t = 500$ is larger.

Figure 2 illustrates the time evolution of the two energy functionals. We observe that they both seem to stabilize at a minimum by the time $t = 50$.




Figure 1: Numerical solutions to system (1)-(3) with initial densities given by $\rho_A(0, x) = .5 + e^{-(x-1)^2}$ and $\rho_B(0, x) = .1 + e^{-(x+1)^2}$. Panel (b) illustrates transient dynamics and panel (c) the long-time behavior of the solution.




Figure 2: Energy decay with time

Figure 3 illustrates similar results as discussed above. A difference is that the initial densities have a similar mass such that $\rho_A(0, x) = .5 + e^{-(x-1)^2}$ and $\rho_B(0, x) = .5 + e^{-(x+1)^2}$ as observed in panel (a) of Figure 3. Thus, the final states of the densities are the same, as seen in panel (c).




Figure 3: Numerical solutions with initial densities $\rho_A(0, x) = .5 + e^{-(x-1)^2}$ and $\rho_B(0, x) = .5 + e^{-(x+1)^2}$

Finally, we illustrate a result in two-dimensions in Figure 4. These results are consistent with Theorem 5.1. On a final note, the numerical schemes seem to break when initial densities have large

mass.




Figure 4: Numerical solutions in two-dimension with initial densities $\rho_A(0, x) = .1 + e^{-|x-2|^2}$ and $\rho_B(0, x) = .1 + e^{-|x+2|^2}$

Acknowledgements

A. Barbaro was supported by the NSF through grant No. DMS-1319462. N. Rodríguez was partially funded by the NSF DMS-1516778. H. Yoldaş was supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 639638). N. Zamponi acknowledges support from the Alexander von Humboldt foundation. The authors gratefully acknowledge the American Institute of Mathematics (AIM), where this project began.

References

- [1] Abdulaziz Alsenafi and Alethea B. T. Barbaro. A convection–diffusion model for gang territoriality. *Physica A: Statistical Mechanics and its Applications*, 510:765–786, 2018.
- [2] Athmane Bakhta and Virginie Ehrlacher. Global existence of bounded weak solutions to degenerate cross-diffusion equations in moving domain. *arXiv preprint*, arXiv: 1508.06449, 2015.
- [3] Judith Berendsen, Martin Burger, and Jan-Frederik Pietschmann. On a cross-diffusion model for multiple species with nonlocal interaction and size exclusion. *Nonlinear Analysis*, 159:10–39, 2017.
- [4] Martin Burger, Marco Di Francesco, Jan-Frederik Pietschmann, and Bärbel Schlake. Nonlinear cross-diffusion with size exclusion. *SIAM Journal on Mathematical Analysis*, 42(6):2842–2871, 2010.

- [5] Stavros N. Busenberg and Curtis C. Travis. Epidemic models with spatial spread due to population migration. *Journal of Mathematical Biology*, 16(2):181–198, 1983.
- [6] Xiuqing Chen, Esther S. Daus, and Ansgar Jüngel. Global existence analysis of cross-diffusion population systems for multiple species. *Archive Rational Mechanics and Analysis*, 227(2):715–747, 2018.
- [7] Esther S. Daus, Pina Milišić, and Nicola Zamponi. Analysis of a degenerate and singular volume-filling cross-diffusion system modeling biofilm growth. *SIAM Journal on Mathematical Analysis*, 51(4):3569–3605, 2019.
- [8] Laurent Desvillettes, Thomas Lepoutre, and Ayman Moussa. Entropy, duality, and cross diffusion. *SIAM Journal on Mathematical Analysis*, 46(1):820–853, 2014.
- [9] Laurent Desvillettes, Thomas Lepoutre, Ayman Moussa, and Ariane Trescases. On the entropic structure of reaction-cross diffusion systems. *Communications in Partial Differential Equations*, 40(9):1705–1747, September 2015.
- [10] Laurent Desvillettes and Ariane Trescases. New results for triangular reaction cross diffusion system. *Journal of Mathematical Analysis and Applications*, 430(1):32–59, 2015.
- [11] Marco Di Francesco, Antonio Esposito, and Simone Fagioli. Nonlinear degenerate cross-diffusion systems with nonlocal interaction. *Nonlinear Analysis*, 169:94–117, 2018.
- [12] Joshua M. Epstein. *Nonlinear dynamics, mathematical biology, and social science*. Reading, MA: Addison-Wesley, 1997.
- [13] Eduard Feireisl and Antonín Novotný. *Singular limits in thermodynamics of viscous fluids. 2nd edition*. Cham: Birkhäuser, 2nd edition edition, 2017.
- [14] Gonzalo Galiano and Virginia Selgas. On a cross-diffusion segregation problem arising from a model of interacting particles. *Nonlinear Analysis: Real World Applications*, 18:34–49, 2014.
- [15] Anita Gerstenmayer and Ansgar Jüngel. Analysis of a degenerate parabolic cross-diffusion system for ion transport. *Journal of Mathematical Analysis and Applications*, 461(1):523–543, 2018.
- [16] Morton E. Gurtin and A. C. Pipkin. A note on interacting populations that disperse to avoid crowding. *Quarterly of Applied Mathematics*, 42:87–94, 1984.
- [17] Ansgar Jüngel. The boundedness-by-entropy method for cross-diffusion systems. *Nonlinearity*, 28(6):1963–2001, 2015.
- [18] Ansgar Jüngel. Cross-diffusion systems with entropy structure. *Proceedings of Equadiff 2017 Conference*, 2017.
- [19] Masaaki Morisita. Habitat preference and evaluation of environment of an animal: experimental studies on the population density of an ant-lion, *Glenuroides japonicus* M’L. (1). *Physiology and Ecology*, 5(1):1–16, 1952.
- [20] Masaaki Morisita. Measuring of habitat value by the ‘environmental density’ method. in Statistical Ecology. Vol. 1. Spatial Patterns and Statistical Distributions, 1. *Pennsylvania State Univ. Press. University Park*, 1:379–401, 1969.
- [21] Nanako Shigesada, Kohkichi Kawasaki, and Ei Teramoto. Spatial segregation of interacting species. *Journal of Theoretical Biology*, 79(1):83–99, 1979.

- [22] Katarzyna Sznajd-Weron and Józef Sznajd. Opinion evolution in closed community. *International Journal of Modern Physics C*, 11(06):1157–1165, 2000.
- [23] Vladimir K. Vanag and Irving R. Epstein. Cross-diffusion and pattern formation in reaction–diffusion systems. *Physical Chemistry Chemical Physics*, 11(6):897–912, 2009.
- [24] Nicola Zamponi and Ansgar Jüngel. Analysis of degenerate cross-diffusion population models with volume filling. *Annales de l’Institut Henri Poincaré C, Analyse non linéaire*, 34(1):1–29, 2017.