



**HAL**  
open science

# Solvability of a transmission problem in $L_p$ -spaces with generalized diffusion equation

Alexandre Thorel

► **To cite this version:**

Alexandre Thorel. Solvability of a transmission problem in  $L_p$ -spaces with generalized diffusion equation. *Electronic Journal of Differential Equations*, 2024, 2024 (78), pp.1-32. 10.58997/ejde.2024.78 . hal-02975666v2

**HAL Id: hal-02975666**

**<https://hal.science/hal-02975666v2>**

Submitted on 2 Dec 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Solvability of a transmission problem in $L^p$ -spaces with generalized diffusion equation

Alexandre Thorel

Université Le Havre Normandie, Normandie Univ, LMAH UR 3821, 76600 Le Havre, France.

alexandre.thorel@univ-lehavre.fr

## Abstract

We study a transmission problem, in population dynamics, between two juxtaposed habitats. In each habitat, we consider a generalized diffusion equation composed by the Laplace operator and a biharmonic term. We, indeed, allow that the coefficients in front of each term could be negative or null. Using semigroups theory and functional calculus, we give some relation between coefficients to obtain the existence and the uniqueness of the classical solution in  $L^p$ -spaces.

**Key Words and Phrases:** Operational differential equations, functional calculus, analytic semigroups, BIP operators, maximal regularity.

**2020 Mathematics Subject Classification:** 35B65, 35J48, 35R20, 47A60, 47D06.

## 1 Introduction

In this work, using analytic semigroups theory, we study a transmission problem for a coupled system of generalized diffusion equations in  $L^p$ -spaces, with  $p \in (1, +\infty)$ . We denote by generalized diffusion equation, an equation of the following form

$$k\Delta^2 u - l\Delta u = g,$$

with  $k, l \in \mathbb{R}$  and  $g$  given. This equation is obtained using the Landau-Ginzburg free energy functional, we refer to [8] or [29] for more details. This work is a natural continuation of the works done in [19] and [35]. Here, we investigate the influence of the Laplace operator and the biharmonic term in the diffusion. In population dynamics, the Laplace operator model the short range diffusion whereas the biharmonic operator represents the long range diffusion. Thus, generalized diffusion is a linear combination of these two operators.

Usually, in most models  $k, l > 0$ , but in many works like for instance [8], [16], [27], [28] or [29], the authors explain that the biharmonic term plays a stabilizing role if  $k > 0$  and a destabilizing role when  $k < 0$ . This is why, in the present paper, we consider  $k \in \mathbb{R} \setminus \{0\}$  and  $l \in \mathbb{R}$ .

Many works have treated generalized diffusion equations and transmission problems associated to it. For instance, we refer to [3], [6], [8], [10], [20], [21], [22], [23], [28], [29] and [30], for the study of such an equation in population dynamics and to [11], [13], [19] and [35] for transmission problems associated to it. Note that [19] and [35], consider applications in population dynamics whereas [11] and [13], consider applications in plate theory.

We define  $\Omega = \Omega_- \cup \Omega_+$ , the  $n$ -dimensional area,  $n \in \mathbb{N} \setminus \{0, 1\}$ , constituted by the two juxtaposed habitats  $\Omega_- := (a, \gamma) \times \omega$  and  $\Omega_+ := (\gamma, b) \times \omega$  with their interface  $\Gamma = \{\gamma\} \times \omega$ , where  $a, \gamma, b \in \mathbb{R}$  with  $a < \gamma < b$  and  $\omega$  being a bounded domain of  $\mathbb{R}^{n-1}$ .

We investigate the study of the following transmission problem

$$(EQ_{pde}) \begin{cases} k_+ \Delta^2 u_+ - l_+ \Delta u_+ = g_+, & \text{in } \Omega_+ \\ k_- \Delta^2 u_- - l_- \Delta u_- = g_-, & \text{in } \Omega_-, \end{cases}$$

where  $k_{\pm} \in \mathbb{R} \setminus \{0\}$ ,  $l_{\pm} \in \mathbb{R}$ ,  $u_{\pm} \in \Omega_{\pm}$  are population density and  $g_{\pm} \in L^p(\Omega_{\pm})$  are given.

Note that the case  $k_{\pm}, l_{\pm} > 0$  has been already treated in [19] and the case  $l_{\pm} = 0$  with  $k_{\pm} > 0$  has been already treated in [35]. Thus, in the the present article, the most important new results concern the other cases. Indeed, as in the two previous cases, the key point of this article is the inversion of determinant operator but unlike the two previous cases, the writing of this determinant operator by the functional calculus must take into account the sign of the constants  $\frac{l_{+}}{k_{+}}$  and  $\frac{l_{-}}{k_{-}}$ . Therefore it is necessary to detail each case and for each of these cases, we give conditions between the different constants  $k_{\pm}$  and  $l_{\pm}$  which make it possible to invert this determinant operator.

Here, we denote by  $(x, y)$  the spatial variables with  $x \in (a, b)$  and  $y \in \omega$ . The above equations are supplemented by the following boundary and transmission conditions

$$(BC_{pde}) \left\{ \begin{array}{l} (1) \left\{ \begin{array}{l} u_{-}(x, \zeta) = 0, \quad x \in (a, \gamma), \quad \zeta \in \partial\omega \\ u_{+}(x, \zeta) = 0, \quad x \in (\gamma, b), \quad \zeta \in \partial\omega \\ \Delta u_{-}(x, \zeta) = 0, \quad x \in (a, \gamma), \quad \zeta \in \partial\omega \\ \Delta u_{+}(x, \zeta) = 0, \quad x \in (\gamma, b), \quad \zeta \in \partial\omega \end{array} \right. \\ (2) \left\{ \begin{array}{l} u_{-}(a, y) = \varphi_{1}^{-}(y), \quad y \in \omega \\ u_{+}(b, y) = \varphi_{1}^{+}(y), \quad y \in \omega \\ \frac{\partial u_{-}}{\partial x}(a, y) = \varphi_{2}^{-}(y), \quad y \in \omega \\ \frac{\partial u_{+}}{\partial x}(b, y) = \varphi_{2}^{+}(y), \quad y \in \omega, \end{array} \right. \end{array} \right.$$

where  $\varphi_{1}^{\pm}$  and  $\varphi_{2}^{\pm}$  are given in suitable spaces, and

$$(TC_{pde}) \left\{ \begin{array}{l} u_{-} = u_{+} \quad \text{on } \Gamma \\ \frac{\partial u_{-}}{\partial x} = \frac{\partial u_{+}}{\partial x} \quad \text{on } \Gamma \\ k_{-} \Delta u_{-} = k_{+} \Delta u_{+} \quad \text{on } \Gamma \\ \frac{\partial}{\partial x} (k_{-} \Delta u_{-} - l_{-} u_{-}) = \frac{\partial}{\partial x} (k_{+} \Delta u_{+} - l_{+} u_{+}) \quad \text{on } \Gamma. \end{array} \right.$$

In  $(BC_{pde})$ , the boundary conditions on the two first lines of (1) means that the individuals could not lie on the boundaries  $(a, b) \times \partial\omega$ , because, for instance, they die or the edge is impassable. The boundary conditions on the two second lines of (1) mean that there is no dispersal in the normal direction. It follows that the dispersal vanishes on  $(a, b) \times \partial\omega$ . In (2), the population density and the flux are given, for instance on  $\{a\} \times \omega$  and on  $\{b\} \times \omega$ . This signifies that the habitats are not isolated. Then, in  $(TC_{pde})$ , the two first transmission conditions mean the continuity of the density and its flux at the interface, while the two second express, in some sense, the continuity of the dispersal and its flux at the interface  $\Gamma$ .

This article is organized as follows.

In section 2, we give our operational problem. Section 3 is devoted to some recall on BIP operators and real interpolation spaces. In section 4, we give our assumptions and main results. Then, in section 5, we state some preliminary results that will be useful to prove our main result. Finally, section 6, which is composed of three parts, is devoted to the proof of our main result.

## 2 Operational formulation

Since we have

$$\Delta u = \frac{\partial^2 u}{\partial x^2} + \Delta_y u,$$

we set

$$\begin{cases} D(A_0) := \{\psi \in W^{2,p}(\omega) : \psi = 0 \text{ on } \partial\omega\} \\ \forall \psi \in D(A_0), \quad A_0\psi = \Delta_g\psi. \end{cases} \quad (1)$$

Thus, using operator  $A_0$ , it follows that each equation of  $(EQ_{pde})$  writes

$$u_{\pm}^{(4)}(x) + (2A_0 - \frac{l_{\pm}}{k_{\pm}} I)u_{\pm}''(x) + (A_0^2 - \frac{l_{\pm}}{k_{\pm}} A_0)u_{\pm}(x) = f_{\pm}(x),$$

where  $u_{\pm}(x) := u_{\pm}(x, \cdot)$ ,  $f_{\pm}(x) := g_{\pm}(x, \cdot)/k_{\pm}$  and  $f_- \in L^p(a, \gamma; L^p(\omega))$ ,  $f_+ \in L^p(\gamma, b; L^p(\omega))$  with  $p \in (1, +\infty)$ .

Moreover, since the boundary conditions (1) of  $(BC_{pde})$  are included in the domain of  $A_0$ , problem  $(EQ_{pde}) - (BC_{pde}) - (TC_{pde})$  becomes

$$\left\{ \begin{array}{l} u_+^{(4)}(x) + (2A_0 - \frac{l_+}{k_+} I)u_+''(x) + (A_0^2 - \frac{l_+}{k_+} A_0)u_+(x) = f_+(x), \quad \text{for a.e. } x \in (\gamma, b) \\ u_-^{(4)}(x) + (2A_0 - \frac{l_-}{k_-} I)u_-''(x) + (A_0^2 - \frac{l_-}{k_-} A_0)u_-(x) = f_-(x), \quad \text{for a.e. } x \in (a, \gamma) \\ u_-(a) = \varphi_1^-, \quad u_+(b) = \varphi_1^+ \\ u'_-(a) = \varphi_2^-, \quad u'_+(b) = \varphi_2^+ \\ u_-(\gamma) = u_+(\gamma) \\ u'_-(\gamma) = u'_+(\gamma) \\ k_-u_-''(\gamma) + k_-A_0u_-(\gamma) = k_+u_+''(\gamma) + k_+A_0u_+(\gamma) \\ k_-u_-^{(3)}(\gamma) + k_-A_0u'_-(\gamma) - l_-u'_-(\gamma) = k_+u_+^{(3)}(\gamma) + k_+A_0u'_+(\gamma) - l_+u'_+(\gamma), \end{array} \right.$$

Then, we will consider a more general case using  $(A, D(A))$ , instead of  $(A_0, D(A_0))$ , with  $-A$  a BIP operator of angle  $\theta_A \in (0, \pi)$  on a UMD space  $X$ , see below for the definitions of BIP operator and UMD spaces, and  $f \in L^p(a, b; X)$ .

More precisely, setting  $r_{\pm} = \frac{l_{\pm}}{k_{\pm}}$ , we study the following transmission problem (P):

$$(P) \left\{ \begin{array}{l} (EQ) \begin{cases} u_+^{(4)}(x) + (2A - r_+ I)u_+''(x) + (A^2 - r_+ A)u_+(x) = f_+(x), \quad x \in (\gamma, b) \\ u_-^{(4)}(x) + (2A - r_- I)u_-''(x) + (A^2 - r_- A)u_-(x) = f_-(x), \quad x \in (a, \gamma) \end{cases} \\ (BC) \begin{cases} u_-(a) = \varphi_1^-, \quad u_+(b) = \varphi_1^+ \\ u'_-(a) = \varphi_2^-, \quad u'_+(b) = \varphi_2^+ \end{cases} \\ (TC) \begin{cases} u_-(\gamma) = u_+(\gamma) \\ u'_-(\gamma) = u'_+(\gamma) \\ k_-u_-''(\gamma) + k_-Au_-(\gamma) = k_+u_+''(\gamma) + k_+Au_+(\gamma) \\ k_-u_-^{(3)}(\gamma) + k_-Au'_-(\gamma) - l_-u'_-(\gamma) = k_+u_+^{(3)}(\gamma) + k_+Au'_+(\gamma) - l_+u'_+(\gamma). \end{cases} \end{array} \right.$$

The transmission conditions  $(TC)$  will be divided into

$$(TC1) \begin{cases} u_-(\gamma) = u_+(\gamma) \\ u'_-(\gamma) = u'_+(\gamma), \end{cases}$$

and

$$(TC2) \begin{cases} k_-u_-''(\gamma) + k_-Au_-(\gamma) = k_+u_+''(\gamma) + k_+Au_+(\gamma) \\ k_-u_-^{(3)}(\gamma) + k_-Au'_-(\gamma) - l_-u'_-(\gamma) = k_+u_+^{(3)}(\gamma) + k_+Au'_+(\gamma) - l_+u'_+(\gamma). \end{cases}$$

Note that  $(TC2)$  is well defined, see Lemma 3.8 below.

We will search a classical solution of problem (P), that is a solution  $u$  such that

$$\begin{cases} u_+ := u|_{(\gamma,b)} \in W^{4,p}(\gamma, b; X) \cap L^p(\gamma, b; D(A^2)), & u_+'' \in L^p(\gamma, b; D(A)), \\ u_- := u|_{(a,\gamma)} \in W^{4,p}(a, \gamma; X) \cap L^p(a, \gamma; D(A^2)), & u_-'' \in L^p(a, \gamma; D(A)), \end{cases} \quad (2)$$

and which satisfies  $(EQ) - (BC) - (TC)$ .

Note that such a solution is not  $W^{4,p}(a, b; X)$  but uniquely  $W^{4,p}(a, \gamma; X)$  in  $\Omega_-$  and  $W^{4,p}(\gamma, b; X)$  in  $\Omega_+$ .

### 3 Definitions and prerequisites

#### 3.1 The class of Bounded Imaginary Powers of operators

**Definition 3.1.** A Banach space  $X$  is a UMD space if and only if for all  $p \in (1, +\infty)$ , the Hilbert transform is bounded from  $L^p(\mathbb{R}, X)$  into itself (see [4] and [5]).

**Definition 3.2.** A closed linear operator  $T_1$  is called sectorial of angle  $\alpha \in [0, \pi)$  if

$$\begin{aligned} i) & \quad \sigma(T_1) \subset \overline{S_\alpha}, \\ ii) & \quad \forall \alpha' \in (\alpha, \pi), \quad \sup \left\{ \|\lambda(\lambda I - T_1)^{-1}\|_{\mathcal{L}(X)} : \lambda \in \mathbb{C} \setminus \overline{S_{\alpha'}} \right\} < +\infty, \end{aligned}$$

where

$$S_\alpha := \begin{cases} \{z \in \mathbb{C} : z \neq 0 \text{ and } |\arg(z)| < \alpha\} & \text{if } \alpha \in (0, \pi), \\ (0, +\infty) & \text{if } \alpha = 0, \end{cases} \quad (3)$$

see [17], p. 19.

**Remark 3.3.** From [18], p. 342, we know that any injective sectorial operator  $T_1$  admits imaginary powers  $T_1^{is}$ ,  $s \in \mathbb{R}$ , but, in general,  $T_1^{is}$  is not bounded.

**Definition 3.4.** Let  $\theta \in [0, \pi)$ . We denote by  $\text{BIP}(X, \theta)$ , the class of sectorial injective operators  $T_2$  such that

$$\begin{aligned} i) & \quad \overline{D(T_2)} = \overline{R(T_2)} = X, \\ ii) & \quad \forall s \in \mathbb{R}, \quad T_2^{is} \in \mathcal{L}(X), \\ iii) & \quad \exists C \geq 1, \forall s \in \mathbb{R}, \quad \|T_2^{is}\|_{\mathcal{L}(X)} \leq Ce^{|s|\theta}, \end{aligned}$$

see [31], p. 430.

#### 3.2 Interpolation spaces

Here we recall some properties about real interpolation spaces in particular cases.

**Definition 3.5.** Let  $T_3 : D(T_3) \subset X \rightarrow X$  be a linear operator such that

$$(0, +\infty) \subset \rho(T_3) \quad \text{and} \quad \exists C > 0 : \forall t > 0, \quad \|t(T_3 - tI)^{-1}\|_{\mathcal{L}(X)} \leq C. \quad (4)$$

Let  $k \in \mathbb{N} \setminus \{0\}$ ,  $\theta \in (0, 1)$  and  $q \in [1, +\infty]$ . We will use the real interpolation spaces

$$(D(T_3^k), X)_{\theta, q} = (X, D(T_3^k))_{1-\theta, q},$$

defined, for instance, in [24], or in [25].

In particular, for  $k = 1$ , we have the following characterization

$$(D(T_3), X)_{\theta, q} := \left\{ \psi \in X : t \mapsto t^{1-\theta} \|T_3(T_3 - tI)^{-1}\psi\|_X \in L_*^q(0, +\infty) \right\},$$

where  $L_*^q(0, +\infty)$  is given by

$$L_*^q(0, +\infty; \mathbb{C}) := \left\{ f \in L^q(0, +\infty) : \left( \int_0^{+\infty} |f(t)|^q \frac{dt}{t} \right)^{1/q} < +\infty \right\}, \quad \text{for } q \in [1, +\infty),$$

and for  $q = +\infty$ , by

$$L_*^\infty(0, +\infty; \mathbb{C}) := \sup_{t \in (0, +\infty)} |f(t)|,$$

see [9] p. 325, or [15], p. 665, Teorema 3, or section 1.14 of [36], where this space is denoted by  $(X, D(T_3))_{1-\theta, q}$ . Note that we can also characterize the space  $(D(T_3), X)_{\theta, q}$  taking into account the Osservazione, p. 666, in [15].

We set also, for any  $k \in \mathbb{N} \setminus \{0\}$

$$(D(T_3), X)_{k+\theta, q} := \left\{ \psi \in D(T_3^k) : T_3^k \psi \in (D(T_3), X)_{\theta, q} \right\},$$

and

$$(X, D(T_3))_{k+\theta, q} := \left\{ \psi \in D(T_3^k) : T_3^k \psi \in (X, D(T_3))_{\theta, q} \right\},$$

see [26], definition 3.2, p. 64.

**Remark 3.6.** The general situation of the real interpolation space  $(X_0, X_1)_{\theta, q}$  with  $X_0, X_1$  two Banach spaces such that  $X_0 \hookrightarrow X_1$ , is described in [24].

**Remark 3.7.** Note that for  $T_3$  satisfying (4),  $T_3^k$  is closed for any  $k \in \mathbb{N} \setminus \{0\}$  since  $\rho(T_3) \neq \emptyset$ ; consequently, if  $k\theta < 1$ , we have

$$(D(T_3^k), X)_{\theta, q} = (X, D(T_3^k))_{1-\theta, q} = (X, D(T_3))_{k-k\theta, q} = (D(T_3), X)_{(k-1)+k\theta, q} \subset D(T_3^{k-1}).$$

For more details see [25], (2.1.13), p. 43 or [15], p. 676, Teorema 6.

We recall the following lemma.

**Lemma 3.8** ([15]). Let  $T_3$  be a linear operator satisfying (4). Let  $u$  such that

$$u \in W^{n, p}(a_1, b_1; X) \cap L^p(a_1, b_1; D(T_3^k)),$$

where  $a_1, b_1 \in \mathbb{R}$  with  $a_1 < b_1$ ,  $n, k \in \mathbb{N} \setminus \{0\}$  and  $p \in (1, +\infty)$ . Then for any  $j \in \mathbb{N}$  satisfying the Poulsen condition  $0 < \frac{1}{p} + j < n$  and  $s \in \{a_1, b_1\}$ , we have

$$u^{(j)}(s) \in (D(T_3^k), X)_{\frac{j}{n} + \frac{1}{np}, p}.$$

This result is proved in [15], p. 678, Teorema 2'.

## 4 Assumptions and statement of results

### 4.1 Hypotheses

In all the sequel,  $r_+, r_- \in \mathbb{R}$ ,  $k_+ k_- > 0$  and  $A$  denotes a closed linear operator in  $X$ . We assume the following hypotheses:

- (H<sub>1</sub>)  $X$  is a UMD space,
- (H<sub>2</sub>)  $[\min(r_+, r_-, 0), +\infty) \subset \rho(A)$ ,
- (H<sub>3</sub>)  $-A \in \text{BIP}(X, \theta_A)$  for some  $\theta_A \in [0, \pi)$ ,
- (H<sub>4</sub>)  $-A \in \text{Sect}(0)$ .

**Remark 4.1.**

1. Due to (H<sub>2</sub>), if at least one parameter  $r_+$  or  $r_-$  is negative or null, then  $0 \in \rho(A)$ .
2. Operator  $A_0$ , defined by (1), satisfies all the previous hypotheses with  $X = L^q(\omega)$ ,  $q \in (1, +\infty)$  and  $r_{\pm} \in \rho(A_0)$ . From [33], Proposition 3, p. 207,  $X$  satisfies (H<sub>1</sub>) and taking  $A_0 + r_{\pm}I$  in [14], Theorem 9.15, p. 241 and Lemma 9.17, p. 242, we deduce that  $A_0$  satisfies (H<sub>2</sub>). Moreover, (H<sub>3</sub>) is satisfied for every  $\theta_A \in [0, \pi)$ , from [32], Theorem C, p. 166-167. Finally, (H<sub>4</sub>) is satisfied thanks to [17], section 8.3, p. 232.
3. In the scalar case, to solve each equation of (EQ), we need to solve the characteristic equations

$$\chi^4 + (2A - r_{\pm})\chi^2 + (A^2 - r_{\pm}A) = 0,$$

thus, in our operational case, we consider the following operators

$$L_- := -\sqrt{-A + r_-I}, \quad L_+ := -\sqrt{-A + r_+I} \quad \text{and} \quad M := -\sqrt{-A}. \quad (5)$$

Due to (H<sub>2</sub>) and (H<sub>3</sub>),  $-A$ ,  $-A + r_-I$  and  $-A + r_+I$  are sectorial operators, so the existence of  $L_-$ ,  $L_+$  and  $M$  is ensured, see for instance [17], e), p. 25 and [1], Theorem 2.3, p. 69.

4. From [17], Proposition 3.1.9, p. 65, we have  $D(L_-) = D(L_+) = D(M)$ . Thus, for  $n, m \in \mathbb{N}$  and  $m \leq n$

$$D(L_{\pm}^n) = D(M^n) = D(L_{\pm}^m M^{n-m}) = D(M^m L_{\pm}^{n-m}).$$

5. From [31], Theorem 3, p. 437 and [1], Theorem 2.3, p. 69, assumptions (H<sub>2</sub>) and (H<sub>3</sub>) imply that  $-A + r_{\pm}I \in \text{BIP}(X, \theta_A)$  and due to [17], Proposition 3.2.1, e), p. 71, that

$$-L_-, -L_+, -M \in \text{BIP}(X, \theta_A/2).$$

Moreover, from [31], Theorem 4, p. 441, we get

$$-(L_- + M), -(L_+ + M) \in \text{BIP}(X, \theta_A/2 + \varepsilon),$$

for any  $\varepsilon \in (0, \pi/2 - \theta_A/2)$ .

Since we have  $0 < \theta_A/2 < \pi/2$ , then due to [31], Theorem 2, p. 437, we deduce that  $L_-$ ,  $L_+$ ,  $M$ ,  $L_- + M$  and  $L_+ + M$  generate bounded analytic semigroups  $(e^{xL_-})_{x \geq 0}$ ,  $(e^{xL_+})_{x \geq 0}$ ,  $(e^{xM})_{x \geq 0}$ ,  $(e^{x(L_-+M)})_{x \geq 0}$  and  $(e^{x(L_++M)})_{x \geq 0}$ .

6. Using the Dore-Venni sums theorem, see [12], we deduce from (H<sub>1</sub>), (H<sub>2</sub>) and (H<sub>3</sub>) that  $0 \in \rho(M) \cap \rho(L_-) \cap \rho(L_+) \cap \rho(L_+ + M) \cap \rho(L_- + M)$ .

7. From (5), we deduce that

$$\forall \psi \in D(M^2), \quad (L_+^2 - M^2)\psi = r_+ \psi \quad \text{and} \quad (L_-^2 - M^2)\psi = r_- \psi. \quad (6)$$

and also that

$$\forall \psi \in D(M), \quad (L_+ - M)\psi = r_+(L_+ + M)^{-1}\psi \quad \text{and} \quad (L_- - M)\psi = r_-(L_- + M)^{-1}\psi. \quad (7)$$

## 4.2 Main results

To solve our operational problem (P), we introduce two problems:

$$(P_+) \begin{cases} u_+^{(4)}(x) + (2A - r_+ I)u_+''(x) + (A^2 - r_+ A)u_+(x) = f_+(x), & \text{for a.e. } x \in (\gamma, b) \\ u_+(\gamma) = \psi_1, \quad u_+(b) = \varphi_1^+ \\ u_+'(\gamma) = \psi_2, \quad u_+'(b) = \varphi_2^+. \end{cases}$$

and

$$(P_-) \begin{cases} u_-^{(4)}(x) + (2A - r_- I)u_-''(x) + (A^2 - r_- A)u_-(x) = f_-(x), & \text{for a.e. } x \in (a, \gamma) \\ u_-(a) = \varphi_1^-, \quad u_-(\gamma) = \psi_1 \\ u_-'(a) = \varphi_2^-, \quad u_-'(\gamma) = \psi_2, \end{cases}$$

**Remark 4.2.** Recall that  $u$  is a classical solution of (P) if and only if there exist  $\psi_1, \psi_2 \in X$  such that

- (i)  $u_-$  is a classical solution of  $(P_-)$ ,
- (ii)  $u_+$  is a classical solution of  $(P_+)$ ,
- (iii)  $u_-$  and  $u_+$  satisfy (TC2).

Therefore, our aim is to state that there exists a unique couple  $(\psi_1, \psi_2)$  which satisfies (i), (ii) and (iii).

**Theorem 4.3.** Let  $f_- \in L^p(a, \gamma; X)$  and  $f_+ \in L^p(\gamma, b; X)$  with  $p \in (1, +\infty)$ . Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+k_- > 0$ . Thus

1. when  $r_+, r_- \in \mathbb{R} \setminus \{0\}$ ,

- if  $r_+ > 0$  and  $r_- > 0$ ,
- if  $r_+ < 0$  and  $r_- < 0$ , such that

$$(l_+ - l_-)(k_+ - k_-) \geq 0,$$

- if  $r_+ > 0$  and  $r_- < 0$ , such that

$$-6l_-k_+ + l_+k_+ + l_-k_- \geq 0,$$

- if  $r_+ < 0$  and  $r_- > 0$ , such that

$$-6l_+k_- + l_+k_+ + l_-k_- \geq 0,$$

2. when  $r_+ \in \mathbb{R} \setminus \{0\}$  and  $r_- = 0$  with  $\frac{k_-}{k_+} \leq 2$ ,

- if  $r_+ > 0$  such that

$$r_+ \geq \frac{(\sqrt{t+1} + \sqrt{t})^2}{t^2} \frac{k_+^2}{4k_-^2}, \quad \text{for } t \in \left(0, \frac{1}{r_+ \|A^{-1}\|_{\mathcal{L}(X)}}\right) \text{ fixed,}$$

- if  $r_+ < 0$  such that

$$r_+ \leq -\frac{27k_+^2}{64k_-^2},$$



3. when  $r_+ = 0$  and  $r_- \in \mathbb{R} \setminus \{0\}$  with  $\frac{k_+}{k_-} \leq 2$ ,

- if  $r_- > 0$  such that

$$r_- \geq \frac{(\sqrt{t+1} + \sqrt{t})^2}{t^2} \frac{k_-^2}{4k_+^2}, \quad \text{for } t \in \left(0, \frac{1}{r_- \|A^{-1}\|_{\mathcal{L}(X)}}\right) \text{ fixed,}$$

- if  $r_- < 0$  such that

$$r_- \leq -\frac{27k_-^2}{64k_+^2},$$

then, there exists a unique classical solution  $u$ , of the transmission problem (P) if and only if

$$\varphi_1^+, \varphi_1^- \in (D(A), X)_{1+\frac{1}{2p}, p} \quad \text{and} \quad \varphi_2^+, \varphi_2^- \in (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p}. \quad (8)$$

**Remark 4.4.** Since the third case is the symmetric of the second one, replacing  $k_+$  by  $k_-$  and  $l_+$  by  $l_-$ , the proof is exactly the same. Thus, we omit it.

**Remark 4.5.** If  $A = A_0$ , to satisfy the first condition set in the second or the third case of Theorem 4.3, since  $\|A^{-1}\|_{\mathcal{L}(X)} \geq \frac{1}{C_\omega}$ , where  $C_\omega$  is the Poincaré constant, it suffices to take  $\omega$  sufficiently small because the more  $\omega$  is small, the more  $C_\omega$  is large, see for instance [7], Corollary 2.2, p. 95 and Corollary 2.3, p. 96, or [2], Remark 3, p. 15.

As a consequence of the previous Theorem, we state the following corollary.

**Corollary 4.6.** Let  $n \geq 2$ ,  $f_+ \in L^p(\Omega_+)$  and  $f_- \in L^p(\Omega_-)$  with  $p \in (1, +\infty)$  and  $p > n$ . Assume that  $\omega$  is a bounded open set of  $\mathbb{R}^{n-1}$  with  $C^2$ -boundary. Let  $k_+, k_-, l_+ > 0$  and  $l_- < 0$  with  $k_+ = k_-$ . Then, there exists a unique solution  $u$  of  $(P_{pde})$ , such that we have  $u_- \in W^{4,p}(\Omega_-)$  and  $u_+ \in W^{4,p}(\Omega_+)$ , if and only if

$$\begin{cases} \varphi_1^\pm, \varphi_2^\pm \in W^{2,p}(\omega) \cap W_0^{1,p}(\omega) \\ \Delta \varphi_1^\pm \in W^{2-\frac{1}{p},p}(\omega) \cap W_0^{1,p}(\omega) \\ \Delta \varphi_2^\pm \in W^{1-\frac{1}{p},p}(\omega) \cap W_0^{1,p}(\omega). \end{cases}$$

The proof is quite similar to the one stated in [19], Corollary 1, p. 2941, or in [20], Corollary 2.7, p. 357. Thus we omit it.

## 5 Preliminary results

In all the sequel, we set

$$c = \gamma - a > 0 \quad \text{and} \quad d = b - \gamma > 0.$$

From Remark 4.2, to solve problem (P) we must first study problems  $(P_+)$  and  $(P_-)$ . To this end, we need the following invertibility result obtained in [20] and [34].

**Lemma 5.1** ([20] and [34]). Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$  and  $(H_4)$  hold. Then operators  $U_{\pm}, V_{\pm} \in \mathcal{L}(X)$  defined by

$$\left\{ \begin{array}{l} U_+ := \begin{cases} I - e^{d(L_++M)} - \frac{1}{r_+} (L_+ + M)^2 (e^{dM} - e^{dL_+}), & \text{if } r_+ \in \mathbb{R} \setminus \{0\} \\ I - e^{2dM} + 2dM e^{dM}, & \text{if } r_+ = 0 \end{cases} \\ V_+ := \begin{cases} I - e^{d(L_++M)} + \frac{1}{r_+} (L_+ + M)^2 (e^{dM} - e^{dL_+}), & \text{if } r_+ \in \mathbb{R} \setminus \{0\} \\ I - e^{2dM} - 2dM e^{dM}, & \text{if } r_+ = 0 \end{cases} \\ U_- := \begin{cases} I - e^{c(L_-+M)} - \frac{1}{r_-} (L_- + M)^2 (e^{cM} - e^{cL_-}), & \text{if } r_- \in \mathbb{R} \setminus \{0\} \\ I - e^{2cM} + 2cM e^{cM}, & \text{if } r_- = 0 \end{cases} \\ V_- := \begin{cases} I - e^{c(L_-+M)} + \frac{1}{r_-} (L_- + M)^2 (e^{cM} - e^{cL_-}), & \text{if } r_- \in \mathbb{R} \setminus \{0\} \\ I - e^{2cM} - 2cM e^{cM}, & \text{if } r_- = 0, \end{cases} \end{array} \right. \quad (9)$$

are invertible with bounded inverses.

From Remark 4.1, statement 4,  $U_{\pm}$  and  $V_{\pm}$  are well defined. For a detailed proof, see [20], Proposition 5.4 with  $k = r_{\pm}$  and [34], Proposition 4.5, p. 645.

## 5.1 Transmission system

### 5.1.1 First case

Assume that  $r_{\pm} \in \mathbb{R} \setminus \{0\}$ . We set

$$\left\{ \begin{array}{l} P_1^+ = k_+(L_+ + M) \left( U_+^{-1}(I + e^{dM})(I - e^{dL_+}) + V_+^{-1}(I - e^{dM})(I + e^{dL_+}) \right) \\ P_2^+ = k_+(L_+ + M) \left( U_+^{-1}(I - e^{dM})(I - e^{dL_+}) + V_+^{-1}(I + e^{dM})(I + e^{dL_+}) \right) \\ P_3^+ = k_+(L_+ + M)L_+ \left( U_+^{-1}(I + e^{dM})(I + e^{dL_+}) + V_+^{-1}(I - e^{dM})(I - e^{dL_+}) \right), \end{array} \right. \quad (10)$$

and similarly

$$\left\{ \begin{array}{l} P_1^- = k_-(L_- + M) \left( U_-^{-1}(I + e^{cM})(I - e^{cL_-}) + V_-^{-1}(I - e^{cM})(I + e^{cL_-}) \right) \\ P_2^- = k_-(L_- + M) \left( U_-^{-1}(I - e^{cM})(I - e^{cL_-}) + V_-^{-1}(I + e^{cM})(I + e^{cL_-}) \right) \\ P_3^- = k_-(L_- + M)L_- \left( U_-^{-1}(I + e^{cM})(I + e^{cL_-}) + V_-^{-1}(I - e^{cM})(I - e^{cL_-}) \right). \end{array} \right. \quad (11)$$

Moreover, we note

$$\begin{aligned} S_1 &= k_+(L_+ + M) \left( U_+^{-1}(I - e^{dL_+})\tilde{\varphi}_2^+ + V_+^{-1}(I + e^{dL_+})\tilde{\varphi}_4^+ \right) \\ &\quad - k_-(L_- + M) \left( U_-^{-1}(I - e^{cL_-})\tilde{\varphi}_2^- + V_-^{-1}(I + e^{cL_-})\tilde{\varphi}_4^- \right), \end{aligned} \quad (12)$$

and

$$\begin{aligned} S_2 &= -k_+(L_+ + M) \left( U_+^{-1}(I + e^{dM})\tilde{\varphi}_1^+ + V_+^{-1}(I - e^{dM})\tilde{\varphi}_3^+ \right) \\ &\quad - k_-(L_- + M) \left( U_-^{-1}(I + e^{cM})\tilde{\varphi}_1^- + V_-^{-1}(I - e^{cM})\tilde{\varphi}_3^- \right) - 2M^{-1}R_1, \end{aligned} \quad (13)$$

with

$$R_1 = -k_+F_+'''(\gamma) + k_+M^2F_+'(\gamma) + l_+F_+'(\gamma) + k_-F_-'''(\gamma) - k_-M^2F_-'(\gamma) - l_-F_-'(\gamma), \quad (14)$$

where  $F_+$  is the unique classical solution of problem

$$\begin{cases} u_+^{(4)}(x) + (2A - r_+ I)u_+''(x) + (A^2 - r_+ A)u_+(x) = f_+(x), & \text{for a.e. } x \in (\gamma, b) \\ u_+(\gamma) = u_+(b) = u_+''(\gamma) = u_+''(b) = 0, \end{cases} \quad (15)$$

and  $F_-$  is the unique classical solution of problem

$$\begin{cases} u_-^{(4)}(x) + (2A - r_- I)u_-''(x) + (A^2 - r_- A)u_-(x) = f_-(x), & \text{for a.e. } x \in (a, \gamma) \\ u_-(a) = u_-(\gamma) = u_-''(a) = u_-''(\gamma) = 0. \end{cases} \quad (16)$$

For an explicit representation formula of the solution of both previous problems, we refer to [20], Theorem 2.2, p. 355-356.

**Remark 5.2.** Since  $F_\pm$  is a classical solution of (15), respectively (16), from Lemma 3.8, it follows that, for  $j = 0, 1, 2, 3$  and  $s = a, \gamma$  or  $b$

$$F_\pm^{(j)}(s) \in (D(M), X)_{3-j+\frac{1}{p}, p}.$$

Now, with our notations, we recall a useful result of [19], Theorem 4.6, p. 2945. This result is proved for  $r_\pm > 0$  but clearly remains true for  $r_\pm < 0$ .

**Theorem 5.3** ([19]). Let  $f_- \in L^p(a, \gamma; X)$  and  $f_+ \in L^p(\gamma, b; X)$ , with  $p \in (1, +\infty)$ . Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$  and  $(H_4)$  hold. Then, the transmission problem  $(P)$  has a unique classical solution if and only if the data  $\varphi_1^+, \varphi_1^-, \varphi_2^+, \varphi_2^-$  satisfy (8) and system

$$\begin{cases} (P_1^- - P_1^+) M\psi_1 + (P_2^+ + P_2^-) \psi_2 = S_1 \\ (P_3^+ + P_3^-) \psi_1 + (P_1^- - P_1^+) \psi_2 = S_2, \end{cases} \quad (17)$$

has a unique solution  $(\psi_1, \psi_2)$  such that

$$(\psi_1, \psi_2) \in (D(A), X)_{1+\frac{1}{2p}, p} \times (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p}. \quad (18)$$

### 5.1.2 Second case

Now, assume that  $r_- = 0$ ; then  $l_- = 0$  and the transmission conditions  $(TC2)$  becomes

$$(TC2') \begin{cases} k_- u_-''(\gamma) + k_- A u_-(\gamma) = k_+ u_+''(\gamma) + k_+ A u_+(\gamma) \\ k_- u_-^{(3)}(\gamma) + k_- A u_-'(\gamma) = k_+ u_+^{(3)}(\gamma) + k_+ A u_+'(\gamma) - l_+ u_+'(\gamma). \end{cases} \quad (19)$$

Our aim here is to establish a similar result to the previous one. To this end, we set

$$\begin{cases} Q_1^- = k_- (U_-^{-1} + V_-^{-1}) (I - e^{2cM}) \\ Q_2^- = k_- (U_-^{-1} (I - e^{cM})^2 + V_-^{-1} (I + e^{cM})^2) \\ Q_3^- = k_- (U_-^{-1} (I + e^{cM})^2 + V_-^{-1} (I - e^{cM})^2), \end{cases} \quad (20)$$

Moreover, we note

$$\begin{aligned} S_3 &= 2k_- M (U_-^{-1} (I - e^{cM}) \tilde{\varphi}_2^- + V_-^{-1} (I + e^{cM}) \tilde{\varphi}_4^-) \\ &\quad - k_+ (L_+ + M) (U_+^{-1} (I - e^{dL_+}) \tilde{\varphi}_2^+ + V_+^{-1} (I + e^{dL_+}) \tilde{\varphi}_4^+), \end{aligned} \quad (21)$$

and

$$\begin{aligned} S_4 = & -2k_-M \left( U_-^{-1} \left( I + e^{cM} \right) \tilde{\varphi}_2^- + V_-^{-1} \left( I - e^{cM} \right) \tilde{\varphi}_4^- \right) \\ & -k_+(L_+ + M) \left( U_+^{-1} \left( I + e^{dM} \right) \tilde{\varphi}_1^+ + V_+^{-1} \left( I - e^{dM} \right) \tilde{\varphi}_3^+ \right) + 2M^{-1}R_2, \end{aligned} \quad (22)$$

with

$$R_2 = -k_- \tilde{F}_-'''(\gamma) + k_- M^2 \tilde{F}_-''(\gamma) + k_+ F_+'''(\gamma) - k_+ M^2 F_+''(\gamma) - l_+ F_+'(\gamma), \quad (23)$$

where  $\tilde{F}_-$  is the classical solution of problem

$$\begin{cases} u_-^{(4)}(x) + 2A u_-''(x) + A^2 u_-(x) = f_-(x), & \text{a.e. } x \in (a, \gamma) \\ u_-(a) = u_-(\gamma) = u_-''(a) = u_-''(\gamma) = 0. \end{cases} \quad (24)$$

**Remark 5.4.** Since  $\tilde{F}_-$  is a classical solution of (16), as in Remark 5.2, from Lemma 3.8, it follows that, for  $j = 0, 1, 2, 3$  and  $s = a, \gamma$  or  $b$

$$\tilde{F}_-^{(j)}(s) \in (D(M), X)_{3-j+\frac{1}{p}, p}.$$

**Theorem 5.5.** Let  $f_- \in L^p(a, \gamma; X)$  and  $f_+ \in L^p(\gamma, b; X)$ , with  $p \in (1, +\infty)$ . Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$  and  $(H_4)$  hold. Then, problem (P) has a unique classical solution if and only if the data  $\varphi_1^+$ ,  $\varphi_1^-$ ,  $\varphi_2^+$ ,  $\varphi_2^-$  satisfy (8) and system

$$\begin{cases} \left( P_1^+ - 2MQ_1^- \right) M\psi_1 - \left( P_2^+ + 2MQ_2^- \right) \psi_2 = S_3 \\ \left( P_3^+ + 2MQ_3^- \right) \psi_1 + \left( 2MQ_1^- - P_1^+ \right) \psi_2 = S_4, \end{cases} \quad (25)$$

has a unique solution  $(\psi_1, \psi_2)$  satisfying (18).

*Proof.* We follow the same steps than the proof of Theorem 4.6, p. 2945 in [19], we only point out the key points. From [20], Theorem 2.5, statement 2, there exists a unique classical solution  $u_+$  of  $(P_+)$  if and only if

$$\varphi_1^+, \psi_1 \in (D(A), X)_{1+\frac{1}{2p}, p} \quad \text{and} \quad \varphi_2^+, \psi_2 \in (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p}. \quad (26)$$

Recall that, from Remark 3.7, we have

$$(D(A), X)_{1+\frac{1}{2p}, p} = (D(M), X)_{3+\frac{1}{p}, p} \quad \text{and} \quad (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p} = (D(M), X)_{2+\frac{1}{p}, p}. \quad (27)$$

This solution is explicitly given in [19], Proposition 2, p. 2943-2944, from which we deduce that

$$k_+ \left( u_+''(\gamma) - M^2 u_+(\gamma) \right) = l_+ \left( I - e^{dL_+} \right) \alpha_2^+ + l_+ \left( I + e^{dL_+} \right) \alpha_4^+,$$

and

$$\begin{aligned} k_+ \left( u_+^{(3)}(\gamma) - M^2 u_+'(\gamma) \right) - l_+ u_+'(\gamma) &= -l_+ M \left( I + e^{dM} \right) \alpha_1^+ - l_+ M \left( I - e^{dM} \right) \alpha_3^+ \\ &\quad + k_+ F_+'''(\gamma) - k_+ M^2 F_+''(\gamma) - l_+ F_+'(\gamma), \end{aligned}$$

where

$$\begin{cases} \alpha_1^+ = \frac{1}{2r_+} (L_+ + M) U_+^{-1} \left[ L_+ (I + e^{dL_+}) \psi_1 - (I - e^{dL_+}) \psi_2 + \tilde{\varphi}_1^+ \right] \\ \alpha_2^+ = -\frac{1}{2r_+} (L_+ + M) U_+^{-1} \left[ M (I + e^{dM}) \psi_1 - (I - e^{dM}) \psi_2 + \tilde{\varphi}_2^+ \right] \\ \alpha_3^+ = \frac{1}{2r_+} (L_+ + M) V_+^{-1} \left[ L_+ (I - e^{dL_+}) \psi_1 - (I + e^{dL_+}) \psi_2 + \tilde{\varphi}_3^+ \right] \\ \alpha_4^+ = -\frac{1}{2r_+} (L_+ + M) V_+^{-1} \left[ M (I - e^{dM}) \psi_1 - (I + e^{dM}) \psi_2 + \tilde{\varphi}_4^+ \right], \end{cases} \quad (28)$$

with

$$\begin{cases} \tilde{\varphi}_1^+ &= -L_+ (I + e^{dL_+}) \varphi_1^+ + (I - e^{dL_+}) (F'_+(b) + F'_+(\gamma) - \varphi_2^+) \\ \tilde{\varphi}_2^+ &= -M (I + e^{dM}) \varphi_1^+ + (I - e^{dM}) (F'_+(b) + F'_+(\gamma) - \varphi_2^+) \\ \tilde{\varphi}_3^+ &= L_+ (I - e^{dL_+}) \varphi_1^+ - (I + e^{dL_+}) (F'_+(b) - F'_+(\gamma) - \varphi_2^+) \\ \tilde{\varphi}_4^+ &= M (I - e^{dM}) \varphi_1^+ - (I + e^{dM}) (F'_+(b) - F'_+(\gamma) - \varphi_2^+), \end{cases} \quad (29)$$

and  $F_+$  is the unique classical solution of problem (15).

In the same way, from [34], Theorem 2.8, statement 2, p. 637, there exists a unique classical solution  $u_-$  of problem  $(P_-)$  if and only if

$$\varphi_1^-, \psi_1 \in (D(A), X)_{1+\frac{1}{2p}, p} \quad \text{and} \quad \varphi_2^-, \psi_2 \in (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p}. \quad (30)$$

Note that from (27), we have

$$\varphi_1^-, \psi_1 \in (D(M), X)_{3+\frac{1}{p}, p} \quad \text{and} \quad \varphi_2^-, \psi_2 \in (D(M), X)_{2+\frac{1}{p}, p}.$$

Moreover, this solution, given in [34], Proposition 4.1, p. 640, is explicitly written in [35], Proposition 4.2, from which it follows that

$$k_- (u_-''(\gamma) - M^2 u_-(\gamma)) = -k_- (2M (I - e^{cM}) \alpha_2^- - 2M (I + e^{cM}) \alpha_4^-),$$

and

$$\begin{aligned} k_- (u_-^{(3)}(\gamma) - M^2 u_-'(\gamma)) &= k_- (2M^2 (I + e^{cM}) \alpha_2^- - 2M^2 (I - e^{cM}) \alpha_4^-) \\ &\quad + k_- F_-^{(3)}(\gamma) - k_- M^2 F_-'(\gamma), \end{aligned}$$

where

$$\begin{cases} \alpha_1^- &:= -\frac{1}{2} U_-^{-1} \left[ (I + (I + cM) e^{cM}) \psi_1 - c e^{cM} \psi_2 + \tilde{\varphi}_1^- \right] \\ \alpha_2^- &:= \frac{1}{2} U_-^{-1} \left[ (I + e^{cM}) M \psi_1 + (I - e^{cM}) \psi_2 + \tilde{\varphi}_2^- \right] \\ \alpha_3^- &:= \frac{1}{2} V_-^{-1} \left[ (I - (I + cM) e^{cM}) \psi_1 + c e^{cM} \psi_2 + \tilde{\varphi}_3^- \right] \\ \alpha_4^- &:= -\frac{1}{2} V_-^{-1} \left[ (I - e^{cM}) M \psi_1 + (I + e^{cM}) \psi_2 + \tilde{\varphi}_4^- \right], \end{cases} \quad (31)$$

with

$$\begin{cases} \tilde{\varphi}_1^- &:= -(I + e^{cM}) \varphi_1^- - c e^{cM} (M \varphi_1^- + \varphi_2^- - \tilde{F}'_-(a) - \tilde{F}'_-(\gamma)) \\ \tilde{\varphi}_2^- &:= -M (I + e^{cM}) \varphi_1^- + (I - e^{cM}) (\varphi_2^- - \tilde{F}'_-(a) - \tilde{F}'_-(\gamma)) \\ \tilde{\varphi}_3^- &:= (I - e^{cM}) \varphi_1^- - c e^{cM} (M \varphi_1^- + \varphi_2^- - \tilde{F}'_-(a) + \tilde{F}'_-(\gamma)) \\ \tilde{\varphi}_4^- &:= M (I - e^{cM}) \varphi_1^- - (I + e^{cM}) (\varphi_2^- - \tilde{F}'_-(a) + \tilde{F}'_-(\gamma)). \end{cases} \quad (32)$$

Note that due to (26), (27), (28) and (29), respectively to (27), (30), (31) and (32), we deduce that

$$\alpha_i^\pm \in D(M), \quad \text{for } i = 1, 2, 3, 4 \quad \text{and} \quad \alpha_2^-, \alpha_4^- \in D(M^2).$$

Thus, system  $(TC2')$ , given by (19), writes

$$\begin{cases} -2k_- M \left( (I - e^{cM}) \alpha_2^- - (I + e^{cM}) \alpha_4^- \right) = l_+ \left( (I - e^{dL_+}) \alpha_2^+ + (I + e^{dL_+}) \alpha_4^+ \right) \\ 2k_- M^2 \left( (I + e^{cM}) \alpha_2^- - (I - e^{cM}) \alpha_4^- \right) = -l_+ M \left( (I + e^{dM}) \alpha_1^+ + (I - e^{dM}) \alpha_3^+ \right) + R_2, \end{cases}$$

where  $R_2$  is given by (23). Thus, it follows that the previous system gives

$$\begin{cases} -2k_- U_-^{-1} M (I - e^{cM}) \left[ (I + e^{cM}) M \psi_1 + (I - e^{cM}) \psi_2 + \tilde{\varphi}_2^- \right] \\ -2k_- V_-^{-1} M (I + e^{cM}) \left[ (I - e^{cM}) M \psi_1 + (I + e^{cM}) \psi_2 + \tilde{\varphi}_4^- \right] \\ +k_+(L_+ + M) U_+^{-1} (I - e^{dL_+}) \left[ M(I + e^{dM}) \psi_1 - (I - e^{dM}) \psi_2 + \tilde{\varphi}_2^+ \right] \\ +k_+(L_+ + M) V_+^{-1} (I + e^{dL_+}) \left[ M(I - e^{dM}) \psi_1 - (I + e^{dM}) \psi_2 + \tilde{\varphi}_4^+ \right] = 0 \\ \\ 2k_- M U_-^{-1} (I + e^{cM}) \left[ (I + e^{cM}) M \psi_1 + (I - e^{cM}) \psi_2 + \tilde{\varphi}_2^- \right] \\ +2k_- M V_-^{-1} (I - e^{cM}) \left[ (I - e^{cM}) M \psi_1 + (I + e^{cM}) \psi_2 + \tilde{\varphi}_4^- \right] \\ +k_+(L_+ + M) U_+^{-1} (I + e^{dM}) \left[ L_+(I + e^{dL_+}) \psi_1 - (I - e^{dL_+}) \psi_2 + \tilde{\varphi}_1^+ \right] \\ +k_+(L_+ + M) V_+^{-1} (I - e^{dM}) \left[ L_+(I - e^{dL_+}) \psi_1 - (I + e^{dL_+}) \psi_2 + \tilde{\varphi}_3^+ \right] = 2M^{-1} R_2, \end{cases}$$

Finally, using (10), (20), (21), (22) and (23), we obtain that the previous system writes as system (25).

Conversely, if we assume that (8) holds and system (25) has a unique solution  $(\psi_1, \psi_2)$  satisfying (18), then considering  $u_{\pm}$  the unique classical solution of  $(P_{\pm})$ , we obtain that  $u$  is the unique classical solution of (P).  $\square$

## 5.2 Functional calculus

In this section, by using functional calculus, we rewrite the operators defined in (9), (10), (11) and (20), to inverse the determinant operator of system (17) and system (25).

To this end, we recall some classical notations. For  $\theta \in (0, \pi)$ , we denote by  $H(S_{\theta})$  the space of holomorphic functions on  $S_{\theta}$  (defined by (3)) with values in  $\mathbb{C}$ . Moreover, we consider the following subspace of  $H(S_{\theta})$ :

$$\mathcal{E}_{\infty}(S_{\theta}) := \{f \in H(S_{\theta}) : f = O(|z|^{-s}) \text{ } (|z| \rightarrow +\infty) \text{ for some } s > 0\}.$$

In other words,  $\mathcal{E}_{\infty}(S_{\theta})$  is the space of polynomially decreasing holomorphic functions at  $+\infty$ . Let  $T$  be an invertible sectorial operator of angle  $\theta_T \in (0, \pi)$ . If  $f \in \mathcal{E}_{\infty}(S_{\theta})$ , with  $\theta \in (\theta_T, \pi)$ , then we can define, by functional calculus,  $f(T) \in \mathcal{L}(X)$ , see [17], p. 45.

Then, we recall a useful result from [20], Lemma 5.3, p. 370.

**Lemma 5.6** ([20]). Let  $P$  be an invertible sectorial operator in  $X$  with angle  $\theta$ , for all  $\theta \in (0, \pi)$ . Let  $G \in H(S_{\theta})$ , for some  $\theta \in (0, \pi)$ , such that

- (i)  $1 - G \in \mathcal{E}_{\infty}(S_{\theta})$ ,
- (ii)  $G(x) \neq 0$  for any  $x \in \mathbb{R}_+ \setminus \{0\}$ .

Then,  $G(P) \in \mathcal{L}(X)$ , is invertible with bounded inverse.

Let  $r \in \mathbb{R}$ ,  $r_m = \max(-r, 0)$ ,  $\delta > 0$  and  $z \in \mathbb{C} \setminus (-\infty, r_m]$ . We set

$$\begin{cases} u_{\delta,r}(z) = \begin{cases} 1 - e^{-\delta(\sqrt{z+r}+\sqrt{z})} - \frac{1}{r}(\sqrt{z+r} + \sqrt{z})^2 (e^{-\delta\sqrt{z}} - e^{-\delta\sqrt{z+r}}), & \text{if } r \in \mathbb{R} \setminus \{0\} \\ 1 - e^{-2\delta\sqrt{z}} - 2\delta\sqrt{z}e^{-\delta\sqrt{z}}, & \text{if } r = 0 \end{cases} \\ v_{\delta,r}(z) = \begin{cases} 1 - e^{-\delta(\sqrt{z+r}+\sqrt{z})} + \frac{1}{r}(\sqrt{z+r} + \sqrt{z})^2 (e^{-\delta\sqrt{z}} - e^{-\delta\sqrt{z+r}}), & \text{if } r \in \mathbb{R} \setminus \{0\} \\ 1 - e^{-2\delta\sqrt{z}} + 2\delta\sqrt{z}e^{-\delta\sqrt{z}}, & \text{if } r = 0 \end{cases} \end{cases}$$

and when  $u_{\delta,r}(z) \neq 0$ ,  $v_{\delta,r}(z) \neq 0$ , we note

$$\begin{aligned} f_{\delta,r,1}(z) &= \begin{cases} \begin{aligned} & (\sqrt{z+r} + \sqrt{z}) \sqrt{z+r} u_{\delta,r}^{-1}(z) (1 + e^{-\delta\sqrt{z}}) (1 + e^{-\delta\sqrt{z+r}}) \\ & + (\sqrt{z+r} + \sqrt{z}) \sqrt{z+r} v_{\delta,r}^{-1}(z) (1 - e^{-\delta\sqrt{z}}) (1 - e^{-\delta\sqrt{z+r}}), \end{aligned} & \text{if } r \in \mathbb{R} \setminus \{0\} \\ (u_{\delta,0}^{-1}(z) + v_{\delta,0}^{-1}(z)) (1 - e^{-2\delta\sqrt{z}}), & \text{if } r = 0, \end{cases} \\ f_{\delta,r,2}(z) &= \begin{cases} \begin{aligned} & -(\sqrt{z+r} + \sqrt{z}) u_{\delta,r}^{-1}(z) (1 + e^{-\delta\sqrt{z}}) (1 - e^{-\delta\sqrt{z+r}}) \\ & -(\sqrt{z+r} + \sqrt{z}) v_{\delta,r}^{-1}(z) (1 - e^{-\delta\sqrt{z}}) (1 + e^{-\delta\sqrt{z+r}}), \end{aligned} & \text{if } r \in \mathbb{R} \setminus \{0\} \\ u_{\delta,0}^{-1}(z) (1 - e^{-\delta\sqrt{z}})^2 + v_{\delta,0}^{-1}(z) (1 + e^{-\delta\sqrt{z}})^2, & \text{if } r = 0, \end{cases} \end{aligned}$$

and

$$f_{\delta,r,3}(z) = \begin{cases} \begin{aligned} & -(\sqrt{z+r} + \sqrt{z}) u_{\delta,r}^{-1}(z) (1 - e^{-\delta\sqrt{z}}) (1 - e^{-\delta\sqrt{z+r}}) \\ & -(\sqrt{z+r} + \sqrt{z}) v_{\delta,r}^{-1}(z) (1 + e^{-\delta\sqrt{z}}) (1 + e^{-\delta\sqrt{z+r}}), \end{aligned} & \text{if } r \in \mathbb{R} \setminus \{0\} \\ u_{\delta,0}^{-1}(z) (1 + e^{-\delta\sqrt{z}})^2 + v_{\delta,0}^{-1}(z) (1 - e^{-\delta\sqrt{z}})^2, & \text{if } r = 0 \end{cases}$$

**Remark 5.7.** Note that, from  $(H_2)$  and  $(H_3)$ , if  $r_{\pm} \neq 0$ , we have

$$\begin{cases} u_{c,r_-}(-A) = U_-, & u_{d,r_+}(-A) = U_+, & v_{c,r_-}(-A) = V_-, \\ v_{d,r_+}(-A) = V_+, & k_- f_{c,r_-,1}(-A) = P_1^-, & k_+ f_{d,r_+,1}(-A) = P_1^+, \\ k_- f_{c,r_-,2}(-A) = P_2^-, & k_+ f_{d,r_+,2}(-A) = P_2^+, & k_- f_{c,r_-,3}(-A) = P_3^-, \\ k_+ f_{d,r_+,3}(-A) = P_3^+, \end{cases}$$

and if  $r_- = 0$ , we obtain

$$\begin{cases} u_{c,0}(-A) = U_-, & v_{c,0}(-A) = V_-, & k_- f_{c,0,1}(-A) = Q_1^-, \\ k_- f_{c,0,2}(-A) = Q_2^-, & k_- f_{c,0,3}(-A) = Q_3^-. \end{cases}$$

**Remark 5.8.** Let  $\delta > 0$ ,  $r \in \mathbb{R}$  and  $x \in (r_m, +\infty)$ . Then, when  $r = 0$ , we have

$$1 - e^{-2\delta\sqrt{x}} \pm 2\delta\sqrt{x}e^{-\delta\sqrt{x}} = 2e^{-\delta\sqrt{x}} (\sinh(\delta\sqrt{x}) \pm \delta\sqrt{x}) > 0,$$

and from [20], Lemma 5.2, p. 369, it clear that  $u_{\delta,r}(x) > 0$  and  $v_{\delta,r}(x) > 0$ . Thus, when  $r \neq 0$ , we deduce that

$$f_{\delta,r,1}(x) > 0 \quad \text{and} \quad f_{\delta,r,2}(x), f_{\delta,r,3}(x) < 0,$$

and when  $r = 0$ , we obtain

$$f_{\delta,0,1}(x), f_{\delta,0,2}(x), f_{\delta,0,3}(x) > 0.$$

Moreover, for  $z \in \mathbb{C} \setminus (-\infty, r_m]$  and  $r \in \mathbb{R} \setminus \{0\}$ , we define

$$g_{\delta,r}(z) = -\sqrt{z+r} \left( \left(1 - e^{-2\delta(\sqrt{z+r}+\sqrt{z})}\right)^2 - \frac{1}{r^2}(\sqrt{z+r} + \sqrt{z})^4 \left(e^{-2\delta\sqrt{z}} - e^{-2\delta\sqrt{z+r}}\right)^2 \right) \\ + \sqrt{z} \left( \left(1 - e^{-\delta(\sqrt{z+r}+\sqrt{z})}\right)^2 + \frac{1}{r}(\sqrt{z+r} + \sqrt{z})^2 \left(e^{-\delta\sqrt{z}} - e^{-\delta\sqrt{z+r}}\right)^2 \right)^2,$$

and for  $r = 0$ , we set

$$g_{\delta,0}(z) = (1 + \sqrt{z}) \left(1 - e^{-2\delta\sqrt{z}}\right)^4 + 4 \left(1 - e^{-2\delta\sqrt{z}}\right)^2 e^{-2\delta\sqrt{z}} - 16 \delta^2 z e^{-4\delta\sqrt{z}}$$

**Lemma 5.9.** Let  $\delta > 0$  and  $x \in (r_m, +\infty)$ . Thus, if  $r \in \mathbb{R} \setminus \{0\}$ , then  $g_{\delta,r}(x) < 0$  and if  $r = 0$ , then  $g_{\delta,0}(x) > 0$ .

*Proof.* Let  $\delta > 0$  and  $r \in \mathbb{R} \setminus \{0\}$ . For all  $x \in (r_m, +\infty)$ , from [19], Lemma 4.4, p. 2950, we obtain the result.

Now, consider that  $r = 0$ . Then

$$g_{\delta,0}(x) = (1 + \sqrt{x}) \left(1 - e^{-2\delta\sqrt{x}}\right)^4 + 4e^{-2\delta\sqrt{x}} \left( \left(1 - e^{-2\delta\sqrt{x}}\right)^2 - 4\delta^2 x e^{-2\delta\sqrt{x}} \right) \\ = (1 + \sqrt{x}) \left(1 - e^{-2\delta\sqrt{x}}\right)^4 \\ + 4e^{-2\delta\sqrt{x}} \left(1 - e^{-2\delta\sqrt{x}} - 2\delta\sqrt{x} e^{-\delta\sqrt{x}}\right) \left(1 - e^{-2\delta\sqrt{x}} + 2\delta\sqrt{x} e^{-\delta\sqrt{x}}\right).$$

Hence, since  $\delta, x > 0$ , we have

$$1 - e^{-2\delta\sqrt{x}} - 2\delta\sqrt{x} e^{-\delta\sqrt{x}} = e^{-\delta\sqrt{x}} \left( e^{\delta\sqrt{x}} - e^{-\delta\sqrt{x}} - 2\delta\sqrt{x} \right) \\ = 2e^{-\delta\sqrt{x}} (\sinh(\delta\sqrt{x}) - \delta\sqrt{x}) > 0.$$

Finally, we deduce that  $g_{\delta,0} > 0$ . □

## 6 Proof of the main results

In both cases, assume that problem (P) has a unique classical solution; thus, from Theorem 5.3, respectively Theorem 5.5, (8) holds. Conversely, assume that (8) holds, then due to Theorem 5.3, respectively Theorem 5.5, we have to prove that system (17), respectively system (25), has a unique solution such that (18) holds.

The proof is divided in three parts for both cases. First, we will make explicit, in the first case, the determinant of system (17) and in the second case, the determinant of system (25). Then, in the two cases, we will show the uniqueness of the solution. To this end, we will invert the determinant thanks to functional calculus. Finally, we will prove, in all cases, that  $\psi_1$  and  $\psi_2$  have the expected regularity.

### 6.1 Calculus of the determinant

#### 6.1.1 First case

Here, we consider  $r_+, r_- \in \mathbb{R} \setminus \{0\}$ . We have to make explicit the determinant of system (17) that we recall here

$$\begin{cases} (P_1^- - P_1^+) M\psi_1 + (P_2^+ + P_2^-) \psi_2 = S_1 \\ (P_3^+ + P_3^-) \psi_1 + (P_1^- - P_1^+) \psi_2 = S_2. \end{cases}$$



We write the previous system as a matrix equation  $\Lambda_1 \Psi = S$ , where

$$\Lambda_1 = \begin{pmatrix} (P_1^- - P_1^+) M & (P_2^+ + P_2^-) \\ (P_3^+ + P_3^-) & (P_1^- - P_1^+) \end{pmatrix}, \quad \Psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad \text{and} \quad S = \begin{pmatrix} S_1 \\ S_2 \end{pmatrix}.$$

To solve system (17), we will study the determinant

$$\det(\Lambda_1) := M (P_1^- - P_1^+)^2 - (P_2^+ + P_2^-) (P_3^+ + P_3^-),$$

of the matrix  $\Lambda_1$ . Thus, we set

$$\det(\Lambda_1) = D_1^+ + D_1^- + D_2, \tag{33}$$

where

$$\begin{cases} D_1^+ &= M (P_1^+)^2 - P_3^+ P_2^+ \\ D_1^- &= M (P_1^-)^2 - P_3^- P_2^- \\ D_2 &= -P_3^+ P_2^- - P_3^- P_2^+ - 2MP_1^+ P_1^- \end{cases}$$

Then, we recall the result of [19] (Lemma 5.1, p. 2953), describing the determinant.

**Lemma 6.1** ([19]). We have

1.  $D_1^+ = -4k_+^2 (L_+ + M)^2 U_+^{-2} V_+^{-2} D^+$ , with

$$\begin{aligned} D^+ &= L_+ \left( (I - e^{2d(L_+ + M)})^2 - \frac{1}{r_+^2} (L_+ + M)^4 (e^{2dM} - e^{2dL_+})^2 \right) \\ &\quad - M \left( (I - e^{d(L_+ + M)})^2 + \frac{1}{r_+^2} (L_+ + M)^2 (e^{dM} - e^{dL_+})^2 \right)^2. \end{aligned}$$

2.  $D_1^- = -4k_-^2 (L_- + M)^2 U_-^{-2} V_-^{-2} D^-$ , with

$$\begin{aligned} D^- &= L_- \left( (I - e^{2c(L_- + M)})^2 - \frac{1}{r_-^2} (L_- + M)^4 (e^{2cM} - e^{2cL_-})^2 \right) \\ &\quad - M \left( (I - e^{c(L_- + M)})^2 + \frac{1}{r_-^2} (L_- + M)^2 (e^{cM} - e^{cL_-})^2 \right)^2. \end{aligned}$$

### 6.1.2 Second case

Here, we consider  $r_+ \in \mathbb{R} \setminus \{0\}$  and  $r_- = 0$ . As previously, we make explicit the determinant of system (25) that we recall here

$$\begin{cases} (P_1^+ - 2MQ_1^-) M \psi_1 - (P_2^+ + 2MQ_2^-) \psi_2 = S_3 \\ (P_3^+ + 2MQ_3^-) \psi_1 + (2MQ_1^- - P_1^+) \psi_2 = S_4, \end{cases}$$

We write this system as a matrix equation  $\Lambda_2 \Psi = \tilde{S}$ , where

$$\Lambda_2 = \begin{pmatrix} (P_1^+ - 2MQ_1^-) M & -(P_2^+ + 2MQ_2^-) \\ (P_3^+ + 2MQ_3^-) & (2MQ_1^- - P_1^+) \end{pmatrix}, \quad \Psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad \text{and} \quad \tilde{S} = \begin{pmatrix} S_3 \\ S_4 \end{pmatrix}.$$

To solve system (17), we will study the determinant

$$\det(\Lambda_2) := -M \left( P_1^+ - 2MQ_1^- \right)^2 + \left( P_3^+ + 2MQ_3^- \right) \left( P_2^+ + 2MQ_2^- \right),$$

of the matrix  $\Lambda_2$ . We set

$$\det(\Lambda_2) = D_3^+ + D_3^- + D_4, \quad (34)$$

where

$$D_3^+ = P_2^+ P_3^+ - M \left( P_1^+ \right)^2 \quad \text{and} \quad D_3^- = 4M^2 \left( Q_2^- Q_3^- - M \left( Q_1^- \right)^2 \right),$$

with

$$D_4 = 2M \left( P_3^+ Q_2^- + P_2^+ Q_3^- + M P_1^+ Q_1^- \right).$$

**Lemma 6.2.** We have

1.  $D_3^+ = k_+^2 (L_+ + M)^2 U_+^{-2} V_+^{-2} D_0^+$ , with

$$\begin{aligned} D_0^+ &= L_+ \left( \left( I - e^{2d(L_+ + M)} \right)^2 - \frac{1}{r_+^2} (L_+ + M)^4 \left( e^{2dM} - e^{2dL_+} \right)^2 \right) \\ &\quad - M \left( \left( I - e^{d(L_+ + M)} \right)^2 + \frac{1}{r_+^2} (L_+ + M)^2 \left( e^{dM} - e^{dL_+} \right)^2 \right)^2. \end{aligned}$$

2.  $D_3^- = 16k_-^2 M^2 U_-^{-2} V_-^{-2} D_0^-$ , with

$$D_0^- = (I - M) \left( I - e^{2cM} \right)^4 + 4 \left( I - e^{2cM} \right)^2 e^{2cM} - 16c^2 M^2 e^{4cM}.$$

*Proof.*

1. We have

$$P_2^+ P_3^+ = k_+^2 (L_+ + M)^2 L_+ U_+^{-2} V_+^{-2} D'_+,$$

where

$$\begin{aligned} D'_+ &= \left( U_+^2 + V_+^2 \right) \left( I - e^{2dM} \right) \left( I - e^{2dL_+} \right) \\ &\quad + U_+ V_+ \left( \left( I + e^{dM} \right)^2 \left( I + e^{dL_+} \right)^2 + \left( I - e^{dM} \right)^2 \left( I - e^{dL_+} \right)^2 \right) \\ &= \left( U_+^2 + V_+^2 \right) \left[ \left( I + e^{d(L_+ + M)} \right)^2 - \left( e^{dM} + e^{dL_+} \right)^2 \right] \\ &\quad + 2U_+ V_+ \left[ \left( I + e^{d(L_+ + M)} \right)^2 + \left( e^{dM} + e^{dL_+} \right)^2 \right] \\ &= \left( U_+ + V_+ \right)^2 \left( I + e^{d(L_+ + M)} \right)^2 - \left( V_+ - U_+ \right)^2 \left( e^{dM} + e^{dL_+} \right)^2. \end{aligned}$$

Moreover, from (9), we obtain that

$$U_+ + V_+ = 2 \left( I - e^{d(L_+ + M)} \right) \quad \text{and} \quad V_+ - U_+ = \frac{2}{r_+} (L_+ + M)^2 \left( e^{dM} - e^{dL_+} \right). \quad (35)$$

Then

$$\begin{aligned} D'_+ &= 4 \left( I - e^{d(L_+ + M)} \right)^2 \left( I + e^{d(L_+ + M)} \right)^2 \\ &\quad - \frac{4}{r_+^2} (L_+ + M)^4 \left( e^{dM} - e^{dL_+} \right)^2 \left( e^{dM} + e^{dL_+} \right)^2 \\ &= 4 \left( I - e^{2d(L_+ + M)} \right)^2 - \frac{4}{r_+^2} (L_+ + M)^4 \left( e^{2dM} - e^{2dL_+} \right)^2. \end{aligned}$$

Furthermore, we have

$$M \left( P_1^+ \right)^2 = k_+^2 (L_+ + M)^2 M U_+^{-2} V_+^{-2} D_+'' ,$$

where

$$\begin{aligned} D_+'' &= \left( V_+ (I + e^{dM}) (I - e^{dL_+}) + U_+ (I - e^{dM}) (I + e^{dL_+}) \right)^2 \\ &= \left[ (U_+ + V_+) \left( I - e^{d(L_+ + M)} \right) + (V_+ - U_+) \left( e^{dM} - e^{dL_+} \right) \right]^2 , \end{aligned}$$

and due to (35), it follows that

$$D_+'' = \left[ 2 \left( I - e^{d(L_+ + M)} \right)^2 + \frac{2}{r_+} (L_+ + M)^2 \left( e^{dM} - e^{dL_+} \right)^2 \right]^2 .$$

Finally, we deduce that

$$\begin{aligned} D_3^+ &= P_2^+ P_3^+ - M \left( P_1^+ \right)^2 \\ &= k_+^2 (L_+ + M)^2 U_+^{-2} V_+^{-2} (L_+ D_+^+ - M D_+'' ) , \end{aligned}$$

and setting  $D_0^+ = L_+ D_+^+ - M D_+''$ , we obtain the expected result.

2. We have

$$Q_2^- Q_3^- = k_-^2 U_-^{-2} V_-^{-2} D_-' ,$$

where

$$\begin{aligned} D_-' &= \left( V_- \left( I - e^{cM} \right)^2 + U_- \left( I + e^{cM} \right)^2 \right) \left( V_- \left( I + e^{cM} \right)^2 + U_- \left( I - e^{cM} \right)^2 \right) \\ &= \left( U_-^2 + V_-^2 \right) \left( I - e^{2cM} \right)^2 + 2U_- V_- \left( I - e^{2cM} \right)^2 + 16U_- V_- e^{2cM} , \end{aligned}$$

and

$$\begin{aligned} M \left( Q_1^- \right)^2 &= k_-^2 M U_-^{-2} V_-^{-2} (U_- + V_-)^2 \left( I - e^{2cM} \right)^2 \\ &= k_-^2 U_-^{-2} V_-^{-2} \left[ M \left( U_-^2 + V_-^2 \right) \left( I - e^{2cM} \right)^2 + 2M U_- V_- \left( I - e^{2cM} \right)^2 \right] . \end{aligned}$$

Thus

$$Q_2^- Q_3^- - M \left( Q_1^- \right)^2 = k_-^2 U_-^{-2} V_-^{-2} D_-'' ,$$

where

$$\begin{aligned} D_-'' &= (I - M) \left( U_-^2 + V_-^2 \right) \left( I - e^{2cM} \right)^2 + 2(I - M) U_- V_- \left( I - e^{2cM} \right)^2 \\ &\quad + 16U_- V_- e^{2cM} \\ &= (I - M) (U_- + V_-)^2 \left( I - e^{2cM} \right)^2 + 16U_- V_- e^{2cM} . \end{aligned}$$

Moreover, from (9), we obtain that

$$U_- + V_- = 2 \left( I - e^{2cM} \right) \quad \text{and} \quad U_- V_- = \left( I - e^{2cM} \right)^2 - 4c^2 M^2 e^{2cM} .$$

Then

$$D''_- = 4(I - M) \left( I - e^{2cM} \right)^4 + 16 \left( I - e^{2cM} \right)^2 e^{2cM} - 64c^2 M^2 e^{4cM}.$$

Therefore, it follows that

$$\begin{aligned} D_3^- &= 4M^2 \left( Q_2^- Q_3^- - M \left( Q_1^- \right)^2 \right) \\ &= 16k_-^2 M^2 U_-^{-2} V_-^{-2} D_0^-, \end{aligned}$$

where  $D_0^- = \frac{1}{4} D''_-$ .

□

## 6.2 Inversion of the determinant

### 6.2.1 First case

Here, we consider  $r_+, r_- \in \mathbb{R} \setminus \{0\}$ . Let  $r = \max(-r_+, -r_-, 0) \geq 0$ . By using functional calculus, we prove that the determinant of system (17), given by (33), is invertible with bounded inverse. Due to Lemma 6.1 and the definition of  $D_2$ , we obtain:

$$D_1^+ = g_1^+(-A), \quad D_1^- = g_1^-(-A) \quad \text{and} \quad D_2 = g_2(-A),$$

where, for  $z \in \mathbb{C} \setminus \mathbb{R}_-$ , we have set

$$\begin{cases} g_1^+(z) &= 4k_+^2 (\sqrt{z+r_+} + \sqrt{z})^2 u_{d,r_+}^{-2}(z) v_{d,r_+}^{-2}(z) g_{d,r_+}(z) \\ g_1^-(z) &= 4k_-^2 (\sqrt{z+r_-} + \sqrt{z})^2 u_{c,r_-}^{-2}(z) v_{c,r_-}^{-2}(z) g_{c,r_-}(z) \\ g_2(z) &= k_+ f_{d,r_+,1}(z) k_- f_{c,r_-,3}(z) + k_- f_{c,r_-,1}(z) k_+ f_{d,r_+,3}(z) \\ &\quad - 2\sqrt{z} k_+ f_{d,r_+,2}(z) k_- f_{c,r_-,2}(z), \end{cases}$$

with  $u_{\delta,r}$ ,  $v_{\delta,r}$ ,  $g_{\delta,r}$  and  $f_{\delta,r,i}$  the complex functions defined in section 5.2. Thus

$$\det(\Lambda_1) = D_1^+ + D_1^- + D_2 = f_1(-A), \tag{36}$$

with  $f_1 = g_1^+ + g_1^- + g_2$ . Note that, for some  $\theta \in (0, \pi)$ , we have  $f_1 \in H(S_\theta)$  and due to Remark 5.8 and Lemma 5.9, for  $x > 0$ , we have

$$f_1(x) = g_1^+(x) + g_1^-(x) + g_2(x) < 0. \tag{37}$$

Let  $C_1, C_2$  be two linear operators in  $X$ . We denote by  $C_1 \sim C_2$  the equality  $C_1 = C_2 + \Sigma$ , where  $\Sigma$  is a finite sum of terms of type  $kL_+^l L_-^m M^n e^{\alpha L_+} e^{\beta L_-} e^{\delta M}$ , where  $k \in \mathbb{R}$ ;  $l, m, n \in \mathbb{N}$ ;  $\alpha, \beta, \delta \in \mathbb{R}_+$  with  $\alpha + \beta + \delta \neq 0$ . Note that  $\Sigma$  is a regular term in the sense:

$$\Sigma \in \mathcal{L}(X) \quad \text{with} \quad \Sigma(X) \subset D(M^\infty) := \bigcap_{k \geq 0} D(M^k).$$

Since we have  $U_\pm \sim I, V_\pm \sim I$ , then by setting  $W = U_- U_+ V_- V_+ \sim I$ , we deduce that

$$\begin{cases} WP_1^+ \sim 2k_+(L_+ + M), & WP_1^- \sim 2k_-(L_- + M) \\ WP_2^+ \sim 2k_+(L_+ + M), & WP_2^- \sim 2k_-(L_- + M) \\ WP_3^+ \sim 2k_+(L_+ + M)L_+, & WP_3^- \sim 2k_-(L_- + M)L_- \end{cases}$$

Thus

$$\begin{aligned}
W^2 \det(\Lambda_1) &= M \left( WP_1^+ \right)^2 - \left( WP_2^+ WP_3^+ \right) + M \left( WP_1^- \right)^2 - \left( WP_2^- WP_3^- \right) \\
&\quad - \left( WP_2^- WP_3^+ + WP_2^+ WP_3^- + 2MWP_1^+ WP_1^- \right) \\
&\sim -4k_+^2 (L_+ + M)^2 (L_+ - M) - 4k_-^2 (L_- + M)^2 (L_- - M) \\
&\quad - 4k_+ k_- (L_+ + M)(L_- + M)(L_+ + L_- + 2M).
\end{aligned}$$

From (7), we have

$$\begin{aligned}
-W^2 \det(\Lambda_1) &\sim 4k_+^2 r_+(L_+ + M) + 4k_-^2 r_-(L_- + M) \\
&\quad + 4k_+ k_- (L_+ + M)(L_- + M)(L_+ + L_- + 2M) \\
&\sim 4k_+ l_+(L_+ + M) + 4k_- l_-(L_- + M) \\
&\quad + 4k_+ k_- (L_+ + M)(L_- + M)(L_+ + L_- + 2M)
\end{aligned}$$

Hence, we note

$$B_1 = 4k_+ l_+(L_+ + M) + 4k_- l_-(L_- + M) + 4k_+ k_- (L_+ + M)(L_- + M)(L_+ + L_- + 2M).$$

Thus, we obtain

$$\det(\Lambda_1) = -W^{-2} \left( B_1 + \sum_{j \in J} k_j L_+^{l_j} L_-^{m_j} M^{n_j} e^{\alpha_j L_+} e^{\beta_j L_-} e^{\delta_j M} \right), \quad (38)$$

where  $J$  is a finite set and for any  $j \in J$ :

$$k_j \in \mathbb{R}; \quad l_j, m_j, n_j \in \mathbb{N}, \quad \alpha_j, \beta_j, \delta_j \in \mathbb{R}_+ \quad \text{with} \quad \alpha_j + \beta_j + \delta_j \neq 0.$$

We set

$$B_2 = I + \frac{l_+}{k_-} (L_- + M)^{-1} (L_+ + L_- + 2M)^{-1} + \frac{l_-}{k_+} (L_+ + M)^{-1} (L_+ + L_- + 2M)^{-1}$$

such that

$$B_1 = 4k_+ k_- (L_+ + M)(L_- + M)(L_+ + L_- + 2M)B_2.$$

**Proposition 6.3.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+ k_- > 0$ . If one of the following assumptions holds

- $\frac{l_+}{k_-} > 0$  and  $\frac{l_-}{k_+} > 0$ ,
- $\frac{l_+}{k_-} < 0$  and  $\frac{l_-}{k_+} < 0$ , such that

$$(l_+ - l_-)(k_+ - k_-) \geq 0, \quad (39)$$

- $\frac{l_+}{k_-} > 0$  and  $\frac{l_-}{k_+} < 0$ , such that

$$-6l_- k_+ + l_+ k_+ + l_- k_- \geq 0, \quad (40)$$

- $\frac{l_+}{k_-} < 0$  and  $\frac{l_-}{k_+} > 0$ , such that

$$-6l_+k_- + l_+k_+ + l_-k_- \geq 0, \quad (41)$$

then,  $b_2(x) > 0$ , for  $x > r \geq 0$  and operator  $B_1$ , defined above, is invertible with bounded inverse.

**Remark 6.4.** Since  $k_+k_- > 0$ , then we have the following equivalences

$$\frac{l_+}{k_-} > 0 \iff r_+ > 0 \quad \text{and} \quad \frac{l_-}{k_+} > 0 \iff r_- > 0.$$

*Proof.* From  $(H_2)$  and  $(H_3)$ , since  $k_+k_- \neq 0$ , it is clear that

$$0 \in \rho(4k_+k_-(L_+ + M)(L_- + M)(L_+ + L_- + 2M)).$$

Thus, it remains to prove that  $0 \in \rho(B_2)$ . To this end, we use Lemma 5.6.

Let  $z \in \mathbb{C} \setminus (-\infty, r]$ . We set

$$\begin{aligned} b_2(z) &= 1 + \frac{l_+}{k_-} \frac{1}{(\sqrt{z+r_-} + \sqrt{z})(\sqrt{z+r_+} + \sqrt{z+r_-} + 2\sqrt{z})} \\ &\quad + \frac{l_-}{k_+} \frac{1}{(\sqrt{z+r_+} + \sqrt{z})(\sqrt{z+r_+} + \sqrt{z+r_-} + 2\sqrt{z})}, \end{aligned} \quad (42)$$

hence  $b_2(-A) = B_2$ . Then, for all  $x > r \geq 0$ , it follows

$$\begin{aligned} b_2(x) &= 1 + \frac{l_+}{k_-} \frac{1}{(\sqrt{x+r_-} + \sqrt{x})(\sqrt{x+r_+} + \sqrt{x+r_-} + 2\sqrt{x})} \\ &\quad + \frac{l_-}{k_+} \frac{1}{(\sqrt{x+r_+} + \sqrt{x})(\sqrt{x+r_+} + \sqrt{x+r_-} + 2\sqrt{x})} \end{aligned}$$

Our aim is to prove that  $b_2(x) > 0$ , for all  $x > r$ . To this end, we set

$$y = x - r > 0,$$

hence

$$\begin{aligned} b_2(y+r) &= 1 + \frac{l_+}{k_-} \frac{1}{(\sqrt{y+r+r_-} + \sqrt{y+r})(\sqrt{y+r+r_+} + \sqrt{y+r+r_-} + 2\sqrt{y+r})} \\ &\quad + \frac{l_-}{k_+} \frac{1}{(\sqrt{y+r+r_+} + \sqrt{y+r})(\sqrt{y+r+r_+} + \sqrt{y+r+r_-} + 2\sqrt{y+r})} \\ &= 1 + \frac{1}{(\sqrt{y+r+r_+} + \sqrt{y+r+r_-} + 2\sqrt{y+r})} b_3(y), \end{aligned}$$

where

$$b_3(y) = \frac{\frac{l_+}{k_-}}{(\sqrt{y+r+r_-} + \sqrt{y+r})} + \frac{\frac{l_-}{k_+}}{(\sqrt{y+r+r_+} + \sqrt{y+r})}.$$

Then

$$b'_3(y) = \frac{-\frac{l_+}{k_-} \left( \frac{1}{2\sqrt{y+r+r_-}} + \frac{1}{2\sqrt{y+r}} \right)}{(\sqrt{y+r+r_-} + \sqrt{y+r})^2} + \frac{-\frac{l_-}{k_+} \left( \frac{1}{2\sqrt{y+r+r_+}} + \frac{1}{2\sqrt{y+r}} \right)}{(\sqrt{y+r+r_+} + \sqrt{y+r})^2}$$

and

$$b_2'(y+r) = \frac{-\left(\frac{1}{2\sqrt{y+r+r_+}} + \frac{1}{2\sqrt{y+r+r_-}} + \frac{1}{\sqrt{y+r}}\right)}{(\sqrt{y+r+r_+} + \sqrt{y+r+r_-} + 2\sqrt{y+r})^2} b_3(y) + \frac{1}{(\sqrt{y+r+r_+} + \sqrt{y+r+r_-} + 2\sqrt{y+r})} b_3'(y),$$

Now, we have to study the following fourth cases.

1. If  $\frac{l_+}{k_-} > 0$  and  $\frac{l_-}{k_+} > 0$ , then it is clear that  $b_3 > 0$  and  $b_2 > 0$ .
2. If  $\frac{l_+}{k_-} < 0$  and  $\frac{l_-}{k_+} < 0$ , then  $b_3' > 0$  and  $b_2' > 0$ . Thus  $b_2(y+r) > b_2(r)$  where

$$b_2(r) = 1 + \frac{1}{(\sqrt{r+r_+} + 2\sqrt{r})} b_3(0) > 1 + \frac{1}{2\sqrt{r}} b_3(0),$$

with

$$b_3(0) = \frac{\frac{l_+}{k_-}}{\sqrt{r+r_-} + \sqrt{r}} + \frac{\frac{l_-}{k_+}}{\sqrt{r+r_+} + \sqrt{r}}.$$

Since  $\sqrt{r+r_+} > 0$  and  $\sqrt{r+r_-} > 0$ , it follows that

$$\frac{\frac{l_+}{k_-}}{\sqrt{r+r_-} + \sqrt{r}} > \frac{\frac{l_+}{k_-}}{\sqrt{r}} \quad \text{and} \quad \frac{\frac{l_-}{k_+}}{\sqrt{r+r_+} + \sqrt{r}} > \frac{\frac{l_-}{k_+}}{\sqrt{r}},$$

hence

$$b_3(0) > \frac{1}{\sqrt{r}} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right).$$

Thus, we obtain

$$b_2(r) > 1 + \frac{1}{2r} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right).$$

Moreover, we have

$$1 + \frac{1}{2r} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right) \geq 0 \iff 2r \geq - \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right),$$

where

$$- \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right) = \begin{cases} -r_- \left( \frac{l_+}{l_-} + \frac{k_-}{k_+} \right), & \text{if } r = -r_- \\ -r_+ \left( \frac{k_+}{k_-} + \frac{l_-}{l_+} \right), & \text{if } r = -r_+. \end{cases}$$

Thus, we obtain that

$$2r \geq - \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right) \iff \begin{cases} l_+k_+ + l_-k_- - 2l_-k_+ \geq 0, & \text{if } r = -r_- \\ l_+k_+ + l_-k_- - 2l_+k_- \geq 0, & \text{if } r = -r_+. \end{cases}$$

Furthermore, since  $k_+k_- > 0$ , if  $r = -r_-$ , then  $-\frac{l_-}{k_-} \geq -\frac{l_+}{k_+}$ , hence  $-l_-k_+ \geq -l_+k_-$

and if  $r = -r_+$ , then  $-\frac{l_+}{k_+} \geq -\frac{l_-}{k_-}$ , hence  $-l_+k_- \geq -l_-k_+$ . It follows that

$$\begin{cases} l_+k_+ + l_-k_- - 2l_-k_+ \geq l_+k_+ + l_-k_- - l_+k_- - l_-k_+, & \text{if } r = -r_- \\ l_+k_+ + l_-k_- - 2l_+k_- \geq l_+k_+ + l_-k_- - l_+k_- - l_-k_+, & \text{if } r = -r_+. \end{cases}$$

Finally, if (39) holds, then we obtain  $b_2 > 0$ .

3. If  $\frac{l_+}{k_-} > 0$  and  $\frac{l_-}{k_+} < 0$ , then since  $k_+k_- > 0$ , we have

$$\frac{l_+}{k_-} > 0 \iff \frac{l_+ k_+}{k_+ k_-} k_-^2 > 0 \iff r_+ > 0 \quad \text{and} \quad \frac{l_-}{k_+} < 0 \iff \frac{l_- k_-}{k_- k_+} k_+^2 < 0 \iff r_- < 0.$$

Thus  $r = -r_- > 0$ ,  $r_+ > 0$  and

$$b_3(y) = \frac{\frac{l_+}{k_-}}{(\sqrt{y} + \sqrt{y+r})} + \frac{\frac{l_-}{k_+}}{(\sqrt{y+r+r_+} + \sqrt{y+r})}.$$

Since  $\frac{l_+}{k_-} > 0$  and  $\sqrt{y} < \sqrt{y+r}$  it follows that

$$\frac{\frac{l_+}{k_-}}{\sqrt{y} + \sqrt{y+r}} > \frac{\frac{l_+}{k_-}}{2\sqrt{y+r}}.$$

In the same way, since  $\frac{l_-}{k_+} < 0$  and  $\sqrt{y+r+r_+} > \sqrt{y+r}$ , we deduce that

$$\frac{\frac{l_-}{k_+}}{\sqrt{y+r+r_+} + \sqrt{y+r}} > \frac{\frac{l_-}{k_+}}{2\sqrt{y+r}},$$

hence

$$b_3(y) > \frac{1}{2\sqrt{y+r}} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right).$$

If  $\frac{l_+}{k_-} + \frac{l_-}{k_+} > 0$ , then  $b_3 > 0$  and  $b_2 > 0$ . If  $\frac{l_+}{k_-} + \frac{l_-}{k_+} < 0$ , then we have

$$\begin{aligned} b_2(y+r) &= 1 + \frac{1}{(\sqrt{y+r+r_+} + \sqrt{y} + 2\sqrt{y+r})} b_3(y) \\ &> 1 + \frac{1}{3\sqrt{y+r}} b_3(y) \\ &> 1 + \frac{1}{6(y+r)} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right). \end{aligned}$$

Moreover, we have

$$1 + \frac{1}{6(y+r)} \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right) \geq 0 \iff 6(y+r) + \frac{l_+}{k_-} + \frac{l_-}{k_+} \geq 0.$$

It is obvious that

$$6(y+r) + \left( \frac{l_+}{k_-} + \frac{l_-}{k_+} \right) \geq 6r + \frac{l_+}{k_-} + \frac{l_-}{k_+},$$

thus, since  $k_+k_- > 0$  and here  $r = -r_- = -\frac{l_-}{k_-}$ , we deduce that the previous inequality becomes

$$-6\frac{l_-}{k_-} + \frac{l_+}{k_-} + \frac{l_-}{k_+} \geq 0 \iff -6l_-k_+ + l_+k_+ + l_-k_- \geq 0.$$

Finally, since  $k_+k_- > 0$ , if (40) holds, then  $b_2 > 0$ .

4. If  $\frac{l_+}{k_-} < 0$  and  $\frac{l_-}{k_+} > 0$ , then here  $r = -r_+$  and in the same way than previously, if (41) holds, then  $b_2 > 0$ .



Since  $r = \max(-r_+, -r_-, 0) \geq 0$  and due to  $(H_2)$  and  $(H_3)$ , we deduce that operator  $-A - rI \in \text{BIP}(X, \theta_A)$  with  $0 \in \rho(-A - rI)$ . Thus, considering  $\tilde{b}_2(z) = b_2(z + r)$ , with  $z + r \in \mathbb{C} \setminus \mathbb{R}_-$ , it follows that  $\tilde{b}_2(-A - rI) = B_2$ . Moreover, for a given  $\theta \in (0, \pi)$ , it is clear that  $1 - b_2, 1 - \tilde{b}_2 \in \mathcal{E}_\infty$ . Finally, applying Lemma 5.6 with  $G = \tilde{b}_2$  and  $P = -A - rI$ , we deduce the result.  $\square$

Due to (38) and Proposition 6.3, it follows that

$$\det(\Lambda_1) = -W^{-2}B_1F_1, \quad (43)$$

where

$$F_1 = I + \sum_{j \in J} k_j B_1^{-1} L_+^{l_j} L_-^{m_j} M^{n_j} e^{\alpha_j L_+} e^{\beta_j L_-} e^{\delta_j M}. \quad (44)$$

For  $z \in \mathbb{C} \setminus (-\infty, r]$ , we set

$$b_1(z) = -4k_+k_-(\sqrt{z+r_+} + \sqrt{z})(\sqrt{z+r_-} + \sqrt{z})(\sqrt{z+r_+} + \sqrt{z+r_-} + 2\sqrt{z})b_2(z), \quad (45)$$

where  $b_2$  is given by (42) and

$$\tilde{f}_1(z) = 1 + \sum_{j \in J} k_j b_1(z)^{-1} (-\sqrt{z+r_+})^{l_j} (-\sqrt{z+r_-})^{m_j} (-\sqrt{z})^{n_j} e^{-\alpha_j \sqrt{z+r_+}} e^{-\beta_j \sqrt{z+r_-}} e^{-\delta_j \sqrt{z}}.$$

Then, due to  $(H_2)$  and  $(H_3)$ , we have  $B_1 = b_1(-A)$  and  $F_1 = \tilde{f}_1(-A)$ . Moreover, from (36) and (43), we obtain

$$f_1(-A) = \det(\Lambda_1) = -W^{-2}B_1\tilde{f}_1(-A).$$

Note that, we have

$$f_1(z) = -u_{d,r_+}^{-2}(z)v_{d,r_+}^{-2}(z)u_{c,r_-}^{-2}(z)v_{c,r_-}^{-2}(z)b_1(z)\tilde{f}_1(z). \quad (46)$$

**Proposition 6.5.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+k_- > 0$ . Thus

- if  $r_+ > 0$  and  $r_- > 0$ ,
- if  $r_+ < 0$  and  $r_- < 0$ , such that (39) holds,
- if  $r_+ > 0$  and  $r_- < 0$ , such that (40) holds,
- if  $r_+ < 0$  and  $r_- > 0$ , such that (41) holds,

then,  $F_1 \in \mathcal{L}(X)$ , given by (44), is invertible with bounded inverse.

*Proof.* For a given  $\theta \in (0, \pi)$ , we have  $f_1, \tilde{f}_1 \in H(S_\theta)$ . Moreover, for  $z \in \mathbb{C} \setminus (-\infty, r]$ , since

$$k_j b_1^{-1}(z) (-\sqrt{z+r_+})^{l_j} (-\sqrt{z+r_-})^{m_j} (-\sqrt{z})^{n_j}, \quad \text{for all } j \in J,$$

are polynomial functions, we deduce that  $1 - \tilde{f}_1 \in \mathcal{E}_\infty(S_\theta)$ .

From (37), Proposition 6.3 and Remark 6.4, we know that  $f_1 < 0$  and  $b_2 > 0$  on  $(r, +\infty)$ . Then, since  $u_{d,r_+}, u_{c,r_-}, v_{d,r_+}, v_{c,r_-} > 0$  on  $(r, +\infty)$  and due to (45) and (46), we deduce that  $\tilde{f}_1 < 0$  on  $(r, +\infty)$ .

Therefore, noting  $\tilde{f}_{r,1}(z) = \tilde{f}_1(z + r)$  and applying Lemma 5.6 with  $G = \tilde{f}_1$  and operator  $P = -A - rI$ , thus we deduce that operator  $F_1 = \tilde{f}_{r,1}(-A - rI) = \tilde{f}_1(-A)$  is invertible with bounded inverse.  $\square$

This result finally leads us to state the following main result of this section.

**Proposition 6.6.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+k_- > 0$ . Thus

- if  $r_+ > 0$  and  $r_- > 0$ ,
- if  $r_+ < 0$  and  $r_- < 0$ , such that (39) holds,
- if  $r_+ > 0$  and  $r_- < 0$ , such that (40) holds,
- if  $r_+ < 0$  and  $r_- > 0$ , such that (41) holds,

then  $\det(\Lambda_1)$  is invertible with bounded inverse.

*Proof.* From (43), Proposition 6.3 and Proposition 6.5, it follows that  $\det(\Lambda_1) = -W^{-2}B_1F_1$ , is invertible with bounded inverse.  $\square$

### 6.2.2 Second case

Let  $r_+ \in \mathbb{R} \setminus \{0\}$  and  $r_- = 0$ . In the same way than previously, using functional calculus, we prove that the determinant of system (25), given by (34), is invertible with bounded inverse. Due to Lemma 6.2, and the definition of  $D_4$ , we obtain:

$$D_3^+ = g_3^+(-A), \quad D_3^- = g_3^-(-A) \quad \text{and} \quad D_4 = g_4(-A),$$

where, for  $z \in \mathbb{C} \setminus \mathbb{R}_-$ , we have set

$$\begin{cases} g_3^+(z) &= 4k_+^2(\sqrt{z+r_+} + \sqrt{z})^2 u_{d,r_+}^{-2}(z) v_{d,r_+}^{-2}(z) g_{d,r_+}(z) \\ g_3^-(z) &= 16k_-^2 z u_{c,0}^{-2}(z) v_{c,0}^{-2}(z) g_{c,0}(z) \\ g_4(z) &= -2\sqrt{z}(k_+ f_{d,r_+,3}(z) k_- f_{c,0,2}(z) + k_+ f_{d,r_+,2}(z) k_- f_{c,0,3}(z)) \\ &\quad + 2zk_+ f_{d,r_+,1}(z) k_- f_{c,0,1}(z), \end{cases}$$

with  $u_{\delta,r}$ ,  $v_{\delta,r}$ ,  $g_{\delta,r}$  and  $f_{\delta,r,i}$  the complex functions defined in section 5.2. Thus

$$\det(\Lambda_2) = D_3^+ + D_3^- + D_4 = f_2(-A), \quad (47)$$

with  $f_2 = g_3^+ + g_3^- + g_4$ . Note that, for some  $\theta \in (0, \pi)$ , we have  $f_2 \in H(S_\theta)$  and due to Remark 5.8 and Lemma 5.9, for  $x > \max(-r_+, 0)$ , we have

$$f_2(x) = g_3^+(x) + g_3^-(x) + g_4(x), \quad \text{where} \quad g_3^+ < 0 \quad \text{and} \quad g_3^-, g_4 > 0. \quad (48)$$

**Lemma 6.7.** Let  $k_+k_- > 0$ . Then

- if  $r_+ > 0$  such that

$$r_+ \geq \frac{(\sqrt{t+1} + \sqrt{t})^2}{t^2} \frac{k_+^2}{4k_-^2}, \quad \text{for } t > 0 \text{ fixed.} \quad (49)$$

for all  $x \geq tr_+$ , we have  $f_2(x) > 0$ .

- if  $r_+ < 0$  such that

$$r_+ \leq -\frac{27k_+^2}{64k_-^2}, \quad (50)$$

for all  $x \geq -r_+$ , we have  $f_2(x) > 0$ .

*Proof.* From (48), we deduce

$$f_2(x) \geq g_3^+(x) + g_4(x) \geq g_3^+(x) + 2k_+k_- x f_{d,r_+,1}(x) f_{c,0,1}(x).$$

Let  $r = \max(-r_+, 0)$ . For  $x \in (r, +\infty)$ , setting  $y = x - r > 0$  and noting

$$h_1(y) = g_3^+(y+r) + 2k_+k_- (y+r) f_{d,r_+,1}(y+r) f_{c,0,1}(y+r),$$

it follows

$$\begin{aligned} h_1(y) &= 4k_+^2 \frac{(\sqrt{y+r+r_+} + \sqrt{y+r})^2}{u_{d,r_+}^2(y+r)v_{d,r_+}^2(y+r)} g_{d,r_+}(y+r) \\ &\quad + 2k_+k_- (y+r) f_{d,r_+,1}(y+r) f_{c,0,1}(y+r). \end{aligned}$$

Since we have

$$0 > g_{d,r_+}(y+r) \geq -\sqrt{y+r+r_+} \left(1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right)^2,$$

then

$$h_1(y) \geq 4 \frac{(\sqrt{y+r+r_+} + \sqrt{y+r})\sqrt{y+r+r_+}}{u_{d,r_+}^2(y+r)v_{d,r_+}^2(y+r)u_{c,0}(y+r)v_{c,0}(y+r)} h_2(y),$$

where

$$\begin{aligned} h_2(y) &= -k_+^2 (\sqrt{y+r+r_+} + \sqrt{y+r}) \left(1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right)^2 u_{c,0}(y+r)v_{c,0}(y+r) \\ &\quad + k_+k_- (y+r) \left(1 - e^{-2c\sqrt{y+r}}\right)^2 v_{d,r_+}(y+r) \left(1 + e^{-d\sqrt{y+r}}\right) \left(1 + e^{-d\sqrt{y+r+r_+}}\right) \\ &\quad + k_+k_- (y+r) \left(1 - e^{-2c\sqrt{y+r}}\right)^2 u_{d,r_+}(y+r) \left(1 - e^{-d\sqrt{y+r}}\right) \left(1 - e^{-d\sqrt{y+r+r_+}}\right) \\ &\geq -k_+^2 (\sqrt{y+r+r_+} + \sqrt{y+r}) \left(1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right)^2 \left(1 - e^{-2c\sqrt{y+r}}\right)^2 \\ &\quad + k_+k_- (y+r) \left(1 - e^{-2c\sqrt{y+r}}\right)^2 h_3(y), \end{aligned}$$

with

$$\begin{aligned} h_3(y) &= v_{d,r_+}(y+r) \left(1 + e^{-d\sqrt{y+r}}\right) \left(1 + e^{-d\sqrt{y+r+r_+}}\right) \\ &\quad + u_{d,r_+}(y+r) \left(1 - e^{-d\sqrt{y+r}}\right) \left(1 - e^{-d\sqrt{y+r+r_+}}\right) \end{aligned}$$

and

$$\begin{aligned} h_3(y) &= 2v_{d,r_+}(y+r) \left(1 + e^{-d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right) + 2u_{d,r_+}(y+r) \left(e^{-d\sqrt{y+r}} + e^{-d\sqrt{y+r+r_+}}\right) \\ &= 2 \left(1 - e^{-d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right) \left(1 + e^{-d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right) \\ &\quad + 2 \frac{(\sqrt{y+r+r_+} + \sqrt{y+r})^2}{r_+} \left(e^{-d\sqrt{y+r}} - e^{-d\sqrt{y+r+r_+}}\right) \left(e^{-d\sqrt{y+r}} + e^{-d\sqrt{y+r+r_+}}\right) \\ &= 2 \left(1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})}\right) \\ &\quad + 2 \frac{(\sqrt{y+r+r_+} + \sqrt{y+r})^2}{r_+} \left(e^{-2d\sqrt{y+r}} - e^{-2d\sqrt{y+r+r_+}}\right). \end{aligned}$$

Moreover, for all  $y > 0$ , since we have

$$\frac{e^{-2d\sqrt{y+r}} - e^{-2d\sqrt{y+r+r_+}}}{r_+} > 0, \quad \text{for } r_+ \in \mathbb{R} \setminus \{0\},$$

we deduce that

$$h_3(y) > 2 \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right),$$

and

$$\begin{aligned} h_2(y) &> -k_+^2(\sqrt{y+r+r_+} + \sqrt{y+r}) \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right)^2 \left( 1 - e^{-2c\sqrt{y+r}} \right)^2 \\ &\quad + 2k_+k_- (y+r) \left( 1 - e^{-2c\sqrt{y+r}} \right)^2 \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right) \\ &> -k_+^2(\sqrt{y+r+r_+} + \sqrt{y+r}) \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right) \left( 1 - e^{-2c\sqrt{y+r}} \right)^2 \\ &\quad + 2k_+k_- (y+r) \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right) \left( 1 - e^{-2c\sqrt{y+r}} \right)^2 \\ &> \left( 1 - e^{-2d(\sqrt{y+r+r_+} + \sqrt{y+r})} \right) \left( 1 - e^{-2c\sqrt{y+r}} \right)^2 h_4(y), \end{aligned}$$

where

$$h_4(y) = 2k_+k_- (y+r) - k_+^2(\sqrt{y+r+r_+} + \sqrt{y+r}).$$

Thus

$$h_4(y) \geq 0 \iff \frac{y+r}{\sqrt{y+r+r_+} + \sqrt{y+r}} \geq \frac{k_+^2}{2k_+k_-}. \quad (51)$$

We set

$$h_5(y) = \frac{y+r}{\sqrt{y+r+r_+} + \sqrt{y+r}}, \quad (52)$$

hence

$$h_5'(y) = \left( \frac{1}{\sqrt{y+r+r_+} + \sqrt{y+r}} \right) \left( 1 - \frac{1}{2} \sqrt{\frac{y+r}{y+r+r_+}} \right).$$

1. If  $r_+ < 0$ , then  $r = -r_+$  and

$$\frac{y+r}{y+r+r_+} = \frac{y+r}{y},$$

moreover

$$h_5'(y) \geq 0 \iff 1 - \frac{1}{2} \sqrt{\frac{y+r}{y}} \geq 0 \iff 4 \geq \frac{y+r}{y} \iff y \geq \frac{r}{3}.$$

Thus, we have

$$h_5(y) \geq h_5\left(\frac{r}{3}\right) = \frac{\frac{4r}{3}}{\sqrt{\frac{r}{3}} + 2\sqrt{\frac{r}{3}}} = \frac{4}{3} \sqrt{\frac{r}{3}} > 0.$$

This yields that for all  $y \geq 0$ , we have

$$h_5(y) \geq \frac{4}{3} \sqrt{\frac{r}{3}} > 0.$$

Therefore, from (51) and (52), we deduce that

$$h_4(y) \geq 0 \iff h_5(y) \geq \frac{k_+^2}{2k_+k_-} \iff \frac{4}{3} \sqrt{\frac{r}{3}} \geq \frac{k_+}{2k_-},$$

hence, since  $r = -r_+ > 0$ , we obtain

$$\frac{4}{3} \sqrt{\frac{r}{3}} \geq \frac{k_+}{2k_-} \iff \sqrt{\frac{r}{3}} \geq \frac{3k_+}{8k_-} \iff \frac{r}{3} \geq \frac{9k_+^2}{64k_-^2} \iff -r_+ \geq \frac{27k_+^2}{64k_-^2}.$$

2. If  $r_+ > 0$ , then  $r = 0$  and

$$\frac{y+r}{y+r+r_+} = \frac{y}{y+r_+} < 1,$$

hence  $h'_5 > 0$  and  $h_5$  is an increasing function. Thus, from (52), since  $r = 0$ , it follows that

$$h_5(y) = \frac{y}{\sqrt{y+r_+} + \sqrt{y}},$$

then

$$h_5(y) \geq \frac{k_+^2}{2k_+k_-} \iff \frac{y}{\sqrt{y+r_+} + \sqrt{y}} \geq \frac{k_+}{2k_-}.$$

Moreover, for  $t > 0$  fixed, we have

$$h_5(tr_+) \geq \frac{k_+}{2k_-} \iff \frac{tr_+}{\sqrt{(t+1)r_+} + \sqrt{tr_+}} \geq \frac{k_+}{2k_-} \iff \frac{t}{\sqrt{t+1} + \sqrt{t}} \sqrt{r_+} \geq \frac{k_+}{2k_-},$$

hence

$$\sqrt{r_+} \geq \frac{\sqrt{t+1} + \sqrt{t}}{t} \frac{k_+}{2k_-} \iff r_+ \geq \frac{(\sqrt{t+1} + \sqrt{t})^2}{t^2} \frac{k_+^2}{4k_-^2}.$$

Finally, if  $r_+ > 0$  such that (49) holds, then since  $y = x$ , for all  $x \geq tr_+$ , we have  $h_2(x) > 0$ ,  $h_1(x) > 0$  and  $f_2(x) > 0$ . Moreover,  $r_+ < 0$  such that (50) holds, then for all  $y > 0$ , we have  $h_2(y) > 0$ ,  $h_1(y) > 0$  and since  $y = x + r_+$ , for all  $x > -r_+$ , it follows that  $f_2(x) > 0$ .  $\square$

Therefore, as in the first case, since we have  $U_{\pm} \sim I$  and  $V_{\pm} \sim I$ , then by setting  $W = U_-U_+V_-V_+ \sim I$ , we deduce that

$$\begin{cases} WP_1^+ \sim 2k_+(L_+ + M), & WQ_1^- \sim 2k_-I \\ WP_2^+ \sim 2k_+(L_+ + M), & WQ_2^- \sim 2k_-I \\ WP_3^+ \sim 2k_+(L_+ + M)L_+, & WQ_3^- \sim 2k_-I. \end{cases}$$

Thus

$$\begin{aligned} W^2 \det(\Lambda_2) &= \left( WP_2^+ WP_3^+ - M (WP_1^+)^2 \right) + 4M^2 \left( WQ_2^- WQ_3^- - M (WQ_1^-)^2 \right) \\ &\quad + 2M \left( WP_3^+ WQ_2^- + WP_2^+ WQ_3^- + MWP_1^+ WQ_1^- \right) \\ &\sim 4k_+^2 (L_+ + M)^2 (L_+ - M) + 16k_-^2 M^2 (I - M) \\ &\quad + 8k_+k_- (L_+ + M)M(L_+ + M + I). \end{aligned}$$

From (7), we have

$$\begin{aligned} W^2 \det(\Lambda_2) &\sim 4k_+^2 r_+ (L_+ + M) + 16k_-^2 M^2 (I - M) \\ &\quad + 8k_+k_- (L_+ + M)M(L_+ + M + I) \\ &\sim 4k_+l_+ (L_+ + M) + 16k_-^2 M^2 (I - M) \\ &\quad + 8k_+k_- (L_+ + M)M(L_+ + M + I) \end{aligned}$$

Hence, we note

$$B_3 = 4k_+l_+ (L_+ + M) + 16k_-^2 M^2 (I - M) + 8k_+k_- (L_+ + M)M(L_+ + M + I).$$

Thus, we obtain

$$\det(\Lambda_2) = W^{-2} \left( B_3 + \sum_{j \in J} k_j L_+^{l_j} M^{m_j} e^{\alpha_j L_+} e^{\beta_j M} \right), \quad (53)$$

where  $J$  is a finite set and for any  $j \in J$ :

$$k_j \in \mathbb{R}; \quad l_j, m_j \in \mathbb{N}, \quad \alpha_j, \beta_j \in \mathbb{R}_+ \quad \text{with} \quad \alpha_j + \beta_j \neq 0.$$

**Proposition 6.8.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$  hold and  $k_+ k_- > 0$ . If  $\frac{k_-}{k_+} \leq 2$ , then

$$0 \in \rho \left( 8k_+ k_- (L_+ + M)^2 M - 16k_-^2 M^3 \right).$$

*Proof.* Since  $k_+ k_- > 0$ , we have  $\frac{k_-}{k_+} > 0$  and

$$8k_+ k_- (L_+ + M)^2 M - 16k_-^2 M^3 = 8k_+ k_- M \left[ L_+^2 + 2L_+ M + M^2 - 2\frac{k_-}{k_+} M^2 \right].$$

From Remark 4.1, statement 5 and [31], Corollary 3, p. 444, we deduce that

$$L_+^2, 2L_+ M, M^2 \in \text{BIP}(X, \theta_A).$$

Thus, if  $\frac{k_-}{k_+} \leq 1$ , then

$$L_+^2 + 2L_+ M + M^2 - 2\frac{k_-}{k_+} M^2 = L_+^2 - \frac{k_-}{k_+} M^2 + 2L_+ M + \left( 1 - \frac{k_-}{k_+} \right) M^2.$$

Moreover, for all  $\psi \in D(M^2) = D(A)$ , due to (6), we have

$$\left( L_+^2 - \frac{k_-}{k_+} M^2 \right) \psi = \left[ - \left( 1 - \frac{k_-}{k_+} \right) A + r_+ I \right] \psi,$$

and from [31], Theorem 3, p. 437 and [1], Theorem 2.3, p. 69, assumptions  $(H_2)$  and  $(H_3)$  imply that

$$- \left( 1 - \frac{k_-}{k_+} \right) A + r_+ I \in \text{BIP}(X, \theta_A),$$

and

$$L_+^2 - \frac{k_-}{k_+} M^2 + 2L_+ M + \left( 1 - \frac{k_-}{k_+} \right) M^2 \in \text{BIP}(X, \theta_A + \varepsilon),$$

for any  $\varepsilon \in (0, \pi - \theta_A)$ . Moreover, since  $0 \in \rho(L_+ M)$ , we deduce from [31], remark at the end of p. 445, that  $0 \in \rho \left( L_+^2 - \frac{k_-}{k_+} M^2 + 2L_+ M + \left( 1 - \frac{k_-}{k_+} \right) M^2 \right)$ . Therefore, since  $0 \in \rho(M)$  and  $k_+ k_- > 0$ , it follows that

$$0 \in \rho \left( 8k_+ k_- (L_+ + M)^2 M - 16k_-^2 M^3 \right).$$

In the same way, if  $1 < \frac{k_-}{k_+} \leq 2$ , then

$$L_+^2 + 2L_+ M + M^2 - 2\frac{k_-}{k_+} M^2 = L_+^2 - M^2 + 2L_+ M - 2 \left( \frac{k_-}{k_+} - 1 \right) M^2 + M^2 - M^2,$$

hence, for all  $\psi \in D(M^2) = D(A)$ , from (6), we obtain

$$\left(L_+^2 + 2L_+M + M^2 - 2\frac{k_-}{k_+}M^2\right)\psi = r_+\psi + 2M\left(L_+ - \left(\frac{k_-}{k_+} - 1\right)M\right)\psi. \quad (54)$$

Moreover, we have

$$\left(L_+ - \left(\frac{k_-}{k_+} - 1\right)M\right)\psi = \left(L_+ + \left(\frac{k_-}{k_+} - 1\right)M\right)^{-1}\left(L_+^2 - \left(\frac{k_-}{k_+} - 1\right)^2M^2\right)\psi,$$

and from [31], Theorem 3, p. 437 and [1], Theorem 2.3, p. 69, assumptions  $(H_2)$  and  $(H_3)$  imply that

$$\left(L_+^2 - \left(\frac{k_-}{k_+} - 1\right)^2M^2\right) = -\left(2 - \frac{k_-}{k_+}\right)A + r_+I \in \text{BIP}(X, \theta_A). \quad (55)$$

Finally, from  $(H_2)$ ,  $(H_3)$ , (54), (55) and [31], Theorem 3, p. 437, we deduce that

$$r_+\psi + 2M\left(L_+ - \left(\frac{k_-}{k_+} - 1\right)M\right) \in \text{BIP}(X, \theta_A),$$

and

$$0 \in \rho\left(r_+\psi + 2M\left(L_+ - \left(\frac{k_-}{k_+} - 1\right)M\right)\right).$$

Therefore, since  $0 \in \rho(M)$  and  $k_+k_- > 0$ , it follows that

$$0 \in \rho\left(8k_+k_-(L_+ + M)^2M - 16k_-^2M^3\right).$$

□

We set

$$\begin{aligned} B_4 &= I + 4k_+l_+(L_+ + M)\left(8k_+k_-(L_+ + M)^2M - 16k_-^2M^3\right)^{-1} \\ &\quad + 16k_-^2M^2\left(8k_+k_-(L_+ + M)^2M - 16k_-^2M^3\right)^{-1} \\ &\quad + 8k_+k_-(L_+ + M)M\left(8k_+k_-(L_+ + M)^2M - 16k_-^2M^3\right)^{-1}, \end{aligned} \quad (56)$$

thus, we have

$$B_3 = \left(8k_+k_-(L_+ + M)^2M - 16k_-^2M^3\right)B_4.$$

Moreover, from (53) and noting  $B_5 = 8k_+k_-(L_+ + M)^2M - 16k_-^2M^3$ , we have

$$\det(\Lambda_2) = W^{-2}B_5F_2, \quad (57)$$

where

$$F_2 = B_4 + \sum_{j \in J} k_j B_5^{-1} L_+^{l_j} M^{m_j} e^{\alpha_j L_+} e^{\beta_j M}. \quad (58)$$

Now, for  $z \in \mathbb{C} \setminus [\max(-r_+, 0), +\infty)$ , we set

$$\tilde{f}_2(z) = b_4(z) + \sum_{j \in J} k_j b_5(z)^{-1} \sqrt{z + r_+}^{l_j} \sqrt{z}^{m_j} e^{-\alpha_j \sqrt{z+r_+}} e^{-\beta_j \sqrt{z}},$$

where  $b_3(z) = b_4(z)b_5(z)$ , with

$$\begin{aligned} b_4(z) &= 1 + 4k_+l_+(\sqrt{z+r_+} + \sqrt{z}) \left( 8k_+k_-(\sqrt{z+r_+} + \sqrt{z})^2\sqrt{z} - 16k_-^2\sqrt{z^3} \right)^{-1} \\ &\quad + 16k_-^2z \left( -8k_+k_-(\sqrt{z+r_+} + \sqrt{z})^2\sqrt{z} + 16k_-^2\sqrt{z^3} \right)^{-1} \\ &\quad + 8k_+k_-(\sqrt{z+r_+} + \sqrt{z})\sqrt{z} \left( 8k_+k_-(\sqrt{z+r_+} + \sqrt{z})^2\sqrt{z} - 16k_-^2\sqrt{z^3} \right)^{-1}, \end{aligned}$$

and

$$b_5(z) = -8k_+k_-(\sqrt{z+r_+} + \sqrt{z})^2\sqrt{z} + 16k_-^2\sqrt{z^3}.$$

Then  $\tilde{f}_2(-A) = F_2$ ,  $b_3(-A) = B_3$ ,  $b_4(-A) = B_4$  and  $b_5(-A) = B_5$ . Thus, from (47) and (57), we deduce that

$$f_2(z) = u_{d,r_+}^{-2}(z)v_{d,r_+}^{-2}(z)u_{c,0}^{-2}(z)v_{c,0}^{-2}(z)b_5(z)\tilde{f}_2(z). \quad (59)$$

**Proposition 6.9.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+k_- > 0$  with  $\frac{k_-}{k_+} \leq 2$ .

Thus

- if  $r_+ > 0$  such that

$$r_+ \geq \frac{(\sqrt{t+1} + \sqrt{t})^2}{t^2} \frac{k_+^2}{4k_-^2}, \quad \text{for } t \in \left( 0, \frac{1}{r_+\|A^{-1}\|_{\mathcal{L}(X)}} \right) \text{ fixed}, \quad (60)$$

- if  $r_+ < 0$  such that (50) holds,

then  $F_2$ , given by (58), is invertible with bounded inverse.

*Proof.* From Proposition 6.8 and (58), we deduce that  $F_2$  is well defined.

- Assume that  $r_+ > 0$  such that (49) holds. Then, from Lemma 6.7 and (59), it follows that  $f_2$  does not vanish on  $(tr_+, +\infty)$ , for  $t > 0$  fixed, which involves that  $u_{d,r_+}^{-2}$ ,  $v_{d,r_+}^{-2}$ ,  $u_{c,0}^{-2}$ ,  $v_{c,0}^{-2}$ ,  $b_5$  and  $\tilde{f}_2$  do not vanish on  $(tr_+, +\infty)$ , for  $t > 0$  fixed. Moreover, due to  $(H_2)$ , there exists  $R = \frac{1}{\|A^{-1}\|_{\mathcal{L}(X)}} > 0$  such that  $B(0, R) \subset \rho(A)$ . Therefore, setting  $\tilde{f}_{tr_+,2}(z) = \tilde{f}_2(z + tr_+)$ , with  $t \in \left( 0, \frac{1}{r_+\|A^{-1}\|_{\mathcal{L}(X)}} \right)$  fixed and applying Lemma 5.6 where we have set  $G = \tilde{f}_{tr_+,2}$  and operator  $P = -A - tr_+I \in \text{BIP}(X, \theta_A)$  (due to  $(H_2)$  and  $(H_3)$ ), we deduce that operator  $F_2 = \tilde{f}_{tr_+,2}(-A - tr_+I) = \tilde{f}_2(-A)$  is invertible with bounded inverse.
- Now, assume that  $r_+ < 0$  such that (50) holds. Then  $\tilde{f}_2$  does not vanish on  $(-r_+, +\infty)$ . Moreover, from  $(H_2)$  and  $(H_3)$ , we have  $-A + r_+I \in \text{BIP}(X, \theta_A)$ . It follows that  $F_2 = \tilde{f}_{-r_+,2}(-A + r_+I) = \tilde{f}_2(-A)$  is invertible with bounded inverse.

□

This result finally leads us to state the following main result of this section.

**Proposition 6.10.** Assume that  $(H_1)$ ,  $(H_2)$ ,  $(H_3)$ ,  $(H_4)$  hold and  $k_+k_- > 0$  with  $\frac{k_-}{k_+} \leq 2$ .

Thus

- if  $r_+ > 0$  such that (60) holds,
- if  $r_+ < 0$  such that (50) holds,

then  $\det(\Lambda_2)$  is invertible with bounded inverse.

*Proof.* From (57), Proposition 6.8 and Proposition 6.9, we obtain that  $\det(\Lambda_2) = W^{-2}B_5F_2$ , is invertible with bounded inverse. □



## 6.3 Regularity

### 6.3.1 First case

Here, we consider  $r_+, r_- \in \mathbb{R} \setminus \{0\}$ . From Theorem 5.3, we have to prove that system (17) has a unique solution  $(\psi_1, \psi_2)$  satisfying (18). The existence and uniqueness of this solution is ensured by Proposition 6.6, so we have

$$\begin{cases} \psi_1 &= (P_1^- - P_1^+) [\det(\Lambda_1)]^{-1} S_1 - (P_2^+ + P_2^-) [\det(\Lambda_1)]^{-1} S_2 \\ \psi_2 &= -(P_3^+ - P_3^-) [\det(\Lambda_1)]^{-1} S_1 + M (P_1^- - P_1^+) [\det(\Lambda_1)]^{-1} S_2. \end{cases} \quad (61)$$

Now, we have to study the regularity of  $[\det(\Lambda_1)]^{-1}$ . Since, in this case, the determinant  $\det(\Lambda_1)$  is the same than the one in [19], we deduce, from [19], Lemma 5.3, p. 2958, that there exists  $R_{\det(\Lambda_1)} \in \mathcal{L}(X)$  such that

$$R_{\det(\Lambda_1)}(X) \subset D(M), \quad [\det(\Lambda_1)]^{-1} = N^{-1} + N^{-1}R_{\det(\Lambda_1)},$$

where  $N = 4k_+k_-(L_- + M)(L_+ + M)(L_+ + L_- + 2M)$ . Then, the rest of the proof is similar to the one given in [19], section 5.3. Therefore, from (12) and (13), it follows that  $S_1, S_2 \in (D(M), X)_{1+\frac{1}{p}, p}$  and thus

$$[\det(\Lambda)]^{-1} S_1, [\det(\Lambda)]^{-1} S_2 \in (D(M), X)_{4+\frac{1}{p}, p}. \quad (62)$$

Moreover, from (61), we have

$$\begin{cases} \psi_1 &= -2(k_+(L_+ + M) - k_-(L_- + M)) [\det(\Lambda_1)]^{-1} S_1 \\ &\quad + 2(k_+(L_+ + M) - k_-(L_- + M)) [\det(\Lambda_1)]^{-1} S_2 + \tilde{S}_1 \\ \psi_2 &= -2(k_+(L_+ + M)L_+ + k_-(L_- + M)L_-) [\det(\Lambda_1)]^{-1} S_1 \\ &\quad - 2(k_+(L_+ + M) - k_-(L_- + M)) [\det(\Lambda_1)]^{-1} S_2 + \tilde{S}_2, \end{cases} \quad (63)$$

where  $\tilde{S}_1, \tilde{S}_2 \in D(M^\infty)$ . Finally, from (27), (62) and (63), we obtain

$$\begin{cases} \psi_1 \in (D(M), X)_{3+\frac{1}{p}, p} = (D(A), X)_{1+\frac{1}{2p}, p} \\ \psi_2 \in (D(M), X)_{2+\frac{1}{p}, p} = (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p} \end{cases}$$

### 6.3.2 Second case

Here, we consider  $r_+ \in \mathbb{R} \setminus \{0\}$  and  $r_- = 0$ . From Theorem 5.5, we have to prove that (25) has a unique solution  $(\psi_1, \psi_2)$  satisfying (18). The existence and uniqueness of this solution is ensured by Proposition 6.10, so we have

$$\begin{cases} \psi_1 &= (2MQ_1^- - P_1^+) [\det(\Lambda_2)]^{-1} S_3 + (P_2^+ + 2MQ_2^-) [\det(\Lambda_2)]^{-1} S_4 \\ \psi_2 &= -(P_3^+ + 2MQ_3^-) [\det(\Lambda_2)]^{-1} S_3 + M (P_1^+ - 2MQ_1^-) [\det(\Lambda_2)]^{-1} S_4. \end{cases} \quad (64)$$

Now, we have to study the regularity of  $[\det(\Lambda_2)]^{-1}$ . From (9), (56), (57), (58) and [20], Lemma 5.1, p. 365, we deduce that there exists  $R_{\det(\Lambda_2)} \in \mathcal{L}(X)$  such that

$$R_{\det(\Lambda_2)}(X) \subset D(M), \quad [\det(\Lambda_2)]^{-1} = B_5^{-1} + B_5^{-1}R_{\det(\Lambda_2)},$$

where we recall that  $B_5 = 8k_+k_-(L_+ + M)^2M - 16k_-^2M^3$ . Moreover, from (8), (27), (29) and (32), we have

$$\tilde{\varphi}_1^+, \tilde{\varphi}_2^-, \tilde{\varphi}_2^+, \tilde{\varphi}_3^+, \tilde{\varphi}_4^-, \tilde{\varphi}_4^+ \in (D(M), X)_{2+\frac{1}{p}, p} \quad \text{and} \quad \tilde{\varphi}_1^-, \tilde{\varphi}_3^- \in (D(M), X)_{3+\frac{1}{p}, p}.$$

Thus, from (21), (22), (23), Remark 5.2 and Remark 5.4, we deduce that

$$R_2 \in (D(M), X)_{\frac{1}{p}, p} \quad \text{and} \quad S_3, S_4 \in (D(M), X)_{1+\frac{1}{p}, p},$$

which implies that

$$[\det(\Lambda_2)]^{-1} S_3, [\det(\Lambda_2)]^{-1} S_4 \in (D(M), X)_{4+\frac{1}{p}, p}. \quad (65)$$

Moreover, due to (10), (20), (64) and [20], Lemma 5.1, p. 365, we have

$$\begin{cases} \psi_1 = -2(k_+(L_+ + M) - 2k_-M) [\det(\Lambda_2)]^{-1} S_3 \\ \quad + 2(k_+(L_+ + M) + 2k_-M) [\det(\Lambda_2)]^{-1} S_4 + \tilde{S}_3 \\ \psi_2 = -2(k_+(L_+ + M)L_+ + 2k_-M) [\det(\Lambda_2)]^{-1} S_3 \\ \quad + 2M(k_+(L_+ + M) - 2k_-M) [\det(\Lambda_2)]^{-1} S_4 + \tilde{S}_4, \end{cases} \quad (66)$$

where  $\tilde{S}_3, \tilde{S}_4 \in D(M^\infty)$ . Finally, from (27), (65) and (66), we obtain

$$\begin{cases} \psi_1 \in (D(M), X)_{3+\frac{1}{p}, p} = (D(A), X)_{1+\frac{1}{2p}, p} \\ \psi_2 \in (D(M), X)_{2+\frac{1}{p}, p} = (D(A), X)_{1+\frac{1}{2}+\frac{1}{2p}, p}. \end{cases}$$

## Acknowledgments

I would like to thank the referee for the valuable comments and corrections which have improved this paper.

## References

- [1] W. ARENDT, S. BU & M. HAASE, *Functional calculus, variational methods and Liapunov's theorem*, Arch. Math., 77 (2001), 65-75.
- [2] M.A. AZIZ-ALAOUI, G. CANTIN & A. THOREL, *Synchronization of Turing patterns in complex networks of reaction-diffusion systems set in distinct domains*, Nonlinearity, 37 (2024), 025011 (32pp).
- [3] M. BENHARRAT, F. BOUCHELACHEM & A. THOREL, *On the solvability of fourth-order boundary value problems with accretive operators*, Semigroup Forum, 107 (2023), 17-39.
- [4] J. BOURGAIN, *Some remarks on Banach spaces in which martingale difference sequences are unconditional*, Ark. Mat., 21 (1983), 163-168.
- [5] D.L. BURKHOLDER, *A geometrical characterisation of Banach spaces in which martingale difference sequences are unconditional*, Ann. Probab., 9 (1981), 997-1011.
- [6] G. CANTIN & A. THOREL, *On a generalized diffusion problem: a complex network approach*, Discrete Contin. Dyn. Syst. - B, 27, 4 (2022), 2345-2365.

- [7] R.S. CANTRELL & C. COSNER, “Spatial Ecology via Reaction-Diffusion Equations”, Wiley, 2003.
- [8] D.S. COHEN & J.D. MURRAY, *A generalized diffusion model for growth and dispersal in population*, Journal of Mathematical Biology, 12 (1981), 237-249.
- [9] G. DA PRATO & P. GRISVARD, *Somme d’opérateurs linéaires et équations différentielles opérationnelles*, J. Math. Pures et Appl., 54 (1975), 305-387.
- [10] A. Ducrot, P. Magal & A. Thorel, *An integrated semigroup approach for age structured equations with diffusion and non-homogeneous boundary conditions*, Nonlinear Differ. Equ. Appl., 28, 49 (2021).
- [11] R. DENK & F. KAMMERLANDER, *Exponential stability for a coupled system of damped-undamped plate equations*, IMA J. Appl. Math., 83 (2018), 302-322.
- [12] G. DORE & A. VENNI, *On the closedness of the sum of two closed operators*, Math. Z., 196 (1987), 189-201.
- [13] A. FAVINI, R. LABBAS, S. MAINGOT, K. LEMRABET & H. SIDIBÉ, *Resolution and optimal regularity for a biharmonic equation with impedance boundary conditions and some generalizations*, Discrete Contin. Dyn. Syst. - A, 33, 11-12 (2013), 4991-5014.
- [14] D. GILBARG & N. TRUDINGER, “Elliptic partial differential equations of second order”, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [15] P. GRISVARD, *Spazi di tracce ed Applicazioni*, Rendiconti di Matematica, Serie VI, 5 (1972), 657-729.
- [16] P. GROS, “La représentation de l’espace dans les modèles de dynamiques de populations, Modèles dynamiques déterministes à temps et espace continu”, Ifremer, 2001.
- [17] M. HAASE, “The Functional Calculus for Sectorial Operators, Operator Theory: Advances and Applications”, Birkhäuser Verlag, Basel-Boston-Berlin, 2006.
- [18] H. KOMATSU, *Fractional Powers of Operators*, Pac. J. Math., 19, 2 (1966), 285-346.
- [19] R. LABBAS, K. LEMRABET, S. MAINGOT & A. THOREL, *Generalized linear models for population dynamics in two juxtaposed habitats*, Discrete Contin. Dyn. Syst. - A, 39, 5 (2019), 2933-2960.
- [20] R. LABBAS, S. MAINGOT, D. MANCEAU & A. THOREL, *On the regularity of a generalized diffusion problem arising in population dynamics set in a cylindrical domain*, J. Math. Anal. Appl., 450 (2017), 351-376.
- [21] R. LABBAS, S. MAINGOT & A. THOREL, *Generation of analytic semigroup for some generalized diffusion operators in  $L^p$ -spaces*, Math. Ann., 384 (2022), 1-49.
- [22] R. LABBAS, S. MAINGOT & A. THOREL, *Solvability of a fourth order elliptic problem in a bounded sector, part I*, Boll. Unione Mat. Ital., 17 (2024), 647-666.
- [23] R. LABBAS, S. MAINGOT & A. THOREL, *Solvability of a fourth order elliptic problem in a bounded sector, part II*, Boll. Unione Mat. Ital., (2024).
- [24] J.-L. LIONS & J. PEETRE, *Sur une classe d’espaces d’interpolation*, Publications mathématiques de l’I.H.É.S., 19 (1964), 5-68.

- [25] A. LUNARDI, “Analytic semigroups and optimal regularity in parabolic problems”, Birkhauser, Basel, Boston, Berlin, 1995.
- [26] A. LUNARDI, “Interpolation theory”, Third edition, Lecture Notes, Scuola Normale Superiore di Pisa (New Series), 16, Edizioni della Normale, Pisa, 2018.
- [27] J.D. MURRAY, “Mathematical Biology II: Spatial Models and Biomedical Applications”, Third Edition, Springer, 2003.
- [28] A. NOVICK-COHEN, *On Cahn-Hilliard type equations*, Nonlinear Analysis, Theory, Methods & Applications, 15, 9 (1990), 797-814.
- [29] F.L. OCHOA, *A generalized reaction-diffusion model for spatial structures formed by motile cells*, BioSystems, 17 (1984), 35-50.
- [30] A. OKUBO & S.A. LEVIN, “Diffusion and ecological problems, Mathematical biology”, second edition, Springer-Verlag, Berlin, 2010.
- [31] J. PRÜSS & H. SOHR, *On operators with bounded imaginary powers in Banach spaces*, Mathematische Zeitschrift, 203 (1990), 429-452.
- [32] J. PRÜSS & H. SOHR, *Imaginary powers of elliptic second order differential operators in  $L^p$ -spaces*, Hiroshima Math. J., 23, 1 (1993), 161-192.
- [33] J.L. RUBIO DE FRANCIA, *Martingale and integral transforms of Banach space valued functions*, Probability and Banach Spaces, (Zaragoza, 1985), in: Lecture Notes in Math., 1221, Springer-Verlag, Berlin (1986), 195-222.
- [34] A. THOREL, *Operational approach for biharmonic equations in  $L^p$ -spaces*, J. Evol. Equ., 20 (2020), 631-657.
- [35] A. THOREL, *A biharmonic transmission problem in  $L^p$ -spaces*, Commun. Pure Appl. Anal., 20, 9 (2021), 3193-3213.
- [36] H. TRIEBEL, “Interpolation Theory, Functions Spaces, Differential Operators”, Amsterdam, N.Y., Oxford, North-Holland, 1978.